1 **Age, sex, and social environmental effects on immune cell composition in a free-ranging**
2 **non-human primate**
3

Mitchell R. Sanchez Rosado[1], Nicole Marzan-Rivera[1], Marina M. Watowich[2], Andrea D.

Negron-Del Valle[3] Petraleigh Pantoja[1,4], Melissa A. Pavez-Fox[5], Erin R. Siracusa[5], Eve B.

Cooper[6], Josue E. Negron-Del Valle[7,8], Daniel Phillips[7,8], Angelina Ruiz-Lambides[4], Cayo

Biobank Research Unit, Melween I. Martinez[4], Michael J. Montague[9], Michael L.

Platt[9,10,11], James P. Higham[6], Lauren J. N. Brent[5], Carlos A. Sariol[1,4], Noah Snyder-Mackler[7,8,12]

Affiliations:

[1]Department of Microbiology & Medical Zoology, University of Puerto Rico-Medical Sciences,

San Juan, PR

[2]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[3]Department of Biology, University of Puerto Rico-Humacao, San Juan, PR

[4]Unit of Comparative Medicine, Caribbean Primate Research Center and Animal Resources

Center, University of Puerto Rico-Medical Sciences Campus, San Juan, PR, USA

[5]Centre for Research in Animal Behaviour, University of Exeter, Exeter EX4 4QG, UK

[6]Department of Anthropology, New York University, New York, NY 10003, USA

[7]School of Life Sciences, Arizona State University, Tempe, AZ, USA

[8]Center for Evolution and Medicine, School of Life Sciences, Arizona State University, Tempe,

AZ, USA

[9]Department of Neuroscience, University of Pennsylvania, Philadelphia, PA 19104, USA

[10]Department of Psychology, University of Pennsylvania, Philadelphia, PA 19104, USA

[11]Department of Marketing, Wharton School, University of Pennsylvania, Philadelphia, PA

19104, USA

[12]School for Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA

26
27
28
29

Abstract

Increasing age is associated with dysregulated immune function and increased inflammation– patterns that are also observed in individuals exposed to chronic social adversity. Yet we still know little about how social adversity impacts the immune system and how it might promote age-related diseases. Here, we investigated how immune cell diversity varied with age, sex and social adversity (operationalized as low social status) in free-ranging rhesus macaques. We found age-related signatures of immunosenescence, including lower proportions of CD20+ B cells, CD20+/CD3+ ratio, and CD4+/CD8+ T cell ratio – all signs of diminished antibody production. Age was associated with higher proportions of CD3+/CD8+ Cytotoxic T cells, CD16+/CD3- Natural Killer cells, CD3+/CD4+/CD25+ and CD3+/CD8+/CD25+ T regulatory cells, and CD14+/CD16+/HLA-DR+ intermediate monocytes, and lower levels of CD14+/CD16-/HLA-DR+ classical monocytes, indicating greater amounts of inflammation and immune dysregulation. We also found an effect of exposure to social adversity (i.e., low social status) that was sex-dependent. High-status males, relative to females, had higher CD20+/CD3+ ratios and CD16+/CD3 Natural Killer cell proportions, and lower proportions of CD8+ Cytotoxic T cells. Further, low status females had higher proportions of cytotoxic T cells than high status females, while the opposite was observed in males. High status males had higher CD20+/CD3+ ratios than low status males. Together, our study identifies immune cell types that differ by age in a human-relevant primate model animal, and demonstrates a novel link between sex-dependent immunity and social adversity.

Introduction

The average human lifespan has almost doubled over the past century [1], accompanied by an increase in the prevalence of many age-related diseases, including cardiovascular disease, autoimmune disease, diabetes, arthritis, and cognitive decline [2-5]. As individuals age, there is a disruption in the homeostatic balance between innate and adaptive immunity linked to both increases in age-related disease and susceptibility to infection. This imbalance is reflective of two age-related alterations, namely increased inflammation ("inflammaging") and a decline in adaptive immune function ("immunosenescence") [6-7]. Both alterations disrupt the balance between pro- and anti-inflammatory mediators that characterize a healthy immune system. For example, with increasing age, innate immune cells such as monocytes become more active and release drivers of inflammation that include proinflammatory cytokines (e.g., TNF-α) [8]. Adaptive immune cells, such as B cells and helper T cells, show age-related declines that directly impact long-term immunity, as exemplified by the lower effectiveness of vaccination in older individuals [9].

Yet these age-related alterations in immunity are not universal in their trajectories across individuals. There is substantial heterogeneity with age; not all individuals age at the same rate or fall victim to the same age-related diseases. For instance, some people become hypertensive in their 30s, while a 60-year-old may never suffer from this condition. Part of this heterogeneity is due to sex differences in the immune system, which alter the prevalence and onset of age-related diseases. For example, in many species, females mount a stronger immune response with increasing age compared to males [10-11]. Females also have a stronger age-related increase in inflammatory cells compared to males [12]. Further, in humans, older men are more susceptible to infections, such as leptospirosis, tuberculosis, and hepatitis A, than are women [13].

Heterogeneity in aging can also arise from life experiences, such as exposure to social adversity, which can influence the onset and progression of disease and, ultimately, mortality [14]. Social adversity, which is often associated with low social status, and other social stressors [15-16], has been linked to accelerated aging as indexed by biomarkers like epigenetic age and telomere attrition [17-19]. There are also broad similarities between the effects of age and social adversity on peripheral immune function [20]. For instance, early life social adversity in humans has been linked to increases in proinflammatory T cells [21] – a characteristic usually seen with increasing

age. Further, various adversities and social stressors are associated with a decrease in naïve CD4
T helper cells and an increase in naïve CD8 T cells [22], pointing to how the social environment
can shape immunity. However, the extent to which social adversity may be associated with
immunity across the lifecourse remains unknown. Social adversity might lead to accelerated aging-
related disease onset and death and/or social advantage may confer protection from the effects of
aging.

Social structures in human populations are complex and multifaceted, including structural
inequities and discrimination, and these factors can vary across cultures. Thus, it can be difficult
to measure how social adversity "gets under the skin" in humans to affect immune and overall
health. Rhesus macaques (*Macaca mulatta*), a non-human primate, are an established animal
model that exhibits aging trajectories similar to humans, such as decreases in mobility, but
compressed into a lifespan 3-4x times shorter [23]. Aging parallels are also manifested at the
molecular level: rhesus macaques and humans show similar age-related alterations in immune cell
DNA methylation and gene expression [24]. Rhesus macaques also share many social factors with
humans including the expression of affiliative behaviors, despotism, among other behaviors [25],
making them an ideal animal model for translational aging research.

In rhesus macaques, exposure to social adversity can be captured by measures of social status (i.e.,
dominance rank). Social status is acquired differently for males and females [26-27]: females
inherit status from their mothers, while males typically disperse from their natal group and enter a
new group where they acquire status through a combination of "queuing" and physical contests
[28-29]. Similar to humans, social status in macaques patterns access to resources, and can impact
health and lifespan [30-32]. For instance, low status macaques experience more conspecific
aggression and are therefore more likely to be injured [33], and high status female macaques can
live longer than those with lower status [34]. In addition, social status affects immunity in rhesus
macaques; one experimental study showed that social status predicts gene expression patterns in
peripheral blood mononuclear cells [35].
Here, we characterized age-related variation in immune cell types, as well as the influence of social
adversity and sex on immune cell composition. To do so, we studied a free-ranging population of
rhesus macaques living on the island of Cayo Santiago, Puerto Rico where we were able to

simultaneously document age, sex, and social status in a semi-naturalistic social setting with
minimal human intervention [36].

Methods

Study population:

Cayo Santiago is an island located off the southeastern coast of Puerto Rico inhabited
by approximately 1,600 rhesus macaques. The population is managed by the Caribbean Primate
Research Center (CPRC) and is the oldest primate field station in the world [37]. Cayo
Santiago provides unique research opportunities for behavioral, physiological, demographic,
morphological and genomic studies. The Cayo Santiago Field Station has a minimal intervention
policy, which means that the animals are not managed medically or reproductively. There are no
predators on the island, and senescent phenotypes are commonly observed in this population
[24,38-41]. The animals are direct descendants of rhesus macaques introduced from India in 1938;
since 1957 these animals have been continuously monitored [42]. The animals are identified with
tattoos and ear notches, and demographic data such as age, sex and pedigree have been collected
for decades. During the annual capture and release period, researchers have the opportunity to
collect biological and morphological samples with the assistance of CPRC veterinary staff. For the
past 15 years, the Cayo Biobank Research Unit has collected detailed behavioral data to combine
with the biological samples collected each year. In combination, these data provide the opportunity
to test the relationships between the social environment, immune function, and aging.

Blood sampling:

We collected whole blood from sedated rhesus macaques over three capture and release periods
(n=96 in October - December 2019, n=153 in October 2020 - February 2021 and n=120 in October
2021 - February 2022). Samples were collected in 6ml K2 EDTA tubes (Beckton, Dickson and
Company, cat #367899). We collected a total of 369 samples (200 from males, 169 from females)
from 230 unique individuals (113 males, 117 females; i.e., some animals were sampled across
multiple years of the study), spanning the natural lifespan of macaques on Cayo Santiago (mean
age = 11.8 years, range 0-28 years; **Figure 1A and 1B**). Fresh blood samples were transported at
4°C to the University of Puerto Rico Medical Sciences campus where flow cytometric analysis
was performed within 6 hours of sample collection.

Antibodies and flow cytometric analysis:

An 8-panel antibody cocktail previously validated in rhesus macaques [43-46], consisting of the

following antibodies, was used: CD20-PacBlue/Clone 2H7 (Biolegend), CD3-PerCP/Clone SP34-

2 (BD), CD4-APC/Clone L200 (BD), CD8-Viogreen/Clone BW135/80 (Miltenyi), CD25-

PE/Clone 4E3 (Miltenyi), CD14-FITC/Clone M5E2 (BD), CD16-PEVio770/Clone REA423

(Miltenyi), HLA-DR-APCVio770/Clone REA805 (Miltenyi).

We performed phenotypic characterization of rhesus macaque peripheral blood mononuclear cells

(PBMCs) using multicolor flow cytometry with direct immunofluorescence (View **S. Figure 1** for

gating strategy and **Table S1** for Ab panel) on all 369 animals. Aliquots of 150 µl of heparinized

whole blood were incubated with a mix of the antibodies described for 30 minutes at 25°C (room

temperature). After incubation, the red blood cells were fixed and lysed with BD FACS fix and

lyse solution (Cat #349202). Cells were washed twice using PBS containing 0.05% BSA at 1,700

RPM for 5 minutes and processed in a MACSQuant Analyzer 10 flow cytometer (Miltenyi Biotec,

CA).

Lymphocytes and monocytes were gated based on their characteristic forward scatter (measures

cells based on their size) and side scatter (measures cells based on their granularity) patterns.

Lymphocytes were then further subdivided according to their cell surface markers. Natural killer

(NK) cells were defined as the CD3- and CD16+ population; B cells were defined as CD20+

population and T cells as the CD3+ population. We further subdivided T cells from the CD3+ gate

into CD4+ and CD8+ populations. CD4+CD25+ and CD8+CD25+ T regulatory cells were further

gated from the CD4+ and CD8+ gates. Monocytes were gated based on the combined expression

of the HLA-DR/CD14 markers for classical monocytes, HLA-DR/CD16 markers for non-classical

monocytes, and HLA-DR/CD14/CD16 for intermediate monocytes (**S. Figure 2**). Flow cytometry

gating was performed using Flowjo version 10.7.1 (FlowJo LLC Ashland, OR).

To obtain an accurate representation of the proportions of cell types, we counted only the stained

events of the cells of interest and calculated their proportions based on the subsets of lymphocytes

and monocytes. To calculate cell ratios, such as CD20 + B cell to CD3+ T cell ratio (CD20+/CD3+

ratio) and CD4+ T cell to CD8+ T cell ratio (CD4+/CD8+ ratio), we divided the calculated

proportion of these cell types in each individual sample (e.g., CD20+ B cell/ CD3+ T cell and

CD4+/CD8+ respectively). (Calculations detailed in **Table S2**).

Quantification of social adversity (social status):

We quantified the social status of a subset of animals for which we had behavioral data (**Figure**

1C, n = 250 total samples, 134 from males and 116 from females & n = 145 unique individuals,

73 males and 72 females). We calculated social status (i.e., dominance rank) using the outcomes

of win-loss dominance interactions between pairs of same sex adult groupmates. Because of the

different routes through which males and females acquire their status, we quantified social status

separately for males and females within each group for each year of the study [47-48].

Observations of adult animals (older than 6 years) were collected from two different social groups,

groups V and F, in the year prior to sample collection. In 2019 and 2021, behavioral data were

collected using a previously established 10-minute focal sampling protocol [26]. Briefly, the

protocol consisted of recording state behaviors (e.g., resting, feeding) and agonistic encounters,

which included recording the identity of the focal animal and their social partner. Win-loss

agonistic interactions included threat and submissive behaviors, along with contact and non-

contact aggression. In 2019 and 2021 we also collected additional agonistic interactions *ad-*

libitum. In 2020, all agonistic interactions were collected *ad-libitum* due to restrictions imposed

on behavioral data collection due to the COVID-19 pandemic. In all years, we used known

maternal relatedness to settle behavioral gaps in the female hierarchy [49]. To control for variation

in group size, social status (i.e., dominance rank) was quantified as the percentage of same-sex

adults that an animal outranked in their group. For all analyses, we grouped animals into one of

two social status categories: high-rank (>80% of same-sex adults outranked) and low-rank (< 79%

of same-sex adults outranked)[50].

Statistical analysis:

All statistical analyses were performed using R statistical software R version 4.2.3 [51].

First, we performed a principal component analysis of the cell composition data for all samples (n

= 369, 230 unique individuals) using the *prcomp* function of the *stats* package. Next, we employed

a linear additive mixed-effects approach, using the *lmer* function in the *lmerTest* package to run

sample projections on principal components as a function of age (in years), sex, and sample period
- to control for the technical variation in the flow cytometer lasers, which changed over the
sampling years (*model 1* - **Table S3**) and individual ID as a random effect. We also modeled
sample projections on principal components as a function of the interaction between age and sex
(age*sex) and sampling period - which will ultimately allow us to identify possible sex-dependent
associations with age - and individual ID as a random effect (*model 2* - **Table S3**).

To evaluate each cell type at a more granular level, we employed the same additive linear mixed-
effects to test the proportion of each cell type and certain cell type ratios (e.g., CD4+/CD8+) as a
function of age, sex, and sample period with individual ID as a random effect (*model 3* - **Table
S3**). Finally, we tested the proportion of each cell type and certain cell type ratios (e.g.,
CD4+/CD8+) as a function of the interaction between age and sex (age*sex) and sampling period
with individual ID as a random effect (*model 4* - **Table S3**).

For the subset of samples in which information on social status was available (n = 250 total
samples, 145 unique individuals), we tested for the additive effect of principal component
projections as a function of social status, age, sex and sample period, with individual ID and social
group as a random effect (*model 5* - **Table S3**). We also tested for the principal component
interaction between status and age (status*age) and between status and sex (status*sex), with
individual ID and social group as a random effect (*model 6* - **Table S3**). We then additively tested
the proportion of each cell type and certain cell type ratios as a function of social status, age, sex,
and sample period (*model 7* - **Table S3***)*. To test whether the relationship between the proportion
of cell types and social status depended on sociodemographic variables, we tested the interaction
between: social status and age (status*age) and for social status and sex (status*sex, *model 8* -
Table S3). Furthermore, since we identified interactions between social status and sex, and since
social status is acquired differently for male and females in rhesus macaques, we fitted a separate
model for males and females to test if there was a main effect of social status within each sex on
the proportions of different cell types. Age and sample period were also included in the model to
control for these covariates (*model 9* - **Table S3**).

The linear models and sample sizes for each are summarized in **Table S3.** For every predictor

variable in the full (n = 369) and status (n = 250) datasets, we corrected for multiple hypothesis

tests using the Benjamini Hochberg FDR approach and considered significant associates at FDR

< 0.10. Because Model 7 was only performed in cell types that showed a significant interaction

between sex and social status (and not all the cell types measured), we did not correct for multiple

testing in this model.

Results

Macaques exhibit age-related variation in immune cell composition and inflammation

Age was significantly positively associated with the first principal component (PC) of immune cell

composition (*model 1* - $\beta_{PC1\ age}$ = 0.14, FDR = 7.2 x 10^{-18}, **Figure 1D)**. This first PC, which

explained 32.4% of the variance in cell composition across all samples, was associated with higher

proportions of inflammation-associated cell types, such as cytotoxic T cells and regulatory T cells,

and lower proportions of cells involved in pathogen clearance, including CD20+ B cells and

classical monocytes (**Figure 1E, Table S4).** Thus, older animals exhibited a pattern of greater

inflammation and immunosenescence than younger individuals did.

cytometry. **B)** The dataset was roughly balanced between males and females and captured the

entire natural lifespan of macaques in this population. **C)** We calculated social status by assigning

dominance ranks to 250 samples using observational data collected in the year before each sample

was collected. Animals were assigned to one of three dominance ranks: high, medium, and low.

The social status dataset is a subset of the original age dataset because behavioral data were not

available for all study animals (i.e., it is not collected for infants and juveniles). **D)** PC1 of immune

cell compositions is significantly associated with age (β_{PC1} = 0.14, FDR = 7.2 x 10^{-18}). **E)** The T

cell compartment is positively associated with PC1 (and thus age), while the B cell compartment

is negatively associated with PC1 of immune cell composition.

We then conducted a more granular analysis of the factors associated with the proportions of

individual cell types. Age was significantly associated with signatures of immunosenescence,

including a decline in adaptive immune cells. This was largely driven by lower proportions of

CD20+ B cells in older individuals (*model 3* - $\beta_{CD20\ age}$ = -0.83 ± 0.09, FDR = 1.3 x 10^{-16}, **Figure**

2A), which resulted in significantly lower CD20+/CD3+ ratios in older individuals (*model 3* -

$\beta_{CD20:CD3\ age}$ = -0.04 ± 0.004, FDR = 4.9 x 10^{-15}, **Figure 2B**). Age was also associated with higher

proportions of inflammation-related cells. The proportion of cytotoxic CD8+ Cytotoxic T cells

was significantly higher in older animals (*model 3* - $\beta_{CD8\ age}$ = 0.60 ± 0.07, FDR = 3.2 x 10^{-14},

Figure 2C), resulting in a strong and significant effect of lower CD4+/CD8+ ratios (*model 3* -

$\beta_{CD4:CD8\ age}$ = -0.06 ± 0.008, FDR = 4.3 x 10^{-14}, **Figure 2D**) and higher proportions of CD3+ T cells

in older individuals (*model 3* - β_{CD3}= 0.67 ± 0.11, FDR = 2.2 x 10^{-8}, **S. Figure 3**).

Next, we examined the less abundant but immunologically important regulatory CD8+ and CD4+

T cell populations (CD25+), which are involved in immune suppression and maintenance of self-

tolerance [52] (i.e., the ability to recognize self-antigens). Both CD4+ and CD8+ T regulatory cells

were significantly more abundant in older animals (*model 3* - CD3+CD4+CD25+: β_{age}= 0.16 ±

0.02, FDR = 3.6 x 10^{-10}, **Figure 2E**; *model 3* - CD3+CD8+CD25+: β_{age} = 0.03 ± 0.005, FDR =

1.8 x 10^{-6}, **Figure 2F**), suggesting a reduced age-related ability to regulate endogenous and

exogenous antigens.

Figure 2: Age-associated differences in adaptive immune cell proportions. A) CD20+ B cells
(β_{CD20} = -0.83 ± 0.09, FDR = 1.3 x 10^{-16}) proportions and **B)** CD20+/CD3+ ratio ($\beta_{CD20:CD3}$ = -0.04
± 0.004, FDR = 4.9 x 10^{-15}) are lower in older individuals. **C)** CD8+ Cytotoxic T cells (β_{CD8} = 0.60
± 0.07, FDR = 3.2 x 10^{-14}) are higher in older individuals, while the **D)** CD4+/CD8+ T cell ratio
($\beta_{CD4:CD8}$ = -0.06 ± 0.008, FDR = 4.3 x 10^{-14}) is lower in older individuals. **E)** CD4+ T regulatory
cells (β = 0.16 ± 0.02, FDR = 3.6 x 10^{-10}) and **F)** CD8+ T regulatory cells are higher in older
individuals compared to younger individuals (β = 0.03 ± 0.005, FDR = 1.8 x 10^{-6}), possibly because
of higher baseline levels of inflammation (i.e., inflammaging).

Innate immune cells also showed significant associations with age. Classical monocytes (HLA-
DR+/CD14+/CD16-), which are involved in phagocytosis and extracellular pathogen clearance
[53], were lower in older individuals (*model 3* - $\beta_{CD14++\ age}$ = -0.31 ± 0.16, FDR = 0.07, **Figure
3A**), while intermediate monocytes (HLA-DR+/CD14+/CD16+), involved in immune cell
recruitment and proinflammatory cytokine secretion [53], were higher in older individuals (*model
3* - $\beta_{CD14+CD16+\ age}$ = 0.21 ± 0.09, FDR = 0.04, **Figure 3B)**. The proportion of CD16+CD3- NK cells
– which have a similar role to CD8+ Cytotoxic T cells presenting natural cytotoxicity but are not
antigen specific – was also significantly higher in older individuals (*model 3* - $\beta_{NK\ age}$ = 0.17 ± 0.07,
FDR = 0.03, **Figure 3C**). Together, these results indicate that older individuals show a decrease
in adaptive immunity along with an increase in inflammation-related innate immune cells
compared to younger individuals, potentially disrupting a "healthy" homeostatic immune system.

Figure 3: Age is associated with variation in innate immune cell proportions. A) CD14+/CD16-/HLADR+ Classical monocytes (β_{CD14++} = -0.31 ± 0.16, FDR = 0.07) are lower and **B)** CD14+/CD16+/HLADR+ intermediate monocytes ($\beta_{CD14+CD16+}$ = 0.21 ± 0.09, FDR = 0.04) are higher in older individuals, while **C)** CD16+ NK cells (β_{NK} = 0.17 ± 0.07, FDR = 0.03) are higher in older individuals.

We did not observe statistically significant main effects of sex (*model 3* in *Methods*) or a sex-age interaction (*model 4* in *Methods*) on immune cell proportions. Nevertheless, a trend toward sex differences was observed in both the proportions of CD8+ Cytotoxic T cells (*model 3* - $\beta_{CD8\ sex}$ = 2.19 ± 0.95, FDR = 0.14) and in the CD4+/CD8+ ratio (*model 3* - $\beta_{CD4:CD8\ sex}$ = -0.24 ± 0.09, FDR = 0.14, **S. Figure 4**), with males having a higher proportion of CD8+ Cytotoxic T cells compared to females, and females having a higher CD4+/CD8+ ratio compared to males, suggesting a stronger adaptive immune response in females, which, in part, is generated by CD4+ T helper cells.

Social status and immune cell composition

There was a significant interaction between social status and sex on PC1 (31% of the variance in cell composition across all samples) of immune cell composition (*model 6* - $\beta_{PC1-sex*status}$ = -1.7 ± 0.63, FDR = 0.04, **Figure 4A)**, documenting the sex-dependent impact of social status on immunity.

When modeling males and females together in an additive modeling framework, we found no significant effects of social status on immune cell proportions (all FDR > 0.10, *model 7* in *Methods*), or between the interaction between social status and age (*model 8* in *Methods*). However, we found many significant interactions between social status and sex on immune cell

composition (*model 8* in *Methods*). Because social status is acquired differently for male and
female rhesus macaques, we also carried out post-hoc analyses of the social status effects within
each sex separately (*model 9* in *Methods*).

The CD20+/CD3+ ratio interacted between social status and sex such that it was higher in males
of high social status, but lower in females with high social status (*model 8* - $\beta_{CD20/CD3\ ratio-sex*status}$ =
-0.26 ± 0.11, FDR = 0.06, **Figure 4B)**, both part of the adaptive immune system. This difference
in the CD20+/CD3+ ratio seems to be partially driven by interaction between sex and social status
on the proportion of CD3+ T cells (*model 8* - $\beta_{CD3-sex*status}$ = 11.8 ± 4.42, FDR = 0.04, **S. Figure**
5), such that CD3+ T cells were higher in low social status males, but this pattern was flipped
in females. There was a within-sex main effect of social status in males in the CD20+/CD3+ ratio,
with high social status males having a significantly higher ratio than low social status males did
(*model 9* - $\beta_{CD20/CD3\ ratio-males/status}$ = -1.3 ± 0.06, p = 0.04, **Figure 4B**). No significant effect of social
status on the CD20+/CD3+ ratio was found in females (*model 9* - $\beta_{CD20/CD3\ ratio-females/status}$ = 0.10 ±
0.13, p = 0.19).

Additionally, there was a significant interaction between social status and sex in the proportion of
CD8+ Cytotoxic T cells (*model 8* - $\beta_{CD8-sex*status}$ = 8.3 ± 2.94, FDR = 0.04, **Figure 4C)**, which were
higher in high social status males, but this relationship again flipped females. Our within-sex
analysis revealed a significant main effect of CD8+ Cytotoxic T cells, in which low social status
males had significantly higher proportions of this cell type than did high social status males (*model*
9 - $\beta_{CD8-males/status}$ = 4 ± 1.8, p = 0.03, **Figure 4C**). The opposite main effect was observed in females,
in which high status females had significantly higher proportions of CD8+ Cytotoxic T cells than
did low social status females (*model 9* - $\beta_{CD8-females/status}$ = -4.8 ± 2.1, p = 0.03, **Figure 4C**).

In the innate arm of the immune system, we detected an interaction between social status and sex
on the proportion of CD3-CD16+ NK cells (*model 8* - $\beta_{NK-sex*status}$ = -7.9 ± 3.2, FDR = 0.05, **Figure**
4D), where the proportion was lower in males of low status, but higher in females of low status.
Social status approached significance in the proportions of CD3-CD16+ NK cells in females
(*model 9* - $\beta_{NK-females/status}$ = 5.6 ± 3, p = 0.06, **Figure 4D**), in which low social status females
displayed higher proportions of this cell type compared to high social status females. We found no

significant main effect of social status on CD3-CD16+ NK cells in males (*model 9* - $\beta_{\text{NK-males/status}}$
= -2.27 ± 3.4, p = 0.18).

380
Figure 4: Sex and social status interact to impact immune cell composition. A) PC1 (31% of
the variation in the dataset, $\beta_{\text{PC1-sex*status}}$ = -1.7, FDR = 0.03) recapitulates the interaction between
sex and social status. **B)** Interaction in the CD20+/CD3+ ratio ($\beta_{\text{CD20/CD3 ratio-sex*status}}$ = -0.25 ± 0.05,
FDR = 0.06) shows that this ratio is higher with higher social status in males while it is lower with
higher social status in females; within-sex analysis show that high social status males have a
significantly higher ratio than low social status males ($\beta_{\text{CD20/CD3 ratio-males/status}}$ = -1.3 ± 0.06, p =
0.04). **C)** CD8+ Cytotoxic T cells show an interaction ($\beta_{\text{CD8-sex*status}}$ = 8.3 ± 2.94, FDR = 0.04)
where this cell type is higher in lower social status in males, while it is higher in high social status
females. Within-sex analysis of CD8+ T cells showed low social status males had significantly
higher proportions of this cell type than did high social status males ($\beta_{\text{CD8-males/status}}$ = 4 ± 1.8, p =
0.03), while the opposite effect was observed in females ($\beta_{\text{CD8-females/status}}$ = -4.8 ± 2.1, p = 0.03).
D) Interaction in the proportion of NK Cells ($\beta_{\text{NK-sex*status}}$ = -7.9 ± 3.2, FDR = 0.05), with high
social status males showing higher proportions, while high social status females show lower
proportions.

Discussion

We examined how social status, age, and sex were related to immune cell distributions in a large sample of adult rhesus macaques living in semi-natural conditions. Overall, we found strong and consistent signatures of age-related immune cell dysregulation. We also identified significant links between social status and sex in cells of innate and adaptive arms of the immune system. Together, this variation is likely to influence immune responses to pathogenic challenges as well as the development of inflammation-related diseases.

Overall, macaques exhibited age-related differences in immune cells similar to those observed in humans, including declines in lymphocytes [54]. Here, we also identified more specific cell types with age-related differences. We detected lower proportions of CD20+ B cells at older ages, which may reflect immunosenescence, as these cells are responsible for antibody production, pathogen clearance, and are key cells in the generation of immune memory. Further, a key factor underlying the limited efficacy of vaccines in older individuals is a weakened B cell response [55]; B cells have also been associated with protection against certain types of cancer, such as lung cancer [56].

Similar to two other studies in captive macaques, we found higher CD8+ T cell proportions at older ages [67-58]. Notably, this differs from findings in humans, where both CD8+ Cytotoxic T cells and their effector responses (i.e., stimulus responsiveness) exhibit lower proportions at older ages [59]. It is possible that this discrepancy is only reflected in the overall CD8+ cytotoxic T cell pool, as it has been reported that certain CD8+ T cell subsets – such as memory subsets – increase in proportion and efficacy with age [60]. Alternatively, given that CD8+ T cell subsets have been associated with inflammation and 'inflammaging' [61], there is a possibility that higher overall CD8+ Cytotoxic T cell pool in rhesus macaques is indicative of higher levels of inflammation. The age-related reduction in CD4+/CD8+ ratio corroborates this hypothesis. In support of increased inflammation with age, we found that older animals had significantly higher CD3-CD16+ NK cell proportions in our dataset. Similar to CD8+ Cytotoxic T cells, CD3-CD16+ NK cells respond to intracellular pathogens, secrete multiple proinflammatory mediators, and are crucial during tumor surveillance and signaling [62]. The higher proportions of CD3-CD16+ NK cells predict a higher incidence of inflammation and/or tissue injury in the older population, which is commonly observed in the Cayo Santiago macaque population [33]. As expected, CD4+CD25+ T regulatory

cells as well as CD8+CD25+ T regulatory cells were associated with age, indicating higher levels
of inflammation in older individuals [63]. These results, together with lower levels of CD20+ B
cell proportions and higher levels of CD8+ T cell and CD3-CD16+ NK cell proportions, further
support the hypothesis that the adaptive immune response in rhesus macaques decreases with age
and inflammation-related cell types increase (i.e., 'inflammaging'). Taken together, these
alterations may drive biological and physiological decline that likely increases the risk of
morbidity and mortality in macaques, as it does in humans.

Monocyte proportions also varied with age. Specifically, we found fewer CD14+ classical
monocytes in older animals. These cells are phagocytic cells that ingest pathogens that they
encounter [64]. This age-related reduction may indicate a reduction in phagocytosis (ingestion of
pathogens by classical monocytes) and thus can possibly increase infections in older individuals.
In addition, older individuals had higher proportions of CD14+/CD16+ intermediate monocytes,
which are strongly associated with inflammation [65]. For instance, an increase in this cell type
has been linked to disorders such as chronic kidney disease [66]. The decrease in classical
monocytes, together with an increase in intermediate monocytes, represents yet another signature
of immunosenescence and inflammaging.

One of the strengths of our study system was the ability to quantify social adversity,
operationalized as social status, and test if and how social status influenced immune variation and
whether the effects of status varied with age and/or sex. We found no main effect of social status
on the proportion of immune cell types. Also, there was no interaction between social status and
age on the proportions of immune cell types. This result was contrary to our expectations because
we expected low social status individuals to experience more variation in immune cell types with
increasing age. Nevertheless, we found several interactions between social status and sex, as well
as a within-sex main effect of social status on immune cell composition, possibly reflecting the
different pathways through which social status is acquired in males and females and thus
highlighting the fact that different sexes experience social adversity differently across the
hierarchy.

The interaction of social status and sex influenced adaptive and innate immune cell types such as

CD20+/CD3+ ratio and CD8+ T cells and CD3-CD16+ NK cells, where the proportions of these

cell types associated with status depended on the sex of the individual. In humans, social stressors,

such as lower socioeconomic status and lower subjective social status, can affect cytokine release

and inflammatory responses in peripheral blood mononuclear cells in a sex-dependent manner [67-

68]. However, studies in humans that have looked at the interaction between social stressors (such

as socioeconomic status) and sex on immune cell proportions have found no significant interaction

between these covariates [22], thus making our study unique in reporting sex-dependent effects of

social status.

We also found a significant main effect of social status in the within-sex analysis on the CD20+ B

cell/CD3+ ratio, with high social status males having significantly higher ratios than low status

males. The decrease in the CD20+/CD3+ ratio seems to be driven by a decrease in the proportions

of CD3+ T cells in low social status males (**S. Figure 5**) compared to high status males. Decreases

in this cell type have been associated with decreases in cell-mediated immunity to bacteria and

viruses [69-70], potentially showing that the T cell response in macaques is negatively affected by

low social status. In addition, CD8+ T cell proportions were higher in low social status males

compared to high social status macaques. Few prior studies have assessed sociality-related immune

cell differences in male rhesus [71], likely because of ethical and husbandry constraints, such as

aggressive behavior between males. One study in male Barbary macaques (*Macaca sylvanus*)

reported that males with strong social bonds had lower levels of fecal glucocorticoids [72], which

is typically associated with reduced inflammation [73]. Additionally, studies in cynomolgus

macaques (*Macaca fascicularis*) have shown that low social status males had a higher probability

of being infected with a virus than did high social status macaques [74]. These findings should be

taken with caution, however, as other studies of macaques (rhesus and other species) found no

differences in infection rate or immune responses between high and low status males [75-76].

Although there are currently no data associating social status (or other social stressors) with CD8+

T cells in rhesus macaques, there are reports in other species that CD8+ T cells can mediate the

release of proinflammatory cytokines during stressful conditions [77]. Our finding of higher

proportions of CD8+ T cells in low social status macaques might indicate higher levels of baseline

cytotoxic T cell activation, potentially affecting the CD4+ T cell response. Testing this idea will

require methods such as cytokine analysis or next generation sequencing.

There was also a main effect of social status on the proportion of CD8+ T cells in females, but in

contrast to males, high social status females had significantly higher proportions of this cell type

compared to low social status females. One study also reported lower proportions of CD8+ T cells

in low social status in non-free-ranging female rhesus macaques [34]. Given that females tend to

have lower proportions of CD8+ T cells than males regardless of age [78-80], a lower proportion

of this cell type in low social status females might indicate lower cytotoxic immunity at baseline.

Female social status also had a main effect on the proportion of CD3-CD16+ NK cells (associated

with immune surveillance, inflammation and innate responses), with low social status females

having significantly higher proportions of this cell type than high social status females. Although

a prior study found that the proportion of CD3-CD16+ NK cells did not vary with social status in

female rhesus macaques, it did find that this cell type was the most sensitive to social status.

Specifically, low social status females showed patterns of gene expression consistent with a

proinflammatory phenotype in this cell type in response to lipopolysaccharide [81]. These results

highlight that low status female rhesus macaque may experience higher levels of basal

inflammation, consistent with other studies in this species [81-83].

In conclusion, our results demonstrate that, at the level of circulating immune cell proportions,

macaques and humans show similar age-related variation in immune cell types. Although we did

not detect any significant main effects of sex or sex-age interaction, it is possible that more specific,

but unmeasured adaptive immune cells, such as the effector and memory subsets of B cells and T

cells, could differ between males and females. In future studies, it will be important to measure

other innate immune cell types, such as dendritic cells and granulocytes, since these cell types are

critical for antigen presentation and the development of adaptive immune response. We found that

the effects of social status differed between males and females, which is likely due to sex-

differences in how rhesus macaques obtain social status. Specifically, females inherit their social

status, which remains relatively stable throughout their lives, while males queue and occasionally

fight to establish and maintain their social status, which may lead to stronger effects of status on

immune cell distribution and function. Overall, our study provides detailed insights into the

impacts of social and demographic variation on immune cell status in a non-human primate model

with unparalleled translatability to humans. Future research should quantify the proportions of

these cell types as a function of age using a longitudinal approach, which will require sampling

individuals over the course of years. Immune stimulation tests would also be informative by testing

whether both the age-associated and status-associated differences in immune cell types translate

to immune function.

Author contributions

M.R.S.R., J.P.H., L.J.N.B., C.A.S., M.J.M, M.L.P., and N.S.-M designed research; M.R.S.R.,

N.M.R., M.M.W., A.D.N.-D, P.P., M.A.P.-F., E.R.S., E.B.C., J.E.N.-D., D.P., A.R.L., M.J.M. and

CBRU performed research; M.R.S.R and N.S.-M. analyzed data; and M.S.R. and N.S.-M. wrote

the manuscript. All authors reviewed and revised the manuscript. CBRU members: Lauren J.N.

Brent, Jampes P. Higham, Noah Snyder-Mackler, Michael J. Montague, Michael L. Platt,

Melween I. Martinez, Susan C. Antón, Amanda D. Melin, Jérôme Sallet

The authors declare no competing interest.

Acknowledgements:

Part of Figure 1, panel A, was created with BioRender.com.

We thank the Caribbean Primate Research Center for their help in collecting data, especially

Giselle Carabllo Cruz, and Nahiri Rivera Barreto; Crisanta Serrano and Stephanie Dorta for

laboratory advice; Corbin Johnson, Laura Newman, Alice Baniel, India Schneider-Crease,

Kenneth Chiou, Trisha Zintel, and other members of the Snyder-Mackler, Sariol, Higham, Platt,

and Brent labs for helpful discussions during the course of this work.

Funding:

This work was supported by the National Institutes of Health (R01-AG060931; R01AG060931-

S1 R00-AG051764; R01-MH118203; R01-MH096875; R56-AG071023; C06-OD026690; P40-

OD012217; NSF-1800558; ERC-864461).

Ethical note:

This work was approved by the Institutional Animal Care and Use Committees of the University

of Puerto Rico, Medical Sciences Campus (IACUC Number: A400117).

The authors declare no competing interest.

References

1. Max Roser, Esteban Ortiz-Ospina and Hannah Ritchie (2013) - "Life Expectancy". Published

online at OurWorldInData.org. Retrieved from: 'https://ourworldindata.org/life-expectancy'

[Online Resource]

2. Strait, J. B., & Lakatta, E. G. (2012). Aging-associated cardiovascular changes and their

relationship to heart. https://doi.org/10.1016/j.hfc.2011.08.011

3. Yung, R. L., & Julius, A. (2008). Epigenetics, aging, and autoimmunity. *Autoimmunity*, *41*(4),

329-335. https://doi.org/10.1080/08916930802024889

4. Halim, M., & Halim, A. (2019). The effects of inflammation, aging and oxidative stress on the

pathogenesis of diabetes mellitus (type 2 diabetes). *Diabetes & metabolic syndrome: clinical*

research & reviews, *13*(2), 1165-1172. https://doi.org/10.1016/j.dsx.2019.01.040

5. Sacitharan, P. K. (2019). Ageing and osteoarthritis. *Biochemistry and cell biology of ageing:*

part II clinical science, 123-159. DOI: 10.1007/978-981-13-3681-2_6

6. Koh, S. H., Choi, S. H., Jeong, J. H., Jang, J. W., Park, K. W., Kim, E. J., ... & Kang, S. (2020).

Telomere shortening reflecting physical aging is associated with cognitive decline and dementia

conversion in mild cognitive impairment due to Alzheimer's disease. *Aging (Albany NY)*, *12*(5),

4407. https://doi.org/10.18632/aging.102893

7. Ferrucci, L., & Fabbri, E. (2018). Inflammageing: chronic inflammation in ageing,

cardiovascular disease, and frailty. *Nature Reviews Cardiology*, *15*(9), 505-

522. https://doi.org/10.1038/s41569-018-0064-2

8. Aw, D., Silva, A. B., & Palmer, D. B. (2007). Immunosenescence: emerging challenges for an

ageing population. *Immunology*, *120*(4), 435-446. 10.1111/j.1365-2567.2007.02555.x

9. Hearps, A. C., Martin, G. E., Angelovich, T. A., Cheng, W. J., Maisa, A., Landay, A. L., ... &

Crowe, S. M. (2012). Aging is associated with chronic innate immune activation and dysregulation

of monocyte phenotype and function. *Aging cell*, *11*(5), 867-875. https://doi.org/10.1111/j.1474-

9726.2012.00851.x

10. Giefing-Kröll, C., Berger, P., Lepperdinger, G., & Grubeck-Loebenstein, B. (2015). How sex

and age affect immune responses, susceptibility to infections, and response to vaccination. *Aging*

cell, *14*(3), 309-321. https://doi.org/10.1111/acel.12326

11. Gubbels Bupp, M. R., Potluri, T., Fink, A. L., & Klein, S. L. (2018). The confluence of sex

hormones and aging on immunity. *Frontiers in immunology*, *9*, 1269.

https://doi.org/10.3389/fimmu.2018.01269

12. Klein, S. L., & Flanagan, K. L. (2016). Sex differences in immune responses. *Nature Reviews*

Immunology, *16*(10), 626-638. https://doi.org/10.1038/nri.2016.90

13. Guerra-Silveira, F., & Abad-Franch, F. (2013). Sex bias in infectious disease epidemiology:

patterns and processes. *PloS one*, *8*(4), e62390. https://doi.org/10.1371/journal.pone.0062390

14. Pantell, M., Rehkopf, D., Jutte, D., Syme, S. L., Balmes, J., & Adler, N. (2013). Social

isolation: a predictor of mortality comparable to traditional clinical risk factors. American journal

of public health, 103(11), 2056-2062. https://doi.org/10.2105/AJPH.2013.301261

15. Cole, S. W. (2014). Human social genomics. *PLoS genetics*, *10*(8), e1004601.

https://doi.org/10.1371/journal.pgen.1004601

16. L. F. Berkman, T. Glass, I. Brissette, T. E. Seeman, From social integration to health:

Durkheim in the new millennium. Soc. Sci. Med. 51, 843–857 (2000). doi: 10.1016/S0277-9536

(00)00065-4; pmid: 10972429

17. Anderson, J. A., Johnston, R. A., Lea, A. J., Campos, F. A., Voyles, T. N., Akinyi, M. Y., ...

& Tung, J. (2021). High social status males experience accelerated epigenetic aging in wild

baboons. *Elife*, *10*, e66128. https://doi.org/10.7554/eLife.66128

18. Fiorito, G., Polidoro, S., Dugué, P. A., Kivimaki, M., Ponzi, E., Matullo, G., ... & Vineis, P.

(2017). Social adversity and epigenetic aging: a multi-cohort study on socioeconomic differences

in peripheral blood DNA methylation. *Scientific reports*, *7*(1), 1-12.

https://doi.org/10.1038/s41598-017-16391-5

19. Ridout, K. K., Levandowski, M., Ridout, S. J., Gantz, L., Goonan, K., Palermo, D., ... & Tyrka,

A. R. (2018). Early life adversity and telomere length: a meta-analysis. *Molecular psychiatry*,

23(4), 858-871. https://doi.org/10.1038/mp.2017.26

20. Snyder-Mackler, N., Somel, M., & Tung, J. (2014). Shared signatures of social stress and aging

in peripheral blood mononuclear cell gene expression profiles. *Aging Cell*, *13*(5), 954-957.

https://doi.org/10.1111/acel.12239

21. Elwenspoek, M. M., Hengesch, X., Leenen, F. A., Schritz, A., Sias, K., Schaan, V. K., ... &

Muller, C. P. (2017). Proinflammatory T cell status associated with early life adversity. *The*

Journal of Immunology, *199*(12), 4046-4055. https://doi.org/10.4049/jimmunol.1701082

22. Klopack, Eric T., Eileen M. Crimmins, Steve W. Cole, Teresa E. Seeman, and Judith E. Carroll.

2022. "Social Stressors Associated with Age-Related T Lymphocyte Percentages in Older US

Adults: Evidence from the US Health and Retirement Study." *Proceedings of the National*

Academy of Sciences of the United States of America 119 (25): e2202780119.

https://doi.org/10.1073/pnas.2202780119

23. Roth, G. S., Mattison, J. A., Ottinger, M. A., Chachich, M. E., Lane, M. A., & Ingram, D. K.

(2004). Aging in rhesus monkeys: relevance to human health interventions. *Science*, *305*(5689),

1423-1426. https://doi.org/10.1126/science.1102541

24. Chiou, K. L., Montague, M. J., Goldman, E. A., Watowich, M. M., Sams, S. N., Song, J., ... &

Snyder-Mackler, N. (2020). Rhesus macaques as a tractable physiological model of human ageing.

Philosophical Transactions of the Royal Society B, *375*(1811), 20190612.

https://doi.org/10.1098/rstb.2019.0612

25. Thierry, B., Singh, M., & Kaumanns, W. (Eds.). (2004). *Macaque societies: a model for the*

study of social organization (Vol. 41). Cambridge University Press.

26. Brent, L. J., MacLarnon, A., Platt, M. L., & Semple, S. (2013). Seasonal changes in the

structure of rhesus macaque social networks. *Behavioral Ecology and Sociobiology*, *67*(3), 349-

359. https://doi.org/10.1007/s00265-012-1455-8

27. Kulik, L., Amici, F., Langos, D., & Widdig, A. (2015). Sex differences in the development of

social relationships in rhesus macaques (Macaca mulatta). *International journal of primatology*,

36(2), 353-376. https://doi.org/10.1007/s10764-015-9826-4

28. Vandeleest, J. J., Winkler, S. L., Beisner, B. A., Hannibal, D. L., Atwill, E. R., & McCowan,

B. (2020). Sex differences in the impact of social status on hair cortisol concentrations in rhesus

monkeys (Macaca mulatta). *American journal of primatology*, *82*(1), e23086.

https://doi.org/10.1002/ajp.23086

29. Datta, S. (1988). The acquisition of dominance among free-ranging rhesus monkey siblings.

Animal Behaviour, *36*(3), 754-772. https://doi.org/10.1016/S0003-3472(88)80159-3

30. Snyder-Mackler, N., Burger, J. R., Gaydosh, L., Belsky, D. W., Noppert, G. A., Campos, F.

A., ... & Tung, J. (2020). Social determinants of health and survival in humans and other animals.

Science, *368*(6493). https://doi.org/10.1126/science.aax9553

31. Kohn, J. N., Snyder-Mackler, N., Barreiro, L. B., Johnson, Z. P., Tung, J., & Wilson, M. E.

(2016). Dominance rank causally affects personality and glucocorticoid regulation in female

rhesus macaques. *Psychoneuroendocrinology*, *74*, 179-188. https://doi.org/10.1016/j.psyneuen.2016.09.005

32. Debray, R., Snyder-Mackler, N., Kohn, J. N., Wilson, M. E., Barreiro, L. B., & Tung, J. (2019). Social affiliation predicts mitochondrial DNA copy number in female rhesus macaques. *Biology letters*, *15*(1), 20180643. https://doi.org/10.1098/rsbl.2018.0643

33. Pavez-Fox, M. A., Kimock, C. M., Rivera-Barreto, N., Negron-Del Valle, J. E., Phillips, D., Ruiz-Lambides, A., ... & Brent, L. J. (2022). Reduced injury risk links sociality to survival in a group-living primate. *Iscience*, *25*(11), 105454. https://doi.org/10.1016/j.isci.2022.105454

34. Blomquist, G. E., Sade, D. S., & Berard, J. D. (2011). Rank-related fitness differences and their demographic pathways in semi-free-ranging rhesus macaques (Macaca mulatta). *International Journal of Primatology*, *32*, 193-208. https://doi.org/10.1007/s10764-010-9461-z

35. Tung, J., Barreiro, L. B., Johnson, Z. P., Hansen, K. D., Michopoulos, V., Toufexis, D., ... & Gilad, Y. (2012). Social environment is associated with gene regulatory variation in the rhesus macaque immune system. *Proceedings of the National Academy of Sciences*, *109*(17), 6490-6495. https://doi.org/10.1073/pnas.1202734109

36. Kessler, M. J., & Rawlins, R. G. (2016). A 75-year pictorial history of the Cayo Santiago rhesus monkey colony. *American Journal of Primatology*, *78*(1), 6-43. doi: 10.1002/ajp.22381

37. Clutton-Brock, T. (2016). *Mammal societies*. John Wiley & Sons. https://doi.org/10.1093/jmammal/gyx078

38. Cerroni, A. M., Tomlinson, G. A., Turnquist, J. E., & Grynpas, M. D. (2000). Bone mineral density, osteopenia, and osteoporosis in the rhesus macaques of Cayo Santiago. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, *113*(3), 389-410. https://doi.org/10.1002/1096-8644(200011)113:3<389::AID-AJPA9>3.0.CO;2-I

39. Lee, D. S., Kang, Y. H., Ruiz-Lambides, A. V., & Higham, J. P. (2021). The observed pattern and hidden process of female reproductive trajectories across the life span in a non-human primate. *Journal of Animal Ecology*, *90*(12), 2901-2914. https://doi.org/10.1111/1365-2656.13590

40. Watowich, M. M., Chiou, K. L., Montague, M. J., Cayo Biobank Research Unit, Simons, N. D., Horvath, J. E., ... & Snyder-Mackler, N. (2022). Natural disaster and immunological aging in a nonhuman primate. *Proceedings of the National Academy of Sciences*, *119*(8), e2121663119. https://doi.org/10.1073/pnas.2121663119

41. Cooper, E. B., Brent, L. J., Snyder-Mackler, N., Singh, M., Sengupta, A., Khatiwada, S., ... &

Higham, J. P. (2022). The Natural History of Model Organisms: The rhesus macaque as a success

story of the Anthropocene. *Elife*, *11*, e78169-e78169. 10.7554/elife.78169

42. Rawlins, R. G., & Kessler, M. J. (Eds.). (1986). *The Cayo Santiago macaques: History,*

behavior, and biology. SUnY Press.

43. Pérez-Guzmán, E. X., Pantoja, P., Serrano-Collazo, C., Hassert, M. A., Ortiz-Rosa, A.,

Rodríguez, I. V., ... & Sariol, C. A. (2019). Time elapsed between Zika and dengue virus infections

affects antibody and T cell responses. *Nature communications*, *10*(1), 1-14.

https://doi.org/10.1038/s41467-019-12295-2

44. Marzan-Rivera, N., Serrano-Collazo, C., Cruz, L., Pantoja, P., Ortiz-Rosa, A., Arana, T., ... &

Sariol, C. A. (2022). Infection order outweighs the role of CD4+ T cells in tertiary flavivirus

exposure. *Iscience*, *25*(8), 104764. https://doi.org/10.1016/j.isci.2022.104764

45. Asiedu, C. K., Goodwin, K. J., Balgansuren, G., Jenkins, S. M., Le Bas-Bernardet, S., Jargal,

U., ... & Thomas, J. M. (2005). Elevated T regulatory cells in long-term stable transplant tolerance

in rhesus macaques induced by anti-CD3 immunotoxin and deoxyspergualin. *The Journal of*

Immunology, *175*(12), 8060-8068. DOI: https://doi.org/10.4049/jimmunol.175.12.8060

46. Morita, D., Hattori, Y., Nakamura, T., Igarashi, T., Harashima, H., & Sugita, M. (2013). Major

T cell response to a mycolyl glycolipid is mediated by CD1c molecules in rhesus macaques.

Infection and immunity, *81*(1), 311-316. https://doi.org/10.1128/IAI.00871-12

47. Van Noordwijk, M. A., & van Schaik, C. P. (2004). Sexual selection and the careers of primate

males: paternity concentration, dominance–acquisition tactics and transfer decision. *Sexual*

selection in primates: New and comparative perspectives, 208-229.

https://doi.org/10.1017/CBO9780511542459.014

48. Kimock, C. M., Dubuc, C., Brent, L. J., & Higham, J. P. (2019). Male morphological traits are

heritable but do not predict reproductive success in a sexually-dimorphic primate. *Scientific*

reports, *9*(1), 1-11. https://doi.org/10.1038/s41598-019-52633-4

49. Brent, L. J. N., Semple, S., Dubuc, C., Heistermann, M., & MacLarnon, A. (2011). Social

capital and physiological stress levels in free-ranging adult female rhesus macaques. *Physiology*

& behavior, *102*(1), 76-83. https://doi.org/10.1016/j.physbeh.2010.09.022

50. Madlon-Kay, S., Brent, L., Montague, M., Heller, K., & Platt, M. (2017). Using machine
learning to discover latent social phenotypes in free-ranging macaques. *Brain sciences*, *7*(7), 91.
https://doi.org/10.3390/brainsci7070091

51. R Core Team (2021). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

52. Vignali, D. A., Collison, L. W., & Workman, C. J. (2008). How regulatory T cells work. *Nature*
reviews immunology, *8*(7), 523-532. https://doi.org/10.1038/nri2343

53. Idzkowska, E., Eljaszewicz, A., Miklasz, P., Musial, W. J., Tycinska, A. M., & Moniuszko,
M. (2015). The role of different monocyte subsets in the pathogenesis of atherosclerosis and acute
coronary syndromes. *Scandinavian journal of immunology*, *82*(3), 163-173.
https://doi.org/10.1111/sji.12314

54. Erkeller-Yuksel, F. M., Deneys, V., Yuksel, B., Hannet, I., Hulstaert, F., Hamilton, C., ... &
Lydyard, P. M. (1992). Age-related changes in human blood lymphocyte subpopulations. *The*
Journal of pediatrics, *120*(2), 216-222. https://doi.org/10.1016/S0022-3476(05)80430-5

55. Frasca, D., Blomberg, B. B., Garcia, D., Keilich, S. R., & Haynes, L. (2020). Age-related
factors that affect B cell responses to vaccination in mice and humans. *Immunological reviews*,
296(1), 142-154. https://doi.org/10.1111/imr.12864

56. Germain, C., Gnjatic, S., Tamzalit, F., Knockaert, S., Remark, R., Goc, J., ... & Dieu-Nosjean,
M. C. (2014). Presence of B cells in tertiary lymphoid structures is associated with a protective
immunity in patients with lung cancer. *American journal of respiratory and critical care medicine*,
189(7), 832-844. https://doi.org/10.1164/rccm.201309-1611OC

57. Asquith, M., Haberthur, K., Brown, M., Engelmann, F., Murphy, A., Al-Mahdi, Z., &
Messaoudi, I. (2012). Age-dependent changes in innate immune phenotype and function in rhesus
macaques (Macaca mulatta). *Pathobiology of Aging & Age-related Diseases*, *2*(1), 18052.
https://doi.org/10.3402/pba.v2i0.18052

58. Zheng, H. Y., Zhang, M. X., Pang, W., & Zheng, Y. T. (2014). Aged Chinese rhesus macaques
suffer severe phenotypic T-and B-cell aging accompanied with sex differences. *Experimental*
gerontology, *55*, 113-119. https://doi.org/10.1016/j.exger.2014.04.004

59. Quinn, K. M., Fox, A., Harland, K. L., Russ, B. E., Li, J., Nguyen, T. H., ... & La Gruta, N. L.
(2018). Age-related decline in primary CD8+ T cell responses is associated with the development

of senescence in virtual memory CD8+ T cells. *Cell reports*, *23*(12), 3512-3524. https://doi.org/10.1016/j.celrep.2018.05.057

60. Li, M., Yao, D., Zeng, X., Kasakovski, D., Zhang, Y., Chen, S., ... & Xu, L. (2019). Age related human T cell subset evolution and senescence. *Immunity & Ageing*, *16*(1), 1-7. https://doi.org/10.1186/s12979-019-0165-8

61. Mogilenko, D. A., Shpynov, O., Andhey, P. S., Arthur, L., Swain, A., Esaulova, E., ... & Artyomov, M. N. (2021). Comprehensive profiling of an aging immune system reveals clonal GZMK+ CD8+ T cells as conserved hallmark of inflammaging. *Immunity*, *54*(1), 99-115. https://doi.org/10.1016/j.immuni.2020.11.005

62. Meza Guzman, L. G., Keating, N., & Nicholson, S. E. (2020). Natural killer cells: tumor surveillance and signaling. *Cancers*, *12*(4), 952. doi:10.3390/cancers12040952

63. Rocamora-Reverte, L., Melzer, F. L., Würzner, R., & Weinberger, B. (2020). The complex role of regulatory T cells in immunity and aging. *Frontiers in Immunology*, *11*. https://doi.org/10.3389/fimmu.2020.616949

64. Kapellos, T. S., Bonaguro, L., Gemünd, I., Reusch, N., Saglam, A., Hinkley, E. R., & Schultze, J. L. (2019). Human monocyte subsets and phenotypes in major chronic inflammatory diseases. *Frontiers in immunology*, *10*, 2035. https://doi.org/10.3389/fimmu.2019.02035

65. Italiani, P., & Boraschi, D. (2014). From monocytes to M1/M2 macrophages: phenotypical vs. functional differentiation. *Frontiers in immunology*, *5*, 514. https://doi.org/10.3389/fimmu.2014.00514

66. Naicker, S. D., Cormican, S., Griffin, T. P., Maretto, S., Martin, W. P., Ferguson, J. P., ... & Griffin, M. D. (2018). Chronic kidney disease severity is associated with selective expansion of a distinctive intermediate monocyte subpopulation. *Frontiers in immunology*, *9*, 2845. https://doi.org/10.3389/fimmu.2018.02845

67. Moieni, M., Muscatell, K. A., Jevtic, I., Breen, E. C., Irwin, M. R., & Eisenberger, N. I. (2019). Sex differences in the effect of inflammation on subjective social status: a randomized controlled trial of endotoxin in healthy young adults. *Frontiers in psychology*, *10*, 2167. https://doi.org/10.3389/fpsyg.2019.02167

68. Gassen, J., White, J. D., Peterman, J. L., Mengelkoch, S., Proffitt Leyva, R. P., Prokosch, M. L., ... & Hill, S. E. (2021). Sex differences in the impact of childhood socioeconomic status on immune function. *Scientific reports*, *11*(1), 9827. https://doi.org/10.17605/OSF.IO/DXPZU

69. Monserrat, J., de Pablo, R., Reyes, E., Díaz, D., Barcenilla, H., Zapata, M. R., ... & Álvarez-
Mon, M. (2009). Clinical relevance of the severe abnormalities of the T cell compartment in septic
shock patients. *Critical care*, *13*(1), 1-8. https://doi.org/10.1186/cc7731

70. Chinen, J., Easley, K. A., Mendez, H., & Shearer, W. T. (2001). Decline of CD3-positive T-
cell counts by 6 months of age is associated with rapid disease progression in HIV-1–infected
infants. *Journal of allergy and clinical immunology*, *108*(2), 265-268. 10.1067/mai.2001.116573

71. Pavez-Fox, M. A., Negron-Del Valle, J. E., Thompson, I. J., Walker, C. S., Bauman, S. E.,
Gonzalez, O., ... & Brent, L. J. (2021). Sociality predicts individual variation in the immunity of
free-ranging rhesus macaques. *Physiology & behavior*, *241*, 113560.
https://doi.org/10.1016/j.physbeh.2021.113560

72. Young, C., Majolo, B., Heistermann, M., Schülke, O., & Ostner, J. (2014). Responses to social
and environmental stress are attenuated by strong male bonds in wild macaques. *Proceedings of*
the National Academy of Sciences, *111*(51), 18195-18200.
https://doi.org/10.1073/pnas.1411450111

73. Adcock, I.M., Mumby, S. (2016). Glucocorticoids. In: Page, C., Barnes, P. (eds) Pharmacology
and Therapeutics of Asthma and COPD. Handbook of Experimental Pharmacology, vol 237.
Springer, Cham. https://doi.org/10.1007/164_2016_98

74. Cohen, S., Line, S., Manuck, S. B., Rabin, B. S., Heise, E. R., & Kaplan, J. R. (1997). Chronic
social stress, social status, and susceptibility to upper respiratory infections in nonhuman primates.
Psychosomatic medicine, *59*(3), 213-221. 10.1097/00006842-199705000-00001

75. McAuliffe, J., Vogel, L., Roberts, A., Fahle, G., Fischer, S., Shieh, W. J., ... & Subbarao, K.
(2004). Replication of SARS coronavirus administered into the respiratory tract of African Green,
rhesus and cynomolgus monkeys. *Virology*, *330*(1), 8-15.
https://doi.org/10.1016/j.virol.2004.09.030

76. Skinner, J. M., Caro-Aguilar, I. C., Payne, A. M., Indrawati, L., Fontenot, J., & Heinrichs, J.
H. (2011). Comparison of rhesus and cynomolgus macaques in a Streptococcus pyogenes infection
model for vaccine evaluation. *Microbial pathogenesis*, *50*(1), 39-47.
10.1016/j.micpath.2010.10.004

77. Clark, S. M., Song, C., Li, X., Keegan, A. D., & Tonelli, L. H. (2019). CD8+ T cells promote
cytokine responses to stress. *Cytokine*, *113*, 256-264. 10.1016/j.cyto.2018.07.015

78. Lee, B. W., Yap, H. K., Chew, F. T., Quah, T. C., Prabhakaran, K., Chan, G. S., ... & Seah, C.

C. (1996). Age-and sex-related changes in lymphocyte subpopulations of healthy Asian subjects:

From birth to adulthood. *Cytometry: The Journal of the International Society for Analytical*

Cytology, *26*(1), 8-15. https://doi.org/10.1002/(SICI)1097-0320(19960315)26:1<8::AID-

CYTO2>3.0.CO;2-E

79. Lisse, I. M., Aaby, P., Whittle, H., Jensen, H., Engelmann, M., & Christensen, L. B. (1997).

T-lymphocyte subsets in West African children: impact of age, sex, and season. *The Journal of*

pediatrics, *130*(1), 77-85. https://doi.org/10.1016/S0022-3476(97)70313-5

80. Uppal, S. S., Verma, S., & Dhot, P. S. (2003). Normal values of CD4 and CD8 lymphocyte

subsets in healthy Indian adults and the effects of sex, age, ethnicity, and smoking. *Cytometry Part*

B: Clinical Cytometry: The Journal of the International Society for Analytical Cytology, *52*(1),

32-36. https://doi.org/10.1002/cyto.b.10011

81. Snyder-Mackler, N., Sanz, J., Kohn, J. N., Brinkworth, J. F., Morrow, S., Shaver, A. O., ... &

Barreiro, L. B. (2016). Social status alters immune regulation and response to infection in

macaques. Science, 354(6315), 10411045. https://doi.org/10.1126/science.aah3580

82. Sanz, J., Maurizio, P. L., Snyder-Mackler, N., Simons, N. D., Voyles, T., Kohn, J., ... &

Barreiro, L. B. (2020). Social history and exposure to pathogen signals modulate social status

effects on gene regulation in rhesus macaques. *Proceedings of the National Academy of Sciences*,

117(38), 23317-23322. https://doi.org/10.1073/pnas.1820846116

83. Snyder-Mackler, N., Sanz, J., Kohn, J. N., Voyles, T., Pique-Regi, R., Wilson, M. E., ... &

Tung, J. (2019). Social status alters chromatin accessibility and the gene regulatory response to

glucocorticoid stimulation in rhesus macaques. *Proceedings of the National Academy of Sciences*,

116(4), 1219-1228. https://doi.org/10.1073/pnas.1811758115

844