Dispersal syndrome and landscape fragmentation in the salt-marsh specialist spider *Erigone longipalpis*

Maxime Dahirel[1,2]*, Marie Wullschleger[1], Tristan Berry[1,3], Solène Croci[4], Julien Pétillon[1,5]

[1]Univ Rennes, UR1, CNRS, ECOBIO (Ecosystèmes, Biodiversité, évolution), UMR 6553, F-35000, Rennes, France

[2]Department of Biology, Ghent University, B-9000, Ghent, Belgium

[3]Syndicat Mixte de Gestion des Milieux Naturels, Réserve Naturelle Nationale de l'Étang Noir, F-40510, Seignosse, France

[4]CNRS, Université de Rennes 2, EPHE-PSL, Université d'Angers, Université de Bretagne Occidentale, Université de Caen Normandie, Université de Nantes, UMR LETG, F-35043, Rennes, France

[5]Institute for Coastal and Marine Research, Nelson Mandela University, Port Elizabeth, South Africa

*corresponding author: maxime.dahirel@yahoo.fr

Abstract

Dispersal and its evolution play a key role for population persistence in fragmented landscapes where habitat loss and fragmentation increase the cost of between-habitat movements. In such contexts, it is important to know how variation in dispersal and other traits is structured, and whether responses to landscape fragmentation are aligned with underlying dispersal-trait correlations, or dispersal syndromes. We therefore studied trait variation in *Erigone longipalpis*, a spider species specialist of (often patchy) salt marshes. We collected spiders in two salt-marsh landscapes differing in habitat availability. We then reared lab-born spiders for two generations in controlled conditions, and measured dispersal and its association with various key traits. *E. longipalpis* population densities were lower in the more fragmented landscape. Despite this, we found no evidence of differences in dispersal, or any other trait we studied, between the two landscapes. While a dispersal syndrome was present at the among-individual level (dispersers were more fecund and faster growing, among others), there was no indication it was genetically driven: among-family differences in dispersal were not correlated with differences in other traits. Instead, we showed that the observed phenotypic covariations were mostly due to within-family correlations. We hypothesize that the dispersal syndrome is the result of asymmetric food access among siblings, leading to variation in development rates and carrying over to adult traits. Our results show we need to better understand the sources of dispersal variation and syndromes, especially when dispersal may evolve rapidly in response to environmental change.

Keywords: habitat loss; Linyphiidae; multivariate mixed model; phenotypic correlations; population density

Introduction

Dispersal is a key trait and process that influences, and links together, ecological and evolutionary dynamics (Clobert et al. 2012; Govaert et al. 2019). Individuals' movement between habitats shapes (meta-)population dynamics (Benton and Bowler 2012) while the resulting gene flow can have negative or positive impacts on local adaptation (Garant et al. 2007). Many "dispersal syndromes", i.e. associations/covariations between dispersal and other traits, have been documented across the tree of life (Ronce and Clobert 2012; Stevens et al. 2014; Jacob et al. 2019). These covariations may result both from genetic correlations, caused by pleiotropy or joint selective responses across traits, or from plastic responses to experienced environmental conditions (Ronce and Clobert 2012). These syndromes reinforce the role of dispersal as a nexus between ecological and evolutionary processes. Indeed, the existence of these syndromes mean that dispersal not only redistributes species, individuals and genetic diversity in space, but does it non-randomly with respect to trait values, which has potential consequences for instance for the distribution of ecosystem functions in landscapes (Massol et al. 2017; Little et al. 2019) or for the dynamics of range expansions (Ochocki et al. 2020).

Some general dispersal syndromes have been proposed on theoretical grounds. These include, for instance, expectations of a positive association between high dispersal and the "fast" end of a pace-of-life life history axis (Réale et al. 2010; Stevens et al. 2012; Wright et al. 2019), or the trade-off between fecundity and dispersal often seen in insects (Guerra, 2011). However, in practice, observed syndromes are much more diverse and sometimes contradict these general predictions, especially at the within-species level (Guerra 2011; Ronce and Clobert 2012; Bonte and Dahirel 2017), making generalisations difficult. In the case of dispersal-pace of life syndromes, one possible reason for divergence among studies may be that within-species variation in life history does not always align nicely along a main pace of life axis, and that its existence should be tested rather than assumed (Royauté et al. 2018).

Habitat loss and habitat fragmentation (*sensu stricto*; the increasing isolation between habitat patches) are usually seen as a particularly strong selective pressure on dispersal, as they increase the (many) costs associated with moving from a relatively suitable habitat to another

(Bonte et al. 2012; Cote et al. 2017). Indeed, a reduction of dispersal is often predicted and

observed in response to fragmentation *sensu lato* (i.e. conflating habitat loss and fragmentation

sensu stricto together, Cote et al. 2017). Therefore, if dispersal is integrated with other traits in

syndromes, we may then expect these other traits to also evolve jointly in response to habitat

change. If such syndromes have a genetic basis and act as constraints on evolution (see e.g.

Royauté et al. 2020), then the direction of these evolutionary changes should be somewhat

predictable from syndrome structure.

Here we present a study of dispersal and dispersal syndromes in *Erigone longipalpis* (Sundevall,

1830) (fam. Linyphiidae) spiders. This species is a strong habitat specialist, only found in wet

habitats like floodplains and especially salt marshes (Harvey et al. 2002; Pétillon et al. 2008). At

the regional and continental scales, these landscapes are patchily distributed along the coastal

line (European Environmental Agency, 2020); in additions, within salt marshes, *E. longipalpis* is

restricted to specific vegetation types that are themselves patchy, in part due to human

management (Pétillon et al. 2007; Leroy et al. 2014). As a consequence, we may expect its

dispersal responses to be strongly influenced by habitat fragmentation, much more than related

generalist species; indeed, any given landscape likely contains less favourable habitat for

specialists than generalists. In addition, *E. longipalpis* is one of the most important invertebrate

predators in these habitats (Leroy et al. 2014), so understanding how dispersal influences its

spatial population dynamics may help understand the dynamics of its potential prey resources

at the same time (Fronhofer et al. 2018). Finally, spiders and especially linyphiids have been key

models in dispersal ecology and evolution (Bonte, 2012; Bonte and Saastamoinen, 2012), in part

because they exhibit stereotypical behaviours associated with dispersal (whether long-distance

dispersal by ballooning or short-distance movements by rappelling); dispersal-related

behaviours can thus easily be tracked in an experimental setting (e.g. De Meester and Bonte,

2010). We used lab-born spiders coming from several patches in two landscapes differing in

fragmentation degree to test the following hypotheses:

• Rapid human-induced habitat loss and fragmentation have negative impacts on *Erigone*

longipalpis populations, and also lead to reduced dispersal propensity;

- Dispersal is associated with other life-history traits in a syndrome. Specifically, high dispersal propensity is associated with "faster" life-histories (e.g. faster development time, higher fecundity), following predictions derived from the pace of life hypothesis (Réale et al. 2010; Wright et al. 2019);

- As syndromes can constrain trait evolution (e.g. Royauté et al. 2020), and dispersal evolves with fragmentation, the direction in which other traits evolve can be predicted from the direction of dispersal evolution and the structure of the among-family (potentially genetic) dispersal syndrome.

Material and Methods

Site selection and field sampling

We studied *Erigone longipalpis* spiders living in salt marshes located on the south side of Mont-Saint-Michel Bay (western France), in an area split in two (hereafter "west" and "east" landscapes) by the mouth of the Couesnon river (**Fig. 1**). In this area, *E. longipalpis* is mostly found in sheep-grazed patches dominated by the common saltmarsh-grass *Puccinellia maritima* (Huds.) Parl., 1850 (fam. Poaceae), in which it is one of the most common spider species (Leroy et al. 2014). About forty years ago, *P. maritima* meadows occupied vast continuous swaths of the salt marshes on both sides of the Couesnon river (Valéry et al. 2017), and were one of the two dominant habitats, along with natural vegetation dominated by *Halimione portulacoides* (L.) Aellen, 1938 (fam. Amaranthaceae). Since then, both *P. maritima* meadows and natural habitats have been continuously reduced and fragmented by the rapid expansion of *Elytrigia acuta* (DC.) Tzvelev, 1973 (syn. *Elymus athericus* (Link) Kerguélen 1983, fam. Poaceae) throughout the salt marshes, possibly due to anthropic eutrophication (Valéry et al. 2017). This habitat loss and fragmentation has been much more extensive east of the Couesnon river, while the western side of the salt marshes still conserves relatively large and continuous *Puccinellia* patches (Valéry and Radureau, 2015; Valéry et al. 2017)(**Fig. 1**, **Supplementary Material 1**).

Using the most recent (2013) maps in Valéry et al. (2017), orthophotographs (Institut national de l'information géographique et forestière, 2017), and following ground-truthing surveys in late March - early April 2018, we selected 13 meadow patches with *Puccinellia* as potential sampling sites, 6 on the western side, 7 on the eastern side. All candidate patches were then visited at least once and up to 5 times, in April and May 2018, by groups of two or three researchers. All sampling sessions took place in sunny to lightly clouded weather between 10am and 5pm. Each patch visit lasted 30 minutes (some patches were visited twice in a day), during which we collected by hand all female brown-black linyphiids that had no distinctive opisthosoma pattern (more accurate species determination being difficult to impossible in field conditions). This procedure allowed us to get population density estimates standardised by research effort (in person-hours) and thus comparable among patches. We specifically sought females, but a few male by-catches were kept in low-density patches during the final visit, to potentially add genetic diversity in our crosses. We then identified spiders to species level under a binocular microscope (by placing them in between a petri dish and some plastic foam, so they were stuck and epigyne/pedipalps visible) following Roberts (1993). Female *Erigone longipalpis*

have a characteristic epigyne (Roberts, 1993) that makes them relatively easy to distinguish

from all other linyphiids found in these salt marshes (based on Leroy et al. 2014), including

other *Erigone* species. Although spiders were caught in all visited sites, *E. longipalpis* were only

found in the patches where *P. maritima* was dominant (i.e. with >50% cover; 5/6 of western and

3/7 of eastern patches sampled). Overall, we caught 38 *E. longipalpis* female spiders; 34 of

those 38 laid at least one eggsac while 26 of those 34 produced spiderlings. We did not manage

to produce spiderlings from one of the eight successfully sampled patches (as the cocoons

produced by the sole sampled female did not hatch); therefore, all experiments described

below were done using lab-born spiders originating from 7 patches (4 western and 3 eastern

patches).

Maintenance of spiders in the lab

We kept all spiders in temperature-controlled cabinets (25 ± 1°C) under controlled light regime

(16:8 L:D). They were all housed in plastic boxes and cups with a moist 1cm layer of plaster of

Paris at the bottom (e.g. Mestre and Bonte 2012). Container size, whether spiders were housed

by clutch or individually, and a few other elements, depended on life stage; see details below.

Independently of life stage, we provided all spiders with *ad libitum* access to *Sinella curviseta*

Brook, 1882 springtails as their main food source (*Erigone* spiders in general and *E. longipalpis* in

particular readily predate springtails, e.g. Irmler and Heydemann 1985; Mestre and Bonte 2012).

Adult spiders were kept individually in cylindrical plastic cups (6 cm diameter, 4 cm height). In

addition to springtails, we gave them a *Drosophila* fly once a week (generally *Drosophila*

melanogaster Meigen, 1830 but in some cases *Drosophila suzukii* (Matsumura, 1931) when *D.*

melanogaster supply was low). We renewed springtails and re-humidified the plaster at the

same time. When females laid an egg sac (see below for mating details for lab-born spiders), the

eggs were left in place and the female, along with as many remaining springtails as possible,

gently transferred to a new cup using a small paintbrush. The cup containing the eggs was then

placed, open, into a larger plastic box (8 × 9 cm, height 5 cm). When eggs started to hatch, all

spiderlings were counted and transferred from the cup to the larger box using paintbrushes, if

they had not already moved by themselves. We "seeded" each box with a large number of

springtails (> 100 adults), and drilled a small hole (≈ 5 mm diameter and depth) in the plaster

that we filled with rehydrated baker's yeast *Saccharomyces cerevisiae* Meyen ex Hansen, 1883.

This provided a food source for springtails, allowing them to reproduce and grow, guaranteeing

sustained resources for spiderlings. Yeast was rehydrated weekly, at the same time as the

plaster, and renewed at the same time if needed. We moved spiders to individual boxes after

the penultimate moult, to prevent uncontrolled mating events between siblings.

The experiment ran for two generations of lab-reared spiders, from April to late September

2018. Due to limited room in temperature-controlled cabinets, only ≈ 200 individually housed

adults could be kept at any one time, in addition to all spiderlings' boxes. Every time a "slot"

opened, we chose a replacement among the next adults to emerge at random (accordingly, the

number of lab-born adult spiders per origin patch in the final dataset is correlated with the

number of field-caught spiders from that patch; **Supplementary Material 2**). A total of 530

adults were maintained, including 293 females. We only tested females in subsequent

experiments, both for logistic reasons and because individual-level correlations between

fecundity and other traits can only be expressed in females.

Dispersal traits

We tested female spiders' dispersal propensity following a protocol inspired by previous studies

(e.g. Bonte et al. 2008; De Meester and Bonte 2010). We used circular plaster platforms

(diameter 5.5 cm) exposed to a slightly upward (≈ 20°) directional air flow (wind speed: 1 m.s^{-1} ±

0.2, expected to be optimal for dispersal initiation, Simonneau et al. 2016) as our test arena.

Four 17.5 cm long wood skewers were planted vertically, allowing spiders to climb to initiate

dispersal behaviours. We placed test platforms in water-filled trays to prevent spider escapes.

Spiders were tested individually for 15 minutes at ≈ 25°C. We thoroughly cleaned arenas with

paper towels and water between tests to remove any remaining silk thread that may influence

dispersal decisions (De Meester and Bonte 2010). Individuals were starved (i.e. all remaining

springtails and, more rarely, flies were removed) 12 to 24h before tests to stimulate dispersal

(Weyman et al. 1994).

Preliminary tests to validate the protocol were done using field caught spiders, 5 days post-
capture. For the experiment proper, we tested lab-born spiders on average 8.9 days post-sexual
maturity, and before mating (SD: 2.1 days, range: 7-21). Spiders present a stereotyped "tiptoe"
behaviour associated with silk production prior to dispersal, whether it is long-distance dispersal
by "ballooning" or short-distance dispersal by "rappelling" (Bonte 2012). No actual ballooning
was observed during the experiment. This is not surprising however, as while pre-dispersal
behaviours are a prerequisite for actual dispersal, they are not always (and in some cases not
often) followed by actual dispersal during short experimental tests in standardised conditions
(Lee et al. 2015). In some cases, spiders were seen tiptoeing and then immediately walking on
the produced thread floating in the wind after attaching it to the stick; these were likely
attempts at short-distance dispersal (rappelling), but may also have been attempts at two-step
ballooning by "rafting" (Bell et al. 2005). We therefore used the overall number of tiptoe
attempts per trial (whether or not they were followed by a rappelling attempt) as our overall
measure of dispersal motivation. Using only the number of rappelling attempts instead leads to
similar results (**Data availability**; r between the two variables = 0.89).

We then replaced each spider in its box, and provided it with a fly and springtails.

Mating

Two hours after the dispersal test, each female was presented with a non-sib male originating
from the same source population for mating. Males were left for 24h in the female's box, before
being removed. Some males were reused several times (number of males used: 190, mean
number of mates per used male: 1.45, range: 1-4).

A few spiders laid cocoons before being offered a potential mate. As a precaution, we did not
present a male to these spiders, to avoid potential mixed paternities. However, no spiderling
emerged from any of these "suspicious" cocoons. Unmated spiders can lay unfertilised and
therefore non-viable eggs when no mate is present (Zschokke and Herberstein 2005); we
suspect all these "early" cocoons resulted from this, rather than undetected mating between
siblings. As a result, meaningful fecundity data were available for 273 out of our 293 females.

Other traits

We noted the dates at which each eggsac was laid and hatched, as well as the date of sexual maturity and death for each female spider kept adult. We used these data to calculate development times (from hatching to sexual maturity) and adult longevities (from sexual maturity to death). We note that due to observation biases (detection of hatching) and gaps due to non-work days, development times and longevities may be recorded with some error; see **Statistical analyses** and provided code (**Data availability**) for how we accounted for this. For all females that actually had access to a mate, we additionally recorded the total number of hatchlings produced as our measure of fecundity. The experiment was stopped on September 30th 2018; 12 female spiders were still alive at that time, and we used a censoring indicator to include them properly in longevity analyses (see **Statistical analyses**).

When test females died, they were kept in 90% ethanol before being photographed under binocular microscope. We measured the cephalothorax width and length of each spider to the nearest 0.1 mm and used them as proxies of body size (e.g. Eichenberger et al. 2009). We measured each spider twice to account for measurement error. Other morphological traits that could have been useful to gauge individual condition (e.g. leg length, opisthosoma size) could not be obtained consistently, due to spider bodies sometimes losing integrity fast after death (maybe eaten by springtails), and were thus not used.

Statistical analyses

We analysed our data using a Bayesian workflow, with R (version 4.1.2; R Core Team 2021) and the *brms* R package (Bürkner 2017) as frontends for the Stan language (Carpenter et al. 2017). In addition, data preparation, posterior model evaluation and plotting were facilitated by the *tidybayes, bayesplot, patchwork,* as well as the *tidyverse* suite of packages (Gabry et al. 2019; Kay, 2019; Pedersen, 2019; Wickham et al. 2019). See **Supplementary Material 3** for a detailed description of the models and submodels; we here present a short summary.

We first used a Poisson GLM to determine whether the number of female *E. longipalpis* collected per patch differed between the two landscapes. This model included an offset/rate

term accounting for the fact sampling effort (in person-hours) varied between patches. We ran
this model twice: once using only visited patches where *Puccinellia maritima* was actually the
dominant plant species (>50% of cover), and once using all visited patches (the second one is
presented as **Supplementary Material 4**).

We then used a multivariate/multiresponse mixed model approach to analyse how our traits of
interest differed between the two landscapes, and whether traits were correlated at the
(within-patch) among-individual level.

• body size was analysed using a Gaussian model on (centred and scaled to 1SD)
cephalothorax width data (similar results were found using cephalothorax length
instead; see code in **Data availability**). The submodel included a fixed effect of
landscape, a random effect of patch of origin and an individual-level random effect. The
latter random effect is used to link size and all the other traits in an individual-level
covariance matrix; it is estimable in our Gaussian model separately from residual error
because size was measured twice per individual. 280 females out of 293 were measured;
the others were too badly damaged for their size to be estimated accurately. We used
in-model missing data imputation to account for these (results are similar if we exclude
them instead);

• in adult spiders that survived up to dispersal tests (N= 291 out of 293), dispersal was
analysed using the number of tiptoe events per individual and a Poisson model with a
log link. This submodel also included a fixed effect of landscape, a random effect of
patch of origin and an individual-level random effect. The individual-level random effects
here act as observation-level random effects (Harrison 2014);

• in individuals that were able to mate and survive at least one day post-mating (N = 272
out of 293), we analysed the number of spiderlings produced per observed day as our
measure of fecundity. The Poisson model was the same as the dispersal one above, with
the addition of an offset/rate term denoting the number of observation days (from
mating to death/end of experiment);

• development time (number of days from hatching to sexual maturity) and adult
longevity (days from sexual maturity to death) were also analysed using Poisson models.
As above, these submodels included a fixed effect of landscape, random effects of patch
and individual identity. However, they also included an "observation type" binary
variable denoting whether or not the observations bounding the interval of interest
(hatching, maturity, death) were made the day after a gap in recording, which could bias
estimates. Finally, the longevity model accounted for the fact some individuals were not
observed until natural death by using a right-censoring indicator (equal to 0 if natural
death was recorded during the experiment, and to 1 if the individual outlived the
experiment or died accidentally).

(Note that the methods used in the brms package allowed us to estimate all among-individual
covariances even if the number of individuals effectively tested varies between traits as
described above.)

Finally, we re-ran the same multivariate model as above, this time by adding (correlated)
random effects of mother identity in addition to the individual identity random effects. This
allowed us to partition individual-level covariation into its among-family and within-family
components, the former providing an upper bound on the variation attributable to heritable
differences. We initially intended to go further and use pedigree data and animal models
(Wilson et al. 2010) to separate genetic (co)variances from maternal effects. However, in all
attempts, models failed to converge in a satisfactory way. We attribute that to our small sample
size and to the sparseness of our pedigree (we have no relatedness information on spiders
descending from different wild-caught mothers, let alone coming from different patches)
(Wilson et al. 2010). We therefore settled on an approach that merely separates within- from
among-families (co)variation, using mother ID random effects; we acknowledge that such an
approach cannot separate genetic variation from maternal and common environment effects,
and can thus only provide upper-bound estimates of heritable variation.

For most parameters, we used weakly informative priors inspired by McElreath (2020):
$Normal(0,1)$ priors for fixed effects, $half - Normal(0,1)$ priors for standard deviations (random

effect and residual), and a LKJ(3) for the correlation matrices. The only exceptions were the

intercepts for development time and adult longevity, for which we shifted the fixed effects prior

means to reflect existing information on development and survival times of *Erigone* spiders (e.g.

Bonte et al. 2008; Mestre and Bonte 2012). Again, see **Supplementary Material 3** for a more

formal description of the models and priors. We ran four chains per model for 6000 iterations

each, and used the first 3000 iterations of each chain as warmup. Convergence and mixing were

satisfactory based on graphical checks and values of the updated $\hat{R}$ statistic (Vehtari et al. 2021).

In addition, all parameters had tail and bulk-effective sample sizes > 400 (Vehtari et al. 2021),

and most were >1000. All parameter posterior summaries below are given as means [95%

Highest Posterior Density Intervals].

Results

Erigone longipalpis spiders had lower population densities on the eastern, more fragmented,

side of the salt marshes (**Fig. 2**, $\beta_{[east-west]}$= -1.04 [-1.68; -0.37]). This result still holds if we

include sampled patches where *Puccinellia* was not dominant and no spiders were found

(**Supplementary Material 4**).

Figure 2. Effect of landscape of origin on the number of spiders found per patch (weighted by sampling effort). White dots corresponding to observed data are shown alongside model posteriors. Only data from habitat favourable to *Erigone longipalpis*, i.e. *Puccinellia maritima*-dominated meadows, are presented here; for the data and model using all visited sites, see **Supplementary Material 4**.

We found no evidence for dispersal differences between spiders from the western and eastern landscapes (**Fig. 3a**, $\beta_{[east-west]}$ = 0.18 [-0.38 ; 0.66]). We also found no meaningful differences in development time, adult longevity or body size (**Fig. 3**, $\beta_{[east-west]}$ = 0.06 [-0.15; 0.30], 0.10 [-0.04; 0.22], 0.26 [-0.33, 0.80], respectively). We found no evidence that observations of

longevity or time to maturity including observation gaps are biased in one direction or the other

(β = 0.07 [-0.04; 0.17] and -0.01 [-0.09; 0.07], respectively). These results were similar whether

we used the model where among-individual variance is unpartitioned (above, **Fig. 3**), or when

we re-run the model, partitioning among-individual variance into its among- and within-family

components (**Supplementary Material 5**). By contrast, there was some indication that spiders

were less fecund in the western more continuous landscape (**Fig. 3**, $\beta_{[east-west]}$= 0.69 [0.02;

1.40], but contrary to all other (non-)effects mentioned in this paragraph, support for this one

was model-structure dependent: once we re-ran the model partitioning among-individual

variance into its among- and within-family components, the 95% compatibility interval

overlapped with 0 ($\beta_{[east-west]}$= 0.67 [-0.26; 1.60]; **Supplementary Material 5**).

Figure 3. Phenotypic traits of lab-born spiders as a function of landscape of origin. Observed data are

displayed as boxplots (one boxplot per patch of origin), alongside model posteriors. Predictions are

based on the model where individual trait co-variance is not partitioned between among- and within-

family components; see **Supplementary Material 5** for a similar figure based on the model where

individual-level variation is partitioned.

Regarding the structure of the dispersal syndrome, the analysis of the first multivariate model
shows evidence of multiple individual-level correlations among traits, many of them directly
involving dispersal (**Table 1a**). In particular, more dispersive individuals tended to develop
faster, to be larger, and to have higher fecundity per day. There was however no clear
correlation between dispersal and adult longevity. Maturity and fecundity were also correlated,
with faster-developing spiders being more fecund. Larger spiders matured earlier but also died
earlier, and were also more fecund.

When partitioning the individual variance-covariance matrix into its among- and within-family
component, we found that among the correlations described above, the ones that involved
dispersal, fecundity and time to maturity were still found at the within-family level (**Table 1b**)
but not the among-family level (**Table 1c**). On the other end, the size-longevity and size-
fecundity correlations were detectable at the among-family, but not within-family level. A
negative size-maturity correlation (i.e. spiders maturing later were smaller) was found across
hierarchical levels (**Table 1b,c**). A positive longevity-time to maturity correlation was only
present at the among-family level (mothers producing slower-developping offspring also tended
to produce long-lived individuals when adult; **Table 1c**).

Table 1. Means and 95% Higher Posterior Density intervals for (a) the overall individual-level correlations
among traits, (b) the within-family correlations, and (c) the among-family correlations. (a) is based on the
model where individual trait co-variance is not partitioned between among- and within-family
components; see **Supplementary Material 6** for a similar table based on the model where individual-
level variation is partitioned (made by adding among- and within-family covariances).

(a) overall within-patch, among-individual correlations

	Time to maturity	Body size	Dispersal	Fecundity
Body size	-0.42 [-0.56; -0.27]			
Dispersal	-0.42 [-0.58; -0.25]	0.18 [0.04; 0.31]		
Fecundity	-0.33 [-0.50; -0.17]	0.25 [0.12; 0.38]	0.23 [0.09; 0.36]	
Adult longevity	0.04 [-0.12; 0.21]	-0.27 [-0.40; -0.14]	0.01 [-0.14; 0.15]	-0.03 [-0.18; 0.12]

(b): within-family component only (individual ID random effect)

	Time to maturity	Body size	Dispersal	Fecundity
Body size	-0.38 [-0.57; -0.18]			
Dispersal	-0.40 [-0.61; -0.19]	0.11 [-0.04; 0.26]		
Fecundity	-0.34 [-0.55; -0.11]	0.08 [-0.07; 0.22]	0.18 [0.01; 0.34]	
Adult longevity	-0.18 [-0.39; 0.05]	-0.03 [-0.18; 0.12]	0.09 [-0.06; 0.25]	0.02 [-0.15; 0.20]

(c): among-family component only (mother ID random effect)

	Time to maturity	Body size	Dispersal	Fecundity
Body size	-0.33 [-0.62; -0.01]			
Dispersal	-0.39 [-0.84; 0.08]	0.29 [-0.16; 0.75]		
Fecundity	-0.24 [-0.53; 0.10]	0.45 [0.17;0.71]	0.24 [-0.23; 0.68]	
Adult longevity	0.41 [0.03; 0.75]	-0.67 [-0.90; -0.41]	-0.31 [-0.79; 0.19]	-0.17 [-0.53; 0.19]

Discussion

The loss and fragmentation of sheep-grazed *Puccinellia maritima* meadows, the main habitat supporting *Erigone longipalpis* spiders (Leroy et al. 2014) has been a long-term trend in the Mont-Saint Michel Bay, going back at least 30 years (Valéry et al. 2017). As this loss is not uniform throughout the region, we have been able to compare spiders from landscapes differing in habitat availability (**Supplementary Material 1**). We show that population densities are markedly lower in the most fragmented (*sensu lato*) landscape (**Fig. 2**). While the effects of fragmentation *sensu stricto* on biodiversity are still debated (Fletcher et al. 2018; Fahrig et al. 2019), the negative effects of habitat loss itself are more clearly supported (whether they are attributed to reduction in patch size or reduction in total habitat amount; Bender et al. 1998; Fahrig 2003; Watling et al. 2020). Our results here add one more piece of evidence, even small, in that direction. The fact that habitat loss exerts such a strong negative pressure on *Erigone longipalpis* would suggest that it may also shapes which phenotypes are advantageous, and therefore lead to phenotypic divergence between fragmented and (relatively) more intact landscapes. In addition, changes in population density itself are expected to shape dispersal and related traits (De Meester and Bonte 2010; Endriss et al. 2019). These changes can in some cases carry over across generations even when they are not underpinned by genetic variation, through parental effects (Bitume et al. 2014).

For the above reasons, it is very surprising to see that differences in habitat availability have actually little to no effect on phenotypic traits, including dispersal (**Fig. 3**). This is especially true given the reported cases of phenotypic changes in response to habitat fragmentation and loss (Cheptou et al. 2017; Cote et al. 2017). Here, at most, *E. longipalpis* spiders from the more habitat-limited landscape may be slightly more fecund, but even that finding is model-dependent and uncertain. It is possible that our study was simply not done at the right scale, whether in space or in time. On the spatial side, our two landscapes are separated from each other by about 5 km (**Fig. 1**), and other suitable salt marshes can be found at similar distances on the other side of the bay (Valéry et al. 2017). Spiders can in principle balloon over much larger distances than that (Bell et al. 2005). As prevailing winds in the region blow from the

(south-)west (DREAL Normandie, 2020), this could mean populations in the eastern fragmented
habitat are regularly supplied with immigrants from the western continuous habitat, which
would make adaptation to local conditions much more complicated (Garant et al. 2007). We
note however that the actual typical dispersal distances in ballooning spiders are likely much
lower than rare long-distance dispersal events would suggest, and closer to a few hundred
meters to a few kilometres at most, based on population genetic studies (Croucher et al. 2011;
Reed et al. 2011), or on the spatial scale of effect of landscape on abundances (Schmidt et al.
2008; Jackson and Fahrig, 2012). Also, the potential for long-distance dispersal and gene flow is
not always, in itself, a barrier for the evolution of dispersal (Cheptou et al. 2008). Alternately, if
long-distance dispersal is rare in *E. longipalpis* (Meijer, 1977), individual differences in tiptoe
frequency may be more reflective of short-distance dispersal by rappelling than of ballooning.
Then it may be that spiders do not experience any fragmentation and habitat loss at the scale at
which they disperse, even in the eastern habitat. Indeed, typical rappelling distances are up to a
few meters (Bonte et al. 2008), and even the small patches in the eastern fragmented landscape
are an order of magnitude bigger (**Supplementary Material 1**). It might be possible to resolve
this uncertainty through further experiments that cover an even broader range of habitat loss
and fragmentation, and are designed to more clearly separate variation in long-distance and
short-distance dispersal. Regarding the temporal aspect, available information suggests that the
dynamics of the two landscapes started to diverge ≈ 30 years ago (Valéry et al. 2017), which
would in principle be enough to drive the rapid evolution of dispersal (Cheptou et al. 2008).
However, landscape transformation happened progressively and not suddenly. The key question
here is then: when did the landscapes became divergent enough for it to meaningfully influence
Erigone longipalpis biology? To the best of our knowledge, the long-term data that are needed
to answer this question are unfortunately not available even for better-studied linyphiids, let
alone *E. longipalpis*. Further studies at larger scales, both in space and time, or including
estimates of actual gene flow among populations/landscapes (Reed et al. 2011), are again
probably needed before concluding definitely on the effect of habitat loss on *Erigone longipalpis*
evolution.

We found substantial evidence that key life-history traits are integrated with dispersal in a

syndrome in this species (**Table 1**). *Erigone longipalpis* seems to present a within-species pace of

life syndrome, with faster-developing spiders also being more fecund on a per day basis (**Table**

1a, Ricklefs and Wikelski 2002; Réale et al. 2010; Healy et al. 2019). Dispersal was aligned with

this syndrome in a way that follows some existing predictions and results (e.g. Réale et al. 2010;

Stevens et al. 2013), with faster-growing individuals being more likely to disperse. When the

among-individual covariance matrix was split into its within- and among-family components (the

latter reflecting genetic variation, among other things), however, we found that dispersal-

related correlations, and the development time-fecundity correlation expected from the pace-

of-life hypothesis were only found at the within-family level (**Table 1b,c**). In other terms, there

was no evidence that families that differed in dispersal propensity consistently differed in other

traits, which may explain why there was no "coordinated" responses to habitat loss between

traits (**Fig. 3**). This result is interesting given a recent meta-analysis suggested there is actually

little to no empirical support for the hypothesis that risky behaviour and fast pace of life should

be positively correlated at the genetic level, even when they are correlated at the individual

level (Royauté et al. 2018). In addition, a key prediction of the pace-of-life syndrome hypothesis,

that fast development is associated with reduced longevity (Réale et al. 2010; Healy et al. 2019)

is not supported at the individual level, but is validated at the family level (**Table 1**). While this

specific association may well be genetically driven, one must remember that we cannot

separate additive genetic from maternal effects in this setup. As mothers that produced

consistently fast-developing individuals also tended to produce larger than average individuals

(see the size-time to maturity correlation **Table 1c**), the idea that family differences are driven

by maternal effects more than genetic differences is worth considering. So, in *Erigone*

longipalpis, different parts of what we hypothesized to be a common genetic pace of life

syndrome actually exist at different levels of organisation, and the hypothesized dispersal-pace

of life syndrome only exists at the within-family level, meaning it is presumably not driven by

genetic variation. Altogether, these results highlight the need to reevaluate the assumptions

behind the pace of life syndrome hypothesis, and more generally behind general hypotheses on

the integration between (dispersal) behaviour and life history, in particular by better accounting

for hierarchical levels (among-species vs. individual vs. genetic correlations), dispersal context

dependency and the role of resource availability (Bonte and Dahirel, 2017; Wright et al. 2019;

Laskowski et al. 2021). This includes for instance testing for context-dependent syndromes; our

present dataset is however too small in our opinion for such tests.

Even though we did not find support for a dispersal-life-history syndrome at the family/genetic

level, this syndrome still exists at the individual level (**Table 1b**). The question that remains,

then, is how such a structured among-individual variation can arise if it is not apparently linked

to genetic covariation or to external environmental factors (as experimental conditions were

meant to be constant). A possible explanation for a non-genetic dispersal-fecundity syndrome

would be that direct costs incurred during dispersal, independently of the cause of dispersal,

limit fecundity afterwards (Bonte et al. 2012). Indeed, reduction of fecundity in direct relation to

previous dispersal/movement are well documented in arthropods (Roff, 1977; Matsumura and

Miyatake, 2018). There is one obvious drawback here: this prediction is the exact opposite of

the result we observed here. Alternatively, food availability during development can be a major

driving force of phenotypic variation in spiders, including the specific traits measured in this

study (Mestre and Bonte 2012; Kleinteich et al. 2015; Quiñones-Lebrón et al. 2021). Although

food was provided essentially *ad libitum*, we suggest that the observed dispersal syndrome can

be explained by within-family variation in early access to resources (stochastic and/or due to

unobserved trait variation, in e.g. functional response) causing variation in development time

and adult size, which in turn cause variation in dispersal, fecundity and longevity. Indeed, in

snails for instance, variation in early access to food (in the form of egg cannibalism) is well-

known and can drive long-lasting within-clutch differences in key traits, all else being equal

(Desbuquois 1997). Sibling cannibalism is also present in linyphiid spiders (Vanacker et al. 2004).

Further studies comparing dispersal syndromes in spiders reared in isolation versus in groups

could help confirm or infirm this hypothesis.

To summarise, despite strong ecological effects of habitat loss, and the existence of a dispersal

syndrome, we found no evidence of evolutionary responses to habitat loss in a specialist spider.

Although confirmations are needed, our results highlight the importance of

environmental/developmental conditions in driving dispersal syndromes at the within-species

level (Bonte and Dahirel, 2017). Even though we found no support for a genetic dispersal syndrome, observed associations between dispersal and traits can still substantially shape the dynamics of *Erigone longipalpis* in these salt marshes. Indeed, given the rarity of suitable habitats, it is possible that a specific subset of phenotypes is bound to be disproportionally lost during dispersal in the more fragmented landscape, which may have ecological consequences both on spider population dynamics and on their interactions with prey. It is even possible that, in the absence of dispersal evolution to compensate for it, the systematic loss of more fecund individuals to dispersal is itself one of the causal mechanisms for the observed ecological effects of habitat loss on this habitat specialist spider.

Acknowledgements

We warmly thank Martin Entling for the starter *Sinella curviseta* springtail population, as well as Youn Henry and Thomas Enriquez for giving us access to a regular supply of *Drosophila* flies. Amélie Beillard kindly helped take care of the springtail populations at critical times. We are also grateful to Armelle Ansart for access to temperature-controlled cabinets, and to Françoise Burel who supported some exploratory analyses linked to this study. We also thank Shakira Quiñones for a friendly review of the manuscript before submission.

Author contributions

Initial idea: MD and JP; funding acquisition: MD, SC, JP; site selection and sampling: MD, MW, SC, JP; behavioural experiment design and rearing protocol: MD, MW, TB; behavioural and life history experiments and population maintenance: MW, MD; data analysis: MD, MW; lead manuscript writer: MD.

Data availability

Data and R scripts to reproduce all analyses presented in this manuscript are available on Github (https://github.com/mdahirel/erigone-dispersal-2018) and Zenodo (https://doi.org/10.5281/zenodo.5761384)

Funding

This project was funded by a research grant of the Observatoire des Sciences de l'Univers de Rennes (UMS OSUR) to MD, SC and JP, and by an internal grant of the UMR Ecobio (Biological Invasions research axis) to MD and JP.

References

Bell JR, Bohan DA, Shaw EM, Weyman GS. 2005. Ballooning dispersal using silk: world fauna, phylogenies, genetics and models. *Bull Entomol Res* 95:69-114.

Bender DJ, Contreras TA, Fahrig L. 1998. Habitat loss and population decline: a meta-analysis of the patch size effect. *Ecology* 79:517–533.

Benton TG, Bowler DE. 2012. Linking dispersal to spatial dynamics. In: Clobert J, Baguette M, Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford, UK: Oxford University Press. p. 251–265.

Bitume EV, Bonte D, Ronce O, Olivieri I, Nieberding CM. 2014. Dispersal distance is influenced by parental and grand-parental density. *Proc R Soc B* 281:20141061.

Bonte D. 2012. Case study II: Spiders as a model in dispersal ecology and evolution. In: Clobert J, Baguette M, Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford: Oxford University Press. p. xxvi–xxviii.

Bonte D, Dahirel M. 2017. Dispersal: a central and independent trait in life history. *Oikos* 126:472–479.

Bonte D, Lukáč M, Lens L. 2008. Starvation affects pre-dispersal behaviour of *Erigone* spiders. *Basic Appl Ecol* 9:308–315.

Bonte D, Saastamoinen M. 2012. Dispersal syndromes in butterflies and spiders. In: Clobert J, Baguette M, Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford: Oxford University Press. p. 161–170.

Bonte D, Travis JMJ, Clercq ND, Zwertvaegher I, Lens L. 2008. Thermal conditions during juvenile
development affect adult dispersal in a spider. *PNAS* 105:17000–17005.

Bonte D, Van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, Lehouck V, Matthysen E,
Mustin K, Saastamoinen M, et al. 2012. Costs of dispersal. *Biol Rev* 87:290–312.

Bürkner P-C. 2017. brms: an R package for Bayesian multilevel models using Stan. *J Stat Softw*
80:1–28.

Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P,
Riddell A. 2017. Stan: a probabilistic programming language. *J Stat Softw* 76:1–32.

Cheptou P-O, Carrue O, Rouifed S, Cantarel A. 2008. Rapid evolution of seed dispersal in an
urban environment in the weed *Crepis sancta*. *PNAS* 105:3796–3799.

Cheptou P-O, Hargreaves AL, Bonte D, Jacquemyn H. 2017. Adaptation to fragmentation:
evolutionary dynamics driven by human influences. *Phil Trans R Soc B* 372:20160037.

Clobert J, Baguette M, Benton TG, Bullock JM, editors. 2012. Dispersal ecology and evolution.
Oxford, UK: Oxford University Press.

Cote J, Bestion E, Jacob S, Travis J, Legrand D, Baguette M. 2017. Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. *Ecography* 40:56–73.

Croucher PJP, Oxford GS, Gillespie RG. 2011. Population structure and dispersal in a patchy
landscape: nuclear and mitochondrial markers reveal area effects in the spider *Theridion*
californicum (Araneae: Theridiidae). *Biol J Linn Soc* 104:600–620.

De Meester N, Bonte D. 2010. Information use and density-dependent emigration in an
agrobiont spider. *Behav Ecol* 21:992–998.

Desbuquois C. 1997. Influence of egg cannibalism on growth, survival and feeding in hatchlings
of the land snail *Helix aspersa* Müller (Gastropoda, Pulmonata, Stylommatophora). *Reprod Nutr*
Dev 37:191–202.

DREAL Normandie. 2020. Profil Environnemental Normandie - Climat. Available from:
http://www.normandie.developpement-durable.gouv.fr/le-profil-environnemental-normandie-
r307.html.

Eichenberger B, Siegenthaler E, Schmidt-Entling MH. 2009. Body size determines the outcome of
competition for webs among alien and native sheetweb spiders (Araneae: Linyphiidae). *Ecol*
Entomol 34:363–368.

Endriss SB, Vahsen ML, Bitume EV, Monroe JG, Turner KG, Norton AP, Hufbauer RA. 2019. The
importance of growing up: juvenile environment influences dispersal of individuals and their
neighbours. *Ecol Lett* 22:45–55.

European Environmental Agency. 2020. Corine Land Cover, 2018, Version 2020_20u1. European
Union, Copernicus Land Monitoring Service. Available from: https://land.copernicus.eu/pan-
european/corine-land-cover.

Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst* 34:487–
515.

Fahrig L, Arroyo-Rodríguez V, Bennett JR, Boucher-Lalonde V, Cazetta E, Currie DJ, Eigenbrod F,
Ford AT, Harrison SP, Jaeger JAG, et al. 2019. Is habitat fragmentation bad for biodiversity? *Biol*
Cons 230:179–186.

Fletcher RJ, Didham RK, Banks-Leite C, Barlow J, Ewers RM, Rosindell J, Holt RD, Gonzalez A,
Pardini R, Damschen EI, et al. 2018. Is habitat fragmentation good for biodiversity? *Biol Cons*
226:9–15.

Fronhofer EA, Legrand D, Altermatt F, Ansart A, Blanchet S, Bonte D, Chaine A, Dahirel M,
Laender FD, Raedt JD, et al. 2018. Bottom-up and top-down control of dispersal across major
organismal groups. *Nature Ecol Evol* 2:1859–1863.

Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. 2019. Visualization in Bayesian
workflow. *J Royal Stat Soc A* 182:389–402.

Garant D, Forde SE, Hendry AP. 2007. The multifarious effects of dispersal and gene flow on
contemporary adaptation. *Funct Ecol* 21:434–443.

Govaert L, Fronhofer EA, Lion S, Eizaguirre C, Bonte D, Egas M, Hendry AP, Martins ADB, Melián
CJ, Raeymaekers JAM, et al. 2019. Eco-evolutionary feedbacks—Theoretical models and
perspectives. *Funct Ecol* 33:13–30.

Guerra PA. 2011. Evaluating the life-history trade-off between dispersal capability and
reproduction in wing dimorphic insects: a meta-analysis. *Biol Rev* 86:813–835.

Harrison XA. 2014. Using observation-level random effects to model overdispersion in count
data in ecology and evolution. *PeerJ* 2:e616.

Harvey PR, Nellist DR, Telfer MG, editors. 2002. Provisional atlas of British spiders (Arachnida,
Araneae), volumes 1 and 2. Huntingdon, UK: Biological Records Centre.

Healy K, Ezard THG, Jones OR, Salguero-Gómez R, Buckley YM. 2019. Animal life history is
shaped by the pace of life and the distribution of age-specific mortality and reproduction.
Nature Ecol Evol 3:1217–1224.

Institut national de l'information géographique et forestière. 2017. BD ORTHO®. Available from:
http://www.geoportail.gouv.fr/accueil.

Irmler U, Heydemann B. 1985. Populationsdynamik und Produktion von *Erigone longipalpis*
(Araneae, Micryphantidae) auf einer Salzwiese Nordwestdeutschlands. *Faun Ökol Mitt* 5:443–
454.

Jackson HB, Fahrig L. 2012. What size is a biologically relevant landscape? *Landscape Ecol*
27:929–941.

Jacob S, Chaine AS, Huet M, Clobert J, Legrand D. 2019. Variability in dispersal syndromes is a
key driver of metapopulation dynamics in experimental microcosms. *Am Nat* 194:613–626.

Kay M. 2019. tidybayes: tidy data and geoms for Bayesian models. Available from:
http://mjskay.github.io/tidybayes/.

Kleinteich A, Wilder SM, Schneider JM. 2015. Contributions of juvenile and adult diet to the
lifetime reproductive success and lifespan of a spider. *Oikos* 124:130–138.

Laskowski KL, Moiron M, Niemelä P. 2021. Integrating behavior in life-history theory: allocation
versus acquisition? *Trends Ecol Evol* 36:132-138.

Lee VMJ, Kuntner M, Li D. 2015. Ballooning behavior in the golden orbweb spider *Nephila pilipes*
(Araneae: Nephilidae). *Front Ecol Evol* 3:2.

Leroy B, Le Viol I, Pétillon J. 2014. Complementarity of rarity, specialisation and functional
diversity metrics to assess community responses to environmental changes, using an example of
spider communities in salt marshes. *Ecol Indic* 46:351–357.

Little CJ, Fronhofer EA, Altermatt F. 2019. Dispersal syndromes can impact ecosystem
functioning in spatially structured freshwater populations. *Biol Lett* 15:20180865.

Massol F, Altermatt F, Gounand I, Gravel D, Leibold MA, Mouquet N. 2017. How life-history
traits affect ecosystem properties: effects of dispersal in meta-ecosystems. *Oikos* 126:532–546.

Matsumura K, Miyatake T. 2018. Costs of walking: differences in egg size and starvation
resistance of females between strains of the red flour beetle (*Tribolium castaneum*) artificially
selected for walking ability. *J Evol Biol* 31:1632–1637.

McElreath R. 2020. Statistical rethinking: a Bayesian course with examples in R and Stan. 2nd
edition. Boca Raton, USA: Chapman and Hall/CRC.

Meijer J. 1977. The immigration of spiders (Araneida) into a new polder. *Ecol Entomol* 2:81–90.

Mestre L, Bonte D. 2012. Food stress during juvenile and maternal development shapes natal
and breeding dispersal in a spider. *Behav Ecol* 23:759–764.

Ochocki BM, Saltz JB, Miller TEX. 2020. Demography-dispersal trait correlations modify the eco-
evolutionary dynamics of range expansion. *Am Nat* 195:231–246.

Pedersen TL. 2019. patchwork: the composer of plots. Available from: https://CRAN.R-
project.org/package=patchwork.

Pétillon J, Georges A, Canard A, Lefeuvre J-C, Bakker JP, Ysnel F. 2008. Influence of abiotic
factors on spider and ground beetle communities in different salt-marsh systems. *Basic Appl*
Ecol 9:743–751.

Pétillon J, Georges A, Canard A, Ysnel F. 2007. Impact of cutting and sheep grazing on ground–
active spiders and carabids in intertidal salt marshes (Western France). *Anim Biodivers Conserv*
30:201–209.

Quiñones-Lebrón SG, Kuntner M, Kralj-Fišer S. 2021. The effect of genetics, diet, and social
environment on adult male size in a sexually dimorphic spider. *Evol Ecol* 35:217–234.

R Core Team. 2021. R: a language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing. Available from: https://www.R-project.org/.

Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio P-O. 2010. Personality and
the emergence of the pace-of-life syndrome concept at the population level. *Phil Trans R Soc B*
365:4051–4063.

Reed DH, Teoh V-H, Stratton GE, Hataway RA. 2011. Levels of gene flow among populations of a
wolf spider in a recently fragmented habitat: current versus historical rates. *Conserv Genet*
12:331–335.

Ricklefs RE, Wikelski M. 2002. The physiology/life-history nexus. *Trends Ecol Evol* 17:462–468.

Roberts MJ. 1993. The spiders of Great Britain and Ireland. Colchester, UK: Harley Books.

Roff D. 1977. Dispersal in Dipterans: its costs and consequences. *J Anim Ecol* 46:443–456.

Ronce O, Clobert J. 2012. Dispersal syndromes. In: Clobert J, Baguette M, Benton TG, Bullock JM,
editors. Dispersal ecology and evolution. Oxford, UK: Oxford University Press. p. 119–138.

Royauté R, Berdal MA, Garrison CR, Dochtermann NA. 2018. Paceless life? A meta-analysis of
the pace-of-life syndrome hypothesis. *Behav Ecol Sociobiol* 72:64.

Royauté R, Hedrick A, Dochtermann NA. 2020. Behavioural syndromes shape evolutionary
trajectories via conserved genetic architecture. *Proc R Soc B* 287:20200183.

Schmidt MH, Thies C, Nentwig W, Tscharntke T. 2008. Contrasting responses of arable spiders to
the landscape matrix at different spatial scales. *J Biogeogr* 35:157–166.

Simonneau M, Courtial C, Pétillon J. 2016. Phenological and meteorological determinants of
spider ballooning in an agricultural landscape. *C R Biol* 339: 408-416.

Stevens VM, Trochet A, Blanchet S, Moulherat S, Clobert J, Baguette M. 2013. Dispersal
syndromes and the use of life-histories to predict dispersal. *Evol Appl* 6:630–642.

Stevens VM, Trochet A, Van Dyck H, Clobert J, Baguette M. 2012. How is dispersal integrated in
life histories: a quantitative analysis using butterflies. *Ecol Lett* 15:74–86.

Stevens VM, Whitmee S, Le Galliard J-F, Clobert J, Böhning-Gaese K, Bonte D, Brändle M,
Matthias Dehling D, Hof C, Trochet A, et al. 2014. A comparative analysis of dispersal syndromes
in terrestrial and semi-terrestrial animals. *Ecol Lett* 17:1039–1052.

Valéry L, Radureau A. 2015. Evolution de la végétation des marais salés de la Baie du Mont-
Saint-Michel : analyse cartographique 1984, 2013. Conservatoire du littoral.

Valéry L, Radureau A, Lefeuvre J-C. 2017. Spread of the native grass *Elymus athericus* in salt
marshes of Mont-Saint-Michel bay as an unusual case of coastal eutrophication. *J Coast Conserv*
21:421–433.

Vanacker D, Deroose K, Pardo S, Bonte D, Maelfait J-P. 2004. Cannibalism and prey sharing
among juveniles of the spider *Oedothorax gibbosus* (Blackwall, 1841) (Erigoninae, Linyphiidae,
Araneae). *Belg J Zool* 134:23–28.

Vehtari A, Gelman A, Simpson D, Carpenter B, Bürkner P-C. 2021. Rank-normalization, folding,
and localization: an improved $\hat{R}$ for assessing convergence of MCMC (with Discussion). *Bayesian*
Anal 16:667-718

Watling JI, Arroyo-Rodríguez V, Pfeifer M, Baeten L, Banks-Leite C, Cisneros LM, Fang R, Hamel-
Leigue AC, Lachat T, Leal IR, et al. 2020. Support for the habitat amount hypothesis from a
global synthesis of species density studies. *Ecol Lett* 23:674–681.

Weyman GS, Sunderland KD, Fenlon JS. 1994. The effect of food deprivation on aeronautic
dispersal behaviour (ballooning) in *Erigone* spp. spiders. *Entomol Exp Appl* 73:121–126.

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry
L, Hester J, et al. 2019. Welcome to the Tidyverse. *J Open Source Softw* 4:1686.

Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, Kruuk LEB, Nussey DH.
2010. An ecologist's guide to the animal model. *J Anim Ecol* 79:13–26.

Wright J, Bolstad GH, Araya-Ajoy YG, Dingemanse NJ. 2019. Life-history evolution under
fluctuating density-dependent selection and the adaptive alignment of pace-of-life syndromes.
Biol Rev 94:230–247.

Zschokke S, Herberstein ME. 2005. Laboratory methods for maintaining and studying web-
building spiders. *J Arachnol* 33:205–213.

habitat type   ■ *Puccinellia maritima* dominant   ■ *Puccinellia maritima* present   □ other unfavourable land

E. longipalpis found?   ○ no   ● yes