Dispersal syndrome and landscape fragmentation in the salt-marsh specialist spider *Erigone longipalpis*

Maxime Dahirel*[1,2], Marie Wullschleger[1], Tristan Berry[1,3], Solène Croci[4], Julien Pétillon[1,5]

[1]Univ Rennes, UR1, CNRS, ECOBIO (Ecosystèmes, Biodiversité, évolution), UMR 6553, F-35000, Rennes, France

[2]Department of Biology, Ghent University, B-9000, Ghent, Belgium

[3]Syndicat Mixte de Gestion des Milieux Naturels, Réserve Naturelle Nationale de l'Étang Noir, F-40510, Seignosse, France

[4]CNRS, Université de Rennes 2, EPHE-PSL, Université d'Angers, Université de Bretagne Occidentale, Université de Caen Normandie, Université de Nantes, UMR LETG, F-35043, Rennes, France

[5]Institute for Coastal and Marine Research, Nelson Mandela University, Port Elizabeth, South Africa

*corresponding author: maxime.dahirel@yahoo.fr

Abstract

Dispersal and its evolution play a key role for population persistence in fragmented landscapes where habitat loss and fragmentation increase the cost of between-habitat movements. In such contexts, it is important to know how variation in dispersal and other traits is structured, and whether responses to landscape fragmentation are aligned with underlying dispersal-trait correlations, or dispersal syndromes. We therefore studied trait variation in *Erigone longipalpis*, a European spider species specialist of (often patchy) salt marshes. We collected spiders in two salt-marsh landscapes differing in habitat availability. We then reared lab-born spiders for two generations in controlled conditions, and measured dispersal and its association with various key traits. *E. longipalpis* population densities were lower in the more fragmented landscape. Despite this, we found no evidence of differences in dispersal, or any other trait we studied, between the two landscapes. While a dispersal syndrome was present at the among-individual level (dispersers were more fecund and faster growing, among others), there was no indication it was genetically driven: among-family differences in dispersal were not correlated with differences in other traits. Instead, we showed that the observed phenotypic covariations were mostly due to within-family correlations. We hypothesize that the dispersal syndrome is the result of asymmetric food access among siblings, leading to variation in development rates and carrying over to adult traits. Our results show we need to better understand the sources of dispersal variation and syndromes, especially when dispersal may evolve rapidly in response to environmental change.

Keywords: context-dependent dispersal; habitat loss; Linyphiidae; phenotypic correlations; population density

Introduction

Dispersal is a key trait and process that influences, and links together, ecological and evolutionary dynamics (Clobert et al. 2012; Govaert et al. 2019). Individuals' movement between habitats shapes (meta-)population dynamics (Benton and Bowler 2012) while the resulting gene flow can have negative or positive impacts on local adaptation (Garant et al. 2007). Many "dispersal syndromes", i.e. associations/covariations between dispersal and other traits, have been documented across the tree of life (Ronce and Clobert 2012; Stevens et al. 2014; Jacob et al. 2019). These covariations may result both from genetic correlations, caused by pleiotropy or joint selective responses across traits, or from plastic responses to experienced environmental conditions (Ronce and Clobert 2012). These syndromes reinforce the role of dispersal as a nexus between ecological and evolutionary processes. Indeed, the existence of these syndromes mean that dispersal not only redistributes species, individuals and genetic diversity in space, but does it non-randomly with respect to trait values, which has potential consequences for instance for the distribution of ecosystem functions in landscapes (Massol et al. 2017; Little et al. 2019) or for the dynamics of range expansions (Ochocki et al. 2020).

Some general dispersal syndromes have been proposed on theoretical grounds. These include, for instance, expectations of a positive association between high dispersal and the "fast" end of a pace-of-life life history axis (Réale et al. 2010; Stevens et al. 2012; Wright et al. 2019), or the trade-off between fecundity and dispersal often seen in insects (Guerra 2011). However, in practice, observed syndromes are much more diverse and sometimes contradict these general predictions, especially at the within-species level (Guerra 2011; Ronce and Clobert 2012; Bonte and Dahirel 2017), making generalisations difficult. In the case of dispersal-pace of life syndromes, one possible reason for divergence among studies may be that within-species variation in life history does not always align nicely along a main pace of life axis, and that its existence should be tested rather than assumed (Royauté et al. 2018).

Habitat loss and habitat fragmentation (*sensu stricto*; the increasing isolation between habitat patches) are usually seen as a particularly strong selective pressure on dispersal, as they increase the (many) costs associated with moving from a relatively suitable habitat to another (Bonte et

al. 2012; Cote et al. 2017). Indeed, a reduction of dispersal is often predicted and observed in

response to fragmentation *sensu lato* (i.e. conflating habitat loss and fragmentation *sensu stricto*

together, Cote et al. 2017). Therefore, if dispersal is integrated with other traits in syndromes, we

may then expect these other traits to also evolve jointly in response to habitat change. If such

syndromes have a genetic basis and act as constraints on evolution (see e.g. Royauté et al. 2020),

then the direction of these evolutionary changes should be somewhat predictable from syndrome

structure.

Here we present a study of dispersal and dispersal syndromes in *Erigone longipalpis* (Sundevall,

1830) (fam. Linyphiidae) spiders. This European species is a strong habitat specialist, only found

in wet habitats like floodplains and especially salt marshes (Harvey et al. 2002; Pétillon et al.

2008). At the regional and European scales, these favourable landscapes (and therefore *E.*

longipalpis) are patchily distributed along the Atlantic coastal line (*sensu lato*; including the

Channel, North and Baltic seas; European Environmental Agency 2020; GBIF Secretariat 2022). In

addition, within salt marshes, *E. longipalpis* is mostly restricted to specific vegetation types that

are themselves patchy, in part due to human management (Pétillon et al. 2007; Leroy et al. 2014).

As a consequence, we may expect its dispersal responses to be strongly influenced by habitat

fragmentation, much more than related generalist species; indeed, any given landscape likely

contains less favourable habitat for specialists than generalists. In addition, *E. longipalpis* is one

of the most important invertebrate predators in these habitats (Leroy et al. 2014), so

understanding how dispersal influences its spatial population dynamics may help understand the

dynamics of its potential prey resources at the same time (Fronhofer et al. 2018). Finally, spiders

and especially linyphiids have been key models in dispersal ecology and evolution (Bonte 2012;

Bonte and Saastamoinen 2012), in part because they exhibit stereotypical behaviours associated

with dispersal (whether long-distance dispersal by ballooning or short-distance movements by

rappelling); dispersal-related behaviours can thus easily be tracked in an experimental setting

(e.g. De Meester and Bonte 2010). We used lab-born spiders coming from several patches in two

adjacent landscapes differing in fragmentation degree to test the following hypotheses:

• Rapid human-induced habitat loss and fragmentation have negative impacts on *Erigone*
longipalpis populations, and also lead to reduced dispersal propensity;

•   Dispersal is associated with other life-history traits in a syndrome. Specifically, high

dispersal propensity is associated with "faster" life-histories (e.g. faster development

time, higher fecundity), following predictions derived from the pace of life hypothesis

(Réale et al. 2010; Wright et al. 2019);

•   As syndromes can constrain trait evolution (e.g. Royauté et al. 2020), and dispersal evolves

with fragmentation, the direction in which other traits evolve can be predicted from the

direction of dispersal evolution and the structure of the among-family (potentially genetic)

dispersal syndrome.

Material and Methods

Site selection and field sampling

We studied *Erigone longipalpis* spiders living in salt marshes located on the south side of Mont-

Saint-Michel Bay (western France), in an area split in two (hereafter "west" and "east"

landscapes) by the mouth of the Couesnon river (**Fig. 1**). In this area, *E. longipalpis* is mostly found

in sheep-grazed patches dominated by the common saltmarsh-grass *Puccinellia maritima* (Huds.)

Parl., 1850 (fam. Poaceae), in which it is one of the most common spider species (Leroy et al.

2014). About forty years ago, *P. maritima* meadows occupied vast continuous swaths of the salt

marshes on both sides of the Couesnon river (Valéry and Radureau 2015; Valéry et al. 2017), and

were one of the two dominant habitats, along with natural vegetation dominated by *Halimione*

portulacoides (L.) Aellen, 1938 (fam. Amaranthaceae). Since then, both *P. maritima* meadows and

natural *H. portulacoides* habitats have been continuously reduced and fragmented by the rapid

expansion of *Elytrigia acuta* (DC.) Tzvelev, 1973 (syn. *Elymus athericus* (Link) Kerguélen 1983, fam.

Poaceae) throughout the salt marshes, possibly due to anthropic eutrophication (Valéry et al.

2017). This habitat loss and fragmentation has been much more extensive east of the Couesnon

river than on the western side (Valéry and Radureau 2015; Valéry et al. 2017). Nowadays,

Puccinellia meadows are much more abundant on the western side (less habitat loss), discrete

Puccinellia patches are also much larger, and meadows are also less fragmented *sensu stricto*,

compared to the eastern side (**Fig. 1**, see metrics **Supplementary Material 1**).

Using the most recent (2013) maps in Valéry et al. (2017), orthophotographs (Institut national de l'information géographique et forestière 2017), and following ground-truthing surveys in late March - early April 2018, we selected 13 meadow patches with *Puccinellia* as potential sampling sites, 6 on the western side, 7 on the eastern side. Based on available information, we believe the current extent of all selected patches was continuously occupied by *P. maritima* during the last forty years (Valéry and Radureau 2015; but note that sampling was infrequent and *P. maritima* may have been present but not dominant). All candidate patches were then visited at least once and up to 5 times, in April and May 2018, by groups of two or three researchers. All sampling sessions took place in sunny to lightly clouded weather between 10am and 5pm. Each patch visit lasted 30 minutes (some patches were visited twice in a day), during which we collected by hand all female brown-black linyphiids that had no distinctive opisthosoma pattern (more accurate species determination being difficult to impossible in field conditions). This procedure allowed us to get population density estimates standardised by research effort (in person-hours) and thus comparable among patches. We specifically sought females, but a few male by-catches were kept

in low-density patches during the final visit, to potentially add genetic diversity in our crosses. We

then identified spiders to species level under a binocular microscope (by placing them in between

a petri dish and some plastic foam, so they were stuck and epigyne/pedipalps visible) following

Roberts (1993). Female *Erigone longipalpis* have a characteristic epigyne (Roberts 1993) that

makes them relatively easy to distinguish from all other linyphiids found in these salt marshes

(based on Leroy et al. 2014), including other *Erigone* species. Although spiders were caught in all

visited sites, *E. longipalpis* were only found in the patches where *P. maritima* was dominant

(i.e. with >50% cover; 5/6 of western and 3/7 of eastern patches sampled). Overall, we caught 38

E. longipalpis female spiders; 34 of those 38 laid at least one eggsac while 26 of those 34 produced

spiderlings. We did not manage to produce spiderlings from one of the eight successfully sampled

patches (as the cocoons produced by the sole sampled female did not hatch); therefore, all

experiments described below were done using lab-born spiders originating from 7 patches (4

western and 3 eastern patches).

Maintenance of spiders in the lab

We kept all spiders in temperature-controlled cabinets (25 ± 1°C) under controlled light regime

(16:8 L:D). They were all housed in plastic boxes and cups with a moist 1cm layer of plaster of

Paris at the bottom (e.g. Mestre and Bonte 2012). Container size, whether spiders were housed

by clutch or individually, and a few other elements, depended on life stage; see details below.

Independently of life stage, we provided all spiders with *ad libitum* access to *Sinella curviseta*

Brook, 1882 springtails as their main food source (*Erigone* spiders in general and *E. longipalpis* in

particular readily predate springtails, e.g. Irmler and Heydemann 1985; Mestre and Bonte 2012).

Adult spiders were kept individually in cylindrical plastic cups (6 cm diameter, 4 cm height). In

addition to springtails, we gave them a *Drosophila* fly once a week (generally *Drosophila*

melanogaster Meigen, 1830 but in some cases *Drosophila suzukii* (Matsumura, 1931) when *D.*

melanogaster supply was low). We renewed springtails and re-humidified the plaster at the same

time. When females laid an egg sac (see below for mating details for lab-born spiders), the eggs

were left in place and the female, along with as many remaining springtails as possible, gently

transferred to a new cup using a small paintbrush. The cup containing the eggs was then placed,

open, into a larger plastic box (8 × 9 cm, height 5 cm). When eggs started to hatch, all spiderlings

were counted and transferred from the cup to the larger box using paintbrushes, if they had not

already moved by themselves. We "seeded" each box with a large number of springtails (> 100

adults), and drilled a small hole (≈ 5 mm diameter and depth) in the plaster that we filled with

rehydrated baker's yeast *Saccharomyces cerevisiae* Meyen ex Hansen, 1883. This provided a food

source for springtails, allowing them to reproduce and grow, guaranteeing sustained resources

for spiderlings. Yeast was rehydrated weekly, at the same time as the plaster, and renewed at the

same time if needed. We moved spiders to individual boxes after the penultimate moult, to

prevent uncontrolled mating events between siblings.

The experiment ran for two generations of lab-reared spiders, from April to late September 2018.

Due to limited room in temperature-controlled cabinets, only ≈ 200 individually housed adults

could be kept at any one time, in addition to all spiderlings' boxes. Every time a "slot" opened,

we chose a replacement among the next adults to emerge at random (accordingly, the number

of lab-born adult spiders per origin patch in the final dataset is correlated with the number of

field-caught spiders from that patch; **Supplementary Material 2**). A total of 530 adults were

maintained, including 293 females. We only tested females in subsequent experiments, both for

logistic reasons and because individual-level correlations between fecundity and other traits can

only be expressed in females.

Dispersal traits

We tested female spiders' dispersal propensity following a protocol inspired by previous studies

(e.g. Bonte, Travis, et al. 2008; De Meester and Bonte 2010; Larrivée and Buddle 2011). We used

circular plaster platforms (diameter 5.5 cm) exposed to a slightly upward (≈ 20°) directional air

flow (wind speed: 1 m.s^{-1} ± 0.2, expected to be optimal for dispersal initiation, Simonneau et al.

2016) as our test arena. Four 17.5 cm long wood skewers were planted vertically, allowing spiders

to climb to initiate dispersal behaviours. We placed test platforms in water-filled trays to prevent

spider escapes.

Spiders were tested individually for 15 minutes at ≈ 25°C. We thoroughly cleaned arenas with

paper towels and water between tests to remove any remaining silk thread that may influence

dispersal decisions (De Meester and Bonte 2010). Individuals were starved (i.e. all remaining

springtails and, more rarely, flies were removed) 12 to 24h before tests to stimulate dispersal

(Weyman et al. 1994).

Preliminary tests to validate the protocol were done using field caught spiders, 5 days post-

capture. For the experiment proper, we tested lab-born spiders on average 8.9 days post-sexual

maturity and before mating (SD: 2.1 days, range: 7-21). This is about a quarter to a fifth of the

typical adult lifespan (see **Results**) and similar to previous studies in other *Erigone* species (7 days

in Bonte, Travis, et al. 2008). Spiders present a stereotyped "tiptoe" behaviour associated with

silk production prior to dispersal, whether it is long-distance dispersal by "ballooning" or short-

distance dispersal by "rappelling" (Bonte 2012). No actual ballooning was observed during the

experiment. This is not surprising however, as while pre-dispersal behaviours are a prerequisite

for actual dispersal, they are not always (and in some cases not often) followed by actual dispersal

during short experimental tests in standardised conditions (Lee et al. 2015). In some cases, spiders

were seen tiptoeing and then immediately walking on the produced thread floating in the wind

after attaching it to the stick; these may be attempts at short-distance dispersal (rappelling), or

attempts at two-step ballooning by "rafting" (Bell et al. 2005). We therefore used the overall

number of tiptoe attempts per trial (whether or not they were followed by a rappelling-like

attempt) as our overall measure of dispersal motivation. Using only the number of rappelling-like

attempts instead leads to similar results (**Data availability**; r between the two variables = 0.89).

We then replaced each spider in its box, and provided it with a fly and springtails.

Mating

Two hours after the dispersal test, each female was presented with a non-sib male originating

from the same source population for mating. Males were left for 24h in the female's box, before

being removed. Some males were reused several times (number of males used: 190, mean

number of mates per used male: 1.45, range: 1-4).

A few spiders laid cocoons before being offered a potential mate. As a precaution, we did not

present a male to these spiders, to avoid potential mixed paternities. However, no spiderling

emerged from any of these "suspicious" cocoons. Unmated spiders can lay unfertilised and

therefore non-viable eggs when no mate is present (Zschokke and Herberstein 2005); we suspect

all these "early" cocoons resulted from this, rather than undetected mating between siblings. As

a result, meaningful fecundity data were available for 273 out of our 293 females.

Other traits

We noted the dates at which each eggsac was laid and hatched, as well as the date of sexual

maturity and death for each female spider kept adult. We used these data to calculate

development times (from hatching to sexual maturity) and adult longevities (from sexual maturity

to death). We note that due to observation biases (detection of hatching) and gaps due to non-

work days, development times and longevities may be recorded with some error; see **Statistical**

analyses and provided code (**Data availability**) for how we accounted for this. For all females that

actually had access to a mate, we additionally recorded the total number of hatchlings produced

as our measure of fecundity. The experiment was stopped on September 30th 2018; 12 female

spiders were still alive at that time, and we used a censoring indicator to include them properly

in longevity analyses (see **Statistical analyses**).

When test females died, they were kept in 90% ethanol before being photographed under

binocular microscope. We measured the cephalothorax width and length of each spider to the

nearest 0.1 mm and used them as proxies of body size (e.g. Eichenberger et al. 2009). We

measured each spider twice to account for measurement error. Other morphological traits that

could have been useful to gauge individual condition (e.g. leg length, opisthosoma size) could not

be obtained consistently, due to spider bodies sometimes losing integrity fast after death (maybe

eaten by springtails), and were thus not used.

Statistical analyses

We analysed our data using a Bayesian workflow, with R (version 4.1.0; R Core Team 2021) and

the *brms* R package (Bürkner 2017) as frontends for the Stan language (Carpenter et al. 2017). In

addition, data preparation, posterior model evaluation and plotting were facilitated by the

tidybayes, bayesplot, patchwork, as well as the *tidyverse* suite of packages (Gabry et al. 2019; Kay

2019; Wickham et al. 2019; Pedersen 2020). See **Supplementary Material 3** for a detailed

description of the models and submodels; we here present a short summary.

We first used a Poisson GLM to determine whether the number of female *E. longipalpis* collected

per patch differed between the two landscapes. This model included an offset/rate term

accounting for the fact sampling effort (in person-hours) varied between patches. We ran this

model twice: once using only visited patches where *Puccinellia maritima* was actually the

dominant plant species (>50% of cover), and once using all visited patches (the second one is

presented as **Supplementary Material 4**).

We then used a multivariate/multiresponse mixed model approach to analyse how our traits of

interest differed between the two landscapes, and whether traits were correlated at the (within-

patch) among-individual level.

• body size was analysed using a Gaussian model on (centred and scaled to 1SD)

cephalothorax width data (similar results were found using cephalothorax length instead;

see code in **Data availability**). The submodel included a fixed effect of landscape, a

random effect of patch of origin and an individual-level random effect. The latter random

effect is used to link size and all the other traits in an individual-level covariance matrix; it

is estimable in our Gaussian model separately from residual error because size was

measured twice per individual. 280 females out of 293 were measured; the others were

too badly damaged for their size to be estimated accurately. We used in-model missing

data imputation to account for these (results are similar if we exclude them instead);

• in adult spiders that survived up to dispersal tests (*N*= 291 out of 293), dispersal was

analysed using the number of tiptoe events per individual and a Poisson model with a log

link. This submodel also included a fixed effect of landscape, a random effect of patch of

origin and an individual-level random effect. The individual-level random effects here act

as observation-level random effects, dealing with overdispersion (Harrison 2014), in

addition to allowing us to build the individual-level covariance matrix;

• in individuals that were able to mate and survive at least one day post-mating (*N* = 272

out of 293), we analysed the number of spiderlings produced per observed day as our

measure of fecundity. The Poisson model was the same as the dispersal one above, but

with the total lifetime fecundity as the response and with the addition of an offset/rate

term denoting the number of observation days (from mating to death/end of experiment);

•   development time (number of days from hatching to sexual maturity) and adult longevity

(days from sexual maturity to death) were also analysed using Poisson models. As above,

these submodels included a fixed effect of landscape, random effects of patch and

individual identity. However, they also included an "observation type" binary variable

denoting whether or not the observations bounding the interval of interest (hatching,

maturity, death) were made the day after a gap in recording, which could bias estimates.

Finally, the longevity model accounted for the fact some individuals were not observed

until natural death by using a right-censoring indicator (equal to 0 if natural death was

recorded during the experiment, and to 1 if the individual outlived the experiment or died

accidentally).

(Note that the methods used in the *brms* package allowed us to estimate all among-individual

covariances even if the number of individuals effectively tested varies between traits as described

above.)

Finally, we re-ran the same multivariate model as above, this time by adding (correlated) random

effects of mother identity in addition to the individual identity random effects. This allowed us to

partition individual-level covariation into its among-family and within-family components, the

former providing an upper bound on the variation attributable to heritable differences. If a

covariation between two traits is only found in the within-family matrix after this partition, then

it likely does not result from genetic correlations. We initially intended to go further and use

pedigree data and animal models (Wilson et al. 2010) to properly separate genetic (co)variances

from maternal effects. However, in all attempts, models failed to converge in a satisfactory way.

We attribute that to our small sample size and to the sparseness of our pedigree (we have no

relatedness information on spiders descending from different wild-caught mothers, let alone

coming from different patches) (Wilson et al. 2010). We therefore settled on an approach that

merely separates within- from among-families (co)variation, using mother ID random effects; we

acknowledge that such an approach cannot separate genetic variation from maternal and

common environment effects, and can thus only provide upper-bound estimates of heritable
variation.

For most parameters, we used weakly informative priors inspired by McElreath (2020):
$\mathrm{Normal}(0,1)$ priors for fixed effects, $\mathrm{half-Normal}(0,1)$ priors for standard deviations (random
effect and residual), and a $\mathrm{LKJ}(3)$ for the correlation matrices. The only exceptions were the
intercepts for development time and adult longevity, for which we shifted the fixed effects prior
means to reflect existing information on development and survival times of *Erigone* spiders (e.g.
Bonte, Travis, et al. 2008; Mestre and Bonte 2012). Again, see **Supplementary Material 3** for a
more formal description of the models and priors. We ran four chains per model for 6000
iterations each, and used the first 3000 iterations of each chain as warmup. Convergence and
mixing were satisfactory based on graphical checks and values of the updated $\hat{R}$ statistic (Vehtari
et al. 2021). In addition, all parameters had tail and bulk-effective sample sizes > 400 (Vehtari et
al. 2021), and most were >1000. All parameter posterior summaries below are given as means
[95% Highest Posterior Density Intervals].

Results

325　*Erigone longipalpis* spiders had lower population densities on the eastern, more fragmented, side
326　of the salt marshes (1.03 [0.50; 1.60] female spiders per person-hour vs 2.84 [1.80; 3.90] on the
327　western side, **Fig. 2**, **Table 1**). This result still holds if we include sampled patches where
328　*Puccinellia* was not dominant and no spiders were found (**Supplementary Material 4**).

Figure 2. Effect of landscape of origin on the number of spiders found per patch (weighted by sampling effort). White dots correspond to observed data, black dots and segments to posterior means and 95% intervals, and the posterior density distributions of the predicted means are displayed in green. Only data from habitat favourable to *Erigone longipalpis*, i.e. *Puccinellia maritima*-dominated meadows, are presented here; for the data and model using all visited sites, see **Supplementary Material 4**.

336 **Table 1.** Effect of landscape of origin on population density and measured traits. The posterior means and
337 95% of the coefficients $\beta_{[east-west]}$ describing the differences between the two landscapes are presented
338 along with their 95% intervals, intervals that do not contain zero are in bold. Note that for population
339 density, time to maturity, dispersal, fecundity and longevity, these coefficients are on the latent log scale,
340 due to the use of Poisson models (see **Methods**). For body size, the values are in SD units.

Response	Posterior difference $\beta_{[east-west]}$
Population size	**-1.04 [-1.68; -0.37]**
Time to maturity	0.06 [-0.15; 0.30]
Body size	0.26 [-0.33, 0.80]
Dispersal	0.18 [-0.38 ; 0.66]
Fecundity	0.69 [0.02; 1.40]
Adult longevity	0.10 [-0.04; 0.22]

We found no evidence for dispersal differences between spiders from the western and eastern
landscapes (4.44 [3.05; 5.90] vs 5.38 [3.10; 7.66] tiptoe events per trial, respectively; **Table 1**,**Fig.**
3a). We also found no meaningful differences in development time (25.30 [21.50; 29.20] vs 26.90
[21.90; 32.00] days), adult longevity (42.00 [37.10; 46.80] vs 46.20 [40.30; 52.50] days) or body
size (0.99 [0.95; 1.02] vs 1.02 [0.97; 1.06] mm)(**Table1**,**Fig. 3**). We found no evidence that
observations of longevity or time to maturity including observation gaps are biased in one
direction or the other (β = 0.07 [-0.04; 0.17] and -0.01 [-0.09; 0.07], respectively). These results
were similar whether we used the model where among-individual variance is unpartitioned
(above, **Fig. 3**), or when we re-run the model, partitioning among-individual variance into its
among- and within-family components (**Supplementary Material 5**). By contrast, there was some
indication that spiders were less fecund in the western more continuous landscape (0.19 [0.12;
0.28] vs 0.38 [0.19; 0.61] spiderlings per day; **Fig.3**, **Table 1**), but contrary to all other (non-)effects
mentioned in this paragraph, support for this one was model-structure dependent: once we re-
ran the model partitioning among-individual variance into its among- and within-family

components, the 95% compatibility interval overlapped with 0 ($\beta_{[east-west]}$= 0.67 [-0.26; 1.60];

Supplementary Material 5).

Figure 3. Phenotypic traits of lab-born spiders as a function of landscape of origin. Observed data are

displayed as boxplots (one boxplot per patch of origin); black dots and segments represent posterior

means and 95% intervals, and are displayed along the posterior density distributions of the predicted

means in green. Predictions are based on the model where individual trait co-variance is not partitioned

between among- and within-family components; see **Supplementary Material 5** for a similar figure based

on the model where individual-level variation is partitioned.

Regarding the structure of the dispersal syndrome, the analysis of the first multivariate model

shows evidence of multiple individual-level correlations among traits, many of them directly

involving dispersal (**Table 2a**). In particular, more dispersive individuals tended to develop faster,

to be larger, and to have higher fecundity per day. There was however no clear correlation

between dispersal and adult longevity. Maturity and fecundity were also correlated, with faster-

developing spiders being more fecund. Larger spiders matured earlier but also died earlier, and

were also more fecund.

When partitioning the individual variance-covariance matrix into its among- and within-family

component, we found that among the correlations described above, the ones that involved

dispersal, fecundity and time to maturity were still found at the within-family level (**Table 2b**) but

not the among-family level (**Table 2c**). On the other end, the size-longevity and size-fecundity

correlations were detectable at the among-family, but not within-family level. A negative size-

maturity correlation (i.e. spiders maturing later were smaller) was found across hierarchical levels

(**Table 2b,c**). A positive longevity-time to maturity correlation was only present at the among-

family level (mothers producing slower-developping offspring also tended to produce long-lived

individuals when adult; **Table 2c**).

 Table 2. Means and 95% Higher Posterior Density intervals for (a) the overall individual-level correlations
383   among traits, (b) the within-family correlations only, and (c) the among-family correlations only. (a) is
384   based on the model where individual trait co-variance is not partitioned between among- and within-
385   family components; see **Supplementary Material 6** for a similar table based on the model where
386   individual-level variation is partitioned (made by adding among- and within-family covariances). Intervals
387   that do not overlap with zero are in bold.

(a) overall within-patch, among-individual correlations

	Time to maturity	Body size	Dispersal	Fecundity
Body size	**-0.42 [-0.56; -0.27]**			
Dispersal	**-0.42 [-0.58; -0.25]**	**0.18 [0.04; 0.31]**		
Fecundity	**-0.33 [-0.50; -0.17]**	**0.25 [0.12; 0.38]**	**0.23 [0.09; 0.36]**	
Adult longevity	0.04 [-0.12; 0.21]	**-0.27 [-0.40; -0.14]**	0.01 [-0.14; 0.15]	-0.03 [-0.18; 0.12]

(b): within-family component only (individual ID random effect)

Body size	**-0.38 [-0.57; -0.18]**			
Dispersal	**-0.40 [-0.61; -0.19]**	0.11 [-0.04; 0.26]		
Fecundity	**-0.34 [-0.55; -0.11]**	0.08 [-0.07; 0.22]	**0.18 [0.01; 0.34]**	
Adult longevity	-0.18 [-0.39; 0.05]	-0.03 [-0.18; 0.12]	0.09 [-0.06; 0.25]	0.02 [-0.15; 0.20]

(c): among-family component only (mother ID random effect)

Body size	**-0.33 [-0.62; -0.01]**			
Dispersal	-0.39 [-0.84; 0.08]	0.29 [-0.16; 0.75]		
Fecundity	-0.24 [-0.53; 0.10]	**0.45 [0.17;0.71]**	0.24 [-0.23; 0.68]	
Adult longevity	**0.41 [0.03; 0.75]**	**-0.67 [-0.90; -0.41]**	-0.31 [-0.79; 0.19]	-0.17 [-0.53; 0.19]

Discussion

The loss and fragmentation of sheep-grazed *Puccinellia maritima* meadows, the main habitat

supporting *Erigone longipalpis* spiders (Leroy et al. 2014) has been a long-term trend in the Mont-

Saint Michel Bay, going back at least 30 years (Valéry et al. 2017). As this loss is not uniform

throughout the region, we have been able to compare spiders from landscapes differing in habitat

availability (**Supplementary Material 1**). We show that population densities are markedly lower

in the most fragmented (*sensu lato*) landscape (**Fig. 2**). While the effects of fragmentation *sensu*

stricto on biodiversity are still debated (Fletcher et al. 2018; Fahrig et al. 2019), the negative

effects of habitat loss itself are more clearly supported (whether they are attributed to reduction

in patch size or reduction in total habitat amount; Bender et al. 1998; Fahrig 2003; Watling et al.

2020). Our results here add one more piece of evidence in that direction. The fact that habitat

loss exerts such a strong negative pressure on *Erigone longipalpis* would suggest that it may also

shapes which phenotypes are advantageous, and therefore lead to phenotypic divergence

between fragmented and (relatively) more intact landscapes. In addition, changes in population

density itself are expected to shape dispersal and related traits (De Meester and Bonte 2010;

Endriss et al. 2019). These changes can in some cases carry over across generations even when

they are not underpinned by genetic variation, through parental effects (Bitume et al. 2014).

For the above reasons, it is very surprising to see that differences in habitat availability have

actually little to no effect on phenotypic traits, including dispersal (**Fig. 3**). This is especially true

given the reported cases of phenotypic changes in response to habitat fragmentation and loss

(Cheptou et al. 2017; Cote et al. 2017). Here, at most, *E. longipalpis* spiders from the more habitat-

limited landscape may be slightly more fecund, but even that finding is model-dependent and

uncertain. It is possible that our study was simply not done at the right scale, whether in space or

in time. On the spatial side, our two landscapes are separated from each other by about 5 km

(**Fig. 1**), and other suitable salt marshes can be found at similar distances on the other side of the

bay (Valéry et al. 2017). Spiders can in principle balloon over much larger distances than that (Bell

et al. 2005). As prevailing winds in the region blow from the (south-)west (DREAL Normandie

2020), this could mean populations in the eastern fragmented habitat are regularly supplied with

immigrants from the western continuous habitat, which would make adaptation to local

conditions much more complicated (Garant et al. 2007). However, the long-distance ballooning

events held as records are unlikely to be representative of typical dispersal distances, and unlikely

to contribute much to gene flow between already established populations. Indeed, several lines

of evidence suggest that the actual typical dispersal distances in ballooning spiders are closer to

a few hundred meters to a few kilometres at most, meaning that our two landscapes could

actually be isolated enough to diverge. These lines of evidence include population genetic studies

directly examining how gene flow or genetic similarity vary with distance (Croucher et al. 2011;

Reed et al. 2011), or indirect studies looking at which spatial scale influence of landscape of spider

abundances is the most visible (Schmidt et al. 2008; a metric expected to be tightly linked to

typical dispersal distances, see e.g. Jackson and Fahrig 2012). Also, the potential for long-distance

dispersal and gene flow is not always, in itself, a barrier for the evolution of dispersal (Cheptou et

al. 2008). Alternately, if long-distance dispersal is rare in *E. longipalpis* (Meijer 1977), the

individual differences in tiptoe frequency that we observed may actually be more reflective of

short-distance dispersal by rappelling than of ballooning. Then it may be that spiders do not

experience any fragmentation and habitat loss at the scale at which they disperse, even in the

eastern habitat. Indeed, typical rappelling distances are up to a few meters (Bonte, Lukáč, et al.

2008), and even the small patches in the eastern fragmented landscape are an order of magnitude

bigger (**Supplementary Material 1**). It might be possible to resolve this uncertainty through

further experiments that cover an even broader range of habitat loss and fragmentation, or by

experimental refinements that would allow ballooning to be more easily observable in the lab,

even if it is rare (for possible ideas, see Narimanov et al. 2021). This would make it easier to clearly

separate variation in long-distance and short-distance dispersal. Regarding the temporal aspect,

available information suggests that the dynamics of the two landscapes started to diverge ≈ 30

years ago (Valéry et al. 2017), which is a minimum of 30 generations ago (although *E. longipalpis*

can develop much faster in continually favourable lab conditions, see **Fig. 3**, they seem to have

an annual life cycle in natural conditions; Irmler and Heydemann 1985). This would in principle be

enough to drive the rapid evolution of dispersal (Cheptou et al. 2008). However, landscape

transformation happened progressively and not suddenly. The key question here is then: when

did the landscapes became divergent enough for it to meaningfully influence *Erigone longipalpis*

biology? To the best of our knowledge, the long-term data that are needed to answer this

question are unfortunately not available even for better-studied linyphiids, let alone *E.*

longipalpis. Further studies at larger scales, both in space and time, or including estimates of

actual gene flow among populations/landscapes (Reed et al. 2011), are again probably needed

before concluding definitely on the effect of habitat loss on *Erigone longipalpis* evolution.

We found substantial evidence that key life-history traits are integrated with dispersal in a

syndrome in this species (**Table 2**). *Erigone longipalpis* seems to present a within-species pace of

life syndrome, with faster-developing spiders also being more fecund on a per day basis (Ricklefs

and Wikelski 2002; **Table 2a**, Réale et al. 2010; Healy et al. 2019). Dispersal was aligned with this

syndrome in a way that follows some existing predictions and results (e.g. Réale et al. 2010;

Stevens et al. 2013), with faster-growing individuals being more likely to disperse. When the

among-individual covariance matrix was split into its within- and among-family components (the

latter reflecting genetic variation, among other things), however, we found that dispersal-related

correlations, and the development time-fecundity correlation expected from the pace-of-life

hypothesis were only found at the within-family level (**Table 2b,c**). In other terms, there was no

evidence that families that differed in dispersal propensity consistently differed in other traits,

which may explain why there was no "coordinated" responses to habitat loss between traits (**Fig.**

3). This result is interesting given a recent meta-analysis suggested there is actually little to no

empirical support for the hypothesis that risky behaviour and fast pace of life should be positively

correlated at the genetic level, even when they are correlated at the individual level (Royauté et

al. 2018). In addition, a key prediction of the pace-of-life syndrome hypothesis, that fast

development is associated with reduced longevity (Réale et al. 2010; Healy et al. 2019) is not

supported at the individual level, but is validated at the family level (**Table 2**). While this specific

association may well be genetically driven, one must remember that we cannot separate additive

genetic from maternal effects in this setup. As mothers that produced consistently fast-

developing individuals also tended to produce larger than average individuals (see the size-time

to maturity correlation **Table 2c**), the idea that family differences are driven by maternal effects

more than genetic differences is worth considering. For instance, in the social spider *Cyrtophora*

citricola, Yip et al. (2021) were able to show that that parental, rather than genetic effects,

explained differences in behaviour among families. More generally, maternal effects on key traits

are regularly observed in spiders (e.g. Mestre and Bonte 2012; Quiñones-Lebrón et al. 2021). So,

in *Erigone longipalpis*, different parts of what we hypothesized to be a common genetic pace of

life syndrome actually exist at different levels of organisation, and the hypothesized dispersal-

pace of life syndrome only exists at the within-family level, meaning it is presumably not driven

by genetic variation. Altogether, these results highlight the need to reevaluate the assumptions

behind the pace of life syndrome hypothesis, and more generally behind general hypotheses on

the integration between (dispersal) behaviour and life history, in particular by better accounting

for hierarchical levels (among-species vs. individual vs. genetic correlations), dispersal context

dependency and the role of resource availability (Bonte and Dahirel 2017; Wright et al. 2019;

Laskowski et al. 2020 Nov). This includes for instance testing for context-dependent syndromes;

our present dataset is however too small in our opinion for such tests.

Even though we did not find support for a dispersal-life-history syndrome at the family/genetic

level (no correlations between dispersal and other traits were retained in the among-family

matrix), this syndrome still exists at the individual level (**Table 2b**). The question that remains,

then, is how such a structured among-individual variation can arise if it is not apparently linked

to genetic covariation or to variation in abiotic conditions (which were held constant among

individuals in the experimental setup). A first possible set of explanations for a non-genetic

dispersal syndrome would be that direct costs incurred during dispersal, independently of the

cause of dispersal, limit fecundity or longevity afterwards (Bonte et al. 2012). Indeed, reduction

of fecundity in direct relation to previous dispersal/movement is well documented in arthropods

(Roff 1977; Matsumura and Miyatake 2018). There are two obvious drawbacks here: first, this

prediction is the exact opposite of the result we observed here and second, incurred costs of

dispersal cannot explain the association between dispersal and development time, as the latter

is expressed before dispersal. Alternatively, a second set of explanations may involve variation in

biotic conditions, more specifically experienced competition. Food availability during

development can be a major driving force of phenotypic variation in spiders, including in dispersal

and in the other traits measured in this study (Mestre and Bonte 2012; Kleinteich et al. 2015;

Quiñones-Lebrón et al. 2021). The same is true for population density during development (De

Meester and Bonte 2010). Although food was provided essentially *ad libitum*, we suggest that the

observed dispersal syndrome can be explained by within-family variation in early access to

resources (stochastic and/or due to unobserved trait variation, in e.g. functional response)

causing variation in development time and adult size, which in turn cause variation in dispersal,

fecundity and longevity. Indeed, in snails for instance, variation in early access to food (in the

form of egg cannibalism) is well-known and can drive long-lasting within-clutch differences in key

traits, all else being equal (Desbuquois 1997). Sibling cannibalism is also present in linyphiid

spiders (Vanacker et al. 2004). At the scale of our dataset, this variability may have been further

amplified by the fact we let spiderlings from the same clutch grow in the same box together,

leading to variability in experienced density between clutches (see fecundity data **Fig. 3** and De

Meester and Bonte 2010). Even without cannibalism, variation in density and competition may

have contributed to non-random juvenile survival with respect to traits, further contributing to

the non-genetic syndrome. Further studies comparing dispersal syndromes in spiders reared in

isolation versus in groups could help confirm or infirm this hypothesis. Several studies have now

shown that dispersal syndromes can be context-dependent and change structure depending on

environmental conditions including temperature (in spiders, Bonte and Dahirel 2017), landscape

matrix harshness (in ciliates, Jacob et al. 2020) or indeed food availability (in flies, Mishra et al.

2018).

To summarise, despite strong ecological effects of habitat loss, and the existence of a dispersal

syndrome, we found no evidence of evolutionary responses to habitat loss in a specialist spider.

Although confirmations are needed, our results highlight the importance of

environmental/developmental conditions in driving dispersal syndromes at the within-species

level (Bonte and Dahirel 2017). Even though we found no support for a genetic dispersal

syndrome, observed associations between dispersal and traits can still substantially shape the

dynamics of *Erigone longipalpis* in these salt marshes. Indeed, given the rarity of suitable habitats,

it is possible that a specific subset of phenotypes is bound to be disproportionally lost during

dispersal in the more fragmented landscape, which may have ecological consequences both on

spider population dynamics and on their interactions with prey. It is even possible that, in the

absence of dispersal evolution to compensate for it, the systematic loss of more fecund

individuals to dispersal is itself one of the causal mechanisms for the observed ecological effects

of habitat loss on this habitat specialist spider.

Acknowledgements

We warmly thank Martin Entling for the starter *Sinella curviseta* springtail population, as well as Youn Henry and Thomas Enriquez for giving us access to a regular supply of *Drosophila* flies. Amélie Beillard kindly helped take care of the springtail populations at critical times. We are also grateful to Armelle Ansart for access to temperature-controlled cabinets, and to Françoise Burel who supported some exploratory analyses linked to this study. We also thank Shakira Quiñones for a friendly review of the manuscript before submission, and three anonymous reviewers for their comments.

Author contributions

Initial idea: MD and JP; funding acquisition: MD, SC, JP; site selection and sampling: MD, MW, SC, JP; behavioural experiment design and rearing protocol: MD, MW, TB; behavioural and life history experiments and population maintenance: MW, MD; data analysis: MD, MW; lead manuscript writer: MD.

Data availability

Data and R scripts to reproduce all analyses presented in this manuscript are available on Github (https://github.com/mdahirel/erigone-dispersal-2018) and Zenodo (https://doi.org/10.5281/zenodo.5761384)

Funding

This project was funded by a research grant of the Observatoire des Sciences de l'Univers de Rennes (UMS OSUR) to MD, SC and JP, and by an internal grant of the UMR Ecobio (Biological Invasions research axis) to MD and JP.

References

Bell JR, Bohan DA, Shaw EM, Weyman GS. 2005. Ballooning dispersal using silk: world fauna, phylogenies, genetics and models. *Bull Entomol Res* 95:69-114.

Bender DJ, Contreras TA, Fahrig L. 1998. Habitat loss and population decline: a meta-analysis of

the patch size effect. *Ecology* 79:517–533.

Benton TG, Bowler DE. 2012. Linking dispersal to spatial dynamics. In: Clobert J, Baguette M,

Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford, UK: Oxford University

Press. p. 251–265.

Bitume EV, Bonte D, Ronce O, Olivieri I, Nieberding CM. 2014. Dispersal distance is influenced by

parental and grand-parental density. *Proc R Soc B* 281:20141061.

Bonte D. 2012. Case study II: Spiders as a model in dispersal ecology and evolution. In: Clobert J,

Baguette M, Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford: Oxford

University Press. p. xxvi–xxviii.

Bonte D, Dahirel M. 2017. Dispersal: a central and independent trait in life history. *Oikos* 126:472–

479.

Bonte D, Lukáč M, Lens L. 2008. Starvation affects pre-dispersal behaviour of *Erigone* spiders.

Basic Appl Ecol 9:308–315.

Bonte D, Saastamoinen M. 2012. Dispersal syndromes in butterflies and spiders. In: Clobert J,

Baguette M, Benton TG, Bullock JM, editors. Dispersal ecology and evolution. Oxford: Oxford

University Press. p. 161–170.

Bonte D, Travis JMJ, Clercq ND, Zwertvaegher I, Lens L. 2008. Thermal conditions during juvenile

development affect adult dispersal in a spider. *PNAS* 105:17000–17005.

Bonte D, Van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, Lehouck V, Matthysen E, Mustin

K, Saastamoinen M, et al. 2012. Costs of dispersal. *Biol Rev* 87:290–312.

Bürkner P-C. 2017. brms: an R package for Bayesian multilevel models using Stan. J Stat Softw

80:1–28.

Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, Brubaker M, Guo J, Li P,

Riddell A. 2017. Stan: a probabilistic programming language. J Stat Softw 76:1–32.

Cheptou P-O, Carrue O, Rouifed S, Cantarel A. 2008. Rapid evolution of seed dispersal in an urban

environment in the weed *Crepis sancta*. *PNAS* 105:3796–3799.

Cheptou P-O, Hargreaves AL, Bonte D, Jacquemyn H. 2017. Adaptation to fragmentation:

evolutionary dynamics driven by human influences. *Phil Trans R Soc B* 372:20160037.

Clobert J, Baguette M, Benton TG, Bullock JM, editors. 2012. Dispersal ecology and evolution.

Oxford, UK: Oxford University Press.

Cote J, Bestion E, Jacob S, Travis J, Legrand D, Baguette M. 2017. Evolution of dispersal strategies

and dispersal syndromes in fragmented landscapes. *Ecography* 40:56–73.

Croucher PJP, Oxford GS, Gillespie RG. 2011. Population structure and dispersal in a patchy

landscape: nuclear and mitochondrial markers reveal area effects in the spider *Theridion*

californicum (Araneae: Theridiidae). Biol J Linn Soc 104:600–620.

De Meester N, Bonte D. 2010. Information use and density-dependent emigration in an agrobiont

spider. Behav Ecol 21:992–998.

Desbuquois C. 1997. Influence of egg cannibalism on growth, survival and feeding in hatchlings of

the land snail *Helix aspersa* Müller (Gastropoda, Pulmonata, Stylommatophora). Reprod Nutr Dev

37:191–202.

DREAL Normandie. 2020. Profil Environnemental Normandie - Climat. Available from:

http://www.normandie.developpement-durable.gouv.fr/le-profil-environnemental-normandie-

r307.html.

Eichenberger B, Siegenthaler E, Schmidt-Entling MH. 2009. Body size determines the outcome of

competition for webs among alien and native sheetweb spiders (Araneae: Linyphiidae). *Ecol*

Entomol 34:363–368.

Endriss SB, Vahsen ML, Bitume EV, Monroe JG, Turner KG, Norton AP, Hufbauer RA. 2019. The

importance of growing up: juvenile environment influences dispersal of individuals and their

neighbours. *Ecol Lett* 22:45–55.

European Environmental Agency. 2020. Corine Land Cover, 2018, Version 2020_20u1. European

Union, Copernicus Land Monitoring Service. Available from: https://land.copernicus.eu/pan-

european/corine-land-cover.

Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Evol Syst* 34:487–

515.

Fahrig L, Arroyo-Rodríguez V, Bennett JR, Boucher-Lalonde V, Cazetta E, Currie DJ, Eigenbrod F,

Ford AT, Harrison SP, Jaeger JAG, et al. 2019. Is habitat fragmentation bad for biodiversity? *Biol*

Cons 230:179–186.

Fletcher RJ, Didham RK, Banks-Leite C, Barlow J, Ewers RM, Rosindell J, Holt RD, Gonzalez A,

Pardini R, Damschen EI, et al. 2018. Is habitat fragmentation good for biodiversity? *Biol Cons*

226:9–15.

Fronhofer EA, Legrand D, Altermatt F, Ansart A, Blanchet S, Bonte D, Chaine A, Dahirel M, Laender

FD, Raedt JD, et al. 2018. Bottom-up and top-down control of dispersal across major organismal

groups. Nature Ecol Evol 2:1859–1863.

Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. 2019. Visualization in Bayesian workflow.

J Royal Stat Soc A 182:389–402.

Garant D, Forde SE, Hendry AP. 2007. The multifarious effects of dispersal and gene flow on

contemporary adaptation. *Funct Ecol* 21:434–443.

GBIF Secretariat. 2022. *Erigone longipalpis* (Sundevall, 1830) *in* GBIF Backbone Taxonomy.

Checklist dataset accessed via GBIF.org on 2022-03-02. https://doi.org/10.15468/39omei.

Govaert L, Fronhofer EA, Lion S, Eizaguirre C, Bonte D, Egas M, Hendry AP, Martins ADB, Melián

CJ, Raeymaekers JAM, et al. 2019. Eco-evolutionary feedbacks—Theoretical models and

perspectives. *Funct Ecol* 33:13–30.

Guerra PA. 2011. Evaluating the life-history trade-off between dispersal capability and

reproduction in wing dimorphic insects: a meta-analysis. *Biol Rev* 86:813–835.

Harrison XA. 2014. Using observation-level random effects to model overdispersion in count data
in ecology and evolution. *PeerJ* 2:e616.

Harvey PR, Nellist DR, Telfer MG, editors. 2002. Provisional atlas of British spiders (Arachnida,
Araneae), volumes 1 and 2. Huntingdon, UK: Biological Records Centre.

Healy K, Ezard THG, Jones OR, Salguero-Gómez R, Buckley YM. 2019. Animal life history is shaped
by the pace of life and the distribution of age-specific mortality and reproduction. *Nature Ecol*
Evol 3:1217–1224.

Institut national de l'information géographique et forestière. 2017. BD ORTHO®. Available from:
http://www.geoportail.gouv.fr/accueil.

Irmler U, Heydemann B. 1985. Populationsdynamik und Produktion von *Erigone longipalpis*
(Araneae, Micryphantidae) auf einer Salzwiese Nordwestdeutschlands. Faun Ökol Mitt 5:443–
454.

Jackson HB, Fahrig L. 2012. What size is a biologically relevant landscape? Landscape Ecol 27:929–
941.

Jacob S, Chaine AS, Huet M, Clobert J, Legrand D. 2019. Variability in dispersal syndromes is a key
driver of metapopulation dynamics in experimental microcosms. *Am Nat* 194:613–626.

Jacob S, Laurent E, Morel-Journel T, Schtickzelle N. 2020. Fragmentation and the context-
dependence of dispersal syndromes: matrix harshness modifies resident-disperser phenotypic
differences in microcosms. *Oikos* 129:158–169.

Kay M. 2019. tidybayes: tidy data and geoms for Bayesian models. Available from:
http://mjskay.github.io/tidybayes/.

Kleinteich A, Wilder SM, Schneider JM. 2015. Contributions of juvenile and adult diet to the
lifetime reproductive success and lifespan of a spider. *Oikos* 124:130–138.

Larrivée M, Buddle CM. 2011. Ballooning propensity of canopy and understorey spiders in a
mature temperate hardwood forest. *Ecol Entomol* 36:144–151.

Laskowski KL, Moiron M, Niemelä P. 2021. Integrating behavior in life-history theory: allocation
versus acquisition? Trends *Ecol Evol* 36:132-138.

Lee VMJ, Kuntner M, Li D. 2015. Ballooning behavior in the golden orbweb spider *Nephila pilipes*
(Araneae: Nephilidae). *Front Ecol Evol* 3:2.

Leroy B, Le Viol I, Pétillon J. 2014. Complementarity of rarity, specialisation and functional
diversity metrics to assess community responses to environmental changes, using an example of
spider communities in salt marshes. *Ecol Indic* 46:351–357.

Little CJ, Fronhofer EA, Altermatt F. 2019. Dispersal syndromes can impact ecosystem functioning
in spatially structured freshwater populations. *Biol Lett* 15:20180865.

Massol F, Altermatt F, Gounand I, Gravel D, Leibold MA, Mouquet N. 2017. How life-history traits
affect ecosystem properties: effects of dispersal in meta-ecosystems. *Oikos* 126:532–546.

Matsumura K, Miyatake T. 2018. Costs of walking: differences in egg size and starvation resistance
of females between strains of the red flour beetle (*Tribolium castaneum*) artificially selected for
walking ability. *J Evol Biol* 31:1632–1637.

McElreath R. 2020. Statistical rethinking: a Bayesian course with examples in R and Stan. 2nd
edition. Boca Raton, USA: Chapman and Hall/CRC.

Meijer J. 1977. The immigration of spiders (Araneida) into a new polder. Ecol Entomol 2:81–90.

Mestre L, Bonte D. 2012. Food stress during juvenile and maternal development shapes natal and
breeding dispersal in a spider. *Behav Ecol* 23:759–764.

Mishra A, Tung S, Shreenidhi PM, Aamir Sadiq M, Shree Sruti VR, Chakraborty PP, Dey S. 2018.
Sex differences in dispersal syndrome are modulated by environment and evolution. *Philos Trans*
R Soc Lond B Biol Sci 373:20170428.

Narimanov N, Bonte D, Mason P, Mestre L, Entling MH. 2021. Disentangling the roles of electric
fields and wind in spider dispersal experiments. *J Arachnol* 49:380–383.

Ochocki BM, Saltz JB, Miller TEX. 2020. Demography-dispersal trait correlations modify the eco-
evolutionary dynamics of range expansion. *Am Nat* 195:231–246.

Pedersen TL. 2020. patchwork: the composer of plots. Available from: https://CRAN.R-
project.org/package=patchwork.

Pétillon J, Georges A, Canard A, Lefeuvre J-C, Bakker JP, Ysnel F. 2008. Influence of abiotic factors
on spider and ground beetle communities in different salt-marsh systems. *Basic Appl Ecol* 9:743–
751.

Pétillon J, Georges A, Canard A, Ysnel F. 2007. Impact of cutting and sheep grazing on ground–
active spiders and carabids in intertidal salt marshes (Western France). *Anim Biodivers Conserv*
30:201–209.

Quiñones-Lebrón SG, Kuntner M, Kralj-Fišer S. 2021. The effect of genetics, diet, and social
environment on adult male size in a sexually dimorphic spider. *Evol Ecol* 35:217–234.

R Core Team. 2021. R: a language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing. Available from: https://www.R-project.org/.

Réale D, Garant D, Humphries MM, Bergeron P, Careau V, Montiglio P-O. 2010. Personality and
the emergence of the pace-of-life syndrome concept at the population level. Phil Trans R Soc B
365:4051–4063.

Reed DH, Teoh V-H, Stratton GE, Hataway RA. 2011. Levels of gene flow among populations of a
wolf spider in a recently fragmented habitat: current versus historical rates. Conserv Genet
12:331–335.

Ricklefs RE, Wikelski M. 2002. The physiology/life-history nexus. Trends *Ecol Evol* 17:462–468.

Roberts MJ. 1993. The spiders of Great Britain and Ireland. Colchester, UK: Harley Books.

Roff D. 1977. Dispersal in Dipterans: its costs and consequences. *J Anim Ecol* 46:443–456.

Ronce O, Clobert J. 2012. Dispersal syndromes. In: Clobert J, Baguette M, Benton TG, Bullock JM,
editors. Dispersal ecology and evolution. Oxford, UK: Oxford University Press. p. 119–138.

Royauté R, Berdal MA, Garrison CR, Dochtermann NA. 2018. Paceless life? A meta-analysis of the
pace-of-life syndrome hypothesis. Behav Ecol Sociobiol 72:64.

Royauté R, Hedrick A, Dochtermann NA. 2020. Behavioural syndromes shape evolutionary
trajectories via conserved genetic architecture. *Proc R Soc B* 287:20200183.

Schmidt MH, Thies C, Nentwig W, Tscharntke T. 2008. Contrasting responses of arable spiders to
the landscape matrix at different spatial scales. *J Biogeogr* 35:157–166.

Simonneau M, Courtial C, Pétillon J. 2016. Phenological and meteorological determinants of
spider ballooning in an agricultural landscape. *C R Biol* 339: 408-416.

Stevens VM, Trochet A, Blanchet S, Moulherat S, Clobert J, Baguette M. 2013. Dispersal
syndromes and the use of life-histories to predict dispersal. *Evol Appl* 6:630–642.

Stevens VM, Trochet A, Van Dyck H, Clobert J, Baguette M. 2012. How is dispersal integrated in
life histories: a quantitative analysis using butterflies. *Ecol Lett* 15:74–86.

Stevens VM, Whitmee S, Le Galliard J-F, Clobert J, Böhning-Gaese K, Bonte D, Brändle M, Matthias
Dehling D, Hof C, Trochet A, et al. 2014. A comparative analysis of dispersal syndromes in
terrestrial and semi-terrestrial animals. *Ecol Lett* 17:1039–1052.

Valéry L, Radureau A. 2015. Evolution de la végétation des marais salés de la Baie du Mont-Saint-
Michel : analyse cartographique 1984, 2013. Conservatoire du littoral.

Valéry L, Radureau A, Lefeuvre J-C. 2017. Spread of the native grass *Elymus athericus* in salt
marshes of Mont-Saint-Michel bay as an unusual case of coastal eutrophication. J Coast Conserv
21:421–433.

Vanacker D, Deroose K, Pardo S, Bonte D, Maelfait J-P. 2004. Cannibalism and prey sharing among
juveniles of the spider *Oedothorax gibbosus* (Blackwall, 1841) (Erigoninae, Linyphiidae, Araneae).
Belg J Zool 134:23–28.

Vehtari A, Gelman A, Simpson D, Carpenter B, Bürkner P-C. 2021. Rank-normalization, folding, and localization: an improved $\hat{R}$ for assessing convergence of MCMC (with Discussion). *Bayesian Anal* 16:667-718

Watling JI, Arroyo-Rodríguez V, Pfeifer M, Baeten L, Banks-Leite C, Cisneros LM, Fang R, Hamel-Leigue AC, Lachat T, Leal IR, et al. 2020. Support for the habitat amount hypothesis from a global synthesis of species density studies. *Ecol Lett* 23:674–681.

Weyman GS, Sunderland KD, Fenlon JS. 1994. The effect of food deprivation on aeronautic dispersal behaviour (ballooning) in *Erigone* spp. spiders. *Entomol Exp Appl* 73:121–126.

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A, Henry L, Hester J, et al. 2019. Welcome to the Tidyverse. *J Open Source Softw* 4:1686.

Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, Kruuk LEB, Nussey DH. 2010. An ecologist's guide to the animal model. *J Anim Ecol* 79:13–26.

Wright J, Bolstad GH, Araya-Ajoy YG, Dingemanse NJ. 2019. Life-history evolution under fluctuating density-dependent selection and the adaptive alignment of pace-of-life syndromes. *Biol Rev* 94:230–247.

Yip EC, Smith DR, Lubin Y. 2021. Causes of plasticity and consistency of dispersal behaviour in a group-living spider. *Anim Behav* 175:99–109.

Zschokke S, Herberstein ME. 2005. Laboratory methods for maintaining and studying web-building spiders. *J Arachnol* 33:205–213.