Differential effects of soil conservation practices on arthropods and crop yields

Running title: Tillage impacts on arthropods and crop yields

Elinor M. Lichtenberg[1,2*], Ivan Milosavljević[2,3], Alistair J. Campbell[4], David W. Crowder[2]

1) Department of Biological Sciences and Advanced Environmental Research Institute,

University of North Texas, Denton, Texas, USA

2) Department of Entomology, Washington State University, Pullman, WA, USA

3) Department of Entomology, University of California, Riverside, CA, USA

4) Laboratório de Entomologia, Embrapa Amazônia Oriental, Belém, Pará, Brazil

*Corresponding author: 1155 Union Circle #310559, Denton, TX 76203-5017; elichten@unt.edu
13

15   **Acknowledgements**

16   We thank K. Fillion, L. Rafferty, G. Smetzler, and D. Elicious for field and lab work, the

17   growers who provided access to field sites, and three anonymous reviewers. Funding was

18   provided by the Western Oilseed Cropping Systems program and USDA Hatch grant 1014754.

Abstract

Many agricultural management tactics, such as reduced tillage, aim to promote biodiversity and ecosystem services. Responses to such tactics can be context dependent, however, and differentially impact (i) functional groups of service-providing organisms and (ii) crop yields. In canola (*Brassica napus* L, *B. rapa* L) crop fields we assessed how soil tillage and landscape context (amount of semi-natural habitat within 1 km of each field) affected arthropod biodiversity and crop yield. We assessed effects of full (multiple tillage passes that leave soil surface bare), intermediate (tilled once and some stubble remains), or no (seed planted directly into last year's stubble) tillage on functional groups with unique diets and reproductive strategies: (i) herbivores, (ii) kleptoparasites, (iii) parasitoids, (iv) pollinators, and (v) predators. Effects of tillage and landscape context on arthropod abundance and diversity varied across functional groups. Pollinators responded strongest to tillage, benefitting from intermediate tillage. Predators and herbivores responded strongly to landscape context, as both were more abundant in landscapes with more semi-natural habitat. Our results suggest natural history differences among functional groups mediate effects of landscape context on biodiversity. However, variation in arthropod communities had little effect on canola crop yield. The effects of soil management practices on aboveground arthropods are complex, and practices thought to increase some aspects of agricultural sustainability may not be beneficial in other contexts. Identifying practices such as intermediate tillage that may increase soil quality and arthropod diversity is key to designing agricultural ecosystems that will effectively benefit both biodiversity and human wellbeing.

KEYWORDS: sustainable agriculture, biodiversity, crop yield, tillage, pollinators, natural enemies, soil habitat

Introduction

Agricultural management tactics such as intercropping and reduced tillage are implemented in an effort to support biodiversity and ecological services in agroecosystems without sacrificing crop yield. Reduced tillage, for example, supports biodiversity by creating soil habitat availability and heterogeneity (de Graaff, Hornslein, Throop, Kardol, & van Diepen, 2019; Tamburini et al., 2020). However, responses of organisms to practices such as tillage often vary among service-providing functional groups (Lefcheck et al., 2015; Mitchell et al., 2015). This has led to calls to better assess impacts of habitat change on multiple ecosystem services such as soil quality, pollination, and crop yield (Bommarco, Kleijn, & Potts, 2013; Tamburini et al., 2020). Given the complexity of agricultural food webs, studies should also assess how organisms in unique functional groups respond to management, and whether there are trade-offs or synergies in managing for different ecosystem services.

Effects of in-field soil management practices on ecosystem service providers likely depend on how animal functional groups interact with soil. For example, tillage can harm ground-nesting bees (Ullmann, Meisner, & Williams, 2016) and predators that shelter among weeds (Cranshaw, 2004), but often has little to no impact on herbivorous pests that feed on crops (Tooker, O'Neal, & Rodriguez-Saona, 2020). Reduced tillage is implemented to limit soil erosion and conserve soil moisture. However, it can also affect soil chemical and physical profiles, and availability of flowering plants that feed beneficial insects (Kennedy & Schillinger, 2006). However, impacts of soil management practices on distinct functional groups may also depend on the landscape context. Semi-natural habitat near farms can facilitate dispersal of organisms to crops (Kremen et al., 2007; Tscharntke et al., 2012) and responses of organisms to particular practices may depend on the landscape context. Indeed, some studies show benefits of sustainable agricultural practices

accrue most strongly in relatively simple landscapes, while others show the greatest benefits in

complex landscapes (Kennedy et al., 2013; Lichtenberg et al., 2017; Scheper et al., 2013;

Tscharntke et al., 2012).

While supporting biodiversity is often a goal of agricultural production systems, crops must

also generate high yield. Because yield captures the total contribution of biotic communities and

farming practices, it can be difficult to ascribe yield to single factors (Tamburini, Bommarco,

Kleijn, van der Putten, & Marini, 2019). Studies are thus needed that use a single analytical

framework to assess how habitat availability and diversification affect biodiversity and yield,

both directly and through indirect interactions among organisms and management practices

(Birkhofer et al., 2015; Byrnes et al., 2014; Weekers et al., 2022). Such studies may be

particularly useful when conducted at field-level scales that involve commercial production and

representative growing practices for a given region.

Here we assessed how soil tillage affected arthropod functional groups in canola (*Brassica*

napus L, *B. rapa* L) crops of the Pacific Northwest USA, and how landscape context and tillage

interacted with arthropods to affect yield. Canola provides floral food for pollinators and natural

enemies, but attract pests like aphids (Aphididae) and flea beetles (Chrysomelidae). We first

asked if effects of tillage on arthropod abundance and diversity varied by functional group. We

hypothesised functional groups with soil dependence, like pollinators and kleptoparasites, would

be most strongly negatively impacted by tillage (Rowen, Regan, Barbercheck, & Tooker, 2020).

Second, we asked if responses of functional groups to tillage depended on landscape context,

given variation in mobility and habitat needs of unique organisms (Lichtenberg et al., 2017;

Marja, Tscharntke, & Batáry, 2022). Third, we asked whether variation in arthropod

communities interacted with tillage and landscape context to affect yield (Delaplane & Mayer,

2000; Morandin & Winston, 2005; Reddy, 2017). Reduced tillage often decreases crop yield

(e.g., Lundin, 2019; Tamburini et al., 2020), but enhanced pest control or pollination could

counteract this. This allowed us to assess how agricultural practices directly and indirectly

affected multiple ecosystem services and biodiversity across landscapes.

Materials and Methods

Study sites

We sampled arthropods in spring canola fields in eastern Washington and northern Idaho

during 2013 and 2014 (Fig. S1; Table S1). This heavily agricultural region has patches of semi-

natural habitat amidst considerable acreage of grains, legumes, and canola (Painter, Hinman, &

Roe, 2006; USDA NASS, 2020). The region's loess soils are sandy loams. As a mass-flowering

crop that blooms for up to a month, canola attracts flower-feeding arthropods such as pollinators

and natural enemies like predators and parasitoids (Delaplane & Mayer, 2000; Morandin &

Winston, 2005). Canola fields are thus an effective model system for studying multiple

ecosystem service providers across unique functional groups.

We selected 15 spring-planted canola fields each year ranging from 0.7 to 142 ha (mean ±

SD = 44.4 ± 41.0). The short canola bloom period and relatively small number of spring canola

fields in the region limited further sampling. Most fields were maintained by local farmers with

four maintained by local universities or seed companies. Farmers used several different canola

varieties, sometimes within one field. Most of these varieties were glyphosate resistant, with one

farmer using Beyond® (herbicide) resistant seed. All seeds were treated with a neonicotinoid

(mainly thiamethoxam), and some sites applied a pyrethroid insecticide once after bloom to

control flea beetles and cabbage seed pod weevils (*Ceutorhynchus obstrictus* [Marsham]). Most

farmers also applied a herbicide treatment (glyphosate) before bloom. Canola variety used, and

glyphosate and pyrethroid application, were independent of tillage regime. Fields were spaced at

least 2 km apart, which was sufficiently far relative to insect flight distances to consider them

spatially independent.

Arthropod sampling

We used two collection techniques to sample diverse arthropods that associate with canola:

(i) traps typically used to sample bees and (ii) sweep nets. All sampling occurred along one field

edge on days with daytime temperatures above 13 ˚C and wind speed below 4.5 m/s, while fields

were in full bloom. Sampling locations were haphazardly selected along accessible field edges,

where we could work without damaging crops and where traps on the ground would not be

covered in vegetation. At 10:00, we conducted 100 continuous sweeps in the canola canopy,

emptied net contents – including all arthropods – into a plastic bag, and stored them on ice. In the

lab, we freeze-killed arthropods, then sorted them to morphospecies. Bees were pinned and

specimens in other taxa were stored in ethanol.

Around 12:00, we set out a line of bee traps that stayed in place for 24 h. This line included

two blue vane traps (SpringStar) and six 96.1mL pan traps painted in colours that attract bees

(two each of white, fluorescent yellow, fluorescent blue; Kearns & Inouye, 1993; Leong &

Thorp, 1999) and partially filled with soapy water. We separated traps by 5 m and located them

~0.5 m from the field edge. Traps and sweep netting began at the same spot. The following day,

we collected trapped arthropods (bees as well as other arthropods) using a strainer, washed off

excess soap, and stored the specimens in ethanol. In the lab we washed and dried bees, and

sorted all specimens to genus and morphospecies based on morphological characteristics (Arnett,

2000; Arnett & Thomas, 2000; Arnett, Thomas, Skelley, & Frank, 2002; Boyle & Philogène,

1983; Derraik et al., 2002; Derraik, Early, Closs, & Dickinson, 2010; Michener, 2000; Stehr,

1987a, 1987b). Both adults and immature forms of all arthropod groups were considered.

Specimens were also assigned to five functional groups using literature and information

from local species (Arnett, 2000; Michener, 2000; Stehr, 1987a, 1987b): (i) pollinators, (ii)

herbivores, (iii) predators, (iv) parasitoids, and (v) kleptoparasites (Table S2). This classification

considered both diet and reproductive strategy. Herbivores included major regional canola pests

(thrips [Thysanoptera], aphids, flea beetles, cabbage seedpod weevils, and *Lygus* bugs Hahn;

Reddy, 2017). Kleptoparasites are regulators of bee communities, and may reflect overall levels

of pollinator biodiversity (Sheffield, Pindar, Packer, & Kevan, 2013). We also considered

predators and parasitoids pooled together as "natural enemies".

Field and landscape variables

For each site we assessed agronomic and weather factors that can affect insects: (i) tillage,

(ii) field size, (iii) growing degree days, and (iv) cumulative precipitation (Aldercotte, Simpson,

& Winfree, 2022; Forcella et al., 2021; Fragoso, Jiang, Clayton, & Brunet, 2021; e.g., Skellern,

Welham, Watts, & Cook, 2017; Smith et al., 2020). We asked farmers directly to categorize each

site's tillage regime. A field had either (i) full (4 to 7 passes producing bare soil with no stubble

before seeding; n = 6), (ii) intermediate (field tilled once with the same equipment as full tillage

fields, leaving some stubble remaining before seeding; n = 13), or (iii) no (soil not tilled and seed

planted directly into the previous year's stubble; n = 11) tillage. These represent the three main

tillage regimes used in the region. All fields were tilled in spring (typically April into early May).

Because canola is planted very shallow, spring tillage in the region is recommended to be limited

to 2 to 5 cm (Brown, Davis, Lauver, & Wysocki, 2009). We determined field size in ArcGIS by

creating a polygon tracing the boundary and subtracting the areas of all uncultivated patches

(typically remnant prairie within and adjacent to fields; originally mapped by hand). We gathered

daily temperature and precipitation values from PRISM (PRISM Climate Group, 2020) and

calculated growing degree days and cumulative precipitation through the day before we sampled

arthropods. These environmental measures can cause inter-annual variation in arthropod

populations (Forcella et al., 2021; Skellern et al., 2017). PRISM estimates climatic conditions at

a given location on a given day (at 4-km resolution) using climatologically-aided interpolation

based on 30-year patterns (PRISM Climate Group, 2020). Growing degree days were calculated

as the cumulative amount of heat above 5 ˚C since January 1 (Dickson, 2014). We calculated

precipitation cumulatively from January 1.

We also calculated the amount of semi-natural habitat within 1 km of each field, a radius

that is ecologically relevant for pollinators, herbivores, and natural enemies (Greenleaf,

Williams, Winfree, & Kremen, 2007; Rusch et al., 2016). We determined land cover using

CropScape data (USDA SARS, 2014a, 2014b, 2015a, 2015b) to calculate area of each land cover

type around each site. Semi-natural habitat included forests, grasslands, shrublands, and

wetlands, while crop habitats were classified as agricultural.

Data analysis: effects of tillage and landscape on arthropod communities

We used model selection to test how functional groups responded to tillage and landscape

context. We calculated three community metrics – (i) abundance, (ii) richness, and (iii) evenness

(E_{var}, Smith & Wilson, 1996) – for each functional group at each site. Evenness captures how

individuals are distributed across taxa and indicates the degree that rare taxa affect ecosystem

functioning (Crowder, Northfield, Strand, & Snyder, 2010; Winfree, W. Fox, Williams, Reilly,

& Cariveau, 2015). Although richness and evenness are more abstract when not all taxa can be

identified to species, we consistently identified to morphospecies across all fields and thus our

richness and evenness values provide important measures of the sampled communities. We ran

linear regressions for each functional group with tillage, proportion semi-natural habitat, their

interaction, year, degree days, cumulative precipitation, and field size as fixed effects. These

variables were not collinear (Table S4). As abundance data were overdispersed, we used

negative binomial regressions (MASS package, Venables & Ripley, 2002). We analysed richness

and evenness with linear models (identity link), but herbivore evenness was square-root-

transformed due to heteroscedasticity. All analyses met model assumptions. We used

information-theoretic model selection to assess model fit for functional group metrics (MuMIn

package, Barton, 2014), and selected models with AICc values within 2 of the lowest value

(Burnham & Anderson, 1998). Low richness, and thus lack of variation, prevented us from

analysing kleptoparasite and parasitoid evenness; all other metrics were assessed. There were 13

total models, one for each functional group and community metric, each with 30 observations

(28 for predator evenness).

We used multivariate PERMANOVA to test whether community composition responded to

tillage and landscape context (Oksanen et al., 2019). We categorized the following predictors as

"high" (above the median) and "low" (below the median): proportion semi-natural habitat,

degree days, cumulative precipitation, and field size. Analyses also included tillage, the

interaction between tillage and proportion semi-natural habitat, and year as predictors. We used

Sørensen dissimilarity indices based on presence/absence matrices with singletons species

removed. We analysed each functional group separately. Small numbers of kleptoparasite and

parasitoid taxa prevented analysing these groups. To assess each term's significance via type II

sums of squares (as in regressions) we first ran each model with the option for marginal effects.

We then tested significance of tillage and the proportion of semi-natural habitat via two

additional regressions that used sequential order with the variable of interest assessed after all

other fixed effects had been accounted for. To visualize potential community differences as

functions of tillage and landscape context, we plotted a separate Principle Coordinate Analysis

for each variable and functional group (Oksanen et al., 2019).

Data analysis: effects of tillage, landscape and arthropod communities on crop yields

We assessed how arthropod biodiversity, tillage, and landscape context affected canola

crop yield; farmers provided yield data for their sites directly. We ran separate linear regressions

(identity link) with abundance, richness, or evenness that also included tillage, proportion semi-

natural habitat, and their interaction, year, degree days, cumulative precipitation, and field size as

fixed effects. For each set of models, we selected the best fit models as those with AICc values

within 2 of the smallest value. There were three yield models, one for each community metric,

each with 30 observations (28 for the evenness regression).

Results

We collected 21,446 individuals across 130 taxa, with 15,256 herbivores (20 taxa, 82% of

individuals were pest species), 154 kleptoparasites (8 taxa), 1,080 parasitoids (6 taxa), 4,217

pollinators (65 taxa), and 739 predators (32 taxa). The most abundant herbivores were thrips,

aphids, sciaroid flies, and chrysomelids (7,184, 4,078, 2,041, and 576 individuals, respectively).

The most common pollinators were two halictid bee morphospecies (813 *Lasioglossum* and 581

Agapostemon individuals). The most common natural enemies were chalcidoid wasps (727

individuals, parasitoid) and melyrid beetles (229 individuals, predator). Landscapes around our

sites ranged from 0% to 36% semi-natural habitat, and this variable was independent of tillage

regime (Kruskal-Wallis test: $X^2_2 = 2.47$, *P* = 0.29).

Effects of tillage and landscape on arthropod communities

Overall, pollinators and kleptoparasites were affected by tillage while herbivores and

predators responded most strongly to landscape context. Pollinator richness was higher in fields

with intermediate tillage than fields with no tillage (Fig. 1a; Tables S5, S6). Kleptoparasite

abundance was higher in fields with intermediate or no tillage than in heavily-tilled fields in one

of four best models (and significant at $\alpha < 0.10$ in a second model; Fig. 1b; Tables S6, S7).

Pollinator abundance and evenness, and kleptoparasite richness, were not affected by tillage or

landscape context (Tables S5, S7, S8). In contrast, more semi-natural habitat promoted herbivore

and predator abundance regardless of tillage regime (Fig. 2; Table S7). Herbivore diversity and

predator richness were unaffected by tillage or landscape context (Tables S5, S8). Parasitoids

were unaffected by tillage or landscape context (Tables S5, S7, S8). Tillage and landscape

context never had an interactive effect. They also did not affect community composition of any

functional group (Table S9, Fig. S2).

Environmental variables had mixed effects on arthropod abundance and diversity (Tables

S5, S7, S8). Precipitation generally had stronger impacts on arthropod communities than degree

days. Greater kleptoparasite abundance and richness, pollinator abundance, and predator richness

occurred at sites with lower precipitation. Parasitoid abundance and pollinator evenness had the

opposite pattern. Kleptoparasite richness, parasitoid abundance, and predator richness were

highest at cooler sites. Larger fields were associated with greater kleptoparasite abundance and

richness but lower pollinator evenness.

Differential responses of arthropod functional groups may indicate trade-offs when aiming

to manage biodiversity, but we found stronger evidence for biodiversity synergies than trade-

offs. Functional groups with known trophic relationships often had correlated metrics (Table S3).

For example, we found positive correlations in abundance and diversity between pollinators and

kleptoparasites, and among parasitoids, predators, and herbivores. Abundance and evenness of

pollinators was also positively correlated with abundance and evenness of natural enemies, as

well as individual natural enemy groups (predators or parasitoids) (Table S3). However,

biodiversity synergies may be agronomic trade-offs. Abundances of beneficial pollinators and

detrimental herbivores also positively correlated (Table S3).

Effects of tillage, landscape and arthropod communities on crop yields

Tillage strongly affected canola yield, and yield was lower in fields with no tillage than

with full or intermediate tillage (Fig. 3a, Tables S6, S10). Landscape context did not impact

canola yield, however. We found little evidence for effects of landscape context or arthropod

communities on yield (Tables S6, S10; S11). More semi-natural habitat in the surrounding

landscape promoted more diverse herbivore communities, which lowered yield (Fig. 3b; Tables

S6, S10). However, arthropod abundance and evenness, and richness of other functional groups,

did not impact yield.

Discussion

Soil management may differentially affect functional groups due to differences in resource

needs or dispersal among taxa (Bommarco et al., 2013; Harmon-Threatt, 2020). Because many

of these arthropods provide ecosystem services or are pests, biodiversity can affect crop yield.

We found that agricultural landscapes can simultaneously support pollinators and predators, but

different functional groups respond to habitat variability at different scales. Pollinators were

most affected by tillage within fields, while landscape context most strongly affected herbivores

and predators. Yet, no arthropod group strongly impacted crop yield. Our results showed that

agriculture practices alter crop yield directly, but not always indirectly by affecting arthropods

(as in Ricketts et al., 2016).

We found that pollinator and kleptoparasite, but not herbivore, predator, or parasitoid taxa

responded to tillage. Reduced tillage may promote pollinators and kleptoparasites by destroying

fewer ground-nesting bee nests (Kennedy & Schillinger, 2006; Ullmann et al., 2016), but we

found intermediate tillage supported a more diverse pollinator community than no tillage. One

potential explanation is that untilled soil contains a thick layer of crop stems that prevent ground

nesting (Stinner & House, 1990). However, our observed impacts of tillage along with the

presence of kleptoparasites suggests that many bee species nest in canola crop fields, which is

often assumed to not occur (Kleijn, Rundlöf, Scheper, Smith, & Tscharntke, 2011).

Effects of tillage on natural enemies and herbivores are more well studied than for bees

(Furlan, Milosavljević, Chiarini, & Benvegnù, 2021; Rowen et al., 2020; Tooker et al., 2020).

Reduced tillage can benefit natural enemies by promoting weeds that provide nectar or surface

mulch that provides shelter and prey (Clark, Luna, Stone, & Youngman, 1993; Stinner & House,

1990). These mechanisms are likely not operating in our system as canola provides abundant

nectar, farmers controlled weeds, and crops in the region do not contribute much mulch

(Hammel, 1996). When reduced tillage promotes herbivores, it mainly does so due to less soil

disturbance or by promoting weeds (Rowen et al., 2020). Surface mulch can also protect aphids

from predators (Hesler & Berg, 2003). None of these mechanism apply here, since we mainly

sampled herbivores that reside near the tops of plants rather than in soil, farmers manged weeds,

and mulch was minimal in our fields. Thus, it is not surprising that tillage did not affect natural

enemies or herbivores. This highlights the necessity of studying mechanisms mediating how

species' life histories relate to food and shelter resources (Carvalheiro, Bartomeus, Rollin,

Timóteo, & Tinoco, 2021).

Heterogeneous landscapes provide opportunities for consumers to exploit patchy resources

(Tscharntke et al., 2012), and landscapes with more semi-natural habitat had more predators and

herbivores. Predators and our main herbivores (aphids and thrips) routinely travel long distances

in search of suitable habitat (Loxdale & Lushai, 1999; Schellhorn, Bianchi, & Hsu, 2014). In

contrast, most pollinators are central place foragers that repeatedly return to a single nest. Indeed,

bees visiting canola flowers tend to travel only a few metres from their nest (Robinson, 2019).

Thus, most of the bees we collected likely were nesting in or near the canola fields we sampled,

and were thus affected more by tillage than landscape context. This result mirrors meta-analyses

that suggest that highly mobile organisms are more likely to respond to landscape-scale habitat

patterns than less mobile organisms (Lichtenberg et al., 2017; Schneider et al., 2014).

While we showed that local and landscape habitat patterns affected arthropod communities,

these communities minimally impacted crop yields, similar to studies that have found no

relationship between multi-diversity and multifunctionality (Birkhofer et al., 2018). We did find

that yield was lower in fields with higher herbivore richness. Inspection of herbivore abundances

at each site (Table S12) suggests two potential drivers. First, sites with higher herbivore richness

could have higher pest abundance (Table S3). Second, sites with higher herbivore richness could

be more likely to contain a specific damaging pest. Our data shows high-herbivore-richness

fields contained large numbers of aphids, a key canola pest (Reddy, 2017). We also found more

chrysomelids, curculionids, meloids, scraptiids, pentatomids, and yponomeutids in sites with

higher herbivore richness. However, the only canola pests in these groups are seedpod weevils

(Curculionidae), which damage a later crop stage than we sampled (Reddy, 2017).

Abundance and diversity of pollinators and predators also did not affect yield. Canola has

high variability in pollinator dependence (Ouvrard & Jacquemart, 2019), and the varieties in our

study may not be pollinator dependent (Perrot, Gaba, Roncoroni, Gautier, & Bretagnolle, 2018),

or the study region is windy enough to ensure pollen dispersal. Variation in pollinator

dependence may explain differences between our results and studies that find increased yield of

oilseed crops with higher pollinator abundance (Catarino, Bretagnolle, Perrot, Vialloux, & Gaba,

2019). Another potential explanation is that we measured yield across the entire field but insects

only along the field perimeter (due to logistics constraints). Benefits from pollinators and

predators may have also been limited by insecticide use (neonicotinoid treated seeds) that may

control pests and may reduce pollinator abundance. It is also possible that pollinators or

predators correlate with common measures of single ecosystem services, such as pollen

deposition or consumption of sentinel pests on a small subset of plants. If such patterns were

present, they did not scale up to the entire field.

Tillage intensity did affect crop yields, similar to studies showing reduced tillage reducing

yield for oilseed rape and other crops (Lundin, 2019; Tamburini et al., 2020). Indeed, multiple

sustainability-oriented farming practices sometimes result in lower yield than their conventional

counterparts (Smith et al., 2020; Tamburini et al., 2020). Despite yield loss seen here, reduced

tillage can provide other benefits such as improving soil infiltration, reducing erosion, decreasing

evaporative water loss from soil, and improving soil quality (Hammel, 1996; Kennedy &

Schillinger, 2006). Reducing tillage can also reduce fuel, labour, and machine maintenance costs

(although it does require up-front investment in purchasing and calibrating specialized seeding

equipment, and sometimes increased fertilizing; Brown et al., 2009). These factors might

ultimately increase yield of other crops or reduce farmers' costs. This highlights the complex

decisions that underlie farm management.

Overall, our study highlights the need to understand how biodiversity patterns and crop

yields are simultaneously affected by multiple mechanisms, including via soil management, at

various scales. We showed that tillage impacted pollinators, while landscape context strongly

affected predators and herbivores. These differences likely reflect natural history differences

among functional groups. However, these habitat impacts on biodiversity minimally impacted

yield. It is often assumed that enhancing biodiversity promotes ecosystem services, although

evidence from arthropod-mediated ecosystem services such as pollination and pest control is

mixed (e.g., Birkhofer et al., 2018; Dainese et al., 2019; Ricketts et al., 2016). Without clear

evidence that a conservation action such as reduced tillage is likely to increase crop yield,

adoption by farmers may remain low (Kleijn et al., 2019). Thus, data-driven management of

agricultural landscapes to simultaneously support natural biodiversity and ecosystem services

and boost crop yield requires much more research to determine the contexts in which given

management practices, and soil diversification practices in particular, do or do not meet this

multi-faceted goal.

Conflict of Interest Statement

We declare no conflict of interest.

Author Contribution

EML and DWC conceived research. EML conducted experiments. IM identified arthropods.

EML analysed data and conducted statistical analyses. EML, AJC and DWC wrote the

manuscript. DWC secured funding. All authors read and approved the final manuscript.

Data Availability Statement

Data and code will be available via Zenodo at a future date.

References

Aldercotte, A. H., Simpson, D. T., & Winfree, R. (2022). Crop visitation by wild bees declines

over an 8-year time series: A dramatic trend, or just dramatic between-year variation?

Insect Conservation and Diversity, *15*(5), 522–533. DOI: 10.1111/icad.12589

Arnett, R. H., Jr. (2000). *American Insects: A Handbook of the Insects of America North of*

Mexico, Second Edition. CRC Press.

Arnett, R. H., Jr, & Thomas, M. C. (2000). *American Beetles: Archostemata, Myxophaga,*

Adephaga, Polyphaga: Staphyliniformia. CRC Press.

Arnett, R. H., Jr., Thomas, M. C., Skelley, P. E., & Frank, J. H. (2002). *American Beetles,*

Volume II: Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press.

Barton, K. (2014). *MuMIn: Multi-model inference*. R package version 1.47.1, https://CRAN.R-

project.org/package=MuMIn

Birkhofer, K., Andersson, G. K. S., Bengtsson, J., Bommarco, R., Dänhardt, J., Ekbom, B., …

Smith, H. G. (2018). Relationships between multiple biodiversity components and

ecosystem services along a landscape complexity gradient. *Biological Conservation*, *218*,

247–253. DOI: 10.1016/j.biocon.2017.12.027

Birkhofer, K., Diehl, E., Andersson, J., Ekroos, J., Früh-Müller, A., Machnikowski, F., … Smith,

H. G. (2015). Ecosystem services—Current challenges and opportunities for ecological

research. *Frontiers in Ecology and Evolution*, *2*, 97. DOI: 10.3389/fevo.2014.00087

Bommarco, R., Kleijn, D., & Potts, S. G. (2013). Ecological intensification: Harnessing

ecosystem services for food security. *Trends in Ecology & Evolution*, *28*(4), 230–238.

DOI: 10.1016/j.tree.2012.10.012

Boyle, R. M. D., & Philogène, B. J. R. (1983). The native pollinators of an apple orchard:

Variations and significance. *The Journal of Horticultural Science*, *58*(3), 355–363. DOI:

10.1080/00221589.1983.11515130

Brown, J., Davis, J. B., Lauver, M., & Wysocki, D. (2009). *United States Canola Association*

Canola Growers Manual. Washington, DC: United States Canola Association.

Burnham, K. P., & Anderson, D. R. (1998). *Model Selection and Multimodel Inference: A*

Practical Information-Theoretic Approach (2nd ed.). New York: Springer-Verlag.

Byrnes, J. E. K., Gamfeldt, L., Isbell, F., Lefcheck, J. S., Griffin, J. N., Hector, A., … Duffy, J.

E. (2014). Investigating the relationship between biodiversity and ecosystem

multifunctionality: Challenges and solutions. *Methods in Ecology and Evolution*, *5*(2),

111–124. DOI: 10.1111/2041-210X.12143

Carvalheiro, L. G., Bartomeus, I., Rollin, O., Timóteo, S., & Tinoco, C. F. (2021). The role of

soils on pollination and seed dispersal. *Philosophical Transactions of the Royal Society*

B: Biological Sciences, *376*(1834), 20200171. DOI: 10.1098/rstb.2020.0171

Catarino, R., Bretagnolle, V., Perrot, T., Vialloux, F., & Gaba, S. (2019). Bee pollination

outperforms pesticides for oilseed crop production and profitability. *Proceedings of the*

Royal Society B: Biological Sciences, *286*(1912), 20191550. DOI:

10.1098/rspb.2019.1550

Clark, M. S., Luna, J. M., Stone, N. D., & Youngman, R. R. (1993). Habitat preferences of

generalist predators in reduced-tillage corn. *Journal of Entomological Science*, *28*(4),

404–416. DOI: 10.18474/0749-8004-28.4.404

Cranshaw, W. (2004). *Garden Insects of North America*. Princeton, NJ: Princeton University

Press.

Crowder, D. W., Northfield, T. D., Strand, M. R., & Snyder, W. E. (2010). Organic agriculture

promotes evenness and natural pest control. *Nature*, *466*(7302), 109–112. DOI:

10.1038/nature09183

Dainese, M., Martin, E. A., Aizen, M. A., Albrecht, M., Bartomeus, I., Bommarco, R., …

Steffan-Dewenter, I. (2019). A global synthesis reveals biodiversity-mediated benefits for

crop production. *Science Advances*, *5*(10), eaax0121. DOI: 10.1126/sciadv.aax0121

de Graaff, M.-A., Hornslein, N., Throop, H. L., Kardol, P., & van Diepen, L. T. A. (2019).

Effects of agricultural intensification on soil biodiversity and implications for ecosystem

functioning: A meta-analysis. In D. L. Sparks (Ed.), *Advances in Agronomy* (Vol. 155,

pp. 1–44). Cambridge, MA: Academic Press. DOI: 10.1016/bs.agron.2019.01.001

Delaplane, K. S., & Mayer, D. F. (2000). *Crop Pollination by Bees*. New York: CABI

Publishing.

Derraik, J. G. B., Closs, G. P., Dickinson, K. J. M., Sirvid, P., Barratt, B. I. P., & Patrick, B. H.

(2002). Arthropod morphospecies versus taxonomic species: A case study with Araneae,

Coleoptera, and Lepidoptera. *Conservation Biology*, *16*(4), 1015–1023. DOI:

https://doi.org/10.1046/j.1523-1739.2002.00358.x

Derraik, J. G. B., Early, J. W., Closs, G. P., & Dickinson, K. J. M. (2010). Morphospecies and

taxonomic species comparison for Hymenoptera. *Journal of Insect Science*, *10*(108), 1–7.

DOI: 10.1673/031.010.10801

Dickson, T. J. (2014). *Growing Season Weather Impacts on Canola Phenological Development*

and Quality (Masters' thesis, University of Manitoba). University of Manitoba,

Winnipeg, Manitoba.

Forcella, F., Patel, S., Lenssen, A. W., Hoerning, C., Wells, M. S., Gesch, R. W., & Berti, M. T.

(2021). Weather and landscape influences on pollinator visitation of flowering winter

oilseeds (field pennycress and winter camelina). *Journal of Applied Entomology*, *145*(4),

286–294. DOI: 10.1111/jen.12854

Fragoso, F. P., Jiang, Q., Clayton, M. K., & Brunet, J. (2021). Patch selection by bumble bees

navigating discontinuous landscapes. *Scientific Reports*, *11*(1), 8986. DOI:

10.1038/s41598-021-88394-2

Furlan, L., Milosavljević, I., Chiarini, F., & Benvegnù, I. (2021). Effects of conventional versus

no-tillage systems on the population dynamics of elaterid pests and the associated

damage at establishment of maize crops. *Crop Protection*, *149*, 105751. DOI:

10.1016/j.cropro.2021.105751

Greenleaf, S. S., Williams, N. M., Winfree, R., & Kremen, C. (2007). Bee foraging ranges and

their relationship to body size. *Oecologia*, *153*, 589–596.

Hammel, J. E. (1996). Water conservation practices for sustainable dryland farming systems in

the Pacific Northwest. *American Journal of Alternative Agriculture*, *11*(2/3), 58–63.

Harmon-Threatt, A. (2020). Influence of nesting characteristics on health of wild bee

communities. *Annual Review of Entomology*, *65*(1), 39–56. DOI: 10.1146/annurev-ento-

011019-024955

Hesler, L. S., & Berg, R. K. (2003). Tillage impacts cereal-aphid (Homoptera: Aphididae)

Infestations in spring small grains. *Journal of Economic Entomology*, *96*(6), 1792–1797.

DOI: 10.1093/jee/96.6.1792

Kearns, C. A., & Inouye, D. W. (1993). *Techniques for Pollination Biologists*. Niwot, CO:

University Press of Colorado.

Kennedy, A. C., & Schillinger, W. F. (2006). Soil quality and water intake in traditional-till vs.

No-till paired farms in Washington's Palouse region. *Soil Science Society of America*

Journal, *70*(3), 940–949. DOI: https://doi.org/10.2136/sssaj2005.0160

Kennedy, C. M., Lonsdorf, E., Neel, M. C., Williams, N. M., Ricketts, T. H., Winfree, R., …

Kremen, C. (2013). A global quantitative synthesis of local and landscape effects on wild

bee pollinators in agroecosystems. *Ecology Letters*, *16*(5), 584–599. DOI:

10.1111/ele.12082

Kleijn, D., Bommarco, R., Fijen, T. P. M., Garibaldi, L. A., Potts, S. G., & Putten, W. H. van

der. (2019). Ecological intensification: Bridging the gap between science and practice.

Trends in Ecology & Evolution, *34*(2), 154–166. DOI: 10.1016/j.tree.2018.11.002

Kleijn, D., Rundlöf, M., Scheper, J., Smith, H. G., & Tscharntke, T. (2011). Does conservation

on farmland contribute to halting the biodiversity decline? *Trends in Ecology &*

Evolution, *26*(9), 474–481. DOI: 10.1016/j.tree.2011.05.009

Kremen, C., Williams, N. M., Aizen, M. A., Gemmill-Herren, B., LeBuhn, G., Packer, L., …

Ricketts, T. H. (2007). Pollination and other ecosystem services produced by mobile

organisms: A conceptual framework for the effects of land-use change. *Ecology Letters*,

10(4), 299–314. DOI: 10.1111/j.1461-0248.2007.01018.x

Lefcheck, J. S., Byrnes, J. E. K., Isbell, F., Gamfeldt, L., Griffin, J. N., Eisenhauer, N., … Duffy,

J. E. (2015). Biodiversity enhances ecosystem multifunctionality across trophic levels

and habitats. *Nature Communications*, *6*(1), 1–7. DOI: 10.1038/ncomms7936

Leong, J. M., & Thorp, R. W. (1999). Colour-coded sampling: The pan trap colour preferences

of oligolectic and nonoligolectic bees associated with a vernal pool plant. *Ecological*

Entomology, *24*(3), 329–335. DOI: https://doi.org/10.1046/j.1365-2311.1999.00196.x

Lichtenberg, E. M., Kennedy, C. M., Kremen, C., Batáry, P., Berendse, F., Bommarco, R., …

Crowder, D. W. (2017). A global synthesis of the effects of diversified farming systems

on arthropod diversity within fields and across agricultural landscapes. *Global Change*

Biology, *23*(11), 4946–4957. DOI: 10.1111/gcb.13714

Loxdale, H. D., & Lushai, G. (1999). Slaves of the environment: The movement of herbivorous

insects in relation to their ecology and genotype. *Philosophical Transactions of the Royal*

Society of London. Series B: Biological Sciences, *354*(1388), 1479–1495. DOI:

10.1098/rstb.1999.0492

Lundin, O. (2019). No-till protects spring oilseed rape (*Brassica napus* L.) against crop damage

by flea beetles (*Phyllotreta* spp.). *Agriculture, Ecosystems & Environment*, *278*, 1–5.

DOI: 10.1016/j.agee.2019.03.014

Marja, R., Tscharntke, T., & Batáry, P. (2022). Increasing landscape complexity enhances

species richness of farmland arthropods, agri-environment schemes also abundance – A

meta-analysis. *Agriculture, Ecosystems & Environment*, *326*, 107822. DOI:

10.1016/j.agee.2021.107822

Michener, C. D. (2000). *The Bees of the World*. Baltimore, MD: Johns Hopkins University Press.

Mitchell, M. G. E., Suarez-Castro, A. F., Martinez-Harms, M., Maron, M., McAlpine, C.,

Gaston, K. J., … Rhodes, J. R. (2015). Reframing landscape fragmentation's effects on

ecosystem services. *Trends in Ecology & Evolution*, *30*(4), 190–198. DOI:

10.1016/j.tree.2015.01.011

Morandin, L. A., & Winston, M. L. (2005). Wild bee abundance and seed production in

conventional, organic, and genetically modified canola. *Ecological Applications*, *15*(3),

871–881. DOI: 10.1890/03-5271

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner, H.

(2019). *vegan: Community Ecology Package*. R package version 2.6-4, https://CRAN.R-

project.org/package=vegan

Ouvrard, P., & Jacquemart, A.-L. (2019). Review of methods to investigate pollinator

dependency in oilseed rape (*Brassica napus*). *Field Crops Research*, *231*, 18–29. DOI:

10.1016/j.fcr.2018.11.006

Painter, K., Hinman, H., & Roe, D. (2006). *Economics of spring canola production in dryland*

eastern Washington (No. EB2009E; p. 22). Pullman, WA: Washington State University.

Perrot, T., Gaba, S., Roncoroni, M., Gautier, J.-L., & Bretagnolle, V. (2018). Bees increase

oilseed rape yield under real field conditions. *Agriculture, Ecosystems & Environment*,

266, 39–48. DOI: 10.1016/j.agee.2018.07.020

PRISM Climate Group. (2020). Oregon State University. http://prism.oregonstate.edu

Reddy, G. V. P. (2017). *Integrated Management of Insect Pests on Canola and Other Brassica*

Oilseed Crops. Boston, MA: CABI.

Ricketts, T. H., Watson, K. B., Koh, I., Ellis, A. M., Nicholson, C. C., Posner, S., … Sonter, L. J.

(2016). Disaggregating the evidence linking biodiversity and ecosystem services. *Nature*

Communications, *7*(1), 1–8. DOI: 10.1038/ncomms13106

Robinson, S. V. J. (2019). *Central-place foraging, crop yield, and population change in bees: A*

study in canola agroecosystems (Doctoral thesis, University of Calgary). University of

Calgary, Calgary, AB

Rowen, E. K., Regan, K. H., Barbercheck, M. E., & Tooker, J. F. (2020). Is tillage beneficial or

detrimental for insect and slug management? A meta-analysis. *Agriculture, Ecosystems &*

Environment, *294*, 106849. DOI: 10.1016/j.agee.2020.106849

Rusch, A., Chaplin-Kramer, R., Gardiner, M. M., Hawro, V., Holland, J., Landis, D., …

Bommarco, R. (2016). Agricultural landscape simplification reduces natural pest control:

A quantitative synthesis. *Agriculture, Ecosystems & Environment*, *221*, 198–204. DOI:

10.1016/j.agee.2016.01.039

Schellhorn, N. A., Bianchi, F. J. J. A., & Hsu, C. L. (2014). Movement of entomophagous

arthropods in agricultural landscapes: Links to pest suppression. *Annual Review of*

Entomology, *49*, 559–581. DOI: 10.1146/annurev-ento-011613-161952

Scheper, J., Holzschuh, A., Kuussaari, M., Potts, S. G., Rundlöf, M., Smith, H. G., & Kleijn, D.

(2013). Environmental factors driving the effectiveness of European agri-environmental

measures in mitigating pollinator loss – a meta-analysis. *Ecology Letters*, *16*(7), 912–920.

DOI: 10.1111/ele.12128

Schneider, M. K., Lüscher, G., Jeanneret, P., Arndorfer, M., Ammari, Y., Bailey, D., … Herzog,

F. (2014). Gains to species diversity in organically farmed fields are not propagated at the

farm level. *Nature Communications*, *5*, 4151. DOI: 10.1038/ncomms5151

Sheffield, C. S., Pindar, A., Packer, L., & Kevan, P. G. (2013). The potential of cleptoparasitic

bees as indicator taxa for assessing bee communities. *Apidologie*, *44*(5), 501–510. DOI:

10.1007/s13592-013-0200-2

Skellern, M. P., Welham, S. J., Watts, N. P., & Cook, S. M. (2017). Meteorological and

landscape influences on pollen beetle immigration into oilseed rape crops. *Agriculture,*

Ecosystems & Environment, *241*, 150–159. DOI: 10.1016/j.agee.2017.03.008

Smith, B., & Wilson, J. B. (1996). A consumer's guide to evenness indices. *Oikos*, *76*(1), 70–82.

DOI: 10.2307/3545749

Smith, O. M., Cohen, A. L., Reganold, J. P., Jones, M. S., Orpet, R. J., Taylor, J. M., …

Crowder, D. W. (2020). Landscape context affects the sustainability of organic farming

systems. *Proceedings of the National Academy of Sciences*, *117*(6), 2870–2878. DOI:

10.1073/pnas.1906909117

Stehr, F. W. (1987a). *Immature Insects* (Vol. 1). Dubuque, IA: Kendall/Hunt Publishing

Company

Stehr, F. W. (1987b). *Immature Insects* (Vol. 2). Dubuque, IA: Kendall/Hunt Publishing

Company

Stinner, B. R., & House, G. J. (1990). Arthropods and other invertebrates in conservation-tillage

agriculture. *Annual Review of Entomology*, *35*(1), 299–318. DOI:

10.1146/annurev.en.35.010190.001503

Tamburini, G., Bommarco, R., Kleijn, D., van der Putten, W. H., & Marini, L. (2019).

Pollination contribution to crop yield is often context-dependent: A review of

experimental evidence. *Agriculture, Ecosystems & Environment*, *280*, 16–23. DOI:

10.1016/j.agee.2019.04.022

Tamburini, G., Bommarco, R., Wanger, T. C., Kremen, C., Heijden, M. G. A. van der, Liebman,

M., & Hallin, S. (2020). Agricultural diversification promotes multiple ecosystem

services without compromising yield. *Science Advances*, *6*(45), eaba1715. DOI:

10.1126/sciadv.aba1715

Tooker, J. F., O'Neal, M. E., & Rodriguez-Saona, C. (2020). Balancing disturbance and

conservation in agroecosystems to improve biological control. *Annual Review of*

Entomology, *65*(1), 81–100. DOI: 10.1146/annurev-ento-011019-025143

Tscharntke, T., Tylianakis, J. M., Rand, T. A., Didham, R. K., Fahrig, L., Batáry, P., …

Westphal, C. (2012). Landscape moderation of biodiversity patterns and processes—

Eight hypotheses. *Biological Reviews*, *87*(3), 661–685. DOI: 10.1111/j.1469-

185X.2011.00216.x

Ullmann, K. S., Meisner, M. H., & Williams, N. M. (2016). Impact of tillage on the crop

pollinating, ground-nesting bee, *Peponapis pruinosa* in California. *Agriculture,*

Ecosystems & Environment, *232*, 240–246. DOI: 10.1016/j.agee.2016.08.002

USDA NASS. (2020). *Crop production historical track records* (p. 241). Washington, DC:

USDA, NASS

USDA SARS. (2014a). *2013 Idaho Cropland Data Layer*. Washington, DC: USDA, NASS

USDA SARS. (2014b). *2013 Washington Cropland Data Layer*. Washington, DC: USDA,

NASS

USDA SARS. (2015a). *2014 Idaho Cropland Data Layer*. Washington, DC: USDA, NASS

USDA SARS. (2015b). *2014 Washington Cropland Data Layer*. Washington, DC: USDA,

NASS

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.). NY:

Springer.

Weekers, T., Marshall, L., Leclercq, N., Wood, T. J., Cejas, D., Drepper, B., … Vereecken, N. J.

(2022). Ecological, environmental, and management data indicate apple production is

driven by wild bee diversity and management practices. *Ecological Indicators*, *139*,

108880. DOI: 10.1016/j.ecolind.2022.108880

Winfree, R., W. Fox, J., Williams, N. M., Reilly, J. R., & Cariveau, D. P. (2015). Abundance of

common species, not species richness, drives delivery of a real-world ecosystem service.

Ecology Letters, *18*(7), 626–635. DOI: 10.1111/ele.12424

Figure 1: Box and whisker plots of (a) pollinator richness (total # taxa sampled at each site

across all sampling methods) and (b) kleptoparasite abundance (total # individuals sampled at

each site across all sampling methods) as a function of tillage intensity. "Int." is intermediate

tillage. Boxes span the 25th to 75th percentiles, whiskers extend 1.5 times the interquartile range

beyond the boxes, and there is a line at the median. Lines with asterisks indicate groups that are

statistically different from each other at $\alpha = 0.05$.

Figure 2: (a) Herbivore and (b) predator abundance (total # individuals sampled at each site

across all sampling methods) increase with landscape-scale habitat availability (proportion of

semi-natural habitat in a 1 km radius around a site). Curves show best-fit lines from negative

binomial regressions.

Figure 3: Canola yield (a) was lowest with no tillage and (b) decreased as herbivore richness

increased. "Int." is intermediate tillage. Boxes span the 25th to 75th percentiles, whiskers extend

1.5 times the interquartile range beyond the boxes, and there is a line at the median. Lines with

asterisks (a) indicate groups that are statistically different from each other at $\alpha = 0.05$. The curve

(b) shows the best-fit line from linear regression.