Active populations and growth of soil microorganisms are framed by mean annual

precipitation in three California annual grasslands

Megan M. Foley[1,2], Steven J. Blazewicz[3], Karis J. McFarlane[3], Alex Greenlon[4], Michaela

Hayer[1,2], Jeffrey A. Kimbrel[3], Benjamin J. Koch[1,2], Victoria Monsaint-Queeney[1,2], Keith

Morrison[3], Ember Morrissey[5], Bruce A. Hungate[1,2], Jennifer Pett-Ridge[3,6*]

[1]Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff AZ, USA

[2]Department of Biological Sciences, Northern Arizona University, Flagstaff AZ, USA

[3]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

CA, USA

[4]Department of Environmental Science, Policy, and Management, University of California,

Berkeley CA, USA

[5]Division of Plant and Soil Sciences, West Virginia University, Morgantown, West Virginia,

USA

[6]Life & Environmental Sciences Department, University of California Merced, Merced,

California, USA

*Corresponding author:

Jennifer Pett-Ridge. Physical and Life Sciences Directorate, Lawrence Livermore National

Laboratory, Livermore, CA, USA. Email address: pettridge2@llnl.gov.

Keywords

taxon-specific microbial growth, soil water, environmental filtering, legacy effect, radiocarbon,

soil carbon persistence

Abstract

Earth system models project altered precipitation regimes across much of the globe; in California, the winter wet season is predicted to extend into spring, and the summer dry period to lengthen. How these precipitation trends will affect microbial traits and soil carbon (C) cycling is a key knowledge gap. Specifically, we do not have a mechanistic understanding of the linkages between soil moisture legacy effects, microbial population dynamics and soil C persistence. Using quantitative stable isotope probing (qSIP), we compared total and growing soil microbial communities across three California annual grasslands that span a rainfall gradient yet developed on similar parent material. We also assessed multiple edaphic variables, including the radiocarbon (^{14}C) age of soil C, and found soil C turnover time increased with annual precipitation, but that soil microbes respired recently-fixed C regardless of site rainfall history. Samples were assayed in the wet season, when we expected environmental conditions would be most similar across sites. We hypothesized that growing communities would be more compositionally similar across the gradient than the total background microbiome. We also predicted that the long-term legacy effect of soil water limitation would be reflected in a lower community growth capacity at the driest site. We found that the proportion of the total community that was detected as growing was 28%, 48% and 58% at the wet, intermediate and dry sites, respectively. The composition of growing communities strongly resembled that of total communities, and growing communities were no more similar across the gradient than total communities, indicating a strong effect of climate of the structure of growing microbial communities. Members of three phyla, Acidobacteria, Actinobacteria, and Proteobacteria, were responsible for ~79% of the cumulative ^{18}O assimilation and 80% of all taxa that we defined as 'growers'. Bacterial growth rates were low at the driest site relative to the intermediate and

wettest sites. Reduced growth at the driest site was observed across major phyla, including the

Actinobacteria, Acidobacteria, Bacteroidetes, Gemmatimonadetes and Proteobacteria. Microbial

communities at the driest site displayed phylogenetic clustering, suggesting that climate history

impacts microbial growth through environmental filtering for slow growing taxa. Taxonomic

identity was a strong predictor of growth, such that the growth rates of a taxon at one site

predicted its growth rates at the others. This cross-site coherence in growth is likely a

consequence of genetically determined physiological traits, and is consistent with the idea that

evolutionary history influences growth rate.

1. Introduction

Rainfall patterns and soil water content are crucial controllers of microbial population dynamics (growth and death) and Earth system models project major changes in the timing and intensity of precipitation events globally (Dore, 2005; Schimel, 2018). Since microorganisms mediate a wide range of ecosystem processes, including the accrual and persistence of soil carbon, understanding how the climate history of a site impacts microbial communities is essential. Precipitation is a key controller of ecosystems with Mediterranean climates (cool, wet winters and hot, dry summers) and it plays an equally important role in shaping microbial community structure and function (Angel et al., 2010; Maestre et al., 2015). Microorganisms in Mediterranean ecosystems must withstand both direct physiological stress during prolonged periods of low soil moisture yet be able to compete for resources when seasonal rains return and plant growth resumes (Barnard et al., 2020).

Fluctuations between wet and dry seasons in Mediterranean climates can drive dramatic changes in soil microbial community structure, but most diversity studies rely on DNA sequencing which cannot distinguish genetic material from growing, dormant, and recently dead organisms (Cruz-Martínez et al., 2009; Barnard et al., 2015; Carini et al., 2017). These fractions of the microbial community –growing, dormant, and dead – contribute to soil processes differently (Sokol et al., 2022). Populations that are actively turning over (growing and dying) have a prominent role in biogeochemical processes since their growth requires uptake and transformation of environmental substrates, while their death deposits cellular materials into the soil-mineral matrix where it may be subsequently recycled into living biomass or persist as soil organic matter (Liang et al., 2017). It widely accepted that microbial growth is a major factor in the formation of soil carbon (Bradford et al., 2013). The factors controlling which microbial taxa

grow in soils and the timing of their growth are not well understood, but active communities

have a heightened sensitivity to changes in environmental conditions compared to the total

community (Angel et al., 2013; Baubin et al., 2019). In ecosystems with Mediterranean climates,

strong seasonal changes in soil water may act as an important determinant of which taxa grow

and when.

Soil water also strongly influences microbial function. Historic exposure to low soil

water potential can have a lasting effect on microbial respiration, enzyme activity, and carbon

use efficiency (Averill et al., 2016; Nijs et al., 2019; Hawkes et al., 2020), and is likely to

influence growth as well. Repeated exposure to limited soil moisture may favor traits that confer

tolerance to water stress and select against traits that enhance organisms' ability to grow quickly

or efficiently (Evans and Wallenstein, 2014; Malik et al., 2020a, 2020b). For example, the

production of extracellular polymeric substances (EPS) can be a substantial carbon sink for

bacteria growing under water stress, and osmolyte regulation in response to water stress can

consume both nutrients and energy (Flemming and Wingender, 2001; Sandhya and Ali, 2015). In

bacterial isolates, moisture stress has been shown to drive a tradeoff between EPS production

and the length of lag phase, pointing to the relevance of growth rate in water stress acclimation

(Lennon et al., 2012). While such traits provide a clear advantage during chronic water stress,

there are likely biochemical costs involved in maintaining them. Thus, microbial taxa from soils

routinely exposed to low soil water potentials may have acquired traits that lead to slow growth

rates even in the absence of immediate water stress. At the community level, the mechanism

underlying changes in growth rates may include a shift in microbial community composition, a

change in the physiology of microbial taxa, or both (Evans and Wallenstein, 2014).

Climatic history can shape microbial function via shifts in composition or changes in

microbial physiology. In this study, we compared total and growing microbial communities and

soil characteristics across three California annual grasslands with Mediterranean-type climates

that span a rainfall gradient and developed upon similar parent materials. First, we characterized

the soils to understand how diverging precipitation regimes have shaped the physiochemical

environment that soil microbes experience within each ecosystem. We also analyzed the soils'

radiocarbon (^{14}C) age and ^{14}C of CO_2 to investigate how soil C persistence varies along the

gradient. We then compared the structure of communities inferred from 16S rRNA sequencing

(the "total community") against the communities of growing microorganisms identified with

quantitative stable isotope probing (qSIP) (the "growing community") at each site during the wet

season, when soil water was not limiting (Hungate et al., 2015; Koch et al., 2018). Hypothesizing

that growing community structure is more strongly influenced by recent environmental

conditions than climate history, we expected that similarity in soil water content during the

weeks preceding sampling would result in growing communities that were compositionally more

similar to each other than total communities. Lastly, we quantitatively analyzed patterns of

microbial population growth using qSIP. We hypothesized that growth would be slowest in the

driest site even during the wet season, reflecting a legacy effect of low soil moisture on microbial

growth. We predicted that shifts in composition, likely due to habitat filtering for slow-growing

drought tolerant taxa, would be evidenced by phylogenetic clustering in community composition

(detected via net relatedness index and nearest taxon index).

2. Materials and Methods

2.1 Site description and sample collection

We characterized soil chemistry and bacterial and archaeal microbial communities in

three coastal California annual grasslands with Mediterranean-type climates. The sites span a

significant rainfall gradient (388 mm yr^{-1} to 2833 mm yr^{-1}), yet have similar plant communities

and developed on coastal terraces with similar parent material (Table 1). The wettest site, Angelo

Coast Range Reserve, lies upstream of the headwaters of the South Fork Eel River where

indigenous peoples including the Cahto, Pomo, Wailaki, Yuki, Weott, and Sinkyone historically

held territory (Power et al., 2015). Annual grasslands at Angelo intersperse mixed-oak

woodlands and old-growth conifer forests that occur on strath terraces which formed as the Eel

River cut through the soft coastal sediment (Seidl et al., 1992; Sullivan et al., 2016). The

intermediate site, Hopland Research and Extension Center, sits in the foothills of the Mayacamas

Mountains and encompasses topographically rugged rangeland on territory historically occupied

by the Pomo Nation. The driest site, Sedgwick Reserve, spans the Santa Ynez Valley and San

Rafael Mountains and is located on land where the Chumash people have historically lived.

The three sites developed on similarly-aged sedimentary rock, primarily sandstone and

shale from the Franciscan formation (Table 1). The parent material at Sedgwick also consists of

alluvium from the Paso Robles formation which contains Franciscan mélange and Monterey

shale (Homyak et al., 2018). Angelo soils are Ultic Haploxeralfs of the Hologan-Hollowtree-

Casabonne complex (Berhe et al., 2012), Hopland soils are Typic Haploxeralfs of the Witherall-

Squawrock complex (Fossum et al., 2021), and Sedgwick soils are Pachic Argixerolls belonging

to the Botella series (Homyak et al., 2018). These taxonomies depict three naturally fertile soils

(Angelo and Hopland are Alfisols while Sedgwick is a Mollisol) that experience a moderately

dry, Mediterranean-type climate (Angelo and Hopland are Xeralfs while Sedgwick is Xeroll).

Naturalized non-native annual grasses dominate vegetation at the three sites. *Avena spp.* and

Bromus spp. dominate at Hopland and Sedgwick (Michael Barbour et al., 2007; Leitner et al.,

2017). Vegetation at Angelo is dominated by non-native annual grasses including *Aira*

caryophyllea, Bromus hordeaceuous, Briza minor and *Avena spp.* and by annual and perennial

forbs interspersed with the native perennial grass *Danthonia californica* (Kotanen, 2004; Berhe

et al., 2012; Sullivan et al., 2016) (Table 1).

We collected soils in February-March 2018. At each site, surface vegetation was removed

and three replicate soil cores (10 cm x 5 cm) were collected 1 m apart for chemical

characterization, radiocarbon analyses, and quantitative stable isotope probing. Soil cores were

transported on ice to Lawrence Livermore National Laboratory where they were homogenized by

hand, stones and large roots were removed, and portions were dried at room temperature, stored

at 4 °C, or frozen at -80 °C for different downstream analyses. All analyses reported here were

conducted on surface soils (0-10 cm). We measured gravimetric soil water content by drying

soils to a constant weight at 105°C. We retrieved daily mean air temperature and total

precipitation data from nearby climate stations and on-site records for the calendar year and

thirty-day period preceding sample collection at each site.

2.2 Soil physical and chemical characterization

Soil chemical and physical characterization was conducted at the Oregon State

University Soil Health Laboratory. Bulk C and nitrogen (N) concentrations were quantified using

an Elementar vario MACRO cube elemental analyzer through dry combustion, separation of

gaseous species, and thermal conductivity detection. Trace elements were quantified on an

Agilent 5110 ICP-OES after soils were extracted with 0.002M barium chloride. Soil texture was

determined by the hydrometer method. To measure pH, 4 g of soil was suspended in 8 mL of

0.01 M CaCl$_2$ and equilibrated for 1 hour prior to analysis (Minasny et al., 2011). Soil pH was

measured using a Mettler Toledo Seven Compact pH/Ion meter. Additional soil cores (0-5 cm)

were collected in June of 2019 to construct soil water retention curves for the three sites. Soil

cores were also collected from the top 8-15 cm of soil to calculate bulk density for the retention

curves. Intact soil cores were prepared and analyzed using Hyprop and WP4C instruments in

accordance with manufacturer's instructions (Pertassek et al., 2011; Meter Group, 2019).

2.3 Quantitative x-ray diffraction

Quantitative x-ray diffraction (qXRD) was performed at Lawrence Livermore National

Laboratory as previously described (Fossum et al., 2021). Samples for qXRD were crushed and

passed through a 500 µm sieve. 3 g of soil was then ground for 5 minutes with 15 mL of

methanol using a MCrone mill. The ground samples were transferred to a plastic tray, air dried,

and homogenized for 3 min using a vortex mixer with 10 mm plastic beads (Bakker et al., 2018).

The soil powders were side loaded into XRD sample mounts. All powdered samples were

analyzed on a Bruker D8 advance XRD. The samples were scanned from 3° to 65° 2θ at 0.011°

steps with a 5 second per step count time. Quantitative mineralogy was determined using the

BGMN Rietveld refinement and the Profex interface software (Doebelin and Kleeberg, 2015).

XRD patterns were refined to fit crystal unit cell parameters, size, site occupancy, and preferred

orientation.

2.4 Natural abundance stable isotopes and radiocarbon analysis of bulk soils and incubations

For each core, subsamples of dried homogenized soil were analyzed for δ^{13}C and δ^{15}N at

the Center for Stable Isotope Biogeochemistry at the University of California, Berkeley

(IsoPrime100 mass spectrometer) or prepared for bulk soil radiocarbon measurement by sealed-

tube combustion to CO_2 in the presence of CuO and Ag. A subsample of fresh soil from each

core was incubated in the lab to measure $\Delta^{14}CO_2$ of microbially respired carbon. After removing

roots from soils by hand picking, soils were left undisturbed for one week at room temperature to

allow any remaining roots to senesce. For each core, approximately 150 g dry mass-equivalent

soil was weighed into 150 mL beakers and placed inside quart-sized mason jars with stopcock-

fitted lids. Soils were pre-incubated for one day before incubation jars were sealed and

headspaces flushed with CO_2-free air. Soils were then incubated at room temperature until

enough CO_2 had accumulated for ^{14}C measurement (9-11 days). At the end of the incubation

period, headspace CO_2 was cryogenically purified for radiocarbon analysis. Aliquots of purified

CO_2 were analyzed for $\delta^{13}C$ at the UC Davis Stable Isotope Laboratory (GVI Optima Stable

Isotope Ratio Mass Spectrometer).

CO_2 from both combusted bulk soil and incubation jars was purified cryogenically at the

Center for Accelerator Mass Spectrometry (CAMS) at Lawrence Livermore National Laboratory

using a vacuum line before being reduced to graphite on iron powder in the presence of H_2

(Vogel et al., 1984). Radiocarbon abundance in graphitized samples was determined on the FN

Van de Graaff Accelerator Mass Spectrometer at Lawrence Livermore National Laboratory's

Center for Accelerator Mass Spectrometer. Radiocarbon values are reported in $\Delta^{14}C$ notation

corrected for mass-dependent fractionation using measured $\delta^{13}C$ values (Stuiver and Polach,

1977, p. 14).

2.5 ^{18}O-water quantitative stable isotope probing (qSIP)

For qSIP incubations, 5 g soil from each field replicate was transferred to 15 ml Nalgene

flatbottom vials. Soils were lightly air dried in a laminar flow hood at room temperature for 24

hours prior to isotope addition. This drying is necessary to allow for sufficient enrichment of the

soil water with the isotope label while also maintaining soil moisture reasonably similar to field

moisture during the qSIP incubations. One milliliter of isotopically enriched water (98.15 at%

^{18}O-H_2O) or natural abundance water (as a control) was pipetted onto the soil slowly and evenly

and then gently mixed with the pipette tip (raising soil moisture content by 3-7%). After the

water addition, vials were immediately sealed inside 500 ml mason jars and incubated at room

temperature in the dark for 8 days. At the end of the incubation, soils were frozen in liquid

nitrogen and then stored at -80°C.

DNA was extracted from all soil samples using a modified protocol adapted from

Barnard et al. (2015). Three replicate extractions were conducted for each sample and then

replicate DNA extracts were combined. For each extraction, soil (0.4 g +/- 0.001 g soil) was

added to 2 ml Lysing Matrix E tube (MP Biomedicals) and extracted twice as follows. 500 µl

extraction buffer (5% CTAB, 0.5 M NaCl, 240 mM K_2HPO_4, pH 8.0) and 500 µl 25:24:1

phenol:chloroform:isoamyl alcohol were added before shaking (FastPrep24, MP Biomedicals: 30

s, 5.5 m $^{s-1}$). After centrifugation (16,100 x g, 5 min), residual phenol was removed using pre-

spun 2 ml Phase Lock Gel tubes (5 Prime, Gaithersburg, MD, USA) with an equal volume of

24:1 chloroform:isoamyl alcohol, mixed and centrifuged (16,100 x g, 2 min). The aqueous

phases from both extractions were pooled, mixed with 7 µl RNAase (10 mg/ml), mixed by

inverting, and incubated at 50 °C for 10 min. 335 µL 7.5 M NH4+ acetate was added, mixed by

inverting, incubated (4 °C, 1 h). and centrifuged (16,100 x g, 15 min. The supernatant was

transferred to a new 1.7 ml tube and 1 µl Glycoblue (15 mg/ml) and 1 ml 40% PEG 6000 in 1.6

M NaCl was added, mixed by vortex, and incubated at room temperature in the dark (2 h). After

centrifugation (16,100 x g, 20 min), the pellet was rinsed with 1 ml ice-cold 70% ethanol, air-

dried, resuspended in 30 µl 1xTE and stored at -80 °C.

Samples were subjected to a cesium chloride density gradient formed by physical density

separation via ultracentrifuge as previously described with minor modifications (Blazewicz et al.,

2014; Sieradzki et al., 2020). For each sample, 5 µg of DNA in 150 µL 1xTE was mixed with

1.00 mL gradient buffer, and 4.60 mL CsCl stock (1.885 g mL^{-1}) with a final average density of

1.730 g mL^{-1}. Samples were loaded into 5.2 mL ultracentrifuge tubes and spun at 20 ℃ for 108

hours at 176,284 RCF$_{avg}$ in a Beckman Coulter Optima XE-90 ultracentrifuge using a VTi65.2

rotor. Automated sample fractionation was performed using Lawrence Livermore National

Laboratory's high-throughput SIP ('HT-SIP') pipeline, which automates fractionation and clean-

up tasks for the density gradient SIP protocol. Ultracentrifuge tube contents were fractionated

into 36 fractions (~200 µL each) using an Agilent Technologies 1260 isocratic pump delivering

water at 0.25 mL min^{-1} through a 25G needle inserted through the top of the ultracentrifuge tube.

Each tube was mounted in a Beckman Coulter fraction recovery system with a side port needle

inserted through the bottom. The side port needle was routed to an Agilent 1260 Infinity fraction

collector. Fractions were collected in 96-well deep well plates. The density of each fraction was

then measured using a Reichart AR200 digital refractometer fitted with a prism covering to

facilitate measurement from 5 µL, as previously described (Buckley et al., 2007). DNA in each

fraction was purified and concentrated using a Hamilton Microlab Star liquid handling system

programmed to automate previously described glycogen/PEG precipitations (Neufeld et al.,

2007). Washed DNA pellets were suspended in 40 µL of 1xTE and the DNA concentration of

each fraction was quantified using a PicoGreen fluorescence assay. The fractions for each sample

were binned into 9 groups based on density (1.6900-1.7099 g/ml, 1.7100-1.7149 g/ml, 1.7150-

1.7199 g/ml, 1.7200-1.7249 g/ml, 1.7250-1.7299 g/ml, 1.7300-1.7349 g/ml, 1.7350-1.7399 g/ml,

1.7400-1.7468 g/ml, 1.7469-1.7720 g/ml), and fractions within a binned group were combined

and sequenced.

For 16S rRNA gene amplicon sequencing, non-fractionated DNA as well as density

fractionated DNA was amplified in triplicate 10-uL reactions using primers 515 F and 806 R

(Apprill et al., 2015; Parada et al., 2016). Each reaction contained 1uL sample and 9uL of

Phusion Hot Start II High Fidelity master mix (Thermo Fisher Scientific) including 1.5mM

additional $MgCl_2$. PCR conditions were 95 C for 2 min followed by 20 cycles of 95 C for 30 S,

64.5 C for 30 S, and 72 C for 1 min. The triplicate PCR products were then pooled and diluted

10X and used as a template in a subsequent dual indexing reaction that used the same primers

including the Illumina flowcell adaptor sequences and 8-nucleotide Golay barcodes (15 cycles

identical to initial amplification conditions). Resulting amplicons were purified with AMPure XP

magnetic beads (Beckman Coulter) and quantified with a PicoGreen assay on a BioTek Synergy

HT plate reader. Samples were pooled at equivalent concentrations, purified with the AMPure

XP beads, and quantified using the KAPA Sybr Fast qPCR kit (Kapa Biosciences). A total of 9

unfractionated, and 162 fraction libraries were sequenced on an Illumina MiSeq instrument at

Northern Arizona University's Genetics Core Facility using a 300-cycle v2 reagent kit. Sequence

data and sample metadata have been deposited in the NCBI Sequence Read Archive under the

accession number PRJNA718849.

Paired-end 151 nt reads were filtered to remove phiX and other contaminants with bbduk

v38.56 (default settings except k=31 and hdist=1) (Bushnell, 2014). Fastq files were then

filtered/trimmed for quality (maxEE=5, truncQ=2) and used to generate amplicon sequence

variants (ASVs) with DADA2 v1.10 and phyloseq v1.26 (McMurdie and Holmes, 2013;

Callahan et al., 2016). Chimeric sequences were determined and removed using

removeBimeraDenovo from DADA2. ASV taxonomy was determined using the RDP 16S rRNA

gene database (training set 16) using RDP classifier v2.11, keeping classifications with greater

than 50% confidence (Wang et al., 2007). A phylogenetic tree was built using Muscle v3.8.31

and FastTree v2.1.10 (Edgar, 2004; Price et al., 2010).

2.6 Quantitative stable isotope probing analysis

Quantitative stable isotope probing (qSIP) measures the change in buoyant density of a

taxon's DNA, or segment of DNA, due to assimilation of a substrate-derived heavy isotope. qSIP

with ^{18}O-H$_2$O identifies actively growing microbial taxa since ^{18}O is incorporated into the

phosphodiester backbone of DNA during replication through the rapid exchange of oxygen

between water and inorganic phosphate species (Schwartz, 2007). The amount of ^{18}O

incorporation into microbial DNA during an incubation with ^{18}O-H$_2$O thus approximates a

microorganisms growth rate.

We quantified excess atom fraction (EAF) ^{18}O of bacterial DNA following a modified

version of the procedure described by Hungate et al. (2015) and Koch et al. (2018) using average

DNA concentration to normalize the relative abundance of taxa within each density fraction

(Papp et al., 2018a; Sieradzki et al., 2020) . We calculated median values and 95% confidence

intervals for EAF ^{18}O by bootstrapping (n=1000) across experimental replicates. qSIP analysis

was limited to taxa that occurred in at least 2 (of 3) experimental replicates and in at least 2 (of 9)

density fractions. These criteria were chosen to reduce the likelihood of falsely interpreting

spurious density shifts as growth. Technical error associated with tube-level differences in CsCl

density gradients was corrected as previously described (Morrissey et al., 2017). All qSIP

calculations are available at https://github.com/mmf289.

When comparing values of ^{18}O enrichment between sites, we chose to use the metric

"fraction of maximum potential enrichment" or "(FME) ^{18}O" to account for slight differences in

the enrichment of the soil water during incubation. FME ^{18}O was computed for each taxon as

FME ^{18}O = ^{18}O-EAF$_{taxon}$/average soil water ^{18}O enrichment during eight-day incubation. For all

analyses, 'growing' microbial taxa were identified as ASVs having 95% confidence intervals for

EAF ^{18}O exceeding zero. We refer to the community of growing microbial taxa at each site as

the "growing microbial community". We refer to the microbial community inferred from 16S

rRNA marker gene sequencing of non-fractionated DNA at each site as the "total microbial

community".

2.7 Statistical analyses

All statistical analyses were conducted in R (R Core Team, 2021). Soil properties,

relative abundances of minerals, and bulk and respired soil radiocarbon values were compared

between sites using one-way ANOVA and Fisher's LSD test. A Bonferroni correction was

applied when multiple comparisons were made.

The relative abundances of taxa in total communities were measured from sequencing of

unfractionated DNA samples. The relative abundances of taxa in growing communities were

computed by removing counts from taxa that were not identified as growing via qSIP from

unfractionated DNA samples and then re-computing relative abundance values. We computed

the proportion of the total community that was growing by dividing the number of ASVs

identified as growing by the total number of ASVs inferred from 16S rRNA sequencing at each

site. We then used Bray-Curtis dissimilarity to assess between-site dissimilarity for both growing

and total communities with vegan: Community Ecology Package (Wagner et al. 2019). Principal

coordinate analysis (PCoA) was used to visualize community structure. We tested for differences

in community composition between sites using ANOSIM on ranked Bray-Curtis dissimilarities.

To test if the degree of community dissimilarity between sites differed for total versus growing

microbial communities, we performed a t-test on mean Bray-Curtis dissimilarity for each

pairwise comparison of sites between total and growing communities. A Bonferroni correction

was applied to account for multiple comparisons.

We used one-way ANOVA and Fisher's LSD to test for differences between sites in

community, phylum, and family mean FME ^{18}O and only included taxa that were identified as

growing at each site in this analysis. A Bonferroni correction was applied when multiple

comparisons were made. We also compared FME ^{18}O of taxa that were uniquely growing at

Sedgwick to the FME ^{18}O of taxa that were growing at Sedgwick and at least one other site using

a t-test. We conducted NRI and NTI analyses using the Picante package to determine

phylogenetic relatedness of total microbial communities at each site.

We used linear regression to assess the relationship of taxon-specific FME ^{18}O between

sites and included only individual ASVs that were co-occurring in at least two sites.

Phylogenetic signal analyses were used to test whether the growth (EAF ^{18}O) of related

organisms resembled each other more than would be expected by chance alone[1]. Phylogeny was

built using the SILVA v128 tree using SATé-enabled phylogenetic placement (Janssen et al.,

2018; Bolyen et al., 2019). We measured Blomberg's K and Pagel's λ using phytools (Revell,

2012) for taxon-specific EAF ^{18}O values within each site and included all taxa in this analysis.

3. Results

3.1 Precipitation and air temperature trends prior to sample collection

  We gathered precipitation and air temperature data to assess the environmental conditions at each site during the year and month prior to sample collection. The three sites form a gradient in MAP and MAT with precipitation decreasing and temperature increasing between sites in the order: Angelo, Hopland, and Sedgwick (Table 1). In the calendar year prior to sample collection, the wet season at Angelo and Hopland lasted until April. Precipitation remained scarce or absent during the summer months and returned again by November. In contrast, the wet season at Sedgwick occurred in only January and February and didn't return until the following January (Supplementary Figure 1). In sum, Angelo was the wettest and coldest site and Sedgwick was hottest and driest site during the year preceding sample collection (Supplementary Table 1). During the 30 days leading up to sample collection, total precipitation differed between our sites by an average of only 46 mm and soil moisture of the field-collected soil varied modestly from 17% to 25% (Supplementary Tables 1 & 2). The average air temperature during the 30 days leading up to sample collection ranged from 3.76 °C at Angelo to 10.35 °C at Sedgwick.

3.2 Soil characterization

  To understand how the range in precipitation regimes correlates with the physiochemical environment that soil microbes experience within each ecosystem, we characterized soil mineralogy via quantitative x-ray diffraction (qXRD), constructed soil water retention curves, and determined soil texture, total C and N, trace elements, pH, and effective cation exchange capacity. Soils at the three sites contain the same minerals in varying proportions (Supplementary Table 3). Of the non-clay minerals at the sites, quartz is proportionally most

abundant by weight followed by plagioclase and K-feldspar. At the wet and intermediate sites,

Angelo and Hopland, muscovite is the most abundant clay mineral followed by chlorite and

kaolinite; at the dry site, Sedgwick, kaolinite is most abundant.

Sedgwick, the driest site, contained the highest proportion of clay particles (Table 2). Soil

water retention curves show that Sedgwick soils also experience lower soil water potentials at

higher moisture contents than the intermediate and wet sites (Supplementary Figure 1). Soil

carbon and nitrogen were low at all sites and exhibited little variation. $\delta^{15}N$ of bulk soil did not

vary between sites and $\delta^{13}C$ was slightly lower at Sedgwick than Hopland or Angelo. Soil C:N

exhibited little variation between sites. pH decreased significantly with increasing MAP.

Effective cation exchange capacity and most trace elements were highest in Sedgwick soil, with

the exception of aluminum which was highest in the Angelo soil (Table 2).

3.3 Bulk soil and respired $\Delta^{14}C$

The $\Delta^{14}C$ of bulk surface soils increased along the gradient from the wettest to the dry

site, indicating that the average age of bulk soil C is oldest at Angelo, the site with the highest

annual rainfall (Figure 1). Bulk soil $\Delta^{14}C$ values were above modern at Sedgwick, indicating the

presence of carbon associated with atmospheric weapons testing in the 20[th] century, while the

more depleted $\Delta^{14}C$ values observed for soils from Angelo suggest the presence of older and

more slowly cycling soil C at this site compared to the other sites. $\Delta^{14}C$ at our intermediate

rainfall site, Hopland, falls between these two extremes. For reference, turnover times—

determined using a single-pool, steady-state, homogenous model (Torn et al., 2009)—were

approximately 200, 400, and 700 years for Sedgwick, Hopland, and Angelo, respectively. In

contrast, the $\Delta^{14}C$ of respired CO_2 was not significantly different for the three soils, indicating

that microorganisms uniformly respired recently fixed C (with an average age of 7 years since

fixation from the atmosphere), despite the differences in the average age of bulk soil carbon.

3.4 Structure of growing populations versus the total community

To test the hypothesis that growing communities would be compositionally more similar

to each other than total communities, we analyzed the structure of growing and total

communities using PCoA and pairwise comparisons of Bray-Curtis dissimilarity. The proportion

of the total community that was detected as growing (ASVs having 95% confidence intervals for

EAF ^{18}O exceeding zero were considered growing) was 28%, 48% and 58% at the Angelo,

Hopland and Sedgwick sites, respectively (Supplementary Table 4). Community structure

differed by site for both the total and growing bacterial communities (Figure 2). Growing

communities were not compositionally more similar to each other than total communities

(Supplementary Figure 3). Sedgwick was the most dissimilar from the other sites, as quantified

by mean pairwise Bray-Curtis Dissimilarity (Supplementary Figure 3).

More than 70% of 16S rRNA gene sequences in the total bacterial community at each site

could be attributed to three phyla: Acidobacteria, Actinobacteria (dominant orders

Rubrobacterales, and Solirubrobacterales), and Proteobacteria (dominant orders Burkholderiales,

Sphingomonadales, and Rhizobiales). These phyla dominated growing communities as well,

comprising more than 80% of all bacterial 16S rRNA gene sequences that were growing at each

site and accounting for at least 79% of cumulative ^{18}O assimilation in each site (Supplementary

Figure 4). While these phyla accounted for a large proportion of total growth, there was no

detectable relationship between the relative abundance of an individual taxon and its growth,

meaning some low abundance taxa exhibited fast growth (Supplementary Figure 5).

3.5 Microbial growth along the precipitation gradient

Using qSIP, we analyzed patterns of microbial population growth to test the hypothesis

that growth would be slowest in the historically driest site –even during the wet season—

reflecting the climate history of the site. We used fraction of maximum potential enrichment

(FME) ^{18}O when comparing values of ^{18}O enrichment between sites to account for small

differences in the enrichment of the soil water during incubation. Site specific variation in

microbial growth was present at the community, phylum, and family levels. The mean

community FME ^{18}O was significantly higher in Hopland and Angelo soils than in Sedgwick

soils (Figure 3). These patterns, in general, also persisted across phyla and family levels:

Actinobacteria and Bacteroidetes had a higher FME ^{18}O in Hopland relative to Sedgwick soils,

and Acidobacteria, Gemmatimonadetes, and Proteobacteria had higher FME ^{18}O in both Angelo

and Hopland soils (Figure 3). While the mean FME ^{18}O of Comamonadaceae was highest in

Sedgwick, this result was not statistically significant (P=0.33, ANOVA). No microbial families

exhibited significantly faster growth in Sedgwick soils relative to the other sites (Supplementary

Figure 6).

We calculated net relatedness index (NRI) and nearest taxon index (NTI) to assess

whether reduced microbial growth at the driest site was driven by shifts in community

composition (i.e. through habitat filtering for slow growing taxa). NRI and NTI analyses indicate

phylogenetic clustering of total microbial communities at all three sites (Supplementary Table 6).

We then grouped the taxa growing at Sedgwick according to whether they grew at Sedgwick

alone or in at least one other site. We found that taxa that were unique to Sedgwick were slower

growing than taxa that were growing in at least one other site (Figure 4).

3.6 Evolutionary constraints on microbial growth

To better understand the factors driving variation in growth observed in our study, we used phylogenetic signal analyses to test whether taxon-specific growth was influenced by a microorganism's evolutionary history. In all three grassland sites, we detected significant phylogenetic signals for taxon-specific growth, as determined by Pagel's λ and Blomberg's K (Supplementary Table 7). Stronger phylogenetic signals were observed in Angelo soil ($\lambda = 0.82$, p<0.001, K=0.30, p=0.001) and Hopland ($\lambda = 0.85$, p<0.001, K=0.10, p=0.004) than Sedgwick ($\lambda = 0.68$, p<0.001, K = 0.10, p=0.010). A significant phylogenetic signal indicates that the growth rates of closely related taxa resemble each other more than taxa drawn randomly from the same phylogenetic tree, however the values we observed are somewhat lower than expected under Brownian motion evolution ($\lambda = 1$ and K=1).

Given the evidence for a phylogenetic signal of in-situ microbial growth, we were curious to know if taxa would consistently perform with high or low activity regardless of the site, so we used linear regression to analyze taxon-specific growth rates across sites. For taxa that occurred at more than one site and were also significantly growing, their growth rate at one site was predictive of growth rates in another site (Figure 5). We compared ^{18}O enrichment of bacterial ASVs that were present and growing in at least two sites and found positive, linear correlations of taxon-specific enrichment between sites. The growth rate of a bacterial taxon in one site explained as much as 48% of the variation in that taxon's growth rate in a second site (Supplementary Table 5). The amount of variation explained by each regression was the lowest when comparing the growth of ASVs between Angelo and Sedgwick, the two most distant sites in our gradient, that also have the largest difference in mean annual precipitation.

4. Discussion

In this study, we used quantitative stable isotope probing (qSIP) to assess total and growing soil microbial communities across three California annual grassland ecosystems with Mediterranean climates that span a rainfall gradient. We obtained samples for this study in early 2018 during the wet season when differences in soil moisture between our sites were minimized and assessed multiple edaphic variables to characterize how variation in rainfall influences the physiochemical environment experienced by soil microorganisms at each site. We also measured the radiocarbon (^{14}C) age of the soils to investigate how soil C persistence varies along the gradient and might be related to microbial growth potential. We hypothesized that actively growing communities would be more compositionally similar across the gradient than the total background microbiome due to the similarity in soil water content during the weeks preceding sampling. We also predicted that the long-term legacy effect of soil water limitation would be reflected in lower soil C, and lower growth capacity at the driest site even when soils were moisture replete.

4.1 Soil physiochemical environment

We characterized soils from the three California grasslands to assess how diverging rainfall regimes have shaped the physiochemical environment that microorganisms experience within each ecosystem. The three sites developed upon sedimentary parent material, primarily Franciscan sandstone and shale (Table 1). The three soils contain the same minerals but in varying proportions, pointing to the shared characteristics of parent material yet diverging weathering regimes (Supplementary Table 3). pH decreased with increasing MAP, which is

likely driven by increased leaching of base cations with higher amounts of precipitation

(Voroney and Heck, 2015). Some soil characteristics did not obviously correlate with rainfall

patterns. Effective cation exchange capacity was highest at the driest site, Sedgwick, which is

likely driven by its high clay content, but does not differ between the wet and intermediate sites.

Additionally, soil carbon and nitrogen content, two important drivers of microbial composition

and function, varied very little between the sites (Table 2), suggesting a similar backdrop for

microbial growth, beyond the substantial differences in MAP.

498
4.2 Growing communities resemble the structure of total communities

Microbial growth drives elemental cycling in soils, so understanding the factors

controlling which taxa grow is critical for soil biogeochemistry. In ecosystems with

Mediterranean climates, strong seasonal changes in soil water may be an important determinant

of growing community structure. These patterns have clear effects on instantaneous microbial

growth rates, but may also have more pervasive, long-term legacy effects (Schimel, 2018). In

this study, we characterized the total and growing soil microbial communities in three California

grassland ecosystems that span a rainfall gradient during the wet season when all soils were

moisture replete (Supplementary Table 2). We hypothesized that growing communities would be

compositionally more similar to each other than total communities due to the convergence in

environmental conditions, namely soil moisture and an actively growing plant community,

during the weeks prior to sample collection. In contrast to this expectation, both growing and

total communities were distinct between sites, and the distinctions between sites were of the

same order of magnitude for total and growing communities (Figure 2, Supplementary Figure 3).

Microbial ecologists have long sought a reliable metric of metabolic activity with

taxonomic resolution. For many years, it was assumed that rRNA extracted from environmental

samples could be used to identify metabolically active microbial populations (Jones and Lennon,

2010; Campbell and Kirchman, 2013; Bowsher et al., 2019). Sequencing of rRNA has quantified

strong compositional differences between active and total soil microbial communities at regional

scales, and has also shown that rRNA-based soil microbial communities are more similar over

space than DNA-based communities (Angel et al., 2013; Baubin et al., 2019; Locey et al., 2020).

Our data suggest that a site's climate history, rather than recent environmental conditions, place a

stronger imprint on the structure of growing soil microbial populations. This is another piece of

evidence suggesting that rRNA is an imperfect indicator of metabolically active populations

(Blazewicz et al. 2013). While heavy-water DNA qSIP quantifies growing microbial

populations, the tight coupling between DNA and RNA synthesis in soils suggests that the

majority of active taxa are likely also growing (Papp et al., 2018b).

We characterized microbial communities at a time of the year when soils were replete

with moisture and when environmental conditions should generally be favorable for microbial

growth. However, the growing taxa in this study were still only a subset (28-58%) of all taxa

detected by 16S rRNA sequencing, indicating a substantial presence of dormant organisms or

relic DNA (Supplementary Table 4). While this finding is consistent with our current

understanding of microbial communities in soils (the majority of microorganisms are believed to

be dormant and relic DNA is estimated to comprise on average 40% of prokaryotic and fungal

DNA) (Blagodatskaya and Kuzyakov, 2013; Carini et al., 2017), it raises important questions

about microbial growth in these ecosystems. For example, are non-growing taxa merely the

genetic remnants of dead microbes or are they perhaps dormant taxa poised for growth at another

time of the year? For example, studies in California annual grasslands have documented

microbial succession during the growing season of *Avena spp.*, showing that different microbial

taxa grow at different points throughout the wet season (Shi et al., 2015, 2016; Nuccio et al.,

2020). It is also possible that the average generation time of some microbial taxa may be so slow

that their growth could not be detected during the week-long incubation we conducted. These

questions highlight the need for experiments clarifying the *in situ* limits and controls of growth

in the soil microbiome, information that is fundamental for identifying what maintains the

tremendous microbial diversity and functional potential in soils.

4.3 Persistent Effect of Climate on Microbial Growth

Microbial growth rate approximates a microorganism's contributions to elemental

cycling, so it is essential to understand whether the climate history of a site can influence

microbial growth rates in nature. We hypothesized that a history of repeated exposure to low soil

water potential at the driest site would result in lower microbial growth even when soil water was

not limiting. Consistent with this expectation, community and phylum mean growth was higher

at the wet and intermediate sites, Hopland and Angelo, than at the driest site, Sedgwick (Figure

3). Variation in microbial growth did not appear to be driven by pH, as we would have expected

increasing substrate availability and near-neutral pH levels to, on average, facilitate faster

bacterial growth rates at Sedgwick (Rousk et al., 2009). Microbial growth rates did not appear to

be driven by differences in soil texture either, as clay content (which is highest at Sedgwick) is

thought to support faster growth rates by providing more habitable surface area or pore space for

soil microorganisms (Bååth, 1996; Uhlířová and Šantrůčková, 2003). While precipitation is the

apparent causal factor for the variation in microbial growth rates we observed, we cannot entirely

rule out temperature, rainfall periodicity, or plant productivity as proximal causes.

We suspect the variation in microbial growth in our soils reflects a legacy effect of the

precipitation regime, particularly at our driest site, Sedgwick. Similar findings were documented

for a rainfall gradient in Texas, where microbial growth decreased with lower MAP when

measured under controlled laboratory conditions (Leizeaga et al., 2021). However, the literature

directly testing the legacy effect of low soil moisture on growth rate is inconsistent. While a

number of short-term field and lab experiments have found that historic exposure to drought

reduces microbial growth rate, field experiments operating on longer time scales (>10 years)

have failed to measure the same effect, possibly due to microbial acclimations to drought in the

long-term studies that do not confer reduced growth rates (Meisner et al., 2013; Rousk et al.,

2013; Hicks et al., 2018; Rahman et al., 2018; Nijs et al., 2019).

Historic exposure to low soil water potential can alter microbial function through shifts in

community composition or by directly altering the physiology of taxa (Evans and Wallenstein,

2014). Our findings suggest that variability in microbial growth rate was driven by alterations in

community composition via environmental filtering for slow growing taxa at our driest site. NRI

and NTI indices show that communities at Sedgwick display phylogenetic clustering, which

indicates environmental filtering at this site (Supplementary Table 6). Within the community of

microbes that were growing at the driest site, most (155/197 ASVs) were found growing at this

site alone, and the ASVs that were uniquely growing at Sedgwick had slower growth, on

average, than taxa that occurred and grew in at least one other site (Figure 4). Thus, we infer that

reduced microbial growth at our driest site appears to be influenced predominantly by relatively

slow-growing taxa that are unique to this site.

Since we measured microbial growth at the ASV level using qSIP and 16S rRNA marker

gene sequencing, we lack direct evidence regarding the traits and genomic potential of the

microbial taxa that exhibited faster growth at our driest site. However, under the conditions

evaluated here (wet season, moisture replete soils), the consistent arrangement of growth rates

among taxa from site to site (Figure 5) most likely reflects the influence of genetic constraints on

growth. This finding is consistent with the idea that evolutionary history influences growth rate

and suggests that the phenotype of *in situ* growth rate for a given taxon spans a range constrained

by its genetics and physiology, constraints that persist across rainfall gradients, edaphic

properties, and biological communities (Morrissey et al., 2019). Repeated exposure to low soil

water potential at Sedgwick may favor the success of microbial taxa whose physiological traits

confer resistance to water stress and contribute to slow growth. While not statistically significant

(P=0.33, ANOVA), the family Comamonadaceae exhibited faster in growth Sedgwick relative to

the other sites (Supplementary Figure 6) and members of this family can accumulate

polyhydroxyalkanoates within the cell for C storage and have genes for carotenoid biosynthesis,

EPS synthesis and hydrolysis, and trehalose synthesis (De Luca et al., 2011; dos Santos et al.,

2014). Future studies combining qSIP with metagenomic sequencing may help elucidate the

nature of genetic constraints on microbial growth in these soils.

4.4 Soil C persistence (Δ^{14}C) in the grassland sites

Bulk soil Δ^{14}C values indicate that soil C age increased with MAP (Figure 1), consistent

with findings from a recent global scale synthesis (Heckman et al., 2022). The effect of MAP on

soil C persistence is driven by multiple factors that can vary by soil fraction and depth (Heckman

et al., 2022). Recent models of soil C cycling emphasize physiochemical stabilization

mechanisms and microbial necromass as a precursor for persistent soil C (Lavallee et al., 2020).

In our study, microbial growth, and presumably necromass production, also increased with

increased MAP (Figure 3). If necromass-C is preferentially stabilized in these soils, this could

result in more persistent soil C over time, leading to an older average age for bulk soil C. In

addition, greater rates of vertical flushing of water at our highest MAP site could also facilitate

weathering and the development of secondary clay, aluminum, and iron oxide minerals that are

key contributors to soil C storage and persistence (Rasmussen et al., 2018; Slessarev et al.,

2022).

Our supposition that soil C is older in the wetter sites due to the preferential stabilization

of necromass-C is consistent with current literature on soil C cycling in grasslands. The

transformation of plant-derived substrates to microbial products is believed to be a key process in

the stabilization of soil C (Cotrufo et al., 2012; Kallenbach et al., 2016) and appears to have a

pronounced importance in grasslands (Cotrufo et al., 2012; Kallenbach et al., 2016; Angst et al.,

2021). In grassland ecosystems, soil C age is linked to SOC chemistry such that products that are

more microbial in nature tend to be older (Heckman et al., 2021). Soil C:N, the only indicator of

soil C quality in this study, did not vary between our sites (Table 2). However, soil C:N is a

broad proxy and a more targeted molecular characterization of soil C is needed to conclude

whether microbial processing of C increases with MAP along the gradient studied here.

5. Conclusions

Soil water is an important controller of microbial growth and Earth system models project

significant changes in the timing and intensity of precipitation events globally (Dore, 2005;

Schimel, 2018). Seasonal changes in soil moisture have clear effects on instantaneous microbial

growth rates and our study demonstrates that soil water also has a more pervasive, long-term

legacy effect on microorganisms. Our data show that a site's historical climate regime, rather

than recent environmental conditions, strongly influences the structure of growing microbial

communities. Our study also demonstrates that a history of exposure to low soil moisture reduces

microbial growth, likely through shifts in community composition. Since changes in microbial

community structure operate over long time scales, ecosystems may experience a pervasive

legacy effect of climate history as precipitation regimes change. Lastly, our study demonstrates

that taxon-specific growth rates are remarkably consistent between three grassland sites from a

wide precipitation gradient, consistent with the idea that evolutionary history influences

microbial growth and points to the genetic and physiological constraints on growth rate in nature.

Acknowledgements

We thank Erin Nuccio, Amrita Bhattacharyya, Christina Ramon, and Aaron Chew for assistance

with field sampling and lab analyses; John Bailey and Allison Smith at Hopland Research and

Extension Center for help gathering meteorological data; and the Oregon State University Soil

Lab for chemical and physical soil analyses. We would like to thank Thomas Bristow, at the

NASA Ames Research Center, for allowing access to his lab space for qXRD analysis, and the

Stable Isotope Laboratory at UC Berkeley for stable isotope analyses. MM Foley was supported

by a National Science Foundation Graduate Research Fellowship while completing this work.

This research was supported by the U.S. Department of Energy, Office of Biological and

Environmental Research, Genomic Science Program 'Microbes Persist' Scientific Focus Area

(#SCW1632) at Lawrence Livermore National Laboratory (LLNL) and a subcontract to Northern

Arizona University. Work conducted at LLNL was conducted under the auspices of the US

Department of Energy under Contract DE-AC52-07NA27344.

Literature Cited

Angel, R., Pasternak, Z., Soares, M.I.M., Conrad, R., Gillor, O., 2013. Active and total prokaryotic communities in dryland soils. FEMS Microbiology Ecology 86, 130–138. doi:10.1111/1574-6941.12155

Angel, R., Soares, M.I.M., Ungar, E.D., Gillor, O., 2010. Biogeography of soil archaea and bacteria along a steep precipitation gradient. The ISME Journal 4, 553–563. doi:10.1038/ismej.2009.136

Angst, G., Mueller, K.E., Nierop, K.G., Simpson, M.J., 2021. Plant-or microbial-derived? A review on the molecular composition of stabilized soil organic matter. Soil Biology and Biochemistry 156, 108189.

Apprill, A., McNally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial Ecology 75, 129–137. doi:10.3354/ame01753

Averill, C., Waring, B.G., Hawkes, C.V., 2016. Historical precipitation predictably alters the shape and magnitude of microbial functional response to soil moisture. Global Change Biology 22, 1957–1964. doi:10.1111/gcb.13219

Bååth, E., 1996. Thymidine incorporation of bacteria sequentially extracted from soil using repeated homogenization-centrifugation. Microbial Ecology 31, 153–166.

Bakker, E., Hubert, F., Wander, M.M., Lanson, B., 2018. Soil development under continuous agriculture at the Morrow plots experimental fields from X-ray diffraction profile modelling. Soil Systems 2, 46.

Barnard, R.L., Blazewicz, S.J., Firestone, M.K., 2020. Rewetting of soil: revisiting the origin of soil CO2 emissions. Soil Biology and Biochemistry 107819. doi:10.1016/j.soilbio.2020.107819

Barnard, R.L., Osborne, C.A., Firestone, M.K., 2015. Changing precipitation pattern alters soil microbial community response to wet-up under a Mediterranean-type climate. The ISME Journal 9, 946–957. doi:10.1038/ismej.2014.192

Baubin, C., Farrell, A.M., Šťovíček, A., Ghazaryan, L., Giladi, I., Gillor, O., 2019. Seasonal and spatial variability in total and active bacterial communities from desert soil. Pedobiologia 74, 7–14. doi:10.1016/j.pedobi.2019.02.001

Berhe, A.A., Suttle, K.B., Burton, S.D., Banfield, J.F., 2012. Contingency in the direction and mechanics of soil organic matter responses to increased rainfall. Plant and Soil 358, 371–383. doi:10.1007/s11104-012-1156-0

Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: Critical review of estimation criteria and approaches. Soil Biology and Biochemistry 67, 192–211. doi:10.1016/j.soilbio.2013.08.024

Blazewicz, S.J., Schwartz, E., Firestone, M.K., 2014. Growth and death of bacteria and fungi underlie rainfall☐induced carbon dioxide pulses from seasonally dried soil. Ecology 95, 1162–1172.

Blomberg, S.P., Garland, T., Ives, A.R., 2003. Testing for Phylogenetic Signal in Comparative Data: Behavioral Traits Are More Labile. Evolution 57, 717–745. doi:10.1111/j.0014-3820.2003.tb00285.x

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M.,

Chase, J., Cope, E.K., Da Silva, R., Diener, C., Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson, D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B.D., Kang, K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D., McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J.T., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, M.S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu, Q., Knight, R., Caporaso, J.G., 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology 37, 852–857. doi:10.1038/s41587-019-0209-9

Bowsher, A.W., Kearns, P.J., Shade, A., 2019. 16S rRNA/rRNA gene ratios and cell activity staining reveal consistent patterns of microbial activity in plant-associated soil. Msystems 4, e00003-19.

Bradford, M.A., Keiser, A.D., Davies, C.A., Mersmann, C.A., Strickland, M.S., 2013. Empirical evidence that soil carbon formation from plant inputs is positively related to microbial growth. Biogeochemistry 113, 271–281. doi:10.1007/s10533-012-9822-0

Buckley, D.H., Huangyutitham, V., Hsu, S.-F., Nelson, T.A., 2007. Stable isotope probing with 15N achieved by disentangling the effects of genome G+ C content and isotope enrichment on DNA density. Applied and Environmental Microbiology 73, 3189–3195.

Bushnell, B., 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner (No. LBNL-7065E). Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States).

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods 13, 581–583. doi:10.1038/nmeth.3869

Campbell, B.J., Kirchman, D.L., 2013. Bacterial diversity, community structure and potential growth rates along an estuarine salinity gradient. The ISME Journal 7, 210–220.

Carini, P., Marsden, P.J., Leff, J.W., Morgan, E.E., Strickland, M.S., Fierer, N., 2017. Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. Nature Microbiology 2, 16242. doi:10.1038/nmicrobiol.2016.242

Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2012. The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter? Global Change Biology 19, 988–995. doi:10.1111/gcb.12113

Cruz-Martínez, K., Suttle, K.B., Brodie, E.L., Power, M.E., Andersen, G.L., Banfield, J.F., 2009. Despite strong seasonal responses, soil microbial consortia are more resilient to long-term changes in rainfall than overlying grassland. The ISME Journal 3, 738–744. doi:10.1038/ismej.2009.16

De Luca, G., Barakat, M., Ortet, P., Fochesato, S., Jourlin-Castelli, C., Ansaldi, M., Py, B., Fichant, G., Coutinho, P.M., Voulhoux, R., Bastien, O., Maréchal, E., Henrissat, B.,

Quentin, Y., Noirot, P., Filloux, A., Méjean, V., DuBow, M.S., Barras, F., Barbe, V.,
Weissenbach, J., Mihalcescu, I., Verméglio, A., Achouak, W., Heulin, T., 2011. The
Cyst-Dividing Bacterium Ramlibacter tataouinensis TTB310 Genome Reveals a Well-
Stocked Toolbox for Adaptation to a Desert Environment. PLoS ONE 6, e23784.
doi:10.1371/journal.pone.0023784

Doebelin, N., Kleeberg, R., 2015. Profex: a graphical user interface for the Rietveld refinement
program BGMN. Journal of Applied Crystallography 48, 1573–1580.

Dore, M.H.I., 2005. Climate change and changes in global precipitation patterns: What do we
know? Environment International 31, 1167–1181. doi:10.1016/j.envint.2005.03.004

dos Santos, D.F.K., Kyaw, C.M., De Campos, T.A., Miller, R.N.G., Noronha, E.F., Bustamante,
M.M. da C., Kruger, R., 2014. The Family Cystobacteraceae, in: Rosenberg, E., DeLong,
E.F., Lory, S., Stackebrandt, E., Thompson, F. (Eds.), The Prokaryotes:
Deltaproteobacteria and Epsilonproteobacteria. Springer Berlin Heidelberg, Berlin,
Heidelberg, pp. 19–40. doi:10.1007/978-3-642-39044-9_304

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Research 32, 1792–1797.

Evans, S.E., Wallenstein, M.D., 2014. Climate change alters ecological strategies of soil bacteria.
Ecology Letters 17, 155–164. doi:10.1111/ele.12206

Flemming, H.C., Wingender, J., 2001. Relevance of microbial extracellular polymeric substances
(EPSs)--Part I: Structural and ecological aspects. Water Science and Technology: A
Journal of the International Association on Water Pollution Research 43, 1–8.

Fossum, C., Estera-Molina, K., Yuan, M., Herman, D., Chu-Jacoby, I., Nico, P., Morrison, K.,
Pett-Ridge, J., Firestone, M., 2021. Belowground allocation and dynamics of recently
fixed plant carbon in a California annual grassland soil. BioRxiv 2021.08.23.457405.
doi:10.1101/2021.08.23.457405

Hawkes, C.V., Shinada, M., Kivlin, S.N., 2020. Historical climate legacies on soil respiration
persist despite extreme changes in rainfall. Soil Biology and Biochemistry 143, 107752.
doi:10.1016/j.soilbio.2020.107752

Heckman, K., Hicks Pries, C.E., Lawrence, C.R., Rasmussen, C., Crow, S.E., Hoyt, A.M., von
Fromm, S.F., Shi, Z., Stoner, S., McGrath, C., 2022. Beyond bulk: Density fractions
explain heterogeneity in global soil carbon abundance and persistence. Global Change
Biology 28, 1178–1196.

Heckman, K.A., Nave, L.E., Bowman, M., Gallo, A., Hatten, J.A., Matosziuk, L.M., Possinger,
A.R., SanClements, M., Strahm, B.D., Weiglein, T.L., Rasmussen, C., Swanston, C.W.,
2021. Divergent controls on carbon concentration and persistence between forests and
grasslands of the conterminous US. Biogeochemistry 156, 41–56. doi:10.1007/s10533-
020-00725-z

Hicks, L.C., Rahman, M.M., Carnol, M., Verheyen, K., Rousk, J., 2018. The legacy of mixed
planting and precipitation reduction treatments on soil microbial activity, biomass and
community composition in a young tree plantation. Soil Biology and Biochemistry 124,
227–235. doi:10.1016/j.soilbio.2018.05.027

Homyak, P.M., Blankinship, J.C., Slessarev, E.W., Schaeffer, S.M., Manzoni, S., Schimel, J.P.,
2018. Effects of altered dry season length and plant inputs on soluble soil carbon.
Ecology 99, 2348–2362. doi:10.1002/ecy.2473

Hungate, B.A., Mau, R.L., Schwartz, E., Caporaso, J.G., Dijkstra, P., van Gestel, N., Koch, B.J.,
Liu, C.M., McHugh, T.A., Marks, J.C., Morrissey, E.M., Price, L.B., 2015. Quantitative

Microbial Ecology through Stable Isotope Probing. Applied and Environmental Microbiology 81, 7570. doi:10.1128/AEM.02280-15

Janssen, S., McDonald, D., Gonzalez, A., Navas-Molina, J.A., Jiang, L., Xu, Z.Z., Winker, K., Kado, D.M., Orwoll, E., Manary, M., 2018. Phylogenetic placement of exact amplicon sequences improves associations with clinical information. Msystems 3, e00021-18.

Jones, S.E., Lennon, J.T., 2010. Dormancy contributes to the maintenance of microbial diversity. Proceedings of the National Academy of Sciences 107, 5881–5886.

Kallenbach, C.M., Frey, S.D., Grandy, A.S., 2016. Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls. Nature Communications 7, 13630.

Koch, B.J., McHugh, T.A., Hayer, M., Schwartz, E., Blazewicz, S.J., Dijkstra, P., Gestel, N., Marks, J.C., Mau, R.L., Morrissey, E.M., Pett-Ridge, J., Hungate, B.A., 2018. Estimating taxon-specific population dynamics in diverse microbial communities. Ecosphere 9, e02090. doi:10.1002/ecs2.2090

Kotanen, P.M., 2004. Revegetation following Soil Disturbance and Invasion in a Californian Meadow: a 10-year History of Recovery. Biological Invasions 6, 245–254. doi:10.1023/B:BINV.0000022145.03215.4f

Lavallee, J.M., Soong, J.L., Cotrufo, M.F., 2020. Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century. Global Change Biology 26, 261–273. doi:10.1111/gcb.14859

Leitner, S., Homyak, P.M., Blankinship, J.C., Eberwein, J., Jenerette, G.D., Zechmeister-Boltenstern, S., Schimel, J.P., 2017. Linking NO and N2O emission pulses with the mobilization of mineral and organic N upon rewetting dry soils. Soil Biology and Biochemistry 115, 461–466. doi:10.1016/j.soilbio.2017.09.005

Leizeaga, A., Hicks, L.C., Manoharan, L., Hawkes, C.V., Rousk, J., 2021. Drought legacy affects microbial community trait distributions related to moisture along a savannah grassland precipitation gradient. Journal of Ecology n/a. doi:10.1111/1365-2745.13550

Lennon, J.T., Aanderud, Z.T., Lehmkuhl, B.K., Schoolmaster, D.R., 2012. Mapping the niche space of soil microorganisms using taxonomy and traits. Ecology 93, 1867–1879. doi:10.1890/11-1745.1

Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial control over soil carbon storage. Nature Microbiology 2, 1–6. doi:10.1038/nmicrobiol.2017.105

Locey, K.J., Muscarella, M.E., Larsen, M.L., Bray, S.R., Jones, S.E., Lennon, J.T., 2020. Dormancy dampens the microbial distance–decay relationship. Philosophical Transactions of the Royal Society B: Biological Sciences 375, 20190243. doi:10.1098/rstb.2019.0243

Maestre, F.T., Delgado-Baquerizo, M., Jeffries, T.C., Eldridge, D.J., Ochoa, V., Gozalo, B., Quero, J.L., García-Gómez, M., Gallardo, A., Ulrich, W., Bowker, M.A., Arredondo, T., Barraza-Zepeda, C., Bran, D., Florentino, A., Gaitán, J., Gutiérrez, J.R., Huber-Sannwald, E., Jankju, M., Mau, R.L., Miriti, M., Naseri, K., Ospina, A., Stavi, I., Wang, D., Woods, N.N., Yuan, X., Zaady, E., Singh, B.K., 2015. Increasing aridity reduces soil microbial diversity and abundance in global drylands. Proceedings of the National Academy of Sciences 112, 15684–15689. doi:10.1073/pnas.1516684112

Malik, A.A., Martiny, J.B.H., Brodie, E.L., Martiny, A.C., Treseder, K.K., Allison, S.D., 2020a. Defining trait-based microbial strategies with consequences for soil carbon cycling under climate change. The ISME Journal 14, 1–9. doi:10.1038/s41396-019-0510-0

Malik, A.A., Swenson, T., Weihe, C., Morrison, E.W., Martiny, J.B.H., Brodie, E.L., Northen,
T.R., Allison, S.D., 2020b. Drought and plant litter chemistry alter microbial gene
expression and metabolite production. The ISME Journal 1–12. doi:10.1038/s41396-020-
0683-6
McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE 8, e61217.
doi:10.1371/journal.pone.0061217
Meisner, A., Bååth, E., Rousk, J., 2013. Microbial growth responses upon rewetting soil dried for
four days or one year. Soil Biology and Biochemistry 66, 188–192.
doi:10.1016/j.soilbio.2013.07.014
Meter Group, 2019. WP4C.
Michael Barbour, Todd Keeler-Wolf, Allan A. Schoenherr, 2007. Terrestrial Vegetation of
California, 3rd Edition. University of California Press, Berkeley.
Minasny, B., McBratney, A.B., Brough, D.M., Jacquier, D., 2011. Models relating soil pH
measurements in water and calcium chloride that incorporate electrolyte concentration.
European Journal of Soil Science 62, 728–732. doi:10.1111/j.1365-2389.2011.01386.x
Morrissey, E.M., Mau, R.L., Hayer, M., Liu, X.-J.A., Schwartz, E., Dijkstra, P., Koch, B.J.,
Allen, K., Blazewicz, S.J., Hofmockel, K., Pett-Ridge, J., Hungate, B.A., 2019.
Evolutionary history constrains microbial traits across environmental variation. Nature
Ecology & Evolution 3, 1064–1069. doi:10.1038/s41559-019-0918-y
Morrissey, E.M., Mau, R.L., Schwartz, E., McHugh, T.A., Dijkstra, P., Koch, B.J., Marks, J.C.,
Hungate, B.A., 2017. Bacterial carbon use plasticity, phylogenetic diversity and the
priming of soil organic matter. The ISME Journal 11, 1890–1899.
doi:10.1038/ismej.2017.43
Neufeld, J.D., Vohra, J., Dumont, M.G., Lueders, T., Manefield, M., Friedrich, M.W., Murrell,
J.C., 2007. DNA stable-isotope probing. Nature Protocols 2, 860–866.
doi:10.1038/nprot.2007.109
Nijs, E.A. de, Hicks, L.C., Leizeaga, A., Tietema, A., Rousk, J., 2019. Soil microbial moisture
dependences and responses to drying–rewetting: The legacy of 18 years drought. Global
Change Biology 25, 1005–1015. doi:10.1111/gcb.14508
Nuccio, E.E., Starr, E., Karaoz, U., Brodie, E.L., Zhou, J., Tringe, S.G., Malmstrom, R.R.,
Woyke, T., Banfield, J.F., Firestone, M.K., Pett-Ridge, J., 2020. Niche differentiation is
spatially and temporally regulated in the rhizosphere. The ISME Journal 14, 999–1014.
doi:10.1038/s41396-019-0582-x
Pagel, M., 1999. Inferring the historical patterns of biological evolution. Nature 401, 877–884.
doi:10.1038/44766
Papp, K., Hungate, B.A., Schwartz, E., 2018a. Microbial rRNA Synthesis and Growth Compared
through Quantitative Stable Isotope Probing with $H_2{}^{18}O$. Applied and Environmental
Microbiology 84, e02441-17. doi:10.1128/AEM.02441-17
Papp, K., Hungate, B.A., Schwartz, E., 2018b. Microbial rRNA Synthesis and Growth Compared
through Quantitative Stable Isotope Probing with H218O. Applied and Environmental
Microbiology 84. doi:10.1128/AEM.02441-17
Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global
field samples. Environmental Microbiology 18, 1403–1414. doi:10.1111/1462-
2920.13023

Pertassek, T., Andre, P., Durner, W., 2011. HYPROP - Data Evaluation Software User's Manual.

Power, M.E., Bouma-Gregson, K., Higgins, P., Carlson, S.M., 2015. The Thirsty Eel: Summer
and Winter Flow Thresholds that Tilt the Eel River of Northwestern California from
Salmon-Supporting to Cyanobacterially Degraded States. Copeia 103, 200–211.
doi:10.1643/CE-14-086

Price, M.N., Dehal, P.S., Arkin, A.P., 2010. FastTree 2 – Approximately Maximum-Likelihood
Trees for Large Alignments. PLOS ONE 5, e9490. doi:10.1371/journal.pone.0009490

Rahman, M.M., Hicks, L.C., Verheyen, K., Rousk, J., Carnol, M., 2018. Effects of drought
legacy and tree species admixing on bacterial growth and respiration in a young forest
soil upon drying and rewetting. Soil Biology and Biochemistry 127, 148–155.
doi:10.1016/j.soilbio.2018.09.026

Rasmussen, C., Heckman, K., Wieder, W.R., Keiluweit, M., Lawrence, C.R., Berhe, A.A.,
Blankinship, J.C., Crow, S.E., Druhan, J.L., Hicks Pries, C.E., 2018. Beyond clay:
towards an improved set of variables for predicting soil organic matter content.
Biogeochemistry 137, 297–306.

Rousk, J., Brookes, P.C., Bååth, E., 2009. Contrasting Soil pH Effects on Fungal and Bacterial
Growth Suggest Functional Redundancy in Carbon Mineralization. Applied and
Environmental Microbiology 75, 1589–1596. doi:10.1128/AEM.02775-08

Rousk, J., Smith, A.R., Jones, D.L., 2013. Investigating the long-term legacy of drought and
warming on the soil microbial community across five European shrubland ecosystems.
Global Change Biology 19, 3872–3884. doi:10.1111/gcb.12338

Sandhya, V., Ali, Sk.Z., 2015. The production of exopolysaccharide by Pseudomonas putida
GAP-P45 under various abiotic stress conditions and its role in soil aggregation.
Microbiology 84, 512–519. doi:10.1134/S0026261715040153

Schimel, J.P., 2018. Life in Dry Soils: Effects of Drought on Soil Microbial Communities and
Processes. Annual Review of Ecology, Evolution, and Systematics 49, 409–432.
doi:10.1146/annurev-ecolsys-110617-062614

Schwartz, E., 2007. Characterization of Growing Microorganisms in Soil by Stable Isotope
Probing with H218O. Appl. Environ. Microbiol. 73, 2541–2546.
doi:10.1128/AEM.02021-06

Seidl, M., Dietrich, W., Schmidt, K., de Ploey, J., 1992. The problem of channel erosion into
bedrock. Functional Geomorphology 23, 101–124.

Shi, S., Nuccio, E., Herman, D.J., Rijkers, R., Estera, K., Li, J., Rocha, U.N. da, He, Z., Pett-
Ridge, J., Brodie, E.L., Zhou, J., Firestone, M., 2015. Successional Trajectories of
Rhizosphere Bacterial Communities over Consecutive Seasons. MBio.
doi:10.1128/mBio.00746-15

Shi, S., Nuccio, E.E., Shi, Z.J., He, Z., Zhou, J., Firestone, M.K., 2016. The interconnected
rhizosphere: High network complexity dominates rhizosphere assemblages. Ecology
Letters 19, 926–936. doi:10.1111/ele.12630

Sieradzki, E.T., Koch, B.J., Greenlon, A., Sachdeva, R., Malmstrom, R.R., Mau, R.L.,
Blazewicz, S.J., Firestone, M.K., Hofmockel, K.S., Schwartz, E., Hungate, B.A., Pett-
Ridge, J., 2020. Measurement Error and Resolution in Quantitative Stable Isotope
Probing: Implications for Experimental Design. MSystems 5.
doi:10.1128/mSystems.00151-20

Slessarev, E.W., Chadwick, O.A., Sokol, N.W., Nuccio, E.E., Pett-Ridge, J., 2022. Rock
weathering controls the potential for soil carbon storage at a continental scale.
Biogeochemistry 157, 1–13. doi:10.1007/s10533-021-00859-8
Sokol, N.W., Slessarev, E., Marschmann, G.L., Nicolas, A., Blazewicz, S.J., Brodie, E.L.,
Firestone, M.K., Foley, M.M., Hestrin, R., Hungate, B.A., 2022. Life and death in the soil
microbiome: How ecological processes influence biogeochemistry. Nature Reviews
Microbiology 1–16.
Stuiver, M., Polach, H.A., 1977. Discussion Reporting of 14C Data. Radiocarbon 19, 355–363.
doi:10.1017/S0033822200003672
Sullivan, M.J.P., A.Thomsen, M., Suttle, K.B., 2016. Grassland responses to increased rainfall
depend on the timescale of forcing. Global Change Biology 22, 1655–1665.
doi:10.1111/gcb.13206
Torn, M., Swanston, C., Castanha, C., Trumbore, S., 2009. Storage and turnover of organic
matter in soil. Biophysico-Chemical Processes Involving Natural Nonliving Organic
Matter in Environmental Systems. Wiley, Hoboken 219–272.
Uhlířová, E., Šantrůčková, H., 2003. Growth rate of bacteria is affected by soil texture and
extraction procedure. Soil Biology and Biochemistry 35, 217–224. doi:10.1016/S0038-
0717(02)00254-7
Voroney, R.P., Heck, R.J., 2015. Chapter 2 - The Soil Habitat, in: Paul, E.A. (Ed.), Soil
Microbiology, Ecology and Biochemistry (Fourth Edition). Academic Press, Boston, pp.
15–39. doi:10.1016/B978-0-12-415955-6.00002-5
Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. Applied and
Environmental Microbiology 73, 5261–5267. doi:10.1128/AEM.00062-07
948

Figure Captions

Figure 1: $\Delta^{14}C$ (‰) of bulk soil and respired CO_2 from three CA annual grasslands. The
horizontal line represents atmospheric $\Delta^{14}C$ in 2018 and serves as a reference for post-bomb
enrichment. Differences between grassland site means (n=3) were assessed by one-way
ANOVAs and Fisher's LSD. $\Delta^{14}C$ (‰) of bulk soils differed between sites ($p<0.05$, Fisher's
LSD) except Hopland did not differ from Sedgwick ($p>0.05$, Fisher's LSD). $\Delta^{14}C$ (‰) of
respired CO_2 did not differ between sites ($p>0.05$, Fisher's LSD).
956

957   **Figure 2:** PCoA of Bray-Curtis dissimilarities for total (n=3) and growing (n=3) microbial
958   communities assessed via 16S rRNA gene abundance and $H_2^{18}O$ stable isotope probing in three
959   CA annual grasslands, sampled during the winter wet season. Community composition varied
960   significantly by site for the total and growing microbial communities ($p < 0.01$, ANOSIM).
961

Figure 3: Community and phylum mean fraction of maximum enrichment ^{18}O (FME) of soil
microbial taxa that are growing (95% C.I. for excess atom fraction $^{18}O > 0$) at three CA annual
grassland sites. Differences between grassland site means were assessed by one-way ANOVAs
and Fisher's LSD. Letters indicate significant differences between sites ($p<0.05$, Fisher's LSD).
FME ^{18}O is computed as the excess atom fraction ^{18}O of a taxon divided by the enrichment of the
soil water during the qSIP incubation.
968

Figure 4: Fraction of maximum enrichment ^{18}O (FME) of taxa growing in Sedgwick soil versus
other annual grassland soils. Taxa plotted on the left were those growing in Sedgwick soils as
well at least one other site. Taxa plotted on the right were found growing in Sedgwick soils
alone. The FME ^{18}O of taxa unique to Sedgwick soils is lower than that of taxa that grow in at
least one other site (p-value = 0.01, t-test). FME ^{18}O is computed as the excess atom fraction ^{18}O
of a taxon divided by the enrichment of the soil water during the qSIP incubation.
975

Figure 5: Linear regressions (black line) of fraction of maximum enrichment ^{18}O (FME) of
growing microbial taxa that co-occurred and grew in at least two sites from a latitudinal and
precipitation gradient of annual grasslands in CA (Sedgwick>>Hopland>>Angelo). Shaded area
shows 95% C.I. for the regression. Detailed regression results are presented in Supplementary
Table 5. FME ^{18}O is computed as the excess atom fraction ^{18}O of a taxon divided by the
enrichment of the soil water during the qSIP incubation.

Tables

Table 1: Characteristics of three California annual grasslands examined in this study.

	Angelo Coast Range Reserve	Hopland Research and Extension Center	Sedgwick Reserve
Location	39°44.25`N, 123°37.82`W	39°00.11`N, 123°04.18`W	34°42.70'N, 120°2.33'W
Elevation (m)	475	180	360
MAT max/min (°C)	20/6	23/7	24/7
MAP (mm yr^{-1})	2160	956	383
Parent Material	Argillite/Sandstone interbedding	Sandstone, Shale, Graywacke, Schist	Alluvium, Calcareous shale, Acid sandstone
Soil Order	Alfisol	Alfisol	Mollisol
Soil Series	Holohan-Hollowtree-Casabonne complex	Witherell-Squawrock complex	Botella
Dominant vegetation	*Aira spp., Bromus spp., Briza spp.*	*Avena spp., Bromus spp., Erodium spp., Festuca spp.*	*Avena spp., Bromus spp.*

985 **Table 2:** Physical and chemical characteristics of surface soils from three northern California
986 annual grasslands: Angelo, Hopland, and Sedgwick (mean ± standard error, n = 3). Differences
987 between site means were assessed by one-way ANOVAs and Fisher's LSD. Bold text indicates
988 ANOVA P-values less than 0.05, and BDL signifies values below detection limit.

	Angelo Avg ± SE	Hopland Avg ± SE	Sedgwick Avg ± SE	ANOVA *P*
δ^{13}C Bulk Soil (‰)	-27.54 ± 0.1	-27.64 ± 0.1	-26.99 ± 0.2	**0.04**
δ^{15}N Bulk Soil (‰)	3.75 ± 0.4	4.23 ± 0.3	4.85 ± 0.2	0.10
pH	5.02 ± 0.02	5.55 ± 0.03	6.99 ± 0.03	**<0.01**
Total base cations (meq kg^{-1} soil)	10.49 ± 0.7	10.79 ± 0.7	27.61 ± 0.9	**<0.01**
ECEC (meq kg^{-1} soil)	10.99 ± 0.4	10.84 ± 0.4	27.61 ± 0.5	**<0.01**
% Sand	28 ± 2	45 ± 1	38 ± 1	**<0.01**
% Silt	42 ± 1	36 ± 1	28 ± 2	**<0.01**
% Clay	27 ± 1	19 ± 1	34 ± 0	**<0.01**
% C	1.66 ± 0.3	1.39 ± 0.2	1.99 ± 0.1	0.22
% N	0.13 ± 0.02	0.11 ± 0.01	0.18 ± 0.01	**0.03**
C:N	12.19 ± 0.6	12.37 ± 0.2	11.04 ± 0.1	0.11
Al (mg kg^{-1})	44.77 ± 1.1	5.17 ± 0.4	*BDL*	**<0.01**
Ca (mg kg^{-1})	1455.44 ± 53.0	1350.63 ± 47.6	2380.67 ± 50.3	**<0.01**
K (mg kg^{-1})	98.98 ± 6.3	114.92 ± 13.9	390.39 ± 29.4	**<0.01**
Mg (mg kg^{-1})	353.39 ± 18.0	448.75 ± 14.9	1760.32 ± 38.3	**<0.01**
Na (mg kg^{-1})	3.44 ± 1.2	*BDL*	7.21 ± 1.2	**0.03**