Ocean-Warming During Embryogenesis Programs a Lasting Transcriptomic Signature in Fishes

Daniel M. Ripley[1], Terence Garner[2], Samantha A. Hook[3], Ana Veríssimo[4,5], Bianka Grunow[6], Timo Moritz[7,8], Peter Clayton[2], Holly A. Shiels[1], Adam Stevens[2]

[1]Division of Cardiovascular Sciences, Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK.

[2]Division of Developmental Biology and Medicine, Faculty of Biology, Medicine, and Health, The University of Manchester, Manchester, UK.

[3]Department of Earth and Environmental Sciences, The University of Manchester, Manchester, UK.

[4]CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

[5]BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661 Vairão, Portugal

[6]Institute of Muscle Biology and Growth, Research Institute for Farm Animal Biology, Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany.

[7]Deutsches Meeresmuseum, Katharinenberg 14–20, 18437 Stralsund, Germany

[8]Institute of Biological Sciences, University of Rostock, Albert-Einstein-Straße 3, 18059 Rostock, Germany

KEYWORDS climate change, long term effect, gene expression, plasticity, developmental programming, *Scyliorhinus canicula*, RNA-seq.

RUNNING HEADER Warming programs the transcriptome of fishes

Abstract

Exposure to elevated temperatures during embryogenesis has profound acute effects on the cardiac performance, metabolism, and growth of fishes. Some temperature-induced effects may be retained into, or manifest in, later-life through a mechanism termed developmental programming. In this study, we incubated *Scyliorhinus canicula* embryos at either 15°C or 20°C before transferring the newly hatched sharks to a common set of conditions (15°C) for 5 months. Lasting transcriptomic differences associated with the developmental environment were identified, and interactions between cardiac genes were investigated using hypernetwork modelling. Development at an elevated temperature caused changes in transcriptomic connectivity and entropy, parameters thought to relate to plasticity and fitness. We then validated these observations through a novel re-analysis of published *Danio rerio* and *Dicentrarchus labrax* muscle tissue datasets. Together, these data demonstrate a persistent, programmed effect of developmental temperature on the co-ordination of gene expression in three species of fishes, which may relate to altered plasticity and fitness in later-life.

1 INTRODUCTION

Developmental programming comprises changes to an organism's phenotype that are induced, or programmed, by the environmental conditions experienced during development (Bateson *et al.*, 2014). These developmental conditions elicit changes in DNA and chromatin methylation patterns, with consequent alterations in gene expression (Fellous *et al.*, 2015). Some of these changes in methylation pattern, and their concomitant influence on gene expression, can become fixed and retained into later life. For instance, changes in developmental temperature cause hypermethylation in threespine stickleback (Metzger & Schulte, 2017), and studies in rainbow trout (*Oncorhynchus mykiss*) demonstrate the ability of early-life hypoxic events (Liu *et al.*, 2017) and dietary composition (Geurden *et al.*, 2014) to alter glucose metabolism and homeostasis in later-life. Finally, the developmental temperature of zebrafish (*Danio rerio*) has been shown to program their gene expression and physiology, culminating in an enhanced thermal tolerance (Schaefer & Ryan, 2006) and acclimation capacity (Schnurr *et al.*, 2014; Scott & Johnston, 2012) in adulthood.

As well as altering mean trait values, the embryonic environment can also program an organism's ability to acclimate to conditions experienced in later-life (Beaman *et al.*, 2016). That is, the capacity to physiologically respond to and mitigate future environmental challenges can be enhanced or hindered by an individual's embryonic conditions (Beaman *et al.*, 2016). Evidence in zebrafish (*Danio rerio*) (Scott & Johnston, 2012; Schnurr *et al.*, 2014) and mosquitofish (*Gambusia holbrooki*) (Seebacher *et al.*, 2014), amongst other species (Beaman *et al.*, 2016), show that the thermal environment experienced during early-life can have lasting effects on the acclimation capacity of multiple physiological traits, including metabolic scope (Seebacher *et al.*, 2014), critical thermal maximum (Healy *et al.*, 2019), and swimming performance (Scott & Johnston, 2012).

Phenotypic changes arising from physiological plasticity are underwritten by changes in an organism's gene expression in response to environmental cues. Such plasticity may be affected by an individual´s embryonic environment, which may either facilitate or hinder changes in gene expression later in life (Beaman *et al.*, 2016). Thus, there is a link between an organism's phenotype, gene expression, and plastic potential, which can be modulated by its embryonic environment (Beaman *et al.*, 2016). Transcriptional entropy is a metric describing the order or predictability of gene-gene interactions within a transcriptional network. The more entropic a network, the less structured and less predictable its behaviour. Transcriptional entropy is known to correlate with the differentiation potency and phenotypic plasticity of single cells (Banerji *et al.*, 2013; Teschendorff & Enver, 2017). The more entropic a cell, the greater its number of possible fates (Teschendorff & Enver, 2017), and thus the more plasticity it possesses. Given that several studies have shown that the environmental temperature experienced during development can alter plasticity in the adult organism (Beaman *et al.*, 2016), we aimed to assess whether developmental temperature also alters gene-level plasticity by measuring the network entropy and connectivity of the transcriptome in later life.

Whilst many animals may experience adverse environmental conditions during embryogenesis, some species, owing to their ecology, are more prone than others. Oviparous elasmobranchs, because of their protracted, sessile developmental period, are susceptible to experiencing sub-optimal

environmental conditions for long durations. Oviparous elasmobranchs lay their eggs inside
collagenous egg cases termed mermaid's purses, which the embryo remains within for several
months to over a year, depending on the species (Benjamins *et al.*, 2021; McLaughin & Ogower,
1971). Throughout embryogenesis the embryo is unable to move from, and thus may be consistently
exposed to, unfavourable environmental conditions such as elevated temperatures. As the world's
oceans continue to warm (IPCC, 2021), the likelihood of oviparous elasmobranch embryos
developing in sub-optimal conditions is increasing. Representative pathway (RCP) 8.5, a worst-case
scenario model, estimates a ~3˚C increase in ocean temperatures by the end of the century if
greenhouse gas emissions continue to rise (Cai *et al.*, 2014). The rising ocean temperatures, coupled
with an increasing frequency and duration of marine heatwaves (Oliver *et al.*, 2018), renders
exposure to temperatures above animal's optima highly likely. Exposure to elevated temperatures is
known to affect the growth and development of elasmobranch embryos (Hume, 2019; Musa *et al.*,
2020; Rosa *et al.*, 2014). However, no studies have investigated how embryogenesis at elevated
temperatures may have lasting effects on elasmobranchs through developmental programming
mechanisms.

Here, we incubate *Scyliorhinus canicula* embryos at 15˚C or 20˚C throughout embryogenesis before
moving them to a common-garden environment (15˚C) upon hatching. Following 5 months in the
common garden environment, we sacrificed the animals and performed RNA-seq on the ventricle
tissue of six individuals to investigate whether the developmental environment could program lasting
changes in the gene expression and organisation of the *S. canicula* ventricular transcriptome. Finally,
to validate our findings, we re-analysed published datasets of temperature-induced developmental
programming in zebrafish (*Danio rerio)* hypaxial fast muscle (Scott & Johnston, 2012) and European
seabass (*Dicentrarchus labrax)* muscle (Anastasiadi *et al.*, 2021) to test the consistency of our
observations across multiple species of fishes.

2 MATERIAL AND METHODS

2.1 Experimental animals

Scyliorhinus canicula embryos were collected from a population of 7 adult individuals held at 15˚C at
the Ozeaneum, Stralsund, Germany, and transported to the University of Manchester, UK. Upon
arrival the health and developmental stage of each embryo was assessed using the Musa scale
(Musa *et al.*, 2018), and the egg cases were submerged in dechlorinated tap water for two minutes to
remove any marine microorganisms. Only healthy embryos at stage 1 were used in the study (Musa
et al., 2018). All regulated procedures received approval from the institution's ethical review board.

Following embryonic staging and two days in quarantine (15˚C), individuals were randomly assigned
to a temperature treatment group (15 ± 0.3˚C or 20 ± 0.3˚C). 15˚C was chosen for the control
condition as it falls within the range of temperatures experienced by *S. canicula* in the wild and is the
holding temperature of the parent population at the Ozeaneum, Stralsund, Germany. The egg cases
in each treatment group were hung vertically in two, well aerated, 55 l static seawater (35ppt salinity)
tanks equipped with internal filters and left to continue their embryonic development.

Upon hatching, fin clips were taken to facilitate identification in later-life through microsatellite
analysis. The fin clips were stored in 98% ethanol at -20˚C, and the sharks were moved into one of
four well aerated 400 l static seawater tanks held at 15˚C ± 0.3˚C. The hatchlings from the 20˚C
treatment group were lowered to 15˚C at a rate of 2.5˚C per day, and sharks from both treatment
groups were mixed and randomly allocated to a tank. The sharks were fed a mixture of squid, crab
and krill three times per week. The sharks from both treatment groups were held at 15˚C ± 0.3˚C for
an average of 136 days prior to tissue sampling. During the entire experiment, saltwater changes
were performed three times weekly to maintain ammonia, nitrite and nitrate below detectable levels.

2.2 Tissue sampling
Sharks at 4-5 months age (mean ± SEM = 136 ± 6.3days) were euthanised with an overdose of
buffered tricane methanesulfonate and a fin clip was taken post-humous. The ventricle was excised
from each individual, placed into RNAlater, and flash frozen in liquid nitrogen before being stored at -
80˚C prior to RNA-sequencing.

2.3 Individual identification using microsatellite analysis
To identify which condition an individual was developed in, microsatellite analysis was performed on
the fin clips taken at birth and at the end of the experiment to match each individual using the
methods described in Hook *et al*. (2019). Fin clips were extracted using a BioLine Isolate Genomic kit
with an extended proteinase K digestion to maximise DNA yield. DNA yield was assessed using a
NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific, USA) and gel electrophoresis. A
one-primer cocktail containing 11 microsatellites and three tail dyes was used for DNA amplification
(Griffiths *et al*., 2011) using the QIAGEN multiplex PCR kit. The thermocycling protocol consisted of
initial denaturation cycle at 95°C for 15 minutes, followed by 35 cycles of 94°C for 30 seconds, 60°C
for 90 seconds and 72°C for 45 seconds, and finalised by one cycle at 72°C for 30 minutes. The
products were visualised on a gel and then genotyped using an ABI sequencer. Positive control
samples were added to each plate genotyped to account for possible allele slippage.

Genotypes were scored using GeneMapper© v4.1 (Applied Biosystems) and validated through
Microchecker (van Oosterhout *et al*., 2004). Duplicates were found between the two batches using
CERVUS (Marshall *et al*., 1998). In cases where some alleles within the genotype did not match,
controls were used as a reference to identify possible allele slippage. Probabilities of the identity
analysis (pID) were taken from CERVUS to confirm match identification between the two batches.
147

2.4 *Danio rerio* and *Dicentrarchus labrax* re-analysis

The computational analysis pipeline (sections 2.5-2.7 below) was also implemented on previously
published RNA-seq datasets of temperature-induced developmental programming in zebrafish (*D.*
rerio) hypaxial fast muscle (Scott & Johnston, 2012) and European seabass (*D. labrax*) muscle tissue
(Anastasiadi *et al*., 2021, Anastasiadi & Piferrer, 2019). This analysis was performed to assess the
robustness and between-species consistency of the temperature-induced effects observed in *S.*
canicula.

Full details of the animals used in the re-analysis can be found in Scott & Johnston 2012 (*D. rerio*,

age = 8-9 months, N = 5-8) and Anastasiadi *et al.* 2021 and Anastasiadi & Piferrer, 2019 (*D. labrax*,

age = 36 months, N = 5) respectively. The *D. rerio* and *D. labrax* data were processed in an identical

manner to the *S. canicula* data. Transcriptome assembly statistics for the three species are shown in

table 1.

2.5 de-novo transcriptome assembly and differential expression analysis

The ventricle tissue from six individuals (N = 3 per treatment group), stored in RNAlater at -80˚C, was

used for generating the transcriptome. The tissue samples were extracted using the Qiagen kit with

homogenisation. Basic sample quality was assessed using a Nanodrop (ThermoFisher) and the high-

quality ventricular RNA samples were sequenced using a 76 base pair long, pair-ended reads on the

Illumina TruSeq system. Reads that mapped to human, bacterial, and viral sequences were removed

using DeconSeq (Schmieder and Edwards, 2011) whilst reads mapping to ribosomal RNA were

removed using sortmeRNA (Kopylova *et al.*, 2012). Adaptor sequences and sequences with a low-

quality score (regions averaging a score <5 over a 4bp sliding window, and leading/trailing sequences

scoring <5) were then removed from the cleaned reads using Trimmomatic (Bolger *et al.*, 2014) prior

to transcriptome assembly. 264 million bases survived the quality control process and were used for

transcriptome assembly in Trinity 2.8.4 (Grabherr *et al.*, 2011; Haas *et al.*, 2013). Default parameters

were used in Trinity, except –normalize_reads, producing 347710 contigs with a N50 of 1422, and a

BUSCO score of 88.8%, with 519 single copy, and 2458 duplicated copy, BUSCOs (BUSCO v5.2.2,

Simao *et al.*, 2015). Open reading frames (ORFs) were predicted from the transcripts using

Transdecoder (https://github.com/TransDecoder/TransDecoder/wiki) with a minimum length threshold

of 100 amino acids. The retained predicted protein sequences were annotated using a BLAST search

against the SwissProt database. The quality-controlled reads were then mapped using Bowtie-2

(Langmead & Salzberg, 2012), and pseudo-aligned to the de novo transcript assemblies using

Kallisto (Bray *et al.*, 2016), allowing the relative abundance of each gene to be calculated. Where

multiple transcripts mapped to a single predicted protein, the median transcript abundance was used.

The abundance estimates were filtered by removing fragments that had zero mapped reads in any of

the samples before differential expression analysis was performed with EdgeR (McCarthy *et al.*, 2012;

Robinson *et al.*, 2010) in R 3.6.0 (R core team, 2019).

2.6 Hypernetwork analysis

The differentially expressed genes (DEGs) were then used for hypernetwork analysis to assess the

network properties of the transcriptomes from the 15°C and 20°C treatment groups (supplementary

figure 1). The expression of each DEG was tested for correlations against all the assembled

transcripts within each treatment group. The correlation values were binarized using a cut-off of one

standard deviation from the mean, such that strong positive or negative correlations were retained

(denoted as 1) and weak correlations were discarded (denoted as 0). The binarized correlation matrix

is square (DEGs x DEGs) and represents the incidence matrix of the hypernetwork. The binarized

correlation matrix was then multiplied by the transpose of itself ($M * M^t$) to return the adjacency matrix

of the hypernetwork, where the values in each cell represent the number of correlations with non-

DEGs shared between any pair of DEGs. These values also represent the dimensionality of the

hyperedge connecting each pair of nodes. Supplementary figure 1 gives an overview of the

hypernetwork analysis. The data were visualised as heatmaps to assess clustering of the genes, per-

gene connectivity, and per-gene Shannon entropy (herein, entropy) using the R packages mixOmics

(Rohart *et al.*, 2017) and BioQC (Zhang *et al.*, 2017). A highly connected, central cluster of genes was

identified using hierarchical clustering on each heatmap and the connectivity and entropy of the genes

in these clusters was calculated. Connectivity was calculated for each gene as the sum of shared

connections made between the given gene and all other DEGs in the hypernetwork. Entropy per-gene

was calculated with the same values in a column-by-column manner using BioQC (Zhang *et al.*, 2017)

in R 3.6.0 (R core team, 2019), and scaled to size of the central cluster by taking $\log_2 N^2$, where N is

the length of the gene list.

Having identified differences in the connectivity and entropy of the hypernetwork models of DEG

interactions, we extended this approach to assess the gene network properties across the whole

transcriptome. To do this, we selected a random set of non-differentially expressed genes (N = 160, to

match the size of the DEGs list) and subjected them to the hypernetwork analysis described above.

Entropy and connectivity were calculated on the central cluster genes per iteration, and entropy

scores were then scaled to gene list size by taking $\log_2 N^2$, where N is the length of the gene list. This

hypernetwork approach was iterated 10,000 times to provide a whole-transcriptome assessment of

connectivity and entropy for animals from each treatment group. Wilcoxon tests were used to contrast

entropy and connectivity between the sharks incubated at 15˚C and 20˚C.

2.7 Over-representation analysis on the central cluster of transcriptional activity

Having generated the hypernetwork adjacency matrices from the DEGs, hierarchical clustering was

used and identified a group of highly connected genes present in the 15°C, but not 20°C, incubated

sharks. This refined set of DEGs share the largest number of correlations across the rest of the

transcriptome and as such, the expression patterns of these genes are related. We identified the

highly connected genes present in the cold incubated animals (yellow clusters, fig. 2a, 2c & 2e) and

subtracted the few highly connected genes from the warm-incubated animals (yellow clusters, fig. 2b,

2d & 2f). This final per-species gene list represents the genes whose highly connected nature with the

transcriptome is disrupted by elevated incubated temperature. Over representation analysis was

performed on these genes in DAVID, using each species' full gene list as a reference, to predict

possible biological function (Huang *et al.*, 2009a; Huang *et al.*, 2009b).

3 RESULTS

228 The expression of 160 genes in the ventricle of *S. canicula*, 125 genes from *D. rerio* muscle, and 341

229 from *D. labrax* muscle tissue, were significantly altered by temperature during embryogenesis (*S.*

230 *canicula*: 90 upregulated and 70 downregulated, p-value range = 1x10^{-3} - 3.3x10^{-17}; *Danio rerio*: 112

231   upregulated, 13 downregulated, p-value range = 1×10^{-3} - 7.2×10^{-7}; *Dicentrarchus labrax*: 293

232   upregulated and 48 downregulated, p-value range = 1×10^{-3} – 9.56×10^{-11}).

255   FIGURE 1 The entropy (A & C) and connectivity (B & D) of the differentially expressed genes (A & B)
       and entire transcriptome (C & D) of *Scyliorhinus canicula* (red, N = 3), *Danio rerio* (green, N = 5-8), and
256   *Dicentrarchus labrax* (blue, N = 5) that experienced embryogenesis at different temperatures. Asterisks
       denote statistically significant differences between temperature treatment groups (Wilcoxon test; * p ≤
0.05, ** p ≤ 0.01, *** p ≤ 0.001, **** p ≤ 0.0001).

Hypernetwork analysis was performed on both the DEGs lists identified in each species, and on

10,000 permuted random subsets of the entire transcriptome. Embryogenesis at an elevated

temperature significantly increased the entropy (Figure 1a; *S. canicula* p < 2.2×10^{-16}, *D. rerio* p <

2.2×10^{-16}, *D. labrax* p < 2.2×10^{-16}) of the DEGs in all three species, as well as reducing connectivity in

S. canicula and *D. rerio*, whilst increasing it in *D. labrax* (Figure 1b; *S. canicula* p < 2.2×10^{-16}, *D. rerio*

p = 4.8×10^{-12}, *D. labrax* p = 2.2×10^{-16})

Connectivity was decreased in all three species across the entire transcriptome by an elevated

incubation temperature (Figure 1d; *S. canicula* p < 2.2×10^{-16}, *D. rerio*, p < 2.2×10^{-16}, *D. labrax* p <

2.2×10^{-16}). However, entropy of the entire transcriptome was only increased in *S. canicula* and *D.*

rerio, whilst the *D. labrax* transcriptome displayed a lower entropy following incubation at an elevated

temperature (Figure 1c; *S. canicula* p < 2.2×10^{-16}, *D. rerio*, p < 2.2×10^{-16}, *D. labrax* p < 2.2×10^{-16}).

Scyliorhinus canicula, *D. rerio*, and *D. labrax* all had a clear cluster of highly connected DEGs (yellow

clusters), when raised under control conditions *(*Figure 2a, c, e), which was not present in the animals

raised at an elevated temperature (Figure 2b, d, f). Over-representation analysis of the genes present

in these highly connected clusters highlighted a role for these genes in development and muscle

contractile performance (Tables 2, 3 & 4), amongst other processes.

FIGURE 2 Heatmaps displaying the shared correlations between pairs of differentially expressed genes to the rest of the transcriptome in *Scyliorhinus canicula* (A & B, N = 3), *Danio rerio* (C & D, N = 5-8), and *Dicentrarchus labrax* (E & F, N = 5) that experienced embryogenesis at either control (A, C, E) or elevated temperature (B, D, F) conditions. The colour key, and a histogram of the data, are shown in the insets above each heatmap.

4 DISCUSSION

Previous studies have shown that the thermal environment experienced during embryogenesis can have persistent effects on the physiology, gene expression, and phenotypic plasticity of fishes (Anastasiadi *et al.*, 2021; Beaman *et al.*, 2016; Schnurr *et al.*, 2014; Scott & Johnston, 2012). Here, we show that elevated embryonic temperature also programs lasting changes to the organisation of transcriptional networks in small-spotted catsharks (*S. canicula*), zebrafish (*D. rerio*), and European seabass (*D. labrax*), and suggest that this may play a role in the previously observed phenomena of developmentally programmed, temperature-induced changes in phenotypic plasticity (Beaman *et al.*, 2016). Development at an elevated temperature resulted in differentially expressed genes (DEGs) with a significantly higher entropy, and altered connectivity, in all three species. The properties of the entire transcriptome were also significantly affected by developmental temperature. Transcriptome wide gene connectivity was significantly decreased by an elevated incubation temperature in all three species. Network entropy increased with developmental temperature in *S. canicula* and *D. rerio,* but decreased marginally in *D. labrax*. The magnitude of this change was greatest in *S. canicula*, followed by *D. rerio*, and then *D. labrax*. Interestingly, this order of effect sizes matches the times since exposure, which were 4-5, 8-9, and 34 months in *S. canicula*, *D. rerio*, and *D. labrax*, respectively. Whilst the DEGs identified in each species/tissue type showed little overlap, warming during embryogenesis produced consistent effects on the properties of both the DEG and transcriptome-wide gene networks. Such tissue independent transcriptomic signatures have been demonstrated during growth in mammals (Stevens *et al.*, 2013), and may relate to warming-induced changes in ectotherm early-life growth pattern (Musa *et al.*, 2020; Hume, 2019). These developmentally programmed changes in transcriptional entropy and connectivity will change the response of the transcriptome to future environmental challenges, which may result in altered phenotypic plasticity, and thus may have implications for both physiological and population-level resilience to future warming events.

Changes in the role of the DEGs in the transcriptional network, which manifest as an increase in entropy and changes in connectivity, may consequently alter the ability of these genes and thus the animal to mount a co-ordinated response to future environmental challenges. Recent studies have suggested that the developmental environment influences the capacity to acclimate to environmental changes in later-life (Beaman *et al.*, 2016). For example, intertidal copepods (*Tigriopus californicus*) that undergo embryogenesis at 25°C have the capacity to raise their critical thermal maximum as adults through temperature acclimation (Healy *et al.*, 2019). However, embryos of the same species incubated at 20°C show no acclimation capacity in critical thermal maximum as adults (Healy *et al.*, 2019). Given that phenotypic plasticity is facilitated by co-ordinated changes in gene expression, these developmentally programmed changes in phenotypic plasticity may relate to the changes in transcriptional organisation observed in our study. Further studies support the link between embryonic temperature and later-life acclimation capacity. Mosquitofish (*Gambusia holbrooki*) produce multiple generations per year, with those born in summer experiencing a warm but constant environment, and

those born in spring experiencing cool, but steadily warming, conditions. Recent work has
demonstrated that mosquitofish born in the more thermally variable spring environment have a
greater capacity to acclimate their metabolic processes than mosquitofish from the more thermally
stable summer conditions (Seebacher *et al.*, 2014). Finally, studies of fruit flies (*Drosophila*
melanogaster) have shown that heat tolerance is influenced not just by acclimation temperature, but
by an interaction between acclimation temperature and embryonic temperature, further supporting the
role of the early-life thermal environment in dictating an individual's response to future temperature
challenges (Willot *et al.*, 2021).

Studies of developmental programming resulting from environmental challenges in fish and reptiles
have demonstrated a protective phenotype in later-life, in contrast to the negative effects often
reported in mammals (Galli *et al.*, 2021; Hellgren *et al.*, 2021; Ruhr *et al.*, 2019; Seeabacher *et al.*,
2014). For example, work on the common snapping turtle (*Chelydra serpentina*) revealed that hypoxia
exposure (50% air saturation) throughout embryogenesis improved the anoxia tolerance of the
cardiomyocytes isolated from juveniles (Ruhr *et al.*, 2019). Similarly, jacky dragons (*Amphibolurus*
muricatus) raised with extended basking times (11 vs. 7 hours of daily heat lamp exposure) show a
higher panting threshold than those from control conditions, implying a greater thermal tolerance (So
& Schwanz, 2018). Such protective phenotypes, programmed by environmental challenges during
embryogenesis, may be associated with the altered transcriptional landscape resulting from
environmental stresses during early life. However, exposure to a stressor during embryogenesis does
not always facilitate resilience to that stressor in adulthood. Cuban brown anole (*Anolis sagrei*) eggs
incubated under cool, warm, and hot temperature fluctuations, and then raised in standard conditions
after hatching, show no differences in thermal tolerance as adults (Gunderson *et al.*, 2020). Thus,
whilst developmental exposure to a stressor often influences the capacity to respond to that same
stressor in later-life, it is not always the case.

One mechanism linking the developmental environment to later-life acclimation capacity is through
the activity of DNA-methyltransferases (DNMTs) (Radford, 2018). DNA methylation by DNMTs can
repress gene expression either directly, whereby the methylation prevents the interaction between a
gene and its DNA binding proteins (Watt & Molloy, 1988), or indirectly, through recognition of the
methyl cytosine by methyl cytosine binding proteins, and the consequent recruitment of transcriptional
corepressors (Boyes & Bird, 1991; Klose & Bird, 2006). Through these mechanisms, DNMTs can alter
gene expression. Changes in embryonic temperature have been shown to alter DNA methylation
patterns in fish (Metzger & Schulte, 2017), with a recent study identifying DNMT3a as the mediator of
developmental thermal plasticity in zebrafish (*Danio rerio*) (Loughland *et al.*, 2021). Thus, by
modulating DNMT3a's activity, changes in embryonic temperature can have persistent effects on the
gene expression and plasticity of fishes (Loughland *et al.*, 2021; Metzger & Schulte, 2017). Whilst the
DEGs identified in our study show little overlap between the species/tissue-types (*S. canicula*;
ventricle, table 2; *D. rerio*, hypaxial fast muscle, table 3; and *D. labrax*, muscle, table 4), the observed
changes in entropy and connectivity span the entire transcriptome. This transcriptome-wide
remodelling suggests that a mechanism basal to gene expression, such as DNA/chromatin

methylation, may be facilitating the developmentally programmed changes in gene network co-
ordination.

As well as implications for an organism's plasticity, changes in transcriptional entropy are known to be
a predictor of biological fitness (Zhu *et al.*, 2020). As stress increases, gene expression patterns
become more random and consequently transcriptional entropy increases (Zhu *et al.*, 2020). The
negative relationship between transcriptional entropy and fitness has been demonstrated in seven
species of bacteria, across a range of contexts, and found to be robust (Zhu *et al.,* 2020). Therefore,
within this framework, the increase in entropy caused by developmental exposure to elevated
temperature in fishes could be interpreted as a decline in fitness. Whilst this interpretation is tentative,
given that the negative relationship between entropy and fitness has only presently been
demonstrated in bacteria (Zhu *et al.*, 2020), studies in animals have shown later-life fitness costs of
increased embryonic temperatures. *Drosophila subobscura* incubated at elevated temperatures
throughout juvenilehood show a lower reproductive performance as adults than those incubated in
control conditions, regardless of the temperatures experienced during later-life (Santos *et al.*, 2021).
Therefore, the link between the embryonic environment and later-life fitness is present in ectotherms,
and future studies should probe the potential importance of transcriptional entropy in mediating these
effects.

Climate change is a major threat facing animals. As the world's oceans and rivers continue to warm
(Cai *et al.*, 2014; Oliver *et al.*, 2018), the physiological and population-level stresses exerted upon
fishes will continue to grow. Embryogenesis is a sensitive period in many animal's life histories, and
the conditions experienced during embryonic development influence their growth, physiology, and
behaviour throughout their life. Consequently, to develop a comprehensive appreciation of how fishes
will respond to climate change stressors, it is crucial that we develop an understanding, and
incorporate into predictive models, the role of developmental programming in dictating an organisms'
physiology and capacity to respond to future environmental challenges.

Acknowledgements

We thank Vicky Taylor and the aquarist team in the Biological Services Facility at the University of
Manchester for their assistance in animal husbandry and tissue sampling for the microsatellite analysis.
We thank Mar Pineda and Pierre Delaroche for their dissection skills. We thank the aquarist team of
the Ozeaneum, Stralsund for shark breeding. Finally, we acknowledge the assistance given by
Research IT and the use of the Computational Shared Facility at The University of Manchester.

Competing Interests

We declare no competing interests.

Funding

This work was supported by the Biotechnology and Biological Sciences Research Council Doctoral
Training Studentship to D.M.R, the Higher Education Innovation Fund through the University of
Manchester's Knowledge and Innovation Hub for Environmental Stability to H.A.S, and the pump
priming fund through the University of Manchester's Cardiovascular Division awarded to H.A.S, A.S,
and P.C.

Author Contributions

D.M.R co-ordinated the experiments, analysed the data, and drafted the manuscript. T.G contributed
to the code and analytical method development. S.A.H and A.V performed the microsatellite analysis,
and S.A.H identified the individual animals. B.G and T.M supplied *Scyliorhinus canicula* eggs. H.A.S,
P.C, and A.S conceived the study, secured funding, and reviewed and revised the manuscript. All
authors contributed to the manuscript and gave their approval for publication.

Data Availability

Data are available through NCBI's Gene Expression Omnibus (GEO, Edgar et al., 2002) at the GEO
accession number GSE189976.

TABLE 1 Experimental details and transcriptome assembly statistics for *Scyliorhinus canicula*, *Danio rerio*, and *Dicentrarchus labrax*.

Parameter	Species		
	Scyliorhinus canicula	*Danio rerio*	*Dicentrarchus labrax*
time since exposure	4-5 months	8-9 months	34 months
developmental temperatures	15˚C and 20˚C	27˚C and 32˚C	17.2˚C and 20.8˚C
exposure period	until hatching (~15 weeks)	until hatching (~3 days)	6-63dpf (57 days)
N50	1422	1049	1601
median contig length	363	342	428
BUSCO score	88.8%	78.1%	50.7%
annotated transcripts	19306	18374	12607

TABLE 2 Over-representation analysis of the differentially expressed genes identified in the central cluster of the cold, but not warm, incubated *Scyliorhinus canicula*

Biological Process	P-value
multicellular organism development	0.040
response to tumour necrosis factor	0.074
positive regulation of telomere maintenance via telomerase	0.092

TABLE 3 Over-representation analysis of the differentially expressed genes identified in the central cluster of the cold, but not warm, incubated *Danio rerio.*

Biological Process	P-value
cellular response to UV	0.061
positive regulation of smooth muscle cell proliferation	0.064
negative regulation of apoptotic process	0.069

TABLE 4 Top 10 results from over-representation of the differentially expressed genes identified in the central cluster of the cold, but not warm, incubated *Dicentrarchus labrax.*

Biological Process	P-value
ventricular cardiac muscle tissue morphogenesis	4.4×10^{-21}
muscle filament sliding	1.7×10^{-20}
regulation of the force of heart contraction	2.0×10^{-19}
cardiac muscle contraction	4.6×10^{-18}
transition between fast and slow fiber	2.5×10^{-15}
regulation of heart rate	3.0×10^{-14}
cardiac muscle hypertrophy in response to stress	1.5×10^{-12}
adult heart development	9.8×10^{-12}
skeletal muscle contraction	2.7×10^{-11}
ATP metabolic process	5.4×10^{-11}

Supplementary Figure 1: A schematic qualitatively describing hypernetwork analysis. **1.)** All the differentially expressed genes (DEGs) identified between the two treatment groups are tested for correlations against all other genes identified within the transcriptome. The relationships are binarized as 1s (correlation) or 0s (no correlation) based on a cut-off of one standard deviation. **2.)** The number of correlations to the rest of the transcriptome that are shared between each possible pair of DEGs is calculated. In this, any pair of DEGs that are correlated with many of the same genes will have a high shared correlations value, whilst any pair of DEGs that have no/few shared correlations will have a low shared correlations value. **3.)** DEGs are hierarchically clustered based on the similarity of their shared correlations allowing the identification of associated clusters of genes. This network can then be represented as a heatmap (Figure 2) and the mathematical properties of the network (for instance, connectivity and entropy) can be calculated per treatment group.

References

Anastasiadi, D. & Piferrer, F. (2019). 'Epimutations in developmental genes underlie the onset of
domestication in farmed european sea bass' *Molecular Biology and Evolution*, 36 (10), pp.
2252-2264.

Anastasiadi, D., Shao, C. W., Chen, S. L. & Piferrer, F. (2021). 'Footprints of global change in marine
life: Inferring past environment based on DNA methylation and gene expression
marks', *Molecular Ecology,* 30(3), pp. 747-760.

Banerji, C. R. S., Miranda-Saavedra, D., Severini, S., Widschwendter, M., Enver, T., Zhou, J. X. &
Teschendorff, A. E. (2013). 'Cellular network entropy as the energy potential in Waddington's
differentiation landscape', *Scientific Reports,* 3.

Bateson, P., Gluckman, P. & Hanson, M. (2014). 'The biology of developmental plasticity and the
Predictive Adaptive Response hypothesis', *Journal of Physiology-London,* 592(11), pp. 2357-
2368.

Beaman, J. E., White, C. R. & Seebacher, F. (2016). 'Evolution of Plasticity: Mechanistic Link
between Development and Reversible Acclimation', *Trends in Ecology & Evolution,* 31(3),
pp. 237-249.

Benjamins, S., Cole, G., Naylor, A., Thorburn, J. A. & Dodd, J. 'First confirmed complete incubation
of a flapper skate (Dipturus intermedius) egg in captivity', *Journal of Fish Biology*.

Bolger, A. M., Lohse, M. & Usadel, B. (2014). 'Trimmomatic: a flexible trimmer for Illumina sequence
data', *Bioinformatics,* 30(15), pp. 2114-2120.

Boyes, J. & Bird, A. (1991). 'DNA METHYLATION INHIBITS TRANSCRIPTION INDIRECTLY VIA A
METHYL-CPG BINDING-PROTEIN', *Cell,* 64(6), pp. 1123-1134.

Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. (2016). 'Near-optimal probabilistic RNA-seq
quantification', *Nature Biotechnology,* 34(5), pp. 525-527.

527
Cai OR, Poloczanska ES, Brewer PG, Sundby S, Hilmi K, Fabry VJ, Jung S (2014) The ocean in
climate: change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects. In VR
Barros, CB Field, DJ Dokken, MD Mastrandrea, KJ Mach, TE Bilir, M Chatterjee, KL Ebi, YO
Estrada, RC Genova *et al.*, eds, Contribution of Working Group II to the Fifth Assessment
Report of the Intergovernmental Panel on Climate Change. Cambridge University Press,
Cambridge, United Kingdom and New York, NY, USA, pp. 1655–1731

Edgar, R., Domrachev, M. & Lash, A. E. (2002). 'Gene expression omnibus: Ncbi gene expression
and hybridization array data repository' *Nucleic Acids Research*, 30 (1), pp. 207-210. DOI:
10.1093/nar/30.1.207 Available at: <Go to ISI>://WOS:000173077100055

Fellous, A., Favrel, P. & Riviere, G. (2015). 'Temperature influences histone methylation and mRNA
expression of the Jmj-C histone-demethylase orthologues during the early development of
the oyster Crassostrea gigas', *Marine Genomics,* 19, pp. 23-30.

Galli, G. L. J., Ruhr, I. M., Crossley, J. & Crossley, D. A. (2021). 'The Long-Term Effects of
Developmental Hypoxia on Cardiac Mitochondrial Function in Snapping Turtles', *Frontiers in
Physiology,* 12.

Geurden, I., Mennigen, J., Plagnes-Juan, E., Veron, V., Cerezo, T., Mazurais, D., Zambonino-
Infante, J., Gatesoupe, J., Skiba-Cassy, S. & Panserat, S. (2014). 'High or low dietary
carbohydrate: protein ratios during first-feeding affect glucose metabolism and intestinal
microbiota in juvenile rainbow trout', *Journal of Experimental Biology,* 217(19), pp. 3396-
3406.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan,
L., Raychowdhury, R., Zeng, Q. D., Chen, Z. H., Mauceli, E., Hacohen, N., Gnirke, A., Rhind,
N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N. & Regev, A.
(2011). 'Full-length transcriptome assembly from RNA-Seq data without a reference
genome', *Nature Biotechnology,* 29(7), pp. 644-U130.

Griffiths, A. M., Casane, D., McHugh, M., Wearmouth, V. J., Sims, D. W. & Genner, M. J. (2011).
'Characterisation of polymorphic microsatellite loci in the small-spotted catshark (Scyliorhinus
canicula L.)', *Conservation Genetics Resources,* 3(4), pp. 705-709.

Gunderson AR, Fargevieille A, Warner DA. 2020 Egg incubation temperature does not influence adult
heat tolerance in the lizard Anolis sagrei. Biol. Lett. 16: 20190716.
http://dx.doi.org/10.1098/rsbl.2019.0716

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B.,
Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F.,
Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., Leduc, R. D., Friedman,
N. & Regev, A. (2013). 'De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis', *Nature Protocols,* 8(8), pp. 1494-
1512.

Healy, T. M., Bock, A. K. & Burton, R. S. (2019). 'Variation in developmental temperature alters
adulthood plasticity of thermal tolerance in Tigriopus californicus', *Journal of Experimental
Biology,* 222(22).

Hellgren, K. T., Premanandhan, H., Quinn, C. J., Trafford, A. W. & Galli, G. L. J. (2021). 'Sex-
dependent effects of developmental hypoxia on cardiac mitochondria from adult murine
offspring', *Free Radical Biology and Medicine,* 162, pp. 490-499.

Hook, S. A., McMurray, C., Ripley, D. M., Allen, N., Moritz, T., Grunow, B. & Shiels, H. A. (2019).
'Recognition software successfully aids the identification of individual small-spotted
catsharks Scyliorhinus canicula during their first year of life', *Journal of Fish Biology,* 95(6),
pp. 1465-1470.

Huang, D. W., Sherman, B. T. & Lempicki, R. A. (2009a). 'Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists', *Nucleic Acids
Research,* 37(1), pp. 1-13.

Huang, D. W., Sherman, B. T. & Lempicki, R. A. (2009b). 'Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources', *Nature Protocols,* 4(1), pp. 44-57.

Hume, J. B. (2019). 'Higher temperatures increase developmental rate & reduce body size at
hatching in the small-eyed skate Raja microocellata: implications for exploitation of an
elasmobranch in warming seas', *Journal of Fish Biology,* 95(2), pp. 655-658.

IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-
Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L.
Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T.
Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press.

Klose, R. J. & Bird, A. P. (2006). 'Genomic DNA methylation: the mark and its mediators', *Trends in
Biochemical Sciences,* 31(2), pp. 89-97.

Kopylova, E., Noe, L. & Touzet, H. (2012). 'SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data', *Bioinformatics,* 28(24), pp. 3211-3217.

Langmead, B. & Salzberg, S. L. (2012). 'Fast gapped-read alignment with Bowtie 2', *Nature*
Methods, 9(4), pp. 357-U54.
596
Liu, J. W., Dias, K., Plagnes-Juan, E., Veron, V., Panserat, S. & Marandel, L. (2017). Long-term
programming effect of embryonic hypoxia exposure and high-carbohydrate diet at first feeding
on glucose metabolism in juvenile rainbow trout. Journal of Experimental Biology, 220(20),
3686-3694.

Loughland, I., Little, A. & Seebacher, F. (2021). 'DNA methyltransferase 3a mediates developmental
thermal plasticity', *Bmc Biology,* 19(1).

Marshall, T. C., Slate, J., Kruuk, L. E. B. & Pemberton, J. M. (1998). 'Statistical confidence for
likelihood-based paternity inference in natural populations', *Molecular Ecology,* 7(5), pp. 639-
655.

McCarthy, D. J., Chen, Y. S. & Smyth, G. K. (2012). 'Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation', *Nucleic Acids Research,* 40(10),
pp. 4288-4297.

McLaughlin, R. H. & Ogower, A. K. (1971). 'LIFE-HISTORY AND UNDERWATER STUDIES OF A
HETERODONT SHARK', *Ecological Monographs,* 41(4), pp. 271-+.

Metzger, D. C. H. & Schulte, P. M. (2017). 'Persistent and plastic effects of temperature on DNA
methylation across the genome of threespine stickleback (Gasterosteus
aculeatus)', *Proceedings of the Royal Society B-Biological Sciences,* 284(1864).

Musa, S. M., Czachur, M. V. & Shiels, H. A. (2018). 'Oviparous elasmobranch development inside
the egg case in 7 key stages', *Plos One,* 13(11).

Musa, S. M., Ripley, D. M., Moritz, T. & Shiels, H. A. (2020). 'Ocean warming and hypoxia affect
embryonic growth,fitness and survival of small-spotted catsharks,Scyliorhinus
canicula', *Journal of Fish Biology,* 97(1), pp. 257-264.

Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V.,
Benthuysen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-
Kirkpatrick, S. E., Scannell, H. A., Straub, S. C. & Wernberg, T. (2018). 'Longer and more
frequent marine heatwaves over the past century', *Nature Communications,* 9.
623
R Core Team (2019). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Radford, E. J. (2018). 'Exploring the extent and scope of epigenetic inheritance', *Nature Reviews*
Endocrinology, 14(6), pp. 345-355.

Robinson, M. D., McCarthy, D. J. & Smyth, G. K. (2010). 'edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data', *Bioinformatics,* 26(1), pp.
139-140.

Rohart, F., Gautier, B., Singh, A. & Le Cao, K. A. (2017). 'mixOmics: An R package for 'omics feature
selection and multiple data integration', *Plos Computational Biology,* 13(11).

Rosa, R., Baptista, M., Lopes, V. M., Pegado, M. R., Paula, J. R., Trubenbach, K., Leal, M. C.,
Calado, R. & Repolho, T. (2014). 'Early-life exposure to climate change impairs tropical shark
survival', *Proceedings of the Royal Society B-Biological Sciences,* 281(1793).

Ruhr, I. M., McCourty, H., Bajjig, A., Crossley, D. A., Shiels, H. A. & Galli, G. L. J. (2019).
'Developmental plasticity of cardiac anoxia-tolerance in juvenile common snapping turtles
(Chelydra serpentina)', *Proceedings of the Royal Society B-Biological Sciences,* 286(1905).

Santos, M. A., Carromeu-Santos, A., Quina, A. S., Santos, M., Matos, M. & Simoes, P. (2021). 'High
developmental temperature leads to low reproduction despite adult temperature', *Journal of
Thermal Biology,* 95.

Schaefer, J. & Ryan, A. (2006). 'Developmental plasticity in the thermal tolerance of zebrafish Danio
rerio', *Journal of Fish Biology,* 69(3), pp. 722-734.

Schmieder, R. & Edwards, R. (2011). 'Fast Identification and Removal of Sequence Contamination
from Genomic and Metagenomic Datasets', *Plos One,* 6(3).

Schnurr, M. E., Yin, Y. & Scott, G. R. (2014). 'Temperature during embryonic development has
persistent effects on metabolic enzymes in the muscle of zebrafish', *Journal of Experimental
Biology,* 217(8), pp. 1370-1380.

Scott, G. R. & Johnston, I. A. (2012). 'Temperature during embryonic development has persistent
effects on thermal acclimation capacity in zebrafish', *Proceedings of the National Academy
of Sciences of the United States of America,* 109(35), pp. 14247-14252.

Seebacher, F., Beaman, J. & Little, A. G. (2014). 'Regulation of thermal acclimation varies between
generations of the short-lived mosquitofish that developed in different environmental
conditions', *Functional Ecology,* 28(1), pp. 137-148.

Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. (2015). 'BUSCO:
assessing genome assembly and annotation completeness with single-copy
orthologs', *Bioinformatics,* 31(19), pp. 3210-3212.

So, C. K. J. & Schwanz, L. E. (2018). 'Thermal plasticity due to parental and early-life environments
in the jacky dragon (Amphibolurus muricatus)', *Journal of Experimental Zoology Part a-
Ecological and Integrative Physiology,* 329(6-7), pp. 308-316.

Stevens, A., Hanson, D., Whatmore, A., Destenaves, B., Chatelain, P. & Clayton, P. (2013). 'Human
growth is associated with distinct patterns of gene expression in evolutionarily conserved
networks', *Bmc Genomics,* 14.

Teschendorff, A. E. & Enver, T. (2017). 'Single-cell entropy for accurate estimation of differentiation
potency from a cell's transcriptome', *Nature Communications,* 8.

Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M. & Shipley, P. (2004). 'MICRO-CHECKER:
software for identifying and correcting genotyping errors in microsatellite data', *Molecular
Ecology Notes,* 4(3), pp. 535-538.

Watt, F. & Molloy, P. L. (1988). 'CYTOSINE METHYLATION PREVENTS BINDING TO DNA OF A
HELA-CELL TRANSCRIPTION FACTOR REQUIRED FOR OPTIMAL EXPRESSION OF
THE ADENOVIRUS MAJOR LATE PROMOTER', *Genes & Development,* 2(9), pp. 1136-
1143.

Willot, Q., Ben, L. & Terblanche, J. S. (2021). 'Interactions between developmental and adult
acclimation have distinct consequences for heat tolerance and heat stress recovery', *Journal
of Experimental Biology,* 224(16).

Zhang, J. T. D., Hatje, K., Sturm, G., Broger, C., Ebeling, M., Burtin, M., Terzi, F., Pomposiello, S. I.
& Badi, L. (2017). 'Detect tissue heterogeneity in gene expression data with BioQC', *Bmc
Genomics,* 18.

Zhu, Z. Y., Surujon, D., Ortiz-Marquez, J. C., Huo, W. W., Isberg, R. R., Bento, J. & van Opijnen, T.
(2020). 'Entropy of a bacterial stress response is a generalizable predictor for fitness and
antibiotic sensitivity', *Nature Communications,* 11(1).