Warming During Embryogenesis Induces a Lasting Transcriptomic Signature in Fishes

Daniel M. Ripley[1], Terence Garner[2], Samantha A. Hook[3], Ana Veríssimo [4,5], Bianka Grunow[6], Timo Moritz[7,8], Peter Clayton[2], Holly A. Shiels[1], Adam Stevens[2*]*

[1]Division of Cardiovascular Sciences, Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK.

[2]Division of Developmental Biology and Medicine, Faculty of Biology, Medicine, and Health, The University of Manchester, Manchester, UK.

[3]Department of Earth and Environmental Sciences, The University of Manchester, Manchester, UK.

[4]CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

[5]BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661 Vairão, Portugal

[6]Institute of Muscle Biology and Growth, Research Institute for Farm Animal Biology, Wilhelm-Stahl-Allee 2, 18196 Dummerstorf, Germany.

[7]Deutsches Meeresmuseum, Katharinenberg 14–20, 18439 Stralsund, Germany

[8]Institute of Biological Sciences, University of Rostock, Albert-Einstein-Straße 3, 18059 Rostock, Germany

CORRESPONDING AUTHORS Daniel.ripley@manchester.ac.uk, Adam.stevens@manchester.ac.uk

KEYWORDS climate change, long term effect, gene expression, plasticity, developmental programming, *Scyliorhinus canicula*, RNA-seq.

RUNNING HEADER Warming programs the transcriptome of fishes

Abstract

Exposure to elevated temperatures during embryogenesis has profound acute effects on the cardiac performance, metabolism, and growth of embryonic fishes. Some temperature-induced effects may be retained into, or manifest in, later-life through a mechanism termed developmental programming. In this study, we incubated small-spotted catshark (*Scyliorhinus canicula*) embryos at either 15˚C or 20˚C before transferring the newly hatched sharks to a common set of conditions (15˚C) for 5 months. Lasting transcriptomic differences associated with the developmental environment were identified, and interactions between genes were investigated using network modelling. Development at an elevated temperature caused an increase in transcriptomic entropy, a parameter thought to relate to plasticity. We then validated this observation through a novel re-analysis of published zebrafish (*Danio rerio*), European seabass (*Dicentrarchus labrax*), and three-spined stickleback (*Gasterosteus aculeatus*) datasets and show that developmental temperatures influence the transcriptional response to future thermal challenges by, in part, altering the organisation of gene networks.

1 INTRODUCTION

Developmental programming comprises changes to an organism's phenotype that are induced, or programmed, by the environmental conditions experienced during development (Bateson *et al.*, 2014). These developmental conditions elicit changes in DNA and chromatin methylation patterns, with consequent alterations in gene expression (Anastasiadi *et al.*, 2017, Fellous *et al.*, 2015). Some of these changes in methylation pattern, and their concomitant influence on gene expression, can be retained into later life. For instance, changes in developmental temperature cause hypermethylation in threespined stickleback (Metzger & Schulte, 2017), and studies in rainbow trout (*Oncorhynchus mykiss*) demonstrate the ability of early-life hypoxic events (Liu *et al.*, 2017) and dietary composition (Geurden *et al.*, 2014) to alter glucose metabolism and homeostasis in later-life. Finally, the developmental temperature of zebrafish (*Danio rerio*) has been shown to program their gene expression and physiology, culminating in an enhanced thermal tolerance (Schaefer & Ryan, 2006) and acclimation capacity (Schnurr *et al.*, 2014; Scott & Johnston, 2012) in adulthood.

As well as altering mean trait values, the embryonic environment can also program an organism's ability to acclimate to conditions experienced in later-life (Beaman *et al.*, 2016). That is, the capacity to physiologically respond to and mitigate future environmental challenges can be enhanced, hindered, or simply altered by an individual's embryonic conditions (Beaman *et al.*, 2016). Evidence in zebrafish (*Danio rerio*) (Scott & Johnston, 2012; Schnurr *et al.*, 2014) and mosquitofish (*Gambusia holbrooki*) (Seebacher *et al.*, 2014), amongst other species (Beaman *et al.*, 2016), show that the thermal environment experienced during early-life can have lasting effects on the acclimation capacity of multiple physiological traits, including metabolic scope (Seebacher *et al.*, 2014), critical thermal maximum (Healy *et al.*, 2019), and swimming performance (Scott & Johnston, 2012).

Phenotypic changes arising from physiological plasticity are underwritten by changes in an organism's gene expression in response to environmental cues. Such plasticity can be affected by an individual´s embryonic environment, which may either facilitate or hinder changes in gene expression later in life (Beaman *et al.*, 2016). Thus, there is a link between an organism's phenotype, gene expression, and plastic potential, which can be modulated by its embryonic environment (Beaman *et al.*, 2016). Shannon entropy (herein entropy) of the transcriptome is a metric describing the order or predictability of gene-gene interactions within a transcriptional network. These gene-gene interactions can either be direct (pairwise) or indirect (higher order interactions) (Sanchez, 2019). The more entropic a network, the less structured and less predictable its behaviour. Transcriptional entropy is known to correlate with the differentiation potency and phenotypic plasticity of single cells (Banerji *et al.*, 2013; Teschendorff & Enver, 2017). The more entropic a cell, the greater its number of possible fates (Teschendorff & Enver, 2017), and thus the more plasticity it possesses. Given that several studies have shown that the environmental temperature experienced during development can alter plasticity in the adult organism (Beaman *et al.*, 2016), we aimed to assess whether developmental temperature also alters gene-level plasticity by measuring the network entropy of the transcriptome in later life.

Whilst many animals may experience adverse environmental conditions during embryogenesis, some species, owing to their ecology, are more susceptible than others. Oviparous elasmobranchs,

because of their protracted, sessile developmental period, are susceptible to experiencing sub-

optimal environmental conditions for long durations. Oviparous elasmobranchs develop inside

collagenous egg cases termed mermaid's purses, which the embryo remains within for several

months to over a year, depending on the species (Benjamins *et al.*, 2021; McLaughin & Ogower,

1971). Throughout embryogenesis the embryo is unable to move from the purse, and thus may be

consistently exposed to unfavourable environmental conditions such as elevated temperatures. As

the world's oceans continue to warm (IPCC, 2021, Cai *et al.*, 2014), and the frequency of marine

heatwaves is increasing (Oliver *et al.*, 2018), the likelihood of oviparous elasmobranchs developing in

sub-optimal conditions increases too. Exposure to elevated temperatures is known to affect the

growth and development of elasmobranch embryos (Hume, 2019; Musa *et al.*, 2020; Rosa *et al.*,

2014, Ripley *et al.*, 2021). However, no studies have investigated how embryogenesis at elevated

temperatures may have lasting effects on elasmobranchs through developmental programming.

Here, we incubate *Scyliorhinus canicula* embryos at 15˚C or 20˚C throughout embryogenesis, before

moving them to a common-garden environment (15˚C) upon hatching. Following 5 months in the

common garden environment, we sacrificed the animals and performed RNA-seq on the ventricle

tissue of six individuals to investigate whether the developmental environment could program lasting

changes in the gene expression and organisation of the *S. canicula* ventricular transcriptome.

Ventricle tissue was chosen for sequencing owing to the ventricle's key role in mediating thermal

performance and tolerance (Pörtner and Knust, 2007). Finally, to validate our findings, we re-analysed

published datasets of temperature-induced developmental programming in zebrafish (*Danio rerio)*

hypaxial fast muscle (Scott & Johnston, 2012), European seabass (*Dicentrarchus labrax)* muscle

(Anastasiadi *et al.*, 2021), and threespine stickleback (*Gasterosteus aculeatus*) muscle (Metzger and

Schulte, 2018) to test the consistency of our observations across multiple species of fishes and other

striated muscle tissues.

2 MATERIAL AND METHODS

2.1 Experimental animals

Details of the experimental design can be found in figure 1. *Scyliorhinus canicula* embryos were

collected from a population of 7 randomly mating adult individuals held at 15˚C at the Ozeaneum,

Stralsund, Germany, and transported to the University of Manchester, UK. Upon arrival the health and

developmental stage of each embryo was assessed using the Musa scale (Musa *et al.*, 2018). Only

healthy embryos at stage 1 were used in the study (Musa *et al.*, 2018). All regulated procedures

received approval from the institution's ethical review board and were performed under the Home

Office License P005EFE9F9 held by HAS.

Following embryonic staging, individuals were randomly assigned to a temperature treatment group

(15 ± 0.3˚C or 20 ± 0.3˚C). For the control condition, 15˚C was chosen as it falls within the range of

temperatures experienced by *S. canicula* in the wild and is the holding temperature of the parent

population at the Ozeaneum, Stralsund, Germany. For the treatment group, +5˚C (20˚C) was chosen

as is it represents the increase in ocean temperature predicted by the end of the century, whilst also being within the current range of temperatures experienced by some *S. canicula* populations in the wild (Pegado *et al.*, 2020). The egg cases in each treatment group were hung vertically in two, well aerated, 55 l static seawater (35ppt salinity) tanks equipped with internal filters and left to continue their embryonic development.

Upon hatching, fin clips were taken to facilitate identification in later-life through microsatellite analysis. The fin clips were stored in 98% ethanol at -20˚C, and the sharks were moved into one of four well aerated 400 l static seawater tanks held at 15˚C ± 0.3˚C. The hatchlings from the 20˚C treatment group were lowered to 15˚C at a rate of 2.5˚C per day, and sharks from both treatment groups were mixed and randomly allocated to a tank. The sharks were fed a mixture of squid, crab and krill three times per week. The sharks from both treatment groups were held at 15˚C ± 0.3˚C for an average of 136 days prior to tissue sampling. During the entire experiment, saltwater changes were performed three times weekly to maintain ammonia, nitrite, and nitrate below detectable levels.

2.2 Tissue sampling

Sharks at 4-5 months age (mean ± SEM = 136 ± 6.3days) were euthanised with an overdose of buffered tricane methanesulfonate and a fin clip was taken post-humous. The ventricle was excised from each individual, placed into RNAlater, and frozen in liquid nitrogen before being stored at -80˚C prior to RNA-sequencing.

2.3 Individual identification using microsatellite analysis

To identify which condition an individual was developed in, microsatellite analysis was performed on the fin clips taken at birth and at the end of the experiment to match each individual using the methods described in Hook *et al.* (2019). Fin clips were extracted using a BioLine Isolate Genomic kit with an extended proteinase K digestion to maximise DNA yield. DNA yield was assessed using a NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific, USA) and gel electrophoresis. A one-primer cocktail containing 11 microsatellites and three tail dyes was used for DNA amplification (Griffiths *et al.*, 2011) using the QIAGEN multiplex PCR kit. The thermocycling protocol consisted of initial denaturation cycle at 95°C for 15 minutes, followed by 35 cycles of 94°C for 30 seconds, 60°C for 90 seconds and 72°C for 45 seconds, and finalised by one cycle at 72°C for 30 minutes. The products were visualised on a gel and then genotyped using an ABI sequencer. Positive control samples were added to each plate genotyped to account for possible allele slippage.

Genotypes were scored using GeneMapper© v4.1 (Applied Biosystems) and validated through Microchecker (van Oosterhout *et al.*, 2004). Duplicates were found between the two batches using CERVUS (Marshall *et al.*, 1998). In cases where some alleles within the genotype did not match, controls were used as a reference to identify possible allele slippage. Probabilities of the identity analysis (pID) were taken from CERVUS to confirm match identification between the two batches. As *S. canicula* lay multiple eggs, microsatellite analysis was also used to confirm that no samples used for RNA-sequencing were taken from full siblings.

2.4 *Danio rerio, Dicentrarchus labrax,* and *Gasterosteus aculeatus* re-analysis

The computational analysis pipeline (sections 2.5 and 2.6 below) was also implemented on previously
published RNA-seq datasets of temperature-induced developmental programming in zebrafish (*D.*
rerio) hypaxial fast muscle (Scott & Johnston, 2012), European seabass (*D. labrax*) muscle tissue
(Anastasiadi *et al.*, 2021), and threespine stickleback (*G. aculeatus*) muscle tissue (Metzger and
Schulte, 2018). This re-analysis was performed to assess the robustness and between-species
consistency of any temperature-induced effects observed in *S. canicula*.

The *D. rerio, D. labrax*, and *G. aculeatus* data were processed in an identical manner to the *S.*
canicula data. Developmental conditions and transcriptome assembly statistics for the four species
are shown in table 1. Full details of the animals used in the re-analysis can be found in Scott &
Johnston 2012 (*D. rerio*, N = 4), Anastasiadi *et al.* 2021 (*D. labrax*, N = 5), and Metzger and Schulte
2018 (*G. aculeatus*, N = 6 respectively).

2.5 de-novo transcriptome assembly and differential expression analysis

The ventricle tissue from six individuals (N = 3 per treatment group), stored in RNAlater at -80°C, was
used for generating the transcriptome. The tissue samples were extracted using the Qiagen kit with
homogenisation. Basic sample quality was assessed using a Nanodrop (ThermoFisher) and the high-
quality ventricular RNA samples were sequenced using a 76 base pair long, pair-ended reads on the
Illumina TruSeq system. Reads that mapped to human, bacterial, and viral sequences were removed
using DeconSeq (Schmieder and Edwards, 2011) whilst reads mapping to ribosomal RNA were
removed using sortmeRNA (Kopylova *et al.*, 2012). Adaptor sequences and sequences with a low-
quality score (regions averaging a score <5 over a 4bp sliding window, and leading/trailing sequences
scoring <5) were then removed from the cleaned reads using Trimmomatic (Bolger *et al.*, 2014) prior
to transcriptome assembly. 264 million bases survived the quality control process and were used for
transcriptome assembly in Trinity 2.8.4 (Grabherr *et al.*, 2011; Haas *et al.*, 2013). Default parameters
were used in Trinity, except –normalize_reads, producing 347710 contigs with a N50 of 1422, and a
BUSCO score of 88.8%, with 519 single copy, and 2458 duplicated copy, BUSCOs (BUSCO v5.2.2,
Simao *et al.*, 2015). Open reading frames (ORFs) were predicted from the transcripts using
Transdecoder (https://github.com/TransDecoder/TransDecoder/wiki) with a minimum length threshold
of 100 amino acids. This filtered dataset was then annotated using a BLAST search against the
SwissProt database. The reads were then pseudo-aligned to the curated transcript assemblies using
Kallisto (Bray *et al.*, 2016), allowing the relative abundance of each transcript to be calculated. Where
multiple transcripts mapped to a single BLAST hit, the sum of the transcripts abundances was used.
The abundance estimates were filtered by removing fragments that had zero mapped reads in any of
the samples before differential expression analysis was performed with EdgeR (McCarthy *et al.*, 2012;
Robinson *et al.*, 2010) in R 3.6.0 (R core team, 2019) using the R-script available in Metzger and

Schulte, 2018. Significance levels were adjusted for multiple testing using a false discovery rate
(FDR) correction.

2.6 Entropy analysis

Entropy of both pairwise and higher-order interaction networks was determined in each treatment
group. See figure 2 for a conceptual overview. The expression of each gene was tested for
correlations against all the assembled transcripts within each treatment group. The correlation values
were binarized such that strong positive (top 10% of R-values) or strong negative (bottom 10% of R-
values) correlations were retained (denoted as 1) and weak correlations were discarded (denoted as
0). This binarized correlation matrix is square and represents the pairwise interaction network.

To generate the higher order interaction networks (hypernetworks), the binarized correlation matrix
was multiplied by the transpose of itself ($M \times M^t$) to return the adjacency matrix of the hypernetwork,
where the values in each cell represent the number of shared correlations any given pair of genes
have to the rest of the transcriptome. These values also represent the dimensionality of the
hyperedge connecting each pair of nodes.

To assess whole-transcriptome properties, we selected subsets of 200 genes from the network (either
the binarized network or the hypernetwork) and calculated the entropy of each gene using the R
packages mixOmics (Rohart *et al.*, 2017) and BioQC (Zhang *et al.*, 2017) in R 3.6.0 (R core team,
2019). The mean gene-level entropy was then calculated for each iteration. This approach was
iterated 1000 times for both the binarized and hyper-networks to provide a whole-transcriptome
assessment of both pairwise (binarized network) and higher order (hypernetwork) entropy for animals
from each treatment group. Hypernetwork entropy was then scaled to one by taking $\log_2 N^2$, where N
is the length of the gene list. Wilcoxon tests were used to contrast entropy between the sharks
incubated at 15˚C and 20˚C. This same approach was applied to re-analysed zebrafish, seabass, and
stickleback datasets.
219

2.7 Gene ontology analysis

Gene ontology analysis, leveraging the pairwise and higher-order network entropy, was used to
identify specific gene-pathways whose organisation was influenced by developmental temperature.
Gene pathways (geneontology.org) of interest were selected *a priori* and each treatment group was
analysed separately. Gene networks (both pairwise and higher-order) were created (as described in
section 2.6) from a subset (N = 20) of genes within each gene ontology term, and an entropy score
was calculated. This approach was iterated 1000 times per gene pathway to generate a distribution of
entropy scores for each gene ontology term. A Bayesian approach was used to assess the
differences between groups; entropy distributions for each gene pathway contrasted between
developmental temperatures and β entropy scores were calculated. These β values represent the
difference in entropy between warm and control incubated fish for a given gene pathway. The
posterior of β entropy values were plotted and the 89% credible intervals calculated. Differences in

pathway entropy were defined as significant where the 89% credible interval did not cross 0, per
standard protocol (McElreath, 2018).

2.8 Zebrafish acclimation analysis

A subset of zebrafish from both developmental temperatures in the Scott and Johnston (2012) paper
were cold acclimated (16°C for 28-30 days) in adulthood prior to RNA-sequencing. We re-analysed
these data (transcriptome assembly, read quantification and annotation as described in sections 2.5
and 2.6 above) to look at the effects of developmental temperature on temperature-induced gene
expression changes in later-life. The proportion of genes whose expression changed due to cold
acclimation was compared using Fisher's exact test, and the magnitude of change in the differentially
expressed genes (DEGs) was contrasted with a Wilcoxon test. Finally, we created a logistic
regression model (R package 'lme4' (Bates *et al.*, 2015)) to assess whether a gene's response
(differentially expressed or not, as defined by a false discovery rate corrected p-value (FDR) cut-off of
0.05) to cold-acclimation was influenced by its pre-acclimation entropy in both the pairwise and
higher-order interaction networks. Pairwise network entropy, higher order network entropy,
developmental condition, direction of the change (positive or negative fold change), and all their
interactions were included in the initial logistic regression model. Model simplification was then
performed to remove redundant predictors using stepAIC from the package 'MASS' (Venables and
Ripley, 2002). Predictors retained into the final model were tested for significance using a Wald test
(package 'aod', Lesnoff and Lancelot, 2012), and their effect size calculated (package 'oddsratio',
Schratz, 2020).
252

3 RESULTS

Developmental warming caused the differential expression of 163 (FDR < 0.05) genes in juvenile *S.*
canicula, compared to 66 (FDR < 0.05), 68 (FDR < 0.05), and 21 (FDR < 0.05) in adult *D. labrax*, *D.*
rerio, and *G. aculeatus* respectively (figure 3). The DEGs showed little overlap between species
(figure 3).

Entropy of the pairwise interaction networks was increased due to embryonic warming in all species
(*S. canicula*; figure 4a, p < 0.0001. *D. labrax*; figure 4b, p < 0.0001. *D. rerio*: figure 4c, p < 0.0001. *G.*
aculeatus; figure 4d, p < 0.0001), whilst cooling caused an entropy decrease in *G. aculeatus* (figure
4d*,* p < 0.0001). Gene ontology revealed warming-induced entropy differences in 11/14 (*S. canicula*),
10/14 (*D. labrax*), 13/14 (*D. rerio*) and 7/14 (*G. aculeatus*) of the pathways investigated (figure 5).
Developmental cooling altered the entropy of 14/14 pathways in *G. aculeatus.*

Entropy of the higher-order interaction networks was also increased by developmental warming in *S.*
canicula (figure 6a, p < 0.0001), *D.labrax* (figure 6b, p < 0.0001) and *D. rerio* (figure 6c, p < 0.0001),
but not in *G. aculeatus*, where entropy of the higher-order interactions was highest in the control
animals, and reduced by both developmental warming and cooling (figure 6d, p < 0.0001 & p <
0.0001). Gene ontology revealed warming-induced entropy differences in 13/14 (*S. canicula*), 9/14 (*D.*

labrax), 10/14 (*D. rerio*) and 4/14 (*G. aculeatus*) of the pathways investigated (figure 7).
Developmental cooling altered the entropy of 5/14 pathways in *G. aculeatus.*

D. rerio that were incubated at an elevated temperature showed a greater number (figure 8a, p <
0.0001) and magnitude (figure 8b, p < 0.0001) of gene expression change following 28-30 days cold
acclimation (16˚C) than *D. rerio* developed in control conditions. Of the cold-induced DEGs, 44.1%
were shared across both developmental groups. Genes that were programmed by the developmental
environment were more likely to respond to cold acclimation in adulthood than genes that were not
developmentally programmed (figure 8c, p < 0.0001). The gene-level entropy of both the pairwise (p =
0.031) and higher order (p = 0.044) interaction networks before cold-acclimation were associated with
the probability of a gene responding to cold-acclimation in later-life (pseudo R^2 = 0.091). An increase
in higher order entropy was associated with a reduced likelihood of a gene's expression changing
(one-unit increase in higher order network entropy: odds ratio & 95% CI – 0.816, 0.671-0.996), whilst
an increase in pairwise network entropy was positively associated with the probability of a gene's
expression changing (one-unit increase in pairwise network entropy: odds ratio & 95% CI – 6.344,
1.184-33.699).

4 DISCUSSION

Previous studies have shown that the thermal environment experienced during embryogenesis can
have persistent effects on the physiology, gene expression, and phenotypic plasticity of fishes
(Anastasiadi *et al.*, 2021; Beaman *et al.*, 2016; Schnurr *et al.*, 2014; Scott & Johnston, 2012, Metzger
and Schulte 2017). Here, we document temperature-induced developmental programming for the first
time in elasmobranchs and show that elevated embryonic temperature programs lasting changes to
the organisation of transcriptional networks in small-spotted catshark (*S. canicula*) ventricular tissue.
Furthermore, we demonstrate the consistency of this observation across species and muscle types
through a novel re-analysis of previously published datasets on zebrafish (*D. rerio*), European
seabass (*D. labrax*) and three-spined stickleback (*G. aculeatus*) muscle tissue. Such transcriptomic
signatures, seen here in both ventricle and muscle tissue, begin to suggest a tissue independent
effect such as those previously demonstrated during growth in mammals (Stevens *et al.*, 2013, Lui *et*
al., 2010, Lui *et al.*, 2008). Finally, we suggest that these entropy changes may play a role in the
previously observed phenomena of developmentally programmed, temperature-induced changes in
phenotypic plasticity (Beaman *et al.*, 2016).

Developmental warming programmed few (21-163) differentially expressed genes in later-life, with
little overlap between species (figure 3). However, a consistent increase in the entropy of pairwise
gene interaction networks was observed across each species with increasing developmental
temperature (figure 4). These pairwise interaction networks represent gene-gene correlations, and the
consistent differences in the pairwise network entropy across species gives strong support that
developmental temperature influences the organisation of these simple gene-gene connections in
fishes. Gene ontology analysis of these pairwise networks revealed that pathways involved in
ribosome assembly, response to hypoxia, DNA repair, protein folding, organismal growth, and insulin

receptor signalling were affected by developmental temperature in all four species, with insulin
receptor signalling showing a consistent direction of change (figure 5). The insulin receptor signalling
pathway is a key modulator of growth and metabolism. Environmental temperature is known to have
both immediate (Musa *et al.*, 2020) and delayed (compensatory growth, Mortensen and Damsgard,
1993) effects on growth, which may relate to changes in insulin receptor pathway signalling (Won and
Borski, 2013).

Entropy of the higher order interactions was also affected by developmental warming, increasing in
the catshark, seabass, and zebrafish, but decreasing in the stickleback (figure 6). Only the DNA repair
and protein folding pathways showed warming-induced changes in all four species, with neither
sharing a consistent between-species direction of change (figure 7). These higher order interactions
represent networks of shared connections between gene pairs, and thus capture more nuanced and
complex structure within the transcriptome than the simple gene-gene connections. Given this, it is
perhaps unsurprising that the effects of developmental warming on the structure of these more
complex (hyper)networks are less consistent across species. Whilst it is tempting to speculate on the
reason that the higher order interaction networks of the stickleback differ to those of the catshark,
zebrafish, and seabass, especially given the stickleback's complex evolutionary history (McKinnon
and Rundle, 2002), the data required to disentangle this question does not yet exist. Nevertheless,
these developmentally programmed changes in transcriptional entropy, of both the pairwise and
higher order interaction networks, may change the response of the transcriptome to future
environmental challenges, potentially resulting in altered phenotypic plasticity. Indeed, our zebrafish
re-analysis shows that fish incubated in warmer conditions exhibit increased transcriptional entropy
and a greater number and magnitude of gene-expression changes in response to later-life thermal
challenges (figure 8). Interestingly, genes that were programmed by the developmental environment
(figure 3) were more likely to change in response to later-life cold-acclimation than those not affected
by developmental warming (figure 8c). Thus, there is a degree of consistency in temperature-sensitive
genes throughout ontogeny in zebrafish, which aligns with findings from previous studies on three
spined stickleback (*G. aculeatus*, Metzger and Schulte, 2018). Furthermore, the pre-cold acclimation
entropy of individual genes in both the pairwise and higher order interaction networks was associated
with the likelihood of a gene's expression changing in response to later-life cold acclimation. Whilst
the R^2 of the logistic regression was modest, both the pairwise and higher order entropy had
significant effects on the probability of gene expression change. Furthermore, the odds ratios and
their associated confidence intervals did not cross 1, demonstrating that there is a significant effect of
entropy on the transcriptomic response to temperature acclimation. Whilst this effect may be small
and noisy at the level of individual genes, the cumulative effects across the whole transcriptome may
be marked. Thus, the thermal environment experienced during embryogenesis may influence an
individual's responsiveness to future temperature challenges via changes in entropy of the pairwise
and higher order gene interactions within the transcriptome.

Recent studies have suggested that the developmental environment influences the capacity to
acclimate to environmental changes in later-life (Beaman *et al.*, 2016). For example, intertidal

copepods (*Tigriopus californicus*) that undergo embryogenesis at 25˚C have the capacity to raise their
critical thermal maximum as adults through temperature acclimation (Healy *et al.*, 2019). However,
embryos of the same species incubated at 20˚C show no acclimation capacity in critical thermal
maximum as adults (Healy *et al.*, 2019). Given that phenotypic plasticity is facilitated by co-ordinated
changes in gene expression, these developmentally programmed changes in phenotypic plasticity
may relate to the changes in transcriptional organisation observed in our study. Further studies
support the link between embryonic temperature and later-life acclimation capacity. Mosquitofish
(*Gambusia holbrooki*) produce multiple generations per year, with those born in summer experiencing
a warm but constant environment, and those born in spring experiencing cool, but steadily warming,
conditions. Recent work has demonstrated that mosquitofish born in the more thermally variable
spring environment have a greater capacity to acclimate their metabolic processes than mosquitofish
from the more thermally stable summer conditions (Seebacher *et al.*, 2014). Finally, studies of fruit
flies (*Drosophila melanogaster*) have shown that heat tolerance is influenced not just by acclimation
temperature, but by an interaction between acclimation temperature and embryonic temperature,
further supporting the role of the early-life thermal environment in dictating an individual's response to
future temperature challenges (Willot *et al.*, 2021).

Studies of developmental programming resulting from environmental challenges in fish and reptiles
often show a protective phenotype in later-life, in contrast to the negative effects typically reported in
mammals (Galli *et al.*, 2021; Hellgren *et al.*, 2021; Ruhr *et al.*, 2019; Seeabacher *et al.*, 2014). For
example, work on the common snapping turtle (*Chelydra serpentina*) revealed that hypoxia exposure
(50% air saturation) throughout embryogenesis improved the anoxia tolerance of the cardiomyocytes
isolated from juveniles (Ruhr *et al.*, 2019). Similarly, jacky dragons (*Amphibolurus muricatus*) raised
with extended basking times (11 vs. 7 hours of daily heat lamp exposure) show a higher panting
threshold than those from control conditions, implying a greater thermal tolerance (So & Schwanz,
2018). Such protective phenotypes, programmed by environmental challenges during embryogenesis,
may be associated with the altered transcriptional landscape resulting from environmental stresses
during early life. However, exposure to a stressor during embryogenesis does not always facilitate
resilience to that stressor in adulthood. Cuban brown anole (*Anolis sagrei*) eggs incubated under cool,
warm, and hot temperature fluctuations, and then raised in standard conditions after hatching, show
no differences in thermal tolerance as adults (Gunderson *et al.*, 2020). Thus, whilst developmental
exposure to a stressor often influences the capacity to respond to that same stressor in later-life, it is
not always the case.

One mechanism linking the developmental environment to later-life acclimation capacity is through
the activity of DNA-methyltransferases (DNMTs) (Radford, 2018). DNA methylation by DNMTs can
repress gene expression either directly, whereby the methylation prevents the interaction between a
gene and its DNA binding proteins (Watt & Molloy, 1988), or indirectly, through recognition of the
methyl cytosine by methyl cytosine binding proteins, and the consequent recruitment of transcriptional
corepressors (Boyes & Bird, 1991; Klose & Bird, 2006). Through these mechanisms, DNMTs can alter
gene expression. Changes in embryonic temperature have been shown to alter DNA methylation

patterns in fish (Metzger & Schulte, 2017), with a recent study identifying DNMT3a as the mediator of

developmental thermal plasticity in zebrafish (*Danio rerio*) (Loughland *et al.*, 2021). Thus, by

modulating DNMT3a's activity, changes in embryonic temperature can have persistent effects on the

gene expression and plasticity of fishes (Loughland *et al.*, 2021; Metzger & Schulte, 2017). Whilst the

DEGs identified in our study show little overlap between the species/tissue-types (figure 3), the

observed changes in entropy span the entire transcriptome. This transcriptome-wide remodelling

suggests that a mechanism upstream of gene expression, such as DNA/chromatin methylation, may

be facilitating the developmentally programmed changes in gene network co-ordination.

As well as implications for an organism's plasticity, changes in transcriptional entropy are known to be

a predictor of biological fitness (Zhu *et al.*, 2020). As stress increases, gene expression patterns

become more random and consequently transcriptional entropy increases (Zhu *et al.*, 2020). The

negative relationship between transcriptional entropy and fitness has been demonstrated in seven

species of bacteria, across a range of contexts, and found to be robust (Zhu *et al.,* 2020). Although

the same relationship remains to be tested in fish, the increase in entropy caused by developmental

exposure to elevated temperature could have similar effects on fitness. *Drosophila subobscura*

incubated at elevated temperatures throughout juvenilehood show a lower reproductive performance

as adults than those incubated in control conditions, regardless of the temperatures experienced

during later-life (Santos *et al.*, 2021). Therefore, the link between the embryonic environment and

later-life fitness is present in ectotherms, and future studies should probe the potential importance of

transcriptional entropy in mediating these effects.

Climate change is a major threat facing animals. As the world's oceans and rivers continue to warm

(Cai *et al.*, 2014; Oliver *et al.*, 2018), the physiological and population-level stresses exerted upon

fishes will continue to grow. Embryogenesis is a sensitive period in many animal's life histories, and

the conditions experienced during embryonic development influence their growth, physiology, and

behaviour throughout their life. Consequently, it is crucial that we expand our understanding of the

mechanisms by which the developmental environment can influence an organisms' physiology and

capacity to respond to future environmental challenges.

Acknowledgements

We thank Vicky Taylor and the aquarist team in the Biological Services Facility at the University of

Manchester for their assistance in animal husbandry. We thank Mar Pineda and Pierre Delaroche for

their dissection skills. We thank Dr William Joyce for thoughtful discussion of the project. We thank the

aquarist team of the Ozeaneum, Stralsund for shark breeding. We thank Milton Tan and Lafage for their

stickleback and seabass silhouettes shared, unchanged, through PhyloPic under the license: Creative

Commons — Attribution 3.0 Unported — CC BY 3.0. Finally, we acknowledge the assistance given by

Research IT and the use of the Computational Shared Facility at The University of Manchester.

Competing Interests

We declare no competing interests.

Funding

426   This work was supported by the Biotechnology and Biological Sciences Research Council Doctoral
427   Training Studentship to D.M.R, the Higher Education Innovation Fund through the University of
428   Manchester's Knowledge and Innovation Hub for Environmental Stability to H.A.S, and the pump
429   priming fund through the University of Manchester's Cardiovascular Division awarded to H.A.S, A.S,
430   and P.C.

Author Contributions

432   D.M.R co-ordinated the experiments, analysed the data, and drafted the manuscript. T.G contributed
433   to the code and analytical method development. S.A.H and A.V performed the microsatellite analysis,
434   and S.A.H identified the individual animals. B.G and T.M supplied *Scyliorhinus canicula* eggs. H.A.S,
435   P.C, and A.S conceived the study, secured funding, and reviewed and revised the manuscript. All
436   authors contributed to the manuscript and gave their approval for publication.

Data Availability

438   Data are available through NCBI's Gene Expression Omnibus (GEO, Edgar et al., 2002) at the GEO
439   accession number GSE189976.

References

Anastasiadi, D., Diaz, N. & Piferrer, F. (2017). 'Small ocean temperature increases elicit stage-
dependent changes in DNA methylation and gene expression in a fish, the European sea
bass', *Scientific Reports,* 7.

Anastasiadi, D., Shao, C. W., Chen, S. L. & Piferrer, F. (2021). 'Footprints of global change in marine
life: Inferring past environment based on DNA methylation and gene expression
marks', *Molecular Ecology,* 30(3), pp. 747-760.

Banerji, C. R. S., Miranda-Saavedra, D., Severini, S., Widschwendter, M., Enver, T., Zhou, J. X. &
Teschendorff, A. E. (2013). 'Cellular network entropy as the energy potential in Waddington's
differentiation landscape', *Scientific Reports,* 3.

Bates, D., Machler, M., Bolker, B. M. & Walker, S. C. (2015). 'Fitting Linear Mixed-Effects Models
Using lme4', *Journal of Statistical Software,* 67(1), pp. 1-48.

Bateson, P., Gluckman, P. & Hanson, M. (2014). 'The biology of developmental plasticity and the
Predictive Adaptive Response hypothesis', *Journal of Physiology-London,* 592(11), pp. 2357-
2368.

Beaman, J. E., White, C. R. & Seebacher, F. (2016). 'Evolution of Plasticity: Mechanistic Link
between Development and Reversible Acclimation', *Trends in Ecology & Evolution,* 31(3),
pp. 237-249.

Benjamins, S., Cole, G., Naylor, A., Thorburn, J. A. & Dodd, J. (2021). 'First confirmed complete
incubation of a flapper skate (Dipturus intermedius) egg in captivity', *Journal of Fish Biology.*

Bolger, A. M., Lohse, M. & Usadel, B. (2014). 'Trimmomatic: a flexible trimmer for Illumina sequence
data', *Bioinformatics,* 30(15), pp. 2114-2120.

Boyes, J. & Bird, A. (1991). 'Dna methylation inhibits transcription indirectly via a methyl-cpg binding-
protein', *Cell,* 64(6), pp. 1123-1134.

Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. (2016). 'Near-optimal probabilistic RNA-seq
quantification', *Nature Biotechnology,* 34(5), pp. 525-527.
467
Cai OR, Poloczanska ES, Brewer PG, Sundby S, Hilmi K, Fabry VJ, Jung S (2014) The ocean in
climate: change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects. In VR
Barros, CB Field, DJ Dokken, MD Mastrandrea, KJ Mach, TE Bilir, M Chatterjee, KL Ebi, YO
Estrada, RC Genova *et al.*, eds, Contribution of Working Group II to the Fifth Assessment
Report of the Intergovernmental Panel on Climate Change. Cambridge University Press,
Cambridge, United Kingdom and New York, NY, USA, pp. 1655–1731

Edgar, R., Domrachev, M. & Lash, A. E. (2002). 'Gene expression omnibus: Ncbi gene expression
and hybridization array data repository' *Nucleic Acids Research*, 30 (1), pp. 207-210. DOI:
10.1093/nar/30.1.207 Available at: <Go to ISI>://WOS:000173077100055

Fellous, A., Favrel, P. & Riviere, G. (2015). 'Temperature influences histone methylation and mRNA
expression of the Jmj-C histone-demethylase orthologues during the early development of
the oyster Crassostrea gigas', *Marine Genomics,* 19, pp. 23-30.

Galli, G. L. J., Ruhr, I. M., Crossley, J. & Crossley, D. A. (2021). 'The Long-Term Effects of
Developmental Hypoxia on Cardiac Mitochondrial Function in Snapping Turtles', *Frontiers in*
Physiology, 12.

Geurden, I., Mennigen, J., Plagnes-Juan, E., Veron, V., Cerezo, T., Mazurais, D., Zambonino-
Infante, J., Gatesoupe, J., Skiba-Cassy, S. & Panserat, S. (2014). 'High or low dietary
carbohydrate: protein ratios during first-feeding affect glucose metabolism and intestinal
microbiota in juvenile rainbow trout', *Journal of Experimental Biology,* 217(19), pp. 3396-
3406.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan,
L., Raychowdhury, R., Zeng, Q. D., Chen, Z. H., Mauceli, E., Hacohen, N., Gnirke, A., Rhind,
N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N. & Regev, A.
(2011). 'Full-length transcriptome assembly from RNA-Seq data without a reference
genome', *Nature Biotechnology,* 29(7), pp. 644-U130.

Griffiths, A. M., Casane, D., McHugh, M., Wearmouth, V. J., Sims, D. W. & Genner, M. J. (2011).
'Characterisation of polymorphic microsatellite loci in the small-spotted catshark (Scyliorhinus
canicula L.)', *Conservation Genetics Resources,* 3(4), pp. 705-709.
496
Gunderson AR, Fargevieille A, Warner DA. 2020 Egg incubation temperature does not influence adult
heat tolerance in the lizard Anolis sagrei. Biol. Lett. 16: 20190716.
http://dx.doi.org/10.1098/rsbl.2019.0716

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B.,
Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F.,
Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., Leduc, R. D., Friedman,
N. & Regev, A. (2013). 'De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis', *Nature Protocols,* 8(8), pp. 1494-
1512.

Healy, T. M., Bock, A. K. & Burton, R. S. (2019). 'Variation in developmental temperature alters
adulthood plasticity of thermal tolerance in *Tigriopus californicus*', *Journal of Experimental*
Biology, 222(22).

Hellgren, K. T., Premanandhan, H., Quinn, C. J., Trafford, A. W. & Galli, G. L. J. (2021). 'Sex-
dependent effects of developmental hypoxia on cardiac mitochondria from adult murine
offspring', *Free Radical Biology and Medicine,* 162, pp. 490-499.

Hook, S. A., McMurray, C., Ripley, D. M., Allen, N., Moritz, T., Grunow, B. & Shiels, H. A. (2019).
'Recognition software successfully aids the identification of individual small-spotted
catsharks *Scyliorhinus canicula* during their first year of life', *Journal of Fish Biology,* 95(6),
pp. 1465-1470.

Huang, D. W., Sherman, B. T. & Lempicki, R. A. (2009a). 'Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists', *Nucleic Acids*
Research, 37(1), pp. 1-13.

Huang, D. W., Sherman, B. T. & Lempicki, R. A. (2009b). 'Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources', *Nature Protocols,* 4(1), pp. 44-57.

Hume, J. B. (2019). 'Higher temperatures increase developmental rate & reduce body size at
hatching in the small-eyed skate *Raja microocellata*: implications for exploitation of an
elasmobranch in warming seas', *Journal of Fish Biology,* 95(2), pp. 655-658.
524
IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-
Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L.
Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T.
Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press.

Klose, R. J. & Bird, A. P. (2006). 'Genomic DNA methylation: the mark and its mediators', *Trends in*
Biochemical Sciences, 31(2), pp. 89-97.

Kopylova, E., Noe, L. & Touzet, H. (2012). 'SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data', *Bioinformatics,* 28(24), pp. 3211-3217.

Langmead, B. & Salzberg, S. L. (2012). 'Fast gapped-read alignment with Bowtie 2', *Nature*
Methods, 9(4), pp. 357-U54.

Lesnoff, M., Lancelot, R. (2012). *aod: Analysis of Overdispersed Data*. R package version 1.3.2.

Liu, J. W., Dias, K., Plagnes-Juan, E., Veron, V., Panserat, S. & Marandel, L. (2017). Long-term
programming effect of embryonic hypoxia exposure and high-carbohydrate diet at first feeding
on glucose metabolism in juvenile rainbow trout. *Journal of Experimental Biology*, 220(20),
3686-3694.

Loughland, I., Little, A. & Seebacher, F. (2021). 'DNA methyltransferase 3a mediates developmental
thermal plasticity', *Bmc Biology,* 19(1).

Lui, J. C., Chen, W. P., Barnes, K. M. & Baron, J. (2010). 'Changes in gene expression associated
with aging commonly originate during juvenile growth', *Mechanisms of Ageing and*
Development, 131(10), pp. 641-649.

Lui, J. C., Finkielstain, G. P., Barnes, K. M. & Baron, J. (2008). 'An imprinted gene network that
controls mammalian somatic growth is down-regulated during postnatal growth deceleration
in multiple organs', *American Journal of Physiology-Regulatory Integrative and Comparative*
Physiology, 295(1), pp. R189-R196.

Marshall, T. C., Slate, J., Kruuk, L. E. B. & Pemberton, J. M. (1998). 'Statistical confidence for
likelihood-based paternity inference in natural populations', *Molecular Ecology,* 7(5), pp. 639-
655.

McCarthy, D. J., Chen, Y. S. & Smyth, G. K. (2012). 'Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation', *Nucleic Acids Research,* 40(10),
pp. 4288-4297.

McElreath, R. (2018). "Statistical Rethinking : A Bayesian Course with Examples in R and Stan,"
Statistical Rethinking: A Bayesian Course with Examples in R and Stan, pp. 1–469.

McLaughlin, R. H. & Ogower, A. K. (1971). 'Life-history and underwater studies of a heterodont
shark', *Ecological Monographs,* 41(4), pp. 271.

McKinnon, J. S. & Rundle, H. D. (2002). 'Speciation in nature: The threespine stickleback model
systems' *Trends in Ecology & Evolution*, 17 (10), pp. 480-488.

Metzger, D. C. H. & Schulte, P. M. (2017). 'Persistent and plastic effects of temperature on DNA
methylation across the genome of threespine stickleback (*Gasterosteus
aculeatus*)', *Proceedings of the Royal Society B-Biological Sciences,* 284(1864).

Metzger, D. C. H. & Schulte, P. M. (2018). 'Similarities in temperature-dependent gene expression
plasticity across timescales in threespine stickleback (*Gasterosteus aculeatus*)', *Molecular
Ecology,* 27(10), pp. 2381-2396.

Mortensen, A. & Damsgard, B. (1993). 'Compensatory growth and weight segregation following light
and temperature manipulation of juvenile atlantic salmon (*Salmo-salar*) and arctic charr
(*Salvelinus-alpinus*)', *Aquaculture,* 114(3-4), pp. 261-272.

Musa, S. M., Czachur, M. V. & Shiels, H. A. (2018). 'Oviparous elasmobranch development inside
the egg case in 7 key stages', *Plos One,* 13(11).

Musa, S. M., Ripley, D. M., Moritz, T. & Shiels, H. A. (2020). 'Ocean warming and hypoxia affect
embryonic growth,fitness and survival of small-spotted catsharks, *Scyliorhinus
canicula*', *Journal of Fish Biology,* 97(1), pp. 257-264.

Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V.,
Benthuysen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-
Kirkpatrick, S. E., Scannell, H. A., Straub, S. C. & Wernberg, T. (2018). 'Longer and more
frequent marine heatwaves over the past century', *Nature Communications,* 9.

Pegado, M. R., Santos, C. P., Raffoul, D., Konieczna, M., Sampaio, E., Maulvault, A. L., Diniz, M. &
Rosa, R. (2020). 'Impact of a simulated marine heatwave in the hematological profile of a
temperate shark (*Scyliorhinus canicula*)', *Ecological Indicators,* 114.

Portner, H. O. & Knust, R. (2007). 'Climate change affects marine fishes through the oxygen
limitation of thermal tolerance' *Science*, 315 (5808), pp. 95-97.

R Core Team (2019). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Radford, E. J. (2018). 'Exploring the extent and scope of epigenetic inheritance', *Nature Reviews
Endocrinology,* 14(6), pp. 345-355.

Ripley, D. M., et al. (2021). 'Ocean warming impairs the predator avoidance behaviour of
elasmobranch embryos' *Conservation Physiology*, 9 pp. 1-7.

Robinson, M. D., McCarthy, D. J. & Smyth, G. K. (2010). 'edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data', *Bioinformatics,* 26(1), pp.
139-140.

Rohart, F., Gautier, B., Singh, A. & Le Cao, K. A. (2017). 'mixOmics: An R package for 'omics feature
selection and multiple data integration', *Plos Computational Biology,* 13(11).

Rosa, R., Baptista, M., Lopes, V. M., Pegado, M. R., Paula, J. R., Trubenbach, K., Leal, M. C.,
Calado, R. & Repolho, T. (2014). 'Early-life exposure to climate change impairs tropical shark
survival', *Proceedings of the Royal Society B-Biological Sciences,* 281(1793).

Ruhr, I. M., McCourty, H., Bajjig, A., Crossley, D. A., Shiels, H. A. & Galli, G. L. J. (2019).
'Developmental plasticity of cardiac anoxia-tolerance in juvenile common snapping turtles
(*Chelydra serpentina*)', *Proceedings of the Royal Society B-Biological Sciences,* 286(1905).

Sanchez, A. (2019). 'Defining higher-order interactions in synthetic ecology: Lessons from physics
and quantitative genetics' *Cell Systems*, 9 (6), pp. 519-520.

Santos, M. A., Carromeu-Santos, A., Quina, A. S., Santos, M., Matos, M. & Simoes, P. (2021). 'High
developmental temperature leads to low reproduction despite adult temperature', *Journal of
Thermal Biology,* 95.

Schaefer, J. & Ryan, A. (2006). 'Developmental plasticity in the thermal tolerance of zebrafish *Danio
rerio*', *Journal of Fish Biology,* 69(3), pp. 722-734.

Schmieder, R. & Edwards, R. (2011). 'Fast Identification and Removal of Sequence Contamination
from Genomic and Metagenomic Datasets', *Plos One,* 6(3).

Schnurr, M. E., Yin, Y. & Scott, G. R. (2014). 'Temperature during embryonic development has
persistent effects on metabolic enzymes in the muscle of zebrafish', *Journal of Experimental
Biology,* 217(8), pp. 1370-1380.

Schratz, P. (2020). package 'oddsratio'.

Scott, G. R. & Johnston, I. A. (2012). 'Temperature during embryonic development has persistent
effects on thermal acclimation capacity in zebrafish', *Proceedings of the National Academy
of Sciences of the United States of America,* 109(35), pp. 14247-14252.

Seebacher, F., Beaman, J. & Little, A. G. (2014). 'Regulation of thermal acclimation varies between
generations of the short-lived mosquitofish that developed in different environmental
conditions', *Functional Ecology,* 28(1), pp. 137-148.

Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. (2015). 'BUSCO:
assessing genome assembly and annotation completeness with single-copy
orthologs', *Bioinformatics,* 31(19), pp. 3210-3212.

So, C. K. J. & Schwanz, L. E. (2018). 'Thermal plasticity due to parental and early-life environments
in the jacky dragon (*Amphibolurus muricatus*)', *Journal of Experimental Zoology Part a-
Ecological and Integrative Physiology,* 329(6-7), pp. 308-316.

Stevens, A., Hanson, D., Whatmore, A., Destenaves, B., Chatelain, P. & Clayton, P. (2013). 'Human
growth is associated with distinct patterns of gene expression in evolutionarily conserved
networks', *Bmc Genomics,* 14.

Teschendorff, A. E. & Enver, T. (2017). 'Single-cell entropy for accurate estimation of differentiation
potency from a cell's transcriptome', *Nature Communications,* 8.

Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M. & Shipley, P. (2004). 'MICRO-CHECKER:
software for identifying and correcting genotyping errors in microsatellite data', *Molecular*
Ecology Notes, 4(3), pp. 535-538.
637
Venables, W. N. & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth Edition. Springer,
New York. ISBN 0-387-95457-0
Watt, F. & Molloy, P. L. (1988). 'Cytosine methylation prevents binding to dna of a hela-cell
transcription factor required for optimal expression of the adenovirus major late
promoter', *Genes & Development,* 2(9), pp. 1136-1143.

Willot, Q., Ben, L. & Terblanche, J. S. (2021). 'Interactions between developmental and adult
acclimation have distinct consequences for heat tolerance and heat stress recovery', *Journal*
of Experimental Biology, 224(16).
646
Won, E .T., Borski, R. J. (2013). 'Endocrine regulation of compensatory growth in fish', *Front in*
Endocrinology, 4(74).

Zhang, J. T. D., Hatje, K., Sturm, G., Broger, C., Ebeling, M., Burtin, M., Terzi, F., Pomposiello, S. I.
& Badi, L. (2017). 'Detect tissue heterogeneity in gene expression data with BioQC', *Bmc*
Genomics, 18.

Zhu, Z. Y., Surujon, D., Ortiz-Marquez, J. C., Huo, W. W., Isberg, R. R., Bento, J. & van Opijnen, T.
(2020). 'Entropy of a bacterial stress response is a generalizable predictor for fitness and
antibiotic sensitivity', *Nature Communications,* 11(1).

Figure 1: An overview of the experimental design for the RNA-seq data collected from **A)** *Scyliorhinus canicula*, and data re-analysed from **B)** *Dicentrarchus labrax* (Anastasiadi *et al.*, 2021), **C)** *Gasterosteus aculeatus* (Metzger and Schulte, 2018), and **D)** *Danio rerio* (Scott and Johnston, 2012).

TABLE 1 Experimental details and transcriptome assembly statistics for *Scyliorhinus canicula*, *Danio rerio* (Scott and Johnston 2012), *Dicentrarchus labrax* (Anastasiadi *et al.*, 2021) and *Gasterosteus aculeatus* (Metzger and Schulte, 2017).

Parameter	Species			
	Scyliorhinus canicula	*Danio rerio*	*Dicentrarchus labrax*	*Gasterosteus aculeatus*
time since exposure	4-5 months	8-9 months	34 months	~9 months
developmental temperatures	15˚C and 20˚C	27˚C and 32˚C	17.2˚C and 20.8˚C	12˚C, 18˚C and 24˚C
exposure period	until hatching (~15 weeks)	until hatching (~3 days)	6-63dpf (57 days)	until hatching (5-23 days)
life stage at sampling	juvenile	adult	adult	adult
tissue sampled	ventricle	hypaxial fast muscle	muscle	muscle
N50	1422	916	1601	459
median contig length	363	325	428	338

A
Pairwise Interaction Networks

	F	G	H	I	J
F	1	0	1	0	1
G	0	1	1	1	1
H	1	1	1	0	1
I	0	1	0	1	0
J	1	1	1	0	1

B
Higher Order Interaction Networks

Peripheral genes ⬤

	F	G	H	I	J
F	2	2	1	0	1
G	2	3	1	0	1
H	1	1	3	1	2
I	0	0	1	1	1
J	1	1	2	1	3

C

Low Entropy Network

- few interconnections
- specific pattern of associations

High Entropy Network

- diverse interconnections
- robust to perturbation

Figure 2: An example **A)** pairwise gene interaction network. Correlations between pairs of focal genes are shown by a solid black line, and denoted by a 1 in the associated matrix. Pairwise gene network entropy is calculated from the pairwise interaction matrix and represents the networks' information content (complexity). **B)** an example higher order interaction network. Focal genes (the two being assessed for their interaction) are associated via their shared connections to peripheral (not in the focal network) genes, denoted by the grey lines. An interaction between two focal genes can occur via one (gene H – gene I) or multiple (gene I – gene J) shared connections to the peripheral genes. The adjacency matrix denotes the number of shared connections to peripheral genes each pair of focal genes has. Higher order network entropy is calculated from this adjacency matrix and represents the networks' information content (complexity). **C)** an example low and high entropy network.

Figure 3: Overlap of genes in *Scyliorhinus canicula* (red), *Dicentrarchus labrax* (blue), *Danio rerio* (green), and *Gasterosteus aculeatus* (yellow) that are differentially expressed (FDR < 0.05) in later-life due to an increased embryonic temperature. *D. labrax, D. rerio, and G. aculeatus* data are re-analysed from Anastasiadi *et al.*, 2021, Scott and Johnston 2012, and Metzger and Schulte 2017 respectively.

Figure 4: Entropy of pairwise gene interaction networks in **A)** *Scyliorhinus canicula*, **B)** *Dicentrarchus labrax,* **C)** *Danio rerio*, and **D)** *Gasterosteus aculeatus* that underwent embryogenesis at different temperatures. *Scyliorhinus canicula*; Wilcoxon test, p < 0.0001, N = 3 per group. *Dicentrarchus labrax*; Wilcoxon test, p < 0.0001, N = 5 per group. *Danio rerio*; Wilcoxon test, p < 0.0001, N = 4 per group. *Gasterosteus aculeatus*; Kruskal-wallis test and Dunn's multiple comparisons test, p < 0.0001 (cold vs. control) and p < 0.0001 (warm vs. control), N = 6 per group. *D. labrax, D. rerio, and G. aculeatus* data are re-analysed from Anastasiadi *et al.*, 2021, Scott and Johnston 2012, and Metzger and Schulte 2017 respectively.

Figure 5: Median and 89% credibility intervals for pairwise network entropy of gene ontology pathways between control and warm incubated **A)** *Scyliorhinus canicula* (N = 3), **B)** *Dicentrarchus labrax* (N = 5)*,* **C)** *Danio rerio* (N = 4), and **D)** *Gasterosteus aculeatus* (N = 6). β entropy represents the difference in entropy between warm and control incubated animals. Pathways that are significantly altered by developmental temperature in all species are highlighted in bold. *D. labrax, D. rerio, and G. aculeatus* data are re-analysed from Anastasiadi *et al*., 2021, Scott and Johnston 2012, and Metzger and Schulte 2017 respectively.

Figure 6: Entropy of higher-order gene interaction networks in **A)** *Scyliorhinus canicula*, **B)** *Dicentrarchus labrax,* **C)** *Danio rerio*, and **D)** *Gasterosteus aculeatus* that underwent embryogenesis at different temperatures. *Scyliorhinus canicula*; Wilcoxon test, p < 0.0001, N = 3 per group. *Dicentrarchus labrax*; Wilcoxon test, p < 0.0001, N = 5 per group. *Danio rerio*; Wilcoxon test, p < 0.0001, N = 4 per group. *Gasterosteus aculeatus*; Kruskal-wallis test and Dunn's multiple comparisons test, p < 0.0001 (cold vs. control) and p < 0.0001 (warm vs. control), N = 6 per group. *D. labrax, D. rerio, and G. aculeatus* data are re-analysed from Anastasiadi *et al.*, 2021, Scott and Johnston 2012, and Metzger and Schulte 2017 respectively.

Figure 7: Median and 89% credibility intervals for hypernetwork entropy of gene ontology pathways between control and warm incubated **A)** *Scyliorhinus canicula* (N = 3), **B)** *Dicentrarchus labrax* (N = 5)*,* **C)** *Danio rerio* (N = 4), and **D)** *Gasterosteus aculeatus* (N = 6). β entropy represents the difference in entropy between warm and control incubated animals. Pathways that are significantly altered by developmental temperature in all species are highlighted in bold. *D. labrax, D. rerio, and G. aculeatus* data are re-analysed from Anastasiadi *et al*., 2021, Scott and Johnston 2012, and Metzger and Schulte 2017 respectively.

Figure 8: The cold-acclimation response of zebrafish (*Danio rerio*) that underwent embryogenesis in either control or warmed conditions. **A)** median ± 95% CI of the modulus fold change of the differentially expressed genes (Wilcoxon test, p < 0.0001, N = 4 per group), **B)** the number and proportion of gene expression changes (Fisher's exact test, p < 0.0001, N = 4 per group), and **C)** the similarity of gene expression changes between developmental groups (Fisher's exact test, p < 0.0001, N = 4 per group). *D. rerio* data are re-analysed from Scott and Johnston 2012.