1 Rapid bacterial and fungal successional dynamics in first year after Chaparral wildfire
2
3

4 M. Fabiola Pulido-Chavez[1,*], James W. J. Randolph[1], Cassandra Zalman[2], Loralee Larios[3], Peter
5 M. Homyak[4], and Sydney I. Glassman[1,*]
6
[1]Department of Microbiology and Plant Pathology, University of California-Riverside, 900
University Ave, Riverside, CA, 92521
[2]Schmid College of Science and Technology, Chapman University, Orange, CA, 92866
[3]Department of Botany and Plant Sciences, University of California-Riverside, 900 University
Ave, Riverside, CA, 92521
[4]Department of Environmental Sciences, University of California-Riverside, 900 University Ave,
Riverside, CA, 92521
14
15 *Corresponding authors: Sydney I Glassman, sydneyg@ucr.edu and M. Fabiola Pulido-Chavez,
16 mpuli011@ucr.ed Department of Microbiology and Plant Pathology, University of California-
17 Riverside

18
19 Keywords: Secondary succession, Wildfire, Temporal turnover, pyrophilous microbes, Illumina
20 MiSeq, Chaparral

Abstract

The rise in wildfire frequency and severity across the globe has increased interest in

secondary succession. However, despite the role of soil microbial communities in controlling

biogeochemical cycling and their role in the regeneration of post-fire vegetation, the lack of

measurements immediately post-fire and at high temporal resolution has limited understanding of

microbial secondary succession. To fill this knowledge gap, we sampled soils at 17, 25, 34, 67,

95, 131, 187, 286, and 376 days after a southern California wildfire in fire-adapted chaparral

shrublands. We assessed bacterial and fungal biomass with qPCR of 16S and 18S and richness

and composition with Illumina MiSeq sequencing of 16S and ITS2 amplicons. Fire severely

reduced bacterial biomass by 47%, bacterial richness by 46%, fungal biomass by 86%, and fungal

richness by 68%. The burned bacterial and fungal communities experienced rapid succession,

with 5-6 compositional turnover periods. Analogous to plants, turnover was driven by "fire-

loving" pyrophilous microbes, many of which have been previously found in forests worldwide

and changed markedly in abundance over time. Fungal secondary succession was initiated by the

Basidiomycete yeast *Geminibasidium*, which traded off against the filamentous Ascomycetes

Pyronema, *Aspergillus*, and *Penicillium*. For bacteria, the Proteobacteria *Massilia* dominated all

year, but the Firmicute *Bacillus* and Proteobacteria *Noviherbaspirillum* increased in abundance

over time. Our high-resolution temporal sampling allowed us to capture post-fire microbial

secondary successional dynamics and suggest that putative tradeoffs in thermotolerance,

colonization, and competition among dominant pyrophilous microbes control microbial

succession with possible implications for ecosystem function.

1. Introduction

The rapid increase in wildfire frequency, severity, and extent in the western United States

(Riley & Loehman, 2016) and around the globe (Abatzoglou et al., 2019) has renewed interest in

secondary succession. Secondary succession, or the trajectory along which an ecosystem

develops following a disturbance, such as a wildfire, has been extensively studied for plants

(Derroire et al., 2016; Donato et al., 2012), but belowground microbial communities have been

comparatively overlooked. Understanding how wildfires alter soil microbial succession may be

necessary to predict post-fire effects on ecosystem recovery, and function since soil microbes

drive post-fire organic matter decomposition (Semenova-Nelsen et al., 2019), nutrient cycling

(Pérez-Valera et al., 2020), and plant regeneration (Dove & Hart, 2017).

Plant succession is a major process affecting the health and function of ecosystems.

During succession, the dominant species may change in an orderly and predictable manner

(Shugart, 2013). For example, early colonizers (fast-growing, short-lived species) are often

replaced by late colonizers (slow-growing, long-lived species). However, wildfire can reset the

successional clock, shifting the course away from stability (Reilly & Spies, 2016) and initiating

secondary succession. Plant secondary succession is often contingent on the surviving vegetation

and seedbanks present in the heterogeneous post-fire landscape (Jain et al., 2008). While early

plant establishment often happens in low competition and nutrient-rich environments (Dalling,

2008), succession is often mediated by the quantity and identity of early colonizers (i.e., dispersal

limitations and priority effects (Kennedy et al., 2009)), and their tradeoffs for space and resources

(Tilman, 1990). Indeed, the trajectory of species replacement suggests that early colonizers with

similar resource use (i.e., overlapping niches) will dominate open space but will inevitably be

replaced by late-stage species with differentiation in resource use (i.e., complementarity),

allowing for species coexistence (Pacala et al., 1996; Turnbull et al., 2013). This later stage

community is often considered stable, characterized by small fluctuations in community

composition. However, whether the patterns recognized in plant successional theory translate to

belowground soil microbes remains unclear.

Research on post-fire soil microbiomes suggests that fires can reset microbial successional

trajectories via fire-induced mortality (Hart et al., 2005), changes in microbial richness and

biomass (Dooley & Treseder, 2012; Pressler et al., 2019), and the replacement of fungal

basidiomycetes and symbiotic mycorrhizal fungi with ascomycetes and saprobic fungi (Cairney

& Bastias, 2007; Fox et al., 2022). For over a century, pyrophilous or "fire-loving" fungi have

been consistently found in post-fire mushroom surveys (McMullan-Fisher et al., 2011; Seaver,

1909). More recently, next-generation sequencing indicates that the pyrophilous Ascomycete

Pyronema can increase 100-fold after prescribed fires (Reazin et al., 2016) and dominate over

60% of the sequences in experimental pyrocosms (Bruns et al., 2020). Furthermore, there is

increasing evidence of pyrophilous bacteria, such as the Proteobacteria *Massilia* (Enright et al.,

2022; Whitman et al., 2019). This evidence suggests that pyrophilous microbes in systems that

have evolved with wildfire (Dove et al., 2021) might have fire adaptations analogous to plants

(Rundel, 2018) and, thus, likely follow successional dynamics akin to plants. Pyrophilous

microbes have traits that allow them to survive fires (e.g., heat resistant spores, sclerotia) (Day et

al., 2020; Petersen, 1970) and the post-fire environment (e.g., xerotolerance, affinity for nitrogen

mineralization, and affinity for aromatic hydrocarbon degradation) (Fischer et al., 2021; Nelson et

al., 2022; Steindorff et al., 2021). Moreover, since heat from fire often penetrates only the top few

cm of soil (Neary et al., 1999; Pingree & Kobziar, 2019), like plants, secondary succession may

be initiated by surviving microbes that make up the spore bank (Baar et al., 1999; Glassman et al.,

2016). Currently, most research evaluating post-fire microbiomes is based on single timepoint

sampling (Dove & Hart, 2017; Pressler et al., 2019); thus, the succession of pyrophilous microbes

is nearly unknown. However, recent research consisting of 2-3 sampling time points suggests that

bacteria and fungi experience rapid post-fire community changes (Ferrenberg et al., 2013; Qin &

Liu, 2021; Whitman et al., 2022), indicating that higher temporal resolution sampling is needed to

understand bacterial and fungal successional trajectories.

113   Predicting soil microbial succession can be complicated by direct and indirect wildfire

effects (Neary et al., 1999). Whereas soil burn severity controls direct fire impacts (Reazin et al.,

2016; Whitman et al., 2019), changes in soil moisture can indirectly impact post-fire microbes

(Placella et al., 2012; Yang et al., 2021), potentially driving succession, especially in arid

environments where precipitation is limited. Previous research has established that bacteria

rapidly respond to soil wet-up (Barnard et al., 2013; Placella et al., 2012), whereas fungi are less

responsive to soil moisture changes (Barnard et al., 2013; Evans & Wallenstein, 2012).

Furthermore, microbial life strategies can determine microbial response to fire. Research shows

that fire decreases the richness of ectomycorrhizal fungi (EMF) (Cowan et al., 2016; Glassman et

al., 2016; Pulido-Chavez et al., 2021) and arbuscular fungi (AMF) (Xiang et al., 2015) but can

temporarily increase saprobic richness (Enright et al., 2022; Semenova-Nelsen et al., 2019). Yet,

the limited studies on how fires affect multiple microbial guilds (Certini et al 2021) lack the

resolution to identify the direct effect of wildfire and the length of time that mycorrhizal fungi

survive after a wildfire, a question essential for post-fire ecosystem recovery.

We propose that microbial ecological succession theory can be developed and tested in

nature by focusing on California chaparral. Chaparral is a shrubland adapted to high-severity fire

and a biodiversity hotspot distributed in Mediterranean climates worldwide (Barro & Conard,

1991; Rundel, 2018). Chaparral plant secondary succession is relatively well understood and

typically initiated by fire (Keeley et al., 2005). Yet, although the dominant chaparral vegetation

forms mycorrhizal associations required for their establishment and survival (Allen et al., 2005),

little is known about chaparral microbial succession. Indeed, only 13% of post-fire microbiome

research occurs in shrublands (Pressler et al., 2019). With drylands covering nearly 41% of

Earth's land surface and expanding with climate change (Feng & Fu, 2013), understanding

secondary successional dynamics in dryland systems such as chaparral is critical (Osborne et al.,

2022). While our focus is on chaparral, which is likely to have fire-adapted microbes due to

chaparrals' evolutionary history with fire (Hanes, 1971), we expect that successional patterns will

be broadly generalizable to other biomes since pyrophilous bacteria and fungi appear to be

phylogenetically conserved (Enright et al., 2022). Indeed, fungal taxa within the Ascomycota

family Pyronemataceae and a few Basidiomycota genera appear to be globally distributed in fire-

disturbed Spanish shrublands (Pérez-Valera et al., 2018) and Pinaceae (Reazin et al., 2016;

Whitman et al., 2019; Xiang et al., 2014), Eucalyptus (Ammitzboll et al., 2022; McMullan-Fisher

et al., 2011), and redwood-tanoak forests (Enright et al., 2022).

Here, we performed the highest resolution temporal sampling of post-fire microbiomes to

date. We sampled soils at nine timepoints over one year following a chaparral wildfire, allowing

us to identify immediate and temporal effects of wildfire on microbial successional dynamics and

test the following hypotheses: (H1) wildfire will decrease bacterial and fungal biomass and

richness, leading to a shift in the community composition one year post-fire; (H2) wildfire will

have a distinct impact on fungal guilds with symbiotic mycorrhizal fungi experiencing the largest

declines; (H3) higher soil burn severity will lead to larger reductions in both bacteria and fungi

while precipitation will impose a larger impact on bacterial than fungal succession; (H4)

succession will be initiated by pyrophilous microbes, which like plants, will change in abundance

over time based on differences in ecophysiological traits such as growth and nutrient acquisition.

2. Methods

2.1 Study Area, plot design, and soil collection

The Holy Fire burned 94 km^2 in the Cleveland National Forest in Southern California

from August 6 to September 13, 2018. On September 30, 2018, we selected nine plots (6 burned

and 3 unburned) (Fig. 1A). Plots were selected for similarity in aspect, slope, elevation, and pre-

fire vegetation dominance by manzanita (Arctostaphylos glandulosa), an ectomycorrhizal host,

and chamise (Adenostoma fasciculatum), an arbuscular and ectomycorrhizal host (Allen et al.,

2005). Plots were placed on average 25 m from forest access roads (10-40 m) to avoid edge

effects, and each contained four 1 m^2 subplots located 5 m from the center in each cardinal

direction (Fig. 1B). Our study site experiences a Mediterranean-type climate with hot, dry

summers and cool, wet winters, with an average yearly temperature of 17°C and total

precipitation of 668 mm (average 51.39 mm). Precipitation data, based on monthly summaries,

was gathered from the El Cariso weather station (raws.dri.edu). Soils are mapped in the Cieneba

and Friant series and are classified as Typic Xerorthents and Lithic Haploxerolls. They are sandy

and gravelly loams with an average pH of 6.8 for the burned plots and 6.2 for the unburned plots

(additional plot information in Table S1). Although soil types differ, soil type did not affect

bacterial or fungal biomass or richness (glmer.nb; $p > 0.05$).

We used BAER soil burn severity maps created within seven days after fire containment

(https://burnseverity.cr.usgs.gov) to locate plots within moderate burn severity (Fig. 1A).

Notably, the BAER coarse-scale measurement (30 m) does not coincide well with the spatial

distribution and turnover rate of soil microbial communities (Bahram et al., 2013). Since ash

depth increases with soil heating (Parson et al., 2010) and fire severity (Bodí et al., 2014), we

used ash depth as a proxy for soil burn severity by averaging three separate measurements of ash

depth (cm) per each 1 m² subplot (Fig. 1C). Since ash can be rapidly redistributed via post-fire

wind and rain (Bodí et al., 2014), we measured ash before any precipitation event (first rains

occurred on Oct 3) or high wind events (average wind Sept 3 -Sept 17 was 2.7 m/s). Hence, our

measure of soil burn severity refers to initial ash depth.

We sampled soils at 17, 25, 34, 67, 95, 131, 187, 286, and 376 days, corresponding to

approximately 2 and 3 weeks, and 1, 2, 3, 4, 6, 9, and 12 months post-fire. At each of the nine

time points, we collected the top 10 cm of mineral soil (A horizons) beneath the ash layer from

each burned subplot. Since wildfire results in the combustion of the organic layer, to ensure

homogeneity in sampling and direct comparison between treatments in the unburned plots, we

removed the litter layer before sampling A horizons (Pulido-Chavez et al., 2021). Soils were

collected with a ~250 mL releasable bulb planter cleaned with ethanol after each use to prevent

cross-contamination, resulting in 36 soil samples (9 plots x 4 subplots) per sampling time point.

Soils were transported in individual Whirl-Paks in a cooler to the University of California-

Riverside (UCR) within hours of sampling, stored overnight at 4°C, and sieved (2mm) at room

temperature in ethanol cleaned sieves within 24 hours of sampling to minimize microbial

community turnover (Phillips, 2021). A subsample was frozen at -80°C for future DNA

extraction. While soils were stored at -80°C from 1 month to 1 year before DNA extractions were

performed, differences in storage time at -80°C do not affect DNA quantity and integrity (Lauber

et al., 2010; Pavlovska et al., 2021).

2.3 DNA extraction, amplification, and sequencing

DNA extractions for all nine timepoints were performed in the summer of 2019. Soils

were weighed (0.25 g) using ethanol-cleaned spatulas and processed using Qiagen DNeasy

PowerSoil Kits following the manufacturer's protocol, with an increase in centrifugation time to

1.5 min after adding the C3 solution (a solution used to precipitate the organic and inorganic

matter) due to large amounts of precipitate still in the solution, then stored at -20˚C. Extracted

DNA was amplified using the primer pair ITS4-fun and 5.8s to amplify the ITS2 region for fungi

(Taylor et al., 2016) and the primer pair 515F-806R to amplify the V4 region of the 16S rRNA

gene for archaea and bacteria (Caporaso et al., 2011) using the Dual-Index Sequencing Strategy

(DIP) (Kozich et al., 2013). Although our 16S primers amplify archaea and bacteria, for

simplicity, we refer to 16S methods and results simply as bacteria since archaea only contributed

<1% of sequencing reads. We conducted polymerase chain reaction (PCR) in two steps. The first

PCR amplified gene-specific primers, and the second PCR ligated the DIP barcodes and adaptors

for Illumina sequencing. For bacteria, we combined 1 μL of 1:10 diluted DNA, 10.5 μL of Ultra-

Pure Sterile Molecular Biology Grade water (Genesee Scientific, San Diego, CA, USA), 12.5 μL

of AccuStart ToughMix (2x concentration; Quantabio, Beverly, MA, USA), and 0.5 μL each of

the 10 μM 515F and 806R primers. Thermocycler conditions for PCR1 were: 94°C for 2 min

followed by 29 cycles of 94°C for 30 s, 55°C for 30 s, 68°C for 1 min, followed by a 2 min

extension at 68°C. For fungi, we combined 5 μL of undiluted DNA, 6.5 μL of ultra-pure water,

12.5 μL of AccuStart ToughMix, and 0.5 μL each of the 10 μM ITS4-fun and 5.8s primers.

Thermocycler conditions for PCR1 were: 94 °C for 2 min, followed by 30 cycles of 94 °C for 30

s, 55 °C for 30 s, 68 °C for 2 min, followed by a 10 min extension at 68°C. PCR1 products were

cleaned with AMPure XP magnetic beads (Beckman Coulter Inc., Brea, CA), following the

manufacturer's protocols. The DIP PCR2 primers containing the barcodes and adaptors for

Illumina sequencing were ligated to the amplicons during the second PCR step in a 25 μL

reaction containing 2.5 μL of the 10 μM DIP PCR2 primers, 6.5 μL of ultra-pure water, 12.5 μL

of Accustart ToughMix, and 1 μL of PCR1 product. Thermocycler conditions for PCR2 for

bacteria and fungi were: 94°C for 2 min followed by 10 cycles of 94°C for 30 s, 60°C for 30 s,

and 72°C for 1 min. Bacterial and fungal PCR products for the 324 samples (9 plots x 4 subplots

x 9 timepoints) were then separately pooled based on gel electrophoresis band strength and

cleaned with AMPure following established methods (Glassman et al., 2018). Each pool

contained 2-3 timepoints from burned and unburned plots and was checked for quality and

quantity with the Agilent Bioanalyzer 2100 before combining bacteria and fungi at a 2:3 ratio

(0.4 units for bacteria to 0.6 units for fungi). Since each library only fit 2-3 timepoints for both

bacteria and fungi, we sequenced the nine timepoints across four libraries with Illumina MiSeq

2x300bp at the UCR Institute for Integrative Genome Biology. In addition to our 324

experimental samples, we added negative DNA extractions and PCR controls from each time

point and mock communities (ZymoBIOMICS microbial community standard, Zymo, Irvine,

CA) to each library for additional quality control and inference.

2.4 Bacterial and fungal biomass

We estimated bacterial and fungal gene copy numbers with quantitative (q) PCR as a

proxy for biomass using the Eub338/Eub518 primers for bacteria (Fierer et al., 2005) and

FungiQuant-F/FungiQuant-R primers for fungi (Liu et al., 2012). We used the small subunit since

it is more conserved and has less length variation making it better suited for qPCR than the ITS2

region (Mayer et al., 2021), which is better at species level identification (Schoch et al., 2012).

We generated standard curves using a 10-fold serial dilution of the standards by cloning the 18S

region of the fungus *Saccharomyces cerevisiae* or the 16S region of the bacteria *Escherichia coli*

into puc57 plasmid vectors constructed by GENEWIZ, Inc. (NJ, USA) as previously established

(Averill & Hawkes, 2016). The 10 µL qPCR reactions were performed in triplicate. Reactions

contained 1 µL undiluted DNA, 1 µL of 0.05M Tris-HCl ph8.3, 1 µL of 2.5mM $MgCl_2$ (New

England BioLabs (NEB); Ipswich, MA), 0.5 µL of 0.5mg/ml BSA, 0.5 µL of 0.25mM dNTP

(NEB), 0.4 µL of both primers at 0.4µM, 0.5 µL of 20X Evagreen Dye (VWR International;

Radnor, PA), 0.1 µL of Taq DNA polymerase (NEB) and 4.6 µL ultra-pure water. We employed

the CFX384 Touch Real-Time PCR Detection System with the following conditions: 94˚C for 5

min, followed by 40 cycles of 94˚C for 20 seconds, 52˚C (for bacteria) or 50˚C (for fungi) for 30

seconds, followed by an extension at 72˚C for 30 seconds. Gene copy numbers were generated

using the equation $10^{(Cq-b)/m}$, using the quantification cycle (Cq), calculated as the average

Cq value per sample in relation to the known/calculated copies in the Cq/threshold cycles, the y-

intercept (b) and the slope (m) generated with CFX Maestro software. Values with an $R^2 > 0.994$

were considered acceptable. Gene copy numbers were normalized per gram of dry soil (Tatti et

al., 2016). Notably, all methods of estimating microbial biomass have limitations (Gao et al.,

2022), including qPCR, due to variation in target copy number (Lofgren et al., 2019; Song et al.,

2014). However, the high sensitivity (Tellenbach et al., 2010) of qPCR for quantifying the

selected marker gene (Smith & Osborn, 2009) makes it ideal for estimating microbial biomass

from environmental samples when interest is not in species specificity but in estimates of total

microbial biomass. Moreover, common biomass quantification methods for fungi are correlated

(Cheeke et al., 2017).

2.5 Bioinformatics

Illumina data was processed with Qiime2 version 2020.8 (Bolyen et al., 2019). The

demultiplexed fastQ files from the four Illumina sequencing runs were processed individually

using cutadapt (Martin, 2011) to remove the primers and ran through DADA2 version 2020.8

with the defaults parameters to filter out and remove chimeric sequences and low-quality regions

and to produce Amplicon Sequence Variants (ASVs) (Callahan et al., 2017). Bacteria forward

reads were trimmed to 170 bp and reverse to 163 bp, while fungal forwards reads were trimmed

to 209 bp and the reverse reads to 201 bp. On average, 72% of forward and reverse sequences

merged for bacteria, and 61% merged for fungi. DADA2 outputs from each library were

combined into one library for downstream processing, including the removal of singletons and

taxonomic assignments. We used SILVA version 132 for bacterial (Yilmaz et al., 2014) and

UNITE version 8.2 for fungal (Abarenkov et al., 2020) taxonomic assignment using Qiime2

Naïve Bayes Blast+ classifier. Bacterial sequences assigned to mitochondria and chloroplasts and

fungal sequences not assigned to Kingdom Fungi were removed. Fungal ASV tables were

exported and parsed through FUNGuild (Nguyen et al., 2016) to assign functional ecological

guilds, including only highly probable confidence rankings for AMF, EMF, saprotrophs, and

pathogens. Moreover, negative and mock controls were inspected to ensure that sequences and/or

ASVs in negative controls were negligible and that mock taxonomy correlated with known

communities. Sequences were submitted to the National Center for Biotechnology Information

Sequence Read Archive under BioProject accession number PRJNA761539.

2.6 Statistical analysis

The ASV tables were rarefied to a sequences/sample depth of 7,115 for bacteria and

11,480 for fungi to account for uneven sequencing depth. Fungal and bacterial alpha diversity

was estimated using BiodiversityR version 2.14-2 (Kindt & Coe, 2005) with the following

metrics: observed species richness, Simpson, Shannon, Chao1, ACE, Simpson's evenness, and

Inverse Shannon. Although estimating richness from diverse soil communities has limitations

(Willis et al., 2017), wildfire-affected soil constitutes low diversity communities, and patterns of

species richness were similar across metrics (Fig. S1). Thus, we focused on richness estimated as

the number of observed ASVs after rarefaction for all downstream analyses. To test if wildfire

decreased biomass and richness and if biomass and richness increased with time since fire (H1),

if wildfire will have a distinct impact on fungal guilds (H2), and to determine if changes in

biomass and richness are associated with precipitation and soil burn severity (H3), for both

bacteria and fungi, we performed backward model selection and fitted nine statistical models with

treatment (burned vs. unburned), time (measured as days since fire), ash depth (cm), total

monthly precipitation (mm), and second order interactions as predictors. We used generalized

mixed effect models (glmer) with a negative binomial distribution in the MASS package version

7.3-57 (Venables & Ripley, 2002) to account for the over-dispersion of the data and the fact that

the conditional variance was higher than the conditional mean (Bliss, 1953; Ross & Preece,

1985). Time since fire and precipitation were scaled and centered. The level of nestedness for all

models was tested by running a null model with different nested levels (plot, subplot, and time

since fire) and no predictors. Model selection was made using Akaike Information Criterion

(AICc) in the MuMIN package version 1.46.0 (Barton, 2020). All richness models contained plot,

subplot, and time since fire as random effects, and all biomass models included plot and time

since fire as random effects. The subplot was not included for biomass analysis as it was not

determined to be the best model (AIC selection). Pseudo R^2, or the variance explained (marginal

and conditional) for all models, was calculated using the r.squaredGLMM function in the MuMIn

Package.

We compared beta diversity by generating distance matrices with the vegan Avgdist

function, calculating the square-root transformed median Bray-Curtis dissimilarity per 100

iterations. We used permutational multivariate analysis of variance (PERMANOVA) (Anderson,

2017) as implemented with the adonis function in vegan version 2.6-2 (Oksanen et al., 2018) to

test for the significant effects of wildfire, time since fire, precipitation, ash depth, and second

order interactions, on bacterial and fungal community composition overall. Moreover, we tested

the significance of wildfire at each independent time point (H1). Results were visualized using

Non-Metric Multidimensional Scaling (NMDS) ordinations.

We employed several methods to quantify succession (H4). First, we visualized

succession by characterizing community composition patterns at the genus level using phyloseq

version 1.38.0 (McMurdie & Holmes, 2013) and grouping the relative abundance of the dominant

ASVs (> 3% sequence abundance used for plot legibility) for each time point in burned and

unburned communities independently. Second, we used the vegan mantel function to determine

the correlation between temporal distance and community composition to determine how

turnover varies in the presence or absence of fire, similar to a study examining Sorghum

microbial succession (Gao et al., 2020). Third, to visualize spatial species turnover (Baselga,

2010), we took advantage of the fact that early successional periods display large variability

(Collins, 1990; Pandolfi, 2008). Thus, we tested the homogeneity of the bacterial and fungal

communities during succession using the vegan betadisper function to calculate the multivariate

dispersion of Bray-Curtis dissimilarities and visualized the results using principal coordinates

analysis (PCoA). Finally, we employed the codyn package version 2.0.5 (Hallett et al., 2016) to

identify the patterns of community dynamics over successional time, including the rate of

directional change, stability, and synchrony (Collins et al., 2000). Succession involves directional

change via the replacement of early to late successional species (Clements, 1916; Platt &

Connell, 2003); thus, we calculated the rate of directional change using the Euclidian distance of

each microbial community. Then we measured species turnover to determine if species

appearances or disappearances drive succession. Since stability increases as diversity increases

(diversity-stability hypothesis), we independently measured community stability in the burned

and unburned plots as the mean abundance divided by the standard deviation at each temporal

time point (Lehman & Tilman, 2000). However, community stability depends on species

covariance over time; thus, we measured synchrony (Loreau & de Mazancourt, 2008), where

perfect species synchrony is a value of 1 and 0 equals asynchrony (Valencia et al., 2020).

Asynchrony can result in covariance in species populations, where tradeoffs among species can

contribute to overall community stability.

All statistical analyses were conducted in R 4.1.1 (R Core Team, 2021), and plots were

created with ggplot2 version 3.3.6 (Wickham, 2016). All statistical codes are available on GitHub

https://github.com/pulidofabs/SecondarySuccession-Chaparral.

3. Results

3.1 Sequencing data

The four Illumina MiSeq runs resulted in 9.8 M bacterial and 24.6 M fungal sequences for

an average of 31,052 bacterial and 78,202 fungal sequences/sample and a total of 33,078 bacterial

and 11,480 fungal ASVs. Compared to the experimental samples, bacterial negative controls, on

average, had 10 ASVs, while fungal negative controls had 8 ASVs (Table S2). Rarefaction

resulted in a total of 24,874 bacterial and 7,445 fungal ASVs, and the removal of all control

samples. Extracting the fungal guilds from the rarefied fungal ASV table resulted in 208 EMF, 70

saprobic, 65 AMF, and 26 pathogenic fungal ASVs.

3.2 Wildfire effects on bacterial and fungal biomass and richness

Fire significantly reduced bacterial and fungal biomass and richness during the first post-

fire year (Tables S3). Fire had a larger effect on biomass than richness for both bacteria and fungi

and larger effects on fungi than bacteria overall (Fig. 2). On average, across the year, fire reduced

bacterial richness by 46% (Fig. S1A) and biomass by 47% (Fig. S2A) and fungal richness by

68% (Fig. S1A) and biomass by 86% (Fig. S2B). The direct, immediate effect of wildfire at 17

days post-fire was stronger for biomass than richness, with bacterial biomass decreasing by 84%

(Fig. 2A) and fungal biomass by 97% (Fig. 2B; Table S3). In contrast, at 17 days post-fire, fungal

richness declined by 45% (Fig. 2D), but bacterial richness temporarily increased by 31% but

declined by 29% at 25 days post-fire (Fig. 2C; Table S3). While the differences in biomass and

richness between the burned and unburned plots lessened with time since fire, one year was

insufficient for bacterial and fungal biomass or richness to recover to unburned levels (Fig. 2).

The effects of fire across time remained larger for fungi than for bacteria, with fungal biomass

remaining 80% and richness 61% lower and bacterial biomass remaining 43% and richness 23%

lower in the burned plots at 1-year post-fire (Fig. 2; Table S3).

3.3 Fire directly and indirectly affects biomass and richness

Fire had direct, negative effects on bacterial and fungal biomass and richness. These direct

effects were modified over time by positive interactions with fire for fungal biomass and with soil

burn severity for fungal richness and bacterial biomass (Fig. 3; Table S4). Time since fire also

directly positively impacted fungal richness (Table S4). Soil burn severity had significant adverse

direct effects on bacterial and fungal richness but no significant direct effects on biomass,

meaning that biomass was equally reduced by fire regardless of severity (Figs. 3C, 3D; Table S4).

While precipitation had no significant direct effects, precipitation interacted with fire to positively

impact bacterial biomass (Fig. 2A), and precipitation interacted with time since fire to negatively

affect bacterial richness (Fig. 2C) but positively affect fungal richness (Fig. 2D; Table S4).

3.4 Wildfire effects on the richness of different fungal guilds

Wildfire led to large and significant reductions in species richness for all fungal guilds

(Fig. S3; Table S5), with the largest initial declines for AMF. On average, across time points,

EMF decreased by 68%, pathogens by 71%, saprobes by 86%, and AMF by 98% in burned

compared to unburned plots (Fig. S3), and one year was insufficient for any guild to return to

unburned levels. In fact, at 376 days, EMF were on average 91%, AMF 89%, pathogens 69%,

and saprobes 61% lower in the burned plots (Fig. S4; Table S5). EMF richness declined with soil

burn severity and due to a fire and time interaction, whereas time since fire had a direct, negative

effect on saprobic richness, which declined over time (Table S5). Finally, an interaction between

fire and precipitation negatively affected both EMF and pathogen richness (Table S5).

3.5 Wildfire changes microbial community composition

Fire significantly affected bacterial and fungal community structure, explaining 13% of

the compositional variation for bacteria and 10% for fungi (Table S6). Time since fire had

smaller impacts on community composition, explaining 4% of the variation for bacteria and 1%

for fungi. Ash depth equally affected bacterial and fungal composition explaining 2% of the

variance for both, whereas precipitation had slightly larger impacts on bacteria than fungi,

explaining 3% of bacterial and 1% of the fungal variation (Table S6). There were also small but

significant interaction effects for both bacteria and fungi (Table S6). Furthermore, community

composition between burned and unburned plots significantly varied at all 9-time points for both

bacteria (Fig. S5) and fungi (Fig. S6), with differences in community composition increasing over

time from 12% to 21% for bacteria (Fig. S5) and 9% to 13% for fungi (Fig. S6) from day 17 to

376 post-fire. The overall changes in burned bacterial (Fig. 4A) and fungal composition (Fig. 4C)

were driven by the previously rare taxa, which increased in abundance over time (Figs. 4B, 4D).

Unlike fungi, burned bacterial communities were consistently dominated over time (57% average

sequence abundance) by a single genus, the Proteobacteria *Massilia* (Fig. 4B). However, three

Firmicutes quickly rose in dominance, with *Bacillus* (13%), an uncultured Clostridiales (26%),

and *Paenibacillus* (4%) dominating at 34- and 67-days post-fire. Yet these Firmicutes rapidly

declined as the Proteobacteria *Noviherbaspirillum* increased in abundance over time, from 1% at

34 days to 24% by the end of the year (Fig. 4B). For burned fungi, two Basidiomycetes that

dominated at 17 days post-fire, the yeast *Geminibasidium* (45%) and the EMF *Inocybe* (16%),

rapidly declined while filamentous Ascomycetes increased over time (Fig. 4D). The genera

Pyronema increased as *Geminibasidium* decreased and dominated from 25 to 95 days post-fire,

peaking at 67 days post-fire with 67% sequence abundance. In addition, the Ascomycetes

Aspergillus increased from 2% to 22% and *Penicillium* from 36% to 49% from 17 to 376 days

post-fire (Fig. 4D).

3.6 Bacterial successional dynamics

Burned bacterial communities experienced rapid and distinct successional trajectories

(Fig. 5A) at roughly double the rate of fungi (Fig 5C; Fig. 5D) and were driven by six major

compositional turnover points at 25, 34, 95, 131, 187, and 376 days (Fig. 5E). In contrast, the

unburned bacterial communities experienced low dominance and remained stable over time (Fig.

6A) with no succession (Figs. 6C) or compositional turnover (Figs. 6E). These patterns are also

reflected in the measures of successional dynamics (Table S7). For example, burned bacterial

communities experienced temporal changes in species composition that were directional and

higher (0.16) than in the unburned communities (0.12; Table S7). Burned bacterial communities

also exhibited lower synchrony (0.03) than unburned communities (0.22), resulting in higher

stability in burned (8.35) compared to unburned bacterial communities (5.36; Table S7).

3.7 Fungal successional dynamics

Burned fungal communities experienced rapid and distinct successional trajectories (Fig.

5B; Fig. 5D) that were driven by five major compositional turnover points at 25, 34, 67, 95, and

131 days post-fire (Fig. 5F). In contrast, the unburned fungal communities experienced low

dominance (Fig. 6B) and remained stable over time with no succession (Fig. 6D) or

compositional turnover (Fig. 6F). Like bacteria, these patterns were similarly reflected in the

measures of successional dynamics (Table S7) such that burned fungal communities had higher

directional change (0.49) than the unburned communities (0.08), reflecting stronger patterns of

succession as more species were introduced into the burned system. Burned fungal communities

exhibited slightly lower synchrony (0.04) than unburned communities (0.05) but much lower

community stability (6.42) than unburned communities (8.58), indicating higher susceptibility to

change in burned fungal communities (Table S7).

3.8 Taxa driving microbial succession

We found that bacterial and fungal succession were driven by taxa that traded off in

abundance based on physiological traits, such as thermotolerance and fast colonization (Table 1).

Early turnover events for bacteria (Fig. 5A) and fungi (Fig. 5B) in burned plots were driven by

the disappearance of taxa, more so for bacteria (days 17-95) than fungi (days 17-25; Table S7).

Indeed, increased dominance by a few taxa, including the Firmicutes *Bacillus* at 25 days post-fire,

Paenibacillus and *Domibacillus* (34-67 days), and the Proteobacteria *Noviherbaspirillum* (95

days), and the constant dominance of *Massilia* (34-286 days post-fire) resulted in the

disappearance of early bacterial species, including RB41, *Conexibacter*, *Candidatus*

Udaeobacter, and *Bacillus* over time (Fig. 5A; Table S7). In contrast, fungal succession was

initiated by the rapid dominance of *Geminibasidium* and EMF genera that dominated the

unburned communities, including *Inocybe*, *Cortinarius*, and *Tomentella* in the Basidiomycota and

Balsamia in the Ascomycota and one previously rare Basidiomycete EMF genus *Mallocybe*, a

subgenus of *Inocybe* (Fig. 5B, 6B). Interestingly, all species of EMF rapidly disappeared from the

community at 25 days post-fire and were replaced by the constant dominance of *Pyronema*,

Penicillium, and *Aspergillus* (Fig. 5B). However, as succession ensued, species appearance drove

late-year successional turnover for bacteria (187-376 days; Fig. 5A) and mid-year successional

turnover for fungi (67-187 days; Fig. 5B; Table S7). For bacteria, the appearance of two

Bacteroidetes, *Pedobacter* and *Adhaeribacter*, and the Actinobacteria *Blastococcus* shaped late-

year (days 187-376) bacterial succession (Fig. 5A). For fungi, mid-year succession (days 95-187)

was driven by the appearance of filamentous fungal Ascomycetes in the Pyronemataceae and the

Sordariaceae genus *Gelasinospora*, and Basidiomycota mushroom-forming taxa (Fig. 5B). For

example, the previously dominant EMF genus *Inocybe* remained in the community but at a much

lower abundance, declining with time since fire from 16% at 17 days to 0.1% at 376 days (Fig.

4D). In contrast, the pyrophilous Basidiomycete saprobes *Coprinellus* and *Tephrocybe*, which

were rare in the unburned communities (<0.01% sequence abundance), appeared in the

community later in the year (Fig. 5B). Late-year fungal succession (days 286-376) was driven by

the increased dominance of *Tephrocybe* and *Aspergillus* and the Ascomycota *Rasamsonia*, and

the decreased abundance of *Pyronema* (Fig. 5B).

4. Discussion

Here, we present the highest resolution temporal sampling of post-fire microbiomes to

date, enabling us to show for the first time that bacteria and fungi experience rapid succession

driven by the dominance of pyrophilous taxa that traded off in abundance over time (H4).

Although our study was in California chaparral, many of the dominant pyrophilous taxa identified

here also dominate other burned environments, including Spanish shrublands (Pérez-Valera et al.,

2018) and pine (Bruns et al., 2020; Reazin et al., 2016), spruce (Whitman et al., 2019),

Eucalyptus (Ammitzboll et al., 2022; McMullan-Fisher et al., 2011), and redwood-tanoak forests

(Enright et al., 2022), indicating the generality of our results to wildfire-affected ecosystems. We

found that wildfire decreased bacterial and fungal biomass and richness, leading to community

composition shifts that persisted the entire year (H1). Fungal guilds were differentially affected

by fire and time, with AMF and saprobes experiencing the largest immediate fire impacts, but

over time, EMF experienced the largest declines (H2). Moreover, microbial richness and biomass

changes were driven by multiple abiotic interactions, including interactions between time since

fire, precipitation, and soil burn severity (H3).

4.1 Wildfire decreased bacterial and fungal biomass and richness

Fire decreased soil bacterial and fungal biomass and richness, corroborating previous

post-fire studies in Mediterranean shrublands (Pérez-Valera et al., 2018) and forests (Dooley &

Treseder, 2012). We noted a larger fire effect on fungal biomass and richness relative to bacteria,

consistent with previous research showing that bacteria are more resistant to fire than fungi

(Certini et al., 2021; Glassman et al., 2021; Pourreza et al., 2014; Pressler et al., 2019). Although

richness and biomass increased over time for both microbial groups, one year was insufficient for

either group to recover to unburned levels, consistent with previous studies indicating that in

shrublands, microbial biomass and richness recovery could take over two decades (Pérez-Valera

et al., 2018). Interestingly, we observed a transient increase in bacterial richness directly post-fire,

which may result from taxa such as the Actinobacteria genera *Soliribrobacter* and *Conexibacter*

(Albuquerque & da Costa, 2014) that may be favored by the increase in soil pH observed here

(Table S1) and in other studies (Neary et al., 1999) or by post-fire increases in nitrogen and

phosphorus availability (Certini et al., 2021).

4.2 Wildfire impacts on fungal guilds

Fire-induced mortality was rapid for AMF and saprobes. Yet, EMF were most affected by

fire over time, corroborating studies across ecosystems that point to the negative impact of fire on

mycorrhizal richness, primarily driven by host mortality (Dove & Hart, 2017; Pulido-Chavez et

al., 2021). Indeed, post-fire, all subplots were devoid of vegetation (0% vegetation cover)

compared to 97% vegetation cover in unburned plots. However, we note that the rapid decline in

AMF could be due to primer bias, as the ITS2 primer does not properly detect all AMF taxa

(Lekberg et al., 2018). Moreover, we show that surviving ectomycorrhizal Basidiomycetes

Cortinarius, *Inocybe*, and *Tomentella*, previously found in burned temperate pine forests (Owen

et al., 2019; Pulido-Chavez et al., 2021) and pine-dominated Mediterranean systems (Gassibe et

al., 2011; Hernández-Rodríguez et al., 2013) also dominated early chaparral fungal succession

and remained in the community for 1-2 months. Since EMF largely disappeared after two

months, this potentially answers the question of how long it takes mycorrhizal fungi to perish

after their hosts' death. Although it is possible that the EMF signal detected could be relic DNA

(Carini et al., 2016), high soil temperatures typical of post-fire systems (Amacher et al., 2001;

Neary et al., 1999) are likely to have degraded relic DNA (Sirois & Buckley, 2019; Torti et al.,

2015), suggesting that these EMF could have survived the fire and clung to a dying host for at

least two months. The survival of some EMF genera over others may be attributable to

exploration type and specificity for current versus stored photosynthates (Gray & Kernaghan,

2020; Pena et al., 2010). For example, *Cortinarius*, a long/medium exploration type, which

requires more carbon for maintenance, only survived for 17 days post-fire. In contrast, *Inocybe*, a

short-contact type (Agerer, 2001; Koide et al., 2014), survived at very low abundance for the

entire year, perhaps because it may be less carbon demanding or adapted to warming (Fernandez

et al., 2017). Our results indicate that early EMF could have survived on their hosts' stored

photosynthates or the lower resources provided by the surviving and resprouting manzanita and

chamise shrubs, while the vast majority of EMF experienced fire-induced mortality.

4.3 Pyrophilous fungi dominate the burned communities

543    Previously detected pyrophilous fungi across several ecosystems also dominated

544 California chaparral, indicating that pyrophilous microbes are not biome-specific. Instead, these

545 fungi may be activated by temperature thresholds reached during high-severity fires, which are

546 common in chaparral and some Pinaceae forests (Agee, 1993; Keeley & Zedler, 2009; Neary et

547 al., 1999). Specifically, our burned communities were dominated by Ascomycota in the genera

548 *Pyronema*, *Penicillium*, and *Aspergillus*, similar to dominant post-fire fungal species in

549 Mediterranean shrublands (Livne-Luzon et al., 2021), boreal spruce (Whitman et al., 2019), and

550 montane pine forests (Bruns et al., 2020; Pulido-Chavez et al., 2021). Both *Pyronema* (*P.*

551 *omphalodes*) and *Aspergillus* (*A. fumigatus*) are known pyrophilous fungi of California chaparral

552 (Dunn et al., 1982), and we identified additional species including *P. domesticum*, *A. udagawae*,

553 and *A. elsenburgensis*. These fungi are adapted to wildfire and produce fire-resistant structures,

554 including dormant spores, sclerotia, and conidia, which are heat-activated (Gottlieb, 1950;

555 Moore, 1962; Rhodes, 2006; Warcup & Baker, 1963). The rapid and efficient germination of

556 *Aspergillus* (in the subgenus *Fumigati*, including *A. udagawae*) induced by high temperatures

557 (Rhodes, 2006) and its ability to use various carbon and nitrogen sources, including ammonium

558 and nitrate (Krappmann & Braus, 2005), may position *Aspergillus* to rapidly dominate post-fire.

559 Furthermore, *Pyronema domesticum* can mineralize pyrogenic organic matter ((Fischer et al.,

560 2021), an abundant substrate in post-fire environments. Moreover, *Geminibasidium*, a recently

561 described thermotolerant Basidiomycete yeast (Nguyen et al., 2013), dominated at 17 days post-

562 fire. While not typically described as pyrophilous, presumably because most pyrophilous fungi

563 are described from mushrooms (McMullan-Fisher et al., 2011), two other studies have found

564 *Geminibasidium* to increase post-fire in pine forests (Pulido-Chavez et al., 2021; Yang et al.,

565   2020), suggesting that *Geminibasidium* is an underrepresented pyrophilous fungus. Together,

566   these results indicate that dominant pyrophilous fungi are more widespread than expected and

567   that their response to fire is likely due to fire adaptive traits and temperature thresholds.

569   ***4.4 Pyrophilous fungi drive fungal secondary succession***

570   Although pyrophilous fungi are widespread across ecosystems, we lacked an

understanding of how soon these pyrophilous fungi appear, their turnover rates, and if they

change in abundance over time (Fox et al., 2022). Successional theory states that early

successional stages are dominated by fast-growing or ruderal (R) organisms (Kinzig & Pacala,

2013) that tradeoff between stress tolerance (S) and competitive (C) life-history strategies over

time (Grime, 1977; Zhang et al., 2018). Recent adaptations of Grime's C-S-R to microbiomes

suggest that pyrophilous microbes survive and thrive post-fire with traits analogous to plants,

including post-fire resource acquisition (C), thermotolerant structures (S), and fast growth (R)

(Enright et al., 2022; Whitman et al., 2019). Our data suggest that chaparral pyrophilous microbes

fit into these trait categories (Table 1) and that tradeoffs among these traits might drive microbial

succession.

581   Early post-fire succession was driven by surviving thermotolerant fungi

(i.e., *Geminibasidium* and *Pyronema*), followed by fast-colonizers (i.e., *Penicillium* and

Aspergillus), which were overtaken by competitive fungi capable of exploiting post-fire

resources. (i.e., *Coprinellus* and *Tephrocybe*). Whereas *Geminibasidium* is thermo- and xero-

tolerant (Nguyen et al., 2013), *Pyronema* produces thermotolerant sclerotia (Moore, 1962),

enabling wildfire survival. Although both species are thermotolerant, differences in the

morphological growth characteristics between the yeast *Geminibasidium* and the filamentous

Pyronema may explain the tradeoff in dominance. For example, unicellular proliferation allowed

Geminibasidium to dominate instantly, but the ability to forage for nutrients and rapidly increase

surface colonization may allow the filamentous *Pyronema* to better dominate the open space.

Interestingly, *Gelasinospora heterospora*, a fungus closely related to heat-activated *Neurospora*

crassa (Dettman et al., 2001; Emerson, 1948), produces pigmented mycelia, and is known to

require heat or chemical treatment for germination (Alexopoulous & Mims, 1952), was highly

abundant at 25, 34, and 95 days post-fire, suggesting it is also pyrophilous. The decline of

thermotolerators coincided with the dominance of fast-growing Ascomycete fungi in the genera

Penicillium and *Aspergillus* (Dix & Webster, 1995; McGee et al., 2006), which both produce

copious asexual conidia (Crow, 1992), allowing for rapid colonization of the open niche,

analogous to early dominant post-fire plants, which are also often asexual (James, 1984). Lastly,

like plants, competition appears to drive later successional dynamics (Tilman, 1990; Zhang et al.,

2018), as suggested by the emergence of pyrophilous fungal decomposers, such as *Coprinellus*

and *Tephrocybe* at 9-12 months, corroborating findings from Eucalyptus forests (Ammitzboll et

al., 2022; McMullan-Fisher et al., 2011). Evidence suggests that *Coprinellus* can degrade

aromatic hydrocarbons (Steindorff et al., 2021), supporting the idea that it is outcompeting earlier

fungal taxa for this abundant and complex carbon source. Moreover, *Tephrocybe anthracophila*

has a high affinity for Ammonium-Nitrogen (Legg, 1992; Suzuki, 2017), allowing access to this

abundant post-fire resource. In conclusion, our results indicate that tradeoffs among fire-adapted

traits analogous to those in plants might drive post-fire microbial succession.

4.5 Pyrophilous bacteria dominate and initiate bacterial secondary succession

Our high temporal sampling enabled us to demonstrate that bacteria experience rapid

succession and community turnover initiated by possible aerobic, heterotrophic bacteria that form

endospores and produce antibiotics, which may improve their competitive abilities in the post-fire

environment. Moreover, we show that previously detected putative pyrophilous bacteria from

Canadian (Whitman et al., 2019), Chinese boreal (Xiang et al., 2014), and California redwood-

tanoak forests (Enright et al., 2022) and Spanish shrublands (Sáenz de Miera et al., 2020), also

dominated California chaparral, suggesting that these pyrophilous bacteria might be widely

distributed. Specifically, our burned communities favored taxa in the phyla Actinobacteria,

Acidobacteria, Firmicutes, and Proteobacteria, similar to previous studies (Ammitzboll et al.,

2022; Enright et al., 2022; Whitman et al., 2019; Xiang et al., 2014).

Bacterial successional dynamics differed from fungi in that the first few weeks were

characterized by high diversity and a lack of dominance, mimicking the distributions observed in

the unburned communities. In complex resource environments, such as those created by wildfire,

niche complementarity is more important than competition because diverse communities can

better exploit resources (Eisenhauer et al., 2013), further contributing to community stability.

Our results support this idea, as the dominant taxa in our study are likely to use different

resources. For example, members of the genus *Paenibacillus* can fix nitrogen (Monciardini et al.,

2003; Slepecky & Hemphill, 2006), *Bacillus* and *Conexibacter* can reduce nitrogen, and *Bacillus*

can further solubilize phosphate (Espinosa-de-los-Monteros et al., 2001; Kalayu, 2019).

However, at 25 days, bacteria experienced a large compositional turnover, which allowed for the

dominance of pyrophilous bacteria with fire-adapted traits. While tradeoffs in abundance were

not as stark for bacteria as for fungi, as indicated by the constant dominance of the Proteobacteria

Massilia, the following most abundant bacteria did experience tradeoffs in dominance against

other bacterial genera, possibly driven by physiological traits, such as thermotolerance and fast

colonization (Table 1).

Firmicutes contain some of the most resistant, thermotolerant endospore-producing

bacteria (Grady et al., 2016; Kaur et al., 2018; Mandic-Mulec et al., 2015), including members of

the genera *Bacillus*, *Paenibacillus*, and *Clostridiales* which decreased in abundance as fast-

colonizing genera in the Phylum Proteobacteria that may be capable of exploiting post-fire

substrates (e.g., *Novihersparillium* and *Massilia*) increased in dominance. Moreover, members of

the Firmicute order Clostridiales can trigger sporulation in response to stress (Paredes-Sabja et

al., 2011), potentially allowing for increased proliferation post-fire. While thermotolerance is

crucial for fire survival, rapidly dominating the open niche and using post-fire resources may

ensure post-fire growth. *Massilia* is a fast reproducing bacteria that associates with the

rhizosphere (Li et al., 2014) and colonizes AMF hyphae (Iffis et al., 2014) and the roots of

various plants (Ofek et al., 2012). Thus, the dominance of *Massilia* in our study and other post-

fire environments (Enright et al., 2022; Whitman et al., 2019) suggests that its diverse ecology

and rapid reproduction rate promote its proliferation within the soil (Toljander et al., 2005).

Moreover, our results, coupled with current knowledge in the field, show that the continued

dominance of *Massilia* and *Noviherbaspirillium* might be due to their ability to exploit post-fire

resources. For example, the genera *Massilia* contains species that can degrade long-chain

hydrocarbons (Ren et al., 2018) and reduce nitrates (Bailey et al., 2014). In contrast,

Noviherbaspirillium, a mid-to late-year dominant bacterial genus, may degrade polycyclic

aromatic hydrocarbon (Baldani et al., 2014; Grady et al., 2016; Woolet & Whitman, 2020). We

conclude that although tradeoffs are not as distinctive for bacteria as they are fungi, potentially

due to interaction among traits such as fast growth under high resources and resource acquisition

under low resources, our results provide evidence that pyrophilous bacteria have mechanisms to

survive the fire that may drive them to trade off in abundance from early to late dominant taxa

over time.

4.6 Microbial successional dynamics differ for bacteria and fungi

Chaparral vegetation successional patterns result from the self-replacement of the

dominant pre-fire species (i.e., auto-succession), meaning the pre-fire dominant species establish

early post-fire due to resprouting individuals, and these species gradually dominate the system

(Hanes, 1971). However, this was not the case for bacterial or fungal succession. Most post-fire

dominant genera were rare or absent pre-fire, and these initial colonizers experienced directional

change over time via species replacement. Our results are consistent with successional concepts

of non-catastrophic disturbances, which show that species that survive a disturbance or recover

quickly will dominate early successional dynamics but will inevitably be replaced by late-stage

species via directional replacement mechanisms (Platt & Connell, 2003). Similar directional

replacement patterns have been revealed after primary succession of fungi in a glacier foreland

(Dong et al., 2016). Although a few genera dominated burned microbial communities, bacteria

and fungi displayed different successional patterns. Bacterial burned communities exhibited

lower synchrony and higher stability compared to unburned communities. The low synchrony

indicates that stability may be conferred due to compensatory dynamics driven by tradeoffs in the

abundance of cooccurring competitors (Tilman, 1999). Conversely, the lower stability in the

unburned communities could be due to the increase in species synchrony reducing compensatory

dynamics. In systems with high environmental variability, compensatory dynamics mediated by

species asynchrony often contribute to stability (Yachi & Loreau, 1999). In contrast, burned

fungal communities displayed lower community stability compared to unburned communities,

while synchrony was similarly low in both. This lower stability may indicate that there was an

overall increase in the variation of fungal taxa potentially due to the large loss of species

observed in the fungal community (Tilman et al., 2014), resulting in priority effects and alternate

communities with different dominant genera (Connell & Slatyer, 1977; Fukami, 2015). However,

further research is needed to explore these microbial successional dynamics, as microbial

communities are a vital driver of the post-fire ecosystem recovery.

4.7 Precipitation and burn severity differentially affected bacterial and fungal communities

Although fire was the most influential driver of microbial change, we found that

interactions between time and precipitation also drove successional dynamics by differentially

affecting microbial biomass and richness. Precipitation interacted with time since fire resulting in

a negative correlation with bacterial richness and a positive correlation with fungal richness, with

bacterial richness declining but fungal richness increasing after rain events. At least one study

that performed a three-year precipitation manipulation experiment showed that increased

precipitation decreased bacterial richness (Yang et al., 2021). In contrast to richness, precipitation

interacted with fire to positively affect bacterial biomass, while fungal biomass remained

unaffected by rainfall. The distinct responses to precipitation could be due to microbial

physiological and morphological differences (Blazewicz et al., 2014; Placella et al., 2012;

Selbmann et al., 2013). For example, fungi have high morphological plasticity, including the

ability to alternate growth forms based on environmental conditions, such as hyphal growth in

low nutrient conditions versus unicellular form in rich nutrient conditions (Selbmann et al.,

2013), which may allow fungi to survive small changes in moisture content, allowing for

continued growth over time. Moreover, fungi can also activate or enhance resistance to drought

(Evans & Wallenstein, 2012; Guhr & Kircher, 2020), allowing them to maintain a constant

reproduction rate regardless of the external environmental conditions, thus explaining the positive

impact on fungal richness. In contrast, bacteria can experience cell death due to rapid wet-up,

while other bacterial groups enter a dormant state, inhibiting growth (Schimel, 2018). Thus, to be

evolutionarily successful, some bacteria may respond rapidly to favorable conditions (e.g., rain)

(Leizeaga et al., 2022) and rapidly replicate to outcompete other microbes for the pulse of

nutrients released during the rain event (Homyak et al., 2014), thus potentially explaining the

peaks in richness observed after significant rainfall at day 131.

Previous research shows that ash depth correlates with fire intensity (Rice, 1993) and

severity (Bodí et al., 2014; Parson et al., 2010). Here, we measured ash depth before the first

rains, or significant wind events could remove ash from the site, and we found that ash depth can

serve as a proxy of soil burn severity on a scale relevant to soil microbes. Unlike the BAER

maps, which classified all plots as moderate burn severity, we identified large variations in soil

burn severity across our subplots (Fig. S7). It is well known that wildfires result in a

heterogeneous landscape (Jain et al., 2008); thus, the discrepancy in soil burn severity in the

BAER maps could be due to the lack of ground verification. Since our severity effects were

similar to those found in other studies, we feel confident that ash depth can serve as a proxy for

soil burn severity. We found that severity negatively affected bacterial and fungal richness but not

biomass. Similar to studies in boreal (Whitman et al., 2019), oak (Pourreza et al., 2014), and

Mediterranean forests (Lucas-Borja et al., 2019), bacterial and fungal richness decreased with soil

burn severity. However, contrary to a meta-analysis (Dooley & Treseder, 2012), neither bacterial

nor fungal biomass was directly impacted by soil burn severity, probably due to microbial

adaptations to the natural high severity fire regime of chaparral. However, since we observed an

interaction effect of time since fire and severity on microbial richness, it is possible that

previously observed severity effects, derived from chronosequence studies where space is

substituted for time and which, on average, occur two years post-fire (Pressler et al., 2019), may

represent an interaction effect of time and severity and not a direct effect of soil burn severity.

730
731
732   **5. Conclusion**

733
Our high-resolution temporal sampling allowed us to detect rapid secondary succession in

bacterial and fungal communities and marked tradeoffs in abundance among pyrophilous

microbes over time. We found that the same pyrophilous bacteria and fungi that respond to fires

in temperate and boreal forests and Mediterranean shrublands also dominate in California

chaparral, allowing us to generalize knowledge of post-fire microbiomes to dryland habitats that

are rapidly increasing with climate change (Feng & Fu, 2013). While pyrophilous microbes have

been described in other systems, their turnover rates were unknown due to limited post-fire

sampling (Fox et al., 2022). We hypothesize that tradeoffs among fire-adapted traits, with

thermotolerators dominating first, followed by fast colonizers, and finally by competitors capable

of capitalizing on post-fire resource acquisition, drive post-fire microbial successional dynamics.

It appears likely that pyrophilous microbes share many analogous traits to plants, enabling us to

adapt successional theory developed for plants to microbes while acknowledging that bacteria

and fungi differ in how they respond to fire according to their physiology. We conclude that post-

fire bacteria and fungi experience rapid successional changes after fire, suggesting that these

dynamics can help increase our ability to predict if and how ecosystems recover after a wildfire,

an increasingly widespread disturbance with climate change.

750
751
Acknowledgments
We thank the Cleveland National Forest and the Trabuco Ranger District, including
District Ranger Darrel Vance and Emily Fudge, Jeffrey Heys, Lauren Quon, Jacob Rodriguez,
and Victoria Stempniewicz, for their guidance and help with permitting and site selection. We
thank Judy A. Chung for her help with fieldwork and molecular work and Aral C. Greene, Dylan
Enright, and Sameer S. Saroa for fieldwork assistance. We thank Amelia Nelson, Elizah
Stephens, Kaleigh A. Russell, and anonymous reviewers for their helpful comments on the
manuscript. This work was supported by UC Riverside, the BLM JFSP Award #012641-002 and
Shipley Skinner Award to MFPC and SIG, the USDA-NIFA Award 2022-67014-36675 to SIG
and PMH, and the DOE BER Award DE-SC0023127 to SIG and PMH.

762
763
764
Reference
Abarenkov, K., Zirk, A., Piirmann, T., Pöhönen, R., Ivanov, F., Nilsson, R. H., & Kõljalg, U.
(2020). UNITE QIIME release for Fungi 2 (Version 04.02.2020). UNITE Community.
https://doi.org/10.15156/BIO/786387
Abatzoglou, J. T., Williams, A. P., & Barbero, R. (2019). Global Emergence of Anthropogenic
Climate Change in Fire Weather Indices. Geophysical Research Letters, 46(1), 326–336.
https://doi.org/10.1029/2018GL080959
Agee, J. K. (1993). Fire Ecology of Pacific Northwest Forests. Island Press.
Agerer, R. (2001). Exploration types of ectomycorrhizae. Mycorrhiza, 11(2), 107–114.
https://doi.org/10.1007/s005720100108
Albuquerque, L., & da Costa, M. S. (2014). The Families Conexibacteraceae, Patulibacteraceae
and Solirubrobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F.
Thompson (Eds.), The Prokaryotes: Actinobacteria (pp. 185–200). Springer.
https://doi.org/10.1007/978-3-642-30138-4_200

Alexopoulous, C. J., & Mims, C. W. (1952). Introductory mycology (3rd ed.). John Wilet &
Sons.
Allen, M., Egerton-Warburton, L., Treseder, K., Cario, C. H., Lindahl, A., Lansing, J., Querejeta,
J., Kårén, O., Harney, S., & Zink, T. (2005). Biodiversity of Mycorrhizal Fungi in
Southern California 1. /paper/Biodiversity-of-Mycorrhizal-Fungi-in-Southern-1-Allen-
Egerton-Warburton/445d4f6878927a5307669f5e73df3174703f4d2a
Amacher, M. C., Johnson, A. D., Kutterer, D. E., & Bartos, D. L. (2001). First-year postfire and
postharvest soil temperatures in aspen and conifer stands (RMRS-RP-27; p. RMRS-RP-
27). U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
https://doi.org/10.2737/RMRS-RP-27
Ammitzboll, H., Jordan, G. J., Baker, S. C., Freeman, J., & Bissett, A. (2022). Contrasting
successional responses of soil bacteria and fungi to post-logging burn severity. Forest
Ecology and Management, 508, 120059. https://doi.org/10.1016/j.foreco.2022.120059
Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In
Wiley StatsRef: Statistics Reference Online (pp. 1–15). American Cancer Society.
https://doi.org/10.1002/9781118445112.stat07841
Averill, C., & Hawkes, C. V. (2016). Ectomycorrhizal fungi slow soil carbon cycling. Ecology
Letters, 19(8), 937–947. https://doi.org/10.1111/ele.12631
Baar, J., Horton, T. R., Kretzer, A. M., & Bruns, T. D. (1999). Mycorrhizal colonization of Pinus
muricata from resistant propagules after a stand-replacing wildfire. New Phytologist,
143(2), 409–418. https://doi.org/10.1046/j.1469-8137.1999.00452.x
Bahram, M., Kõljalg, U., Courty, P.-E., Diédhiou, A. G., Kjøller, R., Põlme, S., Ryberg, M.,
Veldre, V., & Tedersoo, L. (2013). The distance decay of similarity in communities of
ectomycorrhizal fungi in different ecosystems and scales. Journal of Ecology, 101(5),
1335–1344. https://doi.org/10.1111/1365-2745.12120
Bailey, A. C., Kellom, M., Poret-Peterson, A. T., Noonan, K., Hartnett, H. E., & Raymond, J.
(2014). Draft Genome Sequence of Massilia sp. Strain BSC265, Isolated from Biological
Soil Crust of Moab, Utah. Genome Announcements, 2(6), e01199-14, 2/6/e01199-14.
https://doi.org/10.1128/genomeA.01199-14
Baldani, J. I., Rouws, L., Cruz, L. M., Olivares, F. L., Schmid, M., & Hartmann, A. (2014). The
Family Oxalobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F.
Thompson (Eds.), The Prokaryotes (pp. 919–974). Springer Berlin Heidelberg.
https://doi.org/10.1007/978-3-642-30197-1_291
Barnard, R. L., Osborne, C. A., & Firestone, M. K. (2013). Responses of soil bacterial and fungal
communities to extreme desiccation and rewetting. The ISME Journal, 7(11), 2229–2241.
https://doi.org/10.1038/ismej.2013.104
Barro, S. C., & Conard, S. G. (1991). Fire effects on California chaparral systems: An overview.
Environment International, 17(2–3), 135–149. https://doi.org/10.1016/0160-
4120(91)90096-9
Barton, K. (2020). MuMIn: Multi-Model Inference (R package version 1.43.17.).
https://CRAN.R-project.org/package=MuMIn
Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. Global
Ecology and Biogeography, 19(1), 134–143. https://doi.org/10.1111/j.1466-
8238.2009.00490.x

Blazewicz, S. J., Schwartz, E., & Firestone, M. K. (2014). Growth and death of bacteria and fungi
underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. Ecology, 95(5),
1162–1172. https://doi.org/10.1890/13-1031.1
Bliss, C. I. (1953). Fitting the negative binomial distribution to biological data (with "Note on the
efficient fitting of the negative binomial" by R. A. Fisher), Biometrics, 9, (9th ed.).
Biometrics.
Bodí, M. B., Martin, D. A., Balfour, V. N., Santín, C., Doerr, S. H., Pereira, P., Cerdà, A., &
Mataix-Solera, J. (2014a). Wildland fire ash: Production, composition and eco-hydro-
geomorphic effects. Earth-Science Reviews, 130, 103–127.
https://doi.org/10.1016/j.earscirev.2013.12.007
Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,
Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger,
K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A.
M., Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and
extensible microbiome data science using QIIME 2. Nature Biotechnology, 37(8), 852–
857. https://doi.org/10.1038/s41587-019-0209-9
Bruns, T. D., Chung, J. A., Carver, A. A., & Glassman, S. I. (2020). A simple pyrocosm for
studying soil microbial response to fire reveals a rapid, massive response by Pyronema
species. PLOS ONE, 15(3), e0222691. https://doi.org/10.1371/journal.pone.0222691
Cairney, J. W. G., & Bastias, B. A. (2007). Influences of fire on forest soil fungal
communitiesThis article is one of a selection of papers published in the Special Forum on
Towards Sustainable Forestry — The Living Soil: Soil Biodiversity and Ecosystem
Function. Canadian Journal of Forest Research, 37(2), 207–215.
https://doi.org/10.1139/x06-190
Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. The ISME Journal, 11(12),
2639–2643. https://doi.org/10.1038/ismej.2017.119
Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P.
J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of
millions of sequences per sample. Proceedings of the National Academy of Sciences,
108(Supplement_1), 4516–4522. https://doi.org/10.1073/pnas.1000080107
Carini, P., Marsden, P. J., Leff, J. W., Morgan, E. E., Strickland, M. S., & Fierer, N. (2016). Relic
DNA is abundant in soil and obscures estimates of soil microbial diversity. Nature
Microbiology, 2(3), 1–6. https://doi.org/10.1038/nmicrobiol.2016.242
Certini, G., Moya, D., Lucas-Borja, M. E., & Mastrolonardo, G. (2021). The impact of fire on
soil-dwelling biota: A review. Forest Ecology and Management, 488, 118989.
https://doi.org/10.1016/j.foreco.2021.118989
Cheeke, T. E., Phillips, R. P., Brzostek, E. R., Rosling, A., Bever, J. D., & Fransson, P. (2017).
Dominant mycorrhizal association of trees alters carbon and nutrient cycling by selecting
for microbial groups with distinct enzyme function. New Phytologist, 214(1), 432–442.
https://doi.org/10.1111/nph.14343
Clements, F. E. (1916). Plant Succession: An Analysis of the Development of Vegetation.
Carnegie Institution of Washington.

Collins, S. L. (1990). Patterns of Community Structure During Succession in Tallgrass Prairie.
Bulletin of the Torrey Botanical Club, 117(4), 397–408. https://doi.org/10.2307/2996837
Collins, S. L., Micheli, F., & Hartt, L. (2000). A method to determine rates and patterns of
variability in ecological communities. Oikos, 91(2), 285–293.
https://doi.org/10.1034/j.1600-0706.2000.910209.x
Connell, J. H., & Slatyer, R. O. (1977). Mechanisms of Succession in Natural Communities and
Their Role in Community Stability and Organization. The American Naturalist, 111(982),
1119–1144.
Cowan, A. D., Smith, J. E., & Fitzgerald, S. A. (2016). Recovering lost ground: Effects of soil
burn intensity on nutrients and ectomycorrhiza communities of ponderosa pine seedlings.
Forest Ecology and Management, 378, 160–172.
https://doi.org/10.1016/j.foreco.2016.07.030
Crow, J. F. (1992). An advantage of sexual reproduction in a rapidly changing environment. The
Journal of Heredity, 83(3), 169–173.
https://doi.org/10.1093/oxfordjournals.jhered.a111187
Dalling, J. W. (2008). Pioneer Species. In B. Fath (Ed.), Encyclopedia of Ecology (Second
Edition) (pp. 181–184). Elsevier. https://doi.org/10.1016/B978-0-444-63768-0.00534-5
Day, N. J., Cumming, S. G., Dunfield, K. E., Johnstone, J. F., Mack, M. C., Reid, K. A.,
Turetsky, M. R., Walker, X. J., & Baltzer, J. L. (2020). Identifying Functional Impacts of
Heat-Resistant Fungi on Boreal Forest Recovery After Wildfire. Frontiers in Forests and
Global Change, 3. https://doi.org/10.3389/ffgc.2020.00068
Derroire, G., Balvanera, P., Castellanos-Castro, C., Decocq, G., Kennard, D. K., Lebrija-Trejos,
E., Leiva, J. A., Odén, P.-C., Powers, J. S., Rico-Gray, V., Tigabu, M., & Healey, J. R.
(2016). Resilience of tropical dry forests – a meta-analysis of changes in species diversity
and composition during secondary succession. Oikos, 125(10), 1386–1397.
https://doi.org/10.1111/oik.03229
Dettman, J. R., Harbinski, F. M., & Taylor, J. W. (2001). Ascospore Morphology Is a Poor
Predictor of the Phylogenetic Relationships of Neurospora and Gelasinospora. Fungal
Genetics and Biology, 34(1), 49–61. https://doi.org/10.1006/fgbi.2001.1289
Dix, N. J., & Webster, J. (1995). Fungal Ecology. Chapman & Hall.
Donato, D. C., Campbell, J. L., & Franklin, J. F. (2012). Multiple successional pathways and
precocity in forest development: Can some forests be born complex? Journal of
Vegetation Science, 23(3), 576–584. https://doi.org/10.1111/j.1654-1103.2011.01362.x
Dong, K., Tripathi, B., Moroenyane, I., Kim, W., Li, N., Chu, H., & Adams, J. (2016). Soil
fungal community development in a high Arctic glacier foreland follows a directional
replacement model, with a mid-successional diversity maximum. Scientific Reports, 6(1),
26360. https://doi.org/10.1038/srep26360
Dooley, S. R., & Treseder, K. K. (2012). The effect of fire on microbial biomass: A meta-analysis
of field studies. Biogeochemistry, 109(1/3), 49–61.
Dove, N. C., & Hart, S. C. (2017). Fire Reduces Fungal Species Richness and In Situ
Mycorrhizal Colonization: A Meta-Analysis. Fire Ecology, 13(2), 37–65.
https://doi.org/10.4996/fireecology.130237746
Dove, N. C., Klingeman, D. M., Carrell, A. A., Cregger, M. A., & Schadt, C. W. (2021). Fire
alters plant microbiome assembly patterns: Integrating the plant and soil microbial

response to disturbance. New Phytologist, 230(6), 2433–2446.
https://doi.org/10.1111/nph.17248
Dunn, P. H., Wells, W. G. I., Dickey, J., & Wohlgemuth, M. (1982). Role of Fungi in Postfire
Stabilization of Chaparral Ash Beds (PSW-58, pp. 378–381) [Gen. Tech. Rep]. Pacific
Southwest Forest and Range Experiment Station, Forest Service, U.S.U.S. Department of
Agriculture.
Eisenhauer, N., Schulz, W., Scheu, S., & Jousset, A. (2013). Niche dimensionality links
biodiversity and invasibility of microbial communities. Functional Ecology, 27(1), 282–
288. https://doi.org/10.1111/j.1365-2435.2012.02060.x
Emerson, M. R. (1948). Chemical Activation of Ascospore Germination in Neurospora crassa1.
Journal of Bacteriology, 55(3), 327–330.
Enright, D. J., Frangioso, K. M., Isobe, K., Rizzo, D. M., & Glassman, S. I. (2022). Mega-fire in
Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and Selects for
Pyrophilous Taxa that are Phylogenetically Conserved. Molecular Ecology.
https://doi.org/10.1111/mec.16399
Espinosa-de-los-Monteros, J., Martinez, A., & Valle, F. (2001). Metabolic profiles and aprE
expression in anaerobic cultures of Bacillus subtilis using nitrate as terminal electron
acceptor. Applied Microbiology and Biotechnology, 57(3), 379–384.
https://doi.org/10.1007/s002530100749
Evans, S. E., & Wallenstein, M. D. (2012). Soil microbial community response to drying and
rewetting stress: Does historical precipitation regime matter? Biogeochemistry, 109(1/3),
101–116.
Feng, S., & Fu, Q. (2013). Expansion of global drylands under a warming climate. Atmospheric
Chemistry and Physics, 13(19), 10081–10094. https://doi.org/10.5194/acp-13-10081-2013
Fernandez, C. W., Nguyen, N. H., Stefanski, A., Han, Y., Hobbie, S. E., Montgomery, R. A.,
Reich, P. B., & Kennedy, P. G. (2017). Ectomycorrhizal fungal response to warming is
linked to poor host performance at the boreal-temperate ecotone. Global Change Biology,
23(4), 1598–1609. https://doi.org/10.1111/gcb.13510
Ferrenberg, S., O'Neill, S. P., Knelman, J. E., Todd, B., Duggan, S., Bradley, D., Robinson, T.,
Schmidt, S. K., Townsend, A. R., Williams, M. W., Cleveland, C. C., Melbourne, B. A.,
Jiang, L., & Nemergut, D. R. (2013). Changes in assembly processes in soil bacterial
communities following a wildfire disturbance. The ISME Journal, 7(6), 1102–1111.
https://doi.org/10.1038/ismej.2013.11
Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of Soil Microbial
Community Structure by Use of Taxon-Specific Quantitative PCR Assays. Applied and
Environmental Microbiology, 71(7), 4117–4120. https://doi.org/10.1128/AEM.71.7.4117-
4120.2005
Fischer, M. S., Stark, F. G., Berry, T. D., Zeba, N., Whitman, T., & Traxler, M. F. (2021).
Pyrolyzed Substrates Induce Aromatic Compound Metabolism in the Post-fire Fungus,
Pyronema domesticum. Frontiers in Microbiology, 12, 729289.
https://doi.org/10.3389/fmicb.2021.729289
Fox, S., Sikes, B. A., Brown, S. P., Cripps, C. L., Glassman, S. I., Hughes, K., Semenova-Nelsen,
T., & Jumpponen, A. (2022). Fire as a driver of fungal diversity—A synthesis of current
knowledge. Mycologia, 0(0), 1–27. https://doi.org/10.1080/00275514.2021.2024422

Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches, Species
Pools, and Priority Effects. Annual Review of Ecology Evolution and Systematics, 46, 1–
23. https://doi.org/10.1146/annurev-ecolsys-110411-160340
Gao, C., Courty, P.-E., Varoquaux, N., Cole, B., Montoya, L., Xu, L., Purdom, E., Vogel, J.,
Hutmacher, R. B., Dahlberg, J. A., Coleman-Derr, D., Lemaux, P. G., & Taylor, J. W.
(2022). Successional adaptive strategies revealed by correlating arbuscular mycorrhizal
fungal abundance with host plant gene expression. Molecular Ecology, n/a(n/a).
https://doi.org/10.1111/mec.16343
Gao, C., Montoya, L., Xu, L., Madera, M., Hollingsworth, J., Purdom, E., Singan, V., Vogel, J.,
Hutmacher, R. B., Dahlberg, J. A., Coleman-Derr, D., Lemaux, P. G., & Taylor, J. W.
(2020). Fungal community assembly in drought-stressed sorghum shows stochasticity,
selection, and universal ecological dynamics. Nature Communications, 11(1), 34.
https://doi.org/10.1038/s41467-019-13913-9
Gassibe, P. V., Fabero, R. F., Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto, P.
(2011). Fungal community succession following wildfire in a Mediterranean vegetation
type dominated by Pinus pinaster in Northwest Spain. Forest Ecology and Management,
262(4), 655–662. https://doi.org/10.1016/j.foreco.2011.04.036
Glassman, S. I., Levine, C. R., DiRocco, A. M., Battles, J. J., & Bruns, T. D. (2016).
Ectomycorrhizal fungal spore bank recovery after a severe forest fire: Some like it hot.
The ISME Journal, 10(5), 1228–1239. https://doi.org/10.1038/ismej.2015.182
Glassman, S. I., Randolph, J. W., Saroa, S. S., Capocchi, J. K., Walters, K. E., Pulido-Chavez, M.
F., & Larios, L. (2021). Prescribed versus wildfire impacts on exotic plants and soil
microbes in California grasslands (p. 2021.09.22.461426).
https://doi.org/10.1101/2021.09.22.461426
Glassman, S. I., Weihe, C., Li, J., Albright, M. B. N., Looby, C. I., Martiny, A. C., Treseder, K.
K., Allison, S. D., & Martiny, J. B. H. (2018). Decomposition responses to climate
depend on microbial community composition. Proceedings of the National Academy of
Sciences, 115(47), 11994–11999. https://doi.org/10.1073/pnas.1811269115
Gottlieb, D. (1950). The physiology of spore germination in fungi. The Botanical Review, 16(5),
229–257. https://doi.org/10.1007/BF02873609
Grady, E. N., MacDonald, J., Liu, L., Richman, A., & Yuan, Z.-C. (2016). Current knowledge
and perspectives of Paenibacillus: A review. Microbial Cell Factories, 15(1), 203.
https://doi.org/10.1186/s12934-016-0603-7
Gray, L., & Kernaghan, G. (2020). Fungal Succession During the Decomposition of
Ectomycorrhizal Fine Roots. Microbial Ecology, 79(2), 271–284.
https://doi.org/10.1007/s00248-019-01418-3
Grime, J. P. (1977). Evidence for the Existence of Three Primary Strategies in Plants and Its
Relevance to Ecological and Evolutionary Theory. The American Naturalist, 111(982),
1169–1194.
Guhr, A., & Kircher, S. (2020). Drought-Induced Stress Priming in Two Distinct Filamentous
Saprotrophic Fungi. Microbial Ecology, 80(1), 27–33. https://doi.org/10.1007/s00248-
019-01481-w
Hallett, L. M., Jones, S. K., MacDonald, A. A. M., Jones, M. B., Flynn, D. F. B., Ripplinger, J.,
Slaughter, P., Gries, C., & Collins, S. L. (2016). codyn: An r package of community

dynamics metrics. Methods in Ecology and Evolution, 7(10), 1146–1151.
https://doi.org/10.1111/2041-210X.12569
Hanes, T. L. (1971). Succession after Fire in the Chaparral of Southern California. 41(1), 27–52.
https://doi.org/doi:10.2307/1942434
Hart, S. C., DeLuca, T. H., Newman, G. S., MacKenzie, M. D., & Boyle, S. I. (2005). Post-fire
vegetative dynamics as drivers of microbial community structure and function in forest
soils. Forest Ecology and Management, 220(1–3), 166–184.
https://doi.org/10.1016/j.foreco.2005.08.012
Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto, P. (2013). Post-fire fungal
succession in a Mediterranean ecosystem dominated by Cistus ladanifer L. Forest Ecology
and Management, 289, 48–57. https://doi.org/10.1016/j.foreco.2012.10.009
Homyak, P. M., Sickman, J. O., Miller, A. E., Melack, J. M., Meixner, T., & Schimel, J. P.
(2014). Assessing Nitrogen-Saturation in a Seasonally Dry Chaparral Watershed:
Limitations of Traditional Indicators of N-Saturation. Ecosystems, 17(7), 1286–1305.
https://doi.org/10.1007/s10021-014-9792-2
Iffis, B., St-Arnaud, M., & Hijri, M. (2014). Bacteria associated with arbuscular mycorrhizal
fungi within roots of plants growing in a soil highly contaminated with aliphatic and
aromatic petroleum hydrocarbons. FEMS Microbiology Letters, 358(1), 44–54.
https://doi.org/10.1111/1574-6968.12533
Jain, T. B., Gould, W. A., Graham, R. T., Pilliod, D. S., Lentile, L. B., & González, G. (2008). A
Soil Burn Severity Index for Understanding Soil-fire Relations in Tropical Forests.
AMBIO: A Journal of the Human Environment, 37(7), 563–568.
https://doi.org/10.1579/0044-7447-37.7.563
James, S. (1984). Lignotubers and Burls: Their Structure, Function and Ecological Significance
in Mediterranean Ecosystems. Botanical Review, 50(3), 225–266.
Kalayu, G. (2019). Phosphate Solubilizing Microorganisms: Promising Approach as
Biofertilizers. International Journal of Agronomy, 2019, 1–7.
https://doi.org/10.1155/2019/4917256
Kaur, N., Seuylemezian, A., Patil, P. P., Patil, P., Krishnamurti, S., Varelas, J., Smith, D. J.,
Mayilraj, S., & Vaishampayan, P. (2018). Paenibacillus xerothermodurans sp. Nov., an
extremely dry heat resistant spore forming bacterium isolated from the soil of Cape
Canaveral, Florida. International Journal of Systematic and Evolutionary Microbiology,
68(10), 3190–3196. https://doi.org/10.1099/ijsem.0.002967
Keeley, J. E., Fotheringham, C. J., & Baer-Keeley, M. (2005). Determinants of Postfire Recovery
and Succession in Mediterranean-Climate Shrublands of California. Ecological
Applications, 15(5), 1515–1534. https://doi.org/10.1890/04-1005
Keeley, J. E., & Zedler, P. H. (2009). Large, high-intensity fire events in Southern California
shrublands: Debunking the fine-grain age patch model. In Ecological Applications (Vol.
19, Issue 1, p. 26). https://doi.org/10.1890/08-0281.1
Kennedy, P. G., Peay, K. G., & Bruns, T. D. (2009). Root tip competition among ectomycorrhizal
fungi: Are priority effects a rule or an exception? Ecology, 90(8), 2098–2107.
https://doi.org/10.1890/08-1291.1

Kindt, R., & Coe, R. (2005). Tree diversity analysis. A manual and software for common
statistical methods for ecological and biodiversity studies. World Agroforestry Centre
(ICRAF).
Kinzig, A. P., & Pacala, S. (2013). 9. Successional Biodiversity and Ecosystem Functioning. In
The Functional Consequences of Biodiversity: Empirical Progress and Theoretical
Extensions (MPB-33) (pp. 175–212). Princeton University Press.
https://doi.org/10.1515/9781400847303.175
Koide, R. T., Fernandez, C., & Malcolm, G. (2014). Determining place and process: Functional
traits of ectomycorrhizal fungi that affect both community structure and ecosystem
function. New Phytologist, 201(2), 433–439. https://doi.org/10.1111/nph.12538
Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013).
Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing
Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. Applied and
Environmental Microbiology, 79(17), 5112–5120. https://doi.org/10.1128/AEM.01043-13
Krappmann, S., & Braus, G. H. (2005). Nitrogen metabolism of Aspergillus and its role in
pathogenicity. Medical Mycology, 43(s1), 31–40.
https://doi.org/10.1080/13693780400024271
Lauber, C. L., Zhou, N., Gordon, J. I., Knight, R., & Fierer, N. (2010). Effect of storage
conditions on the assessment of bacterial community structure in soil and human-
associated samples. FEMS Microbiology Letters, 307(1), 80–86.
https://doi.org/10.1111/j.1574-6968.2010.01965.x
Legg, A. (1992). Fire-site agarics. Mycologist, 6(3), 145–149. https://doi.org/10.1016/S0269-
915X(09)80604-6
Lehman, C. L., & Tilman, D. (2000). Biodiversity, Stability, and Productivity in Competitive
Communities. The American Naturalist, 156(5), 534–552. https://doi.org/10.1086/303402
Leizeaga, A., Meisner, A., Rousk, J., & Bååth, E. (2022). Repeated drying and rewetting cycles
accelerate bacterial growth recovery after rewetting. Biology and Fertility of Soils.
https://doi.org/10.1007/s00374-022-01623-2
Lekberg, Y., Vasar, M., Bullington, L. S., Sepp, S.-K., Antunes, P. M., Bunn, R., Larkin, B. G.,
& Öpik, M. (2018). More bang for the buck? Can arbuscular mycorrhizal fungal
communities be characterized adequately alongside other fungi using general fungal
primers? New Phytologist, 220(4), 971–976. https://doi.org/10.1111/nph.15035
Li, X., Rui, J., Mao, Y., Yannarell, A., & Mackie, R. (2014). Dynamics of the bacterial
community structure in the rhizosphere of a maize cultivar. Soil Biology and
Biochemistry, 68, 392–401. https://doi.org/10.1016/j.soilbio.2013.10.017
Liu, C. M., Kachur, S., Dwan, M. G., Abraham, A. G., Aziz, M., Hsueh, P.-R., Huang, Y.-T.,
Busch, J. D., Lamit, L. J., Gehring, C. A., Keim, P., & Price, L. B. (2012). FungiQuant: A
broad-coverage fungal quantitative real-time PCR assay. BMC Microbiology, 12(1), 255.
https://doi.org/10.1186/1471-2180-12-255
Livne-Luzon, S., Shemesh, H., Osem, Y., Carmel, Y., Migael, H., Avidan, Y., Tsafrir, A.,
Glassman, S. I., Bruns, T. D., & Ovadia, O. (2021). High resilience of the mycorrhizal
community to prescribed seasonal burnings in eastern Mediterranean woodlands.
Mycorrhiza, 31(2), 203–216. https://doi.org/10.1007/s00572-020-01010-5

Lofgren, L. A., Uehling, J. K., Branco, S., Bruns, T. D., Martin, F., & Kennedy, P. G. (2019).
Genome-based estimates of fungal rDNA copy number variation across phylogenetic
scales and ecological lifestyles. Molecular Ecology, 28(4), 721–730.
https://doi.org/10.1111/mec.14995
Loreau, M., & de Mazancourt, C. (2008). Species Synchrony and Its Drivers: Neutral and
Nonneutral Community Dynamics in Fluctuating Environments. The American Naturalist,
172(2), E48–E66. https://doi.org/10.1086/589746
Lucas-Borja, M. E., Miralles, I., Ortega, R., Plaza-Álvarez, P. A., Gonzalez-Romero, J., Sagra, J.,
Soriano-Rodríguez, M., Certini, G., Moya, D., & Heras, J. (2019). Immediate fire-induced
changes in soil microbial community composition in an outdoor experimental controlled
system. Science of The Total Environment, 696, 134033.
https://doi.org/10.1016/j.scitotenv.2019.134033
Mandic-Mulec, I., Stefanic, P., & van Elsas, J. D. (2015). Ecology of Bacillaceae. Microbiology
Spectrum, 3(2), 3.2.16. https://doi.org/10.1128/microbiolspec.TBS-0017-2013
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
https://journal.embnet.org/index.php/embnetjournal/article/view/200/458
Mayer, M., Rewald, B., Matthews, B., Sandén, H., Rosinger, C., Katzensteiner, K., Gorfer, M.,
Berger, H., Tallian, C., Berger, T. W., & Godbold, D. L. (2021). Soil fertility relates to
fungal-mediated decomposition and organic matter turnover in a temperate mountain
forest. New Phytologist, 231(2), 777–790. https://doi.org/10.1111/nph.17421
McGee, P. A., Markovina, A.-L., Jeong, G. C. E., & Cooper, E. D. (2006). Trichocomaceae in
bark survive high temperatures and fire: Trichocomaceae in bark survive high
temperatures and fire. FEMS Microbiology Ecology, 56(3), 365–371.
https://doi.org/10.1111/j.1574-6941.2006.00079.x
McMullan-Fisher, S. J. M., Tom W. May, Richard M. Robinson, Tina L. Bell, Teresa Lebel, Pam
Catcheside, & Alan York. (2011). Fungi and fire in Australian ecosystems: A review of
current knowledge, management implications and future directions. 59, 70–90.
McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE, 8(4), e61217.
https://doi.org/10.1371/journal.pone.0061217
Monciardini, P., Cavaletti, L., Schumann, P., Rohde, M., & Donadio, S. (2003). Conexibacter
woesei gen. Nov., sp. Nov., a novel representative of a deep evolutionary line of descent
within the class Actinobacteria. International Journal of Systematic and Evolutionary
Microbiology, 53(2), 569–576. https://doi.org/10.1099/ijs.0.02400-0
Moore, E. J. (1962). The Ontogeny of the Sclerotia of Pyronema Domesticum. Mycologia, 54(3),
312–316. https://doi.org/10.1080/00275514.1962.12025004
Neary, D. G., Klopatek, C. C., DeBano, L. F., & Ffolliott, P. F. (1999). Fire effects on
belowground sustainability: A review and synthesis. Forest Ecology and Management,
122(1–2), 51–71. https://doi.org/10.1016/S0378-1127(99)00032-8
Nelson, A. R., Narrowe, A. B., Rhoades, C. C., Fegel, T. S., Daly, R. A., Roth, H. K., Chu, R. K.,
Amundson, K. K., Young, R. B., Steindorff, A. S., Mondo, S. J., Grigoriev, I. V.,
Salamov, A., Borch, T., & Wilkins, M. J. (2022). Wildfire-dependent changes in soil
microbiome diversity and function. Nature Microbiology, 7(9), 1–12.
https://doi.org/10.1038/s41564-022-01203-y

Nguyen, H. D., Nickerson, N. L., & Seifert, K. A. (2013). Basidioascus and Geminibasidium: A
new lineage of heat-resistant and xerotolerant basidiomycetes. Mycologia, 105(5), 1231–
1250. https://doi.org/10.3852/12-351
Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., &
Kennedy, P. G. (2016). FUNGuild: An open annotation tool for parsing fungal
community datasets by ecological guild. Fungal Ecology, 20, 241–248.
https://doi.org/10.1016/j.funeco.2015.06.006
Ofek, M., Hadar, Y., & Minz, D. (2012). Ecology of Root Colonizing Massilia
(Oxalobacteraceae). PLOS ONE, 7(7), e40117.
https://doi.org/10.1371/journal.pone.0040117
Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.
(2018). Vegan:Community Ecology Package-Version 2.5.2. Ordination Methods,
Diversity Analysis, and Other Functions for Community and Vegetation Ecologist.
Osborne, B. B., Bestelmeyer, B. T., Currier, C. M., Homyak, P. M., Throop, H. L., Young, K., &
Reed, S. C. (2022). The consequences of climate change for dryland biogeochemistry.
New Phytologist, 236(1), 15–20. https://doi.org/10.1111/nph.18312
Owen, S. M., Patterson, A. M., Gehring, C. A., Sieg, C. H., Baggett, L. S., & Fulé, P. Z. (2019).
Large, high-severity burn patches limit fungal recovery 13 years after wildfire in a
ponderosa pine forest. Soil Biology and Biochemistry, 139, 107616.
https://doi.org/10.1016/j.soilbio.2019.107616
Pacala, S. W., Canham, C. D., Saponara, J., Silander, J. A., Kobe, R. K., & Ribbens, E. (1996).
Forest Models Defined by Field Measurements: Estimation, Error Analysis and
Dynamics. Ecological Monographs, 66(1), 1–43. https://doi.org/10.2307/2963479
Pandolfi, J. M. (2008). Succession. In S. E. Jørgensen & B. D. Fath (Eds.), Encyclopedia of
Ecology (pp. 3416–3424). Academic Press. https://doi.org/10.1016/B978-008045405-
4.00547-4
Paredes-Sabja, D., Setlow, P., & Sarker, M. R. (2011). Germination of spores of Bacillales and
Clostridiales species: Mechanisms and proteins involved. Trends in Microbiology, 19(2),
85–94. https://doi.org/10.1016/j.tim.2010.10.004
Parson, A., Robichaud, P. R., Lewis, S. A., Napper, C., & Clark, J. T. (2010). Field guide for
mapping post-fire soil burn severity (RMRS-GTR-243). U.S. Department of Agriculture,
Forest Service, Rocky Mountain Research Station. https://doi.org/10.2737
Pavlovska, M., Prekrasna, I., Parnikoza, I., & Dykyi, E. (2021). Soil Sample Preservation
Strategy Affects the Microbial Community Structure. Microbes and Environments, 36(1),
ME20134. https://doi.org/10.1264/jsme2.ME20134
Pena, R., Offermann, C., Simon, J., Naumann, P. S., Geßler, A., Holst, J., Dannenmann, M.,
Mayer, H., Kögel-Knabner, I., Rennenberg, H., & Polle, A. (2010). Girdling Affects
Ectomycorrhizal Fungal (EMF) Diversity and Reveals Functional Differences in EMF
Community Composition in a Beech Forest. Applied and Environmental Microbiology,
76(6), 1831–1841. https://doi.org/10.1128/AEM.01703-09
Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2018). Resilience to fire of
phylogenetic diversity across biological domains. Molecular Ecology, 27(13), 2896–2908.
https://doi.org/10.1111/mec.14729

Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2020). Soil microbiome drives
the recovery of ecosystem functions after fire. Soil Biology and Biochemistry, 149,
107948. https://doi.org/10.1016/j.soilbio.2020.107948
Petersen, P. M. (1970). Danish fireplace fungi-an ecological investigation on fungi on burns.
Dansk Botanisk Arkiv, 27(3), 1–97.
Phillips, L. (2021, March 18). General Field Plot Sampling Protocol for DNA-based analyses
(2018-2020). Protocols.Io. https://www.protocols.io/view/general-field-plot-sampling-
protocol-for-dna-based-btd2ni8e
Pingree, M. R. A., & Kobziar, L. N. (2019). The myth of the biological threshold: A review of
biological responses to soil heating associated with wildland fire. Forest Ecology and
Management, 432, 1022–1029. https://doi.org/10.1016/j.foreco.2018.10.032
Placella, S. A., Brodie, E. L., & Firestone, M. K. (2012). Rainfall-induced carbon dioxide pulses
result from sequential resuscitation of phylogenetically clustered microbial groups.
Proceedings of the National Academy of Sciences, 109(27), 10931–10936.
https://doi.org/10.1073/pnas.1204306109
Platt, W. J., & Connell, J. H. (2003). Natural Disturbances and Directional Replacement of
Species. Ecological Monographs, 73(4), 507–522.
Pourreza, M., Hosseini, S. M., Safari Sinegani, A. A., Matinizadeh, M., & Dick, W. A. (2014).
Soil microbial activity in response to fire severity in Zagros oak (Quercus brantii Lindl.)
forests, Iran, after one year. Geoderma, 213, 95–102.
https://doi.org/10.1016/j.geoderma.2013.07.024
Pressler, Y., Moore, J. C., & Cotrufo, M. F. (2019). Belowground community responses to fire:
Meta-analysis reveals contrasting responses of soil microorganisms and mesofauna.
Oikos, 128(3), 309–327. https://doi.org/10.1111/oik.05738
Pulido-Chavez, M. F., Alvarado, E. C., DeLuca, T. H., Edmonds, R. L., & Glassman, S. I. (2021).
High-severity wildfire reduces richness and alters composition of ectomycorrhizal fungi
in low-severity adapted ponderosa pine forests. Forest Ecology and Management, 485,
118923. https://doi.org/10.1016/j.foreco.2021.118923
Qin, Q., & Liu, Y. (2021). Changes in microbial communities at different soil depths through the
first rainy season following severe wildfire in North China artificial Pinus tabulaeformis
forest. Journal of Environmental Management, 280, 111865.
https://doi.org/10.1016/j.jenvman.2020.111865
R Core Team. (2021). R: A language and environment for statistical computing. R Foundation for
Statistical Computing. https://www.R-project.org/.
Reazin, C., Morris, S., Smith, J. E., Cowan, A. D., & Jumpponen, A. (2016). Fires of differing
intensities rapidly select distinct soil fungal communities in a Northwest US ponderosa
pine forest ecosystem. Forest Ecology and Management, 377, 118–127.
https://doi.org/10.1016/j.foreco.2016.07.002
Reilly, M. J., & Spies, T. A. (2016). Disturbance, tree mortality, and implications for
contemporary regional forest change in the Pacific Northwest. Forest Ecology and
Management, 374, 102–110. https://doi.org/10.1016/j.foreco.2016.05.002
Ren, C., Chen, J., Lu, X., Doughty, R., Zhao, F., Zhong, Z., Han, X., Yang, G., Feng, Y., & Ren,
G. (2018). Responses of soil total microbial biomass and community compositions to

rainfall reductions. Soil Biology and Biochemistry, 116, 4–10.
https://doi.org/10.1016/j.soilbio.2017.09.028
Rhodes, J. C. (2006). Aspergillus fumigatus: Growth and virulence. Medical Mycology, 44(s1),
77–81. https://doi.org/10.1080/13693780600779419
Rice, S. K. (1993). Vegetation establishment in post-fire Adenostoma chaparral in relation to
fine-scale pattern in fire intensity and soil nutrients. Journal of Vegetation Science, 4(1),
115–124. https://doi.org/10.2307/3235739
Riley, K. L., & Loehman, R. A. (2016). Mid-21st-century climate changes increase predicted fire
occurrence and fire season length, Northern Rocky Mountains, United States. Ecosphere,
7(11), e01543. https://doi.org/10.1002/ecs2.1543
Ross, G. J. S., & Preece, D. A. (1985). The Negative Binomial Distribution. Journal of the Royal
Statistical Society. Series D (The Statistician), 34(3), 323–335. JSTOR.
https://doi.org/10.2307/2987659
Rundel, P. W. (2018). California Chaparral and Its Global Significance. In E. C. Underwood, H.
D. Safford, N. A. Molinari, & J. E. Keeley (Eds.), Valuing Chaparral (pp. 1–27). Springer
International Publishing. https://doi.org/10.1007/978-3-319-68303-4_1
Sáenz de Miera, L. E., Pinto, R., Gutierrez-Gonzalez, J. J., Calvo, L., & Ansola, G. (2020).
Wildfire effects on diversity and composition in soil bacterial communities. Science of
The Total Environment, 726, 138636. https://doi.org/10.1016/j.scitotenv.2020.138636
Schimel, J. P. (2018). Life in Dry Soils: Effects of Drought on Soil Microbial Communities and
Processes. Annual Review of Ecology, Evolution, and Systematics, 49(1), 409–432.
https://doi.org/10.1146/annurev-ecolsys-110617-062614
Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W.,
Fungal Barcoding Consortium, Fungal Barcoding Consortium Author List, Bolchacova,
E., Voigt, K., Crous, P. W., Miller, A. N., Wingfield, M. J., Aime, M. C., An, K.-D., Bai,
F.-Y., Barreto, R. W., Begerow, D., … Schindel, D. (2012). Nuclear ribosomal internal
transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi.
Proceedings of the National Academy of Sciences, 109(16), 6241–6246.
https://doi.org/10.1073/pnas.1117018109
Seaver, F. J. (1909). Studies in Pyrophilous Fungi: I. The Occurrence and Cultivation of
Pyronema. Mycologia, 1, 131–193.
Selbmann, L., Egidi, E., Isola, D., Onofri, S., Zucconi, L., de Hoog, G. S., Chinaglia, S., Testa,
L., Tosi, S., Balestrazzi, A., Lantieri, A., Compagno, R., Tigini, V., & Varese, G. C.
(2013). Biodiversity, evolution and adaptation of fungi in extreme environments. Plant
Biosystems - An International Journal Dealing with All Aspects of Plant Biology, 147(1),
237–246. https://doi.org/10.1080/11263504.2012.753134
Semenova-Nelsen, T. A., Platt, W. J., Patterson, T. R., Huffman, J., & Sikes, B. A. (2019).
Frequent fire reorganizes fungal communities and slows decomposition across a
heterogeneous pine savanna landscape. New Phytologist, 224(2), 916–927.
https://doi.org/10.1111/nph.16096
Shugart, H. H. (2013). Encyclopedia of Biodiversity. In Succession, Phenomenon Of (2nd
Edition, pp. 63–70). Academic Press.

Sirois, S. H., & Buckley, D. H. (2019). Factors governing extracellular DNA degradation
dynamics in soil. Environmental Microbiology Reports, 11(2), 173–184.
https://doi.org/10.1111/1758-2229.12725
Slepecky, R. A., & Hemphill, H. E. (2006). The Genus Bacillus—Nonmedical. In M. Dworkin, S.
Falkow, E. Rosenberg, K.-H. Schleifer, & E. Stackebrandt (Eds.), The Prokaryotes:
Volume 4: Bacteria: Firmicutes, Cyanobacteria (pp. 530–562). Springer US.
https://doi.org/10.1007/0-387-30744-3_16
Smith, C. J., & Osborn, A. M. (2009). Advantages and limitations of quantitative PCR (Q-PCR)-
based approaches in microbial ecology. FEMS Microbiology Ecology, 67(1), 6–20.
https://doi.org/10.1111/j.1574-6941.2008.00629.x
Song, Z., Vail, A., Sadowsky, M. J., & Schilling, J. S. (2014). Quantitative PCR for measuring
biomass of decomposer fungi in planta. Fungal Ecology, 7, 39–46.
https://doi.org/10.1016/j.funeco.2013.12.004
Steindorff, A. S., Carver, A., Calhoun, S., Stillman, K., Liu, H., Lipzen, A., He, G., Yan, M.,
Pangilinan, J., LaButti, K., Ng, V., Bruns, T. D., & Grigoriev, I. V. (2021). Comparative
genomics of pyrophilous fungi reveals a link between fire events and developmental
genes. Environmental Microbiology, 23(1), 99–109. https://doi.org/10.1111/1462-
2920.15273
Suzuki, A. (2017). Various Aspects of Ammonia Fungi. In T. Satyanarayana, S. K. Deshmukh, &
B. N. Johri (Eds.), Developments in Fungal Biology and Applied Mycology (pp. 39–58).
Springer. https://doi.org/10.1007/978-981-10-4768-8_4
Tatti, E., McKew, B. A., Whitby, C., & Smith, C. J. (2016). Simultaneous DNA-RNA Extraction
from Coastal Sediments and Quantification of 16S rRNA Genes and Transcripts by Real-
time PCR. Journal of Visualized Experiments : JoVE, 112, 54067.
https://doi.org/10.3791/54067
Taylor, D. L., Walters, W. A., Lennon, N. J., Bochicchio, J., Krohn, A., Caporaso, J. G., &
Pennanen, T. (2016). Accurate Estimation of Fungal Diversity and Abundance through
Improved Lineage-Specific Primers Optimized for Illumina Amplicon Sequencing.
Applied and Environmental Microbiology, 82(24), 7217–7226.
https://doi.org/10.1128/AEM.02576-16
Tellenbach, C., Grünig, C. R., & Sieber, T. N. (2010). Suitability of Quantitative Real-Time PCR
To Estimate the Biomass of Fungal Root Endophytes. Applied and Environmental
Microbiology, 76(17), 5764–5772. https://doi.org/10.1128/AEM.00907-10
Tilman, D. (1990). Constraints and Tradeoffs: Toward a Predictive Theory of Competition and
Succession. Oikos, 58(1), 3–15. https://doi.org/10.2307/3565355
Tilman, D. (1999). The Ecological Consequences of Changes in Biodiversity: A Search for
General Principles. Ecology, 80(5), 1455–1474. https://doi.org/10.1890/0012-
9658(1999)080[1455:TECOCI]2.0.CO;2
Tilman, D., Isbell, F., & Cowles, J. M. (2014). Biodiversity and Ecosystem Functioning. Annual
Review of Ecology, Evolution, and Systematics, 45(1), 471–493.
https://doi.org/10.1146/annurev-ecolsys-120213-091917
Toljander, J. F., Artursson, V., Paul, L. R., Jansson, J. K., & Finlay, R. D. (2005). Attachment of
di¡erent soil bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined
by hyphal vitalityand fungal species. FEMS Microbiol Lett, 7.

Torti, A., Lever, M. A., & Jørgensen, B. B. (2015). Origin, dynamics, and implications of
extracellular DNA pools in marine sediments. Marine Genomics, 24, 185–196.
https://doi.org/10.1016/j.margen.2015.08.007
Turnbull, L. A., Levine, J. M., Loreau, M., & Hector, A. (2013). Coexistence, niches and
biodiversity effects on ecosystem functioning. Ecology Letters, 16, 116–127.
https://doi.org/10.1111/ele.12056
Valencia, E., de Bello, F., Galland, T., Adler, P. B., Lepš, J., E-Vojtkó, A., van Klink, R.,
Carmona, C. P., Danihelka, J., Dengler, J., Eldridge, D. J., Estiarte, M., García-González,
R., Garnier, E., Gómez-García, D., Harrison, S. P., Herben, T., Ibáñez, R., Jentsch, A., …
Götzenberger, L. (2020). Synchrony matters more than species richness in plant
community stability at a global scale. Proceedings of the National Academy of Sciences,
117(39), 24345–24351. https://doi.org/10.1073/pnas.1920405117
Venables, W. N., & Ripley, B. D. (2002). Modern Applied Statistics with S (Fourth edition).
Springer. https://www.stats.ox.ac.uk/pub/MASS4/.
Warcup, J. H., & Baker, K. F. (1963). Occurrence of Dormant Ascospores in Soil. Nature,
197(4874), 1317–1318. https://doi.org/10.1038/1971317a0
Whitman, T., Whitman, E., Woolet, J., Flannigan, M. D., Thompson, D. K., & Parisien, M.-A.
(2019). Soil bacterial and fungal response to wildfires in the Canadian boreal forest across
a burn severity gradient. Soil Biology and Biochemistry, 138, 107571.
https://doi.org/10.1016/j.soilbio.2019.107571
Whitman, T., Woolet, J., Sikora, M., Johnson, D. B., & Whitman, E. (2022). Resilience in soil
bacterial communities of the boreal forest from one to five years after wildfire across a
severity gradient. Soil Biology and Biochemistry, 172, 108755.
https://doi.org/10.1016/j.soilbio.2022.108755
Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
https://ggplot2.tidyverse.org
Willis, A., Bunge, J., & Whitman, T. (2017). Improved detection of changes in species richness
in high diversity microbial communities. Journal of the Royal Statistical Society. Series C
(Applied Statistics), 66(5), 963–977.
Woolet, J., & Whitman, T. (2020). Pyrogenic organic matter effects on soil bacterial community
composition. Soil Biology and Biochemistry, 141, 107678.
https://doi.org/10.1016/j.soilbio.2019.107678
Xiang, X., Gibbons, S. M., Yang, J., Kong, J., Sun, R., & Chu, H. (2015). Arbuscular
mycorrhizal fungal communities show low resistance and high resilience to wildfire
disturbance. Plant and Soil, 397(1–2), 347–356. https://doi.org/10.1007/s11104-015-2633-
z
Xiang, X., Shi, Y., Yang, J., Kong, J., Lin, X., Zhang, H., Zeng, J., & Chu, H. (2014). Rapid
recovery of soil bacterial communities after wildfire in a Chinese boreal forest. Scientific
Reports, 4(1), 3829. https://doi.org/10.1038/srep03829
Yachi, S., & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating
environment: The insurance hypothesis. Proceedings of the National Academy of
Sciences, 96(4), 1463–1468. https://doi.org/10.1073/pnas.96.4.1463
Yang, T., Tedersoo, L., Lin, X., Fitzpatrick, M. C., Jia, Y., Liu, X., Ni, Y., Shi, Y., Lu, P., Zhu,
J., & Chu, H. (2020). Distinct fungal successional trajectories following wildfire between

soil horizons in a cold-temperate forest. New Phytologist, 227(2), 572–587.
https://doi.org/10.1111/nph.16531
Yang, Zhu, K., Loik, M. E., & Sun, W. (2021). Differential responses of soil bacteria and fungi to
altered precipitation in a meadow steppe. Geoderma, 384, 114812.
https://doi.org/10.1016/j.geoderma.2020.114812
Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J.,
Ludwig, W., & Glöckner, F. O. (2014). The SILVA and "All-species Living Tree Project
(LTP)" taxonomic frameworks. Nucleic Acids Research, 42(D1), D643–D648.
https://doi.org/10.1093/nar/gkt1209
Zhang, H., Qi, W., & Liu, K. (2018). Functional traits associated with plant colonizing and
competitive ability influence species abundance during secondary succession: Evidence
from subalpine meadows of the Qinghai–Tibetan Plateau. Ecology and Evolution, 8(13),
6529–6536. https://doi.org/10.1002/ece3.4110

1357
1358
1359
1360

Data Accessibility

Raw sequence reads were submitted to the National Center for Biotechnology Information Sequence Read Archive under BioProject accession number PRJNA761539. All statistical codes are available on GitHub https://github.com/pulidofabs/SecondarySuccession-Chaparral.

Benefit-Sharing

Nagoya Protocol is not applicable as there is no genetic material to report. However, benefits from this research ensue from the sharing of our data and results on public databases as described above.

Author's contribution:

SIG conceived of the study and acquired permits; LL, SIG, and PMH came up with the experimental design; SIG, LL, PMH, MFPC, and JWJR performed fieldwork; MFPC and JWJR performed all molecular work; MFPC performed all bioinformatics, statistical analysis, made figures, and wrote the first draft; all authors contributed to writing and editing of the manuscript.

2
Figure 1. A) Location of the study area and plots (6 burned; 3 unburned) within the Holy Fire
burn scar in the Cleveland National Forest in southern California. Soil burn severity is based on
the USDA BAER classifications made at seven days within fire containment. B) Plot
experimental design, each with four 1m² subplots placed 5m from the center in each cardinal
direction. C) Collection of the top 10 cm of soil beneath the ash with a releasable bulb planter
within the 1 m² subplots at 17 days post-fire.
9
10
11

Figure 2. Change between burned (brown) and unburned communities (blue-green) for A) bacterial and B) fungal estimated biomass per gram of soil and C) bacterial and D) fungal species richness at each of the nine timepoints in the first post-fire year. Points represent the mean biomass (A, B) and richness (C, D), and bars represent the standard error of the mean for burned versus unburned plots at each timepoint. Note that gene copy number and species richness are displayed on a daily scale, whereas precipitation is on a monthly scale representing total precipitation for the month in which each soil sample was collected. Note, we sampled twice in Oct. The significance of burned versus unburned plots per timepoint is denoted with an asterisk and is based on a negative binomial regression.

Figure 3. Change in estimated biomass and richness with soil burn severity measured as ash depth at 17 days post-fire for A) bacterial and B) fungal biomass per gram of soil and C) bacterial and D) fungal richness. The significance of biomass and species richness against soil burn severity (SBS) and its interaction with time since (TSF) is based on a negative binomial regression (Table S4). The blue line represents the model's prediction, and the gray represents the standard error.

33
Figure 4. NMDS plot of A) bacterial and C) fungal community composition with colors denoting
treatment and shape indicating time since fire. The NMDS is based on 3-dimensions and has a
stress value of 0.11 for bacteria and 0.12 for fungi. Change in relative sequence abundance of the
five most abundant B) bacterial and D) fungal genera in the burned plots over time. Note that the
dominance of the top 5 bacterial genera began 34 days post-fire since there was no clear
dominance on days 17 and 25 due to the lag in post-fire bacterial declines.

Bacterial and fungal secondary succession

Figure 5. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in burned plots at all 9 timepoints. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in burned plots. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each timepoint for the burned E) bacterial and F) fungal communities. Note that timepoints farther apart with non-overlapping standard error bars indicate a community turnover, whereas overlapping standard error bars represent a lack of compositional turnover. Note that 17 days represents the base level. Thus, the first turnover was initiated at 25 days. Gray bars represent multiple genera of bacteria (A) and fungi (B) that have relative sequence abundance under 3% at each timepoint per treatment. Note that 47% of the bacterial and 20% of the fungal sequences comprise genera <3% relative abundance.

Figure 6. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in unburned plots at all 9-time points. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in unburned plots displaying no significant succession. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each time point for E) bacteria and F) fungi in the unburned plots. In unburned plots, overlapping standard error bars indicate no community turnover across time. Gray bars represent multiple genera of bacteria (A) and fungi (B) that have relative sequence abundance under 3% at each timepoint per treatment. Note the lack of dominance as denoted by the fact that 79% of the bacterial and 51% of the fungal sequences comprise genera <3% relative abundance.

75 **Table 1.** Pyrophilous bacterial and fungal taxa hypothesized position within Grime's C-S-R
76 succession theory (per traits at the genera level, based on current literature).
77

Kingdom	Phylum	Genus	Grimmes Traits (CSR)	Trait description	Citation
Bacteria	Firmicute (Bacillota)	*Bacillus*	Survival	Aerobic, Heterotrophic, Antibiotic-producing, thermotolerant; endospore; reduces nitrogen; solubilizes phosphate	Grady et al., 2016; Kaur et al., 2018; Mandic-Mulec et al., 2015; J. et al., 2001; Kalayu, 2019; Slepecky and Hemphill in the Prokaryotes; Espinosa-de-los-Monteros et al., 2001
Bacteria	Firmicute (Bacillota)	*Clostridiales*	Survival, Fast growth	Thermotolerant; endospore; trigger sporulation in response to stress, reduces nitrogen	Grady et al., 2016; Kaur et al., 2018; Mandic-Mulec et al., 2015; Paredes-Sabja et al., 2011; Kalayu, 2019
Bacteria	Firmicute (Bacillota)	*Paenibacillus*	Survival	Aerobic, Heterotrophic, antibiotic producing (Bacteriocins), thermotolerant; endospore; fixes nitrogen	Grady et al., 2016; Kaur et al., 2018; Mandic-Mulec et al., 2015; Monciardini et al., 2003; Slepecky & Hemphill, 2006
Bacteria	Actinobacteria (Actinomycetota)	*Conexibacter*	Competition	Aerobic, Heterotrophic, Antibiotic-producing, reduces nitrogen	Espinosa-de-los-Monteros et al., 2001; Kalayu, 2019
Bacteria	Proteobacteria (Pseudomonadota)	*Massilia*	Fast Growth; Competition	Aerobic, fast reproduction, diverse ecology, exploit resources	Li et al., 2014;Toljander et al., 2005
Bacteria	Proteobacteria (Pseudomonadota)	*Novihersparillium*	Fast Growth; Competition	Fast reproduction, exploit resources; likely PyOM degrader	Baldani et al., 2014; Grady et al., 2016; Woolet & Whitman, 2020
Fungi	Basidiomycota	*Geminibasidium*	Survival	thermo- and xero-tolerant yeast	Alexopoulous & Mims, 1952
Fungi	Ascomycota	*Pyronema*	Survival Fast Growth	thermotolerant sclerotia; likely PyOM degrader, filamentous,	Moore, 1962; Fisher et al., 2022
Fungi	Ascomycota	*Gelasinospora heterospora*	Survival	Requires heat or chemical treatment for germination, pigmented mycelia	Alexopoulous & Mims, 1952
Fungi	Ascomycota	*Penicillium*	Fast Growth	copious asexual conidia, fast growth, filamentous	Dix & Webster, 1995; McGee et al., 2006;Crow, 1992
Fungi	Ascomycota	*Aspergillus*	Fast Growth	copious asexual conidia, fast growth, filamentous	Dix & Webster, 1995; McGee et al., 2006;Crow, 1992
Fungi	Basidiomycota	*Coprinellus*	Competition	Likely PyOM degrader, filamentous	Steindorff et al., 2021
Fungi	Basidiomycota	*Tephrocybe anthrocophila*	Competition	Affinity for Ammonium or Nitrogen, filamentous	Suzuki, 2017; Legg, 1992

78
79