1   Bacterial and fungal communities experience rapid succession during the first year following a
2   wildfire in a California chaparral
3
4

5   M. Fabiola Pulido-Chavez[1,*], James W. J. Randolph[1], Cassandra Zalman[2], Loralee Larios[3], Peter
6   M. Homyak[4], and Sydney I. Glassman[1,*]
7
8
[1]Department of Microbiology and Plant Pathology, University of California-Riverside, 900
University Ave, Riverside, CA, 92521
[2]Schmid College of Science and Technology, Chapman University, Orange, CA, 92866
[3]Department of Botany and Plant Sciences, University of California-Riverside, 900 University
Ave, Riverside, CA, 92521
[4]Department of Environmental Sciences, University of California-Riverside, 900 University Ave,
Riverside, CA, 92521
16
17
18
19   *Correspondence: Sydney I Glassman, sydneyg@ucr.edu and M. Fabiola Pulido-Chavez,
20   mpuli011@ucr.ed Department of Microbiology and Plant Pathology, University of California-
21   Riverside

24   **Keywords:** Secondary succession, Chaparral, Wildfire, Bacteria, Fungi, Temporal turnover,
25   pyrophilous microbes

Abstract

Abstract

The rise in wildfire frequency and severity across the globe has increased interest in

secondary succession. However, despite the role of soil microbial communities in the

regeneration of post-fire vegetation and biogeochemical cycling, the lack of measurements

immediately post-fire and at high temporal resolution has resulted in a limited understanding of

microbial secondary succession. To fill this knowledge gap, we collected soils at 17, 25, 34, 67,

95, 131, 187, 286, and 376 days after a Southern California wildfire in fire-adapted chaparral

shrublands. We assessed bacterial and fungal biomass with qPCR of 16S and 18S and richness

and composition with Illumina MiSeq sequencing of 16S and ITS2 amplicons. Fire severely

reduced bacterial biomass by 47% and richness by 46%, but the impacts were stronger for fungi,

with biomass decreasing by 86% and richness by 68%. While bacterial biomass and richness

oscillated in response to precipitation, the declines in biomass and richness for both groups

persisted for the entire post-fire year. In contrast to the unburned plots, fungi and bacteria in the

burned plots experienced rapid succession, with 5-6 compositional turnover periods. Analogous

to plants, fire-adapted, or "fire-loving," pyrophilous microbes drove successional dynamics. The

dominant pyrophilous bacteria and fungi experienced marked trade-offs in abundance over time,

especially for fungi, with an initial peak in the Basidiomycete yeast *Geminibasidium* followed by

trade-offs with the filamentous Ascomycetes *Pyronema, Aspergillus,* and *Penicillium*. For

bacteria, the Proteobacteria genus *Massilia* dominated burned plots throughout the year. Yet, the

following two most dominant bacteria, the Firmicute *Bacillus* and Proteobacteria

Noviherbaspirillum, traded off in abundance over time. Our high temporal sampling allowed us

to capture the dynamics of microbial secondary succession for the first time, which indicates that

the trade-offs among a few dominant pyrophilous microbes control microbial succession and

thus might have important impacts on ecosystem functions.

1. Introduction

The rapid increase in wildfire frequency, severity, and extent in the western United States (Riley & Loehman, 2016) and around the globe (Abatzoglou et al., 2019) has encouraged a resurgence of interest in secondary succession. Secondary succession, or the trajectory along which an ecosystem develops following a disturbance, such as a wildfire, has been extensively studied for plants (Derroire et al., 2016; Donato et al., 2012), but belowground microbial communities have been largely overlooked. Understanding how wildfires alter soil microbial succession is necessary to predict post-fire effects on ecosystem recovery, and function since soil microbes drive post-fire organic matter decomposition (Semenova-Nelsen et al., 2019), nutrient cycling (Pérez-Valera et al., 2020), and plant regeneration via symbiotic mycorrhizal fungi (Dove & Hart, 2017).

Microbial succession is a major process affecting the health and function of all ecosystems. However, wildfire-induced mortality (Hart et al., 2005) can reset the successional clock due to reductions in microbial richness and biomass (Dooley & Treseder, 2012; Pressler et al., 2019) and the replacement of fungal basidiomycetes and symbiotic mycorrhizal fungi with ascomycetes and saprobic fungi (Cairney & Bastias, 2007). Although wildfire impacts on soil microbes are understood, the lack of continuous temporal sampling of disturbed ecosystems has limited our understanding of secondary succession. However, existing studies on primary succession suggest that ecological and physiological differences (i.e., growth rate, stress tolerance, nutrient requirements) (Sun et al., 2017) between bacteria and fungi lead to distinct successional trajectories (Brown & Jumpponen, 2014; Fernández-Martínez et al., 2017), driven by environmental filtering or historical constraints (Schluter & Ricklefs, 1993). In disturbed ecosystems, indirect wildfire effects on the soil environment, including soil burn severity, increased ash deposition, soil pH, and nutrients (Köster et al., 2021; Neaxry et al., 1999), can

restructure microbial communities (Whitman et al., 2019). Moreover, differential responses and

adaptations to soil moisture between bacteria and fungi (Placella et al., 2012) (Evans &

Wallenstein, 2012), suggest that post-fire precipitation can affect soil microbial communities

potentially resulting in distinct successional trajectories, as those observed during primary

succession.

In the context of the secondary succession of early microbial assemblages, historical

constraints, such as fire regime, may reflect microbial turnover since pyrophilous "fire-loving"

microbes have likely evolved under the same fire regime as the ecosystem they inhabit. The

rapid fruiting of pyrophilous fungi after fires has been known for over a century (Seaver, 1909),

and in recent decades next-generation sequencing has revealed that pyrophilous Ascomycota

such as *Pyronema* can increase 100-fold post-fire (Reazin et al., 2016) and dominate over 60%

of sequences in a burned pine forest (Bruns et al., 2020). Furthermore, recent studies provide

evidence of pyrophilous bacteria, such as the Proteobacteria *Massilia* (Enright et al., 2021;

Whitman et al., 2019), suggesting that fire resilience and adaptations might be widely distributed

within soil microbiomes. In fact, pyrophilous fungi have a suite of traits that allow for the

survival to fire (i.e., heat resistant spores, sclerotia) and the post-fire environment (i.e.,

xerotolerance, aromatic C degradation) (Bruns et al., 2020; Whitman et al., 2019). However,

although we know that pyrophilous microbes exist, our limited understanding of their

distribution in systems outside of pine forest (Pressler et al., 2019), their successional patterns

and turnover rates, or the replacement of existing taxa with new species, hinders our ability to

explore how pyrophilous microbes mediate the successional trajectory of vegetation in fire-

disturbed ecosystems.

Secondary succession begins in a nutrient-rich environment, with low competition for

nutrients and moisture (Dalling, 2019). Thus, trade-offs in colonization and resource utilization,

a hallmark of successional theory (Tilman 1990), allow surviving or fast-growing organisms to

control the initial successional stages (Turner et al., 1998). Over time, species with similar

resource use (i.e., overlapping niches) as initial colonizers will be excluded from the community

(Pacala et al., 1996), while differentiation in resource use (i.e., complementarity) will promote

species coexistence. Whether these patterns translate to belowground microbes remains unclear.

Additionally, soil microbial communities have short generation times (Hannula et al., 2019), and

on average, the first sampling of post-fire microbiomes occur two years post-fire (Dove & Hart,

2017; Pressler et al., 2019), thereby missing the fine-scale temporal succession that can occur

directly after a fire. To date, a couple of studies have repeatedly sampled the same post-fire

environment and found rapid changes in microbial composition (Ferrenberg et al., 2013; Qin &

Liu, 2021), suggesting that microbial communities likely experience high turnover rates.

Therefore, to capture the dynamics needed to develop microbial successional theory, it is integral

to sample at a scale relevant to soil microbial communities.

Chaparral is a shrubland adapted to high-severity fire and a biodiversity hotspot

distributed in Mediterranean climates worldwide (Barro & Conard, 1991; Rundel, 2018). The

secondary succession of chaparral vegetation is well understood (Keeley et al., 2005). Fire is

considered the main initiator of secondary succession, thus making chaparral ideal for studying

microbial secondary succession. Yet, although the dominant chaparral vegetation forms

mycorrhizal associations required for establishment and survival (Allen et al., 2005; Van der

Heijden et al., 1998), only13% of post-fire microbial research occurs in shrublands (Pressler et

al., 2019), and none discuss secondary succession. Moreover, ectomycorrhizal fungi experience

high mortality rates during wildfires (Dove & Hart, 2017; Pulido-Chavez et al., 2021), but the

ability of mycorrhizal fungi to survive a fire or the length they can survive without a host is

unknown. With drylands covering nearly 41% of the Earth's terrestrial surface and expanding

with climate change (Feng & Fu, 2013), it is critical to understand post-fire successional

dynamics in dryland systems such as chaparral.

Here, we studied the 2018 Holy Fire, which burned 94 km^2 of Southern California

chaparral, to determine the effects of wildfire on soil microbiomes, their successional

trajectories, and turnover rates. We sampled soils at 9-time points during the first post-fire year,

resulting in the highest resolution temporal sampling of post-fire microbes to date, which enabled

us to measure both immediate and temporal effects of wildfire on microbial dynamics across the

first post-fire year. We then tested the following hypotheses: (**H1**) wildfire will decrease

bacterial and fungal biomass and richness immediately following the fire, leading to a shift in the

community composition; (**H2**) bacterial and fungal biomass and richness will increase with time

since fire, but one-year will not be sufficient time for either community to return to unburned

levels; (**H3**) wildfire will have a distinct impact on fungal guilds with symbiotic mycorrhizal

fungi experiencing the largest declines; (**H4**) soil burn severity and precipitation will both affect

microbial succession, but post-fire precipitation will impose a larger impact on bacterial

communities (**H5**) bacterial, and fungal succession will be initiated by pyrophilous microbes,

which will trade-off in abundance over time.

2. Methods

2.1 Study Area, plot design, and soil collection

The Holy Fire burned 94 km^2 in the Cleveland National Forest across Orange and

Riverside counties in Southern California from August 6 to September 13, 2018. On September

30, 2018, within weeks of the fire being declared contained, we established nine plots (6 burned

and 3 unburned) along a gradient of fire severities (**Fig. 1A**). Plots were selected for similarity in

aspect, slope, elevation, and pre-fire vegetation dominance by manzanita *(Arctostaphylos*

glandulosa), an ectomycorrhizal host, and chamise (*Adenostoma fasciculatum),* an arbuscular

and ectomycorrhizal host (Allen et al., 2005). Plots were placed on average 25 m from forest

access roads to avoid edge effects. Each circular plot contained four 1 m^2 subplots located at 5 m

from the center in each cardinal direction (**Fig. 1B**). Our study site experiences a Mediterranean-

type climate with warm, dry summers and cool, wet winters, with an average yearly temperature

of 17˚C and precipitation of 12.5 mm. Precipitation data, based on monthly summaries, was

gathered from the El Cariso weather station accessed from the RAWS USA Climate Archives

(raws.dri.edu). Additional plot information can be found in the supplements **(Table S1)**.

We sampled soils at 17, 25, 34, 67, 95, 131, 187, 286, and 376 days, corresponding to

approximately 2 and 3 weeks, and 1, 2, 3, 4, 6, 9, and 12 months post-fire. We used BAER soil

burn severity maps (Parson et al., 2010) to roughly locate plots at medium and high severity

(**Fig. 1A**), but their coarse-scale measurement (30m) does not coincide with microbial

communities. Since ash depth increases with soil heating (Parson et al., 2010) and fire-severity

(Bodí et al., 2014), we used ash depth as a proxy for burn severity, where we averaged three

separate measurements of ash depth (cm) per each 1m^2 subplot at day 17, before any

precipitation event could wash away the ash. Thus, our measure of soil burn severity refers to

initial ash depth. At each of the 9 time-points, we collected the top 10cm of soil beneath the ash

or the duff from each of the 36 subplots with a ~250 mL releasable bulb planter cleaned between

samples with 70% ethanol (**Fig. 1C**). High-severity wildfires consume the organic layer; thus, for

consistency, we removed the litter layer in the unburned plots before sampling. Soils were

transported to the University of California-Riverside (UCR) within hours of sampling, stored

overnight at 4˚C, sieved (2 mm) within 24 hours, and a subsample was frozen at -80°C for DNA

extraction.

2.3 DNA extraction, amplification, and sequencing

DNA was extracted from soil using Qiagen DNeasy PowerSoil Kits following the

manufacturer's protocol, except the centrifugation time was increased to 1.5 min after adding the

C3 solution due to a large amount of precipitate. DNA extracts were stored at -20˚C for

subsequent analysis. Extracted DNA was amplified using the primer pair ITS4-fun and 5.8s

(Taylor et al., 2016) to amplify the ITS2 region for fungi and the primer pair 515F-806R to

amplify the V4 region of the 16S rRNA gene for archaea and bacteria (Caporaso et al., 2011)

using the Dual-Index Sequencing Strategy (DIP) (Kozich et al., 2013). Although our 16S primers

amplify archaea and bacteria, for simplicity, we refer to 16S methods and results simply as

bacteria since archaea only contributed <1% of sequencing reads. We conducted polymerase

chain reaction (PCR) in two steps. The first PCR amplified gene-specific primers, and the second

PCR ligated the DIP barcodes for Illumina sequencing. For bacteria, we combined 1 µl of 1:10

diluted DNA, 10.5 µL of Ultra-Pure Sterile Molecular Biology Grade (Genesee Scientific, San

Diego, CA, USA), 12.5 µL of AccuStart ToughMix (2x concentration; Quantabio, Beverly, MA,

USA), and 0.5 µL each of the 10 µM 515F and 806R primers. Thermocycler conditions for

PCR1 were: 94°C for 2 min followed by 29 cycles of 94°C for 30 s, 55°C for 30 s, 68°C for 1

min. For fungi, we combined 5 µl of undiluted fungal DNA, 6.5 µl of ultra-pure water, 12.5 µl

of AccuStart ToughMix, and 0.5 µL each of the 10 µM ITS4-fun and 5.8s primers. Thermocycler

conditions for PCR1 were: 94 °C for 2 min, followed by 30 cycles of 94 °C for 30 s, 55 °C for 30

s, 68 °C for 2 min. PCR1 products were then cleaned with AMPure XP magnetic beads

(Beckman Coulter Inc., Brea, CA, USA) following manufacturers' protocols. The DIP PCR2

primers containing the barcodes and adaptors for Illumina sequencing were ligated to the

amplicons during the second PCR step in a 25 µL reaction containing 2.5 µL of the 10 µM DIP

PCR2 primers, 6.5 µL of ultra-pure water, 12.5 µL of Accustart ToughMix, and 1 µL of PCR1

product. Thermocycler conditions for the second PCR for bacteria and fungi were: 94°C for 2

min followed by 9 cycles of 94°C for 30 s, 60°C for 30 s, and 72°C for 1 min. Bacterial and

fungal PCR products for each of the 324 samples (9 plots x 4 sub-plots x 9 time-points) were

then separately pooled based on gel electrophoresis band strength and cleaned with AMPure

following established methods (Glassman et al., 2018). Each pool was then checked for quality

and quantity with the Agilent Bioanalyzer 2100, pooled at a 2:3 ratio (0.4 units for bacteria to 0.6

units for fungi), and sequenced with Illumina MiSeq 2x300bp at the UCR Institute for

Integrative Genome Biology. In addition to our 324 samples sequenced in four Illumina MiSeq

runs, we added negative DNA extractions and negative PCR controls from each time point and

mock communities (ZymoBIOMICS microbial community standard, Zymo, Irvine, CA) to each

library for additional quality control and inference.

2.4 Bacterial and fungal biomass

Bacterial and fungal biomass was quantified with quantitative (q) PCR. We used the

FungiQuant-F and FungiQuant-R primers to target the 18s rDNA for fungi (Liu et al., 2012) and

the Eub338/Eub518 primers for bacteria (Fierer et al., 2005). To estimate bacterial and fungal

small-subunit rDNA gene abundances, standard curves were generated using a 10-fold serial

dilution of the standards that were generated by cloning the 18S region of *Saccharomyces*

cerevisiae or the 16S region of *Escherichia coli* into puc57 plasmid vectors, which were

constructed by GENEWIZ, Inc. (NJ, USA) as previously established (Averill & Hawkes, 2016).

The 10 µl qPCR reactions were performed in triplicate. Reactions contained 1 µL undiluted

DNA, 1 µl of 0.05M Tris-HCl ph8.3, 1 µL of 2.5mM $MgCl_2$ (New England BioLabs (NEB);

Ipswich, MA), 0.5 µL of 0.5mg/ml BSA, 0.5 µL of 0.25mM dNTP (NEB), 0.4 µL of both

primers at 0.4µM, 0.5 µL of 20X Evagreen Dye (VWR), 0.1 µL of Taq DNA polymerase (NEB)

and 4.6 µL ultra-pure water. We employed the CFX384 Touch Real-Time PCR Detection

System with the following conditions: 94˚C for 5 min, followed by 40 cycles of 94˚C for 20

seconds, 52˚C (for bacteria) or 50˚C (for fungi) for 30 seconds, followed by an extension at 72˚C

for 30 seconds. Gene copy numbers were generated using a regression equation, $10^{\wedge}$ (Cq-b)/m,

where the y-intercept and slope were generated from the standard curve. The Cq value was

calculated as the average Cq value per sample in relation to the known/calculated copies in the

standard Cq (CT); values were generated by CFX Maestro software.

2.5 Bioinformatics

Illumina data was processed with Qiime2 (Bolyen et al., 2019). The demultiplexed fastQ

files from the four Illumina sequencing runs were processed individually using cutadapt (Martin,

2011) to remove the primers and ran through DADA2 to filter out and remove chimeric

sequences and low-quality regions and to produce Amplicon Sequence Variants (ASVs)

(Callahan et al., 2017). DADA2 outputs from each library were merged into one library for

downstream processing, including the removal of singletons and taxonomic assignments. We

used the SILVA reference database for bacteria (Yilmaz et al., 2014) and the UNITE reference

database for fungi (Kõljalg et al., 2005) for taxonomic assignment (accessed on 10/27/2020).

Sequences assigned to mitochondria and chloroplasts for bacteria or not assigned to the Kingdom

Fungi were removed from ASV tables before subsequent analysis. Raw fungal ASV tables were

exported and parsed through FUNGuild (Nguyen et al., 2016) to assign functional ecological

guilds to each fungal ASV. We identified the following fungal guilds and included only high

probable confidence rankings for downstream analysis: arbuscular fungi (AMF), ectomycorrhizal

fungi (EMF), saprotrophs, and pathogens. Moreover, negative and mock controls were inspected

to ensure that sequences in negative controls were negligible and that mock taxonomy correlated

with known communities. Sequences were submitted to the National Center for Biotechnology

Information Sequence Read Archive under BioProject accession number PRJNA761539.

2.6 Statistical analysis

Statistical analyses were conducted in R, 4.1.1 (R Core Team, 2021), and all statistical

codes are available on GitHub https://github.com/pulidofabs/SecondarySuccession-Chaparral.

To account for uneven sequencing depth, ASV tables were rarefied to a

sequences/sample depth of 7,115 for bacteria and 11,480 for fungi. Fungal and bacterial alpha

diversity was estimated using BiodiversityR (Kindt & Coe, 2005) with the following metrics:

observed species richness, Simpson, Shannon, Chao1, and ACE. Since patterns of species

richness were similar across metrics **(Fig. S1)**, we focused on richness estimated as the number

of observed ASVs after rarefaction for all downstream analyses. To test if wildfire decreased

biomass and richness (**H1**), if biomass and richness increased with time since fire **(H2),** if

wildfire will have distinct impact on fungal guilds **(H3)**, and to determine if changes in biomass

and richness were associated with precipitation and soil burn severity **(H4),** for both bacteria and

fungi, we fitted thirteen statistical models with treatment (burned vs. unburned), time (measured

as days since fire), ash depth (cm), total monthly precipitation (mm), and their respective

interactions as predictors. We used generalized mixed effect models (glmer) with a negative

binomial distribution in the MASS package (Venables & Ripley, 2002) to account for the over-

dispersion of the data and the fact that the conditional variance was higher than the conditional

mean (Bliss, 1953; Ross & Preece, 1985). Moreover, time since fire and precipitation were

scaled and centered, and the level of nestedness for all models was tested by running a null

model with different nested levels ("plot," "subplot," "time since fire") and no predictors. Model

selections were made via comparison of Akaike Information Criterion (AICc) in the MuMIN

package (Barton, 2020). All richness models contained "plot," "subplot," and "time since fire" as

random effects, and all biomass models contained "plot" and "time since fire" as random effects.

Pseudo $R^{2,}$ or the variance explained (marginal and conditional) for all models, was calculated

using the r.squaredGLMM function in the MuMIn Package.

We compared beta diversity by generating distance matrices with the Vegan Avgdist

function, calculating the square-root transformed median Bray-Curtis dissimilarity per 100

subsampled iterations. We used permutational multivariate analysis of variance (PERMANOVA)

(Anderson, 2017) as implemented with the Adonis function in the vegan package (Oksanen et

al., 2018) to test for the significant effects of wildfire, time since fire, and their interactions, on

bacterial and fungal community composition overall and at each independent time point (**H1**).

Results were visualized using Non-Metric Multidimensional Scaling (NMDS) ordinations.

We visualized succession by characterizing the community composition patterns at the

genus level using phyloseq (McMurdie & Holmes, 2013) and plotting the relative abundance

profile of the dominant ASVs (greater than 3% sequence abundance) for each time point in

burned and unburned communities independently with ggplot2. We tested for significant changes

in abundance of each taxa between burned and unburned plots and over time using glmer.

Moreover, we employed several methods to quantify succession (**H5**). Similar to a study

examining microbial turnover in Sorghum plants (Gao et al., 2020), we used the mantel function

in vegan to determine the correlation between temporal distance and community composition to

determine how turnover varies in the presence or absence of fire. To visualize spatial species

turnover (Baselga, 2010), we took advantage of the fact that early successional periods display

large variability (Collins, 1990; Pandolfi, 2008). Thus, we tested the homogeneity of the

bacterial and fungal communities during succession using the vegan betadisper function to

calculate the multivariate dispersion of Bray-Curtis dissimilarities and visualized the results

using PCoA. Finally, we employed the codyn package (Hallett et al., 2016) to identify various

factors of diversity metrics to assess succession (Collins et al., 2000). First, to identify whether

the community is experiencing its normal successional trajectory, per the ecological successional

theory, we calculated direct linear change, also known as directional change, using the Euclidian

distance of each microbial community. Moreover, we measured total species turnover or the rate

of species change between two-time points to determine if species appearances or disappearances

drive succession. Lastly, we measured community stability as a measure of resilience, or rate of

return to the original state, which is product synchrony or patterns of how species abundance

covary over time (Loreau & de Mazancourt, 2008)

3. Results

3.1 Sequencing data

The four Illumina MiSeq runs resulted in 9.8 M bacterial and 24.6 M fungal sequences

for an average of 31,052 bacterial and 78,202 fungal sequences/sample and a total of 33,078

bacterial and 11,480 fungal ASVs. Rarefaction resulted in a total of 24,874 bacterial and 7,445

fungal ASVs. Extracting the fungal guilds from the rarefied fungal ASV table resulted in 208

EMF, 70 saprobic, 65 AMF, and 26 pathogenic fungal ASVs.

3.2 Wildfire effects on bacterial and fungal biomass and richness

Fire led to large and significant reductions in bacterial and fungal biomass and richness

during the first post-fire year (**Tables S2; S3)**. Fire had larger impacts on biomass than richness

for both bacteria and fungi and larger impacts on fungi than bacteria overall (**Fig. 2**). On average

across the year, fire reduced richness by 46% for bacteria (**Fig. S1A**) and 68% for fungi (**Fig.**

S1A) and biomass by 47% for bacteria **(Fig. S2A)**, and 86% for fungi **(Fig. S2B)**. The direct,

immediate effect of wildfire at 17 days post-fire was stronger for biomass than richness, with

bacterial biomass decreasing by 84% **(Fig. 2A)** and fungal biomass by 97% **(Fig. 2B).** In

contrast, fungal richness declined by 45% **(Fig. 2D)**, but bacterial richness temporarily increased

by 31% but declined by 29% at 25 days post-fire **(Fig. 2C; Table S2)**. While the differences in

biomass and richness between the burned and unburned sites lessened with time since fire, one

year was insufficient for bacterial and fungal biomass or richness to recover to unburned levels

(Fig. 2). The effects of fire across time remained larger for fungi than for bacteria, with fungal

biomass remaining 80% and richness 61% lower and bacterial biomass remaining 43% and

richness 23% lower in the burned plots at 1-year post-fire **(Fig. 2, Table S2).**

3.3 Drivers of changes in richness and biomass

Negative binomial models showed that the direct impact of fire negatively affected

bacterial and fungal biomass and richness **(Table S3)**, which was lessened by the positive

interaction effect of time since fire and soil burn severity **(Fig. 3; Table S3).** While time since

fire only directly affected fungal richness (**Table S3)**. Soil burn severity had significant adverse

direct effects on bacterial and fungal richness but no significant effects on biomass, meaning that

biomass was equally reduced in the presence of fire regardless of severity **(Fig. 3; Table S3)**.

Precipitation did not directly affect bacterial or fungal biomass or richness, but it interacted with

time since fire affected bacterial negatively **(Fig. 2C)** and positively affected fungal richness

(Fig. 2D; Table S3). However, bacterial biomass responded positively to a fire and precipitation

interaction (**Fig. 2A; Table S3).**

3.4 Wildfire effects on the richness of the different fungal guilds

Wildfire led to large and significant reductions for all fungal guilds **(Table S4, Fig. S3),**

with the largest declines for AMF. On average across time points, EMF decreased by 68%,

pathogens by 71%, saprobes by 86%, and AMF by 98% in burned plots compared to unburned

plots **(Fig. S3).** Moreover, one year was insufficient for any guild to return to unburned levels. In

fact, at 376 days, AMF was on average 89%, pathogens 69%, saprobes 61% lower in the burned

plots (Fig. S4). Interestingly, EMF experienced a decrease in richness over time, resulting in

species richness being 91% lower in the burned plots at 376 days post-fire. **(Fig. S4)**. Although

EMF richness did not significantly differ from the unburned plots at 25 days, fire interacted with

time since fire caused EMF to decrease in the burned plots **(Fig. S4; Table S4)**. EMF richness

further declined with soil burn severity **(Table S4).** Moreover, a fire and precipitation interaction

negatively affected EMF and pathogen richness **(Table S4).** In the unburned plots, mycorrhizal

and pathogen richness remained unchanged over time **(Fig. S4)**, but an interaction between fire

and time resulted in a sharp dip in saprobic richness in the unburned plots at 286 days post-fire,

causing a brief overlap between the burned and unburned plots **(Fig. S4, Table S4).**

3.5 Wildfire changes microbial community composition

Fire significantly affected the community structure of bacteria and fungi, explaining 13%

of the compositional variation for bacteria and 11% for fungi **(Table S5).** Time since fire had

smaller impacts on community composition, explaining 4% of the variation for bacteria and 1%

for fungi. In comparison, a fire and time interaction explained 2% of the variation for bacteria

and 1% for fungi **(Table S5).** Community composition between burned and unburned plots

significantly varied at all 9-time points for both bacteria **(Fig. S5)** and fungi **(Fig. S6),** with

differences increasing over time from 12% to 21% for bacteria **(Fig. S5)** and 9% to 13% for

fungi from day 17 to day 376 post-fire **(Fig. S6)**. The changes in bacterial **(Fig. 4B)** and fungal

composition **(Fig. 4D)** were driven by the dominance of previously rare taxa, which traded off in

abundance over time in the burned plots. Bacterial communities experienced less discernable

trade-offs than fungi due to the consistent dominance of the Proteobacteria *Massilia*, which

composed on average 57% of the sequences throughout the year in the burned plots **(Fig. 4B)**.

However, three Firmicutes quickly dominated, with *Bacillus* (13%), an uncultured Clostridiales

(26%), and *Paenibacillus* (4%) dominating at 34 and 67 days post-fire. The Firmicutes rapidly

declined and traded abundances with the Proteobacteria *Noviherbaspirillum,* which increased in

abundance over time, from 1% at 34 days to 24% by the end of the year **(Fig. 4B).** For fungi, two

Basidiomycetes that dominated at 17 days post-fire, the yeast *Geminibasidium* (45%) and the

EMF *Inocybe* (16%), rapidly declined and traded abundance with filamentous Ascomycetes **(Fig.**

4D). While the Ascomycete *Penicillium* quickly established dominance and constantly occupied

an average sequence abundance of 38% over the year **(Figs. 4D)**, *Pyronema* rapidly dominated,

peaking at 67 days with 67% of the total sequences post-fire. *Pyronema* slowly declined

thereafter as *Aspergillus* slowly increased in abundance over time, from 2% at17 days to 22%

376 days post-fire **(Fig. 4D)**.

3.6 Microbial successional dynamics

Directional change in the community was higher in the burned bacterial and fungal

communities, confirming that succession is occurring **(Table S6).** Moreover, the low synchrony

observed in bacterial and fungal burned communities (Table S6) indicates community stability

due to the dominance of a few genera. Analysis of the community stability collaborated these

results and was higher for bacteria than fungi, and it was higher in the burned than unburned

bacterial communities **(Tables S6).** Indicating that bacterial succession is rapidly proceeding to

the point of potential equilibrium due to increasing bacterial richness and resilience. In contrast,

fungal communities are less stable more susceptible to change. Visual confirmation of these

trends established that both bacterial **(Fig. 5A)** and fungal **(Fig. 5B)** communities experienced

rapid and distinct successional trajectories in the burned plots, with bacteria **(Fig. 5C)**

experiencing successional changes at double the rate of fungi **(Fig. 5D)**. These successional

changes in the burned plots were driven by six major compositional turnover points for bacteria

at 25, 34, 95, 131, 187, and 376 days post-fire **(Fig. 5E)** and five for fungi at 25, 34, 67, 95, and

131 days post-fire **(Fig. 5F).** In contrast, the unburned bacterial **(Fig. 6A)** and fungal **(Fig. 6B)**

communities experienced lower dominance and remained stable over time with no succession

(Figs. 6C, 6D) or compositional turnover (**Figs. 6E, 6F).**

3.7 Taxa driving microbial succession

Early turnover events for bacteria **(Fig. 5A)** and fungi **(Fig. 5B)** were driven by

the disappearance of taxa, more so for bacteria (days 17-95) than fungi (17-25; **Table S6**).

Whereas high diversity and the lack of dominance drove early bacterial succession **(Fig. 5A)**, as

succession proceeded, dominance by a few taxa, including the Firmicutes *Bacillus* at 25 days

post-fire, *Paenibacillus* and *Domibacillus* (34-67 days), and the Proteobacteria

Noviherbaspirillum (95 days), and the constant dominance of *Massilia* (34-286 days post-fire)

resulted in the disappearance of bacterial species over time **(Fig. 5A; Table S6)**. In contrast,

fungal succession was initiated by the rapid dominance of *Geminibasidium* and EMF genera that

dominated the unburned communities, including *Inocybe, Cortinarius*, and *Tomentella* in the

Basidiomycota and *Balsamia* in the Ascomycota and one previously rare Basidiomycete EMF

genus *Mallocybe,* a subgenus of *Inocybe* **(Fig. 5B, 6B).** Interestingly, all species of EMF rapidly

disappeared from the community at 25 days post-fire and were replaced by the constant

dominance of *Pyronema*, *Penicillium,* and *Aspergillus* **(Fig. 5B)**. However, as succession ensued,

species appearance drove late-stage bacterial (187-376 days; **Fig. 5A**) and mid-stage fungal (67-

187 days) succession **(Fig. 5B; Table S6).** For bacteria, the appearance of two Bacteroidetes,

Pedobacter and *Adhaeribacter*, and the Actinobacteria *Blastococcus* shaped mid-stage bacterial

succession (**Fig. 5A**). For fungi, mid-stage succession was driven by the appearance of

Basidiomycota mushroom-forming taxa, including the return of the previously dominant EMF

genus *Inocybe*, and the appearance of the saprobes, *Coprinellus,* and *Tephrocybe,* which were

rare in the unburned communities (<0.01% sequence abundance). The appearance of unknown

filamentous Ascomycota in the Sordariaceae and Pyronemataceae also drove mid-stage fungal

succession. In contrast, late-stage succession was driven by the increased dominance of

Tephrocybe and *Aspergillus* and the disappearance of *Inocybe* and *Rasamsonia,* and the

decreased abundance of *Pyronema* from 286-376 days post-fire **(Fig. 5B).**

4. Discussion

Here, we present the highest resolution temporal sampling of post-fire microbiomes to

date, enabling us to show for the first time that bacteria and fungi experience rapid succession

driven by the dominance of previously rare "fire-loving" pyrophilous taxa that increase post-fire

but experience marked trade-offs in abundance over time. Although our study was in California

chaparral, many of the dominant pyrophilous taxa identified here also dominate other burned

environments, including pine (Bruns et al., 2020; Reazin et al., 2016), spruce forests (Whitman et

al., 2019), redwood tanoak forests (Enright et al., 2021), and Spanish shrublands (Pérez-Valera et

al., 2018), indicating the generality of our results to wildfire-affected ecosystems. We found that

wildfire decreased bacterial and fungal biomass and richness, shifting community composition

(H1) and persisting for the entire year for both bacteria and fungi (**H2**). Fungal guilds were

differentially affected by fire and time, with AMF and saprobes experiencing larger direct fire

impacts than pathogens or EMF. However, over time, mycorrhizal fungi experience larger

declines than pathogens and saprobes (**H3).** Interestingly, AMF immediately succumbed to the

death of their host while EMF survived for 1-2 months, answering the question of how long it

takes mycorrhizal fungi to die after the death of their host. Moreover, soil severity negatively

affected bacterial and fungal richness, while bacterial biomass and fungal richness were

positively influenced by precipitation's interaction with time and fire (**H4).** Finally, we found

strong evidence that fire causes bacteria and fungi to undergo rapid secondary succession

dominated by pyrophilous microbes that trade-off in abundance over time (**H5),** likely driving

post-fire ecosystem recovery.

4.1 Wildfire decreased bacterial and fungal biomass and richness

Soil bacterial and fungal biomass and richness declined in the burned sites, corroborating

previous post-fire studies in Mediterranean shrublands (Pérez-Valera et al., 2018) and forests

(Dooley & Treseder, 2012). We noted a more significant fire effect on fungal biomass and

richness relative to bacteria, consistent with previous research (Certini et al., 2021; Glassman et

al., 2021; Pourreza et al., 2014; Pressler et al., 2019), showing that bacteria are more resistant to

fire than fungi (Certini et al., 2021; Dunn et al., 1982). Although richness and biomass increased

over time for both microbial groups, one year was insufficient for either group to recover to

unburned levels, consistent with previous studies (Pérez-Valera et al., 2018; Treseder et al.,

2004), indicating that microbial biomass and richness could take decades to recover. Moreover,

we observed a transient increase in bacterial richness directly post-fire. This transient increase

may result from taxa such as bacteria in the Actinobacteria genera *Soliribrobacter* and

Conexibacter (Albuquerque & da Costa, 2014) that may be favored by the increase in soil pH

observed here **(Table S1)** and in other studies (Neary et al., 1999; Rousk et al., 2010) or by post-

fire pulses of nitrogen and phosphorus (Certini et al., 2021). We observed a larger direct fire

impact on AMF and pathogens than EMF and saprobes; however, EMF was more impacted over

time. Although studies have noted the decrease in EMF and saprobic richness (Dove & Hart,

2017; Enright et al., 2021; Pulido-Chavez et al., 2021) and AMF richness or colonization (Dove

& Hart, 2017; Enright et al., 2021), generalization of our observations are difficult as the vast

majority of studies focus on 1-2 guilds (Dove & Hart, 2017). Although EMF did not experience

the largest immediate effect of wildfire compared to the other guilds, we speculate that EMF

persisted for a few weeks post-fire because their host roots remained alive. Thus, the increase in

EMF mortality with time is dependent on host mortality, providing evidence for the importance

of surviving host for EMF survival.

4.2 Precipitation and soil burn severity affected soil bacterial and fungal communities

Due to our high-resolution temporal sampling and fine-scale measurement of soil burn

severity, we detected the effect of precipitation and fire severity on microbial recovery.

Precipitation did not affect fungal biomass, interacted with time to negatively affect bacterial but

positively affected fungal richness, and bacterial biomass in the burned plots. These results are

consistent with previous studies (Barnard et al., 2013), which showed that bacterial biomass can

bloom in response to rewetting. This distinct response in richness could be due to mortality and

survival characteristics and physiological and morphological differences between these microbial

groups (Blazewicz et al., 2014; Placella et al., 2012; Selbmann et al., 2013) and their potential

adaptation to normal water conditions in chaparral. For example, fungi can prime themselves for

stressful situations by activating or enhancing resistance to drought (Evans & Wallenstein, 2012;

Guhr & Kircher, 2020), allowing them to maintain a constant reproduction rate regardless of the

external environmental conditions. Moreover, the high morphological plasticity of fungi,

including their ability to alternate growth forms based on environmental conditions, such as

hyphal growth in low nutrient conditions or meristematic growth in stressful conditions

(Selbmann et al., 2013), may allow fungi to survive small changes in moisture content, while

forcing certain bacterial groups to enter a dormant state, inhibiting growth (Schimel, 2018).

Thus, to be evolutionarily successful, bacteria may respond rapidly to favorable conditions, e.g.,

rain, and rapidly replicate to outcompete other microbes for the pulse of nutrients released during

the rain event (Homyak et al., 2014), thus potentially explaining the peaks in biomass and

richness observed after high rain events (day 131).

499   Our results show that initial ash depth, measured before the first rains, can serve as an

index of soil burn severity measured on a scale relevant to the soil microbiome, which can aid in

understanding the initial and long-lasting impacts of fire severity on soil microbiomes. Similar to

studies in boreal (Whitman et al., 2019) and oak forests (Pourreza et al., 2014), bacterial and

fungal richness strongly responded to soil burn severity. However, contrary to a meta-analysis

(Dooley & Treseder, 2012), neither bacterial nor fungal biomass was directly impacted by soil

burn severity. Due to the frequent high-severity fires in this system, we speculate that biomass

might depend on a temperature threshold (fire intensity) rather than fire severity. However, since

we observed an interaction effect of time since fire and severity on biomass, we speculate that

previously observed severity effects, which results from studies that on average sample two years

post-fire (Dove & Hart, 2017; Pressler et al., 2019), may represent an interaction effect of time

and severity and not a direct effect of severity.

.

4.3 Pyrophilous bacteria dominate burned communities and drive succession

513   Wildfire led to a complete turnover of the chaparral soil bacterial communities favoring

putative pyrophilous bacteria that have been detected after fires in temperate and boreal forests,

such as taxa belonging to the phyla Actinobacteria, Acidobacteria, Firmicutes, and

Proteobacteria (Enright et al., 2021; Whitman et al., 2019; Xiang et al., 2014). In particular, the

dominance of the genera *Massilia, Noviherbaspirillium,* and *Paenibacillus* may be attributed to

their ability to survive the wildfire and exploit transient niches in the early post-fire environment.

For example, *Paenibacillus xerothermodurans* is an endospore-forming bacterium that has been

shown to contain heat-inducible transcriptional proteins, which can protect heat-induced protein

denaturation (Kaur et al., 2018). However, *Massilia* does not form endospores but was

nonetheless the most dominant genus post-fire. *Massilia* is a known fast reproducing (Enright et

al., 2021; Whitman et al., 2019) rhizospheric bacteria (Li et al., 2014), capable of colonizing the

roots of various plants (Ofek et al., 2012) and forming associations with arbuscular fungi (Iffis et

al., 2014). Thus, we speculate that a fast reproduction rate and arbuscular association allow

Massilia to promote its ability to dominate post-fire systems by increasing its proliferation within

the soil (Toljander et al., 2005). Lastly, an uncultured Firmicute Clostridiales dominated early

succession. Previous studies suggest that Clostridiales produce resistant spores that can survive

harsh conditions and lack of nutrients by triggering sporulation (Paredes-Sabja et al., 2011). This

suggests that Clostridiales survived the wildfire and quickly sporulated, resulting in early post-

fire dominance. Together these results suggest that post-fire dominant bacteria in wildfire-

adapted systems have coevolved and developed mechanisms to survive the fire resulting in niche

exclusion and dominance in recently burned ecosystems.

Our high temporal resolution sampling enabled us to demonstrate the secondary

succession of bacterial communities and contextualize these dynamics in broader successional

theory. Moreover, we were able to show that putative pyrophilous bacteria experience rapid

turnover, resulting in a marked trade-off in abundance over time, information currently lacking

in soil bacterial communities and post-fire disturbed ecosystems. In bacterial communities,

succession and community turnover occurred rapidly and was initiated by aerobic, heterotrophic

bacteria that form endospores and produce antibiotics, which may improve their competitive

abilities.

Like plant successional patterns (Capitanio & Carcaillet, 2008; Egler, 1954), early

successional stages were characterized by high bacterial diversity and a lack of dominance,

mimicking the distributions observed in unburned communities. Previous studies suggest that in

complex resource environments, such as those created by wildfire, niche complementarity

becomes more important than competition because diverse communities can better exploit

resources (Eisenhauer et al., 2013). Our results support this idea, as the dominant taxa in our

study have been shown to use different forms and types of resources. For example,

*Paenibacillu*s can fix nitrogen (Monciardini et al., 2003; Slepecky & Hemphill, 2006), *Bacillus*

and *Conexibacter* can reduce nitrogen, and *Bacillus* can further solubilize phosphate (J. et al.,

2001; Kalayu, 2019), suggesting that niche partitioning may promote positive interactions

between bacteria in the early successional stages (<25 days). However, as succession progressed,

the bacterial community converged on one dominant functional strategy, where all three genera

that dominated most successional stages (*Massilia, Paenibacillus,* and *Noviherbaspirillum)* have

similar functional roles, including phosphorus solubilization (*Paenibacillus, Massilia*), nitrogen

fixation (*Paenibacillus*), and reduction (*Massilia, Paenibacillus*) (Bailey et al., 2014; Baldani et

al., 2014; Grady et al., 2016; Gu et al., 2016; Wan et al., 2020; Woolet & Whitman, 2020).

Moreover, *Massilia* and *Noviherbaspirillium* are putative polycyclic aromatic hydrocarbon

degraders, a component of pyrogenic organic matter, an abundant post-fire carbon resource

(Baldani et al., 2014; Grady et al., 2016), which may enable them to thrive in the post-fire

environment; suggesting that the diverse functional potential of the dominant taxa may allow

these bacteria to survive in a changing environment.

4.3 Pyrophilous fungi dominate the burned communities and drive succession

Interestingly, previously detected pyrophilous fungi from globally burned ecosystems

also dominate California chaparral. Although our experiment did not study mechanisms

controlling pyrophilous microbial dominance, it is clear that pyrophilous microbes are not

biome-specific. Instead, these fungi may be activated by temperature thresholds, such as

temperatures reached during high severity fire, which are common in chaparral and some

Pinaceae forests (Agee, 1993; Keeley & Zedler, 2009; Neary et al., 1999). Specifically, our

communities were dominated by Ascomycota in the genera *Pyronema, Penicillium*, and

Aspergillus, similar to dominant fungal species in wildfire affected boreal spruce forest

(Whitman et al., 2019), montane pine forests (Fujimura et al., 2005; Pulido-Chavez et al., 2021),

Mediterranean shrublands (Livne-Luzon et al., 2021) and controlled pyrocosm studies (Bruns et

al., 2020) Both *Pyronema (P. omphalodes)* and *Aspergillus* (*A. fumigatus*) are known

pyrophilous fungi of California chaparral (Dunn et al., 1982). We identified additional species

that responded positively to wildfire, including *P. domesticum*, *A. udagawae*, and *A.*

elsenburgensis. These pyrophilous fungi are adapted to wildfire (Moore, 1962; Rhodes, 2006)

and produce fire-resistant structures such as dormant spores, sclerotia, and conidia (Smith et al.,

2015; Warcup & Baker, 1963), which are heat activated (Gottlieb, 1950; Warcup & Baker,

1963). The rapid and efficient germination of *Aspergillus* (in the subgenus *Fumigati*, including

A. udagawae) induced by high temperatures (Rhodes, 2006) and its ability to use various carbon

and nitrogen sources, including ammonium and nitrate (Krappmann & Braus, 2005), may

position *Aspergillus* to rapidly dominate after chaparral fires. Furthermore, *Pyronema*

domesticum, a relatively well-studied pyrophilous fungus, can mineralize pyrogenic organic

matter, a dominant substrate in burned environments (Fischer et al., 2021), thus potentially

allowing it to take advantage of the abundant food source in this system. Finally,

Geminibasidum, a recently described thermotolerant Basidiomycete yeast (H. D. T. Nguyen et

al., 2013), dominated 45% of our sequences at 17 days post-fire. While not typically described as

pyrophilous, presumably because most pyrophilous fungi are described from sporocarps

(McMullan-Fisher et al., 2011), recent studies using next-generation sequencing have found

Geminibasidium to increase post-fire in pine forests (Pulido-Chavez et al., 2021; Yang et al.,

2020). Thus, it is likely that *Geminibasidium* is an underrepresented pyrophilous fungus that

appears after high severity fires in multiple ecosystems. Together, these results suggest that

dominant pyrophilous fungi are more widespread than expected and that their response to fire

might be due to fire adaptations and requirements.

Although knowledge of pyrophilous fungi is growing, we lacked an understanding of

how soon these pyrophilous fungi appear, how they trade in abundance over time, and their

turnover rates (Fox et al., 2022). We show that pyrophilous fungi experience rapid turnover,

trading in abundance over time, and by establishing their dominance in early successional stages

(<25 days), they maintain control of the system for the entire year. Early fungal successional

stages (3 weeks post-fire) were dominated by surviving ectomycorrhizal Basidiomycetes, such as

Cortinarius, Inocybe, and *Tomentella,* which dominated the unburned fungal communities and

that have been previously found in burned pine forests (Gassibe et al., 2011; Glassman et al.,

2016; Owen et al., 2019; Pulido-Chavez et al., 2021) and pine-dominated Mediterranean systems

(Gassibe et al., 2011; Hernández-Rodríguez et al., 2013). However, fast-growing saprobic and

pyrophilous fungi in the genera *Pyronema, Penicillium*, and *Aspergillus* quickly dominated all

successional time points, beginning at 25 days. The results for *Pyronema* are consistent with

pyrocosm studies (Bruns et al., 2020). Although it is possible that the EMF signal detected could

be due to relic DNA (Carini et al., 2016), high UV conditions, such as those in post-fire systems

(Neary et al., 1999), rapidly degrade relic DNA (Torti et al., 2015), suggesting that these EMF

survived the fire and clung to a dying host for at least 2 months, as in the case of *Inocybe*. Thus,

we suggest that this is not relic DNA but a true signal of how long it takes EMF to die once their

host dies. Ectomycorrhizal survival may be attributable to their exploration type and specificity

for current vs. stored photosynthates (Gray & Kernaghan, 2020; Pena et al., 2010). For example,

long/medium exploration types, such as *Cortinarius,* require more carbon for maintenance and

only remain in the community for 17 days post-fire. Whereas *Inocybe*, a short-contact type

(Agerer, 2001; Koide et al., 2014), survived for up to 67 days post-fire. Thus, we speculate that

early EMF could have survived on their hosts' stored photosynthates. However, not all EMF

hosts died, as sprouting was observed as early as 286 days post-fire, suggesting that the EMF that

survived the fire could have survived on the lower resources provided by the resprouting

manzanita and chamise shrubs.

Successional theory states that early successional stages are dominated by fast-growing

species that quickly dominate the open space (Kinzig & Pacala, 2013). Our data suggest that this

trait in the ecological theory applies to fungal succession. The fungal community was dominated

by fast-growing, thermotolerant fungi, *Penicillium*, *Aspergillus,* and *Pyronema* (Dix & Webster,

1995; McGee et al., 2006). It is well known that both *Penicillium* and *Aspergillus* can readily

reproduce via asexual means, thus in their asexual states, they can readily reproduce without

needing a comparable mate (Crow, 1992), allowing rapid colonization of the open niche.

Although stress selection is suggested to dominate sexual reproduction (Grishkan et al., 2003),

our results suggest that in chaparral, a fire-adapted ecosystem, stress adaptations such as the

melanized conidia and thick cell walls of *Aspergillus* and *Penicillium* (Cordero & Casadevall,

2017; Lagashetti et al., 2019), and the formation of dormant spores or conidia might be

characteristic of the historical fire regime of the system, thus allowing these fungi to rapidly

dominate all successional time points during the first post-fire year. Lastly, we noted the

emergence of fungal decomposers, such as *Coprinellus* and *Tephrocybe,* at later successional

stages, supporting previous research (McMullan-Fisher et al., 2011). The onset of fruiting body

development of well-known decomposers, such as *Coprinellus*, suggest that the potential

mortality of early successional forbs and the increased dead woody material in the system might

be providing the organic C required for saprobic communities.

5 Conclusion

Due to our unprecedented high-resolution temporal sampling, we show that bacterial and

fungal communities experience rapid secondary succession and trade-offs in abundance. We

found that the same pyrophilous bacteria and fungi that respond to fires in temperate and boreal

forests and Mediterranean shrublands also dominate in California chaparral, allowing us to

generalize knowledge of post-fire microbiomes to dryland habitats that are rapidly increasing

with climate change (Feng & Fu, 2013). While pyrophilous microbes have been described in

other systems, their turnover rates were unknown due to the lack of repeated post-fire sampling.

We found that microbial communities were dominated by post-fire microbes that were rare or

absent pre-fire and which trade-off in abundance over time, in a manner that may affect post-fire

biogeochemical cycling and vegetation regeneration due to their abilities to transform post-fire

nutrients or associate with regenerating vegetation. Additionally, post-fire dominant bacteria and

fungi in wildfire-adapted systems appear to have many traits analogous to plants, such as fire

adaptations and the ability to grow fast and rapidly exploit post-fire niches, making it easier to

adapt successional theory developed in plants to microbes. We conclude that although wildfire

leads to drastic reductions in microbial species richness, biomass, and community composition,

post-fire bacteria and fungi experience rapid successional changes that align with plant

successional theory. Understanding these dynamics can increase our ability to predict if and how

ecosystems will recover after fire, a disturbance that is increasingly widespread with climate

change.

Acknowledgments

We thank the Cleveland National Forest and the Trabuco Ranger District, including District

Ranger Darrel Vance and Officers Jeffrey Heys, Emily Fudge, Victoria Stempniewicz, and Jacob

F. Rodriguez, for their guidance, help with the permitting and site selection process. We thank

Judy A. Chung for help with fieldwork and molecular work and Aral C. Greene, Dylan Enright,

and Sameer S. Saroa for fieldwork assistance. We thank Amelia Nelson, Elizah Stephens, and

Kaleigh A. Russell, and our anonymous reviewers for their helpful comments on the manuscript.

This work was supported by UC Riverside, the BLM JFSP Award #012641-002, and Shipley

Skinner Award to MFPC and SIG, and the USDA-NIFA Award 2022-67014-36675 to SIG and

PMH.

Conflict of interest

Authors do not declare any conflict of interest or relationship that might influence the author's

objectivity.

Author's contribution: SIG conceived of the study and acquired permits; LL, SIG, and PMH

came up with the experimental design; SIG, LL, PMH, MFPC, and JWJR performed fieldwork;

MFPC and JWJR performed all molecular work; MFPC performed all bioinformatics, statistical

analysis, made all of the figures, and wrote the first draft; all authors contributed to writing and

editing of the manuscript.

683
684 **Figure 1.** A) Location of the study area and plots (6 burned; 3 unburned) within the Holy Fire
685 burn scar in the Cleveland National Forest in Southern California. Soil burn severity is based on
686 the USDA BAER classifications. B) Plot experimental design, each with four 1m^2 subplots
687 placed 5m from the center in each cardinal direction. C) Collection of the top 10 cm of soil
688 beneath the ash with a releasable bulb planter within the 1 m^2 subplots at 17 days post-fire.
689
690

Figure 2. Change in biomass for A) bacteria and B) fungi and change in richness for C) bacteria and D) fungi at each of the 9 time-points in the first year post-fire in burned and unburned plots. Points represent mean differences averaged across burned and unburned plots, and bars represent the standard error of the mean. Precipitation is based on total monthly precipitation (mm). The significance of burned versus unburned plots per time point is denoted with an asterisk and is based on a negative binomial regression.

Figure 3. Change in A) bacterial and B) fungal biomass and C) bacterial and D) fungal richness in relation to soil burn severity measured as ash depth at 17 days post-fire. The significance of biomass and species richness against soil burn severity (SBS) and its interaction with time since (TSF) fire is based on a negative binomial regression.

Figure 4. NMDS plot of A) bacterial and C) fungal community composition with colors denoting treatment and shape indicating time since fire. The NMDS is based on 3-dimensions and has a stress value of 0.11 for bacteria and 0.12 for fungi. Change in relative sequence abundance of the five most abundant B) bacterial and D) fungal genera in the burned plots over time. Note, the dominance of the top 5 bacterial genera began at 34 days post-fire since there wasn't clear dominance at days 17 and 25 due to the lag in post-fire bacterial declines.

Figure 5. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in burned plots at all 9-time points. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in burned plots. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each time point for the burned E) bacterial and F) fungal communities. Note time points farther apart indicate a community turnover, and overlapping points with standard errors represent a lack of compositional turnover.

Figure 6. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in unburned plots at all 9-time points. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in unburned plots displaying no significant succession. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each time point for E) bacteria and F) fungi in the unburned plots. Note overlapping standard error bars indicate no community turnover across time in unburned plots.

Reference

Abatzoglou, J. T., Williams, A. P., & Barbero, R. (2019). Global Emergence of Anthropogenic Climate Change in Fire Weather Indices. *Geophysical Research Letters*, *46*(1), 326–336. https://doi.org/10.1029/2018GL080959

Agee, J. K. (1993). *Fire Ecology of Pacific Northwest Forests*. Island Press.

Agerer, R. (2001). Exploration types of ectomycorrhizae. *Mycorrhiza*, *11*(2), 107–114. https://doi.org/10.1007/s005720100108

Albuquerque, L., & da Costa, M. S. (2014). The Families Conexibacteraceae, Patulibacteraceae and Solirubrobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes: Actinobacteria* (pp. 185–200). Springer. https://doi.org/10.1007/978-3-642-30138-4_200

Allen, M., Egerton-Warburton, L., Treseder, K., Cario, C. H., Lindahl, A., Lansing, J., Querejeta, J., Kårén, O., Harney, S., & Zink, T. (2005). *Biodiversity of Mycorrhizal Fungi in Southern California 1*. /paper/Biodiversity-of-Mycorrhizal-Fungi-in-Southern-1-Allen-Egerton-Warburton/445d4f6878927a5307669f5e73df3174703f4d2a

Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In *Wiley StatsRef: Statistics Reference Online* (pp. 1–15). American Cancer Society. https://doi.org/10.1002/9781118445112.stat07841

Averill, C., & Hawkes, C. V. (2016). Ectomycorrhizal fungi slow soil carbon cycling. *Ecology Letters*, *19*(8), 937–947. https://doi.org/10.1111/ele.12631

Bailey, A. C., Kellom, M., Poret-Peterson, A. T., Noonan, K., Hartnett, H. E., & Raymond, J. (2014). Draft Genome Sequence of Massilia sp. Strain BSC265, Isolated from Biological Soil Crust of Moab, Utah. *Genome Announcements*, *2*(6), e01199-14, 2/6/e01199-14. https://doi.org/10.1128/genomeA.01199-14

Baldani, J. I., Rouws, L., Cruz, L. M., Olivares, F. L., Schmid, M., & Hartmann, A. (2014). The Family Oxalobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes* (pp. 919–974). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-30197-1_291

Barnard, R. L., Osborne, C. A., & Firestone, M. K. (2013). Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *The ISME Journal*, *7*(11), 2229–2241. https://doi.org/10.1038/ismej.2013.104

Barro, S. C., & Conard, S. G. (1991). Fire effects on California chaparral systems: An overview. *Environment International*, *17*(2–3), 135–149. https://doi.org/10.1016/0160-4120(91)90096-9

Barton, K. (2020). *MuMIn: Multi-Model Inference* (R package version 1.43.17.) [Computer software]. https://CRAN.R-project.org/package=MuMIn

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, *19*(1), 134–143. https://doi.org/10.1111/j.1466-8238.2009.00490.x

Blazewicz, S. J., Schwartz, E., & Firestone, M. K. (2014). Growth and death of bacteria and fungi underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. *Ecology*, *95*(5), 1162–1172. https://doi.org/10.1890/13-1031.1

Bliss, C. I. (1953). *Fitting the negative binomial distribution to biological data (with "Note on the efficient fitting of the negative binomial" by R. A. Fisher), Biometrics, 9,* (9th ed.). Biometrics.

Bodí, M. B., Martin, D. A., Balfour, V. N., Santín, C., Doerr, S. H., Pereira, P., Cerdà, A., & Mataix-Solera, J. (2014). Wildland fire ash: Production, composition and eco-hydro-

geomorphic effects. *Earth-Science Reviews*, *130*, 103–127.
https://doi.org/10.1016/j.earscirev.2013.12.007

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,
Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger,
K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A.
M., Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and
extensible microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8), 852–
857. https://doi.org/10.1038/s41587-019-0209-9

Brown, S. P., & Jumpponen, A. (2014). Contrasting primary successional trajectories of fungi
and bacteria in retreating glacier soils. *Molecular Ecology*, *23*(2), 481–497.
https://doi.org/10.1111/mec.12487

Bruns, T. D., Chung, J. A., Carver, A. A., & Glassman, S. I. (2020). A simple pyrocosm for
studying soil microbial response to fire reveals a rapid, massive response by Pyronema
species. *PLOS ONE*, *15*(3), e0222691. https://doi.org/10.1371/journal.pone.0222691

Cairney, J. W. G., & Bastias, B. A. (2007). Influences of fire on forest soil fungal
communitiesThis article is one of a selection of papers published in the Special Forum on
Towards Sustainable Forestry — The Living Soil: Soil Biodiversity and Ecosystem
Function. *Canadian Journal of Forest Research*, *37*(2), 207–215.
https://doi.org/10.1139/x06-190

Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *The ISME Journal*, *11*(12),
2639–2643. https://doi.org/10.1038/ismej.2017.119

Capitanio, R., & Carcaillet, C. (2008). Post-fire Mediterranean vegetation dynamics and
diversity: A discussion of succession models. *Forest Ecology and Management*, *255*(3),
431–439. https://doi.org/10.1016/j.foreco.2007.09.010

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P.
J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of
millions of sequences per sample. *Proceedings of the National Academy of Sciences*,
108(Supplement_1), 4516–4522. https://doi.org/10.1073/pnas.1000080107

Carini, P., Marsden, P. J., Leff, J. W., Morgan, E. E., Strickland, M. S., & Fierer, N. (2016).
Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. *Nature*
Microbiology, *2*(3), 1–6. https://doi.org/10.1038/nmicrobiol.2016.242

Certini, G., Moya, D., Lucas-Borja, M. E., & Mastrolonardo, G. (2021). The impact of fire on
soil-dwelling biota: A review. *Forest Ecology and Management*, *488*, 118989.
https://doi.org/10.1016/j.foreco.2021.118989

Collins, S. L. (1990). Patterns of Community Structure During Succession in Tallgrass Prairie.
Bulletin of the Torrey Botanical Club, *117*(4), 397–408. https://doi.org/10.2307/2996837

Collins, S. L., Micheli, F., & Hartt, L. (2000). A method to determine rates and patterns of
variability in ecological communities. *Oikos*, *91*(2), 285–293.
https://doi.org/10.1034/j.1600-0706.2000.910209.x

Cordero, R. J., & Casadevall, A. (2017). Functions of fungal melanin beyond virulence. *Fungal*
Biology Reviews, *31*(2), 99–112. https://doi.org/10.1016/j.fbr.2016.12.003

Crow, J. F. (1992). An advantage of sexual reproduction in a rapidly changing environment. *The*
Journal of Heredity, *83*(3), 169–173.
https://doi.org/10.1093/oxfordjournals.jhered.a111187

Dalling, J. W. (2019). *Pioneer Species. In:Fath BD (ed) Encyclopedia of ecology.* Elsevier.
https://doi.org/10.1016/B978-0-444-63768-0.00534-5

Derroire, G., Balvanera, P., Castellanos-Castro, C., Decocq, G., Kennard, D. K., Lebrija-Trejos,
E., Leiva, J. A., Odén, P.-C., Powers, J. S., Rico-Gray, V., Tigabu, M., & Healey, J. R.
(2016). Resilience of tropical dry forests – a meta-analysis of changes in species diversity
and composition during secondary succession. *Oikos*, *125*(10), 1386–1397.
https://doi.org/10.1111/oik.03229
Dix, N. J., & Webster, J. (1995). *Fungal Ecology*. Chapman & Hall.
Donato, D. C., Campbell, J. L., & Franklin, J. F. (2012). Multiple successional pathways and
precocity in forest development: Can some forests be born complex? *Journal of*
Vegetation Science, *23*(3), 576–584. https://doi.org/10.1111/j.1654-1103.2011.01362.x
Dooley, S. R., & Treseder, K. K. (2012). The effect of fire on microbial biomass: A meta-
analysis of field studies. *Biogeochemistry*, *109*(1/3), 49–61.
Dove, N. C., & Hart, S. C. (2017). Fire Reduces Fungal Species Richness and In Situ
Mycorrhizal Colonization: A Meta-Analysis. *Fire Ecology*, *13*(2), 37–65.
https://doi.org/10.4996/fireecology.130237746
Dunn, P. H., Wells, W. G. I., Dickey, J., & Wohlgemuth, M. (1982). *Role of Fungi in Postfire*
Stabilization of Chaparral Ash Beds (PSW-58, pp. 378–381) [Gen. Tech. Rep]. Pacific
Southwest Forest and Range Experiment Station, Forest Service, U.S.U.S. Department of
Agriculture.
Egler, F. E. (1954). Vegetation science concepts I. Initial floristic composition, a factor in old-
field vegetation development with 2 figs. *Vegetatio*, *4*(6), 412–417.
https://doi.org/10.1007/BF00275587
Eisenhauer, N., Schulz, W., Scheu, S., & Jousset, A. (2013). Niche dimensionality links
biodiversity and invasibility of microbial communities. *Functional Ecology*, *27*(1), 282–
288. https://doi.org/10.1111/j.1365-2435.2012.02060.x
Enright, D. J., Frangioso, K. M., Isobe, K., Rizzo, D. M., & Glassman, S. I. (2021). Mega-fire in
Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and Selects for
Pyrophilous Taxa and Traits that are Phylogenetically Conserved. *BioRxiv*,
2021.06.30.450634. https://doi.org/10.1101/2021.06.30.450634
Evans, S. E., & Wallenstein, M. D. (2012). Soil microbial community response to drying and
rewetting stress: Does historical precipitation regime matter? *Biogeochemistry*, *109*(1/3),
101–116.
Feng, S., & Fu, Q. (2013). Expansion of global drylands under a warming climate. *Atmospheric*
Chemistry and Physics, *13*(19), 10081–10094. https://doi.org/10.5194/acp-13-10081-
2013
Fernández-Martínez, M. A., Pérez-Ortega, S., Pointing, S. B., Allan Green, T. G., Pintado, A.,
Rozzi, R., Sancho, L. G., & de los Ríos, A. (2017). Microbial succession dynamics along
glacier forefield chronosequences in Tierra del Fuego (Chile). *Polar Biology*, *40*(10),
1939–1957. https://doi.org/10.1007/s00300-017-2110-7
Ferrenberg, S., O'Neill, S. P., Knelman, J. E., Todd, B., Duggan, S., Bradley, D., Robinson, T.,
Schmidt, S. K., Townsend, A. R., Williams, M. W., Cleveland, C. C., Melbourne, B. A.,
Jiang, L., & Nemergut, D. R. (2013). Changes in assembly processes in soil bacterial
communities following a wildfire disturbance. *The ISME Journal*, *7*(6), 1102–1111.
https://doi.org/10.1038/ismej.2013.11
Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of Soil Microbial
Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Applied and*
Environmental Microbiology, *71*(7), 4117–4120.
https://doi.org/10.1128/AEM.71.7.4117-4120.2005

Fischer, M. S., Stark, F. G., Berry, T. D., Zeba, N., Whitman, T., & Traxler, M. F. (2021). *Pyrolyzed substrates induce aromatic compound metabolism in the post-fire fungus,Pyronema domesticum* [Preprint]. Microbiology. https://doi.org/10.1101/2021.03.15.435558

Fujimura, K. E., Smith, J. E., Horton, T. R., Weber, N. S., & Spatafora, J. W. (2005). Pezizalean mycorrhizas and sporocarps in ponderosa pine (Pinus ponderosa) after prescribed fires in eastern Oregon, USA. *Mycorrhiza*, *15*(2), 79–86. https://doi.org/10.1007/s00572-004-0303-8

Gao, C., Montoya, L., Xu, L., Madera, M., Hollingsworth, J., Purdom, E., Singan, V., Vogel, J., Hutmacher, R. B., Dahlberg, J. A., Coleman-Derr, D., Lemaux, P. G., & Taylor, J. W. (2020). Fungal community assembly in drought-stressed sorghum shows stochasticity, selection, and universal ecological dynamics. *Nature Communications*, *11*(1), 34. https://doi.org/10.1038/s41467-019-13913-9

Gassibe, P. V., Fabero, R. F., Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto, P. (2011). Fungal community succession following wildfire in a Mediterranean vegetation type dominated by Pinus pinaster in Northwest Spain. *Forest Ecology and Management*, *262*(4), 655–662. https://doi.org/10.1016/j.foreco.2011.04.036

Glassman, S. I., Levine, C. R., DiRocco, A. M., Battles, J. J., & Bruns, T. D. (2016). Ectomycorrhizal fungal spore bank recovery after a severe forest fire: Some like it hot. *The ISME Journal*, *10*(5), 1228–1239. https://doi.org/10.1038/ismej.2015.182

Glassman, S. I., Randolph, J. W., Saroa, S. S., Capocchi, J. K., Walters, K. E., Pulido-Chavez, M. F., & Larios, L. (2021). *Prescribed versus wildfire impacts on exotic plants and soil microbes in California grasslands* (p. 2021.09.22.461426). https://doi.org/10.1101/2021.09.22.461426

Glassman, S. I., Weihe, C., Li, J., Albright, M. B. N., Looby, C. I., Martiny, A. C., Treseder, K. K., Allison, S. D., & Martiny, J. B. H. (2018). Decomposition responses to climate depend on microbial community composition. *Proceedings of the National Academy of Sciences*, *115*(47), 11994–11999. https://doi.org/10.1073/pnas.1811269115

Gottlieb, D. (1950). The physiology of spore germination in fungi. *The Botanical Review*, *16*(5), 229–257. https://doi.org/10.1007/BF02873609

Grady, E. N., MacDonald, J., Liu, L., Richman, A., & Yuan, Z.-C. (2016). Current knowledge and perspectives of Paenibacillus: A review. *Microbial Cell Factories*, *15*(1), 203. https://doi.org/10.1186/s12934-016-0603-7

Gray, L., & Kernaghan, G. (2020). Fungal Succession During the Decomposition of Ectomycorrhizal Fine Roots. *Microbial Ecology*, *79*(2), 271–284. https://doi.org/10.1007/s00248-019-01418-3

Grishkan, I., Korol, A. B., Nevo, E., & Wasser, S. P. (2003). Ecological stress and sex evolution in soil microfungi. *Proceedings of the Royal Society B: Biological Sciences*, *270*(1510), 13–18. https://doi.org/10.1098/rspb.2002.2194

Gu, H., Lou, J., Wang, H., Yang, Y., Wu, L., Wu, J., & Xu, J. (2016). Biodegradation, Biosorption of Phenanthrene and Its Trans-Membrane Transport by Massilia sp. WF1 and Phanerochaete chrysosporium. *Frontiers in Microbiology*, *7*. https://doi.org/10.3389/fmicb.2016.00038

Guhr, A., & Kircher, S. (2020). Drought-Induced Stress Priming in Two Distinct Filamentous Saprotrophic Fungi. *Microbial Ecology*, *80*(1), 27–33. https://doi.org/10.1007/s00248-019-01481-w

Hallett, L. M., Jones, S. K., MacDonald, A. A. M., Jones, M. B., Flynn, D. F. B., Ripplinger, J.,
Slaughter, P., Gries, C., & Collins, S. L. (2016). codyn: An r package of community
dynamics metrics. *Methods in Ecology and Evolution*, *7*(10), 1146–1151.
https://doi.org/10.1111/2041-210X.12569
Hart, S. C., DeLuca, T. H., Newman, G. S., MacKenzie, M. D., & Boyle, S. I. (2005). Post-fire
vegetative dynamics as drivers of microbial community structure and function in forest
soils. *Forest Ecology and Management*, *220*(1–3), 166–184.
https://doi.org/10.1016/j.foreco.2005.08.012
Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto, P. (2013). Post-fire fungal
succession in a Mediterranean ecosystem dominated by Cistus ladanifer L. *Forest*
Ecology and Management, *289*, 48–57. https://doi.org/10.1016/j.foreco.2012.10.009
Homyak, P. M., Sickman, J. O., Miller, A. E., Melack, J. M., Meixner, T., & Schimel, J. P.
(2014). Assessing Nitrogen-Saturation in a Seasonally Dry Chaparral Watershed:
Limitations of Traditional Indicators of N-Saturation. *Ecosystems*, *17*(7), 1286–1305.
https://doi.org/10.1007/s10021-014-9792-2
Iffis, B., St-Arnaud, M., & Hijri, M. (2014). Bacteria associated with arbuscular mycorrhizal
fungi within roots of plants growing in a soil highly contaminated with aliphatic and
aromatic petroleum hydrocarbons. *FEMS Microbiology Letters*, *358*(1), 44–54.
https://doi.org/10.1111/1574-6968.12533
J., E. de los M., A., M., & F., V. (2001). Metabolic profiles and aprE expression in anaerobic
cultures of Bacillus subtilis using nitrate as terminal electron acceptor. *Applied*
Microbiology and Biotechnology, *57*(3), 379–384.
https://doi.org/10.1007/s002530100749
Kalayu, G. (2019). Phosphate Solubilizing Microorganisms: Promising Approach as
Biofertilizers. *International Journal of Agronomy*, *2019*, 1–7.
https://doi.org/10.1155/2019/4917256
Kaur, N., Seuylemezian, A., Patil, P. P., Patil, P., Krishnamurti, S., Varelas, J., Smith, D. J.,
Mayilraj, S., & Vaishampayan, P. (2018). Paenibacillus xerothermodurans sp. Nov., an
extremely dry heat resistant spore forming bacterium isolated from the soil of Cape
Canaveral, Florida. *International Journal of Systematic and Evolutionary Microbiology*,
68(10), 3190–3196. https://doi.org/10.1099/ijsem.0.002967
Keeley, J. E., Fotheringham, C. J., & Baer-Keeley, M. (2005). Determinants of Postfire
Recovery and Succession in Mediterranean-Climate Shrublands of California. *Ecological*
Applications, *15*(5), 1515–1534. https://doi.org/10.1890/04-1005
Keeley, J. E., & Zedler, P. H. (2009). Large, high-intensity fire events in Southern California
shrublands: Debunking the fine-grain age patch model. In *Ecological Applications* (Vol.
19, Issue 1, p. 26). https://doi.org/10.1890/08-0281.1
Kindt, R., & Coe, R. (2005). *Tree diversity analysis. A manual and software for common*
statistical methods for ecological and biodiversity studies. World Agroforestry Centre
(ICRAF).
Kinzig, A. P., & Pacala, S. (2013). 9. Successional Biodiversity and Ecosystem Functioning. In
The Functional Consequences of Biodiversity: Empirical Progress and Theoretical
Extensions (MPB-33) (pp. 175–212). Princeton University Press.
https://doi.org/10.1515/9781400847303.175
Koide, R. T., Fernandez, C., & Malcolm, G. (2014). Determining place and process: Functional
traits of ectomycorrhizal fungi that affect both community structure and ecosystem
function. *New Phytologist*, *201*(2), 433–439. https://doi.org/10.1111/nph.12538

Kõljalg, U., Larsson, K.-H., Abarenkov, K., Nilsson, R. H., Alexander, I. J., Eberhardt, U., Erland, S., Høiland, K., Kjøller, R., Larsson, E., Pennanen, T., Sen, R., Taylor, A. F. S., Tedersoo, L., & Vrålstad, T. (2005). UNITE: A database providing web-based methods for the molecular identification of ectomycorrhizal fungi. *New Phytologist*, *166*(3), 1063–1068. https://doi.org/10.1111/j.1469-8137.2005.01376.x

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*, *79*(17), 5112–5120. https://doi.org/10.1128/AEM.01043-13

Krappmann, S., & Braus, G. H. (2005). Nitrogen metabolism of *Aspergillus* and its role in pathogenicity. *Medical Mycology*, *43*(s1), 31–40. https://doi.org/10.1080/13693780400024271

Lagashetti, A. C., Dufossé, L., Singh, S. K., & Singh, P. N. (2019). Fungal Pigments and Their Prospects in Different Industries. *Microorganisms*, *7*(12), 604. https://doi.org/10.3390/microorganisms7120604

Li, X., Rui, J., Mao, Y., Yannarell, A., & Mackie, R. (2014). Dynamics of the bacterial community structure in the rhizosphere of a maize cultivar. *Soil Biology and Biochemistry*, *68*, 392–401. https://doi.org/10.1016/j.soilbio.2013.10.017

Liu, C. M., Kachur, S., Dwan, M. G., Abraham, A. G., Aziz, M., Hsueh, P.-R., Huang, Y.-T., Busch, J. D., Lamit, L. J., Gehring, C. A., Keim, P., & Price, L. B. (2012). FungiQuant: A broad-coverage fungal quantitative real-time PCR assay. *BMC Microbiology*, *12*(1), 255. https://doi.org/10.1186/1471-2180-12-255

Livne-Luzon, S., Shemesh, H., Osem, Y., Carmel, Y., Migael, H., Avidan, Y., Tsafrir, A., Glassman, S. I., Bruns, T. D., & Ovadia, O. (2021). High resilience of the mycorrhizal community to prescribed seasonal burnings in eastern Mediterranean woodlands. *Mycorrhiza*, *31*(2), 203–216. https://doi.org/10.1007/s00572-020-01010-5

Loreau, M., & de Mazancourt, C. (2008). Species Synchrony and Its Drivers: Neutral and Nonneutral Community Dynamics in Fluctuating Environments. *The American Naturalist*, *172*(2), E48–E66. https://doi.org/10.1086/589746

Martin, M. (2011). *Cutadapt removes adapter sequences from high-throughput sequencing reads*. https://journal.embnet.org/index.php/embnetjournal/article/view/200/458

McGee, P. A., Markovina, A.-L., Jeong, G. C. E., & Cooper, E. D. (2006). Trichocomaceae in bark survive high temperatures and fire: Trichocomaceae in bark survive high temperatures and fire. *FEMS Microbiology Ecology*, *56*(3), 365–371. https://doi.org/10.1111/j.1574-6941.2006.00079.x

McMullan-Fisher, S. J. M., Tom W. May, Richard M. Robinson, Tina L. Bell, Teresa Lebel, Pam Catcheside, & Alan York. (2011). *Fungi and fire in Australian ecosystems: A review of current knowledge, management implications and future directions*. *59*, 70–90.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLOS ONE*, *8*(4), e61217. https://doi.org/10.1371/journal.pone.0061217

Monciardini, P., Cavaletti, L., Schumann, P., Rohde, M., & Donadio, S. (2003). Conexibacter woesei gen. Nov., sp. Nov., a novel representative of a deep evolutionary line of descent within the class Actinobacteria. *International Journal of Systematic and Evolutionary Microbiology*, *53*(2), 569–576. https://doi.org/10.1099/ijs.0.02400-0

Moore, E. J. (1962). The Ontogeny of the Sclerotia of Pyronema Domesticum. *Mycologia*, *54*(3), 312–316. https://doi.org/10.1080/00275514.1962.12025004

Neary, D. G., Klopatek, C. C., DeBano, L. F., & Ffolliott, P. F. (1999). Fire effects on belowground sustainability: A review and synthesis. *Forest Ecology and Management*, *122*(1–2), 51–71. https://doi.org/10.1016/S0378-1127(99)00032-8

Nguyen, H. D. T., Nickerson, N. L., & Seifert, K. A. (2013). Basidioascus and Geminibasidium: A new lineage of heat-resistant and xerotolerant basidiomycetes. *Mycologia*, *105*(5), 1231–1250. JSTOR.

Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., & Kennedy, P. G. (2016). FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, *20*, 241–248. https://doi.org/10.1016/j.funeco.2015.06.006

Ofek, M., Hadar, Y., & Minz, D. (2012). Ecology of Root Colonizing Massilia (Oxalobacteraceae). *PLOS ONE*, *7*(7), e40117. https://doi.org/10.1371/journal.pone.0040117

Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2018). *Vegan:Community Ecology Package-Version 2.5.2. Ordination Methods, Diversity Analysis, and Other Functions for Community and Vegetation Ecologist*.

Owen, S. M., Patterson, A. M., Gehring, C. A., Sieg, C. H., Baggett, L. S., & Fulé, P. Z. (2019). Large, high-severity burn patches limit fungal recovery 13 years after wildfire in a ponderosa pine forest. *Soil Biology and Biochemistry*, *139*, 107616. https://doi.org/10.1016/j.soilbio.2019.107616

Pacala, S. W., Canham, C. D., Saponara, J., Silander, J. A., Kobe, R. K., & Ribbens, E. (1996). Forest Models Defined by Field Measurements: Estimation, Error Analysis and Dynamics. *Ecological Monographs*, *66*(1), 1–43. https://doi.org/10.2307/2963479

Pandolfi, J. M. (2008). Succession. In S. E. Jørgensen & B. D. Fath (Eds.), *Encyclopedia of Ecology* (pp. 3416–3424). Academic Press. https://doi.org/10.1016/B978-008045405-4.00547-4

Paredes-Sabja, D., Setlow, P., & Sarker, M. R. (2011). Germination of spores of Bacillales and Clostridiales species: Mechanisms and proteins involved. *Trends in Microbiology*, *19*(2), 85–94. https://doi.org/10.1016/j.tim.2010.10.004

Parson, A., Robichaud, P. R., Lewis, S. A., Napper, C., & Clark, J. T. (2010). *Field guide for mapping post-fire soil burn severity* (RMRS-GTR-243; p. RMRS-GTR-243). U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. https://doi.org/10.2737/RMRS-GTR-243

Pena, R., Offermann, C., Simon, J., Naumann, P. S., Geßler, A., Holst, J., Dannenmann, M., Mayer, H., Kögel-Knabner, I., Rennenberg, H., & Polle, A. (2010). Girdling Affects Ectomycorrhizal Fungal (EMF) Diversity and Reveals Functional Differences in EMF Community Composition in a Beech Forest. *Applied and Environmental Microbiology*, *76*(6), 1831–1841. https://doi.org/10.1128/AEM.01703-09

Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2018). Resilience to fire of phylogenetic diversity across biological domains. *Molecular Ecology*, *27*(13), 2896–2908. https://doi.org/10.1111/mec.14729

Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2020). Soil microbiome drives the recovery of ecosystem functions after fire. *Soil Biology and Biochemistry*, *149*, 107948. https://doi.org/10.1016/j.soilbio.2020.107948

Placella, S. A., Brodie, E. L., & Firestone, M. K. (2012). Rainfall-induced carbon dioxide pulses
result from sequential resuscitation of phylogenetically clustered microbial groups.
Proceedings of the National Academy of Sciences, *109*(27), 10931–10936.
https://doi.org/10.1073/pnas.1204306109
Pourreza, M., Hosseini, S. M., Safari Sinegani, A. A., Matinizadeh, M., & Dick, W. A. (2014).
Soil microbial activity in response to fire severity in Zagros oak (Quercus brantii Lindl.)
forests, Iran, after one year. *Geoderma*, *213*, 95–102.
https://doi.org/10.1016/j.geoderma.2013.07.024
Pressler, Y., Moore, J. C., & Cotrufo, M. F. (2019). Belowground community responses to fire:
Meta-analysis reveals contrasting responses of soil microorganisms and mesofauna.
Oikos, *128*(3), 309–327. https://doi.org/10.1111/oik.05738
Pulido-Chavez, M. F., Alvarado, E. C., DeLuca, T. H., Edmonds, R. L., & Glassman, S. I.
(2021). High-severity wildfire reduces richness and alters composition of
ectomycorrhizal fungi in low-severity adapted ponderosa pine forests. *Forest Ecology*
and Management, *485*, 118923. https://doi.org/10.1016/j.foreco.2021.118923
Qin, Q., & Liu, Y. (2021). Changes in microbial communities at different soil depths through the
first rainy season following severe wildfire in North China artificial Pinus tabulaeformis
forest. *Journal of Environmental Management*, *280*, 111865.
https://doi.org/10.1016/j.jenvman.2020.111865
R Core Team. (2021). *R: A language and environment for statistical computing. R Foundation*
for Statistical Computing. https://www.R-project.org/.
Reazin, C., Morris, S., Smith, J. E., Cowan, A. D., & Jumpponen, A. (2016). Fires of differing
intensities rapidly select distinct soil fungal communities in a Northwest US ponderosa
pine forest ecosystem. *Forest Ecology and Management*, *377*, 118–127.
https://doi.org/10.1016/j.foreco.2016.07.002
Rhodes, J. C. (2006). *Aspergillus fumigatus*: Growth and virulence. *Medical Mycology*, *44*(s1),
77–81. https://doi.org/10.1080/13693780600779419
Riley, K. L., & Loehman, R. A. (2016). Mid-21st-century climate changes increase predicted fire
occurrence and fire season length, Northern Rocky Mountains, United States. *Ecosphere*,
7(11), e01543. https://doi.org/10.1002/ecs2.1543
Ross, G. J. S., & Preece, D. A. (1985). The Negative Binomial Distribution. *Journal of the Royal*
Statistical Society. Series D (The Statistician), *34*(3), 323–335. JSTOR.
https://doi.org/10.2307/2987659
Rousk, J., Brookes, P. C., & Bååth, E. (2010). Investigating the mechanisms for the opposing pH
relationships of fungal and bacterial growth in soil. *Soil Biology and Biochemistry*, *42*(6),
926–934. https://doi.org/10.1016/j.soilbio.2010.02.009
Rundel, P. W. (2018). California Chaparral and Its Global Significance. In E. C. Underwood, H.
D. Safford, N. A. Molinari, & J. E. Keeley (Eds.), *Valuing Chaparral* (pp. 1–27).
Springer International Publishing. https://doi.org/10.1007/978-3-319-68303-4_1
Schimel, J. P. (2018). Life in Dry Soils: Effects of Drought on Soil Microbial Communities and
Processes. *Annual Review of Ecology, Evolution, and Systematics*, *49*(1), 409–432.
https://doi.org/10.1146/annurev-ecolsys-110617-062614
Schluter, D., & Ricklefs, R. E. (1993). *Species diversity: An introduction to the problem. Species*
diversity in ecological communities. Chicago University Press.
Selbmann, L., Egidi, E., Isola, D., Onofri, S., Zucconi, L., de Hoog, G. S., Chinaglia, S., Testa,
L., Tosi, S., Balestrazzi, A., Lantieri, A., Compagno, R., Tigini, V., & Varese, G. C.
(2013). Biodiversity, evolution and adaptation of fungi in extreme environments. *Plant*

Biosystems - An International Journal Dealing with All Aspects of Plant Biology, *147*(1), 237–246. https://doi.org/10.1080/11263504.2012.753134

Semenova-Nelsen, T. A., Platt, W. J., Patterson, T. R., Huffman, J., & Sikes, B. A. (2019). Frequent fire reorganizes fungal communities and slows decomposition across a heterogeneous pine savanna landscape. *New Phytologist*, *224*(2), 916–927. https://doi.org/10.1111/nph.16096

Slepecky, R. A., & Hemphill, H. E. (2006). The Genus Bacillus—Nonmedical. In M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer, & E. Stackebrandt (Eds.), *The Prokaryotes: Volume 4: Bacteria: Firmicutes, Cyanobacteria* (pp. 530–562). Springer US. https://doi.org/10.1007/0-387-30744-3_16

Smith, M. E., Henkel, T. W., & Rollins, J. A. (2015). How many fungi make sclerotia? *Fungal Ecology*, *13*, 211–220. https://doi.org/10.1016/j.funeco.2014.08.010

Sun, S., Li, S., Avera, B. N., Strahm, B. D., & Badgley, B. D. (2017). Soil Bacterial and Fungal Communities Show Distinct Recovery Patterns during Forest Ecosystem Restoration. *Applied and Environmental Microbiology*, *83*(14), e00966-17. https://doi.org/10.1128/AEM.00966-17

Taylor, D. L., Walters, W. A., Lennon, N. J., Bochicchio, J., Krohn, A., Caporaso, J. G., & Pennanen, T. (2016). Accurate Estimation of Fungal Diversity and Abundance through Improved Lineage-Specific Primers Optimized for Illumina Amplicon Sequencing. *Applied and Environmental Microbiology*, *82*(24), 7217–7226. https://doi.org/10.1128/AEM.02576-16

Toljander, J. F., Artursson, V., Paul, L. R., Jansson, J. K., & Finlay, R. D. (2005). Attachment of di¡erent soil bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal vitalityand fungal species. *FEMS Microbiol Lett*, 7.

Torti, A., Lever, M. A., & Jørgensen, B. B. (2015). Origin, dynamics, and implications of extracellular DNA pools in marine sediments. *Marine Genomics*, *24*, 185–196. https://doi.org/10.1016/j.margen.2015.08.007

Treseder, K. K., Mack, M. C., & Cross, A. (2004). Relationships Among Fires, Fungi, and Soil Dynamics in Alaskan Boreal Forests. *Ecological Applications*, *14*(6), 1826–1838. https://doi.org/10.1890/03-5133

Turner, M. G., Baker, W. L., Peterson, C. J., & Peet, R. K. (1998). Factors Influencing Succession: Lessons from Large, Infrequent Natural Disturbances. *Ecosystems*, *1*(6), 511–523. https://doi.org/10.1007/s100219900047

Van der Heijden, M. G. A., Klironomos, J. N., Ursic, M., Moutoglis, P., Streitwolf-Engel, R., Boller, T., Wiemken, A., & Sanders, I. R. (1998). Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*, *396*(6706), 69–72. https://doi.org/10.1038/23932

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (Fourth edition). Springer. https://www.stats.ox.ac.uk/pub/MASS4/.

Wan, W., Qin, Y., Wu, H., Zuo, W., He, H., Tan, J., Wang, Y., & He, D. (2020). Isolation and Characterization of Phosphorus Solubilizing Bacteria With Multiple Phosphorus Sources Utilizing Capability and Their Potential for Lead Immobilization in Soil. *Frontiers in Microbiology*, *0*. https://doi.org/10.3389/fmicb.2020.00752

Warcup, J. H., & Baker, K. F. (1963). Occurrence of Dormant Ascospores in Soil. *Nature*, *197*(4874), 1317–1318. https://doi.org/10.1038/1971317a0

Whitman, T., Whitman, E., Woolet, J., Flannigan, M. D., Thompson, D. K., & Parisien, M.-A. (2019). Soil bacterial and fungal response to wildfires in the Canadian boreal forest

across a burn severity gradient. *Soil Biology and Biochemistry, 138*, 107571. https://doi.org/10.1016/j.soilbio.2019.107571

Woolet, J., & Whitman, T. (2020). Pyrogenic organic matter effects on soil bacterial community composition. *Soil Biology and Biochemistry*, *141*, 107678. https://doi.org/10.1016/j.soilbio.2019.107678

Xiang, X., Shi, Y., Yang, J., Kong, J., Lin, X., Zhang, H., Zeng, J., & Chu, H. (2014). Rapid recovery of soil bacterial communities after wildfire in a Chinese boreal forest. *Scientific Reports*, *4*. https://doi.org/10.1038/srep03829

Yang, T., Tedersoo, L., Lin, X., Fitzpatrick, M. C., Jia, Y., Liu, X., Ni, Y., Shi, Y., Lu, P., Zhu, J., & Chu, H. (2020). Distinct fungal successional trajectories following wildfire between soil horizons in a cold-temperate forest. *New Phytologist*, *227*(2), 572–587. https://doi.org/10.1111/nph.16531

Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., & Glöckner, F. O. (2014). The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. *Nucleic Acids Research*, *42*(D1), D643–D648. https://doi.org/10.1093/nar/gkt1209