1   Bacterial and fungal communities experience rapid secondary succession during the first year
2   following wildfire in California chaparral
3
4

5   M. Fabiola Pulido-Chavez[1,*], James W. J. Randolph[2], Cassandra Zalman[3], Loralee Larios[4], Peter
6   M. Homyak[5], and Sydney I. Glassman[6,*]
7
8
[1]Department of Microbiology and Plant Pathology, University of California-Riverside, 900
University Ave, Riverside, CA, 92521, mpuli011@ucr.edu
[2]Department of Microbiology and Plant Pathology, University of California-Riverside,
900 University Ave, Riverside, CA, 92521, jrand005@ucr.edu
[3]Schmid College of Science and Technology, Chapman University, Orange, CA, 92866,
medvedef@chapman.edu
[4]Department of Botany and Plant Sciences, University of California-Riverside, 900 University
Ave, Riverside, CA, 92521, loralee@ucr.edu
[5]Department of Environmental Sciences, University of California-Riverside, 900 University Ave,
Riverside, CA, 92521, phomyak@ucr.edu
[6]Department of Microbiology and Plant Pathology, University of California-Riverside,
900 University Ave, Riverside, CA, 92521, sydneyg@ucr.edu
26 *Correspondence: Sydney I Glassman, sydneyg@ucr.edu and M. Fabiola Pulido-Chavez,
27 mpuli011@ucr.ed Department of Microbiology and Plant Pathology, University of California-
28 Riverside

Keywords: Secondary succession, Chaparral, Wildfire, Bacteria, Fungi, Temporal turnover,
pyrophilous microbes, Illumina MiSeq, next-generation sequencing

Abstract

The rise in wildfire frequency and severity across the globe has increased interest in

secondary succession. However, despite the role of soil microbial communities in the

regeneration of post-fire vegetation and biogeochemical cycling, the lack of measurements

immediately post-fire and at high temporal resolution has resulted in a limited understanding of

microbial secondary succession. To fill this knowledge gap, we sampled soils at 17, 25, 34, 67,

95, 131, 187, 286, and 376 days after a Southern California wildfire in fire-adapted chaparral

shrublands. We assessed bacterial and fungal biomass with qPCR of 16S and 18S and richness

and composition with Illumina MiSeq sequencing of 16S and ITS2 amplicons. Fire severely

reduced bacterial biomass by 47% and richness by 46%, but the impacts were stronger for fungi,

with biomass decreasing by 86% and richness by 68%. While bacterial biomass and richness

oscillated in response to precipitation, the declines in biomass and richness for both groups

persisted for the entire year. In contrast to the unburned plots, burned fungal and bacterial

communities experienced rapid succession, with 5-6 compositional turnover periods. Analogous

to plants, "fire-loving" pyrophilous microbes increased post-fire and drove successional

dynamics. Interestingly, many of the same pyrophilous taxa found in forests worldwide also

dominated post-fire chaparral and traded off markedly in abundance over time. For fungi, an

initial peak in the Basidiomycete yeast *Geminibasidium* traded off with the filamentous

Ascomycetes *Pyronema, Aspergillus*, and *Penicillium*. For bacteria, the Proteobacteria genus

Massilia dominated burned plots throughout the year, but the following most dominant bacteria,

the Firmicute *Bacillus* and Proteobacteria *Noviherbaspirillum*, traded off in abundance over time.

Our unprecedented high-resolution temporal sampling allowed us to capture post-fire microbial

secondary successional dynamics and show that putative tradeoffs in thermotolerance,

colonization, and competition among dominant pyrophilous microbes control microbial

succession with possible implications for ecosystem function.

1. Introduction

The rapid increase in wildfire frequency, severity, and extent in the western United States

(Riley & Loehman, 2016) and around the globe (Abatzoglou et al., 2019) has renewed interest in

secondary succession. Secondary succession, or the trajectory along which an ecosystem

develops following a disturbance, such as a wildfire, has been extensively studied for plants

(Derroire et al., 2016; Donato et al., 2012), but belowground microbial communities have been

comparatively overlooked. Understanding how wildfires alter soil microbial succession may be

necessary to predict post-fire effects on ecosystem recovery, and function since soil microbes

drive post-fire organic matter decomposition (Semenova-Nelsen et al., 2019), nutrient cycling

(Pérez-Valera et al., 2020), and plant regeneration (Dove & Hart, 2017).

Plant succession is a major process affecting the health and function of ecosystems.

During succession, the dominant species may change in an orderly and predictable manner

(Shugart, 2013). Early colonizers (fast-growing, short-lived species) are often replaced by long-

lived stable communities . However, wildfire can reset the successional clock, shifting the course

of succession away from equilibrium (Reilly & Spies, 2016) and initiating secondary succession.

Because secondary succession often begins with low competition in a nutrient-rich environment

(Dalling, 2008), colonization and resource use tradeoffs will often drive successional dynamics

(Tilman, 1990). Indeed, the sequence of species replacement suggests that early colonizers with

similar resource use (i.e., overlapping niches) will dominate open space but will inevitably be

replaced by late-stage species with differentiation in resource use (i.e., complementarity),

allowing for species coexistence (Pacala et al., 1996; Turnbull et al., 2013). However, whether

the patterns recognized in plant successional theory translate to belowground soil microbes

remains unclear.

Research on post-fire soil microbiomes suggests that fires can reset microbial succession

via fire-induced mortality (Hart et al., 2005), reductions in microbial richness and biomass

(Dooley & Treseder, 2012; Pressler et al., 2019), and the replacement of fungal basidiomycetes

and symbiotic mycorrhizal fungi with ascomycetes and saprobic fungi (Cairney & Bastias, 2007;

Fox et al., 2022). Indeed, for over a century, pyrophilous "fire-loving" fungi have been

consistently found in post-fire mushroom surveys (McMullan-Fisher et al., 2011; Seaver, 1909).

More recently, next-generation sequencing indicates that the pyrophilous Ascomycete *Pyronema*

can increase 100 fold after prescribed fires (Reazin et al., 2016) and dominate over 60% of the

sequences in experimental pyrocosms (Bruns et al., 2020). Furthermore, there is increasing

evidence of pyrophilous bacteria, such as the Proteobacteria *Massilia* (Enright et al., 2022;

Whitman et al., 2019). Together, evidence suggests that pyrophilous microbes likely evolved

under the same fire regime as the ecosystems they inhabit (Dove et al., 2021) and likely have fire

adaptations analogous to plants (Rundel, 2018). Pyrophilous microbes have traits that allow for

the survival to fire (i.e., heat resistant spores, sclerotia) (Day et al., 2020; Petersen, 1970) and the

post-fire environment (i.e., xerotolerance, decomposition of nitrogen and aromatic hydrocarbons)

(Fischer et al., 2021; Nelson et al., 2021; Steindorff et al., 2021). However, most research is

based on single time-point sampling (Dove & Hart, 2017; Pressler et al., 2019); thus, the

successional trajectory of pyrophilous microbes is nearly unknown. However, recent research

suggests that bacteria and fungi experience rapid post-fire community changes (Ferrenberg et al.,

2013; Qin & Liu, 2021), leading to different successional trajectories (Ammitzboll et al., 2022).

Although these studies sampled two to three times in the first post-fire year, to adequately

capture the rapid growth and turnover rates of soil microbes and thus the dynamics needed to

develop microbial successional theory (Hannula et al., 2019), it is integral to sample directly

post-fire and at high temporal resolution.

Here, we propose that microbial ecological succession theory can be developed and

tested in nature by focusing on California chaparral. Chaparral is a shrubland adapted to high-

severity fire and a biodiversity hotspot distributed in Mediterranean climates worldwide (Barro

& Conard, 1991; Rundel, 2018). Chaparral plant secondary succession is relatively well

understood and typically initiated by fire (Keeley et al., 2005). Yet, although the dominant

chaparral vegetation forms mycorrhizal associations required for their establishment and survival

(Allen et al., 2005), little is known about chaparral microbial succession. Indeed, only 13% of

post-fire microbiome research occurs in shrublands (Pressler et al., 2019). With drylands

covering nearly 41% of Earth's terrestrial surface and expanding with climate change (Feng &

Fu, 2013), it is critical to understand secondary successional dynamics in dryland systems such

as chaparral. While our focus is on chaparral, which is likely to have fire-adapted microbes due

to its evolutionary history, we expect that successional patterns will be broadly generalizable to

other biomes since pyrophilous bacteria and fungi appear to be phylogenetically conserved

(Enright et al., 2022). Indeed, some fungal taxa within the Ascomycota family Pyronemataceae

and a few Basidiomycota genera appear to be globally distributed in fire-disturbed Spanish

shrublands (Pérez-Valera et al., 2018) and Pinaceae (Hughes et al., 2020; Reazin et al., 2016;

Whitman et al., 2019; Xiang et al., 2014), Eucalyptus (Ammitzboll et al., 2022; McMullan-

Fisher et al., 2011), and Redwood-Tanoak forests (Enright et al., 2022).

However, predicting the successional trajectory of soil microbes is complicated by direct

and indirect wildfire effects (Neary et al., 1999). Whereas soil burn severity drives direct fire

impacts (Reazin et al., 2016; Whitman et al., 2019), changes in soil moisture, pH, and nutrients

can indirectly impact post-fire microbes (Köster et al., 2014; Neary et al., 1999), potentially

driving succession via environmental filtering (Schluter & Ricklefs, 1993). Indeed, research

supports the role of differential responses to soil burn severity (Whitman et al., 2019) and soil

moisture in structuring microbiomes (Evans & Wallenstein, 2012; Placella et al., 2012; Yang et

al., 2021). Bacteria rapidly respond to soil wet-up (Barnard et al., 2013; Placella et al., 2012),

whereas fungi appear to be more resilient to soil moisture changes (Barnard et al., 2013; Evans &

Wallenstein, 2012). Furthermore, microbial life strategies can determine microbial response to

fire. Research shows that fire decreases ectomycorrhizal fungi (EMF) (Cowan et al., 2016;

Glassman et al., 2016; Pulido-Chavez et al., 2021) and arbuscular fungi (AMF) (Xiang et al.,

2015) but can temporarily increase saprobic richness (Enright et al., 2022; Semenova-Nelsen et

al., 2019). Yet, the limited studies on how fires affect multiple microbial guilds (Certini et al

2021) lack the resolution to identify the direct effect of wildfire and the length of time that

mycorrhizal fungi can survive after plant host death, a question essential for post-fire ecosystem

recovery.

Here, we performed the highest resolution temporal sampling of post-fire microbiomes to

date, sampling soil at 9-time points over one year following a chaparral wildfire, allowing us to

identify immediate and temporal effects of wildfire on microbial successional dynamics. We

then tested the following hypotheses: (**H1**) wildfire will decrease bacterial and fungal biomass

and richness, leading to a shift in the community composition, where one year will not be

sufficient time to return to unburned levels; (**H2**) wildfire will have a distinct impact on fungal

guilds with symbiotic mycorrhizal fungi experiencing the largest declines; **(H3)** higher soil burn

severity will lead to larger reductions in both bacteria and fungi while precipitation will impose a

larger impact on bacterial than fungal succession; **(H4)** succession will be initiated by

pyrophilous microbes, which like plants, will trade off in abundance over time based on their

traits.

2. Methods

2.1 Study Area, plot design, and soil collection

The Holy Fire burned 94 km^2 in the Cleveland National Forest in Southern California

from August 6 to September 13, 2018. On September 30, 2018, we established nine plots (6

burned and 3 unburned) along a gradient of fire severities (**Fig. 1A**). Plots were selected for

similarity in aspect, slope, elevation, and pre-fire vegetation dominance by manzanita

(Arctostaphylos glandulosa), an ectomycorrhizal host, and chamise (*Adenostoma fasciculatum),*

an arbuscular and ectomycorrhizal host (Allen et al., 2005). Plots were placed on average 25 m

from forest access roads to avoid edge effects. Each circular plot contained four 1 m^2 subplots

located 5 m from the center in each cardinal direction (**Fig. 1B**). Our study site experiences a

Mediterranean-type climate with warm, dry summers and cool, wet winters, with an average

yearly temperature of 17˚C and precipitation of 12.5 mm. Precipitation data, based on monthly

summaries, was gathered from the El Cariso weather station (raws.dri.edu). Additional plot

information can be found in the supplements (**Table S1**).

We sampled soils at 17, 25, 34, 67, 95, 131, 187, 286, and 376 days, corresponding to

approximately 2 and 3 weeks, and 1, 2, 3, 4, 6, 9, and 12 months post-fire. We used BAER soil

burn severity maps (Parson et al., 2010) to roughly locate plots at medium and high severity

(**Fig. 1A**), but their coarse-scale measurement (30m) does not coincide well with microbial

communities. Since ash depth increases with soil heating (Parson et al., 2010) and fire severity

(Bodí et al., 2014), we used ash depth as a proxy for soil burn severity, where we averaged three

separate measurements of ash depth (cm) per each 1 m² subplot at day 17, before any

precipitation event could wash away the ash. Thus, our measure of soil burn severity refers to

initial ash depth. At each of the 9 time points, we collected the top 10 cm of soil beneath the ash

or duff from each of the 36 subplots with a ~250 mL cleaned releasable bulb planter (**Fig. 1C**).

Soils were transported to the University of California-Riverside (UCR) within hours of sampling,

stored overnight at 4°C, sieved (2 mm) within 24 hours, and a subsample was frozen at -80°C for

DNA extraction.

2.3 DNA extraction, amplification, and sequencing

DNA was extracted from soil using Qiagen DNeasy PowerSoil Kits following the

manufacturer's protocol, except the centrifugation time was increased to 1.5 min after adding the

C3 solution due to large amounts of precipitate. DNA extracts were stored at -20°C for

subsequent analysis. Extracted DNA was amplified using the primer pair ITS4-fun and 5.8s

(Taylor et al., 2016) to amplify the ITS2 region for fungi and the primer pair 515F-806R to

amplify the V4 region of the 16S rRNA gene for archaea and bacteria (Caporaso et al., 2011)

using the Dual-Index Sequencing Strategy (DIP) (Kozich et al., 2013). Although our 16S primers

amplify archaea and bacteria, for simplicity, we refer to 16S methods and results simply as

bacteria since archaea only contributed <1% of sequencing reads. We conducted polymerase

chain reaction (PCR) in two steps. The first PCR amplified gene-specific primers, and the second

PCR ligated the DIP barcodes and adaptors for Illumina sequencing. For bacteria, we combined

1 μl of 1:10 diluted DNA, 10.5 μL of Ultra-Pure Sterile Molecular Biology Grade (Genesee

Scientific, San Diego, CA, USA), 12.5 µL of AccuStart ToughMix (2x concentration; Quantabio,

Beverly, MA, USA), and 0.5 µL each of the 10 µM 515F and 806R primers. Thermocycler

conditions for PCR1 were: 94°C for 2 min followed by 29 cycles of 94°C for 30 s, 55°C for 30 s,

68°C for 1 min followed by a 2 min extension at 68°C. For fungi, we combined 5 µl of

undiluted fungal DNA, 6.5 µl of ultra-pure water, 12.5 µl of AccuStart ToughMix, and 0.5 µL

each of the 10 µM ITS4-fun and 5.8s primers. Thermocycler conditions for PCR1 were: 94 °C

for 2 min, followed by 30 cycles of 94 °C for 30 s, 55 °C for 30 s, 68 °C for 2 min followed by a

10 min extension at 68°C. PCR1 products were cleaned with AMPure XP magnetic beads

(Beckman Coulter Inc., Brea, CA), following the manufacturer's protocols. The DIP PCR2

primers containing the barcodes and adaptors for Illumina sequencing were ligated to the

amplicons during the second PCR step in a 25 µL reaction containing 2.5 µL of the 10 µM DIP

PCR2 primers, 6.5 µL of ultra-pure water, 12.5 µL of Accustart ToughMix, and 1 µL of PCR1

product. Thermocycler conditions for PCR2 for bacteria and fungi were: 94°C for 2 min

followed by 10 cycles of 94°C for 30 s, 60°C for 30 s, and 72°C for 1 min. Bacterial and fungal

PCR products for each of the 324 samples (9 plots x 4 sub-plots x 9 time-points) were then

separately pooled based on gel electrophoresis band strength and cleaned with AMPure

following established methods (Glassman et al., 2018). Each pool was then checked for quality

and quantity with the Agilent Bioanalyzer 2100, pooled at a 2:3 ratio (0.4 units for bacteria to 0.6

units for fungi), and sequenced with Illumina MiSeq 2x300bp at the UCR Institute for

Integrative Genome Biology. In addition to our 324 samples sequenced in four Illumina MiSeq

runs, we added negative DNA extractions and PCR controls from each time point and mock

communities (ZymoBIOMICS microbial community standard, Zymo, Irvine, CA) to each library

for additional quality control and inference.

2.4 Bacterial and fungal biomass

Bacterial and fungal biomass was quantified with quantitative (q) PCR. We used the

FungiQuant-F/FungiQuant-R primers to target the 18s rDNA for fungi (Liu et al., 2012) and the

Eub338/Eub518 primers for bacteria (Fierer et al., 2005). To estimate bacterial and fungal small-

subunit rDNA gene abundances, we generated standard curves using a 10-fold serial dilution of

the standards by cloning the 18S region of *Saccharomyces cerevisiae* or the 16S region of

Escherichia coli into puc57 plasmid vectors constructed by GENEWIZ, Inc. (NJ, USA) as

previously established (Averill & Hawkes, 2016). The 10 μl qPCR reactions were performed in

triplicate. Reactions contained 1 μL undiluted DNA, 1 μl of 0.05M Tris-HCl ph8.3, 1 μL of

2.5mM $MgCl_2$ (New England BioLabs (NEB); Ipswich, MA), 0.5 μL of 0.5mg/ml BSA, 0.5 μL

of 0.25mM dNTP (NEB), 0.4 μL of both primers at 0.4μM, 0.5 μL of 20X Evagreen Dye (VWR

International; Radnor, PA), 0.1 μL of Taq DNA polymerase (NEB) and 4.6 μL ultra-pure water.

We employed the CFX384 Touch Real-Time PCR Detection System with the following

conditions: 94°C for 5 min, followed by 40 cycles of 94°C for 20 seconds, 52°C (for bacteria) or

50°C (for fungi) for 30 seconds, followed by an extension at 72°C for 30 seconds. Gene copy

numbers were generated using a regression equation using the quantification cycle (Cq),

calculated as the average Cq value per sample in relation to the known/calculated copies in the

Cq/threshold cycles (Ct). Thus, the gene copy number was determined using the following

equation, $10^{(Cq-b)/m}$, where the y-intercept and slope were generated from the standard curve.

All values were generated by CFX Maestro software.

2.5 Bioinformatics

Illumina data was processed with Qiime2 (Bolyen et al., 2019). The demultiplexed fastQ

files from the four Illumina sequencing runs were processed individually using cutadapt (Martin,

2011) to remove the primers and ran through DADA2 to filter out and remove chimeric

sequences and low-quality regions and to produce Amplicon Sequence Variants (ASVs)

(Callahan et al., 2017). DADA2 outputs from each library were merged into one library for

downstream processing, including the removal of singletons and taxonomic assignments. We

used the SILVA reference database for bacteria (Yilmaz et al., 2014) and the UNITE reference

database for fungi (Kõljalg et al., 2005) for taxonomic assignment (accessed on 10/27/2020).

Sequences assigned to mitochondria and chloroplasts for bacteria or not assigned to the Kingdom

Fungi were removed from ASV tables before subsequent analysis. Fungal ASV tables were

exported and parsed through FUNGuild (Nguyen et al., 2016) to assign functional ecological

guilds. We only included high probable confidence rankings for the following guilds for

downstream analysis: AMF, EMF, saprotrophs, and pathogens. Moreover, negative and mock

controls were inspected to ensure that sequences in negative controls were negligible and that

mock taxonomy correlated with known communities. Sequences were submitted to the National

Center for Biotechnology Information Sequence Read Archive under BioProject accession

number PRJNA761539.

2.6 Statistical analysis

Statistical analyses were conducted in R 4.1.1 (R Core Team, 2021), and all statistical

codes are available on GitHub https://github.com/pulidofabs/SecondarySuccession-Chaparral.

The ASV tables were rarefied to a sequences/sample depth of 7,115 for bacteria and 11,480 for

fungi to account for uneven sequencing depth. Fungal and bacterial alpha diversity was estimated

using BiodiversityR (Kindt & Coe, 2005) with the following metrics: observed species richness,

Simpson, Shannon, Chao1, and ACE. Since patterns of species richness were similar across

metrics **(Fig. S1)**, we focused on richness estimated as the number of observed ASVs after

rarefaction for all downstream analyses. To test if wildfire decreased biomass and richness and if

biomass and richness increased with time since fire **(H1)**, if wildfire will have a distinct impact

on fungal guilds **(H2)**, and to determine if changes in biomass and richness are associated with

precipitation and soil burn severity **(H3)**, for both bacteria and fungi, we fitted thirteen statistical

models with treatment (burned vs. unburned), time (measured as days since fire), ash depth (cm),

total monthly precipitation (mm), and their respective interactions as predictors. We used

generalized mixed effect models (glmer) with a negative binomial distribution in the MASS

package (Venables & Ripley, 2002) to account for the over-dispersion of the data and the fact

that the conditional variance was higher than the conditional mean (Bliss, 1953; Ross & Preece,

1985). Time since fire and precipitation were scaled and centered. The level of nestedness for all

models was tested by running a null model with different nested levels (plot, subplot, and time

since fire) and no predictors. Model selection was made using Akaike Information Criterion

(AICc) in the MuMIN package (Barton, 2020). All richness models contained plot, subplot, and

time since fire as random effects, and all biomass models included plot and time since fire as

random effects. Pseudo R^2, or the variance explained (marginal and conditional) for all models,

was calculated using the r.squaredGLMM function in the MuMIn Package.

We compared beta diversity by generating distance matrices with the vegan Avgdist

function, calculating the square-root transformed median Bray-Curtis dissimilarity per 100

subsampled iterations. We used permutational multivariate analysis of variance (PERMANOVA)

(Anderson, 2017) as implemented with the adonis function in vegan (Oksanen et al., 2018) to test

for the significant effects of wildfire, time since fire, and their interactions, on bacterial and

fungal community composition overall and at each independent time point (**H1**). Results were

visualized using Non-Metric Multidimensional Scaling (NMDS) ordinations.

We employed several methods to quantify succession (**H4**). We visualized succession by

characterizing community composition patterns at the genus level using phyloseq (McMurdie &

Holmes, 2013) and plotting the relative abundance profile of the dominant ASVs (> 3%

sequence abundance) for each time point in burned versus unburned communities with ggplot2

(Wickham H, 2016). We tested for significant changes in abundance of each taxon between

burned and unburned plots and over time using glmer. Similar to a study examining Sorghum

microbial succession (Gao et al., 2020), we used the vegan mantel function to determine the

correlation between temporal distance and community composition to determine how turnover

varies in the presence or absence of fire. To visualize spatial species turnover (Baselga, 2010),

we took advantage of the fact that early successional periods display large variability (Collins,

1990; Pandolfi, 2008). Thus, we tested the homogeneity of the bacterial and fungal communities

during succession using the vegan betadisper function to calculate the multivariate dispersion of

Bray-Curtis dissimilarities and visualized the results using principal coordinates analysis

(PCoA). Finally, we employed the codyn package (Hallett et al., 2016) to identify various

diversity metrics to assess succession (Collins et al., 2000). To evaluate whether the community

is following its normal successional trajectory, we calculated direct linear change, also known as

directional change, using the Euclidian distance of each microbial community. Next, we

measured species turnover to determine if species appearances or disappearances drive

succession. Lastly, we measured community stability as a measure of resilience, which

incorporates synchrony, or patterns of how species abundance covary over time (Loreau & de

Mazancourt, 2008).

3. Results

3.1 Sequencing data

The four Illumina MiSeq runs resulted in 9.8 M bacterial and 24.6 M fungal sequences

for an average of 31,052 bacterial and 78,202 fungal sequences/sample and a total of 33,078

bacterial and 11,480 fungal ASVs. Rarefaction resulted in a total of 24,874 bacterial and 7,445

fungal ASVs. Extracting the fungal guilds from the rarefied fungal ASV table resulted in 208

EMF, 70 saprobic, 65 AMF, and 26 pathogenic fungal ASVs.

3.2 Wildfire effects on bacterial and fungal biomass and richness

Fire led to significant reductions in bacterial and fungal biomass and richness during the

first post-fire year (**Tables S2; S3**). Fire had more significant impacts on biomass than richness

for both bacteria and fungi and larger effects on fungi than bacteria overall (**Fig. 2**). On average,

across the year, fire reduced richness by 46% for bacteria and 68% for fungi (**Fig. S1A**) and

biomass by 47% for bacteria (**Fig. S2A**) and 86% for fungi (**Fig. S2B**). The direct, immediate

effect of wildfire at 17 days post-fire was stronger for biomass than richness, with bacterial

biomass decreasing by 84% (**Fig. 2A**) and fungal biomass by 97% (**Fig. 2B**). In contrast, fungal

richness declined by 45% (**Fig. 2D**), but bacterial richness temporarily increased by 31% at 17

days but declined by 29% at 25 days post-fire (**Fig. 2C; Table S2**). While the differences in

biomass and richness between the burned and unburned sites lessened with time since fire, one

year was insufficient for bacterial and fungal biomass or richness to recover to unburned levels

(**Fig. 2**). The effects of fire across time remained larger for fungi than for bacteria, with fungal

biomass remaining 80% and richness 61% lower and bacterial biomass remaining 43% and

richness 23% lower in the burned plots at 1-year post-fire (**Fig. 2, Table S2**).

3.3 Drivers of changes in richness and biomass

Negative binomial models showed that the direct impact of fire negatively affected

bacterial and fungal biomass and richness, which was lessened by the positive interaction effect

of soil burn severity with time since fire (**Fig. 3; Table S3**). Furthermore, fire effects on fungal

richness were further attenuated by a direct and positive impact of time since fire **(Table S3)**.

Soil burn severity had significant adverse direct effects on bacterial and fungal richness but no

significant direct effects on biomass, meaning that biomass was equally reduced in the presence

of fire regardless of severity **(Figs. 3C,3D; Table S3)**. While precipitation had no significant

direct effects, precipitation interacted with time since fire to negatively affect bacterial richness

(Fig. 2C) but positively affect fungal richness **(Fig. 2D; Table S3)**. Moreover, bacterial biomass

responded positively to a fire and precipitation interaction (**Fig. 2A; Table S3**).

3.4 Wildfire effects on the richness of different fungal guilds

Wildfire led to large and significant reductions for all fungal guilds (**Table S4, Fig. S3**),

with the largest initial declines for AMF. On average, across time points, EMF decreased by

68%, pathogens by 71%, saprobes by 86%, and AMF by 98% in burned plots compared to

unburned plots (**Fig. S3**). Moreover, one year was insufficient for any guild to return to unburned

levels. In fact, at 376 days, AMF was on average 89%, pathogens 69%, and saprobes 61% lower

in the burned plots (**Fig. S4**). Interestingly, EMF experienced a decrease in richness over time

due to a fire and time interaction, resulting in species richness being 91% lower in the burned

plots at 376 days post-fire (**Fig. S4; Table S4**). EMF richness further declined with soil burn

severity (**Table S4**). Moreover, a fire and precipitation interaction negatively affected EMF and

pathogen richness **(Table S4).**

3.5 Wildfire changes microbial community composition

Fire significantly affected bacterial and fungal community structure, explaining 13% of

the compositional variation for bacteria and 11% for fungi. Time since fire had smaller impacts

on community composition, explaining 4% of the variation for bacteria and 1% for fungi.

Finally, the fire and time interaction explained 2% variation for bacteria and 1% for fungi (**Table**

S5). Community composition between burned and unburned plots significantly varied at all 9-

time points for both bacteria (**Fig. S5**) and fungi (**Fig. S6**), with differences increasing over time

from 12% to 21% for bacteria (**Fig. S5**) and 9% to 13% for fungi (**Fig. S6**) from day 17 to 376

post-fire. The changes in burned bacterial (**Fig. 4A**) and fungal composition (**Fig. 4C**) were

driven by the dominance of previously rare taxa, which traded off in abundance over time (**Figs.**

4B, 4D). Burned bacterial communities experienced less discernable tradeoffs than fungi due to

the consistent dominance of the Proteobacteria *Massilia*, which composed on average 57% of the

sequences throughout the year (**Fig. 4B**). However, three Firmicutes quickly dominated, with

Bacillus (13%), an uncultured Clostridiales (26%), and *Paenibacillus* (4%) dominating at 34-

and 67-days post-fire. The Firmicutes rapidly declined and traded off with the Proteobacteria

Noviherbaspirillum, which increased in abundance over time, from 1% at 34 days to 24% by the

end of the year (**Fig. 4B**). For burned fungi, two Basidiomycetes that dominated at 17 days post-

fire, the yeast *Geminibasidium* (45%) and the EMF *Inocybe* (16%), rapidly declined and traded

off with filamentous Ascomycetes **(Fig. 4D)**, including *Pyronema*, which increased as

Geminibasidium decreased. Indeed, *Pyronema* dominated from days 25 to 95 post-fire, peaking

at 67 days post-fire with 67% sequence abundance. In addition, the Ascomycetes *Aspergillus*

increased from 2% to 22% and *Penicillium* from 36% to 49% from 17 to 376 days post-fire **(Fig.**

4D).

3.6 Microbial successional dynamics

Directional change was higher in burned than unburned bacterial and fungal

communities, suggesting that succession is occurring in burned plots **(Table S6)**. Moreover,

bacterial and fungal burned communities experienced low synchrony **(Table S6),** indicating

community stability due to the dominance of a few genera. Indeed, community stability was

higher in burned than unburned plots and for bacteria than fungi **(Tables S6)**, suggesting that

bacterial succession is proceeding to potential equilibrium due to increasing bacterial richness

and resilience. In contrast, fungal communities were more susceptible to change. Visual

confirmation of these trends established that burned bacterial **(Fig. 5A)** and fungal **(Fig. 5B)**

communities experienced rapid and distinct successional trajectories, with bacteria **(Fig. 5C)**

experiencing successional changes at roughly double the rate of fungi **(Fig. 5D)**. These

successional changes in burned plots were driven by six major compositional turnover points for

bacteria at 25, 34, 95, 131, 187, and 376 days **(Fig. 5E)** and five for fungi at 25, 34, 67, 95, and

131 days post-fire **(Fig. 5F)**. In contrast, the unburned bacterial **(Fig. 6A)** and fungal **(Fig. 6B)**

communities experienced low dominance and remained stable over time with no succession

(Figs. 6C, 6D) or compositional turnover **(Figs. 6E, 6F).**

3.7 Taxa driving microbial succession

Early turnover events for bacteria **(Fig. 5A)** and fungi **(Fig. 5B)** in burned plots were

driven by the disappearance of taxa, more so for bacteria (days 17-95) than fungi (days 17-25;

Table S6). Indeed, increased dominance by a few taxa, including the Firmicutes *Bacillus* at 25

days post-fire, *Paenibacillus* and *Domibacillus* (34-67 days), and the Proteobacteria

Noviherbaspirillum (95 days), and the constant dominance of *Massilia* (34-286 days post-fire)

resulted in the disappearance of early surviving bacterial species, including *RB41*, *Conexibacter*,

Candidatus Udaeobacter, and *Bacillus* over time **(Fig. 5A; Table S6)**. In contrast, fungal

succession was initiated by the rapid dominance of *Geminibasidium* and EMF genera that

dominated the unburned communities, including *Inocybe*, *Cortinarius*, and *Tomentella* in the

Basidiomycota and *Balsamia* in the Ascomycota and one previously rare Basidiomycete EMF

genus *Mallocybe*, a subgenus of *Inocybe* **(Fig. 5B, 6B)**. Interestingly, all species of EMF rapidly

disappeared from the community at 25 days post-fire and were replaced by the constant

dominance of *Pyronema*, *Penicillium*, and *Aspergillus* **(Fig. 5B)**. However, as succession ensued,

species appearance drove late-year successional stages for bacteria (187-376 days; **Fig. 5A**) and

mid-year successional stages for fungi (67-187 days; **Fig. 5B; Table S6**). For bacteria, the

appearance of two Bacteroidetes, *Pedobacter* and *Adhaeribacter*, and the Actinobacteria

Blastococcus shaped late-year (days 187-376) bacterial succession **(Fig. 5A)**. For fungi, mid-year

succession (days 95-187) was driven by the appearance of filamentous fungal Ascomycetes in

the Pyronemataceae and in the Sordariaceae genus *Gelasinospora*, and Basidiomycota

mushroom-forming taxa. For example, the previously dominant EMF genus *Inocybe* returned,

and the pyrophilous Basidiomycete saprobes *Coprinellus* and *Tephrocybe*, which were rare in the

unburned communities (<0.01% sequence abundance), also appeared. Late-year fungal

succession (days 286-376) was driven by the increased dominance of *Tephrocybe* and

Aspergillus, the disappearance of *Inocybe* and the Ascomycota *Rasamsonia,* and the decreased

abundance of *Pyronema* **(Fig. 5B)**.

4. Discussion

Here, we present the highest resolution temporal sampling of post-fire microbiomes to

date, enabling us to show for the first time that bacteria and fungi experience rapid succession

driven by the dominance of pyrophilous taxa that trade off in abundance over time **(H4)**.

Although our study was in California chaparral, many of the dominant pyrophilous taxa

identified here also dominate other burned environments, including Spanish shrublands (Pérez-

Valera et al., 2018) and pine (Bruns et al., 2020; Reazin et al., 2016), spruce (Whitman et al.,

2019), Eucalyptus (Ammitzboll et al., 2022; McMullan-Fisher et al., 2011), and redwood-tanoak

forests (Enright et al., 2022), indicating the generality of our results to wildfire-affected

ecosystems. We found that wildfire decreased bacterial and fungal biomass and richness, leading

to community composition shifts that persisted the entire year (**H1**). Fungal guilds were

differentially affected by fire and time, with AMF and saprobes experiencing the largest

immediate fire impacts, but over time, EMF experienced the largest declines (**H2**). Interestingly,

AMF rapidly succumbed to the death of their host while EMF survived for 1-2 months,

answering the question of how long it takes mycorrhizal fungi to die after their host. Moreover,

microbial richness and biomass changes were driven by multiple abiotic interactions, including

interactions between time since fire, precipitation, and soil burn severity (**H3**).

4.1 Wildfire decreased bacterial and fungal biomass and richness

Fire decreased soil bacterial and fungal biomass and richness, corroborating previous

post-fire studies in Mediterranean shrublands (Pérez-Valera et al., 2018) and forests (Dooley &

Treseder, 2012). We noted a larger fire effect on fungal biomass and richness relative to bacteria,

consistent with previous research showing that bacteria are more resistant to fire than fungi

(Certini et al., 2021; Glassman et al., 2021; Pourreza et al., 2014; Pressler et al., 2019). Although

richness and biomass increased over time for both microbial groups, one year was insufficient

for either group to recover to unburned levels, consistent with previous studies that indicate that

microbial biomass and richness recovery could take decades (Pérez-Valera et al., 2018; Treseder

et al., 2004). Interestingly, we observed a transient increase in bacterial richness directly post-

fire, which may result from taxa such as the Actinobacteria genera *Soliribrobacter* and

Conexibacter (Albuquerque & da Costa, 2014) that may be favored by the increase in soil pH

observed here **(Table S1)** and in other studies (Neary et al., 1999) or by post-fire increases in

nitrogen and phosphorus availability (Certini et al., 2021). While several species of EMF

survived for the first months, fire-induced mortality was rapid for AMF and saprobes. Yet, EMF

was most impacted by fire over time, corroborating studies across ecosystems that point to the

negative impact of fire on mycorrhizal richness, primarily driven by host mortality (Dove &

Hart, 2017; Pulido-Chavez et al., 2021).

4.2 Pyrophilous fungi dominate the burned communities

Previously detected pyrophilous fungi from globally burned ecosystems also dominated

California chaparral, indicating that pyrophilous microbes are not biome-specific. Instead, these

fungi may be activated by temperature thresholds reached during high severity fires, which are

common in chaparral and some Pinaceae forests (Agee, 1993; Keeley & Zedler, 2009; Neary et

al., 1999). Specifically, our burned communities were dominated by Ascomycota in the genera

Pyronema, Penicillium, and *Aspergillus,* similar to dominant fungal species post-fire in

Mediterranean shrublands (Livne-Luzon et al., 2021), boreal spruce (Whitman et al., 2019), and

montane pine forests (Bruns et al., 2020; Fujimura et al., 2005; Pulido-Chavez et al., 2021). Both

Pyronema (P. omphalodes) and *Aspergillus (A. fumigatus*) are known pyrophilous fungi of

California chaparral (Dunn et al., 1982), and we identified additional species including *P.*

domesticum, *A. udagawae*, and *A. elsenburgensis*. These fungi are adapted to wildfire and

produce fire-resistant structures, including dormant spores, sclerotia, and conidia, which are

heat-activated (Gottlieb, 1950; Moore, 1962; Rhodes, 2006; Warcup & Baker, 1963). The rapid

and efficient germination of *Aspergillus* (in the subgenus *Fumigati*, including *A. udagawae*)

induced by high temperatures (Rhodes, 2006) and its ability to use various carbon and nitrogen

sources, including ammonium and nitrate (Krappmann & Braus, 2005), may position *Aspergillus*

to rapidly dominate post- fire. Furthermore, *Pyronema domesticum*, a relatively well-studied

pyrophilous fungus, can mineralize pyrogenic organic matter, PyOM (Fischer et al., 2021),

allowing it to take advantage of an abundant post-fire substrate. Moreover, *Geminibasidium*, a

recently described thermotolerant Basidiomycete yeast (Nguyen et al., 2013), dominated at 17

days post-fire. While not typically described as pyrophilous, presumably because most

pyrophilous fungi are described from mushrooms (McMullan-Fisher et al., 2011), two other

studies have found *Geminibasidium* to increase post-fire in pine forests (Pulido-Chavez et al.,

2021; Yang et al., 2020), suggesting that it is pyrophilous. Together, these results indicate that

dominant pyrophilous fungi are more widespread than expected and that their response to fire is

likely due to fire adaptive traits and temperature thresholds.

4.3 Pyrophilous fungi drive fungal secondary succession

Although it is clear that pyrophilous fungi exist and are widespread, we lacked an

understanding of how soon these pyrophilous fungi appear, their turnover rates, and if they trade

off in abundance over time (Fox et al., 2022). Here, we show that surviving ectomycorrhizal

Basidiomycetes *Cortinarius, Inocybe,* and *Tomentella,* previously found in burned temperate

pine forests (Owen et al., 2019; Pulido-Chavez et al., 2021) and pine-dominated Mediterranean

systems (Gassibe et al., 2011; Hernández-Rodríguez et al., 2013) also dominate early chaparral

fungal succession. Although it is possible that the EMF signal detected could be relic DNA

(Carini et al., 2016), high UV conditions, common in post-fire systems (Neary et al., 1999), are

likely to have degraded relic DNA (Torti et al., 2015), suggesting that these EMF could have

survived the fire and clung to a dying host for at least two months. The survival of these EMF

may be attributable to their exploration type and specificity for current versus stored

photosynthates (Gray & Kernaghan, 2020; Pena et al., 2010). For example, *Cortinarius,* a

long/medium exploration type, which requires more carbon for maintenance, only survived for

17 days post-fire. In contrast, *Inocybe,* a short-contact type (Agerer, 2001; Koide et al., 2014),

survived for up to 67 days post-fire, suggesting that it may be less carbon demanding or adapted

to warming (Fernandez et al., 2017). Our results indicate that early EMF could have survived on

their hosts' stored photosynthates or the lower resources provided by the surviving and

resprouting manzanita and chamise shrubs.

Successional theory states that early successional stages are dominated by fast-growing

or ruderal (R) organisms (Kinzig & Pacala, 2013) that tradeoff between stress tolerance (S) and

competitive (C) life-history strategies over time (Grime, 1977; Zhang et al., 2018). Recent

adaptations of Grime's C-S-R to microbiomes suggest that pyrophilous microbes survive and

thrive post-fire with traits analogous to plants, including post-fire resource acquisition (C),

thermotolerant structures (S), and fast growth (R) (Enright et al., 2022; Whitman et al., 2019).

Our data suggest that pyrophilous microbes fit into these trait categories and that tradeoffs

among these traits might drive microbial succession, such that surviving thermotolerant fungi

(i.e., *Geminibasidium* and *Pyronema*) colonize first, followed by fast-colonizers (i.e., *Penicillium*

and *Aspergillus*), which were overtaken by competitive fungi capable of exploiting post-fire

resources. (i.e., *Coprinellus* and *Tephrocybe)*. Whereas *Geminibasidium* is thermo- and xero-

tolerant (Nguyen et al., 2013), *Pyronema* produces thermotolerant sclerotia (Moore, 1962),

enabling wildfire survival. Although both species are thermotolerant, differences in the

morphological growth characteristics between the yeast *Geminibasidium* and the filamentous

Pyronema may explain the tradeoff in dominance. For example, unicellular proliferation allowed

Geminibasidium to dominate instantly, but the ability to forage for nutrients and rapidly increase

surface colonization may allow the filamentous *Pyronema* to better dominate the open space.

Interestingly, *Gelasinospora heterospora*, a fungus closely related to heat-activated *Neurospora*

crassa (Dettman et al., 2001; Emerson, 1948), that produces pigmented mycelia, and is known to

require heat or chemical treatment for germination (Alexopoulous & Mims, 1952), was highly

abundant at 25, 34, and 95 days post-fire, suggesting it is also pyrophilous. The decline of

thermotolerators coincided with the dominance of fast-growing Ascomycete fungi in the genera

Penicillium and *Aspergillus* (Dix & Webster, 1995; McGee et al., 2006), which both produce

copious asexual conidia (Crow, 1992), allowing for rapid colonization of the open niche,

analogous to early dominant post-fire plants, which are also often asexual (James, 1984). Lastly,

like plants, competition appears to drive later successional dynamics (Tilman, 1990; Zhang et al.,

2018), as suggested by the emergence of pyrophilous fungal decomposers, such as *Coprinellus*

and *Tephrocybe* at 9-12 months, corroborating findings from Eucalyptus forests (Ammitzboll et

al., 2022; McMullan-Fisher et al., 2011). Evidence suggests that *Coprinellus* can degrade

aromatic hydrocarbons (Steindorff et al., 2021), supporting the idea that it is outcompeting

earlier fungal taxa for this abundant and complex carbon source. In conclusion, our results

indicate that tradeoffs among fire-adapted traits analogous to those in plants might drive post-fire

microbial succession.

4.4 Pyrophilous bacteria dominate and initiate bacterial secondary succession

Our high temporal sampling enabled us to demonstrate that bacteria experience rapid

succession and community turnover initiated by aerobic, heterotrophic bacteria that form

endospores and produce antibiotics, which may improve their competitive abilities in the post-

fire environment. Moreover, we show that previously detected putative pyrophilous bacteria

from Canadian (Whitman et al., 2019) and Chinese boreal (Xiang et al., 2014), and California

redwood-tanoak forests (Enright et al., 2022) also dominated California chaparral, suggesting

that these pyrophilous bacteria might be widely distributed. Specifically, our burned

communities favored taxa in the phyla Actinobacteria, Acidobacteria, Firmicutes, and

Proteobacteria, similar to previous studies (Ammitzboll et al., 2022; Enright et al., 2022;

Whitman et al., 2019; Xiang et al., 2014).

Bacterial successional dynamics were slightly different from fungi. The first few weeks

of bacterial succession were characterized by high diversity and a lack of dominance, mimicking

the distributions observed in the unburned communities. In complex resource environments, such

as those created by wildfire, niche complementarity is more important than competition because

diverse communities can better exploit resources (Eisenhauer et al., 2013). Our results support

this idea, as the dominant taxa in our study have been shown to use different resources. For

example, *Paenibacillu*s can fix nitrogen (Monciardini et al., 2003; Slepecky & Hemphill, 2006),

Bacillus and *Conexibacter* can reduce nitrogen, and *Bacillus* can further solubilize phosphate (J.

et al., 2001; Kalayu, 2019). However, at 25 days, bacteria experienced a large compositional

turnover, which allowed for the dominance of pyrophilous bacteria with fire-adapted traits.

While tradeoffs in abundance were not as stark for bacteria as for fungi, as indicated by the

constant dominance of the Proteobacteria *Massilia*, the next most abundant bacteria did

experience tradeoffs from thermotolerant Firmicutes (i.e., *Bacillus, Paenibacillus*, and

Clostridiales) to fast-colonizing Proteobacteria that are also capable of exploiting post-fire

substrates such *Novihersparillium* and *Massilia*. Thermotolerance and heat-stress adaptation

dominated survival traits, allowing Firmicutes to dominate as early as 25 days post-fire.

Firmicutes contain some of the most resistant, thermotolerant endospore producing bacteria

(Grady et al., 2016; Kaur et al., 2018; Mandic-Mulec et al., 2015), as well as the species

Paenibacillus xerothermodurans, which contains heat-inducible transcriptional proteins that

protect against protein denaturation (Paredes-Sabja et al., 2011). Moreover, Clostridiales can

trigger sporulation in response to stress (Paredes-Sabja et al., 2011), allowing for increased

proliferation post-fire. While thermotolerance is crucial for fire survival, rapidly dominating the

open niche and using post-fire resources may ensure post-fire growth. *Massilia* is a fast

reproducing bacteria that associates with the rhizosphere (Li et al., 2014) and colonizes AMF

hyphae (Iffis et al., 2014) and the roots of various plants (Ofek et al., 2012). Thus, the dominance

of *Massilia* in our study and other post-fire environments (Enright et al., 2022; Whitman et al.,

2019) suggests that its diverse ecology and rapid reproduction rate promote its proliferation

within the soil (Toljander et al., 2005). Moreover, our results, coupled with current knowledge in

the field, show that the continued dominance of *Massilia* and *Noviherbaspirillium* might be due

to their ability to exploit post-fire resources. For example, *Bacillus* fix and reduce nitrogen

(Bailey et al., 2014), while *Noviherbaspirillium,* a mid-to late-year dominant bacterial genus,

may degrade polycyclic aromatic hydrocarbon (Baldani et al., 2014; Grady et al., 2016; Woolet

& Whitman, 2020). We conclude that although tradeoffs are not as distinctive for bacteria as

they are fungi, potentially due to interaction among traits such as fast growth under high

resources and resource acquisition under low resources, our results provide evidence that

pyrophilous bacteria have mechanisms to survive the fire that may drive them to tradeoff in

abundance over time.

4.5 Precipitation and soil burn severity differentially affected bacteria and fungi

Although fire was the most significant driver of microbial change, we found that

interactions between time and precipitation also drove successional dynamics by differentially

affecting microbial biomass and richness. Precipitation interacted with time since the fire and

negatively affected bacterial but positively affected fungal richness, with bacterial richness

declining but fungal richness increasing after rain events. At least one study that performed a

three-year precipitation manipulation experiment showed that increased precipitation decreased

bacterial richness (Yang et al., 2021). In contrast to richness, precipitation interacted with fire to

positively affect bacterial biomass, while fungal biomass remained unaffected by rainfall. The

distinct responses to precipitation could be due to microbial physiological and morphological

differences (Blazewicz et al., 2014; Placella et al., 2012; Selbmann et al., 2013). For example,

fungi have high morphological plasticity, including the ability to alternate growth forms based on

environmental conditions, such as hyphal growth in low nutrient conditions versus unicellular

form in rich nutrient conditions (Selbmann et al., 2013), which may allow fungi to survive small

changes in moisture content, allowing for continued growth over time. Moreover, fungi can also

activate or enhance resistance to drought (Evans & Wallenstein, 2012; Guhr & Kircher, 2020),

allowing them to maintain a constant reproduction rate regardless of the external environmental

conditions, thus explaining the positive impact on fungal richness. In contrast, bacteria can

experience cell death due to rapid wet-up, while other bacterial groups enter a dormant state,

inhibiting growth (Schimel, 2018). Thus, to be evolutionarily successful, bacteria may respond

rapidly to favorable conditions (e.g., rain) (Leizeaga et al., 2022) and rapidly replicate to

outcompete other microbes for the pulse of nutrients released during the rain event (Homyak et

al., 2014), thus potentially explaining the peaks in richness observed after significant rainfall at

day 131.

Here, we measured ash depth before the first rains could wash away ash from the site,

showing that ash depth can serve as an index of soil burn severity on a scale relevant to soil

microbes and that severity negatively affected bacterial and fungal richness but not biomass.

Similar to studies in boreal (Whitman et al., 2019) and oak forests (Pourreza et al., 2014),

bacterial and fungal richness decreased with soil burn severity. However, contrary to a meta-

analysis (Dooley & Treseder, 2012), neither bacterial nor fungal biomass was directly impacted

by soil burn severity, probably due to microbial adaptations to the natural high severity fire

regime of chaparral. However, since we observed an interaction effect of time since fire and

severity on microbial biomass, it is possible that previously observed severity effects, derived

from chronosequence studies where space is substituted for time and which, on average, occur

two years post-fire (Pressler et al., 2019), may represent an interaction effect of time and severity

and not a direct effect of soil burn severity.

5 Conclusion

Our unprecedented high-resolution temporal sampling allowed us to detect rapid

secondary succession in bacterial and fungal communities and marked tradeoffs in abundance

among pyrophilous microbes over time. We found that the same pyrophilous bacteria and fungi

that respond to fires in temperate and boreal forests and Mediterranean shrublands also dominate

in California chaparral, allowing us to generalize knowledge of post-fire microbiomes to dryland

habitats that are rapidly increasing with climate change (Feng & Fu, 2013). While pyrophilous

microbes have been described in other systems, their turnover rates were unknown due to limited

post-fire sampling (Fox et al., 2022). We suggest that tradeoffs among fire-adapted traits, with

thermotolerators dominating first, followed by fast colonizers, and finally by competitors capable

of capitalizing on post-fire resource acquisition, drive post-fire microbial successional dynamics.

It seems that pyrophilous microbes share many analogous traits to plants, enabling us to adapt

successional theory developed in plants to microbes while acknowledging that bacteria and fungi

do have differences in how they respond to fire due to their physiology. We conclude that post-

fire bacteria and fungi experience rapid successional changes after fire and believe that

understanding these dynamics can increase our ability to predict if and how ecosystems will

recover after fire, an increasingly widespread disturbance with climate change.

Acknowledgments

We thank the Cleveland National Forest and the Trabuco Ranger District, including

District Ranger Darrel Vance and Emily Fudge, Jeffrey Heys, Lauren Quon, Jacob Rodriguez,

and Victoria Stempniewicz, for their guidance help with permitting and site selection. We thank

Judy A. Chung for help with fieldwork and molecular work and Aral C. Greene, Dylan Enright,

and Sameer S. Saroa for fieldwork assistance. We thank Amelia Nelson, Elizah Stephens,

Kaleigh A. Russell, and our anonymous reviewers for their helpful comments on the manuscript.

This work was supported by UC Riverside, the BLM JFSP Award #012641-002 and Shipley

Skinner Award to MFPC and SIG, and the USDA-NIFA Award 2022-67014-36675 to SIG and

PMH.

Conflict of interest

666 Authors do not declare any conflict of interest or relationship that might influence the author's

667 objectivity.

Author's contribution:

670 SIG conceived of the study and acquired permits; LL, SIG, and PMH came up with the

671 experimental design; SIG, LL, PMH, MFPC, and JWJR performed fieldwork; MFPC and JWJR

672 performed all molecular work; MFPC performed all bioinformatics, statistical analysis, made all

673 of the figures, and wrote the first draft; all authors contributed to writing and editing of the

674 manuscript.

Figure 1. A) Location of the study area and plots (6 burned; 3 unburned) within the Holy Fire burn scar in the Cleveland National Forest in Southern California. Soil burn severity is based on the USDA BAER classifications. B) Plot experimental design, each with four 1m² subplots placed 5m from the center in each cardinal direction. C) Collection of the top 10 cm of soil beneath the ash with a releasable bulb planter within the 1 m² subplots at 17 days post-fire.

Biomass

Richness

Figure 2. Change in biomass for A) bacteria and B) fungi and change in richness for C) bacteria and D) fungi at each of the 9 time-points in the first year post-fire in burned and unburned plots. Points represent mean differences averaged across burned and unburned plots, and bars represent the standard error of the mean. Precipitation is based on total monthly precipitation (mm). The significance of burned versus unburned plots per time point is denoted with an asterisk and is based on a negative binomial regression.

Figure 3. Change in A) bacterial and B) fungal biomass and C) bacterial and D) fungal richness in relation to soil burn severity measured as ash depth at 17 days post-fire. The significance of biomass and species richness against soil burn severity (SBS) and its interaction with time since (TSF) fire is based on a negative binomial regression, where the unburned plots are represented in blue-green and burned in brown.

Figure 4. NMDS plot of A) bacterial and C) fungal community composition with colors denoting treatment and shape indicating time since fire. The NMDS is based on 3-dimensions and has a stress value of 0.11 for bacteria and 0.12 for fungi. Change in relative sequence abundance of the five most abundant B) bacterial and D) fungal genera in the burned plots over time. Note, the dominance of the top 5 bacterial genera began 34 days post-fire since there was no clear dominance at days 17 and 25 due to the lag in post-fire bacterial declines.

Figure 5. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in burned plots at all 9 time points. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in burned plots. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each time point for the burned E) bacterial and F) fungal communities. Note that time points farther apart indicate a community turnover, whereas overlapping points with standard errors represent a lack of compositional turnover. Note, 17 days represents the base level and the first turnover thus initiated at 25 days.

Figure 6. Relative sequence abundance of A) bacterial and B) fungal genera (ectomycorrhizal fungi denoted with an asterisk) in unburned plots at all 9-time points. Mantel correlation between Bray-Curtis dissimilarity and Euclidean temporal distances in C) bacterial and D) fungal community composition in unburned plots displaying no significant succession. Principal components analysis (PCoA) of the mean and standard error Bray-Curtis community dissimilarity at each time point for E) bacteria and F) fungi in the unburned plots. Note that overlapping standard error bars indicate no community turnover across time in unburned plots.

Reference

Abatzoglou, J. T., Williams, A. P., & Barbero, R. (2019). Global Emergence of Anthropogenic Climate Change in Fire Weather Indices. *Geophysical Research Letters*, *46*(1), 326–336. https://doi.org/10.1029/2018GL080959

Agee, J. K. (1993). *Fire Ecology of Pacific Northwest Forests*. Island Press.

Agerer, R. (2001). Exploration types of ectomycorrhizae. *Mycorrhiza*, *11*(2), 107–114. https://doi.org/10.1007/s005720100108

Albuquerque, L., & da Costa, M. S. (2014). The Families Conexibacteraceae, Patulibacteraceae and Solirubrobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes: Actinobacteria* (pp. 185–200). Springer. https://doi.org/10.1007/978-3-642-30138-4_200

Alexopoulous, C. J., & Mims, C. W. (1952). *Introductory mycology* (3rd ed.). John Wilet & Sons.

Allen, M., Egerton-Warburton, L., Treseder, K., Cario, C. H., Lindahl, A., Lansing, J., Querejeta, J., Kårén, O., Harney, S., & Zink, T. (2005). *Biodiversity of Mycorrhizal Fungi in Southern California 1*. /paper/Biodiversity-of-Mycorrhizal-Fungi-in-Southern-1-Allen-Egerton-Warburton/445d4f6878927a5307669f5e73df3174703f4d2a

Ammitzboll, H., Jordan, G. J., Baker, S. C., Freeman, J., & Bissett, A. (2022). Contrasting successional responses of soil bacteria and fungi to post-logging burn severity. *Forest Ecology and Management*, *508*, 120059. https://doi.org/10.1016/j.foreco.2022.120059

Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In *Wiley StatsRef: Statistics Reference Online* (pp. 1–15). American Cancer Society. https://doi.org/10.1002/9781118445112.stat07841

Averill, C., & Hawkes, C. V. (2016). Ectomycorrhizal fungi slow soil carbon cycling. *Ecology Letters*, *19*(8), 937–947. https://doi.org/10.1111/ele.12631

Bailey, A. C., Kellom, M., Poret-Peterson, A. T., Noonan, K., Hartnett, H. E., & Raymond, J. (2014). Draft Genome Sequence of Massilia sp. Strain BSC265, Isolated from Biological Soil Crust of Moab, Utah. *Genome Announcements*, *2*(6), e01199-14, 2/6/e01199-14. https://doi.org/10.1128/genomeA.01199-14

Baldani, J. I., Rouws, L., Cruz, L. M., Olivares, F. L., Schmid, M., & Hartmann, A. (2014). The Family Oxalobacteraceae. In E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes* (pp. 919–974). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-30197-1_291

Barnard, R. L., Osborne, C. A., & Firestone, M. K. (2013). Responses of soil bacterial and fungal communities to extreme desiccation and rewetting. *The ISME Journal*, *7*(11), 2229–2241. https://doi.org/10.1038/ismej.2013.104

Barro, S. C., & Conard, S. G. (1991). Fire effects on California chaparral systems: An overview. *Environment International*, *17*(2–3), 135–149. https://doi.org/10.1016/0160-4120(91)90096-9

Barton, K. (2020). *MuMIn: Multi-Model Inference* (R package version 1.43.17.) [Computer software]. https://CRAN.R-project.org/package=MuMIn

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, *19*(1), 134–143. https://doi.org/10.1111/j.1466-8238.2009.00490.x

Blazewicz, S. J., Schwartz, E., & Firestone, M. K. (2014). Growth and death of bacteria and
fungi underlie rainfall-induced carbon dioxide pulses from seasonally dried soil. *Ecology*,
95(5), 1162–1172. https://doi.org/10.1890/13-1031.1

Bliss, C. I. (1953). *Fitting the negative binomial distribution to biological data (with "Note on*
the efficient fitting of the negative binomial" by R. A. Fisher), Biometrics, 9, (9th ed.).
Biometrics.

Bodí, M. B., Martin, D. A., Balfour, V. N., Santín, C., Doerr, S. H., Pereira, P., Cerdà, A., &
Mataix-Solera, J. (2014). Wildland fire ash: Production, composition and eco-hydro-
geomorphic effects. *Earth-Science Reviews*, *130*, 103–127.
https://doi.org/10.1016/j.earscirev.2013.12.007

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,
Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger,
K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A.
M., Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and
extensible microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8), 852–
857. https://doi.org/10.1038/s41587-019-0209-9

Bruns, T. D., Chung, J. A., Carver, A. A., & Glassman, S. I. (2020). A simple pyrocosm for
studying soil microbial response to fire reveals a rapid, massive response by Pyronema
species. *PLOS ONE*, *15*(3), e0222691. https://doi.org/10.1371/journal.pone.0222691

Cairney, J. W. G., & Bastias, B. A. (2007). Influences of fire on forest soil fungal
communitiesThis article is one of a selection of papers published in the Special Forum on
Towards Sustainable Forestry — The Living Soil: Soil Biodiversity and Ecosystem
Function. *Canadian Journal of Forest Research*, *37*(2), 207–215.
https://doi.org/10.1139/x06-190

Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *The ISME Journal*, *11*(12),
2639–2643. https://doi.org/10.1038/ismej.2017.119

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P.
J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of
millions of sequences per sample. *Proceedings of the National Academy of Sciences*,
108(Supplement_1), 4516–4522. https://doi.org/10.1073/pnas.1000080107

Carini, P., Marsden, P. J., Leff, J. W., Morgan, E. E., Strickland, M. S., & Fierer, N. (2016).
Relic DNA is abundant in soil and obscures estimates of soil microbial diversity. *Nature*
Microbiology, *2*(3), 1–6. https://doi.org/10.1038/nmicrobiol.2016.242

Certini, G., Moya, D., Lucas-Borja, M. E., & Mastrolonardo, G. (2021). The impact of fire on
soil-dwelling biota: A review. *Forest Ecology and Management*, *488*, 118989.
https://doi.org/10.1016/j.foreco.2021.118989

Collins, S. L. (1990). Patterns of Community Structure During Succession in Tallgrass Prairie.
Bulletin of the Torrey Botanical Club, *117*(4), 397–408. https://doi.org/10.2307/2996837

Collins, S. L., Micheli, F., & Hartt, L. (2000). A method to determine rates and patterns of
variability in ecological communities. *Oikos*, *91*(2), 285–293.
https://doi.org/10.1034/j.1600-0706.2000.910209.x

Cowan, A. D., Smith, J. E., & Fitzgerald, S. A. (2016). Recovering lost ground: Effects of soil
burn intensity on nutrients and ectomycorrhiza communities of ponderosa pine seedlings.
Forest Ecology and Management, *378*, 160–172.
https://doi.org/10.1016/j.foreco.2016.07.030

Crow, J. F. (1992). An advantage of sexual reproduction in a rapidly changing environment. *The*
Journal of Heredity, *83*(3), 169–173.
https://doi.org/10.1093/oxfordjournals.jhered.a111187
Dalling, J. W. (2008). Pioneer Species. In B. Fath (Ed.), *Encyclopedia of Ecology (Second*
Edition) (pp. 181–184). Elsevier. https://doi.org/10.1016/B978-0-444-63768-0.00534-5
Day, N. J., Cumming, S. G., Dunfield, K. E., Johnstone, J. F., Mack, M. C., Reid, K. A.,
Turetsky, M. R., Walker, X. J., & Baltzer, J. L. (2020). Identifying Functional Impacts of
Heat-Resistant Fungi on Boreal Forest Recovery After Wildfire. *Frontiers in Forests and*
Global Change, *3*. https://doi.org/10.3389/ffgc.2020.00068
Derroire, G., Balvanera, P., Castellanos-Castro, C., Decocq, G., Kennard, D. K., Lebrija-Trejos,
E., Leiva, J. A., Odén, P.-C., Powers, J. S., Rico-Gray, V., Tigabu, M., & Healey, J. R.
(2016). Resilience of tropical dry forests – a meta-analysis of changes in species diversity
and composition during secondary succession. *Oikos*, *125*(10), 1386–1397.
https://doi.org/10.1111/oik.03229
Dettman, J. R., Harbinski, F. M., & Taylor, J. W. (2001). Ascospore Morphology Is a Poor
Predictor of the Phylogenetic Relationships of Neurospora and Gelasinospora. *Fungal*
Genetics and Biology, *34*(1), 49–61. https://doi.org/10.1006/fgbi.2001.1289
Dix, N. J., & Webster, J. (1995). *Fungal Ecology*. Chapman & Hall.
Donato, D. C., Campbell, J. L., & Franklin, J. F. (2012). Multiple successional pathways and
precocity in forest development: Can some forests be born complex? *Journal of*
Vegetation Science, *23*(3), 576–584. https://doi.org/10.1111/j.1654-1103.2011.01362.x
Dooley, S. R., & Treseder, K. K. (2012). The effect of fire on microbial biomass: A meta-
analysis of field studies. *Biogeochemistry*, *109*(1/3), 49–61.
Dove, N. C., & Hart, S. C. (2017). Fire Reduces Fungal Species Richness and In Situ
Mycorrhizal Colonization: A Meta-Analysis. *Fire Ecology*, *13*(2), 37–65.
https://doi.org/10.4996/fireecology.130237746
Dove, N. C., Klingeman, D. M., Carrell, A. A., Cregger, M. A., & Schadt, C. W. (2021). Fire
alters plant microbiome assembly patterns: Integrating the plant and soil microbial
response to disturbance. *New Phytologist*, *230*(6), 2433–2446.
https://doi.org/10.1111/nph.17248
Dunn, P. H., Wells, W. G. I., Dickey, J., & Wohlgemuth, M. (1982). *Role of Fungi in Postfire*
Stabilization of Chaparral Ash Beds (PSW-58, pp. 378–381) [Gen. Tech. Rep]. Pacific
Southwest Forest and Range Experiment Station, Forest Service, U.S.U.S. Department of
Agriculture.
Eisenhauer, N., Schulz, W., Scheu, S., & Jousset, A. (2013). Niche dimensionality links
biodiversity and invasibility of microbial communities. *Functional Ecology*, *27*(1), 282–
288. https://doi.org/10.1111/j.1365-2435.2012.02060.x
Emerson, M. R. (1948). Chemical Activation of Ascospore Germination in Neurospora crassa1.
Journal of Bacteriology, *55*(3), 327–330.
Enright, D. J., Frangioso, K. M., Isobe, K., Rizzo, D. M., & Glassman, S. I. (2022). Mega-fire in
Redwood Tanoak Forest Reduces Bacterial and Fungal Richness and Selects for
Pyrophilous Taxa that are Phylogenetically Conserved. *Molecular Ecology*.
https://doi.org/10.1111/mec.16399
Evans, S. E., & Wallenstein, M. D. (2012). Soil microbial community response to drying and
rewetting stress: Does historical precipitation regime matter? *Biogeochemistry*, *109*(1/3),
101–116.

Feng, S., & Fu, Q. (2013). Expansion of global drylands under a warming climate. *Atmospheric*
Chemistry and Physics, *13*(19), 10081–10094. https://doi.org/10.5194/acp-13-10081-
2013
Fernandez, C. W., Nguyen, N. H., Stefanski, A., Han, Y., Hobbie, S. E., Montgomery, R. A.,
Reich, P. B., & Kennedy, P. G. (2017). Ectomycorrhizal fungal response to warming is
linked to poor host performance at the boreal-temperate ecotone. *Global Change Biology*,
23(4), 1598–1609. https://doi.org/10.1111/gcb.13510
Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of Soil Microbial
Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Applied and*
Environmental Microbiology, *71*(7), 4117–4120.
https://doi.org/10.1128/AEM.71.7.4117-4120.2005
Fischer, M. S., Stark, F. G., Berry, T. D., Zeba, N., Whitman, T., & Traxler, M. F. (2021).
Pyrolyzed Substrates Induce Aromatic Compound Metabolism in the Post-fire Fungus,
Pyronema domesticum. *Frontiers in Microbiology*, *12*, 729289.
https://doi.org/10.3389/fmicb.2021.729289
Fox, S., Sikes, B. A., Brown, S. P., Cripps, C. L., Glassman, S. I., Hughes, K., Semenova-
Nelsen, T., & Jumpponen, A. (2022). Fire as a driver of fungal diversity—A synthesis of
current knowledge. *Mycologia*, *0*(0), 1–27.
https://doi.org/10.1080/00275514.2021.2024422
Fujimura, K. E., Smith, J. E., Horton, T. R., Weber, N. S., & Spatafora, J. W. (2005). Pezizalean
mycorrhizas and sporocarps in ponderosa pine (Pinus ponderosa) after prescribed fires in
eastern Oregon, USA. *Mycorrhiza*, *15*(2), 79–86. https://doi.org/10.1007/s00572-004-
0303-8
Gao, C., Montoya, L., Xu, L., Madera, M., Hollingsworth, J., Purdom, E., Singan, V., Vogel, J.,
Hutmacher, R. B., Dahlberg, J. A., Coleman-Derr, D., Lemaux, P. G., & Taylor, J. W.
(2020). Fungal community assembly in drought-stressed sorghum shows stochasticity,
selection, and universal ecological dynamics. *Nature Communications*, *11*(1), 34.
https://doi.org/10.1038/s41467-019-13913-9
Gassibe, P. V., Fabero, R. F., Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto,
P. (2011). Fungal community succession following wildfire in a Mediterranean
vegetation type dominated by Pinus pinaster in Northwest Spain. *Forest Ecology and*
Management, *262*(4), 655–662. https://doi.org/10.1016/j.foreco.2011.04.036
Glassman, S. I., Levine, C. R., DiRocco, A. M., Battles, J. J., & Bruns, T. D. (2016).
Ectomycorrhizal fungal spore bank recovery after a severe forest fire: Some like it hot.
The ISME Journal, *10*(5), 1228–1239. https://doi.org/10.1038/ismej.2015.182
Glassman, S. I., Randolph, J. W., Saroa, S. S., Capocchi, J. K., Walters, K. E., Pulido-Chavez,
M. F., & Larios, L. (2021). *Prescribed versus wildfire impacts on exotic plants and soil*
microbes in California grasslands (p. 2021.09.22.461426).
https://doi.org/10.1101/2021.09.22.461426
Glassman, S. I., Weihe, C., Li, J., Albright, M. B. N., Looby, C. I., Martiny, A. C., Treseder, K.
K., Allison, S. D., & Martiny, J. B. H. (2018). Decomposition responses to climate
depend on microbial community composition. *Proceedings of the National Academy of*
Sciences, *115*(47), 11994–11999. https://doi.org/10.1073/pnas.1811269115
Gottlieb, D. (1950). The physiology of spore germination in fungi. *The Botanical Review*, *16*(5),
229–257. https://doi.org/10.1007/BF02873609

Grady, E. N., MacDonald, J., Liu, L., Richman, A., & Yuan, Z.-C. (2016). Current knowledge
and perspectives of Paenibacillus: A review. *Microbial Cell Factories*, *15*(1), 203.
https://doi.org/10.1186/s12934-016-0603-7

Gray, L., & Kernaghan, G. (2020). Fungal Succession During the Decomposition of
Ectomycorrhizal Fine Roots. *Microbial Ecology*, *79*(2), 271–284.
https://doi.org/10.1007/s00248-019-01418-3

Grime, J. P. (1977). Evidence for the Existence of Three Primary Strategies in Plants and Its
Relevance to Ecological and Evolutionary Theory. *The American Naturalist*, *111*(982),
1169–1194.

Guhr, A., & Kircher, S. (2020). Drought-Induced Stress Priming in Two Distinct Filamentous
Saprotrophic Fungi. *Microbial Ecology*, *80*(1), 27–33. https://doi.org/10.1007/s00248-
019-01481-w

Hallett, L. M., Jones, S. K., MacDonald, A. A. M., Jones, M. B., Flynn, D. F. B., Ripplinger, J.,
Slaughter, P., Gries, C., & Collins, S. L. (2016). codyn: An r package of community
dynamics metrics. *Methods in Ecology and Evolution*, *7*(10), 1146–1151.
https://doi.org/10.1111/2041-210X.12569

Hannula, S. E., Kielak, A. M., Steinauer, K., Huberty, M., Jongen, R., De Long, J. R., Heinen,
R., & Bezemer, T. M. (2019). Time after Time: Temporal Variation in the Effects of
Grass and Forb Species on Soil Bacterial and Fungal Communities. *MBio*, *10*(6), e02635-
19. https://doi.org/10.1128/mBio.02635-19

Hart, S. C., DeLuca, T. H., Newman, G. S., MacKenzie, M. D., & Boyle, S. I. (2005). Post-fire
vegetative dynamics as drivers of microbial community structure and function in forest
soils. *Forest Ecology and Management*, *220*(1–3), 166–184.
https://doi.org/10.1016/j.foreco.2005.08.012

Hernández-Rodríguez, M., Oria-de-Rueda, J. A., & Martín-Pinto, P. (2013). Post-fire fungal
succession in a Mediterranean ecosystem dominated by Cistus ladanifer L. *Forest
Ecology and Management*, *289*, 48–57. https://doi.org/10.1016/j.foreco.2012.10.009

Homyak, P. M., Sickman, J. O., Miller, A. E., Melack, J. M., Meixner, T., & Schimel, J. P.
(2014). Assessing Nitrogen-Saturation in a Seasonally Dry Chaparral Watershed:
Limitations of Traditional Indicators of N-Saturation. *Ecosystems*, *17*(7), 1286–1305.
https://doi.org/10.1007/s10021-014-9792-2

Hughes, K. W., Matheny, P. B., Miller, A. N., Petersen, R. H., Iturriaga, T. M., Johnson, K. D.,
Methven, A. S., Raudabaugh, D. B., Swenie, R. A., & Bruns, T. D. (2020). Pyrophilous
fungi detected after wildfires in the Great Smoky Mountains National Park expand
known species ranges and biodiversity estimates. *Mycologia*, *112*(4), 677–698.
https://doi.org/10.1080/00275514.2020.1740381

Iffis, B., St-Arnaud, M., & Hijri, M. (2014). Bacteria associated with arbuscular mycorrhizal
fungi within roots of plants growing in a soil highly contaminated with aliphatic and
aromatic petroleum hydrocarbons. *FEMS Microbiology Letters*, *358*(1), 44–54.
https://doi.org/10.1111/1574-6968.12533

J., E. de los M., A., M., & F., V. (2001). Metabolic profiles and aprE expression in anaerobic
cultures of Bacillus subtilis using nitrate as terminal electron acceptor. *Applied
Microbiology and Biotechnology*, *57*(3), 379–384.
https://doi.org/10.1007/s002530100749

James, S. (1984). Lignotubers and Burls: Their Structure, Function and Ecological Significance
in Mediterranean Ecosystems. *Botanical Review*, *50*(3), 225–266.

Kalayu, G. (2019). Phosphate Solubilizing Microorganisms: Promising Approach as
Biofertilizers. *International Journal of Agronomy*, *2019*, 1–7.
https://doi.org/10.1155/2019/4917256
Kaur, N., Seuylemezian, A., Patil, P. P., Patil, P., Krishnamurti, S., Varelas, J., Smith, D. J.,
Mayilraj, S., & Vaishampayan, P. (2018). Paenibacillus xerothermodurans sp. Nov., an
extremely dry heat resistant spore forming bacterium isolated from the soil of Cape
Canaveral, Florida. *International Journal of Systematic and Evolutionary Microbiology*,
68(10), 3190–3196. https://doi.org/10.1099/ijsem.0.002967
Keeley, J. E., Fotheringham, C. J., & Baer-Keeley, M. (2005). Determinants of Postfire
Recovery and Succession in Mediterranean-Climate Shrublands of California. *Ecological*
Applications, *15*(5), 1515–1534. https://doi.org/10.1890/04-1005
Keeley, J. E., & Zedler, P. H. (2009). Large, high-intensity fire events in Southern California
shrublands: Debunking the fine-grain age patch model. In *Ecological Applications* (Vol.
19, Issue 1, p. 26). https://doi.org/10.1890/08-0281.1
Kindt, R., & Coe, R. (2005). *Tree diversity analysis. A manual and software for common*
statistical methods for ecological and biodiversity studies. World Agroforestry Centre
(ICRAF).
Kinzig, A. P., & Pacala, S. (2013). 9. Successional Biodiversity and Ecosystem Functioning. In
The Functional Consequences of Biodiversity: Empirical Progress and Theoretical
Extensions (MPB-33) (pp. 175–212). Princeton University Press.
https://doi.org/10.1515/9781400847303.175
Koide, R. T., Fernandez, C., & Malcolm, G. (2014). Determining place and process: Functional
traits of ectomycorrhizal fungi that affect both community structure and ecosystem
function. *New Phytologist*, *201*(2), 433–439. https://doi.org/10.1111/nph.12538
Kõljalg, U., Larsson, K.-H., Abarenkov, K., Nilsson, R. H., Alexander, I. J., Eberhardt, U.,
Erland, S., Høiland, K., Kjøller, R., Larsson, E., Pennanen, T., Sen, R., Taylor, A. F. S.,
Tedersoo, L., & Vrålstad, T. (2005). UNITE: A database providing web-based methods
for the molecular identification of ectomycorrhizal fungi. *New Phytologist*, *166*(3), 1063–
1068. https://doi.org/10.1111/j.1469-8137.2005.01376.x
Köster, K., Berninger, F., Lindén, A., Köster, E., & Pumpanen, J. (2014). Recovery in fungal
biomass is related to decrease in soil organic matter turnover time in a boreal fire
chronosequence. *Geoderma*, *235–236*, 74–82.
https://doi.org/10.1016/j.geoderma.2014.07.001
Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013).
Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing
Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and*
Environmental Microbiology, *79*(17), 5112–5120. https://doi.org/10.1128/AEM.01043-
13
Krappmann, S., & Braus, G. H. (2005). Nitrogen metabolism of *Aspergillus* and its role in
pathogenicity. *Medical Mycology*, *43*(s1), 31–40.
https://doi.org/10.1080/13693780400024271
Leizeaga, A., Meisner, A., Rousk, J., & Bååth, E. (2022). Repeated drying and rewetting cycles
accelerate bacterial growth recovery after rewetting. *Biology and Fertility of Soils*.
https://doi.org/10.1007/s00374-022-01623-2

Li, X., Rui, J., Mao, Y., Yannarell, A., & Mackie, R. (2014). Dynamics of the bacterial
1026   community structure in the rhizosphere of a maize cultivar. *Soil Biology and*
1027   *Biochemistry*, *68*, 392–401. https://doi.org/10.1016/j.soilbio.2013.10.017
1028 Liu, C. M., Kachur, S., Dwan, M. G., Abraham, A. G., Aziz, M., Hsueh, P.-R., Huang, Y.-T.,
1029   Busch, J. D., Lamit, L. J., Gehring, C. A., Keim, P., & Price, L. B. (2012). FungiQuant: A
1030   broad-coverage fungal quantitative real-time PCR assay. *BMC Microbiology*, *12*(1), 255.
1031   https://doi.org/10.1186/1471-2180-12-255
1032 Livne-Luzon, S., Shemesh, H., Osem, Y., Carmel, Y., Migael, H., Avidan, Y., Tsafrir, A.,
1033   Glassman, S. I., Bruns, T. D., & Ovadia, O. (2021). High resilience of the mycorrhizal
1034   community to prescribed seasonal burnings in eastern Mediterranean woodlands.
1035   *Mycorrhiza*, *31*(2), 203–216. https://doi.org/10.1007/s00572-020-01010-5
1036 Loreau, M., & de Mazancourt, C. (2008). Species Synchrony and Its Drivers: Neutral and
1037   Nonneutral Community Dynamics in Fluctuating Environments. *The American*
1038   *Naturalist*, *172*(2), E48–E66. https://doi.org/10.1086/589746
1039 Mandic-Mulec, I., Stefanic, P., & van Elsas, J. D. (2015). Ecology of Bacillaceae. *Microbiology*
1040   *Spectrum*, *3*(2), 3.2.16. https://doi.org/10.1128/microbiolspec.TBS-0017-2013
1041 Martin, M. (2011). *Cutadapt removes adapter sequences from high-throughput sequencing*
1042   *reads*. https://journal.embnet.org/index.php/embnetjournal/article/view/200/458
1043 McGee, P. A., Markovina, A.-L., Jeong, G. C. E., & Cooper, E. D. (2006). Trichocomaceae in
1044   bark survive high temperatures and fire: Trichocomaceae in bark survive high
1045   temperatures and fire. *FEMS Microbiology Ecology*, *56*(3), 365–371.
1046   https://doi.org/10.1111/j.1574-6941.2006.00079.x
1047 McMullan-Fisher, S. J. M., Tom W. May, Richard M. Robinson, Tina L. Bell, Teresa Lebel,
1048   Pam Catcheside, & Alan York. (2011). *Fungi and fire in Australian ecosystems: A review*
1049   *of current knowledge, management implications and future directions*. *59*, 70–90.
1050 McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
1051   Analysis and Graphics of Microbiome Census Data. *PLOS ONE*, *8*(4), e61217.
1052   https://doi.org/10.1371/journal.pone.0061217
1053 Monciardini, P., Cavaletti, L., Schumann, P., Rohde, M., & Donadio, S. (2003). Conexibacter
1054   woesei gen. Nov., sp. Nov., a novel representative of a deep evolutionary line of descent
1055   within the class Actinobacteria. *International Journal of Systematic and Evolutionary*
1056   *Microbiology*, *53*(2), 569–576. https://doi.org/10.1099/ijs.0.02400-0
1057 Moore, E. J. (1962). The Ontogeny of the Sclerotia of Pyronema Domesticum. *Mycologia*, *54*(3),
1058   312–316. https://doi.org/10.1080/00275514.1962.12025004
1059 Neary, D. G., Klopatek, C. C., DeBano, L. F., & Ffolliott, P. F. (1999). Fire effects on
1060   belowground sustainability: A review and synthesis. *Forest Ecology and Management*,
1061   *122*(1–2), 51–71. https://doi.org/10.1016/S0378-1127(99)00032-8
1062 Nelson, A. R., Narrowe, A. B., Rhoades, C. C., Fegel, T. S., Daly, R. A., Roth, H. K., Chu, R.
1063   K., Amundson, K. K., Geonczy, S. E., Emerson, J. B., Young, R. B., Steindorff, A. S.,
1064   Mondo, S. J., Grigoriev, I. V., Salamov, A., Borch, T., & Wilkins, M. J. (2021). *Playing*
1065   *with FiRE: A genome resolved view of the soil microbiome responses to high severity*
1066   *forest wildfire* [Preprint]. Microbiology. https://doi.org/10.1101/2021.08.17.456416
1067 Nguyen, H. D., Nickerson, N. L., & Seifert, K. A. (2013). Basidioascus and Geminibasidium: A
1068   new lineage of heat-resistant and xerotolerant basidiomycetes. *Mycologia*, *105*(5), 1231–
1069   1250. https://doi.org/10.3852/12-351

Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., &
Kennedy, P. G. (2016). FUNGuild: An open annotation tool for parsing fungal
community datasets by ecological guild. *Fungal Ecology*, *20*, 241–248.
https://doi.org/10.1016/j.funeco.2015.06.006
Ofek, M., Hadar, Y., & Minz, D. (2012). Ecology of Root Colonizing Massilia
(Oxalobacteraceae). *PLOS ONE*, *7*(7), e40117.
https://doi.org/10.1371/journal.pone.0040117
Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.
(2018). *Vegan:Community Ecology Package-Version 2.5.2. Ordination Methods,*
Diversity Analysis, and Other Functions for Community and Vegetation Ecologist.
Owen, S. M., Patterson, A. M., Gehring, C. A., Sieg, C. H., Baggett, L. S., & Fulé, P. Z. (2019).
Large, high-severity burn patches limit fungal recovery 13 years after wildfire in a
ponderosa pine forest. *Soil Biology and Biochemistry*, *139*, 107616.
https://doi.org/10.1016/j.soilbio.2019.107616
Pacala, S. W., Canham, C. D., Saponara, J., Silander, J. A., Kobe, R. K., & Ribbens, E. (1996).
Forest Models Defined by Field Measurements: Estimation, Error Analysis and
Dynamics. *Ecological Monographs*, *66*(1), 1–43. https://doi.org/10.2307/2963479
Pandolfi, J. M. (2008). Succession. In S. E. Jørgensen & B. D. Fath (Eds.), *Encyclopedia of*
Ecology (pp. 3416–3424). Academic Press. https://doi.org/10.1016/B978-008045405-
4.00547-4
Paredes-Sabja, D., Setlow, P., & Sarker, M. R. (2011). Germination of spores of Bacillales and
Clostridiales species: Mechanisms and proteins involved. *Trends in Microbiology*, *19*(2),
85–94. https://doi.org/10.1016/j.tim.2010.10.004
Parson, A., Robichaud, P. R., Lewis, S. A., Napper, C., & Clark, J. T. (2010). *Field guide for*
mapping post-fire soil burn severity (RMRS-GTR-243; p. RMRS-GTR-243). U.S.
Department of Agriculture, Forest Service, Rocky Mountain Research Station.
https://doi.org/10.2737/RMRS-GTR-243
Pena, R., Offermann, C., Simon, J., Naumann, P. S., Geßler, A., Holst, J., Dannenmann, M.,
Mayer, H., Kögel-Knabner, I., Rennenberg, H., & Polle, A. (2010). Girdling Affects
Ectomycorrhizal Fungal (EMF) Diversity and Reveals Functional Differences in EMF
Community Composition in a Beech Forest. *Applied and Environmental Microbiology*,
76(6), 1831–1841. https://doi.org/10.1128/AEM.01703-09
Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2018). Resilience to fire of
phylogenetic diversity across biological domains. *Molecular Ecology*, *27*(13), 2896–
2908. https://doi.org/10.1111/mec.14729
Pérez-Valera, E., Verdú, M., Navarro-Cano, J. A., & Goberna, M. (2020). Soil microbiome
drives the recovery of ecosystem functions after fire. *Soil Biology and Biochemistry*, *149*,
107948. https://doi.org/10.1016/j.soilbio.2020.107948
Petersen, P. M. (1970). Danish fireplace fungi-an ecological investigation on fungi on burns.
Dansk Botanisk Arkiv, *27*(3), 1–97.
Placella, S. A., Brodie, E. L., & Firestone, M. K. (2012). Rainfall-induced carbon dioxide pulses
result from sequential resuscitation of phylogenetically clustered microbial groups.
Proceedings of the National Academy of Sciences, *109*(27), 10931–10936.
https://doi.org/10.1073/pnas.1204306109

Pourreza, M., Hosseini, S. M., Safari Sinegani, A. A., Matinizadeh, M., & Dick, W. A. (2014).
Soil microbial activity in response to fire severity in Zagros oak (Quercus brantii Lindl.)
forests, Iran, after one year. *Geoderma*, *213*, 95–102.
https://doi.org/10.1016/j.geoderma.2013.07.024
Pressler, Y., Moore, J. C., & Cotrufo, M. F. (2019). Belowground community responses to fire:
Meta-analysis reveals contrasting responses of soil microorganisms and mesofauna.
Oikos, *128*(3), 309–327. https://doi.org/10.1111/oik.05738
Pulido-Chavez, M. F., Alvarado, E. C., DeLuca, T. H., Edmonds, R. L., & Glassman, S. I.
(2021). High-severity wildfire reduces richness and alters composition of
ectomycorrhizal fungi in low-severity adapted ponderosa pine forests. *Forest Ecology*
and Management, *485*, 118923. https://doi.org/10.1016/j.foreco.2021.118923
R Core Team. (2021). *R: A language and environment for statistical computing*. *R Foundation*
for Statistical Computing. https://www.R-project.org/.
Reazin, C., Morris, S., Smith, J. E., Cowan, A. D., & Jumpponen, A. (2016). Fires of differing
intensities rapidly select distinct soil fungal communities in a Northwest US ponderosa
pine forest ecosystem. *Forest Ecology and Management*, *377*, 118–127.
https://doi.org/10.1016/j.foreco.2016.07.002
Reilly, M. J., & Spies, T. A. (2016). Disturbance, tree mortality, and implications for
contemporary regional forest change in the Pacific Northwest. *Forest Ecology and*
Management, *374*, 102–110. https://doi.org/10.1016/j.foreco.2016.05.002
Rhodes, J. C. (2006). *Aspergillus fumigatus*: Growth and virulence. *Medical Mycology*, *44*(s1),
77–81. https://doi.org/10.1080/13693780600779419
Riley, K. L., & Loehman, R. A. (2016). Mid-21st-century climate changes increase predicted fire
occurrence and fire season length, Northern Rocky Mountains, United States. *Ecosphere*,
7(11), e01543. https://doi.org/10.1002/ecs2.1543
Ross, G. J. S., & Preece, D. A. (1985). The Negative Binomial Distribution. *Journal of the Royal*
Statistical Society. Series D (The Statistician), *34*(3), 323–335. JSTOR.
https://doi.org/10.2307/2987659
Rundel, P. W. (2018). California Chaparral and Its Global Significance. In E. C. Underwood, H.
D. Safford, N. A. Molinari, & J. E. Keeley (Eds.), *Valuing Chaparral* (pp. 1–27).
Springer International Publishing. https://doi.org/10.1007/978-3-319-68303-4_1
Schimel, J. P. (2018). Life in Dry Soils: Effects of Drought on Soil Microbial Communities and
Processes. *Annual Review of Ecology, Evolution, and Systematics*, *49*(1), 409–432.
https://doi.org/10.1146/annurev-ecolsys-110617-062614
Schluter, D., & Ricklefs, R. E. (1993). *Species diversity: An introduction to the problem. Species*
diversity in ecological communities. Chicago University Press.
Seaver, F. J. (1909). Studies in Pyrophilous Fungi: I. The Occurrence and Cultivation of
Pyronema. *Mycologia*, *1*, 131–193.
Selbmann, L., Egidi, E., Isola, D., Onofri, S., Zucconi, L., de Hoog, G. S., Chinaglia, S., Testa,
L., Tosi, S., Balestrazzi, A., Lantieri, A., Compagno, R., Tigini, V., & Varese, G. C.
(2013). Biodiversity, evolution and adaptation of fungi in extreme environments. *Plant*
Biosystems - An International Journal Dealing with All Aspects of Plant Biology, *147*(1),
237–246. https://doi.org/10.1080/11263504.2012.753134
Semenova-Nelsen, T. A., Platt, W. J., Patterson, T. R., Huffman, J., & Sikes, B. A. (2019).
Frequent fire reorganizes fungal communities and slows decomposition across a

heterogeneous pine savanna landscape. *New Phytologist*, *224*(2), 916–927.
https://doi.org/10.1111/nph.16096
Shugart, H. H. (2013). Encyclopedia of Biodiversity. In *Succession, Phenomenon Of* (2nd
Edition, pp. 63–70). Academic Press.
Slepecky, R. A., & Hemphill, H. E. (2006). The Genus Bacillus—Nonmedical. In M. Dworkin,
S. Falkow, E. Rosenberg, K.-H. Schleifer, & E. Stackebrandt (Eds.), *The Prokaryotes:*
Volume 4: Bacteria: Firmicutes, Cyanobacteria (pp. 530–562). Springer US.
https://doi.org/10.1007/0-387-30744-3_16
Steindorff, A. S., Carver, A., Calhoun, S., Stillman, K., Liu, H., Lipzen, A., He, G., Yan, M.,
Pangilinan, J., LaButti, K., Ng, V., Bruns, T. D., & Grigoriev, I. V. (2021). Comparative
genomics of pyrophilous fungi reveals a link between fire events and developmental
genes. *Environmental Microbiology*, *23*(1), 99–109. https://doi.org/10.1111/1462-
2920.15273
Taylor, D. L., Walters, W. A., Lennon, N. J., Bochicchio, J., Krohn, A., Caporaso, J. G., &
Pennanen, T. (2016). Accurate Estimation of Fungal Diversity and Abundance through
Improved Lineage-Specific Primers Optimized for Illumina Amplicon Sequencing.
Applied and Environmental Microbiology, *82*(24), 7217–7226.
https://doi.org/10.1128/AEM.02576-16
Tilman, D. (1990). Constraints and Tradeoffs: Toward a Predictive Theory of Competition and
Succession. *Oikos*, *58*(1), 3–15. https://doi.org/10.2307/3565355
Toljander, J. F., Artursson, V., Paul, L. R., Jansson, J. K., & Finlay, R. D. (2005). Attachment of
di¡erent soil bacteria to arbuscular mycorrhizal fungal extraradical hyphae is determined
by hyphal vitalityand fungal species. *FEMS Microbiol Lett*, 7.
Torti, A., Lever, M. A., & Jørgensen, B. B. (2015). Origin, dynamics, and implications of
extracellular DNA pools in marine sediments. *Marine Genomics*, *24*, 185–196.
https://doi.org/10.1016/j.margen.2015.08.007
Treseder, K. K., Mack, M. C., & Cross, A. (2004). Relationships Among Fires, Fungi, and Soil
Dynamics in Alaskan Boreal Forests. *Ecological Applications*, *14*(6), 1826–1838.
https://doi.org/10.1890/03-5133
Turnbull, L. A., Levine, J. M., Loreau, M., & Hector, A. (2013). Coexistence, niches and
biodiversity effects on ecosystem functioning. *Ecology Letters*, *16*, 116–127.
https://doi.org/10.1111/ele.12056
Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (Fourth edition).
Springer. https://www.stats.ox.ac.uk/pub/MASS4/.
Warcup, J. H., & Baker, K. F. (1963). Occurrence of Dormant Ascospores in Soil. *Nature*,
197(4874), 1317–1318. https://doi.org/10.1038/1971317a0
Whitman, T., Whitman, E., Woolet, J., Flannigan, M. D., Thompson, D. K., & Parisien, M.-A.
(2019). Soil bacterial and fungal response to wildfires in the Canadian boreal forest
across a burn severity gradient. *Soil Biology and Biochemistry*, *138*, 107571.
https://doi.org/10.1016/j.soilbio.2019.107571
Wickham H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
https://ggplot2.tidyverse.org
Woolet, J., & Whitman, T. (2020). Pyrogenic organic matter effects on soil bacterial community
composition. *Soil Biology and Biochemistry*, *141*, 107678.
https://doi.org/10.1016/j.soilbio.2019.107678

Xiang, X., Gibbons, S. M., Yang, J., Kong, J., Sun, R., & Chu, H. (2015). Arbuscular
mycorrhizal fungal communities show low resistance and high resilience to wildfire
disturbance. *Plant and Soil*, *397*(1–2), 347–356. https://doi.org/10.1007/s11104-015-
2633-z
Xiang, X., Shi, Y., Yang, J., Kong, J., Lin, X., Zhang, H., Zeng, J., & Chu, H. (2014). Rapid
recovery of soil bacterial communities after wildfire in a Chinese boreal forest. *Scientific*
Reports, *4*(1), 3829. https://doi.org/10.1038/srep03829
Yang, T., Tedersoo, L., Lin, X., Fitzpatrick, M. C., Jia, Y., Liu, X., Ni, Y., Shi, Y., Lu, P., Zhu,
J., & Chu, H. (2020). Distinct fungal successional trajectories following wildfire between
soil horizons in a cold-temperate forest. *New Phytologist*, *227*(2), 572–587.
https://doi.org/10.1111/nph.16531
Yang, Zhu, K., Loik, M. E., & Sun, W. (2021). Differential responses of soil bacteria and fungi
to altered precipitation in a meadow steppe. *Geoderma*, *384*, 114812.
https://doi.org/10.1016/j.geoderma.2020.114812
Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J.,
Ludwig, W., & Glöckner, F. O. (2014). The SILVA and "All-species Living Tree Project
(LTP)" taxonomic frameworks. *Nucleic Acids Research*, *42*(D1), D643–D648.
https://doi.org/10.1093/nar/gkt1209
Zhang, H., Qi, W., & Liu, K. (2018). Functional traits associated with plant colonizing and
competitive ability influence species abundance during secondary succession: Evidence
from subalpine meadows of the Qinghai–Tibetan Plateau. *Ecology and Evolution*, *8*(13),
6529–6536. https://doi.org/10.1002/ece3.4110
1227
1228