The price of defence: toxins, visual signals and oxidative state in an aposematic butterfly

Jonathan D. Blount [1*†], **Hannah M. Rowland**[2,3†], **Christopher Mitchell**[1], **Michael P. Speed**[4], **Graeme D. Ruxton**[5], **John A. Endler**[6], **Lincoln P. Brower**[7]

[1] Centre for Ecology & Conservation, College of Life & Environmental Sciences, University of Exeter, Penryn Campus, Cornwall TR10 9FE, UK

[2] Max Planck Institute for Chemical Ecology, Hans-Knöll-Straße 8, Jena, 07745, Germany

[3] Department of Zoology, University of Cambridge, Cambridge CB2 3EJ, UK

[4] Institute of Integrative Biology, University of Liverpool, Liverpool L69 7ZB, UK

[5] School of Biology, Sir Harold Mitchell Building, Greenside Place, St Andrews, United Kingdom

[6] Centre for Integrative Ecology, School of Life & Environmental Sciences, Deakin University, Waurn Ponds VIC 3216, Australia

[7] Department of Biology, Sweet Briar College, Sweet Briar VA 24595, USA

* Authors for correspondence: j.d.blount@exeter.ac.uk and hrowland@ice.mpg.de

† Joint first authors

Running headline: Oxidative stress and warning signal honesty

SUMMARY

In a variety of aposematic species, the conspicuousness of an individual's warning signal and

the quantity of its chemical defence are positively correlated. This apparent honest signalling

in aposematism is predicted by resource competition models which assume that the

production and maintenance of aposematic defences compete for access to antioxidant

molecules that have dual functions as pigments directly responsible for colouration and in

protecting against oxidative lipid damage. Here we study a model aposematic system—the

monarch butterfly (*Danaus plexippus*) and make use of the variable phytochemistry of its

larval host-plant, milkweeds (Asclepiadaceae), to manipulate the concentration of

sequestered cardenolides. We test two fundamental assumptions of resource competition

models: that (1) the possession of secondary defences is associated with costs in the form of

oxidative lipid damage and reduced antioxidant defences; and (2) that oxidative damage or

decreases in antioxidant defences can reduce the capacity of individuals to produce

aposematic displays. Monarch caterpillars that sequestered the highest concentrations of

cardenolides exhibited higher levels of oxidative lipid damage as adults. The relationship

between warning signals, cardenolide concentrations and oxidative damage differed between

the sexes. In male monarchs conspicuousness was explained by an interaction between

oxidative damage and sequestration: as males sequester more cardenolides, those with high

levels of oxidative damage become less conspicuous, while those that sequester lower levels

of cardenolides equally invest in conspicuous with increasing oxidative damage. There was

no significant effect of oxidative damage or concentration of sequestered cardenolides on

female conspicuousness. Our results demonstrate physiological linkage between the

production of coloration and protection from autotoxicity, that warning signals can be honest

indicators of defensive capability, and that the relationships are different between the sexes.

Introduction

Many species of animals, plants and micro-organisms possess chemical defences that reduce

the likelihood of their being eaten [1]. Aposematic species, such as the monarch butterfly

(*Danaus plexippus*), signal their chemical defences to predators with conspicuous warning

colours [2]. In some species, the conspicuousness of the warning signal and the quantity of

chemical defence are positively correlated, for example in dendrobatid frogs [3-5], marine

opisthobranchs [6], ladybird beetles [7, 8], and paper wasps *Polistes dominula* [9]. The

identification of these relationships has led to renewed interest in the idea that warning

signals may inform predators about levels of prey defence and that such signals are honest

indicators of defensive capability [10]. Several theoretical studies have, however, predicted

the opposite: that well defended prey should reduce investment in signals, because signalling

carries an additional conspicuousness cost, and such prey stand a good chance of surviving

any attack unharmed [11-13]. Negative signal-defence correlations have been reported across

Dendrobatid species of poison dart frogs [14] including the granular poison dart frog

Dendrobates granuliferus [15]. Using a novel theoretical model Blount et al [16] could

explain both the positive and negative correlations between warning colours and chemical

defences if these two traits are linked through the competitive use of a shared resource. One

possible shared resource is energy, which can be limiting for the sequestration or biosynthesis

of toxins [17]. Alternatively, sequestration of toxins could impose metabolic costs in terms of

oxidative lipid damage and reduced antioxidant defences – an idea that has has received

limited empirical attention in regard to warning signals.

Many plant allelochemicals are powerful pro-oxidants, which, when ingested, can cause

oxidative lipid damage [18]. In their resource competition framework, Blount et al [16]

envisaged that pigments used in prey warning signals (e.g., carotenoids, flavonoids, melanins,

and pteridines) might play a dual role imparting colour as well as acting as antioxidants that

prevent self-damage when storing toxins. In their model, if antioxidants are required to

enable high levels of toxicity, signal reliability can be explained if the brightest and most

toxic species gain access to more of the limiting resource than those that are less bright and

less toxic. The prediction of signal honesty breaks down in their model when prey have very

abundant resources, in which case it pays to divert these resources increasingly into toxins

and not into warning signals, because less conspicuous but highly toxic prey encounter

predators less often and have high chances of surviving attacks [16]. Because directly

controlling antioxidant availability is challenging experimentally, whether colour-toxin

relationships are influenced by the resource state of prey is poorly understood [14, 15], and

has not yet been examined experimentally [11, 12, 16, 17, 19].

Here we manipulate diet-derived toxins available to an individual to test whether toxin

sequestration and the production and maintenance of warning signals are linked via oxidative

state. We achieve this by using a model aposematic system, the monarch butterfly, that

selectively metabolises cardenolides present in its host-plants and stores them in its body [20,

21], and by using the known inter- and intraspecific variability in cardenolide concentration

and cardenolide polarity of the monarch's larval milkweed host-plants (Asclepiadaceae; [21,

22]) to manipulate the toxins available to an individual to sequester. We test whether (1) the

quantity of sequestered secondary defence by caterpillars is associated with oxidative lipid

damage and reduced antioxidant defences in adults; and (2) whether oxidative lipid damage

incurred during sequestration affects the capacity of adult butterflies to produce aposematic

displays.

Because there is substantial genetic variation in sequestration ability, and because

evolutionary history and contemporary species interactions may influence patterns of

cardenolide sequestration [23] we used monarchs from the same population. We reared them

on four milkweed species that show within- and between-species, and among-population

variation in cardenolide concentration, diversity, and polarity [24-27]. The polarity of

cardenolides can influence their toxicity [21] and transport [28], so our four plant species

were used to represent a gradient of distinct cardenolide defence, with the aim to create

different costs of sequestration.

We reared monarch caterpillars on either *Ascelpias curassavica*, which is a critical hostplant

worldwide and contains a diversity of cardenolides, including apolar voruscharin which is

detrimental to monarch performance [21, 29-32]; *A. syriaca* (common milkweed) which is

the principal native food plant of the monarch east of the Rocky Mountains and is

characterized by more polar cardenolides though also contains the apolar labriformin [33,

34]; *A. incarnata* (swamp milkweed) which is a preferred hostplant in the Midwest USA and

has low concentrations of cardenolides [35]; or *A. tuberosa* (butterfly milkweed) which is an

occasional native foodplant in the wild with low concentrations of cardenolides [21, 36].

Based on the published records of biological activity and chromatic quantification (e.g.,

HPLC and thin layer chromatography) of cardenolides of milkweeds in different

populations/latitudes [37, 38], we predicted that concentrations of cardenolides in the body

and wings of newly eclosed adults would differ according to the cardenolides of the host-

plant on which they were reared (see methods): *A. curassavica* (*Ac*) $\geq$ *A. syriaca* (*As*) > *A.*

incarnata (*Ai*) $\geq$ *A. tuberosa* (*At*). Because sequestration, modification, and storage of

allelochemicals are believed to be oxidatively stressful for many chemically defended

organisms [18], we predicted that body levels of oxidative lipid peroxidation would differ

amongst individuals according to host-plant, as follows: *Ac$\geq$As>Ai$\geq$At*. In turn, we predicted

that survival to eclosion would differ according to host-plant as follows: *At$\geq$Ai>As$\geq$Ac*, and

that the redness, luminance, and conspicuousness of the warning signals would differ:

Ac$\geq$As>Ai$\geq$At. In addition to testing for treatment effects, we examined associations between

individual levels of cardiac glycosides (CGs), oxidative state, and warning signal redness,

luminance, and conspicuousness. We predicted a positive association between levels of CGs

and and lipid peroxidation, a negative association of CGs with antioxidant defences, and that

highly toxic prey will reduce investment in signals if they have high levels of oxidative

damage because they cannot bear the oxidative cost of investing in both traits [16].

Materials and methods

(a) Milkweed plants

Milkweeds (*A. incarnata, A. tuberosa, A. curassavica, and A. syriaca*) were locally sourced

in Amherst, Virginia, USA. We did not measure cardenolide content in the leaves of the

plants used in the experiment, but used published records about the spatial repeatability of

cardenolide defences that take into account latitudinal clines [37-39]. From these records we

predicted that for milkweeds grown in Mid-Atlantic and Southeastern regions of the United

States, total cardenolide content would vary in the order *A. curassavica* (*Ac*) $\geq$ *A. syriaca* (*As*)

> *A. incarnata* (*Ai*) $\geq$ *A. tuberosa* (*At*). The plants were grown without the use of pesticides as

described previously [40]. All plants were healthy with undamaged leaves at the start of the

experiment.

(b) *Milkweed Feeding Assay and preparation of samples*

Eggs were obtained from wild caught Monarch females in Virginia (USA) and larvae were

reared by LPB on single plants. All larvae had abundant food during the experiment. Larvae

were monitored daily for pupation starting at Day 12.

Upon eclosion, the date and the individual's sex were recorded. Adults were placed in

glassine envelopes and frozen on dry ice until dead, then stored in a freezer at -80°C before

being shipped to the UK on dry ice. Adults were stored at -80°C in JDB's laboratory until

they were analysed.

For analysis, the gut was removed, and the head, thorax and abdomen (hereafter, 'body')

were bisected longitudinally. One half of the body was returned to storage at -80°C for later

analysis of cardenolides. The other half of the body was immediately homogenised in ice-

cold phosphate-buffered saline (PBS; 5 % w/v), for analyses of oxidative stress. The wings

were removed using dissecting scissors and shipped on dry ice to HMR's laboratory for

analyses of colouration (see below). Wings were then returned (on dry ice) to JDB's

laboratory for cardenolide analysis.

(c) Determination of oxidative stress and cardenolide concentration

Determining oxidative state requires a range of assays including antioxidant defences and

oxidative damage. We assayed three such components: total superoxide dismutase (SOD),

which are metalloenzymes that catalyse the dismutation of superoxide into oxygen and

hydrogen peroxide and form a crucial part of intracellular antioxidant defences; total

antioxidant capacity (TAC), which measures the activity of low molecular weight chain-

breaking antioxidants including ascorbate, a-tocopherol, carotenoids and flavonoids, and

therefore gives an overview of the antioxidant status of individuals (including diet-derived

antioxidants); and malondialdehyde (MDA), which is formed by the β-scission of peroxidised

polyunsaturated fatty acids, and therefore is a definitive marker of oxidative damage.

(i)   *Total superoxide dismutase (SOD)*

Total SOD was assayed by measuring the dismutation of superoxide radicals generated by

xanthine oxidase and hypoxanthine. We followed the instructions of the kit (#706002;

Cayman Chemical, Michigan, USA), except that tissue homogenates were further diluted

with the supplied sample buffer (1:100 v/v) to ensure that SOD activity fell within the range

of the standard curve. Samples were assayed in duplicate and are reported as units of SOD

activity (U) per mg tissue.

(ii)   Total antioxidant capacity (TAC)

TAC was assayed by measuring the capacity of tissue homogenate to inhibit the oxidation of

2,2'-azino-di-[3-ethylbenzthiazoline sulphonate] (#709001; Cayman Chemical). The

homogenate was further diluted with sample buffer (1:5 v/v) to bring the absorbance values

within the range of the standard curve. Samples were assayed in duplicate, as per the kit

instructions. Data are reported as nmols of TAC activity (Trolox equivalents) per g tissue.

(iii)   Malondialdehyde (MDA)

MDA was measured using HPLC with fluorescence detection as described previously for

tissue homogenates (e.g. Blount & Pike 2011). All chemicals were HPLC grade, and

chemical solutions were prepared using ultra pure water (Milli-Q Synthesis; Millipore,

Watford, UK). Samples were derived in 2 mL capacity polypropylene screw-top

microcentrifuge tubes. For tissue homogenates and for the standard (1,1,3,3-

tetraethoxypropane, TEP; see below) a 20 µl aliquot was added to 20 µl butylated

hydroxytoluene solution (BHT; 0.05% w/v in 95 % ethanol), 160 µl phosphoric acid solution

(H_3PO_4; 0.44 *M*), and 40 µl thiobarbituric acid solution (TBA; 42 m*M*). Samples were

capped, vortex mixed for 5 s, then heated at 100 °C for 1 h in a dry bath incubator to allow

formation of MDA-TBA adducts. Samples were then cooled on ice for 5 min, before 80 µl *n*-

butanol was added and tubes were vortex mixed for 20s. Tubes were centrifuged at 12 x g

and 4 °C for 3 min, before the upper (*n*-butanol) phase was collected and transferred to an

HPLC vial for analysis. Samples (10 µl) were injected into a Dionex HPLC system (Dionex

Corporation, California, USA) fitted with a 5µm ODS guard column and a Hewlett-Packard

Hypersil 5µ ODS 100 x 4.6 mm column maintained at 37 °C. The mobile phase was

methanol-buffer (40:60, v/v), the buffer being a 50 m*M* anhydrous solution of potassium

monobasic phosphate at pH 6.8 (adjusted using 5 *M* potassium hydroxide solution), running

isocratically over 3.5 min at a flow rate of 1 ml min^{-1}. Data were collected using a

fluorescence detector (RF2000; Dionex) set at 515 nm (excitation) and 553 nm (emission).

For calibration a standard curve was prepared using a TEP stock solution (5 m*M* in 40 %

ethanol) serially diluted using 40 % ethanol. Data are presented as nmols MDA per g tissue.

(iv)   Cardenolide analysis

For cardenolide analysis, samples (individual wings and the remaining half of the body) were

dried in an oven at 60 °C for 16 h. The dried material was ground to a fine powder using a

pestle and mortar, and weighed to the nearest 0.0001 g using a GR-202 electronic balance

(A&D Instruments Ltd., Abington, UK). The samples were placed in screw-top

polypropylene tubes for de-fatting and cardenolide extraction. To remove fats, 2 ml hexane

was added and samples were incubated at 35°C for 30 min, before being centrifuged at 10 x g

and 4ºC for 5 min; the supernatant was collected and discarded. Ethanol (1.9 ml) containing

20 μg digitoxin (#D5878; Sigma-Aldrich, Dorset, UK) as an internal standard were added to

the pellet and it was briefly vortexed to extract cardenolides (Ritland 1994). Samples were

placed in an XUB5 ultrasonic bath (Grant Instruments, Shepreth, UK) at 60°C for 60 min,

followed by further centrifugation (9.6 x g and 4ºC for 5 min). The supernatant was removed

and the remaining pellet dried using a Savant ISS 110 SpeedVac Concentrator at room

temperature (Thermo Scientific, Altrincham, UK). Samples were then dissolved in 0.5 ml

methanol by vortexing, and 50 μl of the solution was injected into a Dionex HPLC system

fitted with a Waters Spherisorb 5μm ODS2 column (150 x 4.6 mm) maintained at 20°C. A

multistep gradient of acetonitrile (ACN) and ultra pure water was used as the mobile phase,

as follows: initial concentration 20 % ACN held for 5 minutes, ramp to 70 % ACN at 20 mins

and held until 25 minutes, ramp to 95 % ACN at 30 mins and held until 35 mins, returning to

20 % ACN at 40 mins, with an end time of 50 mins. The flow rate was 0.7 ml min^{-1}. Peaks

were collected using a PDA-100 photodiode array detector (Dionex) at an absorbance value

of 218 nm. Spectral data between 200-400 nm were collected, and cardenolides were

identified as peaks with a symmetrical absorbance band with a maximum absorbance

between 217-222 nm [41, 42]. Data are presented as μg total cardenolides per 0.1 g tissue.

(d) Measurements and visual modelling of colouration

The reflectance of the fore- and hindwings of each butterfly, and of four leaves from one of

the representative backgrounds of the leaves of *Asclepias syriaca* (obtained from A.

Agrawal), were measured using an Ocean Optics USB2000 spectrophotometer, with

specimens illuminated at 45º to normal by a DH1000 balanced halogen deuterium light

source. The measuring spot diameter was 3 mm, with spectra recorded at 0.34nm intervals

from 300 to 700nm and measured relative to a WS-1 reflectance standard. For each wing, 3-4

non-overlapping measurements were taken per orange segment of the wing (figure s1; total

number of measurement spots per forewing = 13-14; hind wing = 28-29). For the leaves we

took ten non-overlapping measurements. Spectrophotometry data were recorded using Ocean

Optics OOBase32.

Reflectance spectra were reduced to 1nm intervals ranging from wavelengths 300-700 nm

using customised MATLAB code (written by I. C. Cuthill). To determine differences in

chroma between treatments, we modelled the predicted photon catches of a generalized

passerine bird (passerines being the main avian predators of monarchs [43]) for each

spectrum following the Vorobyev-Osorio model [44], using a customized MATLAB code

(provided by I. C. Cuthill, 2006). We used the cone types of a blue tit, *Cyanistes caeruleus*

[45, longwave, LWS, λmax 563nm; mediumwave, MWS, λmax 503 nm; shortwave, SWS,

λmax 448 nm ; ultraviolet, UVS, λmax 371 nm; and double dorsal, DD λmax 563 nm;

46].We calculated redness as the ratio of long wave to medium wave cone responses. We also

calculated a measure of conspicuousness as ΔS between the butterflies and *A. syriaca* leaves

using the Vorobyev-Osorio colour discrimination model, which is based on evidence that

colour discrimination is determined by noise arising in the photoreceptors and is independent

of light intensity. We used a Weber fraction value of 0.05 for the most abundant cone type

[47]. We calculated luminance as the response of the dorsal double cone [48].

(e) Data analyses

Left- and right-wing cardenolide concentration (from here on CG) were significantly

positively correlated (r = 0.96, t = 20.75, df = 38, p = 2.2e-16). There was no significant

correlation between asymmetry in wing CG concentrations and the mean CG of the wings (r

= 0.10, t = 0.64, df = 38, P = 0.52), and so for each individual we calculated a mean wing CG

concentration. Left- and right-wing redness were significantly positive correlated (r = 0.65, t

= 5.30, df = 38, P = 5.2e-06), as were left- and right-wing conspicuousness (r = 0.59, t = 4.41,

df = 37, P = 8.6e-05), and left- and right-wing luminance (r = 0.396, t = 2.66, df = 38, P =

0.0114). For each individual we calculated a mean forewing redness, conspicuousness, and

luminance.

We log transformed malondialdehyde (MDA) and wing and body CG and to normalise the

distribution of the residuals. We analysed the effect of foodplant on total CG content,

bodyCG and wingCG content using a GLM with a Gaussian distribution and identity link

function. We set *A. tuberosa* as the level with which we compared the other plants, because

of its low cardenolide content. Eclosion success or failure was analysed with a binomial

GLM. To analyse how cardenolides and oxidative stress affected colour metrics in the

butterflies we used two methods. First, we used the R drop1 function to remove single model

terms to find the minimal adequate model and compared models using AIC (see

supplementary tables s1-s11). We explored interactions between two continuous variables

using the coplot function in the R package car [49]. Second, we built a generalized linear

model including six covariates *a priori* expected to affect warning signals (three markers of

oxidative state and three measures of sequestration – body, wing, and total CG content). The

model response variables were conspicuousness, redness, or brightness. We standardized all

covariates to a mean of 0 and a standard deviation (SD) of 1 to facilitate comparisons

between models and variables. We ran all possible models containing each covariate (without

interactions) using the R package MuMIn [50] and summed the cumulative AIC support for

the models containing each covariate. MDA, body cardenolides, and total cardenolides had

AIC weights of 0.153, 0.081, and 0.075 (supplementary table s8). All data were analysed

using R version 4.0.4.

Results

CG concentrations

In support of our first hypothesis, concentrations of cardenolides in the body and wings of

newly eclosed adults differed according to which host-plant species they were reared on

(Figure 1a; $F_{(3,37)} = 8.262$, P = 0.0002). Monarchs reared on *Ac* and *As* sequestered

significantly more cardenolides in total than those reared on *At* (estimate = 822.9 ± 184.3, t =

4.47, P = 7.6e-05; estimate = 726.6 ± 187.1, t = 3.88, P =0.0004). Those reared on *Ai* did not

sequester significantly different total amounts of CGs to those on *At* (estimate = 213.9 ±

189.2, t = 1.13, P = 0.27). Males tended to sequester more cardenolides in total than females,

but this was not significant at the alpha 0.05 level (estimate = 243.5 ± 133.4, t = 1.825, P =

0.076).

Body CG concentration tended to differ between food plants, but this was not significant at

the alpha 0.05 level (Figure 1b; $F_{(4,37)}= 2.471$, P = 0.061). Wing CG concentration

significantly varied between host-plants (Figure 1c; $F_{(3,37)}= 26.23$, P <0.001 ; Fig. 1c). Wing

CGs were significantly higher in *Ac* and *As* compared to At (estimate = 1.56 ± 0.27, t =

5.698, P = 1.6e-06; estimate = 1.77 ± 0.29, t = 6.120, P = <0.001), but did not differ

significantly between *At* and *Ai* (p=0.38), nor between *As* and *Ac* (p= 0.79).

Survival to pupation and eclosion

We predicted that sequestration of cardenolides would impose a cost that would be associated

with reduced survivorship to eclosion. Successful eclosion was significantly higher in

monarchs reared on *At* (89%) compared with those reared on *As* (20%; estimate = -2.06 ±

0.751, t = 2.75, P = 0.009), while there was no significant difference between *Ai* (70%) and

At (89%, estimate = -0.696 ± 0.728, t = -0.956, P = 0.34) or *At* (89%) and *Ac* (62%; estimate

= -0.927 ± 0.690, t = -1.344, P = 0.19). Survival to pupation did not differ significantly

according to host-plant species (0% *At*, 23% *Ai*, 33% *As*, 27% *Ac*; Fisher's exact test P =

0.28).

CG and oxidative stress

Malondialdehyde (MDA)

Contrary to our prediction that oxidative lipid damage would differ by hostplant, we found

that the minimal adequate model did not include species of host-plant, and only included sex

and the total concentration of cardenolide. In line with our prediction of a positive association

between concentration of sequestered CGs and markers of oxidative damage, across all

treatment groups individuals that sequestered higher total concentrations of CGs had higher

levels of MDA (estimate = 0.0005 ± 0.0002, t = 2.51, P = 0.016). There was no significant

difference between the sexes (estimate = -0.3097 ± 0.198, t = -1.57, P = 0.13).

Superoxide dismutase activity (SOD)

The minimal adequate model included only body CG concentration, which had no

statistically significant association with SOD (estimate = 0.933 ± 0.616, t = 1.51, P = 0.14).

Total antioxidant capacity (TAC)

The minimal adequate model included foodplant, which did not significantly affect the

concentrations of total antioxidant capacity (TAC: $F_{(3,38)}$= 0.3203, P = 0.81).

Sequestration, oxidative stress and warning signals

Because malondialdehyde was the only marker associated with individual levels of

sequestration, we proceeded to analyse its association with warning signals, but did not

conduct analyses on superoxide dismutase activity or total antioxidant capacity effects on

warning signals. We found a significant three-way interaction between MDA x total CG

concentration x sex on the conspicuousness (chroma ΔS) of monarch forewings (estimate = -

0.009 $\pm$ 0.005, t = -2.05, P = 0.049). Because males were more conspicuous than females, we

separated the analysis by sex. We found a significant two way-interaction between MDA and

total CG concentration on the conspicuousness (chroma ΔS) of male forewings (estimate = -

0.006 $\pm$ 0.002, t = -2.98, P = 0.007) that is represented by a curved surface (Figure 3). Males

with the highest levels of sequestered cardenolides had the most conspicuous warning signals

when oxidative damage was lowest, and reduced their warning signal conspicuousness as

oxidative damage increased, while those that sequestered the lowest levels of cardenolides

invested equally in warning signal conspicuous with increasing oxidative damage. There was

no significant effect of oxidative damage or concentration of sequestered cardenolides on

female forewing conspicuousness (MDA estimate = 0.320 $\pm$ 2.7859, t = 0.12, P = 0.911; CG

total estimate = -0.0005 $\pm$ 0.003, t = -0.14, P = 0.895). Males had significantly redder

forewings than females (estimate = 0.263 $\pm$ 0.063, t = 4.15, P = 0.0002), and as the

concentration of sequestered cardenolides increased the redness of the forewings increased

for both males and females, but this was not significant at the alpha 0.05 level (estimate =

0.0007 $\pm$ 0.0003, t = 1.95, P = 0.059). Monarchs with low levels of oxidative damage

produced redder warning signals with increasing concentrations of sequestered cardenolides,

whereas and monarchs with high levels of oxidative damage produced less red warning

signals with increasing concentrations of sequestered cardenolides, but this was also not

significant at the alpha 0.05 level (MDA x total CG interaction: estimate = -0.0002 $\pm$

0.00009, t = -1.864, P = 0.071). Males had significantly brighter forewings than females

(estimate = 0.027 ± 0.007, t = 4.101, P = 0.0002). Both male and female monarchs with

higher levels of MDA had significantly less bright forewings (estimate = -0.01 ± 0.005, t =

2.04, P = 0.048).

Discussion

We reared individual monarch caterpillars on whole plants from single populations of four

milkweed species with varying phytochemistry [*Asclepias tuberosa*, *A. incarnata*, *A.syriaca*,

and *A.curassavica*; 51]. Monarchs that sequestered higher concentrations of cardenolides

experienced higher levels of oxidative damage than those that sequestered lower

concentrations (measured as the concentration of malondialdehyde; MDA). Although there is

some evidence that cardenolides can be a burden for monarch caterpillars [22, 52], our results

are some of the first to show physiological costs of sequestration for monarchs [see also 21,

39]. Male monarchs that sequestered the highest concentrations of cardenolides had the most

conspicuous warning signals when oxidative damage was lowest, and reduced their warning

signal conspicuousness as oxidative damage increased, while those that sequestered the

lowest levels of cardenolides invested equally in warning signal conspicuous with increasing

oxidative damage. Our results suggest that highly toxic prey reduce investment in signals

because they cannot bear the oxidative cost of investing highly in both traits. This is the first

evidence to support Blount et al's [16] resource allocation trade-off model, and shows that

variable access to resources can result in differential costs of signalling in monarch butterflies

[53].

Some authors [54] have rejected the idea that there can be a mechanistic link between

warning signals and chemical defences that involve intrinsic costs of production or

maintenance. However, condition-dependent expression of colourful signals can arise when

the underlying physiological pathways that produce colouration are dependent on the same

core cellular processes [16, 55]. We found stable levels of antioxidant defences (superoxidase

dismutase and total antioxidant capacity) but increased oxidative damage was associated with

reduced warning signal conspicuousness, which suggests that the monarchs that are better

able to control pro-oxidant levels can sequester more cardenolides and still invest in warning

signals [56]. Similar patterns are observed in *Aristolochia*-feeding pipevine swallowtails

butterflies (*Battus philenor*) that store nitrophenanthrenes and have significantly higher tissue

levels of carotenoids than related species which mimic them in coloration but do not store

toxins [57]. Similarly, the rank order of body concentrations of carotenoids in three species of

lepidoptera matches the rank order of dietary exposure to the prooxidant toxin

furanocoumarin (8-MOP) [Papilio polyxenes > Spodoptera eridania > Trichoplusia ni; 18].

Antioxidants are involved in detoxification processes [58] as well as pigment synthesis

pathways directly [59], or as cofactors of enzymes [60]. Our results allow us to speculate that

antioxidant availability has a role in the biochemistry underlying the variations in warning

colouration and toxicity in aposematic animals. However, warning colours are usually

regulated by more than one mechanism, and we suggest that this area of research warrants

further biochemical study.

It is likely that oxidative state depends on the combination of genetic, environmental and

gene by environment interactions that determine an individual's condition. Monarchs show

patterns of local adaptation to their hostplants [based on larval growth rate 39], and also show

GxE interactions in sequestration ability [23] which may reflect either a lack of evolutionary

history with different species of *Asclepias*, or a physiological trade-off in sequestration

ability. Our results suggest that populations of monarchs that are locally adapted to high

cardenolide milkweeds could be less subject to oxidative damage, either because they have

evolved higher antioxidant efficiency, or because they are able to accumulate more

antioxidants to provide enhanced antioxidant protection. Further research on this topic could

have interesting implications for understanding host shifts and range expansions as well as

our understanding of automimicry (the occurrence of palatable 'cheaters' in a chemically

defended population). Whether automimicry in monarchs arises directly from a deficiency in

an individual's capacity to cope with reactive oxygen species warrants further study [61].

The resource competition model [16] does not investigate or predict sex differences in

resource limitation. However, Blount et al [62] found that ladybird females (the larger sex)

are more susceptible to resource limitation than males, and hence more likely to signal

honestly. In this study we found costs of signalling in male monarchs. Males are slightly

larger than females and are the more active sex; this behavioural difference may increase

their probability of detection and risk of predation, and also increase their sensitivity to

oxidative damage. If predators are sensitive to the combined qualities of toxins and displays

when they attack prey, this could mean they avoid highly conspicuous prey and reject highly

defended but less conspicuous prey, and instead predate moderately defended and

conspicuous individuals, this could maintain the signal variability we observe in males and in

other aposematic prey. The relationship between visual signal strength, chemical defence, and

predator response warrants further study.

Although we did not measure cardenolide content in milkweed plants, our results confirm

previous research showing that the concentration of cardenolides in monarch butterflies

varies depending on host-plant chemistry [63, 64], and follows a standard approach to using

the natural between species and population variability in plant toxicity to test for costs of

sequestration [21, 37, 51]. However, it is methodologically difficult to separate the costs of

sequestration from other potential effects of plant allelochemicals [65]. Future work could

focus on comparative phytochemical profiles to understand resource availability, and

experiments that control nutritional content while manipulating toxin content.

In conclusion, our results support the proposal that oxidative state can be a key physiological

mechanism that links warning colours to sequestration costs [16], and that the contribution of

pigments to antioxidant activity for maintaining redox homeostasis is important [61]. Our

results show that specialist herbivores must balance the benefits of plant secondary

compounds for sequestration with the burden that these same compounds impose [though see

51, 66]. Future research should examine the possibly of natural selection for oxidative

capacity, and whether aposematism evolves in species that have high antioxidant capacity.

These studies could provide new theories for understanding the evolution of aposematism

when coloration and toxicity do not coevolve. Documenting the costs associated with using

secondary defences in natural systems is important for our understanding of the ecology and

evolution of aposematism.

Acknowledgements

We thank Prof. Linda Fink for her support in drafting the manuscript, Prof. Innes Cuthill for

the MATLAB code for visual modelling, and Prof. Anurag Agrawal for leaves of *Ascelpias*

syriaca. JDB was supported by a Royal Society University Research Fellowship. HMR was

supported by a Junior Research Fellowship from Churchill College, Cambridge, and by

funding from the Max Planck Society.

Author contributions

Jonathan D. Blount conceptualization, methodology, resources, data curation, writing -

original draft, project administration, funding acquisition.

Hannah M. Rowland methodology, formal analysis, investigation, data curation, writing -

original draft, visualization.

Christopher Mitchell investigation, data curation.

Michael P. Speed conceptualization, methodology, writing - review & editing.

Graeme D. Ruxton conceptualization, methodology, writing - review & editing.

John A. Endler conceptualization, writing - review & editing, visualization.

Lincoln P. Brower conceptualization, methodology, investigation, resources.

References

[1] Blum, M.S. 1981 *Chemical Defenses of Arthropods*. New York, Academic Press.

[2] Ruxton, G.D., Allen, W.L., Sherratt, T.N. & Speed, M.P. 2018 *Avoiding Attack. The*
Evolutionary Ecology of Crypsis, Aposematism, and Mimicry. Oxford, UK., Oxford
University Press.

[3] Summers, K. & and Clough, M.E. 2001 The evolution of coloration and toxicity in the
poison frog family (Dendrobatidae). *Proceedings of the National Academy of Sciences* **98**,
6227-6232.

[4] Santos, J.C. & Cannatella, D.C. 2011 Phenotypic integration emerges from aposematism
and scale in poison frogs. *Proceedings of the National Academy of Sciences* **108**, 6175-6180.
(doi:10.1073/pnas.1010952108).

[5] Maan, M.E. & Cummings, M.E. 2009 Sexual dimorphism and directional sexual selection
on aposematic signals in a poison frog. *Proceedings of the National Academy of Sciences*
106, 19072-19077. (doi:10.1073/pnas.0903327106).

[6] Cortesi, F. & Cheney, K.L. 2010 Conspicuousness is correlated with toxicity in marine
opisthobranchs. *Journal of Evolutionary Biology* **23**, 1509-1518. (doi:doi:10.1111/j.1420-
9101.2010.02018.x).

[7] Bezzerides, A., L., McGraw, K.J., Parker, R.S. & Husseini, J. 2007 Elytra color as a

signal of chemical defense in the Asian ladybird beetle Harmonia axyridis. *Behavioral*

Ecology and Sociobiology **61**, 1401-1408. (doi:10.1007/s00265-007-0371-9).

[8] Winters, A.E., Stevens, M., Mitchell, C., Blomberg, S.P. & Blount, J.D. 2014 Maternal

effects and warning signal honesty in eggs and offspring of an aposematic ladybird beetle.

Functional Ecology **28**, 1187-1196. (doi:doi:10.1111/1365-2435.12266).

[9] Vidal-Cordero, J.M., Moreno-Rueda, G., López-Orta, A., Marfil-Daza, C., Ros-Santaella,

J.L. & Ortiz-Sánchez, F.J. 2012 Brighter-colored paper wasps (Polistes dominula) have larger

poison glands. *Frontiers in Zoology* **9**, 20. (doi:10.1186/1742-9994-9-20).

[10] Summers, K., Speed, M.P., Blount, J.D. & Stuckert, A.M.M. 2015 Are aposematic

signals honest? A review. *Journal of Evolutionary Biology*, n/a-n/a. (doi:10.1111/jeb.12676).

[11] Leimar, O., Enquist, M. & Sillentullberg, B. 1986 Evolutionary Stability of Aposematic

Coloration and Prey Unprofitability - a Theoretical-Analysis. *American Naturalist* **128**, 469-

490.

[12] Speed, M.P. & Ruxton, G.D. 2007 How bright and how nasty: Explaining diversity in

warning signal strength. *Evolution* **61**, 623-635.

[13] Ruxton, G.D., Speed, M.P. & Broom, M. 2009 Identifying the ecological conditions that

select for intermediate levels of aposematic signalling. *Evolutionary Ecology* **23**, 491-501.

(doi:10.1007/s10682-008-9247-3).

[14] Darst, C.R., Cummings, M.E. & Cannatella, D.C. 2006 A mechanism for diversity in

warning signals: Conspicuousness versus toxicity in poison frogs. *Proc. Natl. Acad. Sci. U. S.*

A. **103**, 5852-5857.

[15] Wang, I.J. 2011 Inversely related aposematic traits: reduced conspicuousness evolves

with increased toxicity in a polymorphic poison-dart frog. *Evolution* **65**, 1637-1649.

(doi:doi:10.1111/j.1558-5646.2011.01257.x).

[16] Blount, J.D., Speed, M.P., Ruxton, G.D. & Stephens, P.A. 2009 *Warning displays may*

function as honest signals of toxicity 871-877 p.

[17] Holloway, G.J., de Jong, P.W., Brakefield, P.M. & de Vos, H. (1991) Chemical defence

in ladybird beetles (Coccinellidae). I. Distribution of coccinelline and individual variation in

defence in 7-spot ladybirds (Coccinella septempunctata). *Chemoecology,* **2,** 7-14.

[18] Ahmad, S. 1992 Biochemical defence of pro-oxidant plant allelochemicals by

herbivorous insects. . *Biochemical Systematics and Ecology* **20**, 269-296.

[19] Holen, O.H. & Svennungsen, T.O. 2012 Aposematism and the Handicap Principle. *The*

American Naturalist **180**, 629-641. (doi:10.1086/667890).

[20] Malcolm, S.B. & Brower, L.P. 1989 Evolutionary and Ecological Implications of

Cardenolide Sequestration in the Monarch Butterfly. *Experientia* **45**, 284-295.

[21] Agrawal, A.A., Böröczky, K., Haribal, M., Hastings, A.P., White, R.A., Jiang, R. &

Duplais, C. 2021 Cardenolides, toxicity, and the costs of sequestration in the coevolutionary

interaction between monarchs and milkweeds. *Proceedings of the National Academy of*

Sciences **118**, e2024463118. (doi:10.1073/pnas.2024463118).

[22] Tao, L., Berns, A.R. & Hunter, M.D. 2014 Why does a good thing become too much?

Interactions between foliar nutrients and toxins determine performance of an insect herbivore.

Functional Ecology **28**, 190-196. (doi:https://doi.org/10.1111/1365-2435.12163).

[23] Freedman, M.G., Choquette, S.-L., Ramírez, S.R., Strauss, S.Y., Hunter, M.D. &

Vannette, R.L. 2021 Population-specific patterns of toxin sequestration in monarch butterflies

from around the world. *bioRxiv,* 2021.2010.2015.464593. (doi:10.1101/2021.10.15.464593).

[24] Brower, L.P. 1969 Ecological chemistry. *Scientific American* **220**, 22-29.

[25] Brower, L.P. 1972 Ecological Chemistry of Palatability Cardiac Glycoside Spectrum in

Monarch Butterflies, Danaus-Plexippus, and Asclepias Milkweeds. *American Zoologist* **12**,

712-713.

[26] Züst, T., Petschenka, G., Hastings, A.P. & Agrawal, A.A. 2019 Toxicity of Milkweed
Leaves and Latex: Chromatographic Quantification Versus Biological Activity of
Cardenolides in 16 Asclepias Species. *J Chem Ecol* **45**, 50-60. (doi:10.1007/s10886-018-
1040-3).

[27] Tao, L., Hoang, K.M., Hunter, M.D. & de Roode, J.C. 2016 Fitness costs of animal
medication: antiparasitic plant chemicals reduce fitness of monarch butterfly hosts. *J Anim*
Ecol **85**, 1246-1254. (doi:10.1111/1365-2656.12558).

[28] Groen, S.C., LaPlante, E.R., Alexandre, N.M., Agrawal, A.A., Dobler, S. & Whiteman,
N.K. 2017 Multidrug transporters and organic anion transporting polypeptides protect insects
against the toxic effects of cardenolides. *Insect Biochemistry and Molecular Biology* **81**, 51–
61.

[29] Seiber, J.N., Tuskes, P.M., Brower, L.P. & Nelson, C.J. 1980 Pharmacodynamics of
Some Individual Milkweed Cardenolides Fed to Larvae of the Monarch Butterfly (Danaus-
Plexippus L). *Journal of Chemical Ecology* **6**, 321-339.

[30] Roeske, C.N., Seiber, J.N., Brower, L.P. & Moffitt, C.M. 1976 Milkweed Cardenolides
and Their Comparative Processing by Monarch Butterflies (Danaus plexippus L.). In
Biochemical Interaction Between Plants and Insects (eds. J.W. Wallace & R.L. Mansell), pp.
93-167. Boston, MA, Springer US.

[31] Faldyn, M.J., Hunter, M.D. & Elderd, B.D. 2018 Climate change and an invasive,
tropical milkweed: an ecological trap for monarch butterflies. *Ecology* **99**, 1031-1038.
(doi:https://doi.org/10.1002/ecy.2198).

[32] Agrawal, A.A. 2017 *Monarchs and Milkweed: A Migrating Butterfly, a Poisonous Plant,*
and their Remarkable Story of Coevolution Princeton, NJ., Princeton University Press.

[33] Geest, E.A., Wolfenbarger, L.L. & McCarty, J.P. 2019 Recruitment, survival, and
parasitism of monarch butterflies (Danaus plexippus) in milkweed gardens and conservation
areas. *Journal of Insect Conservation* **23**, 211-224. (doi:10.1007/s10841-018-0102-8).

[34] Malcolm, S.B., Cockrell, B.J. & Brower, L.P. 1989 Cardenolide Fingerprint of Monarch

Butterflies Reared on Common Milkweed, Asclepias-Syriaca-L. *Journal of Chemical*

Ecology **15**, 819-853.

[35] Petschenka, G. & Agrawal, A.A. 2015 Milkweed butterfly resistance to plant toxins is

linked to sequestration, not coping with a toxic diet. *Proceedings. Biological sciences / The*

Royal Society **282**, 20151865. (doi:10.1098/rspb.2015.1865).

[36] Brower, L.P. 1970 Plant poisons in a terrestrial food chain and implications for mimicry

theory. . In *Biochemical Coevolution (Proceedings of the Twenty-Ninth Annual Biology*

Colloquium). (ed. K.L. Chambers), pp. 69-82. Corvallis, Oregon., Oregon State University

Press, .

[37] Soule, A.J., Decker, L.E. & Hunter, M.D. 2020 Effects of diet and temperature on

monarch butterfly wing morphology and flight ability. *Journal of Insect Conservation* **24**,

961-975. (doi:10.1007/s10841-020-00267-7).

[38] Woods, E.C., Hastings, A.P., Turley, N.E., Heard, S.B. & Agrawal, A.A. 2012 Adaptive

geographical clines in the growth and defense of a native plant. *Ecological Monographs* **82**,

149-168. (doi:https://doi.org/10.1890/11-1446.1).

[39] Freedman, M.G., Jason, C., Ramírez, S.R. & Strauss, S.Y. 2020 Host plant adaptation

during contemporary range expansion in the monarch butterfly. *Evolution* **74**, 377-391.

(doi:https://doi.org/10.1111/evo.13914).

[40] Pocius, V.M., Debinski, D.M., Pleasants, J.M., Bidne, K.G., Hellmich, R.L. & Brower,

L.P. 2017 Milkweed Matters: Monarch Butterfly (Lepidoptera: Nymphalidae) Survival and

Development on Nine Midwestern Milkweed Species. *Environ. Entomol.* **46**, 1098-1105.

(doi:10.1093/ee/nvx137).

[41] Malcolm, S.B. & Zalucki, M.P. 1996 Milkweed latex and cardenolide induction may

resolve the lethal plant defence paradox. In *Proceedings of the 9th International Symposium*

on Insect-Plant Relationships (eds. E. Städler, M. Rowell-Rahier & R. Bauer), pp. 193-196.

Dordrecht, Springer Netherlands.

[42] Rasmann, S., Agrawal, A.A., Cook, S.C. & Erwin, A.C. 2009 Cardenolides, induced
responses, and interactions between above- and belowground herbivores of milkweed
(Asclepias spp.). *Ecology* **90**, 2393-2404. (doi:https://doi.org/10.1890/08-1895.1).

[43] Fink, L.S., Brower, L.P., Waide, R.B. & Spitzer, P.R. 1983 Overwintering Monarch
Butterflies as Food for Insectivorous Birds in Mexico. *Biotropica* **15**, 151-153.

[44] Vorobyev, M. & Osorio, D. 1998 Receptor noise as a determinant of colour thresholds.
Proc. R. Soc. Lond. Ser. B-Biol. Sci. **265**, 351-358.

[45] Keyser, A.J. & Hill, G.E. 2000 Structurally based plumage coloration is an honest signal
of quality in male blue grosbeaks. *Behavioral Ecology* **11**, 202-209.

[46] Hart, N.S., Partridge, J.C., Cuthill, I.C. & Bennett, A.T.D. 2000 Visual pigments, oil
droplets, ocular media and cone photoreceptor distribution in two species of passerine bird:
the blue tit (Parus caeruleus L.) and the blackbird (Turdus merula L.). *J. Comp. Physiol. A-*
Sens. Neural Behav. Physiol. **186**, 375-387.

[47] Stevens, M. & Merilaita, S. 2011 *Animal Camouflage: Mechanisms and Function.*
Cambridge, UK, Cambridge University Press.

[48] Osorio, D. & Vorobyev, M. 2005 Photoreceptor spectral sensitivities in terrestrial
animals: adaptations for luminance and colour vision. *Proceedings of the Royal Society of*
London, Series B **272**, 1745-1752.

[49] Fox, J. & Weisberg, S. 2019 *An R Companion to Applied Regression*. Third edition. ed,
Sage, Thousand Oaks CA.

[50] Bartoń, K. 2020 MuMIn: Multi-Model Inference. (

[51] Erickson, J.M. 1973 The Utilization of Various Asclepias Species by Larvae
of the Monarch Butterfly, Danaus Plexippus. *Psyche* **80**, 028693.
(doi:10.1155/1973/28693).

[52] Zalucki, M.P., Brower, L.P. & Alonso, A. 2001 Detrimental effects of latex and cardiac

glycosides on survival and growth of first-instar monarch butterfly larvae Danaus plexippus

feeding on the sandhill milkweed Asclepias humistrata. *Ecological Entomology* **26**, 212-224.

[53] Zahavi, A. & Zahavi, A. 1997 *The handicap principle: a missing piece of Darwin's*

puzzle. Oxford, UK, Oxford University Press.

[54] Guilford, T. & Dawkins, M.S. 1993 Are warning colors handicaps? *Evolution* **47**, 400.

[55] Hill, G.E. 2011 Condition-dependent traits as signals of the functionality of vital cellular

processes. *Ecology Letters* **14**, 625-634. (doi:https://doi.org/10.1111/j.1461-

0248.2011.01622.x).

[56] Costantini, D., Fanfani, A. & Dell'Omo, G. 2008 Effects of corticosteroids on oxidative

damage and circulating carotenoids in captive adult kestrels (Falco tinnunculus). *Journal of*

Comparative Physiology B **178**, 829-835. (doi:10.1007/s00360-008-0270-z).

[57] Harborne, J.B. & Williams, C.A. 2001 Anthocyanins and other flavonoids. *Natural*

product reports, **18**, 310-333.

[58] Enayati, A.A., Ranson, H. & Hemingway, J. (2005) Insect glutathione transferases and

insecticide resistance. *Insect Mol Biol,* **14,** 3-8.

[59] Ito, S. & Prota, G. (1977) A facile one-step synthesis of cysteinyldopas using mushroom

tyrosinase. *Experientia,* **33,** 1118-1119.

[60] Shamim, G., Ranjan, K.S., Pandey, M.D. & Ramani, R. (2014) Biochemistry and

biosynthesis of insect pigments. *EJE,* **111,** 149-164.

[61] Costantini, D. & Verhulst, S. 2009 Does high antioxidant capacity indicate low oxidative

stress? *Functional Ecology* **23**, 506-509. (doi:https://doi.org/10.1111/j.1365-

2435.2009.01546.x).

[62] Blount, J.D., Rowland, H.M., Drijfhout, F.P., Endler, J.A., Inger, R., Sloggett, J.J.,

Hurst, G.D.D., Hodgson, D.J. & Speed, M.P. 2012 How the ladybird got its spots: effects of

resource limitation on the honesty of aposematic signals. *Functional Ecology* **26**, 334-342.

(doi:10.1111/j.1365-2435.2012.01961.x).

[63] Jones, P.L., Petschenka, G., Flacht, L. & Agrawal, A.A. 2019 Cardenolide Intake,

Sequestration, and Excretion by the Monarch Butterfly along Gradients of Plant Toxicity and

Larval Ontogeny. *Journal of Chemical Ecology* **45**, 264-277. (doi:10.1007/s10886-019-

01055-7).

[64] Brower, L.P., Nelson, C.J., Seiber, J.N., Fink, L.S. & Bond, C. 1988 Exaptation as an

alternative to coevolution in the cardenolide-based chemical defense of monarch butterflies

(*Danaus plexippus L.*) against avian predators. In *Chemical mediation of coevolution* (ed.

K.C. Spencer). London, Harcourt Brace Jovanovich.

[65] Bowers, M.D. 1992 The evolution of unpalatability and the cost of chemical defense in

insects. . In *Insect chemical ecology: an evolutionary approach. Chapman & Hall, New York,*

New York, USA. (eds. B.D. Roitberg & M.B. Isman), pp. 216–244.

[66] Petschenka, G. & Agrawal, A.A. 2015 Milkweed butterfly resistance to plant toxins is

linked to sequestration, not coping with a toxic diet. *Proceedings of the Royal Society B:*

Biological Sciences **282**. (doi:10.1098/rspb.2015.1865).

Figure 1. cardenolide sequestration by monarchs feeding on *Ascelpias tuberosa* (AT), *A.*

Incarnata (AI), *A. Syriaca* (AS), and *A. curassavica* (AC); a) total cardenolide (µg total

cardenolides per 0.1 g tissue) measured in the wings and bodies, b) body cardenolide content,

c) wing cardenolide content. Boxplots show the median, interquartile range, and the whiskers

represent the largest and smallest value within 1.5 times the 25th and 75th percentile. Outliers

are represented by small black dots. The mean cardenolide concentration is represented by

the larger black dot.

Figure 2. Total cardenolide concentration for males (filled triangle) and females (open circle)

and Malondialdehyde (MDA) concentration. Line is the smoothed conditional mean with

95% confidence intervals.

Figure 3. Three-dimensional surface (perspective) plot for the response surface predicting

the relationship between the conspicuousness of the male monarch's forewing, oxidative

damage measured as malondialdehyde (MDA), and total cardenolide concentration.

Figure 1a, b, c.

Figure 2

Figure 3