1

The p97-UBXD8 complex modulates ER-Mitochondria contact sites by modulating membrane

lipid saturation and composition

Rakesh Ganji[1], Joao A. Paulo[2], Yuecheng Xi[3], Ian Kline[3], Jiang Zhu[4], Christoph S. Clemen[5],
Conrad C. Weihl[4], John G. Purdy[3], Steve P. Gygi[2], and Malavika Raman[1#]

[1] Department of Developmental Molecular and Chemical Biology, Tufts University School of

Medicine, Boston MA

[2] Department of Cell Biology Harvard Medical School, Boston MA

[3] Department of Immunobiology, BIO5 Institute, University of Arizona School College of Medicine,

Tucson Arizona

[4] Department of Neurology, Washington University School of Medicine, Saint Louis,

Missouri

[5] Institute of Aerospace Medicine, German Aerospace Center, Cologne, Germany; Center for

Physiology and Pathophysiology, Institute of Vegetative Physiology, Medical Faculty, University

of Cologne, Cologne, Germany.

Address correspondence to Malavika Raman, malavika.raman@tufts.edu

Abstract

The intimate association between the endoplasmic reticulum (ER) and mitochondrial membranes at ER-mitochondria contact sites serves as a platform for several critical cellular processes, in particular lipid synthesis. Enzymes involved in lipid biosynthesis are enriched at contacts and membrane lipid composition at contacts is distinct relative to surrounding membranes. How contacts are remodeled and the subsequent biological consequences of altered contacts such as perturbed lipid metabolism remains poorly understood. Here we show that the p97 AAA-ATPase and its ER-tethered ubiquitin-X domain adaptor 8 (UBXD8) regulate the prevalence of ER-mitochondria contacts. The p97-UBXD8 complex localizes to contacts and loss of this complex increases contacts in a manner that is dependent on p97 catalytic activity. Quantitative proteomics of purified contacts demonstrates alterations in proteins regulating lipid metabolism upon loss of UBXD8. Furthermore, lipidomics studies indicate significant changes in distinct lipid species in UBXD8 knockout cells. We show that loss of p97-UBXD8 results in perturbed contacts due to an increase in membrane lipid saturation via SREBP1 and the lipid desaturase SCD1. Aberrant contacts in p97-UBXD8 loss of function cells can be rescued by supplementation with unsaturated fatty acids or overexpression of SCD1. Perturbation of contacts and inherent lipid synthesis is emerging as a hallmark to a variety of human disorders such as neurodegeneration. Notably, we find that the SREBP1-SCD1 pathway is negatively impacted in the brains of mice with p97 mutations that cause neurodegeneration. Our results suggest that contacts are exquisitely sensitive to alterations to membrane lipid composition and saturation in a manner that is dependent on p97-UBXD8.

Main

Contact sites between the ER and mitochondria allow for compartmentalization of biosynthetic reactions such as lipid synthesis, calcium transport and apoptosis among others[1-3]. ER-mitochondria contact sites (henceforth referred to as contacts unless specified otherwise) form when the membrane of these organelles come into close apposition (observed to be between 5 – 100 nm) without fusion[2,4]. Contacts are stabilized by the interaction between tethering proteins that reside in opposing membranes; thus, the ability to regulate transient interactions between these tethers (through their abundance or post-translational modifications) enables the rapid formation and dissolution of contacts to meet cellular needs[5-9].

Regulated protein degradation via the ubiquitin proteasome system is an efficient means to modulate contacts as numerous ubiquitin-reliant protein quality control mechanisms surrounding the ER and mitochondria can be co-opted to modulate the contact site proteome[10,11]. The p97 AAA-ATPase is an abundant, evolutionarily conserved, ubiquitin-selective unfoldase that regulates multiple protein quality control pathways surrounding the ER and mitochondria. p97 is ideally positioned to regulate contacts by mediating the extraction and degradation of membrane-embedded tethers or contact resident proteins[12-14]. Importantly, specificity in p97-regulated pathways occurs via association with numerous adaptors that recruit p97 to specific substrates[15,16]. We isolated contacts (also known as mitochondria-associated membranes, MAM) from HEK-293T cells and probed for p97 and select adaptor proteins. Percoll gradient centrifugation of crude mitochondria releases associated ER membranes allowing for the purification of ER-mitochondria contacts[17]. Using a known contact site-enriched protein as a marker (fatty acid coenzyme A ligase 4, FACL4) as well as markers for mitochondria (TOMM20) and ER (SEC61β), we found that the ER-localized p97 adaptors UBXD8 and UBXD2 (but not cytosolic UBXN1) are enriched at contacts and that p97 is also present in these fractions (Fig. 1a). To determine the role of p97 complexes at contacts we used a split luciferase reporter wherein the N-terminal fragment of luciferase is targeted to mitochondria and the C-terminal

fragment to the ER using established targeting sequences[18]. Functional luciferase activity is

reconstituted when the two organelles establish close range contacts and luciferase activity can

be measured using the substrate, Enduren. We verified the functionality of the reporter system in

wildtype cells by over-expressing receptor accessory protein 1 (REEP1)[18], a contact tether and

found a robust increase in contacts (Supplementary Fig.1a). To determine what role if any p97-

adaptor complexes may have at contacts, cells were transfected with split luciferase cDNAs and

siRNAs and luminescence was measured. Loss of p97 and UBXD8 resulted in an increase in

contacts, whereas depletion of UBXD2 or another p97 adaptor UBXD7 had no impact (Fig. 1b, c

and Supplementary Fig.1b). Hence even though UBXD2 localizes to contacts it does not regulate

them and we focus on the role of UBXD8 for the remainder of the study. To verify the specificity

of the phenotype, we expressed wildtype p97 or UBXD8 siRNA-resistant cDNAs and were able

to rescue the phenotype (Fig. 1b, c). However, individual ATP catalytic site mutants in p97 were

unable to rescue (Fig. 1c). To assess the role of individual domains in UBXD8, we expressed

point mutants in the ubiquitin associated (UBA) or ubiquitin-X (UBX) domains that serve to bind

ubiquitin and p97 respectively, as well as deletion of the UAS domain that has been reported to

mediate the oligomerization of UBXD8[19]. Loss of these functional domains in UBXD8 prevented

rescue of increased contacts (Fig. 1b, Supplementary Fig. 1c). Two additional assays were used

to verify the role of p97-UBXD8 in regulating contacts. We measured contacts by measuring the

extent of co-localization between the ER and mitochondria by confocal microscopy

(Supplementary Fig. 1d, e), and we performed transmission EM (TEM) studies in wildtype and

UBXD8 knockout (KO) cells generated by CRISPR-Cas9 gene editing (Fig. 1d-f). UBXD8 KO

cells had a significant increase in ER tubules closely apposed to mitochondria (Fig. 1e, f and

Supplementary Fig. 1e). No defects in overall mitochondria number or morphology were apparent

in UBXD8 KO cells (Supplementary Fig. 1f).

The p97-UBXD8 complex is critical for maintaining ER quality control by regulating ERAD and

loss of this complex can cause ER stress due to deficits in ERAD[20]. To determine whether altered

contacts observed upon depletion of p97-UBXD8 is due to increased ER stress, we treated cells

with ER stressors tunicamycin and thapsigargin which cause protein misfolding in the ER.

However, no impact on contacts was observed under these conditions (Supplementary Fig. 2a).

Furthermore, we probed for the ER chaperone BiP which is induced upon ER stress and found

no difference in BiP levels between wildtype and UBXD8 KO cells (Supplementary Fig.2b).

Together with our findings suggest that the p97-UBXD8 complex has a novel role in regulating

contacts independent of its role in ERAD.

To identify pathways altered in UBXD8 KO cells that may contribute to the contact site

defect, we isolated contacts (MAMs) from triplicate wildtype and UBXD8 KO HEK-293T cells by

biochemical fractionation and performed multiplexed, quantitative proteomics on the post-nuclear

supernatant and MAM fractions using tandem mass tags (TMT) (Fig. 2a, b and Supplementary

Fig. 3a-f, Supplementary Table 1)[21,22]. We used two filtering criteria for downstream analysis: $|\log_2$

WT:KO ratio$|$ > 1.0 and $-\log_{10}$ p value >1.5. The abundance of 23 proteins was enriched and 28

proteins was depleted in the MAM fraction of UBXD8 KO cells out of a total of 4499 quantified

(Fig. 2b and Supplementary Fig. 3a). Putative contact site proteins identified in previous studies

were present in our dataset validating our approach (Fig. 2b and Supplementary Fig. 3b,

Supplementary Table 1) [18,23-25]. Furthermore, we identified significant enrichment of known p97-

UBXD8 substrates such as squalene monooxygenase (SQLE), and HMG-CoA reductase

(HMGCR) in UBXD8 KO cells (Fig. 2b and Supplementary Fig. 3g, h). Interestingly, proteins

involved in lipid or cholesterol metabolism and lysosome function were also enriched in the

UBXD8 KO contact proteome ($\log_2$ WT:KO ratio < -0.65 and $-\log_{10}$ p value >1.5) (Fig. 2b-d and

Supplementary Fig.3c). In summary, our quantitative proteomic studies of the contact proteome

in UBXD8 KO cells suggests that perturbation in the abundance of numerous enzymes linked to

lipid biosynthesis may underlie the dysregulation in contacts.

To better understand how cellular lipid metabolism may be impacted by loss of UBXD8

we measured the lipidome of wildtype and UBXD8 KO cells. We identified and quantitatively

measured the relative levels of phospholipids (PLs) using liquid-chromatography high-resolution

tandem mass spectrometry (LC-MS/MS). Our analyses examined the major classes of PLs found

in membranes (and synthesized at contacts) including phosphatidylcholine (PC),

phosphatidylethanolamine (PE), phosphatidylserine (PS), and phosphatidylinositol (PI). We also

examined one-tailed lysophospholipids, which are metabolic byproducts of phospholipids (e.g.,

LPC, LPE, LPS, and LPI). We determined the concentration of 151 PLs in UBXD8 KO cells

relative to their concentration in wildtype cells. In general, most PLs were more abundant in KO

cells (Fig. 2e, f and Supplementary Fig. 4a-c, Supplementary Table 2). Approximately two-thirds

of the two-tailed PC and PE and one-tailed LPC and LPE species were ≥2-fold more abundant in

UBXD8 KO cells (Fig. 2e, f Supplementary Fig. 4a-d). The PC species that increased the most in

the UBXD8 KO cells contained one or two double bonds among the fatty acyl tails (i.e., PC (44:1),

PC (44:2), PC (46:1), PC (46:2), PC (48:2), PC (50:2), and PC (52:2)). MS/MS identification of

the individual tails revealed that each of these lipids contained only saturated or monounsaturated

fatty acids ranging in size from C16:0 to C32:1 and that lipids with one or fewer double bonds in

each tail were increased the most by the loss of UBXD8 (Supplementary Fig. 4f and

Supplementary Table 2). Phospholipids are synthesized via a diacylglycerol (DG) intermediate

and DGs are metabolized to generate PLs and TGs. We therefore extended our lipidomics studies

to measure DG and two additional neutral lipids: triacylglycerol (TG) and cholesteryl esters (CE)

that are major components of lipid droplets whose biogenesis at the ER is regulated by UBXD8[26].

The relative concentration of most DGs were altered by less than 2-fold (Supplementary Fig. 4e).

Of the ten DG species increased by ≥2-fold in UBXD8 KO cells, nine contained only saturated or

monounsaturated tails ranging in size from C14:0 to C32:0. Similar to PC and PE, most TG

species were ≥2-fold more abundant in KO cells relative to control cells (Supplementary Fig. 4e).

The TG species whose abundance was enhanced the most by the loss of UBXD8 were also

enriched in saturated and monounsaturated fatty acyl tails from C14:0 to C32:1. In contrast, most

CE species was unaltered or slightly depleted by the loss of UBXD8 (Supplementary Fig. 4e). In

summary, loss of UBXD8 shifts the lipidome to have a greater abundance of PC, PE, DG, and

TG with saturated and monounsaturated fatty acyl tails demonstrating that UBXD8 is necessary

for regulating lipid concentrations. The synthesis of these phospholipids occurs at contacts and

their altered abundance in UBXD8 KO cells may contribute to defects in contacts[27].

The significantly altered lipidome and related alterations in lipid biosynthetic enzymes

prompted us to ask if these changes stem from the regulation of the sterol regulatory element-

binding proteins 1 and 2 (SREBP1/2) pathway. p97-UBXD8 mediates the activation of ER-

localized SREBPs by mediating ubiquitin-dependent membrane extraction and degradation of the

SREBP negative regulator insulin induced gene 1 (INSIG1), in a sterol or fatty acid-dependent

manner (Supplementary Fig. 5a)[28,29]. A similar pathway for lipid sensing in *S.cerevisiae*, requires

Cdc48p and Ubx2p (orthologs of p97 and UBXD8 respectively),[30] and yeast lacking Ubx2p have

more saturated cellular membranes due to loss of transcriptional activation of Δ^9 desaturase

ole1[30]. We found that loss of p97 or UBXD8 resulted in a significant loss of SREBP1 activation

and accumulation of the inactive ER-tethered form concomitant with stabilization of INSIG1 (Fig.

3a-d). Lipid desaturases are transcriptional targets of SREBP1 and were decreased in abundance

at the transcript level in UBXD8 KO cells (Supplementary Fig.5b). This parallels the decreased

protein abundance of the best characterized desaturases, SCD1 (Δ^9) and FADS1 (Δ^5) in UBXD8

KO or p97 depleted cells (Fig. 3a-d). In contrast, UBXD8 loss did not significantly impact SREBP2

activation or its downstream targets although this may be cell type or context specific (Fig.3a, b

and Supplementary Fig.5c, d). Previous studies have reported that contacts have lipid raft-like

properties and are enriched in cholesterol and sphingolipids[31], thus the localization of cholesterol-

sensing proteins to these sub-domains may be advantageous for rapid pathway activation. We

find that SREBP1 and SCD1 are enriched at contacts isolated by biochemical fractionation (Fig.

3e). Notably, an increase in ER-tethered SREBP1 and a decrease in SCD1 were apparent in the

MAM fractions of UBXD8 KO cells (Fig. 3f). Furthermore, over-expression of mature SREBP1a

and c and to a lesser extent SREBP2, is sufficient to rescue contacts in p97-UBXD8 depleted

cells (Supplementary Fig. 5e, f). Hence, defective SREBP1 activation underlies increased

contacts in p97-UBXD8 depleted cells.

The decrease in the abundance of lipid desaturases and our finding that UBXD8 KO cells

have increased phospholipids with saturated or mono-unsaturated tails, prompted us to determine

whether altering membrane lipid saturation perturbed contacts between the ER and mitochondria.

FADS1 isoforms are localized to both ER and mitochondria[32], therefore we focused on ER-

localized SCD1[33] as it was enriched at contacts (Fig. 3e). We treated wild type HEK-293T cells

with the SCD1 inhibitor MF438[34] and found that contacts between the ER and mitochondria

increased in a manner that could be rescued by supplementing cells with monounsaturated oleic

acid, the product of SCD1, (Supplementary Fig. 5g). We next evaluated whether re-expressing

SCD1 in p97 or UBXD8-depleted cells rescued the increased contact site phenotype.

Overexpression of wildtype SCD1 rescued ER-mitochondria contacts to wild type levels in p97-

UBXD8 depleted cells (Fig. 3g, Supplementary Fig. 5h). In contrast, a catalytically inactive version

of SCD1 was unable to rescue the phenotype (Fig. 3g, Supplementary Fig. 5h). Notably,

overexpression of SCD1 catalytic mutant in wild type cells resulted in an increase in contacts

suggesting that the resulting ordered lipid bilayers may impact contacts (Figure 3g). To further

extend these findings, we asked if simply supplementing p97-UBXD8 depleted cells with

unsaturated oleic acid (18:1), a precursor for the generation of polyunsaturated fatty acids in cells

was sufficient to rescue contacts. Indeed, oleic acid but not saturated palmitic acid (16:0) rescued

the contact defect in p97-UBXD8 depleted cells. Strikingly, palmitic acid alone increased contacts

in wildtype cells (Fig. 3h).

Collectively, these results suggested that contacts are exquisitely sensitive to perturbations in

lipid profiles within cellular membranes and that loss of p97-UBXD8 alters membrane lipid

composition and saturation. We evaluated whether loss of p97-UBXD8 impacts lipid saturation

globally within the cell. We used lipid-binding pyrene probes that insert into both disordered

(unsaturated) and ordered (saturated) lipid bilayers throughout the cell and undergo monomer to

excimer formation in a manner dependent on local membrane order. We measured the ratio of

monomer to excimer fluorescence as an indicator of membrane order in cells treated with the p97

inhibitor CB-5083 or in UBXD8 KO cells. Loss of UBXD8 or inhibition of p97 resulted in more

ordered cellular membranes (monomer: excimer ratio <1) compared to controls in two different

cell lines (Fig.4a, Supplementary Fig.6a). Strikingly this phenotype can be reversed by incubating

cells with oleic but not palmitic acid (Fig.4a, Supplementary Fig.6a). Thus, the local degradation

of INSIG1 and subsequent activation of SREBP1 at contacts impacts lipid composition and

saturation throughout the cell but significantly impact contacts due to their reliance on membranes

for association. These findings prompted us to investigate whether the global increase in

membrane order impacts other ER-organelle contacts. Analysis of TEM data from wildtype and

UBXD8 KO cells demonstrated increased contacts between the ER and plasma membrane (Fig.

4b, c). Whether other ER-organelle contacts are similarly perturbed is under investigation.

Notably, UBXD8 KO cells have an increase in membranous whorls containing concentric

membrane layers, reminiscent of multilamellar bodies that function in lipid storage and secretion

(Supplementary Fig. 6b). Interestingly, these membrane rich structures were also identified in

yeast strains lacking Ubx2p [30] and may arise due to imbalances in lipid levels.

Mutations in p97 cause several primarily neurodegenerative protein aggregation disorders.

These include inclusion body myopathy with Paget's disease of the bone and frontotemporal

dementia (IBMPFD, also known as multi-system proteinopathy 1, MSP-1)[35], amyotrophic lateral

sclerosis (ALS)[36,37], Charcot Marie Type IIB[38], among others. We investigated whether p97

disease-associated mutations altered contacts between the ER and mitochondria and perturbed

the SREBP1-SCD1 pathway. We measured contacts in mouse embryonic fibroblasts

heterozygous or homozygous for p97 R155H (a prevalent mutation observed in patients) and

observed that cells with p97 R155H homozygous mutation had a significant increase in contacts

that could be rescued with oleic acid but not palmitic acid (Fig. 4d).

Next, we evaluated the SREBP-SCD1 pathway in the brains of the two distinct p97 mouse

models[39]. A conditional knockout of p97 (p97 cKO) in the cortex and hippocampus has recently

been shown to develop cortical atrophy, neuronal loss and TDP43 inclusions reminiscent of

frontotemporal dementia[39]. We stained for SREBP1 and SCD1 in the CA1 regions of one-month-

old p97 cKO mice before neurodegeneration phenotypes are observed. Strikingly, p97 cKO mice

had a significant decrease of SREBP1 and SCD1 immunoreactivity in the CA1 region compared

to age-matched controls (Fig. 4e, f). To assess whether these defects were also present in a

pathogenic context, we immunoblotted for SREBP1 and SCD1 in brain lysates from 6- and 13-

month-old p97$^{R155C/WT}$ mice[39,40]. SREBP1 processing was significantly diminished at 13 months in

p97$^{R155C/WT}$ mice relative to controls (Fig. 4g, h). Similarly, loss of SCD1 protein levels was

apparent at 6 months and continued to decline at 13 months in p97$^{R155C/WT}$ mice (Fig. 4g, h).

Collectively, our findings suggest that p97 mutations that cause disease may also have underlying

lipid metabolism deficits that could contribute to disease pathology.

Discussion

Here we have identified an unanticipated role for p97 and its ER-tethered adaptor UBXD8

in regulating ER-mitochondria contacts by perturbing membrane composition and fluidity in

multiple cell types. We propose that altered lipid bilayers that arise upon loss of UBXD8 impact

contacts in multiple ways: (1) preventing their dynamic association and disassociation due to loss

of fluidity, and (2) negatively impacting the lateral movement of tethering proteins within

membranes. UBXD8 is unique among p97 adaptors in its evolutionarily conserved and

multifunctional roles in lipid sensing and metabolism[19,26,30,41-43]. Our proteomic and lipidomic data

from UBXD8 KO cells suggests that widespread changes in lipid metabolism is likely due to the

inability to mobilize SREBP1 from the ER. Similar to the yeast *ubx2* deletion phenotype, we find

decreased abundance of lipid desaturases, particularly SCD1 and complementation of p97-

UBXD8 depleted cells with SCD1 rescues contact defects. However, given the significant

changes in lipid profiles in UBXD8 KO cells, it is likely that SCD1 regulation is not the only

mechanism at play. It remains to be determined whether UBXD8 also facilitates the degradation

of contact tethers. We note that UBXD8 is present in pure mitochondrial fractions (Fig. 1a).

Previous studies have observed dual localization of UBXD8 to ER and mitochondria[44,45] and

recently, the yeast ortholog, Ubx2p was reported to localize to the outer mitochondrial membrane

where it associates with the translocon in the outer mitochondrial membrane (TOMM) complex to

clear stalled polypeptides in a p97-dependent manner[46]. Further studies are required to determine

whether mammalian UBXD8 functions in an analogous manner on mitochondria. Aberrant contact

sites are emerging as a common feature in the pathophysiology of a wide spectrum of human

diseases ranging from diabetes to neurodegeneration[47-49]. We find that p97 mutations that cause

proteinopathies also exhibit increased contacts and display significantly decreased levels of

SREBP1 and SCD1. A recent report found that motor neurons from ALS patients with p97

mutations exhibited more contacts between the ER and mitochondria relative to controls[50]. Thus,

altered organelles contacts and downstream lipid synthesis warrants further investigation in p97

associated diseases.

Figure Legends

Fig. 1. Loss of p97 and UBXD8 results in increased ER-mitochondria contacts.

a, Immunoblot of indicated proteins from subcellular fractionation of HEK-293T cells, PNS: post-

nuclear supernatant, MAM: Mitochondria associated membrane (n > 3 biologically independent

samples). **b**, Top: Domain organization of UBXD8 and indicated mutations, Middle: Split

luciferase assay to measure contacts in HEK-293T cells transfected with siRNAs to UBXD8 and

indicated C-HA/FLAG siRNA-resistant rescue constructs, RLU: relative luminescence unit,

Bottom: Immunoblot of indicated proteins. UBA: ubiquitin associated, UAS: upstream activating

sequence, UBX: ubiquitin X ($n > 3$ biologically independent samples). **c**, Top: Domain

organization of p97 and indicated mutations, Middle: Split luciferase assay to measure contacts

in HEK-293T cells transfected with siRNAs to p97 and indicated N-Myc siRNA-resistant rescue

constructs, RLU: relative luminescence unit, Bottom: Immunoblot of indicated proteins ($n \geq 3$

biologically independent samples). **d**, Immunoblot of UBXD8 in CRISPR-Cas9 edited HEK-293T

and HeLa-Flp-IN-T-Rex cells, KO: knockout. **e**, Representative transmission EM micrographs of

wildtype and UBXD8 KO HEK-293T cells illustrating contacts between ER and mitochondria. **f**,

Quantification of contact length between ER and mitochondria in each genotype from (**e**)

(measurements are from $n = 3$ biological replicates with WT = 50 cells from 65 fields and UBXD8

KO = 53 cells from 71 fields). OMM: Outer mitochondrial membrane. Data are means ± SEM (*,

**, ***$P < 0.05$, 0.01, 0.0001 respectively, One-way ANOVA with Tukey's multiple comparison

test). Scale bar, 100 nm.

Fig. 2. Quantitative proteomics and lipidomics identifies a role for UBXD8 in regulating

lipid metabolism at contacts.

a, Schematic of tandem mass tag (TMT) proteomic workflow from wildtype and UBXD8 KO HEK-

293T cells. PNS: post-nuclear supernatant, MAM: mitochondria associated membrane. **b**,

Volcano plot of the ($-\log_{10}$-transformed P value *versus* the $\log_2$-transformed ratio of wildtype/

UBXD8 KO) proteins identified from MAM fractions of HEK-293T cells. $n = 3$ biologically

independent samples for each genotype. P values were determined by empirical Bayesian

statistical methods (adjusted for multiple comparisons) using the *LIMMA* R package; for

parameters, individual P values and q values, see Supplementary Table 1. **c,** Network of

differentially enriched terms shown as clustered functional ontology categories. Each node

represents a functional ontology term enriched in the TMT data (**a, b**) as scored by Metascape[51].

Networks were generated using Cytoscape v3.8.2. Size of node represents number of genes

identified in each term by gene ontology (GO). Grey and Blue donuts represent percent of genes

identified in each GO term in wildtype or UBXD8 KO respectively. Node outline thickness

represents $-\log_{10}$-transformed P value of each term. The inner circle color of each node indicates

the corresponding functional GO cluster. **d**, Bubble plot representing significantly enriched GO

clusters identified from TMT proteomics of MAM fractions in wildtype (blue) or UBXD8 KO (green)

cells (**a-c**). Size of the circle indicates the number of genes identified in each cluster. **e-f**, Relative

levels of Phosphatidylcholine (PC) (**e**) and Phosphatidylethanolamine (PE) (**f**) in HEK-293T WT

and UBXD8 KO cells. PLs were measured by LC-MS/MS following normalization by total protein

amount. Each dot in the plot represents the level of a PL in UBXD8 KO cells relative to its level in

wildtype cells. The dashed line represents a relative level of 1 (e.g., the level in UBXD8 KO cells

is equal to the level in wildtype cells). (n = 3 biologically independent experiments were performed,

each with duplicate samples). Statistical analysis was performed on the log transformed relative

fold change values (UBXD8 KO relative to WT) using independent t tests and Benjamini-

Hochberg correction in R stats package (p-values are listed in Supplemental Table 1).

Fig. 3. Loss of SREBP1 activation and SCD1 expression upon p97-UBXD8 depletion is

responsible to contact defects.

a-d Immunoblot and the corresponding band intensity quantifications of indicated proteins in the

SREBP pathway in wildtype and UBXD8 KO HEK-293T cells (**a & b**), or p97-siRNA depleted cells

(**c & d**). All samples were transfected with INSIG1-HA/FLAG due to lack of reliable antibodies to

the endogenous protein. ($n ≥ 3$ biologically independent samples). **e**, Immunoblot of indicated

SREBP pathway proteins from subcellular fractionation of HEK-293T cells, PNS: post-nuclear

supernatant, MAM: mitochondria associated membrane. (n = 3 biologically independent

samples). **f**, Immunoblot of indicated SREBP pathway proteins from subcellular fractionation of

wildtype and UBXD8 KO HEK-293T cells, MAMs: mitochondria associated membrane. (n = 3

biologically independent samples). Corresponding fold changes (FC: UBXD8 KO vs WT) of

SREBP1 and SCD1 normalized to FACL4 is shown. **g**, Split luciferase assay in HEK293T cells

transfected with indicated siRNAs and wildtype or catalytically dead SCD1. GFP-HA/FLAG was

transfected as a negative control. RLU: relative luminescence unit. ($n \geq 3$ independent biological

replicates). **h**, Split luciferase assay in HEK293T cells transfected with indicated siRNAs and

treated with either monounsaturated oleic acid or saturated palmitic acid. RLU: relative

luminescence unit. ($n \geq 3$ independent biological replicates). Data are means $\pm$ SEM (*, **,

***$P < 0.05$, 0.01, 0.0001 respectively. Paired t test (**b & d**), or One-way ANOVA with Tukey's

multiple comparison test (**g & h**).

Fig. 4. Supplementation of p97-UBXD8 depleted cells with oleic acid rescues contact

defects.

a, Global membrane fluidity was measured using a pyrene-based lipid probe in wildtype and

UBXD8 KO HEK-293T cells. Wildtype cells were also treated with 5 μM of the p97 inhibitor CB-

5083 for 4 hours. Cells were supplemented with indicated concentrations of oleic acid and palmitic

acid for 4 hours. The fold change (Treated$_{excimer:monomer}$ vs Control$_{excimer:monomer}$) of the ratio of

excimer (Em. Max. 460nm) to monomer (Em max. 400nm) fluorescence is indicated. Fold

changes < 1 indicate more ordered lipid bilayers relative to wildtype untreated control. ($n \geq 3$

biologically independent samples). **b**, Representative transmission EM micrographs of wildtype

and UBXD8 KO HEK-293T cells illustrating contacts between ER and plasma membrane. **c**,

Quantification of contact length between ER and plasma membrane in each genotype from (**b**)

(measurements are from $n = 3$ biological replicates with WT = 50 cells from 65 fields and UBXD8

KO = 53 cells from 71 fields). **d**, Split luciferase assay to measure contacts in mouse embryonic

fibroblasts with heterozygous or homozygous p97 R155H mutation. Cells were supplemented

with indicated concentrations of oleic acid and palmitic acid for 4 hours. ($n \geq 3$ biologically

independent samples). **e,** Representative SREBP1 and SCD1 staining from CA1 regions of 1

month-old control (CAMK2α) and p97 cKO mice (scale bar is 25 μm).   **f,** Quantification of

fluorescence intensity of images in (**e**). Individual points represent mean ROI intensity from each

mouse, 3 or 4 animals per group. **g,** Representative immunoblot for SREBP1 and SCD1 from

cortical brain lysates of 12-month-old control (C57), or 6- and 12-month-old p97[R155C/WT] mice (n=4

for each group). Pan 14-3-3 was used as housekeeping control. **h,** Quantification of (**g**). The ratio

of mature SREBP to total SREBP is shown. SCD1 intensities are normalized to 14-3-3 levels in

each lane. Individual points represent each mouse, 4 animals per group. Data are means ± SEM

(*, **, ***$P < 0.05$, 0.01, 0.0001 respectively. Significance was analyzed by One-way ANOVA with

Newman-Keuls multiple comparison test (**a**) or Tukey's multiple comparison test (**c, d**) or

Dunnett's multiple comparison (**h**) or Student's *t*-test (**f**).

Supplementary Figure Legends

Supplementary Fig. 1. Loss of p97 and UBXD8 results in increased ER-mitochondria

contacts.

a, Split luciferase assay to measure contacts in HEK-293T cells transfected with REEP1 (*n* > 3

biologically independent samples). **b**, Left: Split luciferase assay to measure contacts in HEK-

293T cells transfected with indicated siRNAs. (*n* ≥ 3 biologically independent samples). Right:

Immunoblot of HEK 293T cells transfected with indicated siRNAs. **c,** Affinity purification of

indicated UBXD8-HA/FLAG constructs transiently expressed in HEK-293T cells. Immunoblots of

whole cell lysates and affinity purifications probed with anti-HA, anti-p97, anti-PCNA, and ubiquitin

antibodies (*n* = 3 biologically independent samples). **d**, Top: Representative confocal image of

HeLa-Flp-IN-TRex cells stably expressing Sec61β-eGFP (green, ER) and stained for endogenous

TOMM20, (red, mitochondria). Bottom: ImageJ image analysis pipeline for the quantification of

contacts between ER and mitochondria[52]. **e**, Quantification of ER- mitochondria contacts in cells

transfected with indicated siRNAs (left and middle panel) or in UBXD8 KO HeLa-Flp-IN-TRex

cells (right panel) using assay in (**d**). Bottom panels show immunoblots for knockdown efficiency.

(n = 3 biologically independent samples) N: numbers of cells analyzed in each condition.

(Quartiles represent the upper 75th percentile and the lower 25th percentile. The line inside the

box represents the median. Whiskers indicate distribution of data from minimum to maximum in

a condition.). **f**, Quantification of number of mitochondria per field from transmission electron

microscopy of wildtype and UBXD8 KO cells. (Measurements are from n = 3 biological replicates

with WT = 50 cells in 65 fields and UBXD8 KO = 53 cells in 71 fields). Data are means ± SEM (*,

**, ***P < 0.05, 0.01, 0.0001 respectively. One-way ANOVA with Tukey's multiple comparison test

(**b & e** (left and middle panel)), Paired t-test (**a**) or Unpaired t test with Welch's correction (**e** (right

panel) & **f**).

Supplementary Fig. 2. ER stress does not alter ER-mitochondria contacts.

a, Split luciferase assay to measure contacts in HEK-293T cells treated with 2.5 µM Tunicamycin

(Tu), 1.5 µM Thapsigargin (Tg), for 2 hours. (n = 3 biologically independent samples). Immunoblot

of HEK-293T cells treated with Tu and Tg and probed for the indicated proteins. **b**, Immunoblot

of HEK-293T wildtype and UBXD8 KO cells treated with 1.5 µM Thapsigargin (Tg) for 2 hours,

UT: Untreated. (n = 3 biologically independent samples). Data are means ± SEM (ns: not

significant). One-way ANOVA with Tukey's multiple comparison test (**a**).

Supplementary Fig. 3. Quantitative proteomics of wildtype and UBXD8 KO contact

proteome.

a, Table depicting number of proteins and peptides quantified in post-nuclear supernatant and

mitochondria associated membrane fractions identified by proteomics in wildtype and UBXD8 KO

cells. Number of proteins up- or downregulated at $\log_2$- fold change (FC) (wildtype/ UBXD8 KO)

± 0.65 and ±1 is indicated. **b**, Venn diagram depicting overlap of our dataset with other putative

mitochondria associated membrane proteins identified by proteomics[18,23-25] **c**, Protein-protein

interaction network of differentially expressed proteins from MAM fractions of HEK-293T cells

involved in ERAD, cholesterol biosynthesis and lysosome function shown as clustered functional

categories. Protein associations were determined using STRING database with score ≥ 0.4. Each

node represents a protein belonging to enriched GO clusters as scored by Metascape. Size of

node represents $-\log_{10}$-transformed P value and color of node represents $\log_2$- fold change (FC)

(WT / UBXD8 KO). **d**, Volcano plot of the $-\log_{10}$-transformed P value *versus* the $\log_2$-transformed

ratio of wildtype/ UBXD8 KO proteins identified in the post-nuclear supernatant of HEK-293T cells.

n = 3 (each genotype) biologically independent samples. P values were computed by empirical

Bayesian statistical methods (adjusted for multiple comparisons) available in *Limma* R package;

for parameters, individual P values and q values, see Supplementary Table 1. **e**, Network of

differentially enriched functional ontology terms shown as clustered functional ontology

categories. Each node represents a functional ontology term enriched in the TMT data (**d**) as

scored by Metascape and networks generated using Cytoscape v3.8.2. Size of node represents

number of genes identified in each term by gene ontology (GO). Grey and Blue donuts represent

percent of genes identified in each GO term in wildtype or UBXD8 KO respectively. Node outline

thickness represents $-\log_{10}$-transformed P value of each term. The inner circle color of each node

indicates the corresponding functional GO cluster. **f,** Bubble plot representing significantly

enriched GO clusters identified from TMT proteomics of post-nuclear fractions in wildtype (blue)

or UBXD8 KO (green) cells (**d-e**). Size of the circle indicates the number of genes identified in

each cluster. **g**,**h**, Squalene epoxidase (SQLE) half-life measurements in wildtype and UBXD8

KO HEK 293T cells. FLAG-SQLE was transiently expressed, and cells were treated with 100

μg/mL cycloheximide for the indicated times. Samples were resolved on SDS-PAGE for

immunoblots (**g**) and levels of SQLE were quantified and normalized to loading control PCNA (**h**).

(n = 3 biologically independent samples). Data are means ± SEM (*, **, *** $P <$ 0.05, 0.01, 0.0001

respectively. One-way ANOVA with Tukey's multiple comparison test (**h**).

Supplementary Fig. 4. Loss of UBXD8 alters cellular lipidome with increased abundance of saturated fatty acid tail containing phospholipids

a, Relative levels of phospholipids (PLs), namely PG, phosphatidylglycerol; PI, phosphatidylinositol; and PS, phosphatidylserine in HEK-293T WT and UBXD8 KO cells. PLs were measured by LC-MS/MS following normalization by total protein amount. Each dot in the plot represents the level of a PL in UBXD8 KO cells relative to its level in wildtype cells. The dashed line represents a relative level of 1 (e.g., the level in UBXD8 KO cells is equal to the level in wildtype cells). (n = 3 biologically independent experiments were performed, each with duplicate samples). **b**, Relative levels of DG, Diacylglycerol; TG, Triacylglycerol in HEK-293T WT and UBXD8 KO cells. The DG and TG is quantified and visualized similarly as in **a**. (n = 3 biologically independent experiments were performed, each with duplicate samples). **c**, Relative levels of CE, Cholesteryl esters in HEK-293T WT and UBXD8 KO cells. The CE is quantified and visualized similarly as in **a**. (n = 3 biologically independent samples). **d-e**, Changes in the relative levels of PLs (**d**) including lysophopholipids (LPC, Lysophosphatidyl choline; LPE, Lysophosphatidyl ethanolamine; LPS, Lysophosphatidyl Serine; and LPI, Lysophophatidyl Inositol) and (**e**) Neutral lipids (DGs, TGs, and CEs) in HEK-293T WT and UBXD8 KO cells were quantified. The averaged relative fold changes were log_2 transformed and visualized as a heatmap. (n = 3 biologically independent experiments were performed, each with duplicate samples). **f**, Dot plots representing fold change of PC and PE lipids in UBDX8 KO cells relative to WT cells and fatty acid tails determined by MS/MS. The two tails of the lipid are shown and organized so that tail 1 (y-axis) is the shorter tail and tail 2 (x-axis) is the longer tail. The labels tail 1 and tail 2 do not represent their stereospecific number (sn). Grey boxes indicate the increase in saturated and mono-unsaturated tails in UBXD8 KO cells. Statistical analysis was performed on the log_2 transformed relative fold change values (UBXD8 KO relative to WT) using independent t tests and Benjamini-Hochberg correction in R stats package (p-values are listed in Supplemental Table S3).

Supplementary Fig. 5. Diminished SREBP pathway activation in UBXD8 KO cells

a, Schematic of SREBP pathway activation. In cholesterol-replete conditions, SCAP-INSIGs-SREBPs are in an inactive tripartite complex in the ER membrane. Cholesterol depletion triggers a conformational change in SCAP releasing it from INSIGs and enabling the transport of SCAP-SREBPs to the Golgi. Here SREBPs are cleaved sequentially by site 1 and site 2 proteases to release the active transcription factor. INSIGs are ubiquitylated and extracted for degradation from the membrane by p97-UBXD8. **b-c,** Real-time quantitative PCR of SREBP1 target genes including lipid desaturases (**b**), and SREBP2 target genes (**c**). **d,** Band intensity quantifications of mature SREBP2 in wildtype and UBXD8 KO HEK-293T cells corresponding to **Fig 3a** ($n \geq 3$ biologically independent samples) **e,** Split luciferase assay to measure contacts in HEK-293T cells transfected with siRNAs to UBXD8 or p97 and indicated 2X-FLAG-tagged mature SREBP1a, 1c, and 2 constructs. RLU: relative luminescence unit. ($n = 4$ biologically independent samples). **f**, Immunoblot of indicated proteins in HEK-293T cells transfected with siRNAs to UBXD8 or p97 and indicated 2X-FLAG-tagged mature SREBP1a, 1c, and 2 constructs. Immunoblots were probed with antibodies to SREBP1 and 2 to visualize immature and transfected mature forms. **g**, Split luciferase assay to measure contacts in HEK 293T cells treated with SCD1 inhibitor MF438 at 1 μM for 4 hours. Cells were also treated with oleic acid for 4 hours as indicated. ($n \geq 3$ biologically independent samples). **h,** Immunoblot of indicated proteins in HEK293T cells transfected with indicated siRNAs and wildtype or catalytically dead mutant of SCD1. GFP-HA/FLAG was transfected as a negative control. Related for Fig. **3e**. ($n = 3$ independent biological replicates). Data are means ± SEM (*, **, *** $P < 0.05$, 0.01, 0.0001 respectively. One-way ANOVA with Dunnett's multiple comparison test (**b, c**), Paired t test with Welch's correction (**d**) or One-way ANOVA with Tukey's multiple comparison test (**e & g**).

Supplementary Fig. 6. Loss of UBXD8 perturbs membrane saturation.

a, Global membrane fluidity in cells was measured using pyrene-based lipid probes in wildtype

and UBXD8 KO HeLa-Flp-IN-TRex cells. Wildtype cells were also treated with 5 μM of the p97

inhibitor CB-5083 for 4 hours. Cells were supplemented with indicated concentrations of oleic acid

and palmitic acid for 4 hours. The fold change (Treated$_{excimer:monomer}$ *vs* Control$_{excimer:monomer}$) of ratio

of excimer (Em. Max. 460nm) to monomer (Em max. 400nm) fluorescence is indicated. Fold

changes < 1 indicate more ordered lipid bilayers relative to WT untreated control. (n = 3

biologically independent samples). **b**, Representative transmission EM micrographs of

multilamellar bodies (red arrows) containing membrane whorls in UBXD8 KO HEK-293T cells.

Lower panel shows quantification of multilamellar bodies from images in (**c**). (Measurements are

from n = 3 biological replicates with WT = 50 cells in 65 fields and UBXD8 KO = 120 cells in 122

fields). Data are means ± SEM (*, **, ***P < 0.05, 0.01, 0.0001 respectively, One-way ANOVA with

Tukey's multiple comparison test (**a**). Scale bar, 100 nm.

Materials and Methods

Cell culture, transfections, immunoprecipitations, and treatments.

HEK293T, HeLa Kyoto Mouse embryonic fibroblasts (MEFs), and HeLa-Flp-IN-TREX

(HFTs, gift from Brian Raught, University of Toronto) cells were cultured in Dulbecco's modified

Eagle's medium, supplemented with 10% fetal bovine serum (FBS) and 100 units/mL penicillin

and streptomycin. Cells were maintained in a humidified, 5 % CO_2 atmosphere at 37°C. HeLa

Kyoto wildtype cell line was a generous gift from Ron Kopito (Stanford University).

For siRNA transfections, HEK293T cells were trypsinized and reverse transfected with

siRNAs. HeLa Kyoto, MEF, and HFT cells were trypsinized and seeded into a 12-well or 6-well

dish 24 hours prior to siRNA transfections. In both reverse and forward transfections, the cells

were transfected with 20 nM siRNAs using RNAiMax (Invitrogen) according to the manufacturer's

protocol. Cells were harvested 48-72 hours post transfection. For DNA transfections, HEK293T

cells in 6 well plates were transfected with 0.75 μg pcDNA3-Mit-NRluc91 and pcDNA3-CRluc92-

ER, 0.75 μg of UBXD8-C-HA/FLAG constructs, 1 μg of N-Myc-p97 constructs, 1 μg of p97-C-Myc,

1 μg of SCD1-C-HA/FLAG constructs, 0.2-0.5 μg of GFP-C-HA/FLAG, 0.75 μg of pCIG construct,

0.75 μg of REEP1, or 0.75 μg each of 2X-FLAG-SREBP1a, 2X-FLAG-SREBP1c, 2X-FLAG-

SREBP2 using Polyehtylenimine (PEI) at 1:4 DNA:PEI ratio and typically harvested 36-48 hours

post transfection. HeLa Kyoto, MEF, and HFT cells were transfected with cDNA using

Lipofectamine 2000 (Invitrogen) and the cells were harvested 36-48 hours post transfection. Cells

were lysed in mammalian cell lysis buffer (50 mM Tris-Cl, pH 6.8, 150 mM NaCl, 0.5% Nonidet

P-40, HALT Protease inhibitors (Pierce) and 1 mM DTT). Cells were incubated at 4^{o}C for 10 min

and then centrifuged at 14,000 rpm for 15 min at 4^{o}C. The supernatant was collected, and protein

concentration was estimated using the DC protein assay kit (Biorad). Protein G agarose (Pierce,

Thermo Fisher scientific) and the indicated antibodies were used for immunoprecipitation at 4^{o}C

for 3-5 hours. Beads were washed 3-5 times in 1 ml mammalian cell lysis buffer and resuspended

in 2X SDS sample buffer. Cells were treated with 1 μM of Bortezomib, 5 μM CB-5083, 5 μg/ml of

puromycin, 1 μM SCD1 inhibitor (MF438), 1 mM Oleic acid, or 0.25 mM Palmitic acid for the

indicated times (see figures for details). A full list of constructs used in this study can be found in

Supplementary Table 3.

Generation of CRISPR cell lines.

The CRISPR-Cas9 gene editing system was used to generate UBXD8 knockout cell lines in

HEK293T, and HFT cells. The guide sequence 5' GTTAACCTGCAGGGGTCGTGA 3' was

cloned into the pX459 vector carrying the hSpCas9 and transiently transfected into HEK293T and

HeLa-Flp-IN-TRex cells using Lipofectamine 3000 (Invitrogen) per the manufacturer's protocol.

36 hours post-transfection the cells were selected with 1 μg/ml puromycin for a further 24-36-hrs.

The surviving cells were then serially diluted into 96 well plates for clonal selection and expression

levels were monitored by immunoblotting.

Antibodies, siRNAs and Reagents.

The p97 (10736-1-AP), UBXD8 (16251-1-AP), FACL4 (22401-1-AP), UBXD2 (21052-1-AP),

HRD1 (13473-1-AP), Sec61β (51020-2-AP), Calnexin (10427-2-AP), UBXN1 (16135-1-AP),

SREBP1 (14088-1-AP), SREBP2 (28212-1-AP), FADS1 (10627-1-AP), anti-GFP (66002-1-AP)

and SCD1 (23393-1-AP) antibodies were from Proteintech Inc. The TIMM23 (H-8; sc514463),

TOMM20 (F-10; sc17764), TOMM70 (A-8; sc390545), pan-ubiquitin (P4D1; sc8017), c-Myc

(9E10; sc40), β-Actin (AC-15; sc69879), GAPDH (O411; sc47724), PCNA (PC10; sc56), UBE2J1

(B-6; sc377002), and UBE2S (C-1; sc390917) antibodies were obtained from Santa Cruz

Biotechnologies. LC3B (D11; 3868S), and BiP (C50B12; 3177T) were from Cell Signaling

Technologies. Derlin1 (A302-849A-T), p97 (A300-589A), and TEB4/MARCHIV (A304-171A-T)

were from Bethyl laboratories. The following antibodies UBXD7 (PA5-61972; Invitrogen), anti-HA

(16B12; MMS-101P, Covance), anti-FLAG (M2; F3165 Sigma Aldrich), INSIG1 (ab70784;

Abcam) were used for immunoblotting. HRP conjugated anti-rabbit (W401B) and anti-mouse

(W402B) secondary antibodies were from Promega. CB-5083 was a gift from Cleave Biosciences

and Bortezomib was from Selleckchem. Palmitic acid (100905) is from MP Biomedicals and Oleic

acid (270290050) is from Acros Organics. MF438 (569406) is from Millipore Sigma. All siRNAs

were purchased from Ambion (Thermo Fisher Scientific). UBXD8-0 (s23260), UBXD8-9 (s23259),

UBXD7-7 (s24997). UBXD2-1 (D-014184-03), UBXD2-2 (D-014184-04), HRD1-3 (D-007090-03),

and HRD1-4 (D-007090-04) were purchased from GE Dharmacon. siControl (SIC001) was from

Millipore Sigma. p97 siRNAs (2-HSS111263 and 3-HSS111264) were from Invitrogen (Thermo

Fisher Scientific). p97 rescue constructs were previously published [53] and were resistant to siRNA

2. UBXD8-C-HA/FLAG construct was previously published [53]. The UBXD8 rescue constructs,

including UBA* (^{17}LLQF^{20} mutated to ^{17}AAAA^{20}), ΔUAS (deleted amino acids between 122-277),

and UBX* (^{407}FPR^{409} mutated to ^{407}AAA^{409}), were cloned using overlap PCR followed by Gibson

assembly (NEB) cloning into pHAGE-C-HA/FLAG and were resistant to siRNA # 0. The SCD1-C-

HA/FLAG WT and catalytic dead mutant (His^{160} His^{161} and His^{301} His^{302} mutated to Ala^{160} Ala^{161}

and Ala^{301} Ala^{302}) constructs were cloned using overlap PCR followed by Gateway cloning

(Thermo Fisher Scientific) into pHAGE-C-HA/FLAG.

Mitochondria-associated membranes (MAM) fractionation.

MAMs were isolated as previously described [54,55]. Briefly, HEK293T or HeLa-Flp-IN-T-Rex cells

were seeded into four 150 mm TC dishes. Cells were lysed in Homogenization buffer (225 mM

mannitol, 75 mM sucrose, and 30 mM Tris-Cl, pH 7.4) using a Dounce homogenizer. The lysate

was centrifuged three times at 600xg for 5 minutes to remove unlysed cells and nuclei resulting

in post-nuclear supernatants (PNS). The cleared lysate was centrifuged at 7000xg to separate

crude mitochondrial pellet and supernatant containing microsomes. The supernatant was cleared

by centrifugation at 20,000xg for 30 minutes followed by microsome isolation using high-speed

centrifugation at 100,000xg for 1 hour. The crude mitochondrial pellet was washed twice in

homogenization buffer containing 0.1 mM EGTA at 7000xg and 10,000xg for 10 minutes. MAMs

were isolated from crude mitochondria using 30% Percoll gradient centrifugation at 95,000xg for

1 hr in a swinging-bucket rotor. The banded MAM fraction was washed once with phosphate-

buffered saline (PBS) before lysing in lysis buffer (50 mM Tris-Cl, pH 7.2, 150 mM NaCl, 2%

SDS). The pure mitochondrial fractions were resuspended and washed in mitochondrial

resuspension buffer (250 mM mannitol, 0.5 mM EGTA, 5 mM HEPES pH7.4). Mitochondrial

membranes were solubilized using 0.5% (v/v) Digitonin. Protein concentrations for both soluble

and pellet fractions were determined by DC protein assay kit (Biorad).

Split luciferase assay to measure ER-Mitochondria contacts.

Cells seeded a day prior in a 12-well plates were co-transfected with 0.75 μg pcDNA3-Mit-

NRluc91 and pcDNA3-CRluc92-ER (kind gift from Jeffrey A. Golden, Brigham and Women's

Hospital, Boston) using PEI at a 1:4 (DNA:PEI) ratio, or Lipofectamine 2000 (Invitrogen) as per

manufacturer's protocol. Media was changed after 6 hour and 18 hour later the cells are split into

a clear bottom white 96-well plate with 50-100K cells per well. After 24 hours, 30 μM of live-cell

substrate Enduren (Promega) was added to cells and incubated for 2-3 hours in a 37^{o}C incubator.

The luminescence was measured using a SpectraMax iD3 multi-well plate reader. The

luminescence measurements were normalized to the cell viability in each condition. Cell viability

was measured using Cell Titer-Glo (Promega) according to the manufacturers' instructions.

Relative luminescence units (RLU) for each cell line were normalized to the DMSO treated

samples to derive fold changes in RLU. Mean, standard error of means (SEM) and statistical

significance were calculated by one way ANOVA with indicated post-hoc analysis using

GraphPad Prism 5.01 (www.graphpad.com).

Immunofluorescence and Microscopy.

HFT cells stably expressing Sec61β-eGFP were grown on #1.5 cover slips in a 12 well plate and

transfected with indicated siRNAs using RNAiMax. 48 hours post-transfection, cells were washed

briefly in PBS and fixed with 4% paraformaldehyde at room temperature for 15 min. Cells were

washed in PBS and permeabilized in ice-cold 100% methanol at -20^{o}C for 10 min. Cells were

washed three times in PBS and blocked in 2% BSA in PBS with 0.3% Triton X-100 for 1 hour.

The coverslips were incubated overnight with the indicated antibodies in a humidified chamber.

The cells were washed and incubated for a further hour with appropriate Alexa-Fluor conjugated

secondary antibodies (Molecular Probes) for 1 hour in the dark. Cells were washed with PBS,

nuclei were stained with Hoechst and mounted on slides. All images were collected using a Nikon

A1R scan head with spectral detector and resonant scanners on a Ti-E motorized inverted

microscope equipped with 60× Plan Apo NA 1.4 objective lens. The indicated fluorophores were

excited with either a 405nm, 488nm or 594nm laser line. Images were analyzed using FIJI

(https://t/imagej.net/Fiji). Using a previously described method[52], co-localized pixel analysis to

quantify the ER-mitochondrial contact sites was performed using an ImageJ macro containing

tubeness, colocalization highlighter, and isophotcounter plugins (Supplementary fig 2a).

Lipid depletion and fatty acid supplementation.

Cells were depleted of or supplemented with fatty acids as previously described[56-58]. Briefly, cells

were treated with DMEM containing 0.5% lipid-depleted fetal calf serum (LDFCS; S5394, Sigma-

Aldrich) for 24 hours. 500 mM Oleic acid in DMSO was used as a stock solution to prepare a

working solution of 1 mM in DMEM containing 0.5% LDFCS. Cells were treated for 4 hr. 500 mM

palmitic acid stock solution was prepared in 100% ethanol by heating to 70°C for 20 min. This

stock solution, was used to prepare 0.25 mM palmitic acid solution in DMEM containing 0.5%

LDFCS which was heated in a water-bath at 50°C for 2 hr. The 0.25 mM palmitic acid solution is

cooled down to 37°C before adding to cells. Cell were incubated in palmitic acid solution for 4 hr.

All working solutions were prepared immediately prior to treatment.

Real time polymerase chain reaction (PCR)

Equal number of HEK293T WT or UBXD8 KO cells were seeded into a 6-well plate. The next day,

total RNA was isolated as per manufacturer's instructions using the PureLink RNA Mini kit

(Thermo fisher). The purified RNA was quantified and an equal amount of RNA was used for

cDNA preparation using iScript cDNA synthesis kit (Biorad). GAPDH was used as a housekeeping

gene was used. Primer sequences used in this study can be found in Supplementary Table 3.

Real time PCR was performed using the Powerup SyBr green master mix (Thermo Fisher). Data

analyses were carried out using the $2^{-\Delta\Delta Ct}$ method.

Membrane fluidity measurements

Cells were seeded in clear bottom black 96-well plate and treated with 5 μM CB-5083, lipid

depletion or lipid supplementation as described above. Membrane fluidity was measured using a

membrane fluidity kit (Abcam, ab189819) as per manufacturer's instructions. The assay uses a

lipophilic, membrane embedding pyrenedecanoic acid probe which undergoes a spectral shift in

emission from monomer (Em 400nm) to excimer (Em 460nm) based on local membrane fluidity

upon excitation at 360nm. The excimer to monomer (Em 460nm / Em 400nm) ratio was calculated

for each sample. Then fold changes of ratios (Treated$_{excimer:monomer}$ *vs* Control$_{excimer:monomer}$) were

deduced to provide a relative estimate of membrane fluidity compared to the wildtype untreated

control. A fold change less than 1 indicates ordered membranes relative to control.

Transmission Electron microscopy.

Cells were fixed in 2.5% glutaraldehyde, 3% paraformaldehyde with 5% sucrose in 0.1 M sodium

cacodylate buffer (pH 7.4), pelleted, and post fixed in 1% OsO_4 in veronal-acetate buffer. The

cells were stained *en bloc* overnight with 0.5% uranyl acetate in veronal-acetate buffer (pH 6.0),

then dehydrated and embedded in Embed-812 resin. Sections were cut on a Leica EM UC7 ultra

microtome with a Diatome diamond knife at a thickness setting of 50 nm, stained with 2% uranyl

acetate, and lead citrate. The sections were examined using a FEI Tecnai spirit at 80KV and

photographed with an AMT CCD camera. The images were analyzed manually for the ER-

Mitochondrial and ER-PM contacts using FIJI (https:/t/imagej.net/Fiji). Briefly, the scale of image

was set using Set Scale tool on ImageJ. Followed by measuring the length of ER, PM, or

perimeter of mitochondria using freehand line tool. The percent of contact length was determined

by taking the ratio of the length of ER, or PM (within contact distances of 25-100nm) to the

perimeter of mitochondria. The data were analyzed using GraphPad Prism 5.01 for Windows,

GraphPad Software, San Diego California USA, (www.graphpad.com).

TMT-based proteomics.

Sample preparation, digestion, and TMT labeling.

The PNS and MAM fractions were isolated from HEK293T WT or UBXD8 KO cells. 100 μg protein from each sample was precipitated using 15% (v/v) Trichloroacetic acid (TCA) followed by 100% Acetone washes. The protein pellets were resuspended in 200 mM N-(2-Hydroxyethyl)piperazine-N′-(3-propanesulfonic acid) (EPPS) (pH 8.5) buffer followed by reduction using 5 mM tris(2-carboxyethyl)phosphine (TCEP), alkylation with 14 mM iodoacetamide and quenched using 5 mM dithiothreitol treatments. The reduced and alkylated protein was precipitated using methanol and chloroform. The protein mixture was digested with LysC (Wako) overnight followed by Trypsin (Pierce) digestion for 6 hours at 37°C. The trypsin was inactivated with 30% (v/v) acetonitrile. The digested peptides were labelled with 0.2 mg per reaction of 6-plex TMT reagents (ThermoFisher scientific) (126, 127N, 127C, 128N, 128C, and 129N) at room temperature for 1 hour. The reaction was quenched using 0.5% (v/v) Hydroxylamine for 15 min. A 2.5 μL aliquot from the labeling reaction was tested for labeling efficiency. TMT-labeled peptides from each sample were pooled together at a 1:1 ratio. The pooled peptide mix was dried under vacuum and resuspended in 5% formic acid for 15 min. The resuspended peptide sample was further purified using C18 solid-phase extraction (SPE) (Sep-Pak, Waters).

Off-line basic pH reverse-phase (BPRP) fractionation.

We fractionated the pooled, labeled peptide sample using BPRP HPLC[59]. We used an Agilent 1200 pump equipped with a degasser and a detector (set at 220 and 280 nm wavelength). Peptides were subjected to a 50-min linear gradient from 5% to 35% acetonitrile in 10 mM ammonium bicarbonate pH 8 at a flow rate of 0.6 mL/min over an Agilent 300Extend C18 column (3.5 µm particles, 4.6 mm ID and 220 mm in length). The peptide mixture was fractionated into a total of 96 fractions, which were consolidated into 24 super-fractions[60]. Samples were subsequently acidified with 1% formic acid and vacuum centrifuged to near dryness. Each

consolidated fraction was desalted via StageTip, dried again via vacuum centrifugation, and

reconstituted in 5% acetonitrile, 5% formic acid for LC-MS/MS processing.

Liquid chromatography and tandem mass spectrometry.

Mass spectrometric data were collected on an Orbitrap Lumos mass spectrometer coupled to a

Proxeon NanoLC-1000 UHPLC. The 100 μm capillary column was packed with 35 cm of Accucore

150 resin (2.6 μm, 150Å; ThermoFisher Scientific). The scan sequence began with an MS1

spectrum (Orbitrap analysis, resolution 120,000, 350−1400 Th, automatic gain control (AGC)

target 5 x10^5, maximum injection time 50 ms). Data were acquired for 150 minutes per fraction.

SPS-MS3 analysis was used to reduce ion interference[61,62]. MS2 analysis consisted of collision-

induced dissociation (CID), quadrupole ion trap analysis, automatic gain control (AGC) 1 x10^4,

NCE (normalized collision energy) 35, q-value 0.25, maximum injection time 60 ms), isolation

window at 0.5 Th. Following acquisition of each MS2 spectrum, we collected an MS3 spectrum in

which multiple MS2 fragment ions were captured in the MS3 precursor population using isolation

waveforms with multiple frequency notches. MS3 precursors were fragmented by HCD and

analyzed using the Orbitrap (NCE 65, AGC 3.0 x10^5, isolation window 1.3 Th, maximum injection

time 150 ms, resolution was 50,000).

Data analysis.

Spectra were converted to mzXML via MSconvert[63]. Database searching included all entries from

the Human UniProt Database (downloaded: August 2018). The database was concatenated with

one composed of all protein sequences for that database in the reversed order. Searches were

performed using a 50-ppm precursor ion tolerance for total protein level profiling. The product ion

tolerance was set to 0.9 Da. These wide mass tolerance windows were chosen to maximize

sensitivity in conjunction with Comet searches and linear discriminant analysis[64,65]. TMT tags on

lysine residues and peptide N-termini (+229.163 Da for TMT) and carbamidomethylation of

cysteine residues (+57.021 Da) were set as static modifications, while oxidation of methionine

residues (+15.995 Da) was set as a variable modification. Peptide-spectrum matches (PSMs)

were adjusted to a 1% false discovery rate (FDR)[66,67]. PSM filtering was performed using a linear

discriminant analysis, as described previously[65] and then assembled further to a final protein-level

FDR of 1% [67]. Proteins were quantified by summing reporter ion counts across all matching PSMs,

also as described previously[68]. Reporter ion intensities were adjusted to correct for the isotopic

impurities of the different TMT reagents according to manufacturer specifications. The signal-to-

noise (S/N) measurements of peptides assigned to each protein were summed and these values

were normalized so that the sum of the signal for all proteins in each channel was equivalent to

account for equal protein loading. Finally, each protein abundance measurement was scaled,

such that the summed signal-to-noise for that protein across all channels equaled 100, thereby

generating a relative abundance (RA) measurement.

Downstream data analyses for TMT datasets were carried out using the R statistical package

(v4.0.3) and Bioconductor (v3.12; BiocManager 1.30.10). TMT channel intensities were quantile

normalized and then the data were log-transformed. The log transformed data were analyzed with

limma-based R package where p-values were FDR adjusted using an empirical Bayesian

statistical. Differentially expressed proteins were determined using a $\log_2$ (fold change (WT *vs*

UBXD8 KO)) threshold of > +/- 0.7.

Gene ontology (GO) functional enrichment analyses of proteomics data.

The differentially expressed proteins were further annotated and GO functional enrichment

analysis was performed using Metascape online tool (http://metascape.org)[51]. The GO cluster

network and protein-protein interaction network generated by metascape and the STRING

database (https://string-db.org/), respectively, were imported into Cytoscape software (v3.8.2) to

add required attributes (fold changes, p-values, gene number, and conditions) and prepared for

the visualization. Other proteomic data visualizations were performed using the RStudio software

(v1.4.1103), including hrbrthemes (v0.8.0), viridis (v0.6.1), dplyr (v.1.0.7), and ggplot2 (v 3.3.5).

Lipidomics

Sample preparation, mass spectrometry, and identification.

For each independent lipidomic experiment, HEK293T WT and UBXD8 KO cells were seeded in triplicate. Two of the three samples for each condition were used for lipidomics (i.e., lipids from duplicate samples were extracted and analyzed in parallel to determine technical variation). The third sample was used to determine the total protein concentration. Cells were washed with PBS, scraped into cold 50% methanol, centrifuged, and the cell pellets were frozen. Next, cells were resuspended in cold 50% methanol and transferred to glass vials. Chloroform was added and the mixture was gently vortexed and centrifuged at 1,000x g for 5 min at 4°C. Lipids were transferred to a clean glass vial using a glass Hamilton syringe. Lipids were extracted twice using chloroform prior to being dried under nitrogen gas. Samples were normalized according to protein concentration when resuspended in a 1:1:1 solution of methanol:chloroform:isopropanol prior to mass spectrometry (MS) analysis. The samples were stored at 4°C in an autosampler during data collection.

Lipids were identified and quantitatively measured using ultra high-performance liquid-chromatography high-resolution tandem MS/MS (UHPLC-MS/MS) as recently described[69,70]. Separation of lipids was done by reverse-phase chromatography using a Kinetex 2.6 μm C18 column (Phenomenex 00F-4462-AN) at 60°C using a Vanquish UHPLC system (Thermo Scientific) and two solvents: solvent A (40:60 water-methanol plus 10mM ammonium formate and 0.1% formic acid) and solvent B (10:90 methanol-isopropanol plus 10mM ammonium formate and 0.1% formic acid). UHPLC was performed at a 0.25 ml per min flow rate for 30 min per sample, starting at 25% solvent B and ending at 100% solvent B as described. The column was washed and equilibrated between samples. Samples were run in a semi-random order where WT or UBXD8 KO samples were interspersed with blank samples. Lipids were ionized using a heated electrospray ionization (HESI) source and nitrogen gas and measured using a Q-Exactive Plus

mass spectrometer operating at a MS1 resolution of either 70,000 or 140,000 and a MS2

resolution of 35,000. MS1 Spectra were collected over a mass range of 200 to 1,600 m/z with an

automatic gain control (AGC) setting of 1e6 and transient times of 250 ms (70,000 resolution) or

520 ms (140,000 resolution). MS2 spectra were collected using a transient time of 120 ms and

an AGC setting of 1e5. Each sample was analyzed using negative and positive ion modes. The

mass analyzer was calibrated weekly. SPLASH LIPIDOMIX mass spectrometry standards (Avanti

Polar Lipids) were used in determining extraction efficiencies and lipid quantitation. Quality control

(QC) samples consisting of lipids extracted from the National Institute of Standards and

Technology (NIST) Standard Reference Material 1950 Metabolites in Frozen Human Plasma

which contains plasma pooled from 100 healthy donors were used in this study. In parallel to the

samples, a control that lacked cells was used to determine any contaminants from the lipid

extraction and measurement steps. Any lipids found in the no cell control were removed during

analysis steps.

Lipids were identified and quantified using MAVEN[71], EI-MAVEN (Elucidata), Xcalibur

(ThermoFisher Scientific), and LipidSearch software (ThermoFisher Scientific). UHPLC retention

time, MS^1 peaks, and MS^2 fragments were used to identify lipids. The lipid retention time, MS1

peak shape, isotopic distribution, and MS^2 fragments were visually confirmed for all lipids reported

in this study. Peak area was used to determined lipid abundance. Lipids were included if they

were observed in 3-6 samples in both UBXD8 KO and WT cells. Missing values in a sample were

not imputed. The fold change of each lipid in UBXD8 KO cells relative to its level in WT cells was

used to test for statistical difference between UBXD8 KO and WT cells using independent t-tests

and the Benjamini-Hochberg correction method to control for false statistical discovery. The

following lipid classes were included in the analysis: cholesteryl esters (CE), diacylglycerol (DG),

phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylglycerol (PG),

phosphatidylinositol (PI), phosphatidylserine (PS), and triacylglycerol (TG). Guidelines from the

Lipidomic Standards Initiative were followed for lipid species identification and quantification,

including consideration of isotopic patterns resulting from naturally occurring ^{13}C atoms and

isomeric overlap. The following MS^2 information was used to confirm each lipid species: PC

fragment of 184.073 (positive mode) and tail identification using formic adduct (negative mode);

PE fragment of 196.038 or the tail plus 197.046 (negative mode) and neutral loss (NL) of 141.019

(positive mode); PG fragment of 152.996 plus the identification of the FA tails (negative mode)

and NL 189.04 of [M+NH4]+ adduct (positive mode); PI fragment of 241.012 (negative) and NL

277.056 of [M+NH4]+ adduct (positive mode); PS NL of 87.032 (negative); DG and TG by NL of

FA tails (positive mode); and CE fragment of 369.352 or neutral loss of 368.35 (positive).

Mouse studies

C57BL/6 (stock No.: 000664) and p97$^{R155H/WT}$ (B6;129S-Vcptm1Itl/J, Stock No: 021968) were

purchased from Jackson Laboratory. p97$^{R155C/WT}$ and p97 cKO (VCPFL/FL; CaMKIIa-Cre) were

obtained as reported previously[39,40]. All mice utilized in the study were on a C57BL/6 background.

Both male and female mice were used in this study. Animal procedures were performed in

accordance with protocols approved by the Animal Studies Committee at Washington University

School of Medicine.

Immunohistochemistry of tissue sections

Mice were anesthetized in an isoflurane chamber and perfused with PBS containing herapin. The

whole brain was removed from the skull and fixed in 4% PFA overnight at 4 °C degrees, cut

coronally into 40 micrometer sections, and stored in cryoprotectant solution at 4 °C degree for

staining. Sections were first rinsed 3 times with TBS and then blocked with blocking solution for

30 minutes (5% normal goat serum with 0.1% Triton X-100 in TBS). Sections were stained with

the primary antibody (SREBP1, 1:500, SCD1: 1:250 dilution) in TBS-0.1% Triton X-100 plus 2%

normal goat serum at 4 °C overnight, followed by 3 washes with TBS. Sections were then

incubated with the Alexa 488 and 555 tagged secondary antibodies (1:1000) for 2 hours at room

temperature, followed by counterstaining with DAPI (1:1000) for 20 min. After three washes with

TBS, the sections were mounted on the glass slides. True black (Biotium, NC1125051) was

incubated with the sections for 5 minutes to quench the auto-fluorescence. Slides were cover-

slipped using Prolong Gold mounting medium. Images were acquired using a Nikon Eclipse 80i

fluorescence microscope. All images were taken with the same fluorescent settings and

subsequently adjusted equally for brightness and contrast to ensure accurate pathology

quantification. For CA1 regions, ROIs are drawn according to DAPI staining and the mean

intensity was measured in each ROI by ImageJ. Background intensities (from three regions per

image were subtracted.

Immunoblot

Mouse cortex was lysed in RIPA buffer with protease inhibitor cocktails (PMSF and PIC) followed

by sonication (two cycles of 30 seconds cycles at 50% power). The protein concentration is

estimated by the BCA assay. Samples were loaded into 10% gel and transferred into

nitrocellulose membrane. The membranes were blocked by 5% milk in PBS-0.2% Tween20 and

incubated with the primary antibody in blocking solution overnight at 4 °C degrees. The membrane

was then washed three times with PBS-0.2% Tween20 and incubated with a secondary goat anti-

rabbit HRP antibody (1:5000) for 1 hour. Blot was rinsed three times with PBS-0.2% Tween20

and probed by a fresh mixture of ECL reagents at dark and then exposed by SYNGENE.

Statistical analyses.

For all experiments, N ≥ 3 biological replicates for each condition were examined. Fold changes,

SEM, and statistical analyses were performed using GraphPad Prism version 5.01 for Windows,

GraphPad Software, San Diego California USA (www.graphpad.com). The Statistical tests

performed, SEM, and statistical significance values are mentioned in the figures and

supplementary tables.

Acknowledgements

We thank Peter Juo and Karl Munger for critical reading of the manuscript. We are grateful to

Jeffrey Golden (Brigham and Women's Hospital, Harvard Medical School) for split luciferase

constructs, and the Whitehead Institute Electron Microscopy core for electron microscopy. We

would also like to thank Jacob Klickstein for developing the ImageJ script for ER-mitochondria co-

localization. This work is supported by the NIH grant GM127557 to M.R., funds from the University

of Arizona Health Sciences and BIO5 Institute to J.G.P., and NIH grant AG031867 to C.C.W.

Respective Contributions

R.G and M.R conceived the studies. R.G performed all experiments except for lipidomics studies

(that were performed by Y.X. and J.G.P. and analyzed by Y.X., I.K. and J.G.P.) and mouse

immunohistochemistry and immunoblotting (that were performed by J.Z). J.P and S.P.G assisted

with proteomic studies. M.R wrote the manuscript with input from R.G and J.G.P and C.C.W.

Competing Interests

The authors declare no conflicts of interest.

Request for reagents

Please contact the corresponding author, M.R for reagent requests.

Data availability

All raw proteomic and lipidomic data will be made available on public servers upon acceptance of

the manuscript. Any other data is available from the corresponding author upon request.

Code availability

The ImageJ macro to quantify ER-mitochondria contact sites will be made available on Github

upon manuscript acceptance.

References

1   Rowland, A. A. & Voeltz, G. K. Endoplasmic reticulum-mitochondria contacts: function of
the junction. *Nat Rev Mol Cell Biol* **13**, 607-625, doi:10.1038/nrm3440 (2012).
2   Barazzuol, L., Giamogante, F. & Calì, T. Mitochondria Associated Membranes (MAMs):
Architecture   and   physiopathological   role.   *Cell   Calcium*   **94**,   102343,
doi:10.1016/j.ceca.2020.102343 (2021).
3   Vance, J. E. MAM (mitochondria-associated membranes) in mammalian cells: lipids and
beyond. *Biochim Biophys Acta* **1841**, 595-609, doi:10.1016/j.bbalip.2013.11.014 (2014).
4   Kornmann, B. [ERMES, a multifunctional complex connecting endoplasmic reticulum and
mitochondria]. *Med Sci (Paris)* **26**, 145-146, doi:10.1051/medsci/2010262145 (2010).
5   de Brito, O. M. & Scorrano, L. Mitofusin 2 tethers endoplasmic reticulum to mitochondria.
Nature **456**, 605-610, doi:10.1038/nature07534 (2008).
6   Kornmann, B. & Walter, P. ERMES-mediated ER-mitochondria contacts: molecular hubs
for   the   regulation   of   mitochondrial   biology.   *J   Cell   Sci*   **123**,   1389-1393,
doi:10.1242/jcs.058636 (2010).
7   De Vos, K. J. *et al.* VAPB interacts with the mitochondrial protein PTPIP51 to regulate
calcium homeostasis. *Hum Mol Genet* **21**, 1299-1311, doi:10.1093/hmg/ddr559 (2012).
8   Chandra, D. *et al.* Association of active caspase 8 with the mitochondrial membrane during
apoptosis: potential roles in cleaving BAP31 and caspase 3 and mediating mitochondrion-
endoplasmic reticulum cross talk in etoposide-induced cell death. *Mol Cell Biol* **24**, 6592-
6607, doi:10.1128/mcb.24.15.6592-6607.2004 (2004).
9   Rapizzi, E. *et al.* Recombinant expression of the voltage-dependent anion channel
enhances the transfer of Ca2+ microdomains to mitochondria. *J Cell Biol* **159**, 613-624,
doi:10.1083/jcb.200205091 (2002).
10   Preston, G. M. & Brodsky, J. L. The evolving role of ubiquitin modification in endoplasmic
reticulum-associated   degradation.   *Biochem   J*   **474**,   445-469,   doi:10.1042/bcj20160582
(2017).
11   Song, J., Herrmann, J. M. & Becker, T. Quality control of the mitochondrial proteome. *Nat*
Rev Mol Cell Biol **22**, 54-70, doi:10.1038/s41580-020-00300-2 (2021).
12   Meyer, H. p97 complexes as signal integration hubs. *BMC Biol* **10**, 48, doi:10.1186/1741-
7007-10-48 (2012).
13   Ganji, R., Mukkavalli, S., Somanji, F. & Raman, M. The VCP-UBXN1 Complex Mediates
Triage of Ubiquitylated Cytosolic Proteins Bound to the BAG6 Complex. *Mol Cell Biol* **38**,
doi:10.1128/MCB.00154-18 (2018).
14   Tanaka, A. *et al.* Proteasome and p97 mediate mitophagy and degradation of mitofusins
induced by Parkin. *J Cell Biol* **191**, 1367-1380, doi:10.1083/jcb.201007013 (2010).
15   Alexandru, G. *et al.* UBXD7 binds multiple ubiquitin ligases and implicates p97 in
HIF1alpha turnover. *Cell* **134**, 804-816, doi:10.1016/j.cell.2008.06.048 (2008).
16   Raman, M. *et al.* Systematic proteomics of the VCP-UBXD adaptor network identifies a
role   for   UBXN10   in   regulating   ciliogenesis.   *Nat   Cell   Biol*   **17**,   1356-1369,
doi:10.1038/ncb3238 (2015).
17   Williamson, C. D., Wong, D. S., Bozidis, P., Zhang, A. & Colberg-Poley, A. M. Isolation of
Endoplasmic Reticulum, Mitochondria, and Mitochondria-Associated Membrane and
Detergent Resistant Membrane Fractions from Transfected Cells and from Human
Cytomegalovirus-Infected Primary Fibroblasts. *Curr Protoc Cell Biol* **68**, 3.27.21-23.27.33,
doi:10.1002/0471143030.cb0327s68 (2015).
18   Cho, I. T. *et al.* Ascorbate peroxidase proximity labeling coupled with biochemical
fractionation identifies promoters of endoplasmic reticulum-mitochondrial contacts. *J Biol*
Chem **292**, 16382-16392, doi:10.1074/jbc.M117.795286 (2017).

19 Kim, H. *et al.* UAS domain of Ubxd8 and FAF1 polymerizes upon interaction with long-
chain unsaturated fatty acids. *J Lipid Res* **54**, 2144-2152, doi:10.1194/jlr.M037218 (2013).

20 Xu, Y., Liu, Y., Lee, J. G. & Ye, Y. A ubiquitin-like domain recruits an oligomeric chaperone
to a retrotranslocation complex in endoplasmic reticulum-associated degradation. *J Biol
Chem* **288**, 18068-18076, doi:10.1074/jbc.M112.449199 (2013).

21 Ting, L., Rad, R., Gygi, S. P. & Haas, W. MS3 eliminates ratio distortion in isobaric
multiplexed quantitative proteomics. *Nat Methods* **8**, 937-940, doi:10.1038/nmeth.1714
(2011).

22 McAlister, G. C. *et al.* MultiNotch MS3 enables accurate, sensitive, and multiplexed
detection of differential expression across cancer cell line proteomes. *Anal Chem* **86**,
7150-7158, doi:10.1021/ac502040v (2014).

23 Hung, V. *et al.* Proteomic mapping of cytosol-facing outer mitochondrial and ER
membranes in living human cells by proximity biotinylation. *Elife* **6**,
doi:10.7554/eLife.24463 (2017).

24 Kwak, C. *et al.* Contact-ID, a tool for profiling organelle contact sites, reveals regulatory
proteins of mitochondrial-associated membrane formation. *Proc Natl Acad Sci U S A* **117**,
12109-12120, doi:10.1073/pnas.1916584117 (2020).

25 Cho, K. F. *et al.* Split-TurboID enables contact-dependent proximity labeling in cells. *Proc
Natl Acad Sci U S A* **117**, 12143-12154, doi:10.1073/pnas.1919528117 (2020).

26 Olzmann, J. A., Richter, C. M. & Kopito, R. R. Spatial regulation of UBXD8 and p97/VCP
controls ATGL-mediated lipid droplet turnover. *Proc Natl Acad Sci U S A* **110**, 1345-1350,
doi:10.1073/pnas.1213738110 (2013).

27 Vance, J. E. Phospholipid synthesis in a membrane fraction associated with mitochondria.
J Biol Chem **265**, 7248-7256 (1990).

28 Ikeda, Y. *et al.* Regulated endoplasmic reticulum-associated degradation of a polytopic
protein: p97 recruits proteasomes to Insig-1 before extraction from membranes. *J Biol
Chem* **284**, 34889-34900, doi:10.1074/jbc.M109.044875 (2009).

29 Horton, J. D., Goldstein, J. L. & Brown, M. S. SREBPs: activators of the complete program
of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* **109**, 1125-1131,
doi:10.1172/jci15593 (2002).

30 Surma, M. A. *et al.* A lipid E-MAP identifies Ubx2 as a critical regulator of lipid saturation
and lipid bilayer stress. *Mol Cell* **51**, 519-530, doi:10.1016/j.molcel.2013.06.014 (2013).

31 Hayashi, T. & Fujimoto, M. Detergent-resistant microdomains determine the localization
of sigma-1 receptors to the endoplasmic reticulum-mitochondria junction. *Mol Pharmacol*
77, 517-528, doi:10.1124/mol.109.062539 (2010).

32 Park, W. J. *et al.* A novel FADS1 isoform potentiates FADS2-mediated production of
eicosanoid precursor fatty acids. *J Lipid Res* **53**, 1502-1512, doi:10.1194/jlr.M025312
(2012).

33 Bai, Y. *et al.* X-ray structure of a mammalian stearoyl-CoA desaturase. *Nature* **524**, 252-
256, doi:10.1038/nature14549 (2015).

34 Léger, S. *et al.* Synthesis and biological activity of a potent and orally bioavailable SCD
inhibitor (MF-438). *Bioorg Med Chem Lett* **20**, 499-502, doi:10.1016/j.bmcl.2009.11.111
(2010).

35 Watts, G. D. *et al.* Novel VCP mutations in inclusion body myopathy associated with Paget
disease of bone and frontotemporal dementia. *Clin Genet* **72**, 420-426,
doi:10.1111/j.1399-0004.2007.00887.x (2007).

36 Johnson, J. O. *et al.* Exome sequencing reveals VCP mutations as a cause of familial
ALS. *Neuron* **68**, 857-864, doi:10.1016/j.neuron.2010.11.036 (2010).

37 Abramzon, Y. *et al.* Valosin-containing protein (VCP) mutations in sporadic amyotrophic
lateral sclerosis. *Neurobiol Aging* **33**, 2231 e2231-2231 e2236,
doi:10.1016/j.neurobiolaging.2012.04.005 (2012).

38　Gonzalez, M. A. *et al.* A novel mutation in VCP causes Charcot-Marie-Tooth Type 2
disease. *Brain* **137**, 2897-2902, doi:10.1093/brain/awu224 (2014).
39　Wani, A. *et al.* Neuronal VCP loss of function recapitulates FTLD-TDP pathology. *Cell Rep*
36, 109399, doi:10.1016/j.celrep.2021.109399 (2021).
40　Clemen, C. S. *et al.* The heterozygous R155C VCP mutation: Toxic in humans! Harmless
in mice? *Biochem Biophys Res Commun* **503**, 2770-2777, doi:10.1016/j.bbrc.2018.08.038
(2018).
41　Loregger, A. *et al.* Haploid Mammalian Genetic Screen Identifies UBXD8 as a Key
Determinant of HMGCR Degradation and Cholesterol Biosynthesis. *Arterioscler Thromb*
Vasc Biol **37**, 2064-2074, doi:10.1161/atvbaha.117.310002 (2017).
42　Schrul, B. & Kopito, R. R. Peroxin-dependent targeting of a lipid-droplet-destined
membrane protein to ER subdomains. *Nat Cell Biol* **18**, 740-751, doi:10.1038/ncb3373
(2016).
43　Kolawa, N. *et al.* Perturbations to the ubiquitin conjugate proteome in yeast δubx mutants
identify Ubx2 as a regulator of membrane lipid composition. *Mol Cell Proteomics* **12**, 2791-
2803, doi:10.1074/mcp.M113.030163 (2013).
44　Nahar, S., Chowdhury, A., Ogura, T. & Esaki, M. A AAA ATPase Cdc48 with a cofactor
Ubx2 facilitates ubiquitylation of a mitochondrial fusion-promoting factor Fzo1 for
proteasomal degradation. *J Biochem* **167**, 279-286, doi:10.1093/jb/mvz104 (2020).
45　Chowdhury, A., Ogura, T. & Esaki, M. Two Cdc48 cofactors Ubp3 and Ubx2 regulate
mitochondrial morphology and protein turnover. *J Biochem* **164**, 349-358,
doi:10.1093/jb/mvy057 (2018).
46　Mårtensson, C. U. *et al.* Mitochondrial protein translocation-associated degradation.
Nature **569**, 679-683, doi:10.1038/s41586-019-1227-y (2019).
47　Rieusset, J. Contribution of mitochondria and endoplasmic reticulum dysfunction in insulin
resistance: Distinct or interrelated roles? *Diabetes Metab* **41**, 358-368,
doi:10.1016/j.diabet.2015.02.006 (2015).
48　Xu, L., Wang, X. & Tong, C. Endoplasmic Reticulum-Mitochondria Contact Sites and
Neurodegeneration. *Front Cell Dev Biol* **8**, 428, doi:10.3389/fcell.2020.00428 (2020).
49　Marchi, S. *et al.* Oncogenic and oncosuppressive signal transduction at mitochondria-
associated endoplasmic reticulum membranes. *Mol Cell Oncol* **1**, e956469,
doi:10.4161/23723548.2014.956469 (2014).
50　Hall, C. E. *et al.* Progressive Motor Neuron Pathology and the Role of Astrocytes in a
Human Stem Cell Model of VCP-Related ALS. *Cell Rep* **19**, 1739-1749,
doi:10.1016/j.celrep.2017.05.024 (2017).
51　Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-09234-6 (2019).
52　Filadi, R. *et al.* Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria
coupling. *Proc Natl Acad Sci U S A* **112**, E2174-2181, doi:10.1073/pnas.1504880112
(2015).
53　Raman, M., Havens, C. G., Walter, J. C. & Harper, J. W. A genome-wide screen identifies
p97 as an essential regulator of DNA damage-dependent CDT1 destruction. *Mol Cell* **44**,
72-84, doi:10.1016/j.molcel.2011.06.036 (2011).
54　Wieckowski, M. R., Giorgi, C., Lebiedzinska, M., Duszynski, J. & Pinton, P. Isolation of
mitochondria-associated membranes and mitochondria from animal tissues and cells. *Nat*
Protoc **4**, 1582-1590, doi:10.1038/nprot.2009.151 (2009).
55　Williamson, C. D., Wong, D. S., Bozidis, P., Zhang, A. & Colberg-Poley, A. M. Isolation of
Endoplasmic Reticulum, Mitochondria, and Mitochondria-Associated Membrane and
Detergent Resistant Membrane Fractions from Transfected Cells and from Human
Cytomegalovirus-Infected Primary Fibroblasts. *Curr Protoc Cell Biol* **68**, 3 27 21-23 27 33,
doi:10.1002/0471143030.cb0327s68 (2015).

56 Wang, Z. *et al.* Saturated fatty acids activate microglia via Toll-like receptor 4/NF-kappaB
signalling. *Br J Nutr* **107**, 229-241, doi:10.1017/S0007114511002868 (2012).
57 Davis, O. B. *et al.* NPC1-mTORC1 Signaling Couples Cholesterol Sensing to Organelle
Homeostasis and Is a Targetable Pathway in Niemann-Pick Type C. *Dev Cell* **56**, 260-276
e267, doi:10.1016/j.devcel.2020.11.016 (2021).
58 Lim, C. Y. *et al.* ER-lysosome contacts enable cholesterol sensing by mTORC1 and drive
aberrant growth signalling in Niemann-Pick type C. *Nat Cell Biol* **21**, 1206-1218,
doi:10.1038/s41556-019-0391-5 (2019).
59 Wang, Y. *et al.* Reversed-phase chromatography with multiple fraction concatenation
strategy for proteome profiling of human MCF10A cells. *Proteomics* **11**, 2019-2026,
doi:10.1002/pmic.201000722 (2011).
60 Paulo, J. A. *et al.* Quantitative mass spectrometry-based multiplexing compares the
abundance of 5000 S. cerevisiae proteins across 10 carbon sources. *J Proteomics* **148**,
85-93, doi:10.1016/j.jprot.2016.07.005 (2016).
61 Gygi, J. P. *et al.* Web-Based Search Tool for Visualizing Instrument Performance Using
the Triple Knockout (TKO) Proteome Standard. *J Proteome Res* **18**, 687-693,
doi:10.1021/acs.jproteome.8b00737 (2019).
62 Paulo, J. A., O'Connell, J. D. & Gygi, S. P. A Triple Knockout (TKO) Proteomics Standard
for Diagnosing Ion Interference in Isobaric Labeling Experiments. *J Am Soc Mass*
Spectrom **27**, 1620-1625, doi:10.1007/s13361-016-1434-9 (2016).
63 Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nat*
Biotechnol **30**, 918-920, doi:10.1038/nbt.2377 (2012).
64 Beausoleil, S. A., Villen, J., Gerber, S. A., Rush, J. & Gygi, S. P. A probability-based
approach for high-throughput protein phosphorylation analysis and site localization.
Nature biotechnology **24**, 1285-1292, doi:10.1038/nbt1240 (2006).
65 Huttlin, E. L. *et al.* A tissue-specific atlas of mouse protein phosphorylation and
expression. *Cell* **143**, 1174-1189, doi:10.1016/j.cell.2010.12.001 (2010).
66 Elias, J. E. & Gygi, S. P. Target-decoy search strategy for mass spectrometry-based
proteomics. *Methods Mol Biol* **604**, 55-71, doi:10.1007/978-1-60761-444-9_5 (2010).
67 Elias, J. E. & Gygi, S. P. Target-decoy search strategy for increased confidence in large-
scale protein identifications by mass spectrometry. *Nat Methods* **4**, 207-214,
doi:10.1038/nmeth1019 (2007).
68 McAlister, G. C. *et al.* Increasing the Multiplexing Capacity of TMTs Using Reporter Ion
Isotopologues with Isobaric Masses. *Analytical Chemistry* **84**, 7469-7478,
doi:10.1021/ac301572t (2012).
69 Xi, Y., Harwood, S., Wise, L. M. & Purdy, J. G. Human Cytomegalovirus pUL37x1 Is
Important for Remodeling of Host Lipid Metabolism. *J Virol* **93**, doi:10.1128/JVI.00843-19
(2019).
70 Xi, Y., Lindenmayer, L., Kline, I., von Einem, J. & Purdy, J. G. Human Cytomegalovirus
Uses a Host Stress Response To Balance the Elongation of Saturated/Monounsaturated
and Polyunsaturated Very-Long-Chain Fatty Acids. *mBio* **12**, doi:10.1128/mBio.00167-21
(2021).
71 Melamud, E., Vastag, L. & Rabinowitz, J. D. Metabolomic analysis and visualization
engine for LC-MS data. *Anal Chem* **82**, 9818-9826, doi:10.1021/ac1021166 (2010).
1060
1061

Figure 1.

Figure 2

Figure 3.

Figure 4.

a Global Membrane Fluidity (HEK-293T)

Fold Change (Treated (Excimer:Monomer) *vs* Control (Excimer:Monomer))

1mM Oleic Acid	0.25 mM Palmitic Acid

Ctrl — CB5083 — UBXD8 KO — WT

b HEK293T WT — Nucleus — ER — UBXD8 KO — ER — Nucleus — Scale bar: 100nm

c % PM apposed with ER (*vs* PM length per field)

WT — UBXD8 KO

<25 nm — 26-50 nm — 51-100 nm

Distance between ER-Plasma Membrane

d ER-Mito contacts

Fold Change (RLU)

1 mM Oleic acid	0.25 mM Palmitic acid

p97^wt/wt — p97^R155H/wt Heterozygous — p97^R155H/R155H Homozygous

Mouse Embryonic Fibroblasts

e DAPI — SREBP-1 — DAPI — SCD1

CAMK — p97 cKO

f SREBP1 Intensity — CAMK — p97 cKO

SCD1 Intensity — CAMK — p97 cKO

g p97^WT/R155C

C57 — R155C (6 mos) — R155C (13 mos)

SREBP1 — Immature — mature — 150 kDa — 100 kDa — 75 kDa

14-3-3 — 37 kDa

SCD1 — 50 kDa

14-3-3 — 37 kDa

h Active SREBP1/ total SREBP1

C57 — 6 — 13 — Months — R155C

SCD/ pan 14-3-3

C57 — 6 — 13 — Months — R155C

Supplementary Figure 1.

Supplementary Figure 2.

a

b

Supplementary Figure 3.

a

		Post nuclear supernatant	Mitochondria-associated membrane
	No. of proteins quantified	7556	4499
	No. of peptides quantified	56738	35371
Log₂FC=-1	No. of upregulated proteins in UBXD8 KO	29	23
Log₂FC=+1	No. of upregulated proteins in WT	74	28
Log₂FC=-0.65	No. of upregulated proteins in UBXD8 KO	123	108
Log₂FC=+0.65	No. of upregulated proteins in WT	226	112

Supplementary Figure 6.

Scale bar: 100nm

	Number of fields used for quantification	Total number of whorls	Percentage
HEK293T WT	65	4	6.15%
UBXD8 KO	122	40	32.7%