Testosterone promotes Coxsackievirus B3 pathogenesis in an oral inoculation mouse model

Adeeba H. Dhalech[#], Caleb M. Corn[#], Vrushali Mangale, and Christopher M. Robinson*

Department of Microbiology and Immunology, Indiana University School of Medicine,

Indianapolis, IN, 46202, USA.

[#]These authors contributed equally to this work.

*To whom correspondence should be addressed.

Mailing address: 635 Barnhill Drive, Indianapolis, IN, 46202

Phone: 317-274-7822

Fax: 317-274-3331

Running Title: CVB3 fecal shedding and pathogenesis is enhanced by testosterone

Abstract

Enteroviruses initiate infection in the gastrointestinal tract, and sex is often a biological variable

that impacts infection. The role of sex hormones on enterovirus pathogenesis, however, is

unclear. Previous data indicate that sex hormones can influence intestinal replication of

Coxsackievirus B3 (CVB3), an enterovirus in the Picornavirus family. To determine if

testosterone promotes CVB3 infection, male mice were castrated and provided placebo or

testosterone-filled capsules. We found that testosterone-treated mice shed significantly more

CVB3 in the feces and succumbed to CVB3-induced disease at a higher rate than castrated mice

given a placebo. Treatment of male mice with an androgen receptor antagonist, flutamide,

protected male mice from CVB3-induced lethality, further confirming the role of testosterone in

viral pathogenesis. We also observed higher viral loads in peripheral tissues of testosterone-

treated mice and an increase in the cytokine and chemokine response. Finally, we found that

testosterone treatment in female mice increased fecal CVB3 shedding but had no impact on viral

lethality. Overall, these data indicate that testosterone and androgen receptor signaling can

promote CVB3 replication in the intestine and enhance CVB3 lethality in a sex-dependent

manner.

Importance

Biological sex plays a significant role in the outcome of various infections and diseases. The

impact of sex hormones on intestinal replication and dissemination of Coxsackievirus B3

remains limited. Using an oral inoculation model, we found that testosterone enhances CVB3

shedding and lethality in male mice. Further, testosterone can promote CVB3 shedding in female

mice. This work highlights the role of testosterone in intestinal replication of CVB3 and suggests

that sex hormones can impact the replication of enteric viruses.

Introduction

Sex plays a significant role in human disease. Males are often more susceptible to pathogens, while females are more predisposed to autoimmune diseases (1-4). This disparity is linked to the immune system as sex hormones can significantly affect immune responses. Androgens, such as testosterone, can suppress immunity, delaying the elimination of pathogens (5-9). On the other hand, estrogens can enhance both the cell-mediated and humoral immune response, and there is evidence supporting estrogen's influence on disease severity in cardiovascular diseases and traumatic brain injuries (10-16). However, the mechanism and consequences of sex hormones on infectious diseases remain limited.

Coxsackievirus is a non-enveloped RNA virus in the *Picornaviridae* family and implicated in several human diseases, including viral myocarditis, HFMD, and meningitis (17-19). Infants, children, and immunocompromised individuals are the most susceptible to these diseases, and Coxsackievirus is associated with an 11% fatality rate in neonates (20-22). Coxsackievirus B3 (CVB3) is one of the leading causes of viral myocarditis. There is a strong sex bias in viral myocarditis as it affects more males than females, with a mortality rate of 2:1 in individuals infected under the age of 40 (23, 24). Animal models of CVB3 have provided further evidence of this sex bias. Our laboratory, and others, have shown that sex hormones contribute to CVB3 pathogenesis. Castration of male mice before infection reduces CVB3-induced myocarditis and mortality (25, 26). In CVB3-infected mice, dysregulation of the immune response is hypothesized to contribute to mortality. Testosterone and estradiol influence the CD4[+] T cell response to CVB3 to either promote or protect against viral-induced myocarditis (27-29). Castration of male mice also significantly reduces anti-inflammatory M2 macrophages resulting

in more severe myocarditis following CVB3 infection, suggesting other immune cells contribute

to myocarditis (30). Finally, genes on the Y chromosome also likely contribute to CVB3-induced

disease, indicating that viral myocarditis is multifactorial in males (31, 32). Therefore, many

questions remain as to the mechanism of the sex bias in Coxsackievirus infections.

While animal models have provided vital information on CVB3 pathogenesis, many of these

models mimic systemic infection through intraperitoneal injections of the virus. However,

Coxsackievirus is a member of a group of viruses transmitted through the fecal-oral route,

termed enteroviruses. Very few studies have shown successful oral inoculation of mice with

Coxsackievirus (33-37). Recently we established an oral inoculation model to examine CVB3

replication in the intestine. Using this model, we also observed a sex bias in infection. We found

that male mice support robust intestinal replication and succumb to CVB3 induced disease.

Female mice, however, are mainly resistant to CVB3 with limited viral replication in the

intestine (38). Gonadectomy before CVB3 infection impacted fecal shedding in mice and

protected male mice from CVB3-induced lethality, suggesting that sex hormones contribute to

viral pathogenesis. Here, we show that testosterone enhances CVB3 shedding in male and female

mice. Testosterone also enhanced lethality in male mice, but not in female mice. Overall, these

data highlight the importance of examining sex bias in CVB3 induced disease.

Results

Testosterone and androgen receptor signaling enhances CVB3 lethality in male mice

following oral inoculation. We have previously established an oral inoculation model for CVB3

using C57BL/6 *Ifnar*$^{-/-}$ (deficient for that interferon α/β receptor) mice (38). We found that

castration of male *Ifnar*$^{-/-}$ mice prior to oral infection with CVB3 protected against CVB3-

induced lethality and significantly reduced CVB3 fecal shedding. To determine if testosterone,

specifically, could impact CVB3 pathogenesis, we surgically castrated male *Ifnar*$^{-/-}$ mice to

deplete endogenous testosterone. As a control, we also performed mock castrations on male

mice. One week after surgery, we implanted mice with placebo or exogenous testosterone

capsules. To assess hormone replacement, we determined serum concentrations of testosterone

by ELISA. Castrated mice that received the placebo capsule had a significantly lower serum

testosterone concentration than mock-castrated and castrated male mice that received

testosterone (Fig. 1A). Further, mock castrated and castrated mice receiving exogenous

testosterone had similar amounts of serum testosterone, confirming successful hormone

treatment. One week following hormone treatment, male mice were orally inoculated with $5x10^7$

PFU CVB3 and monitored for survival for 14 days post-inoculation (dpi). We found that, like

our previous study (38), testosterone-depleted mice (castrated + placebo) were protected from

CVB3-induced lethality (Fig 1B). Further, mock castrated mice and testosterone-treated

(castrated + testosterone) mice were significantly more likely to die from CVB3 inoculation than

testosterone-depleted mice. These data indicate that testosterone contributes to CVB3-induced

lethality in male mice.

To confirm the effects of testosterone on male *Ifnar*$^{-/-}$ mice, we hypothesized that treatment with

an androgen receptor antagonist, flutamide, would also confer similar protection against CVB3

infection. To test this hypothesis, we administered flutamide (25µg/kg) by intraperitoneal

injections to male *Ifnar*$^{-/-}$ mice prior to oral inoculation with CVB3 and monitored mice for

survival for 14 dpi. We found that mice treated with flutamide were protected entirely from

CVB3-induced lethality compared to mice provided vehicle control injections (Fig. 1C). Taken

together, these data indicate that testosterone and androgen receptor signaling in orally CVB3

inoculated male *Ifnar*$^{-/-}$ mice is a significant contributor to viral-induced lethality.

Testosterone promotes CVB3 fecal shedding and viral dissemination to peripheral tissues.

CVB3 is spread through the fecal-oral route; therefore, we next examined CVB3 fecal shedding,

indicative of viral replication in the intestine (38). Following oral inoculation, we collected feces

at 1, 2, and 3 dpi and quantified fecal CVB3 using a plaque assay on HeLa cells. We found that

at 2 and 3 dpi, testosterone-depleted mice shed significantly less CVB3 in the feces than mock-

castrated male mice (Fig. 2A and 2B). Further, castrated mice receiving exogenous testosterone

shed similar levels of CVB3 as compared to mock-castrated mice and significantly more CVB3

than testosterone-depleted mice. Since we observed higher fecal shedding in mice with

testosterone, we hypothesized that these mice with testosterone would also have higher viral

titers in the peripheral tissues following oral inoculation. To examine this hypothesis, we

harvested the heart, liver, kidney, spleen, and pancreas at three dpi and quantified CVB3 tissue

titers by a plaque assay. Similar to fecal shedding, we observed higher tissue CVB3 titers for

mock castrated and testosterone-treated mice than testosterone-depleted mice (Fig. 2C). Taken

together, these data suggest that testosterone enhances fecal shedding and promotes viral

dissemination of CVB3 to peripheral tissues following replication in the intestine.

The inflammatory response to CVB3 is enhanced in mice with testosterone. The immune

response plays a pivotal role in tissue damage following CVB3 infection (27, 39-43). Previously,

we observed a systemic immune response to CVB3 in male mice as measured by serum cytokine

and chemokine levels (38). To test whether testosterone altered the magnitude of the cytokine

and chemokine response, we measured the serum concentrations of 32 cytokine and chemokines

at 3 dpi. In mock castrated mice, we saw an overall increase in the serum concentrations of pro-

inflammatory cytokines and chemokines IL-6, IFN-γ, TNF-α, IP-10, MCP-1, and RANTES

compared to uninfected males (Fig 3). The only cytokines and chemokines significantly altered

by testosterone were KC and G-CSF, which were found in greater serum concentrations in mock

castrated and testosterone-treated mice than uninfected mice. In contrast, testosterone-depleted

mice had a decrease in serum cytokine and chemokine concentration levels consistent with lower

levels of viral replication in the peripheral tissues. These data indicate that male $Ifnar^{-/-}$ mice

with testosterone elicit a more robust immune response than testosterone-depleted mice

following oral inoculation of CVB3.

CVB3 does not induce liver damage or cause pancreatitis in $Ifnar^{-/-}$ mice following oral

inoculation. We found that CVB3 tissue titers were the highest in the liver and pancreas

following CVB3 inoculation; therefore, we hypothesized that testosterone could enhance liver

necrosis and/or pancreatitis in $Ifnar^{-/-}$ mice. First, to test this hypothesis, we examined alanine

aminotransferase (ALT) and lipase levels in the serum by ELISA at 3 dpi to determine liver and

pancreas damage, respectively. We observed no significant increase in ALT levels or lipase

levels in the serum of mock or castrated male mice following oral inoculation (Fig. 4A and 4B).

Next, we examined histological sections of the liver and the pancreas from male mice sacrificed

at 3 dpi. Similar to ELISA results, we observed no evidence of hepatic necrosis or inflammation

in the liver (Fig. 4C). Further, we did not observe a loss of acinar cells, beta cells, or

inflammation in the pancreas (Fig. 4D). These data indicate that CVB3-Nancy does not induce

liver damage or pancreatitis after oral inoculation of *Ifnar*[-/-] mice.

Testosterone does not enhance CVB3-induced lethality in female mice but does enhance

fecal shedding and viral dissemination. Previously we found that orally CVB3 inoculated

female *Ifnar*[-/-] mice were protected entirely from CVB3-induced lethality and had limited viral

fecal shedding (38). Since exogenous testosterone treatment restored fecal shedding and lethality

in male mice, we hypothesized that testosterone might enhance pathogenesis in female mice. To

examine this hypothesis, we provided exogenous testosterone or placebo capsules to female mice

before oral inoculation with CVB3. Like our castration studies, female mice with testosterone

had significantly higher serum testosterone concentrations than placebo controls (Fig. 5A). One

week after hormone treatment, female mice were orally inoculated with 5×10^7 PFUs of CVB3

and monitored for survival. In contrast to males, both testosterone- and placebo-treated infected

females were protected from CVB3-induced lethality (Fig. 5B). Next, to examine fecal CVB3

shedding, feces were collected from infected mice at 1, 2, and 3 dpi, processed, and quantified by

a standard plaque assay. We observed significantly more CVB3 in the feces at 3 dpi in female

mice that received testosterone than female mice that received placebo capsules (Fig. 5C).

Finally, we examined CVB3 titers in peripheral tissues at 3 dpi to test if testosterone enhanced

viral dissemination. We found a significant increase in the viral loads of the heart, kidney, and

spleen in testosterone-treated female mice (Fig. 5D). The viral load in the liver was higher in

testosterone-treated mice but did not reach statistical significance. In contrast to the other organs,

the viral load in the pancreas was similar between placebo- and testosterone-treated female mice.

Overall, these data indicate that testosterone can enhance fecal CVB3 shedding and viral

dissemination to the heart, kidney, and spleen; however, testosterone does not contribute to

CVB3-induced lethality in female mice.

Discussion

Sexual dimorphism is commonly observed in various infectious, autoimmune, and

cardiovascular diseases. Males are often more susceptible to diseases caused by bacteria, viruses,

fungi, or parasites. Females, however, are more predisposed to autoimmune diseases like

systemic lupus erythematosus, multiple sclerosis, and rheumatoid arthritis. Additionally, clinical

and epidemiological studies have highlighted the disparity in cardiovascular diseases with an

increased mortality rate in men. CVB3 also displays a sex bias in human infections where males

are twice as likely to have severe sequelae. Previously, we observed a sex bias in CVB3 orally

inoculated male and female *Ifnar*[-/-] mice, consistent with human infections (38). Here we show

that testosterone, the primary sex hormone in males, impacts CVB3 pathogenesis following oral

infection.

Our data indicate that testosterone enhances intestinal replication of CVB3. We observed that

testosterone could promote fecal shedding of CVB3 in both male and female *Ifnar*[-/-] mice (Figs.

2A and 5C). To our knowledge, this is the first time that testosterone has been demonstrated to

influence the intestinal replication of an enteric virus. These data differ from previous studies

using systemic models of CVB3 infection. In this model, castration of wild-type, immune-

competent male mice did not impact viral replication in the heart (30). This difference may be

due to a sex bias in organ-specific replication of CVB3 and dependent on the type I IFN

response. In agreement, we have previously shown that while CVB3 replicates better in the

intestine of wild-type male mice, CVB3 replicated equally in the organs of wild-type male and

female mice following systemic infection. Further, the loss of type 1 IFN also facilitates sex-

dependent replication in the liver and spleen (38). Overall, these data suggest that organ-specific

androgen signaling may play a significant role in CVB3 replication.

Unfortunately, the mechanism for the sex hormone modulation of intestinal replication is

unclear. Previous data indicate that sex hormones, including testosterone, can enhance CVB3

attachment to cardiomyocytes (26); therefore, testosterone may alter the expression of viral

receptors on intestinal cells. Additionally, testosterone can interact with immune cells and

suppress inflammation (30, 44-47). It is possible that testosterone dampens the innate immune

response in the intestine, promoting initial viral replication in the intestine. Our data indicate that

the loss of testosterone broadly reduces the systemic cytokine and chemokine response to CVB3

(Fig. 3). However, it is unclear if this reduction is due to loss of testosterone or limited CVB3

replication and dissemination. Whether testosterone influences CVB3 binding to intestinal

epithelial cells or alters the intestinal immune response during initial viral replication warrants

further study.

Recent data also show that intestinal bacteria can influence the intestinal replication of enteric

viruses, including CVB3 (48-50). We have recently demonstrated that specific intestinal bacteria

in male mice play a role in promoting CVB3 infectivity and viral stability (51). While the

mechanism is unclear for CVB3, a similar enterovirus, poliovirus, has been found to use bacteria

to enhance attachment to the poliovirus receptor (49, 52). Interestingly, sex-specific differences

in intestinal bacteria can account for the sex bias in mouse models of type-1 diabetes (53, 54).

Since bacteria can also metabolize hormones that alter bacterial growth (55), it is intriguing to

speculate that sex hormones may change the overall intestinal ecosystem to directly or indirectly

impact CVB3 replication.

With a systemic infection model of CVB3, previous studies have found that testosterone

enhances lethality in male mice (25), consistent with our data. Castrating male mice, or by using

flutamide, an androgen receptor antagonist, we were also able to rescue mice from CVB3-

induced lethality (Fig. 1B and 1C). Interestingly, testosterone increased CVB3 shedding and

viral dissemination in female mice in our model, but testosterone did not enhance lethality (Fig.

5). In agreement with our data, systemic models of CVB3 in infection have shown that while

testosterone treatment promoted cardiac inflammation in female mice following CVB3 infection,

testosterone did not increase mortality in female mice (25, 27). These data suggest that other sex-

dependent factors may play a role in clearing the viral infection in female mice. Previous studies

have shown that the Y chromosome contributes to CVB3 pathogenesis (31, 32); therefore, genes

on the sex chromosomes may represent one of these critical factors. Future studies are required

to identify Y-specific genes and other potential correlates of immune protection that may

promote or limit CVB3 pathogenesis in mice.

The cause of death in our oral inoculation model remains unknown. CVB3 is the primary cause

of viral myocarditis. However, in C57BL/6 mice, CVB3 also causes hepatic damage and

pancreatitis (56-59). Since we observed high viral loads in the liver and pancreas of testosterone-

treated animals (Fig 2C), we hypothesized that the male mice died of hepatitis or pancreatitis.

However, contrary to our hypothesis, we did not observe inflammation or tissue damage in the

liver or pancreas following infection (Fig 4C and 4D). We confirmed these histology data by

serum ALT and Lipase concentrations at 3 dpi (Fig 4A and 4B). The cause of death is still under

investigation; however, enteric viruses, including poliovirus and Coxsackievirus, can cross the

blood-brain barrier, infect the central nervous system, and cause viral meningitis and encephalitis

(60, 61). Further studies are required to determine viral loads in the central nervous system and

resolve the determinants of lethality in the oral inoculation model.

In conclusion, we found that testosterone promotes intestinal CVB3 replication in orally

inoculated male and female *Ifnar*$^{-/-}$ mice. Further, testosterone enhances viral-induced lethality in

a sex-dependent manner. The exact mechanism of how testosterone aggravates disease is

unclear, but future studies will be necessary to determine how this sex hormone promotes

intestinal CVB3 infection. Overall, these data reinforce the importance of sex as a biological

variable in enteric viral infections.

Materials and Methods

Cells and virus

HeLa cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% calf serum and 1% penicillin-streptomycin at 37˚C with 5% CO_2. The CVB3-Nancy infectious clone was obtained from Marco Vignuzzi (Pasteur Institute, Paris, France) and propagated in HeLa cells as described previously (38). CVB3 was quantified by a standard plaque assay using HeLa cells.

Mouse experiments

All animals were handled according to the Guide for the Care of Laboratory Animals of the National Institutes of Health. All mouse studies were performed at Indiana University School of Medicine using protocols approved by the local Institutional Animal Care and Use Committee in a manner designated to minimalize pain, and any animals that exhibited severe disease were euthanized immediately by CO_2 inhalation. C57BL/6 $PVR^{+/+}$ $Ifnar^{-/-}$ mice were obtained from S. Koike (Tokyo, Japan) (62). One week post hormone treatment, mice were orally infected with 5x10^7 PFU of CVB3 IC Nancy. All adult experimental mice were 10-15 weeks old at the time of infection. Feces from infected mice were collected 1, 2, and 3 days post-infection, processed as previously described, and the fecal virus was quantified by a standard plaque assay (38).

Castration and hormone manipulation

Eight-week to ten-week-old male mice were put under anesthesia, and their testes were surgically removed or mock castrated as a surgical control. Testosterone implants were

constructed using silastic tubing (inner diameter-.078", outer diameter-.125"; Dow Chemical

Company). After placing 7.5mm of Crystalline Testosterone (Sigma Aldrich) in the tubing, the

ends of the tubing were sealed with 2.5 mm of medical adhesive (732 Multi-Purpose Sealant,

Dowsil). After the medical adhesive dried, the implants were incubated at 37°C overnight in

sterile phosphate-buffered saline. To ensure osmoregulation, implants that were found floating

were discarded. Due to the light sensitivity of sex steroid hormones, the testosterone implants

were concealed from light. One-week post castration, castrated mice were administered either

testosterone or placebo capsules subcutaneously under the right shoulder. Mock-castrated mice

were given placebo capsules. Testosterone levels of mouse serum were quantified using a

rat/mouse testosterone ELISA following the manufacturer's instructions (MP Biomedical).

Flutamide (TCI America, Portland, OR) was dissolved in 100% DMSO and diluted in double-

distilled water. As a vehicle control, 100% DMSO was also diluted in double-distilled water

before administration. Male mice were given 25µg/kg of flutamide or vehicle control by

intraperitoneal injection for three consecutive days before oral inoculation with CVB3-Nancy.

Tissue collection and histological analysis

The heart, liver, spleen, kidneys, and pancreas were aseptically collected 3 days post-infection

and homogenized in phosphate-buffered saline using 0.9-2.0 mm stainless steel beads in a Bullet

Blender (Next Advance). Cellular debris was removed by centrifugation at 12,000xg for 10 min

at 4°C, supernatants were collected, and CVB3 was quantified by plaque assay on HeLa cells.

Additionally, a portion of the aseptically removed organs were cut, washed in 1XPBS, put in

histology cassettes, and fixed overnight using 10% Neutral buffered formalin (NBF). The

samples were transferred to 70% Ethanol. The tissues were then paraffin-embedded, sectioned,

and stained with hematoxylin and eosin (H&E) by the Indiana University Histology Core.

Serum collection and analysis

Blood was collected from the inferior vena cava from infected or uninfected male and female

mice 3 days post-infection and incubated at room temperature for 30 mins to initiate coagulation.

Samples were centrifuged at 2000 rpm for 15 mins, and separated serum was collected and

stored at -20 C for downstream analysis. Alanine transaminase (EN0207Mu-1, CusaBio

Technologies) and Pancreatic lipase (E91453Mu-1, Cusabio Technologies) were measured by

ELISA. Serum cytokine levels were measured by the Indiana University Multiplex Analysis

Core using a Millipore Milliplex MAP Mouse Cytokine/Chemokine Magnetic Kit (Millipore

Sigma, Burlington, MA).

Statistical Analysis

Comparisons between control and study groups were analyzed using either a Mann-Whitney U

test or a one-way analysis of variance (ANOVA). A Log-rank test was used for survival curve

analysis. Error bars in the figures represent the standard errors of the means. A p-value <0.05

was considered significant. All analyses were performed using GraphPad Prism 9 (GraphPad

Software, La Jolla, CA).

Acknowledgments

We would like to thank Samantha Scholz, DVM and Keely Szilagyi, DVM for their help and

input with the mouse castration and testosterone experiments. This work is funded by a K01

DK110216, R03 DK124749, a Showalter Trust Award, and a Biomedical Research Grant from

the Indiana Clinical and Translational Sciences Institute to CMR.

References

1. Fuller AC, Kang B, Kang HK, Yahikozowa H, Dal Canto MC, Kim BS. 2005. Gender bias in
Theiler's virus-induced demyelinating disease correlates with the level of antiviral
immune responses. J Immunol 175:3955-63.
2. Shah VA, Chong CY, Chan KP, Ng W, Ling AE. 2003. Clinical characteristics of an outbreak
of hand, foot and mouth disease in Singapore. Ann Acad Med Singapore 32:381-7.
3. Aminu M, Ahmad AA, Umoh JU, de Beer MC, Esona MD, Steele AD. 2007. Adenovirus
infection in children with diarrhea disease in Northwestern Nigeria. Ann Afr Med 6:168-
73.
4. Curiel RE, Miller MH, Ishikawa R, Thomas DC, Bigley NJ. 1993. Does the gender
difference in interferon production seen in picornavirus-infected spleen cell cultures
from ICR Swiss mice have any in vivo significance? J Interferon Res 13:387-95.
5. Hou J, Zheng WF. 1988. Effect of sex hormones on NK and ADCC activity of mice. Int J
Immunopharmacol 10:15-22.
6. Rettew JA, Huet-Hudson YM, Marriott I. 2008. Testosterone reduces macrophage
expression in the mouse of toll-like receptor 4, a trigger for inflammation and innate
immunity. Biol Reprod 78:432-7.
7. McKay LI, Cidlowski JA. 1999. Molecular control of immune/inflammatory responses:
interactions between nuclear factor-kappa B and steroid receptor-signaling pathways.
Endocr Rev 20:435-59.
8. Furman D, Hejblum BP, Simon N, Jojic V, Dekker CL, Thiebaut R, Tibshirani RJ, Davis MM.
2014. Systems analysis of sex differences reveals an immunosuppressive role for
testosterone in the response to influenza vaccination. Proc Natl Acad Sci U S A 111:869-
74.
9. Trigunaite A, Dimo J, Jorgensen TN. 2015. Suppressive effects of androgens on the
immune system. Cell Immunol 294:87-94.
10. Brotfain E, Gruenbaum SE, Boyko M, Kutz R, Zlotnik A, Klein M. 2016. Neuroprotection
by Estrogen and Progesterone in Traumatic Brain Injury and Spinal Cord Injury. Curr
Neuropharmacol 14:641-53.
11. Salem ML, Hossain MS, Nomoto K. 2000. Mediation of the immunomodulatory effect of
beta-estradiol on inflammatory responses by inhibition of recruitment and activation of
inflammatory cells and their gene expression of TNF-alpha and IFN-gamma. Int Arch
Allergy Immunol 121:235-45.
12. Straub RH. 2007. The complex role of estrogens in inflammation. Endocr Rev 28:521-74.
13. Sorachi K, Kumagai S, Sugita M, Yodoi J, Imura H. 1993. Enhancing effect of 17 beta-
estradiol on human NK cell activity. Immunol Lett 36:31-5.
14. Zhou N, Yue Y, Xiong S. 2018. Sex Hormone Contributes to Sexually Dimorphic
Susceptibility in CVB3-Induced Viral Myocarditis via Modulating IFN-gamma(+) NK Cell
Production. Can J Cardiol 34:492-501.
15. Chakrabarti S, Lekontseva O, Davidge ST. 2008. Estrogen is a modulator of vascular
inflammation. IUBMB Life 60:376-82.
16. Murphy E. 2011. Estrogen signaling and cardiovascular disease. Circ Res 109:687-96.

17.   Kosrirukvongs P, Kanyok R, Sitritantikorn S, Wasi C. 1996. Acute hemorrhagic
conjunctivitis outbreak in Thailand, 1992. Southeast Asian J Trop Med Public Health
27:244-9.
18.   He SJ, Han JF, Ding XX, Wang YD, Qin CF. 2013. Characterization of enterovirus 71 and
coxsackievirus A16 isolated in hand, foot, and mouth disease patients in Guangdong,
2010. Int J Infect Dis 17:e1025-30.
19.   Andreoletti L, Leveque N, Boulagnon C, Brasselet C, Fornes P. 2009. Viral causes of
human myocarditis. Arch Cardiovasc Dis 102:559-68.
20.   Khetsuriani N, Lamonte A, Oberste MS, Pallansch M. 2006. Neonatal enterovirus
infections reported to the national enterovirus surveillance system in the United States,
1983-2003. Pediatr Infect Dis J 25:889-93.
21.   Khetsuriani N, Lamonte-Fowlkes A, Oberst S, Pallansch MA, Centers for Disease C,
Prevention. 2006. Enterovirus surveillance--United States, 1970-2005. MMWR Surveill
Summ 55:1-20.
22.   Abedi GR, Watson JT, Nix WA, Oberste MS, Gerber SI. 2018. Enterovirus and
Parechovirus Surveillance - United States, 2014-2016. MMWR Morb Mortal Wkly Rep
67:515-518.
23.   Fairweather D, Cooper LT, Jr., Blauwet LA. 2013. Sex and gender differences in
myocarditis and dilated cardiomyopathy. Curr Probl Cardiol 38:7-46.
24.   Woodruff JF. 1980. Viral myocarditis. A review. Am J Pathol 101:425-84.
25.   Huber SA, Job LP, Auld KR. 1982. Influence of sex hormones on Coxsackie B-3 virus
infection in Balb/c mice. Cell Immunol 67:173-9.
26.   Lyden DC, Olszewski J, Feran M, Job LP, Huber SA. 1987. Coxsackievirus B-3-induced
myocarditis. Effect of sex steroids on viremia and infectivity of cardiocytes. Am J Pathol
126:432-8.
27.   Huber SA, Kupperman J, Newell MK. 1999. Hormonal regulation of CD4(+) T-cell
responses in Coxsackievirus B3-induced myocarditis in mice. J Virol 73:4689-95.
28.   Huber SA. 2008. Coxsackievirus B3-induced myocarditis: infection of females during the
estrus phase of the ovarian cycle leads to activation of T regulatory cells. Virology
378:292-8.
29.   Huber SA, Pfaeffle B. 1994. Differential Th1 and Th2 cell responses in male and female
BALB/c mice infected with coxsackievirus group B type 3. J Virol 68:5126-32.
30.   Frisancho-Kiss S, Coronado MJ, Frisancho JA, Lau VM, Rose NR, Klein SL, Fairweather D.
2009. Gonadectomy of male BALB/c mice increases Tim-3(+) alternatively activated M2
macrophages, Tim-3(+) T cells, Th2 cells and Treg in the heart during acute
coxsackievirus-induced myocarditis. Brain Behav Immun 23:649-57.
31.   Robinson DP, Huber SA, Moussawi M, Roberts B, Teuscher C, Watkins R, Arnold AP, Klein
SL. 2011. Sex chromosome complement contributes to sex differences in coxsackievirus
B3 but not influenza A virus pathogenesis. Biol Sex Differ 2:8.
32.   Case LK, Toussaint L, Moussawi M, Roberts B, Saligrama N, Brossay L, Huber SA,
Teuscher C. 2012. Chromosome y regulates survival following murine Coxsackievirus b3
infection. G3 (Bethesda) 2:115-21.
33.   Bopegamage S, Kovacova J, Vargova A, Motusova J, Petrovicova A, Benkovicova M,
Gomolcak P, Bakkers J, van Kuppeveld F, Melchers WJ, Galama JM. 2005. Coxsackie B

476   virus infection of mice: inoculation by the oral route protects the pancreas from
477   damage, but not from infection. J Gen Virol 86:3271-80.

478   34.   Bopegamage S, Borsanyiova M, Vargova A, Petrovicova A, Benkovicova M, Gomolcak P.
479   2003. Coxsackievirus infection of mice. I. Viral kinetics and histopathological changes in
480   mice experimentally infected with coxsackieviruses B3 and B4 by oral route. Acta Virol
481   47:245-51.

482   35.   Harrath R, Bourlet T, Delezay O, Douche-Aourik F, Omar S, Aouni M, Pozzetto B. 2004.
483   Coxsackievirus B3 replication and persistence in intestinal cells from mice infected orally
484   and in the human CaCo-2 cell line. J Med Virol 74:283-90.

485   36.   Loria RM, Kibrick S, Broitman SA. 1974. Peroral infection with group B coxsackievirus in
486   the adult mouse: protective functions of the gut. J Infect Dis 130:539-43.

487   37.   Loria RM, Kibrick S, Broitman SA. 1974. Peroral infection with group B coxsackievirus in
488   the newborn mouse: a model for human infection. J Infect Dis 130:225-230.

489   38.   Robinson CM, Wang Y, Pfeiffer JK. 2017. Sex-Dependent Intestinal Replication of an
490   Enteric Virus. J Virol 91.

491   39.   Huber SA, Job LP, Woodruff JF. 1981. Sex-related differences in the pattern of
492   Coxsackievirus B-3-induced immune spleen cell cytotoxicity against virus-infected
493   myofibers. Infect Immun 32:68-73.

494   40.   Estrin M, Huber SA. 1987. Coxsackievirus B3-induced myocarditis. Autoimmunity is
495   L3T4+ T helper cell and IL-2 independent in BALB/c mice. Am J Pathol 127:335-41.

496   41.   Henke A, Huber S, Stelzner A, Whitton JL. 1995. The role of CD8+ T lymphocytes in
497   Coxsackievirus B3-induced myocarditis. J Virol 69:6720-8.

498   42.   Frisancho-Kiss S, Davis SE, Nyland JF, Frisancho JA, Cihakova D, Barrett MA, Rose NR,
499   Fairweather D. 2007. Cutting edge: cross-regulation by TLR4 and T cell Ig mucin-3
500   determines sex differences in inflammatory heart disease. J Immunol 178:6710-4.

501   43.   Fairweather D, Rose NR. 2007. Coxsackievirus-induced myocarditis in mice: a model of
502   autoimmune disease for studying immunotoxicity. Methods 41:118-22.

503   44.   Trigunaite A, Dimo J, Jørgensen TN. 2015. Suppressive effects of androgens on the
504   immune system. Cellular Immunology 294:87-94.

505   45.   Araneo BA, Dowell T, Diegel M, Daynes RA. 1991. Dihydrotestosterone exerts a
506   depressive influence on the production of interleukin-4 (IL-4), IL-5, and gamma-
507   interferon, but not IL-2 by activated murine T cells. Blood 78:688-99.

508   46.   Frisancho-Kiss S, Nyland JF, Davis SE, Frisancho JA, Barrett MA, Rose NR, Fairweather D.
509   2006. Sex differences in Coxsackievirus B3-induced myocarditis: IL-12Rbeta1 signaling
510   and IFN-gamma increase inflammation in males independent from STAT4. Brain Res
511   1126:139-47.

512   47.   Kadel S, Kovats S. 2018. Sex Hormones Regulate Innate Immune Cells and Promote Sex
513   Differences in Respiratory Virus Infection. Frontiers in Immunology 9.

514   48.   Robinson CM, Woods Acevedo MA, McCune BT, Pfeiffer JK. 2019. Related Enteric
515   Viruses Have Different Requirements for Host Microbiota in Mice. J Virol 93.

516   49.   Kuss SK, Best GT, Etheredge CA, Pruijssers AJ, Frierson JM, Hooper LV, Dermody TS,
517   Pfeiffer JK. 2011. Intestinal microbiota promote enteric virus replication and systemic
518   pathogenesis. Science 334:249-52.

50.   Baldridge MT, Nice TJ, McCune BT, Yokoyama CC, Kambal A, Wheadon M, Diamond MS,
Ivanova Y, Artyomov M, Virgin HW. 2015. Commensal microbes and interferon-lambda
determine persistence of enteric murine norovirus infection. Science 347:266-9.
51.   Dhalech AH, Fuller TD, Robinson CM. 2021. Specific Bacterial Cell Wall Components
Influence the Stability of Coxsackievirus B3. J Virol 95:e0142421.
52.   Robinson CM, Jesudhasan PR, Pfeiffer JK. 2014. Bacterial lipopolysaccharide binding
enhances virion stability and promotes environmental fitness of an enteric virus. Cell
Host Microbe 15:36-46.
53.   Markle JG, Frank DN, Mortin-Toth S, Robertson CE, Feazel LM, Rolle-Kampczyk U, von
Bergen M, McCoy KD, Macpherson AJ, Danska JS. 2013. Sex differences in the gut
microbiome drive hormone-dependent regulation of autoimmunity. Science 339:1084-
8.
54.   Yurkovetskiy L, Burrows M, Khan AA, Graham L, Volchkov P, Becker L, Antonopoulos D,
Umesaki Y, Chervonsky AV. 2013. Gender bias in autoimmunity is influenced by
microbiota. Immunity 39:400-12.
55.   Kornman KS, Loesche WJ. 1982. Effects of estradiol and progesterone on Bacteroides
melaninogenicus and Bacteroides gingivalis. Infect Immun 35:256-63.
56.   Liu JY, Wang SM, Chen IC, Yu CK, Liu CC. 2013. Hepatic damage caused by coxsackievirus
B3 is dependent on age-related tissue tropisms associated with the coxsackievirus-
adenovirus receptor. Pathog Dis 68:52-60.
57.   Koestner W, Spanier J, Klause T, Tegtmeyer PK, Becker J, Herder V, Borst K, Todt D,
Lienenklaus S, Gerhauser I, Detje CN, Geffers R, Langereis MA, Vondran FWR, Yuan Q,
van Kuppeveld FJM, Ott M, Staeheli P, Steinmann E, Baumgartner W, Wacker F, Kalinke
U. 2018. Interferon-beta expression and type I interferon receptor signaling of
hepatocytes prevent hepatic necrosis and virus dissemination in Coxsackievirus B3-
infected mice. PLoS Pathog 14:e1007235.
58.   Alirezaei M, Flynn CT, Wood MR, Whitton JL. 2012. Pancreatic acinar cell-specific
autophagy disruption reduces coxsackievirus replication and pathogenesis in vivo. Cell
Host Microbe 11:298-305.
59.   Lasrado N, Gangaplara A, Massilamany C, Arumugam R, Shelbourn A, Rasquinha MT,
Basavalingappa RH, Delhon G, Xiang SH, Pattnaik AK, Steffen D, Reddy J. 2021.
Attenuated strain of CVB3 with a mutation in the CAR-interacting region protects
against both myocarditis and pancreatitis. Sci Rep 11:12432.
60.   Rhoades RE, Tabor-Godwin JM, Tsueng G, Feuer R. 2011. Enterovirus infections of the
central nervous system. Virology 411:288-305.
61.   Sin J, Mangale V, Thienphrapa W, Gottlieb RA, Feuer R. 2015. Recent progress in
understanding coxsackievirus replication, dissemination, and pathogenesis. Virology
484:288-304.
62.   Ida-Hosonuma M, Iwasaki T, Yoshikawa T, Nagata N, Sato Y, Sata T, Yoneyama M, Fujita
T, Taya C, Yonekawa H, Koike S. 2005. The alpha/beta interferon response controls
tissue tropism and pathogenicity of poliovirus. J Virol 79:4460-9.
560

Figure Legends

Figure 1. CVB3-induced lethality is enhanced by testosterone. (A) Serum testosterone

concentrations in mock or castrated male C57BL/6 *Ifnar*[-/-] provided either placebo or

testosterone capsules. **$p<0.01$, one-way ANOVA. (B) Survival of mock castrated + placebo,

castrated + placebo, and castrated + testosterone mice after oral inoculation with $5x10^7$ PFU of

CVB3-Nancy. n=8-9 mice per group. **$p<0.01$, ***$p<0.001$, Log-rank test. (C) Survival of male

Ifnar[-/-] mice treated with flutamide or vehicle control after oral inoculation with $5x10^7$ PFU of

CVB3-Nancy. n=8-9 mice per group. *$p<0.05$ Log-rank test.

Figure 2. CVB3 fecal shedding and tissue titers following oral inoculation. Mice were orally

inoculated with $5x10^7$ PFU of CVB3-Nancy. (A) CVB3-Nancy fecal titers in mock castrated,

castrated + placebo, and castrated + mice. (B) CVB3-Nancy fecal titers in mock castrated,

castrated + placebo, and castrated + mice at days 2 and 3 post-inoculation. *$p<0.05$, **$p<0.01$,

Kruskal-Wallis test. (C) CVB3-Nancy tissue titers. Mice were euthanized at 3 dpi, and tissues

were collected in mock castrated (blue), castrated + placebo (gray), and castrated + testosterone

(blue and gray) *Ifnar*[-/-] mice.

Figure 3. The inflammatory response to CVB3 is enhanced in mice with testosterone. Serum

cytokine quantification in uninfected, mock castrated, castrated + placebo, and castrated +

testosterone *Ifnar*[-/-] mice. Infected mice were orally inoculated with $5x10^7$ PFUs of CVB3-

Nancy, and serum was collected 3 dpi.

Figure 4. Oral inoculation of *Ifnar*[-/-] mice with CVB3-Nancy does not induce liver damage or

cause pancreatitis. Serum levels of (A) ALT and (B) pancreatic lipase were quantified by ELISA

at 3 dpi. One-way ANOVA. H&E stained (C) liver and (D) pancreas collected at 3 dpi following

oral inoculation of mock castrated, castrated + placebo, and castrated + testosterone male *Ifnar*[-/-]

mice. Histology is representative of two experiments and at least 4 mice per group.

Figure 5. Testosterone promotes CVB3 shedding and dissemination in female *Ifnar*[-/-] mice, but

not lethality. (A) Serum testosterone concentrations in female C57BL/6 *Ifnar*[-/-] provided either

placebo or testosterone capsules. (B) Survival of female C57BL/6 *Ifnar*[-/-] provided either placebo

or testosterone capsules after oral inoculation with 5×10^7 PFU of CVB3-Nancy. (C) CVB3-

Nancy fecal titers in female C57BL/6 *Ifnar*[-/-] provided either placebo or testosterone. *p<0.05,

Mann-Whitney test. (D) CVB3-Nancy tissue titers in female C57BL/6 *Ifnar*[-/-] provided either

placebo or testosterone. *p<0.05, **p<0.01, Mann-Whitney test.

Figure 1

A.

B.

C.

Figure 2

A.

B.

C.

Figure 3

Figure 4

Figure 5

A.

B.

C.

D.