Long read sequencing reveals sequential complex rearrangements driven by Hepatitis B virus integration

Songbo Wang[1,2], Jiadong Lin[1,2], Xiaofei Yang[2,3], Zihang Li[4], Tun Xu[1,2], Peng Jia[1,2], Tingjie Wang[1,2], Bo Wang[5], Liangshuo Hu[5], Kai Ye[1,2,4,6,7,*]

[1]School of Automation Science and Engineering, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, Shaanxi, China

[2]MOE Key Lab for Intelligent Networks & Networks Security, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, Shaanxi, China.

[3]School of Computer Science and Technology, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, Shaanxi, China

[4]School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, Shaanxi, China

[5]The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaanxi, China.

[6]Faculty of Science, Leiden University, Leiden, The Netherlands

[7]Genome Institute, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, Shaanxi, China.

[*]Correspondence to Kai Ye, E-mail: kaiye@xjtu.edu.cn

Integration of Hepatitis B (HBV) virus into human genome disrupts genetic structures and

cellular functions. Here, we conducted multiplatform long read sequencing on two cell lines

and five clinical samples of HBV-induced hepatocellular carcinomas (HCC). We resolved

two types of complex viral integration induced genome rearrangements and established a

Time-phased Integration and Rearrangement Model (TIRM) to depict their formation

progress by differentiating inserted HBV copies with HiFi long reads. We showed that the

two complex types were initialized from focal replacements and the fragile virus-human

junctions triggered subsequent rearrangements. We further revealed that these

rearrangements promoted a prevalent loss-of-heterozygosity at chr4q, accounting for 19.5%

of HCC samples in ICGC cohort and contributing to immune and metabolic dysfunction.

Overall, our long read based analysis reveals a novel sequential rearrangement progress

driven by HBV integration, hinting the structural and functional implications on human

genomes.

Persistent hepatitis B virus (HBV) infection is reported to be responsible for >50% of all

hepatocellular carcinoma (HCC) worldwide[1,2]. Chronic liver inflammation and cell proliferation

promoted by viral proteins such as HBx have been described as main molecular mechanism of

HBV-induced HCC (HBV-HCC)[3]. Besides, integration of HBV fragments into human genome,

which may activate host genes and cause genome instability, happens in more than half of liver

cancers (36 out of 61 samples[4]) and is considered as a causal mechanism that lead to

carcinogenesis[5].

Several studies utilized short read sequencing (SRS) to survey the genomic and oncogenic

preference of HBV integrations sites[6-8]. However, due to the short read length (typically 100-

200bp), SRS detects single human-virus junction but fails to reveal the full-picture patterns. With

the advantages of long read sequencing (LRS) such as CLR and ONT, the genomic complexity

caused by HBV integration has been classified into either intra- or inter-chromosome

alterations[1,9,10]. One typical intra-chromosome integration directly affects cancer driver genes such

as TERT and MYC[1,11]. For inter-chromosome integrations, translocations, inducing focal gain,

deletion and generation of dicentric chromosome, have been characterized as main patterns.

Nevertheless, these studies have shown the association between viral integration and human

genome structural rearrangements, we still lack a full picture of how virus disrupt the genome due

to small sample number and the high error-prone CLR or ONT, which often lead to detection

artifacts from studies with low-coverage data (less than 20X)[9,10]. Therefore, long and accurate HiFi

long read sequencing would be ideal to characterize HBV associated genome rearrangements and

their underlying mechanisms, for better diagnosis, prognosis and treatment.

In this study, we applied multi-platform LRS on HBV-integrated tumor samples, focused on

identifying comprehensive and novel HBV integration patterns as well as their underlying

mechanisms and impact on human genome. In total 7 samples, consisting of 2 HBV-HCC cell

lines (SNU-182 and SNU-387) and 5 clinical samples, were sequenced with the most advanced

LRS technologies such as CLR, ONT and CCS (**Supplementary Table1**). Collectively, in our

data, we identified 17 full-view HBV integration events (**Methods, Supplementary Fig.1**), which

were classified into three types (**Figure 1a**). Type I was characterized as viral genome fragmented

and inserted into human genome through simple and focal replacement. Type II involved complex

and HBV-induced unbalanced translocations while Type III referred to the most complex event,

chromothripsis. Specifically, these two novel complex patterns as well as their formation

mechanisms were introduced and we proposed a Time-phased Integration and Rearrangement

Model (TIRM for short) to describe their formation progress from Type I (**Figure 1b**). Finally, we

utilized HCC WGS and RNA-seq data from ICGC cohort to explore the prevalence of our

discoveries and their implications on immune and metabolic systems.

In type I integration mode, the focal replacement events (2 out of 17) found in our samples focally

disrupted human genomes by replacing short human sequence with HBV fragments. For instance,

two HBV fragments replaced about 70bp of intergenic region between FAM83F and TNRC6B

(chr22:40,044,376-40,044,446) in cell line SNU-387 and therefore, we could observe two human-

virus junctions (chr22: 40,044,376-HBV: 1832, HBV: 570-chr22: 40,044,446) in one single LRS

read simultaneously (**Supplementary Fig.2**). Although FAM83F and TNRC6B have been

associated with tumorigenesis, their expression was not altered compared with other HCC samples.

Besides, based on the HBV integration patterns observed in LRS data, we deduced 3 potential

focal replacement events (**Supplementary Fig.2**) from published single human-virus junctions in

ICGC HCC dataset[4]. All 3 events located at gene intron regions and did not cause expression

alteration (**Supplementary Fig.2**). Those focal replacements were gene-related integrations but

were different from promoter- or exon-preferred integrations at TERT or KMT2B[1], which caused

significant overexpression or altered transcript. Therefore, these type I replacement events might

not directly contribute carcinogenesis but led to instable genome, especially fragile at around the
human-virus junctions. As a result, the initial step of our TIRM was accomplished and instable
genome promoted subsequent rearrangements.
The type II complex events (11 out of 17) caused large scale copy number variation (CNV) through
unbalanced-translocations. As a result, integrated HBV fragments fused two chromosomes
together. Both two virus-human junctions linking two chromosomes were right at the boundary of
significant CNAs, with one gain of heterozygosity (GOH) in one chromosome and one loss of
heterozygosity (LOH) in the other one. Among the 11 HBV-induced unbalanced translocation
events, we identified a novel pattern of HBV fragments from 8 events, in which two distinct copies
of HBV fragments harboring different single-nucleotide-variants (SNVs) were inserted between
two human chromosomes with different orientations. For instance, in Event3 from SNU-182,
chromosome 4 and 8 were linked together by four HBV fragments, two of which headed forward
while the other two headed reverse **(Figure 2a)**. Both forward and reversed fragments could form
a nearly complete HBV genome. Meanwhile, HiFi long reads facilitated the identification of SNVs,
differentiating forward and reversed fragments**,** indicating that the different-oriented fragments
were originated from two independent integration events during long-term and persistent infection.
Next, we built a phylogeny tree with all HBV fragments in cell line SNU-182 **(Methods, Figure**
2b). We noticed that HBV fragments from different integration events located at different branches
and nodes in the tree, indicating multiple different integration events in one sample at different
time points. Moreover, different-oriented fragments connecting two chromosomes (e.g.
chromosome 4 and 8 in Event3) even located on different branches, confirming our hypothesis
that current Event3 were initialized by two independent HBV integrations events and the inter-
translocation happened as a subsequent rearrangement.
Given the evidences above, we proposed the Time-phased Integration and Rearrangement Model
(TIRM for short, need rename) to illustrate the integration process and its potential formation
mechanism **(Figure 2c)**. Initially, during the persistent infection, HBV integrated into two different
chromosomes at two different time points through two separate DNA double-strand-breaks (DSB)
and its subsequent repair pathways, such as microhomology-mediated end joining (MMEJ)[12]. This
process would form two individual replacement events. Later, similar as the transposable elements
that make the genome unstable, the two integrations broke again and left two exposed DSBs at the
human-virus junctions. Finally, under the micro-homology mediated break-induced replication

(MMBIR)[13], the replication fork jumped from one DSB to the other to form a new chromosome,

resulting in GOH and LOH simultaneously on two different chromosomes. In general, the TIRM

well established the process of the initial integration and the following translocation step by step.

(A more complicated unbalanced translocation was found in Event9 of Patient1, three enhancer-

enriched segments from chr3, chr5 and chr12 were fused with 2 integrated HBV fragments

including open reading frames of HBx and Core **(Supplementary Fig.3)**. We assumed it as a novel

HBV-host DNA contact by hijacking human enhancers to form super enhancers, which have been

defined as spatial clustering of groups of traditional enhancers. As reported in human

papillomavirus 16 (HPV16), the super-enhancer, which is formed by integrated HPV16 segments

and human enhancers, forced the transcription of viral oncogenes[14]. In HBV, the similar enhancer-

contact was described in a 3D way between HBV cccDNA–host DNA[15]. Therefore, we found a

direct evidence that the enhancer-contact could be formed by integration and its subsequent

rearrangement.

The most complex integration pattern, integration-induced chromothripsis, was identified from 2

of our clinical samples. For instance, in Patient2, we observed chromosome-wide CNAs and 30

genomic rearrangements localized at chr14 and chr16 **(Figure 3a)**. Those rearrangements

frequently linked together copy number neutral, loss or amplified regions, indicating that chr14

and chr16 underwent a chromothripsis event. Moreover, HBV integrated into the boundary of the

rightmost amplified region and formed a symmetrical fold-back-inversion structure, which was a

strong evidence for breakage-fusion-bridge (BFB) mechanism[16]. Meanwhile, we also found

another fold-back-inversion structure at the boundary of the highest amplified region

(Supplementary Figure 4). We could infer that the second BFB happened later than the HBV-

mediate BFB by comparing the CNA around the breakpoints because the CN always doubled after

a BFB event.

The above evidences also implied a TIRM, starting with focal virus integration and resulting in

chromothripsis events **(Figure 3b)**. Specifically, at the beginning time point, HBV integrated into

chr14 in the way like replacement as well. Later, at the integration site, DSB happened and was

replicated in S phase. In G2, the two broken ends were glued together by non-homologous end-

joining (NHEJ) and formed a dicentric chromosome, which was cut off during mitosis. After that,

in the next cell cycle, the second BFB happened and the chromosome became so unstable that

chromothripsis eventually happened. Similar TIRM was also identified in Patient1 on chr7, where

the chromothripsis occurred after the HBV-induced BFB **(Supplementary Figure 5)**.

The whole-genome deep LRS provided us the opportunity to investigate the impacts of the two

complex rearrangements on human genome. Notably, we observed a frequent LOH at chr4's long

arm (chr4q-LOH for short), presenting in 5 out of 7 in our data. To reveal its prevalence in HCC,

we collected the HCC WGS data (n=303) as well as their CNA profile[17] across 2 projects (LIRI-

JP and LIHC-US) in ICGC cohort **(Supplementary Table2)**. We found ~19.5% HCC samples

(n=59) included chr4q-LOH breakpoints **(Figure 4a)**, exceeding that in ovarian cancer (11%)[18].

When considering viral conditions, ~12% of the NBNC-HCCs while ~32% of HBV-HCCs showed

chr4q-LOH, suggesting HBV as a promotion factor of chr4q-LOH. We also examined the

correlation between HBV integration and these LOHs. Although not all these chr4q-LOHs were

directly induced by HBV integration (3 out of 7 in our data, 8 out of 59 in ICGC), we considered

HBV could promoted the LOH by integration and subsequent rearrangement.

Next, we examined the enrichment and hotspots of these chr4q-LOH breakpoints. 16 out of 59

focally located at chr4q33-34, leading to frequent chr4q35-LOH (Cluster1, **Supplementary Fig**

6a). To reveal the implications of these LOHs on genes, we collected the corresponding transcript

read count matrix[17] from ICGC cohort (n=118). Previous reports found that genes located at

chr4q35, such as IRF2 and ING2, suffered frequent LOH and considered them as potential HCC

suppressor genes[19]. However, compared with chr4q-neutral samples, two genes located at chr4q35,

ACSL1 and F11, were observed significantly down-regulated (P.adj=0.004 and 0.002, respectively)

(Figure 4b). The roles of ACSL1 and F11 have been associated with immunological processes [20-

22] and low expression of the two genes were correlated with poor survival in HCC patients[23].

Therefore, down-regulation of the two genes implied that the host immune system might under a

disorder, which could also be proved by other 14 down-regulated immune-related genes such as

C1s, C2, and CFH and so on **(Figure 4c, Supplementary Table3)**.

We then performed GO and KEGG enrichment on all 357 down-regulated genes. KEGG item

complement and coagulation cascades was highly enriched (**Figure 4d**). HBV has been reported

to inhibit the expression of complement C3 and C4 to cause immune dysfunction by chronic and

persistent infection[24]. This enrichment suggested that the immune dysfunction might be worsen in

the effort of HBV integration and the subsequent genome instability at chr4q, which finally

contributed to carcinoma progress.

In addition to chr4q35-LOH, we discovered another enriched cluster at chr4q12-13 region, leading

a larger lost from chr4q13 to 3' telomere (37 out of 59, Cluster2). Similar gene expression and

pathway enrichment analysis were performed on Cluster2 samples as well. Except the two

immune-related genes mentioned above, metabolic genes such as UGT2B genes[25] and ADH1B [26]

that located after chr4q13 were significantly down-regulated **(Figure 4b)**. As a consequence, 22

metabolism pathways (12 in Cluster1) were enriched **(Supplementary Fig 6b)**. Since metabolic

alteration clearly characterized HCC[27], HBV-integration might worsen the aberrant metabolism in

the same way as that in immune dysfunction.

Gene analysis among Cluster1 and Cluster2 suggested that HBV integration could indirectly

disturb genes and affect immune and metabolic pathways through subsequent rearrangements. It

has been known that HBV integration was a random event[28]. However, integrating into specific

regions, such as chr4q12-13 and chr4q-33-34, would accelerate genome instability and

tumorigenesis. In general, the initial integration might not be deadly (none of them in exons or

promoters), but the subsequent rearrangement described in TIRM could alter gene copies and

granted cells with selective advantages to HCC.

In this study, utilizing LRS on HBV-HCC samples, we proposed the Time-phased Integration and

Rearrangement Model (TIRM) to demonstrate the formation process of the two complex types of

HBV-induced genome instability and their corresponding mechanisms, such as MMEJ, MMBIR

and BFB. The model characterized the initial stage of the integration-induced rearrangements as a

simple replacement mediated by MMEJ. Then MMBIR and BFB mechanisms were involved to

generate complex events. Moreover, we showed two frequent HBV-induced chr4-LOH regions

that play important roles on immune and metabolic functions, which may grant the cells selective

advantages and finally contribute to carcinoma process.

LRS technology allows us to characterize the full-picture of HBV integration. Although we self-

developed a few of scripts that could precisely identify HBV integration sites and patterns, a

complete algorithm that can not only detect integrations but reconstruct the genomes

rearrangements is still expected to carry on. Moreover, LRS is still incapable of covering entirely

the large-scale chromothripsis and we need reconstruct the event manually. Limited by the number

of our samples, we were incapable to discover the expression-level influence of the integrations

and subsequent rearrangements. Therefore, we collected RNA-seq samples from ICGC cohort.

ICGC HCC samples are sequenced with SRS technology, which may lead to missing integration

sites. That's the cause that we observed ~50% HBV integrations in chr4-LOH breakpoints in our

samples but that in ICGC cohort is low. Current studies, including the proposed TIRM, are based

on the end-state HCC. However, a better understanding of the sequential process and molecular

mechanisms of HBV integration requires a number of samples that undergo different infection and

carcinoma stages. These will also lead to a better understanding of the impact of integration on

both HBV-self and human. We also expect a comprehensive study that reveal the causation of the

integration, which is still obscured for now.

HBV integrations have been discussed and explored with SRS and LRS in the past decade. We

demonstrate that LRS especially the HiFi LRS can comprehensively characterize the novel

integration patterns, mechanisms and impacts on human genome. As a result, long-read HBV

integration studies will be essential to reveal their role in the genetics of HCC. Our work lays down

an important foundation for further large-scale HBV integration studies with LRS, making up the

disadvantages of SRS and investigating the full spectrum of HBV integrations.

    MAIN FIGURES

    Figure 1

Fig. 1 | Introduction of novel discoveries. a, Classification of found integration events. 3 major types of integration have been identified and their corresponding CNAs are shown in the line-plots. The 3 types are ranged from left to right, which accorded with their complexity from simple to complex. The black arrows indicated the formation results. **b**, The Time-phased Integration and Rearrangement Model (TIRM for short). During persistent infection, HBV evolved and might integrated into human genome by replacement at any time point (outside the

box). Later, through a new DSB, the integration site broke again and subsequent rearrangement

happened to form the complex Type2 and Type3 (inside the box).

Figure 2

Fig 2. | HBV-induced unbalanced translocation. a, Observed evidence from SNU-182-Event3,

including CNA and direct single read that covered the entire event. **b**, Phylogeny tree built with

all HBV fragments from SNU-182. Fragments from same event but with opposite orientations

were marked in same color. Fragments' coordinates were indicated by colored arrows. **c**,

Proposed TRIM to explain the formation of unbalanced translocation. First, two independent

replacement integrations happened with unique HBV fragments. Then, under the MMBIR

mechanism, the replication fork jump from one human-virus junction to another. Finally, the

unbalanced translocation was formed. As a result, blue and green chromosomes were linked

together by unique HBV fragments. The green chromosome suffered a LOH while the blue

chromosome suffered a GOH.

Figure 3

Fig 3 | HBV-induced chromothripsis. **a**, CNA and inter/intro-chromosome translocations in

Patient2. Two BFBs and HBV integration site were marked as non-gray arrows. **b**, Proposed

TIRM to explain the formation of chromothripsis. The two BFBs happened at human-virus

junction and human genome, respectively. Then the chromothripsis happened and the retained

fragments are re-connected by NHEJ to form the final new chromosome.

Fig. 4 | Analysis results from ICGC. **a**, Enrichment of LOH segments and breakpoints across ICGC HCC cohort. Liver tumors (HCC) had two significantly enriched breakpoints regions. **b**, Influence of LOH on ACSL1, F11, UGT2B7 and ADH1B. **c**, Volcano plot for differential gene expression in Cluster1. Immune-related genes were labeled. **d**, Scatter plot of KEGG pathway enrichment statistics. Pathways terms were ranged by gene ratio and top 20 were displayed in the figure.

Methods

Cell lines

SNU-182 and SNU-387 were acquired from the National Collection of Authenticated Cell Cultures. The SNU-387 were sequenced on the PacBio Sequel machine, producing the CLR reads while SNU-182 were sequenced on the PacBio Sequel II sequencer, generating HiFi reads. Briefly, libraries for single molecule real-time (SMRT) PacBio genome sequencing were constructed following the standard protocols. The high molecule genomic DNA was sheared to ~20Kbp targeted size, followed by damage repair and end repair, blunt-end adaptor ligation and size selection. The average CLR sequencing coverage was 45 reads per bp while that forThe average HiFi sequencing coverage was 24 reads per bp.

Additionally, the RNA sequencing of SNU-182 and SNU-387 were produced. A total amount of 1 g RNA per sample was used for library preparation, and the library was generated using NEBNext UltraTM RNA Library Prep Kit for Illumina (NEB, USA), following the manufacturer's recommendations. Then, the prepared library was sequenced on the Illumina Novaseq platform, generating the 150bp paired-end reads.

Clinical samples

Five primary HCC samples were collected from the First Affiliated Hospital of Xi'an Jiaotong University. Five collected tumor tissues were sequenced on the Nanopore PremethION sequencer according to the manufactures' instructions. In general, the long DNA fragments were qualified and selected using the PippinHT system (Sage Science, USA), and the A-ligation reaction were conducted with NEBNext Ultra II End Repair/dA-tailing Kit. The adapter in the SQK-LSK109 (Oxford Nanopore Technologies, UK) was used for further ligation reaction and DNA library was measured by Qubit 4.0 Fluorometer (Invitrogen, USA). Finally, about 700ng DNA library was constructed for each tumor tissue and sequenced with PremethION. The average ONT sequencing coverage was 45 reads per bp.

ICGC cohort

Project LIRI-JP and LIHC-US were selected from ICGC cohort (n=303). Filtering out those in black list, we total obtained 303 HCC samples, 118 of which have RNA-seq. The corresponding normal paired data were also collected. We utilized the corresponding profiles of CNA and

transcript read count aggregated by PCAWG working group. The viral condition for each sample

was also obtained from previous studies as well[4,23,29].

Characterization of HBV integration with LRS. HBV integration pattern characterization was

performed by a self-developed pipeline, including: (1) read mapping (2) candidate reads selection;

(3) cluster reads to events; and (4) classification of integration events

(1) Read mapping.

Raw reads from the LRS technologies were aligned to GRCh38 with HBV genome via

Minimap2[30]. The preset parameter of Minimap2, 'map-pb' and 'map-ont' were activated for

PacBio and Oxford Nanopore (ONT) reads, respectively. Besides soft-clip and MD tag were

set by –Y and –MD. Parameter –a and --eqx was set to output SAM format with =/X CIGAR

operators. Then we used SAMtools[31] to sort and index the output.

(2) Candidate reads selection.

We selected reads that span both human and viral genomes by traversing reads that partially

aligned on HBV genome. Minimum map quality was set to 20 to filter imprecise reads. For

each candidate read r_i, we collect all its primary and supplementary alignments and sorted

them by read coordinates. Therefore, r_i can be represented as all its alignments: $r_i =<$

$(chr1, start1, end1, T), (HBV, start2, end2, F), ... (chr3, start2, end2, T) >$, where T and F

denoted the mapping direction.

(3) Cluster reads to events.

We cluster candidate reads to integration events. Different reads but with similar alignments

were merged together based on their coordinates. We set the maximum difference between two

alignments to 50bp. When comparing two reads, we considered their orientations and toke the

minus strand into account. After merging, integration events with less than 5 supporting reads

were filtered out.

(4) Classification of integration events.

Integration events were classified to 3 types, including focal replacement, unbalanced

translocation and chromothripsis. Reads that supporting focal replacement usually included 4

alignments: the middle two aligned to HBV while the left and right segments aligned focally

and closely on the same chromosome. The unbalanced translocations featured as at least 3

alignments, two of which aligned to different chromosomes and the left aligned to HBV. The

chromothripsis always included fold back inversions formed by HBV fragments. As a result,

we could observed a nearly symmetrical arrangement of alignments. Accompanying with the

CNA and inter-/intra-translocations, we determined the chromothripsis event.

Phylogeny tree. We used HiFi sequenced SNU-182 to build a phylogeny tree. Firstly, we

separated apart the aligned HBV fragments by the four events. We also separated the different

oriented segments in one event. Because HBV always fragmented and some sequences might lost,

we then fixed the gap region with reference genome sequences. As a result, we obtained 7

individual HBV mutated genomes. We used MEGA-X[32] (GUI mode) to build a neighbor-joining

tree using maximum composite likelihood method. Both transitions and transversions of

substitutions were included. The test of phylogeny ran bootstrap method 500 replications.

Copy number variation analysis. For LRS, genome-wide profile of CNA was performed by

CNVkit[33] without paired control samples. We ran CNVkit under the default settings except the bin

size was set to 100Kbp, and copy number plots were created with the scatter option. In addition,

we applied the IGVtools[34] to all chromosomes at 100Kbp resolution, generating the binned whole

genome coverage. The binned coverage was saved as TDF format for further visualization. For

SRS (ICGC samples), CNV profiles of all samples were obtained from PWAWG working group.

The published results included total CN, major CN and minor CN of each chromosome segment.

We further implemented a simple script to visualize the results for better understanding.

LOH or GOH in ICGC samples were determined by relative change between two neighboring CN

segments. CN segments with length less that 10Kbp were excluded to avoid sequence coverage

bias. To avoid aneuploidies which were common in cancer, we strictly required that the CN change

must start from neutral CN (CN=2). Therefore, LOH meant CN change from 2 to 1 while GOH

meant CN change from 2 to 3. Besides, we required LOH or GOH to extend to the telomere.

Samples whose chr4 satisfied the above conditions were kept and the corresponding change

breakpoints were kept as well. Chr4-neutral samples' chr4 kept CN=2.

Chromothripsis detection. Chromothripsis featured as massive CNVs and inter-/intra-

chromosome translocations. CNV breakpoints of our samples were obtained from CNVkit

mentioned above. Translocation detection for chromothripsis followed a similar pipeline as that

for HBV integration. We collected all abnormal reads with two alignments aligned in different

chromosomes (inter-translocation) or in same chromosome but aligned two distant region (> 1Mb,

intra-translocation). After clustering single reads and filtering by minimum supporting reads

(default 5), we assigned each translocation breakpoint with CNV breakpoint by imitating the

coordinate's differences within 50bp. Those inter-/intra- translocations without CNV were

excluded. Finally, we visualize CNV and translocations together utilizing karyoploteR[35].

Gene expression and pathway enrichment. ICGC HCC samples were contributed into 4 groups.

Group1 and group2 were samples from chr4q-LOH breakpoint Custer1 and Cluter2, respectively.

Group3 included all chr4-netural HCC samples. Group4 was the set of all normal-paired samples

of group 1, 2 and 3. We performed two rounds of analysis: <group1, group3> and <group2, group3>

to reveal the impaction of different chr4q-LOH clusters on genes. Besides, we also performed

<group3, group4> as a control to understand the gene expression alteration after tumorigenesis.

Gene transcript read count matrix was obtained from PWAWG working group[17]. Differential

transcript expression between each two groups was analyzed by DESeq2[36]. Genes with P.adjust <

0.05 were considered as significantly altered. Besides, genes with log2FoldChange > 1 or <-1 were

set to 'UP' and 'DOWN', respectively for further pathway enrichment. For genes with more than

one transcript, we chose the most significant one and removed the left. The visualization (volcano

plot) was done by R package ggpubr (https://rpkgs.datanovia.com/ggpubr).

KEGG and Go pathway enrichments were perform with clusterProfiler[37]. Up- and down-regulated

genes were enriched separately. The p and q value cutoff were both set to 0.05. Pathway items that

satisfied the p and q value cutoff and had more than 10 genes enriched were kept and visualize by

R package enrichplot (https://yulab-smu.top/biomedical-knowledge-mining-book/).

Acknowledgements

K.Y. and X.Y. are supported by the National Science Foundation of China (32125009, 32070663,

62172325,), the Key Construction Program of the National '985' Project, the World-Class

Universities (Disciplines), and the Characteristic Development Guidance Funds for the Central

Universities and the Fundamental Research Funds for the Central Universities.

Author contributions

K.Y. designed and supervised the research. S.W. and J.L performed HBV integration analysis.

S.W., T.X. and T.W performed the gene expression analysis. B.W. and L.H. performed clinical

surgeries and offered the clinical HCC samples. J.P performed reads alignment. Z.L. performed

Sanger analysis of HBV-human junctions. K.Y., S.W., J.L. and X.Y. wrote the paper. All authors

read and proved the final version of this manuscript.

Competing interesting

The authors declare no competing interests.

REFERENCES

1. Peneau, C. *et al.* Hepatitis B virus integrations promote local and distant oncogenic driver alterations in hepatocellular carcinoma. *Gut* (2021).

2. Guthle, M. & Dollinger, M.M. Epidemiology and risk factors of hepatocellular carcinoma. *Radiologe* **54**, 654-+ (2014).

3. An, P., Xu, J.H., Yu, Y.Y. & Winkler, C.A. Host and Viral Genetic Variation in HBV-Related Hepatocellular Carcinoma. *Frontiers in Genetics* **9**(2018).

4. Zapatka, M. *et al.* The landscape of viral associations in human cancers. *Nature Genetics* **52**, 320-+ (2020).

5. Tarocchi, M., Polvani, S., Marroncini, G. & Galli, A. Molecular mechanism of hepatitis B virus-induced hepatocarcinogenesis. *World Journal of Gastroenterology* **20**, 11630-11640 (2014).

6. Sung, W.K. *et al.* Genome-wide survey of recurrent HBV integration in hepatocellular carcinoma. *Nature Genetics* **44**, 765-U188 (2012).

7. Zhao, L.H. *et al.* Genomic and oncogenic preference of HBV integration in hepatocellular carcinoma. *Nature Communications* **7**(2016).

8. Jiang, Z. *et al.* The effects of hepatitis B virus integration into the genomes of hepatocellular carcinoma patients. *Genome Res* **22**, 593-601 (2012).

9. Zhuo, Z. *et al.* Long-read sequencing reveals the structural complexity of genomic integration of HBV DNA in hepatocellular carcinoma. *NPJ Genom Med* **6**, 84 (2021).

10. Álvarez, E.G. *et al.* Aberrant integration of Hepatitis B virus DNA promotes major restructuring of human hepatocellular carcinoma genome architecture. *Nature Communications* **12**(2021).

11. Li, C.L. *et al.* Androgen Receptor Enhances Hepatic Telomerase Reverse Transcriptase Gene Transcription After Hepatitis B Virus Integration or Point Mutation in Promoter Region. *Hepatology* **69**, 498-512 (2019).

12. Tu, T., Budzinska, M.A., Shackel, N.A. & Urban, S. HBV DNA Integration: Molecular Mechanisms and Clinical Implications. *Viruses* **9**(2017).

13. Zhang, F. *et al.* The DNA replication FoSTeS/MMBIR mechanism can generate genomic, genic and exonic complex rearrangements in humans. *Nat Genet* **41**, 849-53 (2009).

14. Warburton, A. *et al.* HPV integration hijacks and multimerizes a cellular enhancer to generate a viral-cellular super-enhancer that drives high viral oncogene expression. *Plos Genetics* **14**(2018).

15. Yang, B. *et al.* 3D landscape of Hepatitis B virus interactions with human chromatins. *Cell Discov* **6**, 95 (2020).

16. Li, Y.L. *et al.* Constitutional and somatic rearrangement of chromosome 21 in acute lymphoblastic leukaemia. *Nature* **508**, 98-+ (2014).

17. Campbell, P.J. *et al.* Pan-cancer analysis of whole genomes. *Nature* **578**, 82-+ (2020).

18.   Wrzeszczynski, K.O. *et al.* Identification of Tumor Suppressors and Oncogenes from
Genomic and Epigenetic Features in Ovarian Cancer. *Plos One* **6**(2011).
19.   Zhang, H. *et al.* Analysis of loss of heterozygosity on chromosome 4q in
hepatocellular carcinoma using high-throughput SNP array. *Oncol Rep* **23**, 445-55
(2010).
20.   Kalugotla, G. *et al.* Frontline Science: Acyl-CoA synthetase 1 exacerbates lipotoxic
inflammasome activation in primary macrophages. *Journal of Leukocyte Biology*
106, 803-814 (2019).
21.   Roelands, J. *et al.* Long-Chain Acyl-CoA Synthetase 1 Role in Sepsis and Immunity:
Perspectives From a Parallel Review of Public Transcriptome Datasets and of the
Literature. *Frontiers in Immunology* **10**(2019).
22.   Deng, Y.B. *et al.* Long non-coding RNA F11-AS1 inhibits HBV-related
hepatocellular carcinoma progression by regulating NR1I3 via binding to
microRNA-211-5p. *Journal of Cellular and Molecular Medicine* **24**, 1848-1865
(2020).
23.   Lian, Q.Y. *et al.* HCCDB: A Database of Hepatocellular Carcinoma Expression Atlas.
Genomics Proteomics & Bioinformatics **16**, 269-275 (2018).
24.   Zhu, C. *et al.* Hepatitis B virus inhibits the expression of complement C3 and C4, in
vitro and in vivo. *Oncol Lett* **15**, 7459-7463 (2018).
25.   Lu, L.L. *et al.* Drug-Metabolizing Activity, Protein and Gene Expression of UDP-
Glucuronosyltransferases Are Significantly Altered in Hepatocellular Carcinoma
Patients. *Plos One* **10**(2015).
26.   Shih, S., Huang, Y.T. & Yang, H.I. A multiple mediator analysis approach to quantify
the effects of the ADH1B and ALDH2 genes on hepatocellular carcinoma risk.
Genetic Epidemiology **42**, 394-404 (2018).
27.   De Matteis, S. *et al.* Aberrant Metabolism in Hepatocellular Carcinoma Provides
Diagnostic and Therapeutic Opportunities. *Oxid Med Cell Longev* **2018**, 7512159
(2018).
28.   Budzinska, M.A., Shackel, N.A., Urban, S. & Tu, T. Cellular Genomic Sites of Hepatitis
B Virus DNA Integration. *Genes* **9**(2018).
29.   Furuta, M. *et al.* Characterization of HBV integration patterns and timing in liver
cancer and HBV-infected livers. *Oncotarget* **9**, 25075-25088 (2018).
30.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094-3100 (2018).
31.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**(2021).
32.   Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and*
Evolution **35**, 1547-1549 (2018).

33.   Talevich, E., Shain, A.H., Botton, T. & Bastian, B.C. CNVkit: Genome-Wide Copy
474   Number Detection and Visualization from Targeted DNA Sequencing. *Plos*
475   *Computational Biology* **12**(2016).
476   34.   Robinson, J.T. *et al.* Integrative genomics viewer. *Nature Biotechnology* **29**, 24-26
477   (2011).
478   35.   Gel, B. & Serra, E. karyoploteR: an R/Bioconductor package to plot customizable
479   genomes displaying arbitrary data. *Bioinformatics* **33**, 3088-3090 (2017).
480   36.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and
481   dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**(2014).
482   37.   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
483   data. *Innovation (N Y)* **2**, 100141 (2021).
484