1
2
3 **Lessons on fruiting body morphogenesis from genomes and**
4 **transcriptomes of Agaricomycetes**
5

6 László G. Nagy[1]*, Peter Jan Vonk[2], Markus Künzler[3], Csenge Földi[1], Máté Virágh[1], Robin A.
7 Ohm[2], Florian Hennicke[4], Balázs Bálint[1], Árpád Csernetics[1], Botond Hegedüs[1], Zhihao Hou[1],
8 Xiao-Bin Liu[1], Shen Nan[5], Manish Pareek[1], Neha Sahu[1], Benedek Szathmári[1], Torda Varga[1],
9 Hongli Wu[1], Xiao Yang[5], Zsolt Merényi[1]
10

1 Synthetic and Systems Biology Unit, Biological Research Center, Szeged, 6726, Hungary
2 Microbiology, Department of Biology, Faculty of Science, Utrecht University, Padualaan 8,
3584 CH, Utrecht, The Netherlands
3 Eidgenössische Technische Hochschule Zürich, Department of Biology, Institute of
Microbiology, Zürich, Switzerland
4 Project Group Genetics and Genomics of Fungi, Chair Evolution of Plants and Fungi,
Ruhr-University Bochum, 44780, Bochum, North Rhine-Westphalia, Germany
5 Institute of Applied Mycology, Huazhong Agricultural University, 430070 Hubei Province,
PR China
20
21 ** To whom correspondence should be addressed: lnagy@fungenomelab.com*
22

Abstract

Fruiting bodies of mushroom-forming fungi (Agaricomycetes) are among the most complex
structures produced by fungi. Unlike vegetative hyphae, fruiting bodies grow determinately
and follow a genetically encoded developmental program that orchestrates tissue
differentiation, growth and sexual sporulation. In spite of more than a century of research,
our understanding of the molecular details of fruiting body morphogenesis is limited and a
general synthesis on the genetics of this complex process is lacking. In this paper, we aim to
comprehensively identify conserved genes related to fruiting body morphogenesis and distill
novel functional hypotheses for functionally poorly characterized genes. As a result of this
analysis, we report 921 conserved developmentally expressed gene families, only a few
dozens of which have previously been reported in fruiting body development. Based on
literature data, conserved expression patterns and functional annotations, we provide
informed hypotheses on the potential role of these gene families in fruiting body
development, yielding the most complete description of molecular processes in fruiting body
morphogenesis to date. We discuss genes related to the initiation of fruiting, differentiation,
growth, cell surface and cell wall, defense, transcriptional regulation as well as signal
transduction. Based on these data we derive a general model of fruiting body development,
which includes an early, proliferative phase that is mostly concerned with laying out the
mushroom body plan (via cell division and differentiation), and a second phase of growth via
cell expansion as well as meiotic events and sporulation. Altogether, our discussions cover
1480 genes of *Coprinopsis cinerea*, and their orthologs in *Agaricus bisporus, Cyclocybe*
aegerita, Armillaria ostoyae, Auriculariopsis ampla, Laccaria bicolor, Lentinula edodes,
Lentinus tigrinus, Mycena kentingensis, Phanerochaete chrysosporium, Pleurotus ostreatus,

and *Schizophyllum commune*, providing functional hypotheses for ~10% of genes in the
genomes of these species. Although experimental evidence for the role of these genes will
need to be established in the future, our data provide a roadmap for guiding functional
analyses of fruiting related genes in the Agaricomycetes. We anticipate that the gene
compendium presented here, combined with developments in functional genomics
approaches will contribute to uncovering the genetic bases of one of the most spectacular
multicellular developmental processes in fungi.
53
54 Key words: functional annotation; comparative genomics; cell wall remodeling; development;
55 fruiting body morphogenesis; mushroom; transcriptome
56
57

143   1. Introduction

Fungi are one of the five main extant lineages in which complex macroscopic structures
emerged over the course of evolution. The best known macroscopic structures of fungi are
sexual fruiting bodies, which enclose spore-producing cells into a 3-dimensional, protective
environment and facilitate spore dispersal(Nagy et al. 2018). Sexual fruiting bodies are
complex multicellular structures based on the definition of Knoll(Knoll 2011) and show
evidence for repeated evolution across the fungal tree of life(Stajich et al. 2009; Merényi et
al. 2020a). They are found in at least 8 lineages, with best known representatives belonging
to the Pezizomycotina (e.g. morels, truffles), Agaricomycotina (e.g. agarics, boletes),
however, spectacular examples evolved in several other lineages, such as *Endogone* spp. in
the Mucoromycota(Chang et al. 2019b) or the enigmatic *Neolecta* genus in the
Taphrinomycotina(Nagy 2017; Nguyen et al. 2017). Fruiting bodies reached highest
complexity in the Agaricomycetes, which include developmentally integrated morphologies
that follow a spatially and temporally tightly regulated developmental program(Hibbett 2004;
Virágh et al. 2021). In terms of complexity level, certain Agaricomycete fruiting bodies
approach the complexity of simple animals and plants(Taylor and Ellison 2010; Nagy et al.
2018).

Fruiting body morphogenesis has been a subject of intense research, however,
despite tremendous efforts in this field, morphogenesis of Basidiomycota fruiting bodies is
quite poorly known. In comparison, fruiting body development and its underlying genetics are
considerably better explored in the Ascomycota (see e.g.(Pöggeler et al. 2006a)). This may
be due to a mixture of factors, including the complexity of Basidiomycota developmental
programs, the lack of easy-to-use model systems or the relative difficulty of hypothesis
testing via genetic manipulation (e.g. due to low frequency of homologous recombination(De
Jong et al. 2010)). Research on Basidiomycete fruiting body morphogenesis has recently
been relying largely on comparative -omics techniques, which provided insights into the
gene repertoires and the regulation of gene expression during fruiting body development.
This complements the rich body of literature on functionally characterized genes in
Coprinopsis cinerea (summarized in Kües(Kües 2000)), *Schizophyllum commune* and other
species. There are several recent milestones in fruiting body research such as the
publication of key genome sequences(Ohm et al. 2010; Stajich et al. 2010; Morin et al. 2012)
or comparative transcriptomic datasets (e.g. (Morin et al. 2012; Plaza et al. 2014a; Kües and
Navarro-González 2015; Almási et al. 2019; Krizsán et al. 2019; Merényi et al. 2021)). An
emerging driving force of research on fruiting body morphogenesis is the mushroom
industry, which produces sexual fruiting bodies for food or as sources of bioactive
compounds(Royse et al. 2017; Ma et al. 2018). Fruiting body development has been a
subject of several excellent reviews, surprisingly, however, the last gene-centric review of
fruiting body development in mushroom-forming fungi was published in 2000(Whiteford and
Thurston 2000). This paper aims to complement recent reviews on the morphogenesis of
vegetative hyphae(Riquelme et al. 2018), and on the genome of *Neurospora*
crassa(Borkovich et al. 2004) with knowledge on fruiting body morphogenesis in the
Agaricomycetes.
The aim of this paper is to systematically catalog genes and genetic processes
related to fruiting body morphogenesis in the Agaricomycetes, based on both literature
reviews and a meta-analysis of published developmental transcriptomes. We provide a
handbook-like inventory of gene families and their putative functions based on expression
data and functional annotations borrowed from well-researched model systems. Because a
negligible proportion of morphogenesis genes has so far been functionally characterized in
the Agaricomycetes, it is currently impossible to build a comprehensive picture based on
mechanistic studies alone, as has recently been done for hypha morphogenesis(Riquelme et
al. 2018). Instead, we derive functional hypotheses for genes based on comparative
expression data across multiple species. Our starting point is that genes which are
developmentally regulated in the majority of species for which transcriptome data are
available, should belong to the core pathways of fruiting body morphogenesis, whereas
genes that show species-specific expression patterns should be responsible for sculpting
species-specific morphologies or represent transcriptional or technical noise. Finally, we
incorporate novel and previous observations into a new synthesis on processes active
during fruiting body morphogenesis.
201

1.1. What are fruiting bodies and how do they develop

Fruiting bodies, also called basidiomes, basidiocarps or simply 'mushrooms' in lay language,
may be some of the best known structures of fungi. They are sexual reproductive organs
that facilitate spore dispersal. While in simple Basidiomycota spores are born on naked

basidia, mushroom-forming fungi evolved complex 3-dimensional structures that offer
tremendous advantages in spore dispersal efficiency.
Developmental events that take place in fruiting bodies have been investigated
previously in several species, of which *Coprinopsis/Coprinellus* spp. and *S. commune* are
the two most widely used groups of models. The development of fruiting bodies in these
species has been described in previous reviews(Kües 2000; Kües and Liu 2000; Palmer and
Horton 2006; Kües and Navarro-González 2015), therefore, we here only provide a short
introduction, and describe the process in more detail (combining new results) in Chapter 5.
Briefly, fruiting body development is initiated on sexually competent mycelia as the
reprogramming of hyphal branching patterns, which results in the interlacing of hyphae to
form a 3-dimensional fruiting body initial(Kües and Liu 2000; Kües and Navarro-González
2015). Rather than relying on zygotic (e.g. maternal) mRNA cargo as animals and plants do,
fungal hyphae undergo transcriptional reprogramming during fruiting body initiation. This
reprogramming is influenced and triggered by a mixture of - often species-specific - factors,
such as starvation, light or temperature changes (reviewed by(Sakamoto 2018)). Once
appropriate internal and external signals converge on the initiation of fruiting, morphogenesis
starts with an undifferentiated initial (hyphal knot, aggregate), which undergoes tissue
differentiation to form primordia. In subsequent late primordium and young fruiting body
stages rapid growth (stipe elongation and cap expansion) and meiosis happen in a
coordinated manner, which ultimately results in species-specific mature morphologies and
culminates in spore formation and release.
Development of the fruiting bodies is a tightly regulated process that has a high
degree of autonomy. That is, once started, the developmental program is not interrupted by
even large perturbations, such as injuries, or removal of the fruiting body from the supporting
colony at or after certain stages(Moore 2013). In fruiting bodies not only morphogenetic
events, but also several internal processes (e.g. nutritional and sexual) are developmentally
regulated. For example, in *C. cinerea*, stipe elongation starts immediately after meiosis takes
place, after which spore production, maturation and cap expansion follow in a tightly
choreographed chronology(Kües 2000; Moore 2013). For detailed description of the process
and molecular summaries of current knowledge on fruiting body morphogenesis the reader
is referred to recent reviews(Kües 2000; Kües and Navarro-González 2015; Virágh et al.
2021) and the following detailed chapters of this paper.

1.2. The evolution of Agaricomycete fruiting bodies: from simple to complex
morphogenesis

How fruiting body complexity evolved has interested mycologists for decades and has
received considerable attention recently. Therefore we provide only a brief introduction here
and refer the reader to recent reviews on the topic (e.g.(Kües and Navarro-González 2015;
Nagy et al. 2018; Virágh et al. 2021)). Phylogenetic studies have made tremendous progress
in uncovering broad patterns of morphological transformation and trends in fruiting body
evolution (e.g. (Hibbett 2004; Varga et al. 2019; Sánchez-García et al. 2020)). Fruiting body
morphologies show a great diversity in the Basidiomycota, from naked lawns of basidia, to
resupinate, coralloid or highly complex, or the so called pileate-stipitate fruiting bodies of
well-known agarics (e.g. *C. cinerea*) (Fig. 1).
Agaricomycete fruiting bodies might have emerged in or around the last common
ancestor of Dacrymycetes and Agaricomycetes over 400 million years ago(Varga et al.
2019; Prasanna et al. 2020), although there is some uncertainty around ancestral states.

This is partly due to the presence of jelly-like telia reminiscent of fruiting bodies in the
Pucciniomycotina, uncertainties in the branching order of Basidiomycota subphyla and
reductive evolution in early-diverging clades (Bartheletiomycetes, Wallemiomycetes) of the
Agaricomycotina subphylum (reviewed in (Prasanna et al. 2020)). Nevertheless, all
Agaricomycetes share a single, fruiting body-forming ancestor, which suggests there should
be a core, deeply conserved genetic toolkit of fruiting body development. Phylogenetic
comparative studies have provided sound evidence that this ancestor had
corticioid/resupinate/jelly like fruiting bodies (e.g. *Phanerochaete chrysosporium*) and that
more complex forms derived from such morphologies repeatedly(Hibbett 2004; Varga et al.
2019). Within the Agaricomycetes, tissues that protect and enclose basidia became thicker
and more complex, eventually leading to the mushroom fruiting body structures we know
today. From the perspective of this paper, it is important to mention that phylogenetic studies
have also established that fruiting body evolution is highly convergent, that is, many of the
characteristic traits (e.g. cap, gills) evolved independently several times. Therefore, an
evolutionary framework is important for understanding how fruiting bodies evolved and for
establishing homologies among tissues. Gills, caps and stipes, for example, although
superficially similar, are not homologous across Polyporales and Agaricales. This implies
that such, convergent forms (e.g. gills of *Lentinus tigrinus* in the Polyporales and that of *C.*
cinerea in the Agaricales) are not homologous across lineages and their development should
not be expected to follow the same genetic principles. On the other hand, basidia, the sexual
processes that happen therein and the development of 3-dimensional fruiting body tissue,
are homologous across all mushroom-forming fungi (i.e. they have a single origin in the
Basidiomycota), providing an reliable framework for analysis.
These examples illustrate how the diversity of fruiting bodies can complicate the
identification of core components of their morphogenesis. Ideally, we are interested in
developmental processes and underlying genetic components that uniformly characterize
fruiting bodies of all mushroom-forming fungi, i.e. those that can be traced back to the first
fruiting body-forming Agaricomycete ancestor.
Therefore, building 'minimal models', which encompass the minimum set of genes
necessary to build a fruiting body are highly relevant. Such minimal models could be, among
others, species that produce resupinate fruiting bodies (e.g. *Ph. chrysosporium*)(Krizsán et
al. 2019). Recently, *Cryptococcus neoformans* (Tremellomycetes, Basidiomycota) proved to
be a useful model organism, because it produces simple lawns of basidia, sometimes
referred to as fruiting bodies, in which basidium differentiation, meiosis and spore formation
take place(Liu et al. 2018). Therefore, *C. neoformans* 'fruiting bodies' encompass all
processes needed for Basidiomycota sexual reproduction but lack features essential to the
definition of fruiting bodies such as three-dimensional organization, tissue differentiation and
others. Merenyi et al recently sorted developmental genes of *Pleurotus ostreatus* and seven
other complex species into groups that were developmentally regulated both in *C.*
neoformans and mushroom-forming fungi and those in which developmental expression was
observed only in mushroom-forming fungi(Merényi et al. 2021). 'Shared' orthogroups
comprised most mitosis/meiosis related genes, but also some previously thought to be
specific for complex morphogenesis (e.g. fasciclins). This indicates that sexual
morphogenesis, even at the very simple forms that exist in *C. neoformans*, involves a
morphogenetic program. Comparing genes that are developmentally regulated in fruiting
bodies to those regulated in *C. neoformans,* thus, can help constructing more precise
genetic models of complex fruiting body morphogenesis.

1.3. Cellular and regulatory processes involved in fruiting body development

Fruiting body development is an incredibly complex process and this complexity is reflected
in the underlying cellular and genetic processes. Several cellular processes are emerging as
important regulatory mechanisms underlying fruiting body morphogenesis, whereas some
others that proved developmentally relevant in animals and plants seem to be of limited
importance in fungi (e.g. gene body methylation(A et al. 2018; Wen et al. 2019)). This
chapter summarizes the most important - mostly transcription related - processes described
in fruiting bodies recently. Splicing and RNA interference are discussed below under
chapters 4.5.3 and 4.6, respectively.
Probably the most influential and most commonly assayed mechanisms of fruiting
body development are transcriptional reprogramming and gene expression changes. Gene
expression changes naturally associate with developmental processes and are informative
regarding the cellular processes that take place during fruiting. *In silico* functional analyses
of differentially or developmentally expressed genes have been a primary tool for
mycologists to understand fruiting body functions. Developmentally dynamic gene
expression patterns form the primary data source for this paper and the main part of the
paper (Chapter 4) is devoted to discussing genes with conserved expression patterns. A
significant unresolved question is how gene expression changes are regulated in various
stages of fruiting body development. It is not known currently if transcriptional
reprogramming events (e.g. during fruiting body initiation) are mediated by broad regulation
of chromatin accessibility, by regulating the expression and activity of regulatory genes (e.g.
transcription factors), or a mixture of these. A potentially significant regulatory mechanism is
post-translational modification of transcription factors or molecular switches in intracellular
signaling pathways, as demonstrated for the *Hom2* or the *Ras1* gene of *S. commune*(Knabe
et al. 2013; Pelkmans et al. 2017b). Another exciting avenue of future research is the
regulatory potential of selective post-translational protein modification by the
ubiquitin/proteasome system. Although direct evidence for its role in fruiting body
development has not yet been published, indirect evidence was provided by the remarkable
expansion of genes encoding F-box, BTB/POZ and RING-type zinc finger proteins in the
genomes of mushroom-forming fungi(Krizsán et al. 2019). These gene families provide the
substrate-specificity of ubiquitin ligases and have been shown, in other lineages, to be
involved in transcriptional regulation(Kipreos and Pagano 2000).
Natural antisense transcripts (NATs) are RNA transcripts that show complementarity to
transcripts of protein coding genes. NATs occur ubiquitously across eukaryotes and have
been detected in mushroom-forming fungi too(Ohm et al. 2010; Muraguchi et al. 2015;
Merényi et al. 2021). Because NATs do not encode proteins, they often evolve very fast and
are hard to link to specific functions. Nevertheless, their expression is developmentally
dynamic in all species examined to date, suggesting that they may be involved in regulating
development. Muraguchi et al reported that an antisense transcript of the stipe elongation
related AspE gene is transcribed just before stipe elongation(Muraguchi et al. 2015).
However, it should also be kept in mind that many NATs may result from random
transcription initiation and may therefore be transcriptional noise in fruiting body
transcriptomes(Merényi et al. 2021).
Allele-specific expression (ASE) refers to the imbalance of gene expression from two
divergent alleles in diploid or dikaryotic organisms. Fungal dikaryons almost always harbor
two allelically different monokaryons (heterokaryons), which can lead to intermediate, A- or
B-dominated expression. ASE has so far been detected in fruiting body transcriptomes of *A.*

bisporus(Gehrmann et al. 2018) and *P. ostreatus*(Merényi et al. 2021), but may be
widespread in the Agaricomycetes. In *A. bisporus*, Gehrmann et al suggested that ASE may
facilitate the division of labor between nuclei in a dikaryon(Gehrmann et al. 2018). Another
study, based on *P. ostreatus*, suggested that, at macroevolutionary timescales, ASE may be
a neutrally arising phenomenon that can nevertheless generate adaptive gene expression
variation(Merényi et al. 2021). It was found to be mostly characteristic of evolutionarily young
genes, consistent with neutral evolution. Allele-specific expression has agriculturally broad
consequences in hybrid genetics of plants(Bartoš et al. 2019) and could become a similarly
central question in the mushroom industry.
RNA editing has recently been detected in fruiting body transcriptomes of several
Ascomycota species(H et al. 2016; Liu et al. 2017; Teichert et al. 2017) and subsequently
reported in Basidiomycota also(Zhu et al. 2014; Wu et al. 2019). Like alternative splicing,
RNA editing provides a (post-)transcriptional mechanism for diversifying transcripts encoded
by a single gene, by enzymatically modifying certain bases of the primary RNA transcript. A
critical analysis recently found no evidence for RNA editing in fruiting body transcriptomes of
P. ostreatus(Merényi et al. 2021). This study raised the possibility that, because allele-
specific expression and RNA editing can generate similar signatures in RNA-Seq data, ASE
may have been mistaken for RNA editing in some prior Basidiomycota analyses and that
more research is needed to prove a role of RNA editing in mushroom development.
Several other exciting phenomena may be involved in the morphogenesis of
Basidiomycota fruiting bodies, including regulatory processes that may yet to be discovered.
For example, epigenetic regulation represents a potentially widespread but largely
underexplored regulatory mechanism for fruiting body morphogenesis(Vonk and Ohm 2021).
Similarly, upstream open reading frames (uORFs), which are open reading frames located in
the 5' untranslated region of an mRNA, can regulate (e.g. via encoded peptides) or interfere
with the transcription of the gene(HM et al. 2009). uORFs are known to be important in the
asexual development of Ascomycota(Han and Adams 2001) [and have recently been
detected in developmental genes of *C. cinerea* (Hegedus et al in prep)].

1.4. Genomic and transcriptomic resources

Genomic resources are also available for most model species of fruiting body development
(e.g. *C. cinerea* (Schaeff.) Redhead, Vilgalys & Moncalvo*, S. commune* Fr.*, Cyclocybe*
aegerita (V. Brig.) Vizzini)(Ohm et al. 2010; Stajich et al. 2010; Muraguchi et al. 2015; Gupta
et al. 2018; Almási et al. 2019) and those used by the mushroom industry, including
Hericium erinaceus (Bull.) Pers.(Gong et al. 2020; W et al. 2020)*, Ganoderma spp.*(Chen et
al. 2012)*, Flammulina spp.*(Park et al. 2014, 2019; Chen et al. 2020)*, L. edodes* (Berk.)
Pegler(Chen et al. 2016; Shim et al. 2016; Sakamoto et al. 2017a)*, Pleurotus spp*(A et al.
2018; Wang et al. 2018a; Dai et al. 2019; Lee et al. 2021)*, Agaricus spp*(Morin et al. 2012;
O'Connor et al. 2019; Sonnenberg et al. 2020)*, Tricholoma matsutake* (S. Ito & S. Imai)
Singer(Min et al. 2020)*, Sparassis crispa* (Wulfen) Fr.(Kiyama et al. 2018)*, Hypsizygus*
marmoreus (Peck) H.E. Bigelow(Min et al. 2018), among others. More broadly, in the
Agaricomycetes, draft genome sequences have been proliferating at a steady pace, due to
interest in biodiversity, lignocellulose decomposition(Martinez et al. 2004, 2009a; Floudas et
al. 2012; Riley et al. 2014) and mycorrhiza-formation(Martin et al. 2008; Kohler et al. 2015;
Miyauchi et al. 2020). This trend has virtually eliminated genome data being the bottleneck in
biological discovery in fungi. This is further supported by the spread of third generation
sequencing techniques, long-read sequencing data, which allow more and more complete

(telomere-to-telomere) assemblies and a range of analyses that draft genomes did not
permit.
The continuous development of technologies has also influenced the identification of
genes related to fruiting body morphogenesis. While early studies isolated fruiting body-
specific proteins/genes using mutant analyses, hybridization techniques (among others),
these were later substituted by RT-PCR and microarray studies, then by high throughput
sequencing analyses, such as 5'-serial analysis of gene expression (SAGE(WW et al. 2008;
Cheng et al. 2013a)) and most recently RNA-Seq. As of today, RNA-Seq datasets are
available for a wide range of species and conditions. These include *C. cinerea*(Plaza et al.
2014a; Muraguchi et al. 2015; Krizsán et al. 2019; Xie et al. 2020)*, Flammulina spp*(Yan et
al. 2019; Liu et al. 2020b)*, L. edodes*(Park et al. 2017; Sakamoto et al. 2017b; Song et al.
2018b; Y et al. 2018; Yoo et al. 2019; Kim et al. 2020)*, Agaricus spp*(Gehrmann et al. 2018;
Lu et al. 2020; O'Connor et al. 2021)*, Pleurotus ostreatus* (Jacq.) P. Kumm.(Wen et al. 2019;
Merényi et al. 2021)*, H. marmoreus*(Zhang et al. 2015b)*, C. aegerita*(Orban et al. 2021)*,
Phanerochaete chrysosporium* Burds.(Krizsán et al. 2019)*, Lentinus tigrinus* (Bull.)
Fr.(Krizsán et al. 2019)*, Mycena kentingensis* Y.S. Shih, Chi Y. Chen, W.W. Lin & H.W.
Kao(Ke et al. 2020a)*, Rickenella mellea* (Singer & Clémençon) Lamoure(Krizsán et al.
2019)*, Armillaria ostoyae (*Romagn.) Herink(Sipos et al. 2017a)*, Pisolithus microcarpus*
(Cooke & Massee) G. Cunn (Pereira et al. 2017)*, Laccaria bicolor* (Maire) P.D. Orton
(Ruytinx et al in prep)*, S. commune*(Ohm et al. 2010) and *Auriculariopsis ampla* (Lév.)
Maire(Almási et al. 2019)*. Many of these datasets build the foundation of this study, by
providing windows into developmentally expressed genes in different fungal species.
A general observation we made during the analyses we present hereafter is that the
resolution of the transcriptomic data is key to identifying expression dynamics and patterns.
Both tissue-wise and temporal resolution of the data allowed the discovery of more
developmentally relevant expression patterns. We found several cases where tissue-
resolved transcriptomes provided clear signal for conserved, tissue-specific expression of a
gene, whereas in non-resolved transcriptomes the same gene showed a more or less flat
expression curve. This is an inherent nature of bulk RNA-Seq data: spatial expression
patterns average out in samples that contain large populations of multiple cell types.
Beyond gene expression, a broad array of other types of information could be
leveraged for understanding fruiting body development. Notable recent examples include
assays of the overrepresentation of certain gene families in the genomes of focal
species(Krizsán et al. 2019), a population genomics study combined with selection analyses
and gene expression profiling in *L. edodes*(Zhang et al. 2021c) to identify genes involved in
fruiting body development(Zhang et al. 2021c), or ChIP-Seq analyses for assaying promoter
occupancy and thus transcriptional activity(Vonk and Ohm 2021).

429 2. Description of the approach used in this paper

In this study, we address the puzzle of how a fruiting body develops and what novel insight
can be gleaned from systematic comparisons of developmental transcriptomes. We aim at
providing a comprehensive listing of putatively development-related gene families in the
Agaricomycetes. We provide, for each discussed functional group or gene family,
information on their expression, orthology to well-established model organisms
(*Saccharomyces cerevisiae, Schizosaccharomyces pombe, Aspergillus nidulans*,
Neurospora crassa) and to key mushroom-forming fungi as tables. We also annotate several

conserved pathways in *C. cinerea*, based on strict 1-to-1 orthology to known pathway
members in well-researched model systems (see Methods for details).
Each chapter is structured so that it starts with a literature review of the given gene
family followed by a discussion of novel findings (if any) based on our meta-analysis of
transcriptomes. Our main units of investigation are **conserved developmentally expressed**
(CDE) orthogroups, which are groups of single-copy genes in which the majority (>65%[1]) of
the genes were developmentally expressed (see Methods and Merenyi et al 2021(Merényi et
al. 2021)). For simplicity CDE orthogroups are considered to be groups of genes sharing
similar functions (see limitations below) in fruiting body development.
We performed a meta-analysis of 12 developmental transcriptomes (*Agaricus*
bisporus(Gehrmann et al. 2018)*, C. aegerita*(Orban et al. 2021)*, A. ostoyae*(Sipos et al.
2017a)*, A. ampla*(Almási et al. 2019)*, C. cinerea*(Krizsán et al. 2019)*, L. bicolor* (Ruytinx et al
in prep)*, L. edodes*(Zhang et al. 2021d)*, L. tigrinus*(Krizsán et al. 2019)*, M. kentingensis*(Ke
et al. 2020a)*, Ph. chrysosporium*(Krizsán et al. 2019)*, P. ostreatus*(Krizsán et al. 2019)*, S.*
commune(Almási et al. 2019)) (Fig. 1). These species belong to two orders (Agaricales,
Polyporales), which span an estimated 200 million years of evolution(Varga et al. 2019), and
represent diverse ecologies and morphological adaptations, such as simple resupinate (*Ph.*
chrysosporium), coralloid (*Pterula gracilis*), pileate-stipitate (e.g. *C. cinerea*) and cyphelloid
(*A. ampla, S. commune*) forms. Each of these species and their ecological adaptations show
several unique aspects (e.g. fruiting body ecotypes) which could be related to gene
expression data. However, we are here interested in only the shared aspects of fruiting body
development, and therefore use the diversity of the compared species in focusing our
attention to widely conserved fruiting body genes which should help define the core building
blocks required for fruiting body morphogenesis.
The comparative approach we follow is paramount to separate 'wheat from chaff', i.e.
developmentally regulated genes that are relevant for the development of fruiting bodies in
general from genes with taxonomically restricted or species-specific developmental roles.
This builds on the assumption that a gene involved in core fruiting body functions should be
conserved across Agaricomycetes and should show similar expression dynamics in all
species. Therefore, in our approach we focus on conservation of both sequence and
expression. A comparative approach is also desirable because it is now well-known that
gene expression is associated with 'biological noise', which may influence the detection of
differentially expressed or developmentally regulated genes(Merényi et al. 2021) in any
single species, whereas across-species comparisons effectively cancel the effects of gene
expression noise.
This study, like all others, has limitations. First, it should be noted that evidence for
gene function remains, in an overwhelming majority of cases, circumstantial, inferred from
expression patterns or extrapolated from the functions of orthologous genes in model fungi.
At the moment we rely on these types of evidence due to the scarcity of developmental
genetic and in-depth functional studies in most Agaricomycetes. Despite the inferential
nature of the study, many orthologs display very conserved expression patterns across
Agaricomycetes species (e.g. aquaporins in stipe tissues, see below), allowing us to make
confident predictions on the function of such genes. Second, a perfect comparative
transcriptomic study would compare corresponding developmental stages across species. In
the Agaricomycetes establishing homology relationships among individual species is

---

[1] We deviaed from this rule where justified.

complicated (see above). Furthermore, the developmental transcriptomes we used differ in
the sampled time points and resolution (e.g. tissue-wise or bulk) and do not follow a unified
nomenclature for developmental stages (e.g. aggregate in *S. commune* vs. hyphal knot in *C.*
cinerea). Therefore, we here focused broadly on comparing expression dynamics and the
shape of the expression curve, rather than attempting to find 1-to-1 correspondence
between stages of different species. Finally, in the present study, we focus on genes with
developmentally dynamic expression. However, not all morphogenetically relevant genes
show expression dynamics; constantly expressed genes can often be important for
development, for example, if regulated post-transcriptionally (e.g. alternative splicing
isoforms(Gehrmann et al. 2016; Krizsán et al. 2019)) or post-translationally (e.g.
phosphorylation, ubiquitylation etc.(Knabe et al. 2013; Pelkmans et al. 2017a)).
493

494   3. Methods

Bioinformatic analyses of RNA-Seq data
The raw RNA-Seq data of 12 previously published Basidiomycota species were reanalysed
following Merenyi et al(Merényi et al. 2021). The following reference genomes were used: *C.*
cinerea (AmutBmut pab1-1 v1.0(Muraguchi et al. 2015))*, A. ostoyae* (C18/9(Sipos et al.
2017b))*, A. ampla* (NL-1724 v1.0(Almási et al. 2019)), *S. commune* (H4-8a and H4-8b(Ohm
et al. 2010))*, L. bicolor* (v2.0(Martin et al. 2008))*, L. edodes* (Le(Bin) 0899 ss11 v1.0(Zhang
et al. 2021c))*, L. tigrinus* (RLP-9953-sp(Wu et al. 2018)), *Ph. chrysosporium* (RP-78
v2.2(Ohm et al. 2014)), *P. ostreatus* (PC15 v2.0(Riley et al. 2014))*, M. kentingensis*(Ke et al.
2020a)*, A. bisporus* (var. bisporus H97(Morin et al. 2012))*, C. aegerita* (AAE-3(Gupta et al.
2018))*.* To remove adaptors, ambiguous nucleotides and any low quality read ends, reads
were trimmed using bbduk.sh and overlapping read pairs were merged with bbmerge.sh
(part of BBMap/BBTools; http://sourceforge.net/projects/bbmap/) with the following
parameters: qtrim=rl trimq=25 minlen=40. A two-pass STAR alignment(Veeneman et al.
2016) was performed against reference genomes with the same parameters as in our
previous study(Krizsán et al. 2019) except that the maximal intron length was reduced to
3000 nt. Reads were counted and summarized for each exon with FeatureCounts(Liao et al.
2014) taking into account the mode of sequencing (paired-end, single-end). Since the
Cyclocybe transcriptome was obtained with the Quantseq method, reads were counted only
in the -100 - +400 region of the 3' ends of the genes. Read count data were normalized
using EdgeR(Robinson et al. 2010). Expression levels were calculated as fragments per
kilobase of transcript per million mapped reads (FPKM).
Developmentally regulated genes were identified as before(Krizsán et al. 2019;
Merényi et al. 2021), based on a stricter four-fold or a more permissive two-fold change
cutoff between adjacent developmental stages or tissue types. Genes showing a 4-fold
upregulation in the first primordium stage relative to vegetative mycelium were termed 'FB-
init', and considered regulated during fruiting body initiation. It should be noted that the data
we analyze is noisy, based on different chemistries, different labs and protocols, etc., which
introduces a certain amount of data loss in our analyses. Nevertheless, the transition to
fruiting body development is such a dramatic change that we expect similar gene expression
changes to be discoverable across species, despite the heterogeneity of the data. This is
supported by previous comparative transcriptomic studies(Plaza et al. 2014b; Krizsán et al.

2019; Merényi et al. 2020b, 2021) as well as the biologically meaningful conclusions
obtained here.
528
Orthology based on reciprocal best hits
Proteins of each species were searched against the proteomes of other species using the
reciprocal best hit (RBH) module of MMSeqs2(Steinegger and Söding 2017). To remove
spurious RBHs, a bidirectional 50% coverage and at least 1e-6 e-value were required.
Proteins were clustered into single copy clusters using a connected component clustering
algorithm in the igraph package([CSL STYLE ERROR: reference with no printed form.]),
based on reciprocal best hits. This step clustered 74.3% of the proteins in the examined
species. The other 25.7% of proteins form clusters with duplication in any or more species.
For these 1:1 orthogroups were circumscribed using the 'ortholog coding' algorithm of the
COMPARE pipeline(Nagy et al. 2014). In the case of terminal duplications, orthogroup
membership was decided based on connectivity; the protein which showed more hits to
other members of the orthogroup was selected while the other(s) moved to separate
orthogroup(s). Complete orthology relationships of genes across all 12 species are given in
this paper are presented in Supplementary Table 1.
This approach is suitable for identifying moderately to very conserved gene groups,
whereas for highly volatile genes (i.e those that show frequent duplication/loss), a more
liberal approach building on orthologous groups (like(Krizsán et al. 2019)) is more
appropriate. For fast-evolving genes, approaches looking at sequence conservation might
not work. Therefore, in the case of certain gene families, such as defense genes or cell wall
remodeling CAZymes, we relied on primary gene family classification instead of reciprocal
best hit based orthogroups. In the case of CAZymes, we followed the cazy.org
classification(Levasseur et al. 2013).
551
Identification of CDE orthogroups
The central unit of discussion in this paper is referred to as 'conserved developmentally
expressed (CDE) orthogroup', which refers to groups of single-copy genes (one gene per
species) in which the majority of genes is developmentally regulated either at fold change >2
or >4. As a principal rule, we considered an orthogroup to be a CDE orthogroup if >65% of
the genes in it were developmentally regulated, and at least 8 species were represented.
Deviations from this rule were allowed in unique cases (e.g. taxonomically restricted but
important gene family) and are transparently presented in tables and supplementary tables
under each of the following chapters.

4. Results: developmentally expressed gene classes in the
Agaricomycetes

In this paper we synthesized literature reviews with a meta-analysis of developmental
transcriptomes to identify conserved gene families involved in fruiting body development. For
the latter we reanalysed data for 12 species and identified developmentally regulated genes,
i.e. those that showed >2 or >4-fold changes between successive developmental stages or
adjacent tissue types (Supplementary Table 2). Statistics on developmentally expressed
genes are provided in Supplementary Table 2/b. Numbers of developmentally regulated
genes were similar to those reported by previous studies,(Almási et al. 2019; Krizsán et al.

2019; Merényi et al. 2020b, 2021) therefore, we here do not discuss these in detail, rather
focus on those that are conserved across species.
We organized developmentally regulated genes into strict 1-to-1 orthogroups
following the logic of Merenyi et al(Merényi et al. 2021). Orthogroups in which >65% of the
genes were developmentally regulated were designated as conserved developmentally
expressed (CDE) orthogroups at fold change values of >2 or >4. We identified 921 CDE
orthogroups across the twelve Agaricomycetes species; these form the basis of the
discussion presented in the rest of this paper. Statistics on the number of species which had
developmentally regulated genes in these orthogroups are shown on Fig. 2/A. The
transcriptome data of *A. bisporus* was an outlier in several aspects. We currently do not
know if this is due to biological differences of this species or technical reasons, nevertheless
this transcriptome was assigned lower weight and discussed rarely in the following chapters.
By relying on conserved developmentally expressed orthogroups, we here take shared
expression dynamics as a proxy for conservation of function. It should be mentioned that this
approach may be, in several cases, only a crude approximation of real conservation of
function, but facilitates comparative discussion of gene families and their speculated
functions in fruiting body morphogenesis.
We functionally characterized CDE orthogroups and arranged them into broader
groups based on function inferred by orthology to genes in well-known model systems (*S.*
cerevisiae, Schizosaccharomyces pombe, A. nidulans, N. crassa) and/or their Gene
ontology and InterPro annotations. We defined 17 broader functional groups, cell
division/proliferation, defense, transcriptional regulators, signal transduction, cell wall
biosynthesis and remodeling, cell surface proteins, secondary metabolism, cytoskeleton,
basic metabolism and acetyl-CoA production, lipid metabolism, storage carbohydrate
metabolism, ubiquitin/proteasome system, transporters, functionally poorly known genes,
unannotated genes, stress response related genes and ferric reductases (Fig. 2/B). The
main categories were further subdivided into finer groups to facilitate discussions of putative
functions. We categorized developmental genes manually to the best of our knowledge
based on inferred gene function, though the subdivision is probably not impeccable. For
example, overlaps exist between chromatin remodeling and DNA replication related
regulatory factors or between the latter and cytoskeleton related genes (e.g. kinesins
involved in moving chromatids during mitosis).
The two largest categories were functionally poorly characterized and unannotated
genes, these contained 182 and 176 CDE orthogroups, respectively (Fig. 2/B). The first,
'functionally poorly characterized genes' contains orthogroups which we could not
confidently link to any fruiting body functions. It is possible that this category could be
subdivided and with further manual annotation more orthogroups could be explained in the
context of fruiting body development. On the other hand, unannotated genes are those that
contain no existing Pfam or InterPro domain signatures and lack all kinds of functional
annotation. The largest functional group for which a function could be determined was
related to cell proliferation (DNA replication, DNA repair, mitosis, meiosis) followed by cell
wall biosynthesis and remodeling. Transcriptional regulators (including transcription factors,
chromatin-related and RNA-binding regulatory proteins), transporters and Acetyl-CoA
production related genes were also represented by a considerable number of genes. Certain
functional categories, such as 'defense' and 'secondary metabolism' contained only a small
number of CDE orthogroups. In these cases, these low numbers reflect the lack of
conservation of gene sequences, rather than the lack of a role of these gene families in
fruiting bodies.

Our discussion is guided here by conservation of developmentally regulated families.
However, there are gene families that do not form conserved orthogroups, but are
nevertheless very important for fruiting body development. These can be either multigene
families that undergo frequent duplication and thus orthology relationships are intricate (e.g.
hydrophobins), or fast-evolving families in which divergence quickly erases orthology
relationships (e.g. F-Box proteins, defense-related proteins). Therefore, in addition to gene
families/pathways identified through CDE orthogroups, we discuss broader, fast-evolving
gene families based on annotations at the gene family level only (i.e. without orthogroups)
and performed manual annotations to identify broader containing gene families or cellular
pathways. Accordingly, in addition to the 921 CDE orthogroups, we discuss further 558 gene
groups which belong to broad functionalities involved in fruiting body development. For some
processes we were able to reconstruct (nearly) complete pathways, such as in the case of
mitosis/meiosis related genes or fatty acid biosynthesis, whereas in other cases we only
document the widespread developmental regulation and expression profile conservation of a
gene family, but without information on the containing pathway of broader cellular function.
For some gene groups, such as cytoskeletal genes, GPCRs or MAP kinase pathways, we
provide annotations despite their limited expression dynamics during fruiting body
development. Such gene families may be important in fruiting body development despite
their low expression dynamics, therefore, we decided to annotate them in order to facilitate
comparative discussion across mushroom-forming fungi and more thoroughly investigated
Ascomycota model systems.
In the following chapters we discuss each of the more important gene groups in
detail, with emphasis on expression profiles, putative function and role in fruiting body
development. Besides well-known ones, we intentionally focus on lesser-known gene
families with widespread developmental expression, to potentially distill new insights into
fruiting body development. We provide catalogues and references to recent reviews for well-
researched gene families as well (e.g. hydrophobins or laccases).
645

4.1. Cell division, proliferation and growth

4.1.1. Meiotic and mitotic gene expression show two distinct expression peaks

Mitosis and meiosis are key to the development and growth of multicellular organisms.
Mushroom fruiting bodies are no exception: mitosis leads to cell proliferation early in fruiting
body development, whereas meiosis happens in basidia, which are localized to gills, to
produce spores.
Transcriptomic data provided clear evidence for expression peaks of genes related to
mitotic or meiotic cell division, as well as associated processes in DNA replication, repair,
chromosome dynamics/movement etc., during fruiting body development. The enrichment of
meiotic/mitotic genes during fruiting body development and sporulation was noted in *C.*
cinerea(Burns et al. 2010; Muraguchi et al. 2015; X et al. 2015; Krizsán et al. 2019),
Pisolithus microcarpus(Pereira et al. 2017)*, Tricholoma matsutake*(Tang et al. 2020) and *L.*
edodes(Song et al. 2018b), *Agaricus blazei*(Lu et al. 2020) as well as in a six-species
comparison published by Krizsan et al(Krizsán et al. 2019), among others. Similar signal for
meiotic gene upregulation has been detected in Ascomycota fruiting bodies as
well(Rodenburg et al. 2018). *C. cinerea* has been used as a model system of Basidiomycota
meiosis, in particular in so called white cap mutants that have meiosis-associated defects(Lu

et al. 2003). These works revealed many conserved and some unique aspects of the meiotic
process as well as resulted in meiosis being one of the best understood processes in this
species(Seitz et al. 1996; NY et al. 1997; L et al. 1999; Gerecke and Zolan 2000; Nara et al.
2001; Cummings et al. 2002; Lu et al. 2003; SH et al. 2004; Muraguchi et al. 2008a; AM et
al. 2009).
Probably the most detailed such study was performed by Burns et al(Burns et al.
2010), who investigated by microarray the transcriptional events that happen in *C. cinerea*
gills during a time course that encompasses the meiotic events. They found that 2,721
genes exhibited changing probe intensity during the six time points around meiosis. Using
dikaryotic mycelium as reference, they identified 886 genes that were expressed in gill tissue
only; these genes included core meiotic components. Like in other species, genes were
induced in successive waves during the meiotic process; these waves were arranged into 9
clusters based on expression trajectories. The clusters separated genes into functional
groups, e.g. those required for prophase I or sporulation in early and late in the time series,
respectively. Early genes further included ones related to DNA replication, cytoskeleton
organization and regulation. Late induced genes in their study are mostly related to spore
formation and show overlaps with genes related to ascospore formation in *S. cerevisiae*(S et
al. 1998a; Primig et al. 2000) and *Sch. pombe*(Mata et al. 2002). Ribosomal protein coding
gene expression was found to be high up to karyogamy, at which point it starts to decrease
and does not lift up again. This might reflect the shutting down of protein synthesis as the
fruiting body enters the sporulating phase.
Freitas Pereira et al examined sporogenesis in *Pisolithus*, an ectomycorrhizal fungus
in the Boletales, and found that meiotic genes (annotated by KEGG) showed dynamics
throughout the development of basidiospore-containing peridioles(Pereira et al. 2017). In this
species spores are produced in peridioles, lentil-shaped to globular compartments within the
fruiting body. Several genes that were differentially expressed in peridioles of different
maturation stages were annotated as meiotic/cell cycle or cell division related, most of which
showed an expression peak in unconsolidated, young, and mature peridioles. These
observations are consistent with microscopic observations on the sporogenesis of *P.*
microsporus(AN and MD 2010).
We annotated 180 conserved orthogroups as involved in cell proliferation, such as
mitosis, meiosis, DNA replication, DNA repair or the cell cycle, based on expression
conservation 141 of these qualified as CDE orthogroup (Table 1). These included previously
functionally characterized genes such as *ku70*, *rad50* and *Mre11* of *C. cinerea*(AM et al.
2009; Nakazawa et al. 2011), DMC1 (=Lenedo1_1211634) of *L. edodes*(Sakamoto et al.
2009)*, ku80* of *S. commune*(De Jong et al. 2010) and *Msh4* of *P. ostreatus*(Lavrijssen et al.
2020). Most of these genes showed developmental regulation and similar expression profiles
in most of the species. This is not surprising, given that both mitosis and meiosis have well-
documented roles in sculpting fruiting bodies. Nevertheless, to our knowledge, these genes
represent the most comprehensive, although surely not complete, list of cell proliferation
related genes to date. We tentatively subdivided CDE orthogroups into ones related to DNA
replication, DNA repair, meiosis as well as both mitosis and meiosis; these contained 87, 26,
13, 54 genes, respectively (Table 1). These include members of several well-characterized
protein complexes described from S. cerevisiae, such as Ku, SMC, MCM, Ndc80, Gins and
cohesin complexes (Table 1). Orthologs of *C. cinerea* arp9, which was described as a
putative actin-related protein involved in chromatin remodeling(Nakazawa et al. 2016), were
also found to follow similar expression profiles, although at moderate dynamics (Table 1).

These genes showed characteristic expression profiles in most species. The
expression of meiotic genes shows a distinct peak in gill tissues (Fig. 3, Supplementary Fig.
1) or in stages that contain meiotic tissue. Meiotic gene expression precedes early phases of
sporulation and was clearly discernible in *C. neoformans* or *P. ostreatus* in previous
transcriptomic studies(Liu et al. 2018; Merényi et al. 2021). Because the onset of meiosis is
tightly regulated in most species (among mushrooms best known in *C. cinerea*(Kües 2000))
meiotic gene expression also provides a landmark to calibrate developmental chronologies
among species where this may be hard to do because of morphological divergence. It
should be noted that some species develop in ways so that we do not expect a clear peak.
For example, cell proliferation and spore production is not synchronized in *S. commune*,
where it appears that sporulation progresses from the basal, older parts and progresses
towards the edges of the fruiting body(Kües and Liu 2000). In *S. commune*, this may be
explained by the highly derived morphology of the fruiting bodies, consisting of an
assemblage of cyphelloid fruiting bodies. Therefore, we would not expect one clear peak of
meiotic gene expression, unless we sample different zones of the fruiting body separately.
On the other hand, mitotic, DNA replication and repair genes showed a clear peak
associated with primordium stages, and their expression leveled off in mature fruiting bodies
(Fig. 3, Supplementary Fig. 1). This probably reflects intense cell proliferation in primordia,
which establishes the main tissues and developmental modules of the mushroom. The
expression of these genes later decreases to low levels in young fruiting body stipes and
caps (also high in meiotic tissues where one round of mitosis happens) and in mature fruiting
bodies. This peak was also detected in previous studies on *Agaricus blazei*(Lu et al. 2020)
and *P. ostreatus*(Merényi et al. 2021) and probably relates to the biphasic development of
fruiting bodies, where the first phase is concerned with cell proliferation and tissue formation
whereas the second with growth by cell expansion without much change in cell numbers.
Our data suggest that this pattern should be conserved across agarics, but potentially
missing or less clear in species that follow different developmental patterns (e.g. polypores).
Cryptococcus neoformans recently proved to be a particularly useful model for
teasing apart morphogenetic processes(Merényi et al. 2021). This species forms simple
fruiting bodies composed of lawns of aerial hyphae bearing basidia. Liu et al pointed out in
Cryptococcus, that genes co-induced with meiotic genes might be related to basidium
morphogenesis, a process that greatly overlaps with meiosis(Liu et al. 2018). Based on
transcriptome data it may be hard to separate such genes from meiotic genes, because of
the temporal and spatial overlap of their expected expression. Nevertheless, one of the
genes reported by Liu et al, *C. neoformans* Csa1 (*C. cinerea* protein ID: 471238, see Table
TRA below), which has been implicated in basidium development (and to a lesser extent
meiosis(Liu et al. 2018)), is developmentally expressed in 10/10 species in our data and
shows expression peaks that overlap with that of meiotic genes.
An orthogroup that includes the serine-threonine kinases *Sch. pombe* Ran1 and *S.*
cerevisiae Sks1 (represented by *C. cinerea* 456276, Table 1) showed marked peaks in gills
of mature fruiting bodies in all agaricoid species (except *C. cinerea*) and in mature fruiting
bodies in species with no separable gills or caps (*S. commune, A. ampla*). Ran1 is a
negative regulator of *Sch. pombe* meiosis (with Mei2 as target(Caligiuri et al. 1997)),
whereas Sks1 is involved in the adaptation to low glucose concentrations and pseudohyphal
growth in yeast. Despite this apparent strong contradiction, the characteristic expression
peaks in meiotic tissues and developmental stages in mushroom-forming fungi suggest roles
in sporulation-related processes, possibly the repression of meiotic processes during spore
morphogenesis.

In summary, mitotic and meiotic (including associated processes such as DNA
replication and repair) gene expression showed conserved patterns across the examined
mushroom-forming fungi. This probably reflects, on one hand, intense cell proliferation
during primordium development and, on the other hand, meiosis that precedes sporulation in
basidia. These gene expression patterns are easily recognizable in transcriptomic data and
provide landmarks for calibrating developmental series. The downregulation of mitosis
related genes in the second half of fruiting body development coincides with ceasing cell
proliferation and the transition of the mushroom to growth by cell expansion.

766   4.1.2. Ribosomal genes

Ribosomes produce proteins, the necessary building blocks for living cells. They are
composed of four RNA subunits and up to 80 different proteins. The amount of ribosomes,
and thus ribosomal protein expression, in a cell is proportional to its protein synthesis
demand(Kraakman et al. 1993; Jorgensen et al. 2002). Therefore, ribosomal protein
expression has been used as a proxy for the proliferative activity of a cell or a tissue and we
expect it can be similarly informative in mushrooms too.
We observed a preponderance of ribosomal protein genes among CDE orthogroups,
especially at FC>2 and to a lesser extent at FC>4. To look globally at structural proteins of
ribosomes, we identified ribosomal protein genes based on previously reported ribosomal
compositions, excluding mitochondrial ribosomal proteins and ribosome biogenesis-related
genes. The 74 orthogroups identified this way (Table RIB) showed very characteristic
expression dynamics, albeit at low fold change values, in most species. Ribosomal genes
were highly expressed in the early developmental stages of most species followed by
gradual shutting down in young and mature fruiting bodies. A second, sharp expression peak
was observed in gill tissues of *P. ostreatus, A. ostoyae, A. bisporus* and *C. cinerea (*Fig. 4,
Supplementary Fig. 2). Like in the case of mitotic and meiotic genes, we hypothesize that
these two peaks correspond to an early, proliferative stage of development and the protein
synthesis demand of spore production/meiosis. Ribosomal genes seem to be co-expressed
with genes involved in cell division and proliferation (meiotic, DNA replication and repair, see
above) in both primordia and in the gills. This could signify the need for increased protein
production for cell proliferation in primordia and for meiosis and spore production, although
other processes happening in gills requiring a large amount of protein cannot be ruled out.
The characteristic expression of ribosomal genes is a conspicuous pattern and was detected
in partial or complete form in several transcriptomic papers in both Basidiomycota(Cheng et
al. 2013b; Zhou et al. 2014; Zhang et al. 2015b; Song et al. 2018b; Krizsán et al. 2019; Liu
et al. 2020a; Lu et al. 2020) and Ascomycota(Rodenburg et al. 2018; Tong et al. 2020).
The vegetative mycelium generally showed high ribosomal protein-coding gene
expression, though not in all species, which we hypothesize depended on whether samples
for RNA-Seq were harvested from older, inactive or actively growing (e.g. colony edge) parts
of the mycelium.
A ribosome-bound Hsp70 chaperone protein formed a CDE orthogroup
developmentally regulated in 9 species (FC>2) (*C. cinerea* protein ID: 438977) (Table 2).
This gene was also detected previously as upregulated in primordia of *C. cinerea*(Cheng et
al. 2013b). It is orthologous to *S. cerevisiae* Ssb2, a Hsp70-family chaperone that is bound
by the ribosome and escorts the folding of nascent polypeptides(Peisker et al. 2010).
In summary, a characteristic expression trajectory of ribosomal genes is
characteristic of most mushroom-forming fungi, which likely reflects high protein demands of

intensely growing tissue in primordial stages and gills and supports a bi-phasic model of
fruiting body development (see below).

4.1.3. Growth by cell expansion and turgor manipulation - a biphasic model of fruiting
body development

A key difference in fruiting bodies and hyphae is that in the latter growth and expansion
takes place at the tip, whereas in the former, cell expansion can vary between apical and
isotropic. Already de Bary established, by studying *Mycena vulgaris*, that mushroom fruiting
bodies grow mostly by cell expansion not by cell proliferation(Bary et al. 1887). This appears
to be a rather widespread feature of fruiting bodies and shows broad similarity to growth by
cell expansion in plant fruits(Krizsán et al. 2019). More precisely, fruiting body development
seems to be divided into a first proliferative stage, during which cell division, differentiation
as well as tissue formation takes place, and a growth phase, which is primarily concerned
with cell expansion to achieve the mature shapes and sizes of the fruiting bodies(MOORE et
al. 1979; Kües 2000; G et al. 2013; Kües and Navarro-González 2015; Krizsán et al. 2019;
Liu et al. 2021). Of the better investigated species, this two-phased development is probably
most spectacular in *C. cinerea*, which is adapted to very rapidly lifting cap above ground and
expanding it for spore dispersal(Kües 2000).
It should be noted that the two-phase development is not fully universal among fungi
and does not always mean a complete separation of proliferative and expansive phases. For
example, Craig et al(Craig et al. 1977) showed that the top regions of *A. bisporus* stipes
elongated up to 9-fold, whereas cell lengths increased up to 2.2-fold, and from these data
derived that cell division also occurs to achieve this elongation rate. In *S. commune*, it
appears that outer parts of the fruiting body are still growing (by cell proliferation) while older
parts are already sporulating(Kues and Liu 2000).
Whether for the entire fruiting body or only parts of it, growth by cell expansion
seems to be a widespread trait during the second half of development of mushrooms. How
cell expansion is achieved, in general, is poorly known, but it probably requires water supply,
turgor manipulation (by osmolytes), cell wall elasticity and new cell wall material deposition.
Water supply may be realized by the regulated expression of aquaporin genes (see below).
On the other hand, the identity of the osmolytes has not been resolved yet. In the
Ascomycota ions, ion channels, glycerol and sugars were reported as main players of turgor
manipulation(Fischer et al. 2004). In Basidiomycota fruiting bodies, sugars (mannitol,
trehalose) offer themselves as straightforward candidates, but Moore([CSL STYLE ERROR:
reference with no printed form.]) speculated that high molecular weight sugars are unlikely to
be the osmolytes that drive stipe expansion. Trehalose and polyols have been excluded as
osmolytes(Kües 2000; F et al. 2005), whereas potassium oxaloacetate has been
hypothesized, which would be consistent with the upregulation of oxalate metabolism (see
below), but not confirmed(Knoll; Kües 2000). On the other hand, ions would either need to
be transported to the fruiting body from the mycelium, or released locally.
Factors modulating cell wall elasticity and remodeling are better known. During
fruiting body growth, cell wall deposition occurs throughout the cell surface, and not only in
the tip region as in hyphae, which was termed diffuse extension(Kües 2000). In *Agaricus*
bisporus larger proportion of β-1,6-linked glucose side branches in a β-1,3-glucan backbone
was observed in the rapidly elongating top segment of the stipe(Mol and Wessels 1990).
This causes hydrogen-bonds among glucan chains to be weaker in rapidly elongating stipe
compared with mature stipe and vegetative hyphae. It has been hypothesized that these

850 weak bonds are disrupted and reformed by turgor pressure rather than hydrolysing
851 enzymes(Mol et al. 1990). The architecture of cell wall also regulates stipe elongation; in
852 primordial stages chitin microfibrils are arranged isotropically which is remodeled to helical
853 arrangement during development(Kamada et al. 1991). The turgor pressure acting on the
854 cell wall and insertion of new chitin microfibrils allow elongation and in that dynamic
855 rearranges of microfibrils chitin synthases, chitinases and glucanases play the major role in
856 *C. cinerea*(Kamada et al. 1991). Niu et al observed differences in structure of chitin
857 microfibrils in elongating and mature stipe regions(Niu et al. 2015). Extension of the cell wall
858 happens by insertion of newly synthesized microfibrils. In contrast, studies in *Flammulina*
859 and *Coprinopsis* suggested that stipe cell wall extension is mediated by endogenous
860 expansin-like proteins, not turgor pressure or cell wall glycoside hydrolases(Fang et al. 2014;
861 W et al. 2014). Plant expansins mediate Turgor-mediated cell expansion(Baccelli 2015) and
862 several other morphogenetic processes (e.g. organ abscission, fruit softening(Sampedro and
863 Cosgrove 2005)). The auxin-driven expansion of plant cells without changes in cell number
864 is known as acid growth(Rayle and Cleland 1992; Arsuffi and Braybrook 2018), because low
865 pH is required by cell wall modifying proteins (e.g. expansins) to loosen the cell wall. In fungi,
866 it is not known if an analogous process exists, but the acidifying effect of external CO_2 has
867 been hypothesized as a factor in stipe cell elongation(Liu et al. 2021) and there is evidence
868 for expansins participating in the process (see below). Accordingly, high external CO_2 during
869 mushroom cultivation was speculated to cause malformed, long-stipe fruiting bodies by
870 acidifying the extracellular space and activating cell wall modifying proteins(Liu et al. 2021).

871 4.2. Acetyl-CoA production by central carbon metabolism

872 We found evidence for widespread developmental expression of multiple acetyl-coenzyme A
873 (Ac-CoA) producing enzyme families. Ac-CoA is a key intermediate in central carbon
874 metabolism as well as other processes (biosynthesis of lipids and certain amino acids(Chen
875 et al. 2015)). Under normal carbon metabolic conditions, the generated Ac-CoA, which
876 comes from either glycolysis or fatty acid beta-oxidation enters the TCA cycle, where it is
877 oxidized to CO_2 for energy (ATP) production. However, Ac-CoA is required also for the
878 production of biological membranes(Strijbis and Distel 2010; Chen et al. 2015), and our
879 results point in the direction of multiple basic metabolic pathways converging on excess Ac-
880 CoA production for membrane biosynthesis. Combined with gene expression profiles of
881 these genes, we hypothesize that excess Ac-CoA is being channeled into ergosterol and
882 fatty acid production during fruiting body development, as detailed below.
883   For this work we started from Ac-CoA related CDE orthogroups detected in the
884 transcriptomes of the examined species and developmental stages, which belonged to
885 several pathways. We reasoned that if there is an increased demand for Ac-CoA at certain
886 developmental stages of fruiting bodies, then this should be reflected in the regulation of Ac-
887 CoA-producing pathways. Thus, we annotated the TCA cycle, the glyoxal cycle, the fatty
888 acid beta-oxidation, ethanol utilization, glycolytic as well as pyruvate decarboxylation
889 pathways of *C. cinerea*, based on protein orthology to corresponding pathways of *S.
890 cerevisiae* (Fig. 5, Supplementary Fig. 3, Table 3). These led to a model we put forth on Fig.
891 5, with the additional note that it reflects one possible synthesis of the observations into a
892 mechanistic model compatible with what we know about fruiting body development, but that
893 other interpretation may also be viable.
894   First of all, we found that oxaloacetase (oxaloacetate lyase) genes were
895 developmentally expressed in 11 species (FC>4, *C. cinerea* 499734). Oxaloacetases

produce oxalate and acetate from oxaloacetate, which is an intermediate of both the TCA
and the Glyoxylate cycles. In *S. cerevisiae* acetate is converted to Ac-CoA by acetyl-CoA
synthases (Acs1), orthologs of which were developmentally regulated in 11 and 4 species at
fold change of 2 and 4, respectively (represented by *C. cinerea* 500719). We also found a
CDE orthogroup comprising putative orthologs of the *S. cerevisiae* Ac-CoA hydrolase Ach1
(represented by *C. cinerea* 442787, FC>2/4: 7/5). Ach1 is a relatively poorly known gene,
which was recently shown to be responsible for the conversion of Ac-CoA to acetate, which
can then be easily transferred from the mitochondrion to the cytoplasm(Chen et al. 2015).
Cytoplasmic acetate is then converted to cytoplasmic Ac-CoA, a starting material for lipid
synthesis, suggesting that Ach1 mediated Ac-CoA supply for lipid synthesis. Remarkably,
Ach1 orthologs are present in only 8 species and completely missing in *A. bisporus* (missing
even in telomere-to-telomere assemblies(ASM et al. 2020)), *M. kentingensis, Ph.*
chrysosporium, L. tigrinus, Pt. gracilis. This patchiness might indicate that the role of these
genes is dispensable and probably others can take over their roles.
Acetate is also an intermediate of the ethanol utilization pathway, in which two
enzymes form a linear pathway that terminates in the production of Acetyl-CoA(M et al.
2003). Aldehyde dehydrogenases orthologous to AldA of *A. nidulans* and Ald4 of *S.*
cerevisiae formed a CDE orthogroup (*C. cinerea* protein ID: 500663, 9/7); the encoded
enzymes also contribute to acetate production and, in *C. cinerea*, showed peak expression
in young fruiting body caps. Ald4/AldA converts the toxic compound acetaldehyde to
acetate(B et al. 2001). Although the ethanol utilization pathway is mostly discussed in the
context of extracellular ethanol (a nonfermentable carbon source) uptake and degradation,
as well as anaerobic fermentation, acetaldehyde can also be converted to ethanol (Fig. 5), if
it is present in excess amounts in the cells. This may explain the developmental regulation of
orthologs of *S. cerevisiae* Adh3 and *A. nidulans* AlcA/AlcC (represented by *C. cinerea*
500663), which were developmentally expressed in 11/7 species (FC>2/4, Table 3).
Enzymes of the alcohol utilization pathway were recently detected in proteomics data of
fruiting bodies of *Guyanagaster necrorhiza*, where their presence was interpreted as
evidence for anaerobic fermentation within the fruiting body, as part of an intriguing
mutualism with N_2 fixing bacteria(Koch et al. 2021).
The last step of glycolysis is catalyzed by the pyruvate decarboxylase Pdc1 in *S.*
cerevisiae, which converts pyruvate to acetaldehyde. Pdc1 generates cytoplasmic Ac-CoA,
which is essential for the biosynthesis of lipids(Chen et al. 2015). Pdc1 orthologs
(represented by *C. cinerea* 422402) were developmentally expressed in 11/8 species
(FC>2/4), whereas genes encoding other enzymes in glycolysis showed much smaller
dynamics (Supplementary Fig. 3). Together, these observations suggest a widespread
upregulation of acetaldehyde and from that either acetate or ethanol producing enzyme
genes in fruiting bodies.
Upstream of oxaloacetases, enzymes responsible for oxaloacetate production in the
TCA (yeast Mdh1, *C. cinerea* 500663, 11/3) and glyoxylate cycles (yeast Mdh2, *C. cinerea*
449845, 9/3) were also quite widely developmentally regulated. In the glyoxylate cycle we
found orthologs of *S. cerevisiae* Mls1 (malate synthase, *C. cinerea* 530292) and Icl1
(isocitrate lyase, *C. cinerea* 495730) to form CDE orthogroups developmentally regulated in
12/11 and 12/7 species (FC>2/4), respectively. This suggests that the glyoxylate cycle is
strongly induced in fruiting bodies. On the other hand, members of the TCA cycle showed
lower developmental dynamics (Fig. 5).
What happens to the oxalate that is produced by oxaloacetases? It is possible that
the resulting oxalate is a waste product, but it may also be used somehow by the fungus,

e.g. as a developmental signal molecule(Palmieri et al. 2019) or as an electron donor, as
has been observed in wood-decay(Hastrup et al. 2012; Zhu et al. 2016; Grąz et al. 2017),
where it acts as an electron donor for oxidative enzymes. One hypothesis for the role of
oxalate is that oxidative enzymes, which might participate in the modification of the fungal
cell wall during development (see above), utilize it as an electron donor, as do enzymes
involved in wood-decay. On the other hand, if oxalate is a by-product, excess amounts may
need to be broken down actively. We found evidence for the upregulation of oxalate
catabolism in the compared species. The two oxalate degradation pathway in fungi
comprises oxalate decarboxylase, which converts oxalate to formate and CO_2 and formate
dehydrogenase, which breaks formate down to CO_2(Mäkelä et al. 2014). We detected two
CDE orthogroups of oxalate decarboxylases (*C. cinerea* protein ID: 361522, 8/6,
Schco3_2751877, 7/6) and one of formate dehydrogenases (*C. cinerea* protein ID: 452378,
7/6, orthologous to Fdh1 of *S. cerevisiae*)(Table 3), which were upregulated in fruiting bodies
widely, suggesting that fruiting bodies utilize enzymatic mechanisms for the neutralization of
oxalate. An alternative route of oxalate degradation in *S. cerevisiae* is catalyzed by PCS60,
an oxalyl-CoA synthetase(Foster and Nakata 2014). Orthologs of this gene were found in
only a subset of species (represented by *C. cinerea* 474928), but were developmentally
regulated in most of them. In addition, a CDE orthogroup of formate transporters,
orthologous to the *Postia placenta* gene reported in Martinez et al(Martinez et al. 2009b),
was developmentally expressed in 8/5 species (FC>2/4, represented by Schco3_2738153,
missing in *C. cinerea*).
Building on the evidence for intense Ac-CoA production, we annotated and analyzed
the Coenzyme-A producing pathway as well as two other sources of Ac-CoA, the fatty acid
beta-oxidation pathway and the cellular cycle for pyruvate decarboxylation to Ac-CoA. These
pathways were reconstructed and annotated based on corresponding pathways in *S.
cerevisiae*(JM et al. 2012). Members of these pathways did not show remarkable
developmental dynamics (Table 3).
Several genes connected to acetate metabolism and transport also formed CDE
orthogroups. Orthologs of *S. cerevisiae* Ady2, an acetate transporter(S et al. 2004), are
widely developmentally regulated (represented by *C. cinerea* 441597, 7/6), and were
strongly induced in young fruiting body caps of *C. cinerea*. A CDE orthogroup of putative
monocarboxylate (lactate, acetate, succinate, etc.) transporters (*C. cinerea* protein ID:
375404 [7/5])(Table 3) orthologous to *S. cerevisiae* Jen1(M et al. 1999) were also highly
induced in YFB-caps in *C. cinerea*. We also identified putative orthologs of *S. cerevisiae*
Yat1 (represented by *C. cinerea* 443703, 10/5), a carnitine acetyltransferase that shuttles
Ac-CoA across the mitochondrial membrane. A CDE orthogroup of aquaglyceroporins
orthologous to *S. cerevisiae* Fps1, which is involved in the uptake of acetate (*C. cinerea*
protein ID: 373141 7/6), were developmentally regulated in 7/6 species. This gene encodes
a protein that has been associated with either glycerol influx/efflux(Luyten et al. 1995) or
acetate uptake(Mollapour and Piper 2007; Watcharawipas et al. 2018), which leaves some
uncertainty on which role it fulfills in fruiting bodies. Orthologues of *S. cerevisiae* Sym1 also
formed a CDE orthogroup, although only at a fold change cutoff of 2 (*C. cinerea* protein ID:
147126, 9/3) (Table 3). This gene is said to be involved in ethanol utilization and stress
response, but mechanistic details are not known(Reinhold et al. 2012; Gilberti et al. 2018).
Similarly, orthologs of yeast Sfc1, a succinate/fumarate antiporter that transports C4
succinate (which is produced from acetate(Bojunga et al. 1998)), to the mitochondria, are
developmentally regulated in several species (represented by *C. cinerea* 374148, 12/7). The
Sfc1 ortholog is also strongly induced late in the development of *C. neoformans*(Liu et al.

2018). Finally, a CDE orthogroup of putative malate symporters orthologous to *Sch. pombe*
Mae1, which is involved in the uptake of extracellular malic acid was detected as
developmentally regulated in 9/5 species (FC>2/4, represented by *C. cinerea* 534888, Table
3). Whether these genes are related to the above discussed observation of acetate/Ac-CoA
related gene upregulation, or how cell surface transporters may be related remains to be
determined. The induction of these genes does not happen in the same or similar
developmental stage in all species (although there is consistency within species, see the
upregulation of several genes in YFB-caps of *C. cinerea*), nevertheless, their induction
suggests differential acetate concentrations and an important role for acetate during in
fruiting body development.
CO_2 produced by the TCA cycle may be fixed into oxaloacetate by the conversion of
CO_2 to hydrogencarbonate by carbonic anhydrases, then conversion of HCO_3^- and pyruvate
into oxaloacetate by pyruvate carboxylases(Elleuche and Pöggeler 2010) (Pyc1 in *S.*
cerevisiae). We found that several carbonic anhydrases (2-9 genes in Agaricomycetes
genomes) are developmentally regulated in mushroom forming fungi and they form at least
one CDE orthogroup (represented by *C. cinerea* 450047)(Table 3). *C. cinerea* 450047
shows an upregulation in young fruiting bodies, coincident with the induction of Ac-CoA
producing enzymes. This observation raises the possibility that CO_2 produced by the TCA
cycle in fruiting bodies may be fixed into oxaloacetate, which can further be converted to Ac-
CoA (Fig. 5). Of note, carbonic anhydrases are involved in the regulation of sexual
development in the Ascomycota(Elleuche and Pöggeler 2010).
Taken together, we observed developmental regulation of key enzymes in several
pathways that seem to converge on increased acetyl-CoA production, either directly, or
indirectly (through acetate or oxaloacetate). Several connected pathways, including oxalate
degradation, the ethanol degradation pathway, the terminal reaction of glycolysis as well as
evidence from the TCA and glyoxylate cycles were pointing in this direction. The expression
of these genes tended to be upregulated late during development and showed good
consistency within, but not necessarily across, species. A possible interpretation of these
data is that increased Ac-CoA production is needed for growth by cell expansion late in
development, as a starting material for lipid and sterol biosynthesis. This is underpinned by
the fact that biosynthetic pathways of ergosterol and other membrane components are also
upregulated during late developmental stages (e.g. the young fruiting body stage in *C.*
cinerea), probably for membrane assembly needed for rapid cell growth. This interpretation
would be consistent with the two-phased development of several mushroom-forming
species, which consists of a first stage of cell division and differentiation and a second,
during which cell numbers hardly change but their volumes increase considerably via
Turgor-mediated expansion. Such a reprogramming of the fungus's basic metabolism from
using Ac-CoA for energy production (when citrate is in the mitochondrion) to using it for lipid
metabolism (when citrate is in the cytoplasm) seems particularly well-suited for fruiting body
formation, which requires tight regulation of lipid metabolism. It should be mentioned that Ac-
CoA is also required for melanin production(Lee et al. 2019), although it is unlikely to be the
driving force of our observation, because pigment production is not conserved across the
examined species. A previously reported reprogramming of the glyoxylate cycle in
Fomitopsis palustris for the production of oxalate(Munir et al. 2001) seems related, however,
in that case, the main product in demand was hypothesized to be oxalate that facilitates
wood-decay, whereas in our case we hypothesize the product in demand is acetate and
finally Ac-CoA. Both variants of the metabolic pathways nevertheless pass through
oxaloacetate(Munir et al. 2001). The excess Ac-CoA production may rely on, or be

connected to carbohydrate reserves that are deposited in the subhymenial part of the cap of
the mushroom(Ji and Moore 1993; Muraguchi et al. 2015) during primordium development.
However, there may be alternative interpretations and the topic is certainly of interest for
detailed, further research, that integrates more functional data beyond gene expression
observations.
1045

1046 4.3. Lipid metabolism

Lipid metabolism is pivotal to very basic organismal processes such as growth,
differentiation, cell division, energy storage and cell membrane composition, to name a few.
Therefore, in depth understanding of gene expression changes associated with lipid
metabolism, catabolism and transport can be expected to provide novel insights into fruiting
body development. Indeed, our meta-analysis of Agaricomycete transcriptomes revealed
several CDE orthogroups related to lipid metabolism. We discuss these below,
complemented with annotations of major related pathways (but no attempt is made to
comprehensively annotate all pathways) and processes. We base our annotations on data
from well-studied model organisms, such as *S. cerevisiae, A. nidulans* and *Sch. pombe*, with
orthology relationships to Agaricomycetes genes established on the basis of 1-to-1
reciprocal best Blast hits.

1058 4.3.1. Yeast beta oxidation

We reconstructed the beta-oxidation pathway (Table 4) in *C. cinerea* based on the yeast
beta-oxidation pathway(van Roermund et al. 2003) (Fig. 5). The expression of genes
comprising the beta-oxidation pathway in general does not appear to be widely dynamic in
fruiting bodies. We found orthologs of yeast CTA1, a catalase A, to form a CDE orthogroup.
However, this catalase in yeast is only a marginal component of the beta-oxidation pathway,
and may be involved in a wide range of other functions (e.g. signaling). The more
widespread developmental expression of ACS1 orthologs is remarkable (11/4 species at
FC=2/4), for a possible connection with sterol biosynthesis (see above). ACS1 produces
Acetyl-CoA, which is the starting point for ergosterol and membrane lipid biosynthesis.

1068 4.3.2. Ergosterol biosynthesis

We reconstructed the ergosterol biosynthesis pathway based on orthology to known
pathway member genes in *S. cerevisiae*(JM et al. 2012) (Table 5). ERG pathway genes of
Agaricomycetes are generally higher expressed in fruiting bodies than in vegetative mycelia
(VM/P1>1 in 20 out of 21 genes in *C. cinerea*), with expression peaking in young fruiting
bodies of *C. cinerea* (Fig. 6, Table 5). Expression peaks at similar stages in other species
too, in primordium 3/young fruiting body in *P. ostreatus* and young fruiting bodies/mature
fruiting bodies stages in *A. ostoyae* (Supplementary Fig. 4). Similar pictures were obtained
for other species. It should be noted that although the ergosterol biosynthesis pathway
shows clear expression trends, the dynamics of expression is relatively low, mostly between
2<FC<4. Therefore, ergosterol biosynthesis genes mostly do not form CDE orthogroups,
nevertheless, the expression patterns are distinctive so they warrant discussion here.
In these species expression peaks seem to coincide with intense growth by cell
expansion in fruiting bodies. Above we showed that mitotic, ribosomal and histone gene
expression level off in late primordium stages (P2 in *C. cinerea*, see Fig. 3), after which

growth by cell expansion takes place. This might create an increased demand for membrane
constituents during rapid cell expansion/growth by turgor, which is what we might see in the
expression patterns of genes in the ergosterol biosynthesis pathway (Fig. 6, Supplementary
Fig. 4). This hypothesis would also explain why there is no clear peak in *C. neoformans* and
Pt. gracilis, where sudden growth by cell expansion is lacking.
Similarly, a homolog of the Sterol-Regulatory Element Binding Proteins (SREBP(CM
and PJ 2010)), similar to the *S. cerevisiae* transcription factor HMS1 shows a very similar
expression pattern: in *C. cinerea* (*C. cinerea* protein ID: 488426) expression peak in YFB-S
and YFB-C (Fig. 6), whereas in *P. ostreatus* it peaks in mature fruiting body tissues
(PleosPC15_2_174910). This suggests, based on expression data, that these putative
SREBPs might be regulating ergosterol biosynthesis genes also in the Agaricomycetes.
A connection with the above-mentioned Ac-CoA producing basic metabolic
pathways, where certain, but not all, components were found widely upregulated, is possible
here. These pathways produce Acetyl-CoA, which is the starting point of ergosterol
biosynthesis. This raises the possibility that the upregulation of the ethanol utilization
pathway components (see above, Fig. 5) and certain components of the glyoxylate cycle
reflect increased Ac-CoA production which is required as input to ergosterol biosynthesis
and eventually membrane assembly. A competing hypothesis was put forth in *P.*
microcarpus, where upregulation of glyoxylate shunt genes was observed and suggested
that it underlies the production of storage materials for spores(Pereira et al. 2017). Using our
more resolved transcriptome data and based on the observation of ergosterol pathway
expression dynamics, we propose that the connection to the glyoxylate shunt might
represent a link to membrane constituent synthesis, although more data would be needed to
verify this hypothesis.
The upregulation of the ergosterol synthesis pathway was observed in *F. velutipes*,
where most ergosterol biosynthesis genes were upregulated in the young fruiting body
stage(Wang et al. 2019) (approximately in agreement with our definition of young fruiting
bodies) and showing intense growth. Sterol metabolism genes were also found to be more
abundant in fruiting bodies of *H. erinaceus* compared to vegetative mycelia(Zeng et al.
2018). It was reported that orthologs of the lanosterol-alpha-demethylase Erg11 of *S.*
cerevisiae, a component of the ergosterol biosynthesis pathway (corresponding *C. cinerea*
ortholog: 397292), was upregulated in fruiting bodies of *Antrodia cinnamomea*(Lu et al.
2014) and *P. ostreatu*s(Sunagawa and Magae 2005). This is consistent with the
observations made in this study.
We also checked the sphingolipid biosynthesis pathway based on orthology to
pathway members in *S. cerevisiae*. Sphingolipids, along with sterols are important in
membrane rafts and microdomains and form key components of fungal cell membranes.
Although some yeast genes did not have clear orthologs in Agaricomycetes, genes involved
in sphingolipid biosynthesis showed a similar expression profile in *C. cinerea* (Fig. 6, Table
5). This may underscore our hypothesis on the link between growth by cell expansion and
membrane assembly.
A characteristic CDE orthogroup of putative C14-sterol reductase genes was
developmentally expressed in 12/7 species (represented by *C. cinerea* 539004)(Table 5, Fig.
6). These proteins are not part of the ergosterol biosynthesis pathway, but show a high
similarity to *A. nidulans* and *C. albicans* membrane phospholipid metabolism proteins
AN2040 and Erg24, respectively, which are involved in ergosterol metabolism. Genes in this
orthogroup show peak expression in late development, coincident with rapid expansion of
fruiting bodies. In several species they were also upregulated in primordia relative to

vegetative mycelium samples. These expression patterns resemble that of ergosterol
biosynthesis genes, and suggest that this orthogroup may be linked to ergosterol
metabolism.

1134 4.3.3. Sterol transport

We detected several CDE orthogroups that were annotated as related to ergosterol transport
and showed similar upregulation in late development as ergosterol biosynthesis genes did.
We therefore inventoried sterol transport related genes of the families Niemann-Pick type C
protein family (*S. cerevisiae* Npc2), the yeast Ncr1 family, the nonspecific sterol transport
protein (Scp2) family, StAR-related lipid-transfer (START) proteins as well as members of
the Oxysterol Binding Protein Homologue (OSH) protein family(Schulz and Prinz 2007).
Of these, the Ncr1, Npc2, Scp2 and START gene families are single-copy families in
mushroom-forming fungi, whereas the OSH family is represented by 5 members in all
mushroom-forming fungi (except *S. commune* which has 6, Table 6). Npc2 (ortholog *C.*
cinerea 475807), Scp2 (ortholog *C. cinerea* 478741) and START proteins as well as one
member of the OSH family were (*C. cinerea* protein ID: 427232)(Table 6) upregulated during
the second half of development, at or after the young fruiting body stage. This is similar to
the expression patterns seen in ergosterol biosynthesis genes. One member of the OSH
family orthogroup (*C. cinerea* protein ID: 378276) showed a clearly stipe-specific expression,
with peaks in stipes of young fruiting bodies or fruiting bodies of *C. cinerea, A. ostoyae, M.*
kentingensis, A. bisporus and P. ostreatus, albeit mostly with low fold change values
(2<FC<4). Interestingly, the corresponding gene was cap-upregulated in *L. bicolor*
(Lacbi2_456536), whereas in simpler species (*S. commune, A. ampla*), it showed diverse
expression peaks. Ncr1 orthologs (represented by *C. cinerea* 494709) showed consistently
higher expression in fruiting bodies than in vegetative mycelia. Fold changes were generally
low in these gene families, but nevertheless the patterns are phylogenetically very
consistent.
These expression patterns resemble those of ergosterol biosynthesis genes, which
also showed upregulation in mid- to late development, coincident with cell expansion and
growth. This suggests that ergosterol transport, in addition to biosynthesis, is an important
process in the second half of the development of mushroom fruiting bodies. Like synthesis,
transport of sterols might be required to fulfill the needs of rapid cell expansion, which
obviously coincides with cell- and membrane surface expansion too.
Although marginally related, we mention here the CAP domain/Pry1 protein family,
which includes known sterol-binding proteins. The family has been associated with diverse
functionalities, including cell adhesion, pathogenicity (pathogenicity-related PR proteins
belong here), morphogenesis, endocrine and paracrine functions, among others(GM et al.
2008; Schneiter and Pietro 2013). This protein family includes *S. cerevisiae* Pry1, a lipid and
sterol binding protein involved in sterol export(Darwiche et al. 2017). To our best knowledge,
there are no previous reports of this family in fruiting body development, whereas, their role
in morphogenesis in mammals is well-established(GM et al. 2008). We detected, on average
2-5 copies of CAP proteins in the Agaricomycetes, with expansions up to 9 in *Pt. gracilis, L.*
bicolor and *A. ostoyae* as well as a CDE orthogroup of Pry1-related CAP-domain proteins
(represented by *C. cinerea* 470387, FC2/4: 7/6, Table 6, Fig. 6). Members of this orthogroup
showed peak expression late in development, at or after the young fruiting body stage in
most species. The sterol transporting ability and expression patterns suggest that this family
may have a role in sterol transport during the second half of development in Agaricomycetes

fruiting bodies. It should be noted that neither Polyporales species (*Ph. chrysosporium* and
L. tigrinus) had developmentally regulated genes in this family, indicating that their role, if
any, may not be universal.

4.3.4. Phospholipid and fatty acid biosynthesis

The phospholipid and fatty acid biosynthesis pathways are conserved in Agaricomycetes,
with most genes showing clear 1-to-1 orthologs to described pathway members of *S.*
*cerevisiae (*Table 7). Because active cell growth and expansion requires an excess amount
of membrane components (ergosterol, phospholipids), we examined pathways involved in
membrane phospholipid and triacylglycerol synthesis. We reconstructed these pathways
based on well-known reaction steps in *S. cerevisiae* and *Yarrowia lipolytica*(Carman and
Han 2011; Fakas 2017). The first half of the pathway, which starts from glycerol-3-phosphate
and leads to the production of phosphatidic acid, shows upregulation in young and mature
fruiting bodies of *C. cinerea* and *A. ostoyae* (Fig. 7). Phosphatidic acid sits in a branching
point of the pathway, with one branch leading to membrane phospholipids (CDP-DAG
pathway), whereas the other leads to triacylglycerol (TAG), a major storage material. The
CDP-DAG pathway shows a general upregulation in the second half of development (young
and mature fruiting bodies) in *C. cinerea, A. ostoyae, M. kentingensis, A. bisporus*
(Supplementary Fig. 5). These patterns are consistent with increased membrane lipid
biosynthesis in these stages, possibly serving the needs of increasing cell volumes during
rapid growth. On the other hand, TAG pathway members appear upregulated somewhat
later, mostly in mature fruiting bodies in most species (Fig. 7, Supplementary Fig. 5), which
coincides with spore formation and could be related to storage material synthesis in spores.
In *P. ostreatus*, both the CDP-DAG and TAG pathways show remarkably uniform
upregulation in primordium stages (Fig. 7), which is quite different from the pattern seen in
other species. This pattern in *P. ostreatus* is also observable in fatty acid biosynthesis
pathways (see below).
Similar to the CDP-DAG and TAG pathways, fatty acid biosynthesis (Table 7) shows
a signal for upregulation in young fruiting body tissues, with *C. cinerea* orthologs of *S.*
cerevisiae Acc1, Fas2, Elo2 and Faa1 showing peak expression in the YFB stage. Similar
patterns exist in *A. ostoyae, M. kentingensis* and to some extent in *A. bisporus* (Fig. 7). In *P.*
ostreatus, fatty acid biosynthesis genes are co-expressed with the other phospholipid
biosynthesis genes, showing an upregulation in primordia.
Both phospholipid synthesis and fatty acid biosynthesis require Acetyl-CoA as a
starting point. For the former, it is needed for acylation reactions in the conversion of
lysophosphatidic acid to phosphatidic acid (last element of the shared part of the pathway)
whereas in the latter, Ac-CoA is the starting point of fatty acid synthesis(Fakas 2017). This
provides another link with the dynamics of Ac-CoA biosynthesis genes noted above. As in
the case of ergosterol biosynthesis pathway members, phospholipid and fatty acid
metabolism genes showed low expression dynamics (and hence did not form CDE
orthogroups), but their expression patterns were consistent across stages and species.
The expression of mitochondrial membrane components (cardiolipin and
phosphatidylglycerol) is also peaking in young fruiting bodies, mostly in stipe, but not very
characteristic (Fig. 7).

4.3.5. Linoleic acid producing fatty acid desaturases

Several CDE orthogroups of fatty acid desaturases (FAD) were detected among conserved
developmentally expressed genes. Members of these orthogroups show high similarity to *C.*
albicans Fad2 and Fad3 and *A. nidulans* OdeA, which are involved in the synthesis of alpha-
linoleic acid. Linoleic acid is a major membrane component in fungi, which, as an
unsaturated fatty acid, regulates membrane fluidity. It has also been speculated to be the
starting point for the biosynthesis of volatiles, such as 1-octen-3-ol(A et al. 2020). Linoleic
acid showed increased proportions in fruiting bodies and reproductive structures in general
in both the Asco- and Basidiomycota(SHAW 1967; Goodrich-Tanrikulu et al. 1998; H and S
2003; Gessler et al. 2017; Song et al. 2018a). Linoleic acid is also a precursor of oxylipins,
key developmental signaling molecules that are actively produced by reproductive
cells(Combet et al. 2006; Gessler et al. 2017; Orban et al. 2021). Accordingly, in the
Ascomycota, several fatty acid desaturase gene mutations were reported to asexual/sexual
defects(Pöggeler et al. 2006b). The higher proportion of linoleic acid in fruiting bodies of
Agaricomycetes was already noted in 1967(SHAW 1967) with several similar reports(H and
S 2003; Song et al. 2018a) later. The proportion of unsaturated fatty acid (which include
linoleic acid) increases during fruiting body formation(H and S 2003) in *L. edodes*. Three
fatty acid desaturases, two in *L. edodes* (Le.Fad1 and Le.Fad2(H and S 2005)) and one in
C. cinerea (*Cc.odeA*(H and S 2003, 2005; S et al. 2007b)) have been shown to be capable
of producing linoleic acid. Fad genes were found to be differentially expressed in fruiting
bodies of *A. bisporus* previously(Morin et al. 2012).
We found 7-16 fatty acid desaturase genes in the examined Agaricomycetes, many
of which were developmentally regulated, although clear orthology of these genes to
experimentally characterized Ascomycota genes is missing. Each species had at least 2-3
FAD encoding genes upregulated in primordia relative to vegetative mycelium, suggesting
that their functions are key to the initiation of fruiting body development. FAD genes formed
five CDE orthogroups (Table 8). One of these (*C. cinerea* protein ID: 363743) was FB-init in
most species (*C. cinerea, A. ostoyae, A. ampla, M. kentingensis, P. ostreatus, S. commune,*
C. aegerita, L. bicolor), showed high expression in primordium stages, then expression
levels dropped significantly in later developmental stages. The CDE orthogroup containing
Le.FAD1 (Lenedo1_1050661) was upregulated in primordia of most species relative to
mycelium samples (*C. aegerita, A. ampla, S. commune, L. edodes. L. bicolor, L. tigrinus, M.*
kentingensis, but not *A. ostoyae, P. ostreatus* and *A. bisporus,* ortholog missing in *C.*
cinerea), although only slightly (fold change = 1.5-2x, except in *L. edodes* and *Pt. gracilis*
which showed >4-fold upregulation). The orthogroup encompassing *Cc.odeA* (*C. cinerea*
protein ID: 422393) and Le.Fad2 (orthogroup represented by *S. commune* 2511002) was
upregulated in primordia relative to vegetative mycelia of several species (*A. ampla, C.*
aegerita, S. commune, C. cinerea, L. edodes, L. tigrinus and *P. ostreatus,* but not in *A.*
ostoyae, M. kentingensis, Pt. gracilis and *A. bisporus*). Our expression data are consistent
with previous reports of transcripts/protein abundance in case of both Le.FAD1 and Le.FAD2
in *L. edodes*(H and S 2003, 2005). The other two FAD orthogroups showed less
conservation in expression (*C. cinerea* protein ID: 471478, 422976, Fig. 8), including peaks
in later developmental stages, such as young fruiting body caps.
The linoleic acid-producing ability of the proteins encoded by these genes suggests
that the detected CDE orthogroups and expression dynamics relate to linoleic acid
production. It follows from our observations of expression dynamics that linoleic acid
production is upregulated in the primordia of several species. This is consistent with previous

reports of higher linoleic and unsaturated fatty acid proportions in fruiting bodies of
Agaricomycetes (see above). There may be at least three explanations for this observation,
including (i) regulation of membrane fluidity in response to temperature changes, (ii) the
same being genetically encoded in fruiting body development, or (iii) fulfilling the
requirements of oxylipin production. We argue that hypothesis (i) is unlikely, however, we
can't speculate on hypotheses (ii) and (iii).
Regulation of membrane fluidity by increasing the proportion of unsaturated fatty
acids is a common mechanism across organisms(H and S 2005). Therefore, a question
arises whether the higher inferred proportion of linoleic acid and corresponding changes in
gene expression are genetically coded traits of fruiting bodies or simply a consequence of
lower temperature required by several species for fruiting. At least for the transcriptomes we
produced(Sipos et al. 2017a; Almási et al. 2019; Krizsán et al. 2019) we know that
vegetative mycelium samples were harvested at the same temperature as fruiting body
samples for all species, which rules out the possibility that higher linoleic acid proportion is
simply a response to lower fruiting temperatures. Rather, this suggests that higher linoleic
acid concentration is a genetically encoded trait in fruiting bodies. Similar conclusions were
reached in the context of unsaturated fatty acids in *L. edodes*(H and S 2003). From an
evolutionary point of view, such a genetic adaptation could go back to the exposed nature of
fruiting bodies; in the natural environments (soil, wood) the vegetative mycelium is in more
stable conditions than fruiting bodies are, which might therefore need to be adapted to colder
temperatures by increasing the proportion of unsaturated fatty acids in fruiting body tissues.
At the same time, due to their large surfaces and continuous evaporation, fruiting bodies
cool themselves (on average ~2.5C colder than their environments(Dressaire et al. 2016)),
which may explain why genetic encoding of higher linoleic acid proportions is beneficial for
their development. It should be noted that linoleic acid production is also required for the
synthesis of several oxylipin species(Combet et al. 2006), which might be involved in cell-to-
cell communication.

4.3.6. Other potentially membrane lipid related genes

We detected several CDE orthogroups which are not parts of the above-mentioned
pathways, yet might be related to lipid metabolism, membrane organization or assembly. In
many cases the functions of these orthogroups is not clear, but their widespread
developmental expression suggests they might be interesting players in fruiting body
development. Therefore, we discuss them below, with details on potential functions where
available.
Glycerophosphodiester phosphodiesterases - A CDE orthogroup of
glycerophosphodiester phosphodiesterases (represented by *C. cinerea* 20997)(Table 9) was
developmentally expressed in 11 species (FC>4), which makes it one of the most widely
developmentally expressed families. However, it is hard to make functional inferences as
these proteins show only limited similarity to experimentally characterized fungal proteins
(highest to Pgc1 of *S. cerevisiae*). *S. cerevisiae* Pgc1 (Phosphatidylglycerol phospholipase
C) negatively regulates phosphatidylglycerol degradation, leading to increased
concentrations in deletion mutants(Šimočková et al. 2008). The glycerophosphodiester
phosphodiesterase family (IPR030395), is represented by three copies in most examined
Agaricomycetes (some species have 4), of which only one is developmentally expressed in
all species. These form a CDE orthogroup in which most genes are FB-init (*C. aegerita, A.*
ostoyae, A. ampla L. bicolor, L. tigrinus, P. ostreatus, S. commune), whereas expression had

different dynamics in *C. cinerea, L. edodes, M. kentingensis, Ph. chrysosporium*). These
expression patterns suggest that the role of this orthogroup, although unknown at the
moment, may be key to the initiation of fruiting body formation.
We noticed that the detected glycerophosphodiester phosphodiesterases showed
strongly correlated expression with members of another CDE orthogroup (represented by *C.*
cinerea 380149, Fig. 7) in six species (data not shown). This contains putative orthologs of
S. cerevisiae Cpt1, a choline/ethanolamine phosphotransferase that catalyzes the last step
(formation of phosphatidylcholine) of the CPD-choline (Kennedy) pathway in *S. cerevisiae*.
The strongly correlated expression in the case of six species (Pearson r >0.8) suggests that
the two genes are linked and possibly form a linear pathway. However, their role in the
context of fruiting body formation requires further research.
Marvel protein family - We also detected two CDE orthogroups (represented by *C.*
cinerea 444998, 462849) (Table 9) of Marvel proteins showing low similarity to *A. nidulans*
mrvA, *C. albicans* Mrv2 and *S. cerevisiae* Fhn1. The gene family seems to be generally
poorly characterized. In *S. cerevisiae*, it may be involved in nonclassical protein secretion,
might act as a sensor of sphingolipids or be related to membrane organization(M et al.
2010). In metazoans, marvel proteins are related to sterol-rich membrane region
positioning(Sánchez-Pulido et al. 2002). Members of the first orthogroup (*C. cinerea* protein
ID: 444998) were FB-init in several species *(A. ostoyae, M. kentingensis, P. ostreatus, S.*
commune, L. edodes), suggesting that they fulfill key functions in fruiting body initiation. The
second orthogroup (*C. cinerea* protein ID: 462849) showed less characteristic expression
patterns, with upregulation mostly late in development in most species.
Sphingolipid C9-methyltransferases - We found two CDE orthogroups of
sphingolipid C9-methyltransferases (*C. cinerea* protein ID: 395241, 365330)(Table 9) that
were FB-init in nearly all species. One of the orthogroups (*C. cinerea* protein ID: 395241)
contains orthologs of Mts1 of *C. albicans,* which is involved in glucosylceramide
biosynthesis*.* Glucosylceramides are glycosphingolipids that form structural components of
the fungal cell membrane, with diverse roles (cell division, spore germination, pathogenicity,
lipid rafts(Poeta et al. 2014)), although little mechanistic detail to date.
PAP2 enzyme family - A CDE orthogroup of diacylglycerol pyrophosphate
phosphatases (*C. cinerea* protein ID: 380219)(Table 9) was FB-init in most species in which
the gene was present. The encoded proteins show high similarity (but not 1-to-1 orthology)
to *A. nidulans* and *C. albicans* membrane phospholipid metabolism proteins AN1671 and
Dpp3, respectively. The latter is required also for farnesol biosynthesis(Nickerson et al.
2006). These genes belong to the broader *PAP2* family, which produce diacylglycerol and
are key in the synthesis of phospholipids, triacylglycerol and the generation of signaling
molecules with a lipid nature. The upregulation of these genes in fruiting bodies might be
related to increased membrane material production associated with cell proliferation during
early development.
Caleosins - Caleosins are well-characterized in plants and indirect data indicate they
may be involved in oxylipin signaling, lipid metabolism, reproduction(Rahman et al. 2018),
though exact functional knowledge seems to be missing. In fungi, caleosins are involved in
sporulation, pathogenicity and lipid storage(Y et al. 2015). A previous study found that the
caleosin family is present in ~30% of fungal genomes(Rahman et al. 2018), whereas we
found that it is present in most of the examined Agaricomycetes, mostly as a single-copy
gene. Members of a detected caleosin CDE orthogroup (*C. cinerea* protein ID:
376602)(Table 9) were FB-init in most species and consistently had peak expression late in

development, which is compatible with presumed roles in fruiting body initiation and
sporulation, respectively.

4.4. Storage carbohydrate metabolism

Fungal colonies accumulate and mobilize various carbohydrates as reserve material and for
energy storage. Previous research has revealed that there are very specific and tightly
regulated accumulation and mobilization patterns of glycogen, trehalose and mannitol within
fungal fruiting bodies(U et al. 2000). Storage carbohydrate metabolism and distributions in
the mycelium and fruiting bodies have been studied in a number of economically important
mushroom-forming species, including *F. velutipes*(Kitamoto et al. 2001)*, Pleurotus spp.*(Han
et al. 2003; Zhou et al. 2016a; Zhu et al. 2019)*, Pisolithus microcarpus*(AN and MD 2010)*, A.*
bisporus(WELLS et al. 1987; TM et al. 2008; Patyshakuliyeva et al. 2013)*, L. bicolor*(Deveau
et al. 2008) and described to a high precision/resolution in *C. cinerea* in a series of papers
by Moore et al(Kuhad et al. 1987; Ji and Moore 1993; Moore 2013), including links with
cAMP levels (summarized by (U et al. 2000)). Briefly, these studies provided evidence on
tightly regulated glycogen, trehalose and mannitol levels in mycelia and various tissues and
developmental stages of fruiting bodies. They also revealed that the identity and quantities of
major sugars in fruiting bodies differ from species to species; trehalose dominates in *L.*
bicolor(Deveau et al. 2008) and *C. cinerea*(Kues 2000), mannitol in *A. bisporus*(Morton et al.
1985) and *L. edodes*(Tan and Moore), and both present in similar quantities in *P.*
ostreatus(Hong Jai-Sik;Kim 1988; Zhou et al. 2016a).
More recently, genome sequences allowed the reconstruction of carbohydrate
metabolism pathways based on gene orthology information to better studied model systems
such as *S. cerevisiae*. Deveau et al found that most of the genes in known fungal pathways
related to glycogen, trehalose and mannitol metabolism were found in the genome of the
ectomycorrhizal *L. bicolor*(Deveau et al. 2008) and similar patterns can be expected in these
very conserved pathways upon examination of other Agaricomycetes genomes.
Transcriptomic studies further allowed previous biochemical observations to be interpreted in
the light of gene expression patterns, although this has not yet been systematically
performed in any species. In the following sections we review information on the metabolism
of glycogen, trehalose and mannitol gleaned from transcriptomic studies of mushroom
development. Many of these studies highlighted that shared patterns of gene expression
related to storage carbohydrate metabolism exist among mushroom-forming species. For
example, Merenyi et al showed that dynamic glycogen accumulation/breakdown is shared
between *C. neoformans* basidium development and complex fruiting body morphogenesis,
two extremes of Basidiomycota development along the complexity gradient(Merényi et al.
2021). On the other hand, trehalose concentrations were found to react to heat shock in
fruiting bodies of *Flammulina filiformis*(Liu et al. 2016). Our discussion hereafter is based on
well-described pathways of storage carbohydrate metabolism in yeast(François and Parrou
2001; JM et al. 2012).
Glycogen metabolism - Glycogen production and mobilization takes place in various
tissues and developmental stages and follows tight regulation that may differ from species to
species (reviewed in (Ji and Moore 1993; Kues 2000; Xie et al. 2020), see also (Ceccaroli et
al. 2011) for Ascomycota). In the Agaricomycetes, this is best known in *C. cinerea* and *A.*
bisporus, whereas mechanistically glycogen metabolism is best known in *S.*
cerevisiae(François and Parrou 2001), which can be used to reconstruct corresponding
pathways in mushroom-forming fungi. Glycogen synthesis encompasses nucleation by

glycogenins and elongation and branching by glycogen synthases. Glycogenins (orthologs of
S. cerevisiae Glg1) are upregulated in young fruiting body gills and/or mature fruiting body
gills of most compared species (Table 10), respectively. This likely corresponds to intense
glycogen synthesis during basidiospore maturation(Kües 2000). Consistent with this,
orthologs of yeast Gac1, a phosphatase that positively regulates the activity of glycogen
synthases, is upregulated in gills and mature fruiting bodies of multiple species. Gac1 in *S.*
cerevisiae regulates glycogen synthases together with Glc7. The latter has orthologs in
mushroom-forming fungi, but these are not developmentally regulated. In *S. cerevisiae*, a
number of other CBM21 domain proteins (Pig1, Pig2) also participate in the regulation of
glycogen synthase activity, however, these have no clear orthologs in mushroom-forming
fungi. On the other hand, the glycogen branching enzyme Glc3 orthologs (represented by *C.*
cinerea 365699) show more diverse expression peaks, and orthologs are upregulated mostly
in stipe tissues (in *C. cinerea, L. bicolor, P. ostreatus* and *A. ostoyae*) (Fig. 9, Supplementary
Fig. 6), indicating that linear glycogen chain production is transcriptionally decoupled from
the addition of branched side chains to the glycogen backbone.
Genes that encode proteins upstream of glycogen synthesis, for example orthologs
of the uridinephosphoglucose pyrophosphorylase Ugp1 (*C. cinerea* protein ID: 359312) and
the phosphoglucomutase PGM1 (*C. cinerea* protein ID: 497782) of *S. cerevisiae*, are
developmentally regulated in multiple species (Table 10). They show some overlap in
expression patterns, in particular, a tendency for upregulation late in development, but their
pattern of expression is quite unique and show lower expression fold changes.
It should be noted that the glycogen synthase kinase 3 (GSK-3)(Chang et al. 2019a)
gene in *C. cinerea* (*C. cinerea* protein ID: 362899), despite its name, it functionally not
relevant to glycogen metabolism, it is an orthologue of *S. cerevisiae* Rim11, a pleiotropic
kinase involved in cell cycle regulation.
Glycogen mobilization happens through three primary genes in *S.*
cerevisiae(François and Parrou 2001) - all these have orthologs in mushroom-forming fungi
(Table 10). Orthologs of the yeast glycogen phosphorylase Gph1 (represented by *C. cinerea*
495987) mostly show high expression in vegetative mycelia or early development, whereas
those of the glycogen debranching enzyme Gdb1 are upregulated also in mature fruiting
bodies in all species (Fig. 9). This probably indicates differential usage of phosphorolysis
and debranching activities in different stages of development. Sga1 is a sporulation specific
glucoamylase in *S. cerevisiae*. In our species orthologs (represented by *C. cinerea* 473268)
of this gene do not show expression peaks in gills, the tissues in which spore development
takes place. Rather, in *C. cinerea*, this gene is upregulated in young fruiting body cap
tissues. How this upregulation corresponds to patterns of glycogen metabolism remains to
be understood. Large-scale postmeiotic utilization of glycogen accumulated in subhymenial
tissues was reported in *C. cinerea*(Kües 2000), however, subhymenial tissues were part of
gill samples in previous RNA-Seq experiments, which excludes the possibility that Sga1
orthologs were involved in that process. In other species, expression patterns were different,
ranging from upregulation in mature fruiting bodies in *A. ampla*, to peak expression in
vegetative mycelium (*P. ostreatus* and *M. kentingensis*). We also detected a CDE
orthogroup of alpha-amylases (represented by *C. cinerea* 495249), which find *A. nidulans*
AmyA as best hit and are probably involved in glycogen catabolism. In *C. cinerea*, this gene
showed upregulation in primordia and peak expression in young fruiting body caps, similar to
other genes related to glycogen mobilization.
In addition to known members of the glycogen catabolism pathways, we identified
two CBM21 domain containing CDE orthogroups that were developmentally regulated in

11/7 and 11/6 species (*C. cinerea* protein ID: 544130 and 455149) (Table 10). In most
species, this gene had the highest expression in vegetative mycelia, but showed significant
expression dynamics also during fruiting body morphogenesis. This seems to be a
Basidiomycota-specific orthogroup, with no significantly similar proteins in model
Ascomycota. The CBM21 domain mostly occurs in starch-binding proteins (glucoamylase
and α-amylase(Ashikari et al. 1986)) whereas in this orthogroup, the CBM21 domain occurs
alone, providing no specific clues to function. Therefore, our placement of this CDE
orthogroup under this chapter should be viewed tentatively until functional studies prove or
disprove the role of the family in glycogen metabolism.
Beyond these gene families, glycogen utilization patterns in fruiting bodies raise a
number of exciting research avenues, such as the mode of transport within the fruiting body
and the identity of transporters involved in that process.
Trehalose metabolism - Out of the at least five pathways known to be involved in trehalose
synthesis, two have been found in Agaricomycetes(Deveau et al. 2008). The first, canonical
pathway comprises the action of a protein complex formed by four proteins in *S. cerevisiae*
(Tps1 - *C. cinerea* 478544, Tps2 - 473004, Tsl1 and Tps3 - no ortholog(François and Parrou
2001)). Of these, Tsl1 (*C. cinerea* protein ID: 503099) and Tps3 are paralogs that arose from
a recent whole genome duplication and thus there is only one corresponding gene in
filamentous fungi. The three genes showed widespread developmentally regulated
expression in examined Agaricomycetes (Table 10), with upregulation mostly late in
development with notable co-regulation between the three genes, as expected for members
of a protein complex (Fig. 9).
A second, noncanonical pathway involving trehalose phosphorylase (TP) exists in
fungi and was detected in multiple mushroom-forming species(K et al. 1998; Eis and
Nidetzky 1999; Wannet et al. 1999; Deveau et al. 2008; Thammahong et al. 2017).
Trehalose phosphorylase gene formed a CDE orthogroup (*C. cinerea* protein ID: 540817) in
the Agaricomycetes we examined(Merényi et al. 2021) and show upregulation in stipes and
in late stages of development in most species (Table 10), although deviations from this
pattern exist (e.g in *C. cinerea*). For example, the gene showed peak expression in
primordium stages of *A. ampla, S. commune* and *M. kentingensis*(Almási et al. 2019;
Krizsán et al. 2019; Ke et al. 2020b).
For **catabolizing trehalose**, fungi exhibit two types of trehalose hydrolyzing
activities, one termed acid trehalase and the other called neutral trehalase(François and
Parrou 2001). Agaricomycetes species possess clear orthologs of neutral trehalases
orthologous to Nth1 of *S. cerevisiae*. On the other hand, and in contrast to Deveau et al
(Deveau et al. 2008), we did not find proteins in the genomes of Agaricomycetes showing
significant sequence similarity (based on BLAST) to acid trehalase. A single neutral
trehalase gene was found in Agaricomycetes, this formed a CDE orthogroup at FC>2 (*C.
cinerea* protein ID: 437212, Table 10) and showed varied expression patterns in the species
examined.
Orthologs of previously described fungal trehalose transporters were not detectable
in Agaricomycetes genomes, based on sequence similarity, in agreement with a previous
study on *L. bicolor*(Deveau et al. 2008). However, it should be noted that fungal trehalose
transporters are poorly known and current knowledge likely misses a large proportion of
genes with this activity(Thammahong et al. 2017).
Mannitol metabolism - Mannitol accumulates in high concentrations in fruiting
bodies of certain species, possibly as an osmolyte assisting fruiting body
growth(Chakraborty et al. 2004; Zhou et al. 2016b) (see also opposing evidence above). A

NADP-dependent mannitol 2-dehydrogenase was reported from *A. bisporus*(JM and H 1998;
S et al. 2001). In agreement, in *L. bicolor* Deveau et al reported a single mannitol
2-dehydrogenase-dependent pathway of mannitol biosynthesis(Deveau et al. 2008), which
works via the direct reduction of fructose. Using the *A. bisporus* NADP-dependent mannitol
2-dehydrogenase gene(JM and H 1998) as a query, we identified an orthogroup
(represented by *C. cinerea* 429272) of short-chain dehydrogenase/reductase genes that
contained only *A. bisporus, L. bicolor, C. cinerea,* and *P. ostreatus*, but was missing from
other species (formed multiple orthogroups probably due to duplications in *A. bisporus*).
However, members of this orthogroup were developmentally regulated in all species (Table
10); they were upregulated at the initiation of fruiting and showed expression peaks in gills of
C. cinerea and *P. ostreatus*, young and mature caps of *L. bicolor*, as well as a marked
upregulation in the stipe tissues of *A. bisporus*. These observation agree with mannitol
metabolism being enhanced in fruiting bodies of *L. bicolor* (Deveau et al. 2008).
In summary, the genomes of Agaricomycetes contain clear orthologs of
experimentally characterized members of glycogen, trehalose and mannitol metabolism
pathways. Many of the genes in these pathways are developmentally regulated in fruiting
bodies, often showing conserved expression peaks in multiple species, reflecting a
conservation of storage carbohydrate mobilization strategies within fruiting bodies.
Interestingly, in *L. bicolor* trehalose and glycogen metabolism was upregulated in *both*
fruiting bodies and mycorrhizae, providing links between fruiting bodies and ECM(Deveau et
al. 2008; Martin et al. 2008).
1523

1524 4.5. Transcriptional regulators

1525 4.5.1. Transcription factors

Transcription factors are proteins that modulate gene expression by binding to specific DNA
sequences and subsequently activating or inhibiting transcription. Transcription factors play
a major role in the regulation of development across the tree of life, and mushroom-forming
fungi are no exception. They are involved in many stages of development, ranging from the
establishment of a dikaryon to various aspects of mushroom development.
The best studied are the homeodomain transcription factors of the *MatA* mating-type
locus that are essential for the establishment of a fertile dikaryon, which is required for
fruiting body formation in heterothallic basidiomycetes(Raudaskoski and Kothe 2010; Vonk
and Ohm 2018). During mating the homeodomain transcription factors expressed from
compatible *MatA* loci dimerize and activate dikaryon-specific transcriptional pathways,
including the formation of clamp connections. The mechanism of mate recognition is very
similar to that found in the yeast *S. cerevisiae*(Haber 2012) and is present across the fungal
kingdom, although many ascomycetes do not use homeodomain transcription factors for
mating(Houbraken and Dyer 2015). In some Agaricales, like *C. cinerea* and *S. commune*,
the *MatA* locus has undergone expansion, improving the chances of finding a compatible
mate(Freihorst et al. 2016).
Several transcription factors have been identified that act downstream of the *MatA*
homeodomain transcription factors. Deletion of the transcription factors *hom2, tea1* and *wc2*
in *S. commune* resulted in strains that did not show dikaryotic development beyond clamp
formation(Ohm et al. 2011, 2013; Pelkmans et al. 2017a). Generally, the resulting colonies
remained symmetrical and showed no sign of mushroom development. Hom2 is highly

conserved in mushroom-forming fungi and features four putative phosphorylation sites.
Disruption of these phosphorylation sites leads to a constitutively active Hom2 and the
inhibition of vegetative growth and monokaryotic fruiting, indicating that Hom2 is post-
translationally regulated(Pelkmans et al. 2017a). WC2 is part of the blue light sensing white
collar complex first identified in *N. crassa*. In *N. crassa*, WC1 and WC2 interact through PAS
domains and WC1 contains a LOV domain that binds a FAD chromophore when exposed to
blue light. This results in conformation changes in the white collar complex and activates
transcription factor domains on both WC1 and WC2(Corrochano 2019). Orthologs of WC1
(dst1 in *C. cinerea*) and WC2 (Cc.wc2 in *C. cinerea*) in *S. commune* and *C. cinerea* fulfil a
similar role and deletion (*S. commune wc1* and *wc2*, *C. cinerea wc2*) or mutation (*C. cinerea*
wc1) of either *wc1* or *wc2* prevents the formation of mature fruiting bodies and is called a
blind phenotype(Terashima et al. 2005; Kuratani et al. 2010; Nakazawa et al. 2011; Ohm et
al. 2013). In *S. commune* this results in a phenotype where no dikaryotic development is
observed and in increased sensitivity to intense light. However, in *C. cinerea* a blind
phenotype leads to the so called 'dark stipe' phenotype(Kamada et al. 2010). In this
phenotype, the primordial shaft is elongated, but pileus and stipe development cannot be
completed. In *S. commune*, WC1 does not contain a DNA-binding domain, unlike WC1 in *N.*
crassa, making transcriptional activity dependent on WC2(Ohm et al. 2010). Orthologs of
white collar complex members with the same function have been reported in *L. edodes*(Sano
et al. 2009) and *P. ostreatus*(Qi et al. 2020). Furthermore, developmental expression of wc
members was reported for *F. filiformis*(Liu et al. 2020b). Mutants of the gene tea1, which is
downregulated in *Δwc1, Δwc2, Δhom2, Δbri1* and *Δfst4* dikaryons, also did not form
mushrooms(Pelkmans et al. 2017a). It should be noted that in *tea1* deletion mutants,
dikaryotic colonies sporadically produced clusters of mushrooms, indicating that Tea1 is not
essential for fruiting body formation and may act as a "switch" that can be bypassed. The
ARID/BRIGHT DNA binding domain transcription factor Bri1 was initially also thought to play
a major role in fruiting development(Ohm et al. 2011). However, later characterization of *bri1*
deletion strains revealed that inactivation leads to significant growth defects that result in a
delay in fruiting(Pelkmans et al. 2017a). The fungal specific transcription factor *fst4* has a
very similar phenotype where no mushrooms are formed, but deletion strains of this
transcription factor do form asymmetrical colonies after mating, which is the first step in
mushroom development(Kües and Liu 2000; Ohm et al. 2010). In *P. ostreatus*, mutations in
gat1 resulted in a strain that did not fruit(Nakazawa et al. 2019). The missense mutation
inside the zinc-finger binding domain resulted in a dominant gain of function mutation, which
suggests that the mutated transcription factor has an altered DNA-binding specificity. This is
in line with disruption of the ortholog *gat1* in *S. commune*, where the phenotype was not
dominant and resulted in the formation of more, but smaller mushrooms(Ohm et al. 2011). A
general overview of Zn(II)2Cys6 zinc cluster transcription factors, including expression
patterns in *P. ostreatus* fruiting bodies, was published by Hou et al(Hou et al. 2020). In
Ganoderma lucidum, silencing of the c_2h_2-type transcription factor *pacC* also resulted in the
absence of fruiting bodies, as well as altered mycelial growth(Wu et al. 2016). Interestingly,
PacC is conserved across the fungal kingdom and activates genes under alkaline conditions,
and inhibits genes under acidic conditions(Peñalva et al. 2008). Therefore, it is unclear if the
absence of fruiting can be attributed to fruiting-specific transcription or general homeostasis.
In *L. edodes*, PriB was described as a primordium-upregulated transcription factor(Endo et
al. 1994). Finally, silencing of *hada-1* resulted in delayed fruiting, as well as multiple growth
defects, in *Hypsizygus marmoreus*(Zhang et al. 2021b).

1594   Downstream in development, *S. commune c2h2, fst1* and *zfc7* gene disruption
1595 resulted in arrested fruiting body formation, where primordia were formed that did not
1596 develop further into mature fruiting bodies(Ohm et al. 2011; Vonk and Ohm 2021).
1597 Interestingly, overexpression of *c2h2* could increase the rate of fruiting in *A. bisporus*,
1598 indicating that *c2h2* expression is a limiting step in fruiting body development(Pelkmans et
1599 al. 2016). The transcription factors Fst1 and Zfc7 were identified due to enrichment of
1600 histone 3 lysine 4 (H3K4) dimethylation in their genes during fruiting body development after
1601 4 days, which indicates increased genetic accessibility and transcription during this
1602 developmental stage(Vonk and Ohm 2021). Further downstream, the *C. cinerea* high-
1603 mobility group transcription factor Exp1 regulates pileus expansion and autolysis to complete
1604 fruiting(Muraguchi et al. 2008b). Notably, this transcription factor is conserved and
1605 developmentally regulated in all studied mushroom-forming fungi, except for *A. ampla, Pt.*
1606 *gracilis* and *S. commune*, whose fruiting bodies do not feature a pileus.
1607   Instead of promoting fruiting body development, several transcription factors are
1608 involved in promoting vegetative growth or repressing fruiting. These include *F. velutipes lfc1*
1609 and *hmg1*, where silencing increased the rate of fruiting(Wu et al. 2020a; Meng et al. 2021).
1610 Additional transcription factors are *P. ostreatus Pofst3* and *S. commune fst3, gat1* and
1611 *hom1*(Ohm et al. 2011; Qi et al. 2019). These genes are differentially regulated in the
1612 majority of mushroom forming species, and conserved in all of them. When the genes are
1613 deleted (*S. commune*) or silenced (*P. ostreatus)* it results in the formation of more, smaller
1614 mushrooms. Finally, premature stop codons in *pcc1* in *C. cinerea* (natural variation,
1615 truncated from 561 to 210 amino acids, HMG-box not disrupted) and *P. ostreatus*
1616 (CRISPR/Cas9-assisted gene mutagenesis, truncated from 471 to 58 amino acids, HMG-
1617 box disrupted) resulted in monokaryotic fruiting and the formation of pseudoclamps(Murata
1618 et al. 1998; Boontawon et al. 2021). In *C. cinerea* these pseudoclamps were formed during
1619 monokaryotic growth, indicating a relation between the *MatA* homeodomain transcription
1620 factors and Pcc1. In *P. ostreatus* however, dikaryotic strains formed pseudoclamps,
1621 suggesting a role in clamp formation.
1622   Many transcription factors are conserved across fungi or specifically in mushroom-
1623 forming fungi. Both *wc2* and *pacC* are conserved in both ascomycetes and basidiomycetes
1624 and except for *hmg1 (F. velutipes), lfc1 (F. velutipes), pdd1 (F. velutipes), pcc1 (C. cinerea*
1625 and *P. ostreatus)* and *zfc7 (S. commune)* all characterized transcription factors are at least
1626 conserved in mushroom-forming fungi. *Cag1* (*C. cinerea* protein ID: 466216) and *Cc.Tup1*
1627 (*C. cinerea* protein ID: 369655) of *C. cinerea* are further examples of highly conserved
1628 transcription factors(Masuda et al. 2016). They are homologs of the highly conserved
1629 general transcriptional co-repressor Tup1 of *S. cerevisiae*, which can form repressive
1630 complexes with a wide range of sequence-specific transcription factors. Deletion of Cag1
1631 resulted in a cap-growthless phenotype, which, upon closer inspection turned out to be
1632 caused by developmental defects of gills(Masuda et al. 2016). However, the expression
1633 patterns are generally not conserved during fruiting body development(Almási et al. 2019;
1634 Krizsán et al. 2019). For example, the developmental stage with the highest expression level
1635 of the ortholog can vary greatly between species. This can indicate either that transcriptional
1636 regulation of fruiting body development is poorly conserved between species, or that
1637 transcription factors are regulated post-translationally. The *MatA* mating type genes are an
1638 example of post-translational regulation, where the transcription factors are constitutively
1639 expressed, but only become activated when heterodimers are formed between compatible
1640 homeodomain transcription factors. Similarly, in *N. crassa* WC2 is activated by structural
1641 changes in its interaction partner WC1 when blue light is detected. Furthermore, Hom2 in *S.*

commune is constitutively expressed, but post-translationally phosphorylated to become
inactive.
While many expression patterns of transcription factors are not conserved in
mushroom-forming fungi, there are several interesting and conserved transcription factors.
We identified 51 orthogroups of TFs that were developmentally regulated in a significant
number of species (Table 11). Among the functionally characterized transcription factors,
orthologs of dst1 (=WC1, *C. cinerea* protein ID: 369621) and Cc.wc2 (*C. cinerea* protein ID:
8610) of *C. cinerea* formed CDE orthogroups developmentally regulated in 10/4 and 10/6
(FC>2/4) species, respectively. Orthologs of dst1 and Cc.wc2 showed varied expression
profiles in the examined species. An orthogroup of homeodomain transcription factors (*C.*
cinerea protein ID: 497620) is developmentally regulated during fruiting body formation in all
mushroom forming fungi, except for *Pt. gracilis*, which is known for its simplified fruiting body
structure. The detected CDE orthogroups also included exp1 of *C. cinerea* (represented by
C. cinerea 421729), which were developmentally regulated in 9/5 species (FC>2/4). In
accordance with its reported function in *C. cinerea*(Muraguchi et al. 2008b), it showed peak
expression in young fruiting body caps. However, the *A. ostoyae, L. bicolor* and *M.*
kentingensis orthologs were upregulated in stipe tissues. In *P. ostreatus* we found the
highest expression in cap cuticle. Furthermore, a C_2H_2-type transcription factor (*C. cinerea*
protein ID: 494418) is conserved and expressed in all mushrooms except for *A. ampla*, *Pt.*
gracilis and *S. commune*. It may therefore play a similar role to Exp1 in *C.*
cinerea(Muraguchi et al. 2008b)*,* which is also conserved and developmentally regulated in
most fungi except for *A. ampla, Pt. gracilis* and *S. commune*. While many other transcription
factor orthogroups are conserved in all studied mushroom-forming fungi, their expression
patterns are not strongly conserved. Therefore, proteomics or functional analyses are
required to further elucidate the role of transcription factors in fruiting body formation.

4.5.2. RNA binding proteins

Pumilio family – This family comprises evolutionarily conserved sequence-specific
RNA-binding proteins that bind 3' UTR regions of specific mRNAs. This family has diverse
roles in fungi, including the regulation of meiotic and sexual processes (e.g. Mcp2 of *S.*
pombe)(Wilinski et al. 2017; Liu et al. 2018). Pumilio proteins were found to be active during
fruiting body formation in both the Asco- and Basidiomycota(Krizsán et al. 2019; Merényi et
al. 2020b). With a few exceptions, all Agaricomycetes contain six Pumilio protein encoding
genes. These formed two CDE orthogroups (Table 12), one of which (*C. cinerea* protein ID:
378537) contains orthologs of the *Sch. pombe* meiotic coiled-coil protein Mcp2 and shows
expression dynamics reminiscent of mitotic and meiotic genes (see Chapter 4.1.1), and
another, comprising orthologs of *Sch. pombe* Puf3 and *C. neoformans* Pum1 (*C. cinerea*
protein ID: 354109). Both orthogroups are developmentally regulated in a large number of
species (8/7 and 7/6 at FC2/4, respectively). In *C. neoformans*, pum1 has been associated
with basidium development and morphogenesis, suggesting that its orthologs may have
similar roles in Agaricomycetes too.
RRM family - We identified five very conserved CDE orthogroups of RNA recognition
motif (RRM) family proteins. One of these (*C. cinerea* protein ID: 471238) were annotated as
related to mitosis and meiosis and is discussed under that chapter. The RRM family
comprises proteins that bind diverse RNA types and shows considerable diversity in fungi,
with 68-88 genes in the Agaricomycetes. Overall, relatively few members of the family were
developmentally expressed, however, the orthogroups we detected showed a conserved

expression pattern. One of them (*C. cinerea* protein ID: 357717) was developmentally
regulated in all 12 species (Table 12) and was upregulated in the second half of fruiting body
development (young fruiting body and mature fruiting body in *C. cinerea*), in particular in gills
(*A. ostoyae, P. ostreatus, M. kentingensis, C. cinerea*) where gill tissue was sampled
separately, or caps (*L. bicolor, L. edodes*). It was also upregulated considerably in primordia
relative to vegetative mycelia. A less characteristic, but similar and very conserved late
upregulation was observed also in the second CDE orthogroup (*C. cinerea* protein ID:
356106), which contains reciprocal best hits of *Sch. pombe* Vip1.
1696   The third CDE orthogroup (*C. cinerea* protein ID: 193545, 11/11 species at FC=2/4)
contains *C. neoformans* Csa2. Members of this orthogroup showed strong induction in the
first primordium stage in *C. cinerea, A. ostoyae, C. aegerita, L. bicolor, L. edodes, P.*
ostreatus, S. commune, A. ampla, M. kentingensis. In *C. neoformans* it shows a strong peak
24h after the induction of sexual development, which, combined with evidence for regulation
by pum1, led Liu et al to conclude it is involved in basidium development and sporulation(Liu
et al. 2018). It's an orthologue of *Sch. pombe* Mei2, which is related to meiosis. The other
three basidium-related genes reported by Liu et al (Fad1, Fas1, CNA00260)(Liu et al. 2018)
do not have clear orthologs in *Coprinopsis*.
1705   The CDE orthogroup represented by *C. cinerea* 424800 (Table 12) was consistently
1706   upregulated in gills and sporulating stages (in *A. ampla, S. commune*) in all species,
1707   suggesting roles in hymenium and/or spore formation. In model Ascomycota, proteins in this
1708   orthogroup find Pab1 of both *Sch. pombe* and *S. cerevisiae* as their best Blast hits, which
1709   are related to mRNA export and the control of poly(A) tail length.
1710   Finally, the CDE orthogroup comprising *C. cinerea* 368527 includes best reciprocal
hosts of *S. cerevisiae* Nam8, a meiosis-specific splicing factor. Our expression data are
consistent with a meiosis-specific role: the gene is developmentally regulated in 6/4 species
(FC2/4, see above, Table 1) and shows a peak expression in meiotic tissues in all species
(gills, cap+gills or mature fruiting bodies). Based on these data, we hypothesize that this
gene also participates in meiosis-specific splicing events in the basidia during fruiting body
formation.

4.5.3. RNA interference, Small RNA

RNA interference is a conserved eukaryotic gene-silencing mechanism that acts via base
pairing of small noncoding RNAs of ~20-30 nucleotides to DNA/RNA targets. RNAi is key to
developmental and differentiation processes, genome defense against viruses or
transposons. There is very limited knowledge on the role of RNAi in mushroom-forming fungi
(but see(Lau et al. 2018)), whereas the processes are extensively studied in the Ascomycota
(in particular *N. crassa* and *Sch. pombe*(Dang et al. 2011)), where RNAi has well-
documented roles in defense against viruses, transposons, meiotic silencing of unpaired
DNA, translational repression and even inter-organismal pathogenic interactions(Weiberg et
al. 2013).
1727   A fully functional RNAi machinery exists in the Basidiomycota. Based on orthology to
Neurospora proteins, Lau et al annotated dicer-like proteins as Dcl-1 (*C. cinerea* protein ID:
444647), Dcl-2 (*C. cinerea* protein ID: 366047) and Dcl-3 (*C. cinerea* protein ID:
438955)(Lau et al. 2018), whereas Stajich et al annotated the argonaute family proteins
AGO (*C. cinerea* protein ID: 355234), an AGO-like protein (*C. cinerea* protein ID: 473372)
and Qde-2 (*C. cinerea* protein ID: 363313)(Stajich et al. 2010) (Table 13). We identified 3-16
genes encoding argonaute and argonaute-like proteins in the examined Agaricomycetes.

Furthermore, fungal RNA interference involves RNA-dependent RNA polymerases, which
are required for generating double-stranded RNA or for amplifying sRNA signals(Dang et al.
2011). The genomes of the examined Agaricomycetes contain 5-13 RNA-dependent RNA
polymerases; of these, the orthogroup represented by *C. cinerea* 498671 is most similar to
the RDP1 protein of *S. pombe*, which was shown to be integral to RNA
interference(Carmichael et al. 2004). Other RNA-dependent RNA polymerases (Table 11)
might participate in alternative RNAi pathways or signal amplification(S et al. 2012b).
Interestingly, we found no clear orthologs of SAD-1, a meiotic silencing protein(Borkovich et
al. 2004), in the examined genomes.
RNAi components show some expression dynamics during fruiting body
development. Genes encoding Dcl-2, Dcl-3 and AGO-like proteins were upregulated in *C.*
cinerea primordia relative to vegetative mycelium(Lau et al. 2018). Similarly, AGO-like and
Qde-2 were found developmentally regulated in *C. cinerea* by Krizsan et al(Krizsán et al.
2019), although Dcl-1-3 genes were not. All of the seven RNA-dependent RNA polymerases
were developmentally regulated in the data of Krizsan et al. In our meta-analysis, the above-
listed components did not show significant developmental expression dynamics (Table 13).
The only noteworthy orthogroup (represented by *C. cinerea* 502116), was developmentally
regulated in 9/6 species (FC>2/4) and contains Argonaute-like proteins that share highest
similarity with *A nidulans* RsdA and *Sch. pombe* Ago1, the first of which is involved in RNA-
silencing(TM et al. 2008) whereas the second has a large suite of described functions, from
RNA silencing to chromosome segregation and cell cycle arrest(Carmichael et al. 2004). We
found no conserved expression pattern in this orthogroup across the examined species.
The most direct evidence for the role of microRNA-like RNAs (milRNA) in mushroom
development was reported by Lau et al(Lau et al. 2018), who compared small RNA
transcriptome profiles of vegetative mycelium and primordium stages and detected 22
milRNA in *C. cinerea*, several of which were differentially expressed between vegetative
mycelia and primordia. These results point to the potential roles of milRNA in mushroom
development, although more studies are needed to understand its roles.
RNA interference has been exploited as a tool of genetic manipulation in mushroom-
forming fungi, often to compensate for the lack of reliable reverse genetics tools in these
organisms. Gene knockdown has been demonstrated in *S. commune*(de Jong et al. 2006)*,*
C. cinerea(Namekawa et al. 2005; MA et al. 2006; Heneghan et al. 2007)*, L. bicolor*(M et al.
2009a)*, Ph. chrysosporium*(Matityahu et al. 2008)*, P. ostreatus, A. bisporus*(Costa Ana
S.M.B.;Thomas 2009)*, Grifola frondosa*(Sato et al. 2015) and *L. edodes*(Nakade et al. 2011),
among others, demonstrating the utility of knockdown techniques in cases/species when
gene knockout was not feasible (particularly in the pre-CRISPR times).

4.5.4. Chromatin regulators

Chromatin remodeling is a fundamental and general mode of epigenetic gene expression
regulation. Dynamic chromatin states are generated primarily via chromatin remodeling
complexes and histone modifications in fungi, in response to environmental or internal
signals. Importantly, chromatin remodeling often provides the basis for broad changes in
gene expression, by modulating the accessibility of DNA for transcription, as opposed to
sequence-specific transcription factors, which regulate only specific genes. Epigenetic
changes often associate with developmental processes; these receive increasing attention in
multiple lineages, including fungi. Accordingly, studies addressing the role of chromatin
remodeling in mushroom development, although quite few at the moment, revealed

fundamental roles of chromatin states in fruiting-related gene transcription. For example,
deleting *Cc.snf5* (*C. cinerea* protein ID: 365798, see Table 14), a member of the SWI/SNF
chromatin remodeling complex, leads to the failure to initiate fruiting body development in *C.*
cinerea (in addition to defects in dikaryon formation)(Ando et al. 2013). In *C. cinerea* two
further genes identified in mutant analyses, *ich1*(Muraguchi and Kamada 1998) and *Cc.*
rmt1(Nakazawa et al. 2010), may be related to chromatin regulation. Recent ChIP-Seq
analyses of epigenetic states revealed substantial changes in transcriptional activity (with
H3K4me2 as a proxy) during the transition to fruiting(Vonk and Ohm 2021), whereas histone
acetyltransferase complex members were found in quantitative trait loci in *L. edodes*(Zhang
et al. 2021d). The presence of SET domain methyltransferases in the Basidiomycota is also
worth mentioning(Freitag 2017). Although evidence at the moment is scanty, the few
available studies and extrapolation from other organisms suggests that chromatin
remodeling may be a significant factor in fruiting body formation. Therefore, in the following
chapters we annotate and discuss expression patterns of genes related to chromatin
maintenance and remodeling based on homology to known chromatin remodeling
components in *S. cerevisiae* and *N. crassa.*

4.5.4.1. Histone gene expression correlates with fruiting body initiation and sporulation

Histones - Histones are highly conserved proteins that assemble into octameric
nucleosomes and condense DNA into chromatin in a tightly regulated manner. Shuffling of
histone variants can generate nucleosome variants, which, combined with diverse
posttranslational modifications of histones (acetylation, methylation, phosphorylation) can
generate a finely diversified histone code contributing to fine-tuned regulation of chromatin
states, which, in turn, impact transcriptional patterns of genes(Borkovich et al. 2004).
Histone modifications are often used to make readouts of transcriptional states associated
with development(Vonk and Ohm 2021). Agaricomycetes have 14-40 histone and histone-
variant encoding genes. They show a very similar overall expression pattern in all species
that resembles the expression of meiotic/mitotic and ribosomal genes: upregulation in
primordium stages as well as meiotic tissues (gills or caps+gills) coincident with sporulation
(Fig. 10, Supplementary Fig. 7). Though overall very conserved, there are slight variations to
this general pattern: *A. ampla* shows a peak only in primordia, whereas in *C. cinerea* and *L.*
tigrinus a meiosis/sporulation associated peak is better visible, while the primordium-peak is
modest. We detected four CDE orthogroups of histones and histone variants (Table 14).
These include a basidiomycetes-specific histone H2A component (*C. cinerea* protein ID:
363625), which is developmentally expressed in 8/6 species (FC2/4) and show considerable
upregulation in primordia relative to vegetative mycelium in most species. Other CDE
orthogroups also showed primordium- and gill- upregulation. A straightforward interpretation
of the similar expression patterns of histones and cell proliferation related genes is a mutual
link to DNA repair. H2A histone phosphorylation is associated with DNA repair(JA et al.
2000; Borkovich et al. 2004), providing a potential mechanistic explanation for the observed
expression patterns. Somewhat contradicting this is the fact that relatives of the Mec1 kinase
(*C. cinerea* protein ID: 19106), which phosphorylates H2A.Z in *S. cerevisiae*(JA et al. 2000),
are not developmentally regulated in Agaricomycetes. On the other hand, histone
upregulation in primordia and meiotic tissues may also be interpreted as evidence for
intense transcriptional reprogramming, which has been hypothesized for the mycelium-
fruiting body transition(Krizsán et al. 2019; Vonk and Ohm 2021). Alternatively, intense DNA
replication may require increased supply of histones in proliferating tissues.

Histone chaperone - We detected a histone chaperone CDE orthogroup containing
homologs of *S. cerevisiae* Chz1 (represented by *C. cinerea* 498885)(Table 14), which forms
a single-copy gene family in the Agaricomycetes. Members of this orthogroup are involved in
histone replacement during transcription and showed the same expression pattern as
histones (peaking in primordia and meiotic tissues), so we speculate they are involved in the
same cellular processes during fruiting body initiation and sporulation.
We next inventoried histone acetyltransferases, histone deacetylases, histone
methyltransferases, histone ubiquitylases, histone kinases and histone ADP-ribosylases
based on homology to corresponding *N. crassa* genes described by Borkovich et
al(Borkovich et al. 2004) as well as diagnostic conserved domains.
Histone acetyltransferase (HAT) - We found one HAT gene per species in the
examined Agaricomycetes (represented by *C. cinerea* 420113), in contrast to Ascomycota,
that have multiple histone acetyltransferase genes(Borkovich et al. 2004). This family
showed no developmental expression dynamics in our species. However, we detected
several protein N-acetyltransferases, which resemble HATs, but lack orthology to known
HATs and do not contain diagnostic conserved domains of HATs. They share an identical
domain composition with *Sch. pombe* Ard1, a Gcn5-related N-acetyltransferase (GNAT),
which acetylates diverse proteins, including transcriptional regulators. A CDE orthogroup of
GNATs (*C. cinerea* protein ID: 457726) was developmentally expressed in 8/4 species
(FC2/4, Table 14), however, given the lack of clear histone link we only tentatively discuss
this family here, noting that functional analysis is needed to confirm their functional
annotation.
Histone deacetylases (HDACs) - we distinguished two groups of HDACs, one
related to *S. cerevisiae* Hda1 and the Sirtuin family, which were detected in 6-7 (11 in *A.
ostoyae*) and 4-9 copies in the examined Agaricomycetes, respectively. Neither groups
showed considerable developmental expression dynamics, except for one CDE orthogroup
comprising sirtuins (represented by *C. cinerea* 455597), which was developmentally
expressed in 9/9 species (FC2/4, Table 14). This family is orthologous to Hst4 of *Sch.
pombe* and Hst3 of *S. cerevisiae*, which regulate DNA damage response and repair by
regulating histone H3K56 acetylation(Konada et al. 2018). Consistent with their presumed
role, the expression of this gene family followed closely that of mitosis/meiosis and DNA
repair related genes in all species: they showed a small expression peak in primordia and a
larger one in meiotic tissues coinciding with sporulation (Fig. 10).
Histone ADP-ribosylases are poly(ADP-ribose) polymerases (PARPs) and are
among the least understood factors in epigenetic regulation; they attach (poly-)ADP-ribose
moieties to proteins, including histones and are thought to regulate histone turnover and
DNA damage response. In *N. crassa* the family is represented by a single gene,
parp(Borkovich et al. 2004). The examined Agaricomycetes' genomes harbor 3-9 copies of
PARPs, of which two formed CDE orthogroups (*C. cinerea* protein ID: 459336, 384835),
which were developmentally expressed in 8/4 and 6/5 species (FC2/4, Table 14),
respectively. Their expression patterns did not follow that of histones, suggesting that their
actions are independent of those.
Histone methyltransferases and histone kinases showed no noticeable expression
dynamics in fruiting bodies (Table 14), suggesting that they are equally important in fruiting
bodies and vegetative mycelia.
1871

4.5.4.2. Other chromatin related genes

Ich1 **gene family -** The only methyltransferase that might be connected to chromatin
regulation was the gene underlying the 'Ichijiku' mutant, which was first identified by
Muraguchi and Kamada(Muraguchi and Kamada 1998), as a mutant that produced
malformed, barrel-shaped primordia, in which cap development fails. The corresponding *ich1*
gene encodes a nuclear methyltransferase that contains a winged-helix DNA binding domain
and was reported to be upregulated in primordia(Muraguchi et al. 2015). Our results agree
and extend these observations: Ich1 orthologs form a CDE orthogroup (represented by *C.*
cinerea 545797) that is developmentally expressed in 8/6 species (FC2/4)(Table 14) and
shows a marked upregulation in primordia relative to vegetative mycelium in *C. cinerea, A.*
ostoyae (weakly)*, A. ampla, S. commune, A. bisporus, M. kentingensis* and *P. ostreatus.*
The exact mechanism of how Ich1 regulates tissue formation is not known, however, this
gene represents a very interesting target for detailed studies, especially given its putative
role in chromatin regulation and/or methylation. On a related note, *Cc.rmt1*, an arginine
methyltransferase which was suggested to participate in histone methylation and to be
required for hyphal knot and fruiting body formation(Nakazawa et al. 2010), forms an
orthogroup which is developmentally regulated in nine species (at FC>2) (Table 14). The
exact role of *Cc.rmt1* is not known so its role in histone methylation remains preliminary.
Histone isomerases - We found a CDE orthogroup that was widely developmentally
expressed (represented by *C. cinerea* 493226)(Table 14). Ascomycota orthologs of these
genes are histone proline isomerases in the peptidyl-prolyl cis-trans isomerase family, which
contribute to nucleosome assembly and thus regulate gene expression. This orthogroup
shows upregulation in gills of *C. cinerea, A. ostoyae* and *P. ostreatus* (all species in which
gills were sampled separate) and in cap tissues of *L. edodes, M. kentingensis, L. tigrinus.*
These expression patterns are consistent with a role in sporulation. Genes in this orthogroup
also show upregulation in primordia relative to vegetative mycelia, although this is usually
modest (2-3x, 10x in *P. ostreatus*).
HMGA and AT-hook protein - We detected a CDE orthogroup (*C. cinerea* protein
ID: 473747)(Table 14) that includes primarily stipe-upregulated genes (in *C. cinerea, A.*
ostoyae, L. tigrinus, P. ostreatus, M. kentingensis, but not *L. bicolor*) that shows expression
peaks also in YFB cap of *C. cinerea* and FB cap of *P. ostreatus*. We found no expression
dynamics in Schizophyllaceae species, which lack stipes. The orthogroup belongs to the
High mobility group A family, which comprises AT-rich DNA binding proteins with an
important role in chromatin organization. This family appears to be patchily distributed in the
examined species, but nevertheless seems to be important for fruiting body development.
SRA-YDG domain proteins - We detected a CDE orthogroup (*C. cinerea* protein ID:
511209)(Table 14) of proteins containing an SRA-YDG (IPR003105) domain, which occurs
in histone and chromatin binding proteins and may be associated with the cell cycle.
Proteins with this domain seem to be missing in Ascomycota model species (results not
shown), but show expression patterns similar to those of histones. They are upregulated in
primordia and in meiotic stages (gills, caps with gills) in *C. cinerea, A. ostoyae, P. ostreatus*
and *L. edodes.* In the Schizophyllaceae as well as *P. ostreatus*, the gene family is
upregulated in primordia relative to vegetative mycelium samples. Based on their expression
patterns, we hypothesize they are related to cell proliferation, the cell cycle or chromatin
modification, but available data do not allow us to further narrow down their putative function
among these categories.

4.6. Alternative splicing and spliceosome components

Alternative splicing is a post-transcriptional mechanism for regulating transcript identity and abundances. mRNA variants generated by splicing can increase the diversity of proteins transcribed from a single gene, but can also destine transcripts to the nonsense-mediated decay pathway and thus reduce transcript abundance posttranscriptionally(Krizsán et al. 2019). Transcriptomes allow us to describe alternative splicing complexity of organisms at an unprecedented depth. Several studies looked at alternative splicing in fungi(Zhao et al. 2013; Grützmann et al. 2014; Jin et al. 2017), including Agaricomycetes model systems(Lugones et al. 1999a; Gehrmann et al. 2016; Krizsán et al. 2019). The Basidiomycota and, in particular, the Agaricomycetes displayed a higher percentage of spliced genes than did other groups of fungi(Grützmann et al. 2014; Krizsán et al. 2019). Results of these studies highlighted that, of the several possible splicing events, intron retention (IR) is the most common in fungi while exon-skipping, which most amply contributes to generating protein isoforms, accounts for only 0.5-2.5% of fungal splicing events. Most research on splicing focuses on the latter, however, it appears that exon skipping-rich transcriptomes represent an animal innovation(Grau-Bové et al. 2018). IR events have been considered either as transcriptional noise or as a regulatory mechanism(Middleton et al. 2017). IR can introduce premature stop codons in transcripts or new functional elements for RNA binding proteins. Premature stop codons can activate nonsense-mediated decay, a surveillance mechanism that results in the degradation of the mRNA transcript. Recent results suggest that IR may not only regulate transcript levels, but can be a key driver of posttrancriptional processes(Middleton et al. 2017). There is evidence that in the Basidiomycota IR exerts its effect on transcripts independent of the nonsense-mediated decay pathway(Gonzalez-Hilarion et al. 2016).

Functionally characterized examples of developmentally relevant splicing events are known in the Ascomycota. For example, alternative splicing of the blue light receptor wc-1 gene was observed in *Phaeosphaeria* spp. and it was hypothesized that AS may control various light-dependent reactions(Chiu et al. 2010). In *Neurospora wc-1* transcription is controlled by multiple distinct cis-regulatory regions (termed different promoters), which responded differently (not, weakly or strongly) to light and resulted in alternative protein isoforms(Káldi et al. 2006). These observations, although probably just scratching the surface, implicate AS as an important contributor to the development of fruiting bodies.

Alternative transcript isoforms affect a considerable portion of genes during fruiting body development. IR-generated transcript isoforms with a dynamic expression are abundant in fruiting bodies(Gehrmann et al. 2016; Krizsán et al. 2019), affecting 20-46% of expressed genes, pointing towards potential important regulatory roles. In *S. commune*, 69% of the alternative spliced transcripts had altered predicted functional properties or subcellular localization(Gehrmann et al. 2016). Gehrmann et al reported the largest number of splicing events at the fruiting body initiation stage in *S. commune*, whereas Krizsan et al did not find substantial differences in the number of splicing events across developmental stages.

Splicing can influence transcript levels significantly during fruiting body development. A recent study found 425 genes in *S. commune* for which an alternative transcript replaced the highest expressed transcript in some stage of development. Many such events happened at fruiting body initiation (referred to as the aggregate stage), pinpointing the magnitude of transcriptome reprogramming at this developmental transition. A conceptually similar scenario, genes whose expression do not vary considerably during fruiting body

1964 development, but have splicing isoforms that vary in their expression significantly, was
1965 detected for ~160-1300 genes in six Agaricomycetes (Krizsán et al. 2019).
1966   Taken together, recent studies on Agaricomycetes suggest that splicing, in particular
1967 intron retention, is an important process in fruiting body development, contributing to
1968 posttranscriptional regulation of transcript levels or generates functional diversity in proteins
1969 translated from alternative transcripts. The prevalence of IR in fungal transcriptomes
1970 suggests that the main role of splicing in fungi might be posttranscriptional transcript level
1971 regulation rather than the expansion of the proteome space, although mechanistic or
1972 proteome-level studies are missing at the moment.
1973   We here annotated spliceosome components and splicing-associated proteins in *C.*
1974 *cinerea* based on knowledge in *S. cerevisiae* and *Sch. pombe*(Käufer and Potashkin 2000;
1975 JM et al. 2012)*.* We annotated 50 orthogroups associated with splicing, however, none of
1976 them showed remarkable expression dynamics in fruiting bodies (Supplementary Table 3).
1977 These included the gene described as *Le.cdc5* and *Cc.cdc5* in *L. edodes* and *C. cinerea*,
1978 respectively(Miyazaki et al. 2004), a pre-mRNA splicing factor.
1979

1980 4.7. Signal transduction systems and communication

1981 One of the most important multicellular processes.

1982 4.7.1. Oxylipins

In fungi, oxylipins are responsible for a range of signaling functionalities and act as paracrine
hormone-like substances or as so-called 'infochemicals'(Combet et al. 2006; Holighaus and
Rohlfs 2019). Ascomycota species produce diverse oxylipin compounds (reviewed
in(Gessler et al. 2017)) and changes in their levels influence development. For example,
oxylipins can regulate the yeast-to-hypha transition in dimorphic fungi, as well as immune
evasion(Rennemeier et al. 2011) and communication with the host in pathogens, while a
well-known group of oxylipins called psi (precocious sexual inducer) factors regulates sexual
and asexual development in Ascomycota(Tsitsigiannis and Keller 2007). Basidiomycota
oxylipins are less known. The best-known oxylipin of mushroom-forming fungi is 1-octen-3-
ol, an enantiomeric compound that is mainly produced as (R)-(−)-1-octen-3-ol by widely-
known species(Zawirska-Wojtasiak 2004). On the one hand, it acts as an interkingdom-
signaling molecule. It may attract or repel fungivorous arthropods(Fäldt et al. 1999;
Holighaus and Rohlfs 2019) or derogate insect larval development. In combination with the
equally prominent octan-3-one that may attract specialist fungivores, it has also been shown
to attract predators of such fungivores(Fäldt et al. 1999). Furthermore, 1-octen-3-ol has been
shown to retard plant seed germination and vegetative growth(Hung et al. 2014). On the
other hand, 1-octen-3-ol shows a quorum-sensing like action as it inhibits colony growth(Yin
et al. 2015; Kües et al. 2018), conidiation(Fäldt et al. 1999; Yin et al. 2015), fruiting
initiation(Noble et al. 2009; Eastwood et al. 2013) and germination where spores are
crowded(Chitarra et al. 2004; Combet et al. 2006).
The composition of the oxylipin cocktail emitted by intact mushroom tissue changes
in the course of mushroom development while measurement of the most abundant oxylipins
during that process crucially depends on the sampling method: disruption of fungal tissue
has been shown to trigger the release of oxylipins that are not emitted when fungal material
is intact(Fäldt et al. 1999). The changing specific oxylipin bouquet emitted by a developing
mushroom may not solely reflect adaptation to insect attraction for spore dispersal and

repellence of fungivores. It is also possible that, e.g. octan-3-one and 1-octen-3-ol are
produced by the fruiting body to inhibit on-colony spore germination(Combet et al. 2006) or
colonization by mycoparasites(Berendsen et al. 2013). During fruiting body development,
octan-3-one emission either positively(Combet et al. 2009) or negatively(Zhang et al. 2008;
A et al. 2020; Orban et al. 2021) correlates with the amount of 1-octen-3-ol produced by (not
yet) sporulating tissue samples of certain species.
The biosynthesis pathway of 1-octen-3-ol has interested mycologists for a long
time(Combet et al. 2006; Kües et al. 2018; Orban et al. 2021), but remained unresolved to
date. The process starts by oxygenation of polyunsaturated fatty acids (PUFAs) with linoleic
acid as the predominant PUFA in Asco- and Basidiomycota(Combet et al. 2006; Brodhun
and Feussner 2011). It was suggested that lipoxygenases (LOXs), linoleate diol synthases
(LDSs that combine a dioxygenase – DOX – and an isomerase activity), hydroperoxide
lyases (HPLs) and alcohol dehydrogenases (ADHs) participate in 1-octen-3-ol
biosynthesis(Combet et al. 2006). Still, it is possible that 1-octen-3-ol can be produced by
incomplete beta-oxidation or by multiple pathways(Combet et al. 2006; Brodhun and
Feussner 2011). Orban et al. recently attempted to identify genes involved in oxylipin
production based on correlations between 1-octen-3-ol concentrations and gene expression
patterns(Orban et al. 2021) including a gene of an ene-reductase for octan-3-one formation
from ADH-mediated conversion of 1-octen-3-ol.
Two main enzyme classes have been implicated in fungal oxylipin production, LOXs
and LDSs(Combet et al. 2006; Orban et al. 2021). Both act on linoleic acid as the starting
point, from which various compounds are produced by first an oxidation reaction, then by
further modification(Combet et al. 2006). We found LOXs to be highly patchy in
Agaricomycetes, whereas LDSs (basically the Ppo family of proteins) are widely conserved.
Incomplete beta-oxidation has also been suggested as a possible source of oxylipins(Venter
et al. 1997; Brodhun and Feussner 2011), this hypothesis is compatible with the lack of
certain enzymes in many species.
Lipoxygenases (LOXs) - Out of the examined species, we only found lipoxygenases
in *C. aegerita*, *P. ostreatus*, *L. bicolor* and *L. edodes*, which have 1-5 copies. These did not
form conserved orthogroups and their expression patterns mostly did not vary substantially
during development. Exceptions include the gene previously described as
PoLOX1(Plagemann et al. 2013; Y et al. 2013) (PleosPC15_2_1068582). *PoLOX1* displays
a remarkable upregulation (1880 fold, FPKM 94.000) in fruiting bodies relative to vegetative
mycelium. The gene showed consistently higher expression in stipes compared to caps,
which confirms the previous report on the gene(Y et al. 2013). A lipoxygenase gene
(*CaeLOX4*, gene ID AAE3_04864/Genbank accession no. CAA7262624) of *C. aegerita* also
showed a strong upregulation (~150-fold) during late development (post sporulation stage) in
this species, as noted by Orban et al (Orban et al. 2021). Notably, two other LOXs show
strong phase-specific expression increase in fruiting stage tissue: on the one hand,
CaeLOX5 (AAE3_07753) shows an expression pattern quite congruent with the maturation-
associated increase of the 1-octen-3-ol levels in the headspace of the *C. aegerita* culture.
On the other hand, Cae*LOX1* (AAE3_00896) shows transcriptional upregulation before
fruiting body maturation sets in. Such patterns may imply stage-specific action or (partial)
functional redundancy among these genes that must, prospectively, be tested by functional
characterization studies.
Linoleate diol synthases (LDSs) - The other well-known gene family involved in the
production of oxylipins is the one of linoleate diol synthases, which makes up to PpoA-C
family in *A. nidulans*. The main products of the Ppo family are the so-called psi factors,

which regulate diverse developmental processes from yeast-to-hypha transition and cell
aggregation (reviewed in ref(Tsitsigiannis and Keller 2007)). We identified 1-4 LDS genes in
Agaricomycetes with most species having two copies. Although several genes have
reciprocal best hits among Ascomycota (e.g. *C. cinerea* 423716 to PpoC of *A. nidulans*), a
gene tree analysis showed that these are not true orthologs (results not shown) but likely
resulted from a duplication event that predated the split of Asco- and Basidiomycota.
Therefore, it is likely that Basidiomycota linoleate diol synthases do not have the psi factor
producing activity that *A. nidulans* PpoA-C genes and their homologs have(Tsitsigiannis and
Keller 2007).

Linoleate diol synthases grouped into three conserved orthogroups (Table 15), of
which two (*C. cinerea* protein ID: 398037, 423716) were developmentally expressed in a
considerable number of species. One of these (*C. cinerea* protein ID: 423716) includes *U.
maydis* Ssp1 (Ustma2_2_11710), which was found to be specifically expressed in
teliospores(Huber et al. 2002). Genes within these orthogroups seem to show diverse
expression patterns, without noticeable conservation of expression across species. In the
absence of functional information of these genes in Basidiomycota, it is impossible to
speculate on their mechanistic role in fruiting body formation. They are likely to be involved
in oxylipin production(Orban et al. 2021), but further studies are necessary to elucidate their
functions.

Alcohol dehydrogenases – the alcohol dehydrogenase family participates in a
broad range of cellular processes, including oxylipin biosynthesis(Orban et al. 2021). We
identified a CDE orthogroup containing *C. aegerita* Adh1-1 (represented by *C. cinerea*
363127), which showed partial conservation of expression: in most species it was
upregulated during late development (*C. aegerita, A. ostoyae, L. edodes, L. tigrinus, C.
cinerea*). In addition, in *S. commune, Ph. chrysosporium, A. ostoyae* and *L. edodes* it was
also upregulated >4x in primordia relative to vegetative mycelium. In *M. kentingensis* and *A.
ampla* the gene was not developmentally regulated, whereas *L. bicolor* and *P. ostreatus* did
not have orthologs. Based on a previous report of *C. aegerita* Adh1-1(Orban et al. 2021), it is
possible that this orthogroup of alcohol dehydrogenases participate in oxylipin production.
2086

4.7.2. Pheromone and pheromone receptor pathways

The structure and evolution of mating systems, including HD transcription factors (A locus),
pheromone receptors and pheromone precursor genes (B locus) is among the best known
aspects of Agaricomycete biology (e.g.(Kües 2000, 2015; James 2015) refs). The primary
role of pheromone/receptor systems is regulating dikaryotization, i.e. the compatibility of
monokaryons, nuclear movement and other downstream events. Hyphal fusion is believed to
occur ubiquitously in the Basidiomycota, therefore, pheromone/receptor systems may
regulate post-fusion behavior(Raudaskoski and Kothe 2010; Jones and Bennett 2011).
However, pheromone/receptor systems are involved in the regulation of sexual events too.
For example, mating type loci are involved in fruiting body formation in *C. cinerea*(Kues et al.
1998, 2002). Basidiomycete mating systems, pheromone and pheromone receptor genes
have been reviewed recently in detail (e.g.(Raudaskoski and Kothe 2010; Jones and Bennett
2011; James 2015; Kuees 2015)). We here annotate components of the Agaricomycete
mating systems, with special emphasis on copy number variation across the twelve species
examined in this paper and expression dynamics during fruiting body development.

Pheromone precursor genes - Basidiomycetes produce mating pheromones that
are structurally analogous to *S. cerevisiae* a-factor(Jones and Bennett 2011). Mating
pheromones are short peptides that undergo a multi-step maturation process and regulate
intercellular communication and sexual development in a wide range of fungi(Jones and
Bennett 2011). Copy numbers of pheromone precursor genes vary across the
Agaricomycetes and are often found in the vicinity of corresponding pheromone
GPCRs(Raudaskoski and Kothe 2010). Four pheromone precursor genes were found in *L.*
edodes(Wu et al. 2013)*, P. eryngii*(Kim et al. 2014)*, A. bisporus*(Foulongne-Oriol et al.
2021)*, Moniliopthora roreri*(Díaz-Valderrama and Aime 2016)*,* five in *Ph.*
chrysosporium(James et al. 2011)*,* seven in *Pycnoporus cinnabarinus*(Levasseur et al.
2014)*,* two in *P. touliensis*(Gao et al. 2018) and several genes (up to 20) in *C.*
cinerea(Riquelme et al. 2005) and *S. commune*(Specht 1995). Synthetic pheromone
administration affected PriA (a primordium-expressed gene) expression in *L. edodes*,
indicating the involvement of mating pheromones in fruiting body formation(Ha et al. 2018).
One of the pheromone precursor genes was expressed in glucose-limited conditions
following blue light induction of fruiting in *C. cinerea*(Sakamoto et al. 2018).
In our dataset, we found up to nine pheromone-precursor encoding genes, however,
we did not find any in *A. ostoyae, A. bisporus, M. kentingensis, Ph. chrysosporium* and *L.*
tigrinus. The explanation for this may be that they are simply missing from the genome
annotations. Pheromone precursors are short ~50 amino acid peptides and several genome
annotation protocols omit predicted proteins that are <50 or <100 amino acids long. For this
same reason, it is likely that our inventories of pheromone-precursor genes in other
examined species is incomplete. Nevertheless, we found pheromone precursors in *C.*
cinerea, A. ampla, L. bicolor, L. edodes, S. commune and *P. ostreatus*; these genes were
often developmentally regulated, implying that they may participate in developmental
processes, possibly as signal molecules. While knowledge on the role of pheromones in
mating and dikaryon formation is abundant, whether and how they participate in fruiting body
development requires further research.
Pheromone modification - pheromone precursor proteins are post-translationally
modified during their maturation process. This involves prenylated, carboxymethylation as
well as proteolytic cleavage to remove N-terminal sequence fragments(Jones and Bennett
2011).
Post-translational prenylation (mostly farnesylation) on CAAX motifs is performed by
RAM proteins in *S. cerevisiae*(Caldwell et al. 1995)*.* Based on orthology with *S. cerevisiae*
pheromone prenylating enzymes, we identified two genes as putatively involved in
pheromone prenylation. The first is orthologous to *S. cerevisiae RAM1* (*C. cinerea* protein
ID: 491483) and shows very modest expression dynamics (Table 16), whereas the second is
an ortholog of *RAM2* (*C. cinerea* protein ID: 473005), encoding a general prenylation
enzyme that also modifies other peptides, including proteins related to cell wall glucan
biosynthesis. Following prenylation, the *S. cerevisiae* a-factor is cleaved by Rce1 (ortholog:
C. cinerea 377901) and Ste24 (ortholog: *C. cinerea* 419650). Orthologs of these genes
showed constant expression during fruiting body development (Table 16).
On the other hand, we found evidence for the developmental expression of
carboxymethylating enzymes. Two CDE orthogroups of protein-S-isoprenylcysteine O-
methyltransferases (homologous to *S. cerevisiae* Ste14) were detected among conserved
developmentally expressed genes. Ste14 mediates methylation of the isoprenylated C-
terminal CAAX motif in the mating pheromone a-factor(Romano and Michaelis 2001).
Whether these homologs are related to mating pheromone maturation in mushroom-forming

fungi remains unclear, therefore, we only tentatively classified these as signal transduction-
related genes. Two Ste14 homologs of *C. cinerea* were previously found to be expressed
after blue light exposure(Sakamoto et al. 2018). We detected two CDE orthogroups of Ste14
homologs (Table 16). One of the orthogroups (represented by *C. cinerea* 152667) is present
only in five species and upregulated in primordia of *C. cinerea, A. ostoyae, A. bisporus* and
P. ostreatus but interestingly shows the reverse pattern, high in vegetative mycelium and no
expression in any fruiting body stage in *M. kentingensis*. The other orthogroup (represented
by *C. cinerea* 354499) was upregulated in primordia of *C. cinerea, A. ostoyae, Pt. gracilis*
and *S. commune (*but not in *A. ampla* and *P. ostreatus).* More broadly, members of this gene
family were developmentally regulated in several species, although they were not conserved
across species. On the other hand, 1-to-1 orthologs of *S. cerevisiae* Ste14 (represented by
C. cinerea 153902), show constant expression in all species.
As a final step of a-factor maturation, Ste24 and Axl1/Ste23 proteolytically cleave the
N-terminus in two steps(Jones and Bennett 2011). The corresponding Agaricomycete
orthologs did not show developmentally dynamics expression (*C. cinerea* protein ID: 419650
and 358206, respectively)(Table 16).
Pheromone sensing GPCRs - Seven-transmembrane domain G-protein coupled
receptors have been studied in multiple Agaricomycetes(Kües 2000; Raudaskoski and Kothe
2010; Jones and Bennett 2011; Wirth et al. 2021). Pheromone-sensing GPCRs are located
in the B mating type locus in Basidiomycota, together with pheromone precursor genes.
Pheromone sensing GPCRs transduce signals intracellularly to a mitogen-activated protein
kinase (MAPK) cascade (see next chapter). Homology relationships and copy numbers of
Basidiomycete pheromone-sensing GPCRs vary significantly across species, but all
Basidiomycete pheromone GPCRs are homologous to yeast Ste3, which is involved in
mating (reviewed recently(Raudaskoski and Kothe 2010; Fraser et al. 2014)). For example,
four pheromone-like GPCR genes have recently been annotated in *S. commune*, of which
one was suggested to participate in mating, whereas the others were proposed to have a
role in directing growth in vegetative mycelia(Wirth et al. 2021).
We identified 4-19 pheromone GPCR-encoding genes in the examined species.
These genes in general showed low expression dynamics in fruiting bodies (Table 16),
although several genes were upregulated slightly (FC<4) in stipes in certain species.
A-factor export proteins - We identified a CDE orthogroup (represented by *C.*
cinerea 540181)(Table 16) of plasma membrane ATPases which are othologous to *S.*
cerevisiae Ste6, the protein that exports farnesylated a-factor from the cell. This orthogroup
showed partial conservation of expression across the examined species. In *A. ostoyae, L.*
bicolor, L. edodes and *M. kentingensis* it was upregulated in stipe tissues relative to caps,
whereas in *C. cinerea, S. commune, A. ampla* (FC>4) and *C. aegerita* (though only at FC>2)
the gene was upregulated in primordia relative to vegetative mycelium.
2188

4.7.3. MAPK, TOR and cAMP pathways

MAP kinase pathways - The mitogen-activated protein kinase (MAPK) cascades are
evolutionary conserved intracellular signal transduction pathways, which regulate important
cellular processes(Román et al. 2007). The extracellular signal is sensed by membrane
proteins or receptors, which is relayed by sequential phosphorylation and activation of three
kinases (MAPKKK, MAPKK, MAPK) to transcription factors that induce or repress specific
target genes. MAPK cascades usually consist of further components, like various adaptors,

docking and scaffold proteins which are implicated in the spatial and temporal regulation of
MAP kinase signaling(Frawley and Bayram 2020).
While in Ascomycota MAPK members have been extensively examined
(e.g.(Teichert et al. 2014)), relatively little information is available on the expression and role
of MAPK pathways (except the pheromone, see above) in Agaricomycetes. In *Pleurotus*
tuoliensis pheromone-dependent and starvation-dependent MAPK signaling pathway
members were reported to be up-regulated during fruiting(Fu et al. 2017). Le.MAPK, a MAP
kinase of *L. edodes* (Lenedo1_1058304) was reported to be developmentally regulated and
one of its interacting partners was characterized(Szeto et al. 2007). In this paper, we
annotated Agaricomycetes MAPK signaling pathways based on the four main MAPK
pathways known in *S. cerevisiae*: the peptide pheromone pathway (Fus3), the starvation
responding pathway (Kss1), the high osmolarity glycerol stress pathway (Hog1) and the cell
wall integrity (Slt2/Mpk1) pathway (Table 17). We found relatively few clear orthologs in *C.*
cinerea (Table 17). We performed transitive ortholog and domain-based search to find
further putative *S. cerevisiae* orthologs in *C. cinerea*. MAPK pathway components do not
show dynamic expression patterns during fruiting body development.
TOR pathway - The target of rapamycin (TOR) is an evolutionary conserved
serine/threonine kinase that is a key regulator in multiple cellular events(Tatebe and
Shiozaki 2017). In fungi, the TOR pathway plays role in the response to nutrients (especially
sugar and nitrogen) but also involved in stress responses, such as osmotic and oxidative
stresses. TOR signaling pathways are well-studied in yeasts, but data about their functional
analysis in multicellular agaricomycetes is rare. In *G. lucidum* TOR signaling was reported to
play a role in the regulation of chitin and β-1,3-glucan synthesis and hence of cell wall
thickness in an SLT2-MAPK dependent manner(Chen et al. 2019). Orthologs of *Ph.*
chrysosporium were previously annotated(Nguyen et al. 2020). According to a 5'-SAGE
study of *C. cinerea,* TOR pathway members are constantly expressed during the fruiting
body formation, except Rheb (*C. cinerea* protein ID: 493094) a Ras-like small GTPase,
which is able to prevent FKBP12 inhibition of TOR(Cheng et al. 2013a).
Here we annotated TOR pathway components in mushroom-forming fungi and could
find orthologs for all but three yeast genes reported to participate in the pathway (Table 17).
Consistent with previous reports, TOR pathway components showed modest expression
dynamics during fruiting body development. Nevertheless, based on the broad role of the
pathway in fungal physiology it will be interesting to investigate its connection with fruiting
body morphogenesis in the future.
cAMP pathway - Adenylyl cyclase dependent pathway (cAMP-pathway) has diverse roles in
fungi, including cell growth, metabolism, stress resistance and nutrient-sensing (especially
glucose) or starvation in fungi(D'Souza and Heitman 2001). Accordingly, the cAMP pathway
has been intensely researched in mushroom-forming fungi(D'Souza and Heitman 2001;
Tamaki 2007). Genetic manipulation of several cAMP and connected pathway components
in *S. commune* resulted in altered fruiting characteristics, either repression or complete lack
of fruiting body formation(Yamagishi et al. 2002, 2004; Knabe et al. 2013; Pelkmans 2016).
The cAMP pathway has also been linked to morphogenesis in *F. velutipes:* the deletion of a
Gβ-like protein CPC-2 completely impaired fruiting body formation, but the addition of cAMP
or it analog 8-Bromo-cAMP (also a PKA-activator) into the medium restored fruiting in
Fv.cpc2 knockdown mutants(Wu et al. 2020b). Ras and G-protein ⬚ subunits and their
regulators (e.g. *Thn1*) might also be connected to the cAMP pathway(Yamagishi et al. 2002;
Palmer and Horton 2006; Knabe et al. 2013) and were repeatedly reported to affect
morphogenesis of fruiting bodies (e.g.(Schubert et al. 2006)).

In general, cAMP is a known inducer of fruiting in several basidiomycetes, such as *C.*
macrorhizus(Uno and Ishikawa 1973), *C. cinerea*(Swamy et al. 1985)*, S. commune*(Schwalb
1974) and *Ph. chrysosporium*(Gold and Cheng 1979)*.* In the latter, cAMP was speculated to
induce fruiting by overriding the repressive effect of glucose (i.e. mimicked starvation in the
fungus(Gold and Cheng 1979))*.* More recent studies in *S. commune* reported a repressing
effect of cAMP on fruiting, and suggested this was because cAMP is able to mimic high CO_2
levels(Pelkmans 2016). Genes of the cAMP pathway were annotated in *S. commune* and
the pde2 phosphodiesterase (Schco3_2636760), which can degrade cAMP was functionally
analyzed(Pelkmans 2016). The overexpression of pde2 results in fruiting body development
even at elevated CO_2 levels. These results were interpreted as evidence for CO_2 being
sensed via the cAMP pathway(Pelkmans 2016).
We annotated members of the cAMP pathway with *C. cinerea* as reference (Table
17). Consistent with data in *S. commune*(Pelkmans 2016), we could identify most members
of the cAMP pathway in the examined species. Our expression data revealed that cAMP
pathway members show relatively little expression dynamics in mushroom-forming fungi
(Table 17). Nevertheless, this does not preclude an important role of cAMP signaling in
fruiting body formation.
2261

4.7.4. G-protein coupled receptors

G-protein coupled receptors (GPCRs) comprise the largest receptor family in fungi. GPCRs
are seven transmembrane cell surface receptors, which transduce signals from a wide range
of external stimuli by a G-protein heterotrimer(Xue et al. 2008). Fungal GPCRs originally
divided into six classes based on their structural similarity and homology(Xue et al. 2008),
however, this classification was later expanded with additional classes (reviewed in (Brown
et al. 2018)).
Pheromone sensing GPCRs are among the most widely studied in connection with
fruiting body development. In *Flammulina filiformis* six pheromone receptor-like genes were
characterized whereby four were highly expressed in the fruiting body, while two exhibited
the maximal expression level in the mycelia(Meng et al. 2020). Six pheromone-like GPCRs
were characterized also in *S. commune* (*bar3, bbr2, brl1*, *brl2*, *brl3* and *brl4*); it was reported
that the overexpression of *brl2, brl3* and *brl4* enhanced mating or fruiting body formation in
this species(Wirth et al. 2021). We annotated 7 mating-type pheromone receptors in *C.*
cinerea (Table 16). Most of the pheromone receptors are highly expressed during the
fruiting, but do not have orthologs in other species. In addition, five further GPCR genes
were detected, which, similarly to pheromone-sensing GPCRs mostly did not have orthologs
in other species (Table 18). This may suggest that fungal species have a specialised GPCR
repertoire, that is hard to examine using orthology-based approaches.
While GPCRs are not conserved, we detected three CDE orthogroups that may be
related to GPCR signaling and were developmentally regulated across several species. An
RGS (regulator of G-protein signaling) domain containing protein family (*C. cinerea* protein
ID: 479562) was developmentally regulated in 11/7 species (FC>2/4)(Table 18). RGS
proteins are involved in signal transduction by regulating G-protein activity and rapidly
switching on or off G-protein coupled receptors pathways. We also found a CDE orthogroup
containing putative Ran-binding proteins (*C. cinerea* protein ID: 472370), which are
members of the Ras superfamily that regulates all receptor-mediated transport between the
nucleus and the cytoplasm. Finally, we detected two arrestin superfamily orthogroups (*C.*

cinerea protein ID: 473465, *S. commune:* Schco3_2490312) (Table 18). Arrestins are
involved in signal transduction through the inactivation of G protein-coupled receptors, but
some members of the superfamily have been reported in receptor endocytosis, cell cycle
progression and pH sensing as well also(Herranz et al. 2005; Telzrow et al. 2019).

4.7.5. Other signal transduction related families

Kinases - The majority of CDE orthogroups classified as signal transduction were kinases
(Table 15). We found 13 conserved and developmentally expressed kinase orthogroups,
however, most of these proved difficult to functionally characterize, likely due to the high rate
of evolution in kinases and the lack of orthology to well-known genes in model species. One
exception is the CDE orthogroup (*C. cinerea* protein ID: 451611) that contains orthologs of
S. cerevisiae Rck2, which is a member of the high osmolarity glycerol (HOG) MAPK
pathway. The HOG pathway is involved in responding to oxidative and osmotic stress. In
most species, Rck2 orthologs showed early upregulation, with highest expression values in
primordia, though its functional significance is not known. Rck2 orthologs were differentially
expressed between dikaryons and A and B type monokaryons of *C. cinerea* as well(Stajich
et al. 2010), and it was concluded that RCK2 genes in *C. cinerea* must serve different
functions from that of *S. cerevisiae* orthologs. The remaining kinase orthogroups are
functionally uncharacterized and may be interesting targets for functional studies, their
dynamic expression patterns indicate potentially conserved and important functions in
fruiting body development.
Velvet complex - The velvet complex has been identified as a central regulatory hub in
fungal development and secondary metabolism(Bayram et al. 2008). The complex was
characterized in the Ascomycota, where it consists of VeA, LaeA, VelB as well as VelC and
VosA, which assemble into condition-specific (including light-responsive) complexes. It has
been shown that the velvet domain, an ~200 amino acid fold, binds DNA and resembles the
structure of the metazoan transcription factors NF-κB(Ahmed et al. 2013). Proteins
containing a velvet domain exist outside the Pezizomycotina as well, however, these
proteins did not appear to be orthologous to members of the Velvet complex in the
aspergilli(Ojeda-López et al. 2018); accordingly, how they function and whether they form
complexes also in the Basidiomycota and early-diverging fungi is not known.
The first evidence on a potential role of Velvet proteins in Basidiomycota fruiting body
development was provided by Plaza et al(Plaza et al. 2014b). They detected the
upregulation of a velvet domain containing protein in primordia of *C. cinerea* relative to
vegetative mycelia. Following these trails and combining detailed mechanistic knowledge
from the velvet regulon of *A. nidulans*(Krijgsheld et al. 2013), they could show that some,
albeit a minor proportion, of orthologs of the *A. nidulans* velvet-regulated genes are
upregulated in fruiting bodies in *C. cinerea, L. bicolor* and *S. commune*(Plaza et al. 2014b)
also. This suggested some extent of conservation of function in the Dikarya, although the
lack of orthology between Asco- and Basidiomycota proteins casts some uncertainty on how
these data should be interpreted. Velvet protein-encoding genes were reported to be
developmentally regulated in multiple Agaricomycetes species in the comparative analyses
of Krizsan et al(Krizsán et al. 2019).
Our current analysis indicated that the velvet domain occurs in 5-6 proteins (only 4 in
P. ostreatus) encoded in the genomes of mushroom-forming fungi. Surprisingly, the yeast-
like *C. neoformans* also possess 6 genes with a velvet domain, in contrast to Ascomycota
yeasts, which have lost velvet genes(Nagy et al. 2014). Interestingly, Ascomycota on

average have 4 genes that encode velvet-domain containing proteins, indicating an
expansion in the Basidiomycota. Consistent with results of Ojeda-Lopez et al(Ojeda-López
et al. 2018), we found that Basidiomycota velvet proteins lack clear orthology to Ascomycota
genes. Only VelB and VosA of *A. nidulans* have reciprocal best hits in *C. cinerea* (*C. cinerea*
protein ID: 365674 and 374867, respectively).
In terms of expression, observed patterns vary widely among species. For example,
all six genes are developmentally regulated in *C. cinerea*, but none in *A. ostoyae* (both
FC>4), indicating considerable expression plasticity through evolution. We identified three
orthogroups whose members were developmentally regulated in a notable number of
species (Table 19). Of these, genes in one orthogroup (represented by *C. cinerea* 385561)
were mostly upregulated in stipe tissues in the pileate-stipitate species *C. cinerea, A.*
ostoyae, M. kentingensis, *L. edodes* (but only in mature FBs of *L. bicolor*). The gene was not
developmentally regulated in *L. tigrinus* and *P. ostreatus*, possibly indicating the distant
relationship of these species from ones in which the gene is stipe-upregulated. The second
most interesting CDE orthogroup (*C. cinerea* protein ID: 496341) was developmentally
regulated in 10 species (FC>2), but expression peaks varied from primordia (*A. ampla, S.*
commune, A. ostoyae), to young fruiting body caps (*C. cinerea*) and stipes (*M. kentingensis*).
Based on the above data, velvet protein-encoding genes represent an interesting
target for phylogenetic and functional studies of sexual fruiting bodies in the Agaricomycetes;
we expect their analyses will reveal interesting insights into fruiting body development.
2356
STRIPAK complex - The striatin-interacting phosphatase and kinase, STRIPAK (known as
Far complex in(Kück et al. 2019)), is a widely conserved multi-protein complex that can be
found from yeast to metazoans(Reschka et al. 2018). Its function in metazoans is unknown,
whereas in Ascomycota model systems (e.g. *Sordaria macrospora*) it is involved in hyphal
fusion and fruiting body development. It was first discovered in 2004 in fungi(S and U 2004)
and its function has been increasingly clarified in a series of papers since then (see
in(Reschka et al. 2018; Riquelme et al. 2018)). Mechanistically, the STRIPAK complex
provides links between multiple signaling pathways in eukaryotic cells (reviewed in(Kück et
al. 2019)), although many of the details of its interactions with other proteins are not known.
Its role in fruiting body formation in the Basidiomycota has not yet been investigated.
We queried the *C. cinerea* genome with members of the STRIPAK complex
described from *Sordaria* and *S. cerevisiae* (listed in(Reschka et al. 2018)). *C. cinerea* seems
to have a fully functional STRIPAK complex, with all genes having clear orthologs, except for
Far3/7 which only had weak hits. Further, the *C. cinerea* proteins were members of
conserved single-copy Agaricomycetes orthogroups, which indicated that the complex is
conserved in all the examined Agaricomycetes species. Members of this complex did not
show significant expression dynamics (Table 20). The conservation of the complex in the
Agaricomycetes suggests it might have important functions, however, its role in fruiting body
development remains unclear at the moment and requires further research.

4.7.6. Light sensing systems and light receptors

Fungi adapt developmental processes to light signals in time and space to ensure that
sporulating cells can maximize spore dispersal efficiency(Yu and Fischer 2018). The
photobiology and its connection to developmental events are quite well-known in both the
Asco- and Basidiomycota. In the last decade, the molecular architectures and regulatory
networks of sensing and responding to blue, red and green light have been uncovered in

multiple species(Corrochano 2011; Fuller et al. 2015). In this chapter we discuss light
sensing systems in Agaricomycetes and analyze their expression patterns in fruiting bodies.
We note that the best known such system, the white collar complex has transcription factor
activity and is discussed under chapter 4.5.1.
Cryptochromes - Fungal genomes include DASH-type cryptochromes, photolyase
genes that function as blue light photoreceptors and are involved in DNA repair(Fuller et al.
2015). Cryptochromes are regulated by the white collar complex in *N. crassa*(Froehlich et al.
2010) and regulate fruiting body development in certain Ascomycota (e.g. *Cordyceps*
militaris(F et al. 2017)), but to our best knowledge, no such information is available in the
Basidiomycota. Our analysis suggests that cryptochromes are single copy genes in the
Agaricomycetes, except in *L. edodes*, which has two genes. The CDE orthogroup containing
Agaricomycetes cryptochromes (represented by *C. cinerea* 492962)(Table 21) was
developmentally regulated in 9/7 species (FC>2/4), and showed strong induction in
primordia relative to vegetative mycelium in *A. ostoyae, M. kentingensis, L. edodes, P.*
ostreatus, Ph. chrysosporium and at FC>2 in *C. aegerita* and *C. cinerea*. The gene was
constantly expressed in *A. ampla* and *S. commune* and was missing in *L. bicolor.* Based on
these data, cryptochromes may be relevant to fruiting body formation in the Agaricomycetes.
Phytochromes are red- and far-red receptors that influence sexual and asexual
sporulation in fungi. The first circumscribed fungal phytochrome was the *fphA* gene in *A.*
nidulans, which represses sexual development under red light(A et al. 2005). Since then, the
repression of sexual development has been proven for *N. crassa* too and several details of
the process (e.g. interaction with WCC and velvet factors) have been clarified(A and J 2005;
Wang et al. 2016; Yu et al. 2016). In the Basidiomycota, the few reports of phytochromes
include a systematic mutant library in *C. neoformans*, which circumscribed the gene
encoding *TCO3* (Lee et al. 2016), as well as observations of phytochrome upregulation in
browning mycelium of *L. edodes*(Tang et al. 2013) and in rhizomorphs of *A. ostoyae*(Sipos
et al. 2017b). In *U. maydis* red and blue light receptors have been shown to be required for
the development of fruiting body-like structures(Sánchez-Arreguin et al. 2020).
Phytochromes seem to be single-copy genes in the Agaricomycetes examined here.
They formed a conserved orthogroup (represented by *C. cinerea* 470504, Table 21),
however, genes in this orthogroup did not show significant expression dynamics during
fruiting body formation, except in *A. ostoyae* and *C. cinerea*. In *A. ostoyae* the expression of
the phytochrome gene was ~10x higher in stipes than caps in stage 2 primordia and young
fruiting bodies. Similarly, in *C. cinerea* peak expression was observed in elongating stipes in
the young fruiting body stage. However, other species did not display this expression profile,
indicating that phytochrome expression is not conserved across mushroom-forming fungi.
Although expression patterns did not seem to be conserved in fruiting bodies, it is possible
that red light sensing phytochromes are part of a photoreceptor system that signals position
in open air, which is required for sporulation and fruiting body formation for the majority of
species(J et al. 2010; Fuller et al. 2015).
Fungal light sensing systems also include rhodopsins, seven-transmembrane domain
proteins involved in sensing green light(Bieszke et al. 1999). We detected rhodopsins only in
L. tigrinus and *Ph. chrysosporium*, but none of the Agaricales in our dataset.
Genes related to blue light receptors - Blue light is one of the most important
signals for fungi and responses as well as molecular aspects of the fungal blue light
response are among the best known aspects of fungal photobiology in both the Asco- and
Basidiomycota. The most important receptors of blue light are the white collar complex
(WCC), which comprises the photoreceptors/transcription factors WC2 and WC1. These

have been discussed under the chapter on transcription factors (see above). We here only
discuss putative accessory proteins similar to VIVID of *N. crassa*. These proteins harbor a
single PAS domain and act as repressors of the WCC in *N. crassa* and were suggested to
provide a mechanism for adapting to changing light intensities during the day(Fuller et al.
2015). Although evidence for that is missing in the Agaricomycetes, a CDE orthogroup of
PAS domain proteins (represented by *C. cinerea* 500409) was developmentally regulated in
9/7 species (FC>2/4) and showed strong induction in primordia relative to vegetative mycelia
in *A. ostoyae, C. aegerita, C. cinerea, L. bicolor, L. tigrinus, M, kentingensis, Ph.*
chrysosporium but not in *A. ampla* and *P. ostreatus* (and missing from *L. edodes* and *S.*
commune). It will be interesting to see if this or similar genes are involved in light responses
in the Agaricomycetes.
Dst2 family - We found that orthologs of the dst2 gene of *C. cinerea* formed a CDE
orthogroup (represented by *C. cinerea* 361584). Dst2 has been described as a FAD-binding
and Berberine-like domain containing protein that, if mutated, results in a blind dark stipe
phenotype similar to what is observed in mutations on the white collar complex(Kamada et
al. 2010; Kuratani et al. 2010). Based on these observations, it has been hypothesized that
dst2 represents a novel photoreceptor(Kuratani et al. 2010). The typical expression profile of
these genes (seen in *A. ostoyae, C. aegerita, C. cinerea, L. edodes* and *P. ostreatus*)
involved upregulation in primordia relative to mycelium and, in later stages, higher
expression in stipes compared to caps. Variations to this pattern exist, such as only being
FB-init in *L. bicolor* and *S. commune*, or being only stipe upregulated but not FB-init in *M.*
kentingensis and *L. tigrinus*. We hypothesize that dst2 genes might be related to stipe
development and, potentially, photomorphogenesis.
2453

2454 4.8. Cell wall biosynthesis and modification

The cell wall is an organelle that functions as an extracellular matrix in fungi. It provides
integrity, rigidity to cells, defines cell shape, harbors extracellular proteins, represents a
channel of communication with the environment, and provides the first line of physical
defense, to name just a few of its major roles in fungal physiology. Structural roles are
particularly important for the Agaricomycetes that form above ground fruiting bodies that
should be rigid yet plastic enough to withstand environmental forces and support spore
dispersal. In fruiting bodies, the cell wall must serve structural and defense roles but must
also be sufficiently plastic to allow changes in shape and size during development. Although
mechanisms of cell shape determination are not known in mushroom-forming fungi,
regulated cell wall deposition and remodeling by hydrolytic enzymes residing in the wall
itself(Verdín et al. 2019), are undoubtedly of key importance.
There is plenty of evidence for cell wall remodeling and differential cell wall
compositions in fruiting bodies of various species (e.g. (Buser et al. 2010a; Almási et al.
2019; Krizsán et al. 2019; Zhou et al. 2019)), although most of that is based on
transcriptome rather than functional studies. For example, stipe growth may be realized by
the breaking of chitin-glucan cross-links by hydrolytic enzymes(Liu et al. 2021). Similarly,
there are cross-linking (transglycosidases) and lytic (e.g. glycoside hydrolases) enzymes
that rigidify and make the cell wall more plastic, respectively(Verdín et al. 2019).
Mannosylation might also rigidify the cell wall and mannoproteins are often N-glycosylated,
which contributes to water retention and adhesion(Lesage and Bussey 2006). In the
Agaricomycetes, Buser et al reported a fruiting body-specific N-glycan from *C. cinerea*(Buser

et al. 2010b). The regulated interplay of these enzymes might result in the formation of
stage-, tissue-, and species-specific cell wall architectures in fruiting bodies.
In this chapter we review current knowledge on cell wall related genes in mushroom-
forming fungi and analyze CDE orthogroups that are likely linked to cell wall synthesis or
modification. We will not review cell wall composition in fruiting bodies, but refer to earlier
reviews(Kües 2000). Cell wall functioning is quite well-researched in the Ascomycota (e.g. in
S. cerevisiae and *N. crassa*), and to a much smaller extent in the Basidiomycota. We
therefore base much of our functional hypotheses on knowledge in the Ascomycota, noting
that the process may be different and/or significantly more complex in the Basidiomycota.
For example, while some chitin-glucan cross linking enzyme families appear similarly
important both in Asco- and Basidiomycota (e.g. GH16/GH17), there are some (e.g. GH72
and GH76) which are involved in cell wall remodeling in the Ascomycota, but apparently not
in the Basidiomycota. Hereafter we discuss cell wall related CDE orthogroups broken down
into synthetic and remodeling enzymes as well as by substrate (chitin/glucan).
From these analyses a few broad observations emerge, which we discuss here. Most
importantly, the diversity of enzymes that show developmental regulation in fruiting bodies
suggests that the fruiting body cell wall may be significantly more complex than we think,
and we still know quite little about its exact structure and composition. A general pattern that
also arises is that more tissue- and stage-resolved transcriptomes reveal proportionately
more developmentally expressed genes, which indicates that cell wall remodeling and,
consequently, cell wall architectures, likely show high tissue-specificity in fruiting bodies.
Based on our results, it appears that fungal cell wall (FCW) remodeling and crosslinking
genes show significantly more expression dynamics than genes encoding chitin and glucan
synthases. In line with this observation, we found that intracellular components of cell wall
biosynthesis (chaperones, glucan synthase complex, pathways involved in glucan monomer
synthesis) also show little expression dynamics.
Comparisons across species revealed limited conservation in FCW related gene
expression patterns across species, which suggests it diverges rapidly during evolution. This
lends support to the quick evolution of cell wall architectures and corresponding enzyme
profiles in fruiting bodies. Another observation related to this is that FCW-related genes
show complex gene duplication/loss patterns, which complicates the inference of well-
conserved groups of 1-to-1 orthologous genes. Thus, these genes might be more
appropriately analyzed in the context of broader orthogroups (e.g. based on
Orthofinder(Emms and Kelly 2015)) than based on the strict 1-to-1 orthogroups we used.
However, for the sake of uniformity, we stick to the latter, with notes on overall prevalence of
developmental expression in a given family.
An interesting observation is that several FCW active genes upregulated in fruiting
bodies are also expressed during mycorrhiza establishment in ECM fungi such as *L.*
bicolor(Martin et al. 2008; C et al. 2014). Such signals provide clues to shared
morphogenetic mechanisms between ectomycorrhizal development and fruiting bodies.
Finally, an important consideration for cell wall biosynthesis/remodeling genes is
whether they are upregulated in fruiting bodies is the vegetative mycelium sample that was
used as control in the RNA-Seq. Non-growing inactive hyphae (where cell wall synthesis rate
is low) as control of actively growing fruiting bodies can yield more gene expression
differences than an actively growing mycelium sample. Although we do not think this has
significantly influenced our comparisons, because we do not know the sampling conditions
for all species, this possibility should be kept in mind in the interpretation of results.

We attempted to as comprehensively analyze cell wall related gene expression as
possible, at least with consideration of genes that show some expression conservation at the
gene family level. Nevertheless, it is certain that several important, especially species-
specific genes are left out from our inventory. For example, mannosylation, a potentially
interesting process contributing to cell wall functioning, is not discussed here due to the lack
of orthology to experimentally characterized mannosylation components in the Ascomycota.
These aspects will need to be covered in future studies.
2530

4.8.1. Cell wall biosynthesis

4.8.1.1. Chitin biosynthesis (GT2)

Chitin is one of the most important, although usually not the most abundant structural
polymer of the fungal cell wall. This is true for fruiting bodies as well, in which, although cell
wall composition is altered relative to vegetative mycelia(M and C 1981; Mol and Wessels
1990; Kües 2000), chitin is an important component that confers rigidity. Chitin is also a
scaffold for the modification of cell wall structure either by cross-linking, incorporation of
glycoproteins or chemical modifications to the chitin polymers, among others(Verdín et al.
2019). Cell wall chitin assembly involves the action of the UDP-N-acetylglucosamine
biosynthesis pathway (in *S. cerevisiae* Gfa1p, Gna1p, Pcm1p, Qri1p), chitin synthases
(Chs1-3) which elongate the growing chitin polymer as well as chitin deacetylases, which
modify the chitin on the cell surface(Shahinian and Bussey 2000; Klis et al. 2006; Verdín et
al. 2019) (the latter is discussed in chapter 4.8.2.1). While yeasts have up to three chitin
synthases, filamentous fungi have additional chitin synthase genes (on average up to
seven(Verdín et al. 2019)), including highly processive ones that harbor a myosin motor
domain and that were lost in yeasts(N et al. 2006).
We examined chitin synthesis and modification related genes based primarily on
functional annotations and recent reviews(Shahinian and Bussey 2000; Klis et al. 2006;
Verdín et al. 2019). Clear orthologs of the *S. cerevisiae* UDP-N-acetylglucosamine
biosynthesis pathway could be identified in all examined Agaricomycetes species (Table 22).
We identified 8-13 (17 in *L. bicolor*) chitin synthases of the GT2 family, 2-3 chitin synthase
export chaperones as well as orthologs of *S. cerevisiae* SKT5 (activator of chitin synthase 3)
in the examined Agaricomycetes. Interestingly, UDP-N-acetylglucosamine biosynthesis
pathway genes showed more expression dynamics than most chitin synthase genes (Table
22). Among chitin synthases, although many formed conserved orthogroups, only a single
appeared developmentally regulated in a wide array of species. Nevertheless, each species
had developmentally regulated and/or FB-init (sensu Krizsan et al 2019) chitin synthases,
though these did not always belong in conserved orthogroups. The most interesting
orthogroup (represented by *C. cinerea* 359180) was developmentally regulated in 7/6 (at fold
change 2/4) species and showed upregulation and high expression in primordia relative to
vegetative mycelium in several species (*C. cinerea, A. ampla, P. ostreatus, S. commune, C.
aegerita* and *L. edodes*). Comparative genomic data suggests that this orthogroup of chitin
synthases is specific to mushroom-forming fungi or the Basidiomycota, suggesting that it
may be a fruiting body-specific chitin synthase. A second CDE orthogroup of GT2 chitin
synthases (*C. cinerea* protein ID: 519767) was also detected (Table 22).
A chitin synthase regulator described as Csr2 from *C. neoformans*(Banks et al. 2005)
and Skt5 from *S. cerevisiae*(S et al. 1993), is conserved in the Agaricomycetes (orthogroup

represented by *C. cinerea* 538650), but did not show significant expression dynamics (Table
22).

4.8.1.2. β-glucan biosynthesis

4.8.1.2.1. β-1,6-glucan (Kre9/Knh1, GH16, GH63)

In addition to β-1,3-glucan, β-1,6-glucan is also found in fungal cell walls and is
hypothesized to serve gluing purposes by covalently linking β-1,3-glucan to chitin and
mannoproteins(Lesage and Bussey 2006). Its synthesis builds on UDP-glucose produced by
the corresponding pathway in *S. cerevisiae*. We find that most members of the yeast beta-
1,6-glucan biosynthesis pathway have no clear 1-to-1 ortholog in the Agaricomycetes (Kre9,
Knh1, Dfg5, Dcw1, Kre5, Kre1). Only Kre6 (*C. cinerea* protein ID: 447635), Rot1 (*C. cinerea*
protein ID: 241938), Rot2 (*C. cinerea* protein ID: 359282) and CWH41 (*C. cinerea* protein ID:
486333) had clear orthologs. However, at the family level, all known members of the β-1,6-
glucan synthesis machinery are represented in the Agaricomycetes (Table 23), except Kre1,
which seems missing.
Kre9/Knh1 - In yeasts and filamentous Ascomycota, the Kre9/Knh1 and GH16 families
include the main glucan synthases(Lesage and Bussey 2006). Several additional well-
characterized proteins in the endoplasmic reticulum are necessary for β-1,6-glucan
synthesis *(e.g. Big1, Rot1, Rot2, Cwh41).* It should be noted that there is a strong overlap
between genes participating in β-1,6-glucan synthesis and remodeling, therefore, separating
these two processes is harder than it is in the case of chitin or β-1,3-glucan.
The Kre9/Knh1 family of cell surface proteins are involved in β-1,6-glucan assembly
in the cell wall of fungi and have been characterized in a number of Ascomycota
species(Brown and Bussey 1993; Dijkgraaf et al. 1996; S et al. 1998b; Costachel et al.
2005), predominantly yeasts. A member of this family has been independently reported as a
developmentally regulated gene, DRMIP, of *L. edodes*(Szeto et al. 2007)*,* which was
reported to be primarily expressed in primordia. Reports of this family in fruiting body
transcriptomes just started to emerge recently(Almási et al. 2019; Krizsán et al. 2019; Liu et
al. 2020b), and no mechanistic study proved yet a link to cell wall in fruiting bodies. The
family is overrepresented in Agaricomycetes(Krizsán et al. 2019), suggesting it might have
experienced duplications specific to mushroom-forming fungi.
We identified 5-15 Kre9/Knh1 genes in mushroom-forming fungi. Of these 3-4 are
developmentally regulated in each species. Notably, each species has 2-3 genes in this
family that show a considerably higher expression in all fruiting body tissues than in
vegetative mycelia, suggesting that these genes are involved in generating fruiting-body
specific cell walls. For example, in *C. cinerea* we found three FB-init Kre9/Knh1 genes (*C.*
cinerea protein ID: 470185, 359649, 465604), whereas in *L. edodes*, there was one FB-init
gene, in addition to DRMIP, which was highly expressed and two-fold upregulated in all
fruiting body tissues compared to vegetative mycelium. The family displayed more
expression dynamics in more resolved transcriptomes (e.g. *A. ostoyae, C. cinerea*),
suggesting that it has considerable tissue-specificity. In addition, each species showed for
some genes an expression peak in vegetative mycelium, which is consistent with a
partitioning of fruiting body and mycelium-specific roles. Taken together, the Kre9/Knh1
family is likely involved in β-1,6-glucan assembly in the cell wall, with several genes showing
fruiting body-specific expression. We detected five CDE orthogroups of Kre9/Knh1 genes
(represented by *C. cinerea* 359649, 465604, 470186, 417746, *S. commune* 2557059)(Table
23).

GH16 - The other major β-1,6-glucan synthase family is GH16, which includes both glucan
synthases (e.g. Kre6 of *S. cerevisiae*, = *C. cinerea* 447635) and glucan-chitin cross-linking
enzymes that participate in cell wall remodeling(Patel and Free 2019). We discuss this family
in detail under the 'Cell wall remodeling' chapter.
Several ER-localized β-1,6-glucan synthesis related proteins have been described in
yeasts and filamentous Ascomycota(Lesage and Bussey 2006; Orlean 2012; Essen et al.
2020), including Kre5, Big1, Rot1, Rot2 and Cwh41. Rot2 and Cwh41 are also involved in
protein N-glycosylation(Lesage and Bussey 2006). These are conserved in the
Agaricomycetes, mostly forming single-copy families with low expression dynamics during
fruiting body development. The GT24 (glycoprotein α-glucosyltransferase) family includes
the Kre5 gene in *S. cerevisiae* and *C. albicans.* Kre5 has an indirect role in β-1,6-glucan
synthesis and may be involved in the refolding of misfolded N-glycoproteins in the ER, in
particular β-1,6-glucan synthases(Lesage and Bussey 2006). GT24 is present as a single-
copy family in Agaricomycetes (Table 23); their expression is approximately constant in
fruiting bodies. Big1 is an ER membrane protein of unknown function that is required for β-
1,6-glucan synthesis in *S. cerevisiae*(Lesage and Bussey 2006). It is a single-copy gene
family (represented by *C. cinerea* 432992) in the Agaricomycetes, without developmentally
relevant expression patterns (Table 23). Rot1 is a chaperone involved in protein folding (in
general) located in the ER. It is required for cell wall synthesis and N-glycosylation and O-
mannosylation in *S. cerevisiae*(Lesage and Bussey 2006). Rot1 orthologs are single copy
genes (*C. cinerea* protein ID: 241938) in the Agaricomycetes that showed no developmental
dynamics in the species we examined (Table 23). Rot2 is an ER-localized alpha-glucosidase
involved in protein N-glycosylation, required for normal levels of β-1,6-glucan in the cell wall
of *S. cerevisiae*(Lesage and Bussey 2006). It belongs to the GH31 family, which is present in
3-8 copies in Agaricomycetes. They form two CDE orthogroup (FC2/4: 8/6 species,
represented by *C. cinerea* 499280 and FC2/4: 6/5 species, represented by *S. commune*
2543909)(Table 23), which were developmentally regulated in 6/5 and 5/4 species, however,
expression patterns of these orthogroups did not show marked conservation. 1-to-1
orthologs of *S. cerevisiae* Rot2 (represented by *C. cinerea* 359282) showed very small gene
expression dynamics.
The GH63 family includes alpha-glucosidases, such as Cwh41 of *S. cerevisiae,*
involved in cell wall beta-1,6-glucan assembly in the ER. The family is represented by 2
copies in all but one of the examined species (*L. edodes* has 5 genes). GH63 genes form
two conserved orthogroups, of which one (*C. cinerea* protein ID: 486333), containing
orthologs of *S. cerevisiae* Cwh41) did not appear to be interesting, whereas members of the
other (*C. cinerea* protein ID: 494484) showed marked expression peaks coinciding with
spore formation in all species (Table 23). In *C. neoformans*, the gene showed upregulation
at 72h, also coinciding with sporulation. Whether this GH63 orthogroup is needed for spore
wall assembly or other processes that temporally overlap with sporulation cannot be
determined based on these data, nevertheless, the family seems to be interesting in the
context of fruiting body development.

4.8.1.2.2. β-1,3-glucan (GT48)

In *S. cerevisiae* β-1,3-glucan is synthesized by the FKS (Fks1-3) and the GAS (Gas1-4
families as well as a few accessory proteins, such as Smi1 (*C. cinerea* protein ID: 181364)
and Rho1 (*C. cinerea* protein ID: 528086)(Lesage and Bussey 2006; Verdín et al. 2019). Of
the glucan synthases only FKS1 and Gas1 had clear 1-to-1 orthologs is the Agaricomycetes
(*C. cinerea* protein ID: 446030 and 543646, respectively). The glucan synthase complex,

which is formed by FKS1, Rho1, Smi1(Verdín et al. 2019), seems to be conserved in the
Agaricomycetes.
Most Agaricomycetes species have two beta-1,3-glucan synthase genes of the GT48
family (except *M. kentingensis* and *L. tigrinus*, which have 5 and 7 respectively). Somewhat
surprisingly, GT48 genes showed very modest expression dynamics and were mostly not
developmentally regulated. Similarly, the Gas1 family of glucanosyltransferases is present in
1-2 copies in the examined species (*L. bicolor* has 7 and *M. kentingensis* has 4) and they
were not developmentally regulated. Accessory members of the glucan synthase complex
(Smi1, Rho1) showed constant expression in fruiting bodies. Collectively, these data suggest
that *β*-1,3-glucan synthesis is present, but it is not dynamically regulated in fruiting bodies.
2671

4.8.2. Cell wall remodeling

4.8.2.1. Chitin-active enzyme families (GH18, GH20, X325, AA9, AA14, CBM5/12, CBM50,
CE4)

GH18 - Glycoside hydrolase family 18 contains chitinases (both endo- and exochitinases).
Chitinases remodel the chitin structure by cleaving β-1,4 linkage of chitin or modify
oligosaccharide(Verdín et al. 2019) (e.g. cleaving chitobiose from the cell wall). Chitinases
are one of the most straightforward players in fruiting body development and accordingly
have been reported in several previous genomic(Chen et al. 2012; Park et al. 2014),
transcriptomic(Patyshakuliyeva et al. 2013; Sakamoto et al. 2017b; Xie et al. 2018; Krizsán
et al. 2019) and mechanistic(Niu et al. 2016; Zhou et al. 2016c) studies. Despite the clear
prediction of their role in morphogenesis, this has not been mechanistically shown in the
Basidio- or Ascomycota(L et al. 2011), possibly because multiple paralogous genes in each
species' genome complicate functional studies.
In the Basidiomycota, Zhou et al reported chiB1 (*C. cinerea* protein ID: 368217)
(Table 24), a class V exochitinase purified from *C. cinerea* and suggested to be involved in
cap autolysis(Zhou et al. 2016c). Expression data agree with this, the corresponding gene
shows a very strong induction in mature fruiting body caps. Similarly, ChiIII(Niu et al. 2016)
of *C. cinerea* (C. cinerea protein ID: 354859) was reported to have both endo- and
exochitinase activity, and to be expressed mostly in fruiting bodies, with peak expression in
mature fruiting bodies (which is confirmed by our expression data). Most recently, Zhou et al
assayed the role of chitinases in stipe cell wall extension(Zhou et al. 2019) and named six
further chitinases in *C. cinerea*, ChiE1, (*C. cinerea* protein ID: 543586), ChiE2 (*C. cinerea*
protein ID: 520359), ChiEn1 (*C. cinerea* protein ID: 91051), ChiEn2 (*C. cinerea* protein ID:
90984), ChiEn3 (*C. cinerea* protein ID: 470416), and ChiEn4 (*C. cinerea* protein ID: 358869).
Chitinases have also been examined in the context of autolysis of the model mushrooms *C.*
cinerea and *Coprinellus congregatus*(H and HT 2009).
GH18 genes are found in 8-12 copies in Agaricomycetes (up to 20 in Polyporales,
see (Chen et al. 2012)), some of which harbor carbohydrate binding modules (CBMs), but
their function has not been examined. Most chitinases are developmentally expressed in
fruiting bodies. Mycelium-upregulated chitinases are also found in most species; these,
although display peak expression in vegetative mycelia (often ~10x that of fruiting bodies),
often also show >4fold expression dynamics within fruiting bodies. We detected seven CDE
orthogroups *(Table 24), which* included *C. cinerea* ChiB1, ChiE2, ChiEn1 and ChiEn3. In
general, although, the detected chitinase orthogroups were developmentally regulated in
many species, they did not show conserved expression patterns across species, which could

indicate quick divergence of function or the failure of our reciprocal-best-hit based approach
to detect functionally related genes (in the latter case phylogenetic analyses might help).
One orthogroup (represented by *C. cinerea* 520359) shows gill-specific expression peaks in
the species in which separate RNA-Seq data are available for gills (*A. ostoyae, P. ostreatus*)
or late upregulation in *Pt. gracilis, C. aegerita, L. edodes, L. bicolor, Ph. chrysosporium, A.*
ampla and *S. commune*, suggesting it may have something to do with processes that take
place in the hymenium late in development, possibly sporulation.
GH20 and chitin degradation products - A chitin-connected GH family is GH20, which has
β-1,6-N-acetylglucosaminidase activity and thus can cleave chitobiose released by
chitinases further into N-acetylglucosamine (GlcNac). GlcNac may be taken up by the cell for
repurposing of cleaved cell wall components as nutrient source(López-Mondéjar et al. 2009;
de Oliveira et al. 2018) or, to trigger upregulation of chitinolytic genes in a transcriptional
feedback-loop(Langner and Göhre 2015). The latter hypothesis seems more plausible for
cases when the fungus feeds on chitin rather than modifies it for developmental purposes. In
previous reports, GH20 genes have been identified in the post-harvest transcriptome of *L.*
edodes(Sakamoto et al. 2017b). We identified 2-8 GH20 genes in the examined
Agaricomycetes, some of which were developmentally regulated (including the single gene
in *C. neoformans*), but did not form conserved CDE orthogroups. The role of this gene family
in fruiting body development may be worth examining further.
If GlcNac is indeed taken up and metabolized in the cell, then two of the detected
CDE orthogroups may provide clues to its mechanistic bases. An orthogroup of GlcNac
transporters, which contains reciprocal best hits of *C. albicans* NGT1, was developmentally
regulated in 5/4 species (*C. cinerea* protein ID: 464567, Table 24). *C. albicans* NGT1
encodes a specific transporter of GlcNac(FJ and JB 2007; Q et al. 2020). We speculate that
this orthogroup may be involved in the uptake of GlcNac monomers generated by GH20
enzymes extracellularly.
In fungi, the GlcNac catabolic pathway is best described in *C. albicans*(Kumar et al.
2000; Yamada-Okabe et al. 2001), however, only a few of the corresponding genes are
conserved in Agaricomycetes. Only the GlcN6P deaminase (Nag1, *C. cinerea* 356467
FC2/4: 7/4 species) and a GlcNac permease Nag4 (*C. cinerea* protein ID: 544873, FC2/4:
6/4)(Table 24) genes have clear reciprocal best hits in Agaricomycetes, whereas we could
not identify clear orthologs of N-acetylglucosamine kinase (Nag5) or N-acetylglucosamine
deacetylase (Nag2). Our annotations identified a different gene as N-acetylglucosamine
deacetylase, however (*C. cinerea* protein ID: 446427), which was developmentally regulated
in 8/6 species (FC=2/4). We also detected a GlcNac deacetylase orthogroup (*C. cinerea*
protein ID: 446427), which may be involved in intracellular GlcNac catabolism.
In summary, the evidence presented here on GlcNac uptake and catabolism in
mushroom forming-fungi is circumstantial, at best, however, provides clues for further
research on this topic. GlcNac (and any other breakdown intermediate) produced during cell
wall modification may be precious compounds in the fruiting body, which is otherwise
dependent on distal supply of nutrients from the mycelium. This might have prompted the
evolution of efficient repurposing mechanisms in fruiting bodies to mitigate its dependence
on mycelial nutrient supply.
CE4 chitin deacethylases - Chitin deacetylases (CDA) are involved in chitosan production,
a minor but key constituent of the cell wall. Chitosan, a more soluble polymer than chitin,
occurs in both mycelia and fruiting bodies of basidiomycetes, although its role in fruiting body
formation is thought to be limited(M and C 1981; Crestini et al. 1996; Baker et al. 2007). It is
produced by the synergistic activities of chitin synthases and CDAs, where the former

synthesize and the latter deacetylate chitin in various spatial or temporal patterns and to
varying degrees(Sebastian et al. 2020). CDAs belong to the carbohydrate esterase 4 family
(CE4) in the CAZy classification and are homologous to bacterial nodB factors that produce
signal molecules in the rhizobia-legume symbiosis(Buhian and Bensmihen 2018).
Deacetylation of chitin to chitosan makes it resistant to chitinases, which might provide a
mechanism for regulating the extent of chitinase-mediated wall plasticity and thus cell
expansion (e.g. in stipe elongation(Liu et al. 2021)). Fungal CDAs have a potential in
biotechnology, as chitosan is an additive in food, cosmetic and pharmaceutical
products(Morin-Crini et al. 2019). CDAs have poorly known roles in Ascomycota fruiting
body development, with both *N. crassa* CDAs, nevertheless, being highly expressed during
perithecium development, suggesting roles in sexual morphogenesis(Patel and Free 2019).
It should be noted that CE4 also contains acetyl xylan esterases, which are able to break
down the xylan backbone of hemicellulose during wood decay(X and S 2009). Therefore,
some genes of the family in the Agaricomycetes may be related to wood decay too.
CDA genes have been detected in several RNA-Seq studies focusing on fruiting
body development(Morin et al. 2012; Sipos et al. 2017b; Xie et al. 2018; Krizsán et al. 2019).
Three chitin deacetylases have been experimentally characterized in *C. cinerea*, CDA1 (*C.*
cinerea protein ID: 42776), CDA2 (*C. cinerea* protein ID: 502602), and CDA3 (*C. cinerea*
protein ID: 464952)(Table 24), all of which were reported to have high expression in the
elongating region of the stipe(Wang et al. 2018b; Bai et al. 2020). Another chitin
deacetylase, primarily expressed in the fruiting bodies, was investigated in *F. velutipes*(M et
al. 2008).
Searching the genomes of fruiting body forming fungi reveals 5-18 chitin deacetylase
genes, several of which formed conserved single-copy orthogroups. The previously
investigated CDA1-3 genes of *C. cinerea* formed species-specific orthogroups. The majority
of CDA genes in the examined species were developmentally regulated in fruiting bodies,
indicating key functions in fruiting body development. Six out of 17 *C. cinerea* CDA genes
(Table 24) formed CDE orthogroups. These showed diverse expression peaks with no
apparent trend in their expression. However, several genes were upregulated in stipes of
young or mature fruiting bodies. One of them (*C. cinerea* protein ID: 355445) comprise
proteins representing reciprocal best hits of *S. cerevisiae* CDA1, which is involved in the
biosynthesis of the ascospore wall, required for spore wall rigidity. Expression of genes in
this orthogroup did not peak during spore development, suggesting that this specific gene
orthogroup is not involved in basidiospore wall synthesis. On the other hand, the expression
of one and three genes were highest in young fruiting body gills and mature fruiting body
gills and caps, respectively, which might be involved in producing chitosan in spore walls.
On a related note, we could not detect chitosanases, which degrade chitosan, in the
genomes of mushroom-forming fungi, suggesting that chitosan is a terminal cell wall
carbohydrate, not modified or cleaved further.
Carbohydrate binding modules (CBMs) - Carbohydrate binding modules, as their name
suggests, bind various carbohydrates, and thus can frequently be attached to cell wall
modifying enzymes. Chitin binding ability has been shown for at least 9 CBM
families(Sánchez-Vallet et al. 2015). Of the most widely known chitin-binding CBMs (1, 5/12,
14, 18, 19, 50), only CBM1, 5/12, 18 and 50 families are conserved in the Agaricomycetes
we examined. The CBM1 family, which primarily binds cellulose and according to some
reports chitin, does not appear to be involved in fruiting body development: most genes that
encode CBM1 containing proteins have virtually no expression in fruiting bodies. Some such
proteins show significant expression dynamics (e.g. *C. cinerea* 363583, 499390), but are not

conserved across species. The CBM18 family occurs in GH16 proteins, which have been
described above.

CBM5/12 - The CBM5/12 family comprises chitin-binding modules, which assist the
action of cell wall modifying enzymes in fungal development(Hartl et al. 2011). The CBM5
family is sometimes considered a lectin due to its carbohydrate binding ability(Ismaya et al.
2020). Most CBM5/12 modules occur in combination with GH18 domains, where their chitin
binding ability is believed to increase the efficiency of the chitinase(Hartl et al. 2011). We
identified 3-6 CBM5/12 domains in the examined Agaricomycetes. All but one CBM5/12
modules are attached to GH18 chitinases and have thus been discussed above. The single
CBM5/12 module that is not attached to a chitinase occurs on proteins that also harbor a
domain of unknown function (DUF3421) and a DM9 repeat as well. Such proteins occur in
single copy in Agaricomycetes genomes and comprise a CDE orthogroup (*C. cinerea* protein
ID: 413423) that show high expression in vegetative mycelia and mature stipe tissues of
most species (*A. ostoyae, P. ostreatus, M. kentingensis, C. cinerea, L. tigrinus, L. bicolor, L.*
edodes) (Table 24). The protein is functionally uncharacterized. Both the DUF3421 and the
DM9 repeat occur primarily in insects and other animals (but are found also in early-
diverging fungi), but their activity is not known. Their developmental expression and the
presence of a CBM5/12 domain suggests they may be linked to the chitin content in the cell
wall, however, how exactly they function remains unclear.

CBM50 - also known as LysM domain, this family binds chitin and can occur in a
number of chitin-active proteins or on its own(GB et al. 2015). It has been reported to show
developmentally dynamic expression in previous studies of fruiting body
development(Sakamoto et al. 2017b; Sipos et al. 2017b; Krizsán et al. 2019; Huang et al.
2020). LysM domain proteins are widely known as effectors of plant pathogenic fungi, in
which solitary secreted CBM50 modules can mask chitin residues and thus can dampen the
immune response of plants(GB et al. 2015). Solitary CBM50 modules expressed in fruiting
bodies could be involved in defense against insects(Kunzler 2015).

The examined Agaricomycetes contain 3-9 CBM50/LysM domain encoding genes
(except *A. ostoyae,* which has 18, but its expansion may be related to the pathogenic nature
of *Armillaria*(Sipos et al. 2017b)). The CBM50 module occurs almost exclusively on its own
in the examined Agaricomycetes, only *C. aegerita* has proteins that harbored combinations
of CBM50 and chitinase domain signatures. Interestingly, CBM50 encoding genes showed
relatively little expression dynamics within fruiting bodies with 0-2 developmentally
expressed genes per species. This, and reports on the induction of one of the CBM50 genes
of *C. cinerea* in response to bacterial and nematode challenge(Kombrink et al. 2018;
Tayyrov et al. 2018, 2019) could indirectly support a role in defense. However, some genes
had high and constant expression during fruiting body development (e.g. *C. cinerea*
448942), suggesting chitin-linked functions.

AA9 family (Lytic polysaccharide monooxygenases) - The family has been
reported only recently to have fruiting body specific expression patterns(Almási et al. 2019;
Krizsán et al. 2019). AA9 proteins were initially considered to be endoglucanases and
classified in glycoside hydrolase family 61, but turned out to be lytic polysaccharide
monooxygenases (LPMOs) that cleave polysaccharide chains using a distinct oxidative
mechanism(G et al. 2010).

The examined Agaricomycetes possess 7-34 AA9 encoding genes, of which 1-10
were developmentally regulated. A part of this copy number variation can be explained by
lifestyle, as wood-decay species are known to have expanded, while biotrophic or
mycorrhizal species have contracted AA9 repertoires(Miyauchi et al. 2020). AA9 LPMOs

might be active on β-glucan or on both glucan and cellulose as well, providing them with the
activity required to remodel the fungal cell wall(Krizsán et al. 2019). The identified AA9
genes hardly formed conserved orthogroups, which may be a consequence of high
duplication/loss rates.
Several species showed interesting LPMO expression patterns, which were quite
consistent within, but different between species. For example, in *C. cinerea* several genes
had a late upregulation, suggesting that LPMOs might be involved in autolysis. On the other
hand, in *Ph. chrysosporium* many genes showed high expression in vegetative mycelium
and mature fruiting bodies but low in young fruiting bodies, creating U-shaped expression
curves. In *L. tigrinus* several genes showed a marked peak in cap and gill samples of mature
fruiting bodies. The single AA9 gene (Cryne_JEC21_1_562) of *C. neoformans* had a strong
(45-fold) induction in 24h samples, coincident with basidium formation(Liu et al. 2018). In
general, all species displayed several genes with high expression in vegetative mycelium,
but negligible (<5 FPKM) expression in any of the fruiting body tissues, suggesting the
existence of separate vegetative mycelium-specific and fruiting body-specific AA9
subgroups. Yet other AA9 genes displayed uniformly low (<5 FPKM) expression; these
LPMO-s might be specific for the plant cell wall, and thus not induced during development.
Nevertheless, the developmentally dynamic expression of several AA9 genes suggests that
this family is involved in fungal cell wall remodeling during fruiting body development.
AA14 family - We found a conserved orthogroup (represented by *C. cinerea*
362412) that contained members of the recently circumscribed AA14 family of lytic xylan
oxidases(Couturier et al. 2018). These proteins require copper as their native cofactor and
were speculated to unlock xylan-coated cellulose fibers in the plant cell wall. We identified
members of the AA14 family based on BlastP (query *Pycnoporus coccineus* AUM86167 and
AUM86166): we find that blastp with e-value cutoff of 10^{-10} yielded copy numbers that match
perfectly those reported by Couturier et al (Suppl Data Set 1)(Couturier et al. 2018). Fruiting
body forming fungi contained 2-5 copies of the family, whereas yeast-like fungi (*U. maydis*
and *C. neoformans*) contained none. Gene belonging to this family did not display
characteristic expression peaks across species. However, as with other GH and AA families,
their developmental expression in fruiting bodies suggests they may be active on some
component of the fungal cell wall as well or have other extracellular roles.
X325 family - We also detected the recently described X325 family of copper-
containing, LPMO-like proteins(Labourel et al. 2020). As a new gene family, the only
functional knowledge available currently is that it displays a fold similar to fungal LPMOs,
may be involved in copper homeostasis(Garcia-Santamarina et al. 2020), is upregulated in
several mycorrhizal fungi and resides on the surface (via a GPI anchor) of the fungal cell
wall(Labourel et al. 2020). A morphogenetic role for the family has been speculated. We
here report that members of this family form a CDE orthogroup (represented by *C. cinerea*
454418) and are developmentally regulated in a wide range of species (9/6 at FC4/2),
although expression patterns differ between species. Additional developmentally regulated
X325 genes from other species were segregated into other orthogroups. The *L. bicolor* gene
is slightly upregulated in primordia relative to free living mycelium (FC=1,8) and has a peak
expression in young fruiting body stipes.
AA10 and AA11, two well-known chitinolytic LPMO families(Langner and Göhre
2015; Polonio et al. 2021), are missing in the examined Agaricomycetes, so they are unlikely
to be contributors to fruiting body development.

4.8.2.2. Glucan-linked families (GH3, GH5, GH16, GH17, GH30, GH55, GH71, GH128,
GH152, Cdc43, WSC)

Cdc43 - We detected orthologs of *S. cerevisiae* Cdc43 (*C. cinerea* protein ID: 376392) as a
CDE orthogroup in this study (Table 25). Cdc43, together with Ram2, forms the type I
geranylgeranyl transferase of *S. cerevisiae*, which transfers a geranyl-geranyl moiety to
proteins having a CaaX sequence on their C-terminal(AA et al. 1991). One of the targets in
S. cerevisiae is Rho1p, a rho-type GTPase, which is the regulatory subunit of beta-1,3-
glucan synthase(SB et al. 1999). Cdc43 orthologs were developmentally expressed in 11/6
(FC=2/4) (Table 25) species as well as *C. neoformans*. They showed a marked upregulation
from vegetative mycelium to primordia in *A. ampla, A. ostoyae, C. aegerita, L. bicolor, L.
edodes, C. cinerea, S. commune*, and more or less constant expression in *M. kentingensis,
Pt. gracilis* and *P. ostreatus*. On the other hand, orthologs of Ram2 (*C. cinerea* protein ID:
473005) as well as those of the main substrate, Rho1 (*C. cinerea* protein ID: 528086) did not
show strong expression dynamics. We hypothesize that the expression dynamics of Cdc43
mirrors the dynamics of cell wall glucan biosynthesis in fruiting bodies.
GH17 - Enzymes belonging to the GH17 family are known to crosslink β-1,3-glucan in the
cell wall of yeast species and presumably *N. crassa* too(Patel and Free 2019). The family
has been reported in a transcriptomic study of mushroom development(Krizsán et al. 2019)
(as endo-β-1,3-glucanosyltransferase), and it was suggested to be involved in β-1,3-glucan
crosslinking, although direct evidence for this is lacking at the moment. The examined
Agaricomycetes possess 1-4 GH17 encoding genes. *C. cinerea* has four GH17 genes, of
which only one appear to be clear orthologs of Ascomycota genes: *C. cinerea* 389921 is
reciprocal best hit to SCW10 of *S. cerevisiae* a soluble cell wall protein with a poorly known
function. GH17-s formed 2 CDE orthogroups (Table 25). One of the CDE orthogroups
(represented by *C. cinerea* 476698) showed clear tissue-specific expression (cap- or stipe
upregulation), but interestingly, this was not fully consistent across species.
GH16 - The GH16 family shows glycosylhydrolase and glycosyltransferase activity and
contain glucan synthase and crosslinking enzymes, which are active in cell wall synthesis
(e.g. Kre6 and Skn1 of *S. cerevisiae*) and remodeling. GH16s were shown to crosslink β-1,3-
glucan and β-1,6-glucan to chitin fibers in *Candida* spp and *N. crassa*(Patel and Free 2019).
Chitin-glucan crosslinking is key to morphogenesis in fungi(Arroyo et al. 2016). GH16s
comprise the CRH (Congo Red Hypersensitive) family, three members of which (Crh1p,
Crh2p and Crr1p) were originally circumscribed in *S. cerevisiae.* Of these, only Crr1 has a
clear 1-to-1 ortholog in Agaricomycetes (*C. cinerea* protein ID: 449960), the other two do
not. At the transcriptome level, the upregulation of GH16 genes have been reported before
in a comparative study(Krizsán et al. 2019), in *A. bisporus*(Morin et al. 2012;
Patyshakuliyeva et al. 2013)*, Leucocalocybe mongolica*(Duan et al. 2021)*, Pleurotus spp*(Xie
et al. 2018; Ye et al. 2021) as well as during post-harvest development on *L.
edodes*(Sakamoto et al. 2009, 2017b). Mlg1 is a functionally characterized member of the
GH16 family in *L. edodes* assayed for its glucanase activity during postharvest
senescence(Sakamoto et al. 2009).
The species in this study have 21-36 GH16 encoding genes, which is consistent with
a previous report on *L. edodes*(Sakamoto et al. 2017b). About half of the GH16 genes in
each species also carry a 'Beta-glucan synthesis-associated Skn1' domain, which signifies
relationship with the Kre6/Skn1 family, which is involved in β-1,6-glucan synthesis(Garcia-
Rubio et al. 2020). We found five GH16 CDE orthogroups (*C. cinerea* protein ID: 502321,
381870, 411452, 169822, *S. commune* 2686544) (Table 25), that are not orthologous to

functionally characterized GH16 genes in the Ascomycota. Although most genes do not form
conserved orthogroups, possibly because of volatility (frequent duplication/loss), it is
noteworthy that each species has 1-3 genes which show a marked upregulation in the
primordium stage and remain highly expressed in the fruiting body relative to vegetative
mycelium. These might be involved in the formation of 3-dimensional fungal tissues. Another
interesting observation is that eight out of 21 GH16 genes in *C. neoformans* are also
developmentally regulated, with several of them showing peak expression at 24h, coincident
with basidium development.
GH30 - Glycoside hydrolase family 30 (GH30) includes enzymes with diverse activities,
which have been mostly discussed for their role as endo-xylanases in wood decay in the
fungal literature(Wymelenberg et al. 2009; Katsimpouras et al. 2019). However, GH30 also
includes β-1,6-glucanases, which were found to be upregulated in *L. bicolor*
ectomycorrhizae(C et al. 2014), raising the possibility of morphogenetic roles. GH30
members were also upregulated during postharvest development of *L. edodes*(Sakamoto et
al. 2017b). Of the 1-3 (6 in *L. bicolor*) GH30 genes in the examined Agaricomycetes, at most
1-2 are developmentally expressed in fruiting bodies and only a single orthogroup
(represented by *C. cinerea* 381637) was so in several species (Table 25). Genes in this
orthogroup were most often upregulated in stipe tissues relative to pileus in fruiting bodies.
GH30 genes of *C. cinerea* have been proposed to facilitate stipe elongation by in the stipe
base may contribute to replacing β-1,6-glucan to β-1,3-glucan crosslinking in the stipe base,
which may reduce wall extensibility, leading to the cessation of growth(Liu et al. 2021).
Further, the shared upregulation of GH30 genes in multiple fruiting body transcriptomes and
in ectomycorrhizae provide strong evidence for its role in FCW remodeling during
morphogenesis in the Basidiomycota.
GH3 - The GH3 family comprises enzymes with diverse activities, including exo-β-1,3-
glucanase activity, and has been suggested to participate in cell wall remodeling in
Ascomycetes(Patel and Free 2019). The upregulation of GH3 genes in fruiting bodies has
previously been noted in a number of species(Zhou et al. 2018; Krizsán et al. 2019), raising
the possibility that they participate in cell wall remodeling during fruiting body development.
A GH3 family member, that was upregulated in fruiting bodies of *Auricularia heimuer*, was
recently cloned and characterized(Sun et al. 2020). GH3 encoding genes are present in 7-20
copies in examined Agaricomycetes. Three CDE orthogroups belonging to the GH3 family
(*C. cinerea* protein ID: 369876, 373418, *S. commune* Schco3_2629974,) were detected
(Table 25), genes in these orthogroups showed diverse expression patterns in the fruiting
bodies of different species. It should be mentioned that some GH3 enzymes, mostly in
bacteria, possess β-N-acetylhexosaminidase activity, which allows the encoded enzymes to
be involved in chitin utilization, by cleaving N-acetylglucosamine dimers into monomeric
GlcNac(de Oliveira et al. 2018). Most such enzymes are bacterial, with sporadic reports of β-
N-acetylhexosaminidase activity in fungi(S et al. 2014). Therefore, whether GH3
upregulation is linked to chitin utilization or glucan remodeling cannot be determined here.
GH5 - The GH5 family is a very diverse group that includes β-1,4-glucanases, as well as
several other activities. It includes classic cellulases involved in wood decay(Floudas et al.
2012), whereas Ascomycota GH5 genes, such as Exg1, Exg2 and Exg3 of *S. pombe*, or
Exg1 of *S. cerevisiae*, are well-known components of the cell wall glucan remodeling
machinery(Dueñas-Santero et al. 2010). GH5s were shown to not only hydrolyze β-1,3-
glucan, but also to act as transglycosylases, and thus contribute to cell wall
crosslinking(Arroyo et al. 2016). Based on expression patterns, GH5s were speculated to be
active on beta-glucan in fruiting body cell walls also(Y et al. 2005; C et al. 2014; Almási et al.

2019; Krizsán et al. 2019). Krizsan et al reported multiple subfamilies (GH5_7, GH5_9,
GH5_15, GH5_49) with endo-β-1,4-mannanase, endo-β-1,6- and exo-β-1,3-glucanase
activities to be upregulated in fruiting bodies(Krizsán et al. 2019). The family includes the
exo-beta-1,3-glucanase Exg1 from *L. edodes*, which was reported to be fruiting body
specific(Y et al. 2005). Our re-analysis confirms that Exg1 (Lenedo1_1176007) has
consistently higher expression in stipe than in cap tissues.

GH5 is a large family with 16-20(25) copies in the examined Agaricomycetes.
Consistent with their diversity, we detected six CDE orthogroups in our data (*C. cinerea*
protein ID: 546300, 405828, 505989, 496411, 414471, 373700) (Table 25). One of these (*C.*
cinerea protein ID: 546300), contains reciprocal best hits of *Sch. pombe* Exg3, a cytoplasmic
β-1,6-glucanase with unknown cellular function. All species in which this orthogroup is
developmentally regulated showed upregulation in mature fruiting bodies. This could indicate
a link to sporulation, or other late developmental events. Another orthogroup (*C. cinerea*
protein ID: 414471) contains 1-to-1 orthologs of *Sch. pombe* and *S. cerevisiae* Exg1, which
is involved in cell expansion and sporulation, picturing potentially similar roles in
Agaricomycetes too. Members of yet another orthogroup (*C. cinerea* protein ID: 373700), are
consistently higher expressed in all fruiting stages (primordia to mature FB) than in
vegetative mycelia in *A. ostoyae, C. aegerita, A. ampla, C. cinerea, P. ostreatus, S.*
commune and to some extent in *L. bicolor* and *L. edodes* (but not in *A. bisporus* and *Pt.*
gracilis).

GH55 - This family contains exo- or endo-β-1,3-glucanases that have been characterized
functionally in fruiting bodies of two Agaricomycetes. Exg2 was found to be expressed higher
in the stipe than in the cap of *L. edodes*(Sakamoto et al. 2005). The GH55 family has also
been reported to be upregulated during cap expansion of *Volvariella volvacea* and the
corresponding gene was named Exg2(Tao et al. 2013). We detected on average two GH55
genes in the examined Agaricomycetes (5 in *A. ostoyae*). One of these formed a CDE
orthogroup (represented by *C. cinerea* 492834), which included *L. edodes* Exg2 and showed
very strong induction in young fruiting body caps and stipes of *C. cinerea*, caps of *A. ostoyae*
and stipes of *M. kentingensis*. Overall, these data are consistent with the suggested role of
GH55 in cap expansion in *V. volvacea*. In *L. edodes*, the expression of Exg2 was higher in
stipes than in cap tissues. Interestingly, the second GH55 gene of *C. cinerea* (*C. cinerea*
protein ID: 454337) (Table 25) also shows this characteristic induction and high expression
in young fruiting body caps and stipes. In species with simpler morphology (*S. commune, A.*
ampla) the family did not show characteristic expression dynamics. A third GH55 gene
(Armosto1_267056) in *A. ostoyae* showed an induction in primordia relative to vegetative
mycelium.

GH128 - GH128 has been first circumscribed in the context of fruiting body development,
based on a fruiting body expressed β-1,3-glucanase, Glu1, in *L. edodes*(Y et al. 2011).
Since its discovery, some reports have mentioned upregulation of GH128 genes in fruiting
bodies(Krizsán et al. 2019; Duan et al. 2021; Merényi et al. 2021), with a single report also in
connection with wood decay(Mäkinen et al. 2019). On average, the GH128 family has 2-8
members in Agaricomycetes. We detected two CDE orthogroups (represented by *C. cinerea*
463000, 413757) which were expressed in the majority of species (Table 25) and several
GH128 genes which were developmentally expressed in a species-specific way. In *C.*
cinerea, protein 413757 represents the best hit of *L. edodes* GLU1. Available data here and
in previous publications suggest that GH128 might be a primarily fruiting body-related gene
family, worthy of further functional examination.

GH71 - Four orthogroups containing developmentally regulated GH71 alpha-1,3-glucanases
and beta-glucuronidases were detected (*C. cinerea* protein ID: 22054, 464399, 456054,
492754)(Table 25). GH71 genes (also called mutanases) are involved in fungal cell wall
remodeling in a wide range of species, both yeasts and filamentous fungi(Dekker et al. 2004;
C et al. 2014; H et al. 2015). A well-characterized member of this family is the *Sch. pombe*
endo-1,3-alpha-glucanase Agn1, which facilitates cell fission via the degradation of the
septum material(Dekker et al. 2004). In previous studies of mushroom development, this
family was reported only by Krizsan et al(Krizsán et al. 2019), Park et al(Park et al. 2014),
and Morin et al(Morin et al. 2012). In *Pleurotus eryngii*, GH71 genes were found
downregulated in primordia after blue light stimuli(Xie et al. 2018). GH71 genes are present
in 3-12 copies in the examined Agaricomycetes. One of the CDE orthogroups (represented
by *C. cinerea* 456054) is upregulated in primordia relative to vegetative mycelium in 7 of the
10 species (*C. cinerea, A. ostoyae, C. aegerita, L. tigrinus, L. bicolor, Ph. chrysosporium, P.*
ostreatus) in which it is present.
GH152 - The GH152 family, also known as thaumatin-like proteins, which include both
putatively defense-related antimicrobial peptides and β-1,3-glucanases. They show
homology to plant thaumatin-like proteins. Because the literature reports both antimicrobial
and glucanase activity for this family(Kunzler 2015), we do not attempt to subdivide them
and refer to them as GH152/thaumatin proteins. The GH152/thaumatins include an
experimentally characterized gene, Tlg1 in *L. edodes* (Lenedo1_561169), which was
detected in a post-harvest screen of cell wall and lentinan degrading genes(Sakamoto et al.
2006). Lentinan is a glucan-like cell wall component with important medicinal properties in *L.*
edodes. The expression of the gene was detected only after harvest, raising questions as to
what the physiological role of the gene might be under natural circumstances. Our analysis
of the data from Zhang et al did not reveal developmentally dynamic expression for Tlg1
using RNA-Seq data(Zhang et al. 2021c). Structure prediction for an *A. ostoyae*
developmentally expressed protein indicated an acidic cleft in the 3D structure, characteristic
of antimicrobial members of the family(Krizsán et al. 2019). A new report of antifungal
activity of the thaumatin-like protein Le.TLP1 in *L. edodes* further supports putative defense
roles for the GH152 family(Ma et al. 2021).
The family comprises 2-8 (19 in *Pt. gracilis*) copies in examined Agaricomycetes,
many of which are poorly conserved. Therefore, while several genes are developmentally
expressed in each species, only a single CDE orthogroup was detected (represented by *C.*
cinerea 496344)(Table 25). The role of GH152/thaumatins in fruiting body development is
unclear at the moment, it could be involved in defense against microbes, cell wall glucan
remodeling, or both(Kunzler 2015).
Additional beta-glucan active families (GH1, GH6, GH12) were detected in the
broader orthology-based approach of Krizsan et al(Krizsán et al. 2019), which were not
detected in this study, probably because of the strict orthology definition we use.
WSC family - this family of lectin-like proteins comprises a widely conserved but poorly
known group of fungal proteins. The family was originally described as a putative stress
receptor in *S. cerevisiae*(AL et al. 1999) and as a domain in an extracellular β-1,3-
exoglucanase in *Trichoderma harzianum*(Cohen-Kupiec et al. 1999), and later demonstrated
to be a carbohydrate binding module, similar to CBMs(S et al. 2019). The best-characterized
member in the Basidiomycota seems to be involved in beta-glucan binding/FCW remodeling
during plant-fungal interaction in *Serendipita indica*(Wawra et al. 2019). Whether this is true
for other WSC family members too, is not known, but is indicative of a potential role in
adhesion/remodeling at the cell surface.

The examined Agaricomycetes harbored 2-12 copies of WSC-domain encoding
genes. Of these, several showed strong induction in primordia and fruiting bodies in *A.*
ostoyae, C. aegerita, C. cinerea, L. bicolor, L. edodes, L. tigrinus, Ph. chrysosporium, P.
ostreatus and *S. commune* (but not in *A. ampla* and *M. kentingensis*). Genes with a
predicted WSC domain in their encoded proteins were typically primordium-upregulated in all
species. In *C. cinerea* these genes had high expression in early (proliferative) development
(hyphal knots to stage 2 primordia), followed by very low expression in most genes in late
developmental stages. The WSC domain can occur on its own or in combination with other
enzymes (most often GH71 and glyoxal oxidase) and uncharacterized domains, indicating
that it often fulfills a binding role in larger proteins. For example, WSC domain is present on
GH71 proteins that form a 10/8 orthogroup (represented by *C. cinerea* 464399), which we
discussed above.
3099

4.8.2.3. Miscellaneous cell wall remodeling (GH27, GH72, GH76, GH79, GH88, GT18,
pectinesterases)

In addition to chitin- and glucan-active families, several further CAZyme groups showed
widespread developmental regulation in fruiting bodies, for which the substrates in fruiting
bodies are less clear. These make very interesting groups indicating potentially novel cell
wall components, however, they are hard to place in context of fruiting body development as
of now.
GH79 and 88 - An orthogroup containing GH79 β-glucuronidases (*C. cinerea* protein ID:
470689) (Table 26), showed developmental expression in several species, consistent with
reports of Krizsan et al(Krizsán et al. 2019). Low amounts of glucuronic acid was reported in
N. crassa cell walls(Verdín et al. 2019), but the role of GH79 enzymes in cell wall
biosynthesis or remodeling are unknown. On a related note, an orthogroup of GH88
glucuronyl hydrolases was detected (*C. cinerea* protein ID: 384105)(Table 26). GH88 genes
are generally linked to pectin degradation by wood-decay fungi(van den Brink and de Vries
2011); their potential role in fruiting bodies is not known.
GH27 - Similarly, two CDE orthogroups (represented by Armosto1_254810 and
Armosto1_269656) of GH27 enzymes with alpha-galactosidase or α-*N*-
acetylgalactosaminidase activity were detected (Table 26). This family is discussed mostly in
the context of lignocellulose degradation(Kabel et al. 2017), to our knowledge it has not
been reported in fruiting body transcriptomes before. As a tendency, GH27 encoding genes
were often upregulated late in the development of the containing species. Their role in
fruiting body formation is not known.
GT18 - The GT18 family, which contains α-1,3(6)-mannosylglycoprotein β-1,6-N-acetyl-
glucosaminyltransferases showed a patchy distribution in the examined species, being
completely missing from the genomes of *C. cinerea* and *A. ostoyae*. However, it showed a
strong upregulation in primordia relative to vegetative mycelium and constant high
expression in fruiting bodies of *A. bisporus, A. ampla, S. commune, C. aegerita* and *P.*
ostreatus (not in *M. kentingensis and L. bicolor*)(Table 26, *S. commune* protein ID:
Schco3_2633734). This indicates fruiting body specific functions for this family, possibly in
chitin synthesis, although its exact role remains unknown. To our best knowledge, the role of
the GT18 family in modifying the fungal cell wall has not been described in any species.
GH72 - Contrary to our expectations, members of the GH72 family did not appear to be
developmentally relevant in Agaricomycetes. This family contains cell wall transglycosylases

that crosslink glycoproteins to cell wall components in the Ascomycota(Kar et al. 2019), and
therefore we expected it to be developmentally expressed in Basidiomycota as well. Their
expression is considerable but largely constant in Agaricomycetes fruiting bodies. This could
indicate a difference between Asco- and Basidiomycota cell wall composition, or that they
act during fruiting body development irrespective of their constant expression.
GH76 - In the Ascomycota, GH76 endo-1,6-alpha-mannosidases (Dfg5 and Dcw1 in *S.*
cerevisiae) are required for normal cell wall synthesis. Their exact role may not be known
with certainty, but they were hypothesized to be involved in the proper positioning of
glycosylphosphatidylinositol-anchored cell wall proteins(Orlean 2012). To our best
knowledge, the GH76 family has not yet been suggested to participate in fruiting body
development in the Agaricomycetes. It is present 0-10 copies in the Agaricomycetes and
shows predominantly mycelium-specific expression. We only found scattered expression
dynamics in fruiting bodies, such as a peak in stipes of *P. ostreatus*
(PleosPC15_2_1064904), but most genes are not developmentally regulated. Given that the
family is missing in some species completely and the predominantly vegetative mycelium-
specific expression, we hypothesize that the GH76 family does not participate, or only to a
small extent, in cell wall assembly in Agaricomycete fruiting bodies.
Pectinesterases - We detected a CDE orthogroup (represented by Schco3_2614029) that
contains genes generally linked to pectin degradation. It contains pectinesterases that are
developmentally regulated in 6/5 species (FC 2/4) (Table 26). Because pectin is not a known
cell wall component in Agaricomycetes, speculating about the function of these genes is
hard and may require more research.

4.8.2.4. Multicopper oxidases and cupredoxins

Laccases catalyze the oxidation of diverse phenolic and non-phenolic compounds, which
makes them suited to diverse (extra)cellular roles, from lignin degradation, pigment
production to dye decolorization or fruiting body related functions(P 2006). Their biochemical
properties provide laccases with significant potential in industrial applications. Despite very
intense research on laccases (mostly in the context of lignocellulose decomposition), we still
have very limited information on the precise role of laccases in fruiting body development.
Several reports of upregulation of laccase, or multicopper oxidase (MCO, the broader
family in which laccases belong) genes or laccase activity in fruiting bodies of
Agaricomycetes(Zhao and Kwan 1999; Ohga et al. 2000; Chen et al. 2003; Wang and Ng
2005; Madhavan et al. 2014; Sakamoto et al. 2015; Sipos et al. 2017b; Nagy et al. 2018) are
available (reviewed in detail by(Kües et al. 2011)). *Schizophyllum commune* was reported to
have 6 MCO genes (two of which are laccases), of which 3 were reported to be upregulated
in fruiting bodies(Madhavan et al. 2014) (Mco2, Mco3, Lcc1) by RT-PCR, which is consistent
with later RNA-Seq data(Krizsán et al. 2019). Of these, Lcc1 showed the highest similarity to
Fet3-type ferroxidases. Similarly, *P. ostreatus* laccases were reported to be expressed in a
developmental stage-specific manner, with Lacc5 and Lacc12 specifically expressed in
fruiting bodies and primordia, respectively, while other laccases had even or mycelium-
biased expression patterns(Park et al. 2015; Jiao et al. 2018). Stage-specific MCO
expression was reported in *L. bicolor* with specific laccase genes induced in fruiting bodies
(Lcc2), ectomycorrhizae and free-living mycelium(Courty et al. 2009). These studies have
linked laccase expression to two major functions, pigment production in either spores or
fruiting bodies and fungal cell wall modification or cell-cell adhesion(Kües et al. 2011).
Laccases were speculated to polymerize polyphenols in the intercellular spaces and thus
contribute to the binding of neighboring hyphae to each other. The hypothesis of laccases

being related to adhesion comes from Leatham and Stahmann(Leatham and Stahmann
1981), but, to our knowledge, was never experimentally verified. Although not relevant to
fruiting body formation, an interesting inverse relationship between mycelial laccase
expression and flushes of fruiting body development was reported in several species,
possibly related to stopping mycelium growth or nutrient acquisition when nutritional
competence for fruiting is achieved (reviewed in(Kües et al. 2011)).

Developmental studies indicated laccase function in fruiting body formation in
Pleurotus spp(Das et al. 1997). and *L. edodes*(Nakade et al. 2011). In *P. florida* laccase
gene silencing abolished fruiting body development and caused slower vegetative growth,
whereas in *L. edodes* aerial hyphae formation was also reduced and hyphae developed
thinner, non-layered cell walls upon the silencing of the Lcc1 laccase gene. Zhang et al
reported faster vegetative growth and earlier fruiting in laccase (Lcc1) overexpressing strains
of *H. marmoreus* compared to wild type strains(Zhang et al. 2015a), although no clear
hypothesis on how overexpression of a single structural gene could result in these
phenotypes was presented.

We here present annotations for multicopper oxidases, which include laccases,
ascorbate oxidases and ferroxidases(Kües et al. 2011). The 11 genomes of Agaricomycetes
examined here had 5-26 multicopper oxidases, a considerable proportion of which were
developmentally regulated, for example, 13 out of 17 in *C. cinerea* (Table 27). We found
three CDE laccase orthogroups; these comprised *C. cinerea* Lcc8, Lcc2 and their orthologs
as well as reciprocal best hits of the *S. cerevisiae* ferroxidase Fet3. [Of note, we detected
reciprocal best hits of the ferroxidase Fet3 in *C. cinerea*, whereas a previous report, based
on phylogenetic analyses did not identify Fet3-type MCOs in this species(Kües et al. 2011);
we note that this may arise from the limitations of our reciprocal best hit based search and
that accurate classification of MCOs requires the examination of specific catalytic residues.
Orthologs of *C. cinerea* Lcc2 (*C. cinerea* protein ID: 368135) are also involved in
morphogenesis in *L. edodes (as Lcc1)*, and differentially expressed during *Cryptococcus*
basidium formation(Liu et al. 2018; Merényi et al. 2021). We found that Lac4 from *Volvariella*
volvacea, which was implicated in fruiting body development(Chen et al. 2003) showed
highest similarity to the CDE orthogroup that comprised *L. edodes* Lcc1 (represented by *C.*
cinerea 368135). In the examined Agaricomycetes, this group of genes is upregulated in
caps (*P. ostreatus, C. cinerea*), stipes (*M. kentingensis*) or primordia (*A. ampla*), or shows
flat expression curves (other species). The detected ferroxidase orthogroup was present in 7
species, of which it was developmentally regulated in *A. bisporus, A. ostoyae, C. cinerea, L.*
edodes and *P. ostreatus* (FC>2). A Fet3-like gene was also upregulated in *S.*
commune(Madhavan et al. 2014), but this gene did not group into the detected orthogroup in
our analysis. Ferroxidases are membrane proteins that catalyze the oxidation of Fe^{2+} to Fe^{3+}
for iron uptake and have only weak laccase activities(Larrondo et al. 2003). Finally, *C.*
cinerea 440170 is the orthologs of *C. neoformans* Lac1, which has been shown to be
involved in melanin biosynthesis(Lee et al. 2019). This gene shows a strong expression
peak in young fruiting body gills, suggesting that it is involved in producing spore pigments in
C. cinerea.

In summary, experimental and gene expression evidence on the role of multicopper
oxidases in fruiting body development is abundant and -omics data reveal marked
developmental stage-specific expression of laccases. Hypotheses on their role either in
pigment production or cell wall crosslinking/adhesion have been put forth, however, the
available data do not allow us at the moment to select between these hypotheses, until more
developmental studies and gene knockout phenotypes become available.

In addition to *bona fide* MCOs, we identified four CDE orthogroup of proteins which
belong to the Cupredoxin family, like laccases, but do not contain the typical multicopper
oxidase domain. These might be highly diverged MCO genes or distant relatives of true
MCOs with unknown functions. The five orthogroups are conserved across Agaricomycetes
(*C. cinerea* protein ID: 356026, 451850, 441442, 353997, 497434) (Table 27). One of these
(*C. cinerea* protein ID: 356026) is FB-init is *A. ostoyae, C. cinerea* and *P. ostreatus* (not in *A.*
ampla, S. commune and *M. kentingensis*), whereas a second (*C. cinerea* protein ID:
451850) was characteristically late upregulated in most species (*A. ampla, M. kentingensis,*
P. ostreatus, Pt. gracilis). Currently, it is impossible to speculate about the role of these
proteins in fruiting body development.
3238

4.8.2.5. Expansins and cerato-platanins
Fungal expansins (or expansin-like proteins) show similarity to plant expansins, proteins that
modify the plant cell wall but do not show hydrolytic activity towards its polymers. Rather,
they exert their action by loosening the cell wall structure, hence the name loosenins(Quiroz-
Castañeda et al. 2011) and swollenins(M et al. 2002a) for some members of the family.
Fungal expansins with activities towards either the plant or fungal cell wall have been
reported. Certain fungal expansins can bind chitin and facilitating the action of hydrolytic
enzymes by separating cell wall chitin microfibrils. Two experimentally characterized
Basidiomycota expansins are Loos1 from *Bjerkandera adusta*(Quiroz-Castañeda et al. 2011)
and ScExlx1 from *S. commune*(Tovar-Herrera et al. 2015). Both were shown to bind chitin
and facilitate the action of chitinases, therefore, it was concluded that they modify, but not
hydrolyze the chitin polymer structure. Recent studies suggested that expansin-like proteins
are involved in stipe cell elongation in *F. velutipes*(Q et al. 2018) and *C. cinerea*(Niu et al.
2015) and were shown to have a role in stipe elongation, by loosening the chitin polymer
structure (reviewed by(Liu et al. 2021)).
Several expansins show expression patterns compatible with a role in fruiting body
development in previous transcriptome-based studies(Barsottini et al. 2013; Sipos et al.
2017b; Q et al. 2018; Almási et al. 2019; Krizsán et al. 2019). Based on analyses of 201
fungal genomes, Krizsan et al reported that expansins are enriched in the genomes of
Agaricomycetes(Krizsán et al. 2019). The examined mushroom-forming fungi possess 9-27
expansin genes, with *C. cinerea* having 13. Given the dual specificity of expansins, it is likely
that several of these are involved in cellulose degradation. However, a significant proportion
(19-85%) of expansin genes are developmentally regulated in fruiting bodies, indicating roles
in morphogenesis too. Expansins are a highly volatile group of genes that did not group into
conserved orthogroups in our analysis (except two CDE orthogroups, see below), therefore,
comparative analysis of their expression patterns across species is hardly possible. Within
C. cinerea expansins show diverse expression peaks, in gills, cap or stipes with some genes
highest expressed in vegetative mycelia too (Fig. 11). This could indicate tissue- and stage-
specific roles for expansins in fruiting body formation. For example, four genes had a strong
induction in stipes (*C. cinerea* protein ID: 451456, 464389, 359952, 461698), which is
consistent with previous reports of their participation in stipe elongation(Liu et al. 2021).
Three *C. cinerea* genes showed significant increase in expression from the vegetative
mycelium to hyphal knots (*C. cinerea* protein ID: 461698, 286849, 496836). Despite the
volatility of the family, we detected three putative expansin orthogroups (represented by *C.*
cinerea 496836, 467907 and 461698) which were developmentally regulated in 12/10, 11/10
and 10/5 species (at FC>2/4) (Table 28).

Cerato-platanins - Although initially thought to be related to hydrophobins, cerato-
platanins share a common origin with expansins(Barsottini et al. 2013; Gaderer et al. 2014;
Baccelli 2015; Luti et al. 2020). First described from the ascomycete *Ceratocystis platani*,
they were initially and generally considered to be related to pathogenesis(Gaderer et al.
2014; Baccelli 2015; Chen et al. 2017; Luti et al. 2020), however, there is piling evidence, so
far only from transcriptome studies, that they might be important in fruiting body
development as well. In such studies, cerato-platanin genes showed extensive expression
dynamics in fruiting bodies, suggesting an activity on the fungal cell wall(Plaza et al. 2014b;
Sipos et al. 2017b; Almási et al. 2019; Krizsán et al. 2019; Liu et al. 2020b), or in
defense(Kunzler 2015). Structure-based biochemistry studies in *Moniliophthora perniciosa*
showed that certain cerato-platanins are able to bind chitin and are active on the fungal cell
wall(Barsottini et al. 2013). Like expansins, cerato-platanins are strongly enriched in the
genomes of mushroom-forming fungi(Chen et al. 2017; Krizsán et al. 2019). There is,
unfortunately, little information on their mechanistic role in fruiting body development, except
a the study by de Barsottini(Barsottini et al. 2013), which provides proof that cerato-platanins
can be active on chitin and the fungal cell wall, providing a biochemical link to their potential
role in fruiting body development.
Our analyses identified 1-8 cerato-platanins in the genomes of examined mushroom-
forming fungi, which is in line with copy numbers reported recently(Chen et al. 2017). Like
expansins, cerato-platanins are quite volatile and we identified only two conserved CDE
orthogroups (*C. cinerea* protein ID: 439071, 544218) (Table 28), with most genes scattered
in species-specific or smaller groups. The first showed opposing expression patterns in
different species: in *C. cinerea* (protein ID: 544218) it was upregulated in caps relative to
stipes, whereas in *A. ostoyae* and *M. kentingensis* it was upregulated in stipes relative to
caps. In *P. ostreatus* and *M. kentingensis* it showed the highest expression in vegetative
mycelia. In *A. ampla* and *S. commune* it showed a very strong induction (>1000-fold) in
mature fruiting bodies.
Expression data across species indicates that developmentally dynamic cerato-
platanin expression was observed only in certain species, while not in others. For example,
no developmentally regulated cerato-platanins were found in *Ph. chrysosporium, A.*
bisporus, only a single in *M. kentingensis*, but several in *A. ostoyae, C. cinerea, L. tigrinus,*
etc. This indicates that developmental expression of cerato-platanins may be a species-
specific, rather than a conserved feature in mushroom-forming fungi.
3308

4.8.3. Miscellaneous cell wall related families (HTP, DyP, AA3, AA5, GH76, GT8,
Slt2, Cfs1, Eln3, peptidases)
We identified several CDE gene families which are potentially linked to the fungal cell wall,
however, evidence is lacking on their mechanistic roles and/or they could not be confidently
ascribed to either chitin or glucan, and/or are not enzymatic proteins. These are discussed in
this chapter, with noting that the widely observed developmental expression makes these
protein families interesting, however, their assignment to the cell wall is preliminary.
HTPs - Heme-thiolate peroxidases (HTPs) are fungal-specific proteins that were first
described in *C. aegerita* (as haloperoxidase) and use H_2O_2 to modify substrates(R et al.
2004). While their enzymatic properties have been quite thoroughly studied(M et al. 2015a),
their biological roles are poorly known. A remarkable expansion of the family was noted in *A.*
bisporus which, combined with the upregulation of HTPs in *Agaricus* substrates led to the

hypothesis that HTPs are linked to the litter decomposer lifestyle(Morin et al. 2012). In the
context of fruiting bodies, HTPs were reported only in *A. ostoyae*(Sipos et al. 2017b) so far.
We detected two CDE orthogroups of HTPs (represented by *C. cinerea* 479688,
442333)(Table 29) as well as several other species-specific developmentally regulated HTP
genes. The examined Agaricomycetes have 2-10 genes with an expansion to 23 genes in
both *M. kentingensis* and *A. bisporus* as reported earlier(Morin et al. 2012). Most of the
species had at least one HTP encoding gene that is strongly upregulated (up to 50-fold) from
vegetative mycelium to the first primordium stage. These observations establish HTPs as a
potentially important morphogenetic gene family in fruiting body development with currently
unknown function. We tentatively discuss this gene family under the chapter on cell wall
modification, however, it should be mentioned that this is based on an assumption of a
potential role of this family.
DyPs - Dye decolorizing peroxidases (DyPs) are a group of fungal peroxidases that belong
to the broader heme-containing peroxidase family. They are mostly discussed in the context
of plant cell wall degradation by fungi(Floudas et al. 2012); to our knowledge, this is the first
report of its potential role in fruiting body morphogenesis. We detected 0-5 copies (missing in
A. ampla, S. commune and *A. bisporus*) and a patchy distribution in Agaricomycetes.
However, in species that have DyP genes, these are often developmentally expressed. DyP
genes show significant upregulation from vegetative mycelium to the first primordium stages
in *A. ostoyae, C. cinerea, C. aegerita, L. bicolor* and *P. ostreatus*. In *M. kentingensis* all DyP
genes show a marked stipe expression peak, whereas in *Pt. gracilis*, expression curves are
flat. DyP genes grouped into a single CDE orthogroup (represented by *C. cinerea* 407604,
Table 29). While some of the copy number variation is explained by the lignin-degrading
ability of fungi(Almási et al. 2019), the developmental expression of DyP genes suggests
morphogenetic roles as well, possibly in remodeling the cell wall, although this is based on
assumptions distilled from the general activity of the family. It would be interesting to
investigate if, similar to observations during lignin degradation, H_2O_2-generating enzyme
gene expression is co-regulated with peroxidase expression patterns.
AA5_1 (GLOX) - Glyoxal oxidases (GLOX) in the AA5_1 family of auxiliary activities are
H_2O_2 generating enzymes that facilitate the oxidative cleavage of lignin by
peroxidases(Floudas et al. 2012). Surprisingly, GLOXs have also been reported to be
upregulated in fruiting bodies in 6 species of Agaricomycetes(Krizsán et al. 2019) and were
detected in fruiting bodies of *Auricularia polytricha* by protein mass spectrometry(Jia et al.
2017). Since lignin is obviously missing from fruiting bodies, they were speculated to
contribute to cell wall remodeling(Krizsán et al. 2019). We found 1-22 GLOX genes in the
examined Agaricomycetes, the fewest in the Schizophyllaceae (*A. ampla, S. commune*)
possibly because of their reduced morphology or reduced lignin-degrading ability(Almási et
al. 2019). Two CDE orthogroups (*C. cinerea* protein ID: 263775 and 440836 Table 29), and
several species-specific developmentally regulated GLOX genes were found in our data. We
found GLOX genes significantly upregulated in primordia relative to vegetative mycelium
(FB-init) in *A. ostoyae, A. ampla, C. cinerea, L. tigrinus, L. bicolor, S. commune, Ph.*
chrysosporium, P. ostreatus and *C. aegerita* (not in *M. kentingensis, L. edodes* and *A.*
bisporus). Two of the three *C. neoformans* GLOX genes also showed significant
upregulation during basidium development. These data indicate that GLOXs may be
important morphogenetic factors in the Basidiomycota.
AA3_1 - cellobiose dehydrogenases in the AA3_1 family (GMC oxidoreductases) are known
to donate electrons to LPMOs during lignocellulolysis in wood-decay fungi(Floudas et al.
2012). There are only two reports on the potential role of CDHs in fruiting body development,

one is a comparative transcriptomics study(Krizsán et al. 2019) and the other is a report of
the interaction between CDH and laccases in the formation of cinnabarinic acid, the red
fruiting body pigment of *Pycnoporus cinnabarinus*(Temp and Eggert 1999). A CDE
orthogroup of cellobiose dehydrogenase (CDH) genes (*C. cinerea* protein ID: 469460)(Table
29) was developmentally regulated in all species in which it was present. Additional CDH
genes (2-8 genes in Agaricomycetes) were developmentally regulated in a species-specific
manner in several species, although expression levels were low in many cases. The
biological significance of CDH in fruiting body development is currently unknown; based on
previous studies it may be related to cell wall remodeling or pigment production.
Slt2 kinase family - A serine-threonine kinase orthologous to the MAP kinase involved in
the cell wall integrity pathway of *S. cerevisiae* (Slt2) was detected as a CDE orthogroup (*C.*
cinerea protein ID: 413665). Slt2 is a terminal kinase in the pathway that phosphorylates
downstream transcription factors (Rlm1 and Swi4/6 in yeast)(Levin 2005). Although the gene
is developmentally regulated in several species (Table 29), expression patterns are not
consistent across taxa. Given the potentially pleiotropic role of this kinase, it is hard to guess
its role in fruiting bodies, but extrapolations from its role in *S. cerevisiae* would suggest links
to cell wall integrity, morphogenesis or the cell cycle.
Cfs1 family - The orthogroup containing *C. cinerea* Cfs1 (*C. cinerea* protein ID: 447925), a
cyclopropane fatty acid synthase, was developmentally regulated in 10/7 (FC>2/4) species
(Table 29). Cfs1 was identified in a mutant screen of strains with developmental defects; its
mutation leads to failures in the progression of hyphal knot development(Liu 2005) in *C.*
cinerea (stalled at primary hyphal knot). The corresponding gene is highly induced 1h after
blue light exposure(Sakamoto et al. 2018). We discuss this gene under the cell wall,
although some evidence suggests it may be more closely related to plasma membrane
assembly. Structurally, Cfs1 and related proteins are mycolic acid cyclopropane synthases,
a family that occurs in 1-6 copies in the examined Agaricomycetes. Their function is not
known, but presumably they are involved in the synthesis of long fatty acids (mycolic acid is
not known to occur in fungi). In most examined species (*A. ampla, M. kentingensis, C.*
cinerea, P. ostreatus, S. commune), the gene is upregulated in primordia relative to
vegetative mycelium, consistent with its phenotypic role in hyphal knot development.
Eln3 family - This family is based on the *Eln3* gene of *C. cinerea*, which was found in a
REMI mutagenesis screen. The mutant strain displayed defects in stipe elongation and
mutant stipe cells were disorganized and there were large gaps among stipe cells, whereas
in the wild type strain stipe cells align closely with each other(Liu 2005). It has been
hypothesized that *Eln3* is involved in cell-cell connections, possibly adhesion.
We found that *C. cinerea Eln3* (*C. cinerea* protein ID: 380322) formed a small CDE
orthogroup present in 7 species and was developmentally regulated in 6 species (Table 29).
In these species the gene was predominantly FB-init and stipe upregulated, in agreement
with a previously reported role in stipe elongation(Arima et al. 2004). It should be noted that,
in many species, upregulation in primordia is easy to confuse with stipe-specific
upregulation, because primordia often comprise predominantly stipe tissue. In general, we
found 7-8 genes the examined Agaricomycetes (only 4 in *A. bisporus*) with identical domain
composition to *Eln3*. This does not guarantee identical function, nevertheless is indicative of
the diversity of this family in mushrooms. These genes were often FB-init and showed stipe
or cap-specific upregulation depending on the gene and the species. For example, in *C.*
cinerea we found that 5 out of 6 genes were stipe-upregulated. In *A. ostoyae* we found four
cap- and two stipe-upregulated, whereas in *L. bicolor* two cap- and one stipe-upregulated
gene. The widespread developmental regulation of *Eln3* and its homologs suggest

morphogenetic functions, possibly in association with the cell wall. Based on its domain
composition, we hypothesize it represents a cell wall active, possibly new CAZyme family
not yet recognized by Cazy.org.
GT8 - The GT8 family has 1-7 (most frequently 5) copies in the Agaricomycetes. The GT8
family includes both N-acetylglucosaminyltransferases linked to the modification of N-linked
glycans in the Golgi, and glycogenins involved in glycogen metabolism. Buser et al
hypothesized that fruiting body-specific N-glycans generated by GT8 enzymes may
contribute to adhesion in *C. cinerea*(Buser et al. 2010a). GT8 genes formed one CDE
orthogroup, represented by *C. cinerea* 461263 FC2/4: 7/3 species), which is closest to *S.*
cerevisiae Glg1 glycogenin (discussed under carbohydrate metabolism). Of the other GT8
genes *C. cinerea* (*C. cinerea protein ID:* 453097, =*CcGnt1)*(Buser et al. 2010a)*, A. ostoyae,*
L. tigrinus, P. ostreatus, A. ampla and *S. commune* had one FB-init gene each, however,
these did not form CDE orthogroups. CcGnt2 (*C. cinerea* protein ID: 412828) was not
expressed. Other species did not have developmentally regulated genes. Due to the
variability of the family, it is hard to determine whether these are closer to glycogenins or N-
acetylglucosaminyltransferases.
N-glycosylation - Mannosylation and N-glycosylation of proteins in the ER is a key process
for cell wall synthesis. Berends et al provided a detailed view of the N-glycosylation
machinery in *S. commune, C. cinerea* and *L. bicolor* based on orthology to the *S. cerevisiae*
N-glycosylation pathway members and expression data in *S. commune* fruiting bodies and
monokaryotic mycelium(Berends et al. 2009). They conclude that Agaricomycetes have the
gene repertoire to produce oligomannose structures only, in contrast to some Ascomycota
which can produce very complex modifications to proteins. Our expression results broadly
agree with theirs in that we detected very low dynamics of corresponding genes (not shown)
in the fruiting body transcriptomes analyzed here.
3442

3443 4.9. Cell surface and cell wall proteins

Secreted cell surface proteins fulfill diverse functions in the life of fungi, ranging from
signaling, interactions with the environment, adhesion, cell wall modifications, but also
communication with hosts in mutualistic or pathogenic interactions(Pellegrin et al. 2015;
Essen et al. 2020). The fungal cell surface proteome and secretome profile of each species
in a given environment is unique, making constituents of fungal communities recognizable by
their extracellular protein profiles(Campbell et al. 2015). Secreted proteins have been the
subject of intense research, mostly in the context of interactions between fungi and other
organisms, adhesion (e.g. via GPI-anchored adhesins) of fungal cells to various
surfaces(Essen et al. 2020; M et al. 2020b). Fruiting bodies, as environmentally exposed, 3-
dimensional multicellular structures, can be expected to display unique cell-surface
attributes. In fact, cell surface proteins include some of the best-known fruiting body-
associated proteins (e.g. hydrophobins, lectins, see below) or proteins related to defense
(see these in a separate chapter). Conceivably, adhesion and/or cell-to-cell communication,
two of the most important multicellular functions(Knoll 2011), could be realized via the action
of secreted cell surface proteins in fruiting bodies, to just name a few of the diverse ways
mushroom-forming fungi might make use of their cell surface protein repertoire. A significant
open question in fruiting body biology is the nature and role of the extracellular matrix and
the biochemical and communication events that take place in the interhyphal space. Cell

surface proteins are undoubtedly important players in these processes, and therefore
represent interesting targets of future research.
While some cell surface proteins are the best examined fruiting body proteins (e.g.
hydrophobins), the following inventory of conserved developmentally expressed genes
shows that there is plenty of new insight to be learned about the cell surface proteome of
Agaricomycetes fruiting bodies. In this chapter we discuss cell surface protein families in the
context of CDE orthogroups detected in our meta-analysis of fruiting body transcriptomes.
3469

4.9.1. Hydrophobins

Hydrophobins are some of the best-known proteins in fruiting bodies. They confer
hydrophobicity to hyphae, which they achieve by self-assembling into a water-repellent
rodlet layer on the cell surface. Hydrophobins were hypothesized to line air channels in
fruiting bodies(Lugones et al. 1999b), but also contribute to water sealing, which is important
in fruiting bodies as they rely on water transported to them from the vegetative
mycelium(Nehls and Dietz 2014). The family has extensively literature, in general(Wösten
2003; Linder et al. 2005; Bayry et al. 2012), in relation to fruiting body development(Ohm et
al. 2010; Kües and Navarro-González 2015; Kim et al. 2016; Sammer et al. 2016; Y et al.
2019) or on potential uses in bioengineering(Wösten and Scholtmeijer 2015). We therefore
concentrate on novel aspects detected in our data that have not or only rarely been reported
before.
Basidiomycota hydrophobins mostly belong to class I hydrophobins(Sammer et al.
2016). On average 11-44 hydrophobin genes (only 6 in *M. kentingensis*) were detected in
the examined Agaricomycetes, a significant proportion of which were developmentally
regulated. They can be arranged into distinct groups based on expression patterns. Previous
studies reported a near complete turnover of hydrophobin genes from vegetative mycelium
to fruiting body, indicating that two distinct sets of hydrophobins exist(Ma et al. 2007; WW et
al. 2008; Cheng et al. 2013a). We also observed that mycelium-expressed and fruiting body-
expressed hydrophobins form distinct groups. For example, the numbers of hydrophobins
showing mycelium- vs fruiting body specific expression was 6/5, 1/9, 10/4, 23/3 and 5/20 (at
fold change 4/2) in *Ph. chrysosporium, S. commune, P. ostreatus, A. ostoyae* and *C.*
cinerea, respectively. *A. ostoyae* further has two rhizomorph-specifically expressed
hydrophobins, indicating morphogenesis-related diversification of the family. This is in line
with observations in *F. velutipes,* where most hydrophobin genes were found to be
primordium-upregulated(Kim et al. 2016). We also detected hydrophobin genes expressed
during early development, but not later, indicating functions linked to primordium
development, which are turned off during later development. These observations further
substantiate the functional diversification, probably driven by morphogenesis and complexity.
Despite the large diversity and developmental expression of hydrophobins, only 2
CDE orthogroups were detected (*C. cinerea* protein ID: 99661, 419467) (Table 30), which
were present in only a subset (11 and 8 species, respectively) of the examined species. This
is due to the frequent duplication of hydrophobin genes(Mgbeahuruike et al. 2017), which
quickly erases clear orthology relationships. CDE orthogroups did not show conserved
expression patterns. Interestingly, *M. kentingensis* has only 6 hydrophobin genes, none of
which were developmentally expressed in our reanalysis of data from Ke et al(Ke et al.
2020a). A possible explanation for this is that this species produces hygrophanous, rather
than hydrophobic fruiting bodies, which absorb rather than repel water. The relatively low

number of developmentally regulated hydrophobin genes in *A. ostoyae* may also be
explained by the hygrophanous caps of this species. Hygrophanicity is a widespread trait in
the Agaricales and might explain some of the differences in hydrophobin diversity and
expression.
3512   Taken together, hydrophobins, as already established by previous research, are one
3513   of the best-known fruiting body-related structural cell surface proteins in the Agaricomycetes,
3514   with almost all species expressing developmental stage-specific suites of hydrophobin
3515   genes.
3516

4.9.2. Lectins

3517
Lectins will be also discussed in Chapter 10 on defense-related proteins, however, a role in
defense has been proven for only a subset of the diverse lectin repertoires of mushroom-
forming fungi; the remaining non-defense, or functionally unknown lectins are discussed in
this chapter (even though not all lectins may be secreted or their secretion status is unclear
at the moment, (Boulianne et al. 2000)). Lectins are proteins that specifically recognize and
bind various carbohydrate moieties, mostly on the cell surface. This ability makes them
suitable for multiple biological roles, from bacterial agglutination and cytotoxicity against
mushroom grazers(Bleuler-MartÍnez et al. 2011; Kunzler 2015) to carbohydrate binding
modules that fulfill roles related to the fungal cell wall (e.g. remodeling), adhesion, cell-to-cell
communication or other functions in the extracellular matrix. Their carbohydrate binding
ability makes them interesting in a wide range of biomedical fields(Hassan et al. 2015),
including cancer research, which led to the broader lectin family rapidly expanding.
Mushroom fruiting bodies remain one of the richest sources of lectins for applied research.
However, regarding their physiological role for the fungi, boundaries between defense-
related and other lectins are far from clear at the moment(Bleuler-MartÍnez et al. 2011).
Due to interest in their practical applications, there is often more information on the structure
of mushroom lectins than on their roles in the fungus' life. Fruiting bodies express a diverse
set of lectins (reviewed by(A et al. 2013; Kobayashi et al. 2014; Y and H 2014)) and several
of the well-known lectin families were first isolated from mushrooms. For example, the fungal
fruit body lectin family was established based on the *Xerocomus chrysenteron* XCL protein,
which was suggested to have insecticidal activity(C et al. 2004). Besides interest in their
biotechnological applications, lectins have been in the focus of research on fruiting bodies.
Early reports identified specific hyphal knot and primordium upregulated galectins (Cgl1 and
Cgl2) in *C. cinerea*(Boulianne et al. 2000; Bertossa et al. 2004), which were initially
postulated to be cell-cell adhesive or communication related molecules that may bind
basidiolipids(Walser et al. 2005). Simultaneous silencing of the *cgl1* and *cgl2* genes did,
however, not disturb fruiting, arguing against his hypothesis(MA et al. 2006). The lectins
were rather shown to inhibit the development of insect and nematode larvae, in nematodes
via binding to a specific N-glycan epitope, suggesting a function in defense against
predators(Butschi et al. 2010; Bleuler-MartÍnez et al. 2011). Both genes were found
upregulated in primordia, relative to vegetative mycelium in recent RNA-Seq based studies
(cgl1 = *C. cinerea* 473274, cgl2 = *C. cinerea* 488611). Later, Wälti et al described cgl3 (=*C.*
cinerea 539794), which resembles galectins, but binds chitooligosaccharides and had no
effect on larval development of insects and nematodes(Wälti et al. 2008). Beta-glucan
binding lectins of *Serepindita indica* may be involved in fungal cell wall remodeling at the
plant-fungal interface(Wawra et al. 2019). Several other recent reports of individual lectins

from a diverse set of medicinal or edible mushrooms is available but their detailed discussion
is beyond the scope of this paper. Lectins are also involved in Ascomycota fruiting body
(perithecium) formation and interestingly, the gene recently described in *Podospora anserina*
showed highest similarity to *Xerocomus* lectins(Nowrousian and Cebula 2005). Several
lectin families are overrepresented in mushroom-forming fungi(Krizsán et al. 2019),
suggesting key roles in their life cycle.
'Lectomes' evolve fast (like other defense genes) both in terms of sequence and
gene content. Most lectin families a phylogenetically patchily distributed in mushroom
forming fungi, with the probable sole exception of ricin-B lectins, which occur in all examined
species. It has been suggested that the fast gene turnover can contribute to the expression
of defense gene cocktails by the fungus, which is advantageous and that horizontal gene
transfer from bacteria can provide a mechanistic explanation for the phylogenetic patchiness
of several lectin families(Moran et al. 2012; Kunzler 2015).
We here provide an overview of lectin copy numbers in the examined
Agaricomycetes in the context of 8 main families, but note that more recent
classifications(Lebreton et al. 2021) significantly reorganized lectin families and may provide
a more nuanced view on lectin repertoires. Lectin copy numbers and expression patterns
were summarized in the context of 6 Agaricomycetes species by Krizsan et al(Krizsán et al.
2019) with lectins showing the RicinB fold (which includes protease inhibitors too, see above
and(J et al. 2016)) being most diverse and present in all species, whereas other families are
more patchy. Nevertheless, all lectin families had developmentally regulated genes, often
showing strong induction in primordium stages.
As noted earlier (e.g. (Kunzler 2015)), lectins are poorly conserved and hence mostly
did not form conserved orthogroups in our analysis. Only two CDE orthogroups, all
belonging to the RicinB family were detected (*C. cinerea* protein ID: 495335, 353931)(Table
31). Both orthogroups contained secreted proteins. Expression patterns were poorly
conserved within these orthogroups, but we note that each had FB-init genes in several
species, indicating that these secreted Ricin-B family members are involved in the initiation
of fruiting body development.

4.9.3. Other cell surface proteins (HsbA, Con6, PriA, AA7, CFEM, wax synthases,
fasciclins, peptidases, intradiol ring cleavage dioxygenase)

The HsbA family comprises poorly known cell wall galactomannoproteins that includes
HsbA from *Aspergillus oryzae*(S et al. 2006b), a *Magnaporthe* homolog(Soanes et al. 2012)
as well as genes named Mp1 in *Talaromyces marneffei*(Cao et al. 1998) and in *Aspergillus
fumigatus*(KY et al. 2001). Detailed functional information is not available for either of these
genes. The family contains GPI-anchored cell surface proteins that can bind to various
surfaces and was circumscribed in the context of appressorium attachment and
lignocellulose degradation in Ascomycetes. The common denominator between these
circumstances may be the need for adhesion of hyphae to various surfaces, enabling
secreted lytic enzymes to exert their action. In the context of fruiting bodies, the family was
first reported in fruiting bodies of *A. ostoyae*(Sipos et al. 2017b). In our current analysis, we
detected 0-3 members of the family in Agaricomycetes (missing from *L. tigrinus, Ph.
chrysosporium, A. ampla. S. commune*), all of which were developmentally expressed. The
family did not form CDE orthogroups. Expression of members of this family showed
significant dynamics (>100-fold), mostly with gill- or cap-specific expression peaks late
during development. Functional analysis of this gene family in mushroom-forming fungi

would be very interesting. If, as speculated above, the family is related to adhesion, then it
may provide a missing link to adhesion in complex multicellularity in mushrooms.
The Con6 protein family include small hydrophilic peptides that were first discovered as
conidium-specific proteins in *N. crassa* and later shown in *A. nidulans* (ConF and ConJ) to
be responsible for desiccation resistance of asexual spores(Suzuki et al. 2013). The family
also includes the meiotically expressed *Sch. pombe* gene *cum1*. The Con6 family is
conserved in Agaricomycetes, with 1-7 copies in the examined genomes, most of which are
developmentally regulated. They showed two major expression patterns in all species: (i)
strong induction in mature gills or combined gill+cap samples and/or (ii) upregulation in
primordia relative to vegetative mycelia (Fig. 12, Supplementary Fig. 8). For some genes
both for others only one of these expression patterns were observable. The family seems
poorly conserved, nevertheless, they formed three CDE orthogroups (represented by *C.*
cinerea 470234, 186809, *S. commune* 2624349)(Table 32). The gill-specific upregulation,
combined with the conidium-specific expression of the gene family in filamentous
Ascomycota, suggests that Con6 proteins may be involved in producing a spore-specific
protein coating in these species. The orthogroup represented by *C. cinerea* 186809 showed
upregulation in primordia of most species and has low expression in vegetative mycelia and
later developmental stages. The strong upregulation of several Con6-encoding genes in
primordia relative to vegetative mycelia in virtually all species suggests that it has general
roles in fruiting bodies as well, possibly in regulating hydrophobicity/hydrophilicity of the cell
surface, although this requires further research.
Wax synthases are long-chain alcohol O-fatty-acyltransferases that synthesize wax esters,
a typical constituent of the cuticula of plants or the sebum of the animals(Li et al. 2008). To
our best knowledge, only Krizsan et al(Krizsán et al. 2019) indicated developmental
expression for this family previously.
We detected 4-16 wax synthases in the examined Agaricomycetes, the fewest in *A.*
bisporus, but found no CDE orthogroups, suggesting a high duplication rate in the family.
However, each species (except *Pt. gracilis* and *A. bisporus*) had 1-2 FB-init wax synthase
genes, which suggests fruiting body specific roles for these genes. We observed
considerable (>100-fold) upregulation compared to vegetative mycelia in many cases (Fig.
13, Supplementary Fig. 9). Given the widespread upregulation of wax synthases in
mushroom-forming fungi, it is possible that they produce a wax-like layer on the surface of
fruiting body cells. On plants, wax-coating of leaves serves as a defense system that inhibits
penetration of pathogenic fungi and water-sealing(Lewandowska et al. 2020). Products of
fungal wax synthases, although unknown at the moment, could have similar properties in
fruiting bodies, worth investigating in the future.
PriA family - this family comprises secreted cell surface proteins first described as a
primordium-induced protein in *L. edodes* (hence its name)(S et al. 1992). The gene has
been quite extensively studied in *L. edodes*(T and K 2000; T et al. 2004a)*,* where a link to
zinc homeostasis has been suggested. It was also detected as developmentally regulated in
F. velutipes(Liu et al. 2020b)*, P. eryngii*(Fu et al. 2017), *L. edodes*(S et al. 1992) and in six
species by Krizsan et al(Krizsán et al. 2019). The family also includes the virulence-related
secreted adhesin Cfl1 of *C. neoformans*, which is highly induced during sexual development
and has been thoroughly researched in *C. neoformans*(Wang et al. 2012, 2013). Cfl1 has
been suggested to act as a morphogen that induces neighboring cells to phenocopy the
producing cells' morphology(Wang et al. 2013). Cfl1 has 4 homologs in *C. neoformans*,
which have stage-specific expression including Dha1, which is primarily expressed in
basidia(Gyawali et al. 2017). The family is present in 0-4 copies in the Agaricomycetes and

formed two CDE orthogroups. The first of these (*C. cinerea* protein ID: 359866)(Table 32)
showed a very conserved, mycelium biased expression; it was highly expressed in
vegetative mycelia and the first primordium stage, then virtually unexpressed in all later
stages in *A. ampla S. commune, M. kentingensis* and *P. ostreatus*. In *A. ostoyae* expression
dropped immediately after the mycelium sample. The second CDE orthogroup (represented
by *C. cinerea* 419688) contains best hits of *C. neoformans* Dha1 and showed a peak
expression in gills and caps of *C. cinerea, A. bisporus, C. aegerita*, *L. edodes* and late
stages of *S. commune* and *A. ampla.* At the same time, these genes were significantly
upregulated (up to 50-100x) in primordia relative to vegetative mycelium. In *L. edodes*, our
RNA-Seq based observations are consistent with the first reports of the PriA gene as a
primordium-upregulated one(S et al. 1992). The third orthogroup (represented by *S.*
commune 2681702), was present in only 8 species, and was FB-init in *A. ampla, A. ostoyae,*
P. ostreatus, M. kentingensis and *S. commune* (Table 32). In summary, the PriA family
appears to be a very interesting family from the perspective of fruiting body development. In
C. neoformans adhesive and signaling roles have been suggested for the family, although
exact mechanisms of function are still unknown.
Aspartic peptidase A1 - This family is a versatile group of endopeptidases that has
undergone a characteristic expansion in fungi(Revuelta et al. 2014). They were reported to
be involved in pathogenesis, nutrition, cell wall maintenance and a range of other functions.
The best researched family of A1 aspartic peptidases are yapsins, which are involved in cell
wall assembly and remodeling, although their function is not clear; limited evidence suggests
they proteolytically process secreted (cell wall) proteins(Gagnon-Arsenault et al. 2006; Gow
et al. 2017; M et al. 2021). Based on functional knowledge on yapsins, and the abundance
of GPI-anchored A1 aspartic peptidases in fungi, we discuss this family in the cell surface
section, despite almost certainly not all of them are related to the cell wall. The family has
been reported to be upregulated in fruiting bodies by several studies previously(Rahmad et
al. 2014; Song et al. 2018b; Krizsán et al. 2019). Aspartic peptidase A1 family was found as
15-60 copies in Agaricomycetes genomes. We detected 3 CDE orthogroups and several
species-specifically developmentally expressed genes (Table 32), though these did not
display strong expression conservation in the examined species. Nevertheless, the
developmental expression of several members of this family, combined with evidence on
yeast yapsins(Gagnon-Arsenault et al. 2006) and the secreted nature of these proteins
suggest a cell wall related role for aspartic peptidases which may be worth investigating. Of
note, we detected further CDE orthogroups of various peptidase families (M20, M28, M16,
S10), which may also be involved in fruiting body development (Table 32), but our functional
knowledge at the moment is too limited to speculate about their roles. Other peptidase
families that were reported in mushroom-forming fungi include the metalloprotease M43
family, which was reported from *P. ostreatus*(Joh et al. 2004) and has a strong (>700x)
induction upon fruiting in the data published by Merenyi et al(Merényi et al. 2021).
AA7 family - One of the CDE orthogroups (represented by *C. cinerea* 501135)(Table 32)
were top hits of *Fusarium graminearum* ChitO, the only experimentally examined member of
the family known to us(Heuts et al. 2007), which oxidises chitooligosaccharide species. AA7
comprises oligosaccharide oxidising flavo-enzymes which also contain a Berberine-like
domain and produce H_2O_2 during their action. They have been speculated to be involved in
the detoxification of lignocellulose degradation intermediates and in the production of various
secondary metabolites(Levasseur et al. 2013). Based on the activity of their homologs and
their expression in fruiting bodies, it is likely that these AA7 family members are active on
chitin, although their exact roles remain obscure for now.

CFEM domain proteins - We detected 2 CDE orthogroups of CFEM domain containing
proteins (Table 32). Homologous CFEM domain proteins may be involved in defense,
pathogenicity, host-pathogen interactions or iron uptake, as is the rbt5 family of *Candida*
spp.(Gaurav Bairwa et al. 2017) In *C. cinerea* certain mycelium expressed CFEM domain
proteins were reported to be involved in defense(Plaza et al. 2014b). The detected CDE
orthogroups comprise cell surface proteins with unknown functions, though they might be
involved in defense, however, evidence for that is missing at the moment.
Fasciclins are adhesive cell surface proteins that facilitate the attachment of cells to various
surfaces. The few fungal mentions include a report of a role in the adhesion of *Magnaporthe*
oryzae to host surfaces(Liu et al. 2009) as well as a primordium-upregulated gene in *L.*
edodes (*Le.flp1*)(Y et al. 2007). Recently, fasciclins were hypothesized to have been co-
opted for fruiting body development independently in both the Agarico- and
Pezizomycotina(Merényi et al. 2020b). We identified 2-8 fasciclin-encoding genes in the
examined Agaricomycetes. All examined species have developmentally expressed fascicling
genes, although these formed only small CDE orthogroups (Table 32). Several genes
showed an upregulation in primordia relative to vegetative mycelium (as reported in *L.*
edodes), suggesting that this family has widespread roles in mushroom formation, possibly
by facilitating adhesion. We found that the sequence deposited by(Y et al. 2007) (Accession
number: AB287432) as well as ascomycete homologs to which they analyzed are
cupredoxins, not fasciclins, leaving doubts as to the identity of Le.flp1. Nevertheless, this
does not undermine the observation that fasciclin encoding genes are widespread
developmentally expressed genes in mushroom-forming fungi, probably with roles in fruiting
body development.
Intradiol ring cleavage dioxygenases catalyze a ring-opening step in the degradation of
aromatic compounds, and were hypothesized to be involved in xenobiotic detoxification or
the utilization of aromatic compounds as carbon sources(Semana and Powlowski 2019).
However, the physiological role of intradiol ring cleavage dioxygenases in fungi is not known.
The only previous mention of a potential relationship to fruiting bodies reported
developmental regulation in 4 out of 6 species(Krizsán et al. 2019). We identified 2-7 copies
(28 in *A. ostoyae*) in the examined Agaricomycetes, with 1-3 developmentally expressed
genes in most of the species that, however, did not form conserved orthogroups. Members
of the family showed a significant upregulation in primordia relative to vegetative mycelium in
A. ampla, S. commune, C. cinerea, A. ostoyae as well as a strong upregulation (>100fold)
during late development of *C. neoformans*. Two genes show a very distinct upregulation in
young fruiting body caps of *C. cinerea*. No developmentally expressed genes were found in
Ph. chrysosporium and *M. kentingensis*. In summary, intradiol ring cleavage dioxygenases
might be important genes during fruiting body development of several species; they might
degrade aromatic compounds during fruiting, however, the identity of those compounds are
unknown at the moment.
3735

3736 4.10. Defense

Agaricomycete fruiting bodies (mushrooms) are, due to the concentration of biomass,
attractive targets of predators, parasites and pathogens. Accordingly, these structures are
known to be subject to grazing by slugs, to infestations with larvae of phorid and sciarid flies
and to infections by bacteria and molds(Wood et al. 2004; Krivosheina 2008; Lakkireddy et
al. 2020; M et al. 2020a). Since mushrooms are the sexual reproduction organs and a

considerable investment in terms of resources and energy, Agaricomycetes have evolved
several strategies for the protection and defense of these structures(Künzler 2018). Such
strategies include physical barriers(Varga et al. 2021) and chemical defense i.e. the
production of molecules (toxins) impairing the growth, development and/or viability of
antagonists(P 2008; FY and NP 2014; Stadler and Hoffmeister 2015).
Using saprobic Agaricomycete species, whose sexual cycles can be completed
under laboratory conditions, it could be demonstrated that the production of these toxins
occurs axenically i.e. in the absence of the antagonist(Plaza et al. 2014b; Tayyrov et al.
2019). This axenic toxin production is spatiotemporally regulated in that the toxin-encoding
genes are expressed solely during fruiting body formation and in the fruiting body tissue.
Some of these genes belong to the most highly upregulated genes of the entire genome
during fruiting body formation. This expression pattern results in an efficient, constitutive
protection of the fruiting body because at least some toxins are already in place in case the
structure is attacked by an antagonist. It is unclear whether the chemical defense of the
Agaricomycete fruiting body can be fortified by the induction of toxin production in response
to environmental factors e.g. the presence of antagonists. Antagonist-specific induction of
antibacterial and nematotoxic proteins upon challenge with bacteria and fungivorous
nematodes, respectively, has recently been reported for the vegetative mycelium of C.
cinerea(Kombrink et al. 2018; Tayyrov et al. 2019).
Toxins of Agaricomycete fruiting bodies include small molecules (secondary
metabolites)(Kües et al. 2018), peptides (ribosomally or non-ribosomally synthesized) and
proteins, all of which act by binding to and (in case of toxins with enzymatic activity)
converting specific target molecules in the antagonists. In the following, we will summarize
the current knowledge about the structure, function, biosynthesis, regulation and distribution
of Agaricomycete fruiting body toxins. In accordance with the focus of this review on
genomics, we restrict our summary to compounds where the biosynthetic genes have been
identified. We also make references to available gene expression data, but it must be noted
that most toxin-encoding genes evolve fast and thus they do not easily conform to orthology-
based categorizations that are central to the rest of this paper.
Small molecules - The structurally related alkaloids ibotenic acid and muscimol of
the fly agaric *Amanita muscaria* are two well-known examples of defensive secondary
metabolites of Agaricomycete fruiting bodies(Michelot and Melendez-Howell 2003). The
toxicity of these compounds toward flies gave rise to the name of the producer fungus. Their
bioactivity is due to their structural similarity to the neurotransmitters glutamate and GABA
and their binding to respective receptors in the brains of both invertebrates and vertebrates.
The recent discovery of the biosynthetic gene cluster for the compounds revealed that their
biosynthesis initiates from glutamate and makes it possible to identify additional *Amanita*
species producing these compounds on a genetic basis(Obermaier and Müller 2020).
Another psychotropic alkaloid produced by Agaricomycete fruiting bodies is psilocybin
produced by several *Psilocybe* species(J et al. 2017; Torrens-Spence et al. 2018). This
tryptophan derivative is an agonist of serotonergic receptors and is currently tested for
treatment of depressions(Tullis 2021). The transcriptional analysis of the recently identified
biosynthetic genes confirms its fruiting body-specific production(R et al. 2020). The blueing
reaction of psilocybin-producing fruiting bodies upon injury has recently been demonstrated
to be due to enzyme-mediated oxidative oligomerization of psilocybin suggesting that at
least part of the defense function of this compound may be based on optical
deterrence(Lenz et al. 2020). Wound-activation by injury (oxidation) has been reported for

other secondary metabolites of Agaricomycete fruiting bodies(Stadler and Sterner 1998; P
2008) but the genetic basis of these compounds and reactions is often not known.
Defense peptides - The bicyclic octapeptide α-amanitin produced by some *Amanita*,
Galerina, *Conocybe* and *Lepiota* species is probably the most famous toxin of
Agaricomycete fruiting bodies(Walton 2018). The thermal stability and the very low lethal
dose (0.1 mg/kg body weight) of the peptide have led to the common names 'death cap'
(*Amanita phalloides*) or 'destroying angel' (*Amanita bisporigera*) of α-amanitin producers.
Upon ingestion, the peptide is released from the fruiting body tissue and able to passively
enter the intestinal epithelial cells where it binds and inactivates the essential enzyme RNA
polymerase II in the nucleus(X et al. 2018). The fungus protects itself from the toxin by
expressing an insensitive variant of the target enzyme(PA et al. 1978). In 2007, Walton and
coworkers demonstrated that α-amanitin and its relative phallacidin are derived from
ribosomally synthesized precursor peptides with common flanking sequences (MSDIN) by
the action of a dual function prolyloligopeptidase-macrocyclase (POPB)(Hallen et al. 2007; H
et al. 2012). Genome surveys of above Agaricomycete species revealed a large family of
peptide precursor genes differing in the core peptide sequence and, thus, coding for α-
amanitin- and phallacidin-related, but structurally different peptides, referred to as
cycloamanides, whose function is unknown(Pulman et al. 2016; H et al. 2018; He et al.
2020; Zhou et al. 2021). Localization of POPB expression confirmed the fruiting body-
specific expression of α-amanitin and phallacidin(Luo et al. 2010; CH et al. 2018).
Above described secondary metabolites and peptides have received considerable attention,
mainly due to their activity towards humans. During the last two decades of research on
fungal defense, it has become evident, however, that a large part of the chemical defense of
Agaricomycete fruiting bodies, in particular against invertebrate predators and parasites, is
based on protein toxins(M et al. 2002b; J et al. 2016; Tayyrov et al. 2018). These toxins
have received less attention because their activity is usually lost upon boiling and is specific
for a particular group of invertebrates. In addition, based on their fruiting body-specific
expression, some of these proteins have been implicated in fruiting body formation rather
than defense (see the example of *C. cinerea* galectins CGL1 and CGL2 mentioned above),
which has delayed their recognition and functional analysis as toxins(S et al. 2007a; A et al.
2019a). Most of these protein toxins are stored intracellularly and released upon ingestion of
the fruiting body tissue by fungivores(Kunzler 2015; Künzler 2018). Once released, they
interact with the digestive tract of the fungivore. Characterized classes of protein toxins from
Agaricomycete fruiting bodies include lectins(Goldstein and Winter 2007; Bleuler-Martĺnez et
al. 2011; A et al. 2013), protease inhibitors(S et al. 2015; Sabotič et al. 2019), pore-forming
toxins (PFTs)(Anderluh and Maček 2003; JM et al. 2010; Panevska et al. 2020),
ribotoxins(Tayyrov et al. 2019; Ragucci et al. 2021) and biotin-binding proteins(Y et al. 2009;
S et al. 2012a).
Defense related lectins - Fruiting body lectins have long been used as tools in
glycobiology without addressing their biological function(Y and H 2014). In the meantime, it
has been established that these proteins bind to specific glycoepitopes on the epithelial cells
of the digestive tract of the fungivore which can ultimately lead to disruption of the structure
and function of this tissue(Stutz et al. 2015). The mechanism of lectin-mediated toxicity is not
clear yet but, in case of lectins without additional functional domain (hololectins), the
bioactivity appears to depend on the crosslinking of protein-bound glycoepitopes on the
epithelial cell surface(S et al. 2017), whereas in case of lectins with additional functional
domains (chimerolectins), the bioactivity appears to depend on binding to lipid-bound
glycoepitopes and endocytosis(T et al. 2011; Juillot et al. 2016). Fruiting body lectins can

adopt various folds and carbohydrate specificities amongst which the RicinB-like (β-trefoil)
fold and fucoside-binding, respectively, are most common. A prototypic example for such a
fruiting body lectin is the hololectin CCL2 (*C. cinerea* protein ID: 408852) from the model
Agaricomycete *Coprinopsis cinerea* which is present in other Agaricomycete species, adopts
a RicinB-like fold, forms a dimer, binds to a specific fucoside present on nematode N-glycan
cores and is toxic towards bacterivorous and fungivorous nematodes(Schubert et al. 2012; S
et al. 2017; Tayyrov et al. 2018). A summary of diverse entomo- and nematotoxic lectin have
been described from fruiting bodies has been given by Sabotic et al (J et al. 2016). Our data
confirm previous reports on the upregulation (>150-fold) of CCL1 (*C. cinerea* protein ID:
456849) and CCL2 (*C. cinerea* protein ID: 408852) in primordia (Table 33).
The RicinB-like (beta-trefoil) fold is also adopted by protease-inhibitors isolated from
Agaricomycete fruiting bodies(J et al. 2016). These proteins were shown to inhibit serine and
cysteine proteases of various insect larvae. A prototypic example for this type of fruiting body
toxin is Clitocypin, an insecticidal cysteine-protease inhibitor isolated from the
Agaricomycete *Clitocybe nebularis*(J et al. 2011; Šmid et al. 2015). Of the examined
species, the genomes of *C. aegerita* and *L. bicolor* encode 1 and 20 putative clitocypins,
respectively. Our data identified a putatively defense-related RicinB protein orthogroup
(represented by *C. cinerea* 485770)(Table 33). Another example is the six-bladed beta-
propeller lectin Tectonin2 from L. bicolor whose production is induced in fruiting bodies and
ectomycorrhizae(Martin et al. 2008). This lectin forms tetramers harboring 24 binding sites
for O-methylated glycoepitopes and exhibits toxicity towards nematodes and agglutination of
bacteria displaying respective glycoepitopes on their cell surfaces(Wohlschlager et al. 2014;
Sommer et al. 2018). We detected tectonin homologs in the genomes of *L. bicolor* and *C.*
aegerita.
Pore-forming toxins - A common and probably very ancient class of protein toxins
of Agaricomycete fruiting bodies are pore-forming toxins (PFTs)(Peraro and van der Goot
2015). These toxins are divided into α- and β-PFTs depending on whether the predominant
secondary structure motif is an α-helix or β-sheet. Within the producing cell, these proteins
are usually monomeric and soluble, but upon release and binding to a target cell membrane,
they undergo a dramatic conformational change. This change in conformation leads to
oligomerization and insertion of the protein into the membrane. The result of this complex
process is a proteinaceous pore that leads to lysis of the target cell. Among PFTs occurring
in Agaricomycete fruiting bodies, members of the aerolysin and <u>M</u>embrane <u>A</u>ttack <u>C</u>omplex
<u>P</u>er<u>F</u>orin/<u>C</u>holesterol <u>D</u>ependent <u>C</u>ytolysin (MACPF/CDC) β-PFT superfamilies have been
characterized most extensively(Reboul et al. 2016; Cirauqui et al. 2017). Examples for
aerolysins are the hemolytic *Laetiporus sulphureus* lectin (LSL)(JM et al. 2010), a
homologous and recently identified nematotoxic protein, CC1G_11805 (*C. cinerea* protein
ID: 546868) and its two paralogs, from *C. cinerea*(Plaza et al. 2014b) and the hemolytic
flammutoxin from *F. velutipes*(T et al. 2004b). Pri1, a primordium-expressed gene in *C.*
aegerita (Agrae_CAA7262647) also belongs to pore-forming toxins(Fernandez Espinar and
Labarère 1997). The MACPF/CDC superfamily is represented by the insecticidal
pleurotolysin complex, consisting of PlyA and PlyB, from the edible mushroom *P.*
ostreatus(K et al. 2014; M et al. 2015b; N et al. 2015; A et al. 2019a). The main
representatives of the two β-PFT superfamilies differ in the specificity determinant of the
target membrane and the dimensions of the formed membrane pore. Whereas LSL likely
binds to lactose-containing glycoepitopes (as part of glycolipids or GPI-anchored
glycoproteins) on the target membrane via an N-terminal RicinB-type(β-trefoil) lectin domain
and forms homomeric transmembrane β-barrels (pores) of seven or eight protomers with an

inner diameter of ~4 nm(I et al. 2011; Szczesny et al. 2011), pleurotolysin PlyA binds to
sphingophospholipids (sphingomyelins) of cholesterol-rich membranes and directs PlyB to
the formation of transmembrane β-barrels (pores) consisting of 13 heteromeric protomers
and an inner diameter of ~ 8 nm(Novak et al. 2020).
Ribotoxins are ribonucleases cleaving a single phosphodiester bond in the
universally conserved sarcin/ricin loop (SRL) of 23S or 28S rRNA in intact prokaryotic or
eukaryotic ribosomes, respectively(M et al. 2017b). Upon this action, the ribosome is unable
to interact with translation elongation factors and translation is stalled. The SRL is named
after α-sarcin, one of the best characterized ribotoxins secreted by the ascomycetous mold
Aspergillus giganteus(M et al. 2017b). Other fungal ribotoxins have been isolated from
ascomycetous entomopathogens and are generally highly active against insects(A et al.
2009; M et al. 2017a). Recently, the first basidiomycetous ribotoxin, ageritin (Agt1), from
fruiting bodies of the edible mushroom *C. aegerita* was isolated(Landi et al. 2017).
Heterologous expression of the respective cDNA in *Escherichia coli* and phylogenetic
analysis showed that the protein is widespread but patchy among Agaricomycetes and
displays entomotoxic activity similar to its ascomycetous counterparts(A et al. 2019b). In
contrast to its ascomycetous counterparts, however, ageritin does not contain a canonical
signal peptide for secretion suggesting that the protein, as most protein toxins of
Agaricomycete fruiting bodies, is stored intracellularly(Baglivo et al. 2020). In order to avoid
damage to the ribosomes of the producing fungal cell, the protein might be sequestered into
a specific subcellular compartment which might involve N-terminal truncation of the
protein(Ragucci et al. 2021). The agt1 gene (*C. aegerita* CAA7259527) was induced
considerably in primordia relative to vegetative mycelium in *C. aegerita*, based on data
produced by Orban et al (Orban et al. 2021).
Tamavadins - Finally, defense proteins of some Agaricomycete fruiting bodies,
similar to animals and bacteria, include proteins that are able to avidly bind and thus
sequester biotin at equimolar concentrations(Michael Green 1990). The prototype of these
proteins, avidin, is thought to protect the eggs of birds from bacterial infections, but the
protein and its relatives also exhibit broad and strong insecticidal activity(JT et al. 2010). A
family of avidin-related biotin-binding proteins, tamavidin (called Tam1 and Tam2), was
isolated from the edible mushroom *Pleurotus cornucopiae* (Tamogitake mushroom) and
shown to possess very strong entomotoxic and nematotoxic activity(Y et al. 2009; S et al.
2012a). Amongst the fruiting body protein toxins tested against two fungivorous nematode
species, tamavidin was the most effective one(Tayyrov et al. 2018). It is unclear how the
producing cells shield their own biotin against this cytoplasmic protein.
Taken together, this data shows that the chemical armory of Agaricomycete fruiting
bodies is very diverse, in terms of compound classes, molecular targets and antagonist
specificity of the toxins, but three common themes can be recognized: (1) Toxin-encoding
genes are phylogenetically patchy, with each family present in only a subset of species (2)
protein toxins provide a large part of the chemical defense of these structures; and (3)
intracellularly stored toxins directed against nematodes and arthropods are predominant.
Latter finding may reflect the use of Agaricomycete fruiting bodies as habitat by these
organisms(Ruess and Lussenhop 2005; Krivosheina 2008). It has even been postulated that
some mycophagous Drosophila flies exploit fruiting body nematotoxins to purge themselves
from insect-parasitic nematodes(JA et al. 2017).
3930

3931 4.11. Secondary metabolism and natural product biosynthesis

Secondary metabolites are one of the most important classes of molecules in chemical
warfare and developmental signaling in fungi(Spiteller 2015; Keller 2019; Gressler et al.
2021). Based on recent genome surveys it appears that the fungal secondary metabolome is
extremely diverse and is a largely untapped resource of natural products and bioactive
molecules(Keller 2019). This may be particularly true for the Agaricomycetes, despite
several lines of evidence for a large diversity of secondary metabolites produced by
Agaricomycetes(Spiteller 2015; Gressler et al. 2021). From the perspective of fruiting body
formation, it is important to note that secondary metabolites can serve defense or repellent
purposes in fruiting bodies (see chapter on Defense) but can also influence developmental
processes. For example, the deletion of a polyketide synthase gene resulted in malformed
perithecia of *Sordaria macrospora*(D and M 2014). The lack of coprinoferrin production in *C.*
cinerea resulted in the loss of fruiting body formation(Tsunematsu et al. 2019).
Due to differences in basic genome organization between Asco- and Basidiomycota,
the latter possess fewer secondary metabolism gene clusters, yet, according to a recent
study some developmentally regulated genes formed 'clusters' in the genomes of 8 species
and some of these 'clusters' overlapped with predicted biosynthetic gene clusters that might
encode secondary metabolism genes(Merényi et al. 2021).
The number of functionally characterized secondary metabolism genes and linked
natural products of Agaricomycetes is rapidly increasing. For example, *ArmB* and orsellinic
acid in *Armillaria* spp(Lackner et al. 2013), *LpaA* and laetiporic acid in *Laetiporus* sp(Seibold
et al. 2020), or *St.pks1* and strobilurin in *Strobilurus tenacellus*(Nofiani et al. 2018) as well as
numerous terpene synthases have been circumscribed in the past decade (reviewed in more
detail by(Gressler et al. 2021)). Strobilurin derivatives are available commercially and are the
world's most sold fungicides(DW et al. 2002). Siderophore-type secondary metabolites,
coprinoferrin, ferrichrome A and basidioferrin in *C. cinerea, Omphalotus olearius* and
Ceriporiopsis subvermispora, respectively, and their putative synthesizing genes were also
identified. In *C. cinerea cfp1* (*C. cinerea* protein ID: *354233*) was reported to be responsible
for coprinoferrin production(Tsunematsu et al. 2019), whereas in *O. olearius* a cluster
composed of *fso1, omo1* and *ato1*(K et al. 2005). Coprinoferrin proved necessary for fruiting
body development, as the cfp1 mutant did not, only in the presence of exogenously
administered coprinoferrin produce fruiting bodies. Contrary to expectations based on
Ascomycota knowledge, deleting a homolog (*C. cinerea* protein ID: 357031) of *A. nidulans*
LaeA did not compromise coprinoferrin production(Tsunematsu et al. 2019). Despite recent
advances in understanding Agaricomycetes secondary metabolism, given the number of
putative secondary metabolism genes encoded by fungal genomes, we are likely hardly
scratching the surface of natural product diversity in mushroom-forming fungi.
Since many secondary metabolites specifically occur in fruiting bodies of
Agaricomycetes, it is expected that their biosynthetic gene clusters may be developmentally
regulated. However, secondary metabolism related genes evolve fast, which does not make
them easily detectable in the orthology-based framework we use here. Nevertheless, we
identified several secondary metabolism related genes that showed shared and conserved
developmental expression profiles in the examined Agaricomycetes, although the natural
products they produce are unknown in most cases.
Polyketide synthases (PKS) are large (often >1500 amino acid) multi-domain
enzymes that produce diverse secondary metabolites. They are widely distributed in fungi
and are subject to intense research, mostly in the Ascomycota. Polyketide synthases and

the few examples of known natural products of the Basidiomycota have recently been reviewed(Gressler et al. 2021). Basidiomycete PKSs mostly belong to type 1, which is divided into highly-, partially- and non-reducing PKSs. Hybrid nonribosomal peptide synthase-PKS genes have been shown to underlie fungal bioluminescence(Ke et al. 2020a; Gressler et al. 2021).

We identified orthologs of previously described siderophore biosynthesis related genes in the examined Agaricomycetes. Members of the biosynthetic cluster described from *O. olearius*(K et al. 2005) seem to be conserved, with fso1 being orthologous to *C. cinerea* cfp1 (*C. cinerea* protein ID: 354233), whereas L-ornithine N(5)-monooxygenase omo1 and the N-acyltransferase ato1 seem single-copy in Agaricomycetes genomes (represented by *C. cinerea* 492586 and 161175). These genes were not developmentally regulated and had highest expression in vegetative mycelia. We further identified a single CDE orthogroup of polyketide synthases (represented by *C. cinerea* 446258, Table 34), which was developmentally regulated in 11/8 species (FC>2/4). These genes were stipe upregulated (albeit weakly) in most species (*C. cinerea, P. ostreatus, M. kentingensis, A. ostoyae*), although their expression also increased in mature caps of *P. ostreatus, A. ostoyae, M. kentingensis* relative to younger cap stages and in mature fruiting bodies of *C. aegerita*, *A. ampla* and *S. commune* relative to earlier stages. The *C. cinerea* gene shows peak expression in meiotic gill tissues, whereas in *P. ostreatus* and *A. ostoyae*, where gills and caps were sampled separately like in *C. cinerea*, highest expression was observed in caps but not in gills. The *C. cinerea* gene in this orthogroup is the best hit of *Aspergillus fumigatus* PksB, which is a polyketide synthase involved in spore pigment (melanin) production. Based on this similarity, we hypothesize that these genes are also involved in pigment production in Agaricomycetes.

Terpene synthases have been a focus of secondary metabolism studies in mushroom-forming fungi. They encode highly promiscuous enzymes that can generate a large diversity of metabolites(Slot and Gluck-Thaler 2019) and thus provide the basis for fast adaptive evolution. Slot and Gluck-Thaler showed that morphologically complex fungi harbor more terpene synthases than simpler counterparts(Slot and Gluck-Thaler 2019). Similarly, terpene synthases were found to be strongly overrepresented in the genomes of Agaricomycetes compared to other fungi (FDR corrected p-value < 10^{-125})(Krizsán et al. 2019). Twelve sesquiterpene synthases (Fla1-12) were identified in *F. velutipes*(Tao et al. 2016), eleven in *C. aegerita* (Agr1-Agr11)(Zhang et al. 2020, 2021a) and in *Omphalotus olearius* (Omp1-10)(GT et al. 2012) and five (Cop1-Cop6) in *C. cinerea*(Agger et al. 2009), with information on the potential metabolite products in many cases (e.g. viridiflorene/viridiflorol, Δ^{6}-protoilludene, γ-cadinene, germacrene).

Agger et al identified six genes, Cop1 to Cop6, related to terpene biosynthesis in *C. cinerea*(Agger et al. 2009). Of the genes reported by Agger et al, orthologs of Cop2 and Cop4 appear interesting from the perspective of mushroom development. Orthologs of Cop1 show an upregulation in primordia relative to vegetative mycelium in *C. aegerita, A. ampla* (only FC>2), *L. bicolor, Ph. chrysosporium* and were cap-upregulated in *A. ostoyae, P. ostreatus, L. edodes* and *M. kentingensis*. On the other hand, orthologs of Cop4 (*C. cinerea* protein ID: 30510) were FB-init in *C. cinerea, C. aegerita, A. ampla, L. bicolor, L. edodes* and *P. ostreatus* (Table 34). Cop2 (*C. cinerea* protein ID: 443805) is only present in *C. cinerea, A. bisporus* and *P. ostreatus* and is upregulated in primordia relative to vegetative mycelium in all species, suggesting it functions during fruiting body initiation. Products of these genes were reported to as δ-cadinene and germacrene A, but the functions of these natural products remain unknown.

We found 13-28 (63 in *A. ostoyae*) putative terpene synthases in the examined
Agaricomycetes genomes. Of these, several were developmentally regulated, showing one
of two most typical expression profiles: (i) highest expression in vegetative mycelium or (ii) a
strong induction in primordia and other fruiting stages relative to vegetative mycelium. The
latter expression profile is reminiscent of that of typical defense genes (e.g. pore-forming
toxins, see above), but it is compatible with several other putative functions as well. Terpene
synthases formed three CDE orthogroups (Table 34), two of these correspond to *C. cinerea*
Cop1 (*C. cinerea* protein ID: 469738) and Cop4 (*C. cinerea* protein ID: 30510, contains *C.*
aegerita Agr4 also), discussed above. The third CDE orthogroup (represented by *C. cinerea*
488324) is developmentally regulated in 8/6 species, but shows mixed expression profiles.
Taken together, terpene synthases might participate in important fruiting body
functions, including events around the initiation of primordium development, yet their exact
functions, whether the production of defense/repellent metabolites and/or compounds for
communication, remain unknown for the moment.
Phenylalanine ammonia-lyases - We detected two highly conserved CDE
orthogroups of phenylalanine ammonia-lyases (PAL), one of which (*C. cinerea* protein ID:
386671, ortholog of pal1 of *Pleurotus*) is developmentally regulated in 11/9 species, the
other (*C. cinerea* protein ID: 361541, ortholog of pal2 of *Pleurotus*(Hou et al. 2019)) is in 8/5
(out of 9 species in which present) (Table 34). PAL enzymes are best known from plants,
where they are part of the phenylpropanoid pathway, which is involved in the production of
diverse secondary metabolites, associated with defense, lignin and flavonoid
biosynthesis(Dixon et al. 2002). In fungi, phenylalanine ammonia-lyases are discussed
mostly in the context of catabolism of phenylalanine to cinnamic acid(Hyun et al. 2011),
although some reports of the production of diverse compounds have been published(Hyun et
al. 2011). Recently, an association with fruiting body formation has been reported in *F.*
velutipes(Yun et al. 2015) and *P. ostreatus* (Hou et al. 2019).
Our analyses revealed two PAL-encoding genes in most of the examined
Agaricomycetes (up to 4 in *A. ostoyae*). Expression profiles of genes within the two CDE
orthogroups were not conserved, although several genes showed an upregulation in
primordia (e.g. both genes in *C. cinerea*). At the moment the physiological role of PALs is
unknown, to our best knowledge, in any fungus. They could be involved in the production of
defense-related metabolites, as in plants, or other fruiting body related functionalities.
4058

4059 4.12. Cytoskeleton

A plastic cytoskeleton is a key factor for eukaryotic cell shape regulation and has been
named one of the prerequisites for diversification of size, form and function in multicellular
eukaryotes(Knoll 2011). It is involved in the regulation of diverse morphogenetic processes
in fungi, including hyphal growth, conidiation, biofilm formation, dimorphic switching, to name
a few(Riquelme et al. 2018). In line with this, although at a somewhat limited scale, evidence
is available for the role of cytoskeleton rearrangement in fruiting body development.
Evidence from Gene Ontology analyses pointed at differential cytoskeletal gene expression
in *P. microcarpus*(Pereira et al. 2017). Microscopy observations provided evidence for a key
role of cytoskeletal elements during the development of mushroom-forming fungi. For
example, longitudinally oriented microtubules were observed in the sterigmata during spore
formation in *C. cinerea*, along which Golgi vesicles carry carbohydrates to the developing
spore and spore wall(Kües and Liu 2000).

4072    In terms of expression levels, we found only limited dynamics in cytoskeletal proteins,
4073 likely reflecting the need for continuous supply of its components for the cell. Cheng et al
4074 reported an upregulation of tubulin and tubulin-specific chaperone genes in primordium
4075 stages of *C. cinerea*(Cheng et al. 2013a) using 5'-SAGE technology. Although newer data
4076 did not confirm these observations(Muraguchi et al. 2015; Krizsán et al. 2019), both recent
4077 RNA-Seq studies reported upregulation of these genes in stipes of *C. cinerea*.
4078    Following these lines of evidence, we annotated the main cytoskeletal protein groups
4079 in *C. cinerea* based on homology and domain composition and checked their orthologs'
4080 expression patterns across mushroom-forming fungi. We grouped 67 cytoskeletal and
4081 associated protein coding genes into the following categories (Table 35): actin, actin
4082 nucleation, actin filament organization, actin-based motor protein, microtubule, microtubule
4083 heterodimerization, microtubule-based motor protein, regulation of microtubule-based motor
4084 proteins. Copy numbers in these categories agreed with previous annotations, including 2
4085 copies of formins per species and a tubulin gene duplication in the common ancestor of
4086 Basidiomycota(Zhao et al. 2014). Of cytoskeletal genes, only kinesin genes and the
4087 Dynactin complex showed consistent expression dynamics. In both cases, expression peaks
4088 were observed in young fruiting body gills and stipes in *C. cinerea* (Fig. 14, Supplementary
4089 Fig. 10).
4090    Tubulin gene expression (alpha1-, alpha2-, beta-tubulin and tubulin chaperone) was
4091 previously reported to be regulated during early fruiting events of *C. cinerea*, with
4092 upregulation in stage 1 primordia based on 5'-SAGE data(Cheng et al. 2013a). Our data on
4093 the corresponding genes revealed that the three of these, alpha1- (*C. cinerea* protein ID:
4094 176636), beta-tubulin (*C. cinerea* protein ID: 393528) and tubulin chaperone (*C. cinerea*
4095 protein ID: 413977) had constant expression in primordia, but show expression peak in
4096 young fruiting body stipes, whereas alpha2-tubulin (*C. cinerea* protein ID: 359665) was
4097 upregulated in young fruiting body caps. These expression patterns were not consistent
4098 across species (Table 35) and possibly reflect fast stipe elongation of *C. cinerea*.
4099 **Septins** - Septins form a highly conserved family of proteins involved in morphogenesis,
4100 development and in generating cell asymmetries in fungi and in eukaryotes in
4101 general(Gladfelter 2006; Khan et al. 2015). They have been shown to be responsible for
4102 morphogenesis in a number of model fungi, such as bud emergence in *S. cerevisiae*,
4103 septum formation in hyphae or appressorium formation in *Magnaporthe oryzae*, among
4104 others(Gladfelter 2006). Their broad involvement in morphogenesis and limited evidence
4105 from fruiting body forming fungi suggest that septins have important, although to date
4106 underappreciated, roles in fruiting body development. For example, they have been reported
4107 in stipe elongation in *C. cinerea*(Shioya et al. 2013).
4108    We annotated septins based on their domain composition: the combination of two
4109 IPR domain terms (IPR030379, IPR016491) are diagnostic for fungal septins. Most
4110 mushroom-forming fungi have 7 septin genes (*C. cinerea* protein ID: 367587, 416864 [*cdc3*,
4111 = *A. nidulans* AspE, =AspB], 103773, 415573 [top hit of *A. nidulans* AspE], 103809, 455217,
4112 481878), with some having only 6 while a few having 8, which is similar to the septin
4113 repertoires seen in filamentous fungi and yeasts(Douglas et al. 2005).
4114    Septins are highly variable genes, and did not form conserved orthogroups, but
4115 nevertheless maintain some functional conservation in mushroom-forming fungi. Two
4116 orthogroups of septins showed consistent upregulation at the initiation of fruiting body
4117 development, in the very first primordium stages in all species in which they are
4118 present(Krizsán et al. 2019; Merényi et al. 2021). One of these is the top hit of AspE of *A.*
4119 *nidulans*, which is involved in conidiation and spore production(Hernández-Rodríguez et al.

2014). Septin expression seems to be highly induced in stipes in *C. cinerea, M.*
kentingensis, A. ostoyae, but following a different expression tendency in *P. ostreatus*
(mostly in primordia) and *A. bisporus* (mostly in young fruiting bodies). The stipe induced
septins included *Cc.cdc3* of *C. cinerea*, which has been shown to be involved in stipe
elongation(Shioya et al. 2013). The ortholog of *C. cinerea cdc3* was also mostly stipe-
induced in *A. bisporus*(卢园萍 et al. 2019). Muraguchi et al reported a downregulation of
septin expression from the 4_0hrPri primordia (approximately coinciding with stage 2
primordia of Krizsan et al and Kues 2000(Kües 2000; Krizsán et al. 2019)) to the 5_12hrPri
stage of *C. cinerea*(Muraguchi et al. 2015).
The limited diversity of septins raises the question how they can fulfill the diverse
roles they have in fungal morphogenesis. Although most septins had peak expression in
stipes of mushroom-forming fungi (except in certain species) their expression level was
dynamically changing in other tissues and from one developmental stage to the other as
well, suggesting that there is still a long way to uncover their roles in fruiting body
development. Understanding their protein-protein interaction networks is a promising future
avenue of research.

4136 4.13. Ubiquitin-proteasome system

4137 4.13.1. Cop9 signalosome

The Cop9 signalosome is a protein complex conserved across all eukaryotes, which
catalyzes the deactivation of cullin-RING ubiquitin ligase (CRL) complexes by removing the
Nedd8 protein from the cullin subunit of the complex (deneddylation reaction). As such, it is
an important component of various signal transduction pathways and is involved primarily in
multicellular developmental processes. In fungi, it was found dispensable for vegetative
growth, but necessary for sexual fruiting body formation(Busch et al. 2003; Braus et al.
2010), secondary metabolism and it was reported to interact with the Velvet complex(Kress
et al. 2012; Meister et al. 2019). Recognition of the COP9 signalosome (CSN) as an
important player in fruiting body formation started with genetic screens that identified CsnE
and CsnD, which displayed identical defects in fruiting body initiation in *A. nidulans*(Busch et
al. 2003, 2007). The *A. nidulans* Cop9 complex represents an 8-subunit, stereotypical
complex, whereas that of other model fungi (*N. crassa, S. cerevisiae* or *S. pombe*) possess
fewer subunits and are not typical(Braus et al. 2010). Agaricomycetes genomes appear to
encode a typical 8-subunit CSN (Table 36). Neither we nor previous studies found significant
expression dynamics of any subunit of the CSN in mushroom-forming fungi. It is not known
whether the complex is involved in fruiting body development as it is in Ascomycota,
nevertheless, it will be an interesting target for future experiments, especially because it is
involved in light regulation in *N. crassa* and can interact with the Velvet complex, two key
factors that regulate Basidiomycota development also.

4157 4.13.2. Protein ubiquitination

Covalent protein modification by the ubiquitin-proteasome system is a widely conserved
regulatory machinery that functions in several essential processes in eukaryotes and was
recently detected as a potentially very interesting function during fruiting body
development(Krizsán et al. 2019). The regulatory potential of the ubiquitin-proteasome
system stems from its ability to attach the 76-amino acid ubiquitin protein to lysines of target

proteins, which either destines them for degradation by the proteasome or modifies their
activity, cellular localization or DNA-binding ability(LC et al. 2007; Liu and Xue 2011; Finley
et al. 2012). Ubiquitination can modify the activity of transcription factors, which provides an
additional layer of transcriptional regulation(Freiman and Tjian 2003).
The protein ubiquitination machinery is composed of E1 ubiquitin-activating, E2
ubiquitin-conjugating enzymes and E3 ubiquitin ligases, which attach the ubiquitin to target
proteins. While E1 and E2 enzymes are generic components of the system and are present
in limited numbers in eukaryotic genomes (Table 37), ubiquitin ligases provide the substrate
specificity and come in many subtypes and display considerable diversity due to
interchangeable protein components, of which hundreds may exist in the genomes of some
species(Hutchins et al. 2013; Zheng and Shabek 2017). The most diverse such parts are F-
box and BTB/POZ domain proteins, both of which can confer substrate-specificity to the E3
complexes(R et al. 2003) and RING-type zinc finger proteins, which are scaffold proteins
linking target proteins with E2 enzymes(Metzger et al. 2012). These three gene families
were found to be strongly expanded in mushroom-forming fungi, with frequent tissue-specific
developmental regulation seen in RNA-Seq data(Almási et al. 2019; Krizsán et al. 2019)
(see also Table 37). The expansion stems from Agaricomycetes-specific duplication events
and a correlation between fruiting body complexity and F-Box, BTB/POZ and RING-type zinc
finger copy number and expression has been suggested(Krizsán et al. 2019), which could
explain the evolution of increasingly complex mushroom morphologies in the class.
The widespread cellular roles of ubiquitination suggests that it should be instrumental
during fruiting body morphogenesis too. In line with this, the protein ubiquitination system
was found differentially expressed in *F. velutipes* fruiting bodies(M et al. 2006; Liu et al.
2020b) and in caps of the same species subjected to differential CO_2 treatments(Yan et al.
2019). In recent comparative transcriptomics analyses, we provided evidence for the
developmental regulation of several components, mostly those that confer substrate
specificity to E3 ligase complexes, in multiple Agaricomycetes species, including *C. cinerea,*
A. ostoyae, A. ampla, S. commune or *P. ostreatus*, among others(Almási et al. 2019; Krizsán
et al. 2019; Merényi et al. 2021). BTB domain-containing proteins, a family of E3 proteins,
were reported to be upregulated in fruiting bodies in *Antrodia cinnamomea* (Lu et al. 2014).
The combination of a remarkable gene family expansion with developmental regulation
suggests that the ubiquitin-proteasome system could play a pivotal role in fruiting body
development, although mechanistic details of the process remain to be understood.
Ubiquitylation is key in developmental processes of the Ascomycota also, including fruiting
body formation, and in some cases more mechanistic details have been uncovered
(reviewed in(Riquelme et al. 2018)). However, a similar expansion of F-Box domains, as
seen in Agaricomycetes, has not been reported.
Hereafter, we present an annotation of components of the protein ubiquitination
machinery in selected mushroom-forming fungi, with particular emphasis on *C. cinerea*
genes and comparisons across species. In this presentation, we follow the classification of
E3 ubiquitin ligases used by Hutchins et al(Hutchins et al. 2013) and identify protein diversity
of each protein family based on Interpro domain HMM-s. Results of the annotation along
with orthologous genes and conservation levels are shown in Supplementary table 4. A
limitation of this approach is that the presence of a defining domain in the protein does not in
all the cases, indicate a function in the ubiquitination machinery. For example, BTB/POZ
domains, which contribute to expanding the diversity of E3 ligase complexes(R et al. 2003)
can, in rare cases, also occur in proteins not related to ubiquitination (e.g. cytoskeleton or
transcription regulating proteins(Stogios et al. 2005)).

Several components of the ubiquitination machinery are very conserved in
Agaricomycetes genomes, including E1 (on average 3 genes) and E2 enzymes (19-24
genes), Cullin genes (4-10 genes) and deubiquitinating enzyme-encoding genes (23-27
genes) (Supplementary table 4). Most of these genes show no or modest expression
dynamics, which is not surprising given that the ubiquitination machinery is needed
constantly in the cell. Of the E1 enzymes, only *S. cerevisiae* Uba3 orthologs (*C. cinerea*
protein ID: 485679), which belong to the neddylation pathway, are developmentally
regulated to some extent (8/3 species at FC=2/4). In *C. cinerea* 14 of the 23 E2 enzymes
and half (13 of 26) of deubiquitinating enzymes are developmentally regulated, potentially
indicating functional diversification of E2 enzymes and deubiquitinating enzymes during
development (Supplementary table 5). We found that none of the cullin genes in *C. cinerea*
are developmentally regulated. Significantly larger dynamics can be observed in E3 ligase
components, which are discussed in detail below.
RING-type Zinc finger proteins - RING-domain containing proteins, named after
RING (really interesting new gene), are E3 ubiquitin ligases that confer interchangeability to
the ligase complexes(Deshaies and Joazeiro 2009). Eukaryotic genomes contain large
numbers of RING-domain proteins, which, along with other E3 ligases, enable the ubiquitin
system to selectively ubiquitinate proteins. RING-domain proteins are generally poorly
known in filamentous fungi, though it was recently noticed that they are strongly
overrepresented in Agaricomycetes(Krizsán et al. 2019). The *mfbAc* (=Le.MFB1) gene of *L.*
edodes (Lenedo1_1172639) described as a protein with adhesive properties by Kondoh et
al(Kondoh et al. 1995) also belongs to this gene family.
The examined Agaricomycetes possess 81-148 RING-domain containing proteins
(Table 37), consistent with a previous broader sampling across 200 species. On average,
30-50 RING-domain proteins were developmentally regulated in the examined
species(Almási et al. 2019; Krizsán et al. 2019), representing a significant pool of potentially
important regulators. RING-domain proteins are highly variable, with most of them not
showing clear orthology across species. Nevertheless, we found eight CDE orthogroups
which displayed conserved developmental expression, mostly at FC>2 (Table 37). For
example, one of these (represented by *C. cinerea* 378120) showed a very distinct peak in
gills of *C. cinerea, A. ostoyae, P. ostreatus* and caps (containing gills) of *M. kentingensis, L.*
edodes as well as mature fruiting bodies of *S. commune* and *A. ampla*. Another CDE
orthogroup (represented by *C. cinerea* 440772) was upregulated in gill tissues of mature
fruiting bodies in *P. ostreatus, A. ostoyae, M. kentingensis*, *L. bicolor* and mature fruiting
bodies of *S. commune* and *A. ampla*, but not in *A. bisporus, L. edodes* or *C. cinerea*. These
two orthogroups coincide (partly) with meiosis and sporulation, so they could be involved in
these, or other gill-specific processes. Genes in another CDE orthogroup (C. cinerea protein
ID: 543850) showed an abrupt, although modest upregulation at the initiation of fruiting body
development and expression peak in mature fruiting bodies or gills therein in *C. cinerea, A.*
ostoyae, C. aegerita, A. ampla, L. bicolor, L. tigrinus, Ph. chrysosporium, P. ostreatus and *S.*
commune (not in *L. edodes*).
F-box proteins - F-box proteins were recently identified as a potentially key group of
proteins in fruiting body development(Krizsán et al. 2019). They are generally responsible for
providing target specificity to the ubiquitination machinery, by binding and docking substrate
proteins to Cullin E3 ubiquitin ligase complexes. Some F-box proteins interact with multiple
target proteins while others only with a single protein(Jonkers and Rep 2009). Similarly,
some F-Box proteins function independently of the SCF complex(Jonkers and Rep 2009),
allowing for additional, yet poorly known regulatory functions to emerge. For example, F-box

proteins can act as transcriptional co-regulators in plants(Chae et al. 2008) and are present
in large numbers in plant genomes (up to 700-1000 genes(Xu et al. 2009)), providing the
possibility to assemble an enormous diversity of alternative ubiquitin ligase complexes.
The F-box family is highly expanded in mushroom-forming fungi and is composed of
fast-evolving proteins with little similarity to each other. Agaricomycetes possess 69 to
~1200 F-box proteins in their genomes, whereas other filamentous fungi and yeasts possess
up to 90 and 20 such genes(Krizsán et al. 2019), respectively. This represents a strong
enrichment in Agaricomycetes genomes ($P<10^{-140}$), considering that the number of other
proteins is not proportionately higher in this group than in other fungi. F-box protein encoding
genes often showed tissue-specific expression and their numbers showed a positive
correlation with the complexity level of the fruiting bodies(Krizsán et al. 2019).
F-Box proteins are highly variable and form few conserved orthogroups, compared to
their diversity in fungal genomes. However, we found eleven orthogroups that showed some
level of conservation and consistent developmental regulation across species (Table 37).
These included orthologs of *A. nidulans* GrrA (*C. cinerea* protein ID: 358351), which is
needed for meiosis and ascospore maturation(S et al. 2006a; Riquelme et al. 2018), are
conserved across Agaricomycetes and developmentally regulated in five species (FC>2),
indicating that some F-Box proteins are conserved despite the general volatility of the family.
Due to this high variability, F-box proteins are hard to examine systematically using a
comparative transcriptomics approach. Rather, they likely often fulfill species- or clade-
specific roles, which could be studied either with more densely sampled transcriptomic data,
or reverse genetics approaches.
BTB/POZ domain proteins - Proteins that contain the BTB/POZ motif, a protein-
protein interaction (homodimerization) domain, can possess diverse cellular functions, the
most frequent of which are regulation of chromatin structure (thus, transcription) and protein
ubiquitination, as members of E3 ubiquitin ligase complexes(S et al. 1994; R et al. 2003).
They are poorly known in fungi.
In animals, plants as well as fungi outside the Agaricomycetes, BTB domains are
mostly fused to other conserved protein domains(Stogios et al. 2005) (in animals, mostly to
zinc fingers(S et al. 1994)). In contrast, we observed that in the Agaricomycetes, less than
10% of BTB/POZ domain proteins associate with any other domains and virtually never with
DNA binding transcription factor domains, indicating that the gene family expansion in
mushroom-forming fungi is concerned with proteins that only contain the BTB/POZ domain.
This domain composition suggests that the BTB/POZ family is primarily involved in E3
ligases in Agaricomycetes, therefore we discuss this family under the chapter on protein
ubiquitination.
We found 8 CDE orthogroups with some level of conserved developmental
expression in Agaricomycetes (Table 37). Members of two (*C. cinerea* protein ID: 379603,
449824) of these orthogroups are co-expressed in stages or tissues in which meiosis and
sporulation occurs: gills of *A. ostoyae* and *P. ostreatus*, mature fruiting bodies of *Ph.
chrysosporium, A. ampla* and *S. commune,* mature caps of *M. kentingensis, L. bicolor, L.
edodes* (but not developmentally regulated in *C. cinerea*). Another CDE orthogroup (*C.
cinerea* protein ID: 459680) was upregulated in stipes of *C. cinerea, A. ostoyae, M.
kentingensis, L. edodes* (but not in *P. ostreatus* and *A. bisporus*), primordium stages of *A.
ampla* and *S. commune* and during early development (primordia + young fruiting bodies) of
L. bicolor. A fourth CDE orthogroup, which contains reciprocal best hits to the putative E3
ubiquitin ligase protein *btb1* of *Sch. pombe* (*C. cinerea* protein ID: 545078), but contains also

domains that suggest it regulates chromatin condensation and thus gene expression.
Members of this orthogroup were upregulated typically late during development.

4.14. Other gene groups

4.14.1. Transporters

Among the conserved developmentally expressed genes (CDE orthogroups) we detected 49
putatively transport-related gene families. Many of these genes belong to the Major
Facilitator Superfamily or ABC transporters, fast-evolving gene families that are represented
by hundreds of genes in fungal genomes. Transporters have been reported in transcriptomic
studies of several mushroom-forming species, both in fruiting body(Song et al. 2018b;
Krizsán et al. 2019) and ectomycorrhiza formation(Xu et al. 2015). Due to the fast evolution
of these families, orthology to genes of Ascomycota model systems and even within the
Agaricomycetes is lacking in most cases, making it hard to infer their functions.
Nevertheless, some conserved transporter genes were detected, including orthologs of *S.*
cerevisiae Pns1 (choline transporter, *C. cinerea* 355745), *A. nidulans* MepA (ammonium
permease, *C. cinerea* 538158), *Sch. pombe* Kha1 (plasma membrane potassium ion/proton
symporter, *C. cinerea* 449705), *S. cerevisiae* Emp70 (endosomal transport protein, *C.*
cinerea 495380), *S. cerevisiae* Opt1 (plasma membrane oligopeptide transporter, *C. cinerea*
496395), among others (see Table 38). In the absence of functional annotations for the vast
majority of these genes, we list the genes in Table 38, but do not go into detailed analyses,
because it is unlikely that we can arrive at any reasonable speculation about their function.
Notable CDE orthogroups include an MFS transporter *C. cinerea* 465489, which was
developmentally regulated in 9/5 species (FC>2/4) and FB-init in *C. cinerea, A. ostoyae, P.*
ostreatus, S. commune (in the latter at FC>2 only). A phosphate transporter CDE orthogroup
(*C. cinerea* protein ID: 30849) was FB-init in *C. cinerea, C. aegerita, A. ostoyae, A. ampla, S.*
commune, M. kentingensis and, at FC>2 in *L. bicolor (*not in *L. edodes*) (Table 38). A group
of zinc-regulated transceptors (transporter + receptor, *C. cinerea* protein ID: 455616)
orthologous to *S. cerevisiae* Zrt1 was also detected. In summary, transporters can be
expected to fulfill key functions in fruiting bodies, in moving around goods, signal molecules,
modulating osmolarity to name a few, however, information about these in fruiting bodies is
still limited.
Aquaporins are water and solute-transporting membrane channels that are integral to
several different aspects of the fungal lifestyle. They were reported to be overexpressed in
primordia and stipes of *F. filiformis*, relative to other tissues and developmental stages(Liu et
al. 2020b). They were also reported to be upregulated in *L. bicolor* fruiting bodies and
ectomycorrhizae(Xu et al. 2015).
We found 2-11 aquaporin genes in the examined Agaricomycetes and observed
conserved aquaporin expression in two aquaporin orthogroups. One of these (*C. cinerea*
protein ID: 373141) contains putative orthologs of yeast Fps1, an aquaglyceroporin that has
suggested roles in acetate uptake or glycerol transport(Nehls and Dietz 2014), not water
uptake (see under chapter on Acetyl CoA). It is worth noting that aquaglyceroporins were
also differentially regulated in *L. bicolor* fruiting bodies(Xu et al. 2015) and that these
proteins were shown to have a capacity to transport various solutes, including NH_3 and
glycerol. This led Xu et al to hypothesize that these genes could be involved in osmosis
regulation or signaling(Xu et al. 2015), which is different from the conclusion we make above
on acetate transport.

The other orthogroup (represented by *S. commune* 2501677) comprises stipe-
specific aquaporin genes, which show similar expression trajectories in *A. bisporus, A.*
ostoyae, M kentingensis and *P. ostreatus*, as well as in an external *F. filiformis* dataset(Liu et
al. 2020b). In all of these species, the genes showed expression peaks in stipes, mostly in
young and mature fruiting bodies. It is notable that this pattern is seen in pileate-stipitate
species and has been noted in a previous 3-species comparison of transcriptomes(Plaza et
al. 2014b). Species without separate cap and stipe tissues either do not have this gene (*Pt.*
gracilis) or they do, but the gene is not developmentally regulated (*S. commune, Ph.*
chrysosporium). The current genome assembly of *C. cinerea* does not contain an orthologue
in this group, which might be because it became dispensable for the unique developmental
trajectory of this species or because the genome assembly is incomplete at this genomic
region. The marked stipe expression peak suggests a widespread role of water transport
during mushroom development. Further, the increasing expression parallel with
developmental progression could indicate a role in water supply needed for cellular
expansion. On a related note, *C. neoformans* also shows an upregulation of an aquaporin
gene (Cryne_JEC21_1_3081) late in its development.
A consequence of growth by cell expansion is that growth rate is limited by the speed
of water transport, not cell division and CW material deposition(Roper and Seminara 2019).
Therefore, aquaporin-facilitated water transport might be a selected trait in many mushroom-
forming fungi that undergo fast growth. Although unknown at this moment, if fruiting body
expansion is mediated by locally produced osmolytes (e.g. by breaking down glycogen),
then that could produce a gradient that draws water towards the expanding tissues. This
may be further facilitated by water channels formed by aquaporins. Finally, it is possible that
other physical solutions, e.g. inter-hyphal water passageways also help transporting (e.g. via
surface tension) water in the fruiting body.
The shared stipe upregulation of aquaporins can shed light on the origins of the
typical mushroom (pileate-stipitate) morphology. All examined stipitate Agaricales species
showed a stipe-specific aquaporin expression, which supports a single origin of pileate-
stipitate fruiting bodies in the Agaricales. On the other hand, *L. tigrinus,* which evolved
pileate-stipitate fruiting bodies independently in the Polyporales, does not show stipe-specific
aquaporin expression.

4.14.2. Stress response genes

Diverse stress-related genes were detected among CDE orthogroups. Based on functional
annotations and orthology relationships, many of these seem to be related to protection from
oxidative stress, either through direct protection by their protein products (peroxiredoxins) or
via the produced metabolites (ergothioneine) that protect from oxidative stress. We found
evidence for all three pathways of reactive oxygen detoxification, the peroxiredoxin pathway,
cytoplasmic catalases and glutathione peroxidases(Breitenbach et al. 2015), being active in
fruiting bodies. The dynamic expression of these genes may point to the existence of an
oxidative environment in fruiting bodies. However, it is currently unknown in which
tissues/stages and in what cellular contexts this may arise. One possibility is that the action
of cell wall modifying enzymes (e.g. peroxidases), which require an oxidative environment
(H_2O_2), results in excess reactive oxygen species, though this has not been confirmed yet
and remains a hypothesis for now. Below we discuss the main stress-related CDE
orthogroups which we detected in the developmental transcriptomes of Agaricomycetes.

Peroxiredoxins - Peroxiredoxins are antioxidant proteins that can reduce diverse
reactive oxygen species (ROS), including H_2O_2, and are one of the three cellular pathways
of ROS detoxification in fungi. They have also been reported to be involved in signal
transduction. The function of peroxiredoxins are not known in fruiting bodies and we
identified a single report of peroxiredoxin upregulation in fruiting bodies, in
Termitomyces(Rahmad et al. 2014). Another gene from *Antrodia camphorata* was cloned
and biochemically characterized(Wen et al. 2007). We found orthologs of peroxiredoxin Tpx1
of *Sch. pombe* and Tsa1 of *S. cerevisiae* (*C. cinerea* protein ID: 468305) to be
developmentally regulated in 12/8 species (FC>2/4) (Table 39). Tpx1 has been
characterized as the H_2O_2 sensor that relays the redox signal to the transcription factor Pap1
by inducing the formation of intramolecular disulfide bonds in Pap1(Jara et al. 2007).
Orthologs of Tpx1 are upregulated mostly in gill and cap tissues of mature fruiting bodies in
the examined species. Orthologs of the Pap1 transcription factor (represented by *C. cinerea*
378471) are expressed constantly in mushroom-forming fungi. Orthologs of yeast Ahp1
peroxiredoxin (*C. cinerea* protein ID: 444300) were also developmentally regulated in the
majority of species (10/4 species at FC=2/4, Table 39). This gene preferentially eliminates
organic peroxides, rather than H_2O_2 and relays signal to the Cad1 transcription factor in *S.*
cerevisiae(Iwai et al. 2010). In *C. cinerea* this gene shows a very strong induction in young
fruiting body gills, whereas in other species no evidence for a transcript enrichment in gill
tissues was found. Finally, we detected orthologs of *S. cerevisiae* Dot5 (*C. cinerea* protein
ID: 379600) (Table 39), a thioredoxin (thiol-specific peroxidase) that is involved in the
detoxification of peroxides(Cha et al. 2003; Izawa et al. 2003), but was also reported to be
involved in the regulation of telomere length. Thioredoxins, together with peroxiredoxins
(thioredoxin peroxidase) make up the thioredoxin system, which acts similarly to the
glutaredoxin system in ROS scavenging. This orthogroup showed a characteristic
upregulation in primordia relative to vegetative mycelium, then lower expression in late
developmental stages in *C. cinerea, L. edodes, C. aegerita, A. ostoyae* and *A. ampla*, but
not in other species.
Ergothionine biosynthesis - We also detected orthologs of the ergothioneine
biosynthesis protein Egt1 of *S. pombe*. Ergothioneine is a histidine-derived antioxidant
metabolite known to confer beneficial health effects to mushroom fruiting bodies(Weigand-
Heller et al. 2012; Kalaras et al. 2017). For eukaryotic cells, it might be needed for growth
and survival under oxidative stress conditions, for example during conidial survival and
germination(MH et al. 2012; Sheridan et al. 2016). Ergothioneine is produced via a two-step
reaction pathway in *S. pombe*(Pluskal et al. 2014)*,* comprising Egt1 then Egt2. Both are
conserved in *C. cinerea* (*C. cinerea* protein ID: 356399 and 258252, respectively, Table 39),
however, only the first gene is developmentally regulated in the majority of the species, the
other is not. Expression patterns of the Egt1 orthologs varied, in certain species they are FB-
init, while in others it showed stipe-upregulation (*A. ostoyae, L. bicolor, L. edodes, L. tigrinus,*
M. kentingensis, but not *C. cinerea*). How this can be explained is not clear, nevertheless,
our data provide some evidence for the upregulation of ergothioneine biosynthesis in fruiting
bodies. Orthologs of *Sch. pombe* Egt2, which catalyzes the second step of ergothioneine
production (*C. cinerea* protein ID: 258252) were developmentally regulated in a more
restricted set of species.
Dehydro-D-arabinono-1;4-lactone - Two orthogroups that make up the
biosynthesis pathway of dehydro-D-arabinono-1;4-lactone were developmentally expressed
in several species. Dehydro-D-arabinono-1;4-lactone (previously known as D-
erythroascorbate) is a protective metabolite against oxidative stress and is produced by a

pathway comprising Ara1, Ara2 and Alo1 in *S. cerevisiae*(K et al. 2006). Of these, Ara2 and
Alo1 have clear orthologs in Agaricomycetes (represented by *C. cinerea* 365550 and
432313, respectively)(Table 39), whereas Ara1 does not. Both Ara2 and Alo1 orthologs are
strongly upregulated in fruiting bodies. ALO1 orthologs showed a clear upregulation in stipe
tissues of *C. cinerea, A. ostoyae* and *P. ostreatus* (but not in *L. bicolor or L. edodes*),
whereas in *S. commune, A. ampla and L. edodes* it is upregulated in primordia and stays
high throughout fruiting body development. In *C. aegerita*, the gene is strongly upregulated
in the first primordium stage, then expression drops to low levels.
Catalases - Two catalase orthogroups (represented by *C. cinerea* 472224 and
420157)(Table 39) were developmentally regulated in the majority of species. Catalases are
involved in H_2O_2 scavenging and thus protection against reactive oxygen species, but also in
other pathways (such as beta-oxidation, see above), so their assignment to oxidative stress
response is less certain than that of the above-mentioned genes.
Glutathione system - The glutathione system is composed of (i) glutathione, a 3-
amino acid redox-active metabolite(Pócsi et al. 2004; Sato et al. 2009); (ii) glutathione
peroxidases, which reduce H_2O_2; (iii) glutathione reductases, which convert oxidized
glutathione back to its non-oxidized form; as well as (iv) glutaredoxins, which repair
oxidatively damaged proteins. In *S. cerevisiae* glutathione is produced by a two-step reaction
from L-cysteine and L-glutamate by the enzymes GSH1 and GSH2(CM 2001). We
annotated members of this pipeline in *C. cinerea* based on orthology information to
components of the glutathione system in *S. cerevisiae*(Pócsi et al. 2004; Sato et al. 2009)(
Table 39). We identified single-copy orthologs of two proteins in the glutathione synthesis
pathway, glutathione reductase, glutathione peroxidase. Glutaredoxins are present, on
average, as 5-6 distinct genes in Agaricomycetes genomes, except for *Pt. gracilis* and *M.*
kentingensis, which harbor 9 and 11 glutaredoxin genes in their genomes, respectively.
Unlike the other antioxidant systems, the expression patterns in the glutathione system show
only moderate conservation across species, being developmentally regulated in 3-6 species
(Table 39).
Glutathione-S-transferases are also involved in defense against oxidative damage.
They represent a diverse gene family also involved in other cellular detoxification tasks, such
as the neutralization of xenobiotics, drugs or pesticides(M et al. 2009b). Leon-Ramirez et al
hypothesized that in *U. maydis* 'basidiocarps' glutathione-S-transferases are involved in the
protection against oxidizing intracellular conditions generated by some enzymes, such as
NAPDH oxidases(CG et al. 2017).Therefore, we only marginally treat them under this
chapter, noting that any glutathione-S-transferase orthogroup, also by virtue of the lack of
clear orthology to functionally characterized genes from model species, could be related to
any of the possible functions of this family, not necessarily protection from oxidative damage.
Glutathione S-transferases are a diverse gene family in fungi, with 20-49 genes (on average
32) in the genomes of Agaricomycetes. In each of the species, ~50% of glutathione S-
transferase encoding genes (up to 70% in *Coprinopsis*) were developmentally regulated,
indicating that these play important roles in fruiting body development, however, due to the
general lack of conservation of these genes, these did not form CDE orthogroups.
Interestingly, each species had genes that showed a marked upregulation (FC>4) at the
initiation of fruiting body development.

4.14.3. Unannotated genes

Genes that encode proteins with no known conserved domain (Pfam or InterPro) signatures,
referred to as unannotated genes here, are abundant in fungal genomes and represent
some of the most exciting gene families. At the same time, they represent a challenge to link
to fruiting body development, due to the complete lack of functional clues. In general, few
literature reports are available on unannotated genes, possibly due to the difficulty of tackling
their functions. An exception from this is a report of the enrichment of unannotated genes in
mature peridioles of *Pisolithus microcarpus* fruiting bodies(Pereira et al. 2017). An example
of unannotated genes that turn out to be developmentally relevant is the LPMO-like X325
gene family(Labourel et al. 2020), which was circumscribed just recently and appears to be
related to the cell wall (see chapter 4.8.2.1).
We identified 176 CDE orthogroups of unannotated genes (Supplementary Table 6).
These included, for example, *Spc14* and *Spc33* of *S. commune*, which are involved in the
formation of the septal pore cap(Van Peer et al. 2010), are unannotated proteins but form
conserved, single-copy gene families in the Agaricomycetes (represented by *C. cinerea*
419716 and 473658) that was developmentally regulated in 10/6 and 12/5 species,
respectively. In *C. cinerea* alone, we identified 32 FB-init unannotated genes, several of
which showed very significant fold changes (up to 150-200x).
This category of genes included *Ctg1* of *L. edodes* and *Cc.Ctg1* of *C. cinerea*. Ctg1
was originally identified in *L. edodes* as a target of the pre-mRNA splicing factor Cdc5 and
was later shown to influence stipe elongation in *C. cinerea*(Nakazawa et al. 2008). In the
latter study knockdown experiments were unsuccessful, whereas overexpression *C. cinerea*
mutants showed accelerated stipe elongation(Nakazawa et al. 2009). In our data *Ctg1*
orthologs formed a CDE orthogroup (represented by *C. cinerea* 528942) which was
developmentally regulated in 10/8 species (FC>2/4)(Supplementary Table 6). Despite the
conservation of developmental regulation, expression profiles were only partially conserved
across species. For example, the gene was upregulated in mature fruiting bodies in *C.*
aegerita, A. ampla, C. cinerea, L. edodes, P. ostreatus, whereas in other species (e.g. *A.*
ostoyae), it showed upregulation in primordia relative to vegetative mycelium. Overall, CTG1
orthologs seem to be a very exciting group of developmental genes, with no conserved
domains and no mechanistic information on how they might contribute to fruiting body
formation. Its role in stipe elongation needs to be conserved in other species, if expression
profile conservation correlates with phenotype, we may expect other phenotypic effects in
other species.
In summary, unannotated genes may contain very interesting, novel gene families
involved in fruiting body development, yet circumscribing their functions remains challenging
at the moment and will require extensive genetic manipulation techniques.

4.14.4. Functionally poorly known genes

In this group of genes we placed ones that had some (mostly automatic) annotations, but we
could not clearly link them to cellular or fruiting-related processes. Thus, this category differs
from unannotated genes in that unannotated genes are completely devoid of all (even
automated) annotations. Altogether, we assigned 182 CDE orthogroups to this category
(Supplementary Table 7). Some of these genes are discussed briefly below, but we note that
many more interesting, fruiting-related genes are potentially hidden among the 182
orthogroups, however, clarifying their roles is left for future studies.
4533

Cytochrome p450 family - We discuss cytochrome p450s in this category, because they participate in diverse functions, from secondary metabolite biosynthesis (as part of gene clusters, e.g.(Nofiani et al. 2018)) to unknown cellular pathways mediating stipe elongation(Muraguchi and Kamada 2000). We detected 8 CDE orthogroups of cytochrome p450s, most of which are functionally unknown. Among the cytochrome p450s, *C. cinerea* Eln2 should be mentioned. This gene was identified associated with a dominant mutation that confers an elongationless stipe phenotype(Muraguchi and Kamada 2000). Eln2 was reported to be a constitutively expressed cytochrome p450. In accordance with this, we did not find considerable expression dynamics on the orthogroup containing eln2 (represented by *C. cinerea* 381558). However, the developmentally regulated orthogroups may be worthy targets of functional analyses in the future.

GT25 family - This family includes enzymes with diverse activities (lipopolysaccharide β-1,4-galactosyltransferase, β-1,3- or β-1,2-glucosyltransferase). In bacteria, GT25 are responsible for the biosynthesis of the membrane component lipopolysaccharides(Bertani and Ruiz 2018) ('endotoxins'). Most known GT25 enzymes are of bacterial origin, we found no functional study of fungal genes to date. The examined Agaricomycetes had mostly a single gene encoding GT25, which showed sporulation-associated expression peaks in most species: young fruiting body gills in *C. cinerea*, fruiting body gills in *P. ostreatus*, fruiting body cap in *L. edodes* and *C. aegerita*, mature fruiting bodies in *A. ampla* and *S. commune*, etc.. GT25 encoding genes showed very low expression in general, in many cases only detectable in the peak expressing stage/tissue. The broader orthogroup (*C. cinerea* protein ID: 501720) that contains the detected GT25 genes is specific to the Basidiomycota (result not shown), which could indicate bacterial origin into the phylum. The function of these proteins in fruiting bodies is currently unknown, however, expression peaks coincident with sporulation suggest they are related to late developmental events.

UstYa-like mycotoxin biosynthesis proteins - this gene family was first reported to be involved in the synthesis of the ribosomally synthesised peptide toxin ustiloxin in the ascomycete *Ustilaginoidea virens*(T et al. 2015). This family of genes was found to participate in the production of other toxic cyclic peptides, collectively named 'dikaritins', in plant pathogenic fungi (Ding et al. 2016; Vogt and Künzler 2019). Biochemically, they are oxidases of ribosomally produced peptides, including ustiloxins but also alpha-pheromones of *S. cerevisiae* (collectively Kex2-processed repeat proteins, KEPs)(Umemura 2020). Using a simple domain-based search, we detected 2-16 (28 in *A. ostoyae*) genes in the examined Agaricomycetes. Of these, each species had one to multiple genes that showed an induction (>4fold) in primordia relative to vegetative mycelia (except *C. cinerea*, in which two genes were induced in young fruiting body caps), and high expression in all subsequent fruiting body stages, suggesting that their protein product has an increased abundance in fruiting bodies. Although the role of these genes in the Basidiomycota remains unknown for now, we speculate they may be responsible for the production of toxic compounds, possibly with a role in defense.

Carbonic anhydrases (CA) - convert CO_2 to HCO_3^- which is then used in various ways by eukaryotic cells. CAs have been shown to be involved in fruiting body formation: CA knockout strains of the ascomycete *S. macrospora* showed slower vegetative growth and delayed fruiting body formation or defects in ascospore germination(Elleuche and Pöggeler 2009, 2010). The delay in fruiting body formation suggests a lower rate of HCO_3^- buildup necessary for some as yet unknown downstream processes, that, however, can be made up in longer time. Similar roles of CAs have not yet been reported in Basidiomycota, however,

they would be worth investigating as many CAs appeared developmentally regulated in
fruiting bodies. We found 2-9 CA encoding gene in the examined Agaricomycetes and that
each species had at least one developmentally regulated carbonic anhydrase gene,
although clear patterns of expression conservation were not noticeable. We found one CDE
orthogroup (*C. cinerea* protein ID: 450047) that was widely developmentally regulated in
mushroom-forming fungi (8/5 species at FC2/4), showing diverse expression peaks in
different species. It should be noted that CA expression levels potentially react also to
ambient CO_2 concentrations, so some of the expression variation we encountered may be
due to the growth conditions used during fruiting.
CipB gene family - This category of genes included the *cipB* gene reported from *C.*
cinerea(Nakazawa et al. 2009). The *cipB* gene was first identified as an interacting partner of
the splicing factor cdc5 in *L. edodes*(Nakazawa et al. 2008). We found that the orthogroup
that contained *cipB* orthologs (represented by *C. cinerea* 401749) was developmentally
regulated in 10/2 species (FC>2/4), indicating that these genes have widespread although
moderate expression dynamics (Supplementary Table 7). Based on conserved domain
annotations, *cipB* genes are transmembrane ATP synthases located in the mitochondria; its
contribution to fruiting body formation is not known at the moment.

4.14.5. Ferric reductase superfamily proteins and iron metabolism

The Ferric Reductase Domain Superfamily comprises a conserved group of proteins that
include ferric reductases and NAPDH oxidases, both of which are located in the plasma
membrane(Zhang et al. 2013). Ferric reductases are part of a reductive iron uptake system
which, in many model fungi, are involved in maintaining iron and potentially also copper
homeostasis(Gaurav Bairwa et al. 2017). These genes have mostly been studied in the
context of iron acquisition of pathogenic fungi from their hosts. Agaricomycetes don't share
clear orthology with experimentally characterized ferric reductases of the Ascomycota
(reviewed in(Gaurav Bairwa et al. 2017))(cite Hegedus et al 2021). Nevertheless, we
identified eight ferric reductase orthogroups, of which two were developmentally regulated in
multiple species (Table 40), suggesting that ferric reductases have conserved roles in
fruiting body development, although their function in the process is unknown. One of these
was developmentally expressed in 9/5 species (represented by *C. cinerea* 354442, FC>2/4)
contains genes orthologous to *S. cerevisiae* FRE7, which might be involved in copper
homeostasis in *S. cerevisiae*(C et al. 2000). Another orthogroup, which contains orthologs of
A. nidulans NoxA and *N. crassa* nox-1 (represented by *C. cinerea* 260659), was
developmentally regulated in 5/1 species. This gene is a NADPH oxidase, which has been
shown to be involved in developmental signaling and tissue differentiation(Nguyen et al.
2017).
The Ferric reductase superfamily also includes NADPH oxidases (NOXs)(Zhang et
al. 2013), which are reactive oxygen-generating cell membrane proteins involved in
differentiation-related signaling processes and tissue patterning(Blackstone). NOX genes
and their regulator NoxR were reported to have been lost repeatedly in yeast lineages,
underscoring their role in multicellularity(Nguyen et al. 2017). There is evidence for NOXs
regulating sexual, but not asexual or vegetative development in *A. nidulans*(T et al. 2003)
and *G. lucidum* and there is expression evidence in *H. marmoreus*(Zhang et al. 2015b). We
identified two putative NOX orthogroups in Agaricomycetes (*C. cinerea* protein ID: 461481,
260659). A third (represented by *C. cinerea* 152927) was interpreted as a NOX by León-
Ramírez et al(CG et al. 2017), but appears to be a ferric reductase. Despite clear reports of

NOXs being induced in sexual structures of Ascomycota(T et al. 2003), we did not find
evidence for consistent induction of these genes in fruiting bodies, rather, expression
patterns vary among fruiting body forming fungi. NOX genes show modest expression
dynamics in mushroom forming fungi (2<FC<4, Table 40). *C. cinerea* 448679 appears to be
a single-copy homolog of noxR that is conserved among mushroom-forming fungi. Members
of this orthogroup are not developmentally expressed.
4634

4635 5. A synoptic model of fruiting body development

This study aimed to provide a catalog of conserved fruiting body development genes based
on a literature review and a meta-analysis of developmental transcriptomes of 12
mushroom-forming fungi. Based on sequence analyses of genes showing developmentally
expression during fruiting body morphogenesis in these 12 species, we identified 921
conserved developmentally expressed orthogroups and several gene families. To more
broadly explore pathways linked to these genes, as well as genes that are known to be
involved in fruiting but do not show significant expression dynamics, we analyzed further 558
gene families. As a result, altogether 1479 genes of *C. cinerea*, and the orthologs of these in
11 other species are treated in this paper. These allowed us to identify developmental
expression patterns that are shared across species. For most of the conserved
developmental orthogroups, we provided broad functional classifications and speculations
on their probable role in fruiting body development, although elucidating their exact function
and molecular mechanisms of their action remains for future studies. In this chapter we
synthesize these data into an updated, synoptic model of fruiting body development in the
Agaricomycetes (Fig. 15). This model strongly builds on work by previous scholars, who
attempted to provide overviews of developmental and/or molecular mechanisms of
mushroom formation. In a series of papers Moore provided detailed descriptions of
developmental events based on morphological and biochemical analyses(Moore 2005).
Later, Kues summarized knowledge on the life history of *C. cinerea* and details of its
developmental biology(Kües 2000; Kües and Liu 2000). More recently, de Plaza et al, Ohm
et al and Muraguchi et al provided examples of transcription networks active during
development and a molecular overview of *C. cinerea* development based on gene ontology,
respectively(Ohm et al. 2011; Plaza et al. 2014b; Muraguchi et al. 2015). This paper further
elaborates these models with more detail, and a selective presentation of only those
processes that, based on gene expression patterns are conserved across twelve
Agaricomycetes species. We hereafter discuss developmental events in three phases: early
events (primordium formation and cell proliferation), growth by cell expansion and late
events (sexual processes, sporogenesis).

4664 5.1. Early events: primordium formation and cell proliferation

Expression patterns of developmentally regulated gene groups outline several distinct
processes and a temporal sequence of cellular events in fruiting bodies. The first in this
sequence and probably the most interesting is the initiation of fruiting body development,
which involves the rearrangement of hyphal branching patterns, and the light-dependent
formation of a primary, then a compact, secondary hyphal knot(Kües 2000; Liu 2005;
Sakamoto et al. 2018). Hyphal knot formation represents a transition from loosely arranged
hyphae with fractal dimensions to true complex multicellular structures(Nagy et al. 2018).

Fruiting body initiation has been a subject of intense research. Several gene expression
changes have been described in this transition, including light-regulated(Sakamoto et al.
2018), defense (e.g. galectins, pore-forming toxins(Plaza et al. 2014b), lectins(Boulianne et
al. 2000; Walser et al. 2005)), fatty acid biosynthesis(Liu 2005), transcriptional regulation
(TFs, velvet proteins(Ohm et al. 2011; Plaza et al. 2014b)), cell surface (hydrophobins, PriA,
fasciclins(S et al. 1992; Y et al. 2007)), cell wall (Kre9/Knh1, laccases(Kües et al. 2011)) as
well as others(Liu 2005; Hou et al. 2019). We update these observations by several
functionalities in this work.
The largest gene group that shows an upregulation in primordium samples is related
to mitotic cell division (See Fig. 3 in chapter 4.1.1 above), which coincides with tissue
differentiation in the primordium and produces the final number of cells that makes up the
fruiting body. Ribosomal and chromatin-remodeling related gene (e.g. histones) expression
show very similar expression profiles in primordia (see Fig. 4 and 10 in chapter 4.1.2 and
4.5.4.1), suggesting that novel protein synthesis and chromatin remodeling are required cell
proliferation/division in primordia. Chromatin remodeling might be required for mitotic
chromosome condensation and/or for rearranging chromatin accessibility for fruiting-specific
trans regulatory factors. We term this the proliferative phase of development, which, in *C.*
cinerea, encompasses hyphal knots to stage 2 primordia (Fig. 15). This pattern of mitotic,
histone and ribosomal gene expression has been observed in most species, suggesting that
fruiting body development can, in general, be divided into proliferative phase and growth by
cell division. However, some species (e.g. *A. bisporus*(Craig et al. 1977)*,* see chapter 4.1.3)
might not fully conform to this pattern and we expect non-Agaricales species (e.g.
Polyporales, bracket fungi) to not follow this rule.
While the vegetative mycelium is entangled in the substrate and sensitive to bacterial
and fungal competitors, fruiting bodies grow above ground and are, thus, more exposed to
more extreme environmental conditions and biotic stress (animals grazers or infections).
Accordingly, It has been reported that mycelial- and fruiting body-expressed defense gene
repertoires show little overlap, indicating a dramatic switch in defense strategy upon
fruiting(Plaza et al. 2014b). Defense genes generally show little conservation and high
phylogenetic patchiness and did not form conserved orthogroups in our analyses. We,
therefore, discussed them in the context of broader gene families (for detailed reviews
see(Kunzler 2015; Künzler 2018)). Putative secondary metabolism and oxylipin-related
genes might also serve defense and/or antimicrobial purposes.
Probably as a response to differences in environment compared to the substrate,
such as higher temperature or humidity fluctuations, several cell surface protein-encoding
and fatty acid metabolism genes showed induction in primordia. These included
hydrophobins, cerato-platanins, wax synthases and the Con6 family, among others.
Hydrophobins probably help retaining water or provide protection from excess water above-
ground(Lugones et al. 1999b). Wax synthases and Con6 proteins, although haven't been
examined to date, might act in a similar way. We infer changes in cell membrane
composition in primordia from expression changes in genes influencing membrane
composition. For example, fatty acid desaturases related to linoleic acid biosynthesis are
upregulated in fruiting bodies, probably contributing to the higher linoleic acid contents of
fruiting bodies reported before(SHAW 1967; Song et al. 2018a), with potentially to oxylipin
synthesis too(A et al. 2020)). Putative stress-related gene expression changes
(peroxiredoxins and ergothioneine-biosynthesis genes) might also correspond to genetically
encoded mechanisms for coping with adverse above-ground conditions.

Cell wall remodeling emerges as a potentially important process in fruiting body
initiation both in this study and previous ones (e.g.(Liu et al. 2021)). Fruiting body-specific
cell wall architectures exist and are probably produced by a diverse suite of lytic and cell wall
remodeling enzymes that are induced in primordia (Fig. 15). From the array of conserved
orthogroups, it appears that glucan remodeling is more important than chitin modification.
Additionally, oxidative enzyme genes (HTPs, DyP, MCOs, AA5_1) seem to be important in
primordium development, although their functions have not been elucidated, yet. Cell wall
related gene families show conservation at the gene family level, not always at the level of
strict CDE orthogroups. Therefore, the complete array of cell wall remodeling enzymes
induced in primordia is much broader in any species than shown on Fig. 15. Several cell wall
remodeling related gene families described from the Ascomycota were developmentally
regulated in the Agaricomycetes as well (e.g. GH16/17, Kre9/Knh1), whereas some, that are
central to Ascomycota cell wall remodeling showed constant expression (e.g. GH72) or are
missing in the Agaricomycetes (e.g. GH76). Further, it appears that dynamic expression is
mostly observed in cell wall-active enzyme genes, not in upstream (e.g. ER-localized)
components of the cell wall assembly line.
Although the light-dependency of fruiting body initiation differs from species to
species(Sakamoto 2018), we detected some orthogroups that probably correspond to light
reception. These include phytochromes, the *C. cinerea* Dst2 family and potentially PAS
domain containing proteins. The latter is particularly interesting, given that homologs in *N.*
crassa regulate the white collar complex and mediate adaptation to changing light intensities
during the day(Fuller et al. 2015).
The initiation of fruiting body development represents a transition from fractal-like to
complex multicellularity. Accordingly, several complex multicellular processes, such as cell-
to-cell communication, adhesion and the regulation of differentiation(Knoll 2011; Nagy et al.
2018) should turn on in primordia. Like for previous studies, decoding these aspects proved
difficult for us too. We expect regulatory factors, such as transcription factors, kinases,
phosphatases, potentially F-box proteins or non-proteinogenic factors (e.g. small RNA) to
participate in cellular differentiation. We and previous work detected some transcription
factor, kinase, NOX, pheromone, F-box orthogroups upregulated in primordia. However,
these generally did not show significant conservation of expression, so the main regulatory
mechanisms of cellular differentiation in fruiting bodies remain an enigma. It further remains
to be established if differentiation mechanisms described from plants and animals, such as
morphogens, hormone-like substances (e.g. oxylipins) or highly conserved gene regulatory
circuits exist in fungi. Similarly, adhesion-related proteins are mostly unknown at the
moment. Fasciclins and cell wall modification (e.g. by laccases) have been speculated to be
related to adhesion(Nagy et al. 2018; Merényi et al. 2020b), though glucan remodeling
enzymes, wax synthases, the carbohydrate-binding properties of lectins or species-specific
mechanisms could also be key and more research is needed to elucidate what genes glue
hyphae together in 3-dimension fruiting bodies.

5.2. Transition to growth by expansion

4759

At the stage 2 primordium stage (following nomenclature in *C. cinerea*, Fig. 15), cell and
tissue differentiation, cell proliferation has largely completed. This is reflected in the
downregulation of mitotic, ribosomal and chromatin-remodeling genes. At this point, the
fruiting body transitions to a different developmental mechanism, which we term 'growth by
cell expansion'. Growth by cell expansion is a well-known fungal trait and was already noted

by de Bary in 1887(Bary et al. 1887), however, the molecular details have remained
unknown. The phenomenon is marginally similar to fruit ripening in plants, but the
mechanisms are probably completely different, though both involve cell wall loosening,
remodeling and cell expansion.
Based on analyses of developmental transcriptomes, we detected what we believe is
a plausible molecular mechanism for growth by expansion. This involves the upregulation of
membrane biosynthesis genes and the rewiring of basic metabolic pathways for the
production of excess Acetyl-CoA (see chapter 4.2), which is the starting point of the
biosynthesis of membrane components (e.g. ergosterol). Upregulation of genes involved in
sterol transport, ergosterol biosynthesis correlates with the rewiring of basic metabolism. At
the same time, aquaporin gene expression increases in stipes, which probably supports
water transport driving the expansion of cells via turgor. Turgor manipulation requires an
osmolyte, however, despite speculations of trehalose or mannitol serving as osmolytes, its
identity remain unresolved; the expression data analyzed here did not provide unequivocal
evidence for any given osmotically active molecule in fruiting bodies. Trehalose, glycogen
and mannitol metabolism genes are upregulated at this stage, but many of them in gills,
which suggest an involvement in spore storage material production, rather than turgor
manipulation.
For enabling cell volume increases, the fungus deploys a diverse and mostly
species-specific array of cell wall remodeling enzymes. Cell expansion also probably follows
different mechanisms in each tissue. For example, according to the model of stipe
elongation in *C. cinerea* proposed by Liu et al(Liu et al. 2021), CE4 and GH30 genes may
contribute to the rigidification of the cell wall in the stipe base, causing it to be more resistant
to loosening by lytic enzymes than the stipe apex, thereby reducing its elongation potential.
It is reasonable to assume that similar mechanisms underlie the differential expansion
potential of different tissues within the fruiting body. We detected a large suite of conserved
CAZyme orthogroups upregulated in elongating stipes (Fig. 15), whereas only one
conserved orthogroup (GH55) was found in expanding caps. Since caps are composed of
multiple tissues, it is possible that RNA-Seq techniques with higher resolution (e.g. single-
cell RNA-Seq) are needed to uncover conserved tissue-specific cell wall remodeling gene
expression.
What regulatory genes underlie growth by cell expansion will need to be determined
in future studies. We detected conserved orthogroups of several developmentally expressed
regulators (transcription factors, RING-type zinc fingers, F-box, BTB and RNA-binding
proteins) but most showed tissue-specific expression patterns and no expression profile
clearly suggested involvement in regulating cell expansion in general. Interestingly, the Exp1
transcription factor reported to be responsible for cap expansion in *C. cinerea*(Muraguchi et
al. 2008b) was more frequently upregulated in stipes than in caps across the examined
species. We detected a red light-sensing phytochrome orthogroup upregulated in stipes,
which we hypothesize is responsible directing stipe growth away from closed spaces
towards the open air.

5.3. Late events: meiosis and spore production

4806

The young and mature fruiting body (in *C. cinerea* nomenclature) stages are the scene for
most sexual events, including meiosis and spore production. Accordingly, meiotic genes as
well as several genes involved in DNA replication/repair, chromatin remodeling, ribosome
assembly, which showed a peak in the proliferative phase in primordia, show a very sharp

expression peak associated with meiosis (see chapter 4.1.1, 4.1.2 and 4.5.4). We found
several conserved orthogroups of BTB/POZ proteins and RING-type zinc fingers with
expression peaks in gills; these might regulate key events associated with meiosis. A
conserved transcriptional co-repressor, *Cag1* of *C. cinerea* should be mentioned here.
Although this gene and its orthologs did not show significant expression dynamics, its
deletion caused abnormalities in gill development(Masuda et al. 2016).

Genes associated with storage carbohydrate metabolism were upregulated in gills in
most species, which is consistent with a rich body of literature reporting glycogen fluxes in
fruiting bodies(Ji and Moore 1993; Kües 2000). These genes included ones related to
glycogen mobilization, trehalose and mannitol metabolism. We found evidence for glycogen
synthesis (*Gsy1* orthologs), suggesting that this carbohydrate is packaged into spores.

Finally, in *C. cinerea* gills we detected an ortholog of the *A. nidulans* pigment-
producing polyketide synthase (PksB), this gene might be involved in spore melanin
production, although in other species this orthogroup showed different expression dynamics.

6. Outstanding questions related to fruiting body morphogenesis

Despite the advances in understanding cellular events of fruiting body morphogenesis,
several significant questions remain open and await further research. We below outline eight
key questions that we think are instrumental to develop a better understanding of fungal
fruiting body formation and complex multicellularity. This represents a subjective assessment
of significant challenges of fruiting body research, there may be other perspectives on
remaining key challenges to be solved.

1. Although new information on fruiting body morphogenesis accumulates at a steady
pace, several questions related to multicellular organization remain open. Fruiting bodies are
complex multicellular structures and thus display complex developmental programs involving
tissue differentiation and cell type specification. Besides a developmental program, adhesion
and cell-to-cell communication are two key traits for complex multicellularity(Knoll 2011).
While we are starting to understand transcriptional networks governing tissue differentiation
(see chapter 4.5.1), we still know very little about the genes/proteins mediating hypha-hypha
adhesion, communication or whether an extracellular matrix (other than the cell wall) exists
in mushroom-forming fungi. Hypotheses have been put forth on adhesion (via laccases or
fasciclins), however, these have not yet been tested experimentally. Tissue differentiation
results in distinct cell types within the fruiting body, however, our current understanding of
cell type diversity in mushroom-forming fungi is limited. All current estimates for cell type
diversity are based on morphology, which probably represent underestimates; technologies
like single-cell RNA-Seq might provide more insights into cell type classification.

2. Regarding regulatory genes, it is an open question whether 'master' regulators of
fruiting body development or tissue differentiation exists, similarly to animal and plant
developmental regulator systems(Meyerowitz 2002). Developmental processes are often
organized modularly and hierarchically so that the resulting modules have a high level of
autonomy. Such autonomy of mushroom development, to our best knowledge, has rarely
been documented. An exception from this is monokaryotic fruiting, the process of forming
basidiocarps on monokaryotic cultures without mating in certain species (e.g. (Herzog et al.
2016)). The fact that fruiting body initiation can bypass mating suggests that the process has
a certain degree of autonomy, which might be reflected as modularity or hierarchy in the

underlying regulatory networks. Moore et al characterized fruiting body morphogenesis as being made up of developmental subroutines that are somewhat independent of each other and that makes the whole process resilient to a certain degree of developmental imprecision(Loftus et al. 2020).

3. Transcriptional changes during the initiation of fruiting are significant. Most studies with a sufficient number of time points sampled (e.g.(Plaza et al. 2014b; Muraguchi et al. 2015; Sipos et al. 2017b; Almási et al. 2019; Krizsán et al. 2019; Merényi et al. 2021; Zhang et al. 2021d)) (Ruytinx et al in prep) found the largest number of differentially expressed genes at the transition from vegetative mycelium to the hyphal knot/primordium stage. However, whether this transcriptional rewiring event is purely a result of transcription factor activity or whether there are more general changes, for example, in chromatin accessibility or epigenetics, is unknown at the moment. Assays such as ATAC-Seq(Buenrostro et al. 2015) or ChIP-Seq (see e.g.(Vonk and Ohm 2021)) could provide insight into the magnitude of transcriptional versus chromatin-related rearrangement during fruiting body initiation. Similarly, posttranslational modification of proteins may be a significant regulatory mechanism in fruiting body morphogenesis(Pelkmans et al. 2017a). Gene methylation, an important developmental mechanism in animals, seems to be missing in fungi, on the other hand(Bewick et al. 2019).

4. Understanding tissue specificity of genes and encoded proteins is key to uncovering mechanisms of tissue differentiation and predicting gene function. The bulk expression studies and a few studies involving fluorescent protein tagging revealed numerous examples of tissue-specifically expressed genes; however, more systematic studies will be needed to understand tissue-specific expression and the mechanisms that generate it.

5. We identified altogether 176 conserved developmentally expressed orthogroups of unannotated genes. These genes encode proteins that contain no known protein domains, therefore, they lack even automated functional annotations. Unannotated genes are common in fungal genomes and represent an enigma for mycology. Their conservation implies that they have important functions these are, however, unknown in the vast majority of cases (see Spc genes of *S. commune* for an exception, chapter 4.14.3). Therefore, unannotated genes might represent a future treasure trove for developmental biology.

6. Growth by cell expansion is a widely conserved trait of mushroom-forming fungi. Above we provided evidence for this process involving excess Ac-CoA production, the upregulation of membrane biosynthetic processes and cell wall remodeling genes (chapter 4.2, 4.3). However, the identity of the osmolyte that drives turgor increase in fruiting bodies remains unknown and its identification represents a significant challenge for future research.

7. Although not strictly morphogenetic, how the fruiting body is supported by the mycelium is a question with important practical implications in mushroom industry. It is known that the directionality of the transport of goods in the mycelium switches as fruiting body development is initiated(Herman et al. 2020). Also, it has recently been suggested that secondary metabolite biosynthesis during fruiting body development depends also on the mycelium(Orban et al. 2021). Thus, transcriptional and metabolic reprogramming has to affect the entire fungal colony and not only the part where primordia emerge. The feeding and nutrient transport mechanisms as well as regulation of fruiting in relation to nutrient availability are exciting avenues of research, but beyond the primarily morphogenetic scope of the present paper.

8. While useful for discovering shared aspects of fruiting body morphogenesis, the comparative approach used by us, ignores species-specific aspects of morphology,

ecological adaptation, gene repertoires and expression dynamics. Non-conserved aspects of development are certainly important, particularly for understanding the biology of individual cultivated species and will deserve further study in the future. However, given the available information, making functional inferences for taxonomically restricted or species-specific genes is even harder than for conserved genes, therefore, we here restrict ourselves to the discussion of the latter.

7. Conclusions

The development of mushroom fruiting bodies is one of the most complex morphogenetic processes in fungi. A significant body of previous research has focused on understanding the underlying genetics and has made significant progress in cataloging genes related to the initiation, morphogenesis, defense, pigmentation of fruiting bodies, among others. However, available knowledge is still patchy and lacks key details in multiple aspects, such as conserved events in fruiting body morphogenesis and corresponding genes or key regulators of major differentiation steps.

This study aimed at providing a catalog of conserved fruiting body development genes based on a literature review and a meta-analysis of developmental transcriptomes of 12 mushroom-forming fungi. We identified gene orthogroups that show conserved developmental expression across multiple species. Despite some limitations of the genomic and transcriptomic approach used here, this study provided novel insights (such as gene families, orthogroups, conserved expression profiles) into the morphogenesis of mushroom fruiting bodies. We hope these data reveal some aspects of the core genetic program of agaricomycete fruiting body morphogenesis and will provide a roadmap for functional analyses of fruiting body development. We are aware that the summary provided here is still an incomplete survey of developmental genes in fruiting body forming fungi. Comparisons at finer scales, such as within economically relevant genera, denser sampling of time points during development or at better spatial resolution (e.g. single-cell RNA-Seq) could yield additional insights. Joint analyses of gene expression patterns with gene phylogenies or with gene family copy number dynamics, functional genomics assays, the inference of gene co-expression networks or population genomics (e.g. GWAS analyses) among others, could all provide more detailed views on the synergy between evolutionary changes in gene content, expression regulation and morphogenesis. We anticipate that with advances in functional genomics of Agaricomycetes, it will become easier to conduct such studies and shed light on conserved and species-specific aspects of fruiting body morphogenesis.

Acknowledgements

We are grateful to all friends and colleagues who, over the years, inspired our work on fruiting body development through collaborations or discussions. We appreciate the help of Levente Karaffa in making sense of Acetyl-CoA production and basic metabolism. We appreciate the permission of Francis Martin and Jason Tsai to use photographs of *L. bicolor* and *M. kentingensis* in Fig. 1, respectively. We are grateful to Annegret Kohler for providing pre-publication access to the transcriptome of *L. bicolor*. This work was supported by the European Research Council (grant no. 758161 to L.G.N.), the "Momentum" program of the Hungarian Academy of Sciences (contract No. LP2019-13/2019 to L.G.N.), the Hungarian National Research, Development, and Innovation Office (contract No. GINOP-2.3.2-15-

2016-00052), the Szeged Scientists Academy under the sponsorship of the Hungarian Ministry of Innovation and Technology (FEIF/646-4/2021-ITM_SZERZ, to B.Sz., A. Cs. And L.G.N.) and with the professional support of the Doctoral Student Scholarship Program of the Co-operative Doctoral Program of the Ministry of Innovation and Technology financed from the National Research, Development and Innovation Fund (KDP-17-4/PALY-2021 , to Cs.F.). T.V. was supported by the National Talent Programme (NTP-NFTÖ-21-B-0074) of the Hungarian Government. FH gratefully acknowledges funding from the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG) under grant HE 7849/3-1.

References

(PDF) The Igraph Software Package for Complex Network ResearchAvailable from https://www.researchgate.net/publication/221995787_The_Igraph_Software_Package_for_Complex_Network_Research.

2. Stem and cap morphogenesis in Coprinopsis fruit bodies.

A B., K V., R T., J P., D V., N F.-D., R F. 2005. The Aspergillus nidulans phytochrome FphA represses sexual development in red light. Curr. Biol. 15:1833–1838.

A B., R C., M M., L L.-V., L R., AG P., M P., L R. 2018. Transposon-associated epigenetic silencing during Pleurotus ostreatus life cycle. DNA Res. 25:451–464.

A I., J H. 2005. Photosensing fungi: phytochrome in the spotlight. Curr. Biol. 15.

A O., F H., M R. 2020. Volatilomes of Cyclocybe aegerita during different stages of monokaryotic and dikaryotic fruiting. Biol. Chem. 401:995–1004.

A P., V H., M S., M N., Š M., I P., S P., M Z., N R., P M., P V., J R., K S. 2019a. Pore-forming protein complexes from Pleurotus mushrooms kill western corn rootworm and Colorado potato beetle through targeting membrane ceramide phosphoethanolamine. Sci. Rep. 9.

A T., S A., R H., E V., S A., P L., P M., M R., F H., M K. 2019b. Heterologous Production and Functional Characterization of Ageritin, a Novel Type of Ribotoxin Highly Expressed during Fruiting of the Edible Mushroom Agrocybe aegerita. Appl. Environ. Microbiol. 85.

A V., E H.-G., M O., AL M., M B. 2009. Solution structure of hirsutellin A--new insights into the active site and interacting interfaces of ribotoxins. FEBS J. 276:2381–2390.

A V., SM B., A I., Varrot A., Basheer S.M., Imberty A. 2013. Fungal lectins: structure, function and potential applications. Curr. Opin. Struct. Biol. 23:678–685.

AA F., DI J., CC F., MH G., SR J., JA G., F T. 1991. Protein geranylgeranyltransferase of Saccharomyces cerevisiae is specific for Cys-Xaa-Xaa-Leu motif proteins and requires the CDC43 gene product but not the DPR1 gene product. Proc. Natl. Acad. Sci. U. S. A. 88:4448–4452.

Agger S., Lopez-Gallego F., Schmidt-Dannert C. 2009. Diversity of sesquiterpene synthases in the basidiomycete Coprinus cinereus. Mol. Microbiol. 72:1181–1195.

Ahmed Y.L., Gerke J., Park H.-S., Bayram Ö., Neumann P., Ni M., Dickmanns A., Kim S.C., Yu J.-H., Braus G.H., Ficner R. 2013. The Velvet Family of Fungal Regulators Contains a DNA-Binding Domain Structurally Similar to NF-$κ$B. PLOS Biol. 11:e1001750.

AL L., TK L., R B. 1999. Characterization of the Wsc1 protein, a putative receptor in the stress response of Saccharomyces cerevisiae. Genetics. 152:1487–1499.

Almási É., Sahu N., Krizsán K., Bálint B., Kovács G.M., Kiss B., Cseklye J., Drula E., Henrissat B., Nagy I., Chovatia M., Adam C., LaButti K., Lipzen A., Riley R., Grigoriev I. V., Nagy L.G. 2019. Comparative genomics reveals unique wood-decay strategies and fruiting body development in the Schizophyllaceae. New Phytol.

AM M., CS M., OP S., DS M., SN A., ME Z. 2009. Meiotic localization of Mre11 and Rad50 in wild type, spo11-1, and MRN complex mutants of Coprinus cinereus. Chromosoma. 118:471–486.

AN C., MD C. 2010. Basidiosporogenesis, meiosis, and post-meiotic mitosis in the ectomycorrhizal fungus Pisolithusmicrocarpus. Fungal Genet. Biol. 47:477–483.

Anderluh G., Maček P. 2003. Dissecting the Actinoporin Pore-Forming Mechanism.
Structure. 11:1312–1313.
Ando Y., Nakazawa T., Oka K., Nakahori K., Kamada T. 2013. Cc.snf5, a gene encoding a
putative component of the SWI/SNF chromatin remodeling complex, is essential for
sexual development in the agaricomycete Coprinopsis cinerea. Fungal Genet. Biol.
50:82–89.
Arima T., Yamamoto M., Hirata A., Kawano S., Kamada T. 2004. The eln3 gene involved in
fruiting body morphogenesis of Coprinus cinereus encodes a putative membrane
protein with a general glycosyltransferase domain. Fungal Genet. Biol. 41:805–812.
Arroyo J., Farkaš V., Sanz A.B., Cabib E. 2016. 'Strengthening the fungal cell wall through
chitin–glucan cross-links: effects on morphogenesis and cell integrity''.' Cell. Microbiol.
18:1239–1250.
Arsuffi G., Braybrook S.A. 2018. Acid growth: an ongoing trip. J. Exp. Bot. 69:137–146.
Ashikari T., Nakamura N., Tanaka Y., Kiuchi N., Shibano Y., Tanaka T., Amachi T.,
Yoshizumi H. 1986. Rhizopus raw-starch-degrading glucoamylase: Its cloning and
expression in yeast. Agric. Biol. Chem. 50:957–964.
ASM S., N S.-T., B L., RA O., PM H., K S., JJP B., A van P. 2020. Telomere-to-telomere
assembled and centromere annotated genomes of the two main subspecies of the
button mushroom Agaricus bisporus reveal especially polymorphic chromosome ends.
Sci. Rep. 10.
B F., M F., I N. 2001. Ethanol catabolism in Aspergillus nidulans: a model system for
studying gene regulation. Prog. Nucleic Acid Res. Mol. Biol. 69:149–204.
Baccelli I. 2015. Cerato-platanin family proteins: one function for multiple biological roles?
Front. Plant Sci. 5:769.
Baglivo I., Ragucci S., D'Incecco P., Landi N., Russo R., Faoro F., Pedone P. V, Maro A. Di.
2020. Gene Organization, Expression, and Localization of Ribotoxin-Like Protein
Ageritin in Fruiting Body and Mycelium of Agrocybe aegerita. Int. J. Mol. Sci. 2020, Vol.
21, Page 7158. 21:7158.
Bai Y., Wang Y., Liu X., Zhao J., Kang L.Q., Liu Z., Yuan S. 2020. Heterologous expression
and characterization of a novel chitin deacetylase, CDA3, from the mushroom
Coprinopsis cinerea. Int. J. Biol. Macromol. 150:536–545.
Baker L.G., Specht C.A., Donlin M.J., Lodge J.K. 2007. Chitosan, the Deacetylated Form of
Chitin, Is Necessary for Cell Wall Integrity in Cryptococcus neoformans. Eukaryot. Cell.
6:855.
Banks I.R., Specht C.A., Donlin M.J., Gerik K.J., Levitz S.M., Lodge J.K. 2005. A Chitin
Synthase and Its Regulator Protein Are Critical for Chitosan Production and Growth of
the Fungal Pathogen Cryptococcus neoformans. Eukaryot. Cell. 4:1902.
Barsottini M.R. de O., de Oliveira J.F., Adamoski D., Teixeira P.J.P.L., do Prado P.F. V,
Tiezzi H.O., Sforça M.L., Cassago A., Portugal R. V, de Oliveira P.S.L., Zeri A.C. de M.,
Dias S.M.G., Pereira G.A.G., Ambrosio A.L.B. 2013. Functional Diversification of
Cerato-Platanins in Moniliophthora perniciosa as Seen by Differential Expression and
Protein Function Specialization. http://dx.doi.org/10.1094/MPMI-05-13-0148-R.
26:1281–1293.
Bartoš O., Röslein J., Kotusz J., Paces J., Pekárik L., Petrtýl M., Halačka K., Štefková
Kašparová E., Mendel J., Boroń A., Juchno D., Leska A., Jablonska O., Benes V.,
Šídová M., Janko K. 2019. The Legacy of Sexual Ancestors in Phenotypic Variability,
Gene Expression, and Homoeolog Regulation of Asexual Hybrids and Polyploids. Mol.
Biol. Evol. 36:1902–1920.
Bary A. de, Bary A. de, Balfour I.B., Garnsey H.E.F., Garnsey H.E.F. 1887. Comparative
morphology and biology of the Fungi, Mycetozoa and bacteria. Oxford: Clarendon
press,.
Bayram Ö., Krappmann S., Ni M., Bok J.W., Helmstaedt K., Valerius O., Braus-Stromeyer
S., Kwon N.-J., Keller N.P., Yu J.-H., Braus G.H. 2008. VelB/VeA/LaeA Complex
Coordinates Light Signal with Fungal Development and Secondary Metabolism.
Science (80-.). 320:1504–1506.

Bayry J., Aimanianda V., Latge P. 2012. Hydrophobins—Unique Fungal Proteins. PLoS
Pathog. 8:4.
Berends E., Ohm R.A., De Jong J.F., Rouwendal G., Wösten H.A.B., Lugones L.G., Bosch
D. 2009. Genomic and biochemical analysis of N glycosylation in the mushroom-
forming basidiomycete Schizophyllum commune. Appl. Environ. Microbiol. 75:4648–
4652.
Berendsen R.L., Kalkhove S.I.C., Lugones L.G., Baars J.J.P., Wösten H.A.B., Bakker
P.A.H.M. 2013. Effects of the mushroom-volatile 1-octen-3-ol on dry bubble disease.
Appl. Microbiol. Biotechnol. 97:5535–5543.
Bertani B., Ruiz N. 2018. Function and biogenesis of lipopolysaccharides. EcoSal Plus. 8.
Bertossa R.C., Kües U., Aebi M., Künzler M. 2004. Promoter analysis of cgl2, a galectin
encoding gene transcribed during fruiting body formation in Coprinopsis cinerea
(Coprinus cinereus). Fungal Genet. Biol. 41:1120–1131.
Bewick A.J., Hofmeister B.T., Powers R.A., Mondo S.J., Grigoriev I. V., James T.Y., Stajich
J.E., Schmitz R.J. 2019. Diversity of cytosine methylation across the fungal tree of life.
Nat. Ecol. Evol. 3:479–490.
Bieszke J.A., Braun E.L., Bean L.E., Kang S., Natvig D.O., Borkovich K.A. 1999. The nop-1
gene of Neurospora crassa encodes a seven transmembrane helix retinal-binding
protein homologous to archaeal rhodopsins. Proc. Natl. Acad. Sci. U. S. A. 96:8034.
Blackstone N.W. Redox control and the evolution of multicellularity. .
Bleuler-Martínez S., Butschi A., Garbani M., Wîlti M.A., Wohlschlager T., Potthoff E., Sabotia
J., Pohleven J., Lüthy P., Hengartner M.O., Aebi M., Künzler M. 2011. A lectin-
mediated resistance of higher fungi against predators and parasites. Mol. Ecol.
20:3056–3070.
Bojunga N., Kötter P., Entian K.-D. 1998. The succinate/fumarate transporter Acr1p of
Saccharomyces cerevisiae is part of the gluconeogenic pathway and its expression is
regulated by Cat8p. Mol. Gen. Genet. MGG 1998 2605. 260:453–461.
Boontawon T., Nakazawa T., Horii M., Tsuzuki M., Kawauchi M., Sakamoto M., Honda Y.
2021. Functional analyses of Pleurotus ostreatus pcc1 and clp1 using CRISPR/Cas9.
Fungal Genet. Biol. 154:103599.
Borkovich K.A., Alex L.A., Yarden O., Freitag M., Turner G.E., Read N.D., Seiler S., Bell-
Pedersen D., Paietta J., Plesofsky N., Plamann M., Goodrich-Tanrikulu M., Schulte U.,
Mannhaupt G., Nargang F.E., Radford A., Selitrennikoff C., Galagan J.E., Dunlap J.C.,
Loros J.J., Catcheside D., Inoue H., Aramayo R., Polymenis M., Selker E.U., Sachs
M.S., Marzluf G.A., Paulsen I., Davis R., Ebbole D.J., Zelter A., Kalkman E.R.,
O'Rourke R., Bowring F., Yeadon J., Ishii C., Suzuki K., Sakai W., Pratt R. 2004.
Lessons from the genome sequence of Neurospora crassa: tracing the path from
genomic blueprint to multicellular organism. Microbiol. Mol. Biol. Rev. 68:1–108.
Boulianne R.P., Liu Y., Aebi M., Lu B.C., Kües U. 2000. Fruiting body development in
Coprinus cinereus: regulated expression of two galectins secreted by a non-classical
pathwayThe GenBank accession number for the sequence reported in this paper is
AF130360. Microbiology. 146:1841–1853.
Braus G.H., Irniger S., Bayram Ö. 2010. Fungal development and the COP9 signalosome.
Curr. Opin. Microbiol. 13:672–676.
Breitenbach M., Weber M., Rinnerthaler M., Karl T., Breitenbach-Koller L. 2015. Oxidative
Stress in Fungi: Its Function in Signal Transduction, Interaction with Plant Hosts, and
Lignocellulose Degradation. Biomolecules. 5:318.
van den Brink J., de Vries R.P. 2011. Fungal enzyme sets for plant polysaccharide
degradation. Appl. Microbiol. Biotechnol. 2011 916. 91:1477–1492.
Brodhun F., Feussner I. 2011. Oxylipins in fungi. FEBS J. 278:1047–1063.
Brown J.L., Bussey H. 1993. The yeast KRE9 gene encodes an O glycoprotein involved in
cell surface beta-glucan assembly. Mol. Cell. Biol. 13:6346–6356.
Brown N.A., Schrevens S., Van Dijck P., Goldman G.H. 2018. Fungal G-protein-coupled
receptors: Mediators of pathogenesis and targets for disease control. Nat. Microbiol.
Buenrostro J.D., Wu B., Chang H.Y., Greenleaf W.J. 2015. ATAC-seq: A method for

assaying chromatin accessibility genome-wide. Curr. Protoc. Mol. Biol. 2015:21.29.1--
21.29.9.
Buhian W.P., Bensmihen S. 2018. Mini-Review: Nod Factor Regulation of Phytohormone
Signaling and Homeostasis During Rhizobia-Legume Symbiosis. Front. Plant Sci.
0:1247.
Burns C., Stajich J.E., Rechtsteiner A., Casselton L., Hanlon S.E., Wilke S.K., Savytskyy
O.P., Gathman A.C., Lilly W.W., Lieb J.D., Zolan M.E., Pukkila P.J. 2010. Analysis of
the Basidiomycete Coprinopsis cinerea Reveals Conservation of the Core Meiotic
Expression Program over Half a Billion Years of Evolution. PLOS Genet. 6:e1001135.
Busch S., Eckert S.E., Krappmann S., Braus G.H. 2003. The COP9 signalosome is an
essential regulator of development in the filamentous fungus Aspergillus nidulans. Mol.
Microbiol. 49:717–730.
Busch S., Schwier E.U., Nahlik K., Bayram Ö., Helmstaedt K., Draht O.W., Krappmann S.,
Valerius O., Lipscomb W.N., Braus G.H. 2007. An eight-subunit COP9 signalosome
with an intact JAMM motif is required for fungal fruit body formation. Proc. Natl. Acad.
Sci. 104:8089–8094.
Buser R., Lazar Z., Käser S., Künzler M., Aebi M. 2010a. Identification, characterization, and
biosynthesis of a novel N-glycan modification in the fruiting body of the basidiomycete
Coprinopsis cinerea. J. Biol. Chem. 285:10715–10723.
Buser R., Lazar Z., Käser S., Künzler M., Aebi M. 2010b. Identification, Characterization,
and Biosynthesis of a Novel N-Glycan Modification in the Fruiting Body of the
Basidiomycete Coprinopsis cinerea. J. Biol. Chem. 285:10715–10723.
Butschi A., Titz A., Wälti M.A., Olieric V., Paschinger K., Nöbauer K., Guo X., Seeberger
P.H., Wilson I.B.H., Aebi M., Hengartner M.O., Künzler M. 2010. Caenorhabditis
elegans N-glycan Core β-galactoside Confers Sensitivity towards Nematotoxic Fungal
Galectin CGL2. PLoS Pathog. 6:e1000717.
C B., L D., C M.-D., A L., C S.-B., P K., D F., L P., JP S. 2004. A new lectin family with
structure similarity to actinoporins revealed by the crystal structure of Xerocomus
chrysenteron lectin XCL. J. Mol. Biol. 344:1409–1420.
C G., M K., VR I., PO B., DR W. 2000. Identification of the copper regulon in Saccharomyces
cerevisiae by DNA microarrays. J. Biol. Chem. 275:32310–32316.
C V.-F., C C., A K., E M., R B., J P., E D., P C., A W., RP de V., B H., F M. 2014. Genomic
and transcriptomic analysis of Laccaria bicolor CAZome reveals insights into
polysaccharides remodelling during symbiosis establishment. Fungal Genet. Biol.
72:168–181.
Caldwell G.A., Naider F., Becker J.M. 1995. Fungal lipopeptide mating pheromones: A
model system for the study of protein prenylation. Microbiol. Rev. 59:406–422.
Caligiuri M., Connolly T., Beach D. 1997. Ran1 functions to control the Cdc10/Sct1 complex
through Puc1. Mol. Biol. Cell. 8:1117.
Campbell L.T., Padula M.P., Harry E., Carter D.A., Campbell L.T., Padula M.P., Harry E.,
Carter D.A. 2015. You are what you secrete: extracellular proteins and virulence in
Cryptococcus. Microbiol. Aust. 36:93–95.
Cao L., Chan C., Lee C., Wong S.S., Yuen K. 1998. MP1 Encodes an Abundant and Highly
Antigenic Cell Wall Mannoprotein in the Pathogenic Fungus Penicillium marneffei.
Infect. Immun. 66:966.
Carman G.M., Han G.-S. 2011. Regulation of Phospholipid Synthesis in the Yeast
Saccharomyces cerevisiae. Annu. Rev. Biochem. 80:859.
Carmichael J.B., Provost P., Ekwall K., Hobman T.C., JB C., P P., K E., TC H. 2004. ago1
and dcr1, two core components of the RNA interference pathway, functionally diverge
from rdp1 in regulating cell cycle events in Schizosaccharomyces pombe. Mol. Biol.
Cell. 15:1425–1435.
Ceccaroli P., Buffalini M., Saltarelli R., Barbieri E., Polidori E., Ottonello S., Kohler A.,
Tisserant E., Martin F., Stocchi V. 2011. Genomic profiling of carbohydrate metabolism
in the ectomycorrhizal fungus Tuber melanosporum. New Phytol. 189:751–764.
CG L.-R., JL C.-P., D M.-S., A S.-A., ET A.-C., J R.-H. 2017. Transcriptomic analysis of

basidiocarp development in Ustilago maydis (DC) Cda. Fungal Genet. Biol. 101:34–45.

CH Z., JP Z., WQ D., TH L., ZD J. 2018. Molecular cloning and the expression pattern of AePOPB involved in the α-amanitin biosynthesis in Amanita exitialis fruiting bodies. Gene. 662:123–130.

Cha M.-K., Choi Y.-S., Hong S.-K., Kim W.-C., No K.T., Kim I.-H. 2003. Nuclear Thiol Peroxidase as a Functional Alkyl-hydroperoxide Reductase Necessary for Stationary Phase Growth of Saccharomyces cerevisiae*. J. Biol. Chem. 278:24636–24643.

Chae E., Tan Q.K.-G., Hill T.A., Irish V.F. 2008. An Arabidopsis F-box protein acts as a transcriptional co-factor to regulate floral development. Development. 135:1235–1245.

Chakraborty T.K., Basu D., Das N., Sengupta S., Mukherjee M. 2004. The mannitol cycle in Pleurotus ostreatus (Florida). FEMS Microbiol. Lett. 236:307–311.

Chang J., Chan P.L., Xie Y., Ma K.L., Cheung M.K., Kwan H.S. 2019a. Modified recipe to inhibit fruiting body formation for living fungal biomaterial manufacture. PLoS One. 14:e0209812.

Chang Y., Desirò A., Na H., Sandor L., Lipzen A., Clum A., Barry K., Grigoriev I. V., Martin F.M., Stajich J.E., Smith M.E., Bonito G., Spatafora J.W. 2019b. Phylogenomics of Endogonaceae and evolution of mycorrhizas within Mucoromycota. New Phytol. 222:511–525.

Chen D.D., Shi L., Yue S.N., Zhang T.J., Wang S.L., Liu Y.N., Ren A., Zhu J., Yu H.S., Zhao M.W. 2019. The Slt2-MAPK pathway is involved in the mechanism by which target of rapamycin regulates cell wall components in Ganoderma lucidum. Fungal Genet. Biol. 123:70–77.

Chen H., Kovalchuk A., Keriö S., Asiegbu F.O. 2017. Distribution and bioinformatic analysis of the cerato-platanin protein family in Dikarya. http://dx.doi.org/10.3852/13-115. 105:1479–1488.

Chen J., Li J.-M., Tang Y.-J., Ma K., Li B., Zeng X., Liu X.-B., Li Y., Yang Z.-L., Xu W.-N., Xie B.-G., Liu H.-W., Guo S.-X. 2020. Genome-wide analysis and prediction of genes involved in the biosynthesis of polysaccharides and bioactive secondary metabolites in high-temperature-tolerant wild Flammulina filiformis. BMC Genomics 2020 211. 21:1–16.

Chen L., Gong Y., Cai Y., Liu W., Zhou Y., Xiao Y., Xu Z., Liu Y., Lei X., Wang G., Guo M., Ma X., Bian Y. 2016. Genome Sequence of the Edible Cultivated Mushroom Lentinula edodes (Shiitake) Reveals Insights into Lignocellulose Degradation. PLoS One. 11:e0160336.

Chen S., Ma D., Ge W., Buswell J.A. 2003. Induction of laccase activity in the edible straw mushroom, Volvariella volvacea. FEMS Microbiol. Lett. 218:143–148.

Chen S., Xu J., Liu C., Zhu Y., Nelson D.R., Zhou S., Li C., Wang L., Guo X., Sun Y., Luo H., Li Y., Song J., Henrissat B., Levasseur A., Qian J., Li J., Luo X., Shi L., He L., Xiang L., Xu X., Niu Y., Li Q., Han M. V, Yan H., Zhang J., Chen H., Lv A., Wang Z., Liu M., Schwartz D.C., Sun C. 2012. Genome sequence of the model medicinal mushroom Ganoderma lucidum. Nat. Commun. 2012 31. 3:1–9.

Chen Y., Zhang Y., Siewers V., Nielsen J. 2015. Ach1 is involved in shuttling mitochondrial acetyl units for cytosolic C2 provision in Saccharomyces cerevisiae lacking pyruvate decarboxylase. FEMS Yeast Res. 15:1–8.

Cheng C.K., Au C.H., Wilke S.K., Stajich J.E., Zolan M.E., Pukkila P.J., Kwan H.S. 2013a. 5'-Serial Analysis of Gene Expression studies reveal a transcriptomic switch during fruiting body development in Coprinopsis cinerea. BMC Genomics. 14:195.

Cheng C.K., Au C.H., Wilke S.K., Stajich J.E., Zolan M.E., Pukkila P.J., Kwan H.S. 2013b. 5'-Serial Analysis of Gene Expression studies reveal a transcriptomic switch during fruiting body development in Coprinopsis cinerea. BMC Genomics. 14:195.

Chitarra G.S., Abee T., Rombouts F.M., Posthumus M.A., Dijksterhuis J. 2004. Germination of Penicillium paneum conidia is regulated by 1-octen-3-ol, a volatile self-inhibitor. Appl. Environ. Microbiol. 70:2823–2829.

Chiu E.Y.H., Lin Y.H., Wu W., Song Q., Chang P.F.L., Gao L.Y., Chou C.C., Ueng P.P. 2010. Alternative splicing and genetic diversity of the white collar-1 (wc-1) gene in

cereal Phaeosphaeria pathogens. Eur. J. Plant Pathol.

Cirauqui N., Abriata L.A., van der Goot F.G., Dal Peraro M. 2017. Structural,
physicochemical and dynamic features conserved within the aerolysin pore-forming
toxin family. Sci. Reports 2017 71. 7:1–12.

CM B., PJ E. 2010. Sterol regulatory element binding proteins in fungi: hypoxic transcription
factors linked to pathogenesis. Eukaryot. Cell. 9:352–359.

CM G. 2001. Role of the glutathione/glutaredoxin and thioredoxin systems in yeast growth
and response to stress conditions. Mol. Microbiol. 39:533–541.

Cohen-Kupiec R., Broglie K.E., Friesem D., Broglie R.M., Chet I. 1999. Molecular
characterization of a novel $β$-1,3-exoglucanase related to mycoparasitism of
Trichoderma harzianum. Gene. 226:147–154.

Combet E., Henderson J., Eastwood D.C., Burton K.S. 2006. Eight-carbon volatiles in
mushrooms and fungi: properties, analysis, and biosynthesis. Mycoscience. 47:317–
326.

Combet E., Henderson J., Eastwood D.C., Burton K.S. 2009. Influence of sporophore
development, damage, storage, and tissue specificity on the enzymic formation of
volatiles in mushrooms (agaricus bisporus). J. Agric. Food Chem. 57:3709–3717.

Corrochano L.M. 2011. Fungal photobiology: a synopsis. IMA Fungus 2011 21. 2:25–28.

Corrochano L.M. 2019. Light in the Fungal World: From Photoreception to Gene
Transcription and beyond. Annu. Rev. Genet. 53:149–170.

Costa Ana S.M.B.;Thomas D.J.I.⬚;Eastwoo. D.S.B.⬚;Baile. A.M.⬚;Foste. G.D.⬚;Mill.
P.R.⬚;Challe. M.P.. 2009. Quantifiable Downregulation of Endogenous Genes in
Agaricus bisporus Mediated by Expression of RNA Hairpins. J. Microbiol. Biotechnol.
19:271–276.

Costachel C., Coddeville B., Latgé J.-P., Fontaine T. 2005. Glycosylphosphatidylinositol-
anchored Fungal Polysaccharide in *Aspergillus fumigatus*. J. Biol. Chem. 280:39835–
39842.

Courty P.E., Hoegger P.J., Kilaru S., Kohler A., Buée M., Garbaye J., Martin F., Kües U.
2009. Phylogenetic analysis, genomic organization, and expression analysis of multi-
copper oxidases in the ectomycorrhizal basidiomycete Laccaria bicolor. New Phytol.
182:736–750.

Couturier M., Ladevèze S., Sulzenbacher G., Ciano L., Fanuel M., Moreau C., Villares A.,
Cathala B., Chaspoul F., Frandsen K.E., Labourel A., Herpoël-Gimbert I., Grisel S.,
Haon M., Lenfant N., Rogniaux H., Ropartz D., Davies G.J., Rosso M.-N., Walton P.H.,
Henrissat B., Berrin J.-G. 2018. Lytic xylan oxidases from wood-decay fungi unlock
biomass degradation. Nat. Chem. Biol. 2017 143. 14:306–310.

Craig G.D., Gull K., Wood D.A. 1977. Stipe elongation in Agaricus bisporus. Microbiology.
102:337–347.

Crestini C., Kovac B., Giovannozzi-Sermanni G. 1996. Production and isolation of chitosan
by submerged and solid-state fermentation from Lentinus edodes. Biotechnol. Bioeng.
50:207–210.

Cummings W.J., Merino S.T., Young K.G., Li L., Johnson C.W., Sierra E.A., Zolan M.E.
2002. The Coprinus cinereus adherin Rad9 functions in Mre11-dependent DNA repair,
meiotic sister-chromatid cohesion, and meiotic homolog pairing. Proc. Natl. Acad. Sci.
99:14958–14963.

D'Souza C.A., Heitman J. 2001. Conserved cAMP signaling cascades regulate fungal
development and virulence. FEMS Microbiol. Rev. 25:349–364.

D S., M N. 2014. The polyketide synthase gene pks4 is essential for sexual development
and regulates fruiting body morphology in Sordaria macrospora. Fungal Genet. Biol.
68:48–59.

Dai Y., Sun L., Yin X., Gao M., Zhao Y., Jia P., Yuan X., Fu Y., Li Y. 2019. Pleurotus eryngii
Genomes Reveal Evolution and Adaptation to the Gobi Desert Environment. Front.
Microbiol. 0:2024.

Dang Y., Yang Q., Xue Z., Liu Y. 2011. RNA Interference in Fungi: Pathways, Functions,
and Applications. Eukaryot. Cell. 10:1148.

Darwiche R., Mène-Saffrané L., Gfeller D., Asojo O.A., Schneiter R. 2017. The pathogen-
related yeast protein Pry1, a member of the CAP protein superfamily, is a fatty acid-
binding protein. J. Biol. Chem. 292:8304.
Das N., Sengupta S., Mukherjee M. 1997. Importance of laccase in vegetative growth of
Pleurotus florida. Appl. Environ. Microbiol. 63:4120–4122.
Dekker N., Speijer D., Grün C.H., Van Den Berg M., De Haan A., Hochstenbach F. 2004.
Role of the α-glucanase Agn1p in fission-yeast cell separation. Mol. Biol. Cell. 15:3903–
3914.
Deshaies R.J., Joazeiro C.A.P. 2009. RING Domain E3 Ubiquitin Ligases.
https://doi.org/10.1146/annurev.biochem.78.101807.093809. 78:399–434.
Deveau A., Kohler A., Frey-Klett P., Martin F. 2008. The major pathways of carbohydrate
metabolism in the ectomycorrhizal basidiomycete Laccaria bicolor S238N. New Phytol.
180:379–390.
Díaz-Valderrama J.R., Aime M.C. 2016. The cacao pathogen Moniliophthora roreri
(Marasmiaceae) possesses biallelic A and B mating loci but reproduces clonally.
Heredity (Edinb). 116:491–501.
Dijkgraaf G.J.P., Brown J.L., Bussey H. 1996. The KNHl Gene of Saccharomyces cerevisiae
is a Functional Homolog of KREO. 12:683–692.
Ding W., Liu W.-Q., Jia Y., Li Y., van der Donk W.A., Zhang Q. 2016. Biosynthetic
investigation of phomopsins reveals a widespread pathway for ribosomal natural
products in Ascomycetes. Proc. Natl. Acad. Sci. 113:3521–3526.
Dixon R.A., Achnine L., Kota P., Liu C.J., Reddy M.S.S., Wang L. 2002. The
phenylpropanoid pathway and plant defence - A genomics perspective. Mol. Plant
Pathol. 3:371–390.
Douglas L.M., Alvarez F.J., McCreary C., Konopka J.B. 2005. Septin function in yeast model
systems and pathogenic fungi. Eukaryot. Cell. 4:1503–1512.
Dressaire E., Yamada L., Song B., Roper M. 2016. Mushrooms use convectively created
airflows to disperse their spores. Proc. Natl. Acad. Sci. U. S. A.
Duan M., Bao H., Bau T. 2021. Analyses of transcriptomes and the first complete genome of
Leucocalocybe mongolica provide new insights into phylogenetic relationships and
conservation. Sci. Reports 2021 111. 11:1–12.
Dueñas-Santero E., Martín-Cuadrado A.B., Fontaine T., Latgé J.P., del Rey F., de Aldana
C.V. 2010. Characterization of glycoside hydrolase family 5 proteins in
Schizosaccharomyces pombe. Eukaryot. Cell. 9:1650–1660.
DW B., JM C., JR G., AA H., M H., B P.-D. 2002. The strobilurin fungicides. Pest Manag.
Sci. 58:649–662.
Eastwood D.C., Herman B., Noble R., Dobrovin-Pennington A., Sreenivasaprasad S., Burton
K.S. 2013. Environmental regulation of reproductive phase change in Agaricus bisporus
by 1-octen-3-ol, temperature and CO2. Fungal Genet. Biol. 55:54–66.
Eis C., Nidetzky B. 1999. Characterization of trehalose phosphorylase from Schizophyllum
commune. Biochem. J. 341:385.
Elleuche S., Pöggeler S. 2009. $β$-Carbonic Anhydrases Play a Role in Fruiting Body
Development and Ascospore Germination in the Filamentous Fungus Sordaria
macrospora. PLoS One. 4:e5177.
Elleuche S., Pöggeler S. 2010. Carbonic anhydrases in fungi. Microbiology. 156:23–29.
Emms D.M., Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol.
16:157.
Endo H., Kajiwara S., Tsunoka O., Shishido K. 1994. A novel cDNA, priBc, encoding a
protein with a Zn(II)2Cys6 zinc cluster DNA-binding motif, derived from the
basidiomycete Lentinus edodes. Gene. 139:117–121.
Essen L.-O., Vogt M.S., Mösch H.-U. 2020. Diversity of GPI-anchored fungal adhesins. Biol.
Chem. 401:1389–1405.
F T., I G., S V. 2005. Ejection mechanics and trajectory of the ascospores of Gibberella zeae
(anamorph Fuarium graminearum). Fungal Genet. Biol. 42:528–533.

F W., X S., X D., J Z., C D. 2017. DASH-type cryptochromes regulate fruiting body
development and secondary metabolism differently than CmWC-1 in the fungus
Cordyceps militaris. Appl. Microbiol. Biotechnol. 101:4645–4657.
Fakas S. 2017. Lipid biosynthesis in yeasts: A comparison of the lipid biosynthetic pathway
between the model nonoleaginous yeast Saccharomyces cerevisiae and the model
oleaginous yeast Yarrowia lipolytica. Eng. Life Sci. 17:292–302.
Fäldt J., Jonsell M., Nordlander G., Borg-Karlson A.K. 1999. Volatiles of bracket fungi
Fomitopsis pinicola and Fomes fomentarius and their functions as insect attractants. J.
Chem. Ecol. 25:567–590.
Fang H., Zhang W., Niu X., Liu Z., Lu C., Wei H., Yuan S. 2014. Stipe wall extension of
Flammulina velutipes could be induced by an expansin-like protein from Helix aspersa.
Fungal Biol. 118:1–11.
Fernandez Espinar M.-T., Labarère J. 1997. Cloning and sequencing of the Aa-Pri1 gene
specifically expressed during fruiting initiation in the edible mushroom Agrocybe
aegerita, and analysis of the predicted amino-acid sequence. Curr. Genet. 1997 326.
32:420–424.
Finley D., Ulrich H.D., Sommer T., Kaiser P. 2012. The Ubiquitin–Proteasome System of
Saccharomyces cerevisiae. Genetics. 192:319.
Fischer M., Cox J., Davis D.J., Wagner A., Taylor R., Huerta A.J., Money N.P. 2004. New
information on the mechanism of forcible ascospore discharge from Ascobolus
immersus. Fungal Genet. Biol. 41:698–707.
FJ A., JB K. 2007. Identification of an N-acetylglucosamine transporter that mediates hyphal
induction in Candida albicans. Mol. Biol. Cell. 18:965–975.
Floudas D., Binder M., Riley R., Barry K., Blanchette R.A., Henrissat B., Martínez A.T.,
Otillar R., Spatafora J.W., Yadav J.S., Aerts A., Benoit I., Boyd A., Carlson A.,
Copeland A., Coutinho P.M., Vries R.P. de, Ferreira P., Findley K., Foster B., Gaskell
J., Glotzer D., Górecki P., Heitman J., Hesse C., Hori C., Igarashi K., Jurgens J.A.,
Kallen N., Kersten P., Kohler A., Kües U., Kumar T.K.A., Kuo A., LaButti K., Larrondo
L.F., Lindquist E., Ling A., Lombard V., Lucas S., Lundell T., Martin R., McLaughlin
D.J., Morgenstern I., Morin E., Murat C., Nagy L.G., Nolan M., Ohm R.A.,
Patyshakuliyeva A., Rokas A., Ruiz-Dueñas F.J., Sabat G., Salamov A., Samejima M.,
Schmutz J., Slot J.C., John F. St., Stenlid J., Sun H., Sun S., Syed K., Tsang A.,
Wiebenga A., Young D., Pisabarro A., Eastwood D.C., Martin F., Cullen D., Grigoriev I.
V., Hibbett D.S. 2012. The Paleozoic Origin of Enzymatic Lignin Decomposition
Reconstructed from 31 Fungal Genomes. Science (80-.). 336:1715–1719.
Foster J., Nakata P.A. 2014. An oxalyl-CoA synthetase is important for oxalate metabolism
in Saccharomyces cerevisiae. FEBS Lett. 588:160–166.
Foulongne-Oriol M., Taskent O., Kües U., Sonnenberg A.S.M., van Peer A.F., Giraud T.
2021. Mating-Type Locus Organization and Mating-Type Chromosome Differentiation in
the Bipolar Edible Button Mushroom Agaricus bisporus. Genes (Basel). 12:1079.
François J., Parrou J.L. 2001. Reserve carbohydrates metabolism in the yeast
Saccharomyces cerevisiae. FEMS Microbiol. Rev. 25:125–145.
Fraser J.A., Hsueh Y.-P., Findley K.M., Heitman J. 2014. Evolution of the Mating-Type
Locus: The Basidiomycetes. Sex in Fungi. Washington, DC, USA: ASM Press. p. 19–
34.
Frawley D., Bayram Ö. 2020. The pheromone response module, a mitogen-activated protein
kinase pathway implicated in the regulation of fungal development, secondary
metabolism and pathogenicity. Fungal Genet. Biol. 144:103469.
Freihorst D., Fowler T.J., Bartholomew K., Raudaskoski M., Horton J.S., Kothe E. 2016. 13
The Mating-Type Genes of the Basidiomycetes. In: Wendland J., editor. Growth,
Differentiation and Sexuality. Cham: Springer International Publishing. p. 329–349.
Freiman R.N., Tjian R. 2003. Regulating the Regulators: Lysine Modifications Make Their
Mark. Cell. 112:11–17.
Freitag M. 2017. Histone Methylation by SET Domain Proteins in Fungi. Annu. Rev.
Microbiol. 71:413–439.

Froehlich A.C., Chen C.H., Belden W.J., Madeti C., Roenneberg T., Merrow M., Loros J.J.,
Dunlap J.C. 2010. Genetic and molecular characterization of a cryptochrome from the
filamentous fungus Neurospora crassa. Eukaryot. Cell. 9:738–750.
Fu Y., Dai Y., Yang C., Wei P., Song B., Yang Y., Sun L., Zhang Z.-W.W., Li Y. 2017.
Comparative Transcriptome Analysis Identified Candidate Genes Related to Bailinggu
Mushroom Formation and Genetic Markers for Genetic Analyses and Breeding. Sci.
Reports 2017 71. 7:1–12.
Fuller K.K., Loros J.J., Dunlap J.C. 2015. Fungal photobiology: visible light as a signal for
stress, space and time. Curr. Genet. 61:275.
FY L., NP K. 2014. Spatial and temporal control of fungal natural product synthesis. Nat.
Prod. Rep. 31:1277–1286.
G S., AS S., LJ van G. 2013. Development and growth of fruit bodies and crops of the
button mushroom, Agaricus bisporus. Fungal Biol. 117:697–707.
G V.-K., B W., SJ H., Z L., H Z., M S., VG E. 2010. An oxidative enzyme boosting the
enzymatic conversion of recalcitrant polysaccharides. Science. 330:219–222.
Gaderer R., Bonazza K., Seidl-Seiboth V. 2014. Cerato-platanins: A fungal protein family
with intriguing properties and application potential. Appl. Microbiol. Biotechnol.
98:4795–4803.
Gagnon-Arsenault I., Tremblay J., Bourbonnais Y. 2006. Fungal yapsins and cell wall: a
unique family of aspartic peptidases for a distinctive cellular function. FEMS Yeast Res.
6:966–978.
Gao W., Qu J., Zhang J., Sonnenberg A., Chen Q., Zhang Y., Huang C. 2018. A genetic
linkage map of Pleurotus tuoliensis integrated with physical mapping of the de novo
sequenced genome and the mating type loci. BMC Genomics. 19:18.
Garcia-Rubio R., de Oliveira H.C., Rivera J., Trevijano-Contador N. 2020. The Fungal Cell
Wall: Candida, Cryptococcus, and Aspergillus Species. Front. Microbiol. 0:2993.
Garcia-Santamarina S., Probst C., Festa R.A., Ding C., Smith A.D., Conklin S.E., Brander
S., Kinch L.N., Grishin N. V, Franz K.J., Riggs-Gelasco P., Leggio L. Lo, Johansen
K.S., Thiele D.J. 2020. A lytic polysaccharide monooxygenase-like protein functions in
fungal copper import and meningitis. Nat. Chem. Biol. 2020 163. 16:337–344.
Gaurav Bairwa, Jung W.H., W. Kronstad J. 2017. Iron acquisition in fungal pathogens of
humans. Metallomics. 9:215–227.
GB A., L K., OU S., V S.-S. 2015. Molecular diversity of LysM carbohydrate-binding motifs in
fungi. Curr. Genet. 61:103–113.
Gehrmann T., Pelkmans J.F., Lugones L.G., Wösten H.A.B., Abeel T., Reinders M.J.T.
2016. Schizophyllum commune has an extensive and functional alternative splicing
repertoire. Sci. Rep. 6.
Gehrmann T., Pelkmans J.F., Ohm R.A., Vos A.M., Sonnenberg A.S.M.M., Baars J.J.P.P.,
Wösten H.A.B.B., Reinders M.J.T.T., Abeel T. 2018. Nucleus-specific expression in the
multinuclear mushroom-forming fungus Agaricus bisporus reveals different nuclear
regulatory programs. 115:4429–4434.
Gerecke E.E., Zolan M.E. 2000. An mre11 mutant of Coprinus cinereus has defects in
meiotic chromosome pairing, condensation and synapsis. Genetics. 154:1125.
Gessler N.N., Filippovich S.Y., Bachurina G.P., Kharchenko E.A., Groza N. V., Belozerskaya
T.A. 2017. Oxylipins and oxylipin synthesis pathways in fungi. Appl. Biochem. Microbiol.
53:628–639.
Gilberti M., Baruffini E., Donnini C., Dallabona C. 2018. Pathological alleles of MPV17
modeled in the yeast Saccharomyces cerevisiae orthologous gene SYM1 reveal their
inability to take part in a high molecular weight complex. PLoS One. 13.
Gladfelter A.S. 2006. Control of filamentous fungal cell shape by septins and formins. Nat.
Rev. Microbiol. 2006 43. 4:223–229.
GM G., K R., MK O. 2008. The CAP superfamily: cysteine-rich secretory proteins, antigen 5,
and pathogenesis-related 1 proteins--roles in reproduction, cancer, and immune
defense. Endocr. Rev. 29:865–897.
Gold M.H., Cheng T.M. 1979. Conditions for fruit body formation in the white rot

basidiomycete Phanerochaete chrysosporium. Arch. Microbiol. 121:37–41.

Goldstein I.J., Winter H.C. 2007. Mushroom Lectins. Compr. Glycosci. From Chem. to Syst.
Biol. 3–4:601–621.

Gong W., Xie C., Zhou Y., Zhu Z., Wang Y., Peng Y. 2020. A Resequencing-Based
Ultradense Genetic Map of Hericium erinaceus for Anchoring Genome Sequences and
Identifying Genetic Loci Associated With Monokaryon Growth. Front. Microbiol. 0:3129.

Gonzalez-Hilarion S., Paulet D., Lee K.T., Hon C.C., Lechat P., Mogensen E., Moyrand F.,
Proux C., Barboux R., Bussotti G., Hwang J., Coppée J.Y., Bahn Y.S., Janbon G. 2016.
Intron retention-dependent gene regulation in Cryptococcus neoformans. Sci. Rep.

Goodrich-Tanrikulu M., Howe K., Stafford A., Nelson M.A. 1998. Changes in fatty acid
composition of Neurospora crassa accompany sexual development and ascospore
germination. Microbiology. 144:1713–1720.

Gow N.A.R., Latge J.-P., Munro C.A. 2017. The Fungal Cell Wall: Structure, Biosynthesis,
and Function. Microbiol. Spectr. 5.

Grau-Bové X., Ruiz-Trillo I., Irimia M. 2018. Origin of exon skipping-rich transcriptomes in
animals driven by evolution of gene architecture. Genome Biol.

Grąz M., Jarosz-Wilkołazka A., Janusz G., Mazur A., Wielbo J., Koper P., Żebracki K.,
Kubik-Komar A. 2017. Transcriptome-based analysis of the saprophytic fungus
Abortiporus biennis – response to oxalic acid. Microbiol. Res. 199:79–88.

Gressler M., Löhr N.A., Schäfer T., Lawrinowitz S., Seibold P.S., Hoffmeister D. 2021. Mind
the mushroom: natural product biosynthetic genes and enzymes of Basidiomycota. Nat.
Prod. Rep. 38:702–722.

Grützmann K., Szafranski K., Pohl M., Voigt K., Petzold A., Schuster S. 2014. Fungal
alternative splicing is associated with multicellular complexity and virulence: A genome-
wide multi-species study. DNA Res. 21:27–39.

GT W., MB Q., S C., F L.-G., C S.-D. 2012. Draft genome of Omphalotus olearius provides a
predictive framework for sesquiterpenoid natural product biosynthesis in
Basidiomycota. Chem. Biol. 19:772–783.

Gupta D.K., Rühl M., Mishra B., Kleofas V., Hofrichter M., Herzog R., Pecyna M.J., Sharma
R., Kellner H., Hennicke F., Thines M. 2018. The genome sequence of the
commercially cultivated mushroom Agrocybe aegerita reveals a conserved repertoire of
fruiting-related genes and a versatile suite of biopolymer-degrading enzymes. BMC
Genomics. 19:1–13.

Gyawali R., Upadhyay S., Way J., Lin X. 2017. A Family of Secretory Proteins Is Associated
with Different Morphotypes in Cryptococcus neoformans. Appl Env. Microbiol. 83.

H L., HE H.-A., JS S.-C., JD W. 2012. Ribosomal biosynthesis of $α$-amanitin in Galerina
marginata. Fungal Genet. Biol. 49:123–129.

H L., HT C. 2009. Enhanced expression of chitinase during the autolysis of mushroom in
Coprinellus congregatus. J. Microbiol. 47:225–228.

H L., Q C., Y L., X L., R S., HE H.-A., ZL Y. 2018. The MSDIN family in amanitin-producing
mushrooms and evolution of the prolyl oligopeptidase genes. IMA Fungus. 9:225–242.

H L., Q W., Y H., L C., C H., C J., Y L., Y D., Z K., JR X. 2016. Genome-wide A-to-I RNA
editing in fungi independent of ADAR enzymes. Genome Res. 26:499–509.

H M., D S., JE S., V B. 2015. Deciphering the uniqueness of Mucoromycotina cell walls by
combining biochemical and phylogenomic approaches. Environ. Microbiol. 17:1649–
1662.

H S., S K. 2003. A stearoyl-CoA-specific Delta 9 fatty acid desaturase from the
basidiomycete Lentinula edodes. Biosci. Biotechnol. Biochem. 67:2431–2437.

H S., S K. 2005. Cloning and functional characterization of a Delta12 fatty acid desaturase
gene from the basidiomycete Lentinula edodes. Mol. Genet. Genomics. 273:336–341.

Ha B., Kim S., Kim M., Ro H.S. 2018. Activation of the mating pheromone response pathway
of Lentinula edodes by synthetic pheromones. Mycobiology. 46:407–415.

Haber J.E. 2012. Mating-Type Genes and MAT Switching in Saccharomyces cerevisiae.
Genetics. 191:33–64.

Hallen H.E., Luo H., Scott-Craig J.S., Walton J.D. 2007. Gene family encoding the major

toxins of lethal Amanita mushrooms. Proc. Natl. Acad. Sci. 104:19097–19101.

Han S., Adams T.H. 2001. Complex control of the developmental regulatory locus brlA in Aspergillus nidulans. Mol. Genet. Genomics. 266:260–270.

Han S.E., Kwon H. Bin, Lee S.B., Yi B.Y., Murayama I., Kitamoto Y., Byun M.O. 2003. Cloning and characterization of a gene encoding trehalose phosphorylase (TP) from Pleurotus sajor-caju. Protein Expr. Purif. 30:194–202.

Hartl L., Zach S., Seidl-Seiboth V. 2011. Fungal chitinases: diversity, mechanistic properties and biotechnological potential. Appl. Microbiol. Biotechnol. 2011 932. 93:533–543.

Hassan M.A., Rouf R., Tiralongo E., May T.W., Tiralongo J. 2015. Mushroom lectins: specificity, structure and bioactivity relevant to human disease. Int J Mol Sci. 16:7802–7838.

Hastrup A.C.S., III F.G., Lebow P.K., Jensen B. 2012. Enzymatic oxalic acid regulation correlated with wood degradation in four brown-rot fungi. Int. Biodeterior. Biodegrad. , Vol. 75, 2012; pp. 109-114. . 75:109–114.

He Z., Long P., Fang F., Li S., Zhang P., Chen Z. 2020. Diversity of MSDIN family members in amanitin-producing mushrooms and the phylogeny of the MSDIN and prolyl oligopeptidase genes. BMC Genomics 2020 211. 21:1–18.

Heneghan M.N., Costa A.M.S.B., Challen M.P., Mills P.R., Bailey A., Foster G.D. 2007. A comparison of methods for successful triggering of gene silencing in Coprinus cinereus. Mol. Biotechnol. 2007 353. 35:283–296.

Herman K.C., Wösten H.A.B., Fricker M.D., Bleichrodt R.J. 2020. Growth induced translocation effectively directs an amino acid analogue to developing zones in Agaricus bisporus. Fungal Biol. 124:1013–1023.

Hernández-Rodríguez Y., Masuo S., Johnson D., Orlando R., Smith A., Couto-Rodriguez M., Momany M. 2014. Distinct Septin Heteropolymers Co-Exist during Multicellular Development in the Filamentous Fungus Aspergillus nidulans. PLoS One. 9:e92819.

Herranz S., Rodríguez J.M., Bussink H.J., Sánchez-Ferrero J.C., Arst H.N., Peñalva M.A., Vincent O. 2005. Arrestin-related proteins mediate pH signaling in fungi. Proc. Natl. Acad. Sci. U. S. A. 102:12141–12146.

Herzog R., Solovyeva I., Rühl M., Thines M., Hennicke F. 2016. Dikaryotic fruiting body development in a single dikaryon of Agrocybe aegerita and the spectrum of monokaryotic fruiting types in its monokaryotic progeny. Mycol. Prog. 15:947–957.

Heuts D.P.H.M., Janssen D.B., Fraaije M.W. 2007. Changing the substrate specificity of a chitooligosaccharide oxidase from Fusarium graminearum by model-inspired site-directed mutagenesis. FEBS Lett. 581:4905–4909.

Hibbett D.S. 2004. Trends in morphological evolution in homobasidiomycetes inferred using maximum likelihood: A comparison of binary and multistate approaches. Syst. Biol. 53:889–903.

HM H., DE N., J G., MS S. 2009. Evolutionary roles of upstream open reading frames in mediating gene regulation in fungi. Annu. Rev. Microbiol. 63:385–409.

Holighaus G., Rohlfs M. 2019. Volatile and non-volatile fungal oxylipins in fungus-invertebrate interactions. Fungal Ecol. 38:28–36.

Hong Jai-Sik;Kim T.-Y. 1988. Contents of Free-Sugars & Free-Sugaralcohols in Pleurotus ostreatus, Lentinus edods & Agaricus bisporus. Korean J. Food Sci. Technol. 20:459–462.

Hou L., Wang L., Wu X., Gao W., Zhang J., Huang C. 2019. Expression patterns of two pal genes of Pleurotus ostreatus across developmental stages and under heat stress. BMC Microbiol. 19:231.

Hou Z., Chen Q., Zhao M., Huang C., Wu X. 2020. Genome-wide characterization of the Zn(II)2Cys6 zinc cluster-encoding gene family in Pleurotus ostreatus and expression analyses of this family during developmental stages and under heat stress. PeerJ. 8:e9336.

Houbraken J., Dyer P.S. 2015. Induction of the Sexual Cycle in Filamentous Ascomycetes. :23–46.

Huang X., Zhang R., Qiu Y., Wu H., Xiang Q., Yu X., Zhao K., Zhang X., Chen Q., Penttinen

P., Gu Y. 2020. RNA-seq Profiling Showed Divergent Carbohydrate-Active Enzymes (CAZymes) Expression Patterns in Lentinula edodes at Brown Film Formation Stage Under Blue Light Induction. Front. Microbiol. 0:1044.

Huber S., Lottspeich F., Kämper J. 2002. A gene that encodes a product with similarity to dioxygenases is highly expressed in teliospores of Ustilago maydis. Mol. Genet. Genomics 2002 2676. 267:757–771.

Hung R., Lee S., Bennett J.W. 2014. The effects of low concentrations of the enantiomers of mushroom alcohol (1-octen-3-ol) on Arabidopsis thaliana. Mycology. 5:73–80.

Hutchins A.P., Liu S., Diez D., Miranda-Saavedra D. 2013. The Repertoires of Ubiquitinating and Deubiquitinating Enzymes in Eukaryotic Genomes. Mol. Biol. Evol. 30:1172–1187.

Hyun M.W., Yun Y.H., Kim J.Y., Kim S.H. 2011. Fungal and plant phenylalanine ammonia-lyase. Mycobiology. 39:257–265.

I A., I A., B de las R., R M., I R.-C., M M., P G., H T., IJ G., B P.-A., JM M. 2011. High-resolution structural insights on the sugar-recognition and fusion tag properties of a versatile $β$-trefoil lectin domain from the mushroom Laetiporus sulphureus. Glycobiology. 21:1349–1361.

Ismaya W.T., Tjandrawinata R.R., Rachmawati H. 2020. Lectins from the Edible Mushroom Agaricus bisporus and Their Therapeutic Potentials. Molecules. 25.

Iwai K., Naganuma A., Kuge S. 2010. Peroxiredoxin Ahp1 Acts as a Receptor for Alkylhydroperoxides to Induce Disulfide Bond Formation in the Cad1 Transcription Factor *. J. Biol. Chem. 285:10597–10604.

Izawa S., Kuroki N., Inoue Y. 2003. Nuclear thioredoxin peroxidase Dot5 in Saccharomyces cerevisiae: roles in oxidative stress response and disruption of telomeric silencing. Appl. Microbiol. Biotechnol. 2004 641. 64:120–124.

J F., F B., D H. 2017. Enzymatic Synthesis of Psilocybin. Angew. Chem. Int. Ed. Engl. 56:12352–12355.

J P., J B., L V., A L., A C., B S., J K., J S. 2011. Basidiomycete Clitocybe nebularis is rich in lectins with insecticidal activities. Appl. Microbiol. Biotechnol. 91:1141–1148.

J R.-R., M H., C K., S M., R F. 2010. Fungi, hidden in soil or up in the air: light makes a difference. Annu. Rev. Microbiol. 64:585–610.

J S., RA O., M K. 2016. Entomotoxic and nematotoxic lectins and protease inhibitors from fungal fruiting bodies. Appl. Microbiol. Biotechnol. 100:91–111.

JA C., RP O., SJ P., EA H. 2017. Olfactory Preferences of the Parasitic Nematode Howardula aoronymphium and its Insect Host Drosophila falleni. J. Chem. Ecol. 43:362–373.

JA D., NF L., SP J. 2000. A role for Saccharomyces cerevisiae histone H2A in DNA repair. Nature. 408:1001–1004.

James T.Y. 2015. Why mushrooms have evolved to be so promiscuous: Insights from evolutionary and ecological patterns. Fungal Biol. Rev. 29:167–178.

James T.Y., Lee M., van Diepen L.T.A. 2011. A Single mating-type locus composed of homeodomain genes promotes nuclear migration and heterokaryosis in the white-rot fungus phanerochaete chrysosporium. Eukaryot. Cell. 10:249–261.

Jara M., Vivancos A.P., Calvo I.A., Moldón A., Sansó M., Hidalgo E. 2007. The Peroxiredoxin Tpx1 Is Essential as a H2O2 Scavenger during Aerobic Growth in Fission Yeast. https://doi.org/10.1091/mbc.e06-11-1039. 18:2288–2295.

Ji J., Moore D. 1993. Glycogen metabolism in relation to fruit body maturation in Coprinus cinereus. Mycol. Res. 97:283–289.

Jia D., Wang B., Li X., Peng W., Zhou J., Tan H., Tang J., Huang Z., Tan W., Gan B., Yang Z., Zhao J. 2017. Proteomic Analysis Revealed the Fruiting-Body Protein Profile of Auricularia polytricha. Curr. Microbiol. 2017 748. 74:943–951.

Jiao X., Li G., Wang Y., Nie F., Cheng X., Abdullah M., Lin Y., Cai Y. 2018. Molecules | Free Full-Text | Systematic Analysis of the Pleurotus ostreatus Laccase Gene (PoLac) Family and Functional Characterization of PoLac2 Involved in the Degradation of Cotton-Straw Lignin | HTML. Mol. 2018, Vol. 23, Page 880. 23:880.

Jin L., Li G., Yu D., Huang W., Cheng C., Liao S., Wu Q., Zhang Y. 2017. Transcriptome

analysis reveals the complexity of alternative splicing regulation in the fungus Verticillium dahliae. BMC Genomics. 18:130.

JM C., EL H., C A., R B., G B., ET C., KR C., MC C., SS D., SR E., DG F., JE H., BC H., K K., CJ K., SR M., RS N., J P., MS S., M S., S W., ED W. 2012. Saccharomyces Genome Database: the genomics resource of budding yeast. Nucleic Acids Res. 40.

JM M., H T., D S., IJ G. 2010. Laetiporus sulphureus lectin and aerolysin protein family. Adv. Exp. Med. Biol. 677:67–80.

JM S., H M. 1998. Cloning and characterization of NADP-mannitol dehydrogenase cDNA from the button mushroom, Agaricus bisporus, and its expression in response to NaCl stress. Appl. Environ. Microbiol. 64:4689–4696.

Joh J.-H.H., Kim B.-G.G., Kong W.-S.S., Yoo Y.-B.B., Kim N.-K.K., Park H.-R.R., Cho B.-G.G., Lee C.-S.S. 2004. Cloning and developmental expression of a metzincin family metalloprotease cDNA from oyster mushroom Pleurotus ostreatus. FEMS Microbiol. Lett. 239:57–62.

Jones S.K., Bennett R.J. 2011. Fungal Mating Pheromones: Choreographing the Dating Game. Fungal Genet. Biol. 48:668–676.

de Jong J.F., Deelstra H.J., Wösten H.A.B., Lugones L.G. 2006. RNA-Mediated Gene Silencing in Monokaryons and Dikaryons of Schizophyllum commune. Appl. Environ. Microbiol. 72:1267.

De Jong J.F., Ohm R.A., De Bekker C., Wösten H.A.B., Lugones L.G. 2010. Inactivation of ku80 in the mushroom-forming fungus Schizophyllum commune increases the relative incidence of homologous recombination. FEMS Microbiol. Lett. 310:91–95.

Jonkers W., Rep M. 2009. Lessons from fungal F-Box proteins. Eukaryot. Cell. 8:677–695.

Jorgensen P., Nishikawa J.L., Breitkreutz B.J., Tyers M. 2002. Systematic identification of pathways that couple cell growth and division in yeast. Science (80-.). 297:395–400.

JT C., NP M., EP B., LA M. 2010. The use of biotin-binding proteins for insect control. J. Econ. Entomol. 103:497–508.

Juillot S., Cott C., Madl J., Claudinon J., van der Velden N.S.J., Künzler M., Thuenauer R., Römer W. 2016. Uptake of Marasmius oreades agglutinin disrupts integrin-dependent cell adhesion. Biochim. Biophys. Acta. 1860:392.

K A., K F., TA S., E H., R K., K G. 2006. NAD+-specific D-arabinose dehydrogenase and its contribution to erythroascorbic acid production in Saccharomyces cerevisiae. FEBS Lett. 580:6428–6434.

K O., M B., G V., M D.S., K S., P M. 2014. Fungal MACPF-like proteins and aegerolysins: bi-component pore-forming proteins? Subcell. Biochem. 80:271–291.

K S., H Y., Y O., S F., E T., S H. 1998. Production of trehalose synthase from a basidiomycete, Grifola frondosa, in Escherichia coli. Appl. Microbiol. Biotechnol. 50:193–198.

K W., K E., L A., T A., H A. 2005. Characterization of the ferrichrome A biosynthetic gene cluster in the homobasidiomycete Omphalotus olearius. FEMS Microbiol. Lett. 249:157–163.

Kabel M.A., Jurak E., Mäkelä M.R., de Vries R.P. 2017. Occurrence and function of enzymes for lignocellulose degradation in commercial Agaricus bisporus cultivation. Appl. Microbiol. Biotechnol. 2017 10111. 101:4363–4369.

Kalaras M.D., Richie J.P., Calcagnotto A., Beelman R.B., MD K., JP R., A C., RB B. 2017. Mushrooms: A rich source of the antioxidants ergothioneine and glutathione. Food Chem. 233:429–433.

Káldi K., González B.H., Brunner M., K K., BH G., M B. 2006. Transcriptional regulation of the Neurospora circadian clock gene wc-1 affects the phase of circadian output. EMBO Rep. 7:199–204.

Kamada T., Sano H., Nakazawa T., Nakahori K. 2010. Regulation of fruiting body photomorphogenesis in Coprinopsis cinerea. Fungal Genet. Biol.

Kamada T., Takemaru T., Prosser J.I., Gooday G.W. 1991. Right and left handed helicity of chitin microfibrils in stipe cells inCoprinus cinereus. Protoplasma 1991 1651. 165:64–70.

Kar B., Patel P., Ao J., Free S.J. 2019. Neurospora crassa family GH72
glucanosyltransferases function to crosslink cell wall glycoprotein N-linked
galactomannan to cell wall lichenin. Fungal Genet. Biol. 123:60–69.
Katsimpouras C., Dedes G., Thomaidis N.S., Topakas E. 2019. A novel fungal GH30
xylanase with xylobiohydrolase auxiliary activity. Biotechnol. Biofuels 2019 121. 12:1–
14.
Käufer N.F., Potashkin J. 2000. Analysis of the splicing machinery in fission yeast: a
comparison with budding yeast and mammals. Nucleic Acids Res. 28:3003.
Ke H.-M., Lee H.-H., Lin C.-Y.I., Liu Y.-C., Lu M.R., Hsieh J.-W.A., Chang C.-C., Wu P.-H.,
Lu M.J., Li J.-Y., Shang G., Lu R.J.-H., Nagy L.G., Chen P.-Y., Kao H.-W., Tsai I.J.
2020a. Mycena genomes resolve the evolution of fungal bioluminescence. Proc. Natl.
Acad. Sci. 117:31267–31277.
Ke H.-M., Lee H.-H., Lin C.-Y.I., Liu Y.-C., Lu M.R.M.J.M.R., Hsieh J.-W.A., Chang C.-C.,
Wu P.-H., Lu M.R.M.J.M.R., Li J.-Y., Shang G., Lu R.J.-H., Nagy L.G., Chen P.-Y., Kao
H.-W., Tsai I.J. 2020b. Mycena genomes resolve the evolution of fungal
bioluminescence. Proc. Natl. Acad. Sci. 117:31267–31277.
Keller N.P. 2019. Fungal secondary metabolism: regulation, function and drug discovery.
Nat. Rev. Microbiol. 17:167.
Khan A., McQuilken M., Gladfelter A.S. 2015. Septins and Generation of Asymmetries in
Fungal Cells. http://dx.doi.org/10.1146/annurev-micro-091014-104250. 69:487–503.
Kim H. Il, Lee C.S., Park Y.J. 2016. Further characterization of hydrophobin genes in
genome of Flammulina velutipes. Mycoscience. 57:320–325.
Kim J.Y., Kim D.Y., Park Y.J., Jang M.J. 2020. Transcriptome analysis of the edible
mushroom Lentinula edodes in response to blue light. PLoS One.
Kim K.-H., Kang Y.M., Im C.H., Ali A., Kim S.Y., Je H.-J., Kim M.-K., Rho H.S., Lee H.S.,
Kong W.-S., Ryu J.-S. 2014. Identification and Functional Analysis of Pheromone and
Receptor Genes in the B3 Mating Locus of Pleurotus eryngii. PLoS One. 9:e104693.
Kipreos E.T., Pagano M. 2000. The F-box protein family. Genome Biol. 2000 15. 1:1–7.
Kitamoto Y., Kobayashi A., Mori N., Ohga S. 2001. Metabolic function of glycogen
phosphorylase and trehalose phosphorylase in fruit-body formation of Flammulina
velutipes. Mycoscience. 42:143–147.
Kiyama R., Furutani Y., Kawaguchi K., Nakanishi T. 2018. Genome sequence of the
cauliflower mushroom Sparassis crispa (Hanabiratake) and its association with
beneficial usage. Sci. Reports 2018 81. 8:1–11.
Klis F.M., Boorsma A., De Groot P.W.J. 2006. Cell wall construction in Saccharomyces
cerevisiae. Yeast. 23:185–202.
Knabe N., Jung E.M., Freihorst D., Hennicke F., Horton J.S., Kothe E. 2013. A central role
for ras1 in morphogenesis of the basidiomycete Schizophyllum commune. Eukaryot.
Cell. 12:941–952.
Knoll A.H. 2011. The Multiple Origins of Complex Multicellularity. Earth Planet. Sci. 39:217–
239.
Knoll F. Untersuehungen über Längenwachstum und Geotropismus der Fruehtkörperstiele
von Coprinus stiriaeus. .
Kobayashi Y., Tateno H., Ogawa H., Yamamoto K., Hirabayashi J. 2014. Comprehensive
List of Lectins: Origins, Natures, and Carbohydrate Specificities. Methods Mol. Biol.
1200:555–577.
Koch R.A., Yoon G.M., Aryal U.K., Lail K., Amirebrahimi M., LaButti K., Lipzen A., Riley R.,
Barry K., Henrissat B., Grigoriev I. V., Herr J.R., Aime M.C. 2021. Symbiotic nitrogen
fixation in the reproductive structures of a basidiomycete fungus. Curr. Biol. 31:3905-
3914.e6.
Kohler A., Kuo A., Nagy L.G., Morin E., Barry K.W., Buscot F., Canbäck B., Choi C.,
Cichocki N., Clum A., Colpaert J., Copeland A., Costa M.D., Doré J., Floudas D., Gay
G., Girlanda M., Henrissat B., Herrmann S., Hess J., Högberg N., Johansson T., Khouja
H.-R.R., Labutti K., Lahrmann U., Levasseur A., Lindquist E.A., Lipzen A., Marmeisse
R., Martino E., Murat C., Ngan C.Y., Nehls U., Plett J.M., Pringle A., Ohm R.A., Perotto

S., Peter M., Riley R., Rineau F., Ruytinx J., Salamov A., Shah F., Sun H., Tarkka M.,
Tritt A., Veneault-Fourrey C., Zuccaro A., Tunlid A., Grigoriev I. V, Hibbett D.S., Martin
F., Martin F. 2015. Convergent losses of decay mechanisms and rapid turnover of
symbiosis genes in mycorrhizal mutualists. Nat. Genet. 47:410–415.
Kombrink A., Tayyrov A., Essig A., Stöckli M., Micheller S., Hintze J., van Heuvel Y., Dürig
N., Lin C., Kallio P.T., Aebi M., Künzler M. 2018. Induction of antibacterial proteins and
peptides in the coprophilous mushroom Coprinopsis cinerea in response to bacteria.
ISME J. 2018 133. 13:588–602.
Konada L., Aricthota S., Vadla R., Haldar D. 2018. Fission Yeast Sirtuin Hst4 Functions in
Preserving Genomic Integrity by Regulating Replisome Component Mcl1. Sci. Reports
2018 81. 8:1–17.
Kondoh O., Muto A., Kajiwara S., Takagi J., Saito Y., Shishido K. 1995. A fruiting body-
specific cDNA, mfbAc, from the mushroom Lentinus edodes encodes a high-molecular-
weight cell-adhesion protein containing an Arg-Gly-Asp motif. Gene. 154:31–37.
Kraakman L.S., Griffioen G., Zerp S., Groeneveld P., Thevelein J.M., Mager W.H., Planta
R.J. 1993. Growth-related expression of ribosomal protein genes in Saccharomyces
cerevisiae. Mol. Gen. Genet. MGG 1993 2391. 239:196–204.
Kress M.R. von Z., Harting R., Bayram Ö., Christmann M., Irmer H., Valerius O., Schinke J.,
Goldman G.H., Braus G.H. 2012. The COP9 signalosome counteracts the
accumulation of cullin SCF ubiquitin E3 RING ligases during fungal development. Mol.
Microbiol. 83:1162–1177.
Krijgsheld P., Bleichrodt R., van Veluw G.J., Wang F., Müller W.H., Dijksterhuis J., Wösten
H.A.B. 2013. Development in aspergillus. Stud. Mycol.
Krivosheina N.P. 2008. Macromycete fruit bodies as a habitat for dipterans (Insecta,
Diptera). Entomol. Rev. 2008 887. 88:778–792.
Krizsán K., Almási É., Merényi Z., Sahu N., Virágh M., Kószó T., Mondo S., Kiss B., Bálint
B., Kües U., Barry K., Cseklye J., Hegedüs B., Henrissat B., Johnson J., Lipzen A.,
Ohm R.A., Nagy I., Pangilinan J., Yan J., Xiong Y., Grigoriev I. V, Hibbett D.S., Nagy
L.G. 2019. Transcriptomic atlas of mushroom development reveals conserved genes
behind complex multicellularity in fungi. Proc. Natl. Acad. Sci. U. S. A. 116:7409–7418.
Kück U., Radchenko D., Teichert I. 2019. STRIPAK, a highly conserved signaling complex,
controls multiple eukaryotic cellular and developmental processes and is linked with
human diseases. Biol. Chem. 400:1005–1022.
Kuees U. 2015. From two to many: multiple mating types in Basidiomycetes. Fungal Biol.
Rev. 29:126–166.
Kues U. 2000. Life history and developmental processes in the basidiomycete Coprinus
cinereus. Microbiol Mol Biol Rev. 64:316–353.
Kües U. 2000. Life History and Developmental Processes in the Basidiomycete Coprinus
cinereus. Microbiol. Mol. Biol. Rev.
Kües U. 2015. From two to many: Multiple mating types in Basidiomycetes. Fungal Biol.
Rev. 29:126–166.
Kues U., Granado J.D., Hermann R., Boulianne R.P., Kertesz-Chaloupkova K., Aebi M.
1998. The A mating type and blue light regulate all known differentiation processes in
the basidiomycete Coprinus cinereus. Mol Gen Genet. 260:81–91.
Kües U., Khonsuntia W., Subba S., Dörnte B. 2018. Volatiles in Communication of
<Emphasis Type="Italic">Agaricomycetes</Emphasis>. Physiol. Genet.:149–212.
Kues U., Liu Y. 2000. Fruiting body production in basidiomycetes. Appl. Microbiol.
Biotechnol.
Kües U., Liu Y. 2000. Fruiting body production in basidiomycetes. Appl. Microbiol.
Biotechnol. 54:141–152.
Kües U., Navarro-González M. 2015. How do Agaricomycetes shape their fruiting bodies? 1.
Morphological aspects of development. Fungal Biol. Rev. 29:63–97.
Kües U., Rühl M., U K., M R. 2011. Multiple multi-copper oxidase gene families in
basidiomycetes - what for? Curr. Genomics. 12:72–94.
Kues U., Walser P.J., Klaus M.J., Aebi M. 2002. Influence of activated A and B mating-type

pathways on developmental processes in the basidiomycete Coprinus cinereus. Mol Genet Genomics. 268:262–271.

Kuhad R.C., Rosin I. V, Moore D. 1987. A possible relation between cyclic-AMP levels and glycogen mobilization in Coprinus cinereus. Trans. Br. Mycol. Soc. 88:229–236.

Kumar M.J., Jamaluddin M.S., Natarajan K., Kaur D., Datta A. 2000. The inducible N-acetylglucosamine catabolic pathway gene cluster in Candida albicans: Discrete N-acetylglucosamine-inducible factors interact at the promoter of NAG1. Proc. Natl. Acad. Sci. 97:14218–14223.

Kunzler M. 2015. Hitting the sweet spot-glycans as targets of fungal defense effector proteins. Molecules. 20:8144–8167.

Künzler M. 2018. How fungi defend themselves against microbial competitors and animal predators. PLOS Pathog. 14:e1007184.

Kuratani M., Tanaka K., Terashima K., Muraguchi H., Nakazawa T., Nakahori K., Kamada T. 2010. The dst2 gene essential for photomorphogenesis of Coprinopsis cinerea encodes a protein with a putative FAD-binding-4 domain. Fungal Genet. Biol.

KY Y., CM C., KM C., PC W., XY C., AS L., L C. 2001. Characterization of AFMP1: a novel target for serodiagnosis of aspergillosis. J. Clin. Microbiol. 39:3830–3837.

L A.-F., C C., C L., V A., V S.-S., E M., JP L. 2011. Functional analysis of the fungal/plant class chitinase family in Aspergillus fumigatus. Fungal Genet. Biol. 48:418–429.

L L., EE G., ME Z. 1999. Homolog pairing and meiotic progression in Coprinus cinereus. Chromosoma. 108:384–392.

Labourel A., Frandsen K.E.H., Zhang F., Brouilly N., Grisel S., Haon M., Ciano L., Ropartz D., Fanuel M., Martin F., Navarro D., Rosso M.-N., Tandrup T., Bissaro B., Johansen K.S., Zerva A., Walton P.H., Henrissat B., Leggio L. Lo, Berrin J.-G. 2020. A fungal family of lytic polysaccharide monooxygenase-like copper proteins. Nat. Chem. Biol. 2020 163. 16:345–350.

Lackner G., Bohnert M., Wick J., Hoffmeister D. 2013. Assembly of Melleolide Antibiotics Involves a Polyketide Synthase with Cross-Coupling Activity. Chem. Biol. 20:1101–1106.

Lakkireddy K., Khonsuntia W., Kües U. 2020. Mycoparasite Hypomyces odoratus infests Agaricus xanthodermus fruiting bodies in nature. AMB Express. 10.

Landi N., Pacifico S., Ragucci S., Iglesias R., Piccolella S., Amici A., Di Giuseppe A.M.A., Di Maro A. 2017. Purification, characterization and cytotoxicity assessment of Ageritin: The first ribotoxin from the basidiomycete mushroom Agrocybe aegerita. Biochim. Biophys. Acta - Gen. Subj. 1861:1113–1121.

Langner T., Göhre V. 2015. Fungal chitinases: function, regulation, and potential roles in plant/pathogen interactions. Curr. Genet. 2015 622. 62:243–254.

Larrondo L.F., Salas L., Melo F., Vicuña R., Cullen D. 2003. A Novel Extracellular Multicopper Oxidase from Phanerochaete chrysosporium with Ferroxidase Activity. Appl. Environ. Microbiol. 69:6257–6263.

Lau A.Y.-T., Cheng X., Cheng C.K., Nong W., Cheung M.K., Chan R.H.-F., Hui J.H.L., Kwan H.S. 2018. Discovery of microRNA-like RNAs during early fruiting body development in the model mushroom Coprinopsis cinerea. bioRxiv.

Lavrijssen B., Baars J.P., Lugones L.G., Scholtmeijer K., Sedaghat Telgerd N., Sonnenberg A.S.M., van Peer A.F. 2020. Interruption of an MSH4 homolog blocks meiosis in metaphase I and eliminates spore formation in Pleurotus ostreatus. PLoS One. 15:e0241749.

LC T., X W., A A., Z C., J T.-F., H Y., NP P., BS C., C C.-C., H E.-B., P T., SG C., HJ K., T M., X J., PP P. 2007. Ubiquitination regulates PTEN nuclear import and tumor suppression. Cell. 128:141–156.

Leatham G.F., Stahmann M.A. 1981. Studies on the Laccase of Lentinus edodes: Specificity, Localization and Association with the Development of Fruiting Bodies. Microbiology. 125:147–157.

Lebreton A., Bonnardel F., Dai Y.-C., Imberty A., Martin F.M., Lisacek F. 2021. A Comprehensive Phylogenetic and Bioinformatics Survey of Lectins in the Fungal

Kingdom. J. Fungi. 7:453.
Lee D., Jang E.H., Lee M., Kim S.W., Lee Y., Lee K.T., Bahna Y.S. 2019. Unraveling
melanin biosynthesis and signaling networks in cryptococcus neoformans. MBio. 10.
Lee K.T., So Y.S., Yang D.H., Jung K.W., Choi J., Lee D.G., Kwon H., Jang J., Wang L.L.,
Cha S., Meyers G.L., Jeong E., Jin J.H., Lee Y., Hong J., Bang S., Ji J.H., Park G.,
Byun H.J., Park S.W., Park Y.M., Adedoyin G., Kim T., Averette A.F., Choi J.S.,
Heitman J., Cheong E., Lee Y.H., Bahn Y.S. 2016. Systematic functional analysis of
kinases in the fungal pathogen Cryptococcus neoformans. Nat. Commun. 7.
Lee Y.Y., de Ulzurrun G.V.D., Schwarz E.M., Stajich J.E., Hsueh Y.P. 2021. Genome
sequence of the oyster mushroom Pleurotus ostreatus strain PC9. G3 Genes,
Genomes, Genet. 11.
Lenz C., Wick J., Braga D., García-Altares M., Lackner G., Hertweck C., Gressler M.,
Hoffmeister D. 2020. Injury-Triggered Blueing Reactions of Psilocybe "Magic"
Mushrooms. Angew. Chemie Int. Ed. 59:1450–1454.
Lesage G., Bussey H. 2006. Cell Wall Assembly in Saccharomyces cerevisiae. Microbiol.
Mol. Biol. Rev. 70:317.
Levasseur A., Drula E., Lombard V., Coutinho P.M., Henrissat B. 2013. Expansion of the
enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes.
Biotechnol. Biofuels. 6.
Levasseur A., Lomascolo A., Chabrol O., Ruiz-Dueñas F.J., Boukhris-Uzan E., Piumi F.,
Kües U., Ram A.F.J., Murat C., Haon M., Benoit I., Arfi Y., Chevret D., Drula E., Kwon
M.J., Gouret P., Lesage-Meessen L., Lombard V., Mariette J., Noirot C., Park J.,
Patyshakuliyeva A., Sigoillot J.C., Wiebenga A., Wösten H.A.B., Martin F., Coutinho
P.M., de Vries R.P., Martínez A.T., Klopp C., Pontarotti P., Henrissat B., Record E.
2014. The genome of the white-rot fungus Pycnoporus cinnabarinus: A basidiomycete
model with a versatile arsenal for lignocellulosic biomass breakdown. BMC Genomics.
15:486.
Levin D.E. 2005. Cell Wall Integrity Signaling in Saccharomyces cerevisiae. Microbiol. Mol.
Biol. Rev. 69:262.
Lewandowska M., Keyl A., Feussner I. 2020. Wax biosynthesis in response to danger: its
regulation upon abiotic and biotic stress. New Phytol. 227:698–713.
Li F., Wu X., Lam P., Bird D., Zheng H., Samuels L., Jetter R., Kunst L., F L., X W., P L., D
B., H Z., L S., R J., L K., Li F., Wu X., Lam P., Bird D., Zheng H., Samuels L., Jetter R.,
Kunst L. 2008. Identification of the Wax Ester Synthase/Acyl-Coenzyme
A:Diacylglycerol Acyltransferase WSD1 Required for Stem Wax Ester Biosynthesis in
Arabidopsis. Plant Physiol. 148:97–107.
Liao Y., Smyth G.K., Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 30:923–930.
Linder M.B., Szilvay G.R., Nakari-Setälä T., Penttilä M.E. 2005. Hydrophobins: the protein-
amphiphiles of filamentous fungi. FEMS Microbiol. Rev. 29:877–896.
Liu C., Bi J., Kang L., Zhou J., Liu X., Liu Z., Yuan S. 2021. The molecular mechanism of
stipe cell wall extension for mushroom stipe elongation growth. Fungal Biol. Rev.
35:14–26.
Liu H., Li Y., Chen D., Qi Z., Wang Q., Wang J., Jiang C., Xu J.-R. 2017. A-to-I RNA editing
is developmentally regulated and generally adaptive for sexual reproduction in
Neurospora crassa. Proc. Natl. Acad. Sci. 114:E7756--E7765.
Liu J., Shang X., Liu J., Tan Q. 2016. Changes in trehalose content, enzyme activity and
gene expression related to trehalose metabolism in Flammulina velutipes under heat
shock. Microbiology. 162:1274–1285.
Liu L., He G.J., Chen L., Zheng J., Chen Y., Shen L., Tian X., Li E., Yang E., Liao G., Wang
L. 2018. Genetic basis for coordination of meiosis and sexual structure maturation in
cryptococcus neoformans. Elife.
Liu T.B., Chen G.Q., Min H., Lin F.C. 2009. MoFLP1, encoding a novel fungal fasciclin-like
protein, is involved in conidiation and pathogenicity in Magnaporthe oryzae. J. Zhejiang
Univ. Sci. B. 10:434–444.

Liu T.B., Xue C. 2011. The ubiquitin-proteasome system and F-box proteins in pathogenic
fungi. Mycobiology. 39:243–248.
Liu X.-B. Bin, Xia E.-H.H., Li M., Cui Y.-Y.Y., Wang P.-M.M., Zhang J.-X.X., Xie B.-G.G., Xu
J.-P.P., Yan J.-J.J., Li J., Nagy L.G., Yang Z.L. 2020a. Transcriptome data reveal
conserved patterns of fruiting body development and response to heat stress in the
mushroomforming fungus Flammulina filiformis. PLoS One. 15:e0239890.
Liu X. Bin, Xia E.H., Li M., Cui Y.Y., Wang P.M., Zhang J.X., Xie B.G., Xu J.P., Yan J.J., Li
J., Nagy L.G., Yang Z.L. 2020b. Transcriptome data reveal conserved patterns of
fruiting body development and response to heat stress in the mushroomforming fungus
Flammulina filiformis. PLoS One.
Liu Y. 2005. An Essential Gene for Fruiting Body Initiation in the Basidiomycete Coprinopsis
cinerea Is Homologous to Bacterial Cyclopropane Fatty Acid Synthase Genes.
Genetics. 172:873–884.
Loftus M.G., Sánchez C., Moore D., Robson G., Trinci T. 2020. A 21st century miniguide to
sporophore morphogenesis and development in Agaricomycetes and their
biotechnological potential. .
López-Mondéjar R., Catalano V., Kubicek C.P., Seidl V. 2009. The $β$-N-
acetylglucosaminidases NAG1 and NAG2 are essential for growth of Trichoderma
atroviride on chitin. FEBS J. 276:5137–5148.
Lu B.C., Gallo N., Kües U., BC L., N G., U K. 2003. White-cap mutants and meiotic
apoptosis in the basidiomycete Coprinus cinereus. Fungal Genet. Biol. 39:82–93.
Lu M.-Y.J., Fan W.-L., Wang W.-F., Chen T., Tang Y.-C., Chu F.-H., Chang T.-T., Wang S.-
Y., Li M.-Y., Chen Y.-H., Lin Z.-S., Yang K.-J., Chen S.-M., Teng Y.-C., Lin Y.-L., Shaw
J.-F., Wang T.-F., Li W.-H. 2014. Genomic and transcriptomic analyses of the medicinal
fungus Antrodia cinnamomea for its metabolite biosynthesis and sexual development.
Proc. Natl. Acad. Sci. U. S. A. 111:E4743.
Lu Y.P., Liao J.H., Guo Z.J., Cai Z.X., Chen M.Y. 2020. Genome Survey and Transcriptome
Analysis on Mycelia and Primordia of Agaricus blazei. Biomed Res. Int. 2020.
Lugones L.G., Scholtmeijer K., Klootwijk R., Wessels J.G.H. 1999a. Introns are necessary
for mRNA accumulation in Schizophyllum commune. Mol. Microbiol. 32:681–689.
Lugones L.G., Wösten H.A.B., Birkenkamp K.U., Sjollema K.A., Zagers J., Wessels J.G.H.
1999b. Hydrophobins line air channels in fruiting bodies of Schizophyllum commune
and Agaricus bisporus. Mycol. Res. 103:635–640.
Luo H., Hallen-Adams H.E., Scott-Craig J.S., Walton J.D. 2010. Colocalization of Amanitin
and a Candidate Toxin-Processing Prolyl Oligopeptidase in Amanita Basidiocarps.
Eukaryot. Cell. 9:1891.
Luti S., Sella L., Quarantin A., Pazzagli L., Baccelli I. 2020. Twenty years of research on
cerato-platanin family proteins: clues, conclusions, and unsolved issues. Fungal Biol.
Rev. 34:13–24.
Luyten K., Albertyn J., Skibbe W.F., Prior B.A., Ramos J., Thevelein J.M., Hohmann S.
1995. Fps1, a yeast member of the MIP family of channel proteins, is a facilitator for
glycerol uptake and efflux and is inactive under osmotic stress. EMBO J. 14:1360.
M C., S P., RP A., C G., C L. 1999. The lactate-proton symport of Saccharomyces cerevisiae
is encoded by JEN1. J. Bacteriol. 181:2620–2623.
M F., J K., B F. 2003. Relationships between the ethanol utilization (alc) pathway and
unrelated catabolic pathways in Aspergillus nidulans. Eur. J. Biochem. 270:3555–3564.
M H., H K., MJ P., R U. 2015a. Fungal unspecific peroxygenases: heme-thiolate proteins
that combine peroxidase and cytochrome p450 properties. Adv. Exp. Med. Biol.
851:341–368.
M H., SM T., SJ M., RA H., KL H., CT B., E O. 2020a. Bacterial Brown Pit, a New Disease of
Edible Mushrooms Caused by Mycetocola sp. Plant Dis. 104:1445–1454.
M K., S D., F M., AG P. 2009a. RNA silencing in the model mycorrhizal fungus Laccaria
bicolor: gene knock-down of nitrate reductase results in inhibition of symbiosis with
Populus. Environ. Microbiol. 11:1878–1896.
M L., B Ž., D H., K L., R T., V M. 2021. Systematic Comparison of Cell Wall-Related Proteins

of Different Yeasts. J. fungi (Basel, Switzerland). 7:1–19.

M L., G G., V S., A K., R R., W T., J M., M O. 2010. C terminus of Nce102 determines the
structure and function of microdomains in the Saccharomyces cerevisiae plasma
membrane. Eukaryot. Cell. 9:1184–1192.

M M., AA N., M D., JP J., E G. 2009b. The fungal glutathione S-transferase system.
Evidence of new classes in the wood-degrading basidiomycete Phanerochaete
chrysosporium. Cell. Mol. Life Sci. 66:3711–3725.

M N.-L., C G.M. 1981. The cell walls of Agaricus bisporus and Agaricus campestris fruiting
body hyphae. Can. J. Microbiol. 27:779–787.

M N., N K., M S., P M., G A., K S. 2015b. Fungal aegerolysin-like proteins: distribution,
activities, and applications. Appl. Microbiol. Biotechnol. 99:601–610.

M O., P M., F B., JG G., Á M.-D.-P., L G.-O. 2017a. Characterization of a new toxin from the
entomopathogenic fungus Metarhizium anisopliae: the ribotoxin anisoplin. Biol. Chem.
398:135–142.

M O., R L.-G., JC L.-R., Á M.-D.-P., M O., M M.-L., J L., JG G., L G.-O. 2017b. Fungal
Ribotoxins: A Review of Potential Biotechnological Applications. Toxins (Basel). 9.

M S., M P., S H., J P., B S., E N., A B., M W., M P. 2002a. Swollenin, a Trichoderma reesei
protein with sequence similarity to the plant expansins, exhibits disruption activity on
cellulosic materials. Eur. J. Biochem. 269:4202–4211.

M S., P C., M B., M B., M M.-N., JP L., I M. 2020b. GPI Anchored Proteins in Aspergillus
fumigatus and Cell Wall Morphogenesis. Curr. Top. Microbiol. Immunol. 425:167–186.

M W., V T., L P., L C., D F. 2002b. Proteins as active compounds involved in insecticidal
activity of mushroom fruitbodies. J. Econ. Entomol. 95:603–607.

M Y., M K., S I., G T., M O., M S. 2008. Isolation and characterization of a gene coding for
chitin deacetylase specifically expressed during fruiting body development in the
basidiomycete Flammulina velutipes and its expression in the yeast Pichia pastoris.
FEMS Microbiol. Lett. 289:130–137.

M Y., S S., K S., G T., S I., M O., M S., Yamada M., Sakuraba S., Shibata K., Taguchi G.,
Inatomi S., Okazaki M., Shimosaka M. 2006. Isolation and analysis of genes specifically
expressed during fruiting body development in the basidiomycete Flammulina velutipes
by fluorescence differential display. FEMS Microbiol. Lett. 254:165–172.

Ma A., Shan L., Wang N., Zheng L., Chen L., Xie B. 2007. Characterization of a Pleurotus
ostreatus fruiting body-specific hydrophobin gene, Po.hyd. J. Basic Microbiol. 47:317–
324.

Ma G., Yang W., Zhao L., Pei F., Fang D., Hu Q. 2018. A critical review on the health
promoting effects of mushrooms nutraceuticals. Food Sci. Hum. Wellness. 7:125–133.

MA W., C V., RM B., A G., M A., M K., Wälti M.A., Villalba C., Buser R.M., Grünler A., Aebi
M., Künzler M. 2006. Targeted gene silencing in the model mushroom Coprinopsis
cinerea (Coprinus cinereus) by expression of homologous hairpin RNAs. Eukaryot. Cell.
5:732–744.

Ma X., Fan X., Wang G., Xu R., Yan L., Zhou Y., Gong Y., Xiao Y., Bian Y. 2021. Enhanced
Expression of Thaumatin-like Protein Gene (LeTLP1) Endows Resistance to
Trichoderma atroviride in Lentinula edodes. Life. 11:863.

Madhavan S., Krause K., Jung E.-M., Kothe E. 2014. Differential regulation of multi-copper
oxidases in Schizophyllum commune during sexual development. Mycol. Prog. 2014
134. 13:1199–1206.

Mäkelä M.R., Sietiö O.-M., Vries R.P. de, Timonen S., Hildén K. 2014. Oxalate-Metabolising
Genes of the White-Rot Fungus Dichomitus squalens Are Differentially Induced on
Wood and at High Proton Concentration. PLoS One. 9:e87959.

Mäkinen M., Kuuskeri J., Laine P., Smolander O.-P., Kovalchuk A., Zeng Z., Asiegbu F.O.,
Paulin L., Auvinen P., Lundell T. 2019. Genome description of Phlebia radiata 79 with
comparative genomics analysis on lignocellulose decomposition machinery of phlebioid
fungi. BMC Genomics 2019 201. 20:1–22.

Martin F., Aerts A., Ahrén D., Brun A., Danchin E.G.J., Duchaussoy F., Gibon J., Kohler A.,
Lindquist E., Pereda V., Salamov A., Shapiro H.J., Wuyts J., Blaudez D., Buée M.,

Brokstein P., Canbäck B., Cohen D., Courty P.E., Coutinho P.M., Delaruelle C., Detter J.C., Deveau A., DiFazio S., Duplessis S., Fraissinet-Tachet L., Lucic E., Frey-Klett P., Fourrey C., Feussner I., Gay G., Grimwood J., Hoegger P.J., Jain P., Kilaru S., Labbé J., Lin Y.C., Legué V., Le Tacon F., Marmeisse R., Melayah D., Montanini B., Muratet M., Nehls U., Niculita-Hirzel H., Secq M.P.O.-L., Peter M., Quesneville H., Rajashekar B., Reich M., Rouhier N., Schmutz J., Yin T., Chalot M., Henrissat B., Kües U., Lucas S., Van de Peer Y., Podila G.K., Polle A., Pukkila P.J., Richardson P.M., Rouzé P., Sanders I.R., Stajich J.E., Tunlid A., Tuskan G., Grigoriev I. V. 2008. The genome of Laccaria bicolor provides insights into mycorrhizal symbiosis. Nature. 452:88–92.

Martinez D., Challacombe J., Morgenstern I., Hibbett D., Schmoll M., Kubicek C.P., Ferreira P., Ruiz-Duenas F.J., Martinez A.T., Kersten P., Hammel K.E., Wymelenberg A. Vanden, Gaskell J., Lindquist E., Sabat G., BonDurant S.S., Larrondo L.F., Canessa P., Vicuna R., Yadav J., Doddapaneni H., Subramanian V., Pisabarro A.G., Lavín J.L., Oguiza J.A., Master E., Henrissat B., Coutinho P.M., Harris P., Magnuson J.K., Baker S.E., Bruno K., Kenealy W., Hoegger P.J., Kües U., Ramaiya P., Lucas S., Salamov A., Shapiro H., Tu H., Chee C.L., Misra M., Xie G., Teter S., Yaver D., James T., Mokrejs M., Pospisek M., Grigoriev I. V., Brettin T., Rokhsar D., Berka R., Cullen D. 2009a. Genome, transcriptome, and secretome analysis of wood decay fungus Postia placenta supports unique mechanisms of lignocellulose conversion. Proc. Natl. Acad. Sci. 106:1954–1959.

Martinez D., Challacombe J., Morgenstern I., Hibbett D., Schmoll M., Kubicek C.P., Ferreira P., Ruiz-Duenas F.J., Martinez A.T., Kersten P., Hammel K.E., Vanden Wymelenberg A., Gaskell J., Lindquist E., Sabat G., Splinter BonDurant S., Larrondo L.F., Canessa P., Vicuna R., Yadav J., Doddapaneni H., Subramanian V., Pisabarro A.G., Lavín J.L., Oguiza J.A., Master E., Henrissat B., Coutinho P.M., Harris P., Magnuson J.K., Baker S.E., Bruno K., Kenealy W., Hoegger P.J., Kües U., Ramaiya P., Lucas S., Salamov A., Shapiro H., Tu H., Chee C.L., Misra M., Xie G., Teter S., Yaver D., James T., Mokrejs M., Pospisek M., Grigoriev I. V, Brettin T., Rokhsar D., Berka R., Cullen D. 2009b. Genome, transcriptome, and secretome analysis of wood decay fungus *Postia placenta* supports unique mechanisms of lignocellulose conversion. Proc. Natl. Acad. Sci. 106:1954–1959.

Martinez D., Larrondo L.F., Putnam N., Sollewijn Gelpke M.D., Huang K., Chapman J., Helfenbein K.G., Ramaiya P., Detter J.C., Larimer F., Coutinho P.M., Henrissat B., Berka R., Cullen D., Rokhsar D. 2004. Genome sequence of the lignocellulose degrading fungus Phanerochaete chrysosporium strain RP78. Nat. Biotechnol. 22:695–700.

Masuda R., Iguchi N., Tukuta K., Nagoshi T., Kemuriyama K., Muraguchi H. 2016. The Coprinopsis cinerea Tup1 homologue Cag1 is required for gill formation during fruiting body morphogenesis. Biol. Open. 5:1844–1852.

Mata J., Lyne R., Burns G., Bähler J. 2002. The transcriptional program of meiosis and sporulation in fission yeast. Nat. Genet. 2002 321. 32:143–147.

Matityahu A., Hadar Y., Dosoretz C.G., Belinky P.A. 2008. Gene silencing by RNA interference in the white rot fungus Phanerochaete chrysosporium. Appl. Environ. Microbiol. 74:5359–5365.

Meister C., Thieme K.G., Thieme S., Köhler A.M., Schmitt K., Valerius O., Braus G.H. 2019. COP9 Signalosome Interaction with UspA/Usp15 Deubiquitinase Controls VeA-Mediated Fungal Multicellular Development. Biomol. 2019, Vol. 9, Page 238. 9:238.

Meng L., Chou T., Jiang S., Wang L., Zhu M., Mukhtar I., Xie B., Wang W. 2020. Characterization and expression pattern analysis of pheromone receptor-like genes in Winter Mushroom Flammulina filiformis. Arch. Microbiol. 202:2671–2678.

Meng L., Lyu X., Shi L., Wang Q., Wang L., Zhu M., Mukhtar I., Xie B., Wang W. 2021. The transcription factor FvHmg1 negatively regulates fruiting body development in Winter Mushroom Flammulina velutipes. Gene. 785:145618.

Merényi Z., Prasanna A.N., Wang Z., Kovács K., Hegedüs B., Bálint B., Papp B., Townsend J.P., Nagy L.G. 2020a. Unmatched level of molecular convergence among deeply

6043   divergent complex multicellular fungi. Mol. Biol. Evol. 37:2228–2240.
6044 Merényi Z., Prasanna A.N., Wang Z., Kovács K., Hegedüs B., Bálint B., Papp B., Townsend
6045   J.P., Nagy L.G. 2020b. Unmatched Level of Molecular Convergence among Deeply
6046   Divergent Complex Multicellular Fungi. Mol. Biol. Evol. 37:2228–2240.
6047 Merényi Z., Virágh M., Gluck-Thaler E., Slot J.C., Kiss B., Varga T., Geösel A., Hegedüs B.,
6048   Bálint B., Nagy L.G. 2021. Gene age predicts the transcriptional landscape of sexual
6049   morphogenesis in multicellular fungi. bioRxiv.:2021.06.04.447176.
6050 Metzger M.B., Hristova V.A., Weissman A.M. 2012. HECT and RING finger families of E3
6051   ubiquitin ligases at a glance. J. Cell Sci. 125:531–537.
6052 Meyerowitz E.M. 2002. Comparative genomics. Plants compared to animals: The broadest
6053   comparative study of development. Science (80-.). 295:1482–1485.
6054 Mgbeahuruike A.C., Kovalchuk A., Asiegbu F.O. 2017. Comparative genomics and
6055   evolutionary analysis of hydrophobins from three species of wood-degrading fungi.
6056   http://dx.doi.org/10.3852/13-077. 105:1471–1478.
6057 MH B., V B.-P., D M., L E. 2012. The Neurospora crassa mutant NcΔEgt-1 identifies an
6058   ergothioneine biosynthetic gene and demonstrates that ergothioneine enhances
6059   conidial survival and protects against peroxide toxicity during conidial germination.
6060   Fungal Genet. Biol. 49:160–172.
6061 Michael Green N. 1990. [5] Avidin and streptavidin. Methods Enzymol. 184:51–67.
6062 Michelot D., Melendez-Howell L.M. 2003. Amanita muscaria: chemistry, biology, toxicology,
6063   and ethnomycology. Mycol. Res. 107:131–146.
6064 Middleton R., Gao D., Thomas A., Singh B., Au A., Wong J.J.-L., Bomane A., Cosson B.,
6065   Eyras E., Rasko J.E.J., Ritchie W. 2017. IRFinder: assessing the impact of intron
6066   retention on mammalian gene expression. Genome Biol. 2017 181. 18:1–11.
6067 Min B., Kim S., Oh Y.-L., Kong W.-S., Park H., Cho H., Jang K.-Y., Kim J.-G., Choi I.-G.
6068   2018. Genomic discovery of the hypsin gene and biosynthetic pathways for terpenoids
6069   in Hypsizygus marmoreus. BMC Genomics 2018 191. 19:1–12.
6070 Min B., Yoon H., Park J., Oh Y.-L., Kong W.-S., Kim J.-G., Choi I.-G. 2020. Unusual genome
6071   expansion and transcription suppression in ectomycorrhizal Tricholoma matsutake by
6072   insertions of transposable elements. PLoS One. 15:e0227923.
6073 Miyauchi S., Kiss E., Kuo A., Drula E., Kohler A., Sánchez-García M., Morin E.,
6074   Andreopoulos B., Barry K.W., Bonito G., Buée M., Carver A., Chen C., Cichocki N.,
6075   Clum A., Culley D., Crous P.W., Fauchery L., Girlanda M., Hayes R.D., Kéri Z., LaButti
6076   K., Lipzen A., Lombard V., Magnuson J., Maillard F., Murat C., Nolan M., Ohm R.A.,
6077   Pangilinan J., Pereira M. de F., Perotto S., Peter M., Pfister S., Riley R., Sitrit Y.,
6078   Stielow J.B., Szöllősi G., Žifčáková L., Štursová M., Spatafora J.W., Tedersoo L.,
6079   Vaario L.M., Yamada A., Yan M., Wang P., Xu J., Bruns T., Baldrian P., Vilgalys R.,
6080   Dunand C., Henrissat B., Grigoriev I. V, Hibbett D., Nagy L.G., Martin F.M. 2020. Large-
6081   scale genome sequencing of mycorrhizal fungi provides insights into the early evolution
6082   of symbiotic traits. Nat. Commun.
6083 Miyazaki Y., Jojima T., Ono T., Yamazaki T., Shishido K. 2004. A cDNA homologue of
6084   Schizosaccharomyces pombe cdc5 + from the mushroom Lentinula edodes:
6085   Characterization of the cDNA and its expressed product. Biochim. Biophys. Acta - Gene
6086   Struct. Expr. 1680:93–102.
6087 Mol P.C., Vermeulen C.A., Wessels J.G.H. 1990. Diffuse extension of hyphae in stipes of
6088   Agaricus bisporus may be based on a unique wall structure. Mycol. Res. 94:480–488.
6089 Mol P.C., Wessels J.G.H. 1990. Differences in wall structure between substrate hyphae and
6090   hyphae of fruit-body stipes in Agaricus bisporus. Mycol. Res. 94:472–479.
6091 Mollapour M., Piper P.W. 2007. Hog1 Mitogen-Activated Protein Kinase Phosphorylation
6092   Targets the Yeast Fps1 Aquaglyceroporin for Endocytosis, Thereby Rendering Cells
6093   Resistant to Acetic Acid. Mol. Cell. Biol. 27:6446.
6094 Moore D. 2005. Principles of Mushroom Developmental Biology. .
6095 Moore D. (David). 2013. Coprinopsis□: an autobiography. :208.
6096 MOORE D., ELHITI M.M.Y., BUTLER R.D. 1979. MORPHOGENENSIS OF THE
6097   CARPOPHORE OF COPRINUS CINEREUS. New Phytol. 83:695–722.

Moran Y., Fredman D., Szczesny P., Grynberg M., Technau U. 2012. Recurrent horizontal
transfer of bacterial toxin genes to eukaryotes. Mol. Biol. Evol. 29:2223–2230.
Morin-Crini N., Lichtfouse E., Torri G., Crini G. 2019. Applications of chitosan in food,
pharmaceuticals, medicine, cosmetics, agriculture, textiles, pulp and paper,
biotechnology, and environmental chemistry. Environ. Chem. Lett. 2019 174. 17:1667–
1692.
Morin E., Kohler A., Baker A.R., Foulongne-Oriol M., Lombard V., Nagy L.G., Ohm R.A.,
Patyshakuliyeva A., Brun A., Aerts A.L., Bailey A.M., Billette C., Coutinho P.M., Deakin
G., Doddapaneni H., Floudas D., Grimwood J., Hildén K., Kües U., Labutti K.M.,
Lapidus A., Lindquist E.A., Lucas S.M., Murat C., Riley R.W., Salamov A.A., Schmutz
J., Subramanian V., Wösten H.A.B., Xu J., Eastwood D.C., Foster G.D., Sonnenberg
A.S.M., Cullen D., de Vries R.P., Lundell T., Hibbett D.S., Henrissat B., Burton K.S.,
Kerrigan R.W., Challen M.P., Grigoriev I. V., Martin F. 2012. Genome sequence of the
button mushroom Agaricus bisporus reveals mechanisms governing adaptation to a
humic-rich ecological niche. Proc. Natl. Acad. Sci. U. S. A. 109:17501–17506.
Morton N., Hammond J.B.W., Dickerson A.G. 1985. The Agaricus bisporus mannitol
pathway during sporophore growth. Trans. Br. Mycol. Soc. 85:671–675.
Munir E., Yoon J.J., Tokimatsu T., Hattori T., Shimada M. 2001. A physiological role for
oxalic acid biosynthesis in the wood-rotting basidiomycete Fomitopsis palustris. Proc.
Natl. Acad. Sci. 98:11126–11130.
Muraguchi H., Abe K., Nakagawa M., Nakamura K., Yanagi S.O., H M., K A., M N., K N., SO
Y. 2008a. Identification and characterisation of structural maintenance of chromosome
1 (smc1) mutants of Coprinopsis cinerea. Mol. Genet. Genomics. 280:223–232.
Muraguchi H., Fujita T., Kishibe Y., Konno K., Ueda N., Nakahori K., Yanagi S.O., Kamada
T., H M., T F., Y K., K K., N U., K N., SO Y., T K., Muraguchi H., Fujita T., Kishibe Y.,
Konno K., Ueda N., Nakahori K., Yanagi S.O., Kamada T. 2008b. The exp1 gene
essential for pileus expansion and autolysis of the inky cap mushroom Coprinopsis
cinerea (Coprinus cinereus) encodes an HMG protein. Fungal Genet. Biol. 45:890–896.
Muraguchi H., Kamada T. 1998. The ich1 gene of the mushroom Coprinus cinereus is
essential for pileus formation in fruiting. Development. 125:3133–3141.
Muraguchi H., Kamada T. 2000. A mutation in the eln2 gene encoding a cytochrome P450 of
Coprinus cinereus affects mushroom morphogenesis. Fungal Genet. Biol. 29:49–59.
Muraguchi H., Umezawa K., Niikura M., Yoshida M., Kozaki T., Ishii K., Sakai K., Shimizu
M., Nakahori K., Sakamoto Y., Choi C., Ngan C.Y., Lindquist E., Lipzen A., Tritt A.,
Haridas S., Barry K., Grigoriev I. V, Pukkila P.J. 2015. Strand-specific RNA-seq
analyses of fruiting body development in Coprinopsis cinerea. PLoS One. 10.
Murata Y., Fujii M., Zolan M.E., Kamada T. 1998. Molecular analysis of pcc1, a gene that
leads to A-regulated sexual morphogenesis in Coprinus cinereus. Genetics. 149:1753–
1761.
N L., SC K., I F., RH L., CF R., TT C.-D., BA S., O K., DA T., SM E., I V., JA T., T H., K O.,
EM H., RK T., JC W., M T., HR S., MA D. 2015. Conformational changes during pore
formation by the perforin-related protein pleurotolysin. PLoS Biol. 13.
N T., S Y., A O., H H., Takeshita N., Yamashita S., Ohta A., Horiuchi H. 2006. Aspergillus
nidulans class V and VI chitin synthases CsmA and CsmB, each with a myosin motor-
like domain, perform compensatory functions that are essential for hyphal tip growth.
Mol. Microbiol. 59:1380–1394.
Nagy L.G. 2017. Evolution: Complex Multicellular Life with 5,500 Genes. Curr. Biol.
27:R609–R612.
Nagy L.G., Kovács G.M., Krizsán K. 2018. Complex multicellularity in fungi: evolutionary
convergence, single origin, or both? Biol. Rev.
Nagy L.G., Ohm R.A., Kovács G.M., Floudas D., Riley R., Gácser A., Sipiczki M., Davis J.M.,
Doty S.L., de Hoog G.S., Lang B.F., Spatafora J.W., Martin F.M., Grigoriev I. V, Hibbett
D.S. 2014. Latent homology and convergent regulatory evolution underlies the repeated
emergence of yeasts. 5:4471.
Nakade K., Watanabe H., Sakamoto Y., Sato T. 2011. Gene silencing of the Lentinula

edodes lcc1 gene by expression of a homologous inverted repeat sequence. Microbiol. Res. 166:484–493.

Nakazawa T., Ando Y., Hata T., Nakahori K., T N., Y A., T H., K N. 2016. A mutation in the Cc.arp9 gene encoding a putative actin-related protein causes defects in fruiting initiation and asexual development in the agaricomycete Coprinopsis cinerea. Curr. Genet. 62:565–574.

Nakazawa T., Ando Y., Kitaaki K., Nakahori K., Kamada T. 2011. Efficient gene targeting in δCc.ku70 or δCc.lig4 mutants of the agaricomycete coprinopsis cinerea. Fungal Genet. Biol. 48:939–946.

Nakazawa T., Kaneko S., Miyazaki Y., Jojima T., Yamazaki T., Katsukawa S., Shishido K. 2008. Basidiomycete Lentinula edodes CDC5 and a novel interacting protein CIPB bind to a newly isolated target gene in an unusual manner. Fungal Genet. Biol. 45:818–828.

Nakazawa T., Kaneko S., Murata H., Kamada T., Shishido K. 2009. The homologue of Lentinula edodes ctg1, a target for CDC5 and its interacting partner CIPB, from Coprinopsis cinerea is involved in fruitingbody morphogenesis of C. cinerea. Mycoscience. 50:331–342.

Nakazawa T., Morimoto R., Wu H., Kodera R., Sakamoto M., Honda Y. 2019. Dominant effects of gat1 mutations on the ligninolytic activity of the white-rot fungus Pleurotus ostreatus. Fungal Biol. 123:209–217.

Nakazawa T., Tatsuta Y., Fujita T., Nakahori K., Kamada T. 2010. Mutations in the Cc.rmt1 gene encoding a putative protein arginine methyltransferase alter developmental programs in the basidiomycete Coprinopsis cinerea. Curr. Genet. 56:361–367.

Namekawa S.H., Iwabata K., Sugawara H., Hamada F.N., Koshiyama A., Chiku H., Kamada T., Sakaguchi K. 2005. Knockdown of LIM15/DMC1 in the mushroom Coprinus cinereus by double-stranded RNA-mediated gene silencing. Microbiology. 151:3669–3678.

Nara T., Hamada F., Namekawa S., Sakaguchi K. 2001. Strand Exchange Reaction in Vitro and DNA-Dependent ATPase Activity of Recombinant LIM15/DMC1 and RAD51 Proteins from Coprinus cinereus. Biochem. Biophys. Res. Commun. 285:92–97.

Nehls U., Dietz S. 2014. Fungal aquaporins: cellular functions and ecophysiological perspectives. Appl. Microbiol. Biotechnol. 2014 9821. 98:8835–8851.

Nguyen D.V., Roret T., Fernandez-Gonzalez A., Kohler A., Morel-Rouhier M., Gelhaye E., Sormani R. 2020. Target of Rapamycin pathway in the white-rot fungus Phanerochaete chrysosporium. PLoS One. 15.

Nguyen T.A., Cissé O.H., Yun Wong J., Zheng P., Hewitt D., Nowrousian M., Stajich J.E., Jedd G. 2017. Innovation and constraint leading to complex multicellularity in the Ascomycota. 8:14444.

Nickerson K.W., Atkin A.L., Hornby J.M., KW N., AL A., JM H. 2006. Quorum sensing in dimorphic fungi: farnesol and beyond. Appl. Environ. Microbiol. 72:3805–3813.

Niu X., Liu C.-C., Xiong Y.-J., Yang M.-M., Ma F., Liu Z.-H., Yuan S. 2016. The Modes of Action of ChiIII, a Chitinase from Mushroom Coprinopsis cinerea, Shift with Changes in the Length of GlcNAc Oligomers. J. Agric. Food Chem. 64:6958–6968.

Niu X., Liu Z., Zhou Y., Wang J., Zhang W., Yuan S. 2015. Stipe cell wall architecture varies with the stipe elongation of the mushroom Coprinopsis cinerea. Fungal Biol. 119:946–956.

Noble R., Dobrovin-Pennington A., Hobbs P.J., Pederby J., Rodger J.P.A. 2009. Volatile C8 compounds and pseudomonads influence primordium formation of Agaricus bisporus. Mycologia. 101:583–591.

Nofiani R., de Mattos-Shipley K., Lebe K.E., Han L.-C., Iqbal Z., Bailey A.M., Willis C.L., Simpson T.J., Cox R.J. 2018. Strobilurin biosynthesis in Basidiomycete fungi. Nat. Commun. 2018 91. 9:1–11.

Novak M., Krpan T., Panevska A., Shewell L.K., Day C.J., Jennings M.P., Guella G., Sepčić K. 2020. Binding specificity of ostreolysin A6 towards Sf9 insect cell lipids. Biochim. Biophys. Acta - Biomembr. 1862:183307.

Nowrousian M., Cebula P. 2005. The gene for a lectin-like protein is transcriptionally

activated during sexual development, but is not essential for fruiting body formation in
the filamentous fungus Sordaria macrospora. BMC Microbiol. 5:64.
NY S., JM L., GJ V., HH O., JD P., ME Z. 1997. Isolation and characterization of rad51
orthologs from Coprinus cinereus and Lycopersicon esculentum, and phylogenetic
analysis of eukaryotic recA homologs. Curr. Genet. 31:144–157.
O'Connor E., Doyle S., Amini A., Grogan H., Fitzpatrick D.A. 2021. Transmission of
mushroom virus X and the impact of virus infection on the transcriptomes and
proteomes of different strains of Agaricus bisporus. Fungal Biol. 125:704–717.
O'Connor E., McGowan J., McCarthy C.G.P., Amini A., Grogan H., Fitzpatrick D.A. 2019.
Whole genome sequence of the commercially relevant mushroom strain Agaricus
bisporus var. bisporus ARP23. G3 Genes, Genomes, Genet. 9:3057–3066.
Obermaier S., Müller M. 2020. Ibotenic Acid Biosynthesis in the Fly Agaric Is Initiated by
Glutamate Hydroxylation. Angew. Chemie Int. Ed. 59:12432–12435.
Ohga S., Cho N.-S., Thurston C.F., Wood D.A. 2000. Transcriptional regulation of laccase
and cellulase in relation to fruit body formation in the mycelium of Lentinula edodes on
a sawdust-based substrate. Mycoscience. 41:149–153.
Ohm R.A., Aerts D., Wösten H.A.B., Lugones L.G. 2013. The blue light receptor complex
WC-1/2 of *Schizophyllum commune*is involved in mushroom formation and protection
against phototoxicity. Environ. Microbiol. 15:943–955.
Ohm R.A., de Jong J.F., de Bekker C., Wösten H.A.B., Lugones L.G. 2011. Transcription
factor genes of Schizophyllum commune involved in regulation of mushroom formation.
Mol. Microbiol. 81:1433–1445.
Ohm R.A., Jong J.F. de, Lugones L.G., Aerts A., Kothe E., Stajich J.E., Vries R.P. de,
Record E., Levasseur A., Baker S.E., Bartholomew K.A., Coutinho P.M., Erdmann S.,
Fowler T.J., Gathman A.C., Lombard V., Henrissat B., Knabe N., Kües U., Lilly W.W.,
Lindquist E., Lucas S., Magnuson J.K., Piumi F., Raudaskoski M., Salamov A.,
Schmutz J., Schwarze F.W.M.R., vanKuyk P.A., Horton J.S., Grigoriev I. V, Wösten
H.A.B., de Jong J.F., Lugones L.G., Aerts A., Kothe E., Stajich J.E., de Vries R.P.,
Record E., Levasseur A., Baker S.E., Bartholomew K.A., Coutinho P.M., Erdmann S.,
Fowler T.J., Gathman A.C., Lombard V., Henrissat B., Knabe N., Kües U., Lilly W.W.,
Lindquist E., Lucas S., Magnuson J.K., Piumi F., Raudaskoski M., Salamov A.,
Schmutz J., Schwarze F.W.M.R., vanKuyk P.A., Horton J.S., Grigoriev I. V, Wösten
H.A.B. 2010. Genome sequence of the model mushroom Schizophyllum commune.
Nat. Biotechnol. 2010 289. 28:957–963.
Ohm R.A., Riley R., Salamov A., Min B., Choi I.G., Grigoriev I. V. 2014. Genomics of wood-
degrading fungi. Fungal Genet. Biol. 72:82–90.
Ojeda-López M., Chen W., Eagle C.E., Gutiérrez G., Jia W.L., Swilaiman S.S., Huang Z.,
Park H.S., Yu J.H., Cánovas D., Dyer P.S. 2018. Evolution of asexual and sexual
reproduction in the aspergilli. Stud. Mycol.
de Oliveira E.S., Junges Â., Sbaraini N., Andreis F.C., Thompson C.E., Staats C.C., Schrank
A. 2018. Molecular evolution and transcriptional profile of GH3 and GH20 β-N-
acetylglucosaminidases in the entomopathogenic fungus *Metarhizium anisopliae*.
Genet. Mol. Biol. 41:843–857.
Orban A., Weber A., Herzog R., Hennicke F., Rühl M. 2021. Transcriptome of different
fruiting stages in the cultivated mushroom Cyclocybe aegerita suggests a complex
regulation of fruiting and reveals enzymes putatively involved in fungal oxylipin
biosynthesis. BMC Genomics 2021 221. 22:1–23.
Orlean P. 2012. Architecture and Biosynthesis of the Saccharomyces cerevisiae Cell Wall.
Genetics. 192:775.
P B. 2006. Fungal laccases - occurrence and properties. FEMS Microbiol. Rev. 30:215–242.
P S. 2008. Chemical defence strategies of higher fungi. Chemistry. 14:9100–9110.
PA H., AC V., JF A. 1978. The insensitivity of mushroom nuclear RNA polymerase activity to
inhibition by amatoxins. Arch. Microbiol. 118:317–319.
Palmer G.E., Horton J.S. 2006. Mushrooms by magic: making connections between signal
transduction and fruiting body development in the basidiomycete fungus Schizophyllum

commune. FEMS Microbiol. Lett. 262:1–8.
Palmieri F., Estoppey A., House G.L., Lohberger A., Bindschedler S., Chain P.S.G., Junier
P. 2019. Oxalic acid, a molecule at the crossroads of bacterial-fungal interactions. Adv.
Appl. Microbiol.:1–29.
Panevska A., Skočaj M., Modic Š., Razinger J., Sepčić K. 2020. Aegerolysins from the
fungal genus Pleurotus – Bioinsecticidal proteins with multiple potential applications. J.
Invertebr. Pathol.:107474.
Park M., Kim M., Kim S., Ha B., Ro H.-S.S. 2015. Differential expression of laccase genes in
Pleurotus ostreatus and biochemical characterization of laccase isozymes produced in
Pichia pastoris. Mycobiology. 43:280.
Park S.G., il Yoo S., Ryu D.S., Lee H., Ahn Y.J., Ryu H., Ko J., Hong C.P. 2017. Long-read
transcriptome data for improved gene prediction in Lentinula edodes. Data Br. 15:454–
458.
Park Y.-J., Baek J.H., Lee S., Kim C., Rhee H., Kim H., Seo J.-S., Park H.-R., Yoon D.-E.,
Nam J.-Y., Kim H.-I., Kim J.-G., Yoon H., Kang H.-W., Cho J.-Y., Song E.-S., Sung G.-
H., Yoo Y.-B., Lee C.-S., Lee B.-M., Kong W.-S. 2014. Whole Genome and Global
Gene Expression Analyses of the Model Mushroom Flammulina velutipes Reveal a
High Capacity for Lignocellulose Degradation. PLoS One. 9:e93560.
Park Y.-J., Lee C.-S., Kong W.-S. 2019. Genomic Insights into the Fungal Lignocellulolytic
Machinery of Flammulina rossica. Microorganisms. 7.
Patel P.K., Free S.J. 2019. The Genetics and Biochemistry of Cell Wall Structure and
Synthesis in Neurospora crassa, a Model Filamentous Fungus. Front. Microbiol.
0:2294.
Patyshakuliyeva A., Jurak E., Kohler A., Baker A., Battaglia E., De Bruijn W., Burton K.S.,
Challen M.P., Coutinho P.M., Eastwood D.C., Gruben B.S., Mäkelä M.R., Martin F.,
Nadal M., van den Brink J., Wiebenga A., Zhou M., Henrissat B., Kabel M., Gruppen H.,
De Vries R.P. 2013. Carbohydrate utilization and metabolism is highly differentiated in
Agaricus bisporus. BMC Genomics. 14.
Van Peer A.F., Wang F., Van Driel K.G.A., De Jong J.F., Van Donselaar E.G., Müller W.H.,
Boekhout T., Lugones L.G., Wösten H.A.B., Peer A.F. Van, Wang F., Driel K.G.A. Van,
Jong J.F. De, Donselaar E.G. Van, Müller W.H., Boekhout T., Lugones L.G., Wösten
H.A.B. 2010. The septal pore cap is an organelle that functions in vegetative growth
and mushroom formation of the wood-rot fungus Schizophyllum commune. Environ.
Microbiol. 12:833–844.
Peisker K., Chiabudini M., Rospert S. 2010. The ribosome-bound Hsp70 homolog Ssb of
Saccharomyces cerevisiae. Biochim. Biophys. Acta - Mol. Cell Res. 1803:662–672.
Pelkmans J.F. 2016. Environmental signaling and regulation of mushroom formation. .
Pelkmans J.F., Patil M.B., Gehrmann T., Reinders M.J.T., Wösten H.A.B., Lugones L.G.
2017a. Transcription factors of schizophyllum commune involved in mushroom
formation and modulation of vegetative growth. Sci. Rep. 7:1–11.
Pelkmans J.F., Patil M.B., Gehrmann T., Reinders M.J.T.T., Wösten H.A.B.B., Lugones L.G.
2017b. Transcription factors of schizophyllum commune involved in mushroom
formation and modulation of vegetative growth. Sci. Rep. 7:1–11.
Pelkmans J.F., Vos A.M., Scholtmeijer K., Hendrix E., Baars J.J.P., Gehrmann T., Reinders
M.J.T., Lugones L.G., Wösten H.A.B. 2016. The transcriptional regulator c2h2
accelerates mushroom formation in Agaricus bisporus. Appl. Microbiol. Biotechnol.
100:7151–7159.
Pellegrin C., Morin E., Martin F.M., Veneault-Fourrey C. 2015. Comparative Analysis of
Secretomes from Ectomycorrhizal Fungi with an Emphasis on Small-Secreted Proteins.
Front. Microbiol. 6:1278.
Peñalva M.A., Tilburn J., Bignell E., Arst H.N. 2008. Ambient pH gene regulation in fungi:
making connections. Trends Microbiol. 16:291–300.
Peraro M.D., van der Goot F.G. 2015. Pore-forming toxins: ancient, but never really out of
fashion. Nat. Rev. Microbiol. 2015 142. 14:77–92.
Pereira M. de F., Campos A.N. da R., Anastacio T.C., Morin E., Brommonschenkel S.H.,

Martin F., Kohler A., Costa M.D. 2017. The transcriptional landscape of
basidiosporogenesis in mature Pisolithus microcarpus basidiocarp. BMC Genomics
2017 181. 18:1–16.
Plagemann I., Zelena K., Arendt P., Ringel P.D., Krings U., Berger R.G. 2013. LOXPsa1, the
first recombinant lipoxygenase from a basidiomycete fungus. J. Mol. Catal. B Enzym.
87:99–104.
Plaza D.F., Lin C.-W.W., van der Velden N.S.J., Aebi M., Kunzler M., Künzler M., Kunzler
M., Künzler M. 2014a. Comparative transcriptomics of the model mushroom
Coprinopsis cinerea reveals tissue-specific armories and a conserved circuitry for
sexual development. BMC Genomics. 15:492.
Plaza D.F., Lin C.W., van der Velden N.S., Aebi M., Kunzler M. 2014b. Comparative
transcriptomics of the model mushroom Coprinopsis cinerea reveals tissue-specific
armories and a conserved circuitry for sexual development. BMC Genomics. 15:492.
Pluskal T., Ueno M., Yanagida M. 2014. Genetic and Metabolomic Dissection of the
Ergothioneine and Selenoneine Biosynthetic Pathway in the Fission Yeast, S. pombe,
and Construction of an Overproduction System. PLoS One. 9:97774.
Pócsi I., Prade R.A., Penninckx M.J. 2004. Glutathione, Altruistic Metabolite in Fungi. Adv.
Microb. Physiol. 49:1–76.
Poeta M. Del, Nimrichter L., Rodrigues M.L., Luberto C. 2014. Synthesis and Biological
Properties of Fungal Glucosylceramide. PLoS Pathog. 10.
Pöggeler S., Nowrousian M., Kück U. 2006a. 16 Fruiting-Body Development in
Ascomycetes. .
Pöggeler S., Nowrousian M., Kück U. 2006b. Fruiting-Body Development in Ascomycetes.
In: Kües U., Fischer R., editors. Growth, Differentiation and Sexuality. Berlin,
Heidelberg: Springer Berlin Heidelberg. p. 325–355.
Polonio Á., Fernández-Ortuño D., de Vicente A., Pérez-García A. 2021. A haustorial-
expressed lytic polysaccharide monooxygenase from the cucurbit powdery mildew
pathogen Podosphaera xanthii contributes to the suppression of chitin-triggered
immunity. Mol. Plant Pathol. 22:580–601.
Prasanna A.N., Gerber D., Kijpornyongpan T., Aime M.C., Doyle V.P., Nagy L.G. 2020.
Model choice, missing data, and taxon sampling impact phylogenomic inference of
deep basidiomycota relationships. Syst. Biol.
Primig M., Williams R.M., Winzeler E.A., Tevzadze G.G., Conway A.R., Hwang S.Y., Davis
R.W., Esposito R.E. 2000. The core meiotic transcriptome in budding yeasts. Nat.
Genet. 2000 264. 26:415–423.
Pulman J.A., Childs K.L., Sgambelluri R.M., Walton J.D. 2016. Expansion and diversification
of the MSDIN family of cyclic peptide genes in the poisonous agarics Amanita
phalloides and A. bisporigera. BMC Genomics 2016 171. 17:1–14.
Q H., X H., I M., L G., R H., L F., J Y., Y T., B C., B X. 2018. Identification and Expression
Patterns of fvexpl1, an Expansin-Like Protein-Encoding Gene, Suggest an Auxiliary
Role in the Stipe Morphogenesis of Flammulina velutipes. J. Microbiol. Biotechnol.
28:622–629.
Q Z., L X., S Y., Q Z., X W., L W., Z H., Y Y. 2020. NGT1 Is Essential for N-
Acetylglucosamine-Mediated Filamentous Growth Inhibition and HXK1 Functions as a
Positive Regulator of Filamentous Growth in Candida tropicalis. Int. J. Mol. Sci. 21:1–
18.
Qi Y., Chen H., Zhang M., Wen Q., Qiu L., Shen J. 2019. Identification and expression
analysis of Pofst3 suggests a role during Pleurotus ostreatus primordia formation.
Fungal Biol. 123:200–208.
Qi Y., Sun X., Ma L., Wen Q., Qiu L., Shen J. 2020. Identification of two Pleurotus ostreatus
blue light receptor genes (PoWC-1 and PoWC-2) and in vivo confirmation of complex
PoWC-12 formation through yeast two hybrid system. Fungal Biol. 124:8–14.
Quiroz-Castañeda R.E., Martínez-Anaya C., Cuervo-Soto L.I., Segovia L., Folch-Mallol J.L.
2011. Loosenin, a novel protein with cellulose-disrupting activity from Bjerkandera
adusta. Microb. Cell Factories 2011 101. 10:1–9.

R D., J F., S D., M G., D H. 2020. S-Adenosyl-l-Methionine Salvage Impacts Psilocybin Formation in "Magic" Mushrooms. Chembiochem. 21:1364–1371.

R G., S W., S A., J Y., DA W. 2003. BTB/POZ domain proteins are putative substrate adaptors for cullin 3 ubiquitin ligases. Mol. Cell. 12:783–790.

R U., J N., K S., J S., M H. 2004. Novel haloperoxidase from the agaric basidiomycete Agrocybe aegerita oxidizes aryl alcohols and aldehydes. Appl. Environ. Microbiol. 70:4575–4581.

Ragucci S., Landi N., Russo R., Valletta M., Pedone P.V., Chambery A., Maro A. Di. 2021. Ageritin from Pioppino Mushroom: The Prototype of Ribotoxin-Like Proteins, a Novel Family of Specific Ribonucleases in Edible Mushrooms. Toxins 2021, Vol. 13, Page 263. 13:263.

Rahmad N., Al-Obaidi J.R., Rashid N.M., Zean N., Yusoff M.H.Y., Shaharuddin N., Jamil N.A., Saleh N. 2014. Comparative proteomic analysis of different developmental stages of the edible mushroom Termitomyces heimii. Biol. Res. 47:30.

Rahman F., Hassan M., Hanano A., Fitzpatrick D.A., McCarthy C.G.P., Murphy D.J. 2018. Evolutionary, structural and functional analysis of the caleosin/peroxygenase gene family in the Fungi. BMC Genomics 2018 191. 19:1–24.

Raudaskoski M., Kothe E. 2010. Basidiomycete mating type genes and pheromone signaling. Eukaryot Cell. 9:847–859.

Rayle D.L., Cleland R.E. 1992. The Acid Growth Theory of auxin-induced cell elongation is alive and well. Plant Physiol. 99:1271–1274.

Reboul C.F., Whisstock J.C., Dunstone M.A. 2016. Giant MACPF/CDC pore forming toxins: A class of their own. Biochim. Biophys. Acta - Biomembr. 1858:475–486.

Reinhold R., Krüger V., Meinecke M., Schulz C., Schmidt B., Grunau S.D., Guiard B., Wiedemann N., Laan M. van der, Wagner R., Rehling P., Dudek J. 2012. The Channel-Forming Sym1 Protein Is Transported by the TIM23 Complex in a Presequence-Independent Manner. Mol. Cell. Biol. 32:5009.

Rennemeier C., Schwab M., Lermann U., Albert C., Kämmerer U., Frambach T., Morschhäuser J., Dietl J., Staib P. 2011. Seminal plasma protects human spermatozoa and pathogenic yeasts from capture by dendritic cells. Hum. Reprod. 26:987–999.

Reschka E.J., Nordzieke S., Valerius O., Braus G.H., Pöggeler S., J. Reschka E., Nordzieke S., Valerius O., Braus G.H., Pöggeler S. 2018. A novel STRIPAK complex component mediates hyphal fusion and fruiting-body development in filamentous fungi. Mol. Microbiol. 110:513–532.

Revuelta M. V, van Kan J.A.L., Kay J., ten Have A. 2014. Extensive Expansion of A1 Family Aspartic Proteinases in Fungi Revealed by Evolutionary Analyses of 107 Complete Eukaryotic Proteomes. Genome Biol. Evol. 6:1480–1494.

Riley R., Salamov A.A., Brown D.W., Nagy L.G., Floudas D., Held B.W., Levasseur A., Lombard V., Morin E., Otillar R., Lindquist E.A., Sun H., LaButti K.M., Schmutz J., Jabbour D., Luo H., Baker S.E., Pisabarro A.G., Walton J.D., Blanchette R.A., Henrissat B., Martin F., Cullen D., Hibbett D.S., Grigoriev I. V. 2014. Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. Proc. Natl. Acad. Sci. U. S. A. 111:9923–9928.

Riquelme M., Aguirre J., Bartnicki-García S., Braus G.H., Feldbrügge M., Fleig U., Hansberg W., Herrera-Estrella A., Kämper J., Kück U., Mouriño-Pérez R.R., Takeshita N., Fischer R. 2018. Fungal Morphogenesis, from the Polarized Growth of Hyphae to Complex Reproduction and Infection Structures. Microbiol. Mol. Biol. Rev. 82:e00068-17.

Riquelme M., Challen M.P., Casselton L.A., Brown A.J. 2005. The origin of multiple B mating specificities in Coprinus cinereus. Genetics. 170:1105–1119.

Robinson M.D., McCarthy D.J., Smyth G.K. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 26:139–140.

Rodenburg S.Y.A., Terhem R.B., Veloso J., Stassen J.H.M., Van Kan J.A.L. 2018. Functional analysis of mating type genes and transcriptome analysis during fruiting body development of botrytis cinerea. MBio. 9.

van Roermund C.W.T., Waterham H.R., Ijlst L., Wanders R.J.A. 2003. Fatty acid metabolism
in Saccharomyces cerevisiae. Cell. Mol. Life Sci. C. 2003 609. 60:1838–1851.
Román E., Arana D.M., Nombela C., Alonso-Monge R., Pla J. 2007. MAP kinase pathways
as regulators of fungal virulence. Trends Microbiol. 15:181–190.
Romano J.D., Michaelis S. 2001. Topological and mutational analysis of Saccharomyces
cerevisiae Ste14p, founding member of the isoprenylcysteine carboxyl
methyltransferase family. Mol. Biol. Cell. 12:1957–1971.
Roper M., Seminara A. 2019. Mycofluidics: The Fluid Mechanics of Fungal Adaptation.
https://doi.org/10.1146/annurev-fluid-122316-045308. 51:511–538.
Royse D.J., Baars J., Tan Q. 2017. Current Overview of Mushroom Production in the World.
Edible and Medicinal Mushrooms. .
Ruess L., Lussenhop J. 2005. Trophic Interactions of Fungi and Animals. :581–598.
S B.-M., K S., R S., M C., JM M., M H., M S., A V., M K. 2017. Dimerization of the fungal
defense lectin CCL2 is essential for its toxicity against nematodes. Glycobiology.
27:486–500.
S B.-M., S S., M A., M K. 2012a. Biotin-binding proteins in the defense of mushrooms
against predators and parasites. Appl. Environ. Microbiol. 78:8485–8487.
S B., J P., I V., K R., F P., T T., P M., A S., K S. 2007a. Ostreolysin enhances fruiting
initiation in the oyster mushroom (Pleurotus ostreatus). Mycol. Res. 111:1431–1436.
S C., FE N., A V., S T.-M., RM R.-V. 2012b. Two distinct RNA-dependent RNA polymerases
are required for initiation and amplification of RNA silencing in the basal fungus Mucor
circinelloides. Mol. Microbiol. 83:379–394.
S C., J D., M E., J M., D B., PO B., I H. 1998a. The transcriptional program of sporulation in
budding yeast. Science. 282:699–705.
S H., J S., H M., U B., J S. 2001. The crystallographic structure of the mannitol 2-
dehydrogenase NADP+ binary complex from Agaricus bisporus. J. Biol. Chem.
276:27555–27561.
S K., K Y., K H., H M., T S., T K., K S., Kajiwara S., Yamaoka K., Hori K., Miyazawa H.,
Saito T., Kanno T., Shishido K. 1992. Isolation and sequence of a developmentally
regulated putative novel gene, priA, from the basidiomycete Lentinus edodes. Gene.
114:173–178.
S K., N J., B M., S B., RA P., GH B. 2006a. The Aspergillus nidulans F-box protein GrrA
links SCF activity to meiosis. Mol. Microbiol. 61:76–88.
S K., T S., S I., Y H., T O. 1993. A mutant allele skt5 affecting protoplast regeneration and
killer toxin resistance has double mutations in its wild-type structural gene in
Saccharomyces cerevisiae. Biosci. Biotechnol. Biochem. 57:1391–1393.
S N., M L., H B. 1998b. Isolation of Candida glabrata homologs of the Saccharomyces
cerevisiae KRE9 and KNH1 genes and their involvement in cell wall beta-1,6-glucan
synthesis. J. Bacteriol. 180:5020–5029.
S O., H M., T T., Y Y., F H., K G., T N., K A. 2006b. Novel hydrophobic surface binding
protein, HsbA, produced by Aspergillus oryzae. Appl. Environ. Microbiol. 72:2407–
2413.
S O., Y T., A W., M I. 2019. Carbohydrate-binding property of a cell wall integrity and stress
response component (WSC) domain of an alcohol oxidase from the rice blast pathogen
Pyricularia oryzae. Enzyme Microb. Technol. 125:13–20.
S P., F D., S B., C J., M C. 2004. Ady2p is essential for the acetate permease activity in the
yeast Saccharomyces cerevisiae. Yeast. 21:201–210.
S P., U K. 2004. A WD40 repeat protein regulates fungal cell differentiation and can be
replaced functionally by the mammalian homologue striatin. Eukaryot. Cell. 3:232–240.
S Y., S S., Q Y., X F., Z J., X Y. 2014. Biochemical characterization of the first fungal
glycoside hydrolyase family 3 $β$-N-acetylglucosaminidase from Rhizomucor miehei.
J. Agric. Food Chem. 62:5181–5190.
S Z., D G., GG P., JL C., FA L. 1994. The BTB domain, found primarily in zinc finger
proteins, defines an evolutionarily conserved family that includes several
developmentally regulated genes in Drosophila. Proc. Natl. Acad. Sci. U. S. A.

91:10717–10721.
S Z., E S., K I., S S. 2007b. Identification of a novel bifunctional delta12/delta15 fatty acid
desaturase from a basidiomycete, Coprinus cinereus TD#822-2. FEBS Lett. 581:315–
319.
S Ž., J P., J K., J S. 2015. $β$-Trefoil structure enables interactions between lectins and
protease inhibitors that regulate their biological functions. J. Biochem. 158:83–90.
Sabotič J., Renko M., Kos J. 2019. $β$-Trefoil Protease Inhibitors Unique to Higher Fungi.
Acta Chim. Slov. 66:28–36.
Sakamoto Y. 2018. Influences of environmental factors on fruiting body induction,
development and maturation in mushroom-forming fungi. Fungal Biol. Rev.
Sakamoto Y., Minato K., Nagai M., Mizuno M., Sato T. 2005. Characterization of the
Lentinula edodes exg2 gene encoding a lentinan-degrading exo-$β$-1,3-glucanase.
Curr. Genet. 2005 483. 48:195–203.
Sakamoto Y., Nakade K., Sato S., Yoshida K., Miyazaki K., Natsume S., Konno N. 2017a.
Lentinula edodes Genome Survey and Postharvest Transcriptome Analysis. Appl.
Environ. Microbiol. 83.
Sakamoto Y., Nakade K., Sato S., Yoshida K., Miyazaki K., Natsume S., Konno N. 2017b.
Lentinula edodes genome survey and postharvest transcriptome analysis. Appl.
Environ. Microbiol. 83.
Sakamoto Y., Nakade K., Sato T. 2009. Characterization of the post-harvest changes in
gene transcription in the gill of the Lentinula edodes fruiting body. Curr. Genet. 55:409–
423.
Sakamoto Y., Nakade K., Yoshida K., Natsume S., Miyazaki K., Sato S., van Peer A.F.,
Konno N. 2015. Grouping of multicopper oxidases in Lentinula edodes by sequence
similarities and expression patterns. AMB Express. 5.
Sakamoto Y., Sato S., Ito M., Ando Y., Nakahori K., Muraguchi H. 2018. Blue light exposure
and nutrient conditions influence the expression of genes involved in simultaneous
hyphal knot formation in Coprinopsis cinerea. Microbiol. Res. 217:81–90.
Sakamoto Y., Watanabe H., Nagai M., Nakade K., Takahashi M., Sato T. 2006. Lentinula
edodes tlg1 encodes a thaumatin-like protein that is involved in lentinan degradation
and fruiting body senescence. Plant Physiol. 141:793–801.
Sammer D., Krause K., Gube M., Wagner K., Kothe E. 2016. Hydrophobins in the Life Cycle
of the Ectomycorrhizal Basidiomycete Tricholoma vaccinum. PLoS One. 11:e0167773.
Sampedro J., Cosgrove D.J. 2005. The expansin superfamily. Genome Biol. 2005 612. 6:1–
11.
Sánchez-Arreguin J.A., Cabrera-Ponce J.L., León-Ramírez C.G., Camargo-Escalante M.O.,
Ruiz-Herrera J. 2020. Analysis of the photoreceptors involved in the light-depending
basidiocarp formation in Ustilago maydis. Arch. Microbiol. 202:93–103.
Sánchez-García M., Ryberg M., Khan F.K., Varga T., Nagy L.G., Hibbett D.S. 2020. Fruiting
body form, not nutritional mode, is the major driver of diversification in mushroom-
forming fungi. Proc. Natl. Acad. Sci. U. S. A. 117:32528–32534.
Sánchez-Pulido L., Martín-Belmonte F., Valencia A., Alonso M.A. 2002. MARVEL: a
conserved domain involved in membrane apposition events. Trends Biochem. Sci.
27:599–601.
Sánchez-Vallet A., Mesters J.R., Thomma B.P.H.J. 2015. The battle for chitin recognition in
plant-microbe interactions. FEMS Microbiol. Rev. 39:171–183.
Sano H., Kaneko S., Sakamoto Y., Sato T., Shishido K. 2009. The basidiomycetous
mushroom Lentinula edodes white collar-2 homolog PHRB, a partner of putative blue-
light photoreceptor PHRA, binds to a specific site in the promoter region of the L.
edodes tyrosinase gene. Fungal Genet. Biol.
Sato I., Shimizu M., Hoshino T., Takaya N. 2009. The Glutathione System of Aspergillus
nidulans Involves a Fungus-specific Glutathione S-Transferase. J. Biol. Chem.
284:8042.
Sato M., Kurahashi A., Nishibori K., Fujimori F. 2015. Development of a transformation
system for the edible mushroom Grifola frondosa: Demonstrating heterologous gene

expression and RNAi-mediated gene silencing. Mycoscience. 56:364–372.

SB I., H Q., M A., T W., Y O. 1999. Prenylation of Rho1p is required for activation of yeast 1,
3-beta-glucan synthase. J. Biol. Chem. 274:38119–38124.

Schneiter R., Pietro A. Di. 2013. The CAP protein superfamily: function in sterol export and
fungal virulence. Biomol. Concepts. 4:519–525.

Schubert D., Raudaskoski M., Knabe N., Kothe E. 2006. Ras GTPase-activating protein
Gap1 of the homobasidiomycete Schizophyllum commune regulates hyphal growth
orientation and sexual development. Eukaryot. Cell. 5:683–695.

Schubert M., Bleuler-Martinez S., Butschi A., Wälti M.A., Egloff P., Stutz K., Yan S., Wilson
I.B.H., Hengartner M.O., Aebi M., Allain F.H.T., Künzler M. 2012. Plasticity of the $β$-
trefoil protein fold in the recognition and control of invertebrate predators and parasites
by a fungal defence system. PLoS Pathog. 8.

Schulz T.A., Prinz W.A. 2007. Sterol Transport In Yeast and the Oxysterol Binding Protein
Homologue (OSH) Family. Biochim. Biophys. Acta. 1771:769.

Schwalb M.N. 1974. Effect of adenosine 3′,5′-cyclic monophosphate on the morphogenesis
of fruit bodies of Schizophyllum commune. Arch. Microbiol. 96:17–20.

Sebastian J., Rouissi T., Brar S.K. 2020. Fungal chitosan: prospects and challenges. Handb.
Chitin Chitosan.:419–452.

Seibold P.S., Lenz C., Gressler M., Hoffmeister D. 2020. The Laetiporus polyketide synthase
LpaA produces a series of antifungal polyenes. J. Antibiot. 2020 7310. 73:711–720.

Seitz L.C., Tang K., Cummings W.J., Zolan M.E. 1996. The Rad9 Gene of Coprinus
Cinereus Encodes a Proline-Rich Protein Required for Meiotic Chromosome
Condensation and Synapsis. Genetics. 142:1105.

Semana P., Powlowski J. 2019. Four aromatic intradiol ring cleavage dioxygenases from
aspergillus niger. Appl. Environ. Microbiol. 85.

SH N., FN H., K S. 2004. [Latest frontiers of meiosis research in Coprinus cinereus].
Seikagaku. 76:1450–1454.

Shahinian S., Bussey H. 2000. $β$-1,6-Glucan synthesis in Saccharomyces cerevisiae. Mol.
Microbiol. 35:477–489.

SHAW R. 1967. Fatty Acids of Fruiting Bodies of Basidiomycetes. Nat. 1967 2135071.
213:86–87.

Sheridan K.J., Lechner B.E., Keeffe G.O., Keller M.A., Werner E.R., Lindner H., Jones G.W.,
Haas H., Doyle S. 2016. Ergothioneine Biosynthesis and Functionality in the
Opportunistic Fungal Pathogen, Aspergillus fumigatus. Sci. Reports 2016 61. 6:1–17.

Shim D., Park S.G., Kim K., Bae W., Lee G.W., Ha B.S., Ro H.S., Kim M., Ryoo R., Rhee
S.K., Nou I.S., Koo C.D., Hong C.P., Ryu H. 2016. Whole genome de novo sequencing
and genome annotation of the world popular cultivated edible mushroom, Lentinula
edodes. J. Biotechnol. 223:24–25.

Shioya T., Nakamura H., Ishii N., Takahashi N., Sakamoto Y., Ozaki N., Kobayashi M.,
Okano K., Kamada T., Muraguchi H., T S., H N., N I., N T., Y S., N O., M K., K O., T K.,
H M. 2013. The Coprinopsis cinerea septin Cc.Cdc3 is involved in stipe cell elongation.
Fungal Genet. Biol. 58–59:80–90.

Šimočková M., Holič R., Tahotná D., Patton-Vogt J., Griač P. 2008. Yeast Pgc1p (YPL206c)
controls the amount of phosphatidylglycerol via a phospholipase C-type degradation
mechanism. J. Biol. Chem. 283:17107–17115.

Sipos G., Prasanna A.N., Walter M.C., O'Connor E., Bálint B., Krizsán K., Kiss B., Hess J.,
Varga T., Slot J., Riley R., Bóka B., Rigling D., Barry K., Lee J., Mihaltcheva S., LaButti
K., Lipzen A., Waldron R., Moloney N.M., Sperisen C., Kredics L., Vágvölgyi C.,
Patrignani A., Fitzpatrick D., Nagy I., Doyle S., Anderson J.B., Grigoriev I. V., Güldener
U., Münsterkötter M., Nagy L.G. 2017a. Genome expansion and lineage-specific
genetic innovations in the forest pathogenic fungi Armillaria. Nat. Ecol. Evol. 2017 112.
1:1931–1941.

Sipos G., Prasanna A.N., Walter M.C., O'Connor E., Bálint B., Krizsán K., Kiss B., Hess J.,
Varga T., Slot J., Riley R., Bóka B., Rigling D., Barry K., Lee J., Mihaltcheva S., LaButti
K., Lipzen A., Waldron R., Moloney N.M., Sperisen C., Kredics L., Vágvölgyi C.,

Patrignani A., Fitzpatrick D., Nagy I., Doyle S., Anderson J.B., Grigoriev I. V, Güldener
U., Münsterkötter M., Nagy L.G. 2017b. Genome expansion and lineage-specific
genetic innovations in the forest pathogenic fungi Armillaria. Nat. Ecol. Evol. 1:1931–
1941.
Slot J.C., Gluck-Thaler E. 2019. Metabolic gene clusters, fungal diversity, and the generation
of accessory functions. Curr. Opin. Genet. Dev. 58–59:17–24.
Šmid I., Rotter A., Gruden K., Brzin J., Buh Gašparič M., Kos J., Žel J., Sabotič J. 2015.
Clitocypin, a fungal cysteine protease inhibitor, exerts its insecticidal effect on Colorado
potato beetle larvae by inhibiting their digestive cysteine proteases. Pestic. Biochem.
Physiol. 122:59–66.
Soanes D.M., Chakrabarti A., Paszkiewicz K.H., Dawe A.L., Talbot N.J. 2012. Genome-wide
Transcriptional Profiling of Appressorium Development by the Rice Blast Fungus
Magnaporthe oryzae. PLOS Pathog. 8:e1002514.
Sommer R., Makshakova O.N., Wohlschlager T., Hutin S., Marsh M., Titz A., Künzler M.,
Varrot A. 2018. Crystal Structures of Fungal Tectonin in Complex with O-Methylated
Glycans Suggest Key Role in Innate Immune Defense. Structure. 26:391-402.e4.
Song C.H., Cho K.Y., Nair N.G., Vine J. 2018a. Growth Stimulation and Lipid Synthesis in
Lentinus Edodes. https://doi.org/10.1080/00275514.1989.12025782. 81:514–522.
Song H.-Y., Kim D.-H., Kim J.-M. 2018b. Comparative transcriptome analysis of dikaryotic
mycelia and mature fruiting bodies in the edible mushroom Lentinula edodes. Sci. Rep.
8.
Sonnenberg A.S.M., Sedaghat-Telgerd N., Lavrijssen B., Ohm R.A., Hendrickx P.M.,
Scholtmeijer K., Baars J.J.P., van Peer A. 2020. Telomere-to-telomere assembled and
centromere annotated genomes of the two main subspecies of the button mushroom
Agaricus bisporus reveal especially polymorphic chromosome ends. Sci. Reports 2020
101. 10:1–15.
Specht C.A. 1995. Isolation of the Bα and Bβ mating-type loci of Schizophyllum commune.
Curr. Genet. 28:374–379.
Spiteller P. 2015. Chemical ecology of fungi. Nat. Prod. Rep. 32:971–993.
Stadler M., Hoffmeister D. 2015. Fungal natural products—the mushroom perspective. Front.
Microbiol. 0:127.
Stadler M., Sterner O. 1998. Production of bioactive secondary metabolites in the fruit
bodies of macrofungi as a response to injury. Phytochemistry. 49:1013–1019.
Stajich J.E., Berbee M.L., Blackwell M., Hibbett D.S., James T.Y., Spatafora J.W., Taylor
J.W. 2009. The Fungi. Curr. Biol. 19:R840--R845.
Stajich J.E., Wilke S.K., Ahrén D., Au C.H., Birren B.W., Borodovsky M., Burns C., Canbäck
B., Casselton L.A., Cheng C.K., Deng J., Dietrich F.S., Fargo D.C., Farman M.L.,
Gathman A.C., Goldberg J., Guigó R., Hoegger P.J., Hooker J.B., Huggins A., James
T.Y., Kamada T., Kilaru S., Kodira C., Kües U., Kupfer D., Kwan H.S., Lomsadze A., Li
W., Lilly W.W., Ma L.-J., Mackey A.J., Manning G., Martin F., Muraguchi H., Natvig
D.O., Palmerini H., Ramesh M.A., Rehmeyer C.J., Roe B.A., Shenoy N., Stanke M.,
Ter-Hovhannisyan V., Tunlid A., Velagapudi R., Vision T.J., Zeng Q., Zolan M.E.,
Pukkila P.J. 2010. Insights into evolution of multicellular fungi from the assembled
chromosomes of the mushroom Coprinopsis cinerea (Coprinus cinereus). Proc. Natl.
Acad. Sci. U. S. A. 107:11889–94.
Steinegger M., Söding J. 2017. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. Nat. Biotechnol. 35:1026–1028.
Stogios P.J., Downs G.S., Jauhal J.J.S., Nandra S.K., Privé G.G. 2005. Sequence and
structural analysis of BTB domain proteins. Genome Biol. 2005 610. 6:1–18.
Strijbis K., Distel B. 2010. Intracellular Acetyl Unit Transport in Fungal Carbon Metabolism.
Eukaryot. Cell. 9:1809.
Stutz K., Kaech A., Aebi M., Künzler M., Hengartner M.O. 2015. Disruption of the C. elegans
Intestinal Brush Border by the Fungal Lectin CCL2 Phenocopies Dietary Lectin Toxicity
in Mammals. PLoS One. 10:e0129381.
Sun J., Ma Y., Xu X., Liu Z., Zou L. 2020. Molecular cloning and bioinformatics analyses of a

GH3 beta-glucosidase from Auricularia heimuer. http://mc.manuscriptcentral.com/tbeq. 34:850–859.

Sunagawa M., Magae Y. 2005. Isolation of genes differentially expressed during the fruit body development of Pleurotus ostreatus by differential display of RAPD. FEMS Microbiol. Lett. 246:279–284.

Suzuki S., Sarikaya Bayram Ö., Bayram Ö., Braus G.H. 2013. conF and conJ contribute to conidia germination and stress response in the filamentous fungus Aspergillus nidulans. Fungal Genet. Biol. 56:42–53.

Swamy S., Uno I., Ishikawa T. 1985. Regulation of cyclic AMP metabolism by the incompatibility factors in Coprinus cinereus. J. Gen. Microbiol. 131:3211–3217.

Szczesny P., Iacovache I., Muszewska A., Ginalski K., van der Goot F.G., Grynberg M. 2011. Extending the Aerolysin Family: From Bacteria to Vertebrates. PLoS One. 6:e20349.

Szeto C.Y., Leung G.S., Kwan H.S., CY S., GS L., HS K. 2007. Le.MAPK and its interacting partner, Le.DRMIP, in fruiting body development in Lentinula edodes. Gene. 393:87–93.

T H., T S., H E. 2004a. Isolation of genes specifically expressed in the fruit body of the edible basidiomycete Lentinula edodes. Biosci. Biotechnol. Biochem. 68:468–472.

T I., K S. 2000. Decreased zinc ion accumulation by the basidiomycete Lentinus edodes over-expressing L. edodes PriA gene. FEMS Microbiol. Lett. 193:111–115.

T L.-O., H R.-R., J A. 2003. Reactive oxygen species generated by microbial NADPH oxidase NoxA regulate sexual development in Aspergillus nidulans. Mol. Microbiol. 50:1241–1255.

T T., N N., M U., T K., G T., M M., K A. 2015. Ustiloxins, fungal cyclic peptides, are ribosomally synthesized in Ustilaginoidea virens. Bioinformatics. 31:981–985.

T T., Y M., K C., K O., N K., N S.-T., N D., Y H., K T. 2004b. Protein sequence analysis, cloning, and expression of flammutoxin, a pore-forming cytolysin from Flammulina velutipes. Maturation of dimeric precursor to monomeric active form by carboxyl-terminal truncation. J. Biol. Chem. 279:54161–54172.

T W., A B., K Z., SC V., U A. dem K., P G., S B.-M., MO H., M A., M K. 2011. Nematotoxicity of Marasmius oreades agglutinin (MOA) depends on glycolipid binding and cysteine protease activity. J. Biol. Chem. 286:30337–30343.

Tamaki H. 2007. Glucose-stimulated cAMP-protein kinase a pathway in yeast Saccharomyces cerevisiae. J. Biosci. Bioeng. 104:245–250.

Tan Y.H., Moore D. High concentrations of mannitol in the shiitake mushroom Lentinula edodes. .

Tang L., Jian H., Song C., Bao D., Shang X., Wu D., Tan Q., Zhang X. 2013. Transcriptome analysis of candidate genes and signaling pathways associated with light-induced brown film formation in Lentinula edodes. Appl. Microbiol. Biotechnol. 97:4977–4989.

Tang X., Ding X., Hou Y. 2020. Comparative analysis of transcriptomes revealed the molecular mechanism of development of Tricholoma matsutake at different stages of fruiting bodies. Food Sci. Biotechnol. 2020 297. 29:939–951.

Tao Q., Ma K., Yang Y., Wang K., Chen B., Huang Y., Han J., Bao L., Liu X.-B., Yang Z., Yin W.-B., Liu H. 2016. Bioactive Sesquiterpenes from the Edible Mushroom Flammulina velutipes and Their Biosynthetic Pathway Confirmed by Genome Analysis and Chemical Evidence. J. Org. Chem. 81:9867–9877.

Tao Y., Xie B., Yang Z., Chen Z., Chen B., Deng Y., Jiang Y., van Peer A.F. 2013. Identification and expression analysis of a new glycoside hydrolase family 55 exo-beta-1,3-glucanase-encoding gene in Volvariella volvacea suggests a role in fruiting body development. Gene. 527:154–160.

Tatebe H., Shiozaki K. 2017. Evolutionary conservation of the components in the tor signaling pathways. Biomolecules. 7.

Taylor J.W., Ellison C.E. 2010. Mushrooms: Morphological complexity in the fungi. Proc. Natl. Acad. Sci. 107:11655–11656.

Tayyrov A., Schmieder S.S., Bleuler-Martinez S., Plaza D.F., Künzlera M. 2018. Toxicity of

potential fungal defense proteins towards the fungivorous nematodes Aphelenchus avenae and Bursaphelenchus okinawaensis. Appl. Environ. Microbiol. 84.

Tayyrov A., Stanley C.E., Azevedo S., Künzler M. 2019. Combining microfluidics and RNA-sequencing to assess the inducible defensome of a mushroom against nematodes. BMC Genomics 2019 201. 20:1–13.

Teichert I., Dahlmann T.A., Kuck U., Nowrousian M., Kück U., Nowrousian M. 2017. RNA Editing During Sexual Development Occurs in Distantly Related Filamentous Ascomycetes. Genome Biol Evol. 9:855–868.

Teichert I., Steffens E.K., Schnaß N., Fränzel B., Krisp C., Wolters D.A., Kück U. 2014. PRO40 Is a Scaffold Protein of the Cell Wall Integrity Pathway, Linking the MAP Kinase Module to the Upstream Activator Protein Kinase C. PLoS Genet. 10:e1004582.

Telzrow C.L., Nichols C.B., Castro-Lopez N., Wormley F.L., Alspaugh J.A. 2019. A fungal arrestin protein contributes to cell cycle progression and pathogenesis. MBio. 10.

Temp U., Eggert C. 1999. Novel interaction between laccase and cellobiose dehydrogenase during pigment synthesis in the white rot fungus Pycnoporus cinnabarinus. Appl. Environ. Microbiol. 65:389–395.

Terashima K., Yuki K., Muraguchi H., Akiyama M., Kamada T. 2005. The dst1 gene involved in mushroom photomorphogenesis of Coprinus cinereus encodes a putative photoreceptor for blue light. Genetics.

Thammahong A., Puttikamonkul S., Perfect J.R., Brennan R.G., Cramer R.A. 2017. Central Role of the Trehalose Biosynthesis Pathway in the Pathogenesis of Human Fungal Infections: Opportunities and Challenges for Therapeutic Development. Microbiol. Mol. Biol. Rev. 81.

TM H., JW B., MD A., Y R.-D., C S., NP K. 2008. RNA silencing gene truncation in the filamentous fungus Aspergillus nidulans. Eukaryot. Cell. 7:339–349.

Tong X., Zhang H., Wang F., Xue Z., Cao J., Peng C., Guo J. 2020. Comparative transcriptome analysis revealed genes involved in the fruiting body development of Ophiocordyceps sinensis. PeerJ.

Torrens-Spence M.P., Liu C.-T., Pluskal T., Chung Y.K., Weng J.-K. 2018. Monoamine Biosynthesis via a Noncanonical Calcium-Activatable Aromatic Amino Acid Decarboxylase in Psilocybin Mushroom. ACS Chem. Biol. 13:3343–3353.

Tovar-Herrera O.E., Batista-García R.A., Sánchez-Carbente M.D.R., Iracheta-Cárdenas M.M., Arévalo-Niño K., Folch-Mallol J.L. 2015. A novel expansin protein from the white-rot fungus Schizophyllum commune. PLoS One. 10.

Tsitsigiannis D.I., Keller N.P. 2007. Oxylipins as developmental and host-fungal communication signals. Trends Microbiol.

Tsunematsu Y., Takanishi J., Asai S., Masuya T., Nakazawa T., Watanabe K. 2019. Genomic Mushroom Hunting Decrypts Coprinoferrin, A Siderophore Secondary Metabolite Vital to Fungal Cell Development. Org. Lett. 21:7582–7586.

Tullis P. 2021. How ecstasy and psilocybin are shaking up psychiatry. Nature. 589:506–509.

U K., Kues U., Kües U. 2000. Life history and developmental processes in the basidiomycete Coprinus cinereus. Microbiol. Mol. Biol. Rev. 64:316–353.

Umemura M. 2020. Peptides derived from Kex2-processed repeat proteins are widely distributed and highly diverse in the Fungi kingdom. Fungal Biol. Biotechnol. 7:11.

Uno I., Ishikawa T. 1973. Purification and identification of the fruiting-inducing substances in Coprinus macrorhizus. J. Bacteriol. 113:1240–1248.

Varga T., Földi C., Bense V., Nagy L.G. 2021. Developmental innovations promote species diversification in mushroom-forming fungi. bioRxiv.:2021.03.10.434564.

Varga T., Krizsán K., Földi C., Dima B., Sánchez-García M., Sánchez-Ramírez S., Szöllősi G.J., Szarkándi J.G., Papp V., Albert L., Andreopoulos W., Angelini C., Antonín V., Barry K.W., Bougher N.L., Buchanan P., Buyck B., Bense V., Catcheside P., Chovatia M., Cooper J., Dämon W., Desjardin D., Finy P., Geml J., Haridas S., Hughes K., Justo A., Karasiński D., Kautmanova I., Kiss B., Kocsubé S., Kotiranta H., LaButti K.M., Lechner B.E., Liimatainen K., Lipzen A., Lukács Z., Mihaltcheva S., Morgado L.N., Niskanen T., Noordeloos M.E., Ohm R.A., Ortiz-Santana B., Ovrebo C., Rácz N., Riley

R., Savchenko A., Shiryaev A., Soop K., Spirin V., Szebenyi C., Tomšovský M., Tulloss R.E., Uehling J., Grigoriev I. V., Vágvölgyi C., Papp T., Martin F.M., Miettinen O., Hibbett D.S., Nagy L.G. 2019. Megaphylogeny resolves global patterns of mushroom evolution. Nat. Ecol. Evol.

Veeneman B.A., Shukla S., Dhanasekaran S.M., Chinnaiyan A.M., Nesvizhskii A.I. 2016. Two-pass alignment improves novel splice junction quantification. Bioinformatics. 32:43–49.

Venter P., Kock J.L.F., Kumar G.S., Botha A., Coetzee D.J., Botes P.J., Bhatt R.K., Falck J.R., Schewe T., Nigam S. 1997. Production of 3R-hydroxy-polyenoic fatty acids by the yeast Dipodascopsis uninucleata. Lipids. 32:1277–1283.

Verdín J., Sánchez-León E., Rico-Ramírez A.M., Martínez-Núñez L., Fajardo-Somera R.A., Riquelme M. 2019. Off the wall: The rhyme and reason of Neurospora crassa hyphal morphogenesis. Cell Surf. 5:100020.

Virágh M., Merényi Z., Csernetics Á., Földi C., Sahu N., Liu X.-B., Hibbett D.S., Nagy L.G. 2021. Evolutionary Morphogenesis of Sexual Fruiting Bodies in Basidiomycota: Toward a New Evo-Devo Synthesis. Microbiol. Mol. Biol. Rev. 86.

Vogt E., Künzler M. 2019. Discovery of novel fungal RiPP biosynthetic pathways and their application for the development of peptide therapeutics. Appl. Microbiol. Biotechnol. 2019 10314. 103:5567–5581.

Vonk P.J., Ohm R.A. 2018. The role of homeodomain transcription factors in fungal development. Fungal Biol. Rev. 32:219–230.

Vonk P.J., Ohm R.A. 2021. H3K4me2 ChIP-Seq reveals the epigenetic landscape during mushroom formation and novel developmental regulators of Schizophyllum commune. Sci. Reports 2021 111. 11:1–15.

W G., Y W., C X., Y Z., Z Z., Y P. 2020. Whole genome sequence of an edible and medicinal mushroom, Hericium erinaceus (Basidiomycota, Fungi). Genomics. 112:2393–2399.

W Z., X W., Y Z.Z., Z L., W Z., X N., Y Z.Z., S P., Y Z.Z., S Y., Zhang W., Wu X., Zhou Y., Liu Z., Zhang W., Niu X., Zhao Y., Pei S., Zhao Y., Yuan S. 2014. Characterization of stipe elongation of the mushroom Coprinopsis cinerea. Microbiology. 160:1893–1902.

Walser P.J., Kües U., Aebi M., Künzler M. 2005. Ligand interactions of the Coprinopsis cinerea galectins. Fungal Genet. Biol.

Wälti M.A., Walser P.J., Thore S., Grünler A., Bednar M., Künzler M., Aebi M. 2008. Structural Basis for Chitotetraose Coordination by CGL3, a Novel Galectin-Related Protein from Coprinopsis cinerea. J. Mol. Biol. 379:146–159.

Walton J. 2018. Ecology and Evolution of the <Emphasis Type="Italic">Amanita</Emphasis> Cyclic Peptide Toxins. Cycl. Pept. Toxins Amanita Other Poisonous Mushrooms.:93–130.

Wang H.X., Ng T.B. 2005. Purification of a laccase from fruiting bodies of the mushroom Pleurotus eryngii. Appl. Microbiol. Biotechnol. 2005 695. 69:521–525.

Wang L., Gao W., Wu X., Zhao M., Qu J., Huang C., Zhang J. 2018a. Genome-Wide Characterization and Expression Analyses of Pleurotus ostreatus MYB Transcription Factors during Developmental Stages and under Heat Stress Based on de novo Sequenced Genome. Int. J. Mol. Sci. 19.

Wang L., Tian X., Gyawali R., Lin X. 2013. Fungal adhesion protein guides community behaviors and autoinduction in a paracrine manner. Proc Natl Acad Sci U S A. 110:11571–11576.

Wang L., Zhai B., Lin X. 2012. The Link between Morphotype Transition and Virulence in Cryptococcus neoformans. PLoS Pathog. 8:e1002765.

Wang R., Ma P., Li C., Xiao L., Liang Z., Dong J. 2019. Combining transcriptomics and metabolomics to reveal the underlying molecular mechanism of ergosterol biosynthesis during the fruiting process of Flammulina velutipes. BMC Genomics 2019 201. 20:1–12.

Wang Y., Niu X., Guo X., Yu H., Liu Z., Zhang Z., Yuan S. 2018b. Heterologous expression, characterization and possible functions of the chitin deacetylases, Cda1 and Cda2, from mushroom Coprinopsis cinerea. Glycobiology. 28:318–332.

Wang Z., Li N., Li J., Dunlap J.C., Trail F., Townsend J.P. 2016. The fast-evolving phy-2

gene modulates sexual development in response to light in the model fungus
Neurospora crassa. MBio. 7.
Wannet W.J.B., Aben E.M.J., Van Der Drift C., Van Griensven L.J.L.D., Vogels G.D., Op
Den Camp H.J.M. 1999. Trehalose phosphorylase activity and carbohydrate levels
during axenic fruiting in three Agaricus bisporus strains. Curr. Microbiol. 39:205–210.
Watcharawipas A., Watanabe D., Takagi H. 2018. Sodium Acetate Responses in
Saccharomyces cerevisiae and the Ubiquitin Ligase Rsp5. Front. Microbiol. 9.
Wawra S., Fesel P., Widmer H., Neumann U., Lahrmann U., Becker S., Hehemann J.-H.,
Langen G., Zuccaro A. 2019. FGB1 and WSC3 are in planta-induced $β$-glucan-
binding fungal lectins with different functions. New Phytol. 222:1493–1506.
Weiberg A., Wang M., Lin F.M., Zhao H., Zhang Z., Kaloshian I., Huang H. Da, Jin H. 2013.
Fungal small RNAs suppress plant immunity by hijacking host RNA interference
pathways. Science (80-.). 342:118–123.
Weigand-Heller A.B.J., Kris-Etherton P.M., Beelman R.B. 2012. The bioavailability of
ergothioneine from mushrooms (Agaricus bisporus) and the acute effects on antioxidant
capacity and biomarkers of inflammation. Prev. Med. (Baltim). 54:S75--S78.
WELLS T.K., HAMMOND J.B.W., DICKERSON A.G. 1987. VARIATIONS IN ACTIVITIES
OF GLYCOGEN PHOSPHORYLASE AND TREHALASE DURING THE PERIODIC
FRUITING OF THE EDIBLE MUSHROOM AGARICUS BISPORUS (LANGE) IMBACH.
New Phytol. 105:273–280.
Wen J., Zhang Z., Gong L., Xun H., Li J., Qi B., Wang Q., Li X., Li Y., Liu B. 2019.
Transcriptome Changes during Major Developmental Transitions Accompanied with
Little Alteration of DNA Methylome in Two Pleurotus Species. Genes (Basel). 10.
Wen L., Huang H.-M., Juang R.-H., Lin C.-T. 2007. Biochemical characterization of 1-Cys
peroxiredoxin from Antrodia camphorata. Appl. Microbiol. Biotechnol. 2006 736.
73:1314–1322.
Whiteford J.R., Thurston C.F. 2000. The molecular genetics of cultivated mushrooms. Adv.
Microb. Physiol. 42:1–23.
Wilinski D., Buter N., Klocko A.D., Lapointe C.P., Selker E.U., Gasch A.P., Wickens M. 2017.
Recurrent rewiring and emergence of RNA regulatory networks. Proc. Natl. Acad. Sci.
114:E2816--E2825.
Wirth S., Freihorst D., Krause K., Kothe E. 2021. What role might non-mating receptors play
in schizophyllum commune? J. Fungi. 7.
Wohlschlager T., Butschi A., Grassi P., Sutov G., Gauss R., Hauck D., Schmieder S.S.,
Knobel M., Titz A., Dell A., Haslam S.M., Hengartner M.O., Aebi M., Kun☐zler M. 2014.
Methylated glycans as conserved targets of animal and fungal innate defense. Proc.
Natl. Acad. Sci. U. S. A. 111:E2787–E2796.
Wood W.F., Clark T.J., Bradshaw D.E., Foy B.D., Largent D.L., Thompson B.L. 2004.
Clitolactone: A Banana Slug Antifeedant from Clitocybe flaccida. Mycologia. 96:23.
Wösten H.A.B. 2003. Hydrophobins: Multipurpose Proteins.
http://dx.doi.org/10.1146/annurev.micro.55.1.625. 55:625–646.
Wösten H.A.B., Scholtmeijer K. 2015. Applications of hydrophobins: current state and
perspectives. Appl. Microbiol. Biotechnol. 2015 994. 99:1587–1597.
Wu B., Gaskell J., Zhang J., Toapanta C., Ahrendt S., Grigoriev I. V, Blanchette R.A.,
Schilling J.S., Master E., Cullen D., Hibbett D.S. 2019. Evolution of substrate-specific
gene expression and RNA editing in brown rot wood-decaying fungi. ISME J. 2019 136.
13:1391–1403.
Wu B., Xu Z., Knudson A., Carlson A., Chen N., Kovaka S., LaButti K., Lipzen A., Pennachio
C., Riley R., Schakwitz W., Umezawa K., Ohm R.A., Grigoriev I. V, Nagy L.G., Gibbons
J., Hibbett D. 2018. Genomics and Development of *Lentinus tigrinus*☐: A White-Rot
Wood-Decaying Mushroom with Dimorphic Fruiting Bodies. Genome Biol. Evol.
10:3250–3261.
Wu F.-L., Zhang G., Ren A., Dang Z.-H., Shi L., Jiang A.-L., Zhao M.-W., FL W., G Z., A R.,
ZH D., L S., AL J., MW Z. 2016. The pH-responsive transcription factor PacC regulates
mycelial growth, fruiting body development, and ganoderic acid biosynthesis in

Ganoderma lucidum. Mycologia. 108:1104–1113.
Wu L., van Peer A., Song W., Wang H., Chen M., Tan Q., Song C., Zhang M., Bao D. 2013.
Cloning of the Lentinula edodes B mating-type locus and identification of the genetic
structure controlling B mating. Gene. 531:270–278.
Wu T., Hu C., Xie B., Wei S., Zhang L., Zhu Z., Zhang Z., Li S. 2020a. A putative
transcription factor LFC1 negatively regulates development and yield of winter
mushroom. Appl. Microbiol. Biotechnol. 2020 10413. 104:5827–5844.
Wu T., Zhang Z., Hu C., Zhang L., Wei S., Li S. 2020b. A WD40 Protein Encoding Gene
Fvcpc2 Positively Regulates Mushroom Development and Yield in Flammulina
velutipes. Front. Microbiol. 11:498.
WW C., KT N., RS S., CH A., HS K. 2008. Gene expression studies of the dikaryotic
mycelium and primordium of Lentinula edodes by serial analysis of gene expression.
Mycol. Res. 112:950–964.
Wymelenberg A. Vanden, Gaskell J., Mozuch M., Kersten P., Sabat G., Martinez D., Cullen
D. 2009. Transcriptome and Secretome Analyses of Phanerochaete chrysosporium
Reveal Complex Patterns of Gene Expression. Appl. Environ. Microbiol. 75:4058.
X C., JH H., YY L., PT W.L., HS K., Cheng X., Hui J.H.L., Lee Y.Y., Wan Law P.T., Kwan
H.S. 2015. A "developmental hourglass" in fungi. Mol. Biol. Evol. 32:1556–1566.
X L., L F., C W., P C. 2018. Cryo-EM structure of a mammalian RNA polymerase II
elongation complex inhibited by $α$-amanitin. J. Biol. Chem. 293:7189–7194.
X L., S D. 2009. Molecular characterization of a new acetyl xylan esterase (AXEII) from
edible straw mushroom Volvariella volvacea with both de-O-acetylation and de-N-
acetylation activity. FEMS Microbiol. Lett. 295:50–56.
Xie C., Gong W., Zhu Z., Yan L., Hu Z., Peng Y. 2018. Comparative transcriptomics of
Pleurotus eryngii reveals blue-light regulation of carbohydrate-active enzymes
(CAZymes) expression at primordium differentiated into fruiting body stage. Genomics.
110:201–209.
Xie Y., Chang J., Kwan H.S. 2020. Carbon metabolism and transcriptome in developmental
paths differentiation of a homokaryotic Coprinopsis cinerea strain. Fungal Genet. Biol.
143:103432.
Xu G., Ma H., Nei M., Kong H. 2009. Evolution of F-box genes in plants: Different modes of
sequence divergence and their relationships with functional diversification. Proc. Natl.
Acad. Sci. 106:835–840.
Xu H., Navarro-Ródenas A., Cooke J.E.K., Zwiazek J.J., H X., A N.-R., JE C., JJ Z., Xu H.,
Navarro-Ródenas A., Cooke J.E.K., Zwiazek J.J. 2015. Transcript profiling of
aquaporins during basidiocarp development in Laccaria bicolor ectomycorrhizal with
Picea glauca. Mycorrhiza 2015 261. 26:19–31.
Xue C., Hsueh Y.P., Heitman J. 2008. Magnificent seven: Roles of G protein-coupled
receptors in extracellular sensing in fungi. FEMS Microbiol. Rev.
Y F., A O.-U., T G., Y P., NO K. 2015. Involvement of a caleosin in lipid storage, spore
dispersal, and virulence in the entomopathogenic filamentous fungus, Beauveria
bassiana. Environ. Microbiol. 17:4600–4614.
Y K., H K. 2014. Fungal lectins: a growing family. Methods Mol. Biol. 1200:15–38.
Y M., S K., M S., K S., T Y., M N., K B., Miyazaki Y., Kaneko S., Sunagawa M., Shishido K.,
Yamazaki T., Nakamura M., Babasaki K. 2007. The fruiting-specific Le.flp1 gene,
encoding a novel fungal fasciclin-like protein, of the basidiomycetous mushroom
Lentinula edodes. Curr. Genet. 51:367–375.
Y S., K N., N K., Sakamoto Y., Nakade K., Konno N. 2011. Endo-$β$-1,3-glucanase GLU1,
from the fruiting body of Lentinula edodes, belongs to a new glycoside hydrolase family.
Appl. Environ. Microbiol. 77:8350–8354.
Y S., T I., T S., Sakamoto Y., Irie T., Sato T. 2005. Isolation and characterization of a fruiting
body-specific exo-beta-1,3-glucanase-encoding gene, exg1, from Lentinula edodes.
Curr. Genet. 47:244–252.
Y T., M T., J S., S U., Y K., N O., M I., T Y. 2009. Tamavidins--novel avidin-like biotin-binding
proteins from the Tamogitake mushroom. FEBS J. 276:1383–1397.

Y T., R C., J Y., Y L., Z T., H S., B X. 2019. A hydrophobin gene, Hyd9, plays an important role in the formation of aerial hyphae and primordia in Flammulina filiformis. Gene. 706:84–90.

Y T., S T., T K., T J. 2013. Molecular characterization of a lipoxygenase from the basidiomycete mushroom Pleurotus ostreatus. Biosci. Biotechnol. Biochem. 77:38–45.

Y W., X Z., W L. 2018. De novo transcriptomic analysis during Lentinula edodes fruiting body growth. Gene. 641:326–334.

Yamada-Okabe T., Sakamori Y., Mio T., Yamada-Okabe H. 2001. Identification and characterization of the genes for N-acetylglucosamine kinase and N-acetylglucosamine-phosphate deacetylase in the pathogenic fungus Candida albicans. Eur. J. Biochem. 268:2498–2505.

Yamagishi K., Kimura T., Suzuki M., Shinmoto H. 2002. Suppression of fruit-body formation by constitutively active G-protein α-subunits ScGP-A and ScGP-C in the homobasidiomycete Schizophyllum commune. Microbiology. 148:2797–2809.

Yamagishi K., Kimura T., Suzuki M., Shinmoto H., Yamaki K.J. 2004. Elevation of intracellular cAMP levels by dominant active heterotrimeric G protein alpha subunits ScGP-A and ScGP-C in homobasidiomycete, Schizophyllum commune. Biosci. Biotechnol. Biochem. 68:1017–1026.

Yan J.-J.J., Tong Z.-J.J., Liu Y.-Y.Y., Li Y.-N.N., Zhao C., Mukhtar I., Tao Y.-X.X., Chen B.-Z.Z., Deng Y.-J.J., Xie B.-G.G. 2019. Comparative transcriptomics of Flammulina filiformis suggests a high CO2 concentration inhibits early pileus expansion by decreasing cell division control pathways. Int. J. Mol. Sci. 20:5923.

Ye S.-Q., Zou Y., Zheng Q.-W., Liu Y.-L., Li R.-R., Lin J.-F., Guo L.-Q. 2021. TMT-MS/MS proteomic analysis of the carbohydrate-active enzymes in the fruiting body of Pleurotus tuoliensis during storage. J. Sci. Food Agric. 101:1879–1891.

Yin G., Padhi S., Lee S., Hung R., Zhao G., Bennett J.W. 2015. Effects of Three Volatile Oxylipins on Colony Development in Two Species of Fungi and on Drosophila Larval Metamorphosis. Curr. Microbiol. 71:347–356.

Yoo S., Lee H.-Y., Markkandan K., Moon S., Ahn Y.J., Ji S., Ko J., Kim S.-J., Ryu H., Hong C.P. 2019. Comparative transcriptome analysis identified candidate genes involved in mycelium browning in Lentinula edodes. 20.

Yu Z., Armant O., Fischer R. 2016. Fungi use the SakA (HogA) pathway for phytochrome-dependent light signalling. Nat. Microbiol. 2016 15. 1:1–7.

Yu Z., Fischer R. 2018. Light sensing and responses in fungi. Nat. Rev. Microbiol. 2018 171. 17:25–36.

Yun Y.H., Koo J.S., Kim S.H., Kong W.S. 2015. Cloning and expression analysis of phenylalanine ammonia-lyase gene in the mycelium and fruit body of the edible mushroom Flammulina velutipes. Mycobiology. 43:327–332.

Zawirska-Wojtasiak R. 2004. Optical purity of (R)-(-)-1-octen-3-ol in the aroma of various species of edible mushrooms. Food Chem. 86:113–118.

Zeng X., Ling H., Yang J., Chen J., Guo S. 2018. Proteome analysis provides insight into the regulation of bioactive metabolites in Hericium erinaceus. Gene. 666:108–115.

Zhang C., Chen X., Lee R.T.C., Rehka T., Maurer-Stroh S., Rühl M. 2021a. Bioinformatics-aided identification, characterization and applications of mushroom linalool synthases. Commun. Biol. 4:1–11.

Zhang C., Chen X., Orban A., Shukal S., Birk F., Too H.-P., Rühl M. 2020. Agrocybe aegerita Serves As a Gateway for Identifying Sesquiterpene Biosynthetic Enzymes in Higher Fungi. ACS Chem. Biol. 15:1268–1277.

Zhang J., Chen H., Chen M., Ren A., Huang J., Wang H., Zhao M., Feng Z. 2015a. Cloning and functional analysis of a laccase gene during fruiting body formation in Hypsizygus marmoreus. Microbiol. Res. 179:54–63.

Zhang J., Hao H., Liu H., Wang Q., Chen M., Feng Z., Chen H. 2021b. Genetic and functional analysis of the Zn(II)2Cys6 transcription factor HADA-1 in Hypsizygus marmoreus. Appl. Microbiol. Biotechnol. 2021 1057. 105:2815–2829.

Zhang J., Ren A., Chen H., Zhao M., Shi L., Chen M., Wang H., Feng Z. 2015b.

Transcriptome Analysis and Its Application in Identifying Genes Associated with Fruiting Body Development in Basidiomycete Hypsizygus marmoreus. PLoS One. 10:e0123025.

Zhang J., Shen N., Li C., Xiang X., Liu G., Gui Y., Patev S., Hibbett D.S., Barry K., Andreopoulos W., Lipzen A., Riley R., He G., Yan M., Grigoriev I. V., Shan Kwan H., Kit Cheung M., Bian Y., Xiao Y. 2021c. Population genomics provides insights into the genetic basis of adaptive evolution in the mushroom-forming fungus Lentinula edodes. J. Adv. Res.

Zhang L., Gong W., Li C., Shen N., Gui Y., Bian Y., Kwan H.S., Cheung M.K., Xiao Y. 2021d. RNA-Seq-based high-resolution linkage map reveals the genetic architecture of fruiting body development in shiitake mushroom, Lentinula edodes. Comput. Struct. Biotechnol. J. 19:1641–1653.

Zhang X., Krause K.-H., Xenarios I., Soldati T., Boeckmann B. 2013. Evolution of the Ferric Reductase Domain (FRD) Superfamily: Modularity, Functional Diversification, and Signature Motifs. PLoS One. 8:e58126.

Zhang Z.-M., Wu W.-W., Li G.-K. 2008. A GC--MS Study of the Volatile Organic Composition of Straw and Oyster Mushrooms During Maturity and its Relation to Antioxidant Activity. J. Chromatogr. Sci. 46:690–696.

Zhao C., Waalwijk C., de Wit P.J.G.M., Tang D., van der Lee T. 2013. RNA-Seq analysis reveals new gene models and alternative splicing in the fungal pathogen Fusarium graminearum. BMC Genomics 2013 141. 14:1–16.

Zhao J., Kwan H.S. 1999. Characterization, molecular cloning, and differential expression analysis of laccase genes from the edible mushroom Lentinula edodes. Appl. Environ. Microbiol. 65:4908–4913.

Zhao Z., Liu H., Luo Y., Zhou S., An L., Wang C., Jin Q., Zhou M., Xu J.R. 2014. Molecular evolution and functional divergence of tubulin superfamily in the fungal tree of life. Sci. Rep. 4.

Zheng N., Shabek N. 2017. Ubiquitin Ligases: Structure, Function, and Regulation. https://doi.org/10.1146/annurev-biochem-060815-014922. 86:129–157.

Zhou J., Kang L., Liu C., Niu X., Wang X., Liu H., Zhang W., Liu Z., Latgé J.-P., Yuan S. 2019. Chitinases Play a Key Role in Stipe Cell Wall Extension in the Mushroom Coprinopsis cinerea. Appl. Environ. Microbiol. 85.

Zhou S., Li X., Lüli Y., Li X., Chen Z.H., Yuan P., Yang Z.L., Li G., Luo H. 2021. Novel Cyclic Peptides from Lethal Amanita Mushrooms through a Genome-Guided Approach. J. Fungi. 7:204.

Zhou S., Ma F., Zhang X., Zhang J. 2016a. Carbohydrate changes during growth and fruiting in Pleurotus ostreatus. Fungal Biol. 120:852–861.

Zhou S., Ma F., Zhang X., Zhang J., S Z., F M., X Z., J Z., Zhou S., Ma F., Zhang X., Zhang J., S Z., F M., X Z., J Z. 2016b. Carbohydrate changes during growth and fruiting in Pleurotus ostreatus. Fungal Biol. 120:852–861.

Zhou S., Zhang J., Ma F., Tang C., Tang Q., Zhang X. 2018. Investigation of lignocellulolytic enzymes during different growth phases of Ganoderma lucidum strain G0119 using genomic, transcriptomic and secretomic analyses. PLoS One. 13:e0198404.

Zhou Y., Chen L., Fan X., Bian Y. 2014. De Novo Assembly of Auricularia polytricha Transcriptome Using Illumina Sequencing for Gene Discovery and SSR Marker Identification. PLoS One. 9:e91740.

Zhou Y., Kang L., Niu X., Wang J., Liu Z., Yuan S. 2016c. Purification, characterization and physiological significance of a chitinase from the pilei of Coprinopsis cinerea fruiting bodies. FEMS Microbiol. Lett. 363:1–7.

Zhu W., Hu J., Li Y., Yang B., Guan Y., Xu C., Chen F., Chi J., Bao Y. 2019. Comparative Proteomic Analysis of Pleurotus ostreatus Reveals Great Metabolic Differences in the Cap and Stipe Development and the Potential Role of Ca2+ in the Primordium Differentiation. Int. J. Mol. Sci. 2019, Vol. 20, Page 6317. 20:6317.

Zhu Y., Luo H., Zhang X., Song J., Sun C., Ji A., Xu J., Chen S. 2014. Abundant and Selective RNA-Editing Events in the Medicinal Mushroom Ganoderma lucidum.

Genetics. 196:1047.

Zhu Y., Zhuang L., Goodell B., Cao J., Mahaney J. 2016. Iron sequestration in brown-rot
fungi by oxalate and the production of reactive oxygen species (ROS). Int. Biodeterior.
Biodegradation. 109:185–190.

卢园萍, 郭仲杰, 蔡志欣, 施肖堃, 陈美元, 廖剑华, 王泽生, Yuan-Ping L.U., Zhong-Jie G.U.O.,
Zhi-Xin C.A.I., Xiao-Kun S.H.I., Mei-Yuan C., Jian-Hua L., Ze-Sheng W. 2019.
双孢蘑菇Septin基因Ab.Cdc3的克隆与分析. 菌物学报. 37:1635–1642.

7040

Figures

7042

Fig. 1. Species used in this study. Top row left to right: *Agaricus bisporus, Cyclocybe aegerita, Armillaria ostoyae, Auriculariopsis ampla;* Middle row: *Coprinopsis cinerea, Laccaria bicolor, Lentinula edodes, Lentinus tigrinus;* Bottom row: *Mycena kentingensis, Phanerochaete chrysosporium, Pleurotus ostreatus, Schizophyllum commune*

7048
7049

7050
7051 **Fig. 2.** The distribution of conserved developmentally regulated orthogroups across species.
7052 A The number of orthogroups as a function of the number of species in which the orthogroup
7053 is developmentally regulated at fold change >2 (top) and fold change >4 (bottom). Horizontal
7054 axis represent the number of species. B Numbers of genes in each of the functional
7055 categories discussed in the paper.
7056

Fig. 3 Expression heatmap of DNA replication, repair, mitosis and meiosis related genes in *C. cinerea*. A model expression trajectory, illustrating the two peaks characteristic of most examined species, is shown above the left panel. Well-delimited complexes mentioned in the paper are shown separately. Genes are denoted by Protein IDs. Blue and red colors represent low and high expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7068
Fig. 4. Expression heatmap of ribosomal protein encoding genes in *C. cinerea*. Genes are
denoted by Protein IDs. Blue and red colors represent low and high expression, respectively.
Developmental stages are abbreviated as follows: VM – vegetative mycelium, H – hyphal

7072   knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap,
7073   YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting
7074   body cap (including gills), FB.S – mature fruiting body stipe.
7075
7076

7077
7078   **Fig. 5.** Basic metabolic pathways of *C. cinerea.* Genes in the pathways are denoted by the
7079   *S. cerevisiae* gene name, followed by the protein ID of the *C. cinerea* ortholog and, in
7080   parentheses the number of species in which the gene was developmentally regulated at fold
7081   change >2/>4. Pathways are colored differently for clarity.
7082

7083
Fig. 6. Expression heatmap of ergosterol and sphingolipid biosynthesis related genes in *C.*
cinerea. Genes are denoted by Protein IDs. Blue and red colors represent low and high
expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative
mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C –
young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe,
FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7090

7091
Fig. 7. Expression heatmap of membrane phospholipid and fatty acid biosynthetic genes in
C. cinerea. Genes are denoted by Protein IDs. Blue and red colors represent low and high
expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative
mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C –
young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe,
FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.
7098

7099

7100
Fig. 8 Expression heatmap of putatively linoleic acid biosynthesis related fatty acid
desaturase genes in *C. cinerea*. Genes are denoted by Protein IDs. Blue and red colors
represent low and high expression, respectively. Developmental stages are abbreviated as
follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2
primordium, YFB.C – young fruiting body cap, YFB.L – young fruiting body gills, YFB.S –
young fruiting body stipe, FB.C – mature fruiting body cap (including gills), FB.S – mature
fruiting body stipe.
7108

Fig. 9. Expression heatmap of storage carbohydrate metabolism genes in *C. cinerea*. Genes are denoted by Protein IDs. Blue and red colors represent low and high expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7117
Fig. 10 Expression heatmap of chromatin remodeling related genes (histones, histone
chaperones, histone deacethylases) in *C. cinerea*. Genes are denoted by Protein IDs. Blue
and red colors represent low and high expression, respectively. Developmental stages are
abbreviated as follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1
primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap, YFB.L – young
fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting body cap
(including gills), FB.S – mature fruiting body stipe.
7125

Fig. 11. Expression heatmap of expansin-encoding genes in *C. cinerea*. Genes are denoted by Protein IDs. Blue and red colors represent low and high expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

Fig. 12. Expression heatmap of Con6 family cell surface protein-encoding genes in *C. cinerea*. Genes are denoted by Protein IDs. Blue and red colors represent low and high expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7141

7142
Fig. 13. Expression heatmap of putative wax synthase encoding genes in *C. cinerea*. Genes
are denoted by Protein IDs. Blue and red colors represent low and high expression,
respectively. Developmental stages are abbreviated as follows: VM – vegetative mycelium,
H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C – young fruiting
body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe, FB.C –
mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7149
Fig. 14. Expression heatmap of kinesin genes and members of the Dynactin complex in *C.*
cinerea. Genes are denoted by Protein IDs. Blue and red colors represent low and high
expression, respectively. Developmental stages are abbreviated as follows: VM – vegetative
mycelium, H – hyphal knot, P1 – stage 1 primordium, P2 – stage 2 primordium, YFB.C –
young fruiting body cap, YFB.L – young fruiting body gills, YFB.S – young fruiting body stipe,
FB.C – mature fruiting body cap (including gills), FB.S – mature fruiting body stipe.

7156

7157
Fig. 15: The most important, conserved genes belonging to each phase are shown - genes
with limited expression conservation or species-specific expression are not depicted.
[footnote: *pigment patterns vary by species] TP: trehalose phosphorylase, PKS: pyloketide
synthase; OSH: Oxysterol Binding Protein Homologue; AQP: aquaporins; EXP: expansins;
MCO: multicopper oxidase; RRM: RNA recognition motif family proteins; PAP2:
diacylglycerol pyrophosphate phosphatases; FAD: fatty acid desaturase; GD-PDE:
Glycerophosphodiester phosphodiesterases; S-C9-M: sphingolipid C9-methyltransferases;
IRC: intradiol ring cleavage dioxygenase; PAL: phenylalanine ammonia-lyase; SM:
secondary metabolism; HTP: heme-thiolate peroxidase; CP: cerato-platanin. genes
associated with cap autolysis are not shown on this Figure, because it is a species-specific
trait of *C. cinerea*. Note that light induction schemes are not shown on this Figure, as this
varies from species to species.
7170

Table Legends

7171

7172 Supplementary Table 1. Table summarizing ortholog protein ID for all discussed orthogroups
7173 in twelve Agaricomycetes.
7174
7175 Supplementary Table 2. Summary of read numbers and alignments in the analyzed fruiting
7176 body transcriptomes.
7177
Supplementary Table 3. Summary of developmental expression dynamics of CDE
orthogroups of genes related to alternative splicing across 12 species. Protein ID of a
representative protein is given follows by the number of species in which the orthogroup is
developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative
function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent,
1-gene present but not developmentally regulated, 2 - developmentally regulated at fold
change >2, 4- developmentally regulated at fold change >4.

7185
7186   Supplementary table 4. Copy number distributions of genes related to protein ubiquitination
7187   and the proteasome in twelve examined Agaricomycetes.
7188
7189   Supplementary table 5. Genes related to protein ubiquitination and the proteasome in *C.*
7190   *cinerea.*
7191
Supplementary Table 6. Summary of developmental expression dynamics of CDE
orthogroups of unannotated genes across 12 species. Protein ID of a representative protein
is given follows by the number of species in which the orthogroup is developmentally
regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and
ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene
present but not developmentally regulated, 2 - developmentally regulated at fold change >2,
4 - developmentally regulated at fold change >4.
7199
Supplementary Table 7. Summary of developmental expression dynamics of CDE
orthogroups of functionally poorly known genes across 12 species. Protein ID of a
representative protein is given follows by the number of species in which the orthogroup is
developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative
function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent,
1-gene present but not developmentally regulated, 2 - developmentally regulated at fold
change >2, 4- developmentally regulated at fold change >4.

Table 1. Summary of developmental expression dynamics of CDE orthogroups of cell division-related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| ID | Fold change | | Putative function | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
S.c. 2052819	6	5	DNA endonuclease involved in DNA excision repair	–	0	4	4	0	0	1	2	4	0	0	4	4
A.o. 268345	7	5	DNA recombination and repair protein Rad51-like	–	0	4	0	4	2	0	4	0	4	2	4	0
365932	11	3	DNA recombination and repair protein Rad51-like	–	2	1	4	2	2	2	2	4	2	2	2	4
S.c. 2463645	7	4	DNA repair protein Rad52/59/22	–	0	4	2	0	0	2	4	2	0	0	4	4
468197	7	4	Postreplication repair E3 ubiquitin-protein ligase RAD18	–	2	1	0	4	1	1	1	4	2	2	4	4
493516	9	6	DNA repair and recombination protein RDH54	YBR073W	4	1	1	4	1	4	4	4	4	2	2	2
439328	5	3	DNA repair protein RAD59	YDL059C	4	0	0	4	2	0	2	0	4	1	0	0
175000	10	4	DNA mismatch repair protein MSH6	YDR097C	4	2	2	4	1	1	2	4	4	2	2	2
354321	10	5	DNA repair and recombination protein RAD54	YGL163C	2	2	1	4	1	4	4	4	4	2	4	2
60974	9	6	ATP-dependent DNA helicase SRS2	YJL092W	4	1	2	4	1	1	4	4	4	4	2	2
438569	8	5	Nucleoside diphosphate kinase	YKL067W	4	1	4	4	2	1	4	2	2	1	4	1
405038	11	6	Flap endonuclease 1	YKL113C	4	2	4	2	2	2	4	4	4	1	2	4
531259	10	5	DNA repair protein RAD5	YLR032W	4	4	2	2	2	1	4	2	4	1	2	4
492739	11	7	Double-strand break repair protein MRE11	YMR224C	2	2	4	4	4	1	4	4	4	2	4	2
438045	12	5	Ribonuclease H2 subunit A	YNL072W	4	2	2	4	2	2	2	4	4	2	4	2
444936	7	3	DNA repair protein RAD50	YNL250W	2	0	4	4	1	1	2	2	1	1	4	2
541864	11	7	DNA mismatch repair protein MSH2	YOL090W	4	4	4	4	1	2	4	2	4	2	4	2
493340	8	3	Serine/threonine-protein kinase RAD53	YPL153C	4	2	1	4	1	2	2	2	4	1	2	1
360650	11	6	AP endonuclease 1	–	4	4	2	2	4	2	4	2	1	4	4	2
444409	9	5	BRCT domain containing protein	–	4	1	1	2	0	2	2	2	4	4	4	4
463086	9	5	DNA damage checkpoint protein 1	–	2	4	4	2	1	2	4	1	4	1	4	2
461486	8	2	DNA repair metallo-beta-lactamase	–	4	1	2	2	0	1	4	2	2	2	2	1
436861	8	2	DNA repair protein crb2	–	4	0	2	4	2	1	2	2	2	1	2	1
369412	8	3	Kinetochore-associated protein DSN1	–	2	2	1	4	1	0	2	2	0	2	4	4
359386	9	6	Ribonuclease H2 subunit B	–	4	2	4	0	1	1	2	4	4	2	4	4

359391	11	7	UV-damage endonuclease	–	4	2	4	4	4	4	4	2	1	2	4	2
498107	10	9	Cell division control protein 6	–	4	2	4	4	1	4	4	4	4	1	4	4
442457	10	4	Checkpoint protein hus1	–	2	1	2	2	2	1	4	4	4	2	4	2
402634	12	5	Chromatin assembly factor 1 subunit A	–	2	4	2	4	2	2	4	4	4	2	2	2
504757	10	5	Chromosome segregation in meiosis protein 3	–	2	4	2	2	2	1	1	4	4	2	4	4
451918	12	7	Kinesin	–	2	2	2	2	2	4	4	4	4	4	4	4
404203	9	7	N-acetyltransferase ECO1	–	2	4	1	4	2	1	4	4	4	1	4	4
509021	6	3	DNA helicase	–	4	1	2	4	1	1	2	0	2	1	4	1
498847	8	4	Origin recognition complex associated protein	–	2	1	4	2	1	1	4	2	4	2	4	1
465815	7	3	Polynucleotide 3'-phosphatase	–	4	1	2	2	2	1	2	1	4	1	4	1
373872	10	6	Topoisomerase 1-associated factor 1	–	4	2	1	4	2	2	4	4	4	2	1	4
358128	7	2	Replication factor A protein 1	YAR007C	4	1	1	2	1	1	2	2	2	2	4	1
435970	11	5	Serine/threonine-protein kinase KIN3	YAR018C	2	4	2	2	1	2	4	4	2	4	2	4
362687	9	3	DNA polymerase alpha subunit B	YBL035C	2	2	1	2	2	1	2	4	2	1	4	4
436862	9	5	Kinesin-like protein KIP1	YBL063W	2	4	4	2	1	1	4	4	2	1	2	4
364090	10	6	Condensin complex subunit 2	YBL097W	4	1	2	4	0	2	4	2	4	2	4	4
224847	11	7	Replication factor C subunit 5	YBR087W	4	4	4	2	1	2	4	4	4	2	4	2
374046	11	8	Proliferating cell nuclear antigen	YBR088C	4	4	4	4	2	2	4	4	4	1	4	2
439635	10	8	Pachytene checkpoint protein 2	YBR186W	4	4	1	4	2	1	4	4	4	2	4	4
439612	12	10	Deoxyuridine 5'-triphosphate nucleotidohydrolase	YBR252W	4	4	2	4	4	4	4	4	2	4	4	4
445594	11	4	Cell division control protein 7	YDL017W	4	2	2	2	2	2	2	4	4	1	4	2
445949	7	4	DNA polymerase delta catalytic subunit	YDL102W	4	1	2	2	0	1	4	4	1	1	4	2
467336	12	8	MutS protein homolog 5	YDL154W	4	4	4	4	4	2	4	2	2	2	4	4
368409	10	5	DNA ligase 1	YDL164C	2	2	2	4	2	1	4	4	4	1	4	2
356564	10	7	DDK kinase regulatory subunit DBF4	YDR052C	4	2	0	4	2	4	4	4	4	1	4	2
469713	11	6	Condensin complex subunit 3	YDR325W	1	4	2	4	2	2	4	4	2	2	4	4
425950	11	6	Ribonucleoside-diphosphate reductase large chain 1	YER070W	4	2	2	4	2	1	4	4	4	2	2	4
436798	12	8	Uncharacterized ATP-dependent helicase IRC5	YFR038W	4	2	4	4	2	4	4	4	4	2	4	2
358096	12	9	ATP-dependent DNA helicase MER3	YGL251C	2	4	4	4	2	4	2	4	4	4	4	4
521062	11	4	Minichromosome maintenance protein 10	YIL150C	2	2	2	4	2	2	4	2	2	1	4	4
411896	8	3	ATP-dependent DNA helicase MPH1	YIR002C	4	4	0	2	1	2	4	2	2	1	2	0
448596	10	4	DNA primase small subunit	YIR008C	2	4	2	2	2	1	4	4	2	1	2	4
386329	12	10	Ribonucleoside-diphosphate reductase small chain 1	YJL026W	2	4	4	4	4	4	4	4	4	2	4	4
368398	11	7	Mitotic spindle checkpoint component MAD2	YJL030W	4	4	2	2	2	2	4	4	0	4	4	4
373206	12	9	Replication factor C subunit 2	YJR068W	4	4	4	4	2	2	4	4	4	2	4	4
516390	7	3	Structural maintenance of chromosomes protein 4	YLR086W	4	0	2	2	1	2	0	4	4	0	2	1
500550	9	7	Cell division control protein 45	YLR103C	4	4	2	4	1	1	4	4	2	1	4	4

471179	12	9	G2/mitotic-specific cyclin-4	YLR210W	4	4	2	4	4	2	4	4	4	2	4	4
448884	9	6	DNA topoisomerase 3	YLR234W	4	4	1	4	1	2	4	2	4	2	4	1
399877	8	1	FACT complex subunit POB3	YML069W	2	1	2	2	2	2	2	2	1	1	4	1
360590	11	8	Chromosome transmission fidelity protein 18	YMR078C	4	4	4	2	4	2	4	4	4	1	2	4
173502	11	7	ATP-dependent helicase SGS1	YMR190C	2	4	2	4	2	2	4	4	4	1	4	4
449605	10	2	DNA polymerase epsilon catalytic subunit A	YNL262W	2	2	2	2	1	1	4	2	2	2	2	4
471912	11	6	Replication factor C subunit 3	YNL290W	4	2	4	4	2	2	4	2	4	1	4	2
498594	12	6	Thymidylate synthase	YOR074C	4	4	4	2	2	4	4	2	4	2	2	2
403599	10	3	ATP-dependent helicase ULS1	YOR191W	4	2	2	2	2	2	4	4	2	1	2	1
448855	6	3	Replication factor C subunit 1	YOR217W	4	1	2	2	1	1	1	2	4	1	4	1
356401	10	7	ATP-dependent DNA helicase CHL1	YPL008W	4	4	2	4	2	1	4	4	4	2	1	4
498104	11	7	DNA polymerase alpha-binding protein	YPR135W	4	4	2	1	2	2	4	4	4	2	4	4
467729	12	0	DNA polymerase epsilon subunit B	YPR175W	2	2	2	2	2	2	2	2	2	2	2	2
448702	7	2	ATP-dependent DNA ligase	–	2	2	4	4	1	0	2	1	0	1	2	2
537653	10	3	E3 ubiquitin-protein ligase substrate receptor MMS22	–	2	4	4	2	0	2	2	4	2	1	2	2
49522	10	5	Mitochondrial morphogenesis protein SLD7	–	4	1	2	2	2	4	2	1	4	4	4	2
292223	11	5	Alpha-actinin-binding protein	–	4	4	2	4	2	2	4	2	4	1	2	2
426459	9	2	Chromosome transmission fidelity protein 8	–	4	2	2	2	2	1	1	4	2	1	2	2
367776	10	5	DNA helicase	–	4	4	2	2	0	2	4	4	4	1	2	2
374638	10	5	DNA helicase	–	4	4	2	4	2	2	2	1	4	0	4	2
447030	10	6	DNA helicase	–	4	4	2	2	4	1	4	2	1	2	4	4
460017	10	5	DNA polymerase delta subunit 3	–	4	2	2	4	0	2	4	2	4	2	4	1
464859	12	10	DNA replication associated protein	–	4	4	2	4	2	4	4	4	4	4	4	4
447452	10	6	DNA replication factor	–	2	2	2	4	1	2	4	4	4	1	4	4
471732	11	8	Formin	–	4	4	4	2	2	4	4	4	2	1	4	4
365191	11	5	Replication factor A protein 3	–	2	2	1	2	2	4	4	4	4	2	4	2
464852	9	4	Telomere binding protein	–	4	4	0	4	2	1	2	2	2	1	2	4
539797	10	7	TPX2 domain-containing protein	–	4	4	2	1	4	2	2	4	4	0	4	4
394344	10	1	Cohesin subunit SCC3	–	2	2	2	2	2	1	2	2	2	1	4	2
442128	7	1	Sister chromatid cohesion protein 1	–	4	1	2	2	1	1	2	2	2	1	2	1
542870	8	3	Structural maintenance of chromosomes protein 1	YFL008W	4	0	0	2	1	1	4	2	2	2	4	2
406915	11	2	Structural maintenance of chromosomes protein 3	YJL074C	2	2	2	2	1	2	2	4	2	2	4	2
445120	10	1	DNA replication complex GINS protein SLD5	–	4	2	1	2	2	2	2	2	2	2	2	1
380189	8	3	DNA replication complex GINS protein PSF1	YDR013W	4	4	2	2	0	2	1	4	1	2	2	1
421683	10	4	DNA replication complex GINS protein PSF2	YJL072C	4	2	2	2	1	1	2	4	4	2	2	4
439626	9	5	DNA replication complex GINS protein PSF3	YOL146W	4	2	2	2	1	1	4	1	4	2	4	4
193478	9	3	DNA replication licensing factor MCM2	YBL023C	2	2	1	2	2	1	4	4	2	1	2	4

ID			Protein	ORF												
435526	11	5	DNA replication licensing factor MCM7	YBR202W	2	2	4	2	2	1	4	4	2	2	4	4
464525	9	5	DNA replication licensing factor MCM3	YEL032W	2	4	2	2	2	1	4	4	1	1	4	4
421772	10	4	DNA replication licensing factor MCM6	YGL201C	2	2	1	4	1	2	4	4	2	2	4	2
401934	10	6	Minichromosome maintenance protein 5	YLR274W	4	4	1	2	2	2	4	4	2	1	4	4
402513	9	3	DNA replication licensing factor MCM4	YPR019W	4	2	0	4	2	1	2	4	2	1	2	2
450001	11	5	Kinetochore protein NUF2	–	4	2	2	4	2	2	4	4	2	1	4	2
393128	11	2	Kinetochore protein NDC80	YIL144W	2	2	2	4	2	1	2	4	2	2	2	2
167797	10	2	Kinetochore protein SPC24	–	2	4	2	2	2	1	2	2	4	1	2	2
453581	9	6	Kinetochore protein SPC25	–	4	2	2	4	1	1	4	4	4	1	4	2
366401	12	5	E3 SUMO-protein ligase MMS21	–	2	4	2	2	2	2	4	4	2	2	4	2
492711	10	6	Non-structural maintenance of chromosomes element 1	–	4	4	0	4	2	1	2	2	4	2	4	4
412693	9	2	Non-structural maintenance of chromosome element 4	YDL105W	2	2	2	0	2	1	2	4	2	1	4	2
25048	8	5	Structural maintenance of chromosomes protein 5	YOL034W	4	2	0	4	1	2	1	2	4	1	4	4
499664	10	4	Non-structural maintenance of chromosome element 3	–	2	2	1	4	2	4	2	2	4	2	4	1
446459	9	6	Meiotic nuclear division protein 1	–	4	1	2	4	0	2	4	2	4	1	4	4
A.o. 269495	7	7	Meiotic recombination protein DMC1	–	0	4	0	4	4	4	4	0	4	4	0	0
468689	10	8	DNA helicase	–	4	2	0	4	4	4	4	0	4	2	4	4
371359	12	12	PHO85 cyclin-2	YDL127W	4	4	4	4	4	4	4	4	4	4	4	4
502170	12	11	MutS protein homolog 4	YFL003C	4	4	2	4	4	4	4	4	4	4	4	4
238535	12	10	Homologous-pairing protein 2	YGL033W	2	4	2	4	4	4	4	4	4	4	4	4
368527	10	7	Protein NAM8	YHR086W	4	2	1	4	4	4	4	2	4	2	4	1
479000	11	8	Meiotic sister chromatid recombination protein 1	YML128C	4	2	2	4	4	4	4	2	4	1	4	4
374758	9	4	Spindle assembly checkpoint kinase	YPL209C	2	2	4	4	1	2	4	4	1	2	2	1
S.c. 85873	9	7	Nuclear fusion protein KAR5	–	0	4	2	4	1	2	4	4	0	4	4	4
474563	6	4	Holliday junction resolvase	–	4	0	0	4	2	2	0	1	0	4	4	0
469496	6	2	Meiosis specific protein	–	4	1	2	1	2	1	1	2	4	1	2	1
378537	12	12	Meiotic coiled-coil protein	–	4	4	4	4	4	4	4	4	4	4	4	4
185247	7	0	Actin-like protein ARP9	–	2	1	2	2	0	1	2	2	1	2	2	1
447320	10	7	Altered inheritance of mitochondria protein 44	–	4	4	2	4	1	2	1	4	4	2	4	4
465387	8	4	Checkpoint protein RAD24	–	4	2	1	4	0	2	2	4	4	1	2	1
S.c. 2570476	4	1	DNA polymerase IV	–	2	1	0	2	2	1	1	0	0	1	4	1
461421	8	2	G1/S-specific cyclin CLN3	–	1	2	2	4	1	1	2	2	4	1	2	2
404166	11	7	Holliday junction resolvase YEN1	–	4	4	2	2	4	2	2	4	4	4	4	1
182474	10	7	M-phase inducer phosphatase	–	4	4	2	2	1	2	4	4	4	1	4	4
215276	11	6	Mediator of replication checkpoint protein 1	–	4	2	2	4	4	2	2	4	4	2	4	0
472431	9	7	Meiosis-specific protein SPO11	–	4	4	1	4	2	4	4	1	0	2	4	4

ID			Description	S.c.												
461802	11	2	Protein SFI1	–	2	2	2	4	1	2	2	2	2	2	4	2
269110	5	4	DNA helicase	–	4	4	0	4	0	0	4	0	0	0	0	2
365598	11	5	Serine/threonine-protein kinase Haspin homolog ALK1	–	2	4	2	2	2	1	4	4	4	2	2	4
466770	10	4	Serine/threonine-protein kinase MPS1	–	4	2	2	2	2	2	4	4	2	1	4	1
469725	10	4	Spindle assembly checkpoint component MAD3	–	2	4	2	2	2	1	4	4	2	2	4	1
S.c. 2503158	8	7	Uracil-DNA glycosylase	–	0	4	4	4	2	4	4	0	4	0	0	4
442763	9	3	Kinetochore-associated protein MTW1	YAL034W-A	4	1	2	1	2	1	2	4	4	2	2	2
476052	8	3	Origin recognition complex subunit 2	YBR060C	4	1	1	2	2	2	4	4	1	2	2	1
377937	7	5	Anaphase-promoting complex subunit 11	YDL008W	4	1	4	4	0	2	4	1	2	1	4	1
518718	5	0	General transcription and DNA repair factor IIH subunit TFB1	YDR311W	2	1	2	1	2	1	1	1	1	1	2	2
479908	8	2	Tyrosine-protein phosphatase CDC14	YFR028C	2	2	2	4	1	1	2	1	2	1	4	2
394368	9	5	Structural maintenance of chromosomes protein 2	YFR031C	4	0	2	2	1	2	4	4	4	1	2	4
74935	8	5	Spindle pole body component SPC105	YGL093W	4	2	2	4	2	1	0	4	4	1	4	1
545958	12	7	APC/C activator protein CDC20	YGL116W	2	4	2	2	2	4	4	4	4	4	4	2
538972	9	3	Checkpoint serine/threonine-protein kinase BUB1	YGR188C	2	4	2	2	1	1	4	4	1	2	2	2
439189	10	5	Cell division control protein 6	YJL194W	2	4	2	4	2	1	2	4	4	2	4	1
497872	5	0	Anaphase-promoting complex subunit CDC16	YKL022C	2	2	1	2	1	1	2	1	2	1	1	1
354027	9	3	DNA primase large subunit	YKL045W	2	4	1	2	1	1	2	4	4	2	2	2
402736	5	1	Inner kinetochore subunit MIF2	YKL089W	1	1	1	2	2	1	1	1	4	1	2	2
439911	8	4	Condensin complex subunit 1	YLR272C	4	1	2	2	1	1	4	4	2	2	4	1
360249	8	6	Uracil-DNA glycosylase	YML021C	4	1	4	2	4	1	4	2	4	0	4	1
403152	10	6	Cell cycle serine/threonine-protein kinase CDC5/MSD2	YMR001C	4	2	1	2	1	2	4	4	4	4	4	2
449882	11	4	DNA topoisomerase 2	YNL088W	2	2	4	2	2	1	4	4	2	1	4	2
441663	9	5	DNA polymerase alpha catalytic subunit A	YNL102W	4	4	1	2	0	1	4	4	2	2	4	2
439045	11	8	PHO85 cyclin-1	YNL289W	4	4	4	4	4	4	4	2	1	2	4	2
375607	11	6	Cell division control protein 31	YOR257W	4	2	2	2	1	4	4	4	4	2	4	2
533614	12	11	G2/mitotic-specific cyclin-2	YPR119W	4	4	2	4	4	4	4	4	4	4	4	4
380995	10	8	Kinesin-like protein KAR3	YPR141C	4	2	4	4	0	4	4	4	4	2	4	1
438441	10	6	Anaphase spindle elongation protein 1	–	2	2	1	2	0	4	4	4	4	2	4	4
114016	9	6	Chromo/chromo shadow domain	–	4	2	4	4	2	1	2	1	4	1	4	4
522601	7	3	DASH complex subunit Dam1	–	4	1	1	4	2	2	2	4	2	1	0	1
472683	7	3	DNA replication regulator SLD2	–	4	1	0	2	1	2	4	2	0	1	4	2
380283	7	4	DNA replication regulator SLD3	–	2	1	2	4	0	2	4	1	4	1	4	0
463767	7	1	Dynactin subunit 3	–	2	1	1	2	2	2	0	2	4	1	2	1
466211	10	5	Kinetochore protein	–	2	1	2	2	2	2	4	4	4	1	4	4
468013	3	2	Mitotic-spindle organizing protein 1	–	4	1	0	2	0	1	1	0	0	1	4	1

ID	>2	>4	Putatively function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
537412	9	1	Nuclear rim protein 1	–	2	2	2	1	1	1	2	2	2	2	4	2
431070	12	10	Anaphase-promoting complex subunit 4	–	4	4	2	4	4	4	4	4	4	2	4	4
365564	9	7	CCCh-type zinc finger protein	–	4	4	2	4	4	4	4	0	0	0	4	2
540942	8	1	DNA damage response (DDR) and DNA repair protein	–	4	1	2	2	1	1	2	2	2	2	2	1
A.o. 259630	4	3	DNA helicase	–	0	0	0	4	0	0	4	0	0	2	4	0
446333	9	4	DNA helicase	–	2	4	0	4	4	1	4	2	0	2	2	2
450108	7	0	DNA helicase	–	2	1	2	2	1	1	2	2	1	2	2	1
460394	12	10	Rad21/Rec8-like protein	–	2	4	2	4	4	4	4	4	4	4	4	4
A.o. 254302	8	5	Ribonuclease H2; subunit C	–	0	4	0	2	2	1	4	4	0	2	4	4
374534	9	1	Separase	–	2	1	1	2	2	1	4	2	2	2	2	2
496196	5	4	SET domain protein	–	4	1	1	4	4	1	1	1	4	1	2	0
366767	10	5	RecQ genome instability protein	–	4	2	1	4	2	1	4	2	2	4	4	2
493093	12	6	Sporulation-specific protein 12	–	2	4	4	2	2	2	4	4	4	2	4	2
443396	10	3	Ku80 protein	YMR106C	4	2	2	1	2	1	4	2	4	2	2	2
443149	7	3	Ku70 protein	YMR284W	4	1	4	2	1	1	2	2	2	1	4	1

Table 2. Summary of developmental expression dynamics of CDE orthogroups of ribosomal genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | S. cerevisiae ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
423325	10	3	60S ribosomal protein L28e	–	4	2	2	2	2	4	2	2	4	1	2	1
405648	10	2	60S ribosomal protein L19-B	YBL027W	4	2	2	2	2	4	2	2	2	1	2	1
412936	10	2	40S ribosomal protein S8-A	YBL072C	4	2	2	2	1	2	2	2	2	2	4	1
478794	11	3	60S ribosomal protein L23-A	YBL087C	4	2	2	2	2	4	2	2	4	2	2	1
366589	9	5	60S ribosomal protein L4-A	YBR031W	4	2	4	2	0	4	4	2	4	1	2	1
497971	7	5	40S ribosomal protein S11-B	YBR048W	4	2	4	0	2	4	0	1	4	1	4	1
464129	8	4	40S ribosomal protein S6-B	YBR181C	4	2	4	2	1	4	2	1	4	1	2	1
430785	8	2	60S ribosomal protein L21-A	YBR191W	4	2	2	2	2	1	4	1	0	2	2	1
467322	9	5	40S ribosomal protein S14-A	YCR031C	4	2	4	2	2	4	2	1	4	1	4	1
493413	9	1	Putatively involved intra-nuclear transport of pre-60S ribosomal subunits	YCR072C	4	2	2	2	1	2	2	2	2	1	2	1
403137	11	4	60S ribosomal protein L31-A	YDL075W	4	2	4	2	2	4	2	2	4	2	2	1
398174	9	3	40S ribosomal protein S16-B	YDL083C	4	2	2	2	2	2	2	1	4	1	4	1
412254	8	2	60S ribosomal protein L35-B	YDL136W	4	2	2	2	2	4	2	1	0	1	2	1
493053	10	4	Putatively involved in	YDL208W	4	2	2	4	2	2	2	2	4	1	4	1

pseudouridylation of rRNA

64051	9	4	40S ribosomal protein S13	YDR064W	4	2	4	2	1	4	2	2	2	1	4	1
381793	10	4	60S acidic ribosomal protein P2-beta	YDR382W	4	2	2	2	2	4	2	2	4	1	4	1
459549	8	2	60S ribosomal protein L12-B	YDR418W	4	2	2	2	1	4	2	1	2	1	2	0
404670	11	4	40S ribosomal protein S17-B	YDR447C	4	2	4	2	2	4	2	1	4	2	2	2
442748	8	2	40S ribosomal protein S18-A	YDR450W	4	2	2	2	0	4	2	1	0	1	2	2
499692	10	5	60S ribosomal protein L27-B	YDR471W	4	2	2	2	2	4	4	1	4	2	4	1
429262	9	4	60S ribosomal protein L34-A	YER056C-A	4	2	2	2	2	4	4	1	0	2	4	1
439980	9	3	40S ribosomal protein S24-A	YER074W	4	0	0	2	2	4	2	2	2	1	4	2
329287	10	5	40S ribosomal protein S26-B	YER131W	4	2	4	2	2	4	2	2	4	1	4	1
426937	9	2	60S ribosomal protein L2-A	YFR031C-A	4	2	2	2	2	4	2	1	0	2	2	1
S.c. 2742667	10	4	60S ribosomal protein L7A	YGL076C	0	2	4	2	2	4	4	2	4	2	2	1
441325	10	3	60S ribosomal protein L28	YGL103W	4	2	2	2	2	4	2	1	4	2	2	1
475518	10	4	40S ribosomal protein S2	YGL123W	4	2	4	2	2	4	2	2	4	1	2	1
437986	9	4	60S ribosomal protein L1-B	YGL135W	4	2	2	2	2	4	2	1	4	1	4	1
480276	8	3	40S ribosomal protein S25-A	YGR027C	4	2	4	2	2	4	2	1	0	1	2	1
475978	10	3	60S ribosomal protein L26-B	YGR034W	4	2	2	0	2	4	4	2	2	2	2	1
130885	9	3	60S ribosomal protein L11-B	YGR085C	4	2	4	2	2	4	2	2	0	1	2	1
495378	5	0	40S ribosomal protein S23-A	YGR118W	2	1	2	2	1	2	1	1	0	2	1	1
379934	9	3	60S ribosomal protein L24-B	YGR148C	4	2	2	2	2	4	2	0	4	1	2	1
494449	10	4	40S ribosomal protein S0-A	YGR214W	4	2	4	2	2	4	2	2	4	1	2	1
467085	10	3	60S ribosomal protein L14-B	YHL001W	4	2	2	2	2	4	2	2	4	1	2	1
165767	9	3	40S ribosomal protein S20	YHL015W	4	2	2	2	2	4	2	2	0	1	4	1
438384	6	1	40S ribosomal protein S27-B	YHR021C	4	1	1	2	0	2	2	1	2	1	2	1
214528	6	6	Protein involved in rRNA-processing	YHR087W	4	1	4	4	1	1	4	4	1	1	4	0
441187	9	4	60S ribosomal protein L42-B	YHR141C	4	2	1	2	2	4	2	1	4	2	4	1
441652	7	3	40S ribosomal protein S4-B	YHR203C	4	2	0	2	0	4	2	1	4	1	2	1
497565	8	2	60S ribosomal protein L39	YJL189W	4	1	2	2	0	4	2	2	0	2	2	1
375577	9	5	40S ribosomal protein S22-A	YJL190C	4	2	4	2	2	4	4	1	4	1	2	1
385517	8	4	60S ribosomal protein L43-B	YJR094W-A	4	2	0	0	0	4	2	2	4	2	4	1
446338	10	3	40S ribosomal protein S5	YJR123W	4	2	4	2	2	4	2	1	2	2	2	1
472703	8	2	60S ribosomal protein L8-B	YLL045C	2	2	4	2	2	2	2	1	0	1	4	1
374411	7	2	60S ribosomal protein L22-A	YLR061W	2	2	4	2	2	1	2	1	0	1	4	1
471911	7	4	Ubiquitin-40S ribosomal protein S31	YLR167W	4	0	4	2	1	4	2	1	4	1	2	1
447700	7	3	60S ribosomal protein L37-A	YLR185W	4	0	2	2	0	4	2	1	4	1	2	1
353298	7	2	40S ribosomal protein S28-B	YLR264W	4	2	2	2	0	4	2	1	0	2	0	0
398047	7	3	60S ribosomal protein L38	YLR325C	4	2	2	0	0	4	2	1	0	2	4	1
367451	7	3	60S acidic ribosomal protein P0	YLR340W	4	2	1	2	1	2	4	1	2	1	4	1
398594	9	3	40S ribosomal protein S29-A	YLR388W	4	2	2	2	0	4	2	2	0	2	4	1
385868	10	4	60S ribosomal protein L6-B	YLR448W	4	2	4	2	2	4	2	2	0	2	4	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
474164	9	5	40S ribosomal protein S1-B	YML063W	4	2	4	2	2	4	4	1	2	1	4	1
424304	8	4	60S ribosomal protein L13-B	YMR142C	4	2	0	2	2	4	0	4	2	1	4	1
456295	8	2	60S ribosomal protein L36-A	YMR194W	4	2	2	2	0	4	2	2	0	1	2	1
375811	8	3	60S ribosomal protein L20-A	YMR242C	4	2	4		0	4	2	1	2	1	2	1
412005	9	3	60S ribosomal protein L9-B	YNL067W	4	2	2	2	2	4	2	1	2	1	4	1
416494	5	2	40S ribosomal protein S3	YNL178W	4	2	0	0	1	4	2	1	2	1	0	1
438977	9	0	Ribosome-bound, Hsp70-type chaperone	YNL209W	2	2	2	2	2	1	2	2	2	1	2	1
462840	8	4	60S ribosomal protein L18-B	YNL301C	4	2	4	2	1	4	2	1	2	1	4	1
21836	6	0	Putatively involved in nucleolar processing of pre-18S ribosomal RNA	YNL308C	2	1	2	2	2	1	1	1	2	1	2	1
381804	8	2	40S ribosomal protein S15	YOL040C	4	2	2	2	0	4	2	1	2	1	2	1
465651	10	4	40S ribosomal protein S19-A	YOL121C	4	2	4	2	2	4	2	2	4	1	2	1
426850	7	4	60S ribosomal protein L3	YOR063W	4	2	2	0	2	4	0	1	4	1	4	1
401275	9	6	40S ribosomal protein S12	YOR369C	4	2	1	4	0	4	4	2	4	2	4	1
430789	10	4	40S ribosomal protein S9-A	YPL081W	4	2	4	2	2	4	2	2	4	1	2	1
467768	8	5	60S ribosomal protein L5	YPL131W	4	2	4	2	2	4	4	1	0	1	4	1
356089	10	4	60S ribosomal protein L33-A	YPL143W	4	2	2	0	2	4	2	2	4	2	4	1
539347	5	3	60S ribosomal protein L29	YFR032C-A	4	2	0	0	0	4	4	1	0	2	0	1
394207	7	3	60S ribosomal protein L30	YGL030W	4	2	0	2	0	4	2	1	4	1	2	1
539473	7	2	60S ribosomal protein L16-A	YIL133C	4	2	2	0	2	4	2	1	0	1	2	1

Table 3. Summary of developmental expression dynamics of CDE orthogroups of genes related to Acetyl-CoA production and basic metabolism across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
500663	9	7	Aldehyde dehydrogenase 5, mitochondrial	YER073W	4	4	4	4	1	1	2	4	2	1	4	4
500719	11	4	Acetyl-coenzyme A synthetase 2	YLR153C	2	2	4	4	1	2	2	4	2	2	2	4
273601	11	7	Alcohol dehydrogenase 3, mitochondrial	YMR083W	4	2	4	4	4	4	2	4	2	1	4	2
421282	7	4	Aldehyde dehydrogenase	YMR170C	4	0	2	4	1	2	4	0	4	1	2	0
450047	8	5	Carbonic anhydrase	–	4	2	4	2	1	0	1	2	4	1	4	4
355051	9	5	Ceramide synthase	–	4	2	4	2	2	1	4	2	1		4	1
82352	9	6	Uncharacterized TLC domain-containing protein YPR114W	YPR114W	4	1	2	4	4	2	1	4	4	1	4	2
503618	9	3	3-methyl-2-oxobutanoate hydroxymethyltransferase	YBR176W	4	2	2	2	4	2	2	2	0	1	4	1
443766	8	1	Dephospho-CoA kinase CAB5	YDR196C	2	2	1	0	2	2	2	1	4	2	2	1

474787	6	1	Pantothenate kinase CAB1	YDR531W	4	1	2	0	0	2	2	1	1	1	2	2
461914	6	2	Phosphopantetheine adenylyltransferase	YGR277C	1	1	0	2	4	0	2	1	2	1	4	2
282840	5	2	2-dehydropantoate 2-reductase	YHR063C	4	1	1	2	2	1	1	1	2	1	4	1
500145	6	3	Pantoate--beta-alanine ligase	YIL145C	4	1	0	1	4	1	2	1	2	1	2	4
374749	8	1	Coenzyme A biosynthesis protein 3	YKL088W	2	2	4	2	1	2	2	2	0	1	2	1
453548	8	4	Pyruvate kinase 1	YAL038W	4	2	4	4	1	1	2	2	1	1	4	2
499160	9	5	Glucose-6-phosphate isomerase	YBR196C	4	1	4	2	2	4	2	2	4	1	4	1
365495	5	2	Glucokinase-1	YCL040W	4	2	1	1	2	2	1	1	1	1	4	1
442066	7	4	Phosphoglycerate kinase	YCR012W	4	1	2	4	1	1	2	4	2	1	4	1
173684	9	5	Triosephosphate isomerase	YDR050C	4	1	4	2	2	2	4	4	2	1	4	1
402119	8	3	Hexokinase-2	YGL253W	2	2	4	4	1	2	2	1	2	1	4	1
245737	9	4	Enolase 1	YGR254W	4	1	4	2	2	2	2	4	2	1	4	1
264276	7	5	Glyceraldehyde-3-phosphate dehydrogenase 1	YJL052W	4	1	4	2	0	2	4	4	1	1	4	1
546961	8	4	Fructose-bisphosphate aldolase	YKL060C	4	1	4	2	2	1	4	2	2	1	4	1
422402	11	8	Pyruvate decarboxylase isozyme 1	YLR044C	4	2	4	4	4	2	4	4	4	1	4	2
436876	9	6	Fructose-1,6-bisphosphatase	YLR377C	4	1	4	4	2	2	2	4	0	1	4	4
462738	8	1	ATP-dependent 6-phosphofructokinase subunit beta	YMR205C	2	1	4	2	2	1	2	2	2	1	2	1
495730	12	7	Isocitrate lyase	YER065C	2	4	2	4	4	2	2	4	4	4	2	4
530292	12	11	Malate synthase 2	YIR031C	4	4	4	4	4	2	4	4	4	4	4	4
501208	6	0	Homocitrate dehydratase, mitochondrial	YJL200C	2	2	2	0	2	1	2	2	1	1	0	1
374148	12	7	Succinate/fumarate mitochondrial transporter	YJR095W	4	4	2	4	4	2	2	4	2	2	2	4
493553	10	5	Phosphoenolpyruvate carboxykinase	YKR097W	4	2	0	4	2	2	1	4	2	2	4	4
449845	9	3	Malate dehydrogenase	–	1	1	4	2	2	2	2	4	2	1	4	2
442593	10	4	ATP citrate synthase	–	4	2	2	4	2	4	2	2	2	1	4	1
443703	10	5	Mitochondrial carnitine O-acetyltransferase	YAR035W	4	1	4	4	2	2	1	4	4	2	2	2
442787	7	5	Acetyl-CoA hydrolase	YBL015W	4	4	0	4	1	4	2	0	0	0	2	4
441597	7	6	Accumulation of dyads protein 2	YCR010C	4	1	4	4	0	4	4	1	2	1	4	1
373141	10	8	Aquaglycerol porin AQY3	YFL054C	4	4	4	4	2	2	4	0	4	0	4	4
357888	10	2	Mitochondrial oxaloacetate transport protein	YKL120W	2	1	2	2	4	2	2	1	2	2	4	2
375404	11	6	Carboxylic acid transporter protein homolog	YKL217W	4	2	4	4	2	4	2	2	4	1	4	2
147126	9	3	Protein SYM1	YLR251W	2	2	4	2	1	1	4	4	2	1	2	2
470219	9	3	Carnitine O-acetyltransferase, mitochondrial	YML042W	2	1	0	2	4	4	2	2	1	2	4	2
474886	9	3	Citrate/oxoglutarate carrier protein	YMR241W	4	1	4	2	0	2	2	2	2	1	4	2
543155	7	6	Acetate kinase	–	4	2	0	4	4	0	0	4	0	1	4	4
534888	9	5	Malate transporter	–	2	4	4	4	2	0	2	2	0	1	4	4
S.c. 2738153	8	5	Formate transporter	–	0	1	4	4	4	0	4	2	2	1	4	2
474928	8	5	Oxalate-CoA ligase	–	4	0	2	4	2	1	4	4	2	1	4	1
452378	9	7	Formate dehydrogenase 1	YOR388C	4	4	1	4	4	0	4	2	4	1	4	2

Protein ID	FC>2	FC>4	Putative function	S.c. ortholog												
361522 S.c.	10	8	Oxalate decarboxylase	–	4	4	4	2	2	4	4	4	1	1	4	4
2751877	6	5	Oxalate decarboxylase	–	0	4	4	0	0	0	0	2	0	4	4	4
499734	11	11	Oxaloacetase	–	4	4	4	4	4	0	4	4	4	4	4	4
385900	11	2	Pyruvate dehydrogenase E1 component subunit beta, mitochondrial	YBR221C	4	2	2	2	2	2	2	2	2	2	4	1
501813	9	3	Pyruvate dehydrogenase E1 component subunit alpha, mitochondrial	YER178W	4	1	4	2	4	2	2	2	2	1	2	1
440087	10	3	Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex, mitochondrial	YNL071W	4	2	4	2	2	2	2	2	2	1	4	1
273609	9	6	NADP-dependent malic enzyme	–	4	4	4	4	2	2	2	0	4	0	4	1
364080	7	2	Pyruvate carboxylase 2	YBR218C	4	1	2	2	1	2	2	2	1	1	4	1
369183	10	1	Dihydrolipoyllysine-residue succinyltransferase, mitochondrial	YDR148C	2	2	2	2	1	2	2	2	2	1	4	2
492830	8	1	Dihydrolipoyl dehydrogenase, mitochondrial	YFL018C	2	2	2	2	1	2	2	2	1	1	4	1
454395	10	3	Succinate--CoA ligase subunit beta, mitochondrial	YGR244C	4	2	2	1	2	2	2	2	2	1	4	4
354219	5	2	Protein phosphatase 2C homolog 7, mitochondrial	YHR076W	1	1	2	1	2	1	1	1	2	1	4	4
545855	7	1	2-oxoglutarate dehydrogenase, mitochondrial	YIL125W	2	2	1	2	0	2	1	1	2	1	2	4
380777	7	6	NAD-dependent malic enzyme, mitochondrial	YKL029C	4	1	4	4	4	4	2	1	1	1	4	1
409635	11	3	Malate dehydrogenase, mitochondrial	YKL085W	2	2	4	2	2	4	2	2	2	2	4	1
483615	6	0	Succinate dehydrogenase cytochrome b subunit, mitochondrial	YKL141W	1	1	2	0	1	2	2	1	2	2	2	1
400441	6	0	Succinate dehydrogenase flavoprotein subunit, mitochondrial	YKL148C	2	2	2	2	1	1	1	1	1	2	2	1
363708	8	0	Succinate dehydrogenase iron-sulfur subunit, mitochondrial	YLL041C	2	1	2	2	2	2	2	1	2	1	2	1
367547	9	0	Mitochondrial inner membrane protein SHH4	YLR164W	2	1	2	2	1	2	2	2	2	2	2	1
503237	6	0	Aconitate hydratase, mitochondrial	YLR304C	2	2	2	2	0	1	1	2	2	1	1	1
474027	8	2	Isocitrate dehydrogenase subunit 1, mitochondrial	YNL037C	2	2	4	2	4	1	2	1	1	1	2	2
497099	11	4	Citrate synthase, mitochondrial	YNR001C	2	2	4	4	2	2	2	2	4	1	4	2
449179	6	1	Succinate dehydrogenase assembly factor 2, mitochondrial	YOL071W	2	1	0	1	1	2	2	4	0	2	2	1
444503	9	4	Isocitrate dehydrogenase subunit 2, mitochondrial	YOR136W	4	2	4	2	4	2	2	2	0	1	4	1
360292	10	3	Succinate--CoA ligase subunit alpha, mitochondrial	YOR142W	4	2	2	2	2	2	1	2	2	1	4	4
464081	6	2	Fumarate hydratase, mitochondrial	YPL262W	1	1	2	2	2	1	4	2	1	1	4	1

Table 4. Summary of developmental expression dynamics of CDE orthogroups of fatty acid beta-oxidation related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also

given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Species Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
437115	7	4	General amino acid permease AGP2	YBR132C	4	2	4	1	4	2	4	0	1	0	1	2
336239	6	1	Acyl-coenzyme A oxidase	YGL205W	1	1	1	2	2	1	2	2	1	1	4	2
438072	10	5	Acyl-CoA-binding protein	YGR037C	4	2	4	4	2	2	2	4	2	1	4	1
365907	6	2	3-ketoacyl-CoA thiolase, peroxisomal	YIL160C	4	1	2	2	4	2	0	1	2	1	1	1
231560	5	1	Peroxisomal hydratase-dehydrogenase-epimerase	YKR009C	2	1	1	4	2	1	2	1	2	1	1	1
354296	6	2	Aspartate aminotransferase	YLR027C	2	2	4	0	0	2	1	1	1	1	4	2
433737	8	3	Peroxisomal 2,4-dienoyl-CoA reductase SPS19 [(3E)-enoyl-CoA-producing]	YNL202W	4	1	4	2	2	4	1	1	2	1	2	2
374386	7	1	Peroxisomal membrane protein PMP27	YOL147C	2	1	2	2	2	2	1	1	2	1	4	1
436931	6	2	Delta(3,5)-Delta(2,4)-dienoyl-CoA isomerase	YOR180C	2	1	2	1	0	4	1	2	0	1	4	2
401814	5	2	Peroxisomal long-chain fatty acid import protein 2	YPL147W	4	1	1	2	4	1	1	1	2	1	2	1
395825	8	3	Peroxisomal adenine nucleotide transporter 1	YPR128C	4	1	2	2	1	4	2	2	2	1	4	1
437261	6	4	Long-chain-fatty-acid--CoA ligase 1	YOR317W	2	1	4	4	2	1	4	1	0	1	4	1

Table 5. Summary of developmental expression dynamics of CDE orthogroups of ergosterol and sphingolipid biosynthesis related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Species Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
488426	9	3	Transcription factor	–	4	1	0	4	2	2	4	2	2	2	2	1
Late pathway																
366787	9	4	Ergosterol biosynthetic protein 28	YER044C	2	1	2	2	4	2	4	4	2	1	4	1
366442	11	6	Sterol-4-alpha-carboxylate 3-dehydrogenase, decarboxylating	YGL001C	4	2	4	4	2	4	2	4	2	2	4	1
544612	10	7	Delta(24(24(1)))-sterol reductase	YGL012W	4	2	4	2	0	4	4	4	2	4	4	1
496056	10	7	Methylsterol monooxygenase	YGR060W	4	1	4	2	4	4	2	4	2	4	4	1
489517	9	6	Squalene monooxygenase	YGR175C	4	1	2	2	0	4	4	4	4	2	4	1

Protein ID	FC>2	FC>4	Function	S. cerevisiae	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
397292	11	7	Lanosterol 14-alpha demethylase	YHR007C	4	2	4	2	2	4	4	4	2	4	4	1
413057	6	3	NADPH--cytochrome P450 reductase	YHR042W	4	1	4	2	4	1	1	2	1	1	2	1
420341	8	5	Lanosterol synthase	YHR072W	4	2	4	2	0	1	4	4	1	2	4	1
494314	7	4	Delta(7)-sterol 5(6)-desaturase	YLR056W	4	2	4	1	2	2	4	1	1	4	1	1
462226	8	4	3-keto-steroid reductase	YLR100W	4	1	4	2	0	1	4	2	2	1	4	2
373683	10	7	Sterol 24-C-methyltransferase	YML008C	4	1	4	2	4	4	2	4	4	4	2	1
357271	10	6	Cytochrome P450 61	YMR015C	4	0	4	2	2	4	2	4	2	4	4	1
17387	11	7	C-8 sterol isomerase	YMR202W	4	2	4	2	2	4	2	4	4	4	4	1
535364	10	3	Delta(14)-sterol reductase	YNL280C	4	2	2	2	1	2	4	2	2	2	4	1

Mevalonate pathway

Protein ID	FC>2	FC>4	Function	S. cerevisiae	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
358591	7	2	Cystathionine beta-lyase	YGL184C	4	4	1	1	1	1	2	2	2	2	2	1
43445	9	6	Squalene synthase	YHR190W	4	2	4	4	0	4	2	4	2	1	4	1
459666	9	5	Farnesyl pyrophosphate synthase	YJL167W	4	2	2	1	2	2	4	4	4	1	4	1
360799	8	2	3-hydroxy-3-methylglutaryl-coenzyme A reductase 2	YLR450W	4	1	2	2	2	2	1	2	1	2	4	1
439474	12	5	Hydroxymethylglutaryl-CoA synthase	YML126C	4	2	4	2	4	4	2	4	2	2	2	2
355838	12	10	Phosphomevalonate kinase	YMR220W	4	4	4	4	4	4	4	4	2	4	4	2
438368	8	2	Diphosphomevalonate decarboxylase	YNR043W	2	2	2	1	2	1	2	2	1	4	4	1
355558	7	0	Acetyl-CoA acetyltransferase	YPL028W	2	2	1	2	1	2	1	2	1	2	2	1
460395	6	2	Isopentenyl-diphosphate Delta-isomerase	YPL117C	4	1	1	2	1	1	2	4	1	2	2	1
539004	12	7	3-oxo-5-alpha-steroid 4-dehydrogenase	–	4	4	2	4	2	4	4	2	2	2	4	4

Sphingolipid biosynthesis

Protein ID	FC>2	FC>4	Function	S. cerevisiae	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
388282	9	5	Alkaline ceramidase YPC1	YBR183W	4	2	4	4	2	2	1	4	2	1	4	1
429678	6	1	Serine palmitoyltransferase 2	YDR062W	4	1	2	1	2	2	1	2	1	1	2	1
372474	7	1	Sphingosine-1-phosphate lyase	YDR294C	2	1	2	4	2	1	2	2	1	1	2	1
379742	7	0	Sphingolipid C4-hydroxylase SUR2	YDR297W	2	1	1	2	0	2	2	1	2	1	2	2
456289	8	0	Ceramide synthase LAG1	YHL003C	1	2	0	2	2	1	2	2	2	1	2	2
420050	6	3	Dihydrosphingosine 1-phosphate phosphatase LCB3	YJL134W	4	1	1	4	1	2	2	1	1	1	4	2
441179	7	2	Ceramide very long chain fatty acid hydroxylase SCS7	YMR272C	4	1	2	2	2	4	2	1	1	1	2	1
546344	6	3	Serine palmitoyltransferase 1	YMR296C	4	1	2	1	4	1	2	1	2	1	4	1
1859	6	2	Sphingoid long chain base kinase 4	YOR171C	4	1	1	1	1	2	4	2	2	1	2	1

Table 6. Summary of developmental expression dynamics of CDE orthogroups of sterol transport related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			S. cerevisiae ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
470387	7	6	Prefoldin subunit 2	–	4	4	4	4	1	2	4	0	1	0	4	0
475807	7	5	Phosphatidylglycerol/phosphatidylinositol transfer protein	YDL046W	2	1	4	4	0	4	1	1	4	1	4	2
378276	6	1	Oxysterol-binding protein homolog 7	YHR001W	4	1	2	2	1	1	1	1	2	1	2	2
494709	7	1	NPC intracellular sterol transporter 1-related protein 1	YPL006W	4	2	1	2	2	1	2	1	1	1	2	2
427232	10	2	Oxysterol-binding protein homolog 4	YPL145C	4	2	2	4	2	1	2	2	2	1	2	2
478741	6	2	SCP2 sterol transport protein family	–	2	1	2	4	1	2	0	1	2	1	4	1

Table 7. Summary of developmental expression dynamics of CDE orthogroups of membrane phoshpolipid and fatty acid biosynthesis genes genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4-developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			S. cerevisiae ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
Cardiolipin biosynthesis pathway																
539990	4	0	Phosphatidylglycerophosphatase GEP4, mitochondrial	YHR100C	2	2	1	0	1	1	1	2	0	1	2	1
471105	9	3	CDP-diacylglycerol--glycerol-3-phosphate 3-phosphatidyltransferase	YCL004W	2	1	2	2	2	1	2	1	4	4	4	2
522776	6	1	Cardiolipin synthase	YDL142C	4	2	1	0	1	2	2	1	0	1	2	2
CPD-DAG pathway																
442684	8	1	Phosphatidate cytidylyltransferase	YBR029C	2	2	2	1	2	1	1	2	1	4	2	...
436995	8	1	CDP-diacylglycerol-serine O-phosphatidyltransferase	YER026C	4	1	2	2	0	2	2	2	1	1	2	2
444758	11	4	Phosphatidylethanolamine N-methyltransferase	YGR157W	4	2	2	4	2	2	2	4	1	2	4	2
379032	7	1	Phosphatidylserine decarboxylase proenzyme 2	YGR170W	2	1	2	2	2	1	1	2	2	1	4	1
285011	9	6	Phosphatidyl-N-methylethanolamine N-methyltransferase	YJR073C	4	1	4	0	4	2	2	4	4	2	4	1
377789	7	2	Phosphatidylserine decarboxylase proenzyme 1, mitochondrial	YNL169C	4	1	2	4	0	1	2	2	1	1	2	2
436992	6	2	CDP-diacylglycerol-inositol 3-phosphatidyltransferase	YPR113W	4	1	2	2	2	2	1	1	1	1	4	1
Kennedy pathway																
449161	7	3	Choline-phosphate cytidylyltransferase	YGR202C	4	1	0	4	0	1	4	2	0	2	2	2
411802	3	1	Choline kinase	YLR133W	4	1	1	1	1	1	1	1	0	1	2	2
380149	10	5	Cholinephosphotransferase 1	YNL130C	4	2	4	2	2	4	2	2	1	1	4	4

Shared part of the pathway

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
76420	1	1	Long-chain-fatty-acid--CoA ligase 2	YER015W	4	0	0	0	0	0	0	0	0	0	0	0
439965	6	4	Elongation of fatty acids protein 2	YCR034W	4	1	4	1	4	2	1	2	0	1	4	1
462371	5	0	1-acyl-sn-glycerol-3-phosphate acyltransferase	YDL052C	2	1	1	1	2	1	2	1	2	1	2	1
379857	9	3	Acetyl-CoA carboxylase	YNR016C	4	2	1	4	0	2	2	4	1	2	2	2
414377	7	6	Glycerol-3-phosphate dehydrogenase [NAD(+)] 2, mitochondrial	YOL059W	4	1	4	4	2	1	4	1	4	1	4	1
372300	9	4	Lysophospholipid acyltransferase	YOR175C	4	2	4	2	2	2	2	1	4	1	4	1
538984	9	5	Fatty acid synthase subunit alpha	YPL231W	4	2	1	4	2	2	4	4	1	1	4	2
72617	4	4	Fatty acid desaturase	–	4	1	1	4	0	1	1	4	0	0	0	4
4671	6	0	Glycerol-3-phosphate O-acyltransferase 1	YBL011W	2	1	2	2	1	1	2	1	2	1	2	1
450732	8	3	NADPH-dependent 1-acyldihydroxyacetone phosphate reductase	YIL124W	2	1	2	4	2	1	4	1	2	2	4	1
422393	10	6	Fatty acid desaturase	–	4	2	4	2	4	4	2	4	1	1	2	4
437261	6	4	Long-chain-fatty-acid--CoA ligase 1	YOR317W	2	1	4	4	2	1	4	1	0	1	4	1

TAG pathway

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
437679	4	0	Phosphatidic acid phosphohydrolase 1	–	2	1	1	2	1	1	1	2	1	1	1	2
493617	3	1	Phospholipid:diacylglycerol acyltransferase	YNR008W	4	1	1	2	1	1	1	1	1	1	2	1
374579	7	2	Sterol O-acyltransferase 2	–	2	1	2	2	0	2	1	1	4	1	4	2
479811	6	2	Diacylglycerol O-acyltransferase 1	YOR245C	1	1	1	4	1	2	2	1	2	1	4	2

Table 8. Summary of developmental expression dynamics of CDE orthogroups of linoleic acid biosynthesis related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | S. cerevisiae ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
S.c. 2511002	10	3	Fatty acid desaturase	–	0	2	4	2	2	2	4	4	1	2	2	2
363743	9	7	Fatty acid desaturase	–	4	4	2	4	4	4	0	4	0	0	2	4
422976	8	8	Fatty acid desaturase	–	4	0	4	4	4	4	4	0	4	0	4	0
471478	9	7	Fatty acid desaturase	–	4	2	4	4	4	0	4	2	1	0	4	4
493933	11	9	Fatty acid desaturase	–	2	4	4	4	4	1	4	4	4	4	2	4
422393	10	6	Fatty acid desaturase	–	4	2	4	2	4	4	2	4	1	1	2	4

Table 9. Summary of developmental expression dynamics of CDE orthogroups of other lipid metabolism related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
20997	12	11	Glycerophosphodiester phosphodiesterase	–	4	4	4	4	4	4	4	4	4	2	4	4
380219	6	6	Diacylglycerol pyrophosphate phosphatases	–	4	4	0	4	0	0	0	0	4	0	4	4
376602	9	9	Caleosin	–	4	4	1	0	4	4	0	4	4	4	4	4
365330	12	9	Sphingolipid C9-methyltransferase	–	4	2	2	4	4	4	4	4	2	4	4	4
395241	11	8	Sphingolipid C9-methyltransferase	–	4	2	4	4	2	4	4	4	4	2	4	1
444998	8	6	Marvel domain protein	–	4	1	0	4	1	1	4	2	4	2	4	4
462849	10	5	Marvel domain protein	–	4	1	0	2	2	2	4	2	4	4	2	4

Table 10. Summary of developmental expression dynamics of CDE orthogroups of storage carbohydrate metabolism genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
360850	9	8	Alcohol dehydrogenase	–	4	1	4	4	4	4	4	2	4	1	4	1
369180	10	7	Sorbitol dehydrogenase	–	2	4	4	4	0	4	2	4	4	1	2	4
455149	11	6	CBM21 domain containing protein	–	4	2	2	4	4	2	2	2	4	1	4	4
441344	7	2	Glucoamylase	–	2	4	0	2	1	0	0	2	4	2	2	1
365699	8	5	1,4-alpha-glucan-branching enzyme	YEL011W	4	1	2	4	2	4	4	1	0	1	4	2
496146	2	0	Serine/threonine-protein phosphatase controlling glycogen metabolism	YER133W	2	1	1	0	1	2	1	1	1	1	1	1
457685	9	5	Glycogen synthase	YFR015C	4	2	2	4	1	2	4	1	4	1	4	2
473268	6	3	Glucoamylase	YIL099W	4	0	2	4	0	1	2	0	4	0	2	0
461263	7	3	Glycogenin	YJL137C	4	1	2	4	2	1	2	1	1	1	4	2
359312	9	5	UTP-glucose-1-phosphate uridylyltransferase	YKL035W	4	2	2	4	2	4	4	1	2	1	4	1

ID	>2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
497782	7	5	Phosphoglucomutase 2	YMR105C	4	1	4	4	0	2	4	2	0	1	4	1
538513	2	1	Cyclin-dependent protein kinase	YPL031C	2	1	1	1	1	1	1	1	1	1	4	1
495987	9	7	Glycogen phosphorylase	YPR160W	4	1	4	4	2	4	4	4	2	1	4	1
405214	10	6	Glycogen debranching enzyme	YPR184W	2	2	2	4	4	1	4	4	4	1	4	2
495249	10	7	Glycoside hydrolase family 13 alpha-amylase	–	4	4	4	4	4	4	2	2	0	2	4	1
544130	11	7	CBM21 domain containing protein	–	4	2	4	4	4	2	4	4	2	1	4	2
429272	4	4	Mannitol dehydrogenase	–	4	0	4	0	0	4	0	0	0	0	4	0
478544	8	2	Alpha,alpha-trehalose-phosphate synthase	YBR126C	2	1	2	2	1	2	2	2	4	1	4	1
437212	10	2	Neutral alpha-trehalase	YDR001C	2	2	2	2	4	2	1	2	2	1	2	4
473004	8	4	Trehalose-phosphatase	YDR074W	4	1	2	2	4	2	1	4	2	1	4	1
503099	7	3	Trehalose synthase complex regulatory subunit TSL1	YML100W	4	1	2	2	1	4	1	2	2	1	4	1
376383	9	6	Trehalase	–	4	2	2	0	4	4	1	4	4	1	4	2
540817	11	6	Trehalose phosphorylase	–	4	4	2	4	4	1	2	2	2	2	4	4

Table 11. Summary of developmental expression dynamics of CDE orthogroups of transcriptional regulator genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			*S. cerevisiae* ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
421729	9	5	Transcription factor, *C. cinerea* exp1	–	4	1	4	4	4	2	2	2	4	1	2	1
395717	7	7	Transcription factor	–	4	1	0	4	4	4	4	4	4	1	0	1
404567	9	7	Transcription factor	–	4	2	0	0	0	4	4	4	4	2	4	4
437393	11	9	Transcription factor	–	4	4	4	4	4	4	4	2	4	4	0	2
446227	9	8	Transcription factor	–	4	4	4	4	4	1	4	4	1	2	1	4
491825	10	4	Transcription factor	–	4	2	2	4	2	1	2	2	4	1	4	2
494418	10	8	Transcription factor	–	4	0	4	4	4	2	4	4	2	4	0	0
497620	12	9	Transcription factor	–	4	4	2	2	4	4	4	4	2	4	4	4
S.c. 2520583	7	6	Transcription factor	–	0	4	2	4	4	0	0	4	4	0	0	4
8610	10	6	Transcription factor, *C. cinerea* Cc.wc2	–	4	2	4	4	2	2	4	2	4	1	0	4
148855	10	6	Transcription factor, *L. edodes* PriB	–	4	4	1	2	2	2	4	2	4	1	4	4
493627	10	9	Transcription factor, *S. commune* Hom1	–	4	4	4	4	1	2	4	4	4	1	4	4
449111	7	6	Transcription factor	YDR043C	4	1	4	0	4	0	4	4	0	2	4	1
495715	12	12	Transcription factor, *S. commune* Tea1	YEL007W	4	4	4	4	4	4	4	4	4	4	4	4

ID			Description	Name	V1	V2	V3	V4	V5	V6	V7	V8	V9	V10	V11	V12
458586	10	3	Transcription-related RNA-binding protein	YGR159C	2	2	2	4	2	2	4	2	2	1	4	1
482097	10	7	Transcription factor	YPR015C	2	4	4	2	4	4	0	2	4	1	4	4
155160	9	6	Transcription initiation factor	–	4	2	2	4	4	4	4	2	4	1	0	0
449345	9	5	Transcription factor	–	4	2	1	4	4	4	2	2	2	1	4	1
463573	9	5	Transcription factor, *S. commune* Fst3	–	4	1	2	4	2	4	2	2	4	1	4	1
365344	9	6	Transcription factor, *S. commune* Fst4	–	4	2	2	4	4	4	2	1	4	1	4	1
211795	10	5	Transcription factor	–	4	4	2	4	2	4	2	2	4	1	2	1
498166	10	6	Possibly related to transcription elongation factors	–	2	4	4	2	1	1	4	4	4	2	2	4
508285	11	4	Potentially involved in in pre-mRNA 3' end processing	–	4	2	2	4	0	2	2	2	4	2	2	4
497197	7	6	Rho termination factor	–	4	1	0	4	0	2	4	4	4	4	0	0
142396	10	7	Transcription factor	–	4	1	4	2	4	4	4	4	4	2	0	2
351135	7	5	Transcription factor	–	4	2	4	4	0	4	2	1	0	1	4	1
357193	9	7	Transcription factor	–	4	1	4	4	2	1	4	4	4	1	4	2
359157	10	6	Transcription factor	–	4	2	2	4	4	2	2	4	4	1	4	1
360688	11	9	Transcription factor	–	4	4	4	4	2	4	4	4	4	1	2	4
367840	8	7	Transcription factor	–	4	1	4	4	1	4	4	2	4	1	4	1
374546	8	5	Transcription factor	–	4	2	4	4	0	4	0	2	2	1	4	1
411205	9	7	Transcription factor	–	4	4	4	4	2	4	1	2	4	1	4	1
416057	10	4	Transcription factor	–	2	2	2	4	2	1	4	4	2	2	0	4
440363	12	7	Transcription factor	–	4	2	2	4	4	4	4	2	4	2	4	2
441820	6	5	Transcription factor	–	4	1	4	4	2	4	1	1	1	1	4	1
442154	10	7	Transcription factor	–	4	4	4	4	2	1	2	4	2	4	0	4
443979	10	8	Transcription factor	–	4	4	4	4	0	2	4	4	0	2	4	4
451915	7	6	Transcription factor	–	4	0	2	0	4	4	0	4	4	4	0	0
452897	9	8	Transcription factor	–	4	4	2	4	4	4	0	0	1	4	4	4
465631	7	6	Transcription factor	–	4	2	4	4	0	4	4	0	4	0	0	1
492675	7	4	Transcription factor	–	4	2	4	2	1	4	1	0	1	1	4	2
500056	10	6	Transcription factor	–	4	2	2	4	2	4	1	4	4	1	4	2
S.c. 2614371	7	7	Transcription factor	–	0	4	0	4	0	0	4	0	4	4	4	4
S.c. 2705300	6	5	Transcription factor	–	0	4	0	4	0	2	4	4	0	4	0	1
438877	9	5	Transcription related protein	–	2	4	2	4	4	2	4	2	1	1	4	1
471238	10	10	RNA binding protein; distantly related to yeast PAB1	–	4	4	0	4	4	0	4	4	4	4	4	4
369621	10	4	Transcription factor, *C. cinerea* dst1	–	4	2	4	4	2	1	2	2	2	1	4	2
538904	8	6	Transcription factor, *S. commune* Gat1	–	4	2	4	4	1	4	4	1	4	1	2	1
355613	9	3	Transcription factor, *S. commune* Hom2	–	4	1	4	2	4	2	2	2	2	1	1	2
413307	7	5	Transcription related Mynd-type zinc finger	–	4	4	4	4	0	4	1	2	0	1	2	1

Table 12. Summary of developmental expression dynamics of CDE orthogroups of RNA-binding protein encoding genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
357717	12	12	RNA binding domain protein	–	4	4	4	4	4	4	4	4	4	4	4	4
354109	8	6	mRNA-binding Pumilio protein	YLL013C	4	4	1	4	0	2	2	4	0	1	4	4
444698	10	3	RNA binding domain protein	–	2	2	2	2	0	2	4	1	4	2	4	2
541323	11	10	Armadillo domain containing RNA binding protein	–	4	4	4	4	4	2	4	4	4	1	4	4
356106	10	5	RNA binding domain protein	–	4	2	2	4	4	2	2	2	4	1	4	1
424800	12	11	RNA binding domain protein	–	4	4	2	4	4	4	4	4	4	4	4	4
193545	11	11	RNA binding domain protein	–	4	4	4	4	4	4	4	4	4	4	0	4

Table 13. Summary of developmental expression dynamics of CDE orthogroups of RNA interference related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
155983	6	2	RNA-dependent RNA polymerase	–	4	1	2	2	1	2	4	2	1	1	1	1
498671	4	1	RNA-dependent RNA polymerase	–	4	1	1	1	2	1	2	1	2	1	0	1
438955	4	2	Dicer-like protein	–	2	1	1	2	4	4	1	1	1	1	1	1
464843	2	1	RNA-dependent RNA polymerase	YDL128W	4	0	0	0	0	2	0	0	0	0	0	0
366047	6	1	Dicer-like protein	YLR332W	1	1	1	2	2	4	2	1	2	1	2	1
473372	8	0	AGO-like protein	–	2	2	1	2	1	2	2	2	2	1	2	1
495272	6	4	Argonaute-like protein	–	4	1	0	2	2	1	4	1	4	1	4	1
502116	9	6	Argonaute-like proteins with high similarity to *A. nidulans* rsdA and *S. pombe* ago1	–	4	2	1	4	2	4	4	4	4	1	2	1
363313	7	1	Orthologs of QDE-2	–	2	2	1	2	1	1	2	2	4	1	2	1
440860	5	1	RNA-dependent RNA polymerase	–	4	1	2	2	1	1	1	1	2	1	2	1
496192	5	1	RNA-dependent RNA polymerase	–	2	0	0	2	2	4	2	1	0	1	1	0
498410	5	1	RNA-dependent RNA polymerase	–	4	1	2	2	1	0	2	1	1	1	2	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
355234	1	1	AGO ortholog	–	4	0	0	0	0	0	0	0	0	0	0	0
368472	1	0	Argonaute-like protein	–	2	0	0	0	0	0	0	0	0	0	0	0
465906	1	0	Argonaute-like protein	–	2	0	0	0	0	0	0	0	0	0	0	0
444647	3	0	Dicer-like protein	–	2	1	1	1	1	1	1	1	2	1	2	1
124704	1	1	RNA-dependent RNA polymerase	–	4	0	0	0	0	0	0	0	0	0	0	0
444494	1	1	RNA-dependent RNA polymerase	–	4	0	0	0	0	0	0	0	0	0	0	0

Table 14. Summary of developmental expression dynamics of CDE orthogroups of chromatin-related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			*S. cerevisiae* ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
58294	10	0	Actin-related protein 4	–	2	2	2	2	2	1	2	2	2	2	2	1
485296	9	0	Protein arginine N-methyltransferase 1	YBR034C	2	1	2	2	2	2	2	2	2	1	2	1
498885	11	8	Histone H2A.Z-specific chaperone CHZ1	YER030W	4	0	2	4	2	2	4	4	4	4	4	4
493226	10	7	Histone isomerase	YML074C	4	2	4	4	4	2	2	4	4	1	4	1
455597	9	9	NAD-dependent histone deacetylase HST3	YOR025W	4	4	4	4	4	1	0	4	4	1	4	4
545797	11	9	*C. cinerea* Ich1 gene	–	4	4	4	4	1	2	4	4	4	4	2	4
392307	11	5	HIRA-interacting protein 3	–	2	2	4	4	2	2	4	2	4	1	4	2
384835	8	6	Poly(ADP-ribose) polymerase	–	4	2	0	4	0	0	2	4	4	1	4	4
459336	9	5	Poly(ADP-ribose) polymerase	–	2	2	2	4	1	4	4	0	4	1	4	2
473747	8	6	High mobility group protein	–	4	1	4	4	2	2	0	4	4	1	4	0
66389	10	4	Suppressor protein STM1	YLR150W	4	2	4	2	1	4	2	2	4	2	2	1
264120	2	0	SET domain protein	YJL168C	1	1	1	1	1	1	2	1	1	1	2	1
170659	1	0	SET domain protein	–	2	0	0	0	0	0	0	0	0	0	0	0
456684	3	0	SET domain protein	–	2	1	1	1	0	2	0	2	0	0	1	1
497003	1	0	SET domain protein	–	2	1	0	1	1	0	1	1	1	1	0	1
365798	2	0	*C. cinerea* Snf5 family chromatin remodeling protein	–	2	1	1	1	2	1	1	1	1	1	1	1
252667	8	2	Histone acetyltransferase type B subunit 2	YEL056W	2	2	4	2	1	1	2	2	4	1	2	1
518613	10	4	Histone acetyltransferase complex member	YLL022C	2	2	2	2	0	2	4	4	4	1	4	2
420113	7	4	Histone acetyltransferase type B catalytic subunit	YPL001W	4	1	2	1	1	2	4	1	4	1	4	2
457726	8	4	Histone acethytransferase	–	4	1	4	4	2	1	2	0	2	0	4	2
430366	6	0	Histone deacetylase	YGL194C	2	1	1	2	2	1	1	2	2	1	2	1
415916	4	0	Histone deacethylase	YNL021W	2	1	1	2	2	1	2	1	1	1	1	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
354601	6	2	Histone deacetylase	YNL330C	4	2	2	2	1	1	2	1	1	1	4	1
52602	5	3	Histone deacetylase	YPL116W	4	1	1	4	1	1	4	2	0	2	1	1
364572	5	0	Histone deacetylase	–	1	1	2	2	1	1	2	1	2	1	2	1
465141	8	2	Histone deacetylase	–	2	4	4	2	2	1	2	1	2	1	1	2
363625	8	6	Agaricales-specific histone group	–	4	0	0	0	4	4	2	4	4	2	4	0
154410	11	6	Centromere-specific histone variant	–	2	4	2	2	2	4	2	4	4	1	4	4
182362	11	5	Histone	–	4	4	2	2	2	2	2	2	4	0	4	4
473916	8	6	Histone	–	2	0	4	4	1	4	0	2	4	0	4	4
393767	9	5	Histone H2B	–	1	1	4	4	2	2	4	4	0	2	2	4
361866	5	4	Histone H2B.2	YBL002W	2	0	0	4	0	0	4	4	4	0	0	0
418275	10	7	Histone H2A.2	YBL003C	2	4	4	4	2	4	4	0	0	2	4	4
401831	9	6	Histone H4	YBR009C	1	4	4	2	1	4	2	4	4	1	2	4
546355	4	1	Histone H3	YBR010W	2	0	1	2	1	1	1	2	4	1	1	1
284515	11	2	Histone H2A.Z	YOL012C	2	2	4	2	2	2	2	2	4	2	2	1
467558	10	9	Histone H1	YPL127C	4	4	4	4	4	4	4	4	1	2	0	4
320458	0	0	Histone	–	1	0	0	0	0	0	0	0	0	0	0	0
324852	0	0	Histone	–	1	0	0	0	0	0	0	0	0	0	0	0
380437	1	1	Histone	–	4	0	0	0	0	0	0	0	0	0	0	0
385598	1	1	Histone	–	4	0	0	0	0	0	0	0	0	0	0	0
446837	1	0	Histone	–	2	0	0	0	0	0	0	0	0	0	0	0
492876	1	0	Histone	–	2	0	0	0	0	0	0	0	0	0	0	0
503008	1	0	Histone	–	2	0	0	0	0	0	0	0	0	0	0	0
452576	4	0	AMP-activated protein serine/threonine kinase alpha subunit Ssp2	YDR477W	2	1	1	2	1	1	2	1	2	1	1	1
360033	4	1	5'-AMP-activated protein kinase subunit gamma	YGL115W	1	1	4	1	1	1	2	2	1	1	2	1
377394	8	1	Histone-lysine N-methyltransferase, H3 lysine-79 specific	YDR440W	2	1	1	4	2	1	2	2	2	1	2	2
511209	9	8	SRA-YDG domain protein	–	2	4	4	4	4	4	4	0	0	0	4	4

Table 15. Summary of developmental expression dynamics of CDE orthogroups of signal transduction related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			*S. cerevisiae* ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
S.c. 2643671	7	6	Protein kinase related to mating	–	1	2	4	2	4	4	4	1	1	1	4	4
363127	8	7	Alcohol dehydrogenase	YMR318C	4	1	0	4	4	0	4	4	2	4	0	4

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
398037	9	7	Linoleate diol synthase	–	4	1	4	2	2	4	4	0	4	4	0	4
423716	10	5	Linoleate diol synthase kinase	–	4	4	2	4	0	2	4	2	1	2	4	2
357261	8	7	Serine-threonine kinase	–	4	4	0	4	0	2	0	4	0	4	4	4
397547	7	5	Serine-threonine kinase	–	4	0	0	4	4	0	0	2	4	2	0	4
405337	12	11	Serine-threonine kinase	–	4	4	2	4	4	4	4	4	4	4	4	4
441896	10	8	Serine-threonine kinase	–	4	0	2	4	4	4	2	4	4	4		0
456276	10	8	Serine-threonine kinase	–	2	4	2	4	4	4	4	1	4	1	4	4
493180	9	4	Serine-threonine kinase	–	4	2	2	4	1	2	1	2	2	1	4	4
538093	11	8	Serine-threonine kinase	–	2	4	2	4	2	4	4	4	0	4	4	4
462137	11	9	Serine-threonine kinase	YDR247W	2	2	4	4	4	4	4	4	4	1	4	4
449305	10	4	P38 mitogen-activated protein kinase HOG1	YLR113W	4	1	2	2	2	4	4	2	2	1	4	2
451611	10	6	Serine-threonine kinase	YLR248W	4	2	1	4	4	4	2	2	2	1	4	4
448193	10	6	Serine-threonine kinase	–	2	4	1	4	1	2	4	4	4	2	4	2
359429	6	5	Serine-threonine kinase	–	4	0	0	4	1	4	2	0	4	0	4	0
448807	6	5	Serine-threonine kinase	–	4	0	4	2	4	0	0	4	4	1	0	0
470972	8	6	Serine-threonine kinase	–	4	4	0	2	0	4	1	4	2	4	0	4

Table 16. Summary of developmental expression dynamics of CDE orthogroups of pheromone precursor genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog *in S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			S. cerevisiae ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
153902	2	0	Protein-S-isoprenylcysteine O-methyltransferase	YDR410C	2	1	0	1	2	1	1	1	1	1	1	1
116060	1	1	Mating pheromone precursor	–	4	0	0	0	0	0	0	0	0	0	0	0
446846	1	1	Mating pheromone precursor	–	4	0	0	0	0	0	0	0	0	0	0	0
446955	1	1	Mating pheromone precursor	–	4	0	0	0	0	0	0	0	0	0	0	0
446956	1	1	Mating pheromone precursor	–	4	0	0	0	0	0	0	0	0	0	0	0
446958	0	0	Mating pheromone precursor	–	1	0	0	0	0	0	0	0	0	0	0	0
446959	1	1	Mating pheromone precursor	–	4	0	0	0	0	0	0	0	0	0	0	0
446966	0	0	Mating pheromone precursor	–	1	0	0	0	0	0	0	0	0	0	0	0
356637	1	1	Pheromone sensing GPCR Ste3 homolog	–	4	0	0	0	0	0	0	0	0	0	0	0
394943	4	2	Pheromone sensing GPCR Ste3 homolog	–	4	0	0	4	2	1	0	0	0	0	2	0
477564	5	3	Pheromone sensing GPCR Ste3 homolog	–	4	0	2	0	4	0	0	2	0	0	4	0
377901	7	3	CAAX prenyl protease 2	–	4	1	2	1	1	2	4	1	4	1	2	2

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
203704	6	3	Pheromone sensing GPCR Ste3 homolog	–	4	2	0	4	4	2	1	0	1	1	0	2
465723	8	4	Pheromone sensing GPCR Ste3 homolog	–	2	1	4	4	2	1	4	2	1	1	4	2
486965	5	3	Pheromone sensing GPCR Ste3 homolog	–	2	0	0	4	0	4	0	2	1	1	4	0
491483	3	2	Protein farnesyltransferase subunit beta	YDL090C	4	1	1	1	4	1	1	1	1	1	2	1
437675	5	0	Pheromone-processing carboxypeptidase KEX1	YGL203C	2	1	1	1	2	2	2	1	1	1	2	1
545529	5	1	Dipeptidyl aminopeptidase B	YHR028C	1	1	4	2	1	1	1	2	2	1	2	1
419650	6	1	CAAX prenyl protease 1 ste24	YJR117W	4	2	2	1	2	1	1	2	1	1	2	1
473005	8	2	Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha	YKL019W	4	2	1	4	2	2	2	2	1	1	2	1
540181	10	4	Alpha-factor-transporting ATPase	YKL209C	4	4	2	2	2	4	2	2	4	1	1	2
358206	8	2	A-factor-processing enzyme	YLR389C	2	1	1	2	2	4	4	1	2	1	2	2
448165	5	2	Putative kexin	YNL238W	1	2	2	4	1	1	1	1	4	1	1	2
106286	7	2	Pheromone sensing GPCR Ste3 homolog	–	4	1	2	4	2	0	2	2	2	0	0	1
152667	9	9	Homolog of *S. cerevisiae* Ste14	–	4	0	4	4	4	4	4	4	1	4	4	0

Table 17. Summary of developmental expression dynamics of CDE orthogroups of conserved signal transduction pathway genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| ID | Fold change | | Putative function | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
cAMP																
465055	4	0	Phosphatidylinositol phosphate phosphatase	–	2	1	2	1	1	1	2	1	2	1	1	1
462760	6	0	GTPase involved in G-protein signaling in adenylate cyclase activation	–	2	1	2	2	1	2	1	2	1	1	2	1
539854	10	3	Low-affinity cyclic AMP phosphodiesterase	YGL248W	4	4	2	2	1	2	4	2	2	1	2	2
494851	7	2	Phosphatidylinositol phosphate phosphatase	YKL212W	2	2	4	2	1	1	2	1	2	1	4	1
175104	5	1	High-affinity cyclic AMP phosphodiesterase	YOR360C	1	1	4	2	1	2	1	1	1	1	2	2
45965	7	2	Protein kinase A catalytic subunit 2	YPL203W	4	2	2	2	1	1	1	2	2	1	4	1
361063	6	1	Protein kinase A catalytic subunit 1	–	2	1	4	1	2	1	1	2	2	1	2	1
497554	2	0	Adenylate cyclase	–	1	1	1	2	1	1	1	1	2	1	1	1
449966	1	1	Carbonic anhydrase	–	4	0	0	0	0	0	0	0	0	0	0	0
Cell wall stress																
453815	1	0	1-phosphatidylinositol 4-kinase	–	2	1	1	1	1	1	1	1	1	1	1	1

STT4

ID			Description	ORF												
464132	4	2	1-phosphatidylinositol-4-phosphate 5-kinase	–	4	1	1	1	2	2	1	1	1	1	4	1
494847	2	0	Mitogen-activated protein kinase kinase kinase BCK1	–	1	0	0	1	1	2	1	0	2	1	0	1
186221	4	0	Mitogen-activated protein kinase kinase MKK1	–	1	1	2	2	1	1	1	1	1	1	2	2
466997	3	0	Rho family guanine nucleotide exchange factor ROM2	–	2	1	1	1	0	1	2	1	2	1	1	1
419097	5	1	Classical protein kinase C alpha type	YBL105C	2	1	2	1	1	2	1	1	1	1	4	1
373470	7	2	Cyclin-dependent serine/threonine-protein kinase CDC28	YBR160W	2	1	1	0	1	4	2	2	1	1	4	2
418410	5	1	3-phosphoinositide dependent protein kinase-3	YDR490C	2	2	1	1	2	1	1	1	2	1	4	1
528086	6	0	Rho family GTPase RHO1, Ras homolog gene family, member A	YPR165W	2	1	2	2	1	1		1	2	1	2	1
446030	7	1	1,3-beta-D-glucan synthase	YGR032W	2	1	2	4	1	2		1	2	1	2	1
498153	6	2	Tyrosine protein phosphatase PTP3	YER075C	2	2	1	4	1	1	2	1	2	1	4	1
413665	8	5	Mitogen-activated serine/threonine-protein kinase SLT2	YHR030C	2	1	4	4	1	4	2	2	4	1	4	1

High osmolarity pathway

ID			Description	ORF												
474858	4	0	Mitogen-activated protein kinase kinase kinase, osomolarity two-component system, response regulator SSK2	–	1	1	1	2	2	2	1	1	1	1	2	1
494483	4	0	Rho family guanine nucleotide exchange factor CDC24, cell division control protein 24	–	2	1	1	2	1	2	1	1	2	1	1	1
446453	6	0	Type 2C protein phosphatase PTC1	YDL006W	2	1	2	2	1	1	2	1	2	1	0	2
356186	6	1	High osmolarity two-component System, phosphorelay intermediate protein YPD1	YDL235C	4	0	0	2	0	1	2	2	2	1	0	2
484600	10	4	G-protein beta-like WD40 protein	YMR116C	4	2	4	2	2	1	2	2	4	2	4	1
498153	6	2	Tyrosine protein phosphatase PTP3	YER075C	2	2	1	4	1	1	2	1	2	1	4	1
419006	5	0	Pheromone receptor transcription factor	YMR043W	2	1	1	2	1	2	2	2	0	1	1	0

Pheromone pathway

ID			Description	ORF												
16197	1	0	GTPase-activating protein SST2	–	2	1	1	1	1	1	1	1	1	1	1	1
370279	4	0	Mitogen-activated protein kinase kinase	–	1	2	2	1	2	1	1	2	1	1	1	1
353910	1	0	Mitogen-activated protein kinase kinase kinase STE11	–	1	1	0	2	1	1	1	1	1	1	0	1
446910	3	0	Serine/threonine protein kinase SKM1/CLA4, Serine/threonine-protein kinase CLA4	–	1	1	1	2	2	1	1	1	0	1	2	1
416655	3	0	Rho family GTPase CDC42	–	2	1	1	0	1	1	2	1	1	1	2	1
437458	3	0	Scaffold protein BEM1; phosphatidylinositol-3-phosphate-binding protein BEM1, bud emergence protein 1	–	2	1	2	1	1	1	1	1	2	1	1	1
455590	2	0	The RA domain of Ste50 adaptor protein is required for delivery of Ste11 to the plasma membrane in the filamentous growth signaling pathway (???)	–	2	1	1	1	0	1	1	1	1	1	2	1

370180	4	0	Tyrosine/serine/threonine protein phosphatase MSG5	–	2	1	1	1	2	1	1	1	2	1	2	1
445505	5	1	Mitogen-activated serine/threonine-protein kinase FUS3	YBL016W	1	2	2	1	1	1	2	4	1	1	2	1
429295	6	0	Mitogen-activated protein kinase kinase kinase kinase STE20, p21-activated kinase 1	YHL007C	2	1	2	2	0	2	1	1	2	1	2	1
84056	4	2	Guanine nucleotide-binding protein subunit gamma	YJR086W	2	1	2	0	1	1	1	1	4	1	4	1
449416	5	0	Guanine nucleotide-binding protein GI/GS/GT subunit beta-1	YOR212W	2	1	2	2	1	1	1	1	2	1	2	1
498153	6	2	Tyrosine protein phosphatase PTP3	YER075C	2	2	1	4	1	1	2	1	2	1	4	1
419006	5	0	Pheromone receptor transcription factor	YMR043W	2	1	1	2	1	2	2	2	0	1	1	0
376911	8	1	Homeodomain family transcription factor STE12	–	4	1	2	2	1	2	2	2	2	1	0	2

Starvation pathway

356859	4	2	SHO1 osmosensor	–	4	1	2	1	1	2	1	1	1	1	4	1
197636	2	1	Signaling mucin MSB2	–	2	0	1	0	1	0	0	1	4	0	0	0
389009	7	0	14-3-3 family protein BMH1	YDR099W	1	2	2	1	2	2	1	2	2	1	2	1
498153	6	2	Tyrosine protein phosphatase PTP3	YER075C	2	2	1	4	1	1	2	1	2	1	4	1
376911	8	1	Homeodomain family transcription factor STE12	–	4	1	2	2	1	2	2	2	2	1	0	2

TOR

407528	6	3	SEA complex subunit Seh1	YGL100W	4	1	2	2	1	1	1	4	4	1	2	1
365555	4	0	Protein kinase A regulatory subunit	YIL033C	2	1	2	2	1	1	1	2	1	1	1	1
434690	10	4	Phosphoinositide PI4,5P(2) binding Protein, forms a complex with Slm2p	YIL105C	2	1	4	4	2	2	4	2	2	1	4	2
502431	5	1	COPII-coated vesicle component Sec13	YLR208W	1	1	2	1	2	2	1	2	0	1	4	1
466474	8	6	FK506-binding protein 1	YNL135C	2	4	4	0	4	2	4	4	1	1	4	1
81628	6	0	Gtr1/RagA G protein Gtr1	–	2	1	2	2	1	2	2	1	1	1	2	1
417412	2	0	Component of a membrane-bound complex containing the Tor2p kinase	–	2	1	1	1	1	1	1	1	1	1	2	1
498449	1	0	DEP domain containing 5; subunit of SEACIT (GATOR1)	–	2	1	1	1	0	1	1	1	1	1	1	1
438057	4	1	Gtr1/RagA G protein Gtr2	–	2	1	1	2	1	1	1	1	1	1	4	2
467184	4	1	Guanyl-nucleotide exchange factor, HOPS complex subunit Vam6	–	1	1	2	0	1	1	4	1	2	1	2	1
440799	4	0	Nitrogen permease regulator 2-like protein; subunit of SEACIT (GATOR1)	–	1	1	1	1	1	2	2	1	2	1	2	1
366098	2	0	Nitrogen permease regulator 3-like protein; subunit of SEACIT (GATOR1)	–	2	1	1	1	1	2	1	1	1	1	1	1
374354	4	1	Protein required for the transport of Gap1p	–	4	2	1	2	1	1	1	1	1	1	2	1
493094	1	1	Ras homolog	–	4	0	0	0	0	0	0	0	0	0	0	0
517273	3	0	Subunit of SEACAT, a subcomplex of the SEA complex; Mtc1p	–	1	1	1	2	1	2	1	1	1	1	2	1
367284	0	0	Subunit of SEACAT, a subcomplex of the SEA complex; Sea4p	–	1	1	1	1	1	1	1	1	1	1	1	1
440869	2	0	Subunit of TORC1	–	1	1	1	1	1	2	1	1	2	1	1	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
495775	3	1	Subunit of TORC2, RICTOR homolog	–	4	1	1	2	1	1	1	1	0	1	2	1
439934	2	0	Target of rapamycin	–	2	1	1	2	0	1	1	1	1	1	1	1
23387	3	0	Tuberous sclerosis 2 protein homolog	–	2	1	1	2	1	1	1	1	1	1	2	1

Table 18. Summary of developmental expression dynamics of CDE orthogroups of G-protein coupled receptor genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
283463	6	0	G-protein coupled receptor	YKL039W	2	1	2	2	1	1	2	1	2	1	2	1
263790	4	3	G-protein coupled receptor	–	4	1	0	4	1	4	0	0	0	0	0	2
485193	6	0	G-protein coupled receptor	–	2	2	1	2	2	1	1	2	1	1	2	1
379008	0	0	G-protein coupled receptor	–	1	0	0	0	0	0	0	0	0	0	0	0
489342	1	1	G-protein coupled receptor	–	4	0	0	0	0	0	0	0	0	0	0	0
229833	11	4	G-protein involved in signal transduction	–	2	1	4	2	2	4	2	2	4	2	4	2
479562	11	7	Regulator of G-protein signaling	–	4	4	4	4	2	2	4	1	2	2	4	4
S.c. 2490312	9	6	Putative arrestin	–	0	4	2	4	4	4	1	2	0	4	4	2
473465	10	5	Putative arrestin	–	4	4	4	2	1	2	2	2	4	1	4	2
472370	11	7	Ras superfamily protein	–	4	2	2	4	4	4	1	4	4	2	4	2

Table 19. Summary of developmental expression dynamics of CDE orthogroups of velvet protein encoding genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
365674	4	1	Velvet factor	–	4	1	1	1	1	1	2	1	2	1	2	1
355684	3	3	Velvet factor	–	4	0	1	0	4	4	0	0	0	0	0	0
374867	6	1	Velvet factor	–	2	1	2	1	2	2	2	1	4	0	1	1
385561	8	5	Velvet factor	–	4	4	1	0	0	4	2	4	4	1	2	2

| 496341 | 10 | 4 | Velvet factor | – | 4 | 2 | 0 | 2 | 2 | 4 | 2 | 1 | 4 | 2 | 4 | 2 |

Table 20. Summary of developmental expression dynamics of CDE orthogroups of STRIPAK complex protein genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| ID | Fold change | | Putative function | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
51761	1	0	Forkhead-associated domain	–	2	1	1	1	1	1	1	1	1	1	1	1
439971	1	0	Protein phosphatase PP2A regulatory subunit A	YAL016W	1	1	1	1	1	1	1	1	1	1	2	1
280222	5	0	Serine/threonine-protein kinase KIC1	YHR102W	1	1	1	2	2	2	1	2	1	1	2	1
375927	4	0	Factor arrest protein 11	YNL127W	2	1	1	2	1	1	2	1	1	1	2	1
463106	5	2	Serine/threonine-protein phosphatase PP2A-like PPG1	YNR032W	4	1	4	1	1	1	2	2	1	1	2	1
499433	8	2	MOB kinase activator	–	4	2	2	1	2	1	1	4	2	2	2	1
108726	5	0	Striatin	–	2	1	2	2	1	1	1	2	1	1	2	1

Table 21. Summary of developmental expression dynamics of CDE orthogroups of light sensing related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| ID | Fold change | | Putative function | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
492962	9	7	DASH-type cryptochrome	YOR386W	4	1	2	4	4	0	4	2	4	4	4	1
470504	7	2	Phytochrome	–	4	1	0	4	2	0	2	1	2	1	2	2
361584	11	9	*C. cinerea* dst2, putative AA7 family member	–	4	2	4	4	4	4	4	4	4	1	4	2

Table 22. Summary of developmental expression dynamics of CDE orthogroups of chitin biosynthesis genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4.

ID	Fold change >2	Fold change >4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
371191	9	4	Glucosamine 6-phosphate N-acetyltransferase	–	4	2	4	4	1	2	2	2	2	1	4	1
538650	6	0	Chitin synthase regulator SKT5	YBL061C	2	1	2	2	2	1	1	2	2	1	1	1
542039	8	1	Chitin synthase 3	YBR023C	2	2	2	4	2	2	1	1	2	1	2	1
357695	7	5	UDP-N-acetylglucosamine pyrophosphorylase	YDL103C	4	2	4	4	4	1	1	2	1	1	4	1
478844	4	1	Phosphoacetylglucosamine mutase involved in chitin precursor synthesis	YEL058W	4	1	1	2	2	1	1	1	1	1	2	1
440525	9	4	Glutamine-fructose-6-phosphate aminotransferase	YKL104C	4	1	4	4	2	2	2	2	2	1	4	1
439486	3	1	Chitin synthase 1	YNL192W	4	1	1	1	1	2	1	1	1	1	2	1
473500	4	2	Cell wall related glycosyltransferase	YPL227C	2	1	4	1	1	1	1	2	1	1	4	1
382141	7	1	Dolichol-phosphate mannosyltransferase	YPR183W	2	1	2	1	1	4	1	2	2	1	2	2
372414	4	2	Cell wall related glycosyltransferase	–	2	4	0	1	2	0	4	1	0	1	0	1
15443	1	1	Chitin synthase	–	4	0	0	0	0	0	0	0	0	0	0	0
234699	3	0	Chitin synthase	–	1	1	2	0	1	1	1	1	2	1	1	2
357091	5	1	Chitin synthase	–	1	1	1	2	2	2	1	1	2	1	4	1
359180	12	10	Chitin synthase	–	4	4	4	4	2	4	4	4	2	4	4	4
493091	8	2	Chitin synthase	–	2	1	4	2	2	2	2	1	2	1	4	1
374670	3	0	Chitin synthase with myosin motor domain	–	2	1	0	2	1	1	1	1	1	1	2	1
497826	5	0	Chitin synthase with myosin motor domain	–	2	1	2	2	1	2	1	1	2	1	1	1
519767	11	6	Glycosyltransferase 2 family protein	–	4	2	4	4	2	4	2	4	2	1	4	2

Table 23. Summary of developmental expression dynamics of CDE orthogroups of glucan biosynthesis related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	Fold change >4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
417746	12	4	Kre9/Knh1 family putative beta-1,6-glucan assembly protein	–	4	2	2	4	2	2	2	2	4	2	4	2
181364	4	0	Smi1 family protein regulating chitin deposition and cell wall assembly	–	2	1	0	2	2	1	1	1	1	1	2	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
359282	9	2	Glucosidase 2 subunit alpha	YBR229C	2	1	4	4	2	2	2	2	2	1	2	1
486333	7	0	Mannosyl-oligosaccharide glucosidase	YGL027C	2	1	2	2	2	1	1	2	1	1	2	2
494484	8	6	Glycoside hydrolase 63 family alpha-glucosidase	YMR196W	4	1	4	4	2	1	4	1	4	1	4	2
241938	5	1	Protein ROT1 involvedin regulating cell wall 1,6-beta-glucan level	YMR200W	4	1	2	2	1	1	1	1	2	1	1	2
543646	1	0	1,3-beta-glucanosyltransferase GAS1	YMR307W	2	0	0	0	0	0	0	0	0	0	0	0
447635	3	0	Beta-glucan synthesis-associated protein KRE6	YPR159W	2	1	1	2	1	1	1	1	0	1	2	1
S.c. 2584405	7	6	Beta-1,6-glucan synthase, GH16 family protein	–	0	4	0	2	0	0	4	4	4	1	4	4
432992	6	1	Big1 family protein putetivaly required for beta-1,6-glucan synthesis	–	4	1	2	2	2	2	1	1	2	1	1	1
499280	8	6	Glycoside hydrolase 31 family alpha-glucosidase	–	4	0	1	4	0	4	2	4	4	2	4	0
S.c. 2543909	6	5	Glycoside hydrolase 31 family alpha-glucosidase	–	0	4	2	4	0	0	4	4	0	1	4	1
526754	10	3	Glycoside hydrolase 9 family protein	–	2	2	2	2	1	1	4	4	4	2	2	2
359649	5	4	Kre9/Knh1 family putative beta-1,6-glucan assembly protein	–	4	1	4	4	1	4	2	1	0	1	0	1
465604	10	3	Kre9/Knh1 family putative beta-1,6-glucan assembly protein	–	4	1	2	2	2	4	2	2	2	1	4	2
470186	12	10	Kre9/Knh1 family putative beta-1,6-glucan assembly protein	–	4	2	4	4	4	4	4	2	4	4	4	4
S.c. 2557059	9	9	Kre9/Knh1 family putative beta-1,6-glucan assembly protein	–	0	4	0	4	4	0	4	4	4	4	4	4
528086	6	0	Rho family GTPase RHO1, Ras homolog gene family, member A	YPR165W	2	1	2	2	1	2	1	1	2	1	2	1
446030	7	1	1,3-beta-D-glucan synthase	YGR032W	2	1	2	4	1	2	2	1	2	1	2	1

Table 24. Summary of developmental expression dynamics of CDE orthogroups of chitin remodeling genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			*S. cerevisiae* ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
360173	7	2	CE4 chitin deacetylase	–	4	1	2	4	0	1	2	2	0	1	2	2
368217	10	9	Glycoside hydrolase 18, chitinase	–	4	4	4	4	4	0	4	4	2	4	0	4
S.c. 85210	9	7	Glycoside hydrolase 18, chitinase	–	0	4	4	2	4	0	4	4	4	2	4	1
543586	7	2	Glycoside hydrolase 18, chitinase	YDR371W	4	1	2	2	2	1	2	2	1	1	4	1

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
544873	6	4	Polyamine transporter 2	YGR138C	4	1	0	4	0	2	4	2	0	1	4	1
354859	10	4	Endochitinase	YLR286C	4	2	1	2	4	2	2	4	4	2	2	1
355445	11	8	CE4 chitin deacetylase 1	YLR307W	4	2	2	4	4	1	4	4	4	4	4	2
413423	11	7	CBM5/12 domain protein	–	4	2	4	4	4	2	2	4	4	1	4	2
375733	7	6	CE4 chitin deacetylase	–	4	1	4	4	2	4	1	1	4	1	4	1
377259	10	4	CE4 chitin deacetylase	–	4	4	4	1	4	2	0	2	2	2	2	2
379321	7	5	CE4 chitin deacetylase	–	4	0	4	0	4	2	2	0	4	0	4	0
42776	2	1	CE4 chitin deacetylase	–	4	0	0	0	2	0	0	0	0	0	0	0
440521	7	7	CE4 chitin deacetylase	–	4	4	4	4	4	4	0	0	0	0	4	1
464952	1	1	CE4 chitin deacetylase	–	4	0	0	0	0	0	0	0	0	0	0	0
502602	1	1	CE4 chitin deacetylase	–	4	0	0	0	0	0	0	0	0	0	0	0
358869	8	3	Glycoside hydrolase 18, chitinase	–	4	2	2	0	2	2	4	0	0	1	4	2
470416	9	6	Glycoside hydrolase 18, chitinase	–	4	2	4	4	2	4	0	2	4	0	4	1
488885	10	7	Glycoside hydrolase 18, chitinase	–	4	4	4	4	4	1	4	2	1	4	2	2
90984	1	1	Glycoside hydrolase 18, chitinase	–	4	0	0	0	0	0	0	0	0	0	0	0
91051	9	8	Glycoside hydrolase 18, chitinase	–	4	1	4	4	0	4	4	4	2	4	4	1
446427	8	6	Glycoside hydrolase 20 family putative β-1,6-N-acetylglucosaminidase	–	4	1	4	4	4	0	4	2	0	2	4	1
464567	9	6	Glycoside hydrolase 20 family putative β-1,6-N-acetylglucosaminidase	–	4	1	4	4	4	2	2	4	2	1	4	1
520359	10	7	Glycoside hydrolase family 18 chitinase	–	4	4	0	4	2	4	4	2	0	2	4	4
356467	7	4	GlcN6P deaminase (Nag1 family) involved in cell wall biogenesis	–	4	1	4	0	4	0	2	4	2	2	1	1
454418	9	6	X325 family LPMO-like protein	–	1	2	4	1	2	4	4	4	4	1	4	2

Table 25. Summary of developmental expression dynamics of CDE orthogroups of glucan remodeling related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			S. cerevisiae ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
S.c. 2686544	9	9	Beta-1,6-glucan synthase, GH16 family protein	–	0	4	0	4	0	4	4	4	4	4	4	4
546300	10	10	Beta-glucanase; GH5 family	–	1	4	4	4	0	4	4	4	4	4	4	4
464399	10	8	GH71 alpha-1,3-glucanase/beta glucuronidase family protein	–	4	4	4	4	4	4	2	0	4	0	4	2
373418	9	7	Glycoside hydrolase 3 family protein	–	4	1	4	4	4	2	4	0	4	0	2	4
449960	6	2	Probable glycosidase CRH2	YEL040W	4	2	2	1	2	1	1	1	4	2	1	1

Protein ID	FC>2	FC>4	Putative function	S. cerevisiae ortholog												
376392	11	6	Geranylgeranyl transferase type-1 subunit beta	YGL155W	4	4	4	4	2	4	4	2	2	1	2	2
414471	11	8	Glucan 1,3-beta-glucosidase I/II	YLR300W	4	1	2	4	4	4	4	2	4	4	4	2
389921	10	4	Family 17 glucosidase SCW10	YMR305C	4	4	2	4	4	2	2	2	2	1	2	1
169822	5	5	Beta-1,6-glucan synthase, GH16 family protein	–	4	0	4	4	1	4	1	0	4	0	0	0
381870	5	5	Beta-1,6-glucan synthase, GH16 family protein	–	4	4	1	4	4	1	0	0	0	4	1	0
411452	6	5	Beta-1,6-glucan synthase, GH16 family protein	–	4	1	0	4	0	4	4	4	2	1	0	1
502321	11	9	Beta-1,6-glucan synthase, GH16 family protein	–	4	4	4	4	2	4	4	4	4	1	4	2
373700	9	5	Beta-glucanase; GH5 family	–	4	2	4	4	2	2	2	0	0	0	4	4
496411	7	7	Beta-glucanase; GH5 family	–	4	4	4	1	4	0	4	4	1	1	4	1
505989	11	8	Beta-glucanase; GH5 family	–	4	2	4	4	1	4	4	4	2	4	4	2
22054	9	6	GH71 alpha-1,3-glucanase/beta glucuronidase family protein	–	4	4	2	4	1	2	4	2	0	0	4	4
492754	6	5	GH71 alpha-1,3-glucanase/beta glucuronidase family protein	–	4	0	0	4	0	4	0	1	4	2	4	0
463000	6	5	Glycoside hydrolase 128 family protein, putative β-1,3-glucanase	–	4	1	0	4	1	1	4	1	4	1	2	4
413757	7	5	Glycoside hydrolase 128 family, putative β-1,3-glucanase	–	4	1	4	0	2	4	2	0	4	0	4	1
476698	8	7	Glycoside hydrolase 17 family protein with putative β-1,3-glucan crosslinking activity	–	4	1	4	2	4	4	4	4	4	1	1	1
369876	10	4	Glycoside hydrolase 3 family protein	–	4	2	4	4	2	2	2	4	2	1	1	2
405828	9	8	Glycoside hydrolase 3 family protein	–	4	4	4	4	4	0	2	4	4	1	4	1
S.c. 2629974	8	6	Glycoside hydrolase 3 family protein	–	0	4	2	4	4	0	4	4	2	1	4	1
454337	1	1	Glycoside hydrolase family 55 putative β-1,3-glucanase	–	4	0	0	0	0	0	0	0	0	0	0	0
381637	11	9	Glycoside hydrolase family protein with putative glucan crosslinking activity	–	4	4	4	4	4	4	2	4	4	4	2	1
494762	10	3	PL8 family chondroitin lyase	–	4	2	2	0	2	2	0	4	2	2	4	2
543210	6	6	Alginate lyase	–	4	0	4	0	4	0	0	4	4	1	4	0
492834	11	9	Pectate lyase family protein	–	4	2	4	4	4	4	4	4	4	4	2	1
496344	7	7	Thaumatin/GH152 family protein	–	4	1	0	4	0	4	4	4	1	4	0	4
456054	10	8	WSC family protein , putative b-1,3-exoglucanase	–	4	0	4	4	4	4	4	2	4	2	4	0
528086	6	0	Rho family GTPase RHO1, Ras homolog gene family, member A	YPR165W	2	1	2	2	1	2	1	1	2	1	2	1

Table 26. Summary of developmental expression dynamics of CDE orthogroups of miscellaneous cell wall remodeling genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A.*

ID	Fold change		Putative function	*S. cerevisiae* ortholog	Species											
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
384105	9	6	Glucunoryl hydrolase; GH88	–	2	4	4	4	4	2	1	4	2	1	4	1
A.o. 254810	7	5	Glycoside hydrolase; family 27	–	0	4	0	4	2	0	4	0	4	2	4	0
A.o. 269656	8	6	Glycoside hydrolase; family 27	–	0	4	4	1	4	2	4	4	4	0	0	2
S.c. 2614029	7	6	Pectinesterase	–	0	4	4	1	1	2	1	4	1	4	4	4
470689	9	8	Beta-glucuronidase	–	4	0	4	4	2	4	4	4	4	1	4	0
S.c. 2633734	5	4	GT18 α-1,3(6)-mannosylglycoprotein β-1,6-N-acetyl-glucosaminyltransferase	–	0	4	4	0	1	2	0	0	1	0	4	4

Table 27. Summary of developmental expression dynamics of CDE orthogroups of multicopper oxidase genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change		Putative function	*S. cerevisiae* ortholog	Species											
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
408308	0	0	*C. cinerea* laccase lcc1	–	1	0	0	0	0	0	0	0	0	0	0	0
378517	0	0	*C. cinerea* laccase lcc15	–	1	0	0	0	0	0	0	0	0	0	0	0
443425	1	1	*C. cinerea* laccase lcc5	–	4	0	0	0	0	0	0	0	0	0	0	0
467038	0	0	*C. cinerea* laccase lcc6	–	1	0	0	0	0	0	0	0	0	0	0	0
89107	0	0	*C. cinerea* laccase lcc7	–	1	0	0	0	0	0	0	0	0	0	0	0
368135	10	4	*C. cinerea* laccase lcc15	–	4	4	4	2	2	2	0	2	2	0	4	2
353997	9	4	Cupredoxin	–	4	1	4	4	2	2	1	2	4	1	2	2
501548	5	3	Iron transport multicopper oxidase FET3	YMR058W	4	0	4	4	1	0	2	0	0	1	2	0
266892	2	1	*C. cinerea* laccase lcc3	–	2	0	0	0	0	0	4	0	0	0	0	0
440170	2	2	*C. cinerea* laccase lcc8	–	4	0	4	0	0	0	0	0	0	0	0	0
A.o. 257754	7	7	Cupredoxin	–	0	0	1	4	4	1	4	4	4	4	4	0
356026	9	7	Cupredoxin	–	4	1	0	4	4	4	4	2	4	2	4	1
409847	5	3	Cupredoxin	–	4	4	0	0	2	4	0	0	0	0	1	2
441442	9	4	Cupredoxin	–	4	1	1	4	0	2	4	2	4	2	2	2
451850	6	5	Cupredoxin	–	4	2	4	4	1	0	0	0	4	1	4	0
497434	6	5	Cupredoxin	–	4	0	0	1	4	4	4	0	4	0	2	0

Table 28. Summary of developmental expression dynamics of CDE orthogroups of expansin genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
439071	6	5	Cerato-platanin	–	4	4	0	4	4	0	2	0	0	0	0	4
544218	8	8	Cerato-platanin	–	4	0	0	4	0	4	4	4	4	1	4	4
461698	10	5	Expansin	–	4	2	4	2	4	4	4	1	0	2	2	2
467907	11	10	Expansin	–	4	4	2	4	0	4	4	4	4	4	4	4
496836	12	10	Expansin	–	4	4	4	4	4	2	4	4	4	2	4	4

Table 29. Summary of developmental expression dynamics of CDE orthogroups of miscellaneous cell wall related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
380322	6	4	*C. cinerea* Eln3	–	4	0	4	4	1	4	2	0	2	0	0	0
407604	5	5	Dye decolorising peroxidase	–	4	0	0	4	4	4	0	0	0	0	4	0
447925	10	7	*C. cinerea* Cfs1	–	4	4	4	2	1	1	4	4	4	2	4	2
469460	8	6	Cellobiose dehydrogenase	–	4	0	0	4	2	1	4	4	4	2	4	0
442333	11	7	Chloroperoxidase	–	4	2	4	4	4	2	4	2	4	1	4	2
479688	10	7	Chloroperoxidase	–	4	4	0	4	2	2	4	2	4	1	4	4
263775	9	7	Glyoxal oxidase	–	4	0	2	4	4	2	0	4	4	4	4	0
440836	10	10	Glyoxal oxidase	–	4	4	1	4	4	0	4	4	4	4	4	4
413665	8	5	Mitogen-activated serine/threonine-protein kinase SLT2	YHR030C	2	1	4	4	1	4	2	2	4	1	4	1

Table 30. Summary of developmental expression dynamics of CDE orthogroups of hydrophobin genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 -

developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
99661	7	7	Hydrophobin	–	1	4	0	0	4	1	4	4	0	4	4	4
419467	6	6	Hydrophobin	–	4	4	4	4	0	4	1	0	0	1	0	4

Table 31. Summary of developmental expression dynamics of genes belonging to conserved lectin orthogroups across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
353931	8	7	RicinB lectin protein	–	4	4	0	4	4	0	4	0	0	2	4	4
495335	9	9	RicinB lectin protein	–	4	1	4	4	4	4	0	4	4	1	4	4

Table 32. Summary of developmental expression dynamics of CDE orthogroups of cell surface protein encoding genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	*S. cerevisiae* ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
493867	6	1	Fasciclin	YOR100C	2	2	2	2	1	1	1	1	1	1	2	4
186809 S.c.	6	5	Con6 family cell surface protein	–	4	2	0	4	0	0	4	1	0	1	4	4
2624349	5	5	Con6 family cell surface protein	–	0	4	0	0	0	0	4	0	4	0	4	4
361136	7	4	Fasciclin	–	4	1	2	4	2	2	0	4	0	1	4	1
437384	7	5	Fasciclin	–	1	4	4	1	4	2	4	4	0	1	2	0
419688	7	5	PriA family protein	–	4	2	4	0	4	1	4	1	0	2	0	4
470234	6	6	Con6 family cell surface protein	–	1	4	4	0	0	0	4	0	4	0	4	4

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
359866	9	7	PriA family protein	–	1	4	4	2	4	2	4	4	4	1	4	1
S.c. 2681702	8	7	PriA family protein	–	0	4	2	4	0	4	4	0	4	0	4	4
501135	9	9	AA7 chitooligosaccharide oxidase	–	4	1	4	4	1	4	4	4	4	0	4	4
513409	9	7	AA7 chitooligosaccharide oxidase	–	4	1	4	4	4	4	4	1	2	1	4	2
A.o. 269613	6	5	Aspartic peptidase	–	0	0	1	4	4	4	0	4	0	2	4	0
442743	10	10	Aspartic peptidase	–	4	4	4	4	4	0	4	4	0	4	4	4
S.c. 75642	9	9	Aspartic peptidase	–	0	4	0	4	4	4	4	4	4	4	0	4
367835	10	8	Peptidase M20	–	4	2	4	4	0	2	4	4	4	0	4	4
S.c. 2547616	8	7	Serine carboxypeptidase	–	0	2	1	4	4	4	4	4	4	0	4	1
S.c. 2645761	8	7	Serine carboxypeptidase	–	0	4	4	4	4	1	2	4	4	4	1	1
462300	10	7	Serine carboxypeptidase S28	–	4	2	4	4	2	4	1	2	4	4	4	1

Table 33. Summary of developmental expression dynamics of CDE orthogroups of defense-related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
185637	9	5	CFEM domain containing protein	–	4	1	4	4	2	2	2	2	4	1	4	1
439390	9	6	CFEM domain containing protein	–	4	4	4	4	2	2	2	4	0	1	1	4
485770	7	5	RicinB lectin protein	–	4	2	4	0	0	1	4	2	0	0	4	4

Table 34. Summary of developmental expression dynamics of CDE orthogroups of terpene synthase genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change >2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
361541	8	5	Phenylalanine ammonia-lyase	–	4	1	4	2	4	4	0	0	2	0	4	2
386671	11	9	Phenylalanine ammonia-lyase	–	4	2	4	4	4	4	2	4	4	4	4	1
446258	11	8	Putative polyketide synthase	YER061C	4	4	4	4	4	1	2	2	2	4	4	4
394772	4	2	Putative terpene synthase	–	4	0	1	4	1	2	1	1	1	1	2	0
443805	3	2	Putative terpene synthase	–	4	0	2	0	0	0	0	1	0	0	4	0
495900	1	1	Putative terpene synthase	–	1	0	0	0	4	0	0	1	1	0	1	0
124670	5	2	Putative terpene synthase	–	4	0	1	4	1	2	0	2	1	0	2	0
30510	9	8	Putative terpene synthase	–	4	4	4	4	4	4	4	2	0	1	4	1
469738	10	8	Putative terpene synthase	–	1	1	4	4	4	4	4	2	4	4	4	2
201523	1	1	Putative terpene synthase	–	4	0	0	0	0	0	0	0	0	0	0	0

Table 35. Summary of developmental expression dynamics of CDE orthogroups of cytoskeleton-related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

	Fold change			*S. cerevisiae* ortholog	Species											
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
130811	3	1	LisH domain protein	–	4	1	1	2	0	2	1	1	0	1	0	1
504216	9	1	Misato tubulin	–	2	4	2	2	2	1	2	1	2	1	2	2
455153	4	0	Actin cytoskeleton-regulatory complex protein PAN1	YIR006C	2	1	2	1	1	2	1	1	1	1	2	1
496271	3	0	Adenyl cyclase associated protein	YNL138W	2	1	1	1	1	1	1	2	1	1	2	1
356686	6	1	Predicted Rho GTPase	–	2	1	2	1	1	2	2	1	4	1	2	1
381381	11	8	Agaricales-specific Rho GTPase Activating Protein	–	4	4	2	4	4	4	2	2	4	1	4	4
472493	4	0	Potential dynactin complex member	–	2	1	1	2	1	1	1	2	2	1	1	1
61699	5	1	Actin	YFL039C	1	2	2	2	2	1	1	1	1	1	4	1
258718	5	1	Myosin	YHR023W	2	1	1	1	1	2	2	2	1	1	4	1
473079	4	2	Myosin	YMR109W	4	1	2	1	1	2	1	1	1	1	4	1
419142	2	0	Myosin	YOR326W	2	1	1	1	1	1	1	1	1	1	2	1
461274	2	0	Protein SLA2	–	1	1	2	1	1	1	2	1	1	1	1	1
13965	2	0	Actin depolymerization related protein	YCR088W	2	1	1	1	1	1	1	1	1	1	2	1
505644	6	2	Cofilin-homolog protein	YDR063W	2	1	4	0	2	1	2	2	1	1	4	1
448482	6	3	Fimbrin	YDR129C	4	1	4	0	2	1	1	2	2	1	4	1
59959	3	0	Twinfilin	YGR080W	2	1	1	2	1	1	1	1	1	1	2	1
366601	4	0	F-actin capping protein	YIL034C	2	1	2	2	1	1	1	1	0	1	2	1
466954	1	0	Tropomyosin	YIL138C	2	0	0	0	0	0	0	0	0	0	0	0
443986	4	1	Actin capping protein	YKL007W	4	1	2	1	1	1	1	1	2	1	2	1
370292	2	0	Cofilin	YLL050C	1	1	1	2	1	1	1	1	0	1	2	1

484047	8	5	Profilin	YOR122C	4	1	4	4	2	1	2	4	2	1	4	0
444584	5	0	Actin filament binding protein	YPL242C	2	1	2	2	1	2	1	1	2	1	1	1
126636	3	0	Coactosin	–	2	1	1	2	1	1	1	1	0	1	2	1
355823	0	0	Villin/Gelsolin	–	1	0	0	0	0	0	0	0	0	0	0	0
376343	4	0	Villin/Gelsolin	–	1	1	2	2	1	2	1	2	1	1	0	1
358108	3	1	Arp2/3 complex member	YBR234C	4	1	2	1	1	1	1	1	1	1	2	1
130742	3	0	Arp2/3 complex member	YDL029W	2	1	2	1	0	1	2	1	1	1	1	1
498732	4	1	Arp2/3 complex regulating protein	YDL146W	4	1	1	1	1	1	2	1	1	2	2	1
114966	6	1	Arp2/3 complex member	YIL062C	2	1	4	1	2	2	2	1	0	1	2	1
57120	6	0	Formin	YIL159W	2	1	2	2	1	2	1	1	2	1	2	1
415488	3	0	Arp2/3 complex member	YJR065C	1	2	2	0	1	1	1	1	1	1	2	1
192562	2	0	Arp2/3 complex member	YKL013C	1	1	2	1	1	1	1	1	0	1	2	1
493018	2	0	Verprolin	YLR337C	2	1	0	1	0	1	1	1	0	0	2	1
421678	4	1	Arp2/3 complex member	YLR370C	1	1	2	1	1	1	2	1	2	1	4	1
148867	2	0	Formin	YNL271C	2	1	1	1	1	1	2	1	0	1	1	1
546441	1	0	Arp2/3 complex member	YNR035C	2	1	1	1	0	1	1	1	1	1	1	1
465398	6	0	WASP protein	YOR181W	2	1	1	2	0	2	1	1	2	1	2	2
544540	8	4	WASP protein	–	4	1	2	4	2	2	4	4	2	1	0	1
176636	5	2	Alpha tubulin	–	4	1	2	0	1	1	1	2	0	2	1	4
393528	6	1	Beta-tubulin	YFL037W	2	2	2	1	1	1	2	2	0	4	1	1
359665	4	1	Alpha tubulin	YML085C	4	2	0	0	1	1	2	1	1	1	2	1
519006	3	1	Beta tubulin	–	2	1	4	1	1	1	2	1	1	1	1	1
494087	3	1	Kinesin	–	4	1	1	1	1	1	2	1	2	1	1	1
3150	6	0	Kinesin	–	2	1	2	1	2	2	1	1	2	1	2	1
447928	3	0	Kinesin	–	2	1	1	1	1	1	1	1	2	1	2	1
461075	12	9	Kinesin	–	4	4	4	4	2	4	4	4	4	2	2	4
367694	9	5	Kinesin	YDR424C	4	2	2	0	2	4	2	4	1	1	4	4
410146	12	7	Kinesin	YPL155C	2	4	4	2	2	4	4	4	4	2	2	4
494863	10	4	Kinesin	–	4	2	2	4	2	2	4	4	2	1	2	1
113001	6	0	Microtubule plus end tracking protein	YER016W	2	1	1	2	1	1	2	2	2	1	2	1
527694	4	1	Tubulin specific chaperone.	–	2	1	2	2	1	1	1	1	1	1	4	1
441578	6	1	Gamma-tubulin comlex member	YHR172W	2	1	1	2	0	2	2	1	2	1	4	1
361158	7	1	Gamma tubulin	YLR212C	4	1	1	2	2	1	2	2	2	1	2	1
156038	5	2	Gamma-tubulin comlex member	YNL126W	4	1	1	2	1	1	2	2	1	1	4	1
357668	5	0	Gamma-tubulin comlex member	–	1	2	1	2	1	1	1	1	2	2	2	1
358462	6	1	Gamma-tubulin comlex member	–	4	2	1	2	2	1	2	2	1	1	1	1
463163	4	1	Gamma-tubulin comlex member	–	2	1	1	1	1	2	4	1	1	1	2	1
474549	4	0	Dynactin complex member protein	–	2	1	1	1	1	1	2	2	1	1	2	1
447285	9	2	Dynactin complex member protein	–	2	1	1	2	2	2	4	2	4	1	2	2
454161	4	0	Dynactin complex member protein	–	2	1	1	2	1	1	2	1	0	1	2	1
488749	3	0	Dynactin complex member protein	–	2	1	1	1	1	1	1	1	1	1	2	2
437053	3	1	Dynactin complex member protein	YOR269W	2	1	1	1	1	1	1	1	4	1	2	1
415573	9	8	Septin	–	4	4	4	4	4	4	0	4	4	1	0	2

ID	>2	>4	Function	Ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
455217	10	7	Septin	YCR002C	4	2	4	4	4	2	4	4	1	1	4	2
367587	10	6	Septin	YHR107C	4	2	4	4	4	2	4	2	2	1	4	1
481878	6	3	Septin	YJR076C	2	1	4	4	2	1	2	0	0	0	4	1
416864	8	4	Septin	YLR314C	4	2	4	4	2	1	2	2	1	1	4	1
103773	5	1	Septin	–	4	1	1	2	1	1	2	2	2	1	1	1
103809	5	2	Septin	–	4	1	2	1	1	2	2	1	4	1	1	1
413977	4	2	Tubulin binding cofactor A	–	2	1	1	1	1	1	1	2	4	1	4	1
362068	4	0	Tubulin binding cofactor D	–	2	1	1	2	1	1	2	1	2	1	1	1

Table 36. Summary of developmental expression dynamics of CDE orthogroups of Cop9 signalosome genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change		Putative function	*S. cerevisiae* ortholog	Species											
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
357329	9	5	Proteasome component	–	4	2	4	2	1	2	4	4	1	1	4	2
502637	6	0	Defective-in-cullin neddylation protein	–	1	0	1	2	2	2	1	1	2	1	2	2
496080	4	2	Nedd8-like ubiquitin	–	2	1	4	1	0	2	1	1	0	1	4	1
438017	7	1	COP9 signalosome complex subunit 5	YDL216C	4	1	2	2	1	2	2	2	2	1	1	1
497773	3	0	Ubiquitin carboxyl-terminal hydrolase 12	YJL197W	1	1	1	1	0	2	2	1	1	1	2	1
494197	4	0	Cop9 signalosome complex subunit 12	YJR084W	2	2	1	1	1	1	1	1	1	1	2	2
450685	3	0	NEDD8-conjugating enzyme UBC12	YLR306W	2	1	1	1	1	2	1	1	2	1	1	1
503638	7	0	NEDD8-activating enzyme E1 regulatory subunit	YPL003W	2	2	2	2	1	1	1	2	2	1	2	1
485679	8	3	NEDD8-activating enzyme E1 catalytic subunit	YPR066W	4	2	4	2	1	1	2	2	4	1	2	1
237692	9	4	Ubiquitin carboxyl-terminal hydrolase	–	4	1	4	2	1	2	4	2	2	2	4	1
439373	5	0	COP9 signalosome complex subunit 1	–	2	1	1	2	1	1	1	2	2	1	2	1
204652	3	0	COP9 signalosome complex subunit 3	–	2	1	1	1	1	1	2	1	2	1	1	1
444512	3	0	COP9 signalosome complex subunit 4	–	2	1	2	1	1	1	1	2	1	1	0	1
390754	4	0	COP9 signalosome protein	–	1	1	2	2	1	1	2	1	1	1	2	1
35248	4	0	COP9 signalosome subunit 6 protein	–	1	1	1	2	1	1	2	2	1	1	2	1
437024	4	1	Csn8 subunit protein	–	2	1	1	1	1	1	2	1	2	1	4	1
442909	2	0	Proteasome component	–	1	1	1	1	2	1	2	1	1	1	1	1
255029	2	0	COP9 signalosome complex subunit 2	–	2	0	1	1	1	1	1	1	2	1	1	1

Table 37. Summary of developmental expression dynamics of CDE orthogroups of protein ubiquitination and the proteasome related genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
354939	11	5	BTB/POZ domain containing protein	–	4	1	2	4	4	2	2	4	2	2	4	2
379603	9	8	BTB/POZ domain containing protein	–	4	4	1	4	0	4	4	2	0	4	4	4
383774	7	5	BTB/POZ domain containing protein	–	4	2	0	4	4	2	0	0	0	0	4	4
405851	7	5	BTB/POZ domain containing protein	–	4	0	0	2	2	4	4	0	4	0	4	0
446662	5	5	BTB/POZ domain containing protein	–	4	0	4	1	4	4	0	1	1	1	4	0
449824	8	8	BTB/POZ domain containing protein	–	4	4	1	4	0	4	4	1	4	1	4	4
451372	9	4	BTB/POZ domain containing protein	–	2	4	4	2	1	1	2	2	4	1	2	4
459680	11	8	BTB/POZ domain containing protein	–	4	2	4	4	4	4	4	4	4	1	2	2
374447	10	7	BTB/POZ domain containing protein	–	4	4	4	4	2	1	0	2	2	4	4	4
545078	5	1	BTB/POZ domain containing protein	–	2	2	1	2	0	2	1	1	4	1	1	1
358351	5	0	Orthologs of *A. nidulans* GrrA	–	2	1	1	2	2	1	1	1	2	1	2	1
447658	10	5	F-box protein	–	4	4	1	4	1	2	4	2	4	2	2	2
503496	9	5	F-box protein	–	4	1	4	4	2	2	4	4	2	1	2	1
145676	10	6	F-box protein	–	4	2	4	2	4	4	2	4	1	2	2	1
359613	10	6	F-box protein	–	2	2	4	4	4	2	4	4	4	1	2	1
470790	10	6	F-box protein	–	4	2	4	4	2	4	4	2	4	1	1	2
166096	9	7	F-box protein	–	4	0	0	4	2	4	4	4	4	2	4	0
356986	9	6	F-box protein	–	4	1	4	4	2	4	2	2	4	1	4	1
377171	7	2	F-box protein	–	2	2	2	4	1	1	1	2	4	1	1	2
382134	7	4	F-box protein	–	4	1	1	4	2	1	4	2	0	1	4	2
438595	7	7	F-box protein	–	4	1	0	4	1	4	4	0	4	1	4	4
445665	8	7	F-box protein	–	4	1	4	4	4	4	2	4	1	0	1	1
451935	7	6	F-box protein	–	1	4	1	4	0	4	0	4	0	4	4	2
471789	6	6	F-box protein	–	4	1	4	4	4	4	4	0	1	0	0	1
490594	7	3	F-box protein	–	4	1	1	2	2	1	4	1	2	1	4	2
501200	12	9	F-box protein	–	4	4	4	4	4	4	4	2	2	2	4	4
440772	11	7	RING-type zinc finger domain containing protein	–	4	4	2	4	2	2	2	0	4	4	4	4

ID	>2	>4	Putative function	S. cerevisiae ortholog	Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
364816	10	3	RING-type zinc finger domain containing protein	–	2	1	4	2	2	2	4	2	2	4	2	1
378120	8	8	RING-type zinc finger domain containing protein	–	4	4	0	4	4	1	4	1	4	0	4	4
455937	9	2	RING-type zinc finger domain containing protein	–	4	2	2	2	0	4	2	2	2	1	2	0
466739	10	7	RING-type zinc finger domain containing protein	–	4	2	2	4	4	1	4	4	4	2	4	1
467805	10	7	RING-type zinc finger domain containing protein	–	2	1	4	4	4	2	4	2	4	4	4	1
543850	11	9	RING-type zinc finger domain containing protein	–	4	4	2	4	4	2	4	4	0	4	4	4

Table 38. Summary of developmental expression dynamics of CDE orthogroups of transporter encoding genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

ID	Fold change		Putative function	S. cerevisiae ortholog	Species											
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
538158	8	5	Ammonium/urea transporter	–	4	1	0	4	0	4	2	2	4	1	4	2
404532	9	5	Major facilitator superfamily protein	–	4	1	0	2	4	2	4	0	4	2	4	2
492885	9	5	Major facilitator superfamily protein	–	4	1	2	2	1	1	2	4	4	2	4	4
S.c. 2537163	9	5	Multi antimicrobial extrusion protein	–	0	2	0	4	4	2	2	4	4	1	4	2
233439	9	6	Tetrapeptide transporter; OPT1/isp4 family	–	2	4	2	4	4	1	4	4	1	2	4	1
30849	11	6	10TM phosphate transporter	–	4	4	2	4	4	2	2	2	4	1	4	2
371400	11	9	Major facilitator superfamily protein	–	4	1	4	4	4	4	4	4	2	4	4	2
464027	7	6	Major facilitator superfamily protein	–	4	1	4	0	4	4	2	0	4	1	4	1
468563	6	6	Major facilitator superfamily protein	–	4	4	0	4	1	0	4	1	0	4	4	1
501434	9	9	Major facilitator superfamily protein	–	4	0	4	4	4	4	4	4	4	1	4	1
436779	10	0	Nuclear pore protein 84/107	–	2	2	2	2	1	1	2	2	2	2	2	2
S.c. 2283957	8	6	Sugar/inositol transporter	–	0	2	0	4	4	4	4	4	0	1	4	2
355160	10	9	Threonine/Serine exporter; ThrE	–	4	4	1	4	4	4	4	2	4	1	4	4
432854	10	6	Membrane transport protein family	YBR287W	4	2	2	4	4	2	4	4	1	1	2	4
449861	10	4	GABA-specific permease	YDL210W	2	1	4	2	4	1	2	4	4	2	2	2
428934	10	6	Sugar transporter STL1	YDR536W	4	2	4	4	4	4	4	1	2	1	2	2
422892	10	2	Purine-cytosine permease FCY2	YER056C	2	2	4	2	4	2	1	2	2	2	2	1
455616	5	5	Zinc-regulated transporter 1	YGL255W	4	1	4	4	0	4	4	1	1	1	1	1

ID			Name	YGene	1	2	3	4	5	6	7	8	9	10	11	12
465413	9	6	Urea transporter	YHL016C	4	1	0	4	4	4	4	2	4	1	2	2
496395	9	7	Oligopeptide transporter 1	YJL212C	4	4	2	4	4	1	4	2	4	1	4	0
502591	9	6	Polyamine transporter 1	YLL028W	4	2	4	1	1	2	4	4	4	0	2	4
360410	9	8	Thiamine pathway transporter THI73	YLR004C	4	1	4	4	1	4	4	4	4	1	4	2
495380	7	1	Transmembrane 9 superfamily member 1	YLR083C	4	2	2	2	1	1	2	1	0	1	2	2
223186	11	3	Inorganic phosphate transporter PHO84	YML123C	4	1	2	2	2	4	2	2	2	2	4	2
393558	10	6	Vacuolar basic amino acid transporter 1	YMR088C	4	4	2	4	2	1	4	2	1	4	2	4
502991	10	6	Major facilitator superfamily protein	YMR279C	4	4	4	4	2	2	4	2	0	1	4	2
421680	7	3	Calcium channel YVC1	YOR087W	2	2	2	4	1	1	4	1	1	1	2	4
542873	10	5	ATP-dependent permease PDR12	YPL058C	2	2	2	4	2	4	1	2	4	1	4	4
462297	10	2	Sulfate transporter YPR003C	YPR003C	2	1	2	4	2	2	2	2	4	1	2	2
S.c. 2501677	9	6	Aquaporin	–	0	4	4	4	2	1	4	2	4	1	4	2
A.o. 253413	6	6	Major facilitator superfamily protein	–	0	0	4	4	0	4	4	4	0	0	4	0
493842	9	6	Major facilitator superfamily protein	–	4	1	4	4	4	2	2	1	4	1	4	2
495288	8	6	Major facilitator superfamily protein	–	4	4	2	4	4	4	0	4	1	1	2	1
496145	9	7	Sugar/inositol transporter	–	4	4	1	4	2	1	4	4	4	2	4	1
498282	10	6	Major facilitator superfamily protein	–	4	2	2	4	1	4	4	2	2	4	4	1
474705	10	6	ABC transporter	–	4	2	0	4	2	4	4	1	2	2	4	4
357414	6	5	Major facilitator superfamily protein	–	4	1	4	4	1	0	0	4	2	1	4	1
420930	8	7	Major facilitator superfamily protein	–	4	4	4	4	2	4	4	1	0	1	4	1
463045	9	6	Major facilitator superfamily protein	–	4	2	4	4	2	4	0	2	0	1	4	4
465489	9	5	Major facilitator superfamily protein	–	4	1	4	4	0	1	2	2	4	2	4	2
488859	9	5	Mitochondrial substrate/solute carrier	–	4	2	4	2	4	1	2	1	4	1	4	2
451236	10	2	Secretory carrier membrane proteins (SCAMP)	–	2	2	1	4	2	2	4	2	2	1	2	2
374240	10	8	Sodium/calcium exchanger	–	4	1	0	4	2	4	4	2	4	4	4	4
450674	9	7	Sugar phosphate transporter	–	4	2	4	4	2	1	4	4	4	1	4	1
537526	10	7	Sugar phosphate transporter	–	4	4	2	4	4	2	2	1	4	1	4	4
358584	7	5	Sugar/inositol transporter	–	4	4	2	4	1	0	1	4	1	1	4	2
487841	11	8	Sugar/inositol transporter	–	4	4	4	4	2	4	4	2	2	1		4
502428	7	5	Sugar/inositol transporter	–	4	2	2	4	1	1	1	4	4	1	4	1
S.c. 2627703	6	5	Sugar/inositol transporter	–	0	2	4	4	0	0	4	0	4	0	0	4
397645	9	6	High glucose sensor RGT2	YDL138W	4	4	4	2	0	1	2	2	4	1	4	4
463015	8	7	Hexose transporter HXT10	YFL011W	4	4	0	4	4	0	4	4	0	1	4	2

Table 39. Summary of developmental expression dynamics of CDE orthogroups of stress response genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| ID | Fold change | | Putative function | *S. cerevisiae* ortholog | Species | | | | | | | | | | | |
	>2	>4			Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
475934	7	7	12 kDa heat shock protein	YFL014W	4	4	0	4	0	0	4	4	1	1	4	4
457314	8	5	ATP-dependent molecular chaperone HSC82	YMR186W	4	1	4	4	2	1	4	1	2	1	4	2
492695	6	5	Highly conserved plasma membrane protein	–	4	1	4	4	0	4	0	2	0	0	0	4
496981	9	7	Small heat shock protein HSP20	–	4	2	2	4	1	4	4	4	4	1	4	1
23471	11	5	Endoplasmic reticulum chaperone BiP	YJL034W	4	2	4	2	4	2	2	4	2	2	4	1
367102	11	6	Small heat shock protein HSP20	–	2	2	4	4	2	4	4	4	4	2	2	1
420157	8	5	Catalase	–	4	2	2	2	4	0	0	4	0	4	4	1
472224	11	7	Peroxisomal catalase A	YDR256C	4	2	4	4	2	2	2	4	4	1	4	4
432313	11	11	D-arabinono-1;4-lactone oxidase	YML086C	4	4	4	4	4	4	4	4	4	1	4	4
365550	7	2	D-arabinono-1;4-lactone oxidase	YMR041C	2	2	2	4	0	1	2	2	0	1	4	1
356399	11	8	Ergothioneine biosynthesis protein Egt1	–	4	2	4	4	4	4	4	2	2	1	4	4
355021	9	7	Peroxiredoxin	YBL064C	4	2	0	4	2	4	4	4	4	1	4	1
258252	5	4	Ergothioneine biosynthesis protein Egt2	–	4	1	4	1	4	2	1	4	1	1	1	1
28769	8	2	Glutathione peroxidase-like peroxiredoxin 2	YBR244W	4	1	4	2	2	2	2	2	0	1	2	1
415694	7	0	Monothiol glutaredoxin-6	YDL010W	2	1	2	1	1	1	1	2	2	2	2	2
441808	7	3	Glutaredoxin	YDR513W	2	2	2	4	1	4	4	1	1	1	2	1
378207	5	1	Monothiol glutaredoxin-4	YER174C	1	1	1	2	4	2	1	1	2	1	1	2
412620	5	2	Glutamate--cysteine ligase	YJL101C	4	1	4	2	1	2	2	1	1	1	1	1
370027	5	1	Glutathione synthetase	YOL049W	4	1	1	2	1	1	2	2	1	1	2	1
356216	7	1	Monothiol glutaredoxin-5, mitochondrial	YPL059W	2	1	2	1	0	2	2	2	2	1	4	1
499716	5	3	Gluthatione reductase	YPL091W	1	1	4	2	1	4	1	2	0	1	4	1
39451	7	3	Glutaredoxin	–	4	2	4	2	1	1	2	2	4	1	0	1
378471	4	0	Peroxiredoxin	–	1	1	1	2	2	1	1	2	0	1	2	1
379600	9	6	Peroxiredoxin DOT5	YIL010W	4	4	1	4	4	2	4	2	1	1	4	2
444300	10	4	Peroxiredoxin AHP1	YLR109W	4	1	4	4	2	2	2	2	2	1	4	2
468305	12	8	Peroxiredoxin TSA1	YML028W	4	2	4	4	2	4	2	4	4	4	4	2
437202	8	6	Peroxiredoxin	–	4	1	2	4	2	4	4	0	4	0	4	1
449056	9	7	Peroxiredoxin	–	4	2	4	4	2	1	4	4	1	1	4	4
S.c. 263240	5	3	Peroxiredoxin	–	0	2	0	0	0	0	4	4	0	0	4	2

Table 40. Summary of developmental expression dynamics of CDE orthogroups of ferric reductase genes across 12 species. Protein ID of a representative protein is given follows by the number of species in which the orthogroup is developmentally regulated at fold change 2 and 4 (FC>2 and FC>4, respectively). Putative function and ortholog in *S. cerevisiae* (if any) are also given. Abbreviations: 0-gene absent, 1-gene present but not developmentally regulated, 2 - developmentally regulated at fold change >2, 4- developmentally regulated at fold change >4. Species names are abbreviated as: Cc – *C. cinerea*, Aa – *A. ampla*, Ab – *A. bisporus*, Ao – *A. ostoyae*, Ca – *C. aegerita*, Lb – *L. bicolor*, Le – *L. edodes*, Lt – *L. tigrinus*, Mk – *M. kentingensis*, Pc – *Ph. chrysosporium*, Po – *P. ostreatus*, Sc – *S. commune*.

| | Fold change | | | S. cerevisiae ortholog | Species | | | | | | | | | | | |
ID	>2	>4	Putative function		Cc	Aa	Ab	Ao	Ca	Lb	Le	Lt	Mk	Pc	Po	Sc
354442	9	5	Ferric reductase	–	4	0	4	4	0	4	4	2	2	2	2	1
152927	9	5	Ferric reductase	YOR384W	4	1	1	2	4	2	4	2	4	4	2	1
260659	5	1	Ferric reductase	–	1	1	4	2	2	1	1	1	2	1	2	1
448679	3	1	Ferric reductase	–	1	1	2	1	1	1	1	1	4	1	2	1
461481	4	0	Ferric reductase	–	2	1	2	2	2	1	1	1	0	1	1	1