Exploring the Connection between Autophagy and Heat-Stress Tolerance in Drosophila melanogaster

Quentin Willot[1,3], Andre du Toit[2], Sholto de Wet[2], Elizabeth J. Huisamen[1], Ben Loos[2*], John S. Terblanche[1*]

1. Centre for Invasion Biology, Department of Conservation Ecology & Entomology, Stellenbosch University, Stellenbosch 7600, South Africa
2. Department of Physiological Sciences, Stellenbosch University, Stellenbosch 7600, South Africa
3. Current address: Zoophysiology, Department of Biology, Aarhus University, 8000, Aarhus-C, Denmark

Corresponding author: Quentin Willot (qwillot@bio.au.dk).

* These authors have contributed equally to this work

Abstract

Mechanisms aimed at recovering from heat-induced damage are closely associated with the ability of ectotherms to survive exposition to stressful temperatures. Among these mechanisms the respective contribution of autophagy, a ubiquitous stress-responsive catabolic process, has more recently come to light. By increasing the turnover of cellular structures as well as the clearance of long-lived protein and protein aggregates, the induction of autophagy has been linked to increased tolerance to range of abiotic stressors in diverse ectothermic organisms. Since our understanding of the relationship between autophagy and heat-tolerance currently remains limited in insect models, we hypothesized that (1) heat-stress would cause an increase of autophagy in *Drosophila melanogaster* tissues and (2) rapamycin exposure would trigger a detectable autophagic response in flies and increase their heat-tolerance. In line with our hypothesis, we report that flies exposed to heat-stress present signs of protein aggregation and appears to trigger an autophagy-related homoeostatic response as a result. We further show that rapamycin feeding causes the systemic effect associated with TOR inhibition, induces autophagy at least locally in the fly gut, and increase the heat-stress tolerance of individuals. This points toward a likely substantial contribution of this autophagy to cope with stressful temperatures in insects.

Keywords

 Autophagy, rapamycin, heat-tolerance, insects, heat-stress

1. Introduction

Without limited ability to regulate their internal body temperatures, terrestrial ectotherms' physiology remains largely contingent on external temperatures and the degree of climatic variability they experience [1, 2]. They thus rely on a wide array of molecular-level processes to cope with fluctuation of environmental conditions and increase their resilience to physiological stress [3, 4]. Stirred by the projected issues for ectotherms related to climate change and exposure to more frequent episodes of climatic extremes [5, 6], the exploration of these molecular basis in invertebrates has received renewed attention [1, 7]. Most species are indeed capable of transient to permanent shifts in heat hardiness following prior exposure to stressful temperatures, a process commonly referred to as thermal acclimation, which increase the odds of survival to subsequent stress exposure [6, 8, 9]. Several adaptative and acclimation responses at the molecular-level have been correlated with enhanced tolerance to temperatures-induced stress, including, but not restricted to, the overexpression of heat-shock proteins (Hsps) [10-12], increased clearance of cellular protein aggregates [13, 14], modulation of cell-membrane phospholipids [15-17], or improved capacity for ion balance defence [18-20].

Within the molecular toolbox available to invertebrates to cope with environmental and energetic stressors, the contribution of autophagy recently enjoyed renewed attention [21-23]. Essentially, autophagy is a bulk lysosomal degradation pathway conserved in all eukaryotic cells aimed at clearing cytosolic components, targeting mainly organelles, long-lived proteins, and protein aggregates [24-26]. In doing so, autophagy serves a "house-keeping" function and eliminates old or damaged cellular components that may lead to the disruption of homeostasis [24], which in a context of temperature-induced stress is paramount for organisms [27, 28]. Three types of autophagic pathways can be recognized depending on how intracellular material reaches the lysosome [29]. Briefly, during macroautophagy, large cytoplasmic components or large protein aggregates can be captured within an autophagosome, the functional unit of autophagy, which then merges with the lysosomal system for digestion of its content [29]. Lysosomes can also directly capture

smaller components and organelles through microautophagy [30]. Finally, denatured

peptides can selectively be targeted to reach the lysosome via the assistance of Hsp70-

family proteins, a process referred to as chaperone-mediated autophagy (CMA) [31]. In all

cases, these autophagic pathways lead to the degradation of cellular targets. From a

regulatory perspective, in *Drosophila melanogaster*, autophagy induction is under the control

of the Target of Rapamycin (TOR) pathway [25, 32-35]. TOR inhibition in response to stress

or exposure to rapamycin causes an cellular increase in autophagic flux and limits the

synthesis of proteins that are not involved in the stress response [32]. This both directly

mitigates the potential damages caused by abiotic stress while improving cellular metabolic

robustness and survival by maximizing ATP and metabolite availability [36]. Modulation of

autophagy, that relate to a more global responses from the TOR pathway, has therefore

ultimately linked to enhanced survival to various stressors in animals, including insects, such

as starvation, desiccation, toxics, heavy metals or oxidative damage [22, 23, 37-43]. In the

specific case of heat-stress, it has been demonstrated that activation of the transcription

factor HSF-1 which controls the cellular heat-shock response, and transcription of heat-

inducible Hsps, also induces autophagy in multiple tissues in *Caenorhabditis elegans* [14].

This helps reduce the accumulation of denatured protein and increases chances of survival

[14]. A link between rapamycin exposure and increased heat-tolerance in *D. melanogaster*

has also been more recently suggested [44]. Taken together, this provides strong

circumstantial evidence for the upregulation of autophagy to potentially be another pivotal

aspect underpinning heat tolerance in insects.

Here, we thus aimed at exploring the potential mechanisms bridging rapamycin

exposure, autophagy, heat-stress, and increased heat-tolerance in an insect model. We

report that a rapamycin-enriched diet does induce the systemic effect associated with TOR

inhibition in *D. melanogaster*, including impaired larval growth and increased formation of

lysosomal acidic compartments in the fly gut. Second, we show that exposition to heat-stress

further relates to accumulation of autophagy-related clearance proteins in *Drosophila* tissues,

where a combination of rapamycin-exposure and heat-stress causes maximal abundance of

GABARAPl1/2, Ref(2)P and Hsp70 in flies. Finally, we link rapamycin-exposure to enhanced

tolerance to, and recovery from, heat stress. Putting autophagy within the context of heat-

stress, we show that its induction can indeed play an important role nested within the cellular

mechanisms enabling increased insect resilience to temperatures.

2. Methods

2.1 Fly Stocks and rearing

Wild *Drosophila* flies were bait-trapped in April 2019 around the Stellenbosch University

campus, Western Cape, South Africa. Single female flies were isolated into 250 ml rearing

flask supplied with 50ml of Bloomington cornmeal diet [45] and left to lay eggs for one day.

Between 2 and 5 of the first emerging adults were used for species determination following

the key of Markow TAO'Grady P [46]. Stocks were kept at a density of ~100 flies per flask, at

a constant 25°C and a 12:12 light-dark cycle until experiments.

2.2 Rapamycin Treatment and heat-shock

For larval developmental assays, rapamycin (LC Laboratories) was dissolved in molecular

biology grade absolute ethanol (Sigma-Alldricht) and added to the Bloomington cornmeal

diet [45] at a concentration known to induce systemic effects in flies (50 or 200µM, [37]. For

control (0µM), ethanol alone was added. For adults involved in dissection, microscopy,

western blotting and heat-knockdown/recovery assays, eight days old flies were fed liquid

food (5% yeast extract, 5% sucrose, 1% ethanol, 1% brilliant blue FCF) with the appropriate

concentration of rapamycin (0, 50 or 200µM) for two days. Blue food commercial dye

(Brilliant blue FCF, Robertson's Blue Food Colouring) was used to visually confirm feeding.

For western blot analysis, rapamycin-treated adults were further either heat-shocked (37°C,

1h) or kept at 25°C to created 4 conditions within a full-factorial design: (1) Non rapamycin-

exposed/non heat-shocked flies (control, rapa-/HS-), (2) rapamycin-exposed/non heat-

shocked flies (rapa+/HS-), (3) rapamycin non-exposed/heat-shocked flies (rapa-/HS+), and

(4) rapamycin-exposed/heat-shocked flies (rapa+/HS+). Flies were immediately flash-frozen

after treatment and stored at -80°C for downstream analysis.

2.3 Developmental assays

Single 10-day old female flies were transferred into new vials containing the Bloomington cornmeal diet [45], supplemented with 0, 50 or 200μM rapamycin, and left to lay eggs for 24h. Development of larvae in the medium at 23°C was monitored daily, and number of pupations and eclosions recorded.

2.4 Heat-knockdown and recovery assays

For heat-knockdown experiments at 37°C, groups of 10 flies exposed to either 0, 50 or 200 μM rapamycin for 2 days were isolated into glass vials containing a moist cotton ball. Vials were then submerged in a programmable water bath (GD120 series, Grant Instrument Ltd) and the temperature inside the vials was monitored using 0.075 mm diameter thermocouples (Type T Thermocouple (Copper/Constantan), OMEGA Engineering, CT, USA) connected to a digital thermometer (Fluke 52-II Dual Input Digital Thermometer, WA, USA). Flies were exposed to a static 37°C heat-stress and checked every 15 minutes for loss of righting response, defined as the onset muscle coordination loss, when flies would fall on their back after a gentle vial shake and display total absence of movements. Between 110 and 130 flies were used for each experimental condition (0, 50 or 200 rapamycin-exposed) to extract knockdown curves at 37°C. For heat-knockdown experiments at 41°C, because the onset of muscle coordination would occur much faster, single flies were isolated into glass vials, exposed to 41°C, and monitored continuously. As soon as the loss of righting response occurred, each heat knocked-down fly was immediately placed back at room temperature (22°C) and monitored until recovery, defined as the time at which a fly could stand on its legs again, without stumbling (lack of coordinated movement) or falling over from external stimulation (gentle vial shaking). Between 45 and 51 flies were used to reconstruct knockdown/recovery curves at 41°C. Times to knockdown at 41°C and times to recovery were finally plotted against each other and the dataset screened for potential correlations.

2.5 Dissection, staining and confocal microscopy

To visualize the induction of autophagy in *D. melanogaster*, the gut of control or rapamycin-treated flies were dissected out in 4°C chilled PBS and then stained in a 10 µM Lysotracker Red:1µg/ml Hoechst solution (LysoTracker™ Red DND-99, Hoechst 33342 Solution, Thermofisher Scientific) for 3 minutes. Each preparation was washed three times with cold PBS, fixed overnight in 4% PFA at 4°C and mounted within the next 2 hours. Imaging was performed using a Zeiss LSM 780 ELYRA P.S.1 confocal microscope using a 20x objective. In each case, the midgut region between the proventriculus and Malpighian tubules were acquired. Laser power, digital gain and optical settings were kept constant between the images. Images were then binarized using a set threshold and pixel of high values quantified with the ImageJ software 1.50 [47]. The ratio of positive pixels per counting area was pooled for each individual and presented as an average made over between 7 and 13 individuals for each condition.

2.6 Western blotting

Following rapamycin/heat-shock treatments, whole fly bodies were collected, homogenized, and sonicated using a MixSonic (S-4000, ThermoFisher Scientific, Johannesburg, South Africa). The lysate was then centrifuged at 10 000 RPM for 20 min at 4°C (Spectrafuge, 24D, Labnet) to pellet nuclei and large cell fragments. The supernatant was collected into 1.5 mL Eppendorf tubes (#P2TUB003C-0001.5, Lasec, Cape Town, South Africa) and stored at -80°C. 15 mg of protein was mixed with Laemmli's buffer (1 mL Tis-HCl (0.5 M; pH 6.8), 1.6 mL 10% sodium dodecyl sulphate (SDS, #MKCG7687, Sigma-Aldrich, Johannesburg, South Africa), 0.8 mL glycerol (#1045422, Merck, Johannesburg, South Africa) and 0.4 mL of 0.05% bromophenol blue (#1041675, Merck) in 3.8 mL dH2O) in a 2:1 ratio. Samples were subsequently boiled at 95°C for 5 min. Proteins were separated on a Fast Cast TGX Stain-Free gel (#456-8084, Bio- Rad, Johannesburg, South Africa) consisting of a 12% resolving and 4% stacking component, respectively (1610174, Bio-Rad) and transferred to a PVDF membrane (#170-84156, Bio-Rad) using the Trans-Blot Turbo (#170-4155, Bio-Rad).

Membranes were blocked for 2 h using 5% non-fat milk made up in 1x TBS-T (137 mM

sodium chloride, 20 mM Tris, 0.1% Tween-20 at pH 7.6). Membranes were washed 3 times

for 5 min using TBS-T and incubated with primary anti-bodies overnight at 4°C. Primary anti-

bodies anti-Hsp70 (ab2787, Abcam, Germany), anti-Ref(2)P (ab178440, Abcam, Germany),

and anti GABARAP1+ GABARAP2 (ab109364 Abcam, Germany) were used at a 1:5000

dilution in TBS-T. Membranes were washed three times with TBS-T for 5 min and incubated

with peroxidase-linked anti mouse or rabbit IgG at a 1:10,000 dilution in TBS-T

(#7076/#7074S, Cell Signaling, Germany) for one hour at room temperature. Finally,

membranes were visualized using Clarity Western ECL substrate (#170-5061, Bio-Rad),

prepared at a 1:1 ratio and acquired using the ChemiDoc MP System (Bio-Rad).

2.7 Statistical analysis

Statistical analyses were performed using GraphPad Prism 9.01. Kruskal-Wallis tests were

followed by a Dunn's multiple comparisons post hoc test. One-way ANOVAs were performed

followed by a Fisher's LSD post-hoc test. Significance for log-rank tests analyses of Kaplan-

Meier survival and development curves, linear regressions, Kruskal-Wallis, and ANOVA tests

were accepted at $p<0.05$.

3. Results

3.1 Rapamycin feeding delays Drosophila larval development

Delayed larval development because of TOR inhibition is a documented effect of rapamycin

exposure in *D. melanogaster* (Zhang *et al.,* 2000). We thus performed larval growth assays

to confirm that, in our experimental setup, a rapamycin-enriched diet would induce the

expected systemic effects associated with TOR inhibition. Larvae were left to develop in

Bloomington cornmeal diet (Lewis, 1960) supplemented with either 0 (control), 50 or 200 µM

of Rapamycin. The number of pupations and imagoes hatching was recorded as a function

time at 23°C. 50 µM rapamycin exposure significantly delayed the onset of first larval

pupations by about 4 days and hatching by 6 days (Fig.3; Log-ranked test, p<0.001). Survival

from first instar to imago was estimated at 98% in control conditions (N=453) and dropped to

36% with 50 µM supplemented rapamycin food (N=129). No pupation events were observed

at 200 µM (all larvae remained either in second or third instars until mortality reached 100%

after 20 days).

3.2. Rapamycin feeding induces modulation of autophagy in the Drosophila *midgut*

Adult flies were fed control or rapamycin supplemented food at 50 and 200µM. Midguts of

flies were dissected and stained with Lysotracker prior to imaging. On average, there was a

significant increase in the number of Lysotracker-positive compartments observed in the

midgut of flies after rapamycin treatment, suggestive of drug uptake in the digestive track and

local autophagy induction (Fig.2A, B). Quantification of the average number of pixels above

intensity threshold in midgut areas of flies revealed of a dose-dependent response as a

function of the rapamycin concentration in their food (Fig. 2C).

3.3. Exposition to heat-stress and rapamycin modulates autophagy in fly tissues

A qualitative confirmation of engagement of the autophagosome/lysosome system was

performed on fly issues through western blot analysis (Fig 2A, N=4). Flies were either kept

on control conditions (rapa-/HS-) or exposed to rapamycin and/or heat-shock (rapa+/HS-,

rapa-/HS+, rapa+/HS+), and blotting was performed against: (1) GABARAPL1/2, which are

Atg8 protein family members involved in autophagosome formation [48], (2) Ref(2)P, the

Drosophila melanogaster ortholog of p62/sequestosome-1, involved in directing aggregated

protein cargoes towards autophagic/proteasomal degradation pathways by acting as

receptor [49, 50] and (3) HSP70-family proteins that are hallmark of proteotoxic stress and

involved both in the peptide-refolding machinery and CMA [31, 51]. GABARAPL1/2 levels

reached near-statistical significance following heat-stress in rapamycin-treated conditions

(p=0.053) (Fig. 3A). Ref(2)P abundance as compared to control was significantly increased

following heat-stress and the combination of heat-stress and rapamycin exposure (p<0.005).

Finally, the abundance of Hsp70 was not impacted by rapamycin alone but increased

following heat shock and the combination of rapamycin and heat-shock as compared to

control (*p*<0.05; Fig. 3A). Representative western blots are shown (Fig. 3B).

3.4. Rapamycin treatment delays time to knockdown in flies at 37°C

Eight-day old adult flies were fed a rapamycin-enriched diet at either 0 (control), 50 or 200

µM for two days, and then subsequently exposed to a static 37°C stress to induce a time-

dependent heat knockdown. The number of knocked down flies as a function of time was

recorded every 15 minutes (Fig. 4). Rapamycin exposure delayed the time for flies to enter

heat knockdown (Log-ranked test, p<0.001). 50 µM rapamycin exposure in food was already

sufficient to induce a plateau response.

3.5. Rapamycin treatment decreases time to recovery post knockdown

Eight-day old adult flies were fed a rapamycin-enriched diet at either 0 (control), 50 or 200

µM for two days and subsequently exposed to a static 41°C stress to induce a fast heat-

knockdown. Flies were monitored continuously and put back at room temperature (23°C) for

recovery after the loss of muscle coordination. The impact of rapamycin exposure on heat

knockdown at 41°C and its linked recoveries are presented in Fig. 5. For heat knockdown at

41°C, no effect of rapamycin exposure was observed (Fig. 5A). However, rapamycin

exposure reduced the time to recovery post knockdown in flies (Fig. 5B, Log-ranked test,

p<0.001). Finally, for each individual fly, single times to knockdown at 41°C were plotted

against single times to recovery, and significant regressions were screened both as a

function of rapamycin exposure (Fig. S1A) and on the global dataset (Fig. S1B; Table S1).

Only a weak correlation between time to heat knockdown and heat recovery was found for

flies fed with 50 µM rapamycin (R^2=0.22, p <0.001, Fig. S1A) and on the global dataset

(R^2=0.05, p<0.01; Fig. S1B). No significant correlations were recorded for other conditions

(Table S1).

4. Discussion

A number of markers to track autophagy have been validated in *Drosophila* so far, among which the Atg-8 family protein can potentially track all stages of the autophagy pathway by providing key information on autophagosome abundance [48, 52, 53] and the accumulation of ref(2)p which directs ubiquitinated proteins/proteins-aggregates towards the proteasome and/or selective autophagy [49, 50, 52, 53]. While these two markers target complementary, independent functions within the scope of autophagic activities [54], these are limited in obtaining reliable overviews of the autophagic rate (or flux) because of the inherent dynamic nature of this process [52, 55]. For example, the number of autophagosomes in tissues can increase either due to enhanced synthesis or due to accumulation brought by lysosomal dysfunction [56], and since autophagosome synthesis can be balanced by increased degradation, enhanced autophagic rate might result in constant Atg-8 level estimates at a snapshot despite the autophagy flux being upregulated towards a new steady state flux [52, 53, 57]. To circumvent these problems, the use of lysosomal inhibitors such bafilomycin or chloroquine are often used *in vitro* [52, 53, 55, 57]. However, these are poorly compatible with studies involving intact larvae or adult flies because of dire secondary effects that the general inhibition of lysosomal degradation has on the physiology of whole organisms [52]. Likewise, in live *D. melanogaste*r larvae and flies, inhibition of the TOR signalling pathway by rapamycin exposure bears global effects on metabolism via e.g., increased autophagic turnover, and limited *de-novo* protein synthesis. These effects have been documented from hindered larval growth and delayed imago eclosions [33], to increased lysotracker-positive punctate in the fly gut which is indicative of local autophagy induction, and increased survival to starvation and paraquat [37]. Thus, to link autophagy induction with enhanced survival to heat-stress within the boundaries of our own experimental system, it was imperative to first confirm that a rapamycin-rich diet would indeed result in the expected effects associated with the drug-exposure in both larvae and adults. Consistent with previous findings, we confirm that exposure to rapamycin does indeed significantly delay *D. melanogaster* larval development (Fig. 1), and that a 2-days exposure to rapamycin-enriched diet causes a dose-

effect response in the number of lysotracker-positive punctate in the fly midgut (Fig. 2).

Taken together, this provides strong circumstantial evidence that rapamycin-feeding as

performed during our treatments indeed causes the documented effects associated with

TOR inhibition and promotes autophagy, at the very least, locally in the fly gut.

Based on these results, we attempted to detect whether rapamycin-feeding would

cause autophagy induction at the systemic level in the fly, by tracking autophagic markers in

tissues through western-blotting (Fig. 3). We could not detect a statistical significant effect of

rapamycin exposure alone on GABARAPL1/2 levels (Atg-8 family member) [48] nor on

Ref(2)P abundance (Fig. 3A) in fly tissues. One likely potential explanation for this could be

found in the limited temporal resolution offered by our experimental design, as flies were

exposed for 2-days to the drug prior to experiments. Markers such has Ref(2)P have been

shown to decrease rapidly following autophagy induction but restored during such prolonged

stress durations [58], and it could well be that a 2-days rapamycin exposure period might

simply fail to track changes in marker abundance within their relevant timeframe. However,

the near-statistical increased of GABARAPL1/2 abundance (p=0.053), and the increased

abundance of Ref(2)P and Hsp70 (p<0.05) in either heat-stressed and/or heat-

stressed/rapamycin exposed flies offer interesting perspectives (Fig. 3). First, increased

GABARAPL1/2 levels following heat-stress and especially following heat-stress and

rapamycin exposure could potentially reflect autophagosome accumulation through

moderate autophagy induction [48]. Second, because Ref(2)p binds to ubiquitin-positive

aggregates before degradation through the autophagic/proteasome pathway [49, 50], its

accumulation can either reflect reduced degradation rate of protein aggregates or, more

likely under the circumstances, an expected increase in protein aggregation associated with

heat-stress [59-61]. Finally, and while we initially used well established Hsp70-family proteins

as an internal marker for successful heat-stress exposure in flies [10], its peaking levels

through the combination of heat-stress and rapamycin exposure could point toward an

attempt for chaperone-mediated autophagy to recycle denatured peptides before they

precipitate in toxic aggregates [31, 62]. While the full extent of these interpretations remains

merely speculative without further testing, and while we could not statistically confirm

autophagy induction in fly tissues following rapamycin-feeding alone probably because of

the practical time-resolution limitation that a 2-day drug exposure offers, our results

nonetheless suggest heat-stress does induce an expected proteotoxic stress in fly tissues

and, to a degree, that an autophagy-related homoeostatic response might be triggered as a

result thereof.

Finally, we show that rapamycin-exposure increases heat-tolerance in flies (Fig. 4).

Albeit to the best of our knowledge no direct link between rapamycin exposure, autophagy

induction, and heat-stress survival had been made in insect models prior to this work, this

echoes with previous findings involving rapamycin feeding to cause a slight but significant

increase of the critical thermal limits of *D. melanogaster* flies [44], and the overall

involvement of autophagy in surviving an array of deleterious conditions in ectotherms [21-

23, 37-43, 63, 64]. Solid support for a role of autophagy in mitigating heat-induced damages

and increasing survival in invertebrates is also recorded from *Caenorhabditis elegans* where

a causal link between autophagy induction, heat-acclimation and increased upper heat

tolerance was made [14]. In perspective, we thus first show in *D. melanogaster* flies that 50

µM rapamycin exposure is already sufficient to induce a plateau-response stress survival

under a 37°C stress exposure (Fig. 4, such delayed heat-knockdown was not observed in

experiments conducted at 41°C which likely underpins that knockdown occurred too fast in

this condition for us to detect a statistically significant response, Fig. 5A). Second, we were

able to detect a benefit of rapamycin exposure in reducing recovery time post heat-

knockdown (Fig. 5B). Although we provide evidence of a good correlation between these

phenotypic effects and autophagy induction in response to rapamycin and heat-stress

exposure in flies, we can only theorize on the responsible underlying molecular mechanisms

causing the observed heat-tolerance increase. TOR inhibition and autophagy induction have

been linked with a number of non-mutually exclusive processes related to cellular metabolic

resilience and repair mechanisms, including increased clearance of toxic protein aggregates

[14, 65], increase clearance of old and/or denatured peptides [62], optimization energy

production and cellular metabolic state to maximize survival [36] and finally, recycling of

damaged/surplus organelles such as mitochondria [23, 24]. Several of these could be jointly

involved in the observed phenotypic increase in heat-tolerance in flies, and further work

would be necessary to test and disentangle the respective contribution of each of them within

the framework of autophagy induction.

In conclusion, our results show that both rapamycin and heat-stress support degrees of

autophagic induction in *D. melanogaster*, and that rapamycin exposure correlates with an

increase in heat-tolerance and speed of recovery post-knockdown in adults. This highlights

the induction of autophagy as potentially key mechanism to increase resilience to

temperatures in insects. A deeper exploration of autophagy's role in relation to other stress

responsive pathways remains however necessary and could provide important perspectives

on the broader integrated cellular stress-response to temperatures in ectotherms.

Authors' contribution

Q.W., B.L. and J.S.T. conceived and planned the study. Q.W. and E.H. collected samples.

Q.W., A.dT and S.dW performed lab experiments, Q.W. and S.dW analysed the data. All

authors contributed to drafting the article, approved the final published version, and agree to

be held accountable for all aspects of the work

Acknowledgment

We thank Lize Engelbrecht and the team from the Central Analytical Facility of Stellenbosch

University for their help with confocal imagery analysis.

Preprint

A earlier version of this manuscript can be found in bioRxiv [66].

Competing interests

We declare we have no competing interests.

Funding

This work was supported by a Claude Leon Foundation post-doctoral fellowship (to Q.W.)

and a running cost funding (to J.T.) from the DSI-NRF Center of Excellence for Invasion

Biology.

References

1 González☐Tokman, D., Córdoba☐Aguilar, A., Dáttilo, W., Lira☐Noriega, A.,
Sánchez☐Guillén, R. A., Villalobos, F. 2020 Insect responses to heat: physiological
mechanisms, evolution and ecological implications in a warming world. *Biological Reviews*.
95, 802-821.

2 Woods, H. A., Dillon, M. E., Pincebourde, S. 2015 The roles of microclimatic diversity and
of behavior in mediating the responses of ectotherms to climate change. *Journal of Thermal*
Biology. **54**, 86-97.

3 Chown, S. L., Chown, S., Nicolson, S. 2004 *Insect physiological ecology: mechanisms and*
patterns. Oxford University Press.

4 Chown, S. L., Gaston, K. J. 2010 Body size variation in insects: a macroecological
perspective. *Biological Reviews*. **85**, 139-169.

5 Jørgensen, L. B., Ørsted, M., Malte, H., Wang, T., Overgaard, J. 2022 Extreme escalation of
heat failure rates in ectotherms with global warming. *Nature*. 1-6.

6 Weaving, H., Terblanche, J. S., Pottier, P., English, S. 2022 Meta-analysis reveals weak but
pervasive plasticity in insect thermal limits. *Nature communications*. **13**, 1-11.

7 Somero, G. N. 2020 The cellular stress response and temperature: Function, regulation, and
evolution. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*.
333, 379-397.

8 Sørensen, J. G., Kristensen, T. N., Overgaard, J. 2016 Evolutionary and ecological patterns
of thermal acclimation capacity in Drosophila: is it important for keeping up with climate
change? *Current Opinion in Insect Science*. **17**, 98-104.

9 Terblanche, J. S., Hoffmann, A. A. 2020 Validating measurements of acclimation for
climate change adaptation. *Current opinion in insect science*. **41**, 7-16.

10 Feder, M. E., Hofmann, G. E. 1999 Heat-shock proteins, molecular chaperones, and the
stress response: evolutionary and ecological physiology. *Annual review of physiology*. **61**,
243-282.

11 King, A. M., MacRae, T. H. 2015 Insect heat shock proteins during stress and diapause.
Annu. Rev. Entomol. **60**, 59-75.

12 Rinehart, J. P., Li, A., Yocum, G. D., Robich, R. M., Hayward, S. A., Denlinger, D. L.
2007 Up-regulation of heat shock proteins is essential for cold survival during insect
diapause. *Proceedings of the National Academy of Sciences*. **104**, 11130-11137.

13 Colinet, H., Nguyen, T. T. A., Cloutier, C., Michaud, D., Hance, T. 2007 Proteomic
profiling of a parasitic wasp exposed to constant and fluctuating cold exposure. *Insect*
biochemistry and molecular biology. **37**, 1177-1188.

14 Kumsta, C., Chang, J. T., Schmalz, J., Hansen, M. 2017 Hormetic heat stress and HSF-1
induce autophagy to improve survival and proteostasis in C. elegans. *Nature communications*.
8, 1-12.

15 Overgaard, J., Sørensen, J. G., Petersen, S. O., Loeschcke, V., Holmstrup, M. 2005
Changes in membrane lipid composition following rapid cold hardening in Drosophila
melanogaster. *Journal of insect physiology*. **51**, 1173-1182.

16 Overgaard, J., Tomčala, A., Sørensen, J. G., Holmstrup, M., Krogh, P. H., Šimek, P.,
Koštál, V. 2008 Effects of acclimation temperature on thermal tolerance and membrane
phospholipid composition in the fruit fly Drosophila melanogaster. *Journal of insect*
physiology. **54**, 619-629.

17 Slotsbo, S., Sørensen, J. G., Holmstrup, M., Kostal, V., Kellermann, V., Overgaard, J.
2016 Tropical to subpolar gradient in phospholipid composition suggests adaptive tuning of
biological membrane function in drosophilids. *Functional Ecology*. **30**, 759-768.

18 Andersen, M. K., Folkersen, R., MacMillan, H. A., Overgaard, J. 2017 Cold acclimation
improves chill tolerance in the migratory locust through preservation of ion balance and
membrane potential. *Journal of Experimental Biology*. **220**, 487-496.

19 Koštál, V., Renault, D., Mehrabianova, A., Bastl, J. 2007 Insect cold tolerance and repair
of chill-injury at fluctuating thermal regimes: role of ion homeostasis. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology. **147**, 231-238.

20 Overgaard, J., Gerber, L., Andersen, M. K. 2021 Osmoregulatory capacity at low
temperature is critical for insect cold tolerance. *Current Opinion in Insect Science*. **47**, 38-45.

21 Tachibana, S.-I., Matsuzaki, S., Tanaka, M., Shiota, M., Motooka, D., Nakamura, S., Goto,
S. G. 2020 The autophagy-related protein GABARAP is induced during overwintering in the
bean bug (Hemiptera: Alydidae). *Journal of economic entomology*. **113**, 427-434.

22 Teets, N. M., Denlinger, D. L. 2013 Autophagy in Antarctica: combating dehydration
stress in the world's southernmost insect. *Autophagy*. **9**, 629-631.

23 Lebenzon, J. E., Denezis, P. W., Mohammad, L., Mathers, K. E., Turnbull, K. F., Staples,
J. F., Sinclair, B. J. 2022 Reversible mitophagy drives metabolic suppression in diapausing
beetles. *Proceedings of the National Academy of Sciences*. **119**, e2201089119.

24 Kroemer, G., Mariño, G., Levine, B. 2010 Autophagy and the integrated stress response.
Molecular cell. **40**, 280-293.

25 Maruzs, T., Simon-Vecsei, Z., Kiss, V., Csizmadia, T., Juhász, G. 2019 On the fly: recent
progress on autophagy and aging in Drosophila. *Frontiers in Cell and Developmental*
Biology. **7**, 140.

26 Mizushima, N. 2007 Autophagy: process and function. *Genes & development*. **21**, 2861-
2873.

27 Bowler, K. 2018 Heat death in poikilotherms: Is there a common cause? *Journal of*
thermal biology. **76**, 77-79.

28 Goldberg, A. L. 2003 Protein degradation and protection against misfolded or damaged
proteins. *Nature*. **426**, 895-899.

29 Parzych, K. R., Klionsky, D. J. 2014 An overview of autophagy: morphology, mechanism,
and regulation. *Antioxidants & redox signaling*. **20**, 460-473.

30 Wang, L., Klionsky, D. J., Shen, H.-M. 2022 The emerging mechanisms and functions of
microautophagy. *Nature Reviews Molecular Cell Biology*. 1-18.

31 Kaushik, S., Cuervo, A. M. 2018 The coming of age of chaperone-mediated autophagy.
Nature reviews Molecular cell biology. **19**, 365-381.

32 Wullschleger, S., Loewith, R., Hall, M. N. 2006 TOR signaling in growth and metabolism.
Cell. **124**, 471-484.

33 Zhang, H., Stallock, J. P., Ng, J. C., Reinhard, C., Neufeld, T. P. 2000 Regulation of
cellular growth by the Drosophila target of rapamycin dTOR. *Genes & development*. **14**,
2712-2724.

34 Li, R., Xiao, Y., Li, K., Tian, L. 2022 Transcription and Post-translational Regulation of
Autophagy in Insects. *Frontiers in Physiology*. **13**, 825202-825202.

35 Chang, Y.-Y., Neufeld, T. P. 2009 An Atg1/Atg13 complex with multiple roles in TOR-
mediated autophagy regulation. *Molecular biology of the cell*. **20**, 2004-2014.

36 Loos, B., Engelbrecht, A.-M., Lockshin, R. A., Klionsky, D. J., Zakeri, Z. 2013 The
variability of autophagy and cell death susceptibility: Unanswered questions. *Autophagy*. **9**,
1270-1285.

37 Bjedov, I., Toivonen, J. M., Kerr, F., Slack, C., Jacobson, J., Foley, A., Partridge, L. 2010
Mechanisms of life span extension by rapamycin in the fruit fly Drosophila melanogaster.
Cell metabolism. **11**, 35-46.

38 Moore, M. N., Allen, J. I., Somerfield, P. J. 2006 Autophagy: role in surviving
environmental stress. *Marine Environmental Research*. **62**, S420-S425.

39 Minois, N., Carmona-Gutierrez, D., Bauer, M., Rockenfeller, P., Eisenberg, T.,
Brandhorst, S., Sigrist, S., Kroemer, G., Madeo, F. 2012 Spermidine promotes stress

resistance in Drosophila melanogaster through autophagy-dependent and-independent pathways. *Cell death & disease*. **3**, e401-e401.

Mizunoe, Y., Kobayashi, M., Sudo, Y., Watanabe, S., Yasukawa, H., Natori, D., Hoshino, A., Negishi, A., Okita, N., Komatsu, M. 2018 Trehalose protects against oxidative stress by regulating the Keap1–Nrf2 and autophagy pathways. *Redox biology*. **15**, 115-124.

Zhang, D.-W., Xiao, Z.-J., Zeng, B.-P., Li, K., Tang, Y.-L. 2019 Insect behavior and physiological adaptation mechanisms under starvation stress. *Frontiers in physiology*. **10**, 163.

Babczyńska, A., Nowak, A., Kafel, A., Łozowski, B., Rost-Roszkowska, M., Tarnawska, M., Augustyniak, M., Sawadro, M., Molenda, A. 2020 Autophagy: a necessary defense against extreme cadmium intoxication in a multigenerational 2D experiment. *Scientific reports*. **10**, 1-12.

Valko, A., Perez-Pandolfo, S., Sorianello, E., Brech, A., Wappner, P., Melani, M. 2022 Adaptation to hypoxia in Drosophila melanogaster requires autophagy. *Autophagy*. **18**, 909-920.

Rohde, P. D., Bøcker, A., Jensen, C. A. B., Bergstrøm, A. L., Madsen, M. I. J., Christensen, S. L., Villadsen, S. B., Kristensen, T. N. 2021 Genotype and Trait Specific Responses to Rapamycin Intake in Drosophila melanogaster. *Insects*. **12**, 474.

Lewis, E. 1960 A new standard food medium. *Drosophila information service*. **34**, 1-55.

Markow, T. A., O'Grady, P. 2005 *Drosophila: a guide to species identification and use*. Elsevier.

Schneider, C. A., Rasband, W. S., Eliceiri, K. W. 2012 NIH Image to ImageJ: 25 years of image analysis. *Nature methods*. **9**, 671-675.

Schaaf, M. B., Keulers, T. G., Vooijs, M. A., Rouschop, K. M. 2016 LC3/GABARAP family proteins: autophagy (un) related functions. *The FASEB journal*. **30**, 3961-3978.

Liu, W. J., Ye, L., Huang, W. F., Guo, L. J., Xu, Z. G., Wu, H. L., Yang, C., Liu, H. F. 2016 p62 links the autophagy pathway and the ubiqutin–proteasome system upon ubiquitinated protein degradation. *Cellular & molecular biology letters*. **21**, 1-14.

DeVorkin, L., Gorski, S. M. 2014 Monitoring autophagic flux using Ref (2) P, the Drosophila p62 ortholog. *Cold Spring Harbor Protocols*. **2014**, pdb. prot080333.

Rosenzweig, R., Nillegoda, N. B., Mayer, M. P., Bukau, B. 2019 The Hsp70 chaperone network. *Nature reviews molecular cell biology*. **20**, 665-680.

Mauvezin, C., Ayala, C., Braden, C. R., Kim, J., Neufeld, T. P. 2014 Assays to monitor autophagy in Drosophila. *Methods*. **68**, 134-139.

53 Lőrincz, P., Mauvezin, C., Juhász, G. 2017 Exploring autophagy in Drosophila. *Cells*. **6**,
22.

54 de Wet, S., Du Toit, A., Loos, B. 2021 Spermidine and rapamycin reveal distinct
autophagy flux response and cargo receptor clearance profile. *Cells*. **10**, 95.

55 Klionsky, D. J., Abdel-Aziz, A. K., Abdelfatah, S., Abdellatif, M., Abdoli, A., Abel, S.,
Abeliovich, H., Abildgaard, M. H., Abudu, Y. P., Acevedo-Arozena, A. 2021 Guidelines for
the use and interpretation of assays for monitoring autophagy. *autophagy*. **17**, 1-382.

56 Lim, J., Lachenmayer, M. L., Wu, S., Liu, W., Kundu, M., Wang, R., Komatsu, M., Oh, Y.
J., Zhao, Y., Yue, Z. 2015 Proteotoxic stress induces phosphorylation of p62/SQSTM1 by
ULK1 to regulate selective autophagic clearance of protein aggregates. *PLoS genetics*. **11**,
e1004987.

57 Loos, B., Toit, A. d., Hofmeyr, J.-H. S. 2014 Defining and measuring autophagosome
flux—concept and reality. *Autophagy*. **10**, 2087-2096.

58 Sahani, M. H., Itakura, E., Mizushima, N. 2014 Expression of the autophagy substrate
SQSTM1/p62 is restored during prolonged starvation depending on transcriptional
upregulation and autophagy-derived amino acids. *Autophagy*. **10**, 431-441.

59 Willot, Q., Gueydan, C., Aron, S. 2017 Proteome stability, heat hardening and heat-shock
protein expression profiles in Cataglyphis desert ants. *Journal of Experimental Biology*. **220**,
1721-1728.

60 Parsell, D. A., Kowal, A. S., Singer, M. A., Lindquist, S. 1994 Protein disaggregation
mediated by heat-shock protein Hspl04. *Nature*. **372**, 475-478.

61 Tyedmers, J., Mogk, A., Bukau, B. 2010 Cellular strategies for controlling protein
aggregation. *Nature reviews Molecular cell biology*. **11**, 777-788.

62 Cuervo, A. M., Wong, E. 2014 Chaperone-mediated autophagy: roles in disease and aging.
Cell research. **24**, 92-104.

63 Des Marteaux, L. E., McKinnon, A. H., Udaka, H., Toxopeus, J., Sinclair, B. J. 2017
Effects of cold-acclimation on gene expression in Fall field cricket (Gryllus pennsylvanicus)
ionoregulatory tissues. *BMC genomics*. **18**, 1-17.

64 Gerken, A. R., Eller, O. C., Hahn, D. A., Morgan, T. J. 2015 Constraints, independence,
and evolution of thermal plasticity: probing genetic architecture of long-and short-term
thermal acclimation. *Proceedings of the National Academy of Sciences*. **112**, 4399-4404.

65 Lamark, T., Johansen, T. 2012 Aggrephagy: selective disposal of protein aggregates by
macroautophagy. *International journal of cell biology*. **2012**,

66 Willot, Q., du Toit, A., Terblanche, J. S., Loos, B. 2021 Rapamycin induces autophagy and

increases heat tolerance in Drosophila melanogaster. *bioRxiv*.

Figures and tables

Figure 1. Number of *Drosophila melanogaster* pupation and hatching events as a function of

time at 23°C, for larvae fed either control (0 µM) or a 50 µM rapamycin supplemented diet.

Rapamycin exposure delayed both the onset of pupation and hatching by several days.

Different lowercase letters indicate significant differences between treatments (Log-rank

tests, P<0.001).

Figure 2. Representative confocal fluorescence images of adult *Drosophila* midguts stained

with LysoTracker Red (scale bar is 50 µM). **A.** Midgut of control flies. **B.** Midgut of rapamycin

treated flies (200 µM). **C.** Average number of pixels above intensity threshold over midgut

images of 7 to 13 individuals per conditions. Different lowercase letters indicate significant

differences between treatments (Kruskal-Wallis one-way analysis of variance followed by

Dunn's multiple comparisons post hoc test, $p < 0.01$).

Figure 3. Western blots analysis (N=4) following 200 µM rapamycin exposure of fly tissues directed against GABARAP, Ref(2)P and Hsp70. **A.** Relative abundance to control of GABARAPL1/2, Ref(2)P and Hsp70-family proteins following rapamycin exposure and/or heat-shock. GABARAP reached near-statistical significance following heat-stress in rapamycin treated conditions (p=0.053). Ref(2)P levels were significantly up-regulated following both heat-stress and heat-stress/rapamycin exposure conditions ($p<0.05$). Finally, Hsp70 levels were not impacted by rapamycin exposure alone, but increased following heat-shock to reach peak level through a combination of rapamycin exposure and heat-shock ($p<0.05$). Asterix represent significant differences relative to control (One-way analysis of

565   variance followed by a Fisher's LSD post-hoc test, $p<0.05$). **B.** Representative western blots

566   are shown.

Figure 4. Survival of *Drosophila melanogaster* flies that were exposed to a rapamycin-
enriched diet to a 37°C static heat stress (N=115 to 130 per condition). Rapamycin exposure
significantly delayed time-to heat knockdown in flies, and 50 µM exposure was sufficient to
induce a plateau response. Different lowercase letters indicate significant differences
between treatments (Log-rank tests, P<0.001).

576 **Figure 5.** knockdown and recovery curves of *D. melanogaster* flies that were exposed to a

577 rapamycin-enriched diet as a function of time at 41°C. **Fig. 5A.** Rapamycin exposure did not

578 delay time to heat knockdown in flies at 41°C**. Fig. 5B.** Rapamycin exposure decreased time

579 to recovery from heat knockdown of flies. Different lowercase letters indicate significant

580 differences between treatments (Log-rank tests, *p*<0.001).

582