Neurexins in serotonergic neurons regulate serotonin transmission and complex mouse behaviors

Amy Cheung[1-3], <u>Kohtarou Konno</u>[4], Yuka Imamura Kawasawa[5], Aya Matsui[6], Manabu Abe[7], Kenji Sakimura[7], Toshikuni Sasaoka[8], Takeshi Uemura[9,10], <u>Masahiko Watanabe</u>[4], and Kensuke Futai[1,2*]

1. Department of Neurobiology, University of Massachusetts Chan Medical School, 364 Plantation Street, Worcester, MA 01605-2324, USA

2. Brudnick Neuropsychiatric Research Institute, University of Massachusetts Chan Medical School, 364 Plantation Street, Worcester, MA 01605-2324, USA

3. Medical Scientist Training Program, University of Massachusetts Chan Medical School, 364 Plantation Street, Worcester, MA 01605-2324, USA

4. Department of Anatomy, Faculty of Medicine, Hokkaido University, Sapporo, Japan

5. Departments of Pharmacology and Biochemistry & Molecular Biology, Institute for Personalized Medicine, Pennsylvania State University College of Medicine, 500 University Drive, Hershey, PA 17033, USA

6. Vollum Institute, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR, 97239-3098, USA

7. Department of Animal Model Development, Brain Research Institute, Niigata University, Niigata 951-8585, Japan

8. Department of Comparative and Experimental Medicine, Brain Research Institute, Niigata University, Niigata 951-8585, Japan

9. Division of Gene Research, Research Center for Advanced Science, Shinshu University, Nagano 390-8621, Japan

10. Institute for Biomedical Sciences, Interdisciplinary Cluster for Cutting Edge Research, Shinshu University, Nagano 390-8621, Japan

* Corresponding author: Kensuke.Futai@umassmed.edu
Tel: 1-774-455-4318

31 Text Pages with 4 Figures, 1 Table, 1 Meta data supplement, and 6 Supplemental Figures
Number of Words: Abstract 174

Key Words:
neurexin, serotonin, serotonin transporter, serotonergic transmission, raphe nuclei, hippocampus, social behavior, mouse

Abstract

Extensive serotonin (5-HT) innervation throughout the brain corroborates 5-HT's modulatory role in numerous cognitive activities. Volume transmission is the major mode for 5-HT transmission but mechanisms underlying 5-HT signaling are still largely unknown. Abnormal brain 5-HT levels and function have been implicated in autism spectrum disorder (ASD). Neurexin (Nrxn) genes encode presynaptic cell adhesion molecules important for the regulation of synaptic neurotransmitter release, notably glutamatergic and GABAergic transmission. Mutations in Nrxn genes are associated with neurodevelopmental disorders including ASD. However, the role of Nrxn genes in the 5-HT system is poorly understood. Here, we generated a mouse model with all three Nrxn genes disrupted specifically in 5-HT neurons to study how Nrxns affect 5-HT transmission. Loss of Nrxns in 5-HT neurons reduced the number of serotonin neurons in the early postnatal stage, impaired 5-HT release, and decreased 5-HT release sites and serotonin transporter expression. Furthermore, 5-HT neuron-specific Nrxn knockout reduced sociability and increased depressive-like behavior. Our results highlight functional roles for Nrxns in 5-HT neurotransmission, 5-HT neuron survival, and the execution of complex behaviors.

Introduction

Serotonin (5-hydroxytryptamine, 5-HT) neurons in the raphe nuclei (RN) project their axons throughout the brain and modulate social interactions, stress responses, and valence among other processes. Abnormalities in 5-HT signaling have been extensively reported in neuropsychiatric disorders including depression, anxiety disorders, schizophrenia (SCZ), and autism spectrum disorder (ASD) (Lesch & Waider, 2012). 5-HT reaches postsynaptic specializations through volume transmission or at synapses and synaptic triads (Belmer et al., 2017). While much work has focused on deciphering receptor and reuptake dynamics in 5-HT signaling, the functional component important for 5-HT release remains undefined.

Nrxn genes (*Nrxn1-3*) encode alpha-, beta-, and gamma- (*α/βNrxn1-3*, *γNrxn1*) isoforms, and regulate synapse specification and function (Sudhof, 2017). Copy number variations and mutations in Nrxns are associated with ASD and SCZ (Sudhof, 2017). Numerous studies of α and βNrxn KO mice demonstrate impaired excitatory and inhibitory synaptic transmission (Sudhof, 2017). While Nrxns regulate fast synaptic transmission, no studies have examined the role of Nrxns in central neuromodulatory systems like the 5-HT system. Therefore, elucidating the impact of Nrxns in 5-HT transmission will allow a better understanding of pathophysiological mechanisms underlying neuropsychiatric disorders.

In this study, we investigated the functions of Nrxns in the 5-HT system by assessing signaling properties and behavior in 5-HT neuron-specific Nrxn triple knockout (TKO) mice. We demonstrated that the loss of Nrxn genes reduced 5-HT release and the number of 5-HT neurons and release sites in the mouse brain. Moreover, the lack of Nrxns in 5-HT neurons altered social behavior and depressive-like phenotypes. Our findings highlight Nrxns as functional regulators of 5-HT signaling and function.

Results

Expression of Nrxn isoforms in 5-HT neurons and validation of the Fev/RFP/NrxnTKO mouse

line

To characterize Nrxn genes expressed in 5-HT neurons, we analyzed scRNAseq data from a published

database consisting of over 900 single-cell 5-HT neuron datasets (~1 million reads/cell) which

generated 11 different 5-HT neuron clusters from the principal dorsal raphe nucleus (DRN), caudal

DRN (cDRN), and median raphe nucleus (MRN) (**Figure 1A**, **Figure 1-Meta data Supplement 1**) (Ren

et al., 2019). Transcriptional expression of six Nrxn isoforms (α-, βNrxn1/2/3) in 11 clusters indicated

that all 5-HT neuron clusters express at least one α- and βNrxn isoform (**Figure 1B, C**). These results

suggest that Nrxn proteins are expressed in 5-HT neurons in all RN regions.

To test the roles of Nrxns in 5-HT neurons, we generated 5-HT neuron-specific *Nrxn* triple

knockout (TKO) mice by crossing Fev-Cre, an ETS family transcription factor promoting 5-HT neuron-

specific Cre expression, tdTomato (RFP) reporter, and triple Nrxn1/2/3 floxed lines (Fev/RFP/NrxnTKO)

(Scott et al., 2005; Uemura et al., 2022). Cre-negative littermates and Fev/RFP mice were used as

control (Cntl) cohorts. The specific deletion of Nrxns in 5-HT neurons was confirmed by single-cell RT-

qPCR and -dPCR (**Figure 1D, Figure 1-Figure Supplement 1**). The Fev/RFP/NrxnTKO line was fertile

and viable and did not demonstrate obvious differences in gross appearance.

Reduced 5-HT release in Fev/RFP/NrxnTKO mice

While numerous studies indicate that Nrxns regulate fast neurotransmitter release including that of

glutamate and GABA, no studies have tested Nrxn function in central neuromodulatory systems. To

directly examine the role of Nrxn on 5-HT release, we measured 5-HT transients in the DRN and

hippocampus using fast-scan cyclic voltammetry (FSCV) (**Figure 2**). 5-HT release was recorded with a

voltage ramp delivered through carbon-fiber electrodes (**Figure 2A**). Current-voltage (CV) plots

displayed the expected currents for 5-HT oxidation and reduction peak potentials, indicating specificity

for 5-HT (**Figure 2B**).

5-HT transients were recorded in the DRN where 5-HT neurons are highly clustered in littermate

control and Fev/RFP/NrxnTKO mice (**Figure 2C-F**). Electrical stimulation was applied at two different

stimulus strengths to evoke 5-HT release in acute brain slices containing the DRN. To confirm that

electrically evoked FSCV transients were mediated by 5-HT release, the selective serotonin reuptake

inhibitor fluoxetine (FLX) and action potential inhibiting sodium channel blocker tetrodotoxin (TTX) were

applied (**Figure 2C, D, F**) (Carboni & Di Chiara, 1989). Importantly, 5-HT peak amplitude was

significantly reduced in Fev/RFP/NrxnTKO mice (**Figure 2E**). FLX caused a similar increase in 5-HT

transient area in each genotype, indicating that Nrxn TKO did not change transporter activity (**Figure**

2F). Next, we performed FSCV recordings in the dorsal hippocampal CA3 region to determine whether

differences in 5-HT release could be detected in a distal region receiving 5-HT fiber projections (**Figure**

2G-J). We observed robust suppression of 5-HT currents in Fev/RFP/NrxnTKO mice and no genotype-

specific differences in response to FLX. Taken together, these findings indicate that Nrxns are

important for 5-HT release.

Reduced 5-HT fiber density and active zone number in Fev/RFP/NrxnTKO mice

Next, we analyzed whether Nrxns are important for 5-HT innervation in brain regions that receive 5-HT

projections by analyzing RFP-positive fibers between Fev/RFP (Cntl) and Fev/RFP/NrxnTKO mice

(**Figure 3A-I**) (Awasthi et al., 2021). We found that RFP+ fibers were reduced in the hippocampus,

DRN and MRN of Fev/RFP/NrxnTKO mice relative to controls. Interestingly, no differences were seen

in the projections to the nucleus accumbens suggesting that 5-HT inputs are not globally altered. In

addition, serotonin transporter (SERT) expression in the brain was also reduced in brainstem and

hippocampus (**Figure 3-Figure Supplement 1**). These findings suggest that Nrxns selectively mediate

5-HT fiber area depending on the innervated circuit.

It has been reported that Nrxn TKO in cerebellar granule cells cause cell death (Uemura et al.,

2022), which raises the possibility that the reduced 5-HT fiber density is due to 5-HT neuron cell death.

To address this possibility, we compared the population of RFP-expressing 5-HT-positive 5-HT neurons

in the DRN and MRN between Fev/RFP (Cntl) and Fev/RFP/NrxnTKO mice at different ages (**Figure 3L-M**). The RFP/5-HT ratio between Cntl and Fev/RFP/NrxnTKO mice was comparable at P7 but reduced by increasing age specifically in Fev/RFP/NrxnTKO mice suggesting the postnatal loss of Cre-positive 5-HT neurons in Fev/RFP/NrxnTKO mice. To further elucidate the mechanism underlying reduced 5-HT release in Fev/RFP/NrxnTKO mice, we confirmed the expression of RIM, a major active zone protein, in synaptophysin-positive 5-HT terminals in ultra-thin sections (**Figure 3-Figure Supplement 2**). Next, we performed confocal imaging comparing the expression of RIM1/2 in 5-HT fibers in the hippocampus between Cntl and Fev/RFP/NrxnTKO mice (**Figure 3N, O**). The density of RIM1/2 in Fev/RFP/NrxnTKO mice was lower than that in Cntl. These results suggest that Nrxn TKO reduces 5-HT release by decreasing the number of 5-HT neurons and release sites.

Impaired social behavior in Fev/RFP/NrxnTKO mice

We investigated the behavior of adult Fev/RFP/NrxnTKO mice in a variety of assays. Basic activities, evaluated by locomotor activity, rotarod performance, and open field, did not differ between Fev/RFP/NrxnTKO mice and Cre-negative littermate controls (**Figure 4-Figure Supplement 1**). To examine the role of Nrxns in 5-HT system-related behaviors, we next assessed social behavior. Cntl and Fev/RFP/NrxnTKO underwent a direct social interaction test to examine naturally occurring interactions between a subject mouse and a juvenile stimulus mouse. In trial 1, the stimulus mouse was unfamiliar to the subject mouse. After 24 hours, the subject mouse was re-exposed to the same stimulus mouse (trial 2) (**Figure 4A**). Social investigation was measured across both trials and as a reduction in time that the subject mouse spent investigating the stimulus mouse in trial 2. We found that Fev/RFP/NrxnTKO mice spent less time exploring the stimulus mouse in trial 1 (**Figure 4B**) and differed in their investigation of the stimulus mouse across trials (**Figure 4C**) compared with littermate Ctnl mice. These results suggest that Fev/RFP/NrxnTKO mice have deficits in sociability. To further study sociability in the Fev/RFP/NrxnTKO mice, we performed a three-chamber social interaction test (3CST), in which a stimulus mouse was confined to a cylinder to limit direct interaction (**Figure 4-**

Figure Supplement 2A). Both littermate Cntl and Fev/RFP/NrxnTKO mice showed similar preferences

for a stimulus mouse than for a second empty cylinder in a different chamber (days 1 and 2) and for a

novel stimulus mouse than for the previously encountered juvenile conspecific (day 3) (**Figure 4-Figure**

Supplement 2B, C). Cntl and Fev/RFP/NrxnTKO mice showed different investigation behavior on day

1. However, there were no differences in exploration time between groups which contrasts the

sociability deficits observed in the direct social interaction test. Interestingly, one of the depression tests,

the forced swim test but not tail suspension test, revealed increased immobility behavior in

Fev/RFP/NrxnTKO compared with littermate Cntl mice (**Figure 4D-E**). Other tests addressing learning

and memory and repetitive behaviors displayed no abnormalities in Fev/RFP/NrxnTKO mice (**Figure 4-**

Figure Supplement 3). These results demonstrate that the absence of Nrxns in 5-HT neurons impairs

social behavior and moderately influences depression-related behavior.

Discussion

Nrxns regulate the release of fast neurotransmitters such as glutamate and GABA by coupling Ca^{2+}

channels to presynaptic release machinery (Sudhof, 2017). The study of cerebellins, secreted protein

binding partners of Nrxns, indirectly implicates Nrxns in serotonergic system development (Seigneur et

al., 2021). While cerebellin knockout was shown to alter DRN 5-HTergic circuits, Nrxn roles in central

neuromodulatory systems have never been addressed. Here we provide evidence that Nrxns control

neuromodulatory 5-HT release. We found that the DRN and hippocampus displayed > 40% reduction in

5-HT release in Fev/RFP/NrxnTKO mice. Fev expression begins in the embryonic stage and is found in

a subpopulation of 5-HT neurons (~ 60% of 5-HT neurons) (Scott et al., 2005) (**Figure 3M**). Therefore,

the impact of Nrxn TKO during development and non-Cre expressing 5-HT neurons should be noted.

However, consistent with the robust functional deficit observed in Fev/RFP/NrxnTKO mice, the

structural deficit identified by RFP-positive fiber and release site densities were significant (> 50%),

suggesting that the roles of Nrxns in the 5-HT system are cell survival and the formation of functional

components important for 5-HT release.

Given the predominance of non-junctional specializations, we speculate that Nrxns reside at 5-

HT release sites that lack a direct postsynaptic target. The ability of Nrxns to couple with release

machinery triggering 5-HT vesicle exocytosis and their roles in postsynaptic differentiation at synapses

are yet to be explored. Decreased 5-HT neuron number and release sites suggest that 5-HTergic Nrxns

contribute to cell survival and fiber formation. Indeed, Nrxns are important for cerebellar granule cell

survival through regulating the autocrine neurotrophic-factor (NTF) secretory machinery (Uemura et al.,

2022). Sparse pan-Nrxn deletion has also been shown to blunt inferior olive neuron climbing fiber

projections in the cerebellum while complete removal of Nrxns at climbing fiber synapses did not alter

climbing fiber axons but impaired synaptic transmission (Chen et al., 2017). It is interesting that the loss

of Cre+ 5-HT Nrxn TKO neurons is limited to the early postnatal stage. This may suggest that matured

5-HT neurons use the NTF supply from surrounding non-5-HT and Cre-negative 5-HT neurons. The

possibility of Cre expression altering properties related to 5-HT neuron function in place of Nrxn

deletion cannot be excluded. Further investigation is essential to understand the molecular

mechanisms underlying cell death in Nrxn TKO 5-HT neurons.

Overall behavioral phenotypes of Fev/RFP/NrxnTKO mice are milder than null Nrxn KO mouse

lines (Born et al., 2015; Dachtler et al., 2014; Etherton et al., 2009; Grayton et al., 2013). The observed

deficits in sociability in the direct social interaction test and in the depressive-related forced swim test

are relevant to ASD which often presents with co-occurring conditions. In contrast, the 3CST and tail

suspension tests were normal in Fev/RFP/NrxnTKO mice. The 3CST limits direct interaction with a

stimulus mouse and it is possible that the mode and novelty, as there were genotype differences

observed on initial encounter, of social interaction are critical to Fev/RFP/NrxnTKO mice. Additionally,

the forced swim test was performed following the tail suspension test, and it is possible that

Fev/RFP/NrxnTKO are more susceptible to stress rather than despair-associated coping responses.

The Fev-Cre line limits Cre expression in up to 60% of 5-HT neurons (**Figure 3M**), indicating a need for

a more 5-HT neuron-specific Cre line to fully understand the roles of 5-HTergic Nrxns in animal

behaviors.

Our results reveal that Nrxns expressed in midbrain 5-HT neurons are important for <u>cell survival</u>

and maintaining the presynaptic molecular function of 5-HT release sites. Further   studies   are

necessary to decipher Nrxn-mediated 5-HT release machinery, examine the consequences of Nrxn

deletion in RN-innervated circuits in other brain regions, and address whether 5-HT therapeutics can

improve behavioral deficits.

Materials and Methods

Animals

All experiments were conducted under approved animal protocols from the Institutional Animal Care

and Use Committee (IACUC) at the University of Massachusetts Chan Medical School. 5-HT neuron-

specific tdTomato mice (Fev/RFP) were generated by crossing $^{lox-STOP-lox}$tdTomato (Jax #007905) and

Fev^{cre} mice (ePet^{Cre}: Jax #012712) (Scott et al., 2005). <u>Both Fev-Cre and RFP lines were backcrossed</u>

<u>with C57BL/6J line for at least 10 generations.</u> Fev/RFP mice were crossed with Nrxn1$^{f/f}$/2$^{f/f}$/3$^{f/f}$ mouse

line (Uchigashima, Konno, et al., 2020; Uemura et al., 2022) to generate 5-HT neuron-specific triple

Nrxn knockout mouse line (Fev$^{Cre/lox-STOP-lox/lox-STOP-lox}$tdTomato/Nrxn1$^{f/f}$/2$^{f/f}$/3$^{f/f}$: Fev/RFP/NrxnTKO). The

Fev/RFP/NrxnTKO line was maintained by breeding Fev/RFP/NrxnTKO mice with <u>littermate</u> Cre-

negative ($^{lox-STOP-lox}$tdTomato/Nrxn1$^{f/f}$/2$^{f/f}$/3$^{f/f}$: Cntl) mice. Unless specified, Cre-negative littermates were

used as controls. Male mice were used in all experiments. For social behavioral experiments, juvenile

mice used as stimuli were 4- to 6- week-old male mice on a C57BL/6J background.

Mice were group housed (2-5 per cage) and maintained in ventilated cages with *ad libitum*

access to food and water on a standard 12-hour light/12-hour dark cycle (lights ON at 7 A.M.) in a

temperature-controlled (20-23°C) facility. One to two weeks prior to experimentation, mice were

acclimated to a reversed light/dark cycle (lights ON at 7 P.M.).

Single-cell RNA extraction and RT-qPCR <u>and -dPCR</u>

The whole procedure was done based on our recently developed protocol (Mao et al., 2018;

Uchigashima, Konno, et al., 2020; Uchigashima, Leung, et al., 2020). Briefly, cytosol from RFP+ 5-HT

neurons in the dorsal and median raphe nuclei were harvested from Fev/RFP (Cntl) and

Fev/RFP/NrxnTKO mice using the whole-cell patch-clamp approach. A SMART-Seq® HT Kit (TAKARA

Bio) was used to prepare the amplified cDNA templates following the manufacturer's instructions

(Uchigashima, Konno, et al., 2020; Uchigashima, Leung, et al., 2020). To assess Nrxn expression in

individual 5-HT neurons of control and Fev/RFP/NrxnTKO mice, the following TaqMan gene expression

assays (Applied Biosystems) with a FAM^{TM} dye on their probes were used: *Nrxn1* (Mm03808857_m1),

Nrxn2 (Mm01236856_m1), *Nrxn3* (Mm00553213_m1), *Tph2* (Mm00557715_m1) and *Gapdh*

(Mm99999915_g1). Additionally, a custom PrimeTime Std® qPCR Assay, which provides the same

type of quantitative PCR assay as TaqMan assays by utilizing hydrolysis probes in conjunction with

gene specific primer pairs, was designed for the TATA-box binding protein (TBP) housekeeping gene

with a HEX™ (VIC®) dye on its probe (Integrated DNA Technologies, Inc.). The assay consisted of a

forward primer (5'-GGGAGAATCATGGACCAGAACA-3'), a reverse primer (5'-

GGTGTTCTGAATAGGCTGTGG-3'), and a probe (/5HEX/CCTTCCACC/Zen/T_TATGCTCAGGGC

TT/3IABkFQ/). All the PCR reactions and analyses were performed blind to genotype. For the real-time

quantitative PCR (qPCR) analysis, StepOnePlus qPCR system (Applied Biosystems) and the relative

expression of *Nrxns* or *Tph2* were calculated as: Relative expression = $2^{Ct,Gapdh}/2^{Ct,Nrxns \text{ or } Tph2}$; Ct,

threshold cycle for target gene amplification, and presented as fold changes relative to that of Cntl. For

the digital PCR (dPCR) analysis, a QuantStudio 3D Digital PCR System and its accompanying

consumables were used and data were analyzed with QuantStudio 3D AnalysisSuite Cloud Software

(Thermo Fisher). The absolute copy number of Nrxn2, Nrxn3, and TBP were measured and the relative

abundance of Nrxn2 and Nrxn3 was calculated by normalizing their copy numbers to that of TBP.

Transcriptome analysis

Single-cell RNA-seq data was obtained from a recent publication (Ren et al., 2019). See Ren et al.,

2019 for the single-cell isolation and sequencing.

Data processing and Clustering: Datasets were downloaded from NCBI Gene Expression Omnibus

(GSE135132).   Reads   were   aligned   to   a   mouse   reference   transcriptome

(Mus_musculus.GRCm38.cdna.all.fa) using kallisto (Bray et al., 2016). Tximport R package (Soneson

et al., 2015) was used to summarize the reads to the gene level. Each isoform was summarized

manually to account for inclusion of spliced exons in the α or β Nrxn isoforms. The manually curated

transcript IDs are provided in **Table 1**. Gene count data were analyzed using Seurat R package v4.0.1

(Hao et al., 2021). After excluding cells with low sequencing depth (50,000 reads) and low number of

detected genes (cut-off was set at 7,500 genes), the remaining 945 cells were assigned to clusters

according to Ren et al. (Ren et al., 2019). Counts were normalized for each cell using the natural

logarithm of 1 + counts per 10,000 [ln(1+counts/10k)]. Cells were visualized using a 2-dimensional t-

distributed Stochastic Neighbor Embedding (tSNE) and violin plots. The R code is provided as codeR

(**Source Code File 1**).

269
Immunoblotting

Membrane fractions (P2 fraction) were used for immunoblottings. The whole experiments and analyses

were performed blind to genotype. Adult mouse brains (8-10 weeks old) were homogenized in ice-cold

buffer (5 mM HEPES (pH 7.4), 1 mM $MgCl_2$, 0.5 mM $CaCl_2$, 1 mM NaF, and protease inhibitors (Roche

cOmplete protease inhibitor: 05892970001) with a Teflon homogenizer (12 strokes). The homogenate

extract was centrifuged at low speed (1,400 x *g* for 10 min). The supernatant was centrifugated at

13,800 x *g* for 10 min. The supernatant was removed, and the P2 was resuspended in modified Radio

Immunoprecipitation Assay (RIPA) buffer (50 mM Tris-HCl pH 8.0, 150 mM NaCl, 0.1% Triton X-100,

0.5% sodium deoxycholate, 0.1 % SDS, 1 mM sodium orthovanadate, 1 mM NaF, Protease inhibitor

tablet). Homogenates (15 µg) were mixed with Sample Buffer (4% SDS, 20% glycerol, 0.004%

bromophenol blue, 0.125 M Tris-Cl, pH 6.8, 2.5% 2-mercaptoethanol) to undergo SDS-PAGE. Proteins

were transferred to PVDF membranes (0.2 µm, BioRad) and all remaining steps were performed at

room temperature. Membranes were blocked in 5% skim milk and 5% Bovine Serum Albumin in Tris-

Buffered Saline (TBS) for 1 hour and then washed in TBS with 0.1% Tween 20 detergent (TBST) for 10

min. Membranes were incubated with primary antibodies prepared in 1% skim milk and 1% Bovine

Serum Albumin in TBST for 2 hours (mouse anti-SERT 1:7000, MAb Technologies, ST51-2; rabbit

anti-Tph2 1:1000, Abcam, ab184505, RRID: AB_2892828; mouse anti-βActin 1:5000, Sigma, A1978,

RRID: AB_476692). After washing with TBST, membranes were incubated with secondary antibodies

for 1 hour (mouse and rabbit HRP, 1:1000, Millipore). Immunoblot signals were detected using an ECL

detection kit (PerkinElmer Life Sciences) and the Biorad Chemdoc system (Li-Cor). Quantification was

performed by Image J software.

Immunohistochemistry

All mice (8-10 weeks old) were transcardially perfused with ice-cold 4% paraformaldehyde (PFA) / 0.1

M phosphate buffer (PB, pH 7.4) under isoflurane anesthesia. Brains were dissected and post-fixed at

4°C in PFA for 2 hours, then cryo-protected in 30% sucrose / 0.1 M PB. Coronal 40 µm-thick brain

sections were cut on a cryostat (CM3050 S, Leica Biosystems). All immunohistochemical incubations

were carried out at room temperature. Sections were permeabilized for 10 min in 0.1% Tween 20 / 0.01

M phosphate-buffered saline (PBS, pH 7.4), blocked for 30 min in 10% normal donkey serum and

incubated overnight in anti-SERT (guinea pig, 1 µg/ml, Frontier Institute, HTT-GP-Af1400, RRID:

AB_2571777), anti-5-HT (goat, 1:1000, Abcam, ab66047, AB_1142794), anti-RIM1/2 (rabbit, 1:1000,

Synaptic Systems, Rb 140203, AB_887775), or anti-RFP (Rabbit, 1:1000, Rockland, 21513) antibodies.

The following day, sections were washed extensively then incubated in donkey anti-guinea pig-

Alexa488, goat-Alexa488, and mouse-Alexa405 antibodies for 2 hours at a dilution of 1:500 (Jackson

ImmunoResearch Laboratories). Sections were then mounted on slides (ProLong Gold, Invitrogen,

P36930) and viewed for acquisition and analysis. Immunohistochemistry for neurotransmitter release

machinery was confirmed using consecutive ultra-thin (100 nm) sections. Modified PFA (4% PFA and

0.1% glutaraldehyde in PB)-fixed brains were embedded in durcupan (Sigma) and consecutive ultra-

thin sections were prepared by Ultracut ultramicrotome (Leica Microsystems). After etching with

saturated sodium ethanolate solution for 1–5 s, ultrathin sections on slides were treated with

ImmunoSaver (Nisshin EM) at 95°C for 30 min. Sections were permeabilized for 10 min in 0.1% Triton

X-100 / 0.01 M PBS (pH 7.4), blocked for 30 min in 10% normal donkey serum, and incubated

overnight in anti-SERT (goat, 1 µg/ml, Frontier Institute, MSFR103270, RRID: AB_2571776), RIM1/2

(Rabbit, Synaptic Systems, 140203, RRID: AB_887775, 1:1000), synaptophysin (guinea pig, 1 µg/ml,

Frontier Institute, MSFR105690, RRID: AB_2571843) and anti-RFP (guinea pig, Frontier Institute,

MSFR101410, RRID: AB_2571648) antibodies. The following day, sections were washed extensively

then incubated with a mixture of Alexa 488-, Cy3-, or Alexa 647-labeled species-specific secondary

antibodies for 2 hr at a dilution of 1:200 (Invitrogen; Jackson ImmunoResearch, West Grove, PA).

Sections were then mounted on slides (ProLong Gold, Invitrogen, P36930) and viewed for acquisition

and analysis.

Imaging

Image acquisition and analysis were performed blind to genotype. RFP density analysis: We analyzed

5-HT fiber innervation to the nucleus accumbens core and shell (NAcc, NAcSh; Bregma 1.18 ± 0.3 mm),

stratum oriens of the CA1, CA2, and CA3 subregions of the dorsal hippocampus (dCA1, dCA2, dCA3;

Bregma -1.46 ± 0.4 mm), stratum oriens of the CA1 subregion of the ventral hippocampus (vCA1; -3.16

± 0.4 mm), dorsal raphe nucleus (DRN; Bregma -4.56 ± 0.4 mm), and median raphe nucleus (MRN;

Bregma -4.5 ± 0.4 mm). To assess the density of RFP fiber inputs, four stained sections from each of

four Fev/RFP (Cntl) and Fev/RFP/NrxnTKO brains containing the nucleus accumbens, hippocampus,

or raphe nuclei were imaged (1024 x 1024 pixels) using a laser scanning confocal microscope

(LSM700, Zeiss) with a 63x oil-immersion objective (NA 1.4) at an optical zoom of 1.6 and Zen black

image acquisition software (Zeiss). For each brain, six randomly chosen 100x fields of view within the

region of interest were acquired with 21 z-stack steps at 0.35 µm spacing providing a 7 µm z-depth to

generate maximum intensity projections (MIPs) of the z-stacks. Images from all brains for a particular region were acquired using identical settings, and data analyses were performed using ImageJ as previously described (Werneburg et al., 2020). The six images from each region per animal were averaged to generate a mean for that region in each animal, with n=4 animals per genotype. A consistent threshold range was determined by subjecting images, blinded to genotype, to background subtraction and manual thresholding for each MIP within one experiment (IsoData segmentation method, 15-225). Using the analyze particles function, the thresholded images were used to calculate the total area of RFP fiber inputs.

RFP positive cell density analysis: We analyzed the relative expression of RFP- and 5-HT- positive neurons in the DRN (Bregma -4.6 ± 0.3 mm) and MRN (Bregma -4.5 ± 0.4 mm). The ratio was derived by dividing the number of RFP- and 5-HT- positive neurons (n=2-5 mice per cohort). To assess the density of RFP-positive 5-HT neurons, three stained sections from each of four or five Fev/RFP (Cntl) and Fev/RFP/NrxnTKO brains containing the DRN or MRN were imaged (1024 x 1024 pixels) using a laser scanning confocal microscope (LSM700, Zeiss) with a 20x water-immersion objective (NA 1.0) at an optical zoom of 0.5 and Zen black image acquisition software (Zeiss). For each section, two z-stack steps at 10-15 μm spacing were used to image the different 5-HT neuron populations. The ratios obtained from four to six images from each brain were averaged to generate a mean for the MRN and DRN of each animal. Images from the DRN and MRN were acquired using identical settings, and data analyses were performed using ImageJ as previously described (Uchigashima, Konno, et al., 2020).

RIM1/2 density analysis: We analyzed RIM1/2 expression in 5-HT terminals projected in the CA1 pyramidal layer. To assess the density of RIM1/2 puncta in RFP- and SERT- positive fibers, four stained sections from each of four Fev/RFP (Cntl) and Fev/RFP/NrxnTKO brains containing the dorsal hippocampus were imaged (1024 x 1024 pixels) using a laser scanning confocal microscope (Olympus, FV1200) with a 60x oil-immersion objective (NA 1.4). For each brain, 10-15 consecutive images were acquired with 0.2 μm steps and 18-33 RPF-/SERT-positive fibers were chosen. Because confocal microscopes can reach z-axis resolutions of 500 nm (Schermelleh et al., 2010), we measured the RIM

signal found in two or more consecutive RFP-/SERT-positive terminals. RPF-/SERT-positive fibers in

Cntl (81 fibers) and Fev/RFP/NrxnTKO (78) mice (n=3 mice per genotype), respectively, were

subjected to imaging analysis using MetaMorph software (Molecular Devices, Foster City, CA).

Electrophysiology

Slice preparation: Mice were anesthetized with isoflurane and decapitated. Brains were removed and

quickly cooled in ice-cold, pre-oxygenated (95% O_2/5% CO_2) aCSF containing the following (in mM):

126 NaCl, 2.5 KCl, 1.2 NaH_2PO_4, 1.2 $MgCl_2$, 2.4 $CaCl_2$, 25 $NaHCO_3$, 20 HEPES, 11 D-glucose, 0.4

ascorbic acid, pH adjusted to 7.4 with NaOH. Coronal slices (400 μm) containing the dorsal

hippocampus or DRN were prepared in ice-cold aCSF using a vibratome (VT1200 S, Leica Biosystems).

Slices were recovered in oxygenated aCSF at room temperature (22-24°C) for at least 1 hour before

use. Slices were then transferred to a recording chamber perfused at a rate of 1 ml/min with room

temperature, oxygenated aCSF.

Fast scan cyclic voltammetry: 5-HT measurements were performed in the radiatum of dorsal CA3 and

DRN. All experiments and analyses were performed blind to genotype. To detect 5-HT release, carbon

fiber electrodes were prepared as previously described (Hashemi et al., 2009; Matsui & Alvarez, 2018).

Carbon fiber electrodes consisted of 7-μm diameter carbon fibers (Goodfellow) inserted into a glass

pipette (A-M Systems, cat# 602500) with ~150-200 μm of exposed fiber. The exposed carbon fibers

were soaked in isopropyl alcohol for 30 min to clean the surface. Next, the exposed fibers were coated

with Nafion solution (Sigma) to improve detection sensitivity by inserting the carbon fiber into Nafion

solution dropped in a 3 mm diameter circle of twisted reference Ag/AgCl wire for 30 sec with constant

application of +1.0 V potential. The carbon fiber electrodes were air dried for 5 min and then placed in a

70°C oven for 10 min. A modified 5-HT voltage ramp was used, in which the carbon fiber electrode was

held at +0.2 V and scanned to +1.0 V, down to -0.1 V, and back to +0.2 V at 1,000 V/s delivered every

100 ms. Prior to recording, the electrodes were conditioned in aCSF with a voltage ramp delivered at

60 Hz for 10 min.

5-HT release was evoked with electrical stimulation (30 pulses, 30 Hz, 150 or 250 μA, 1 ms)

from an adjacent custom-made bipolar tungsten electrode every 10 min. The stimulating electrode was

placed ~100-200 μm away from the carbon fiber electrode (John et al., 2006). Recordings were

performed using a Chem-Clamp amplifier (Dagan Corporation) and Digidata 1550B after low-pass filter

at 3 kHz and digitization at 100 kHz. Data were acquired using pClamp10 (Molecular Devices) and

analyzed with custom written VIGOR software using Igor Pro 8 (32-bit; Wavemetrics) running mafPC

(courtesy of M.A. Xu-Friedman). Carbon fiber electrodes were calibrated with 1 μM 5-HT (Serotonin

HCl, Sigma) at the end of the experiment to convert peak current amplitude of 5-HT transients to

concentration. Three consecutive traces were averaged from each recording condition for analysis.

Background subtracted peak 5-HT transients and area under the curve were determined by subtracting

the current remaining after TTX (tetrodotoxin citrate, Hello Bio) application from the maximum current

measured. Dopamine HCl and fluoxetine HCl were obtained from Sigma.

Behavioral assays

All behavioral experiments were performed on male mice aged 8 weeks or older. Animals were

habituated to the testing room for at least 30 minutes before each experiment and all tests were

conducted under dim red-light conditions and white noise to maintain a constant ambient sound unless

otherwise noted. All experiments and analyses were performed blind to genotype. Animals were used

in only one behavioral paradigm for the direct social interaction test and fear conditioning. Mice

underwent tests for locomotion, anxiety, repetitive behaviors, and depression in the following order:

locomotor activity, open field, elevated plus-maze, grooming, marble burying, tail suspension test,

forced swim test. At least two days of rest were given in between all tests except for the tail suspension

test and forced swim test, during which mice were allowed to rest for at least 7 days in between.

Another cohort of mice completed the object interaction test followed by at least 2 days of rest before

undergoing rotarod. Behavioral testing apparatuses were cleaned with 0.1% Micro-90 (International

Products Corporation) between each mouse.

Locomotor activity: Locomotor activity of each mouse was tracked in photobeam activity chambers

(San Diego Instruments) for 90 minutes. Total horizontal movement was measured in 5-minute bins.

Rotarod: Motor coordination and balance were evaluated on a rotarod apparatus (San Diego

Instruments) with an accelerating rotarod test. In each trial, mice were habituated to a rod rotating at 6

rpm for 30 sec, then the rotation was increased to 60 rpm over 5 min. The latency to fall was measured

over five trials with an interval of 10 min between each trial. Any mice that remained on the apparatus

after 5 min were removed and their time was scored as 5 min.

Open field: Mice were placed in the center of an open arena (41 x 38 x 30.5 cm) facing the furthest wall

and allowed to freely explore the arena for 10 min. Time spent in the center of the arena (20.5 x 19 cm)

was automatically tracked with EthoVision XT 11.5 (Noldus).

Elevated plus-maze: The apparatus (Med Associates) consists of four arms, two enclosed with black

walls (19 cm high) and two open (35 x 6 cm), connected by a central axis (6 x 6 cm) and elevated 74

cm above the floor. Mice were placed in the intersection of the maze facing the furthest open arm and

allowed to freely explore the maze for 5 min. Time spent in the open and closed arms (index of anxiety-

like behavior) and total entries into the open and closed arms (index of locomotor activity) were

automatically measured with MED-PC IV software.

Direct social interaction test: The test was adapted from Hitti and Siegelbaum, 2014 (Hitti & Siegelbaum,

2014). Each mouse was placed individually into a standard mouse cage and allowed to habituate for 5

minutes followed by the introduction of a novel male juvenile mouse. The activity was monitored for 10

min and social behavior initiated by the subject mouse was measured by an experimenter sitting

approximately 2 meters from the testing cage with a silenced stopwatch. Scored behaviors were

described previously (Kogan et al., 2000): direct contact with the juvenile including grooming and

pawing, sniffing including the anogenital area and mouth and close following (within 1 cm) of the

juvenile. After 24 hours, the 10 min test was run again with the previously encountered mouse. Any

aggressive encounters observed between animals led to exclusion of the subject mouse from analysis.

Three-chamber social interaction test: Using a standard three-chamber design (Moy et al., 2004), the

apparatus consisted of a neutral central zone (18 x 40.5 x 22 cm) connecting two identical

compartments (each 19.5 x 40.5 x 22 cm) (Molas et al., 2017). Each of the outer compartments housed

a caged cylinder (8 cm diameter, 18 cm height; 1 cm between each vertical rod) to allow limited, but

direct physical contact between the subject and stimulus animals. Subject mice were placed in the

central zone and habituated to an empty apparatus for 5 min then briefly removed to place a juvenile

mouse under one of the two caged cylinders while the other caged cylinder remained empty

(counterbalanced). The subject mouse was then returned to the apparatus to freely explore all three

compartments for 5 min/day for three consecutive days using the same juvenile placed in the same

compartment. On Day 3, a novel juvenile mouse was placed in the empty caged cylinder. On days 1

and 2, the preference ratio was calculated as: (total social stimulus investigation – total nonsocial

stimulus investigation) / (total investigation). On day 3, the preference ratio was calculated as: (total

novel stimulus investigation – total familiar stimulus investigation) / (total investigation).

Tail suspension test: Mice were placed in a rectangular TST apparatus (28 x 28 x 42 cm) and

suspended by their tails which were wrapped in red lab tape at around 3/4 the distance from the base.

Movement was monitored for 6 min and the last 4 min were scored for immobility behavior (absence of

righting attempt).

Forced swim test: Mice were placed in a plexiglass cylinder (20 cm diameter, 40 cm height) containing

22±1°C water at a depth of 20 cm to prevent them from escaping or touching the bottom. Immobility,

measured as floating in the absence of movement except for those necessary to keep the head above

water, was measured during the last 4 min of a 6 min session. Following the test, mice were gently

dried with a clean paper towel and placed in a fresh cage on top of a heating pad for around 10-15

minutes after which they were returned to their home cage (Yankelevitch-Yahav et al., 2015).

Fear conditioning: The fear conditioning paradigm was adapted from *Herry et al.* (Herry et al., 2008).

Animals were not habituated in the testing room to avoid untimely association with auditory cues. Using

the ANY-maze fear conditioning system (Ugo Basile SRL), mice were placed in a fear conditioning

cage (17 x 17 x 25 cm) in a sound-attenuating box. The paradigm was performed under no light

conditions using two different contexts (context A and B). Mice underwent four phases with 24 hours in

between each session: habituation, acquisition, auditory recall, and contextual recall. On day 1 (context

A), mice were habituated to five 30 sec presentations of the CS+ and CS- (white noise) at 80 dB sound

pressure level. The inter-cue interval was pseudorandomized and each session with the CS+ or CS-

was 10 min. The presentation order of the CS+ and CS- trials were counterbalanced across animals.

On day 2 (context A), discriminative fear conditioning was performed by pairing the CS+ with a US (1

sec foot shock, 0.75 mA, 5 CS+/US pairings; intertrial interval: 22-125 s). The onset of the US

coincided with the last second of the CS+. On day 3, auditory recall was measured in context B with 5

presentations of CS+ and CS-. On day 4, the contextual recall was measured in context A for 10 min.

ANY-maze software was used to analyze freezing behavior (no movement detected for 1 sec), which

was scored automatically with an infrared photobeam assay in the fear conditioning cage.

Object interaction test: The test was adapted from *Molas et al.* (Molas et al., 2017). The apparatus

consisted of a custom-made white Plexiglass T-shaped maze (three arms, each 9 x 29.5 x 20 cm,

connected through a central 9 x 9 cm zone). Mice were placed in the start arm to habituate to the

apparatus for 5 min. Following habituation, they were presented with identical inanimate objects located

at opposite ends of the T-maze arms for 5 min/day on two consecutive days. On day 3, one of the

inanimate objects was replaced with a novel inanimate object placed in the same location

(counterbalanced) for 5 min. The preference ratio was calculated from day 3 data as: (total novel

stimulus investigation – total familiar stimulus investigation) / (total investigation).

Marble burying: 15 sterilized 1.5-cm glass marbles evenly spaced 2 cm apart in three rows of five were

placed in a standard mouse cage with a layer of bedding at a depth of 5-6 cm. A mouse was placed in

the cage for 30 min, then returned to its home cage. The number of marbles buried (2/3 of their depth

covered with bedding) was counted.

Grooming: Self-grooming behavior was scored as previously described (McFarlane et al., 2008; Yang

et al., 2007). Mice were habituated for 5 min in an empty mouse cage with no bedding, then grooming

behavior was observed for 10 min by an experimenter sitting approximately 2 meters from the testing

cage. Cumulative time spent grooming during the 10 min session was recorded using a silenced

stopwatch.

Statistical analyses

Results are represented as mean ± SEM. Statistical significance was set at $p < 0.05$ and evaluated

using paired and unpaired two-tailed Student's t-tests and two-way ANOVA or two-way repeated

measures ANOVA with Šidák's post hoc testing for normally distributed data. Mann Whitney U tests

were used for nonparametric data. Analyses were carried out with GraphPad Prism (GraphPad

Software). No statistical methods were used to determine sample sizes prior to the experiments.

Sample sizes were comparable to many studies using the similar experimental approaches and animal

models. All observed data points were used for statistics.

Acknowledgments

This work was supported by grants from the National Institutes of Health (R01NS085215 and

R01MH130582 to K.F., T32 GM107000 and F30MH122146 to A.C.), the Global Collaborative Research

Project of Brain Research Institute, Niigata University (G2905 to K.F.), and Riccio Neuroscience Fund

to K.F. The authors thank Ms. Naoe Watanabe for skillful technical assistance. We thank Drs. Veronica

Alvarez, Jacqueline N. Crawley, Gilles Martin, Motokazu Uchigashima, and David Weaver for

comments on an earlier draft of the manuscript.

508
509

References

Awasthi, J. R., Tamada, K., Overton, E. T. N., & Takumi, T. (2021, May 1). Comprehensive topographical map of the serotonergic fibers in the male mouse brain. *J Comp Neurol, 529*(7), 1391-1429. https://doi.org/10.1002/cne.25027

Belmer, A., Klenowski, P. M., Patkar, O. L., & Bartlett, S. E. (2017, Apr). Mapping the connectivity of serotonin transporter immunoreactive axons to excitatory and inhibitory neurochemical synapses in the mouse limbic brain. *Brain Struct Funct, 222*(3), 1297-1314. https://doi.org/10.1007/s00429-016-1278-x

Born, G., Grayton, H. M., Langhorst, H., Dudanova, I., Rohlmann, A., Woodward, B. W., Collier, D. A., Fernandes, C., & Missler, M. (2015). Genetic targeting of NRXN2 in mice unveils role in excitatory cortical synapse function and social behaviors. *Front Synaptic Neurosci, 7*, 3. https://doi.org/10.3389/fnsyn.2015.00003

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016, May). Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol, 34*(5), 525-527. https://doi.org/10.1038/nbt.3519

Carboni, E., & Di Chiara, G. (1989). Serotonin release estimated by transcortical dialysis in freely-moving rats. *Neuroscience, 32*(3), 637-645. https://doi.org/10.1016/0306-4522(89)90285-6

Chen, L. Y., Jiang, M., Zhang, B., Gokce, O., & Sudhof, T. C. (2017, May 3). Conditional Deletion of All Neurexins Defines Diversity of Essential Synaptic Organizer Functions for Neurexins. *Neuron, 94*(3), 611-625 e614. https://doi.org/10.1016/j.neuron.2017.04.011

Dachtler, J., Glasper, J., Cohen, R. N., Ivorra, J. L., Swiffen, D. J., Jackson, A. J., Harte, M. K., Rodgers, R. J., & Clapcote, S. J. (2014, Nov 25). Deletion of alpha-neurexin II results in autism-related behaviors in mice. *Transl Psychiatry, 4*, e484. https://doi.org/10.1038/tp.2014.123

Etherton, M. R., Blaiss, C. A., Powell, C. M., & Sudhof, T. C. (2009, Oct 20). Mouse neurexin-1alpha deletion causes correlated electrophysiological and behavioral changes consistent with cognitive impairments. *Proc Natl Acad Sci U S A, 106*(42), 17998-18003. https://doi.org/10.1073/pnas.0910297106

Grayton, H. M., Missler, M., Collier, D. A., & Fernandes, C. (2013). Altered social behaviours in neurexin 1alpha knockout mice resemble core symptoms in neurodevelopmental disorders. *PLoS One, 8*(6), e67114. https://doi.org/10.1371/journal.pone.0067114

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., 3rd, Zheng, S., Butler, A., Lee, M. J., Wilk, A. J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J., Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., Rogers, A. J., McElrath, J. M., Blish, C. A., Gottardo, R., Smibert, P., & Satija, R. (2021, Jun 24). Integrated analysis of multimodal single-cell data. *Cell, 184*(13), 3573-3587 e3529. https://doi.org/10.1016/j.cell.2021.04.048

Hashemi, P., Dankoski, E. C., Petrovic, J., Keithley, R. B., & Wightman, R. M. (2009, Nov 15). Voltammetric detection of 5-hydroxytryptamine release in the rat brain. *Anal Chem, 81*(22), 9462-9471. https://doi.org/10.1021/ac9018846

Herry, C., Ciocchi, S., Senn, V., Demmou, L., Muller, C., & Luthi, A. (2008, Jul 31). Switching on and off fear by distinct neuronal circuits. *Nature, 454*(7204), 600-606. https://doi.org/10.1038/nature07166

Hitti, F. L., & Siegelbaum, S. A. (2014, Apr 3). The hippocampal CA2 region is essential for social memory. *Nature, 508*(7494), 88-92. https://doi.org/10.1038/nature13028

John, C. E., Budygin, E. A., Mateo, Y., & Jones, S. R. (2006, Jan). Neurochemical characterization of the release and uptake of dopamine in ventral tegmental area and serotonin in substantia nigra of the mouse. *J Neurochem, 96*(1), 267-282. https://doi.org/10.1111/j.1471-4159.2005.03557.x

Kogan, J. H., Frankland, P. W., & Silva, A. J. (2000). Long-term memory underlying hippocampus-dependent social recognition in mice. *Hippocampus, 10*(1), 47-56. https://doi.org/10.1002/(SICI)1098-1063(2000)10:1<47::AID-HIPO5>3.0.CO;2-6

Lesch, K. P., & Waider, J. (2012, Oct 4). Serotonin in the modulation of neural plasticity and networks: implications for neurodevelopmental disorders. *Neuron, 76*(1), 175-191. https://doi.org/10.1016/j.neuron.2012.09.013

Mao, W., Salzberg, A. C., Uchigashima, M., Hasegawa, Y., Hock, H., Watanabe, M., Akbarian, S., Kawasawa, Y. I., & Futai, K. (2018, Jun 12). Activity-Induced Regulation of Synaptic Strength through the Chromatin Reader L3mbtl1. *Cell Rep, 23*(11), 3209-3222. https://doi.org/10.1016/j.celrep.2018.05.028

Matsui, A., & Alvarez, V. A. (2018, Jun 26). Cocaine Inhibition of Synaptic Transmission in the Ventral Pallidum Is Pathway-Specific and Mediated by Serotonin. *Cell Rep, 23*(13), 3852-3863. https://doi.org/10.1016/j.celrep.2018.05.076

McFarlane, H. G., Kusek, G. K., Yang, M., Phoenix, J. L., Bolivar, V. J., & Crawley, J. N. (2008, Mar). Autism-like behavioral phenotypes in BTBR T+tf/J mice. *Genes Brain Behav, 7*(2), 152-163. https://doi.org/10.1111/j.1601-183X.2007.00330.x

Molas, S., Zhao-Shea, R., Liu, L., DeGroot, S. R., Gardner, P. D., & Tapper, A. R. (2017, Sep). A circuit-based mechanism underlying familiarity signaling and the preference for novelty. *Nat Neurosci, 20*(9), 1260-1268. https://doi.org/10.1038/nn.4607

Ren, J., Isakova, A., Friedmann, D., Zeng, J., Grutzner, S. M., Pun, A., Zhao, G. Q., Kolluru, S. S., Wang, R., Lin, R., Li, P., Li, A., Raymond, J. L., Luo, Q., Luo, M., Quake, S. R., & Luo, L. (2019, Oct 24). Single-cell transcriptomes and whole-brain projections of serotonin neurons in the mouse dorsal and median raphe nuclei. *Elife, 8*. https://doi.org/10.7554/eLife.49424

599   Schermelleh, L., Heintzmann, R., & Leonhardt, H. (2010, Jul 26). A guide to super-resolution
600   fluorescence microscopy. *J Cell Biol, 190*(2), 165-175. https://doi.org/10.1083/jcb.201002018
601

602   Scott, M. M., Wylie, C. J., Lerch, J. K., Murphy, R., Lobur, K., Herlitze, S., Jiang, W., Conlon, R. A.,
603   Strowbridge, B. W., & Deneris, E. S. (2005, Nov 8). A genetic approach to access serotonin
604   neurons for in vivo and in vitro studies. *Proc Natl Acad Sci U S A, 102*(45), 16472-16477.
605   https://doi.org/10.1073/pnas.0504510102
606

607   Seigneur, E., Wang, J., Dai, J., Polepalli, J., & Sudhof, T. C. (2021, Dec). Cerebellin-2 regulates a
608   serotonergic dorsal raphe circuit that controls compulsive behaviors. *Mol Psychiatry, 26*(12),
609   7509-7521. https://doi.org/10.1038/s41380-021-01187-x
610

611   Soneson, C., Love, M. I., & Robinson, M. D. (2015). Differential analyses for RNA-seq: transcript-level
612   estimates improve gene-level inferences. *F1000Res, 4*, 1521.
613   https://doi.org/10.12688/f1000research.7563.2
614

615   Sudhof, T. C. (2017, Nov 2). Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural
616   Circuits. *Cell, 171*(4), 745-769. https://doi.org/10.1016/j.cell.2017.10.024
617

618   Uchigashima, M., Konno, K., Demchak, E., Cheung, A., Watanabe, T., Keener, D. G., Abe, M., Le, T.,
619   Sakimura, K., Sasaoka, T., Uemura, T., Imamura Kawasawa, Y., Watanabe, M., & Futai, K.
620   (2020, Dec 23). Specific Neuroligin3-alphaNeurexin1 signaling regulates GABAergic synaptic
621   function in mouse hippocampus. *Elife, 9*. https://doi.org/10.7554/eLife.59545
622

623   Uchigashima, M., Leung, M., Watanabe, T., Cheung, A., Le, T., Pallat, S., Dinis, A. L. M., Watanabe,
624   M., Kawasawa, Y. I., & Futai, K. (2020, Jun 19). Neuroligin3 splice isoforms shape inhibitory
625   synaptic function in the mouse hippocampus. *J Biol Chem, 295*(25), 8589-8595.
626   https://doi.org/10.1074/jbc.AC120.012571
627

628   Uemura, T., Suzuki-Kouyama, E., Kawase, S., Kurihara, T., Yasumura, M., Yoshida, T., Fukai, S.,
629   Yamazaki, M., Fei, P., Abe, M., Watanabe, M., Sakimura, K., Mishina, M., & Tabuchi, K. (2022,
630   Apr 5). Neurexins play a crucial role in cerebellar granule cell survival by organizing autocrine
631   machinery for neurotrophins. *Cell Rep, 39*(1), 110624.
632   https://doi.org/10.1016/j.celrep.2022.110624
633

634   Werneburg, S., Jung, J., Kunjamma, R. B., Ha, S. K., Luciano, N. J., Willis, C. M., Gao, G., Biscola, N.
635   P., Havton, L. A., Crocker, S. J., Popko, B., Reich, D. S., & Schafer, D. P. (2020, Jan 14).
636   Targeted Complement Inhibition at Synapses Prevents Microglial Synaptic Engulfment and
637   Synapse Loss in Demyelinating Disease. *Immunity, 52*(1), 167-182 e167.
638   https://doi.org/10.1016/j.immuni.2019.12.004
639

640   Yang, M., Zhodzishsky, V., & Crawley, J. N. (2007, Dec). Social deficits in BTBR T+tf/J mice are
641   unchanged by cross-fostering with C57BL/6J mothers. *Int J Dev Neurosci, 25*(8), 515-521.
642   https://doi.org/10.1016/j.ijdevneu.2007.09.008
643

644   Yankelevitch-Yahav, R., Franko, M., Huly, A., & Doron, R. (2015, Mar 2). The forced swim test as a
645   model of depressive-like behavior. *J Vis Exp*(97). https://doi.org/10.3791/52587
646
647

Figure 1 Nrxn expression in the raphe nuclei and confirmation of Nrxn deletion in Fev/RFP/NrxnTKO mice.

Single-cell transcriptomic analysis of Nrxn expression in dorsal raphe nucleus (DRN), caudal DRN (cDRN), and median raphe nucleus (MRN) 5-HT neurons. (A) Single-cell t-SNE plot of 999 Tph2-positive neurons representing 5-HT neurons analyzed from a recent publication (Ren et al., 2019). 11 transcriptomic clusters were correlated with anatomical location. DRN, cDRN, and MRN 5-HT neurons have six, one, and four different subclusters, respectively. (B) Single-cell t-SNE plots for the expression of six Nrxn isoforms in distinct 5-HT neuron populations. Cells are colored by log-normalized expression of each transcript, and the color legend reflects the expression values of ln(CPM+1). (C) Violin plots of six Nrxn splice isoforms in 11 clusters. Although there are cluster-dependent Nrxn expression patterns (e.g., α and βNrxn3 expression in MRN4), each Nrxn isoform is expressed in cells across all clusters. (D) Validation of the Fev/RFP/NrxnTKO mouse line. Expression of Nrxn genes in tdTomato-positive 5-HT neurons were compared between Fev/RFP (Cntl) and Fev/RFP/NrxnTKO mice. qPCRs against Nrxn 1, 2, 3, Tph2, and Gapdh (internal control) were performed for single-cell cDNA libraries prepared from tdTomato-positive neurons. Number of neurons: Fev/RFP (n = 23 neurons, 4 mice) and Fev/RFP/NrxnTKO (22, 4). Data are reported as mean ± SEM. n.s., not significant, * p < 0.05, *** p < 0.001, **** p < 0.0001; Mann-Whitney U test.

Figure 2

Figure 2 The lack of Nrxns in 5-HT neurons reduces evoked 5-HT release in the dorsal raphe nucleus and hippocampus.

(A) Voltage ramp protocol for detecting 5-HT with FSCV. Rapid cycling (2.2 ms) of the voltage ramp produced a current based on voltage-dependent oxidation (blue, oxi) and reduction (red) representing real-time 5-HT release. (B) Representative current traces when measuring 5-HT (1 μM, blue) or dopamine (DA; 1 μM, orange) using carbon-fiber microelectrodes calibrated on a 5-HT voltage ramp. Insets: Background-subtracted CV plots of the electrochemical current when 5-HT (left) or DA (right) was applied. (C, D) Electrically evoked 5-HT transients detected in the dorsal raphe nucleus (DRN) of littermate control (Cntl) (n = 6 slices, 3 mice) (C) and Fev/RFP/NrxnTKO (6, 3) (D) mice during perfusion with aCSF (left), SERT blocker fluoxetine (FLX, 10 μM, middle), and action potential-blocking TTX (1 μM, right). Top: representative CV plots of the electrochemical current in aCSF, FLX, and TTX. Middle: average 5-HT transients under each condition. Bottom: background-subtracted 3D voltammograms (false color scale) as a function of time (x axis, 20 s) and voltage applied (y axis). Note that TTX completely abolished peak transients indicating that FSCV measurements were mediated by action potential-induced 5-HT release. The expected oxidation peaks for 5-HT in the CV plots acquired from Cntl and Fev/RFP/NrxnTKO mice were similar. FLX application prolonged 5-HT transient area which verified that the FSCV transients detected 5-HT signals. Cntl and Fev/RFP/NrxnTKO mice showed similar responses to FLX. (E) Peak amplitude of 5-HT transients evoked using a 150 μA or 250 μA stimulation train (30 pulses, 30 Hz, 1 ms) in the DRN of Cntl (gray) and Fev/RFP/NrxnTKO (green) mice. 5-HT release was significantly different between genotypes (two-way repeated measures ANOVA: genotype main effect, $F_{1,10}$ = 12.02, ## p = 0.006; stimulation strength main effect, $F_{1,10}$ = 26.89, p = 0.0004; stimulus strength x genotype interaction, $F_{1,10}$ = 1.936, p = 0.2648). (F) Normalized area of 5-HT transients recorded in the DRN before and after FLX application. 5-HT transient area before and after FLX was significantly different (paired t-test: Cntl, t_5 = 11.04, p = 0.0001; Fev/RFP/NrxnTKO, t_5 = 4.111, p = 0.0093). (G, H) Electrically evoked 5-HT transients detected in the hippocampus of Cntl (n = 3) (G) and Fev/RFP/NrxnTKO (n = 3) (H) mice during perfusion with aCSF (left), SERT blocker FLX (10 μM, middle), and action potential-blocking TTX (1 μM, right). Top: representative CV plots of the electrochemical current in aCSF, FLX, and TTX. Middle: average 5-HT transients under each condition. Bottom: background-subtracted 3D voltammograms (false color scale) as a function of time (x axis, 20 s) and voltage applied (y axis). (I) Peak amplitude of 5-HT transients evoked using a 150 μA or 250 μA stimulation train in the hippocampus of Cntl and Fev/RFP/NrxnTKO mice. 5-HT release at each stimulation strength was significantly different between groups (two-way repeated measures ANOVA with Šidák's post hoc test following significant stimulation strength x genotype interaction, $F_{1,9}$ = 6.982, p = 0.0268; stimulation strength main effect, $F_{1,9}$ = 11.24, p = 0.0085; genotype main effect, $F_{1,9}$ = 24.56, p = 0.0008). (J) Normalized area of 5-HT transients recorded in the hippocampus before and after FLX application. 5-HT transient area before and after FLX differed in Cntl mice (paired t-test: Cntl, t_4 = 5.591, p = 0.005; Fev/RFP/NrxnTKO, t_5 = 1.244, p = 0.2686). n.s., not significant, * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001; Cntl vs. Fev/RFP/NrxnTKO: ## p < 0.01; aCSF vs. FLX: #### p < 0.0001.

Figure 3

Figure 3 The absence of Nrxns in 5-HT neurons decreases 5-HT fiber density, neuron number, and release sites.

(A-G) Representative 100x images of RFP-positive fibers in the nucleus accumbens core (NAcc) (A), nucleus accumbens shell (NAcSh) (B), dorsal hippocampal CA1-3 subregions (dCA1, 2, 3) (C-E), ventral hippocampal CA1 (vCA1) (F), dorsal raphe nucleus (DRN) (G), and median raphe nucleus (MRN) (H). (I) Quantification of the area of RFP-positive fibers revealed that 5-HT innervation was altered in specific brain regions in mice lacking Nrxns (n = 4 mice/genotype; for each mouse, 6 fields of view were averaged for each region). * $p < 0.05$, *** $p < 0.001$; unpaired two-tailed Student's t-test. Scale bars, 10 μm. (J-M) Relative proportion of RFP-expressing 5-HT neurons in the DRN and MRN between Fev/RFP (Cntl, top row) and Fev/RFP/NrxnTKO (TKO, bottom row) mice at different ages: postnatal day 7 (P7) (J), 8 weeks (K), and > 14 months old (L). (M) Quantification of RFP+ neurons as a fraction of 5-HT+ (green) neurons at three different postnatal ages. Note that the RFP+/5-HT ratio in TKO mice decreased with aging, suggesting that TKO in 5-HT neurons causes postnatal cell death. The numbers of Cntl and TKO mice were (postnatal age, number of mice): Cntl: P7 and 8w, 4 and 12-13M, 2; TKO: P7 and 8w, 4 and 12-13M, 3. * $p < 0.05$, two-way ANOVA. (N, O) (N) Three consecutive triple immunofluorescence images with 200 nm step for RFP, RIM1/2, and SERT obtained from Fev/RFP (Cntl, left) and TKO (right) brains. (O) Summary of RIM1/2 signal density between Cntl and TKO hippocampal CA3 region. *** $p < 0.0001$; unpaired two-tailed Student's t-test. Scale bars, 10 μm (A-H), 100 μm (J-L) and 2 μm (N).

Figure 4 The absence of Nrxns in 5-HT neurons impairs social behavior.
(A) Direct social interaction test using the same juvenile stimulus across two trials. (B) Littermate control (Cntl) (gray, n = 13) and Fev/RFP/NrxnTKO (green, n = 13) mice differed in their investigation of the juvenile stimulus across the two trials. Both groups spent less time exploring the juvenile stimulus during trial 2 than in trial 1 (two-way repeated measures ANOVA: trial main effect, $F_{1,24}$ = 7.855, ## p = 0.0099; genotype main effect, $F_{1,24}$ = 2.086, p = 0.1616; significant trial x genotype interaction, $F_{1,24}$ = 4.344, p = 0.0479). Šidák's post hoc test identified a significant genotype difference in investigation time in Trial 1 (** p = 0.0041). (C) The difference score of the interaction time across trials was reduced in Fev/RFPNrxnTKO mice (unpaired two-tailed Student's t-test: t_{24} = 2.084, * p = 0.0479). (D) Left, Fev/RFP/NrxnTKO mice displayed increased immobile time compared with Cntl in the forced swim test (unpaired two-tailed Student's t-test: t_{22} = 2.317, * p = 0.0302). Right, no difference in the number of immobile episodes was observed (unpaired two-tailed Student's t-test: t_{22} = 1.301, p = 0.2068). (E) Left, Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice showed no difference in time immobile in the tail suspension test (unpaired two-tailed Student's t-test: t_{22} = 1.070, p = 0.2964). Right, there was no difference between genotypes in the number of immobile episodes (unpaired two-tailed Student's t-test: t_{22} = 1.001, p = 0.3279).

Figure 1-Figure Supplement 1. Validation of the Fev/RFP/NrxnTKO mouse line. The expression of Nrxn2 and Nrxn3 genes in tdTomato-positive 5-HT neurons were compared between Fev/RFP (Cntl) and Fev/RFP/NrxnTKO mice by dPCRs. Endpoint multiplex PCR was performed for each sample-targeted gene and the absolute copy numbers of Nrxn 2 and Nrxn3 (recorded by FAMTM signal) were normalized to that of TBP (recorded by HEX™ (VIC®) signal). Number of neurons: Fev/RFP (n = 3, 3 mice) and Fev/RFP/NrxnTKO (3, 3). Data are reported as mean ± SEM. Mann-Whitney U test.

Figure 3-Figure Supplement 1. The absence of Nrxns in 5-HT neurons reduces SERT expression.

Two brain regions brainstem and hippocampus were dissected from Fev/RFP/NrxnTKO (TKO) and littermate control (Cntl) brains (n = 6/genotype) and immunoblotted with anti-SERT and β-actin (βAct) antibodies. The graph shows SERT immunoblot intensity divided by βAct. Data are reported as mean ± SEM. * p < 0.05, ** p < 0.01; unpaired two-tailed Student's t-test.

Figure 3-Figure Supplement 2 The expression of RIM1/2 in 5-HT fibers in 100 nm ultra-thin sections.

Triple immunofluorescence images for RIM1/2, synaptophysin (Syn), and SERT obtained from four consecutive 100 nm ultra-thin sections (1st slice: A-C, 2nd: D-F, 3rd: G-I, 4th: J-L) in Cntl dorsal hippocampal CA3 region. The boxed area in low magnification images (A, D, G, and J) is enlarged in the lower panels (B, C, E, F, H, I, K, and L). Arrows indicate RIM1/2 immunofluorescent puncta associated with Syn and SERT. Scale bars, 2 μm.

Figure 4-Figure Supplement 1 5-HT neuron-specific Nrxn TKO does not alter basic behavioral activities.

(A) Left, Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice showed no differences in horizontal locomotor activity (two-way repeated measures ANOVA: genotype main effect, $F_{1,22}$ = 0.4666, p = 0.5017; time x genotype interaction, $F_{17,374}$ = 0.9467, p = 0.5188). Right, cumulative locomotor activity did not differ between Cntl and Fev/RFP/NrxnTKO mice over the 90 min period (unpaired two-tailed Student's t-test: t_{22} = 0.6831, p = 0.5017). (B) Cntl (n = 16) and Fev/RFP/NrxnTKO (n = 13) mice showed no differences in latency to fall over 5 trials of accelerating rotarod (two-way repeated measures ANOVA: genotype main effect, $F_{1,27}$ = 0.2204, p = 0.6425; trial x genotype interaction, $F_{4,108}$ = 0.2161, p = 0.929; trial main effect: $F_{(2.943,79.47)}$ = 37.52, p<0.0001). (C) Left, Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice showed no differences in time spent in the center of the open field arena (unpaired two-tailed Student's t-test: t_{22} = 0.04652, p = 0.9633). Right, there were no differences for the distance traveled during open field (unpaired two-tailed Student's t-test: t_{22} = 0.1053, p = 0.9171). (D) Left, the time Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice spent in the open arms during the elevated-plus maze was similar between genotypes (unpaired two-tailed Student's t-test: t_{22} = 1.158, p = 0.2592). Right, Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice both demonstrated similar total arm entries (unpaired two-tailed Student's t-test: t_{22} = 0.3625, p = 0.7204). n.s.: not significant.

Figure 4-Figure Supplement 2 Social behavior in the three-chamber social interaction test is preserved in Fev/RFP/NrxnTKO mice.

(A) Protocol for three-chamber social interaction test. (B) Littermate Cntl (n = 13) and Fev/RFP/NrxnTKO (n = 13) preferred the social stimulus over the nonsocial stimulus on day 1 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,24}$ = 77.08, p < 0.0001; stimulus x genotype interaction, $F_{1,24}$ = 0.6324, p = 0.4243) and day 2 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,24}$ = 83.16, p < 0.0001; stimulus x genotype interaction, $F_{1,24}$ = 0.03609, p = 0.8509). Both groups preferred the novel stimulus over the familiar stimulus on day 3 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,24}$ = 18.3, p = 0.0003; stimulus x genotype interaction, $F_{1,24}$ = 0.4758, p = 0.4969). Genotype differences were observed only on day 1 (two-way repeated measures ANOVA on day 1: genotype main effect, $F_{1,24}$ = 4.798, # p = 0.0384; day 2: genotype main effect, $F_{1,24}$ = 0.05209, p = 0.8214; day 3: genotype main effect, $F_{1,24}$ = 0.4063, p = 0.5299). (C) The preference ratio for a social stimulus differed across days (two-way repeated measures ANOVA: day main effect, $F_{1.951,46.82}$ = 20.67, p < 0.0001), but no differences were seen between groups (two-way repeated measures ANOVA: genotype main effect, $F_{1,24}$ = 0.1992, p = 0.6594; day x genotype interaction, $F_{2,48}$ = 0.5837, p = 0.5617).

Figure 4-Figure Supplement 3

Figure 4-Figure Supplement 3 5-HT neuron-specific Nrxn TKO mice display normal learning and memory and repetitive behaviors.

(A) Protocol for fear conditioning. Fear conditioning was performed over four days. Mice were presented with two different auditory cues, one paired (yellow speaker) with a foot shock (yellow thunderbolt) and one unpaired (green speaker). ITI: intertrial interval. (B) Cntl (n = 14) and Fev/RFP/NrxnTKO (n = 11) mice both showed differences in freezing behavior across the presentations associated with the foot shock (CS+) (two-way repeated measures ANOVA: stimulus main effect, $F_{2.607,59.97} = 45.06$, #### $p < 0.0001$). No differences were seen between groups (two-way repeated measures ANOVA: genotype main effect, $F_{1,23} = 0.00339$, $p = 0.9541$; stimulus x genotype interaction, $F_{4,92} = 0.2652$, $p = 0.8996$). (C) Among Cntl and Fev/RFP/NrxnTKO, freezing to the CS+ was increased relative to the CS- (stimulus with no association between auditory cue and foot shock) (two-way repeated measures ANOVA: stimulus main effect, $F_{1,23} = 85.22$, #### $p < 0.0001$), but no differences were detected between groups (two-way repeated measures ANOVA: genotype main effect, $F_{1,23} = 1.67$, $p = 0.2091$; stimulus x genotype interaction, $F_{1,23} = 0.03121$, $p = 0.8613$). (D) Fev/RFP/NrxnTKO mice showed no altered freezing behavior during contextual recall (unpaired two-tailed Student's t-test: $t_{23} = 0.01457$, $p = 0.9885$). (E) Protocol for object interaction test. (F) Cntl (n = 15) and Fev/RFP/NrxnTKO (n = 12) showed no preference for identical objects on day 1 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,25} = 2.606$, $p = 0.119$; genotype main effect, $F_{1,25} = 0.02422$, $p = 0.8776$; stimulus x genotype interaction, $F_{1,25} = 0.1923$, $p = 0.6648$) and day 2 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,25} = 3.18$, $p = 0.0867$; genotype main effect, $F_{1,25} = 0.27$, $p = 0.6079$; stimulus x genotype interaction, $F_{1,25} = 0.1825$, $p = 0.6729$) and preferred the novel object over the familiar object on day 3 (two-way repeated measures ANOVA: stimulus main effect, $F_{1,25} = 129.6$, $p < 0.0001$; genotype main effect, $F_{1,25} = 0.1452$, $p = 0.7064$; stimulus x genotype interaction, $F_{1,25} = 0.03164$, $p = 0.8603$). (G) The preference ratio for the objects differed across days (two-way repeated measures ANOVA: day main effect, $F_{1.932,48.3} = 47.7$, $p < 0.0001$), but no differences were seen between groups (two-way repeated measures ANOVA: genotype main effect, $F_{2,50} = 0.008519$, $p = 0.9272$; day x genotype interaction, $F_{2,50} = 0.1781$, $p = 0.8374$). (H) Cntl (n = 12) and Fev/RFP/NrxnTKO (n = 12) mice showed no differences in the number of marbles buried (unpaired two-tailed Student's t-test: $t_{22} = 1.491$, $p = 0.1502$). (I) Left, no differences in grooming time were observed between groups (unpaired two-tailed Student's t-test: $t_{22} = 0.04104$, $p = 0.9676$). Right, the number of grooming episodes was similar between groups (unpaired two-tailed Student's t-test: $t_{22} = 0.5443$, $p = 0.5917$).

Table 1. Nrxn transcript IDs used for quantification.

ENSMUST00000072671.13	αNrxn1
ENSMUST00000160844.9	αNrxn1
ENSMUST00000174331.7	αNrxn1
ENSMUST00000159778.7	βNrxn1
ENSMUST00000174337.7	βNrxn1
ENSMUST00000161402.9	αNrxn1
ENSMUST00000054059.14	αNrxn1
ENSMUST00000172466.7	βNrxn1
ENSMUST00000160800.8	αNrxn1
ENSMUST00000113462.7	αNrxn2
ENSMUST00000236635.1	αNrxn2
ENSMUST00000113461.7	αNrxn2
ENSMUST00000235714.1	αNrxn2
ENSMUST00000137166.7	αNrxn2
ENSMUST00000167734.7	αNrxn3
ENSMUST00000190626.6	αNrxn3
ENSMUST00000167103.7	αNrxn3
ENSMUST00000057634.13	αNrxn3
ENSMUST00000238943.1	βNrxn3
ENSMUST00000110133.8	βNrxn3
ENSMUST00000110130.3	βNrxn3
ENSMUST00000167887.7	αNrxn3