1

3 **A striatal circuit balances learned fear in the presence and absence of sensory cues**

7 Michael Kintscher[1], Olexiy Kochubey[1], Ralf Schneggenburger[1*]

11 [1] Laboratory for Synaptic Mechanisms, Brain Mind Institute, School of Life Science, Ecole

12 Polytechnique Fédérale de Lausanne (EPFL), Switzerland.

14 * Corresponding author; e-mail ralf.schneggenburger@epfl.ch

**Abstract**

During fear learning, defensive behaviors like freezing need to be finely balanced in the presence or absence of threat-predicting cues (conditioned stimulus, CS). Nevertheless, the circuits underlying such balancing are largely unknown. Here, we investigate the role of the ventral tail striatum (vTS) in auditory-cued fear learning of male mice. *In-vivo* $Ca^{2+}$ imaging showed that robust sub-populations of direct (D1R+) and indirect pathway neurons (Adora+) in the vTS responded to footshocks, and to the initiation of movements after freezing; moreover, a sub-population of D1R+ neurons increased its responsiveness to an auditory CS during fear learning. *In-vivo* optogenetic silencing shows that footshock-driven activity of D1R+ neurons contributes to fear memory formation, whereas Adora+ neurons modulate freezing in the absence of a learned CS. Circuit tracing identified the posterior insular cortex (pInsCx) as an important cortical input to the vTS, and recording of optogenetically-evoked EPSCs revealed long-term plasticity with opposite outcomes at the cortical synapses onto D1R+ - and Adora+ neurons. Thus, direct- and indirect pathways neurons of the vTS show differential signs of plasticity after fear learning, and balance defensive behaviors in the presence and absence of learned sensory cues.

## Introduction

Fear learning is an evolutionary conserved behavior, critically important for animals to detect signs of danger in an ever-changing environment. As such, fear learning is necessary for the survival of animals (Phelps and LeDoux, 2005; Janak and Tye, 2015). Nevertheless, learned defensive behaviors need to be finely regulated, so that animals can return to their normal behaviours after the cessation of threat -predicting sensory cues (Zanette et al., 2011). Furthermore, a pathological overexpression of defensive behaviours is a hallmark of several anxiety-related disorders in humans (Dunsmoor et al., 2011). Therefore, it is important to understand the neuronal circuits that balance the expression of learned defensive behaviors during and after the presence of threat-predicting sensory cues.

The study of the neuronal mechanisms of fear learning has been strongly facilitated by employing auditory-cued fear learning in model animals like rodents (Davis, 1992; LeDoux, 2000). In fear learning, subjects learn to associate an initially neutral sensory cue, the conditioned stimulus (CS; often an auditory stimulus), with an aversive or painful outcome like a footshock (the unconditioned stimulus, US). After associative learning, subjects will develop a defensive behavior when the CS is later presented alone (Fanselow, 2018). The defensive behavior that is mostly studied in the context of fear learning in rodents is behavioral arrest, also called freezing (Fanselow, 1994). Studies spanning several decades have firmly established that the amygdala has an important role in fear learning (see Davis, 1992; LeDoux, 2000; Duvarci and Paré, 2014; Tovote et al., 2015, for reviews). The lateral amygdala (LA) is viewed as an input structure to the amygdalar complex (LeDoux et al., 1990), which connects to both the basal amygdala (BA) and the central amygdala, where further integration and processing takes place; finally, (Romanski et al., 1993; Quirk et al., 1995; Amano et al., 2011; Grewe et al., 2017) the execution of learned freezing depends on a central amygdala to midbrain (periaqueductal gray) projection (LeDoux et al., 1988;

Tovote et al., 2016). Nevertheless, it is likely that further neuronal circuits beyond these amygdalar

circuits are involved in fear learning.

The striatum is part of the basal ganglia motor system, a neuronal system with important roles in

the control of movement, action selection and reward - based learning (for reviews, see Hikosaka et

al., 2000; Graybiel, 2005; Grillner et al., 2005; Redgrave et al., 2010; Nelson and Kreitzer, 2014;

Klaus et al., 2019). The principal neurons of the striatum are inhibitory projection neurons of two

types. First, striato-nigral neurons project in a direct pathway to output structures like the

Substantia nigra pars reticularis and others; these neurons selectively express D1-dopamine

receptors (D1R). Second, striato-pallidal neurons project in a more indirect pathway towards the

basal ganglia output structures; these neurons selectively express D2-dopamine receptors, and also

adenosine-2A receptors (Adora) (Gerfen et al., 1990; Schiffmann and Vanderhaeghen, 1993).

Different sub-areas of the striatum have different roles in motor control and motor learning. The

dorsal striatum is involved in the learning of motor sequences and in the selection of appropriate

actions (Nelson and Kreitzer, 2014; Klaus et al., 2019), as well as in habit formation (Redgrave et

al., 2010; Burguiere et al., 2015), whereas the ventral striatum is important for reward-based

learning (Humphries and Prescott, 2010; Cox and Witten, 2019).

Recently, based on brain - wide studies of the cortical inputs to the mouse striatum, a further sub-

area of the striatum was identified; a posterior area called tail striatum (Hintiryan et al., 2016;

Hunnicutt et al., 2016; see Valjent and Gangarossa, 2021 for review). Interestingly, early *in-vivo*

recordings in the LA have found CS - and US-responsive neurons also in the tail striatum adjacent

to the LA (Romanski et al., 1993). Recent work has shown that the tail striatum is, similarly as

other striatal areas, composed of D1R-expressing (DR1+) and D2R- and Adora-expressing neurons

(Gangarossa et al., 2019). Furthermore, *in-vivo* imaging studies have shown that dopaminergic

axons in the tail striatum code for salient sensory stimuli, but not for rewarding stimuli (Menegas et

al., 2018). However, the role of the tail striatum in fear learning has remained elusive.

Here, we use *in-vivo* miniature microscope Ca$^{2+}$ imaging, as well as *in-vivo* and *ex-vivo* optogenetic

approaches and circuit tracing, to investigate the role of D1R+, and Adora+ neurons located in the

ventral part of the tail striatum (vTS) in auditory-cued fear learning.

**Results**

**Coding for tones and movement by D1R+ vTS neurons**

We started by imaging the activity of vTS neurons during a three-day fear learning paradigm

(Figure 1A). To access the vTS, a deep brain area close to the basolateral - and central nuclei of the

amygdala, we used miniature-microscope Ca$^{2+}$ imaging (Figure 1B, C; Ghosh et al., 2011). In a

first series of Ca$^{2+}$ imaging experiments, we used *Drd1a$^{Cre}$* mice, to target the expression of

GCaMP6m to neurons of the direct pathway, using a Cre-dependent AAV vector (Figure 1B, C;

Materials and methods). The *Drd1a$^{Cre}$* mouse chosen for this purpose (line EY217 from GenSAT;

see Materials and methods) shows expression in the vTS, but no expression was observed in the

adjacent cortical or claustrum structures (Figure 1D; Figure 1 - figure supplement 1; see also

Gerfen et al., 2013). This allowed us to target striatal neurons selectively with the employed mouse

line.

The fear conditioning protocol consisted of three sessions given on subsequent days (Figure 1A).

On the first day, mice experienced a habituation session during which six 30-s CS stimulation

blocks consisting of 7 kHz tone beeps were applied (see Materials and methods). During this

session, only a small sub-population of D1R+ neurons (6/176) showed a response to the tone beeps

(Figure 1E, I). One day later, each of the six CS blocks was followed by a 1-s footshock, to which

45/163 imaged D1R+ neurons responded with robust Ca$^{2+}$ signals (Figure 1F, H). During this

training session, an increased number of D1R+ neurons responded to tone beeps (CS) (Figure 1J, L;

Chi-square test, $X^2_{df=2} = 8.004$, p = 0.018), although the average $Ca^{2+}$ response of those neurons

that responded was similar to the one observed during the habituation session (Figure 1I, J). During

the training - and recall session, we also observed that tone beeps were in 10 - 15% of the cases

followed by movement transitions; for the calculation of the number of CS - responders, these trials

were removed (see Figure 1 - figure supplement 2). Finally, during a fear memory recall session on

day 3, the CS was presented alone in a different context. During this session, the number of tone -

responsive neurons was comparable to the one during the training session (Figure 1L);

nevertheless, the average Z-scored $Ca^{2+}$ signal was increased above the response amplitude on the

training session (Figure 1J, K; 0.366 ± 0.071 versus 0.093 ± 0.034 for n = 19 and 17 responders,

respectively; 95% CI: [0.2164; 0.516]; [0.0212; 0.1649]; U = 61, p = 0.001, Mann-Whitney test).

Thus, *in-vivo* $Ca^{2+}$ imaging showed that a significant fraction of D1R+ neurons in the vTS responds

to footshocks, and furthermore, that D1R+ neurons increase their responsiveness to the CS.

During fear learning, rodents acquire a defensive behavior in response to a CS, in the form of

freezing (LeDoux, 2000; Fanselow, 2018). The freezing bouts of mice typically lasted a few

seconds, and were interrupted by movement re-initiation (see e.g. Figure 1 E-G, *top*). We asked

whether transitions from freezing to movement would drive activity in D1R+ vTS neurons; we first

restricted this analysis to times when no tones (CS) were presented. Aligning GCaMP6m

fluorescence traces to the movement onset revealed $Ca^{2+}$ events in a sub- population of D1R+

neurons; we call these "movement - ON" responses. The number of responding neurons decreased

across the three-day fear learning protocol, although this trend did not reach statistical significance

(Figure 1M-O; Figure 1S, open symbols; p = 0.117; $X^2_{df=2} = 4.292$, Chi-square test). We next

analyzed movement-ON responses during the 30-s CS presentations (Figure 1P-R). We found that

a substantial number of movement - ON transitions were preceded by a tone; for the calculation of

the percentage of movement - ON responders, these events were removed (Figure 1 - figure

supplement 3). The analysis showed that the number of D1R+ neurons that responded to a

movement - ON transition during the time of the CS presentations increased during the training

session, and was maintained at an elevated level during the recall session (Figure 1S, filled

symbols; p = 0.0155; $X^2_{df=2}$ = 8.340, Chi-square test). Furthermore, the number of neurons

responding to a movement - ON transition was always higher during the CS, as compared to no -

CS periods (Figure 1S). Taken together, this data suggests that subpopulations of D1R+ neurons, in

addition to responding to footshocks- and to tone stimulation, also code for movement onset.

We next analyzed more comprehensively how the $Ca^{2+}$-event frequency depends on the movement

state of the animal *and* on the presence or absence of a CS. For this, we first deconvolved the

fluorescence traces to obtain times of $Ca^{2+}$ events and their amplitudes (Figure 1T; Materials and

methods; Pnevmatikakis et al., 2016; Giovannucci et al., 2019). This allowed us to compute the

amplitude-weighted frequency of $Ca^{2+}$ events during the four combinations of movement states /

tone presentations, for each day of the fear learning protocol (Figure 1U; Figure 1 - figure

supplement 4). The data was significantly different across conditions (p < 0.0001; KW = 138.33,

Kruskal-Wallis test). Pairwise comparisons of these conditional $Ca^{2+}$ event frequencies within each

session showed that on most sessions, the activity of D1R+ neurons was significantly higher when

the mice moved, than when they froze, with the exception of the no-CS times on the training day

(Figure 1T; Dunn's multiple comparisons test; for p-values, see legend to Figure 1 -figure

supplement 4). The $Ca^{2+}$ event frequency on the training day - given that mice moved - was

significantly changed by the CS (Figure 1U; p = 0.0163; Mean rank difference = 241.9, Kruskal-

Wallis test followed by Dunn's multiple comparisons test). Thus, *in-vivo* $Ca^{2+}$ imaging during a

three-day fear learning protocol shows that the activity of the D1R+ neurons in the vTS is higher

during movements than during the freezing state of the mice. Moreover, fear learning increases the

number of neurons with a phasic response to tones (CS), as well as the number of neurons with a

movement - ON response during the CS presentation (Figure 1L, S).

To determine the location of the imaged neurons and to compare them across mice, we aligned the

center of the GRIN lens to a mouse brain atlas in each mouse, and generated a common cell map

based on the cell coordinates relative to the lens center (Materials and methods). This revealed a

hotspot of footshock- and tone-responding D1R+ neurons in the posterior - ventral region of the tail

striatum medial to the LA (Figure 2A-C; Figure 2 - figure supplement 1). D1R+ neurons with

movement - ON responses during the recall sessions were located in a similar area (Figure 2C).

Venn plots of the overlay of the various response types showed that during the habituation session,

neurons with responses to the CS and movement did not strongly overlap, and represented a small

proportion of the imaged neurons (Figure 2D). During the training session, the neurons with tone -

and movement - ON responses increased in numbers as shown above (Figure 1L, S), and about half

of each sub-population also showed a footshock response. During the recall session, the

populations of both tone - and movement responders stayed constant with respect to the training

day, and these response types now overlapped substantially (Figure 2D). Taken together, *in-vivo*

$Ca^{2+}$ imaging shows that a subpopulation of D1R+ vTS neurons, located in a posterior - ventral

hotspot of the tail striatum, responds to footshocks. During the course of fear learning, these

neurons increasingly code for an aversively-motivated CS and for movement - ON transitions,

suggesting that these representations in populations of D1R+ neurons of the vTS undergo plasticity

driven by fear learning.

**Coding for footshocks and movement by Adora+ vTS neurons**

The other large population of principal neurons in the vTS are Adora+ neurons which, in analogy to

other striatal areas, represent neurons of an "indirect" pathway through the basal ganglia (Gerfen et

al., 1990; Gangarossa et al., 2019). We next investigated the *in-vivo* activity of this population of

vTS neurons throughout the three-day fear learning protocol, using an *Adora2a^{Cre}* mouse line to

target the expression of GCaMP6m to Adora+ neurons in the vTS (Figure 3A; Figure 3 - figure

supplement 1; Materials and methods). About forty percent of the Adora+ vTS neurons responded

to footshocks presented during the training session (79 / 201 neurons; Figure 3C, 3E). A small

subpopulation (13/173 or ~ 8%) responded to tone beeps during the CS (Figure 3F). Contrasting

with the D1R+ neurons, the percentage of tone (CS) - responsive neurons did not change during

fear learning (Figure 3F - H; Figure 3I; $X^2_{df=2}$ = 1.756, p = 0.415, Chi-square test). Thus, a large

sub - population of Adora+ neurons in the vTS responds to footshocks, but the number of Adora+

neurons that responds to tones (CS) remains unchanged during fear learning.

We next analyzed whether the activity of Adora+ neurons in the vTS was modulated by the

movement state of the mice. In the absence of tones, a moderate sub-population of Adora+ neurons

showed movement - ON responses during the habituation session (27/173 or ~ 16 %); this number

decreased over the course of fear learning (Figure 3J - L; Figure 3P, open symbols; $X^2_{df=2}$ = 7.557,

p = 0.0229; Chi-square test). In contrast, during the CS, there was a larger number of Adora+

neurons that showed movement - ON responses; their number was unchanged over the three-day

fear learning protocol (Figure 3M - O; Figure 3P, closed symbols; $X^2_{df=2}$ = 0.67, p = 0.717; Chi-

square test). These experiments show that a substantial sub-population of Adora+ neurons in the

vTS codes for movement onset, but this representation was unchanged by fear learning (Figure 3M

- P).

Similarly as for the D1R+ neurons, we next analyzed the activity of Adora+ neurons as a function

of the four combinations of movement state (movement versus freezing) and CS presentation

(presence, or absence of a CS) (Figure 3Q, R). A Kruskal-Wallis test showed that the amplitude -

weighted frequency of $Ca^{2+}$ events differed across categories (Figure 3R; KW = 184.6, p < 0.0001).

During all three behavior sessions, and irrespective of the presence - or absence of a CS, the

activity of Adora+ neurons was larger during movement than during freezing (Figure 3R; see

legend to Figure 3 - figure supplement 2 for the corresponding p - values; post-hoc Dunn's multiple

comparison test). On the other hand, the presence or absence of a CS did not significantly modulate

the activity of Adora+ neurons, irrespective of whether the mice moved, or froze (Figure 3R; see

legend to Figure 3 - figure supplement 4 for p - values; post-hoc Dunn's multiple comparison test).

This analysis thus corroborates our finding that the activity of Adora+ neurons is only little

modulated by tones and that it is more strongly modulated by movement, but that neither of the two

representations are modulated in a plastic fashion by fear learning. The spatial distribution of

Adora+ neurons responding to footshocks, tones and movement-ON transitions was overall similar

to the one of D1R+ neurons (Figure 3 - figure supplement 3). Taken together, *in-vivo* $Ca^{2+}$ imaging

shows that a substantial percentage of Adora+ neurons in the vTS responds to footshocks, and to

movement - ON transitions, and a smaller sub-population of these neurons responds to tones

(Figure 3S). However, Adora+ neurons do not change their responses to tones and movement - ON

transitions during fear learning.

**D1R+ and Adora+ vTS neurons do not instruct freezing or movement**

The *in-vivo* $Ca^{2+}$ imaging experiments showed that sub-populations of neurons within the two main

types of principal neurons in the vTS code for footshocks, for the CS and for movement - ON

transitions; furthermore, D1R+ neurons increased their representation of tones and movements with

fear learning (Figures 1-3). We next wished to investigate how D1R+ and Adora+ vTS neurons

might contribute to fear learning. A classical model of basal ganglia function postulates that D1R+

neurons in the direct pathway initiate movement, whereas Adora+ (or D2R+) neurons in the

indirect pathway suppress movements (Kravitz et al., 2010; Nelson and Kreitzer, 2014; but see

Klaus et al., 2019). Thus, one possible straightforward hypothesis is that activity of Adora+

neurons of the vTS instructs an arrest of movement, or freezing, and vice-versa, that D1R+ neurons

might instruct movement re-initiation. We wished to test this hypothesis by optogenetic activation

of either D1R+ - or Adora+ vTS neurons, at times when naive mice (that had not undergone fear

learning) are engaged in regular exploratory behavior. For this, we used the channelrhodopsin

variant Chronos (Klapoetke et al., 2014), which was expressed bi-laterally and Cre-dependently in

D1R+, or Adora+ neurons of the vTS, using AAV1:hSyn:FLEX:Chronos-eGFP in the respective

Cre-mouse line (Figure 4A). Three to four weeks later, the mice were allowed to explore the fear

conditioning chamber, and trains of optogenetic stimuli were applied at pre-determined intervals,

irrespective of whether the mice moved or paused from movement (pulse duration 1 ms, repeated at

25 Hz for 2s; each train given six times for each mouse). Optogenetic stimulation did not lead to

changes in the movement activity of the $Drd1a^{Cre}$ mice, nor of the $Adora2a^{Cre}$ mice (Figure 4B, C;

N = 8 and 8 mice; $t_6 = 0.521$, $p = 0.621$, and $t_7 = 0.529$, $p = 0.163$ respectively, paired t-test). In

additional experiments with Adora+ neurons, we employed longer light pulses (2 and 5 ms), but we

similarly did not observe effects on the movement state of the mice (Figure 4 - figure supplement

1). These experiments suggest that in naive mice, the activity of neither D1R+ - nor of Adora+

neurons is sufficient to modulate the movement activity of the mice.

**Footshock-driven activity of D1R+ vTS neurons contributes to fear learning**

We found that many D1R+ neurons respond to footshocks, and that following fear learning, these

neurons show an increased response to the CS and to movement - ON transitions (Figures 1, 2).

This suggests that footshock responses might drive a plasticity of CS - representation, and of the

movement - ON representation in D1R+ vTS neurons. To investigate the role of the footshock -

evoked activity of D1R+ vTS neurons in fear learning, we next silenced the activity of D1R+ vTS

neurons during footshock presentation, and observed the effects of this manipulation on freezing

behavior during the training day, and one day later during fear memory recall.

For optogenetic silencing, we expressed the light-sensitive proton pump Archaerhodopsin (Arch;

Chow et al., 2010) in a Cre-dependent manner bilaterally in the vTS of $Drd1a^{Cre}$ mice, and

implanted optic fibers over each vTS (Figure 5A; Figure 5 - figure supplement 1). Mice in a control

group were injected with an AAV vector driving the expression of eGFP (Materials and methods).

Four weeks later, both groups of mice underwent auditory-cued fear learning, and on the training

day yellow laser light (561 nm) was applied for 3s, starting 1s before each footshock, with the aim

to suppress the footshock-driven activity of D1R+ neurons (Figure 5B). We found in *ex-vivo*

experiments that yellow light strongly hyperpolarizes Arch-expressing D1R+ and Adora+ vTS

neurons, suppresses action potential (AP) firing, and does not evoke rebound APs when the light is

switched off (Figure 5 - figure supplement 2).

During the habituation day of the fear learning protocol the mice showed little freezing, as expected

(Figure 5C). On the training day, during which mice received a 1s footshock after each tone block,

mice in both groups showed increasing levels of freezing throughout the training period,

interrupted by low freezing activity immediately following the footshocks, caused by increased

shock-evoked running and escape behavior (Figure 5D; CS periods and footshocks are indicated by

blue and yellow vertical lines, respectively). To quantify freezing behavior, we averaged the

percent freezing during the CS, and during four no-CS epochs (see Figure 5D, light blue vertical

bars, and lower grey bars, respectively). This showed that freezing levels were not different

between the Arch- and the control group, neither for the CS- nor for the no-CS epochs (Figure 5E,

*left*, for the CS epochs: $F_{1,20} = 0.965$, p = 0.338; Figure 5E *right*, for the no-CS epochs: $F_{1,20} =$

0.583, p = 0.454; two-way repeated measures - ANOVA). Thus, optogenetic silencing of the

footshock-evoked activity of D1R+ vTS neurons did not change the freezing behavior of mice

during the training day.

On the third day, we tested for fear memory recall by applying tone stimulation (CS) alone in a

different context (see Materials and methods). The time-resolved freezing analysis revealed a

gradual increase of freezing levels when mice entered the conditioning chamber (to ~ 20%; Figure

5F, arrow). This baseline level of freezing has been observed before (see e.g. Cummings and Clem,

2020), and likely represents a residual contextual fear memory, despite the change of the context

between the training day and the recall day (Materials and methods). The 30-s tone blocks (CS)

caused a vigorous increase in freezing in eGFP-expressing control mice, whereas in the Arch -

expressing mice, the CS was less efficient in driving freezing (Figure 5F, black- and green average

traces, N = 10 eGFP- and N = 12 Arch -expressing $Drd1a^{Cre}$ mice). Averaging and statistical

analysis revealed a significant difference in CS-driven freezing between control - and Arch -

expressing $Drd1a^{Cre}$ mice (Figure 5G, closed data points; $F_{1,20}$ = 4.853, p = 0.039, two-way

repeated measures - ANOVA). On the other hand, freezing during the no - CS epochs was

unchanged between the two groups (Figure 5H, open data points; p = 0.541, two-way repeated

measures - ANOVA). Thus, optogenetic inhibition of D1R+ neurons in the vTS during the

footshocks on the training day causes a diminished auditory-cued recall of fear memory one day

later. This suggests that footshock - driven activity of D1R+ vTS neurons contributes to auditory-

cued fear learning.

**Adora+ vTS neurons suppress learned fear in the absence of a CS**

We next investigated the role of footshock-driven activity in the Adora$^+$ neurons of the vTS for

auditory-cued fear learning. For this, we silenced the activity of Adora$^+$ vTS neurons during the

footshocks given on the training day (Figure 6A, B; Figure 6 - figure supplement 1), in an approach

analogous to the one used for the D1R$^+$ neurons. During the habituation day, we observed low

freezing as expected (Figure 6C). On the training day, mice in both the eGFP - expressing control

group and in the Arch group showed a gradual increase in freezing with successive CS - US

pairings, interrupted only by low freezing activity immediately following the footshocks, as in the

$Drd1a^{Cre}$ mice (Figure 6D). The analysis of freezing during the CS - and no-CS epochs revealed no

differences in the freezing levels between the Arch - and the eGFP groups on the training day

(Figure 6E; $F_{1,\,17}$ = 0.415, p = 0.528 and $F_{1,17}$ = 1.087, p = 0.312 respectively; two-way repeated

measures ANOVA). Thus, similar as for the D1R$^+$ neurons, footshock-driven activity in Adora$^+$

vTS neurons is not necessary for the freezing behavior that develops during the training day.

On the fear memory recall day, the dynamics of the freezing behavior differed between the eGFP

and the Arch group (Figure 6F, G). While the eGFP - expressing control mice displayed an

increased freezing during each CS epoch followed by a relaxation to lower freezing levels, mice in

the Arch group showed a delayed relaxation of freezing following the CS epochs (Figure 6F; black

and red data respectively). To quantify these effects, we analyzed the average time spent freezing

during the CS and during a late no-CS epoch (Figure 6F, light blue bars, and lower grey bars). This

showed a trend towards an *increased* freezing during the no-CS epochs in the Arch group as

compared to eGFP controls, although this difference did not reach statistical significance (Figure

6G, right; $F_{1,17} = 4.40$, p = 0.0512; two-way repeated measures ANOVA, N = 8 and 11 eGFP - and

Arch mice, respectively). On the other hand, freezing during the CS epochs was unchanged

between the Arch - and the control group (Figure 6G, *left*; $F_{1,17} = 0.249$, p = 0.624; two-way

repeated measures ANOVA). Thus, silencing the footshock - driven activity of Adora+ neurons in

the vTS induces a trend towards higher freezing during fear memory recall in the *absence* of the

CS. These findings, together with the results obtained from silencing D1R$^+$ vTS neurons (see

above), suggest that direct- and indirect pathway neurons of the vTS have separate, but functionally

synergistic roles in fear learning. In fact, the action of both sub-systems together increases the

difference between the strength of a defensive behavior in the presence, and absence of a learned

sensory cue.

**Brain-wide screening of presynaptic inputs to the vTS**

Our *in-vivo* Ca$^{2+}$ imaging and optogenetic experiments have revealed a differential role of D1R+ -

and Adora+ vTS neurons in auditory-cued fear learning. To start investigating the role of synaptic

afferents to vTS neurons in these plasticity processes, we next used monosynaptic retrograde rabies

virus tracing to identify the presynaptic neuron pools that provide input to D1R$^+$ and Adora$^+$

neurons in the vTS (Figure 7A; Wickersham et al., 2007; Wall et al., 2013)).

*Drd1a<sup>Cre</sup>*, or *Adora2a<sup>Cre</sup>* mice were injected into the vTS with an AAV-helper virus driving the

Cre-dependent expression of TVA, eGFP and oG, to render the infected neurons competent for

later EnvA-pseudotyped rabies virus uptake (Figure 7A, see Materials and methods for the specific

viruses). Three weeks later, a pseudotyped delta-G rabies vector driving the expression of dsRed

was injected at the same coordinates; control experiments confirmed the specificity of the Cre-

dependent expression of TVA - expressing vector, and the absence of ectopic expression of the

rabies vector (Figure 7 - figure supplement 1). In this way, starter cells at the injection site in the

vTS could be identified by GFP- and dsRed co-labelling (red) (Figure 7B, cells appearing yellow in

the overlay). Trans-synaptically labelled neurons outside of the striatum were analyzed based on

their expression of dsRed. We placed all presynaptic neurons found in N = 2 *Drd1a<sup>Cre</sup>* and N = 2

*Adora2a<sup>Cre</sup>* mice into brain-wide models, with cells from the two mice arbitrarily positioned on

different brain sides (Figure 7E, F; red and blue dots, respectively). In both *Drd1a<sup>Cre</sup>* and

*Adora2a<sup>Cre</sup>* mice, we found the highest number and density of dsRed-expressing cells in the dorsal

(granular and dysgranular) part of the posterior insular cortex (pInsCx), and in the adjacent ventral

part of the secondary somatosensory cortex (S2; see also Figure 7 Table Suppl. 1 for a list of all

brain structures with detected presynaptic neurons; Figure 7C, D, G, H). A sizeable number of

back-labelled neurons was also found in various areas of the primary somatosensory cortex (S1), in

the globus pallidus externa (GPe), and in the thalamic parafascicular nucleus (PF; Figure 7C, D, G,

H; Figure 7 - figure supplement 2). The GPe and the PF stood out by having high densities of

presynaptic neurons to both D1R- and Adora-MSNs, even though their absolute numbers were not

large (Figure 7G, H; black bars). Although the brain-wide distribution of presynaptic input neurons

to D1R$^+$ and Adora$^+$ neurons in the vTS was overall similar, we detected differences on a finer

scale. Thus, we found that structures known to process auditory - and multimodal sensory

information, like the auditory cortex (AUD), the temporal association cortex (TeA), the posterior

triangular thalamic nucleus (PoT), the posterior intralaminar thalamic nucleus (PiL) and the

basolateral amygdala (BLA), were more strongly back-labelled in *Adora2a^{Cre}* mice than in

*Drd1a^{Cre}* mice (Figure 7G, H; Figure 7 - figure supplement 3) (LeDoux, 2000; Weinberger, 2007;

Sacco and Sacchetti, 2010; Dalmay et al., 2019; Barsy et al., 2020). Finally, we analyzed the a-p

distribution of neurons in the insular cortex and S2 that provide input to both D1R+ - and Adora+

neurons in the vTS (Figure 7I, J). Taken together, rabies-virus mediated circuit tracing shows that

the vTS receives its main cortical input from the S2 and the pInsCx, followed by primary

somatosensory areas. Thalamic areas like PF and VPM, and basal ganglia like GPe, as well as

limbic areas like the CeA and BLA especially for the Adora+ neurons, also contain sizeable

numbers of neurons that provide input to the vTS.

**The posterior insular cortex provides strong excitatory drive to the vTS**

Retrograde rabies-virus labelling showed that the pInsCx, and the adjacent ventral S2 are the

primary cortical input areas to the vTS (Figure 7). We next wished to functionally validate the

connection from the pInsCx to the vTS, using optogenetically-assisted circuit-mapping (Petreanu et

al., 2007; Little and Carter, 2013; Gjoni et al., 2018). For this, we focussed on the connection from

the pInsCx to the vTS, and injected a viral vector driving the expression of Chronos (Klapoetke et

al., 2014), primarily targeting the pInsCx (Figure 8A, B; AAV8:hSyn:Chronos-eGFP, note that

some spill-over of virus into the neighboring S2 cannot be excluded). We used *Drd1a^{Cre}* x

*Rosa26^{LSL-tdTomato}*, or *Adora2a^{Cre}* x *Rosa26^{LSL-tdTomato}* mice, to identify each type of principal neurons

by its tdTomato fluorescence in recordings three - six weeks later. Blue light pulses (1 ms) at

maximal light intensity evoked robust optogenetically-evoked EPSCs of 8.8 ± 1.6 nA in D1R+ vTS

neurons (95% CI: [5.28 nA; 12.2 nA]; n = 12 cells), and of 2.5 ± 0.5 nA in Adora+ vTS neurons

(95% CI: [1.49 nA; 3.58 nA], n = 19 cells); the EPSC amplitudes were significantly different

between the two types of principal neurons (U = 40, p = 0.002, Mann Whitney test; Figure 8C-E).

At both connections, gradually increasing the stimulus light intensity led to a smooth increase of

the EPSC amplitude, which shows that many axons from pInsCx neurons converge onto each type

of principal neuron in the vTS (Figure 8C, D). At high light intensities, the optogenetically-evoked

EPSC amplitudes saturated, suggesting that a maximal number of input axons was activated

(Figure 8C, D). The paired-pulse ratio (PPR) did not differ between the two neuron types (Figure

8F; $t_{28} = 1.288$, $p = 0.208$; unpaired t-test; 95% CI: [0.637; 0.827] and [0.711; 0.960] for D1R+ and

Adora+ neurons). On the other hand, the ratio of direct excitation over feedforward inhibition was

significantly smaller in Adora+ vTS neurons as compared to D1R+ neurons (Figure 8 - figure

supplement 1; $U = 31$, $p = 0.0054$, Mann Whitney test; 95% CI: [2.19; 4.10] and [1.05; 2.40] for

D1R+ and Adora+ neurons). Taken together, optogenetically-assisted circuit mapping shows that

both D1R+ and Adora+ neurons of the vTS receive robust excitatory inputs from the pInsCx. This,

together with the monosynaptic rabies tracing (Figure 7), identifies the pInsCx as providing an

important cortical inputs to the vTS.

**Fear learning causes opposite plasticity at cortical synapses on D1R+ and Adora+ neurons**

The *in-vivo* imaging data showed that many D1R+ and Adora+ neurons in the vTS robustly

respond to footshocks, and that during the course of fear learning, D1R+ neurons increase their

responsiveness to the CS and to movement - ON transitions (Figures 1-3). Furthermore, silencing

each vTS neuron population during the footshocks led to characteristic impairments of freezing in

the presence, and absence of a learned CS (Figures 5, 6). These findings suggest that plasticity

takes place at synapses that drive D1R+ and Adora+ vTS neurons, including at the cortical input

synapses. We therefore next measured the AMPA/NMDA ratio and PPR following fear learning, to

probe for postsynaptic or presynaptic forms of LTP induced by fear learning at each connection

(Yin et al., 2009; Shan et al., 2014; Rothwell et al., 2015; Lucas et al., 2016; see Palchaudhuri et

al., 2022 for a review).

For this purpose, we used optogenetically-assisted circuit mapping to measure optogenetically -

evoked EPSCs at each connection. We injected *Drd1a$^{Cre}$* x *Rosa26$^{LSL-tdTomato}$* mice, and in a

separate series of experiments *Adora2a^Cre* x *Rosa26^LSL-tdTomato* mice, with an AAV vector to drive

the expression of Chronos in neurons of the pInsCx. Three to six weeks later, the mice were

subjected to auditory-cued fear learning (Figure 9A). Following the fear-memory recall session on

day 3, which was performed to validate that mice had successfully learned the CS, mice were

sacrificed, and optogenetically-evoked EPSCs were recorded. A control group of mice underwent

the same protocols, but no footshocks were applied ("CS only" group; Figure 9A). Optogenetically

evoked EPSCs at the pInsCx to D1R$^+$ vTS connection showed a significant increase of the

AMPA/NMDA ratio in the CS + US group, as compared to the CS-only group, suggesting that at

this connection, a postsynaptic form of LTP had occurred during fear learning (Figure 9B, D; $t_{21}$ =

3.833, p = 0.001, unpaired t-test; 95% CI: [1.687; 2.764] and [3.128; 4.719] for CS-only and

CS+US). The PPR, however, was unchanged in D1R+ neurons (Figure 9C, D; $t_{21}$ = 0.033, p =

0.974, unpaired t-test; 95% CI: [0.448; 0.765] and [0.429; 0.791] for CS-only and CS+US).

Conversely, at the pInsCx to Adora+ vTS connection, the AMPA/NMDA ratio was unchanged

(Figure 9E, G; $t_{22}$ = 1.445, p = 0.162, unpaired t-test; 95% CI: [1.538; 2.896] and [0.946; 2.251] for

CS-only and CS+US), but instead, the PPR was increased in the CS+US group as compared to the

CS-only group (Figure 9F, G; U = 18, p = 0.001, Mann Whitney test; 95% CI: [0.193; 0.582] and

[0.574; 0.938] for CS-only and CS+US). The latter finding suggests that fear learning induces a

presynaptic form of long-term depression at the pInsCx to Adora+ vTS synapse. Thus, auditory-

cued fear learning drives long-term plasticity with opposite outcomes at cortical synapses onto

D1R$^+$ and Adora$^+$ neurons in the vTS. These differential forms of long-term plasticity might

contribute to the complementary roles of the two types of vTS principal neurons in fear learning.

**Discussion**

Recent studies in mice have identified the posterior (tail) striatum as an anatomically, and connectionally separate part of the striatum, and evidence suggests that dopamine axons in the tail striatum code for salient sensory events (Hintiryan et al., 2016; Hunnicutt et al., 2016; Menegas et al., 2018; see Valjent and Gangarossa, 2021 for a review). Here, we have investigated the role of the tail striatum in auditory-cued fear learning, in part motivated by earlier studies which showed a convergence of US - and CS signaling in the tail striatum located close to the LA (Romanski et al., 1993). We have targeted the ventral half of the tail striatum close to the amygdala, an area which we refer to as "ventral tail striatum", vTS. This region might include the amygdala-striatum transition zone ("AStria"; located immediately adjacent to the amygdala), but our targeting was not limited to the AStria.

Using Cre-driver mouse lines to target D1R - expressing and Adora - expressing neurons of the direct - and indirect pathways (Gerfen et al., 2013), we studied the role of each type of vTS principal neuron in auditory-cued fear learning. We found that both classes of vTS principal neurons were strongly activated by footshock stimuli, and coded for the movement state of the mice; smaller sub-populations of both neuron types also responded to tones in naive mice. Interestingly, the number of tone-responsive D1R+ neurons, but not of Adora+ neurons, increased during auditory-cued fear learning, not dis-similar to the acquisition of tone responsiveness of amygdalar neurons (Quirk et al., 1995; Amano et al., 2011; Grewe et al., 2017). To investigate the role of footshock-driven activity in vTS principal neuron, which most likely drives plasticity in these neurons, we optogenetically silenced neuronal activity at the time of footshock delivery. In $Drd1a^{Cre}$ mice, this manipulation caused decreased auditory-cue driven freezing one day later, which leads us to conclude that footshock - driven plasticity in vTS D1R+ neurons contributes to the formation of a cue - driven fear memory. This conclusion is corroborated by our finding that fear learning induces a postsynaptic form of LTP at a cortical input synapse to D1R+ neurons in the

vTS. On the other hand, silencing Adora+ neurons during the footshock led to a trend towards

increased freezing in the absence of a learned CS, which suggests that indirect pathway neurons of

the vTS have access to learned contextual freezing. Interestingly, the finding that the vTS Adora+

subsystem acts to suppress freezing in the *absence* of a danger - signaling CS, is reminiscent to

aspects of safety learning (Christianson et al., 2012). Furthermore, a previous study has correlated

the *in-vivo* activity of vTS neurons with safety learning, albeit without distinguishing between

direct- and indirect pathway neurons (Rogan et al., 2005). Taken together, we find that neurons in

the direct - and indirect pathway of the vTS have different roles in fear learning, but together they

act synergistically to increase the difference in the strength of a defensive behavior in the presence,

and absence of a learned sensory cue.

*In-vivo* $Ca^{2+}$ imaging also revealed that sizable sub-populations of both types of vTS principal

neurons code for movement - onset, and that in D1R+ neurons, these movement - ON responses

were increased after fear learning (Figures 1, 3). Movement - correlated activity has been observed

in striatal neurons (Markowitz et al., 2018), as well as in several types of sensory cortices and in a

more brain-wide fashion (Niell and Stryker, 2010; Keller et al., 2012; Stringer et al., 2019). At

present, the role of the movement - related activity in the vTS remains unknown. Our optogenetic

activation experiments did not reveal significant effects on the movement - or freezing behavior of

mice, in line with previous findings in the posterior striatum (Guo et al., 2018), but different from

results in the dorsomedial striatum  (Kravitz et al., 2010); the latter striatal area seems to have a

more immediate role in movement control than the tail striatum. Therefore, it is unlikely that the

movement - ON driven activity that we found here in D1R+ or Adora+ neurons of the vTS

contributes to the control of movement. Rather, one could speculate that the movement - related

activity in the vTS represents a signal related to the bodily state of the animal. In this regard, the

strong excitatory input from the pInsCx to the vTS which we uncovered here seems relevant

(Figures 7, 8). Recent studies have shown that the pInsCx displays state-dependent signaling

relating to physiological needs, or to the aversive state of the animal (Livneh et al., 2017; Gehrlach et al., 2019); it is possible that the activity of the pInsCx is additionally modulated by movement - ON transitions of the animal. Thus, future work might investigate the origins, and functions of the movement - related signals in the vTS.

Previous studies in mice analyzed anterograde tracer experiments from cortex to striatum in a brain-wide fashion; this data has also revealed specific cortical input structures to the tail striatum (Hintiryan et al., 2016; Hunnicutt et al., 2016). We have used rabies - virus mediated back-labelling from genetically identified neurons in the vTS, to identify presynaptic input neurons to the vTS more focally, and specifically for direct- and indirect pathway neurons (Figure 7; see also Wall et al., 2013 for a similar approach in the more anterior dorsal striatum). Our results confirm previous studies regarding the innervation of the vTS by auditory cortical areas, and by amygdalar structures (Hunnicutt et al., 2016). Beyond this, our data identifies the dorsal pInsCx and the S2 as major cortical input structures to the vTS, both for D1R+ and Adora+ neurons. We confirmed the functional relevance of the putative connections by optogenetically-assisted circuit mapping, which revealed unusually large EPSCs at the input from the pInsCx to the vTS (note that the ventral S2 was not covered in our *ex-vivo* optogenetic experiments). Based on the graded input - output curves when varying the light intensity, these EPSCs likely reflect converging inputs from many pInsCx neurons (see also Litvina and Chen, 2017; Gjoni et al., 2018). The strong inputs from pInsCx and presumably ventral S2, suggest that in addition to auditory - and visual information (Yamamoto et al., 2012; Guo et al., 2018; Menegas et al., 2018), somatosensory signals are processed in the vTS. For example, it is possible that the pInsCx transmits somatosensory or nociceptive information about a footshock during aversively motivated learning to the vTS; similarly, movement - related signals might be transferred at this connection, as discussed above. Moreover, it was shown that the pInsCx processes auditory information (Rodgers et al., 2008; Sawatari et al., 2011; Gogolla et al., 2014), thus tone information might additionally be transmitted at this connection. Indeed, our

finding of an enhanced AMPA/NMDA ratio following fear learning at the connection from the

pInsCx to D1R+ vTS neurons gives rise to the hypothesis that learned auditory signals, or

movement - related signals, are transmitted at this connection, a hypothesis which could be tested

in future work.

Subtle, but functionally important differences in the brain areas that project to D1R+ versus Adora+

vTS neurons might determine their different *in-vivo* responses, and, in conjunction with the

different output projections of direct- and indirect pathway neurons, the differential role of each

type of vTS neuron during fear learning. Their differential roles in fear learning are likely

additionally shaped by different effects of dopamine in D1R+ versus Adora+ neurons (Gerfen et

al., 1990; Tritsch and Sabatini, 2012; Hjorth et al., 2020). Recent work has shown that a separate

set of midbrain dopamine neurons innervates the tail striatum, and that these dopamine projections

are activated by salient, but not by rewarding stimuli (Menegas et al., 2015; Menegas et al., 2018).

It remains to be tested whether aversive footshock stimuli similarly cause the release of dopamine

in the vTS. Classical studies showed that dopamine release in the amygdala contributes to fear

learning (Lamont and Kokkinidis, 1998; Guarraci et al., 1999; Nader and LeDoux, 1999), and a

recent study showed that footshock-driven activity of VTA dopamine neurons that project to the

BA contributes to fear learning (Tang et al., 2020). Therefore, dopamine signaling in the vTS,

likely from a different source than dopamine release in the BA (Menegas et al., 2015; Tang et al.,

2020), might contribute to diversifying the roles of vTS direct- and indirect pathway neurons in

fear learning.

Fear learning shows sex-specific differences in rats (Maren et al., 1994; Pryce et al., 1999; Gruene

et al., 2015; see Lebron-Milad and Milad, 2012 for review). Thus, we could not investigate female -

and male mice without differentiating between them; rather, we limited our study to male mice (see

Materials and methods). In rodents, males in general show higher freezing than females, whereas

females show a higher degree of fast bouts of escape reactions (called "darting"), which can also

take the form of learned responses (Gruene et al., 2015). Future work might investigate whether the

direct- or indirect pathway neurons in the vTS contribute to sex-specific defensive behaviors like

darting in females, and whether sex-specific differences exist in the circuit wiring and plasticity

mechanisms of the vTS.

**Conclusions and Outlook**

In summary, we show a role for the vTS in auditory - cued fear learning. We find that direct- and

indirect pathway neurons differentially modulate the degree of learned defensive behaviors in the

presence, and absence of learned sensory cues. This uncovers a role of the vTS in balancing cue-

specific reaction on the one hand, with a more generalized fear response on the other hand.

Adaptive fear discrimination is critically important for animal survival, but is dys-regulated in

anxiety disorders (Sangha et al., 2020). Indeed, it was found that during fear generalization, PTSD

patients exhibited stronger responses in the striatum, amygdala and insular cortex amongst other

areas (Morey et al., 2020). Because fear expression in the absence of aversive sensory cues is

downregulated by Adora+ neurons in the vTS, it is an intriguing possibility to harness the

differential pharmacology of direct- and indirect pathway neurons together with *in-vivo*

manipulations of plasticity (Creed et al., 2015), in an attempt to mitigate the effects of fear

generalization.

**Key Resources Table**

| Reagent type (species) or resource | Designation | Source or reference | Identifiers | Additional information |
|---|---|---|---|---|
| strain, strain background (*M. musculus*) | C57BL/6J | The Jackson laboratory | Jax:000664 | |
| genetic reagent (*M. musculus*) | *Drd1a^Cre* | PMID:17855595; PMID:24360541 | MGI:4366803; MMRRC:030778-UCD; RRID:MMRRC_030778-UCD | STOCK Tg(Drd1-cre)EY217Gsat/Mmucd |
| genetic reagent (*M. musculus*) | *Adora2a^Cre* | PMID:17855595; PMID:24360541 | MGI:4361654; MMRRC:036158-UCD; RRID:MMRRC_036158-UCD | B6.FVB(Cg)-Tg(Adora2a-cre)KG139Gsat/Mmucd |
| genetic reagent (*M. musculus*) | *Rosa26^LSL-tdTomato* | PMID:20023653 | Jax:007909; MGI:3809523; RRID:IMSR_JAX:007909 | B6.Cg-Gt(ROSA)26Sortm9(CAG-tdTomato)Hze/J |
| recombinant DNA reagent | AAV1:hSyn:FLEX:GCaMP6m (viral vector) | Viral vector facility, University of Zürich | ZurichVVF:v290-1 | (6.8e12/ml) |
| recombinant DNA reagent | AAV1:hSyn:FLEX:Chronos-eGFP (viral vector) | University of North Carolina vector core | UNC:Boyden-AAV-Syn-Chronos-GFP | (2.8e12/ml) |
| recombinant DNA reagent | AAV1:CAG:FLEX:Arch-eGFP (viral vector) | University of North Carolina vector core | UNC:Boyden-AAV-CAG-FLEX-Arch-GFP | (2.05e12/ml) |
| recombinant DNA reagent | AAV1:CBA:FLEX:Arch-eGFP (viral vector) | University of Pennsylvania vector core | UPenn:AV-1-PV2432;Addgene:22222-AAV1 | (5.48e12/ml) |
| recombinant DNA reagent | AAV1:CAG:FLEX:eGFP (viral vector) | University of North Carolina vector core | UNC:Boyden-AAV-CAG-FLEX-GFP | (4.4e12/ml) |
| recombinant DNA reagent | AAV8:hSyn:Chronos-eGFP (viral vector) | University of North Carolina vector core | UNC:Boyden-AAV-Syn-Chronos-GFP | (6.5e12/ml) |
| recombinant DNA reagent | AAV8:CAG:FLEX:tdTomato (viral vector) | University of North Carolina vector core | UNC:Boyden-AAV-CAG-FLEX-tdTomato | (6.5e12/ml) |

| | | | | |
|---|---|---|---|---|
| recombinant DNA reagent | AAV1:hSyn:FLEX:TVA-2a-eGFP-2a-oG (viral vector) | Viral vector facility, University of Zürich | ZurichVVF:v243-1; Addgene:85225 | (5.3e12/ml) |
| recombinant DNA reagent | EnvA:deltaG–RV:dsRed (viral vector) | PMID:21867879 | Addgene:32638 | |
| commercial assay or kit | Fluroshield mounting medium with DAPI | Sigma Aldrich | Sigma:F6057-20ML | |
| chemical compound, drug | gabazine | Abcam | Abcam:ab120042; Sigma:SR95531 | |
| software, algorithm | VideoFreeze | Med Associates Inc | Med Associates:VideoFreeze | |
| software, algorithm | EthoVision XT | Noldus Information Technologies | Noldus:EthoVisionXT13; RRID:SCR_000441 | version 13 |
| software, algorithm | ezTrack | PMID:31882950 | RRID:SCR_021496 | https://github.com/denisecailab/ezTrack |
| software, algorithm | Igor Pro | Wavemetrics Inc | RRID:SCR_000325 | version 7.08, 64 bit |
| software, algorithm | Inscopix Data Processing Software | Inscopix Inc | Inscopix:IDPS | |
| software, algorithm | CaImAn | PMID:30652683 | RRID:SCR_021152 | https://caiman.readthedocs.io/ |
| software, algorithm | SHARP-Track | doi:10.1101/447995 | | https://github.com/cortex-lab/allenCCF |
| software, algorithm | Brainrender | PMID: 33739286 | RRID:SCR_022328 | https://edspace.american.edu/openbehavior/project/brainrender/ |
| software, algorithm | GraphPad Prism | GraphPad Software | RRID:SCR_002798 | version 9 |
| software, algorithm | NeuroMatic | PMID:29670519 | RRID:SCR_004186 | pluging for IgorPro |
| other | steretotaxic frame for small animals | David Kopf Instruments | David Kopf Instruments:Model 942 | used with Model 921 mouse adapter |
| other | hydraulic one-axis manipulator | Narishige | MO-10 | for virus injections; see Materials and methods |
| other | 600 µm / 7.3 mm ProView(TM) GRIN lens | Inscopix Inc | Inscopix:1050-004413 | used with nVista3.0 system |

| other | nVista imaging system | Inscopix Inc | Inscopix:nVista3.0; RRID:SCR_017407 | for $Ca^{2+}$ imaging of neurons in freely moving mice; see Materials and methods |
|---|---|---|---|---|
| other | optic fiber implants | Thorlabs Inc | Thorlabs:FT200EMT | 200 µm core / 0.39 NA / 230 µm outer diameter |
| other | ceramic ferrule | Thorlabs Inc | Thorlabs:CFLC230 | 230 µm bore / 1.25 mm outer diameter |
| other | Blue light curing dental cement | Ivoclar Vivadent AG | Ivoclar Vivadent:Tetric EvoFlow | for securing implants at the skull surface |
| other | light curing adhesive | Kulzer GmbH | Kulzer:iBond Total Etch | for treatment of skull before application of dental cement |
| other | fear conditioning apparatus | Med Associates Inc | Med Associates:MED-VFC-OPTO-M | see Materials and methods |
| other | electric footshock stimulator | Med Associates Inc | Med Associates:ENV-414S | used within the fear conditioning apparatus |
| other | 561 nm solid-state laser | Changchun New Industries Optoelectronics Technology (CNI) | CNI:MGL-FN-561-AOM | fiber coupled, maximum output 100 mW; for *in-vivo* activation of Arch |
| other | 473 nm solid-state laser | Changchun New Industries Optoelectronics Technology (CNI) | CNI:MBL-FN-473-150mW | fiber coupled, maximum output 150 mW; for *in-vivo* activation of Chronos, see Materials and methods |
| other | vibrating microtome VT1200S | Leica Microsystems | RRID:SCR_020243 | for preparation of brain slices; see Materials and methods |
| other | patch-clamp amplifier EPC10/2 | HEKA Elektronik | RRID:SCR_018399 | for whole-cell patch-clamp recordings; see Materials and methods |
| other | fluorescent microscope BX51WI | Olympus | RRID:SCR_018949 | - to visualize neurons for whole-cell patch-clamp; see Materials and methods |
| other | high-power LED, blue | Cree Inc | Cree:XPEBRY-L1-0000-00P02 | 460 nm; to excite Chronos in slices; see Materials and methods |
| other | high-power LED, green | Cree Inc | Cree:XPEBGR-L1-0000-00D02 | 530 nm; to activate Arch in slices; see Materials and methods |
| other | LED driver | Mightex Systems | Mightex Systems:BLS-1000-2 | |

| other | silicone photodetector | Thorlabs Inc | Thorlabs:DET36A/M | to measure the time-course of LED light pulse in slice experiments; see Figure 8C, D and Materials and methods |
|---|---|---|---|---|
| other | slide scanning fluorescent microscope | Olympus | Olympus:VS120-L100; RRID:SCR_018411 | for imaging post-hoc histology sections; see Materials and methods |
| other | sliding microtome Microm HM450 | ThermoFisher Scientific | RRID:SCR_015959 | to prepare histological brain sections; see Materials and methods |
| other | confocal microscope | Leica SP8 | RRID:SCR_018169 | for imaging post-hoc histology sections; see Materials and methods |

**Materials and methods**

*Animals*

The experiments were performed with different lines of genetically modified mice (*Mus musculus*) of male sex. The rationale for investigating exclusively male mice was as follows. The aims of the study were to investigate with optogenetic methods whether the vTS has a role in fear learning; to image the *in-vivo* responses of D1R+ - and Adora+ neurons of the vTS during fear learning; to identify the main cortical inputs to both types of vTS principal neurons; and to study signs of long-term plasticity at cortical input synapses to both D1R+ - and Adora+ neurons after fear learning. It has been shown, mainly using rats, that sex-specific differences exist in the strength and types of learned defensive behaviors (Maren et al., 1994; Pryce et al., 1999; Gruene et al., 2015). Therefore, including mice both sexes in the study without differentiating between them would have most likely increased the variability of the results. Thus, it would have been necessary to include male *and* female mice in *separate* groups, which would have doubled the number of experimental groups, and experimental animals used in the study. We therefore decided to perform the initial study in male mice (see also Discussion).

The experiments were performed under authorizations for animal experimentation by the veterinary office of the Canton of Vaud, Switzerland (authorizations VD3274 and VD3518). The following mouse lines were used: 1) *Drd1a$^{Cre}$* (B6.129-Tg(Drd1a-cre)EY217Gsat/Mmucd; see Gong et al., 2007; MMRRC: 030778-UCD; 2) *Adora2a$^{Cre}$* (B6.FVB(Cg)-Tg(Adora2a-cre)KG139Gsat/Mmucd; see Gerfen et al., 2013; MMRRC: 036158-UCD; 3) Cre-dependent tdTomato reporter line, *Rosa26$^{LSL-tdTomato}$* (B6.Cg-Gt(ROSA)26Sor<tm9(CAG-tdTomato)Hze>/J; JAX stock #007909; also called "Ai9"; Madisen et al., 2010). All mice strains were back-crossed for at least five generations to a C57BL/6Jbackground. Mice were weaned at 21 days postnatally (P21), and groups of male mice were housed together under a 12/12 h light/dark cycle (7:00 am, light on), with food and water ad libitum. Surgery was performed at P42 - P56; mice were separated into single cages one

day before surgery. For behavioral experiments (Figure 5, 6), mice from 1 - 2 litters were randomly

assigned to control - (GFP - expressing) or effect group (Arch- expressing). Behavioral testing was

performed during the light cycle.

*Viral vectors and injection coordinates*

For *in-vivo* Ca$^{2+}$-imaging experiments (Figures 1 -3), we injected AAV1:hSyn:FLEX:GCaMP6m

(200nl; 6.8 x10$^{12}$ vg[vector genomes]/ml; cat. v290-1; viral vector facility University of Zürich,

Switzerland), into the left vTS of *Drd1a$^{Cre}$* or *Adora2a$^{Cre}$* mice, using the following coordinates:

medio-lateral (ML) 3.2 mm; anterior-posterior (AP) -0.8 to -1.0 mm; dorso-ventral (DV) -4.3 mm

(from bregma). For the optogenetic activation experiments (Figures 4), we injected

AAV1:hSyn:FLEX:Chronos-eGFP (200nl; 2.80x10$^{12}$ vg/ml; University of North Carolina - UNC

vector core, Chapel Hill, NC, USA) bi-laterally into the vTS of *Drd1a$^{Cre}$* or *Adora2a$^{Cre}$* mice. For

the optogenetic inhibition experiments (Figures 5, 6), we injected AAV1:CAG:FLEX:Arch-eGFP

(200nl; 2.05x10$^{12}$ vg/ml; UNC vector core) bi-laterally into the vTS of *Drd1a$^{Cre}$* or *Adora2a$^{Cre}$*

mice. In earlier experiments AAV1:CBA:FLEX:Arch-eGFP (200nl;5.48x10$^{12}$ vg/ml; AV-1-

PV2432; University of Pennsylvania vector core, Philadelphia, PA, USA, now at Addgene, 22222 -

AAV1) was used; both Arch constructs correspond to the initially described "Arch" (Chow et al.,

2010). Mice in the control group received AAV1:CAG:FLEX:eGFP (200nl; 4.4 x10$^{12}$ vg/ml; UNC

vector core).

For the *ex-vivo* optogenetically-assisted circuit-mapping experiments (Figures 8, 9), *Drd1a$^{Cre}$* x

*Rosa26$^{LSL-tdTomato}$* mice, or *Adora2a$^{Cre}$* x *Rosa26$^{LSL-tdTomato}$* mice were injected unilaterally into the

pInsCx with AAV8:hSyn:Chronos-eGFP (200nl; 6.5x10$^{12}$ vg/ml; UNC vector core), at the

following stereotaxic coordinates: ML 4.2 mm; AP -0.55 mm, DV -3.8 mm (from bregma). In

some experiments, *Drd1a$^{Cre}$*- or *Adora2a$^{Cre}$* mice were used, and an AAV8:CAG:FLEX:tdTomato

(200nl; 6.5x10$^{12}$ vg/ml; UNC vector core) was additionally injected into the vTS (coordinates as

above), to visualize *Drd1a^Cre*- or *Adora2a^Cre* - positive neurons for subsequent patch-clamp

recordings.

*Surgery for virus injection, GRIN lens - , and optical fiber implantation*

The surgery procedures for stereotactic injection of viral vectors alone, or combined with fiber

implantation were as described in (Tang et al., 2020). In short, a mouse was anesthetized with

isoflurane (induction with 3%, maintained at 1%) and the head was fixed in a Model 940

stereotactic injection frame (David Kopf Instruments, Tujunga, CA, USA) using non-rupture ear

bars (Zygoma Ear cups, Kopf Instruments Model 921). Local anesthesia was applied

subcutaneously using a mix (50 µl) of lidocaine (1mg/ml), bupivacaine (1.25mg/ml) and

epinephrine (0.625 µg/ml). After the skull was exposed, small craniotomies were drilled above the

indicated coordinates for injection and fiber insertion. For fiber implantation, an additional

craniotomy was made close to Lambda to insert an anchoring micro screw. Virus suspension was

injected using pulled glass pipettes and an oil hydraulic micromanipulator (MO-10, Narishige,

Tokyo, Japan) with an injection speed of ~ 60 nl/min. For *in-vivo* $Ca^{2+}$-imaging (Figures 1 - 3), a

GRIN lens (600 µm / 7.3 mm ProView™ integrated lenses; cat. 1050-004413; Inscopix Inc, Palo

Alto, CA, USA) was implanted 350 µm above the virus injection site. Prior to GRIN lens

implantation, a 25G medical injection needle was slowly inserted and retracted to facilitate the later

insertion of the blunt-ended lens. To reduce deformation of brain tissue due to continuous vertical

pressure, we lowered the GRIN lens with alternating down (150 µm) and up (50 µm) movements

until the last 200 µm before the final position. For *in-vivo* optogenetic experiments (Figures 4 - 6),

optical fiber implants were implanted bilaterally above the virus injection sites. The fiber implants

were custom-made of a 200 µm core / 0.39 NA / 230 µm outer diameter optic fiber (FT200EMT;

Thorlabs Inc, Newton, NJ, USA) inserted into 1.25 mm outer diameter ceramic ferrules (CFLC230;

Thorlabs) as described in (Sparta et al., 2012). The surface of skull, and the GRIN lens (or optical

fiber implants) were treated with a light curing adhesive iBond Total Etch (Kulzer GmbH, Hanau,

Germany). Blue light curing dental cement (Tetric EvoFlow, Ivoclar Vivadent, Schaan,

Liechtenstein) was then applied to the skull to hold the GRIN lens (or optical fiber implants) in

place. The open end of the GRIN lens at the integrated docking platform was sealed using a Kwik-

Sil silicone compound (World Precision Instruments, Sarasota, FL, USA). After stitching, the skin

was covered with Bepanthen® Plus cream (Bayer AG, Leverkusen, Germany), the drinking water

was supplemented with 1 mg/ml paracetamol and the animals were monitored for the following 6

days to ensure proper post-surgical recovery.

*Behavior*

Auditory-cued fear learning was tested three to four weeks after the surgery. Mice were handled by

the experimenter and habituated to the procedure of attaching a dummy miniature-microscope

(Inscopix Inc.) to the GRIN lens platform (in case of $Ca^{2+}$-imaging experiments), or optical patch

cords (in case of optogenetic experiments), for 10-15 min on five consecutive days. An auditory-

cued fear learning paradigm was performed in a conditioning chamber of a Video Fear

Conditioning Optogenetics Package for Mouse (MED-VFC-OPTO-M, Med Associates Inc.,

Fairfax, VT, USA) under control of VideoFreeze® software (Med Associates Inc). On day 1

(habituation), a mouse at a time was connected to the nVista3.0 mini-microscope (Inscopix Inc.) or

to the optic fiber patch cords, and the animal was placed in the conditioning chamber. The latter

was a rectangular chamber with a metal grid floor, cleaned with 70% ethanol. During the ensuiring

habituation session, six tone blocks (CS), each consisting of 30 tone beeps (7 kHz, 80 dB, 100 ms

duration, repeated at 1 Hz for 30 s), were applied 90 s apart. During a training session on day 2, the

mouse was placed in the same chamber and presented with six CS blocks pseudo-randomly spaced

60-120 s apart, each followed by a 1s foot shock US (0.6 mA, AC) delivered by a stimulator (ENV-

414S, Med Associates Inc). During a fear memory recall session on day 3, the mouse was placed in

a conditioning chamber within a different context, consisting of a curved wall and a smooth acrylic

floor, cleaned with perfumed general-purpose soap, and four CS blocks were applied.

For optogenetic silencing experiments with Arch (Figures 5, 6), light was delivered during 3s

starting 1s before the footshock via 200 μm core / 0.22 NA optic fiber patch cords (Doric Lenses,

Canada) from a 561 nm solid state laser (MGL-FN-561-AOM, 100 mW, Changchun New

Industries Optoelectronics Technology, Changchun, China). The laser was equipped with an AOM

and an additional mechanical shutter (SHB05T; Thorlabs). For optogenetic activation experiments

with Chronos (Figure 4), 1-ms light pulses were delivered at 25 Hz, 2s duration from a 473 nm

solid-state laser (MBL-FN-473-150mW, Changchun New Industries Optoelectronics Technology).

The intensity of each laser was adjusted before the experiment to deliver 10 mW light power at the

fiber tip.

The behaviour of animals was recorded at video rate (30 Hz) by the VideoFreeze® software (Med

Associates Inc). Based on the behavioural videos, a movement trace was generated using ezTrack

software (Pennington et al., 2019; see e.g. Figure 1F-H, red trace). The experimenter, and the

person analyzing the data were blinded to the assignment of each mouse to the control - or test

group. The movement index trace from ezTrack was used to compute a binary freezing trace using

custom procedures in IgorPro 7 (WaveMetrics Inc, Lake Oswego, OR, USA). The animal was

considered to be immobile (freezing state) if the movement index was below a threshold of 40

arbitrary units (without cable attachment) or 120 a.u. (with cable attachment), for a minimum

duration of 0.5 s. The binned trace of percent time spent freezing (10s bin size; see e.g. Figures 4G,

J; 5F, H; 6E, G) was calculated as a time-average of the freezing state from the binary trace, and

then averaged across mice in each group.

*Microendoscopic Ca$^{2+}$-imaging data acquisition and analysis*

We used the nVista 3.0 system (Inscopix Inc) for imaging the activity of neurons expressing

GCamp6m in the vTS over the three-day fear conditioning paradigm. Fluorescent images were

acquired at 30 Hz sequentially from three focal planes, resulting in an effective sampling rate of 10 Hz per plane. The intensity of the excitation LED in the nVista3.0 miniature microscope was set to $1 - 1.5$ mW/mm$^2$, and the camera gain was adjusted to achieve pixel values within the dynamic range of the CCD. The TTL pulses delivered from the behavioral setup (Med Associates) were digitized by the nVista 3.0 system to obtain synchronization between the mouse behavior and Ca$^{2+}$-imaging data.

The initial processing of *in-vivo* Ca$^{2+}$-imaging data was done using the Inscopix Data Processing Software (IDPS; Inscopix Inc). This included: 1) deinterleaving of the videos into the frames taken at individual focal planes; 2) spatial filtering; 3) motion correction; 4) export of the processed videos as TIFF image stacks; 5) export of the timestamps for each acquired frame, and for the experimental events such as CS and US timing. Next, the TIFF stacks were processed (except a movement correction step that was done by IDPS) using a Python-based package CaImAn (Giovannucci et al., 2019). The package is specifically optimized for the analysis of wide-field microendoscopic fluorescent Ca$^{2+}$-imaging data using CNMF-E, an adaptation of constrained nonnegative matrix factorization algorithm (Zhou et al., 2018). Detection of ROIs for each focal plane was performed by CaImAn in an unsupervised manner, using the same set of analysis parameters across the three experimental days. This resulted in a set of background-corrected fluorescence traces for each automatically detected neuron, and also included a deconvolution step (Pnevmatikakis et al., 2016). The deconvolution returned the times, and amplitude values of Ca$^{2+}$ events (see Figure 1T, grey vertical bars). The amplitudes of the deconvolved events were cumulated for each imaged neuron and each time epoch, to derive the amplitude-weighted frequency of Ca$^{2+}$ events in one of four defined time epochs (during freezing, and in the absence of a CS [Frz_noCS]; during movement, and in the absence of a CS [Mov_noCS]; during freezing, and in the presence of a 30-s CS block [Frz_CS]; and during movement, and in the presence of a 30-s CS block [Mov_CS]).

Following the CaImAn analysis, the data were analyzed using custom routines in IgorPro 7

(WaveMetrics) as follows. Fluorescence intensity traces for each i-th cell ($F_i(t)$), were

standardized by calculating Z-score traces as $Z_i(t) = \frac{F_i(t) - \langle F_i(t) \rangle}{\sigma(F_i(t))}$, where $\langle F_i(t) \rangle$ and $\sigma(F_i(t))$ are

the mean and the standard deviation of the fluorescence intensity, respectively, calculated from the

whole trace. Accordingly, the deconvolution traces were also normalized by the standard deviation

of the respective fluorescent trace $\sigma(F_i(t))$.

Prior to further analysis, any duplicate cells arising from different focal planes were identified with

a semi-automated routine. In brief, the candidate duplicate cells were automatically short-listed

based on the lateral proximity of their centers ($< 20$ μm lateral distance) and high temporal cross-

correlation coefficient ($> 0.7$) between their Z-score traces. Rejection of one the duplicate cells

featuring lower intensity signal (i.e. the cells more out of focus than the other) was validated

manually.

The Z-score traces from retained cells were analyzed by their temporal alignments to the onsets of

CS, US and movement -ON events as described in the Results (Figures 1, 3 and their supplements).

Neurons were classified as "responders" to a given event if the time-averaged Z-score value in the

relevant time range after the event onset exceeded the chosen threshold. For US events, the range

was 0 to 1 second and the Z-score threshold was 1.0. For the CS and movement - ON events, the

range was 0 to 0.5 seconds and the threshold set to 0.2.

*Rabies tracing*

For rabies tracing experiments (Figure 7) *Drd1a^{Cre}* or *Adora2a^{Cre}* mice were injected into the vTS

(see above for coordinates) with a tricistronic vector; AAV1:hSyn:FLEX:TVA-2a-eGFP-2a-oG

(250 nl; 5.3 x10$^{12}$ vg/ml; cat. v243-1; viral vector facility University of Zürich) to render cells

competent for EnvA-pseudotyped rabies virus uptake (Wickersham et al., 2007; Wall et al., 2013).

In earlier experiments, we used a mix of AAVs (AAV8:hSyn:FLEX:TVA-2a-oG and

AAV8:EF1α:FLEX:H2B-GFP-2a-oG; 250 nl; 1:1) for the same purpose. Three weeks later, the

rabies vector EnvA:deltaG–RV-dsRed was injected at the same coordinates (250 nl; viral vector

core Salk Institute for Biological Studies, La Jolla, CA, USA; Osakada et al., 2011). The animals

were sacrificed 7 days later, and a histological analysis was performed on every second 40 µm

thick coronal section of the entire brain, from the level of the prefrontal cortex up to the end of the

cerebellum. The resulting images were analysed in a semi-automatic fashion, i.e. dsRed positive

neurons were marked manually, brain sections were registered to the Allen brain atlas and marked

neurons were automatically mapped onto the resulting brain regions using Matlab-based software

(SHARP-Track; Shamash et al., 2018). The positioning of all long-range projecting cells was

analysed and plotted on a 3D brain model using the python-based Brainrender software (Claudi et

al., 2020).

*Electrophysiology*

For whole-cell patch-clamp electrophysiology in slices, mice that had undergone surgery for AAV

vector injection (see above) were sacrificed 3 - 6 weeks later. Mice were deeply anesthetized with

isoflurane, and decapitated. The brain was quickly removed from the skull and placed in ice-cold

preparation solution; the subsequent procedures followed the general method of (Ting et al., 2014).

The preparation solution contained (in mM): 110 N-methyl-D-glutamine, 2.5 KCl, 1.2 $NaH_2PO_4$,

20 HEPES, 25 Glucose, 5 Na- ascorbate, 2 Thiourea, 3 sodium pyruvate, 10 $MgCl_2$, 0.5 $CaCl_2$,

saturated with carbogen gas ($O_2$ 95%/$CO_2$ 5%), pH of 7.4 adjusted with HCl. Coronal slices (300

µm) containing the vTS were cut using a Leica VT1200S slicer (Leica Microsystems, Wetzlar,

Germany). Slices were stored for 7 min at 36°C in the preparation solution and were then placed in

a chamber containing a storage solution, composed of (in mM): 92 NaCl, 2.5 KCl, 30 $NaHCO_3$, 1.2

$NaH_2PO_4$, 20 HEPES, 25 glucose, 5 sodium ascorbate, 2 Thiourea, 3 Na-pyruvate, 2 $MgCl_2$ and 2

CaCl$_2$, pH 7.4 at room temperature, saturated with carbogen (Ting et al., 2014). Whole-cell patch-

clamp recordings were performed with an extracellular solution containing (in mM): 125 NaCl, 2.5

KCl, 25 NaHCO$_3$, 1.2 NaH$_2$PO$_4$, 25 glucose, 0.4 Na-ascorbate, 3 Myo-Inositol, 2 Na-pyruvate, 1

MgCl$_2$ and 2 CaCl$_2$, pH7.4, saturated with carbogen gas. The set-up was equipped with an EPC10/2

patch-clamp amplifier (HEKA Elektronik GmbH, Reutlingen, Germany), and an upright

microscope (BX51WI; Olympus, Tokyo, Japan) with a 60x / 0.9 NA water-immersion objective

(LUMPlanFl, Olympus).

Patch-clamp experiments (Figure 8, 9) were performed using a Cs$^+$-based intracellular solution (in

mM): 140 Cs$^+$-gluconate, 10 HEPES, 8 TEA-Cl, 5 Na-phosphocreatine, 4 Mg-ATP, 0.3 Na-GTP, 5

EGTA, pH 7.2 adjusted with CsOH. AMPA/NMDA - ratio experiments were done in the presence

of 5μM GABA$_A$ receptor antagonist gabazine (SR-95531; Abcam, Cambridge, UK). Experiments

to test the archaerhodopsin (Arch) properties (Figure 5 - figure supplement 2) were performed

using a K$^+$-based solution with (in mM): 8 KCl, 145 K-gluconate, 10 HEPES, 3 Na-

phosphocreatine, 4 Mg ATP, 0.3 Na-GTP, 5 EGTA, pH 7.2 adjusted with KOH. All

electrophysiological experiments were conducted at near-physiological temperature (34°C) using

an inlet heater SHM-6, a heated recording chamber RC-26GL/PM-1 and a thermostatic control unit

TC-344B (all from Warner Instruments, Holliston, MA, USA). All chemicals, unless indicated,

were from Sigma-Aldrich (St. Louis, MO, USA).

For activation of the excitatory and inhibitory opsins in slice experiments, and for visualization of

fluorophores in brain slices, high-power LEDs (CREE XP-E2, 460 nm and 530 nm; Cree Inc.,

Durham, NC, USA) were custom-coupled into the epifluorescence port of the microscope.

Illumination was controlled by the EPC 10/2 amplifier DAC board connected to the LED driver

(BLS-1000-2, Mightex Systems, Toronto, Canada). Irradiance was measured by a photodiode

(DET36A/M, Thorlabs) coupled into the illumination light path, whose readings were calibrated by

the light power measured under the 60x objective using a power-meter model 1918-R equipped

with a 818-UV detector (NewPort, Irvine, CA, USA). Electrophysiological recordings were

analyzed in IgorPro (WaveMetrics) using the NeuroMatic plug-in (Rothman and Silver, 2018).

*Histology*

For anatomical analysis of optic fiber - or GRIN lens positions, mice were transcardially perfused

with a 4% paraformaldehyde (PFA) solution. The brains were post-fixed in PFA overnight and then

transferred to 30% sucrose in phosphate - buffered solution for dehydration. Coronal brain sections

of 40 µm thickness were prepared using a HM450 sliding microtome (Thermo Fisher Scientific,

Waltham, MA, USA). Slices were mounted on Superfrost® Plus slides (Thermo Fisher Scientific)

and embedded in Fluoroshield mounting medium containing DAPI to stain cell nuclei (Sigma-

Aldrich). Slices were imaged with a slide scanning fluorescent microscope VS120-L100 (Olympus)

with a 10x/0.4 NA objective, or with a confocal microscope (Leica SP8). Brain atlas overlays are

taken from (Franklin and Paxinos, 2016). The majority of brain structure names and their

abbreviations follows (Franklin and Paxinos, 2016); their correspondence to the Allen brain atlas

are given in Figure 7, Table Suppl. 1.

*Statistical analysis*

Statistical analysis was performed in GraphPad Prism 9 (GraphPad, San Diego, CA, USA). Before

choosing the main statistical test, the distribution of the data was tested for normality using a

Shapiro-Wilk test. When normality was confirmed, we used a paired or unpaired version of the

two-tailed Student's t-tests for two-sample datasets, as indicated. For the comparison of relative

datasets, we used a one-sample t-test. When the data was not normally distributed, we used two-

tailed non-parametric tests: a Wilcoxon matched-pairs signed-rank test for paired comparisons, or

Mann-Whitney U test for unpaired comparisons of two-sample datasets, as indicated.

For datasets with more than two samples (*in-vivo* $Ca^{2+}$ imaging data; Figures 1, 3), the data showed

skewed distributions which did not pass the normality test, therefore we used a non-parametric

version of a one-way ANOVA, a Kruskal-Wallis test. If this test detected significant differences, it

was followed by Dunn's post-hoc test for multiple comparisons (called Dunn's MC test). For

datasets influenced by two factors such as the optogenetic silencing/control group and the time of

the experiment (Figures 5, 6), we used a repeated-measures two-way ANOVA (RM-ANOVA)

separately for the training and fear retrieval days. If RM-ANOVA reported significance, it was

followed by Šidák's post-hoc tests for the respective factor.

The change in number of neurons responding *in-vivo* to different events across training (Figures

1L, 1S, 3I, 3P) was statistically assessed using a Chi-square test. For the optogenetic silencing

experiments, we determined the sample size *a priori* using a G\*Power software (Faul et al., 2007)

for an RM-ANOVA test (two groups, 4 replicate measurements for the fear retrieval day) assuming

an average change in freezing of 20% (from 60 to 40%) and a standard deviation of 15% (resulting

in an effect size of 0.667), significance level $\alpha$=0.05, power 1-$\beta$ = 0.85, and the correlation between

repeated measures of 0.5. The resulting total sample size was N = 16 (N = 8 mice per group), with a

critical $F_{crit}$= 4.6. There was no sample size estimation made for other experiments.

Each specific statistical test is mentioned in the Results text, and the input data along with

statistical test summary (such as p-values, values of test statistics, degrees of freedom, etc.) and the

main descriptive statistics are given in the statistics Tables for each relevant figure. The data are

expressed as mean ± SEM. Statistical significance, if applicable, is indicated in the Figures using

asterisks as $p \leq 0.05$ (\*), $p \leq 0.01$ (\*\*) and $p \leq 0.001$ (\*\*\*).

**Data availability**

The essential raw data leading to the conclusions of this paper will be made available at

10.5281/zenodo.4003654.

**Acknowledgements**

We would like to thank Shriya Palchaudhuri and Denys Osypenko for helpful discussions regarding

the analysis of freezing behavior, and Elena Mombelli for help with anatomical experiments.

Images were acquired at the bio-optical imaging platform of EPFL (BiOP). This work was

supported by an EMBO fellowship to M.K. (# ALTF 224-2015), and by grants from the Swiss

National Science foundation (SNSF; grant number 31003A_176332 / 1) and from the National

Competence Center for Research (NCCR) of the SNSF "Synapsy - The Synaptic Bases of Mental

disease", project P28 (both to R.S.).

## References

Amano, T., Duvarci, S., Popa, D., and Paré, D. (2011). The fear circuit revisited: contributions of the basal amygdala nuclei to conditioned fear. The Journal of neuroscience : the official journal of the Society for Neuroscience *31*, 15481-15489.

Barsy, B., Kocsis, K., Magyar, A., Babiczky, A., Szabó, M., Veres, J.M., Hillier, D., Ulbert, I., Yizhar, O., and Mátyás, F. (2020). Associative and plastic thalamic signaling to the lateral amygdala controls fear behavior. Nature neuroscience *23*, 625-637.

Burguiere, E., Monteiro, P., Mallet, L., Feng, G., and Graybiel, A.M. (2015). Striatal circuits, habits, and implications for obsessive-compulsive disorder. Current opinion in neurobiology *30*, 59-65.

Chow, B.Y., Han, X., Dobry, A.S., Qian, X., Chuong, A.S., Li, M., Henninger, M.A., Belfort, G.M., Lin, Y., Monahan, P.E., and Boyden, E.S. (2010). High-performance genetically targetable optical neural silencing by light-driven proton pumps. Nature *463*, 98-102.

Christianson, J.P., Fernando, A.B., Kazama, A.M., Jovanovic, T., Ostroff, L.E., and Sangha, S. (2012). Inhibition of fear by learned safety signals: a mini-symposium review. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 14118-14124.

Claudi, F., Tyson, L.A., and Branco, T. (2020). Brainrender. A python based software for visualisation of neuroanatomical and morphological data. bioRxiv *https://doi.org/10.1101/2020.02.23.961748*.

Cox, J., and Witten, I.B. (2019). Striatal circuits for reward learning and decision-making. Nature reviews. Neuroscience *20*, 482-494.

Creed, M., Pascoli, V.J., and Lüscher, C. (2015). Addiction therapy. Refining deep brain stimulation to emulate optogenetic treatment of synaptic pathology. Science *347*, 659-664.

Cummings, K.A., and Clem, R.L. (2020). Prefrontal somatostatin interneurons encode fear memory. Nature neuroscience *23*, 61-74.

Dalmay, T., Abs, E., Poorthuis, R.B., Hartung, J., Pu, D.L., Onasch, S., Lozano, Y.R., Signoret-Genest, J., Tovote, P., Gjorgjieva, J., and Letzkus, J.J. (2019). A critical role for neocortical processing of threat memory. Neuron *104*, 1180-1194 e1187.

Davis, M. (1992). The role of the amygdala in fear and anxiety. Annual review of neuroscience *15*, 353-375.

Dunsmoor, J.E., Prince, S.E., Murty, V.P., Kragel, P.A., and LaBar, K.S. (2011). Neurobehavioral mechanisms of human fear generalization. NeuroImage *55*, 1878-1888.

Duvarci, S., and Paré, D. (2014). Amygdala microcircuits controlling learned fear. Neuron *82*, 966-980.

Fanselow, M.S. (1994). Neural organization of the defensive behavior system responsible for fear. Psychonomic bulletin & review *1*, 429-438.

Fanselow, M.S. (2018). The Role of Learning in Threat Imminence and Defensive Behaviors. Curr Opin Behav Sci *24*, 44-49.

Faul, F., Erdfelder, E., Lang, A.G., and Buchner, A. (2007). G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods *39*, 175-191.

Franklin, K.B.J., and Paxinos, G. (2016). The Mouse Brain in stereotaxic coordinates. Elsevier / Academic Press *4th edition*.

Gangarossa, G., Castell, L., Castro, L., Tarot, P., Veyrunes, F., Vincent, P., Bertaso, F., and Valjent, E. (2019). Contrasting patterns of ERK activation in the tail of the striatum in response to aversive and rewarding signals. J Neurochem *151*, 204-226.

Gehrlach, D.A., Dolensek, N., Klein, A.S., Roy Chowdhury, R., Matthys, A., Junghänel, M., Gaitanos, T.N., Podgornik, A., Black, T.D., Reddy Vaka, N., *et al.* (2019). Aversive state processing in the posterior insular cortex. Nature neuroscience *22*, 1424-1437.

Gerfen, C.R., Engber, T.M., Mahan, L.C., Susel, Z., Chase, T.N., Monsma, F.J., Jr., and Sibley, D.R. (1990). D1 and D2 dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons. Science *250*, 1429-1432.

Gerfen, C.R., Paletzki, R., and Heintz, N. (2013). GENSAT BAC cre-recombinase driver lines to study the functional organization of cerebral cortical and basal ganglia circuits. Neuron *80*, 1368-1383.

Ghosh, K.K., Burns, L.D., Cocker, E.D., Nimmerjahn, A., Ziv, Y., Gamal, A.E., and Schnitzer, M.J. (2011). Miniaturized integration of a fluorescence microscope. Nat Methods *8*, 871-878.

Giovannucci, A., Friedrich, J., Gunn, P., Kalfon, J., Brown, B.L., Koay, S.A., Taxidis, J., Najafi, F., Gauthier, J.L., Zhou, P.*, et al.* (2019). CaImAn an open source tool for scalable calcium imaging data analysis. eLife *8*.

Gjoni, E., Zenke, F., Bouhours, B., and Schneggenburger, R. (2018). Specific synaptic input strengths determine the computational properties of excitation-inhibition integration in a sound localization circuit. J Physiol *596*, 4945-4967.

Gogolla, N., Takesian, A.E., Feng, G., Fagiolini, M., and Hensch, T.K. (2014). Sensory integration in mouse insular cortex reflects GABA circuit maturation. Neuron *83*, 894-905.

Gong, S., Doughty, M., Harbaugh, C.R., Cummins, A., Hatten, M.E., Heintz, N., and Gerfen, C.R. (2007). Targeting Cre recombinase to specific neuron populations with bacterial artificial chromosome constructs. The Journal of neuroscience : the official journal of the Society for Neuroscience *27*, 9817-9823.

Graybiel, A.M. (2005). The basal ganglia: learning new tricks and loving it. Current opinion in neurobiology *15*, 638-644.

Grewe, B.F., Gründemann, J., Kitch, L.J., Lecoq, J.A., Parker, J.G., Marshall, J.D., Larkin, M.C., Jercog, P.E., Grenier, F., Li, J.Z.*, et al.* (2017). Neural ensemble dynamics underlying a long-term associative memory. Nature *543*, 670-675.

Grillner, S., Hellgren, J., Menard, A., Saitoh, K., and Wikstrom, M.A. (2005). Mechanisms for selection of basic motor programs - roles for the striatum and pallidum. Trends in neurosciences *28*, 364-370.

Gruene, T.M., Flick, K., Stefano, A., Shea, S.D., and Shansky, R.M. (2015). Sexually divergent expression of active and passive conditioned fear responses in rats. eLife *4*, e11352.

Guarraci, F.A., Frohardt, R.J., and Kapp, B.S. (1999). Amygdaloid D1 dopamine receptor involvement in Pavlovian fear conditioning. Brain Res *827*, 28-40.

Guo, L., Walker, W.I., Ponvert, N.D., Penix, P.L., and Jaramillo, S. (2018). Stable representation of sounds in the posterior striatum during flexible auditory decisions. Nature communications *9*, 1534.

Hikosaka, O., Takikawa, Y., and Kawagoe, R. (2000). Role of the basal ganglia in the control of purposive saccadic eye movements. Physiological reviews *80*, 953-978.

Hintiryan, H., Foster, N.N., Bowman, I., Bay, M., Song, M.Y., Gou, L., Yamashita, S., Bienkowski, M.S., Zingg, B., Zhu, M.*, et al.* (2016). The mouse cortico-striatal projectome. Nature neuroscience *19*, 1100-1114.

Hjorth, J.J.J., Kozlov, A., Carannante, I., Frost Nylen, J., Lindroos, R., Johansson, Y., Tokarska, A., Dorst, M.C., Suryanarayana, S.M., Silberberg, G.*, et al.* (2020). The microcircuits of striatum in silico. Proceedings of the National Academy of Sciences of the United States of America *117*, 9554-9565.

Humphries, M.D., and Prescott, T.J. (2010). The ventral basal ganglia, a selection mechanism at the crossroads of space, strategy, and reward. Progress in neurobiology *90*, 385-417.

Hunnicutt, B.J., Jongbloets, B.C., Birdsong, W.T., Gertz, K.J., Zhong, H., and Mao, T. (2016). A comprehensive excitatory input map of the striatum reveals novel functional organization. eLife *5*.

Janak, P.H., and Tye, K.M. (2015). From circuits to behaviour in the amygdala. Nature *517*, 284-
292.

Keller, G.B., Bonhoeffer, T., and Hubener, M. (2012). Sensorimotor mismatch signals in primary
visual cortex of the behaving mouse. Neuron *74*, 809-815.

Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K., Morimoto,
T.K., Chuong, A.S., Carpenter, E.J., Tian, Z.*, et al.* (2014). Independent optical excitation of
distinct neural populations. Nat Methods *11*, 338-346.

Klaus, A., Alves da Silva, J., and Costa, R.M. (2019). What, if, and when to move: Basal ganglia
circuits and self-paced action initiation. Annual review of neuroscience *42*, 459-483.

Kravitz, A.V., Freeze, B.S., Parker, P.R., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer, A.C.
(2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia
circuitry. Nature *466*, 622-626.

Lamont, E.W., and Kokkinidis, L. (1998). Infusion of the dopamine D1 receptor antagonist SCH
23390 into the amygdala blocks fear expression in a potentiated startle paradigm. Brain Res
*795*, 128-136.

Lebron-Milad, K., and Milad, M.R. (2012). Sex differences, gonadal hormones and the fear
extinction network: implications for anxiety disorders. Biol Mood Anxiety Disord *2*, 3.

LeDoux, J.E. (2000). Emotion circuits in the brain. Annual review of neuroscience *23*, 155-184.

LeDoux, J.E., Cicchetti, P., Xagoraris, A., and Romanski, L.M. (1990). The lateral amygdaloid
nucleus: sensory interface of the amygdala in fear conditioning. J Neuroscience *10*, 1062-1069.

LeDoux, J.E., Iwata, J., Cicchetti, P., and Reis, D.J. (1988). Different projections of the central
amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. The
Journal of neuroscience : the official journal of the Society for Neuroscience *8*, 2517-2529.

Little, J.P., and Carter, A.G. (2013). Synaptic mechanisms underlying strong reciprocal
connectivity between the medial prefrontal cortex and basolateral amygdala. The Journal of
neuroscience : the official journal of the Society for Neuroscience *33*, 15333-15342.

Litvina, E.Y., and Chen, C. (2017). Functional Convergence at the Retinogeniculate Synapse.
Neuron *96*, 330-338 e335.

Livneh, Y., Ramesh, R.N., Burgess, C.R., Levandowski, K.M., Madara, J.C., Fenselau, H., Goldey,
G.J., Diaz, V.E., Jikomes, N., Resch, J.M.*, et al.* (2017). Homeostatic circuits selectively gate
food cue responses in insular cortex. Nature *546*, 611-616.

Lucas, E.K., Jegarl, A.M., Morishita, H., and Clem, R.L. (2016). Multimodal and site-specific
plasticity of amygdala parvalbumin interneurons after fear learning. Neuron *91*, 629-643.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter,
R.D., Hawrylycz, M.J., Jones, A.R.*, et al.* (2010). A robust and high-throughput Cre reporting
and characterization system for the whole mouse brain. Nature neuroscience *13*, 133-140.

Maren, S., De Oca, B., and Fanselow, M.S. (1994). Sex differences in hippocampal long-term
potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP
and contextual learning. Brain Res *661*, 25-34.

Markowitz, J.E., Gillis, W.F., Beron, C.C., Neufeld, S.Q., Robertson, K., Bhagat, N.D., Peterson,
R.E., Peterson, E., Hyun, M., Linderman, S.W.*, et al.* (2018). The Striatum Organizes 3D
Behavior via Moment-to-Moment Action Selection. Cell *174*, 44-58 e17.

Menegas, W., Akiti, K., Amo, R., Uchida, N., and Watabe-Uchida, M. (2018). Dopamine neurons
projecting to the posterior striatum reinforce avoidance of threatening stimuli. Nature
neuroscience *21*, 1421-1430.

Menegas, W., Bergan, J.F., Ogawa, S.K., Isogai, Y., Umadevi Venkataraju, K., Osten, P., Uchida,
N., and Watabe-Uchida, M. (2015). Dopamine neurons projecting to the posterior striatum form
an anatomically distinct subclass. eLife *4*, e10032.

Morey, R.A., Haswell, C.C., Stjepanovic, D., Mid-Atlantic, M.W., Dunsmoor, J.E., and LaBar, K.S. (2020). Neural correlates of conceptual-level fear generalization in posttraumatic stress disorder. Neuropsychopharmacology *45*, 1380-1389.

Nader, K., and LeDoux, J.E. (1999). Inhibition of the mesoamygdala dopaminergic pathway impairs the retrieval of conditioned fear associations. Behavioral neuroscience *113*, 891-901.

Nelson, A.B., and Kreitzer, A.C. (2014). Reassessing models of basal ganglia function and dysfunction. Annual review of neuroscience *37*, 117-135.

Niell, C.M., and Stryker, M.P. (2010). Modulation of visual responses by behavioral state in mouse visual cortex. Neuron *65*, 472-479.

Osakada, F., Mori, T., Cetin, A.H., Marshel, J.H., Virgen, B., and Callaway, E.M. (2011). New rabies virus variants for monitoring and manipulating activity and gene expression in defined neural circuits. Neuron *71*, 617-631.

Palchaudhuri, S., Osypenko, D., and Schneggenburger, R. (2022). Fear Learning: An Evolving Picture for Plasticity at Synaptic Afferents to the Amygdala. Neuroscientist, 10738584221108083.

Pennington, Z.T., Dong, Z., Feng, Y., Vetere, L.M., Page-Harley, L., Shuman, T., and Cai, D.J. (2019). ezTrack: An open-source video analysis pipeline for the investigation of animal behavior. Sci Rep *9*, 19979.

Petreanu, L., Huber, D., Sobczyk, A., and Svoboda, K. (2007). Channelrhodopsin-2-assisted circuit mapping of long-range callosal projections. Nature neuroscience *10*, 663-668.

Phelps, E.A., and LeDoux, J.E. (2005). Contributions of the amygdala to emotion processing: from animal models to human behavior. Neuron *48*, 175-187.

Pnevmatikakis, E.A., Soudry, D., Gao, Y., Machado, T.A., Merel, J., Pfau, D., Reardon, T., Mu, Y., Lacefield, C., Yang, W.*, et al.* (2016). Simultaneous denoising, deconvolution, and demixing of calcium imaging data. Neuron *89*, 285-299.

Pryce, C.R., Lehmann, J., and Feldon, J. (1999). Effect of sex on fear conditioning is similar for context and discrete CS in Wistar, Lewis and Fischer rat strains. Pharmacol Biochem Behav *64*, 753-759.

Quirk, G.J., Repa, C., and LeDoux, J.E. (1995). Fear conditioning enhances short-latency auditory responses of lateral amygdala neurons: parallel recordings in the freely behaving rat. Neuron *15*, 1029-1039.

Redgrave, P., Rodriguez, M., Smith, Y., Rodriguez-Oroz, M.C., Lehericy, S., Bergman, H., Agid, Y., DeLong, M.R., and Obeso, J.A. (2010). Goal-directed and habitual control in the basal ganglia: implications for Parkinson's disease. Nature reviews. Neuroscience *11*, 760-772.

Rodgers, K.M., Benison, A.M., Klein, A., and Barth, D.S. (2008). Auditory, somatosensory, and multisensory insular cortex in the rat. Cerebral cortex *18*, 2941-2951.

Rogan, M.T., Leon, K.S., Perez, D.L., and Kandel, E.R. (2005). Distinct neural signatures for safety and danger in the amygdala and striatum of the mouse. Neuron *46*, 309-320.

Romanski, L.M., Clugnet, M.C., Bordi, F., and LeDoux, J.E. (1993). Somatosensory and auditory convergence in the lateral nucleus of the amygdala. Behavioral neuroscience *107*, 444-450.

Rothman, J.S., and Silver, R.A. (2018). NeuroMatic: An integrated open-source software toolkit for acquisition, analysis and simulation of electrophysiological data. Frontiers in neuroinformatics *12*, 14.

Rothwell, P.E., Hayton, S.J., Sun, G.L., Fuccillo, M.V., Lim, B.K., and Malenka, R.C. (2015). Input- and output-specific regulation of serial order performance by corticostriatal circuits. Neuron *88*, 345-356.

Sacco, T., and Sacchetti, B. (2010). Role of secondary sensory cortices in emotional memory storage and retrieval in rats. Science *329*, 649-656.

Sangha, S., Diehl, M.M., Bergstrom, H.C., and Drew, M.R. (2020). Know safety, no fear. Neuroscience and biobehavioral reviews *108*, 218-230.

Sawatari, H., Tanaka, Y., Takemoto, M., Nishimura, M., Hasegawa, K., Saitoh, K., and Song, W.J. (2011). Identification and characterization of an insular auditory field in mice. The European journal of neuroscience *34*, 1944-1952.

Schiffmann, S.N., and Vanderhaeghen, J.J. (1993). Adenosine A2 receptors regulate the gene expression of striatopallidal and striatonigral neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *13*, 1080-1087.

Shamash, P., Carandini, M., Harris, K., and Steinmetz, N. (2018). A tool for analyzing electrode tracks from slice histology. bioRxiv *http://dx.doi.org/10.1101/447995*.

Shan, Q., Ge, M., Christie, M.J., and Balleine, B.W. (2014). The acquisition of goal-directed actions generates opposing plasticity in direct and indirect pathways in dorsomedial striatum. The Journal of neuroscience : the official journal of the Society for Neuroscience *34*, 9196-9201.

Sparta, D.R., Stamatakis, A.M., Phillips, J.L., Hovelso, N., van Zessen, R., and Stuber, G.D. (2012). Construction of implantable optical fibers for long-term optogenetic manipulation of neural circuits. Nat Protoc *7*, 12-23.

Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D. (2019). Spontaneous behaviors drive multidimensional, brainwide activity. Science *364*, 255.

Tang, W., Kochubey, O., Kintscher, M., and Schneggenburger, R. (2020). A VTA to basal amygdala dopamine projection contributes to signal salient somatosensory events during fear learning. The Journal of neuroscience : the official journal of the Society for Neuroscience *40*, 3969-3980.

Ting, J.T., Daigle, T.L., Chen, Q., and Feng, G. (2014). Acute brain slice methods for adult and aging animals: application of targeted patch clamp analysis and optogenetics. Methods in molecular biology *1183*, 221-242.

Tovote, P., Esposito, M.S., Botta, P., Chaudun, F., Fadok, J.P., Markovic, M., Wolff, S.B., Ramakrishnan, C., Fenno, L., Deisseroth, K.*, et al.* (2016). Midbrain circuits for defensive behaviour. Nature *534*, 206-212.

Tovote, P., Fadok, J.P., and Lüthi, A. (2015). Neuronal circuits for fear and anxiety. Nature reviews. Neuroscience *16*, 317-331.

Tritsch, N.X., and Sabatini, B.L. (2012). Dopaminergic modulation of synaptic transmission in cortex and striatum. Neuron *76*, 33-50.

Valjent, E., and Gangarossa, G. (2021). The tail of the striatum: From anatomy to connectivity and function. Trends in neurosciences *44*, 203-214.

Wall, N.R., De La Parra, M., Callaway, E.M., and Kreitzer, A.C. (2013). Differential innervation of direct- and indirect-pathway striatal projection neurons. Neuron *79*, 347-360.

Weinberger, N.M. (2007). Auditory associative memory and representational plasticity in the primary auditory cortex. Hearing research *229*, 54-68.

Wickersham, I.R., Lyon, D.C., Barnard, R.J., Mori, T., Finke, S., Conzelmann, K.K., Young, J.A., and Callaway, E.M. (2007). Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. Neuron *53*, 639-647.

Yamamoto, S., Monosov, I.E., Yasuda, M., and Hikosaka, O. (2012). What and where information in the caudate tail guides saccades to visual objects. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 11005-11016.

Yin, H.H., Mulcare, S.P., Hilario, M.R., Clouse, E., Holloway, T., Davis, M.I., Hansson, A.C., Lovinger, D.M., and Costa, R.M. (2009). Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill. Nature neuroscience *12*, 333-341.

Zanette, L.Y., White, A.F., Allen, M.C., and Clinchy, M. (2011). Perceived predation risk reduces the number of offspring songbirds produce per year. Science *334*, 1398-1401.

1116    Zhou, P., Resendez, S.L., Rodriguez-Romaguera, J., Jimenez, J.C., Neufeld, S.Q., Giovannucci, A.,
1117       Friedrich, J., Pnevmatikakis, E.A., Stuber, G.D., Hen, R.*, et al.* (2018). Efficient and accurate
1118       extraction of in vivo calcium signals from microendoscopic video data. eLife *7*.

1119

 **Figure Legends**

 **Figure 1**. **Miniature Microscope Ca$^{2+}$-imaging of D1R+ neurons in the vTS reveals coding for**

 **tones and movement during fear learning.**

(**A**) Outline of the fear learning protocol. (**B**) Experimental scheme of the injection of an AAV

vector and placement of the GRIN lens in the vTS of *Drd1a$^{Cre}$* mice. (**C**) Posthoc fluorescence

microscopy image from an injected *Drd1a$^{Cre}$* mouse expressing GCaMP6m (green channel; blue

channel: DAPI). The black region indicates the position of the GRIN lens. The putative imaging

area is depicted with a red rectangle. Scalebar, 500 µm. (**D**) tdTomato expression in *Drd1a$^{Cre}$* x

*Rosa26$^{LSL-tdTomato}$* mice indicates localization of D1R+ neurons in the vTS, but not in neighboring

cortical - nor amygdalar structures. (**E - G**) Movement traces - and freezing-state of an example

mouse (red traces and light blue areas, respectively), Z-scored Ca$^{2+}$ traces for three example

neurons (black traces); and color-coded Z-scored Ca$^{2+}$ traces for all neurons in one example mouse

(bottom). Data are from the fourth CS presentation of day 1 (E), the fourth CS-US presentation of

day 2 (F), and the second CS presentation on day 3 (G). (**H**) Color-coded Z-scored Ca$^{2+}$ responses

to footshocks in all imaged D1R+ neurons from N = 8 mice. Responses with Z > 1 in a time

interval of [0; 1s] were considered as significant (see white dashed line). The bottom panel shows

the average ± S.E.M. of Ca$^{2+}$ traces for all responders (n = 45 neurons; black trace ± grey shades),

and the average ± S.E.M. across all neurons (n = 163; dashed black trace ± grey shades). (**I - K**) Z-

scored Ca$^{2+}$ traces aligned to tone beeps (CS) during the habituation day, training day, and recall

day (I, J, K, respectively). Responses with an average Z-score > 0.2 in the time interval [0; 0.5s]

were considered significant. The black traces in the lower panel follow the same logics as in (H).

(**L**) Percentage of tone - responsive neurons for each day. (**M - O**) Color-coded Z-scored Ca$^{2+}$

traces aligned to the movement - ON events, analyzed at times in-between CS blocks, for the

habituation -, training - and recall days, as indicated (*top* panel). The traces in the bottom panel

were analyzed as in (H). (**P - R**) Ca$^{2+}$ traces aligned to the movement - ON events, analyzed during

the 30s tone blocks (CS), for the habituation -, training - and recall days (P, Q, and R respectively).

**(S)** Percentage of movement - ON responding neurons during the CS, and in the absence of a CS

(closed, and open symbols, respectively). (**T**) Example traces of, from *top* to *bottom*, times of tone

beeps (blue trace); movement index (red trace); and $Ca^{2+}$ traces from four example neurons in one

mouse; times of freezing are highlighted by light blue. The detection of $Ca^{2+}$ - events and their

amplitudes by a deconvolution analysis is indicated by vertical bars (see Materials and methods).

**(U)** The amplitude - weighted frequency of $Ca^{2+}$ events (average ± S.E.M.) is plotted separately for

the four combinations of CS / no CS epochs, and movement / freezing states of the mice, for the

habituation, training, and fear memory recall day. The presence of a CS is indicated by the blue

bars. For statistical parameters, see Results text.

1155

See also:

Figure supplement 1. Localization of Cre-expressing cells in the vTS of *Drd1a$^{Cre}$* x *Rosa26$^{LSL-}$*

*$^{tdTomato}$* mice

Figure supplement 2. Correction of movement - ON responses for CS - responses

Figure supplement 3. Correction of movement - ON responses for CS - responses

Figure supplement 4. Individual data points and statistical comparison for the $Ca^{2+}$ event

frequencies for four different conditions.

1163

**Figure 2. Spatial localization of the imaged D1R+ vTS neurons, and overlap of neurons**

**coding for sensory events and movement state.**

(**A-C**) Maps with the position of all imaged D1R+ vTS neurons, plotted as projection on the

horizontal plane. Shown are the footshock-responses (A), the tone (CS) responses as imaged on

day3 (B), and the movement - onset responses imaged on day 3 (C). Neurons are drawn in red

when their response was considered significant (average Z-score > 1 for footshock responses;

average Z-score > 0.2 for tone, and movement - ON responses; note different scales of the circles).

For a coarse orientation, outlines of the amygdalar nuclei (basolateral amygdala; "BLA" and central

amygdala, "CeA") and cortex ("Ctx") are shown. (**D**) Venn diagrams showing the overlap of

neuronal populations within the D1R+ vTS neurons that respond to tones (green), to movement -

ON events (blue), and to footshock stimulation (red; on day 2 only). Note the increased number of

neurons responding to tones and movement - onset during fear memory recall, and the overlap of

these sub-populations (right panel).

1177

1178 See also:

1179 Figure supplement 1. Maps with the position of the imaged D1R+ neurons, plotted as a projection

1180 onto the coronal plane.

1181

**Figure 3. Adora+ neurons in the vTS code for footshocks and movement onset, but CS coding is less present.**

(**A**) Experimental scheme of the injection of an AAV vector and placement of the GRIN lens in *Adora2a$^{Cre}$* mice. (**B-D**) Movement traces - and freezing-state of an example mouse (red traces and light blue areas, respectively), Z-scored Ca$^{2+}$ traces for three example neurons (black traces); and color-coded Z-scored Ca$^{2+}$ traces for all neurons in one example mouse (bottom). Data are from the second CS presentation of day 1 (D), the fourth CS-US presentation of day 2 (E), and the second CS presentation on day 3 (F). (**E**) Z-scored Ca$^{2+}$ responses to footshocks in all imaged Adora+ neurons from N = 8 mice; responses with average Z > 1 in the interval of [0; 1s] were considered as significant (traces above dashed white line). The bottom panel shows the average ± S.E.M. of Ca$^{2+}$ traces for all responders (n = 79 neurons; black trace ± grey shades), and the average ± S.E.M. across all neurons (n = 201; dashed black trace ± grey shades). (**F - H**). Color-coded Z-scored Ca$^{2+}$ traces aligned to tone beeps (CS) during the habituation day, training day, and recall day (F, G, H, respectively). Responses with average Z-score > 0.2 in the time interval of [0; 0.5s] were considered significant; the black traces in the lower panel were calculated as in (**E**). (**I**) Percentage of tone - responsive neurons for each day. (**J - L**). Ca$^{2+}$ responses to movement - ON events, analyzed outside the CS blocks, for the habituation -, training - and recall days (J, K, and L, respectively). The top - and bottom panels follow the same logics as in (E). (**M - O**) Ca$^{2+}$ responses to movement - ON events, analyzed during the 30s tone blocks (CS), for the habituation, training, and recall days as indicated. (**P**) Percentage of movement - ON responders in the presence - and absence of a CS (closed, and open symbols). (**Q**) Illustration of the Ca$^{2+}$ deconvolution approach for four example neurons in a Adora+ mouse. From *top* to *bottom*, times of tone beeps (blue trace); movement index (red trace); and Ca$^{2+}$ traces from four example neurons; times of freezing are highlighted by light blue. Vertical grey bars indicate the timing and amplitude of the detected Ca$^{2+}$ events. (**R**) The amplitude-weighted frequency of Ca$^{2+}$ events (average ± S.E.M.), analyzed separately for the four combinations of CS / no CS times, and movement / freezing states of the

1208    mice, for the three fear learning days. The presence of a CS is indicated by the blue bars. The p-

1209    values for the indicated statistical comparisons are reported in the Results text. (**S**) Venn diagrams

1210    showing the overlap of neuronal populations within the Adora+ vTS neurons that respond to tones

1211    (green), to movement - onset transitions (blue), and to footshock stimulation.

1212

See also:

Figure supplement 1. Localization of Cre-expressing cells in the vTS of *Adora2a$^{Cre}$* x *Rosa26$^{LSL-}$*

*$^{tdTomato}$* mice

Figure supplement 2. Individual data points and statistical comparison for the Ca$^{2+}$ event

frequencies for four different conditions

Figure supplement 3. Maps with the position and Ca$^{2+}$ signal intensity of the imaged Adora+ vTS

neurons.

1220

1221

1222

1223

**Figure 4. Optogenetic stimulation of D1R+ and Adora+ vTS neurons has no direct effect on movement.**

(**A**) Scheme showing the bilateral placement of optic fibers over each vTS, and the injection of an AAV vector driving the Cre-dependent expression of Chronos. (**B**) $Adora2a^{Cre}$ mice expressing Chronos in Adora+ neurons of the vTS were allowed to explore the fear conditioning chamber, and trains of blue light stimuli (50 pulses of 1 ms length, 25 Hz, duration of 2s) were applied. The average movement indices for N = 8 mice are shown, either centered around the time of light stimulation (red trace), or taken from a control time interval when no light was applied ("ctrl TP"). (**C**) Analogous experiment to the one in (B), now performed for $Drd1a^{Cre}$ mice expressing Chronos in a Cre - dependent manner in the vTS (N= 7). Note that optogenetic stimulation of neither Adora+ neurons (B), nor of D1R+ neurons led to notable changes in the movement of mice (see Results for the statistical parameters).

See also:

Figure supplement 1. Additional optogenetic stimulation experiments with longer pulse widths

**Figure 5. Optogenetic inhibition of D1R+ vTS neurons during footshock reduces the formation of an auditory-cued fear memory.**

(**A**) Scheme showing the bi-lateral injection in $Drd1a^{Cre}$ mice of an AAV vector driving the expression of either Arch or eGFP (for controls), and the placement of optic fibers over each vTS. (**B**) Behavioral paradigm outlining the 3-day fear learning protocol, and the application of yellow light during the footshock presentation on the training day. (**C**) Freezing levels during 30s CS presentations on the habituation day for the control mice (expressing eGFP; black circles), and for Arch - expressing mice (green circles). (**D**) Time - binned average percent freezing traces for control mice (black data) and Arch - expressing mice (green data; bin width is 10 s in both cases). Light blue boxes indicate the 30s CS presentation periods. Grey bars (bottom) indicate epochs of "no CS" analysis. Vertical orange lines indicate the times of footshock presentation. (**E**) Quantification of freezing during the training day for the control mice and Arch - expressing mice (black and green data, respectively). Freezing was averaged during the 30s CS presentations (filled circles, *left*) and for the no CS epochs (open circles, *right*). (**F**) Time-resolved freezing during the cued retrieval on day 3. Note the increases in freezing driven by each CS presentation in control mice (black trace, average of N = 10 eGFP expressing mice), which were smaller in amplitude in Arch - expressing mice (N = 12, green trace). Light blue areas and grey bars (bottom) indicate times of CS presentation, and epochs of "no CS" analysis, respectively. (**G**) Quantification of freezing during the 30s CS presentation (*left*) and during the no CS epochs (*right*). Note the significant reduction of freezing during the CS in the Arch group (see Results for statistical parameters).

See also:

Figure supplement 1. Post - hoc histology of Arch expression and optic fiber placement in $Drd1a^{Cre}$ mice

1269    Figure supplement 2. Properties of light-activated archaerhodopsin in D1R+ - and Adora+ vTS

1270    neurons

1271

**Figure 6**. **Optogenetic inhibition reveals a role of Adora+ vTS neurons in suppressing freezing**

**in the absence of a learned CS.**

(**A**) Scheme of the experimental approach, during which Arch (or eGFP, for controls) was

expressed Cre-dependently in *Adora2a^{Cre}* mice in the vTS, and optic fibers were placed in the vTS

above the injection sites. (**B**) Behavioral paradigm (upper panel), and scheme of the application of

yellow light during the footshock stimulus (US) on the training day (lower panel). (**C**) Freezing

levels during CS presentations on the habituation day for mice expressing eGFP (control group;

black data) and for Arch - expressing mice (red data). (**D**) Time-resolved analysis of freezing for

control mice (black trace, average of N = 8 mice) and Arch - expressing mice (red trace, N = 11

mice). The light blue areas and grey bars (bottom) indicate the time of CS presentation, and the

time windows for "no CS" analysis. (**E**) Quantification of freezing during the CS (left), and during

the 30s "no CS" epochs (right), for control mice (black) and for Arch - expressing mice (red). (**F**)

Time-resolved freezing during the fear memory recall day. The light blue areas and grey bars

(bottom) indicate the time of CS presentation, and the analysis window for "no CS" analysis. (**G**)

Average percent of time spent freezing, analyzed during the 30s CS presentations (left), and during

the no CS epochs (right) of the fear memory retrieval day, for both groups of mice. There was a

trend towards an enhanced freezing in the Arch group at times when no CS was present (right; p =

0.0512; two-way repeated measures ANOVA; see Results for further statistical parameters).

1290

1291  See also:

1292  Figure supplement 1. Post - hoc histology of Arch expression and optic fiber placement in

1293  *Adora2a^{Cre}* mice

1294

**Figure 7. Retrograde transsynaptic tracing of brain-wide inputs to D1R+ and Adora+ vTS neurons.**

(**A**) Scheme of the experimental protocol for rabies-virus mediated transsynaptic tracing. (**B**) *Left*, Example confocal images of the injection site in a *Drd1a^{Cre}* mouse, *Right*, confocal images at a higher magnification taken from the boxed area of the left image. Green, red and blue channels are eGFP-labelled cells expressing the helper viruses (green), rabies-virus expressing cells (red) and DAPI labelling (blue, only shown in the overlay images). Scale bars, 200 µm (*left*) and 50 µm (*right*). (**C, D**) Example widefield epifluorescence images of rabies virus - labelled presynaptic neurons in coronal sections, from a *Drd1a^{Cre}* mouse (C) and an *Adora2a^{Cre}* mouse (D). The abbreviations of brain areas are shown in Figure 7, Suppl. Table 1. Scalebars, 250 µm. (**E, F**) Localizations of presynaptic neurons plotted in a 3D brain model, both for *Drd1a^{Cre}* (E) and *Adora2a^{Cre}* (F) mice. Each brain shows the results of two animals (blue and red dots) with one dataset being artificially mirrored on the other hemisphere for each genotype. For orientation, some brain areas are highlighted (S2 – green, InsCtx – yellow, PF – red, BLA – pink). (**G, H**) Quantification of labelled neurons for each brain area. Upwards and downward data signify cell density and absolute number of cells, quantified for N = 2 mice of each genotype. Data from single animals are plotted as filled grey circles and triangles. (**I, J**) Distribution of presynaptic neurons along the anterior - posterior axis in the insular cortex (I), and in the secondary somatosensory cortex (J). The dashed line indicates the border between the anterior and posterior insular cortex according to (Franklin and Paxinos, 2016).

See also:

Figure supplement 1. Control experiments for the specificity of helper-virus, and rabies virus expression

Figure supplement 2. Example images for transsynaptically retrogradely labelled neurons in the BLA and PF

1321     Figure supplement 3. Analysis of the number of back-labelled neurons in *Adora2a*$^{Cre}$ mice *relative*

1322     to *Drd1a*$^{Cre}$

1323

1324

**Figure 8. pInsCx provides strong excitatory inputs onto D1R+ - and Adora+ neurons in the vTS .**

**(A)** Experimental scheme of injection of an AAV vector driving the expression of Chronos into the pInsCx, and subsequent slice electrophysiology in the vTS. **(B)** *Left*, example fluorescence images of the injection site in the cortex expressing Chonos-eGFP (scalebar, 500 µm); *middle,* overview brightfield image with the position of the patch pipette (black dotted lines) in the vTS; *right*, higher magnification images of a recorded example cell (brightfield, *top*; and tdTomato fluorescence, *bottom*). **(C, D)** *Left*, EPSCs recorded by stimulating with 1 ms light pulses of increasing intensities (blue, photodiode-recorded trace), and *right*, the resulting input-output curve of EPSC amplitude versus light intensity, with data from all recorded cells overlaid. The example cell shown on the left is highlighted in black in the right panel. Data is shown for D1R+ vTS neurons (C, n = 12 recordings) and Adora+ vTS neurons (D, n = 19 recordings). **(E)** Quantification of the maximal amplitude of EPSCs (*left*), and of the EPSC delay (*right*) measured in D1R+ - and Adora+ vTS neurons (n = 12 and n = 19 recordings, respectively). **(F)** Example traces (*left*, and *middle* panel) and quantification of PPR of optogenetically - evoked EPSCs in D1R+ neurons (green; n = 12 recordings) and Adora+ neurons (red; n = 19 recordings).

See also:

Figure supplement 1. Ratio of direct excitation to feedforward inhibition differs at the synapse from pInsCx to D1R+ vTS neurons versus Adora+ neurons

**Figure 9. Fear learning induces long-term plasticity with opposite outcomes at pInsCx synapses onto D1R+ - and Adora+ vTS neurons.**

(**A**) Timeline of the *ex-vivo* optogenetic assessment of plasticity following fear learning. See Results for details. (**B - D**) Measurements of AMPA/NMDA - ratios and PPR in two experimental groups ("CS-only" versus "CS+US pairing") in *Drd1a^Cre* x *Rosa26^LSL-tdTomato* mice, to target the recording of direct pathway vTS neurons. (**B**) NMDA-EPSCs recorded at +50 mV (top traces), and AMPA-EPSCs recorded at -70 mV (bottom traces), recorded in example D1R+ neurons from each experimental group. Vertical dashed line indicates the time of analysis of NMDA-EPSC. (**C**) AMPA-EPSCs (- 70 mV) recorded for the measurement of PPR (50 ms inter-stimulus interval), shown for one example D1R+ neuron from each experimental group. (**D**) Quantification of AMPA/NMDA ratios and PPR recorded in each experimental group of *Drd1a^Cre* x *Rosa26^LSL-tdTomato* mice (CS only, n = 11 recordings; CS+US pairing, n = 12). Note the significantly increased AMPA/NMDA ratio in the CS+US pairing group as compared to the CS-only group (p = 0.001; unpaired t-test), whereas the PPR was unchanged (p = 0.97, unpaired t-test). (**E - G**) Measurements of AMPA/NMDA - ratios and PPR in two experimental groups ("CS-only" versus "CS+US pairing") in *Adora2a^Cre* x *Rosa26^LSL-tdTomato* mice. (**E**) NMDA-EPSCs (top traces), and AMPA-EPSCs (bottom traces) recorded in example Adora+ neurons from each experimental group. (**F**) Example traces for the measurement of PPR of AMPA-EPSCs at - 70 mV, recorded in example Adora+ neurons of each experimental group. (**G**) Quantification of AMPA/NMDA ratios, and paired-pulse ratios recorded in each experimental group of the *Adora2a^Cre* x *Rosa26^LSL-tdTomato* mice (CS only, n = 12 recordings; CS+US pairing, n = 12). Note the significantly increased PPR in the CS+US pairing group as compared to the CS-only group (p = 0.001; Mann-Whitney test). For further statistical parameters, see Results text.

**Figure Legends for Figure Supplements**

**Figure 1 - figure supplement 1**

**Localization of Cre-expressing cells in the vTS of *Drd1a$^{Cre}$* x *Rosa26$^{LSL\text{-}tdTomato}$* mice**

*Drd1a$^{Cre}$* mice were bred with the *Rosa26$^{LSL\text{-}tdTomato}$* reporter line, to drive the expression of tdTomato

in Cre - positive neurons. Shown is every 6th section (40 µm thickness) at three relevant a-p levels

on the level of the posterior striatum (A, B, C; Franklin and Paxinos, 2016). The mouse brain atlas

images of (Franklin and Paxinos, 2016) were overlaid at the indicated Bregma levels. The lower

panels show magnifications, as indicated by dashed boxes in the upper panels. Note that in the

*Drd1a$^{Cre}$* mouse line used for our study (EY217 BAC transgenic mouse line of the GenSAT project,

see Materials and methods), no Cre expression is driven in cortex -, amygdala -, and claustrum

structures laterally and ventrally to the vTS (see also Gerfen et al., 2013), and images for the EY217

BAC transgenic mouse line on the GenSAT website; http://www.gensat.org/). Rather, Cre -

expression is limited to neurons in the vTS (labelled as "CPu", for "caudate putamen", in the mouse

brain atlas overlays). The meaning of the abbreviation of other brain areas is like in (Franklin and

Paxinos, 2016); see also Figure 7 - supplementary table 1.

1385

1386

1387

**Figure 1 - figure supplement 2.**

**Correction of CS responses for movement - ON responses**

**(A - C)** $Ca^{2+}$ responses were aligned to tone beeps as in Figure 1I - K. In addition, a histogram of the movement - ON events during these times was computed (third panel, black histograms). This revealed that on the training day (B) and on the recall day (C), about 10 - 15% of the tones were followed at 0 - 400 ms by a movement (see arrows), indicating that under these conditions, some tones induced movement initiation of the mice. Because movement - ON transitions can by themselves induce an activity of D1R+ neurons (Figure 1M - O), we removed those events that were followed by a movement (around 85 - 90% of all events were *retained*; see exact values at the bottom, averaged across the N = 8 imaged *Drd1a^{Cre}* mice). This led to movement - corrected averaged $Ca^{2+}$ traces that were slightly changed as compared to the average $Ca^{2+}$ traces without omission of events (second panel, red traces; compare with original traces, black). The movement - ON histograms after correction for synchronous movement - ON transitions is also shown (bottom panel, red histograms). The plot of color-coded Z-scored $Ca^{2+}$ responses (top panel) refers to the data *after* correction for movement - ON events. The number of CS - responding neurons were analyzed from this corrected data (top panel, white dashed lines; numbers of responders indicated in red), and used for the analysis in Figure 1L.

**Figure 1 - figure supplement 3.**

**Correction of movement - ON responses for CS - responses**

**(A - C)** $Ca^{2+}$ responses were aligned to movement - ON transitions during the CS (same data as in

Figure 1P - R). In addition, histograms of the CS events relative to the movement - ON events were

computed (third panel, black histograms). This revealed that many movement - ON transitions were

preceded by a tone event (by ~ 200 - 400 ms; arrows), again suggesting that tones likely cause

movement - ON events with a certain probability, a phenomenon especially visible during the

training - recall sessions (see also Figure 1 - figure supplement 2). Because tones can themselves

cause activity in D1R+ neurons (Figure 1), we removed those events which were preceded at 0 -

400 ms by a tone. Despite the relatively high number of coinciding events, there were no marked

changes between the corrected - and non-corrected $Ca^{2+}$ responses (second panel from top,

compared red [corrected] and black [non-corrected] traces). The number of movement - ON

responders was calculated after this correction (see top panel white dashed lines, and red numbers),

and used for the analysis in Figure 1S.

1420

1421

1422

1423

1424

**Figure 1 - figure supplement 4**

**Individual data points and statistical comparison for the Ca$^{2+}$ event frequencies for four different conditions.**

The amplitude - weighted Ca$^{2+}$ event frequency of the data in Figure 1U are re-plotted together with their underlying individual data points from each imaged neuron (n = 176, 163, and 177 on the habituation -, training - and recall day, respectively). Here, different to the meaning of symbols in Figure 1U, the *medians* of each distribution (horizontal bars) and the *25% - 75% inter-quantile range* (vertical bars) are indicated. Abbreviations: Frz_noCS: freezing, in the absence of CS; Mov_noCS: movement, in the absence of CS; Frz_CS: freezing, during the CS; Mov_CS: movement, during the CS.

The p-values of the statistical comparisons are (post-hoc Dunn's multiple comparison tests, following Kruskal-Wallis test; KW = 138.3; p < 0.0001):

1) Habituation day:

Frz_noCS vs Mov_noCS, p < 0.0001; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p > 0.99; Frz_CS vs Mov_CS, p < 0.0001.

2) Training day:

Frz_noCS vs Mov_noCS, p > 0.99; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p = 0.0163; Frz_CS vs Mov_CS, p < 0.0001.

3) Recall day:

Frz_noCS vs Mov_noCS, p = 0.0167; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p > 0.99; Frz_CS vs Mov_CS, p < 0.0001.

See also Statistics table for Figure 1, for more details on p - values and further statistical parameters.

**Figure 2 - figure supplement 1**

**Maps with the position of the imaged D1R+ neurons, plotted as a projection onto the coronal**

**plane.**

Same data as in Figure 2 A - C, but here plotted as a projection onto the coronal plane, to illustrate

the dorso-ventral distribution of imaged neurons. Note that the apparent localization of neurons in

BLA and CeA is an artifact of the projection of the 3D data, and that different a-p locations of these

neurons has to be taken into account (see Figure 2A - C).

1458

1459

**Figure 3 - figure supplement 1**

**Localization of Cre-expressing cells in the vTS of *Adora2a^{Cre}* x *Rosa26^{LSL-tdTomato}* mice**

*Adora2a^{Cre}* mice were bred with the *Rosa26^{LSL-tdTomato}* reporter line, to drive the expression of

tdTomato in Cre - positive neurons. Shown is every 6th section (40 µm thickness) at three relevant

a-p levels on the level of the posterior striatum (A, B, C). Mouse brain atlas images, at their

indicated Bregma levels, were overlaid (Franklin and Paxinos, 2016). Note the localization of

Adora+ neurons in the vTS (labelled "CPu" for caudate putamen), and the presence of axons in the

globus pallidus, external segment (GPe).

1468

**Figure 3 - figure supplement 2**

**Individual data points and statistical comparison for the $Ca^{2+}$ event frequencies for four**

**different conditions.**

The amplitude - weighted $Ca^{2+}$ event frequency of the data in Figure 3R are re-plotted together with

their underlying individual data points from each imaged neuron (n = 173, 201, and 160 on the

habituation -, training - and recall day, respectively). Here, the *medians* of each distribution

(horizontal bars) and the *25% - 75% interquantile ranges* (vertical bars) are indicated.

Abbreviations: Frz_noCS: freezing, in the absence of CS; Mov_noCS: movement, in the absence of

CS; Frz_CS: freezing, during the CS; Mov_CS: movement, during the CS.

The p-values of the statistical comparisons are (post-hoc Dunn's multiple comparison tests,

following Kruskal-Wallis test; KW = 184.6; p < 0.0001):

1) Habituation day:

Frz_noCS vs Mov_noCS, p < 0.0001; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p >

0.99; Frz_CS vs Mov_CS, p < 0.0001.

2) Training day:

Frz_noCS vs Mov_noCS, p < 0.0001; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p >

0.99; Frz_CS vs Mov_CS, p < 0.0001.

3) Recall day:

Frz_noCS vs Mov_noCS, p = 0.0004; Frz_noCS vs Frz_CS, p > 0.99; Mov_noCS vs Mov_CS, p >

0.99; Frz_CS vs Mov_CS, p = 0.012.

See also Statistics table for Figure 1, for more details on p - values and further statistical

parameters.

1491

1492

1493

1494

**Figure 3 - figure supplement 3**

**Maps with the position and Ca$^{2+}$ signal intensity of the imaged Adora+ vTS neurons.**

Data from N = 8 *Adora2a$^{Cre}$* mice is plotted for the US - responses on the training day, tone (CS)

responses on the recall day, and movement - ON responses on the recall day (A, B, and C

respectively). The maps in the upper - and lower panels are plotted as projections onto the

horizontal - and coronal plane, respectively.

**Figure 4 - figure supplement 1**

**Additional optogenetic stimulation experiments with longer pulse widths**

(**A**) Scheme of the experimental approach. In this series of experiments, N = 6 *Adora2a$^{Cre}$* mice

were prepared for bilateral optogenetic stimulation experiments in the vTS in the same manner as

in Figure 4, but longer-durations blue light pulses were used (2 - and 5 ms).

(**B**) Example movement traces of one mouse. Trains of blue light stimuli (2 ms pulse width;

repeated at 25 Hz for 2 s) were triggered by the experimenter; however, the chance that a light

stimulus occurred was determined randomly by the Ethovision XT 13 software at a 50% chance

level.

(**C**) Overlay of movement traces in the same mouse as in (B), separated for events in which a light

stimulus occurred (pink individual traces, and red average trace), and events in which no stimulus

occurred (grey, and black traces). Note that light stimuli (red trace) did not change the movement

index of the mice in any obvious way.

(**D**) Quantification for N = 6 mice. There was no significance difference between the movement

index during trials with light as compared to trials without light, neither for pulses with 2 ms

duration, nor for pulses with 5 ms duration ($t_5 = 0.331$, $p = 0.75$ and $t_5 = 0.27$, $p = 0.80$; paired t-

test).

**Figure 5 - figure supplement 1**

**Post - hoc histology of Arch expression and optic fiber placement in *Drd1a^{Cre}* mice**

Following experiments (Figure 5), the brains of *Drd1a^{Cre}* mice were fixed by transcardial perfusion

with paraformaldehyde (PFA), and 40 µm coronal sections were prepared. The fluorescence of

Arch-eGFP (green channel) and DAPI (blue) was imaged in a slide scanner (see Materials and

methods). The section containing the end point of the optic fiber (see black non-fluorescent hues in

each image) on each brain side for each mouse is shown (N = 12 Arch - expressing *Drd1a^{Cre}* mice

of Figure 5). On each image, the mouse brain atlas image of the estimated corresponding a-p

position was superimposed (see indicated bregma values below each panel; Franklin and Paxinos,

2016). Codes above each panel refer to the mouse identification number.

**Figure 5 - figure supplement 2**

**Properties of light-activated archaerhodopsin in D1R+ - and Adora+ vTS neurons.**

(A) *Drd1a$^{Cre}$* x *Rosa26$^{LSL-tdTomato}$* mice, or *Adora2a$^{Cre}$* x *Rosa26$^{LSL-tdTomato}$* mice were injected with AAV1:CAG:FLEX:Arch-eGFP AAV vector into the vTS , and patch-clamp recordings of Arch - expressing neurons were performed 3 weeks later in slices of the vTS at 34°C. Example of light-induced currents evoked by 300 ms exposures with increasing light (530 nm) intensities in an Adora+ neuron expressing Arch. Maximal light intensity was 15 mW/mm$^2$; the holding potential was -70 mV.

(B) Quantification of the maximal plateau current measured at the end of the light pulse (grey dashed box in A) for D1R+ - (n = 7) and Adora+ vTS neurons (n = 13) expressing Arch.

(C) Example traces recorded in current-clamp showing that light exposure (3 s) was sufficient to block APs generated by a 1s current injection (no light, black trace; with light, red trace). This observation was made in all neurons recorded under current-clamp.

(D-F) Repeated light exposures (300 ms pulses at 1 Hz, 30 times) were applied to test for the efficiency of repeated hyperpolarizations, and for possible rebound APs after light offset. Hyperpolarization (E) and rebound potentials (F) were quantified for D1R+ (n = 6) and Adora+ vTS neurons (n = 13).

**Figure 6 - figure supplement 1**

**Post - hoc histology of Arch expression and optic fiber placement in *Adora2a^Cre* mice**

The post-hoc histological results for N = 11 Arch - expressing *Adora2a^Cre* mice from Figure 6 is shown, in the same display as Figure 5 - figure supplement 1 for *Drd1a^Cre* mice. Note that the ventral half of the tail striatum (thus, the vTS) was targeted at an a-p position of roughly -0.9 mm relative to Bregma (the striatum is labelled as "CPu" in the overlaid brain atlas images of Franklin and Paxinos, 2016).

**Figure 7 - figure supplement 1**

**Control experiments for the specificity of helper-virus, and rabies virus expression**

**(A - D)** The helper virus, AAV8:hSyn:FLEX:TVA-2a-eGFP-2a-oG was injected into the vTS of a *Drd1a^Cre* mouse (A, B), or into the vTS of a C57BL/6J mouse as a negative control (C, D). This resulted in the Cre - dependent expression of eGFP in the *Drd1a^Cre* mouse (B, and insets at higher magnification), whereas the same injection in the C57BL/6J mouse did not induce detectable expression of eGFP (D), thus documenting the specificity of the Cre - dependent expression of the helper viral constructs in *Drd1a^Cre* mice.

**(E, F)** The pseudotyped glycoprotein G-deficient rabies virus, which can drive the expression of dsRed in TVA- and oG expressing cells, was injected into the vTS of a *Drd1a^Cre* mouse, which had *not* undergone a previous injection with the helper AAV8 virus. Choleratoxin (CTB)-Alexa647 was co-injected to document successful targeting of the vTS (F, red channel). No dsRed expression was observed (F, green channel), thus documenting that successful transduction with deltaG-RV, and therefore, retrograde transsynaptic activity of the rabies virus, requires previous Cre-dependent expression of TVA.

**Figure 7 - figure supplement 2**

**Example images for transsynaptically retrogradely labelled neurons in the BLA and PF.**

The data are from the same experiments as shown in Figure 7. The images on the left sides of each panel illustrate a weak density of labelled neurons in the amygdalar region of both a *Drd1a^Cre* (A) and an *Adora2a^Cre* (B) mouse (left images, CeA, BLA, and LA, see also Figure 7 G, H for the quantifications). The images on the right sides illustrate labeling in a confined region of the PF, indicating a notably high *density* of back-labelled neurons in the PF (see also Figure 7 G, H for the quantifications, and Figure 7, Figure Suppl. 1 - Table for the abbreviations of brain areas).

**Figure 7 - figure supplement 3**

**Analysis of the number of back-labelled neurons in *Adora2a^Cre* mice *relative* to *Drd1a^Cre***

The data in Figure 7G, H was re-sorted, to highlight differences in back-labelling between *Drd1a^Cre* and *Adora2a^Cre* mice. Brain areas with the highest relative backlabelling for Adora+ starter neurons in the vTS (as compared to D1R+ starter cells) are shown from *left* to *right*; the relative difference in backlabelling is indicated by numbers. Note that Adora+ vTS neurons, as compared to D1R+ neurons, showed larger proportions of back-labelled presynaptic neurons in the lateral amygdala (LA), auditory cortex (AUD), temporal association cortex (TeA), basolateral amygdala (BLA), central amygdala (CeA), and posterior triangular thalamic nucleus (PoT); interestingly, these areas process auditory- and multimodal sensory information relevant for associative learning (see references in Results). Note that the back-labelling was also higher in the PiL (posterior intralaminar thalamic nucleus), but because there were no back-labelled neurons in *Drd1a^Cre* mice, the PiL was not entered to this list. On the other hand, a higher proportion of neurons providing input to D1R+, as compared to Adora+ vTS neurons, was found in the S1, in the somatosensory thalamus (VPM), motor cortex (MO) and piriform cortex (PIR).

**Figure 7 - Supplementary Table 1.**

Definition of brain area abbreviations, and correspondence of the names across the Allen Brain

Atlas and the Mouse Brain Atlas of Franklin and Paxinos (2016).

**Figure 8 - figure supplement 1**

**Ratio of direct excitation to feedforward inhibition differs at the synapse from pInsCx to**

**D1R+ vTS neurons versus Adora+ neurons**

Optogenetically-evoked EPSCs were measured at -70 mV, and subsequently recordings were made

at the reversal potential for the EPSCs (+5 to +15 mV across experiments). This revealed robust

optogenetically-evoked inhibitory postsynaptic currents (IPSCs). The *left* panel shows example

traces (green: for a $Drd1a^{Cre}$ x $Rosa26^{LSL-tdTomato}$ mouse; red, for a $Adora2a^{Cre}$ x $Rosa26^{LSL-tdTomato}$

mouse). The middle panel shows the quantification of the E/I ratio, which was significantly larger

in D1R+ neurons than in Adora+ neurons (U = 31, p = 0.0054, Mann Whitney test; n = 12 D1R+

and n = 14 Adora+ vTS neurons). The right panel shows the quantification of latency between the

start of the light pulse and the respective PSC (*green data points*, for n = 12 D1R+ neurons; red

data points, for n = 14 Adora+ vTS neurons). For each neuron type, the delays for the EPSCs

(direct excitation) were significantly shorter than the delays for IPSCs; we therefore interpret the

latter as feedforward inhibition (D1R+ neurons: $t_{11}$ = 5.831, p = 0.0001, paired t-test; Adora+

neurons: w = 153, p < 0.0001, Wilcoxon matched-pairs signed-rank test). Data are shown as

average ± SEM.

Figure 1

**A** Footshock response (Day2)

**B** Tone response (Day3)

**C** Move-ON response (Day3, during CS)

**D**

Habituation

Tone (6)   2   Move (17)

Total = 176

Training

Tone (17)   Move (34)

US (45)

Total = 163

Recall

Tone (19)   14   Move (30)

Total = 177

Figure 2

Figure 3

**A**

AAV1:hsyn:FLEX:Chronos-eGFP
in *Drd1a^Cre* or *Adora2a^Cre*

injection site : ventral tail striatum

**B** *Adora2a^Cre*

light

Movement (A.U.)

—— light
—— ctrl TP

time (s)

**C** *Drd1a^Cre*

light

Movement (A.U.)

—— light
—— ctrl TP

time (s)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

*Drd1a^{Cre}* x *Rosa26^{LSL-tdTomato}*

Figure1 - Figure supplement 1

**CS response**

**A** Habituation

**B** Training

**C** Recall

Figure 1 - figure supplement 2

Move-ON response during CS

A Habituation    B Training    C Recall

average Ca²⁺(Z)
CS events
CS events with correction
time (s)

Move-ON events retained after correction
52.9 ± 4.0%    42.6 ± 5.0%    43.7 ± 4.8%

Figure 1 - figure supplement 3

A

Figure 1 - figure supplement 4

Figure 2 - figure supplement 1

*Adora2a$^{Cre}$ x Rosa26$^{LSL-tdTomato}$*

Figure 3 - figure supplement 1

A

Figure 3 - figure supplement 2

A  Footshock response (Day2)

B  Tone response (Day3)

C  Move-ON response (Day3, during CS)

Figure 3 - figure supplement 3

A

AAV1:hsyn:FLEX:Chronos-eGFP

*Adora2a^{Cre}*

B FV9017

□ manual trigger
■ light on at 50% chance

movement index (a.u.)

time (s)

C FV9017

25 Hz/5 ms/10 mW
OR no light

—— trials with light
—— trials w/o light

movement index (a.u.)

time (s)

D

■ ■ trials with light
■ ■ trials w/o light

av. movement index (a.u.)

2 ms    5 ms
pulse duration

Figure 4 - figure supplement 1

Figure 5 - figure supplement 1

A

200 pA
0.1 s

15
mW/mm²

B

max. current (nA)

Drd1a^Cre    Adora2a^Cre

C

w/o light
light on

25mV
0.5 s

50 pA

D

-80 mV

10 mV
0.1 s

E

Hyperpolarization (mV)

-20

-40

-60

Drd1a^Cre    Adora2a^Cre

F

Rebound potential (mV)

-5

-10

Drd1a^Cre    Adora2a^Cre

Figure 5 - figure supplement 2

Figure 6 - figure supplement 1

Figure 7 - figure supplement 1

**A** *Drd1a^Cre*

BLA

PF

Bregma -1.46mm

Bregma -2.30

**B** *Adora2a^Cre*

BLA

PF

Bregma -1.58

Bregma -2.30

Figure 7 - figure supplement 2

**A**

Figure 7 - figure supplement 3

Figure 8 - figure supplement 1

# Figure 7 – supplemental table 1

| Abbreviation | Allen Brain atlas | Franklin and Paxinos |
|---|---|---|
| AI | Agranular insular area | Agranular insular cortex (dorsal, ventral and posterior) |
| AUD | Auditory areas (dorsal and ventral) | Auditory cortex (dorsal, primary and ventral) |
| BLA | Basolateral amygdalar nucleus (anterior, posterior and ventral) | Basolateral amygdaloid nucleus (anterior, posterior) |
| CeA | Central amygdalar nucleus (capsular, medial and lateral) | Central amygdaloid nucleus (capsular, medial and lateral) |
| Ect | Ectorhinal area | Ectorhinal cortex |
| ENT | Entorhinal area | Entorhinal cortex |
| EP | Endopiriform nucleus | Claustrum (dorsal, ventral and endopiriform) |
| GPe | Globus pallidus, external segment | Globus pallidus |
| GPi | Globus pallidus, internal segment | Entopeduncular nucleus |
| InsCtx | Gustatory areas and Visceral area | Insular cortex (granular and dysgranular) |
| LA | Lateral amygdalar nucleus | Lateral amygdaloid nucleus |
| MO | Primary motor area | Primary motor cortex |
| PF | Parafascicular nucleus | Parafascicular thalamic nucleus |
| PIL | Posterior intralaminar thalamic nucleus | Posterior intralaminar thalamic nucleus |
| PIR | Piriform area | Piriform cortex |
| PO | Posterior complex of the thalamus | Posterior thalamic nuclear group |
| PoT | Posterior triangular thalamic nucleus | Posterior thalamic nuclear group, triangular part |
| S1 barrel | Primary somatosensory area, barrel field | Primary somatosensory cortex, barrel field |
| S1 mouth | Primary somatosensory area, mouth | Primary somatosensory cortex, upper lip/barrel field |
| S1 nose | Primary somatosensory area, nose | Primary somatosensory cortex, upper lip |
| S1 unassigned | Primary somatosensory area, unassigned | Primary somatosensory cortex, dysgranular zone |

| S1 upper limb | Primary somatosensory area, upper limb | Primary somatosensory cortex, shoulder region |
|---|---|---|
| S2 | Supplemental somatosensory area | Secondary somatosensory cortex |
| SNR | Substantia nigra, reticular part | Substantia nigra, reticular part |
| TeA | Temporal association areas | Temporal association cortex |
| VPM | Ventral posteromedial nucleus of the thalamus | Ventral posteromedial thalamic nucleus |