1 **Title page**
2
3 **Title:**
4 Immobilization of *C. elegans* on cultivation plates by thermoelectric cooling for high-
5 throughput subcellular-resolution microscopy
6
7 **Running title:**
8 Cooling immobilization for microscopy
9

10 **Author names and affiliations:**
11 Yao L. Wang[1,◊], Erik L. Jaklitsch[1,◊], Noa W. F. Grooms[1], Leilani G. Schulting, Samuel
12 H. Chung[1,*]
13 [◊]equal contribution
14 [1]Department of Bioengineering, Northeastern University, Boston, Massachusetts, USA
15
16 *Corresponding author
17 E-mail: schung@northeastern.edu
18

19 **Keywords:** animal immobilization, subcellular resolution microscopy, fluorescence
20 microscopy, Peltier, cooling, *Caenorhabditis elegans*
21

**Author contributions:**
Several people contributed to the work described in this paper. SHC conceived the idea.
YLW, ELJ, and NWFG carried out experiments, and all authors analyzed the data. SHC
supervised the experiments and the development of the manuscript. ELJ wrote the first
draft of the manuscript; all authors subsequently took part in the revision process and
approve the final copy of the manuscript.

## Abstract

Imaging, visual screens, and optical surgery are frequently applied to the nematode *Caenorhabditis elegans* at subcellular resolution for *in vivo* biological research. However, these approaches remain low-throughput and require significant manual effort. To improve throughput and enable automation in these techniques, we implement a novel cooling method to immobilize *C. elegans* directly on their cultivation plate. Previous studies cooled animals in microfluidics or flooded wells to 1-4 °C. Counterintuitively, we find that cooling to 5-7 °C immobilizes animals more effectively than lower temperatures. At 6 °C, animal movement consists of bouts of submicron nose tip movement occurring at a sufficiently low magnitude and frequency to permit clear imaging. We demonstrate the ability to perform subcellular-resolution fluorescence imaging, including 64× magnification 3D image stacks and 2-min long timelapse recordings of the ASJ neuron without blurring from animal motion. We also observe no long-term side effects from cooling immobilization on animal lifespan or fecundity. We believe our cooling method enables high-throughput and high-resolution microscopy with no chemical or mechanical interventions.

## Introduction

The nematode *C. elegans* is a widely-utilized model organism for studying many

biological processes. Similar to microorganisms, *C. elegans* are typically cultivated on

agarose plates, and they reproduce rapidly by self-fertilization, making them potentially

well-suited for studies requiring a large sample size. Their transparency and a wide

array of labelling techniques allows straightforward visualization of their internal

anatomy. Many studies in *C. elegans* employ high-resolution imaging of subcellular

structures, which necessitates animal immobilization to prevent image blur. Proper

immobilization is especially crucial for techniques involving multiple images in space or

time, such as 3D image stacks (*i.e.*, z-stacks) and timelapse imaging. Typically,

immobilization involves manual manipulation of individual animals and mounting them

on slides. These time- and labor-intensive procedures restrict experiments to small

sample sizes.

Though many strategies have been developed to immobilize *C. elegans*, there are

distinct drawbacks for each, mainly centered around their inefficiency or accessibility

(see Table S1). In brief, chemical methods utilize agents such as sodium azide or

levamisole to immobilize by disrupting physiology (Reeve, 1988). Mechanical methods,

such as polystyrene beads and hydrogel mixtures, physically restrain animals by

pressure, viscosity, or friction (Kim*, et al.*, 2013). Microfluidic and on-chip technologies

constrain animals by pressure, compression, or suction against a membrane/wall or

fluid (Guo*, et al.*, 2008; Zeng*, et al.*, 2008). Finally, extreme temperatures can also

immobilize animals and most techniques cool animals in flooded wells or microfluidics

(Rohde and Yanik, 2011). Despite the many advances in novel immobilization

strategies, several significant drawbacks remain: Some of the above strategies require

fabrication of slides and significant manual pre- and post-processing of individual

animals. They may increase physiological or physical stress on animals (Manjarrez and

Mailler, 2020). Some strategies require complex fabrication and setups or require

significant user training. Several methods may also allow small animal movements that

blur images, particularly in the nose and pharynx. Notably, none of the current

strategies allow for direct imaging on cultivation plates, necessitating time-consuming

animal transfer.

Here we address these shortcomings by developing an approach that readily and

reversibly immobilizes entire animal populations with minimal user effort and integrates

into standard *C. elegans* experimental workflows. We detail the successful operation of

a thermoelectric cooling stage and associated methods that can immobilize *C. elegans*

en masse directly on cultivation plates for high-resolution imaging. We describe our

cooling instrument, which is designed to hold the entire cultivation plate stably at a

desired temperature. We characterize animal movement at a variety of target

temperatures and cooling rates. Surprisingly, we found that cooling to relatively higher

temperatures (specifically 6 °C) minimizes animal movement, though previously-

reported cooling strategies utilized temperatures between 2-4 °C (Rohde and Yanik,

2011; Chung*, et al.*, 2008). Animal movement while cooled consists of bouts of

submicron nose tip movement occurring infrequently enough to permit high-resolution

brightfield imaging during timescales on the order of minutes. Cooling for short-term

immobilization does not impact *C. elegans* fecundity or lifespan. Finally, using our novel

immobilization method, we obtain subcellular-resolution z-stacks and timelapse

fluorescence images of *C. elegans* neurons without blurring from animal motion.

## Results

Cooling stage apparatus and temperature distribution

Our design goal is to uniformly cool an entire 55-mm diameter agar cultivation plate to a

target temperature, immobilizing animals for high-resolution fluorescence imaging while

also preserving transillumination and access to animals. Agar plates are commonly

used for cultivating, observing, and manipulating *C. elegans*, so our instrument has

broad applicability. As shown in Fig. 1ab, our instrument comprises a thin stage made of

an 80-mm diameter sapphire window, a thermally-conductive copper body, a

thermoelectric Peltier heat pump, and a closed-loop liquid cooler. The sapphire window

is transparent for transillumination and has a high thermal conductivity. We designed

notch cut-outs on the colder side of the sapphire window to improve temperature

uniformity across the window. The Peltier is a thermoelectric, solid-state device that

pumps heat between its two sides depending on applied voltage.

The stage rapidly cools agar plates resting on the sapphire window. Heat leaves the

plate to the sapphire window and then transfers to the copper body and Peltier. The

Peltier pumps heat to the closed-loop cooler for removal. Thermal simulation of our

design predicted a uniform thermal distribution (< 1°C temperature difference from

target temperature, $T_{set}$) across the central 40 mm-diameter area of the cultivation plate

(Fig. 1b thermal simulation top view). We constructed, assembled, and preliminarily

tested the device (Fig. 1c) by achieving and holding a variety of temperatures on the

agar surface. We measured the thermal distribution of the agar plate and cooling stage,

demonstrating an even distribution across the plate's central area and ensuring that the

animals are held at the same temperature for uniform immobilization (Fig. 1d). Thus, our

device features include optical transparency for transillumination, high thermal

conductivity for cooling efficiency, and uniform temperature distribution across the agar

surface.

General observations of cooling immobilization on agarose plates

Because our study is the first, to our knowledge, to immobilize animals on agar plates

by cooling, we began by carefully characterizing the impact of temperature on animal

movement. At temperatures ≥ 8 °C, most animals are not immobilized and crawl out of

the imaging field of view within minutes. Similar to prior studies of animals in liquid

solution, cooling to 1-7 °C slows *C. elegans* movement and stops crawling on plates

immediately. By eye, the part of the animal that moves the most under cooling

immobilization is the nose. Pharyngeal pumping is strongly suppressed, the tail also

shows some movement, and the body moves the least. Movie S1 shows typical

examples of immobilized animal movement at 6 °C, where we immobilized three N2

wild-type worms on a cultivation plate. The video rate is sped up 50× to show a 1200-s

timespan in a 24-s fast frame rate video. The body parts are immobilized to varying

levels, consistent with the independent control of these parts (Alkema*, et al.*, 2005) and

our experience with sodium azide anesthesia (Chung*, et al.*, 2013a). Because the nose

tip exhibits the most movement and contains our target neuron, we verify our cooling

immobilization by measuring movement of the nose tip throughout our work described

below.

Optimal cooling temperature for immobilization

We next determined the optimal target temperature, $T_{set}$, to immobilize animals.

Previous reports utilized target temperatures of 2-4 °C (Rohde and Yanik, 2011; Chung*,*

*et al.*, 2008) but presented minimal characterization of movement, quantitative or

qualitative. In our experience, some animal movement, however slow or small, persists

at all temperatures. We have even observed movement of animals in supercooled

buffer held well below 0 °C. By cooling plates to 1-7 °C using our instrument, we

investigated animal immobilization in response to a wide temperature range.

Surprisingly, we found that a relatively higher temperature, 6 °C, is the optimal

temperature for immobilization. Figure 2 shows average nose tip velocity for each

minute at specified times after reaching $T_{set}$. At 1-3 °C, nose movement velocity is slow

initially but does not greatly decrease over the hour after reaching $T_{set}$ (Fig. 2a). We

confirmed that this persistent velocity trend continued for at least 2 h after reaching $T_{set}$

(see below for details). At these velocities (0.5-1.0 μm/s), we could not obtain clear

high-resolution images, even with our advanced microscope and sensitive camera. 4 °C

represents a transition temperature, where velocity is high initially but decreases to

speeds comparable to those at 1-3 °C. At 5-7 °C, velocity is high initially but, decreases

to very low levels after ~50 min at $T_{set}$. This immobilization appears to take effect

progressively over time, in contrast to the immediate immobilization at lower

temperatures and earlier times. The velocities at 50-60 min are significantly ($p \ll 0.001$)

below those at 1-3 °C and near or below our resolution limit. This minimal velocity (< 0.5

µm/s) permits clear high-resolution microscopy. Because the final velocities are similar

between 5-7 °C, we selected 6 °C as the $T_{set}$ for further study.

Different cooling rates produce similar immobilization

We next examined the impact of the cooling rate on immobilization by testing three

methods to cool animals down to 6 °C at different rates.  We term these methods "slow",

"fast", and "abrupt". Figure 3 shows plate cooling profiles for slow and fast cooling

methods. Slow cooling uses a typical 4 °C refrigerator to cool down a lidded cultivation

plate to 6 °C in about 50-60 mins. After this first hour, we transfer this plate to the

cooling stage that is pre-cooled, to maintain animals at 6 °C. In fast cooling, we place a

plate directly onto the cooling stage to bring the plate from 20 °C to 6 °C. The cooling

stage can maintain colder temperatures than the refrigerator; thus, it can cool down

animals to 6 °C within 15 mins. Abrupt cooling requires more user effort: we pick

animals from their original 20 °C plate directly to a pre-cooled 6 °C plate. Based on

strong thermal conductivity and small animal size, we expect animals are cooled to 6 °C

in seconds.

As previously stated, immobilization at higher $T_{set}$ is strongest starting ~50 min after

beginning of cooling, and the first hour of slow cooling is in the refrigerator. Thus, we

measured velocity for an extended length of time, beginning when the animals reached

$T_{set}$ and up to 2 h from the start of cooling. We compared abrupt, fast, and slow cooling

to 6 °C and fast cooling to 1 °C. After cooling to 6 °C, we observe no significant

differences in velocity during the 80-120 min range between slow and fast cooling ($p$ =

0.08) or between slow and abrupt cooling ($p$ = 0.32), indicating that the cooling rate

does not require careful tuning to optimize immobilization. By contrast, in agreement

with our prior results (Fig. 2), fast cooling to 1 °C allows significantly more movement

during the 80-120 min range than fast cooling to 6 °C ($p$ < 0.001). Of the three cooling

methods, slow cooling involves the least amount of user time and effort (details in

Materials and methods). While certain experiments may require fast or abrupt cooling,

we favor the slow cooling method and utilize it for the remainder of this study.

Intermittent nose movements under cooling.

The goal of our device development is to immobilize animals for imaging, especially for

high-resolution microscopy. As such, we ascertained if animals were sufficiently

immobilized for imaging by our cooling stage. We measured nose movements each

second to better quantify immobilization behavior. Figure 5a shows second-by-second

measurements of typical nose movement under 6 °C slow cooling. Similar to sleep and

lethargus (Raizen*, et al.*, 2008; Iwanir*, et al.*, 2013), nose movement under cooling

conditions is intermittent, mostly comprised of seconds-long "bouts" of increased activity

separated by "windows" with little to no movement. We characterized the distance

moved during bouts and the length of the windows. Only 1.4% of movements surpass 1

µm, and 95% of measurements yielded movement below the resolution limit (0.5 µm)

(Fig. 5b). Movements below 1 µm do not produce visible blur even at the highest

magnification of our microscope (64×) because the motion is largely confined to the tip.

Also, most of the experiment duration is contained within longer windows: 96% of the

duration is part of windows that are longer than 0.5 min, and 90% of the time is part of

windows that are longer than 1 min (Fig. 5c). This level of movement is sufficiently low

to image by eye and is also amenable to high-resolution, blur-free brightfield imaging by

a camera, which can be performed at most timepoints. Moreover, technology

improvements have produced increasingly brighter fluorescent labels, enhanced optics,

and efficient cameras. If these tools allow rapid and more sensitive image acquisition,

the immobilization requirements may be relaxed. For instance, partial immobilization,

achievable at earlier times, may be sufficient for imaging with improved technologies,

allowing high-speed repetitive immobilizations for longitudinal imaging. In summary, our

cooling immobilization approach reduces animal movement sufficiently to clearly image

by eye and by camera.

Touch stimulus temporarily reactivates movement

We next investigated the impact of touch stimulus on movement. Gentle and harsh

touch to *C. elegans* alters the existing behavior of the animal and provokes an escape

response (Chalfie*, et al.*, 2014; Chalfie and Sulston, 1981; Alkema*, et al.*, 2005).

Likewise, touching animals in lethargus or sleep also reactivates them (Raizen*, et al.*,

2008; Iwanir*, et al.*, 2013). We immobilized animals using the slow cooling method to 6

°C and stimulated animals by moving them with a pick at 90 min (*i.e.*, harsh touch). As

expected, touch stimulus reactivates immobilized animals (Fig. 6). Compared to the 10

min before touch stimulus, velocity for the following 10 min is modestly elevated ($p =$

0.006) before returning to the immobilization baseline during the subsequent 10 min.

Thus if animals are disturbed while immobilized, they may require a period of time

before returning to full immobilization.

Animals persist in immobilized state after temperature rises

We envision that, after a typical experiment involving cooling immobilization, plates can

be moved back to their incubator to return animals to their cultivation temperature. We

observed by eye the resumption of movement during warming on an inactive cooling

stage. Because animals become strongly immobilized when cooled below 7 °C, we

expected that movement would return at a similar temperature when warming animals

to their cultivation temperature, 20°C. Intriguingly, following slow cooling immobilization

and maintenance at 6 °C for 1 h, no animals resumed crawling before the plate

temperature reached ~ 10 °C. Even after the plate temperature reached 20 °C, at least

50% of animals remained immobilized until stimulated by gentle touch, whether

externally applied or from contact with other animals moving on the plate. When

stimulated by touch, animals transition within a few seconds directly from immobilized to

crawling that appears normal by eye. Thus, while animals resume apparently normal

behavior following cooling immobilization, the transition from immobilized to crawling

may be more complex.

Cooling immobilization does not reduce lifespan or fecundity

Prior studies involving cooling of *C. elegans* indicate that cooling for shorter periods of

time does not appear to impact the animal health (Rohde and Yanik, 2011), but longer

periods (< 4 h) cause an immediate or progressive decline in health (Ohta*, et al.*, 2014;

Robinson and Powell, 2016). We assayed viability and fecundity of animals immobilized

by cooling for 1 h. First, we immobilized animals at 6 °C (slow cooling) and quantified

their lifespan and number of offspring (Fig. 7). Lifespan and fecundity for control and

cooled animals are not significantly different (black, blue data). Second, we immobilized

animals at 1 °C (fast cooling) to test a harsher immobilization condition. Again, lifespan

and fecundity for control and cooled animals are not significantly different (light gray,

purple data). We conclude that cooling immobilization does not significantly affect

animal health.

Timelapse imaging and 3D image stack

Illuminating animals with high intensity light stimulates movement in some animals. We

describe our observations in the Discussion section. By eye, with our setup and animal

strains, we note increased movement primarily when the illumination is focused to a

size less than the animal length. Some animals remain motionless under illumination

intensities sufficient for fluorescence imaging. By imaging animals immobilized on our

cooling stage, we demonstrate high-resolution timelapse and z-stack images of

individual neurons without movement-induced blurring. We obtained timelapse

recordings (2 min length, 1 frame/s, 100-ms exposure) of an ASJ neuron expressing

green fluorescent protein (GFP). By cross-correlation we tracked the movement of

neuron structures within four areas in the field of view that focused on portions of the

neuron (Fig. 8a). The distance the neuron structures moved from their position in the

first frame is shown in Fig. 8b and is usually less than 0.2 μm. Thus, the distance moved

during 2 min of imaging is below our resolution limit of 0.5 μm.

Incompletely immobilized animals produce images with blurred features due to

motion during or between images. Blurring must be minimized or eliminated to obtain

clear images, particularly for samples with fine features, such as the thin fibers of

neurons. We took a z-stack of the ASJ neuron comprising 41 images with 1 μm step

size taken at 1 frame every 4 s. As shown by the maximum projection of the z-stack in

Fig. 8c, while there is haze around neuron structures from inherent out-of-focus and

scattered light in widefield imaging (Wang*, et al.*, 2021), there is no image blur from

motion. Specifically, the neuronal fibers are clearly resolved, even at high resolution.

This clear image indicates minimal animal movement during 160 s of imaging, and it

underscores the effectiveness of our instrument and approach for immobilization.

**Discussion**

**Envisioned workflows**

We envision two potential workflows utilizing our device to enable convenient, reversible

immobilization of many animals for high-resolution imaging. In this study, we

demonstrated the first workflow, where entire cultivation plates are cooled to 6 °C.

Because this method requires ~50 min to maximally immobilize animals, plates are

"precooled" by placing them in a refrigerator for ~1 h, and subsequent cooling on the

stage maintains the immobilization temperature. To maximize efficiency in multi-plate

experiments, a plate is processed on the stage while simultaneously precooling the next

plate that will be processed. Following imaging of immobilized animals, plates are

returned directly to their incubator to bring them back to their cultivation temperature.

We suggest that plates be tapped or otherwise disturbed to trigger movement in animals

rather than allowing them to persist in the immobilized state. The second workflow,

involving a microchamber device for longitudinal imaging, is further described in the

"Future Developments and Applications" section below.

**Advantages of cooling stage and approach**

Our cooling stage and approach have six technical capabilities, including some

significant improvements over prior immobilization methods. First, large *C. elegans*

populations are readily and reversibly immobilized by cooling. This immobilization

allows observation, imaging, and manipulation of many animals over long time periods.

Second, for some applications or genetic backgrounds, immobilization by cooling may

be preferable to other methods due to secondary effects, such as stress response

(Manjarrez and Mailler, 2020). For instance, chemical agents such as sodium azide

induce stress proteins and thermotolerance in *C. elegans* (Massie*, et al.*, 2003). Third,

our stage permits high-resolution fluorescence imaging of animals directly on cultivation

plates. While *in vivo* microscopy on cultivation plates has been a mainstay technique for

decades, it is more often performed by eye and at lower resolution. Our approach

enables the study of subcellular processes and cell morphology without manipulating

individual animals. Fourth, optical transparency of the sapphire window allows easy

observation and imaging by brightfield transillumination, which is a primary method used

for many imaging applications in *C. elegans*. Fifth, we designed our cooling stage for

straightforward installation into common upright microscope setups. The compact stage

is very thin and is easily introduced into and removed from existing setups. Our cooling

approach is also amenable to typical *C. elegans* experimental workflows. Finally, most

immobilization techniques for high-resolution imaging do not easily allow animal

reorientation while imaging. Our cooling stage maintains access to animals, which can

be manipulated by a standard platinum pick throughout imaging, at the cost of time to

reestablish full immobilization (see "Touch stimulus" section above). These six technical

capabilities enable broad application of our device and approach to many problems that

require high-resolution *in vivo* microscopy on large numbers of animals.

Our cooling stage and approach also has three practical, user-friendly features.

First, our approach requires minimal user effort. Most other reversible immobilization

techniques require transfer and manipulation of individual animals to complex setups or

slides, which must also be fabricated. Our approach obviates this work by immobilizing

animals on the cultivation plate and requires only seconds of user effort. For imaging

experiments that involve animal recovery, we estimate that half the experimental time

and a third of the user effort is expended in non-imaging activities such as animal

handling, immobilization, and recovery (Chung*, et al.*, 2013b). Thus, our approach

greatly accelerates the pace of experiments, enabling the straightforward application of

high-resolution imaging to challenging procedures, such as screens. Second, our

cooling stage and approach are very simple and straightforward, and it requires minimal

training to operate. Some immobilization techniques involving complex setups require

significant training and fine-tuning. In contrast our cooling stage has only a single

voltage adjustment to control the immobilization temperature. Finally, our device is

accessible and inexpensive: we assembled it from easily attainable and affordable

components, totaling a cost of ~$500. In summary, our device and approach has

significant technical capabilities in a very user-friendly design.

**Key technical developments and novel findings**

In our study, we develop a novel, compact thermoelectric stage that cools entire

cultivation plates to immobilize populations of *C. elegans* for imaging. Our stage cools

plates to a uniform temperature distribution across the agar surface that is stable over

long time periods. We describe animal movement under many cooling conditions in far

greater detail than previously reported. We demonstrate that cooling has minimal impact

on lifespan and fecundity. Our cooling stage successfully immobilizes animals

sufficiently to allow high-resolution brightfield imaging. Despite some movement arising

from high-intensity illumination for high-resolution fluorescence imaging, we

demonstrate the capacity of our immobilization approach to perform time-lapse and z-

stack fluorescence imaging. These techniques can be challenging as animal movement

during the extended acquisition time blurs the images. In summary, our cooling stage

represents a novel and effective way to reversibly immobilize large populations of *C.*

*elegans* for high-resolution brightfield and fluorescence imaging.

During our study, we also encountered some interesting and unusual animal

behavior. First, previous studies immobilized animals between 2-4 °C (Rohde and

Yanik, 2011; Chung*, et al.*, 2008). By careful and systematic assay, we find that optimal

immobilization of *C. elegans* occurs at a counterintuitively higher temperature of 5-7 °C

after animals have been cooled for ~50 min. In contrast, animals cooled to 1-3 °C

maintain a relatively stable level of movement that is significantly higher than at 6 °C.

Animals held at 1 °C only slightly reduce their movement even after 2 h. The higher

immobilization temperature is actually preferable as it is easier to maintain, has lower

risk of freezing animals, and is presumably a more benign environment. Second, after

cooling immobilization, many animals persist in the immobilized state even after

warmed to cultivation temperatures but resume normal movement rapidly following

gentle touch. Because animals immobilized by cooling also exhibit bouts of movement

and respond to strong stimuli by temporary increased movement, we speculate that our

immobilized state may be similar to sleep. Further studies are required to determine if

cooling immobilization exhibits other hallmarks of sleep, or if it is more akin to partial

anesthesia.

**Limitations of cooling stage and approach**

There are several limitations of our approach. First, animals raised at 15 °C continue to

crawl at our standard immobilization temperature of 6 °C, indicating that cultivation

temperature modulates cooling immobilization. This is not unexpected as prior studies

have shown that raising *C. elegans* at colder temperatures improves cold tolerance

(Murray*, et al.*, 2007).  However, it does require additional fine-tuning of our approach if

animals are not cultivated at 20 °C. Likewise we also speculate that some temperature-

sensitive mutant strains of *C. elegans* may require adjustment of immobilization

temperature, as we have optimized our approach for the wild-type background. Second,

the speed of immobilization is slower than most other methods, but we expect that the

easy step of pre-cooling plates overcomes this weakness for most applications. Third,

imaging fine structures at specific locations exhibiting greater movement, such as the

dendritic cilia at the nose tip, could be challenging.

Finally, high-intensity illumination for fluorescence imaging at high-resolution

stimulates movement in some animals. Our current investigations are aimed at

quantifying and minimizing this effect. By eye, the nose appears to be more sensitive

than the tail or the body to light. The location of GFP expression may affect light

sensitivity as FLP-fluorescent animals could be illuminated with >10× more light than

ASJ-fluorescent animals before stimulating movement. Also, green light illumination (red

fluorescence imaging) triggers much less movement than blue light (green fluorescence

imaging). These gradations of sensitivity are consistent with prior findings that neurons

in the nose, and particularly the ASJ, contribute strongly to wavelength-dependent light

sensitivity in *C. elegans* (Ward*, et al.*, 2008). Despite this sensitivity, we are still able to

demonstrate high-resolution fluorescence imaging at the subcellular level without

motion blur.

**Future developments and applications**

Our cooling stage and associated equipment could be developed in several ways to

greatly enhance its capabilities. A microfluidic or microchamber (Bringmann, 2011)

device utilized with our cooling approach could permit longitudinal imaging of individual

animals. A thin device would enable rapid, repeated cooling for brief imagings with

minimal impact on animal health (Rohde and Yanik, 2011). Such a device would be

useful for tracking development or neuroregeneration following injury. Our study has

also focused on imaging L4 and young adult animals; however, imaging and laser

surgery on larval stages has been instrumental for understanding animal development

and the roles of individual neurons. Larval stages are significantly more challenging to

process for microscopy than later life stages, particularly the recovery step. If the

procedure for cooling immobilization were optimized for larval stages, that would greatly

facilitate larval imaging and laser surgery. In our study, we also completed procedures

manually, but one advantage of our approach is that it is amenable to automation.

Commercial plate handlers exist to move plates to instruments and could be easily

adapted to load plates onto our cooling stage. The cooling immobilization and imaging

could also be automated through multiple coolers and translation stages. There is also

great potential for integrating our approach with image processing and segmentation

algorithms to enable automated recognition of cellular components and targeting for

laser surgery. The developments listed above would significantly extend the capabilities

and accelerate the throughput of our platform allowing its application to a wide array of

questions that are currently unapproachable.

**Materials and methods**

**Device design and components:** We utilized Solidworks to design and model the

thermal distribution of the cooling stage. Components: Peltier heat pump (40 × 40 mm

CP854705-2, from CUI Devices); 99.9% pure copper stage block (100 × 140 × 3 mm,

machined); sapphire window (80 mm diameter, 3 mm thickness, from Altos Photonics);

Thermal paste to connect parts (Arctic MX-4); liquid CPU Cooler (Hydro Series H100x,

from Corsair); copper liquid cooling heatsink (40 × 40 × 10 mm, replacing the original

bulky heatsink); custom installation bracket fabricated by FDM (Fused Deposition

Modeling) 3D printing of polylactide filament.

**Plate temperature measurements:** We measured the thermal distribution of the agar

plate using a calibrated thermal camera (Teledyne FLIR C2). We embedded a wire

thermocouple (Proster digital thermocouple) 1 mm into the agarose on plates to confirm

camera readings and track agarose temperature over time.

**Nematode handling:** We maintained *C. elegans* on Bacto agar plates under standard

conditions at 20 °C (Brenner, 1974). We utilized two strains in our study: a wild-type N2

strain and a fluorescent strain with *daf-11(sa195)V; ofIs1[ptrx-1::trx-1::gfp]* background.

We utilized N2 for all experiments except for imaging, where we utilized the fluorescent

strain. We synchronized large populations of animals for experiments by placing ~30

adults on plates to lay eggs for 4 h. We transferred animals between plates using a pick.

For the touch stimulus experiment, we used a pick to move animals at least 1 cm and

then back to their original location.

**Cooling methods: slow, fast, and abrupt**

In general, the parameters of the cooling methods, such as voltages and times, that

lead to the desired performance, will depend on specific parameters of the cultivation

plates and stage, such as the amount of agar used to make the plates, the efficiency of

the Peltier stage, and the material conductivities.

**Slow cooling:** In our study, we performed slow cooling to a target temperature ($T_{set}$) of

6 °C only. We first moved a 20 °C plate with to a 4 °C refrigerator for 1 h, with plate's lid

on. Then, we moved the plate directly to a cold stage (pre-cooled for 10 min with 4.8 V

across the Peltier), with plate's lid off.

**Fast cooling:** We first moved a 20 °C plate with animals to a cold stage (cooled for 10

min at 12 V), with plate lid off. When the plate temperature reached $T_{set}$ + 5 °C, we

turned the voltage to a lower value (4.8 V for $T_{set}$ = 6 °C and 7.1 V for $T_{set}$ = 1 °C) to

maintain plates at $T_{set}$. This method cools down plates to different target temperatures in

~15 mins.

**Abrupt cooling:** In our study, we performed abrupt cooling to 6 °C only. We moved

animals from 20 °C plates to a pre-cooled plate on a cold stage.

**Tip-Tracking:** We first captured videos of animals under the stated conditions at 20

frames per second (FPS). Then, we extracted 1 frame per second from those videos

using a custom MATLAB script. We manually tracked the nose tip position in those

frames using the Multi-point Tool in ImageJ and calculated velocities for each second in

MATLAB. We averaged the individual velocities of 60 s to produce average velocities

for each minute.

**Lifespan and fecundity analysis:**

We collected N2 wild-type animals in the L4 lifestage on two plates. We left the control

plate in the cultivation incubator. The cooled plate was processed by the slow cooling to

6 °C and fast cooling to 1 °C methods described above. We held animals at $T_{set}$ for 1 h.

We then moved the cooled plate back to the cultivation incubator.

**Lifespan:** A few hours after treatment, we transferred animals to separate plates to

avoid crowding. We observed animals twice daily following established procedure

(Larsen*, et al.*, 1995) to assess mortality. We transferred animals to new plates when

needed to avoid starvation.

**Fecundity:** A few hours after treatment, we transferred animals individually (*i.e.*,

"singled") to plates. Three days later, we moved the singled animals to new plates and

counted the number of hatched offspring for each animal. We repeated this process

until there were no new offspring.

**Timelapse and z-stack imaging:** We used a Leica M205 FA fluorescence

stereomicroscope with 5x, 0.5 numerical aperture (NA) objective to obtain z-stack and

timelapse images. With other optics in the microscope body, we imaged green

fluorescent protein at 64x magnification, with theoretical 0.5 μm transverse and 4 μm

axial resolution. We performed both timelapse and z-stack fluorescence imaging on the

*C. elegans* ASJ neuron. For timelapse imaging, we used 1 FPS with 100-ms exposure

time for 2 minutes to obtain 120 frames in total. For z-stack imaging, we used a 3-axis

stage (Thorlabs MBT616D) to obtain slices over a 40-μm z range, with 1-μm z step,

100-ms exposure time, and 4-s time interval between slices.

**Statistics and interpretation of results:** For comparing nose tip velocities, we

calculated $p$-values using the $t$ test: For different temperature settings or different

cooling rates, we utilized the paired, two-tailed $t$ test. For examining touch stimulus, we

utilized the unequal variance, two-tailed $t$ test.

For assessing the impact of cooling on viability and fecundity, we calculated $p$-value by

using log-rank test and the unpaired, equal variance, two tailed $t$ test, respectively. Data

are represented as average ± standard deviation (SD). In figures, we indicate values

that differ significantly at $p < 0.05$ (*) and $p < 0.001$ (**) levels.

**Acknowledgements**

We thank the members of the Chung Laboratory for feedback on the manuscript. We
acknowledge Wenchao Zhu (Mechanical Engineering, Northeastern Univ.) for technical
support in thermal distribution measurement. We acknowledge Xin Sun
(Bioengineering, Northeastern Univ.) for technical support in 3D printing.

**Figure captions**

**Figure 1. Cooling stage apparatus.** (a) Top-down view of cooling immobilization

stage. Agar cultivation plates placed on the sapphire window. (b) Thermal modeling of

stage and agar plate. Peltier heat pump and liquid cooling system secured to the copper

base. (c) Photograph of cooling stage and agar plate. (d) Thermal image of unlidded

agar plate and stage cooled to $T_{set} = 6$ °C obtained by infrared camera. Temperatures at

indicated points.

**Figure 2. Relatively higher temperatures immobilize animals more effectively.**

Average nose tip velocity each minute for (a) low temperature (1-4 °C) and (b) high

temperature (5-7 °C) under fast cooling immobilization. $n = 6$ animals. $p < 0.001$ (**)

comparing final 10 min of movement for 1-4 °C and 5-7 °C.

**Figure 3. Plate cooling profiles.** Plate temperatures over time for different cooling

methods and $T_{set}$.

**Figure 4. Different cooling rates to 6 °C eventually immobilize similarly.** Average

velocity of nose tip movement for abrupt, fast, and slow cooling methods to 6 °C and

fast cooling to 1 °C. Note that fast and slow cooling velocity measurement begins after

cooling initiation to allow animals to reach $T_{set}$. $n = 6$ animals for each condition. $p <$

0.001 (**) and not significant (ns) comparisons of movement during 80-120 min.

**Figure 5. Movement under cooling comprises intermittent bouts with long interstitial windows.** (a) Movement measured each second reveals intermittent bouts of increased activity. Movements of < 1 μm do not produce significant image blur. (b) Cumulative distribution function of data in part A. Most movements are small: 99% of movements are < 1 μm. (c) Complementary cumulative distribution function of data in part A. Longer windows comprise most of experiment duration: 96% of duration is part of windows > 0.5 min long.

**Figure 6. Touch stimulus briefly reactivates movement.** Nose tip velocity before and after harsh touch at 90 min are significantly different. Animals return to immobilization in 10 min. $n = 6$ animals. $p < 0.05$ (*).

**Figure 7. Cooling immobilization does not reduce *C. elegans* lifespan and fecundity.** Animal lifespan (a) and fecundity (b) after slow cooling to 6 °C or fast cooling to 1 °C, followed by 1 h at $T_{set}$. $n = 110$ (lifespan), 55 (fecundity) per condition. $p < 0.05$ (*) and $p < 0.001$ (**).

**Figure 8. High-resolution imaging under 6 °C cooling immobilization.** Fluorescent *C. elegans* ASJ neuron imaged at 640× magnification. (a) Maximum projection of 2-min timelapse image sequence shows no motion blur. Relative movement of structures in areas 1-4 tracked by cross-correlation. (b) Minimal movement of structures in areas 1-4 during timelapse. (c) Maximum projection of 3D image stack acquired manually. Structures show no motion blur.

555 **Movie S1. 20-min timelapse recording of immobilized animals.** 50× speed, 10×

556 magnification. Movement at times (s): Left animal nose tip- 99, 182, 392, 822, 1045; tail-

557 359, 723. Center animal tail- 133, 297. Right animal nose tip- 0, 292, 469, 488.

**References**

Alkema, M.J., Hunter-Ensor, M., Ringstad, N., and Horvitz, H.R. Tyramine Functions
Independently of Octopamine in the Caenorhabditis elegans Nervous System. Neuron
46, 247-260 (2005). DOI: 10.1016/j.neuron.2005.02.024.

Brenner, S. The genetics of Caenorhabditis elegans. Genetics 77, 71-94 (1974).

Bringmann, H. Agarose hydrogel microcompartments for imaging sleep- and wake-like
behavior and nervous system development in Caenorhabditis elegans larvae. J
Neurosci Methods 201, 78-88 (2011). DOI:
https://doi.org/10.1016/j.jneumeth.2011.07.013.

Chalfie, M., Hart, A.C., Rankin, C.H., and Goodman, M.B. Assaying mechanosensation.
WormBook : the online review of C elegans biology,
10.1895/wormbook.1891.1172.1891 (2014). DOI: 10.1895/wormbook.1.172.1.

Chalfie, M., and Sulston, J. Developmental genetics of the mechanosensory neurons of
Caenorhabditis elegans. Dev Biol 82, 358-370 (1981).

Chung, K.H., Crane, M.M., and Lu, H. Automated on-chip rapid microscopy,
phenotyping and sorting of C. elegans. Nat Methods 5, 637-643 (2008). DOI:
10.1038/nmeth.1227.

Chung, S.H., Schmalz, A., Ruiz, R.C.H., Gabel, C.V., and Mazur, E. Femtosecond
Laser Ablation Reveals Antagonistic Sensory and Neuroendocrine Signaling that
Underlie C-elegans Behavior and Development. Cell Reports 4, 316-326 (2013a). DOI:
10.1016/j.celrep.2013.06.027. PMCID: PMC4780052.

Chung, S.H., Sun, L., and Gabel, C.V. In vivo Neuronal Calcium Imaging in C. elegans.
Journal of Visualized Experiments, e50357 (2013b). DOI: 10.3791/50357. PMCID:
PMC3653678.

Guo, S.X., Bourgeois, F., Chokshi, T., Durr, N.J., Hilliard, M.A., Chronis, N., and Ben-
Yakar, A. Femtosecond laser nanoaxotomy lab-on-a-chip for in vivo nerve regeneration
studies. Nat Methods 5, 531-533 (2008).

Iwanir, S., Tramm, N., Nagy, S., Wright, C., Ish, D., and Biron, D. The Microarchitecture
of C. elegans Behavior during Lethargus: Homeostatic Bout Dynamics, a Typical Body
Posture, and Regulation by a Central Neuron. Sleep 36, 385-395 (2013). DOI:
10.5665/sleep.2456.

Kim, E., Sun, L., Gabel, C.V., and Fang-Yen, C. Long-Term Imaging of Caenorhabditis
elegans Using Nanoparticle-Mediated Immobilization. PLOS ONE 8, e53419 (2013).
DOI: 10.1371/journal.pone.0053419.

Larsen, P.L., Albert, P.S., and Riddle, D.L. Genes that regulate both development and
longevity in Caenorhabditis elegans. Genetics 139, 1567-1583 (1995). DOI:
10.1093/genetics/139.4.1567.

Manjarrez, J.R., and Mailler, R. Stress and timing associated with Caenorhabditis
elegans immobilization methods. Heliyon 6, e04263 (2020). DOI:
https://doi.org/10.1016/j.heliyon.2020.e04263.

Massie, M.R., Lapoczka, E.M., Boggs, K.D., Stine, K.E., and White, G.E. Exposure to
the metabolic inhibitor sodium azide induces stress protein expression and
thermotolerance in the nematode Caenorhabditis elegans. Cell Stress Chaperones 8, 1-
7 (2003).

Murray, P., Hayward, S.A.L., Govan, G.G., Gracey, A.Y., and Cossins, A.R. An explicit
test of the phospholipid saturation hypothesis of acquired cold tolerance in
Caenorhabditis elegans. Proceedings of the National Academy of Sciences 104, 5489-
5494 (2007). DOI: 10.1073/pnas.0609590104.

Ohta, A., Ujisawa, T., Sonoda, S., and Kuhara, A. Light and pheromone-sensing
neurons regulates cold habituation through insulin signalling in Caenorhabditis elegans.
Nat Commun 5 (2014). DOI: 10.1038/ncomms5412.

Raizen, D.M., Zimmerman, J.E., Maycock, M.H., Ta, U.D., You, Y.J., Sundaram, M.V.,
and Pack, A.I. Lethargus is a Caenorhabditis elegans sleep-like state. Nature 451, 569-
572 (2008). DOI: 10.1038/nature06535.

Reeve, E.C.R. The Nematode Caenorhabditis elegans . Edited by William B. Wood and
the Community of C. elegans Researchers. New York: Cold Spring Harbor Laboratory,
1988. 667 pages. US $97.00. ISBN 0 87969 307 X. Genetics Research 52, 243-244
(1988).

Robinson, J.D., and Powell, J.R. Long-term recovery from acute cold shock in
Caenorhabditis elegans. BMC Cell Biol 17, 11 (2016). DOI: 10.1186/s12860-015-0079-
z.

Rohde, C.B., and Yanik, M.F. Subcellular in vivo time-lapse imaging and optical
manipulation of Caenorhabditis elegans in standard multiwell plates. Nat Commun 2, 7
(2011). DOI: 10.1038/ncomms1266.

Wang, Y.L., Grooms, N.W.F., Civale, S.C., and Chung, S.H. Confocal imaging capacity
on a widefield microscope using a spatial light modulator. PLOS ONE 16, e0244034
(2021). DOI: 10.1371/journal.pone.0244034. PMCID: PMC7886194.

Ward, A., Liu, J., Feng, Z.Y., and Xu, X.Z.S. Light-sensitive neurons and channels
mediate phototaxis in C-elegans. Nat Neurosci 11, 916-922 (2008).

Zeng, F., Rohde, C.B., and Yanik, M.F. Sub-cellular precision on-chip small-animal
immobilization, multi-photon imaging and femtosecond-laser manipulation. Lab Chip 8,
653-656 (2008).

**A**

copper block

sapphire

agar plate Ø55

3 mm thick copper, sapphire

Ø80

lengths in mm

**B**

top view

T(°C)

-5

-10

-15

side view

closed-loop chiller

Peltier heat pump

animals on agar

PS dish

**C**

**D**

sapphire

agar plate

copper

5.8

6.1      6.0      6.5

6.2

T(°C)

-5

-10

-15