Rec8 cohesin-mediated loop-axis chromatin architecture is required for meiotic recombination

Takeshi Sakuno[1][*][#], Sanki Tashiro[2][#], Hideki Tanizawa[2], Osamu Iwasaki[2], Da-Qiao Ding[3], Tokuko Haraguchi[1], Ken-ichi Noma[2, 4][*] and Yasushi Hiraoka[1][*]

[1] *Graduate School of Frontier Biosciences, Osaka University, Suita 565-0871, Japan;*

[2] *Institute of Molecular Biology, University of Oregon, Eugene, OR 97403, USA;*

[3] *Advanced ICT Research Institute Kobe, National Institute of Information and Communications Technology, Kobe 651-2492, Japan;*

[4] *Institute for Genetic Medicine, Hokkaido University, Sapporo 060-0815, Japan*

\# These authors contribute equally to this work.

*Corresponding authors: Takeshi Sakuno (sakuno@fbs.osaka-u.ac.jp) and Yasushi Hiraoka (hiraoka@fbs.osaka-u.ac.jp), Graduate School of Frontier Biosciences, Osaka University, Suita 565-0871, Japan; Ken-ichi Noma (noma@uoregon.edu), Institute of Molecular Biology, University of Oregon, Eugene, OR 97403, USA.

Summary

During meiotic prophase, cohesin-dependent axial structures are formed in the synaptonemal complex (SC). However, the functional correlation between these structures and cohesion remains elusive. Here, we examined the formation of the cohesin-dependent axial structure in fission yeast, which forms atypical SCs composed of linear elements (LinEs) resembling the lateral elements of SC but lacking the central elements. The results demonstrated that Rec8 cohesin is crucial for the formation of the loop-axis structure within the atypical SC. Furthermore, the Rec8-mediated loop-axis structure is formed in the absence of LinEs and provides a structural platform for aligning homologous chromosomes. We also identified a *rec8* mutant that lost the ability to assemble the loop-axis structure without losing cohesion. Remarkably, this mutant showed defects in the LinE assembly, resulting in a significant reduction in meiotic recombination. Collectively, our results demonstrate an essential role for the Rec8-dependent loop-axis structure in LinE assembly, facilitating meiotic recombination.

Keywords

meiosis, cohesin, Rec8, chromosome, axis-loop structure, recombination, fission yeast, linear element, miscarriage, developmental disorders

Introduction

Meiosis is a unique type of cell cycle for sexual reproduction, which halves the chromosome number by two successive nuclear divisions after a single round of DNA replication to produce haploid gametes. In the first meiotic division (meiosis I), homologous chromosomes (homologs) are segregated to opposite poles. This type of division is called reductional segregation and is essentially different from mitosis or meiosis II, in which sister chromatids are segregated to opposite poles (Petronczki et al., 2003). Crossovers formed by reciprocal recombination constitute a physical link between homologs, ensuring reductional segregation (Hunter, 2015; Zickler and Kleckner, 1999). Failure to establish such interhomolog links leads to the mis-segregation of chromosomes during meiosis I, resulting in aneuploidy of gametes. This is a major cause of human miscarriage or developmental disorders caused by chromosome trisomy, such as Down syndrome (Nagaoka et al., 2012; Webster and Schuh, 2017).

During meiotic prophase, homologs become closely associated or aligned through an elaborate proteinaceous structure called the synaptonemal complex (SC), followed by crossover recombination. The SC consists of two lateral elements, each along the axis joining a pair of sister chromatids, with one central element connecting the lateral elements (Grey and de Massy, 2021; Kleckner, 2006). The process of chromatin loop formation is thought to be essentially mediated by meiotic cohesin (Klein et al., 1999; Petronczki *et al.*, 2003). The cohesin complex is responsible for sister chromatid cohesion. The complex consists of two structural maintenance of chromosome (SMC) proteins, a kleisin subunit, and a stromal antigen (SA) subunit. Replacement of kleisin commonly occurs in most organisms, in which the somatic kleisin Rad21/Scc1 is replaced by Rec8

during meiosis (Ishiguro, 2019; Mehta et al., 2013; Nasmyth, 2001). Rec8 cohesin is responsible for sister chromatid cohesion and provides a physical link between homologs as a result of crossover (Moore and Orr-Weaver, 1998; Petronczki *et al.*, 2003). Additionally, Rec8 cohesin is required for the assembly of lateral elements in diverse eukaryotes (Cai et al., 2003; Fujiwara et al., 2020; Fukuda et al., 2014; Golubovskaya et al., 2006; Hopkins et al., 2014; Ishiguro et al., 2014; Klein *et al.*, 1999; Llano et al., 2014; Pasierbek et al., 2001; Winters et al., 2014; Xu et al., 2005). A recent study using high-throughput chromosome conformation capture (Hi-C) (Lieberman-Aiden et al., 2009) in the budding yeast *Saccharomyces cerevisiae* clearly revealed that Rec8 cohesin is essential to construct meiosis-specific chromatin loop structures (Schalbetter et al., 2019). As studied in somatic cells, cohesins can form higher-order structures by pulling together distant regions of the chromosome, for which a loop extrusion model has been proposed (Davidson and Peters, 2021; Fudenberg et al., 2016; Kim et al., 2019; Mizuguchi et al., 2014; Rao et al., 2017; Sanborn et al., 2015). Pds5 and Wpl1/Wapl are cohesin-associating factors conserved in yeasts and vertebrates. These are considered to dissociate cohesin from chromatin (Challa et al., 2019; Peters and Nishiyama, 2012), thus affecting higher-order chromosome structures (Challa et al., 2016; Crawley et al., 2016; Ding et al., 2016a; Ding et al., 2006; Haarhuis et al., 2017; Hong et al., 2019; Silva et al., 2020; Tedeschi et al., 2013; Viera et al., 2020).

The fission yeast *Schizosaccharomyces pombe* forms an atypical SC composed of a structure called the linear element (LinE). The LinE resembles the lateral element of the SC, but lacks central element components, such as *S. cerevisiae* Zip1 (Bähler et al., 1993; Davis et al., 2008; Estreicher et al., 2012; Loidl, 2006; Lorenz et al., 2004; Olson et al.,

1978). Full alignment of homologs in *S. pombe* is finally achieved by chiasmata formation

following crossover recombination (Loidl, 2006). The recognition and alignment of

homologs in *S. pombe* relies on oscillatory movement of the nucleus (horsetail

movement) in a bouquet orientation with all telomeres clustered to the spindle pole body

(SPB) (Chikashige et al., 1994; Ding et al., 1998; Ding et al., 2019; Hiraoka, 2020; Loidl,

2006; Yamamoto and Hiraoka, 2001). More recently, a mechanism was proposed in which

noncoding RNA on chromatin contributes to the initial recognition of correct homolog

pairs (Chikashige *et al.*, 1994; Ding *et al.*, 1998; Ding *et al.*, 2019; Hiraoka, 2020; Loidl,

2006; Yamamoto and Hiraoka, 2001).

Importantly, it has been shown that the phosphorylation of Rec11, a homolog of the

meiosis-specific SA3 subunit of Rec8 cohesin in *S. pombe*, by casein kinase 1δ/ε (CK1),

is required for the assembly of LinEs (Phadnis et al., 2015; Sakuno and Watanabe, 2015).

The Rec8 cohesin complex containing phosphorylated Rec11 specifically interacts with

Rec10, a homolog of the lateral element factor Red1/SYCP2. Previous studies in mouse

spermatocytes have shown that depletion of the SYCP2 or SYCP3 structural components

of axial elements causes defects in axial element formation despite normal cohesin-axis

(Fujiwara *et al.*, 2020; Fukuda et al., 2010; Pelttari et al., 2001). Accordingly, the

structural property of chromatin to trail the horsetail movement is maintained in LinE-

defective mutants of *S. pombe* (Ding *et al.*, 2006; Sakuno and Watanabe, 2015). Based on

these observations, it should be emphasized here that the Rec8-dependent chromosomal

structure is strictly independent of lateral elements (or LinEs in *S. pombe*). However, the

significance of the Rec8-dependent axis-loop structure in meiotic recombination remains

unclear.

In this study, we performed Hi-C analysis to describe the meiotic chromosome structures of *S. pombe* in detail. We then sought to clarify the functional significance of the Rec8-dependent loop-axis structure by identifying *rec8* mutants that specifically show defects in axis formation, but not in cohesion. The collective data demonstrate that the Rec8-dependent loop-axis structure itself provides the scaffold to assemble the LinE axis, facilitating the execution of reciprocal recombination.

Results

Induction of highly synchronized meiosis under physiological conditions

To achieve Hi-C analysis of chromosomal configurations during meiotic prophase, we first developed an experimental system to prepare *S. pombe* cell populations undergoing highly synchronized meiotic progression under physiological conditions. For this purpose, we employed an *h-/h- pat1-as2/pat1-as2* diploid strain harboring one copy of the *mat*-Pc gene cassette (Cipak et al., 2012; Funaya et al., 2012; Yamamoto and Hiraoka, 2003). Meiosis can be induced using this ATP analog-sensitive allele of the Pat1 kinase (*pat1-as2*) upon inactivation of the Pat1 kinase by addition of the ATP analog, instead of the temperature-sensitive allele (*pat1-114*), in which raising temperature has detrimental effects on meiosis (Bähler et al., 1991; Cipak *et al.*, 2012). In addition, MDR-sup alleles (Aoi et al., 2014; Kawashima et al., 2012), which allow efficient cellular uptake of ATP analogs, were also exploited. As shown in Figure S1A, the meiotic progression of wild type diploid cells harboring Rec8-green fluorescent protein (GFP) was highly synchronous (~70%), as indicated by the timing of nuclear division in meiosis I. Pre-meiotic DNA replication was almost completed approximately 1.5 hours after the

induction of meiosis by the addition of an ATP analog after G1-arrest via nitrogen

starvation (Figure S1A, left panel).

Hi-C analysis recapitulates meiotic bouquet chromosomes in *S. pombe*

For Hi-C analysis, cells were collected at 0 h (G1) and 2.5 h (meiotic prophase) after

meiosis induction, along with cells in vegetative growth (VG; mainly in mitotic G2 phase).

Notably, even at 0 h, a significant amount of Rec8-GFP protein was expressed and loaded

onto chromatin. The amount of Rec8 peaked at approximately 2.5–3 h after meiotic

induction (Figures S1B and S1C). Hi-C contacts of all three chromosomes in VG cells

detected obvious centromere clustering (indicated by green arrowheads in Figures 1B and

1C), suggesting the organization of the Rabl orientation (Figure 1A, left). In contrast, as

cells progressed from G1 to meiotic prophase, centromere clustering became vague, while

telomere clustering became evident (Figures 1B and 1C; blue arrowhead). Remarkably,

as cells entered the meiotic prophase, X-shaped inter- and intra-chromosomal contacts

appeared (Figures 1B and 1C). These X-shaped contact patterns represent chromosomal

arms bundled at the telomere (Figure 1A; right), as previously demonstrated in *S. pombe*

meiotic prophase (Chikashige et al., 1997; Ding *et al.*, 1998). The degree of alignment

along the intra-chromosomal arms from the telomere was quantified by the "alignment

index" (Figure S1D). The alignment index of chromosome 1 increased from 1.6 (onset of

meiosis at 0 h) to 2.5 (meiotic prophase at 2.5 h) compared with 0.55 in VG (Figure 1D).

A similar structure was observed during meiotic prophase in mouse spermatocytes by Hi-

C analysis, although it was not as evident as in *S. pombe* (Patel et al., 2019).

Rec8 cohesin produces a chromosome shape suitable for alignment

Next, we examined three factors that play a central role in meiotic recombination. Rec10

is a major component of LinE. Rec12 is a Spo11 homolog that induces DNA double-

strand breaks (Keeney, 2001; Sharif et al., 2002). Rec8 is a meiotic cohesin. To

investigate their contributions to meiotic chromosome structures, we confirmed

synchronous progression of meiosis in these mutants and performed Hi-C analyses

(Figure S2). *rec10Δ* and *rec12Δ* cells showed slight defects in formation of the X-shaped

contacts (Figures 2A and 2B), with an alignment index of 2.8 (*rec10Δ*), 2.5 (*rec12Δ*), and

2.5 (*wild type*, *wt*). In contrast, *rec8Δ* cells showed prominent defects with almost no X-

shaped contacts (Figures 2A and 2B; alignment index 1.5). These findings suggest that

Rec8 regulates the alignment of homologs in a manner distinct from LinE formation to

modulate recombination. The bulk of the chromosomes do not follow the horsetail

movement at meiotic prophase in *rec8Δ*, while only telomere regions follow the leading

edge of the moving nucleus (Ding et al., 2016b; Ding *et al.*, 2006). These observations

imply that the traction force generated by the horsetail movement is not transmitted to the

chromosomes in *rec8Δ*, leading to the loss of alignment along the chromosomes.

Therefore, Rec8 cohesin likely provides the chromosome with appropriate structural

properties to counteract the pulling force, independent of LinEs (Figure 3F).

Rec8 cohesin is crucial to form meiosis-specific chromatin loops

Magnification of Hi-C data (2 kbp binning) in chromosome arm regions revealed that

punctate Hi-C interactions occurred at the onset of meiosis (0 h) and became more

prominent as meiosis progressed (2.5 h) in the wild type (arrows in Figures 3A and S3A).

The positions at the bases of these interactions markedly coincided with the previously

identified Rec8 accumulation sites (Ito et al., 2014) (Figures 3B and S3B). Consistently,

as meiosis progressed in the wild type, *P(s)* (the contact frequency versus distance)

indicated a "shoulder" at approximately 50 kbp, representing an increase in interactions

within a region shorter than that length (Figure 3C). Crucially, punctate Hi-C interactions

observed in the wild type were mostly lost in *rec8Δ*, but were retained in *rec10Δ* or *rec12Δ*

(Figures 3A and S3A). In addition, the shoulder in *P(s)* was lost in the *rec8Δ* mutant, but

not in *rec10Δ* or *rec12Δ* (Figure 3D). These data suggest the linear compaction of

chromosomes and the emergence of a Rec8-dependent *cis*-loop structure upon entry into

meiosis in the wild type. In the Hi-C data of 5 kbp binning, topologically associating

domain (TAD)-like signals were observed in wild type VG cells (dotted lines in Figure

3E), representing relatively long-range interactions manifest in mitotic chromosomes

(Kim et al., 2016; Mizuguchi *et al.*, 2014; Patel *et al.*, 2019; Tanizawa et al., 2017).

Interestingly, as Rec8-dependent loop structures appeared during meiosis, these TAD-like

signals disappeared. Thus, *S. pombe* chromosomes undergo a structural transition through

Rec8-dependent formation of chromatin loop-axis structures upon entry into meiosis

(Figure 3F).

Wpl1 in *S. pombe* regulates the length of chromatin loops in meiotic prophase

In Rad21/Scc1 cohesin-dependent chromatin organization with TADs in somatic cells,

Wapl depletion lengthens the chromatin loops and thickens the chromatin axis (Haarhuis

et al., 2017; Tedeschi *et al.*, 2013). However, it is unknown whether Rec8-dependent

chromatin loops are affected by Wapl during meiosis. As shown in Figure 4A, the meiotic

chromosome axial structures observed by Rec8-GFP were more prominent in *wpl1Δ* than

in the wild type, suggesting that the axial structure becomes thicker in *S. pombe* meiosis.

To examine the chromosome structure in *wpl1Δ*, we performed Hi-C analysis as in the

wild type (Figure S4A). In the Hi-C contact maps in *wpl1Δ* (Figure 4B, upper), the

background was lower than that in the wild type (compare with Figure 1B and the lower

panel of Figure 4B). These findings suggest that the inter- and intra-chromosomal long-

range interactions, including X-shaped contacts, were reduced in *wpl1Δ*, which has a

thicker chromatin axis, compared to the wild type (see Figure S4E).

In the amplified view of Hi-C data (2 kbp binning) in *wpl1Δ*, the number of grid-like

punctate signals increased and the size of the diagonals of each grid increased upon entry

into meiosis (Figures 4C and S4B). These findings imply that the nearby chromatin loops

are in contact with each other at their bases with longer loops and thicker axes, leading to

the linear compaction of chromosomes (Figure S4C). Consistently, compared to that of

the wild type, a rightward shift of the shoulder in the *P(s)* (blue dotted line for *wt* and

green dotted line for *wpl1Δ* in Figure 4D) and an increase in longer loop formation (Figure

4E) were observed in *wpl1Δ*. Note that, because axial signals of Rec8-GFP were still

thicker in the absence of LinEs (*wpl1Δ rec10Δ* double mutant, Figure S4D), Rec8 loop-

axis structures formed independently of LinEs, even under the condition of *wpl1Δ*. These

results strongly indicate that Wpl1/Wapl can antagonize the construction of chromosomal

axes restricting chromatin loop extension, even when mediated by meiotic Rec8 cohesin

in *S. pombe* (Figure S4C).

Identification of *rec8* mutants showing defects in axis formation

The foregoing results demonstrate that Rec8 cohesin is crucial for the formation of the

meiosis-specific axis-loop structure in *S. pombe* meiosis. Importantly, ectopic expression

of Rad21 cohesin in meiosis partly compensates for the cohesion defect in the absence of

Rec8 (Kitajima et al., 2003; Yokobayashi et al., 2003). However, Rad21 signals were

obscure along the chromatin axis (Figure S5A). Thus, besides cohesion, Rec8 may have

an additional function that is more robust than Rad21 in chromatin axis formation.

Therefore, it is possible to identify the mutant of *rec8* that specifically shows defects in

axis formation without affecting its cohesion function. To explore this, we examined the

S. pombe strain in which Rec8 and Rec11 were ectopically expressed as the sole source

of cohesin during mitosis, replacing Rad21 and Psc3 (Figure 5A). Using this strain as a

screening host, randomly mutagenized *rec8* cDNAs fused with GFP-coding DNA were

transformed into the host strain. Since Rec8 is the only available kleisin protein in this

strain, transformants will show a lethal or slow-growth phenotype if there is a *rec8*

mutation that causes loss of cohesion. Moreover, using the h^{90} mating-type and *mei4Δ*

background, each transformant arrests at the meiotic prophase on agar plates of

sporulation medium (Horie et al., 1998). We also introduced *wpl1Δ* in the host strain to

produce the prominent axis of Rec8-GFP (Figure 4A) so that defects in axis formation

can be easily evaluated by microscopy observation (Figure 5A). Deletion of $mei4^+$

represses the expression of recombination-related factors such as Rec12, resulting in

almost no recombination (Miyoshi et al., 2012; Murakami-Tonami et al., 2007; Young et

al., 2004). However, in the present study this did not seem to affect the Rec8 axis

formation, at least in the *wpl1Δ* background (Figure S5B).

We searched approximately 3000 viable transformants of the *rec8* mutant library for

mutants that showed axis formation defects with normal cohesion function in

fluorescence microscopy. We identified a mutant that displayed normal growth (Figure

S5C) but lost the Rec8 axis structure (Figure 5B). In this mutant, phenylalanine at amino

acid residue 204 of Rec8 was substituted with serine (*rec8-F204S*). Strikingly, sister

chromatid cohesion was preserved in the *rec8-F204S* mutant, as assessed at the *cut3* gene

locus (Figure 5C) and other chromosomal loci (Figure S5D). In the cohesion assays, we

used *rec8-S552P* or *rec8Δ*, which showed the cohesion defect, as a control strain (Figures

5C and S5D). The *rec8-S552P* mutant showed normal growth during mitosis (Figure

S5C), but displayed defects in axis formation and cohesion during meiosis, probably due

to low protein expression or low stability of the Psm1 subunit in the cohesin complex

(Figure S5E and S5F). Although cohesion defects are expected to occur during mitosis of

the P$_{adh1}$-derived *rec8-S552P* strain (Figure S5C), growth would be ensured by their

overexpression. Since the backup of Rad21 is likely to function only in *rec8Δ* cells, the

rec8-S552P mutant was more deficient in cohesion than *rec8Δ* (Figures 5C and S5D).

The cohesin complex seemed to be formed normally in the *rec8-F204S* mutant with

comparable expression of Rec11 (Figures S5G and S5H). The F204 of Rec8 seemed to

be conserved among Rec8 species in other organisms (Figure 5D), implying the

significance of this residue in regulating Rec8 functions. Furthermore, the analysis of

mutants in which F204 was replaced by amino acids other than serine indicated that the

hydrophobicity of this position of Rec8 may be important for its axis-forming function,

independent of its cohesion function (Figure S5I).

Thereafter, to delineate the details of axis formation defects caused by *rec8-F204S*, we

observed chromosome morphology during horsetail movement in meiotic prophase. In

the *rec8-F204S* mutant, only the nuclear leading edge followed the horsetail movement

with the bulk of chromosomes remaining, similar to *rec8Δ*, in which axial assembly and

chromosome compaction were lost (Figure 5E, Supplemental Movies 1, and 2). To

determine the degree of chromosome compaction more directly, we measured the distance

between the telomere and the *ade8* gene locus during meiotic prophase using a GFP-

tagged telomere protein (Taz1) and a *lacO*/lacI-GFP tag at the *ade8* gene locus (Figure

5F). As in *rec8Δ*, the telomere-*ade8* distance was longer in the *rec8-F204S* mutant than

in the wild type, suggesting that the chromatin of the *rec8-F204S* mutant is flexible and

is abnormally stretched by the traction of horsetail movement.

***rec8-F204S* mutant is defective in chromatin loop formation**

Next, we performed Hi-C analysis in the *rec8-F204S* mutant as in the *rec8-wt* (Figure

S6A). In the Hi-C contacts of three chromosomes at meiotic prophase (2.5 h), formation

of X-shaped contacts representing the bouquet orientation were apparently compromised

in the *rec8-F204S* mutant, similar to *rec8Δ* (Figure 6A). The alignment index of

chromosome 1 at meiotic prophase (2.5 h) decreased in *rec8-F204S* (1.6) compared with

the wild type (*rec8-wt*, 2.6) (Figure 6B). The findings were consistent with the fact that

chromatin is too flexible to resist the traction of horsetail movement in the *rec8-F204*S

mutant (Figures 5E and 5F). Moreover, in the amplified 2 kbp binning view of Hi-C data,

a clear decrease in the punctate Hi-C interactions was observed in *rec8-F204S*, although

it was not as prominent as in *rec8Δ* (Figure 6C). In addition, the shoulder in *P(s)* became

flattened in *rec8-F204S* compared with *rec8-wt*, both at 0 h and 2.5 h after meiosis

induction (Figure 6D). Consistently, chromatin loops with a length of approximately 20

kbp that formed in a Rec8-dependent manner were diminished at meiotic prophase in

rec8-F204S (Figure 6E). These findings suggest that in the *rec8-F204S* mutant, as in

rec8Δ, the Rec8-dependent meiosis-specific short chromatin loop structures are lost,

resulting in a concomitant loss of the structural property of the chromosome required for

proper alignment. Next, we performed a chromatin immunoprecipitation (ChIP) assay to

determine the localization pattern of Rec8 on chromatin. The accumulation of Rec8-

F204S protein apparently decreased compared with wild type Rec8 at the loci of known

cohesin-enriched regions. In contrast, the localization in other chromosome arm regions

seemed to be normal (Figure S6B). Rec8-enriched regions mostly coincided with each

base of the chromatin loops (Figures 2B and S2B). In the case of *rec8-F204S*, the amount

of Rec8 of in the non-enriched region appeared normal. Thus, the reduced Rec8

probability at known enriched regions possibly reflects the noticeable loss of chromatin

loop formation (Figure S6C).

Rec8-dependent axis-loop structure is required for LinE formation and

recombination

To delineate the significance of the Rec8-dependent axis-loop structure in terms of

meiotic recombination, we examined the formation of LinEs during meiotic prophase. In

wild type cells, the LinE component Rec10, marked by mCherry, localized within the

entire nucleus, similar to Rec8 (Figures 7A and S7A; $CK1^+$, $rec8^+$, $rec11$-wt). In *rec8-*

F204S cells, however, Rec10-mCherry formed aberrant dotty or filamentous aggregates

within the nucleus (Figures 7A and S7A), as with *rec8Δ* or *rec11-5A*, in which LinE was

completely lost (Sakuno and Watanabe, 2015). Interestingly, Rec11 phosphorylation by

CK1, which is required for the assembly of LinEs (Sakuno and Watanabe, 2015), occurred

normally in the *rec8-F204S* mutant, similar to the wild type (Figure 7B). We also

expressed mimetic mutations of CK1-mediated phosphorylation of Rec11 (*rec11-5D*),

which can suppress the defects caused by the loss of Rec11 phosphorylation via CK1

(Sakuno and Watanabe, 2015). Strikingly, the expression of Rec11-5D protein did not

suppress defects in LinE formation or meiotic recombination in *rec8-F204S* (Figures S7A

and S7B). These results strongly suggest that axis-loop formation along sister chromatids

via Rec8 cohesin is still essential for LinE assembly, even in the presence of active CK1-

dependent regulation. Consistent with the LinE formation defect, *rec8-F204S* cells

showed a decreased frequency of crossover recombination at several loci (Figure 7C).

Intragenic recombination between the genomic *ade6-M26* allele and plasmid harboring

the *ade6-469* allele was also significantly reduced in *rec8-F204S,* similar to *rec8Δ* (Figure

S7C). Collectively, these results demonstrate the essential role of the Rec8-dependent

loop-axis structure, which may function as a scaffold for LinE assembly, thus facilitating

meiotic recombination.

Discussion

We demonstrate the Rec8-dependent formation of axis-loop structures during meiosis in

S. pombe, an asynaptic organism. The Rec8 cohesin functions as a scaffold for the

formation of the LinE axis, thereby ensuring meiotic recombination during meiosis.

Pairing of homologous chromosomes in *S. pombe* meiosis

Our Hi-C analysis revealed the emergence of an intriguing X-shaped contact map

representing tight contacts between chromosome arm regions near the telomere in *S.*

pombe meiotic prophase. This is consistent with the previous microscopy observation of

telomere clusters at the SPB (the bouquet orientation of chromosomes) during meiotic

prophase (Chikashige *et al.*, 1994; Ding *et al.*, 1998). The Hi-C data indicate that contacts

also occur between non-homologous arms, not only between homologous arms. It should

be noted that chromosomes recognize their homologous partners to pair within the bundle

of all chromosomes. Therefore, there must be mechanisms to eliminate non-homologous pairing, while simultaneously promoting homologous pairing. Such selective forces between chromosomes can be provided by the oscillatory movement of telomere-bundled chromosomes (Chacón et al., 2016; Takao et al., 2019). Additionally, noncoding RNAs play a robust role in recognizing homologous pairs (Ding *et al.*, 2019; Ding et al., 2012). Importantly, Rec8 dominates noncoding RNA-mediated robust pairing (Ding *et al.*, 2016a; Ding *et al.*, 2016b; Hiraoka, 2020). Therefore, Rec8-mediated chromatin structures are an absolute requirement in the pairing homologous chromosomes during meiosis.

Loop-axis structure of chromatin in *S. pombe* meiosis

Upon entry into meiosis in *S. pombe*, chromosomes form a loop between the Rec8 binding sites (Figures 3B and S3B), which is also evident in *S. cerevisiae* meiosis (Muller et al., 2018; Schalbetter *et al.*, 2019). Although this Rec8 geometry is undetectable via Hi-C in mammalian meiosis (Luo et al., 2020; Patel *et al.*, 2019; Wang et al., 2019), it is argued that the chromosomal position of Rec8 or loop location varies from one cell to another in mammals (Patel *et al.*, 2019). Importantly, the Rec8 axis-loop structure formed normally without LinEs in *S. pombe* (*rec10Δ* in Figure 3A). Therefore, we emphasize that the Rec8-dependent axis-loop structure is distinct from LinEs and not a part of them. This relationship seems to be valid between cohesin-dependent axis-loop structures and lateral elements in other SC-forming organisms (Fujiwara *et al.*, 2020; Fukuda *et al.*, 2010; Pelttari *et al.*, 2001). With the emergence of Rec8-dependent loop structures, TAD-like signals observed in mitotic interphase cells mostly disappeared during meiotic prophase (Figures 3A and 3E), similar to mammalian spermatocytes (Luo *et al.*, 2020; Patel *et al.*,

2019; Wang *et al.*, 2019). The emergence of a loop structure with the disappearance of TADs is probably due to the switch of kleisin subunits from Rad21 to Rec8 that occurs upon the commitment to meiosis.

Notably, in the meiotic prophase of *wp1l*Δ cells, the Rec8 axis became thicker, while each chromatin loop became longer (Figures 4A, 4C, 4D, and 4E), as has been described in mammalian somatic interphase cells (Haarhuis *et al.*, 2017). In the case of *S. cerevisiae*, *WPL1* deletion was also reported to cause chromosomal compaction and recombination defects with an increase in Rec8 amounts on chromatin (Challa *et al.*, 2016). If the axial structures become thicker and the chromosomes become more compact, one would intuitively expect that the alignment of homologs would be easier. However, the opposite was true: X-shaped interactions tended to be reduced in the Hi-C analysis of *wpl1*Δ cells, and alignment of chromosome arms also became defective in *wpl1*Δ (Figure S4F), similar to *pds5*Δ (Ding *et al.*, 2016b). Pds5 associates with the cohesin complex and is required for the chromosomal localization of Wpl1 (Ding *et al.*, 2006; Goto et al., 2017; Lin and O'Connell, 2017), explaining the similar phenotypes of *pds5*Δ and *wpl1*Δ, such as thicker Rec8 axis formation and the alignment of chromosome arms during meiosis (Figures 4A, S4E, and S4F). These alignment defects are presumably caused by the overall stiffness of the chromosomes with thicker axes in *wpl1*Δ. In fact, examination of the chromosomes during horsetail movements in *wpl1*Δ rarely revealed torsional turning (Figure S4G, Supplemental Movies 1, and 3). Thus, our findings imply an interesting mechanism by which *S. pombe* maintains proper chromosome stiffness through the function of the Rec8 cohesin.

Possible mechanisms for formation of the Rec8 chromatin axis

The Rec8-dependent emergence of punctate Hi-C contacts only during meiosis suggests mechanisms by which the chromatin loop structure is formed differently from mitotic Rad21 cohesin. Fluorescence microscopy revealed that the Rec8 axis in the *wpl1Δ* background is reproduced in the mutant of Rec10, Hop1 (a homolog of HORMAD), or Mek1 (Figure S5J). All are required for LinE formation (Lorenz et al., 2006). The same was true in the Rec12 mutant and regulators of Rec12, such as Rec7, Rec15, and Mei4 (Figures S5B and S5J). Our Hi-C analysis demonstrated that the chromatin loop structure was also maintained in *rec10Δ* and *rec12Δ* (Figures 3A and S3A). Therefore, critical regulators of Rec8 axis formation remain to be identified among the factors expressed in a meiosis-specific manner and are required for recombination.

Mechanisms of chromatin loop formation have been proposed in mammalian somatic cells. *cis*-Looping of distal chromatin or TAD formation is thought to be established by the action of the Rad21 cohesin complex anchored on chromatin via the CCCTC-binding factor (CTCF) (Li et al., 2020; Nora et al., 2017; Rao et al., 2014; Wutz et al., 2017). More recently, a loop extrusion model was proposed for the formation of chromatin loops by SMC protein complexes, including cohesin (Davidson and Peters, 2021; Fudenberg *et al.*, 2016; Goloborodko et al., 2016; Sanborn *et al.*, 2015). In this model, the SMC protein complex interacts with DNA, reels, and extrudes it through its own ring structure to form a DNA loop (Davidson et al., 2019; Ganji et al., 2018; Kim et al., 2020; Kschonsak et al., 2017; Terakawa et al., 2017). At least *in vitro*, human Rad21 cohesin is able to extrude DNA, but it requires the cohesin loading factors Scc2 and Scc4 (Davidson *et al.*, 2019). Interestingly, CTCF is a conserved factor only in vertebrates (Hore et al., 2008). Thus,

there are no homologs of CTCF in yeasts. In contrast, Scc2 and Scc4 are conserved from

yeasts to humans (Mis4 and Ssl3, respectively, in *S. pombe*). Interestingly, chromatin loop

structures were already observed in *S. pombe* at meiosis entry (0 h) when the Rec8 cohesin

is loaded by Mis4 and Ssl3 before pre-meiotic DNA replication (Figures 3A and S1A).

Furthermore, the *rec8-F204S* mutant isolated in this study showed a significant defect in

the formation of loop structures, despite maintaining the cohesion function. Strikingly,

phenylalanine at position 204 in Rec8 is located in the domain where kleisin interacts

with Scc2/Mis4, as previously shown by an *in vitro* binding assay (Kikuchi et al., 2016).

Thus, as an alternative scenario for an appealing molecular mechanism in the

establishment of meiosis-specific chromatin loop structures, Rec8 and Mis4 with Ssl3

may also play an important role. Mis4 with Ssl3 may replace the function of CTCF in

anchoring cohesin to chromatin. It is also plausible that some changes in chromatin may

have occurred upon entry into meiosis, such as the modification of histones suitable for

the formation of axes with chromatin loops by the Rec8 cohesin.

Rec8 axis as a scaffold for LinE assembly

In the *rec8-F204S* mutant, which was unable to form a loop structure, LinE formation

failed to occur and the recombination frequency was markedly reduced (Figures 7C, S7B,

and S7C). However, phosphorylation of Rec11, which is required for LinE formation,

was intact and the distribution of Rec8-F204S along the chromosomes was largely normal

(Figures 7B and S6B). Based on the observations of the localization of Rec10 (LinE) by

ChIP and fluorescence microscopy, Rec10 and Rec8 overlap in some regions of the

chromosome, but not in many other chromosome regions (Ding et al., 2021; Miyoshi *et*

al., 2012; Sakuno and Watanabe, 2015). It is likely that the vast majority of Rec10 and

other LinE factors initially localize to chromosomes through interaction with Rec8

cohesin, and then are redistributed along the chromosome appropriately. We speculate

that the oligomerization of LinE factors, such as Rec10, is necessary for the formation of

LinE, which requires a certain amount of Rec8 to be present in a continuous axial

structure. The Rec8 axis acts as a scaffold to trigger the axial formation of LinEs.

Otherwise, LinE factors associated with the chromosome are not stabilized (Figure 7D).

As a result, LinE factors are released from the chromosome, and finally form aggregates

in the nucleus (Figures 7A, S7A, and 7D), due to their propensity to adhere to each other,

as in the polycomplex formed by mutants with defects in SC formation during meiosis of

S. cerevisiae (Zickler and Kleckner, 1999). Crucially, in addition to Rec11-5D mutant

protein, expression of the Rec11-Rec10 fusion protein, which can suppress the defects

caused by the loss of Rec11 phosphorylation (Sakuno and Watanabe, 2015), did not

suppress the LinE formation defect of *rec8-F204S* (Figures S7A, S7B and S7D). This

result strongly supports the view that the interaction between Rec8 and Rec10 is not

stabilized in the absence of the Rec8 axis along the chromatin. LinEs in *S. pombe* are

crucial in the assembly of complexes on chromatin that induce double-strand breaks on

DNA. This explains why a Rec8-dependent axis-loop structure is required for meiotic

recombination. Importantly, F204 in *S. pombe* Rec8 appears to be conserved among some

eukaryotes, including mammals. Although the relevant residue in mouse is leucine, we

demonstrated that the phenylalanine to leucine substitution in *S. pombe* Rec8 showed

almost no defects in axis formation (Figures 5D and S5I). Rec8 and Rec11/SA3 are

required for lateral element formation in mammals, as well as for axial element formation

in *S. pombe* (Fukuda *et al.*, 2014; Hopkins *et al.*, 2014; Llano *et al.*, 2014; Sakuno and

Watanabe, 2015; Winters *et al.*, 2014). Thus, there may be a mechanism by which

meiosis-specific cohesin-dependent axial structures can function as a scaffold in the

formation of lateral elements in mammals.

In conclusion, the current results obtained in *S. pombe* identify crucial functions of Rec8-

dependent assembly of chromatin structures that facilitate the execution of meiotic

recombination essential for faithful propagation of genetic materials into the progenies

during meiosis. The diverse and essential functions of the Rec8 cohesin in supporting

meiotic recombination may provide a basis for understanding the molecular mechanisms

of meiotic chromosome axis maturation and regulation of recombination in mammals,

potentially related to miscarriages and congenital abnormalities in humans (Nagaoka *et

al.*, 2012; Webster and Schuh, 2017).

Acknowledgments

We thank Drs. Masamitsu Sato, Shigehiro Kawashima and Juraj Gregan for providing

useful strains, Dr. Kayoko Tanaka for the helpful advice to induce highly synchronous

meiosis and all members of our laboratory for their valuable support and discussion. This

work was supported by JSPS KAKENHI grants JP26711020 (to T.S.), JP19K06503 (to

D.Q.D.), JP18H05528 (to T.H.), JP20H00454 and JP18H05533 (to Y.H.), and also by the

National Institutes of Health/National Institute of General Medical Sciences

(R01GM124195 to K.N.).

Author contributions

T.S., S.T., T.H., N.K., and Y.H. conceived and designed the experiments; T.S. conducted

almost all of the experiments except for the Hi-C, under supervision of T.H. and Y.H; S.T.

and O.I. performed Hi-C related analyses, under supervision of N.K; Bioinformatics-

related works were implemented by H.T; D.-Q.D. provided the data about the pairing

state in fission yeast meiosis; T.S., T.H., K.N., and Y.H. wrote the manuscript with the

input from all the authors.

Declaration of interest

The authors declare no competing interests.

 Figure legends

Figure 1. Hi-C analysis firmly recapitulates the structure of meiotic bouquet

chromosomes

(A) Chromosome arrangement in mitosis and meiosis in *S. pombe*. In mitosis (left),

centromeres are clustered at the spindle pole body (SPB) and display the Rabl orientation.

The grey lines indicate sister chromatids. In meiotic prophase (right), grey and black lines

indicate a homolog pair of chromosomes. Chromosomes are bundled by the telomeres

clustered at the SPB via telomeres, representing the bouquet orientation. (B) Hi-C contact

maps at 20 kbp resolution for three chromosomes prepared at vegetative growth (VG), 0

h, and 2.5 h after meiosis entry. Schematic views of each chromosome are shown on the

top and the left. Green arrowheads represent the centromere clustering. Blue arrowheads

in meiosis 0 h and 2.5 h represent the telomere clustering as depicted by the left lower

diagonal lines. The color scale on the right represents the percent ranking of contact

scores. (C) Heat maps showing the difference of contacts detected by Hi-C. Left,

subtraction of VG data from 0 h data. Right, subtraction of 0 h data from 2.5 h data. The

color scale on the right represents the percent ranking of difference scores. (D) Degree of

alignment of the two chromosome arms of telomere-bundled chromosome 1 is quantified

as the alignment index (Figure S1D) and is shown as the value along with a black vertical

line in the Hi-C contact map.

Figure 2. Rec8 cohesin shapes chromosomes into a structure suitable for its

alignment

(A) Hi-C contact maps of three chromosomes for the indicated strains, prepared from the

cells 0 h and 2.5 h after meiosis entry. The color scale on the right represents the percent

ranking of contact scores. (B) The alignment index for the indicated strains at meiotic

prophase (2.5 h after meiosis induction) is shown as the value along with a black vertical

line in the Hi-C contact map of chromosome 1.

Figure 3. Rec8 cohesin is crucial to construct the meiosis-specific chromatin loops

(A) Amplification of the Hi-C contact map in the 200 kbp region of chromosome 1 plotted

at 2 kbp resolution for the indicated strains at meiosis (0 h) and 2.5 h (at VG only for wild

type). The black arrow depicts the loop structure. (B) A merged view of the Hi-C contact

map and ChIP-chip data of Rec8 at the 200 kbp region of chromosome 1 shown in (A).

ChIP-chip data were adopted from the previously described source (Ito *et al.*, 2014). (C)

Contact probability versus genomic distance, *P(s)*, for VG (red), meiosis 0 h (blue) and

2.5 h (orange) of the wild type. The "shoulder" marked by the dotted line at approximately

50 kbp represents an increase in interactions within a region shorter than that length. (D)

Contact probability versus genomic distance, *P(s)*, for wild type (red), *rec8Δ* (blue),

rec10Δ (orange), and *rec12Δ* (green) 2.5 h after meiotic induction. (E) Amplification of

Hi-C contact maps at the 500 kbp region of chromosome 1and 2 plotted at 5 kbp

resolution for wild type strain at VG, meiosis 0 h and 2.5 h. The dotted line depicts the

TAD-like structure. (F) Schematic diagram of chromosome conformation at meiotic

prophase in *S. pombe*. All chromosomes are tightly aligned from the telomeres to the

centromeres. Rec8 establishes the meiosis-specific chromatin loop structure, irrespective

of LinEs or recombination.

Figure 4. Wpl1 regulates chromatin loop length during meiotic prophase in *S. pombe*

(A) Serial Z sections (0.3 μm each) of Rec8-GFP signals at meiotic prophase in wild type

and *wpl1Δ*. Axis-like signals of Rec8 are clearly observed in *wpl1Δ*. (B) Upper panel

displays Hi-C contact maps of three chromosomes at 20 kbp resolution in *wpl1Δ* at

vegetative growth (VG) and 0 h, and 2.5 h after meiosis entry, with schematic views of

each chromosome. The color scale on the right represents the percent ranking of contact

scores. The lower panel displays heat maps showing the difference of contacts detected

in each genome region by Hi-C in wild type and *wpl1Δ* at VG, and meiosis 0 h and 2.5 h.

The color scale on the right represents the percent ranking of difference scores. (C)

Amplification of Hi-C contact maps showing the 200 kbp region of chromosome 2 plotted

at 2 kbp resolution for the indicated strains at VG and meiosis 0 h and 2.5 h. (D) *P(s)*s for

meiosis 0 h and 2.5 h of wild type (red and blue) and *wpl1Δ* (orange and green). The blue

or green dotted line represent a shoulder of wild type or *wpl1Δ* at 2.5 h after meiosis entry,

respectively. (E) The number of loops per chromatin distance in *wpl1Δ* and wild type was

calculated. The value obtained by dividing *wpl1Δ* data by wild type data is shown as the

$\log_2$ ratio.

Figure 5. Isolation of *rec8* mutant showing defect in axis formation with intact

cohesion function

(A) The left panel shows a schematic view of cohesin complexes in fission yeast in

mitosis and meiosis. In mitosis, Rad21-Psc3 cohesin complex ubiquitously locates along

the whole chromosomes. In meiosis, Rec8-Psc3 cohesin complex exists only at the

centromeres and Rec8-Rec11 cohesin complex at the chromosomal arm regions. The

right panel shows the scheme for isolation of *rec8* mutants that specifically show defects

in axis formation, but normal cohesion in meiotic prophase. The promoter of the *rec8*

gene is replaced by the *adh1* promoter, and the promoter of the *rec11* gene is replaced by

the *adh41* (a weaker version of *adh1*) promoter, to allow their expression both in mitosis

and meiosis. After transformation of a library of GFP-fused *rec8* mutants with a selective

marker *bsdR* (not shown here), the obtained blasticidin-resistant colonies were

subsequently induced to meiosis to observe the axial Rec8-GFP signals. Growth rate of

the blasticidin-resistant clone acts as an indicator of the functionality of Rec8-Rec11

cohesin. (B) Representative pictures of Rec8-GFP with a cell shape (white dotted line)

during meiotic prophase in wild type and the isolated *rec8-F204S* mutant in this screening.

(C) Cohesion defects in the chromosome arm region (*cut3*$^+$ locus, left arm of chromosome

2) were examined in the indicated cells arrested at prophase I by the *mei4Δ* mutation.

After fixation by methanol, the number of heterozygous *cut3*-GFP signals was counted

(n > 150). A score of 2 or 1.5 *cut3*-GFP signals represents the complete or partial loss of

sister chromatid cohesion, respectively. The *rec8-S552P* identified in this screening

displays a remarkable defect in cohesion that is stronger than that of *rec8Δ* (see the text).

Error bars show standard deviations (SD) from three independent experiment. P values

were obtained from Chi-square test for statistical significance. (D) The upper panel is a

schematic diagram of *S. pombe* Rec8. Domains for the known interaction with cohesin-

related factors are shown. The position of F204 is also indicated by the red arrowhead.

The lower panel depicts the alignment of Rec8 homologs in *Shizosaccharomyces pombe*

(Sp), *S. octosporus* (So), human (h), mouse (m) and frog (*Xenopus leaves*) (x). (E) Time-

lapse observation (5 min intervals) of horsetail nuclear movement in the indicated cells

stained with histone H2B-mCherry. The right cartoon represents a model of the chromatin

state for each strain. In *rec8Δ*, Rad21 partially compensates the cohesion defect. (F) The

upper left panel shows a schematic drawing of the distance of *ade8-lacO* inserts from the

telomere of chromosome II. The upper right panels represent images of GFP signals for

the indicated strains. Brighter GFP signals at the edge of nucleus represent telomeres

(Taz1-GFP) and weaker GFP signals represent the *ade8* locus. The lower graph displays

a plot of the distance between the telomere and the *ade8* locus in each cell during meiotic

prophase for the indicated strains with average distance (n=55).

Figure 6. *rec8-F204S* mutant displays defective chromatin loop formation

(A) Hi-C contact maps of three chromosomes at 20 kbp resolution in wild type *rec8*-3×Pk

and *rec8-F204S*-3×Pk cells at 0 h and 2.5 h after meiosis entry, with schematic views of

each chromosome. The color scale on the right represents the percent ranking of contact

scores. (B) The alignment index for indicated strains at meiotic 0 h and 2.5 h after meiosis

induction is shown as the value along with a black vertical line in the Hi-C contact map

of chromosome 1. (C) Amplification of Hi-C contact maps of the 200 kbp region of

chromosome 1 (upper) and chromosome 2 (lower) at 2 kbp resolution, for the indicated

strain at meiosis 0 h and 2.5 h. (D) Contact probability versus genomic distance, *P(s)*, for

meiosis 0 h and 2.5 h of *rec8-wt* (red and blue) and *rec8-F204S* (orange and green). A

shoulder observed in meiosis 2.5 h of *rec8-wt* was shown with dotted line. (E) Number

of each loop per length quantified using genome-wide loop caller HiCCUPS for the

indicated strains at meiosis 0 h (left) and 2.5 h (right).

Figure 7. Rec8-dependent chromatin loop formation and meiotic recombination

(A) Representative pictures of Rec10-mCherry with cell shape (white dotted line) during

meiotic prophase for the indicated strains. The yellow arrows depict the aggregates of

LinE. (B) Western blot analysis. Whole-cell extracts were prepared from meiotic

prophase (2.5 h after meiosis induction) expressing Rec8-3×Flag in wild type and *rec8-*

F204S, with Psm1-3×HA. Rec11 phosphorylation by CK1 was detected by anti-

phosphorylated Rec11 (at T7). The monoclonal anti-tubulin (DM1A, Abcam) antibody

was used as a loading control. (C) Intergenic meiotic recombination for *ura1-lys3* (n >

700), *cox7-vps38* (n > 500), and *vps38-leu1* (n > 500) in the indicated zygotes. Error bars

show standard deviations (SD) from three independent experiment. (D) Model for the

Rec8 cohesin-dependent regulation of LinE assembly and meiotic recombination. After

the commitment in meiosis, Rec8 cohesin shown in blue is loaded onto the chromosomes

via cohesin loader complex (Mis4-Ssl3) shown in orange. Once loaded, Rec8 cohesin

promotes the loop formation that is antagonized by Wpl1 shown by purple. By meiotic

prophase, in parallel with the continuous loading of Rec8 and phosphorylation of Rec11

by CK1 (indicated by red triangles), chromatin looping with axis formation by Rec8

cohesin occurs to form chromatin to recruit LinE factors. See the detailed description in

the Discussion.

STAR Methods

***S. pombe* strains**

All media and growth conditions were as described previously (Moreno et al., 1991). Complete medium supplemented with uracil, lysine, adenine, histidine, and leucine (YE5S), minimal medium (EMM2), and sporulation-inducing medium (SPA and SSA) were used unless otherwise stated. The strains used in this study are listed in Table S1. Deletion of the endogenous genes ($rec12^+$, $rec11^+$, $rec7^+$, $rec15^+$, $mek1^+$, $rec8^+$, $rec10^+$, $hop1^+$, $wpl1^+$, $pds5^+$, and $mei4^+$) and tagging ($rec10^+$, $rec8^+$, $psm1^+$, $rec25^+$, $rad21^+$, and $rec11^+$) using GFP, mCherry, 3×hemagglutinin (HA) tag, 3×Pk (V5 tag), or 3×Flag was performed according to the PCR-based gene targeting method for *S. pombe* using the $ura4^+$, *kanMX6 (kanR), hphMX6 (hygR), natMX6 (natR)*, and *bsdR* genes as selection markers (Bahler et al., 1998; Rabitsch et al., 2001; Sato et al., 2005). Histone H2B tagging by mCherry for time-lapse observation has been described previously (Ruan et al., 2015). The CK1 shut off mutant, *rec11-5A* mutant, *rec11-5D* mutant, and the strain expressing Rec10-GFP-Rec11 fusion protein have been described previously (Sakuno and Watanabe, 2015). For the assay of crossing over recombination frequency, the *hygR* cassette was inserted 1215 bp upstream from the start codon of the $cox7^+$ gene and the *natR* cassette was inserted 1000 bp upstream from the start codon of $vps38^+$ gene using the PCR-based gene targeting method. To generate strains of *rec8* point mutants with tagging of GFP or 3×Pk tag, the endogenous $rec8^+$ locus was cloned with its promoter and terminator into pUC119. Using the NEBuilder system (New England Biolabs), a DNA fragment encoding GFP or 3×Pk tag with *kanR or bsdR* was introduced immediately before the stop codon of the *rec8* gene cloned in pUC119. Next, target sequences of $rec8^+$ were subsequently mutated using the KOD One Master Mix Blue (TOYOBO) to generate an amino acid

substitution. In case of *rec8-S552P*, a C-terminal portion of *rec8*$^+$ open reading frame

including the mutation site and marker gene with the *rec8*$^+$ terminator was amplified by

PCR and transformed into the wild type *rec8*$^+$ locus. Alternatively, in the case of N-

terminal mutations such as *rec8-F204S*, the promoter to terminator of the *rec8*$^+$ gene,

including the mutation and marker gene, was amplified by PCR and transformed into the

Δ*rec8::ura4*$^+$ strain. Correct replacement was confirmed using PCR and direct

sequencing. The strain in which the mitotic cohesin, Rad21 and Psc3, were exchanged

for meiotic cohesin (Rec8 and Rec11) has been described previously (Kitajima *et al.*,

2003). To construct the *pat1-as2* diploid with MDR-sup, haploid strains (*h*$^-$ *pat1-as2*

MDR-sup and *h*$^-$ *pat1-as2 MDR-sup mat-Pc*) were used (they were generously provided

by Masamitsu Sato, Waseda University). After introducing tagging or disruption of a gene

of interest in both haploids, protoplast fusion was performed as described previously

(Flor-Parra et al., 2014), except for the use of Lysing Enzyme (Sigma-Aldrich) instead of

Lallzyme or Zymolyase.

Preparation of meiotic cells

For microscopic observation of Rec8-GFP, Rec11-GFP, Rec10-mCherry, and Rec25-

mCherry, *h*^{90} cells cultured on YE plates at 26°C overnight were streaked on SSA plates,

or suspended in 20 mg/ml of leucine and spotted on SPA or SSA plates. The cells were

incubated at 26°C for 6–10 h to observe the cells at the horsetail stage. In case of

lacO/lacI-GFP observation for cohesion assays (Figure 5C and S5D), *h*$^+$ and *h*$^-$ cells were

mixed with 20 mg/ml of leucine and spotted on SSA or SPA plates. For the

immunoprecipitation of Rec8-GFP with Psm1-3HA (Figures S5F), *h*$^+$/*h*$^-$ diploid cells

with *mei4*Δ were used to arrest the cells at the meiotic prophase. In the case of diploid

cells, the cells were grown in EMM2 liquid medium containing 5 mg/ml NH$_4$Cl at a

density of 5×10^6 cells/ml at 25°C and then resuspended in EMM2 (1% glucose) medium

lacking NH$_4$Cl at a density of 1×10^7 cells/ml at 25°C for 15 h. For the

immunoprecipitation of Rec8-3×Flag with Psm1 (Figure S5G), ChIP assay (Figure S6B),

detection of phosphorylated Rec11 (Figure 7B), and Hi-C analyses, we used h^-/h^- diploid

cells with *pat1-as2* and *MDR-sup* alleles. Cells grown in YE liquid medium at a density

of 5–10×10^6 cells/ml at 25°C were collected by centrifugation and washed twice with

EMM2 (1% glucose) medium lacking NH$_4$Cl (EMM2-N). Cells were then resuspended

in EMM2-N with 50 mg/ml leucine at a density of 5×10^6 cells/ml and incubated at 25℃

for 6.5 h to induce G1 arrest. By adding the same volume of EMM2-N with 50 mg/ml

leucine and 15 µM of 1-NM-PP1 (Toronto Research Chemicals Inc.) (5–8×10^6 cells/ml),

cells can enter meiosis. They were further incubated for 2.5 h or more. Before harvesting

the meiotic cells, 1 mM phenylmethylsulfonyl fluoride (PMSF; Sigma-Aldrich) was

added to reduce protein degradation.

Rec8 mutant screening

The *rec8$^+$* cDNA mutant library used for the screening was constructed as follows. The

DNA region encoding *3×HA-ura4$^+$*of the plasmid VP068 (pUC119 base, *Padh1-*

*rec8$^+$*cDNA-*3×HA-ura4$^+$-Trec8*) (Yokobayashi *et al., 2003*) was replaced by the DNA

region encoding GFP-BsdR to produce VP068-GFP-bsd. Error-prone PCR of *rec8$^+$*

cDNA in VP068-GFP-BsdR by Ex taq (TaKaRa Bio) was performed using several

concentrations of MnCl$_2$ (final 0.05 mM, 0.15 mM, 0.25 mM, and 0.3 mM) and an

additional 1 mM dGTP or dATP with 0.2 mM of dNTP mix (total 8 conditions). By using

the NEBuilder system, amplified *rec8$^+$*cDNA pools (from eight conditions) were then

cloned into the fragment, which was amplified by inverse PCR of VP068-GFP-BsdR

without the *rec8*[+] cDNA region. In total, approximately 5300 *Escherichia coli*

transformants (representing library size) were generated. All colonies were recovered

from the plates, and plasmids (pRec8 library) were extracted. The pRec8 library was

estimated to contain approximately six mutations/kb. Using the pRec8 library as a

template, the region from 620 bp upstream of the *adh1*[+] promoter to 600 bp downstream

of the *rec8*[+] terminator was amplified by PCR. These fragments were used to transform

the screening host strains. More than 95% of blasticidin-resistant colonies obtained after

transformation showed correct integration, as confirmed by PCR. Rec8-GFP signals

during meiotic prophase were observed among the blasticidin-resistant colonies, which

showed relatively normal growth.

Fluorescence microscopy

Fluorescence images were obtained using an Axioplan2 microscope (Carl Zeiss)

equipped with a Quantix cooled CCD camera (Photometrics) and a DeltaVision

microscope system (GE Healthcare Inc.) equipped with a CoolSNAP HQ2 cooled CCD

camera (Photometrics) and a 60× Plan-ApoN SC oil immersion objective lens (NA=1.40;

Olympus). The brightness of the images was adjusted using Fiji software (Schindelin et

al., 2012) without changing the gamma settings. The pairing frequency was quantified as

described previously (Ding et al., 2004).

For time-lapse live-cell imaging of Rec8-GFP and H2B-mCherry, cells were

cultured overnight in liquid EMM2 at 26°C to reach stationary phase, and cells were

spotted onto SPA plates to induce meiosis, and incubated at 26°C for 4–5 h. Zygotes

suspended in 1% glucose EMM2 (1% glucose) without nitrogen were sonicated briefly

and immobilized on a coverslip coated with lectin in a 35 mm glass-bottomed culture dish

(MatTek) and observed at 26°C in liquid EMM2 (1% glucose) without nitrogen using a

DeltaVision microscope in an air-conditioned room. Images were collected at

approximately nine focal sections with 0.45 μm interval at each time point. The images

were collected every 5 min at three focal sections to minimize the excitation light during

long-term observation. Images were subsequently acquired using Z-sectioning and

projected according to the SoftWoRx software program (Applied Precision).

Immunoprecipitation

The cell extracts ($\sim 8 \times 10^7$ cells) were prepared by bead-beating cells in IP buffer (50 mM

HEPES-KOH at pH 8.0, 150 mM NaCl, 1 mM dithiothreitol, 2.5 mM $MgCl_2$, 20%

glycerol, 1 mM PMSF, 0.8% Triton X-100, 0.1% sodium deoxycholate, and 5× protease

inhibitor cocktail (P8215; Sigma-Aldrich). Fifty units of DNase I (TaKaRa Bio) were

added to the cell extract and sonicated (Branson Inc.). After collecting the supernatant as

whole-cell extracts by centrifugation at 15,00 rpm for 30 min, immunoprecipitation was

performed by incubating these extracts for 1 h at 4°C with anti-GFP polyclonal antibodies

(Rockland) or anti-Flag polyclonal antibodies (anti-DDDDK, MBL), and an additional 2

h at 4°C with 20 μl of a slurry of IgG-conjugated dynabeads (Invitrogen). After washing

the beads five times with IP buffer, the whole-cell extracts and immunoprecipitates were

subjected to western blotting using monoclonal anti-GFP (JL-8, Roche), Flag-M2

monoclonal antibody (Sigma-Aldrich), monoclonal anti-HA (3F10, Roche), and

monoclonal anti-tubulin (DM1A, Abcam) antibodies.

ChIP assay

The procedures were carried out as described previously (Yokobayashi and Watanabe, 2005). Anti-FLAG M2 monoclonal antibody (Sigma-Aldrich) and IgG conjugated dynabeads (Invitrogen) were used for immunoprecipitation. DNA prepared from the whole-cell extracts or immunoprecipitated fractions was analyzed via quantitative PCR (qPCR) with the StepOnePlus Real-Time PCR system (Thermo Fisher Scientific) using Power SYBR Green PCR Master Mix$^{©}$ Green I Master (Thermo Fisher Scientific). The primers used for PCR have been described previously (Sakuno and Watanabe, 2015). Immunoprecipitation with control mouse IgG antibody (Abcam) was performed in each experiment to account for nonspecific binding in the ChIP fractions and verify the significance of immunoprecipitation. The percent IP was quantified using three independent qPCR experiments.

Recombination assay

For intergenic recombination, haploid strains containing appropriate markers were mated and sporulated at 26°C on SSA plates for 30 h. The spores were isolated using β-glucuronidase solution (Wako) and plated on YE. The colonies were then replicated in selective media [EMM2-uracil, EMM2-lysine, EMM2-lysine, YE5S containing hygromycin B (FUJIFILM) or YE5S containing clonNAT (WERNER BioAgents)] and YE5S containing phloxine B (Wako) to select haploid cells. In the *ura1-lys3* interval, the degree of lysine auxotrophy among the uracil prototrophic colonies (n>700, replicated three times) was assessed. In the *cox7^{+}-vps38^{+}* interval, the degree of hygromycin B-resistant (*hygR*-containing) colonies among the clonNAT-resistant (*natR*-containing) colonies was assessed (n>500, and replicated three times). In the *vps38^{+}-leu1^{+}* interval, the degree of leucine prototrophic colonies among the clonNAT-resistant (*natR*-

containing) colonies was assessed (n>500, and replicated three times). For intragenic

recombination, h^{90} haploid strains containing the *ade6-M26* allele and plasmids harboring

the *ade6-469* allele according to the *ura4$^+$* marker (Ponticelli and Smith, 1992) were

sporulated at 26°C on SPA plates for 30 h. The spores were isolated by treatment with

glusulase and plated on YE5S containing phloxine B (to count all viable haploid spores)

and EMM2-adenine plates. The frequency of intragenic recombination was quantified as

the degree of adenine prototrophic colonies among viable haploid spores (n>7400, and

replicated three times).

Flow cytometry

A total of $1×10^7$ cells were fixed with 70% ethanol and incubated with 50 μg/ml RNase

A in 500 μl of 50 mM sodium citrate (pH 7.5) for 4 h at 37°C, and then stained with 0.5

μg/ml propidium iodide. DNA content of $1.5×10^5$ cells was measured on a FACS Calibur

instrument (BD Biosciences).

In situ Hi-C

In situ Hi-C experiments were performed as described previously (Tanizawa *et al.*, 2017)

with minor modifications. Asynchronous or meiotic cell cultures of $4×10^8$ cells were

fixed with 3% paraformaldehyde at 26°C for 10 min. The fixed cells were disrupted using

glass beads and Mini-Beadbeater-16 (BioSpec Products) in lysis buffer (50 mM

HEPES/KOH pH 7.5, 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium

deoxycholate, 1× Complete [Roche, 11836170001], and 1 mM PMSF). The lysed cells

were collected by centrifugation and permeabilized by resuspension with 1.11× NEBuffer

2 (B7002, NEB) containing 0.1% sodium dodecyl sulfate (SDS) at 62°C for 7 min. After

SDS was quenched by adding 1/9 volume of 10% Triton X-100, the permeabilized cells

were treated with 25 units of MboI (R0147, NEB) at 37°C overnight. The enzyme was

inactivated by incubation at 62°C for 20 min. After centrifugation, the pellet containing

the restricted DNA fragments was used in the following procedure. A half volume of fill-

in mix [1× NEBuffer 2, 150 µM biotin-14-dATP (NU-835-BIO14, Jena Bioscience), 150

µM dCTP, 150 µM dGTP, 150 µM dTTP, 0.4 units/µL Klenow fragment (M0210, NEB)]

was added to the pellet and incubated at 37°C for 45 min. Subsequently, three volumes

of ligation mix [1.33× T4 DNA ligase buffer (B0202, NEB), 1.11% Triton X-100, 133

µg/ml bovine serum albumin (B9001, NEB), 5.56 units/µl T4 DNA ligase (M0202,

NEB)] was added and incubated at room temperature (~25°C) for 4 h. After ligation, the

pellet was collected by centrifugation, resuspended in proteinase K solution [10 mM Tris-

HCl pH 8.0, 200 mM NaCl, 1 mM EDTA, 0.5% SDS, 0.5 mg/ml proteinase K (25530049,

Invitrogen)], and incubated at 65°C overnight. DNA was purified by phenol-chloroform

treatment followed by ethanol precipitation, and then fragmented by sonication with a

model UCD-200 Bioruptor (Diagenode). After the biotin-labeled DNA was pulled down

with Dynabeads Streptavidin T1 (65601, Invitrogen), the beads were washed twice with

TNB (5 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 M NaCl, 0.05% Tween-20) at 55°C for

2 min. To ligate the Illumina sequencing adaptors to the biotinylated DNA fragments, the

beads were successively treated with end repair mix [1× T4 DNA ligase buffer, 1 mM

dATP, 0.5 mM each dNTP, 0.5 units/µl T4 polynucleotide kinase (M0201, NEB), 0.12

units/µl T4 DNA polymerase (M0203, NEB), 0.05 units/µl Klenow fragment] at 25°C for

30 min, dA-tailing mix [1× NEBuffer 2, 0.5 mM dATP, 0.25 units/µl Klenow exo-

(M0212, NEB)] at 37°C for 30 min, and Illumina adaptor ligation mix [0.075x End repair

reaction buffer (E6050, NEB), 0.32× Ligation master mix (E7595, NEB), 0.011× Ligation

enhancer, and 45 nM NEBNext adaptor for Illumina] at 20°C for 15 min followed by

0.032 units/µl USER treatment at 37°C for 15 min, with two washes with TNB at 55°C

for 2 min after each reaction. The resultant adaptor-ligated Hi-C library was amplified by

PCR with five to seven cycles with Q5 master mix (M0544, NEB) and index primers

(E7335/E7500/E7710/E7730, NEB) before 37 bp paired-end sequencing using the

NextSeq 500 device (Illumina).

In situ Hi-C data analysis

In situ Hi-C data were processed as described previously (Tanizawa *et al.*, 2017). The

paired reads were separately aligned to the fission yeast genome (version ASM294v2.19)

using Bowtie2 (version 2.2.9) with the iterative alignment strategy. The aligned reads

were assigned to MboI fragments. Redundant paired reads derived from PCR biases,

aligned to repetitive sequences, or with low mapping quality (MapQ < 30) were removed.

Paired reads potentially derived from self-ligation and undigested products were also

discarded.

The fission yeast genome was divided into non-overlapped 20-kb, 5-kb, or 2-

kb bins. Raw contact matrices were constructed by counting paired reads assigned to two

bins, and Hi-C biases in the contact matrices were corrected using the ICE method

(Imakaev et al., 2012). The ICE normalization was repeated 30 times. ICE-normalized

contact scores were used for all heat maps, where the maximum intensity corresponded

to the top 5% score. For subtraction heat maps, the top 5% absolute values of difference

scores (T) were extracted. The T and -T scores were drawn as the maximum and minimum

intensity.

Loop detection

Knight and Ruiz (KR) normalized matrices (Rao *et al.*, 2014) in .hic format was

calculated from raw contact matrices using the 'pre' function of Juicer_tools (version

1.22.01) (Durand et al., 2016). Loop calling was performed for KR-normalized matrices

using HiCCUPS ('hiccups' function of Juicer_tools) with the following parameters: r 2000

-k KR -f .01 -p 3 -i 7 -d 10000".

864   **References**

866   Aoi, Y., Sato, M., Sutani, T., Shirahige, K., Kapoor, T.M., and Kawashima, S.A. (2014).
867   Dissecting the first and the second meiotic divisions using a marker-less drug-
868   hypersensitive fission yeast. Cell Cycle *13*, 1327-1334. 10.4161/cc.28294.
869

870   Bahler, J., Wu, J., Longtine, M.S., Shah, N.G., McKenzie III, A., Steever, A.B., Wach, A.,
871   Philippsen, P., and Pringle, J.R. (1998). Heterologous modules for efficient and versatile
872   PCR-based gene targeting in *Schizosaccharomyces pombe*. Yeast *14*, 943-951.
873

874   Bähler, J., Schuchert, P., Grimm, C., and Kohli, J. (1991). Synchronized meiosis and
875   recombination in fission yeast: observations with pat1-114 diploid cells. Curr Genet *19*,
876   445-451. 10.1007/BF00312735.
877

878   Bähler, J., Wyler, T., Loidl, J., and Kohli, J. (1993). Unusual nuclear structures in meiosis
879   prophase of fission yeast : a cytological analysis. J. Cell Biol. *121*, 241-256.
880

881   Cai, X., Dong, F., Edelmann, R.E., and Makaroff, C.A. (2003). The Arabidopsis SYN1
882   cohesin protein is required for sister chromatid arm cohesion and homologous
883   chromosome pairing. Journal of cell science *116*, 2999-3007.
884

885   Chacón, M.R., Delivani, P., and Tolić, I.M. (2016). Meiotic Nuclear Oscillations Are
886   Necessary to Avoid Excessive Chromosome Associations. Cell Rep *17*, 1632-1645.
887   10.1016/j.celrep.2016.10.014.
888

889   Challa, K., Lee, M.S., Shinohara, M., Kim, K.P., and Shinohara, A. (2016). Rad61/Wpl1
890   (Wapl), a cohesin regulator, controls chromosome compaction during meiosis. Nucleic
891   Acids Res *44*, 3190-3203. 10.1093/nar/gkw034.
892

893   Challa, K., Shinohara, M., and Shinohara, A. (2019). Meiotic prophase-like pathway for
894   cleavage-independent removal of cohesin for chromosome morphogenesis. Curr Genet
895   *65*, 817-827. 10.1007/s00294-019-00959-x.
896

897   Chikashige, Y., Ding, D.Q., Funabiki, H., Haraguchi, T., Mashiko, S., Yanagida, M., and
898   Hiraoka, Y. (1994). Telomere-led premeiotic chromosome movement in fission yeast.
899   Science *264*, 270-273. 10.1126/science.8146661.
900

901   Chikashige, Y., Ding, D.Q., Imai, Y., Yamamoto, M., Haraguchi, T., and Hiraoka, Y.
902   (1997). Meiotic nuclear reorganization: switching the position of centromeres and
903   telomeres in the fission yeast Schizosaccharomyces pombe. EMBO J *16*, 193-202.
904   10.1093/emboj/16.1.193.
905

Cipak, L., Hyppa, R.W., Smith, G.R., and Gregan, J. (2012). ATP analog-sensitive Pat1
protein kinase for synchronous fission yeast meiosis at physiological temperature. Cell
Cycle *11*, 1626-1633. 10.4161/cc.20052.

Crawley, O., Barroso, C., Testori, S., Ferrandiz, N., Silva, N., Castellano-Pozo, M., Jaso-
Tamame, A.L., and Martinez-Perez, E. (2016). Cohesin-interacting protein WAPL-1
regulates meiotic chromosome structure and cohesion by antagonizing specific cohesin
complexes. Elife *5*, e10851. 10.7554/eLife.10851.

Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.M. (2019). DNA
loop extrusion by human cohesin. Science *366*, 1338-1345. 10.1126/science.aaz3418.

Davidson, I.F., and Peters, J.M. (2021). Genome folding through loop extrusion by SMC
complexes. Nat Rev Mol Cell Biol *22*, 445-464. 10.1038/s41580-021-00349-7.

Davis, L., Rozalen, A.E., Moreno, S., Smith, G.R., and Martin-Castellanos, C. (2008).
Rec25 and Rec27, novel linear-element components, link cohesin to meiotic DNA
breakage and recombination. Curr Biol *18*, 849-854. 10.1016/j.cub.2008.05.025.

Ding, D.Q., Chikashige, Y., Haraguchi, T., and Hiraoka, Y. (1998). Oscillatory nuclear
movement in fission yeast meiotic prophase is driven by astral microtubules, as revealed
by continuous observation of chromosomes and microtubules in living cells. J Cell Sci
111 (Pt 6), 701-712.

Ding, D.Q., Haraguchi, T., and Hiraoka, Y. (2016a). A cohesin-based structural platform
supporting homologous chromosome pairing in meiosis. Curr Genet *62*, 499-502.
10.1007/s00294-016-0570-x.

Ding, D.Q., Matsuda, A., Okamasa, K., and Hiraoka, Y. (2021). Linear elements are stable
structures along the chromosome axis in fission yeast meiosis. Chromosoma.
10.1007/s00412-021-00757-w.

Ding, D.Q., Matsuda, A., Okamasa, K., Nagahama, Y., Haraguchi, T., and Hiraoka, Y.
(2016b). Meiotic cohesin-based chromosome structure is essential for homologous
chromosome pairing in Schizosaccharomyces pombe. Chromosoma *125*, 205-214.
10.1007/s00412-015-0551-8.

Ding, D.Q., Okamasa, K., Katou, Y., Oya, E., Nakayama, J.I., Chikashige, Y., Shirahige,
K., Haraguchi, T., and Hiraoka, Y. (2019). Chromosome-associated RNA-protein
complexes promote pairing of homologous chromosomes during meiosis in
Schizosaccharomyces pombe. Nat Commun *10*, 5598. 10.1038/s41467-019-13609-0.

Ding, D.Q., Okamasa, K., Yamane, M., Tsutsumi, C., Haraguchi, T., Yamamoto, M., and
Hiraoka, Y. (2012). Meiosis-specific noncoding RNA mediates robust pairing of
homologous chromosomes in meiosis. Science *336*, 732-736. 10.1126/science.1219518.

Ding, D.Q., Sakurai, N., Katou, Y., Itoh, T., Shirahige, K., Haraguchi, T., and Hiraoka, Y.
(2006). Meiotic cohesins modulate chromosome compaction during meiotic prophase in
fission yeast. J Cell Biol *174*, 499-508. jcb.200605074 [pii]
10.1083/jcb.200605074.

Ding, D.Q., Yamamoto, A., Haraguchi, T., and Hiraoka, Y. (2004). Dynamics of
homologous chromosome pairing during meiotic prophase in fission yeast. Dev Cell *6*,
329-341. 10.1016/s1534-5807(04)00059-0.
960

961 Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S., Huntley, M.H., Lander, E.S., and
962 Aiden, E.L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution
963 Hi-C Experiments. Cell Syst *3*, 95-98. 10.1016/j.cels.2016.07.002.
964

965 Estreicher, A., Lorenz, A., and Loidl, J. (2012). Mug20, a novel protein associated with
966 linear elements in fission yeast meiosis. Curr Genet *58*, 119-127. 10.1007/s00294-012-
967 0369-3.
968

969 Flor-Parra, I., Zhurinsky, J., Bernal, M., Gallardo, P., and Daga, R.R. (2014). A Lallzyme
970 MMX-based rapid method for fission yeast protoplast preparation. Yeast *31*, 61-66.
971 10.1002/yea.2994.
972

973 Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny, L.A.
974 (2016). Formation of Chromosomal Domains by Loop Extrusion. Cell Rep *15*, 2038-2049.
975 10.1016/j.celrep.2016.04.085.
976

977 Fujiwara, Y., Horisawa-Takada, Y., Inoue, E., Tani, N., Shibuya, H., Fujimura, S.,
978 Kariyazono, R., Sakata, T., Ohta, K., Araki, K., et al. (2020). Meiotic cohesins mediate
979 initial loading of HORMAD1 to the chromosomes and coordinate SC formation during
980 meiotic prophase. PLoS Genet *16*, e1009048. 10.1371/journal.pgen.1009048.
981

982 Fukuda, T., Daniel, K., Wojtasz, L., Toth, A., and Höög, C. (2010). A novel mammalian
983 HORMA domain-containing protein, HORMAD1, preferentially associates with
984 unsynapsed meiotic chromosomes. Exp Cell Res *316*, 158-171.
985 10.1016/j.yexcr.2009.08.007.
986

987 Fukuda, T., Fukuda, N., Agostinho, A., Hernandez-Hernandez, A., Kouznetsova, A., and
988 Hoog, C. (2014). STAG3-mediated stabilization of REC8 cohesin complexes promotes
989 chromosome synapsis during meiosis. The EMBO journal *33*, 1243-1255.
990 10.1002/embj.201387329.
991

992 Funaya, C., Samarasinghe, S., Pruggnaller, S., Ohta, M., Connolly, Y., Müller, J.,
993 Murakami, H., Grallert, A., Yamamoto, M., Smith, D., et al. (2012). Transient structure
994 associated with the spindle pole body directs meiotic microtubule reorganization in S.
995 pombe. Curr Biol *22*, 562-574. 10.1016/j.cub.2012.02.042.
996

997 Ganji, M., Shaltiel, I.A., Bisht, S., Kim, E., Kalichava, A., Haering, C.H., and Dekker, C.
998 (2018). Real-time imaging of DNA loop extrusion by condensin. Science *360*, 102-105.
999 10.1126/science.aar7831.
1000

1001 Goloborodko, A., Imakaev, M.V., Marko, J.F., and Mirny, L. (2016). Compaction and
1002 segregation of sister chromatids via active loop extrusion. Elife *5*. 10.7554/eLife.14864.

Golubovskaya, I.N., Hamant, O., Timofejeva, L., Wang, C.J., Braun, D., Meeley, R., and Cande, W.Z. (2006). Alleles of afd1 dissect REC8 functions during meiotic prophase I. Journal of cell science *119*, 3306-3315. 10.1242/jcs.03054.

Goto, Y., Yamagishi, Y., Shintomi-Kawamura, M., Abe, M., Tanno, Y., and Watanabe, Y. (2017). Pds5 Regulates Sister-Chromatid Cohesion and Chromosome Bi-orientation through a Conserved Protein Interaction Module. Curr Biol *27*, 1005-1012. 10.1016/j.cub.2017.02.066.

Grey, C., and de Massy, B. (2021). Chromosome Organization in Early Meiotic Prophase. Front Cell Dev Biol *9*, 688878. 10.3389/fcell.2021.688878.

Haarhuis, J.H.I., van der Weide, R.H., Blomen, V.A., Yáñez-Cuna, J.O., Amendola, M., van Ruiten, M.S., Krijger, P.H.L., Teunissen, H., Medema, R.H., van Steensel, B., et al. (2017). The Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. Cell *169*, 693-707.e614. 10.1016/j.cell.2017.04.013.

Hiraoka, Y. (2020). Phase separation drives pairing of homologous chromosomes. Curr Genet *66*, 881-887. 10.1007/s00294-020-01077-9.

Hong, S., Joo, J.H., Yun, H., Kleckner, N., and Kim, K.P. (2019). Recruitment of Rec8, Pds5 and Rad61/Wapl to meiotic homolog pairing, recombination, axis formation and S-phase. Nucleic Acids Res *47*, 11691-11708. 10.1093/nar/gkz903.

Hopkins, J., Hwang, G., Jacob, J., Sapp, N., Bedigian, R., Oka, K., Overbeek, P., Murray, S., and Jordan, P.W. (2014). Meiosis-specific cohesin component, Stag3 is essential for maintaining centromere chromatid cohesion, and required for DNA repair and synapsis between homologous chromosomes. PLoS genetics *10*, e1004413. 10.1371/journal.pgen.1004413.

Hore, T.A., Deakin, J.E., and Marshall Graves, J.A. (2008). The evolution of epigenetic regulators CTCF and BORIS/CTCFL in amniotes. PLoS Genet *4*, e1000169. 10.1371/journal.pgen.1000169.

Horie, S., Watanabe, Y., Tanaka, K., Nishiwaki, S., Fujioka, H., Abe, H., Yamamoto, M., and Shimoda, C. (1998). The Schizosaccharomyces pombe mei4+ gene encodes a meiosis-specific transcription factor containing a forkhead DNA-binding domain. Mol Cell Biol *18*, 2118-2129. 10.1128/MCB.18.4.2118.

Hunter, N. (2015). Meiotic Recombination: The Essence of Heredity. Cold Spring Harb Perspect Biol *7*. 10.1101/cshperspect.a016618.

Imakaev, M., Fudenberg, G., McCord, R.P., Naumova, N., Goloborodko, A., Lajoie, B.R., Dekker, J., and Mirny, L.A. (2012). Iterative correction of Hi-C data reveals hallmarks of chromosome organization. Nat Methods *9*, 999-1003. 10.1038/nmeth.2148.

Ishiguro, K., Kim, J., Shibuya, H., Hernandez-Hernandez, A., Suzuki, A., Fukagawa, T., Shioi, G., Kiyonari, H., Li, X.C., Schimenti, J., et al. (2014). Meiosis-specific cohesin mediates homolog recognition in mouse spermatocytes. Genes Dev *28*, 594-607. gad.237313.113 [pii]10.1101/gad.237313.113.

Ishiguro, K.I. (2019). The cohesin complex in mammalian meiosis. Genes Cells *24*, 6-30. 10.1111/gtc.12652.

Ito, M., Kugou, K., Fawcett, J.A., Mura, S., Ikeda, S., Innan, H., and Ohta, K. (2014). Meiotic recombination cold spots in chromosomal cohesion sites. Genes Cells *19*, 359-373. 10.1111/gtc.12138.

Kawashima, S.A., Takemoto, A., Nurse, P., and Kapoor, T.M. (2012). Analyzing fission yeast multidrug resistance mechanisms to develop a genetically tractable model system for chemical biology. Chem Biol *19*, 893-901. 10.1016/j.chembiol.2012.06.008.

Keeney, S. (2001). Mechanism and control of meiotic recombination initiation. Curr Top Dev Biol *52*, 1-53. 10.1016/s0070-2153(01)52008-6.

Kikuchi, S., Borek, D.M., Otwinowski, Z., Tomchick, D.R., and Yu, H. (2016). Crystal structure of the cohesin loader Scc2 and insight into cohesinopathy. Proc Natl Acad Sci U S A *113*, 12444-12449. 10.1073/pnas.1611333113.

Kim, E., Kerssemakers, J., Shaltiel, I.A., Haering, C.H., and Dekker, C. (2020). DNA-loop extruding condensin complexes can traverse one another. Nature *579*, 438-442. 10.1038/s41586-020-2067-5.

Kim, K.D., Tanizawa, H., Iwasaki, O., and Noma, K. (2016). Transcription factors mediate condensin recruitment and global chromosomal organization in fission yeast. Nat Genet *48*, 1242-1252. 10.1038/ng.3647.

Kim, Y., Shi, Z., Zhang, H., Finkelstein, I.J., and Yu, H. (2019). Human cohesin compacts DNA by loop extrusion. Science *366*, 1345-1349. 10.1126/science.aaz4475.

Kitajima, T.S., Yokobayashi, S., Yamamoto, M., and Watanabe, Y. (2003). Distinct cohesin complexes organize meiotic chromosome domains. Science *300*, 1152-1155.

Kleckner, N. (2006). Chiasma formation: chromatin/axis interplay and the role(s) of the synaptonemal complex. Chromosoma *115*, 175-194. 10.1007/s00412-006-0055-7.

Klein, F., Mahr, P., Galova, M., Buonomo, S.B.C., Michaelis, C., Nairz, K., and Nasmyth, K. (1999). A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis. Cell *98*, 91-103.

Kschonsak, M., Merkel, F., Bisht, S., Metz, J., Rybin, V., Hassler, M., and Haering, C.H. (2017). Structural Basis for a Safety-Belt Mechanism That Anchors Condensin to Chromosomes. Cell *171*, 588-600.e524. 10.1016/j.cell.2017.09.008.

1097

1098 Li, Y., Haarhuis, J.H.I., Sedeño Cacciatore, Á., Oldenkamp, R., van Ruiten, M.S.,
1099 Willems, L., Teunissen, H., Muir, K.W., de Wit, E., Rowland, B.D., and Panne, D. (2020).
1100 The structural basis for cohesin-CTCF-anchored loops. Nature *578*, 472-476.
1101 10.1038/s41586-019-1910-z.

1102

1103 Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling,
1104 A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive
1105 mapping of long-range interactions reveals folding principles of the human genome.
1106 Science *326*, 289-293. 10.1126/science.1181369.

1107

1108 Lin, S.J., and O'Connell, M.J. (2017). DNA Topoisomerase II modulates acetyl-regulation
1109 of cohesin-mediated chromosome dynamics. Curr Genet *63*, 923-930. 10.1007/s00294-
1110 017-0691-x.

1111

1112 Llano, E., Gomez, H.L., Garcia-Tunon, I., Sanchez-Martin, M., Caburet, S., Barbero, J.L.,
1113 Schimenti, J.C., Veitia, R.A., and Pendas, A.M. (2014). STAG3 is a strong candidate gene
1114 for male infertility. Human molecular genetics *23*, 3421-3431. 10.1093/hmg/ddu051.

1115

1116 Loidl, J. (2006). S. pombe linear elements: the modest cousins of synaptonemal
1117 complexes. Chromosoma *115*, 260-271. 10.1007/s00412-006-0047-7.

1118

1119 Lorenz, A., Estreicher, A., Kohli, J., and Loidl, J. (2006). Meiotic recombination proteins
1120 localize to linear elements in Schizosaccharomyces pombe. Chromosoma *115*, 330-340.
1121 10.1007/s00412-006-0053-9.

1122

1123 Lorenz, A., Wells, J.L., Pryce, D.W., Novatchkova, M., Eisenhaber, F., McFarlane, R.J.,
1124 and Loidl, J. (2004). S. pombe meiotic linear elements contain proteins related to
1125 synaptonemal complex components. J Cell Sci *117*, 3343-3351. 10.1242/jcs.01203.

1126

1127 Luo, Z., Wang, X., Jiang, H., Wang, R., Chen, J., Chen, Y., Xu, Q., Cao, J., Gong, X., Wu,
1128 J., et al. (2020). Reorganized 3D Genome Structures Support Transcriptional Regulation
1129 in Mouse Spermatogenesis. iScience *23*, 101034. 10.1016/j.isci.2020.101034.

1130

1131 Mehta, G.D., Kumar, R., Srivastava, S., and Ghosh, S.K. (2013). Cohesin: functions
1132 beyond sister chromatid cohesion. FEBS Lett *587*, 2299-2312.
1133 10.1016/j.febslet.2013.06.035.

1134

1135 Miyoshi, T., Ito, M., Kugou, K., Yamada, S., Furuichi, M., Oda, A., Yamada, T., Hirota,
1136 K., Masai, H., and Ohta, K. (2012). A central coupler for recombination initiation linking
1137 chromosome architecture to S phase checkpoint. Mol Cell *47*, 722-733.
1138 10.1016/j.molcel.2012.06.023.

1139

1140 Mizuguchi, T., Fudenberg, G., Mehta, S., Belton, J.M., Taneja, N., Folco, H.D.,
1141 FitzGerald, P., Dekker, J., Mirny, L., Barrowman, J., and Grewal, S.I.S. (2014). Cohesin-
1142 dependent globules and heterochromatin shape 3D genome architecture in S. pombe.
1143 Nature *516*, 432-435. 10.1038/nature13833.

Moore, D.P., and Orr-Weaver, T.L. (1998). Chromosome segregation during meiosis: building an unambivalent bivalent. Curr Top Dev Biol *37*, 263-299.

Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission yeast *Schizosaccharomyces pombe*. Methods Enzymol. *194*, 795-823.

Muller, H., Scolari, V.F., Agier, N., Piazza, A., Thierry, A., Mercy, G., Descorps-Declere, S., Lazar-Stefanita, L., Espeli, O., Llorente, B., et al. (2018). Characterizing meiotic chromosomes' structure and pairing using a designer sequence optimized for Hi-C. Mol Syst Biol *14*, e8293. 10.15252/msb.20188293.

Murakami-Tonami, Y., Yamada-Namikawa, C., Tochigi, A., Hasegawa, N., Kojima, H., Kunimatsu, M., Nakanishi, M., and Murakami, H. (2007). Mei4p coordinates the onset of meiosis I by regulating cdc25+ in fission yeast. Proc Natl Acad Sci U S A *104*, 14688-14693. 10.1073/pnas.0702906104.

Nagaoka, S.I., Hassold, T.J., and Hunt, P.A. (2012). Human aneuploidy: mechanisms and new insights into an age-old problem. Nat Rev Genet *13*, 493-504. 10.1038/nrg3245.

Nasmyth, K. (2001). Disseminating the genome: joining, resolving, and separating sister chromatids during mitosis and meiosis. Annu. Rev. Genet. *35*, 673-745.

Nora, E.P., Goloborodko, A., Valton, A.L., Gibcus, J.H., Uebersohn, A., Abdennur, N., Dekker, J., Mirny, L.A., and Bruneau, B.G. (2017). Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains from Genomic Compartmentalization. Cell *169*, 930-944.e922. 10.1016/j.cell.2017.05.004.

Olson, L.W., Eden, U., Egel-Mitani, M., and Egel, R. (1978). Asynaptic meiosis in fission yeast? Hereditas *89*, 189-199.

Pasierbek, P., Jantsch, M., Melcher, M., Schleiffer, A., Schweizer, D., and Loidl, J. (2001). A Caenorhabditis elegans cohesion protein with functions in meiotic chromosome pairing and disjunction. Genes & development *15*, 1349-1360.

Patel, L., Kang, R., Rosenberg, S.C., Qiu, Y., Raviram, R., Chee, S., Hu, R., Ren, B., Cole, F., and Corbett, K.D. (2019). Dynamic reorganization of the genome shapes the recombination landscape in meiotic prophase. Nat Struct Mol Biol *26*, 164-174. 10.1038/s41594-019-0187-0.

Pelttari, J., Hoja, M.R., Yuan, L., Liu, J.G., Brundell, E., Moens, P., Santucci-Darmanin, S., Jessberger, R., Barbero, J.L., Heyting, C., and Höög, C. (2001). A meiotic chromosomal core consisting of cohesin complex proteins recruits DNA recombination proteins and promotes synapsis in the absence of an axial element in mammalian meiotic cells. Mol Cell Biol *21*, 5667-5677. 10.1128/MCB.21.16.5667-5677.2001.

Peters, J.M., and Nishiyama, T. (2012). Sister chromatid cohesion. Cold Spring Harb Perspect Biol *4*. 10.1101/cshperspect.a011130.

Petronczki, M., Siomos, M.F., and Nasmyth, K. (2003). Un menage a quatre: the molecular biology of chromosome segregation in meiosis. Cell *112*, 423-440.

Phadnis, N., Cipak, L., Polakova, S., Hyppa, R.W., Cipakova, I., Anrather, D., Karvaiova, L., Mechtler, K., Smith, G.R., and Gregan, J. (2015). Casein Kinase 1 and Phosphorylation of Cohesin Subunit Rec11 (SA3) Promote Meiotic Recombination through Linear Element Formation. PLoS Genet *11*, e1005225. 10.1371/journal.pgen.1005225.

Ponticelli, A.S., and Smith, G.R. (1992). Chromosomal context dependence of a eukaryotic recombinational hot spot. Proc Natl Acad Sci U S A *89*, 227-231. 10.1073/pnas.89.1.227.

Rabitsch, K.P., Toth, A., Galova, M., Schleiffer, A., Schaffner, G., Aigner, E., Rupp, C., Penkner, A.M., Moreno-Borchart, A.C., Primig, M., et al. (2001). A screen for genes required for meiosis and spore formation based on whole-genome expression. Current biology : CB *11*, 1001-1009.

Rao, S.S., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., and Aiden, E.L. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. Cell *159*, 1665-1680. 10.1016/j.cell.2014.11.021.

Rao, S.S.P., Huang, S.C., Glenn St Hilaire, B., Engreitz, J.M., Perez, E.M., Kieffer-Kwon, K.R., Sanborn, A.L., Johnstone, S.E., Bascom, G.D., Bochkov, I.D., et al. (2017). Cohesin Loss Eliminates All Loop Domains. Cell *171*, 305-320.e324. 10.1016/j.cell.2017.09.026.

Ruan, K., Yamamoto, T.G., Asakawa, H., Chikashige, Y., Masukata, H., Haraguchi, T., and Hiraoka, Y. (2015). Meiotic nuclear movements in fission yeast are regulated by the transcription factor Mei4 downstream of a Cds1-dependent replication checkpoint pathway. Genes Cells *20*, 160-172. 10.1111/gtc.12207.

Sakuno, T., and Watanabe, Y. (2015). Phosphorylation of cohesin Rec11/SA3 by casein kinase 1 promotes homologous recombination by assembling the meiotic chromosome axis. Dev Cell *32*, 220-230. 10.1016/j.devcel.2014.11.033.

Sanborn, A.L., Rao, S.S., Huang, S.C., Durand, N.C., Huntley, M.H., Jewett, A.I., Bochkov, I.D., Chinnappan, D., Cutkosky, A., Li, J., et al. (2015). Chromatin extrusion explains key features of loop and domain formation in wild-type and engineered genomes. Proc Natl Acad Sci U S A *112*, E6456-6465. 10.1073/pnas.1518552112.

Sato, M., Dhut, S., and Toda, T. (2005). New drug-registant cassettes for gene disruption and epitope tagging in Schizosaccharomyces pombe. Yeast *22*, 583-591.

Schalbetter, S.A., Fudenberg, G., Baxter, J., Pollard, K.S., and Neale, M.J. (2019). Principles of meiotic chromosome assembly revealed in S. cerevisiae. Nat Commun *10*,

1238   4795. 10.1038/s41467-019-12629-0.

1239

1240   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
1241   Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
1242   platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

1243

1244   Sharif, W.D., Glick, G.G., Davidson, M.K., and Wahls, W.P. (2002). Distinct functions of
1245   S. pombe Rec12 (Spo11) protein and Rec12-dependent crossover recombination
1246   (chiasmata) in meiosis I; and a requirement for Rec12 in meiosis II. Cell Chromosome *1*,
1247   1. 10.1186/1475-9268-1-1.

1248

1249   Silva, M.C.C., Powell, S., Ladstätter, S., Gassler, J., Stocsits, R., Tedeschi, A., Peters,
1250   J.M., and Tachibana, K. (2020). Wapl releases Scc1-cohesin and regulates chromosome
1251   structure and segregation in mouse oocytes. J Cell Biol *219*. 10.1083/jcb.201906100.

1252

1253   Takao, K., Takamiya, K., Ding, D., Haraguchi , T., Hiraoka, Y., Nishimori, H., and Awazu,
1254   A. (2019). Torsional Turning Motion of Chromosomes as an Accelerating Force to Align
1255   Homologous Chromosomes during Meiosis. Journal of the Physical Society of Japan *88*
1256   (*023801*).

1257

1258   Tanizawa, H., Kim, K.D., Iwasaki, O., and Noma, K.I. (2017). Architectural alterations
1259   of the fission yeast genome during the cell cycle. Nat Struct Mol Biol *24*, 965-976.
1260   10.1038/nsmb.3482.

1261

1262   Tedeschi, A., Wutz, G., Huet, S., Jaritz, M., Wuensche, A., Schirghuber, E., Davidson,
1263   I.F., Tang, W., Cisneros, D.A., Bhaskara, V., et al. (2013). Wapl is an essential regulator
1264   of chromatin structure and chromosome segregation. Nature *501*, 564-568.
1265   10.1038/nature12471.

1266

1267   Terakawa, T., Bisht, S., Eeftens, J.M., Dekker, C., Haering, C.H., and Greene, E.C. (2017).
1268   The condensin complex is a mechanochemical motor that translocates along DNA.
1269   Science *358*, 672-676. 10.1126/science.aan6516.

1270

1271   Viera, A., Berenguer, I., Ruiz-Torres, M., Gómez, R., Guajardo, A., Barbero, J.L., Losada,
1272   A., and Suja, J.A. (2020). PDS5 proteins regulate the length of axial elements and
1273   telomere integrity during male mouse meiosis. EMBO Rep *21*, e49273.
1274   10.15252/embr.201949273.

1275

1276   Wang, Y., Wang, H., Zhang, Y., Du, Z., Si, W., Fan, S., Qin, D., Wang, M., Duan, Y., Li,
1277   L., et al. (2019). Reprogramming of Meiotic Chromatin Architecture during
1278   Spermatogenesis. Mol Cell *73*, 547-561.e546. 10.1016/j.molcel.2018.11.019.

1279

1280   Webster, A., and Schuh, M. (2017). Mechanisms of Aneuploidy in Human Eggs. Trends
1281   Cell Biol *27*, 55-68. 10.1016/j.tcb.2016.09.002.

1282

1283   Winters, T., McNicoll, F., and Jessberger, R. (2014). Meiotic cohesin STAG3 is required
1284   for chromosome axis formation and sister chromatid cohesion. The EMBO journal *33*,

1256-1270. 10.1002/embj.201387330.

Wutz, G., Várnai, C., Nagasaka, K., Cisneros, D.A., Stocsits, R.R., Tang, W., Schoenfelder, S., Jessberger, G., Muhar, M., Hossain, M.J., et al. (2017). Topologically associating domains and chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins. EMBO J *36*, 3573-3599. 10.15252/embj.201798004.

Xu, H., Beasley, M.D., Warren, W.D., van der Horst, G.T., and McKay, M.J. (2005). Absence of mouse REC8 cohesin promotes synapsis of sister chromatids in meiosis. Developmental cell *8*, 949-961.

Yamamoto, A., and Hiraoka, Y. (2001). How do meiotic chromosomes meet their homologous partners?: lessons from fission yeast. Bioessays *23*, 526-533. 10.1002/bies.1072.

Yamamoto, A., and Hiraoka, Y. (2003). Monopolar spindle attachment of sister chromatids is ensured by two distinct mechanisms at the first meiotic division in fission yeast. EMBO J *22*, 2284-2296. 10.1093/emboj/cdg222.

Yokobayashi, S., and Watanabe, Y. (2005). The kinetochore protein Moa1 enables cohesion-mediated monopolar attachment at meiosis I. Cell *123*, 803-817.

Yokobayashi, S., Yamamoto, M., and Watanabe, Y. (2003). Cohesins determine the attachment manner of kinetochores to spindle microtubules at meiosis I in fission yeast. Mol Cell Biol *23*, 3965-3973. 10.1128/MCB.23.11.3965-3973.2003.

Young, J.A., Hyppa, R.W., and Smith, G.R. (2004). Conserved and nonconserved proteins for meiotic DNA breakage and repair in yeasts. Genetics *167*, 593-605. 10.1534/genetics.103.023762.

Zickler, D., and Kleckner, N. (1999). Meiotic chromosomes: integrating structure and function. Annu Rev Genet *33*, 603-754. 10.1146/annurev.genet.33.1.603.

Figure 1. Hi-C analysis firmly recapitulates the structure of meiotic bouquet chromosomes.

Figure 2. Rec8 cohesin shapes chromosomes into a structure suitable for its alignment

A

Meiosis 0 hr | Meiosis 2.5 hr

rec10Δ

rec12Δ

rec8Δ

95%
50%
0%

B

Meiosis 2.5hr

rec10Δ — M-index =2.8

rec12Δ — M-index =2.5

rec8Δ — M-index =1.5

Distance (Mb) / Position in Chr 1 (Mb)

Figure 3. Rec8 cohesin is crucial to construct the meiosis-specific chromatin loops.

Figure 4. Wpl1 regulates chromatin loop length during meiotic prophase in *S. pombe*.

A

B

C

Chr 1, 4.05~4.25Mb, 2kbp bins

D **E**

Figure 5. Isolation of *rec8* mutant showing defect in axis formation with intact cohesion function

Figure 6. *rec8-F204S* mutant displays defective chromatin loop formation

Figure 7. Rec8-dependent chromatin loop formation and meiotic recombination

A

Rec10-mCherry

wt

rec8-F204S

rec8Δ

B

C