# Using ultraconserved elements to reconstruct the termite tree of life

**Running title**: UCEs in termites

Simon Hellemans[1,*], Menglin Wang[1], Nonno Hasegawa[1], Jan Šobotník[2], Rudolf H. Scheffrahn[3], Thomas Bourguignon[1,2,*]

[1]Okinawa Institute of Science & Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa 904-0495, Japan.

[2]Faculty of Tropical AgriScience, Czech University of Life Sciences, Kamýcka 129, 16521 Prague, Czech Republic.

[3]Fort Lauderdale Research and Education Center, Institute for Food and Agricultural Sciences, 3205 College Avenue, Davie, Florida 33314 USA.

*Corresponding authors: simon.hellemans@gmail.com (SH); Thomas.Bourguignon@oist.jp (TB)

**ORCID references of authors**

Simon Hellemans: 0000-0003-1266-9134

Menglin Wang: 0000-0003-2206-9503

Nonno Hasegawa: 0000-0001-5149-9129

Jan Šobotník: 0000-0002-8581-637X

Rudolf H. Scheffrahn: 0000-0002-6191-5963

Thomas Bourguignon: 0000-0002-4035-8977

**Author contributions**

SH and TB conceptualized the experiments. JS and RS collected the samples. MW performed

lab experiments and generated data. SH, MW and NH analyzed the data. SH and TB wrote the

manuscript.

**Competing interests**

We declare we have no competing interests.

**Funding**

This work was supported by the Czech Science Foundation (project No. 15-07015Y), the

Internal Grant Agency of the Faculty of Tropical AgriSciences, CULS (20213112), the Japan

Society for the Promotion of Science (JSPS) through a postdoctoral fellowship for foreign

researchers awarded to SH (19F19819), and the Okinawa Institute of Science and Technology

through core unit funding.

## Acknowledgments

41   We thank Yves Roisin, Saran Traoré, and Guy Josens for providing specimens as well as

42   Crystal Clitheroe for helping with DNA extraction and library preparation. We also thank the

43   Scientific Computation and Data Analysis Section (SCDA) of the Okinawa Institute of Science

44   and Technology Graduate University, Okinawa, Japan, for providing access to the OIST

45   computing cluster.

## Electronic Supplementary Materials

48   Additional information is available for this manuscript.

## Data Archiving

The data that support the findings of this study will be deposited in GenBank (see

Supplementary Table S1 for details). Bait sequences, as well as the extracted UCE sequences,

will be deposited on Dryad (Supplementary Data 1-9).

**Abstract**

The phylogenetic history of termites has been investigated using mitochondrial genomes and

transcriptomes. However, both sets of markers have specific limitations. Mitochondrial

genomes represent a single genetic marker likely to yield phylogenetic trees presenting

incongruences with species trees, and transcriptomes can only be obtained from well-preserved

samples. In contrast, ultraconserved elements (UCEs) include a great many independent

markers that can be retrieved from poorly preserved samples. Here, we designed termite-

specific baits targeting 50,616 UCE loci. We tested our UCE bait set on 42 samples of termites

and three samples of *Cryptocercus*, for which we generated low-coverage highly-fragmented

genome assemblies and successfully extracted *in silico* between 3,426 to 42,860 non-duplicated

UCEs per sample. Our maximum likelihood phylogenetic tree, reconstructed using the 5,934

UCE loci retrieved from upward of 75% of samples, was congruent with transcriptome-based

phylogenies, demonstrating that our UCE bait set is reliable and phylogenetically informative.

Combined with non-destructive DNA extraction protocols, our UCE bait set provides the tool

needed to carry out a global taxonomic revision of termites based on poorly preserved

specimens such as old museum samples. The Termite UCE database is maintained at:

https://github.com/oist/TER-UCE-DB/.

**1. Introduction**

Termites are the most ancient lineage of social insects, with a fossil record dating back to the Early Cretaceous, ~135 million years ago (Mya) (Thorne *et al.*, 2000; Grimaldi & Engel, 2005; Engel *et al.*, 2009). All modern termites share a common ancestor estimated at 140-150 Mya by time-calibrated phylogenetic trees (Bourguignon *et al.*, 2015; Bucek *et al.*, 2019). However, the bulk of the modern termite species diversity belongs to the Termitidae, a lineage that originated during the early Eocene, ~50 Mya, as indicated by time-calibrated phylogenetic trees (Bourguignon *et al.*, 2015, 2017; Bucek *et al.*, 2019) and by the fossil record (Engel *et al.*, 2011). While the backbone of the phylogenetic tree of termites is now largely resolved, most termite species are still awaiting to be placed on the tree of life.

Our understanding of the phylogenetic history of termites was mostly based on mitochondrial markers and fossils to calibrate estimated times of divergence until Bucek *et al*. (2019) published a phylogenetic tree of termites based on transcriptome data. The first phylogenetic trees of termites were based on a couple of PCR-amplified mitochondrial markers, sometimes combined with nuclear 18S or 28S sequences and/or morphological characters, which hardly contributed any phylogenetic signal (e.g., Lo *et al.*, 2004; Inward *et al.*, 2007; Legendre *et al.*, 2008). These phylogenies provided a good overview of the relationships among the main termite lineages but lacked the robustness of phylogenetic trees inferred from full mitochondrial genomes (e.g., Cameron *et al.*, 2012; Bourguignon *et al.*, 2015, 2017). Full mitochondrial genomes, which became easy to sequence with the rise of second-generation sequencing technologies, resolve both shallow and deep divergences in the evolutionary history of termites and other insect lineages (Cameron, 2014), making them a marker of choice for phylogenetic reconstructions. However, mitochondrial genomes form a single marker, as all mitochondrial genes are linked and maternally inherited as a single package. Consequently, mitochondrial phylogenies are sometimes discordant with species phylogenies, especially for

closely related species and short internal branches that diverged in periods of time too brief for

alleles to coalesce (Whitfield & Lockhart, 2007; Degnan & Rosenberg, 2009). One example of

such discordance is provided by Sphaerotermitinae, the unambiguous sister group of

Macrotermitinae according to transcriptomic data; but supported as sister to non-

macrotermitine non-foraminitermitine Termitidae by mitochondrial genome phylogenies

(Figure 1; Bucek *et al.*, 2019). Phylogenies based on multiple independent nuclear markers are

needed to resolve the evolutionary history of organisms accurately.

Transcriptomes, the snapshot of genes expressed by an organism during tissue sampling,

include many independent nuclear markers that can be used to build robust phylogenetic trees.

Transcriptome-based phylogenies, reconstructed using up to ~4,000 single-copy orthologous

nuclear genes spanning over 7.7 million nucleotide positions, have provided a robust picture of

the ancient evolutionary history of termites (Bucek *et al.*, 2019). The sequencing of

transcriptomes is now affordable, but, unfortunately, RNA is unstable and can only be extracted

from samples that have been adequately preserved and stored, preventing the use of most

samples collected before the genomic era began and making the approach impractical for large-

scale studies. One alternative is to mine the conserved genetic markers present in whole-

genome shotgun sequencing datasets, such as some datasets generated to sequence

mitochondrial genomes.

Ultraconserved Elements (UCEs) are highly conserved nuclear regions whose functions

remain largely unknown (Bejerano *et al.*, 2004; Faircloth *et al.*, 2012). UCEs are found across

all regions of animal genomes, including the exonic, intronic, and intergenic regions.

Phylogenetic trees inferred from UCEs have contributed to our understanding of the

evolutionary history of various animal lineages spanning across the animal tree of life (e.g.,

Faircloth *et al.*, 2012; Ryu *et al.*, 2012; Smith *et al.*, 2014; White & Braun, 2019; Zhang *et al.*,

2019). Unlike transcriptomes, UCEs can readily be obtained from museum samples through

baiting conserved elements and their phylogenetically-informative flanking regions from

fragmented genome assemblies (Blaimer *et al.*, 2016; Faircloth, 2017; Derkarabetian *et al.*,

2019). No UCE bait set has been designed for termites so far. We filled this gap as follows: (*i*)

we designed *in silico* baits to capture UCEs; (*ii*) we compared phylogenetic trees reconstructed

using all possible combinations of mitochondrial genomes, nuclear ribosomal RNA genes, and

UCEs; and (*iii*) we showed that UCEs obtained from low-coverage shotgun genome assemblies

are an alternative to transcriptome-based phylogenies for the reconstruction of multi-gene

phylogenies. Finally, we set up a Termite UCE Database, thereby ensuring a long-term re-

usability of published data.

**2. Material and Methods**

*Biological samples and sequencing*

We used sequence data from 42 samples of termites and three samples of *Cryptocercus*, the

wood-feeding subsocial cockroach genus forming the sister group of termites. The sequencing

data of 14 species were retrieved from previous studies (for details, see Table S1). The

sequencing data from the remaining 31 species were obtained from samples preserved in 80%

ethanol stored at room temperature or from samples preserved in RNA-later® and stored at

temperatures ranging between -20°C and -80°C until DNA extraction. DNA was extracted

using the DNeasy Blood & Tissue extraction kit (Qiagen). Libraries were prepared using the

NEBNext® Ultra™ II FS DNA Library Preparation Kit (New England Biolabs) and the Unique

Dual Indexing Kit (New England Biolabs), with reagent volumes reduced to one-fifteenth of

recommended volumes. For samples preserved in 80% ethanol, libraries were prepared without

the enzymatic fragmentation step as the DNA of these samples is typically highly fragmented.

Libraries were pooled in equimolar concentration and paired-end sequenced using the Illumina

HiSeq X or Novaseq platforms at a read length of 150 bp.

*UCE loci identification and* in silico *bait design*

The identification of UCE loci was carried out using PHYLUCE *v*1.6.6 (Faircloth, 2016)

following the recommendations of the tutorial (https://phyluce.readthedocs.io/en/stable/) and

outlined workflow (Faircloth, 2017). Four publicly available genomes belonging to distantly

related termite species were used to design baits: *Zootermopsis nevadensis* (Archotermopsidae),

*Cryptotermes secundus* (Kalotermitidae), *Coptotermes formosanus* (Rhinotermitidae), and

*Macrotermes natalensis* (Termitidae). Genome completeness was assessed using BUSCO

*v*4.1.2 (Simão *et al.*, 2015) and QUAST *v*5.0.2 (Gurevich *et al.*, 2013). The genome of *M.*

*natalensis* was chosen as the base genome for bait design due to its comparatively higher quality

(for details, see Table S1).

Repetitive elements, retroelements, transposons, and small RNAs were masked from

genome assemblies using RepeatMasker *v*4.1.1 (Smit *et al.*, 2015) with the command line "-

species eukaryota -div 50". Assemblies were converted in the 2-bit format using the

faToTwoBit tool of the BLAT suite of programs (Kent, 2002). We simulated 100 bp error-free

paired-end sequencing reads from the three genome assemblies other than that of *M. natalensis*

using art_illumina Q *v*2.5.8 (Huang *et al.*, 2012) with the command line "--fcov 2 --mflen 200

--sdev 150". In order to identify orthologous loci representing putative UCEs, the reads

simulated from the three termite genome assemblies were mapped independently on the

genome assembly of *M. natalensis* with a 0.05 substitution rate onto the base genome using the

permissive raw-read aligner Stampy *v*1.0.32 (Lunter & Goodson, 2011). The three alignment

maps were handled with SAMtools *v*1.9 (Li *et al.*, 2009) and converted into BED files with

bedtools *v*2.29.2 (Quinlan & Hall, 2010). In each BED file, putative conserved regions

overlapping by at least one nucleotide were merged using bedtools. Conserved sequences

shorter than 80 bp or containing over 25% of masked nucleotides were discarded using the

phyluce program phyluce_probe_strip_masked_loci_from_set. The putative orthologous loci

found across the four termite genomes were combined into a database using

phyluce_probe_get_multi_merge_table (Supplementary Data 1). A total of 175,535 loci shared

by the four termite genomes were identified and extracted using

phyluce_probe_query_multi_merge_table and

phyluce_probe_get_genome_sequences_from_bed, respectively. Extracted UCE sequences

shorter than 180 bp were buffered to 180 bp by including 5' and 3' flanking regions in equal

amounts with phyluce_probe_get_genome_sequences_from_bed (Supplementary Data 2).

A preliminary set of 120 bp baits was designed from the base genome of *M. natalensis*

using phyluce_probe_get_tiled_probes. Baits targeted a region of 180 bp and overlapped in its

center by 60 bp (at 2X tiling density). UCEs with ambiguous base calls and GC-content above

70% or below 30% were discarded from the bait set. Duplicates, defined as sequences having

50% identity over half of their length, were also removed from the bait set using LASTZ (Harris,

2007) implemented in the programs phyluce_probe_easy_lastz and

phyluce_probe_remove_duplicate_hits_from_probes_using_lastz. In order to further identify

and remove non-specific baits, we aligned the bait set (Supplementary Data 3) to the four

genomes with phyluce_probe_run_multiple_lastzs_sqlite using a minimum identity threshold

of 80% and minimum coverage of 83%. Sequences shorter than 180 bp were buffered to 180

bp by including 5' and 3' flanking regions in equal amounts and extracted from the alignments

using phyluce_probe_slice_sequence_from_genomes. The loci shared by the four termite

genomes were identified using phyluce_probe_get_multi_fasta_table and

phyluce_probe_query_multi_fasta_table (Supplementary Data 4). The final UCE bait set was

designed with phyluce_probe_get_tiled_probe_from_multiple_inputs, and duplicates were

removed using LASTZ as described above (397,910 baits targeting 50,616 loci; Supplementary

Data 5). This final set of loci was tentatively annotated using the GFF file (NCBI Annotation

Release 100) from the *Z. nevadensis* genome assembly (GCF_000696155).

*Genome assembling and mining of phylogenetic markers*

Adapters and low-quality bases were trimmed from raw reads using fastp *v*0.20.1 (Chen *et al.*,

2018), resulting in a total of 4.55 to 448.64 millions of paired-end reads per sample (for details,

see Table S1). Trimmed reads were assembled using metaSPAdes *v*3.13 (Nurk *et al.*, 2017).

The quality and completeness of assemblies were assessed with QUAST and BUSCO (Table

S1). Mitochondrial genome scaffolds were identified in metaSPAdes assemblies and annotated

using MitoFinder *v*1.4 (Allio *et al.*, 2020). Nuclear ribosomal RNA genes (5S, 5.8S, 18S, and

28S) were extracted from metaSPAdes assemblies using barrnap *v*0.9

(https://github.com/tseemann/barrnap). UCE loci were extracted from metaSPAdes assemblies

using the final set of termite baits we designed and the PHYLUCE suite of programs with

parameter values set as recommended in the tutorial and previously published studies (Faircloth

*et al.*, 2015; Faircloth, 2017; Quattrini *et al.*, 2018). Briefly, baits were aligned to the

metaSPAdes assemblies at a minimum similarity threshold of 50% with

phyluce_probe_run_multiple_lastzs_sqlite. Sequences of the metaSPAdes assemblies

matching baits were extracted with the flanking 200 bp situated at both the 5' and 3' ends using

phyluce_probe_slice_sequence_from_genomes. Extracted sequences were mapped back to the

baits using phyluce_assembly_match_contigs_to_probes with a minimum identity of 80% over

67% of bait length to remove duplicates and sequences matching multiple UCE loci

(Supplementary Data 6; Contribution #1 to the Termite UCE Database available at:

https://github.com/oist/TER-UCE-DB/). The average coverage of UCE loci per sample was

obtained using the mapping workflow of PHYLUCE *v*1.7.1.

*Sequence alignment*

The 13 mitochondrial protein-coding genes, two mitochondrial rRNA genes, 22 mitochondrial

tRNA genes, four nuclear rRNA genes, and UCEs were aligned using MAFFT *v*7.305 (Katoh

& Standley, 2013). For mitochondrial protein-coding genes, we translated DNA sequences into

the corresponding amino acid sequences using the transeq function from EMBOSS *v*6.6.0 (Rice

*et al.*, 2000) and aligned protein sequences with MAFFT. Protein alignments were back-

translated into codon alignments using PAL2NAL *v*14 (Suyama *et al.*, 2006). The other four

types of genes, the mitochondrial rRNA and tRNA genes, nuclear rRNA genes, and UCEs,

were aligned as DNA sequences. UCE loci were aligned using MAFFT implemented in

phyluce_align_seqcap_align, and internal trimming was performed under default parameters

with Gblocks (Castresana, 2000; Talavera & Castresana, 2007) implemented in

phyluce_align_get_gblocks_trimmed_alignments_from_untrimmed. Loci absent in more than

25% of taxa were filtered out with phyluce_align_get_only_loci_with_min_taxa. The final

UCE supermatrix was exported using phyluce_align_format_nexus_files_for_raxml

(Supplementary Data 7: alignments; Supplementary Data 8: corresponding reduced bait set).

Mitochondrial and nuclear gene alignments were concatenated using FASconCAT-G_v1.04.pl

(Kück & Longo, 2014).

*Phylogenetic analyses*

We ran one separate phylogenetic analysis for the mitochondrial genome alignment, the nuclear

rRNA alignment, and the UCE alignment. We also ran one phylogenetic analysis for the

combined UCE and mitochondrial genome alignments. The mitochondrial genome alignment

was separated into five distinct partitions: combined rRNAs, combined tRNAs, and combined

first, second, and third codon positions of protein-coding genes. Nuclear rRNA gene and UCE

alignments were given a single partition each. Phylogenetic trees were reconstructed in a

maximum likelihood (ML) framework using IQ-TREE *v*1.6.12 with 1,000 ultrafast bootstrap

replicates (UFB) to assess branch supports (Nguyen *et al.*, 2015; Chernomor *et al.*, 2016; Hoang

*et al.*, 2018). The best-fit nucleotide substitution model was selected for each partition with the

Bayesian Information Criterion using ModelFinder implemented in IQ-TREE

(Kalyaanamoorthy *et al.*, 2017). We calculated a global bootstrap support (GBS) value for each

tree by averaging bootstrap values of all nodes. To assess concordance among UCEs, we carried

out a multi-gene coalescence analysis with ASTRAL-III *v*5.7.7 (Zhang *et al.*, 2018) using

individual gene trees obtained with IQ-TREE as described above. We allowed polytomies to

reduce gene tree biases. Branch supports calculated with ASTRAL represent local posterior

probabilities (LPP), which are based on gene tree quartet frequencies (Sayyari & Mirarab, 2016).

Topological conflicts between individual gene trees and the ASTRAL species tree were

assessed with PhyParts (Smith *et al.*, 2015) and visualized with PhyPartsPieCharts

(https://github.com/mossmatters/phyloscripts/tree/master/phypartspiecharts).

## 3. Results

In silico *data mining*

The mitochondrial genomes were retrieved from all 42 termite metaSPAdes assemblies. We

also retrieved the four nuclear rRNA genes from 84% of the samples (see Table S1).

Our termite UCE bait set targeted a total of 50,616 loci distributed across 1,094 scaffolds

of the *Z. nevadensis* genome assembly (GCF_000696155). Of these 50,616 loci, 6,325 (12.5 %)

were found in the non-coding regions of 787 scaffolds (Supplementary Data 9). The remaining

44,291 loci were found in the coding regions (33,932 in exons) of 8,636 annotated genes

distributed across 886 scaffolds. Of the 8,636 genes, 6,538 (75.7 %) contained more than one

ultraconserved loci.

From the 50,616 targeted loci, we extracted between 3,426 and 42,860 non-duplicated

UCE loci from 42 termite metaSPAdes assemblies (Table S1). The number of non-duplicated

UCE loci extracted from the assemblies of *Cryptocercus* roaches varied between 13,480 and

16,331. The average coverage of UCE loci per sample was between 8.38 to 134.71x (Table S1).

The final supermatrix, complete at 75% and containing loci present in at least 33 of the 45 taxa,

was composed of 5,934 loci spanning over 1,677,394 nucleotide positions, 591,343 of which

were parsimony-informative. The 45 taxa were represented by 939 to 5,928 loci.

*Phylogenetic reconstructions*

Many deep and shallow relationships within termites were poorly resolved by the nuclear rRNA

phylogenetic tree (GBS = 72.12) (Figure S1). Because of its poor performance, we excluded

the rRNA alignment from the analysis performed on combined marker classes. The

phylogenetic reconstruction based on mitochondrial genomes resolved most relationships (GBS

= 87), except for several nodes within the Serritermitidae, the Rhinotermitidae, and the

Termitinae (Figure S2), as previously reported (Bourguignon *et al.*, 2015). The phylogenetic

analysis performed exclusively on UCEs provided the most robust phylogenetic tree among the

analyses performed on separate marker classes (Figure S3; GBS = 98.59). Combining UCEs

and mitochondrial genomes marginally improved the resolution of the phylogenetic

reconstruction (Figure 1; Figure S4; GBS = 99.02). The combined phylogenetic reconstruction

resolved all nodes with high supports, except for the position of the rhinotermitid *Termitogeton*

*planus* (UFB = 52 and 65, respectively). The phylogenetic analysis with ASTRAL revealed

minimal discordance among the 5,934 UCE markers (Figure S5; final normalized quartet score

of 0.89) (LPP = 1), except for five of the 42 nodes that presented conflicts among UCE markers.

Within the Rhinotermitidae, the nodes corresponding to the split of *T. planus* and

*Prorhinotermes simplex* displayed moderate concordance among UCE markers (LPP of 0.89

and 0.83, respectively). Within the Termitidae, the nodes corresponding to the split of

*Neocapritermes utiariti*, *Pericapritermes* sp. 4, and *Nitiditermes + Cavitermes* showed

moderate to high levels of discordance (LPP of 0.66, 0.98, and 0.39, respectively). PhyParts

analyses on a subset of 1,000 gene trees revealed some levels of topological discordances

(Figure S6). Nodes with discordance were mostly dominated by a plethora of topologies rather

than by a single alternative and uninformative gene trees.

**4. Discussion**

We reconstructed phylogenetic trees for 42 species of termites and three species of

*Cryptocercus* using three classes of markers: nuclear rRNA genes, mitochondrial genomes, and

UCEs. The performance of the three types of phylogenetic markers decreased along the

sequence: UCEs, mitochondrial genomes, and nuclear rRNA genes. The phylogenetic tree

inferred from the latter class of markers, the nuclear rRNA genes, was poorly resolved and did

not recover well-established relationships, such as the sister position of *Mastotermes* in respect

to all other termites. The phylogenetic tree inferred from mitochondrial genomes was robust

but failed to retrieve Sphaerotermitinae as sister to Macrotermitinae, as previously reported

(Bourguignon *et al.*, 2015; Bucek *et al.*, 2019). The best phylogenetic tree was that

reconstructed using the 75%-occupancy matrix comprised of 5,934 UCE loci (Figure 1). This

phylogenetic tree was almost fully resolved and largely congruent with the phylogenetic trees

inferred from transcriptomic data (Bucek *et al.*, 2019). Therefore, our results indicate that the

termite UCE bait set we designed performs very well when reconstructing phylogenetic

relationships among termite species. The addition of mitochondrial genome data, which, as

UCEs, can be recovered from shotgun genome assemblies, slightly improved the global

bootstrap support of the termite phylogenetic tree.

The analysis with ASTRAL revealed a few cases of discordance among UCE markers

for lineages of Rhinotermitidae and Termitidae whose phylogenetic position was also

unresolved with transcriptomic data (Bucek *et al.*, 2019). We used 5,934 UCE loci, a large

number of markers that inevitably led to topological discordances between individual UCE

trees and the species tree. These discordances are possibly caused by the lack of phylogenetic

signal present in a single UCE marker and by population-level processes, such as incomplete

lineage sorting and introgression, which frequently occurs during the emergence of new

lineages (Degnan & Rosenberg, 2009; Blom *et al.*, 2017; Parins-Fukuchi *et al.*, 2021). While

recent studies indicated that the phylogenetic resolution can be improved by removing loci

localized in the exons of a single gene (Hedin *et al.*, 2019; Van Dam *et al.*, 2021), our unfiltered

analysis performed as well as analyses performed on transcriptome-based data. The actual

relationships among termite lineages with unresolved positions remain unclear, possibly

reflecting intricate evolutionary history that cannot be satisfactorily resolved by molecular

phylogenetic techniques.

We ran our analyses on samples for which we generated low coverage genome

assemblies. The final bait set targeting a total of 50,616 orthologous loci was obtained from

four termite genomes, belonging to four families. This high number of UCE loci is comparable

to that found in other groups of arthropods (e.g., Buenaventura *et al.*, 2021). Overall, 67% of

loci in the termite bait set were found in exonic regions, further indicating that the UCEs of

arthropods are mostly found in exons (Hedin *et al.*, 2019; Van Dam *et al.*, 2021). We retrieved

numerous UCE sequences for all samples, including many that produced highly fragmented

assemblies with low BUSCO scores (for details, see Table S1). All samples were accurately

placed on the phylogenetic tree reconstructed with the 5,934 loci present in the 75%-occupancy

supermatrix. Therefore, our UCE bait set has the potential to be used for mining

phylogenetically informative genetic data from assemblies obtained from shotgun sequencing

experiments. We established a centralized termite UCE database (https://github.com/oist/TER-

UCE-DB/), which we plan to use to reference all UCE data extracted with the presently

designed bait set, thereby ensuring the long-term re-usability of the available data.

While producing low-coverage genomes is more costly than targeting UCEs through

synthetized baits, low-coverage genome data can be used to investigate a broad range of

questions in addition to phylogenetic reconstruction. Used in combination with non-destructive

DNA extraction protocols, our UCE baits could also be used to obtain sequence data from

material that cannot be damaged, such as specimens from type series. This approach was

successfully applied to centuries-old museum specimens of Opiliones, carpenter bees, and

weevils (Blaimer *et al.*, 2016; Van Dam *et al.*, 2017; Derkarabetian *et al.*, 2019). We recently

obtained the full mitogenome of a Syntype of the termite *Archotermopsis wroughtoni* collected

at the end of the 19[th] century using shotgun sequencing data (Wang *et al.*, 2021). Termite UCEs

could be extracted using the same procedures. Termite taxonomy, which is led by a shrinking

pool of experts and is largely based on soldier and worker gut morphology, could benefit from

the use of the many UCE markers designed in this study (Eggleton, 1999; Korb *et al.*, 2019).

UCE baiting from whole-genome shotgun sequencing is the perfect tool to carry out a global

taxonomic revision of termites.

**References**

Allio, R., Schomaker-Bastos, A., Romiguier, J., Prosdocimi, F., Nabholz, B., Delsuc, F. 2020.

MitoFinder: efficient automated large-scale extraction of mitogenomic data in target

enrichment phylogenomics. *Molecular Ecology Resources* **20**: 892–905.

Bejerano, G., Pheasant, M., Makunin, I., Stephen, S., Kent, W.J., Mattick, J.S., Haussler, D.

2004. Ultraconserved elements in the human genome. *Science* **304**: 1321–1325.

Blaimer, B.B., Lloyd, M.W., Guillory, W.X., Brady, S.G. 2016. Sequence capture and

phylogenetic utility of genomic ultraconserved elements obtained from pinned insect

specimens. *PLoS ONE* **11**: e0161531.

Blom, M.P.K., Bragg, J.G., Potter, S., Moritz, C. 2017. Accounting for uncertainty in gene tree

estimation: summary-coalescent species tree inference in a challenging radiation of

Australian lizards. *Systematic Biology* **66**: 352–366.

Bourguignon, T., Lo, N., Cameron, S.L., Šobotník, J., Hayashi, Y., Shigenobu, S., Watanabe,

D., Roisin, Y., Miura, T., Evans, T.A. 2015. The evolutionary history of termites as

inferred from 66 mitochondrial genomes. *Molecular Biology and Evolution* **32**: 406–421.

Bourguignon, T., Lo, N., Šobotník, J., Ho, S.Y.W., Iqbal, N., Coissac, É., Lee, M., Jendryka,

M.M., Sillam-Dussès, D., Křížková, B., Roisin, Y., Evans, T.A. 2017. Mitochondrial

phylogenomics resolves the global spread of higher termites, ecosystem engineers of the

tropics. *Molecular Biology and Evolution* **34**: 589–597.

Bucek, A., Šobotník, J., He, S., Shi, M., McMahon, D.P., Holmes, E.C., Roisin, Y., Lo, N.,

Bourguignon, T. 2019. Evolution of termite symbiosis informed by transcriptome-based

phylogenies. *Current Biology* **29**: 3728-3734.e4.

Buenaventura, E., Lloyd, M.W., Perilla López, J.M., González, V.L., Thomas-Cabianca, A.,

Dikow, T. 2021. Protein-encoding ultraconserved elements provide a new phylogenomic

perspective of Oestroidea flies (Diptera: Calyptratae). *Systematic Entomology* **46**: 5–27.

Cameron, S.L. 2014. Insect mitochondrial genomics: implications for evolution and phylogeny.

*Annual Review of Entomology* **59**: 95–117.

Cameron, S.L., Lo, N., Bourguignon, T., Svenson, G.J., Evans, T.A. 2012. A mitochondrial

genome phylogeny of termites (Blattodea: Termitoidae): robust support for interfamilial

relationships and molecular synapomorphies define major clades. *Molecular*

*Phylogenetics and Evolution* **65**: 163–173.

Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in

phylogenetic analysis. *Molecular Biology and Evolution* **17**: 540–552.

Chen, S., Zhou, Y., Chen, Y., Gu, J. 2018. Fastp: an ultra-fast all-in-one FASTQ preprocessor.

*Bioinformatics* **34**: i884–i890.

Chernomor, O., Von Haeseler, A., Minh, B.Q. 2016. Terrace aware data structure for

phylogenomic inference from supermatrices. *Systematic Biology* **65**: 997–1008.

Degnan, J.H., Rosenberg, N.A. 2009. Gene tree discordance, phylogenetic inference and the

multispecies coalescent. *Trends in Ecology and Evolution* **24**: 332–340.

Derkarabetian, S., Benavides, L.R., Giribet, G. 2019. Sequence capture phylogenomics of

historical ethanol-preserved museum specimens: unlocking the rest of the vault. *Molecular*

*Ecology Resources* **19**: 1531–1544.

Eggleton, P. 1999. Termite species description rates and the state of termite taxonomy. *Insectes*

*Sociaux* **46**: 1–5.

Engel, M.S., Grimaldi, D.A., Krishna, K. 2009. Termites (Isoptera): their phylogeny,
classification, and rise to ecological dominance. *American Museum Novitates* **3650**: 1–27.

Engel, M.S., Grimaldi, D.A., Nascimbene, P.C., Singh, H. 2011. The termites of Early Eocene
Cambay amber, with the earliest record of the Termitidae (Isoptera). *ZooKeys* **148**: 105–
123.

Faircloth, B.C. 2017. Identifying conserved genomic elements and designing universal bait sets
to enrich them. *Methods in Ecology and Evolution* **8**: 1103–1112.

Faircloth, B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic
loci. *Bioinformatics* **32**: 786–788.

Faircloth, B.C., Branstetter, M.G., White, N.D., Brady, S.G. 2015. Target enrichment of
ultraconserved elements from arthropods provides a genomic perspective on relationships
among Hymenoptera. *Molecular Ecology Resources* **15**: 489–501.

Faircloth, B.C., McCormack, J.E., Crawford, N.G., Harvey, M.G., Brumfield, R.T., Glenn, T.C.
2012. Ultraconserved elements anchor thousands of genetic markers spanning multiple
evolutionary timescales. *Systematic Biology* **61**: 717–726.

Grimaldi, D.A., Engel, M.S. 2005. *Evolution of the Insects*. Cambridge University Press,
Cambridge.

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G. 2013. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**: 1072–1075.

Harris, R.S. 2007. Improved pairwise alignment of genomic DNA. The Pennsylvania State
University.

Hedin, M., Derkarabetian, S., Alfaro, A., Ramírez, M.J., Bond, J.E. 2019. Phylogenomic

analysis and revised classification of atypoid mygalomorph spiders (Araneae,

Mygalomorphae), with notes on arachnid ultraconserved element loci. *PeerJ* **7**: e6864.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S. 2018. UFBoot2:

improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**:

518–522.

Huang, W., Li, L., Myers, J.R., Marth, G.T. 2012. ART: a next-generation sequencing read

simulator. *Bioinformatics* **28**: 593–594.

Inward, D.J.G., Vogler, A.P., Eggleton, P. 2007. A comprehensive phylogenetic analysis of

termites (Isoptera) illuminates key aspects of their evolutionary biology. *Molecular*

*Phylogenetics and Evolution* **44**: 953–967.

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., Von Haeseler, A., Jermiin, L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*

**14**: 587–589.

Katoh, K., Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–

780.

Kent, W.J. 2002. BLAT—The BLAST-like alignment tool. *Genome Research* **12**: 656–664.

Korb, J., Kasseney, B.D., Cakpo, Y.T., Casalla Daza, R.H., Gbenyedji, J.N.K.B., Ilboudo, M.E.,

Josens, G., Koné, N.A., Meusemann, K., Ndiaye, A.B., Okweche, S.I., Poulsen, M., Roisin,

Y., Sankara, F. 2019. Termite taxonomy, challenges and prospects: West Africa, a case

example. *Insects* **10**: 32.

Kück, P., Longo, G.C. 2014. FASconCAT-G: extensive functions for multiple sequence

alignment preparations concerning phylogenetic studies. *Frontiers in Zoology* **11**: 81.

Legendre, F., Whiting, M.F., Bordereau, C., Cancello, E.M., Evans, T. a., Grandcolas, P. 2008.

The phylogeny of termites (Dictyoptera: Isoptera) based on mitochondrial and nuclear

markers: Implications for the evolution of the worker and pseudergate castes, and foraging

behaviors. *Molecular Phylogenetics and Evolution* **48**: 615–627.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:

2078–2079.

Lo, N., Kitade, O., Miura, T., Constantino, R., Matsumoto, T. 2004. Molecular phylogeny of

the Rhinotermitidae. *Insectes Sociaux* **51**: 365–371.

Lunter, G., Goodson, M. 2011. Stampy: a statistical algorithm for sensitive and fast mapping

of Illumina sequence reads. *Genome Research* **21**: 936–939.

Nguyen, L.T., Schmidt, H.A., Von Haeseler, A., Minh, B.Q. 2015. IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*

*Biology and Evolution* **32**: 268–274.

Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P.A. 2017. metaSPAdes: a new versatile

metagenomic assembler. *Genome Research* **27**: 824–834.

Parins-Fukuchi, C., Stull, G.W., Smith, S.A. 2021. Phylogenomic conflict coincides with rapid

morphological innovation. *Proceedings of the National Academy of Sciences of the United*

*States of America* **118**: e2023058118.

Quattrini, A.M., Faircloth, B.C., Dueñas, L.F., Bridge, T.C.L., Brugler, M.R., Calixto-Botía,

I.F., DeLeo, D.M., Forêt, S., Herrera, S., Lee, S.M.Y., Miller, D.J., Prada, C., Rádis-

Baptista, G., Ramírez-Portilla, C., Sánchez, J.A., Rodríguez, E., McFadden, C.S. 2018.

Universal target-enrichment baits for anthozoan (Cnidaria) phylogenomics: new

approaches to long-standing problems. *Molecular Ecology Resources* **18**: 281–295.

Quinlan, A.R., Hall, I.M. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**: 841–842.

Rice, P., Longden, L., Bleasby, A. 2000. EMBOSS: the European Molecular Biology Open

Software Suite. *Trends in Genetics* **16**: 276–277.

Ryu, T., Seridi, L., Ravasi, T. 2012. The evolution of ultraconserved elements with different

phylogenetic origins. *BMC Evolutionary Biology* **12**: 236.

Sayyari, E., Mirarab, S. 2016. Fast coalescent-based computation of local branch support from

quartet frequencies. *Molecular Biology and Evolution* **33**: 1654–1668.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. V., Zdobnov, E.M. 2015.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**: 3210–3212.

Smit, A.F.A., Hubley, R., Green, P. 2015. RepeatMasker Open-4.0 2013-2015.

Smith, B.T., Harvey, M.G., Faircloth, B.C., Glenn, T.C., Brumfield, R.T. 2014. Target capture

and massively parallel sequencing of ultraconserved elements for comparative studies at

shallow evolutionary time scales. *Systematic Biology* **63**: 83–95.

Smith, S.A., Moore, M.J., Brown, J.W., Yang, Y. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals and

plants. *BMC Evolutionary Biology* **15**: 150. BMC Evolutionary Biology.

Suyama, M., Torrents, D., Bork, P. 2006. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Research* **34**: W609–

W612.

Talavera, G., Castresana, J. 2007. Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. *Systematic Biology* **56**:

564–577.

Thorne, B.L., Grimaldi, D.A., Krishna, K. 2000. Early fossil history of the termites. In:

*Termites: Evolution, Sociality, Symbioses, Ecology* (T. Abe, D. E. Bignell, & M. Higashi,

eds), pp. 77–93. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Van Dam, M.H., Henderson, J.B., Esposito, L., Trautwein, M. 2021. Genomic characterization

and curation of UCEs improves species tree reconstruction. *Systematic Biology* **70**: 307–

321.

Van Dam, M.H., Lam, A.W., Sagata, K., Gewa, B., Laufa, R., Balke, M., Faircloth, B.C., Riedel,

A. 2017. Ultraconserved elements (UCEs) resolve the phylogeny of Australasian smurf-

weevils. *PLoS ONE* **12**: e0188044.

Wang, M., Hellemans, S., Šobotník, J., Arora, J., Buček, A., Sillam-Dussès, D., Clitheroe, C.,

Lu, T., Lo, N., Engel, M.S., Roisin, Y., Evans, T.A., Bourguignon, T. 2021. Historical

biogeography of early diverging termite lineages (Isoptera: Teletisoptera). *bioRxiv*

2021.12.02.471008.

White, N.D., Braun, M.J. 2019. Extracting phylogenetic signal from phylogenomic data:

higher-level relationships of the nightbirds (Strisores). *Molecular Phylogenetics and*

*Evolution* **141**: 106611.

Whitfield, J.B., Lockhart, P.J. 2007. Deciphering ancient rapid radiations. *Trends in Ecology*

*and Evolution* **22**: 258–265.

Zhang, C., Rabiee, M., Sayyari, E., Mirarab, S. 2018. ASTRAL-III: polynomial time species

tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**: 153.

Zhang, Y.M., Williams, J.L., Lucky, A. 2019. Understanding UCEs: a comprehensive primer

on using ultraconserved elements for arthropod phylogenomics. *Insect Systematics and*

*Diversity* **3**: 3.

**Figure captions**

**Figure 1**: (**A**) Maximum likelihood phylogenetic tree of termites reconstructed with IQ-TREE

using 5,934 UCE loci and complete mitochondrial genomes. Only UCE loci present in more

than 75% of species were used. Support values are indicated for non-fully resolved nodes:

ultrafast bootstrap (UFB; summarized from the phylogenetic trees reconstructed using UCE

only and UCE + mitochondrial DNA displayed in Figures S3 and S4, respectively) and

ASTRAL-III local posterior probabilities (LPP; phylogenetic tree displayed in Figure S5)

values. (**B**) Family-level summary topology of termites supported by both UCEs (this study)

and transcriptomic data (Bucek *et al*., 2019), with the indication of alternative topologies

inferred from mitochondrial genome data alone (Bourguignon *et al*., 2015, 2017). Unsupported

splits were summarized as polytomies (branches in red).

# A

*Cryptocercus punctulatus*
*Cryptocercus hirtus*
*Cryptocercus meridianus*
*Mastotermes darwiniensis* — Mastotermitidae
*Porotermes quadricollis* — Stolotermitidae
*Hodotermopsis sjostedti* — Hodotermopsidae
*Zootermopsis nevadensis* — Archotermopsidae
*Kalotermes flavicollis*
*Glyptotermes pubescens*
*Neotermes castaneus* — Kalotermitidae
*Cryptotermes secundus*
*Cryptotermes cavifrons*
*Stylotermes halumicus* — Stylotermitidae
*Glossotermes occulatus* — Serritermitidae
*Rhinotermes hispidus* — Rhinotermitinae
*Dolichorhinotermes longilabius*
*Termitogeton planus* — Termitogetoninae
*Prorhinotermes simplex* — Prorhinotermitinae
*Reticulitermes flavipes* — *Reticulitermes* + *Heterotermes* + *Coptotermes* group
*Heterotermes tenuis*
*Coptotermes formosanus*
*Coptotermes testaceus*
*Sphaerotermes sphaerothorax* — Sphaerotermitinae
*Macrotermes natalensis* — Macrotermitinae
*Odontotermes formosanus*
*Labritermes buttelreepeni* — Foraminitermitinae
*Foraminitermes rhinoceros*
*Allognathotermes hypogeus* — Apicotermitinae
*Anoplotermes banksi*
*Astalotermes* sp. D
*Microcerotermes* sp. — Microcerotermitinae
*Embiratermes neotenicus* — Syntermitinae
*Labiotermes labralis*
*Cornitermes pugnax*
*Neocapritermes utiariti* — Termitinae
*Leptomyxotermes doriae* — Nasutitermitinae
*Nasutitermes corniger*
*Constrictotermes cyphergaster*
*Subulitermes* sp. 1-1
*Promirotermes redundans* — Termitinae
*Cavitermes tuberosus*
*Nitiditermes* sp. — Cubitermitinae
*Pericapritermes* sp. 4 — Termitinae
*Dentispicotermes* n. sp. 2
*Amitermes californicus*

(Rhinotermitidae)
Termitidae

● 90≤UFB≤99
● 90>UFB
▲ 1>LPP

⌐ Polyphletic ⌐

0.03

# B

Mastotermitidae
Stolotermitidae
Hodotermopsidae
Archotermopsidae
Kalotermitidae
Stylotermitidae
Serritermitidae — Rhinotermitinae
Rhinotermitinae — Prorhinotermitinae
Termitogetoninae — Termitogetoninae
Prorhinotermitinae — Serritermitinae
*Reticulitermes*+*Heterotermes* +*Coptotermes*-group
Sphaerotermitinae — Macrotermitinae
Macrotermitinae — Foraminitermitinae
Foraminitermitinae — Sphaerotermitinae
Apicotermitinae
Microcerotermitinae
Syntermitinae
Termitinae
Nasutitermitinae
Termitinae
Cubitermitinae
Termitinae