Acute oral toxicity of zinc phosphide: an assessment for wild house mice in Australia

Lyn A. Hinds, Steve Henry, Nikki Van de Weyer, Freya Robinson, Wendy A. Ruscoe and

Peter R. Brown.

CSIRO Health and Biosecurity, GPO Box 1700, Canberra, Australia, 2601

Corresponding author: Lyn.Hinds@csiro.au

Lyn A. Hinds: 0000-0002-4125-2357

Steve Henry: 0000-0002-4372-8769

Nikki Van de Weyer: 0000-0002-8659-4369

Freya Robinson: 0000-0002-8598-2466

Wendy A. Ruscoe: 0000-0002-7763-0356

Peter R. Brown: 0000-0001-5894-8329

Abstract

BACKGROUND The efficacy of zinc phosphide (ZnP) for broadacre control of wild house

mice in Australia is being reported by growers as increasingly variable. Have mice become

less sensitive over time or are they taking a sub-lethal dose and developing aversion? In this

laboratory study the sensitivity of groups of wild caught and an outbred laboratory strain of

mice was assessed using oral gavage of a range of ZnP concentrations. The willingness of

mice to consume ZnP-coated grains was then determined.

RESULTS Each mouse group had very similar LD50 values (72 to 79 mg ZnP per kg body

weight) which are significantly higher than previously reported. Time to death post-gavage

ranged between 2.5 to 48 h. ZnP coated grains (50 g ZnP per kg grain) presented in the

absence of alternative food were consumed and 94 percent of wild mice died. Mice provided

with alternative food and ZnP coated wheat grains (either 25 or 50 g ZnP per kg grain)

consumed toxic and non-toxic grains, and mortality was lower (33 to 55 percent). If a

sublethal amount of ZnP coated grain was consumed, aversion occurred mostly in the

presence of alternative food.

CONCLUSIONS The sensitivity of wild house mice to ZnP in Australia is significantly lower

than previously assumed. Under laboratory conditions ZnP coated grains coated with a new

higher dose (50 g ZnP per kg grain) were readily consumed. Consumption of toxic grain

occurred when alternative food was available but was decreased. It is important to assess

the efficacy of the higher ZnP dose per grain under natural field conditions, especially when

background food is low.

Keywords: *Mus musculus domesticus*, acute rodenticide; LD_{50}, LD_{90}, outbred laboratory

mice, efficacy, aversion

1   INTRODUCTION

Zinc phosphide (Zn_3P_2, hereafter ZnP) has been used for the control of several commensal

and field rodents for several decades.[1] It has the advantage of being an acute rodenticide

with low risk of secondary poisoning and an absence of long-term residues,[2,3] but it is not

rodent specific. Many other mammals (ungulates, carnivores, lagomorphs) and birds

(particularly land and waterfowl and granivorous passerines) are sensitive to ZnP and fish

are highly susceptible.[4-6] The mode of action of ZnP is primarily due to the release of

phosphine gas and its subsequent uptake into the blood stream and major organs. When

food enters the stomach it generally triggers an increase in the release of gastric acid.[7,8] After

consumption of ZnP-food baits the hydrolysis of ZnP results in a rapid release of phosphine

which causes acute toxicity in various organs including the lungs, liver and heart. Animals

show adverse changes in respiratory patterns, hypoxia and loss of control of bodily

movements (ataxia). In the absence of food in the stomach undegraded ZnP can be

absorbed systemically, and animals may survive beyond 48 h but succumb to liver and/or

renal failure 5 - 14 days later. ZnP is thought to disrupt cytochrome C oxidase, leading to

formation of highly reactive oxygen compounds, which result in tissue injury, particularly

those with a high oxygen demand (brain, lungs, liver, and kidney).[8]

The ZnP oral toxicity (LD_{50} - lethal dose that kills 50% animals) for mammals ranges

from 5.6 mg/kg for nutria (*Myocaster coypus*)[4] and 9.6 mg/kg for brushtail possums

(*Trichosurus vulpecula*)[9] to 60-70 mg/kg for sheep, *Ovis aries.*[5] For several rodent species

the LD_{50} is in the range of 20-40 mg ZnP/kg body weight.[4,6] Specifically, for wild house mice,

Mus musculus, the LD_{50} is reported as 32.7 mg/kg body weight, whilst for Swiss Webster

albino mice, an outbred strain of laboratory mice, the value is 54 mg/kg body weight,[10]

though Bell[11] reported a value of 25 mg/kg for white mice.

The wild house mouse (*Mus musculus domesticus*) is a major pest in grain

production areas of Australia.[12,13] Mouse outbreaks occur irregularly and cause significant

economic losses and social impacts.[12-18] ZnP was first registered in 2000 as an in-crop acute

rodenticide for mouse management[19,20] and initially was applied to grain crops at 1 kg/ha

(registered rate of 25 g ZnP/kg wheat; ~3 toxic grains/m^2; each grain is coated with

approximately 1 mg - the equivalent of an LD_{90} dose for wild mice) during periods of

increasing or high mouse numbers. Its efficacy was highest when applied when food was

scarce, especially prior to crop/cereal flowering.[13,18] After aerial application there was a 51%

reduction in adjusted trap success for mice, while after perimeter baiting less than 25%

reduction was observed.[21] In some situations where outbreaks are frequent, it has become

general practice for farmers to apply ZnP as a preventative control strategy when a crop is

being sown, even if mouse numbers are low and not predicted to increase (Henry S and

Brown PR unpublished observations). Furthermore, several applications of ZnP bait have

been required during the crop growing season leading some farmers to question whether

there has been a change in the effectiveness of ZnP over the last 10-15 years (Henry S and

Brown PR unpublished observations). This may also be influenced by the broadscale

adoption of conservation agriculture practices which provide increased cover and reduced

soil disturbance.[22] These observations have raised questions such as: (1) have mice become

tolerant to the toxin? (2) has presentation of the toxin on cereal grain led to higher levels of

bait aversion? and (3) does the presence of moderate background food resources reduce

uptake of ZnP-coated grains which are spread at approximately 3 grains/m^2?

The effectiveness of ZnP-coated grains has previously been assessed in the

laboratory in the presence of alternative food.[23] In that study, wild mice showed a strong

preference for consumption of cereal grains over lentils, and this influenced their uptake of

toxic ZnP-coated barley grains. The mortality rate for mice presented with a choice of lentils

and toxic ZnP-coated cereal grains was 86%, whereas for mice presented with a choice of

cereal grains and toxic ZnP-coated cereal grains the mortality rate was only 50%. One

possible interpretation here is that the presence or absence of food in the stomach of mice

directly impacts the efficacy of the ZnP. Another possible change in the two decades since

ZnP has been registered for control of mice in crops in Australia is that regular and repeated

bait applications may have led to the selection of mice that are now less sensitive or more

tolerant to ZnP. Potentially, mice in areas where ZnP has been applied on a regular basis

may be more tolerant to ZnP than mice that have never been exposed to the toxin.

The current laboratory study was designed to re-examine the acute toxicity of ZnP in

fed and unfed house mice. Our first objective was to determine the LD$_{50}$ for the wild house

mouse in Australia using oral gavage of known concentrations of ZnP in vegetable oil. To

achieve this, wild mice were trapped from an area where ZnP had been used most years,

and from a location where ZnP had never been used. Their responses were compared with

those of an outbred laboratory strain of mice. Considering the results obtained we then

assessed whether wild caught house mice would consume wheat grains coated with a higher

dose of ZnP than is currently in use.

2   METHODS

2.1   Animals

Wild house mice were trapped using single live capture traps (Longworth, Abingdon, UK).
Traps were set with wheat grains and non-absorbent Dacron in the late afternoon and
checked from dawn the following morning. Trapped mice were weighed, measured, sexed
and assessed for general condition before being transferred to small cages (30.5 x 13.5 x
12.0 cm, length x width x depth) containing wood shavings, nesting material, a cardboard
tube, apple pieces for moisture, a mounted water bottle and food pellets (standard
maintenance diet rat and mouse pellets) (Gordon's Specialty Stockfeeds, NSW, Australia).
Mice were transported to the animal facility at CSIRO, ACT, Australia, in an air-conditioned
vehicle. On arrival, they were transferred to individual, clear, mouse-specific cages (26 x 40 x
20 cm, length x width x depth) which contained bedding and nesting material (wood shavings
and tissue paper), water and food *ad libitum*, cardboard tube shelters and popsicle sticks for
environmental enrichment. The room temperature was maintained at 22 ± 2° C. During the
two-week acclimation period mice were checked daily for alertness, and hydration, and
weighed twice per week.

2.2   Experimental design

Experiment 1: Determination of the LD_{50} and LD_{90} of wild and laboratory house mice

Wild house mice were trapped from a farm location on Yorke Peninsula, South Australia,
where ZnP baiting occurred regularly (at least once per year for 9 of the last 12 years). This

group is hereafter referred to as wild mice (Exposed). The second group of wild house mice

was trapped from around a horse stable in Murrumbateman, NSW, where ZnP baiting has

never occurred. This group is hereafter referred to as wild mice (Naïve). The third group of

mice, an outbred strain of laboratory mice, was obtained from the Animal Resources Centre,

Western Australia. Within each experimental group equal numbers of males and females

were used when available.

We used oral gavage of a range of concentrations of ZnP in vegetable oil using the

broad principles of the OECD Up Down protocol (UDP).[24,25] Starting doses were based on

the reported LD_{50} values.[10] For the outbred laboratory strain the starting dose was 50.22 mg

ZnP/kg body weight, for the wild caught mice it was 30 mg ZnP/kg body weight. If any dosed

mice survived, a higher dose was administered to new individuals, while if they died a lower

dose was administered to new individuals. This process was repeated until a dose response

curve was established. Doses given ranged between 30 - 149 mg ZnP/g body weight.

Specific doses based on body weight were given to animals lightly anaesthetised with

Isoflurane (4%) in oxygen, using a 20 g curved gavage needle attached to a 1 ml syringe

between 08:00 and 09:00 h each morning. Once the gavage needle was in position the

plunger of the syringe was gently depressed to administer the individual's calculated dose

(90-300 µl) into the stomach. The gavage needle was gently removed, and the animal

maintained in a vertical position for a few seconds until it showed the swallow reflex and

signs of recovery from the anaesthetic. Each mouse was placed in a glass observation tank

(30 x 30 x 30 cm) covered with a wire mesh lid. The oral gavage procedure took less than

20-30 seconds and anaesthetic recovery was rapid and uneventful.

Animals were observed regularly (every 30 min) for 24 h post gavage and more

frequently (every 15 min) if signs of toxicity were observed. Observation of their behaviour

included: general alertness (normal to lethargic) and movement (normal exploratory activity

to abnormal/unbalanced and a reluctance to move); body position (normal to sternal or

lateral recumbency); clarity of eyes (bright, clear and open to dull, half or fully closed);

respiratory pattern (normal, shallow; gasping; slow, rapid). If mice exhibited lateral or sternal

recumbency and had lost their righting reflex, as well as showing high respiration rate and

lethargy, they were humanely killed in accordance with animal ethics requirements for this

project.

Mice allocated to a treatment dose were divided into a fed or unfed group. For the

fed group, food was available *ad libitum*, while for the unfed group, food was removed 12 h

prior to oral gavage with a ZnP dose. Food was returned to unfed individuals at 8 h post

gavage. Dosed mice were checked through to 48 h post gavage as per the UDP[25] and then

humanely killed using isoflurane then cervical dislocation.

Experiment 2: Determination of uptake and efficacy of ZnP-coated grain (LD$_{90}$ dose defined

in Experiment 1)

Wild caught house mice were trapped from around Parkes, NSW, and transported

back to the laboratory. They were acclimatised for 7-10 days in individual cages and

provided with a maintenance diet of laboratory mouse pellets and wheat grains prior to the

experiment, as described above. Three days before the start of the experiment, laboratory

chow was removed, and only standard wheat grains (hereafter referred to as alternative

food) were offered.

We established three experimental groups. Firstly, we compared intake and mortality

of one group of animals provided with a choice of alternative food ("Fed") and ZnP-coated

grains (50 g ZnP/kg wheat grain) (hereafter referred to as F50), with another group in which

only ZnP-coated grains (50 g ZnP/kg grain) ("UnFed"; hereafter UF50) were provided for 4 h

before the addition of alternative food.

Secondly, a third group of animals was provided with a choice of alternative food and

ZnP-coated grains (25 g ZnP/kg grain) (hereafter F25). We did this to confirm results

obtained for animals receiving a choice of ZnP-coated grains (25 g ZnP/kg grain) and

alternative food as described in [23] and to compare this F25 group with the above-mentioned

F50 group. Two sequential independent trials (Trial 1 and Trial 2), each using ten animals

per treatment group, were undertaken over a two-week period. The preparation of batches of

ZnP-coated wheat is described below.

Individual mice were presented with two small dishes, one containing wheat grains

(≥15% body weight, (g)) and one containing ZnP-coated wheat grains (n = 10). This choice

was offered for three consecutive days. Each day, consumption of each food type was

recorded, and the dish positions reversed in individual mouse boxes. Each dish was secured

to the floor of the cage with Blu Tack (Bostick Pty Ltd, Thomastown, Victoria, Australia) to

minimise spillage.

Mice were monitored at 30 min intervals from the time of introduction of the ZnP-

coated wheat grains for 24 h (Days 1 - 3) and then 2-3 times per day between Days 4 - 6.

Mice that were still alive on Day 6 were humanely killed as described above.

On Days 1 - 3, between 08:00 - 09:00 h, food was removed, cages were cleaned, and

animals were weighed and checked for any signs of distress and returned to the cage. Then

at 16:00 h, for the fed groups (F50 and F25), alternative food (≥15% body weight) and their

respective ZnP-coated grains (n = 10) were provided in separate dishes and observations

commenced. At this time, 16:00 h, the unfed group (UF50) was provided with ZnP-coated

grains (n = 10) only and observations commenced. Their alternative food (≥15% body

weight) was added between 20:00 and 22:00 h.

To potentially increase uptake of ZnP-coated grains, in the second trial no alternative

food was provided to any groups until approximately 6 h after provisioning of ZnP-coated

grains on Days 2 and 3.

2.3 Zinc Phosphide

<u>2.3.1 Preparation of concentrations of ZnP in oil and coating of ZnP onto grains</u>

For Experiment 1, the ZnP powder (93% purity; ≥ 19% active phosphor (P) basis, powder;

Lot # STBJ1895; Sigma Aldrich Castle Hill NSW, Australia) was weighed and added to 100%

pure vegetable oil (Crisco, Goodman Fielder, Moorebank, NSW, Australia) to achieve a

known concentration (range: 3.22 - 20 mg/ml) for oral gavage on that day. The powdered

ZnP in oil forms a suspension of ZnP particles and is not a homogenous solution. Therefore,

the prepared suspension was rapidly vortexed before each individual dose was quickly

loaded into the 1 ml syringe for gavaging. Each day, two aliquots of each concentration were

removed, held at 4° C, and subsequently submitted for independent analysis (see below).

For Experiment 2, two batches of ZnP-coated wheat grains were prepared in-house.

following the method used by commercial bait producers who pre-mix ZnP powder and

sterilised grain before adding oil. To achieve the rate of 25 g ZnP/kg grain, 250 g wheat was

weighed, and 7.8 g ZnP powder (corrected for 80% purity) was added to a glass bottle. The

bottle was capped and continuously rotated until all the powder was distributed on the grains

after which 6.7 ml vegetable oil was added. The rotating process was repeated until all the

grains were evenly covered with both powder and oil.

To achieve the rate of 50 g ZnP/kg grain, the above procedure was repeated using

125 g wheat grains, 7.8 g ZnP powder and 3.0 ml vegetable oil.

Coated grains were then spread onto trays on aluminium foil and placed in a drying

oven at 40° C overnight. ZnP-coated grains were then returned to a labelled airtight glass

bottle and stored in a dark cupboard.

ZnP-coated grains (n= 8-10) from each batch were individually analysed to

independently confirm the expected coating rate of approximately 1 and 2 mg ZnP per grain,

respectively.

<u>2.3.2 Analysis of concentrations of ZnP in oil and ZnP on coated grains</u>

Analysis was undertaken by ACS Laboratories (Australia) (37 Stubbs St, Kensington,

Victoria, 3031), a commercial, independent analytical service provider accredited by the

National Association of Testing Authorities (NATA), Australia (NATA Accreditation No

16973). One aliquot of each ZnP concentration in oil was subsampled (in duplicate) and

analysed by microwave digestion followed by inductively coupled plasma optical emission

spectroscopy (ICP-OES). Approximately 200 µl subsamples were weighed directly into

Teflon digestion tubes to which 5 ml of concentrated nitric acid and 1 ml of 30% hydrogen

peroxide were added. Grains coated with ZnP were also individually analysed by microwave

digestion followed by ICP-OES. Each grain was weighed directly into a Teflon digestion tube

to which 5 ml of concentrated nitric acid and 1 ml of 30% hydrogen peroxide were added.

Each digestion tube was then microwaved for 65 minutes using a temperature gradient from

0-200° C in a Milestone Connect Ethos Lean Compact Microwave Digestor. Once cooled the

contents of the digestion tube were diluted to 50 ml in deionised water, followed by a further

10-fold dilution in deionised water for ICP analysis using an Agilent 5110 ICP-OES (Dual

View). The operating conditions for the instrument were: 1.20 kW RF power; plasma

15.0L/min; auxillary 1.50 L/min; AVS7 sample injection system; injection pump rate 15.7

ml/min; detection - intensity, Zn 206.2 nm/c/s.

The recovery of Zinc measured across a range of known concentrations of ZnP in oil

(7.3, 15.5, 21.4 mg/ml) (quality controls prepared by ACS laboratories) was 100.4 ± 5.4%

(n=4), 95.4 ± 7.9% (n=3) and 101.3 ± 3.5% (n=6) (Mean ± SEM) respectively. The

concentrations of ZnP in oil used for oral gavage were confirmed to be within 96.3 ± 3.7%

(Mean ± SEM) (n=16 samples) of that expected. The independent analysis of the two

batches of ZnP-coated grains confirmed the expected average coating rate per grain. On

average individual grains from the batch mixed at 25 g ZnP/kg grain were coated with 1.06 ±

0.11 mg ZnP (Mean ± SEM; n=8) (1.7, 1.0, 0.9, 1.3, 0.7, 0.9, 1.0, 1.0 mg ZnP/grain), those

from the batch mixed at 50 g ZnP/kg grain were coated with 2.4 ± 0.37 mg ZnP (Mean ±

SEM; n=10) (1.6, 2.9, 2.2, 1.9, 4.8, 3.6, 1.3, 2.0, 1.3 mg ZnP/grain).

2.4 Statistical analysis

The results of the oral gavage were assessed using generalized linear models (GLM),

performed in R.[26] A GLM, was performed using the '*lme4*' package[27] specifying a binominal

error structure to investigate differences in the proportion of mice that died within each

mouse group and dose. To investigate the additional effects that group, sex, fed or unfed,

and their interactions had on the proportion (p) of animals that died, model Akaike

information criterion (AIC) values were compared between the fitted models. The estimated

LD_{50} (where p = 0.5) and LD_{90} (p = 0.9) values and standard error (SE) for each group were

calculated from the best fit model using the *dose.p* function in the *MASS* package.[28]

Estimated LD_{50} and LD_{90} were completed for each mouse group separately and plotted using

the '*plot*' function. Means are presented ± 1 standard error (SEM) throughout.

3 RESULTS

3.1 Experiment 1: Determination of the LD_{50} and LD_{90} of wild and laboratory house mice

The onset of clinical signs of toxicity varied from 2 h to > 6 h post gavage. The time to death

(humane killing) in mice ranged from 2.5 h to 48 h (Table 1). Animals initially showed no

signs of toxicity for several hours but then succumbed within 30-60 min. The first sign of

developing toxicity was a change from a normal resting posture to a hunched posture,

increasing prominence of the guard hairs (pilo-erection) and rapid, shallow breathing. This

was followed by an increased reluctance to move when disturbed, and/or moving with an

unsteady gait (ataxia), and the eyes were half-closed. Animals then showed lateral or sternal

recumbency and loss of righting reflex at which point they were humanely killed. In general,

the higher the oral gavage dose the more rapid the onset of signs of toxicity and the shorter

time to death (Table 1) (Table S1 shows individual timing).

For the laboratory outbred mice (average body weight = 32.4 ± 0.7 g), the time to first

onset of signs of toxicity was delayed compared to the two groups of wild house mice

(average body weight for naïve and exposed mice = 14.5 ± 0.7 and 12.3 ± 0.5 g

respectively). The time to death post gavage was also variable both within and between

mouse groups, ranging from 6.7 - 48 h in outbred lab mice, 2.5 - 24.5 h in naïve wild mice,

and to 3.5 - 32.7 h in exposed wild mice (Table 1).

INSERT Table 1 hereabouts

The proportion of mice that died was dose-dependent (Fig. 1). The proportion of

animals that died at each treatment dose was not different between lab and wild groups, sex,

or fed and unfed status, therefore adding these factors did not improve the model fit (ΔAIC=

6.43; Table S2).

INSERT Fig. 1 hereabouts

The estimated LD_{50} values for each mouse group ranged between 72 and 79 mg

ZnP/kg body weight (Fig. 1; Table 2; $P>0.05$). The estimated LD_{90} values also were not

significantly different between the different mouse groups, ranging between 102 and 120 mg

ZnP/kg body weight; Table 2; $P>0.05$).

INSERT Table 2 hereabouts

None of the Outbred Laboratory mice dosed at 74.4 mg ZnP/kg body weight died

(Fig. 1a; Table S1). If this treatment dose was not included in the model estimating the LD_{50}

and LD_{90} values for outbred mice, the new estimates were 72.8 ± 7.1 and 109.6 ± 14.6 mg

ZnP/kg body weight respectively which are not significantly different from the original

estimates (Table 2).

3.2   Experiment 2: Uptake of ZnP-coated wheat grains

Across both Trials and all treatment groups, mice consumed ZnP-coated grains (18 of 20 mice in UF50, 15 of 20 mice in F50, and 19 of 20 mice in F25) (Table 3), predominantly on the first day of presentation (Tables S3 and S4). Three of the 60 mice (one in each of F50 T1, UF50 T1 and UF50 T2) never consumed any ZnP-coated grains on any of the three days and have been excluded from the analyses.

INSERT Tables 3 hereabouts

For Treatment UF50, nine of 18 animals (50%) were observed to consume ZnP-coated grains within 30-60 mins of their presentation. For the remaining nine animals, six consumed ZnP-coated grains before alternative food was added, while three animals consumed ZnP-coated grains between 1 and 3 h after the addition of alternative food.

For the F50 and F25 treatments in which there was simultaneous presentation of the ZnP-coated grains with alternative food, two of 15 F50 animals (13.3%) and six of 19 F25 animals (31.5%) consumed ZnP-coated grains within the first 60 min. The remaining animals in these fed groups (86.5% and 68.5% respectively) consumed ZnP-coated grains between 2.5 and 19 h after presentation.

Across all treatments, after first recorded consumption of ZnP-coated grain, the shortest time to the onset of signs of toxicity and subsequent humane intervention was 30 min and 1 h respectively. The longest interval to intervention after first recorded ZnP-coated grain consumption was 34 h. The average time to intervention for the three treatments was similar: 8.3 h for UF50,10.2 h for F50 and 13.8 h for F25 (Table 3).

For the two trials of UF50 treatments in which alternative food was not available for the first 4-6 h after presentation of ZnP-coated grains, 17 of 18 mice that consumed ZnP-coated grains died (~94%). By comparison, for the two trials of the F50 treatment where alternative food was available, 15 of 20 mice consumed ZnP-coated grains on Day 1 and

seven of those died (~47%) (Table 3). Similarly, over the two trials of the F25 treatment, on

Day 1, eight of 19 mice that consumed ZnP-coated grains died (42%) (Table 3). Irrespective

of treatment, the number of ZnP-coated grains consumed varied from <0.5 to 4, usually 1-2

grains (Table 3).

For Trial 2, on Days 2 and 3, the F50 and F25 groups did not receive alternative food

for 6 h after presentation of ZnP-coated grains (i.e. same unfed treatment as the UF50

group). For F25 on Day 2, three of the seven remaining mice consumed ZnP-coated grains

(n = 5 - 8 coated grains) before the addition of alternative food and died, but no further

consumption or deaths occurred on Day 3 (n = 4 mice). For F50 on Day 2, four of the eight

remaining mice consumed grains (0.5 - 2 ZnP-coated grains) and died, and on Day 3, one of

the remaining four mice consumed 2.5 ZnP-coated grains and died (Table S4).

Some animals (n = 16/52) which consumed all or part of a ZnP-coated grain on Day 1

(Table 4), did not show any signs of toxicity, did not die, and did not consume ZnP-coated

grains on either Days 2 or 3. This aversion was higher in the groups where background

wheat grains were presented simultaneously with the ZnP-coated grains (F25, n=8, aversion

= 42%; F50, n=7, aversion = 46.7%; UF50, n=1, aversion = 6%) which reduced the quantity

of toxic grain taken (Table 4).

4   DISCUSSION

This assessment of the acute oral toxicity of ZnP has clearly demonstrated that wild house

mice in Australia are significantly less sensitive (LD_{50} of 72-75 mg ZnP/kg body weight) than

previously reported for wild mice in the USA (LD_{50} of 32.68 mg ZnP/kg body weight).[10,11] The

latter value was used by the Australian Pesticides and Veterinary Medicines Authority

(APVMA) to define the dose rate (25 g ZnP/kg grain, whereby ~1 mg ZnP per grain delivers

an LD_{90} to a 15 g mouse) for registration of ZnP for broadscale management of house mice

in cereal crops,[19,20] Our findings indicate that a higher loading of ~2 mg ZnP/grain (equivalent

to an LD_{90} loading) would be required to achieve a lethal outcome after the first consumption

of a single ZnP-coated grain.

In our oral gavage study the outbred strain of laboratory mice also showed a similar

LD_{50} of 79.18 mg/kg to that of the wild mice, whereas Li and Marsh[10], using Swiss Webster

outbred mice, reported an average LD_{50} value of 53.34 mg/kg. They also observed a

significant difference between genders reporting values for males and females of 73.23

mg/kg and 38.86 mg/kg body weight respectively. We detected no gender differences in LD_{50}

estimates within any mouse group, and the LD_{50} values were all within the same range for

the three different mouse groups (72-79 mg/kg). Given the similarity in the LD_{50} values for

naïve and exposed wild mice this also suggests that in the last 20 years regular field

application of ZnP has not led to selection either for non-responders or for tolerant

individuals. There have been no previous reports in the literature indicating any development

of tolerance or genetic resistance to ZnP.[29]

Nevertheless, we observed high variability in the acute response for both outbred

laboratory mice and wild mice after oral gavage intake of ZnP in each of our experiments.

For example, at the estimated LD_{50} of ~74 mg ZnP/kg body weight, 60-70% exposed and

naïve wild mice died whereas none of the outbred laboratory mice died at this dose. In their

original study,[10] 80% wild mice dosed at 35.62 mg ZnP/kg body weight died whereas none of

our wild mice (either exposed or naïve) died after dosing at 30 mg ZnP/kg body weight.

The earliest time at which our study animals were humanely killed after their oral

gavage dose of ZnP or after consumption of ZnP-coated grains was one hour but could be

up to 48h. This time-to-death range and the development of various signs of acute toxicity

are similar to that observed in other studies in mice and rats.[23,30] This reflects the expected

effects of phosphine gas production in the gut and/or direct uptake of ZnP across the gut and

subsequent failure of major internal organs such as the liver, kidney, brain and heart.[8]

When wild mice were presented with ZnP-coated wheat grains (either ZnP50 or

ZnP25), there was also considerable variation in responses after consumption including a

rapid development of aversion as previously reported for several species including wild

mice[23], common voles, *Microtus arvalis*[31] and prairie voles (*Microtus ochragaster* Wagner).[32]

Any wild mouse taking a sub-lethal dose on first exposure, rarely consumed ZnP-coated

grains subsequently and this aversive response was higher when there was a choice of

alternative food (42 - 47% mice became averse after consuming ZnP25- or ZnP50-coated

grains). Aversion was very low (6%) when ZnP50-coated grains were the only food choice

present for >4 hours but this is because 94% mice died after their first consumption of ZnP50

grains. We therefore conclude that increasing the ZnP dose per grain does not decrease its

palatability with aversion likely due to the sub lethal adverse effects of the toxin which is

much more likely to occur with the ZnP25-coated grains.

In vitro digestion studies have shown that the production of phosphine gas was

reduced where wheat grain was digested prior to the addition of ZnP-coated grains.[33] It was

concluded that this reduction was most likely due to the inhibition of degradation of the ZnP

in the presence of digested wheat grains and gastric enzymes. If this occurred in mice, the

presence of food in the stomach could affect the outcome of ZnP-coated grain consumption.

However, we have no direct evidence of this occurring in either of our experiments. Our

observations suggest the variation in responses is due to a combination of different factors

rather than a single factor - we suggest that the outcome for individual mice might vary due

to quantity of ZnP consumed at first presentation (the number of grains consumed and the

loading per grain), the presence of alternative food, their physiological response in the gut

and their body weight.[30]

The amount of ZnP coating individual wheat grains was independently analysed and,

on average, matched the expected dose rate (either 25 g ZnP/kg grain or 50 g ZnP/kg grain).

However, there was variation between grains in their ZnP loading which could have affected

mortality rates and aversion particularly when consumption of these ZnP coated grains

occurred in the presence of other food. Similar variations were observed for a commercially

prepared batch of grains used in another study[23] and may be due to varying amounts of ZnP

powder being trapped in the crease of grains of different sizes and shapes during coating.

In Australia poisoning of birds after ZnP baiting has rarely been recorded [13, 18, 21]. If

there is a higher ZnP coating per grain, does this represent a greater risk to non-target

species, such as granivorous birds? For the currently registered rate, granivorous birds

would need to detect and consume many ZnP-coated grains before a lethal dose

accumulated (see Table 26[29]). Whether doubling the dose of ZnP coating per grain alters the

potential for non-target impacts in field situations is not known and must be carefully

examined. However, given that ZnP-coated grains can only be applied in-crop in Australia

non-target poisoning of wild birds should remain infrequent.[21,33] It is also unlikely that non-

target species would be at greater risk of secondary poisoning as ZnP does not accumulate

in tissues and the greatest risk would be if the predator consumed the digestive tract which

may contain undigested ZnP.[2]

As noted earlier, given the increased LD_{50} value for ZnP in wild mice, the coating per

grain which represents a lethal dose (LD_{90}) for most 15 g mice needs to be approximately 2

mg ZnP per grain. This increased coating rate means that the majority of ZnP-coated grains

presented to mice would likely be a lethal dose, effectively reducing the chances of mice

encountering a sub lethal dose and developing aversion. When wild mice were presented

with grains coated with ~2 mg ZnP they consumed these and died and especially when no

alternative food was available. In the laboratory trials, mice were caged individually and had

easy access to alternative food as well as the ZnP-coated grains giving them ample

opportunity to be selective in their choice of food. This contrasts with what occurs in a field

situation where there is intraspecific competition for food and shelter and there is the threat

of predation. The relationship between available alternative food and the efficacy of ZnP50

needs to be assessed experimentally in a field situation.

The higher values for ZnP LD_{50} provide us with a plausible explanation for the

variable efficacy observed by farmers when the currently ZnP registered rate of 25 g ZnP/kg

grain is applied. Our findings force us to conclude that ZnP bait applied at the currently

prescribed rate of 1 kg/ha (25 g ZnP/kg grain) means that mice are not being presented with

three lethal doses per square metre in the field (2-3 grains/m^2). Instead, to be assured of

receiving a lethal dose, mice would need to detect and consume at least two ZnP25-coated

grains. This immediately raises questions about the capacity of mice to find a second or third

ZnP25-coated grain prior to the onset of symptoms of toxicity in the complex environment in

paddocks where conservation agriculture (CA) is practised.[22] In our laboratory assessment

where food was provided at the same time as the ZnP-coated grain, the proportion of mice

dying was lower regardless of the ZnP dose per grain as mice ate fewer ZnP-coated grains

(Table 3). The artificial nature of the laboratory comparison indicates that further comparative

studies in CA cropping systems are required. We predict that the amount of alternative food

available in the field at the time of ZnP baiting will influence the detection and uptake of ZnP-

coated grains. In addition, the overall efficacy of a ZnP baiting program in field situations is

likely to be affected by the growth stage of the crop being protected. In CA systems where

available alternative food could be >200 grains/m^2, this will likely affect ZnP baiting outcomes

when the toxic grains are spread at only 2-3 grains/m^2. Assessment of the efficacy of the

higher dose (50 g ZnP/kg grain) compared to the registered dose at a population level under

field conditions is needed to verify the findings of these laboratory-based trials.

5   CONCLUSIONS

This study has conclusively demonstrated that the LD_{50} for Australian wild house mice

is more than two-fold higher than previously reported in the literature for wild mice. This

indicates that the loading of ZnP on an individual grain should be doubled to achieve a lethal

dose per grain and avoid bait aversion developing from sub-lethal effects of the toxin. Grains

coated with the equivalent of an LD_{90} quantity (approximately 2 mg), were readily consumed

by mice and mortality was greater than 90%, particularly in the absence of alternative food.

Results were less conclusive when alternative non-toxic grains were provided at the time of

presentation of ZnP-coated grains. Assessment of the efficacy of the higher dose (50 g

ZnP/kg grain) compared to the currently registered dose (25 g ZnP/kg grain) at a population

level under field conditions is needed to validate the findings of these laboratory-based trials.

ACKNOWLEDGEMENTS

This work was supported by the Grains Research and Development Corporation (GRDC)

through project CSP1804-012RTX, and CSIRO Health and Biosecurity. We sincerely thank

farmers for allowing us to trap mice on their properties for this experiment. We thank Leigh

Nelson and Ken Young (GRDC) for ongoing support. Richard Duncan, University of Canberra,

provided helpful advice on statistical analyses. All experiments and procedures were approved

by the CSIRO Wildlife and Large Animal Ethics Committee (Approval No 2019-23) and conform

to the Australian Code of Practice for the Care and Use of Animals for Scientific Purposes.

CONFLICT OF INTEREST DECLARATION

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author

upon reasonable request.

REFERENCES

1. Eisemann JD, Petersen BE, Fagerstone KA, Efficacy of zinc phosphide for controlling Norway rats, roof rats, house mice, *Peromyscus* spp., prairie dogs and ground squirrels: a literature review (1942 - 2000). *Proceedings 10th Wildlife Damage Conference,* USDA National Wildlife Research Center - Staff Publications **212:**335-349 (2003).

2. Sterner RT, Mauldin RE, Regressors of whole-carcass zinc phosphide/phosphine residues in voles: indirect evidence of low hazards to predators and/scavengers. *Archives of Environmental Contamination and Toxicology* **28**:519-523 (1995).

3. Tkadlec E, Rychnovsky B, Residues of Zn_3P_2 in the common vole (*Microtus arvalis*) and secondary poisoning hazards to predators. *Folia Zoologica* **39:**147-56 (1990).

4. Hood GA, Zinc Phosphide − a new look at an old rodenticide for field rodents. *Proceedings of 5th Vertebrate Pest Conference*: University of California Agriculture and Natural Resources.pp 85-92 (1972).

5. Gupta RC, Non-anticoagulant rodenticides. . In: Gupta RC, editor. *Veterinary Toxicology: Basic and Chemical Principles* Academic Press pp 548-560 (2007).

6. Eason C, Ross J, Blackie H, Fairweather A, Toxicology and ecotoxicology of zinc phosphide as used for pest control in New Zealand. *New Zealand Journal of Ecology* **37:**1-11 (2013).

7. Casteel SW, Bailey EM, Jr. A review of zinc phosphide poisoning. *Veterinary and Human Toxicology* **28:**151-154 (1986).

8. Khan SA, Schell MM, Zinc Phosphide. https://www.msdvetmanual.com/toxicology/rodenticide-poisoning/zinc-phosphide (2014).

9. Morgan D, Eason C, Wickstrom M, Zinc phosphide as a poison for control of possums (*Trichosurus vulpecula*): the need for encapsulation. In: Pelz H-J CD, Feare CJ editor. *Advances in vertebrate pest management II*. Filander, Fürth: Verlag; pp. 337-344 (2001)

10. Li J, Marsh RE, LD_{50} determination of zinc phosphide toxicity for house mice and albino laboratory mice. In: Crabb AC, Marsh RE, editors *Proceedings of the 13th Vertebrate Pest Conference*: University of California, Davis, pp. 91-94 (1988).

11. Bell HB, The hazards of secondary poisoning from zinc phosphide to selected vertebrate species. *Master's Thesis* University of Tennessee. 33 pp(1972).

12. Singleton GR, Brown PR, Pech RP, Jacob J, Mutze GJ, Krebs CJ, One hundred years of eruptions of house mice in Australia – a natural biological curio. *Biological Journal of the Linnean Society* **84:**617-627 (2005).

13. Caughley J, Bomford M, Parker B, Sinclair R, Griffiths J, Kelly D, Managing vertebrate pests: rodents. Canberra: Bureau of Resource Sciences and Grains Research Development Corporation, 130 pp (1998a).

14. Brown PR, Huth NI, Banks PB, Singleton GR, Relationship between abundance of rodents and damage to agricultural crops. *Agriculture, Ecosystems & Environment* **120:**405-15 (2007).

15. Brown PR, Singleton GR, Pech R, Hinds LA, Krebs CJ, Rodent outbreaks in Australia: mouse plagues in cereal crops. In: Singleton GR, Belmain SR, Brown PR, Hardy B, editors. *Rodent Outbreaks: Ecology and Impacts*. Los Baños, Philippines: International Rice Research Institute, pp. 225-238 (2010).

16. Brown PR, Arthur AD, Jones DA, Davies MJ, Grice D, Pech RP, Multiple ecological processes underpin the eruptive dynamics of small mammals: House mice in a semi-arid agricultural environment. Ecology and Evolution **10:**3477-3490 (2020).

17. Caughley J, Monamy V, Heiden K, Impact of the 1993 Mouse Plague. Canberra, Australia GRDC, 73 p (1994).

18. Caughley J, Donkin C, Strong K, Managing mouse plagues in rural Australia. In: Baker RO, Crabb AC editors *Proceedings of the 18th Vertebrate Pest Conference*; University of California, Davis, pp. 160-165(1998b).

19. Staples L, Smith M, Pontin K, Hunt W, Use of zinc phosphide to overcome rodent
infestations. In Wright EJ, Webb MC, Highley E editors. *Stored grain in Australia 2003*
Proceedings of the Australian Postharvest Technical Conference, Canberra, ACT
Australia: CSIRO Stored Grain Research Laboratory, pp 110-115 (2003).
20. APVMA, Evaluation of the new active zinc phosphide in the product MouseOff zinc
phosphide bait. https://apvma.gov.au/node/14101. Canberra, Australia: National
Registration Authority for Agricultural and Veterinary Chemicals. (2000).
21. Brown PR, Chambers LK, Singleton GR, Pre-sowing control of house mice (*Mus*
domesticus) using zinc phosphide efficacy and potential non-target effects. *Wildlife*
Research **29**:27-37 (2002).
22. Ruscoe WA, Brown PR, Henry S, Van de Weyer N, Robinson F, Hinds LA, *et al*.,
Conservation agriculture practices have changed habitat use by rodent pests:
implications for management of feral house mice. *Journal of Pest Science*
https://doi.org/10.1007/s10340-021-01370-7. (2021).
23. Henry S, Brown PR, Van de Weyer N, Robinson F, Hinds LA, Effects of background
food on alternative grain uptake and zinc phosphide efficacy in wild house mice. *Pest*
Management Science https://doi.org/10.1002/ps.6720 (2021).
24. OECD, OECD Guidelines for testing of chemicals: Acute oral toxicity – Up-and-Down
procedure. Paris, France, 26 pp. (2001).
25. Bruce RD, An up-and-down procedure for acute toxicity testing. *Fundamental and*
Applied Toxicology **5:**151-157 (1985).
26. R Core Team, *R: A language and environment for statistical computing*.
https://www.R-project.org/ R Foundation for Statistical Computing; Vienna (2020).
27. Bates D, Mächler M, Bolker B, Walker S, Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67:**1-48 (2015).
28. Venables WN, Ripley BD, Modern Applied Statistics with S. 4th ed. New York,
Springer, 498 p. (2002).
29. McLeod L, Saunders G, Pestcides used in the management of vertebrate pests in
Australia: a review. NSW Dept Primary Industries 176 p. (2013).
30. Mason G, Littin KE. The humaness of rodent pest control. *Animal Welfare* **12**:1-37
(2003).
31. Jacob J, Budde M, Leukers A. Efficacy and attractiveness of zinc phosphide bait in
common voles (*Microtus arvalis*). *Pest Management Science* **66:**132-6 (2010).
32. Horak KE, Hofmann NM, Kimball BA, Assessment of zinc phosphide bait shyness
and tools for reducing flavor aversions. *Crop Protection* **112**:214-219 (2018).
33. Paroz GN, Parker RW, Hannan-Jones MA, Szetu JL, Storage stability and
environmental fate of zinc phosphide grain bait: Grains Research & Development
Corporation, 123 p. (1999).
593
594
595
596

Figures

Figure 1: Proportion of mice dying after oral gavage with different ZnP concentrations (mg
ZnP/kg body weight). Calculated dose response curves for (a) Outbred laboratory mice (b)
Naïve wild house mice and (c) Exposed wild house mice. Horizontal red dashed line
represents 50% mortality, vertical solid line equates to LD_{50} value, vertical dashed lines
represent standard error for the LD_{50} estimate. n>4 animals per test dose, with a mix of
males and females.

(a) Outbred laboratory mice

(b) Naïve wild house mice

(c) Exposed wild house mice

Table 1: Mortality (%) and time to death (h) for different mouse groups after oral gavage of different concentrations of ZnP in oil.

Dose (mg ZnP/kg)	*Body weight (g)	n	Mortality (%)	Time to death (h post gavage) (range)
Outbred laboratory mice				
50.22	29.9 ± 1.8	8	25.0	25.0 - 25.3
65.1	34.9 ± 1.6	6	33.3	31.0 - 48.0
74.4	34.8 ± 1.5	5	0.0	-
83.7	32.2 ± 1.8	8	62.5	9.3 - 29.5
102.3	31.9 ± 1.8	8	87.5	7.5 - 28.0
148.8	32.0 ± 2.1	4	100.0	6.7 - 10.0
Mean ± SEM	32.4 ± 0.7 (n = 39)			18.5 ± 2.4 (n = 20)
Wild mice (Naïve)				
29.95	12.9 ± 1.8	8	0.0	-
50.22	15.3 ± 1.8	8	37.5	6.75 - 23.8
74.4	15.3 ± 1.2	7	71.4	6.3 - 19.0
102.3	13.3 ± 1.8	8	62.5	3.17 - 8.4
148.8	15.4 ± 1.8	8	100.0	2.5 - 24.5
Mean ± SEM	14.5 ± 0.7 (n = 39)			9.8 ± 1.5 (n = 21)
Wild mice (exposed)				
30.41	16.3 ± 0.8	8	0.0	
50.22	10.9 ± 0.7	7	14.3	24.0
65.1	11.1 ± 0.9	4	25.0	19.8
74.4	11.6 ± 0.8	8	62.5	3.5 - 23.3
83.7	9.9 ± 0.8	6	50.0	6.7 - 32.7
102.3	12.4 ± 0.8	11	90.9	4.0 - 24.0
Mean ± SEM	12.3 ± 0.5(n = 44)			11.8 ± 2.0 (n = 20)

610 *Body weight (Mean ± SEM) of dosed animals on day of treatment, includes males and females.

612 **Table 2:** Estimated LD_{50} and LD_{90} values (mg ZnP/kg body weight; Mean ± SEM) for different groups of mice.

Mouse group	Total number mice dosed	LD_{50} (mg ZnP/kg body weight)	LD_{90} (mg ZnP/kg body weight)
Outbred Laboratory strain	39	79.18 ± 6.24	114.58 ± 14.6
Naïve wild	39	72.11 ± 9.09	119.50 ± 18.01
Exposed wild	44	75.22 ± 4.39	102.27 ± 9.34

Table 3: Responses of wild mice on Day 1 to exposure and uptake of ZnP-coated grains for each treatment over two trials (T1, T2). Body
weights (Mean ± SEM (n)) of each group at start of each treatment round, number of animals consuming ZnP-coated grains on Day 1, mortality
(%) after consumption, time to death (range, Mean ± SEM), and number of ZnP-coated grains consumed by individual mice.

Treatment	Body weight (g)	Animals consuming toxic grains - Day 1	Animals dying after consumption (%)	Time to death (h) (range) Mean ± SEM	Number of ZnP-coated grains consumed on first occasion by individual animals
UF50 T1	15.0 ± 0.7 (10)	9 of 10	8 of 9 (88.9)	1 - 22	*1*, **2**, **<0.5**, **2**, **1**, **2.5**, **2.3**, **1**, **1**
UF50 T2	14.7 ± 0.5 (10)	9 of 10	9 of 9 (100)	3 - 34	**0.5**, **3**, **3**, **1**, **2**, **2**, **1**, **<0.5**, **4**
			17 of 18 (94.4)	8.3 ± 2.1	
F50 T1	14.1 ± 0.9 (10)	9 of 10	5 of 9 (55)	2 - 14	*0.5, 1, 0.5, <0.5*, **0.5**, **2**, **1.5**, **1**, **3.5**
F50 T2	15.6 ± 0.6 (10)	6 of 10	2 of 6 (33)	2.5 - 17.5	*1, <0.5, 1*, 0.5, **1**, **1**
			7 of 15 (46.7)	10.2 ± 1.9	
F25 T1	15.3 ± 0.7 (10)	10 of 10	5 of 10 (50)	5 - 24	*<0.5, 0.5, 0.5, 3*, **1**, **2**, **1.5**, **<0.5**, **4**, **2**
F25 T2	14.6 ± 0.8 (10)	9 of 10	3 of 9 (33)	3.5 - 27	*1, 0.5, 1, 1*, 0.5, 2, **1**, **1.7**, **1**
			8 of 19 (42.1)	13.8 ± 2.2	

Numbers in bold are for those animals which died after consumption of Zn-P coated grains.
Numbers in italics are those animals which became averse after first uptake of ZnP-coated grains on Day 1 only.
UF50 = UnFed, 50 g ZnP/kg grain; F50 = Fed, 50 g ZnP/kg grain; F25 = Fed, 25 g ZnP/kg grain

Table 4. Mice showing aversion after first consumption of Zn-P coated grains across Trials 1 and 2

Treatment	n	Animals (n) that did not die after taking ZnP-coated grains on Day 1 and did not consume thereafter	Averse (%)
UF50 T1, T2	9, 9	1, 0	6
F50 T1, T2	9, 6	4, 3	46.7
F25 T1, T2	10, 9	4, 4	42.1
Totals	52	16	

Table S1: Time to death (h) after oral gavage with different doses of ZnP in oil for laboratory mice (Swiss Outbred), wild mice (Naïve) and wild
 mice (Exposed).

Dose (mg/kg)	n	No. humanely killed	Time to death (h post gavage)			
			Female (fed)	Female (unfed)	Male (fed)	Male (unfed)
Outbred laboratory mice						
50.22	8	2	25.0		25.3	
65.1	6	2	31		48	
74.4	5	0				
83.7	8	5	9.25,16.0		20.3, 29.5	17.5
102.3	8	7	8, 19.5	7.5	19, 23.5	10, 28
148.8	4	4	7	8.5	6.7	10
Wild mice (Naïve)						
32.2	8	0				
50.22	8	3	6.75	11.17		23.83
74.4	7	5		6.33	7.25, 9.5, 15.5	19
102.3	8	5	3.17, 3.42		3.25, 7.75	8.42
148.8	8	8	2.5	7, 22.33	4, 4.17, 8	8.17, 24.5
Wild mice (exposed)						
30.41	8	0				
50.22	7	1				24.0
65.1	4	1	19.83			
74.4	8	5	5.67, 20.16	3.5	14.58	23.33
83.7	6	3	7.33, 32.67	6.67		
102.3	11	10	5, 5.33, 6.75	4, 8.33	4.67, 5.25, 8	6, 24.0

Table S2: Summary of proportion of animals that died, modelled with dose, group, sex and fed or unfed as varying fixed effect factors. The top model included the oral gavage dose only (bold).

Model	df	AIC	ΔAIC
Dose	**2**	**91.54**	0
Dose + Group	6	97.23	5.69
Dose + Group + Sex + Fed/Unfed	8	97.97	6.43

641 **Table S3.** Number of mice consuming ZnP-coated grains and dying each day over 3 days, and overall mortality (%) for Trial 1 (T1) of each
642 treatment.
643 F25 = Fed, 25 g ZnP/kg grain; F50 = Fed, 50 g ZnP/kg grain; UF50 = UnFed, 50 g ZnP/kg grain. For F25 and F50 ZnP-coated grains and
644 alternative standard wheat grains were added simultaneously. For UF50, alternative food was added 4 h later.

Treatment		Day 1			Day 2			Day 3		Mortality over 3 days (%)
	n	No. taking	No. dead (%)	n	No. taking	No. dead	n	No. taking	No. dead	
UF50 T1	*9	9	8 (88.9)	1	0	0	1	0	0	88.9
F50 T1	*9	9	5 (55.5)	4	0	0	4	0	0	55.6
F25 T1	10	10	5 (50)	5	1	1	4	0	0	60.0

645 *One animal in each group did not consume any ZnP-coated grains and was excluded from analysis.
646
647

648 **Table S4.** Number of mice consuming ZnP-coated grains and dying each day over 3 days, and overall mortality (%) for Trial 2 (T2) of each
649 treatment. F25 = Fed, 25 g ZnP/kg grain; F50 = Fed, 50 g ZnP/kg grain; UF50 = UnFed, 50 g ZnP/kg grain. [#] For Days 2 and 3, all groups were
650 presented with ZnP-coated grains 6 h before addition of alternative standard wheat grains.

| Treatment | Day 1 | | | Day 2[#] | | | Day 3[#] | | | No. dead, |
	n	No. taking	No. dead (%)	n	No. taking	No. dead	n	No. taking	No. dead	days 2-3 (%)
UF50 T2	*9	5	4 (80)	5	5	5	-	-	-	5/5 (100)
F50 T2	10	6	2 (33)	8	4	4	4	1	1	5/8 (62.5)
F25 T2	10	9	3 (30)	7	3	3	4	0	0	3/7 (42.8)

651 *One animal did not consume any ZnP-coated grain and was excluded from analysis.
652