<u>Title</u>: *Salmonella* Typhimurium induces NAIP/NLRC4- and NLRP3/ASC-independent,

caspase-1/4-dependent inflammasome activation in human intestinal epithelial cells

<u>Running title</u>: Human epithelial inflammasome responses to *Salmonella*

<u>Authors</u>: Nawar Naseer[a], Renate Bauer[bd], Jenna Zhang[a], Igor E. Brodsky[c], Isabella

Rauch[b], and Sunny Shin[a#]

<u>Author affiliations</u>:

[a]Department of Microbiology, University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA 19104

[b]Department of Molecular Microbiology and Immunology, Oregon Health & Science

University, Portland, OR 97239

[c]Department of Pathobiology, University of Pennsylvania School of Veterinary Medicine,

Philadelphia, PA 19104

[d]Department of Biosciences, Paris Lodron University of Salzburg, Salzburg, Austria

#: Email correspondence to Sunny Shin: sunshin@pennmedicine.upenn.edu

Abstract

Salmonella enterica serovar Typhimurium is a gram-negative pathogen that causes

diseases ranging from gastroenteritis to systemic infection and sepsis. *Salmonella* uses

type III secretion systems (T3SSs) to inject effectors into host cells. While these

effectors are necessary for bacterial invasion and intracellular survival, intracellular

delivery of T3SS products also enables detection of *Salmonella* by cytosolic immune

sensors. Upon detecting translocated *Salmonella* ligands, these sensors form

multimeric complexes called inflammasomes, which activate caspases that lead to

proinflammatory cytokine release and pyroptosis. In particular, the *Salmonella* T3SS

needle, inner rod, and flagellin proteins activate the NAIP/NLRC4 inflammasome in

murine intestinal epithelial cells (IECs), which leads to restriction of bacterial replication

and extrusion of infected IECs into the intestinal lumen, thereby preventing systemic

dissemination of *Salmonella*. While these processes are studied quite well in mice, the

role of the NAIP/NLRC4 inflammasome in human IECs remains unknown.

Unexpectedly, we found the NAIP/NLRC4 inflammasome is dispensable for early

inflammasome responses to *Salmonella* in both human intestinal epithelial cell lines and

organoids. Additionally, the NLRP3 inflammasome and the adaptor protein ASC are not

required for inflammasome activation in Caco-2 cells. Instead, we observed a partial

requirement for caspase-1, and a necessity for caspase-4 and GSDMD pore-forming

activity in mediating inflammasome responses to *Salmonella* in Caco-2 cells. These

findings suggest that unlike murine IECs, human IECs do not rely on NAIP/NLRC4, and

also do not use NLRP3/ASC. Instead, they primarily use caspases-1 and -4 to mediate

early inflammasome responses to SPI-1-expressing *Salmonella*.

 Introduction

Enteric bacterial pathogens such as *Salmonella* enterica serovar Typhimurium

(hereafter referred to as *Salmonella*) are leading causes of global morbidity and

mortality from diarrheal diseases (1). Contracted upon ingestion of contaminated food or

water, *Salmonella* colonizes the intestinal tract, where it uses evolutionarily conserved

molecular syringes called type III secretion systems (T3SS) to inject effectors, or

virulence factors, into the host cell cytosol (2). *Salmonella* contains two T3SS: the SPI-1

T3SS is expressed early in infection and enables *Salmonella* to invade host cells, while

the SPI-2 T3SS is expressed at later timepoints during infection and allows *Salmonella*

to replicate within host cells such as intestinal epithelial cells (IECs) (2). IECs thus serve

as both the targets of, as well as the first line of physical and innate immune defense

against enteric pathogens like *Salmonella*. Most studies of *Salmonella*'s interactions

with the innate immune system have been conducted in mice. However, key differences

in innate immune genes encoded by mice and humans make it unclear whether mice

fully recapitulate how humans respond to *Salmonella*. Here, we interrogated how

human IECs sense and respond to *Salmonella* infection.

The mammalian immune system can recognize invading intracellular pathogens

through cytosolic sensors such as nucleotide-binding domain, leucine-rich repeat (NLR)

receptors. Upon detecting a bacterial ligand or activity, these receptors oligomerize to

form multimeric signaling complexes called inflammasomes (3). Inflammasomes recruit

and activate cysteine proteases, such as caspase-1 and caspase-8 (3, 4). Some

inflammasomes require an adaptor protein called apoptosis-associated speck-like

protein containing a CARD (ASC) to mediate their interaction with caspases (3). Active

caspases can process the proinflammatory cytokines IL-1α, IL-1β, and IL-18 (3), and

the pore-forming protein GSDMD (5). This leads to GSDMD-dependent release of the

processed cytokines and an inflammatory form of cell death known as pyroptosis (3).

Release of these proinflammatory cytokines alerts nearby cells of the infection, while

pyroptosis can eliminate the pathogen's replicative niche within the infected host cell.

Various cellular insults during infection can trigger activation of different

inflammasomes. Inflammasome activation is critical for control of *Salmonella* infection in

mice (6). In both murine and human macrophages and murine IECs, *Salmonella*

infection activates a family of inflammasome sensors termed NAIPs, which detect the

Salmonella SPI-1 T3SS needle, inner rod, and flagellin proteins (7–12). Activation of the

NAIP/NLRC4 inflammasome specifically in murine IECs restricts bacterial replication,

causes extrusion of infected cells from the intestinal epithelial layer, and prevents

dissemination of *Salmonella* to systemic sites (13–15). In contrast to mice, humans

encode a single NAIP inflammasome sensor, which promiscuously recognizes the

T3SS needle, inner rod, and flagellin proteins in human macrophages (10, 16, 17). The

role of the NAIP/NLRC4 inflammasome in human IECs during *Salmonella* infection

remains unknown.

Salmonella infection can also induce the NLRP3 inflammasome, which can be

activated by a variety of stimuli during infection, including potassium efflux (18). In

murine macrophages, the NLRP3 inflammasome is thought to be important for late

timepoints during *Salmonella* infection (19). In murine macrophages and murine and

human intestinal epithelial cells, the caspase-11 (mice) or caspase-4/5 (humans)

inflammasome is activated late in *Salmonella* infection (19–23). Caspase-4/5/11 detect

cytosolic LPS and form the noncanonical inflammasome, which can secondarily activate

the NLRP3 inflammasome (24, 25). In human macrophages, *Salmonella* infection

triggers recruitment of both NLRC4 and NLRP3 to the same macromolecular complex

(26). However, whether the NLRP3 inflammasome plays a functional role during

Salmonella infection of human IECs has not been previously tested.

Human IECs infected with *Salmonella* undergo caspase-4 inflammasome

activation at late time points following infection (10hpi), when there is a considerable

population of replicating cytosolic bacteria (20, 22). However, early inflammasome

responses to SPI-1-expressing *Salmonella* have not been previously investigated. In

this study, we have found that human IECs undergo inflammasome activation in

response to SPI-1-expressing *Salmonella*. However, T3SS ligands and flagellin were

not sufficient to activate the inflammasome in human Caco-2 cells or intestinal

organoids. Additionally, using a combination of pharmacological inhibitors and

CRISPR/Cas9 technology, we found that the NAIP inflammasome, the NLRP3

inflammasome, and the adaptor protein ASC are all dispensable for early

inflammasome responses to SPI-1-expressing *Salmonella* in Caco-2 cells. Instead, we

observed that caspase-1 is partially required, whereas caspase-4 is necessary for

inflammasome activation in Caco-2 cells in response to *Salmonella* infection. Our

findings delineate the role of several inflammasomes in human IECs *Salmonella*

infection. Importantly, these findings indicate how widely inflammasome responses to

infection can vary between species as well as cell types.

Materials and Methods

<u>Ethics statement</u>

All studies involving human peripheral blood mononuclear cells (PBMCs) and human

intestinal organoids were performed in compliance with the requirements of the US

Department of Health and Human Services and the principles expressed in the

Declaration of Helsinki. Both human PBMCs and organoids are considered to be a

secondary use of deidentified human specimens and are exempt via Title 55 Part 46,

Subpart A of 46.101 (b) of the Code of Federal Regulations. All experiments performed

with murine organoids were done so in compliance with the regulatory standards of, and

were approved by the Oregon Health & Science University Institutional Animal Care and

Use Committee.

<u>Bacterial strains and growth conditions</u>

Salmonella enterica serovar Typhimurium SL1344 WT and Δ*sipB* (27) isogenic

strains were routinely grown shaking overnight at 37°C in Luria-Bertani (LB) broth with

streptomycin (100 μg/ml). Cells were infected with *Salmonella* grown under SPI-1-

inducing conditions (28).

Listeria monocytogenes WT and isogenic strains on the 10403S background

were routinely grown shaking overnight at 30°C in brain heart infusion (BHI) broth (29).

S. Typhimurium ligands PrgJ or PrgI were translationally fused to the truncated N-

terminus of ActA and under the control of the *actA* promoter (29). The *Listeria* strain

expressing *S.* Typhimurium SsaI was constructed using a codon-optimized gene

fragment (IDT) cloned into the pPL2 vector and introduced into *Listeria* as previously

described (29, 30).

<u>Bacterial infections</u>

Where indicated, cells were primed with 100ng/ml or 400ng/ml of Pam3CSK4

(Invivogen) for 3 hours prior to infection. To induce SPI-1 expression, overnight cultures

of *Salmonella* were diluted into LB broth containing 300 mM NaCl and grown for 3 hours

standing at 37°C (28). Overnight cultures of *Listeria* were diluted and grown shaking for

3 hours in BHI. All cultures were pelleted at 6,010 × *g* for 3 minutes, washed once with

PBS, and resuspended in PBS. Cells were infected with *Salmonella* at a multiplicity of

infection (MOI) of 60 or *Listeria* at the indicated MOI for each experiment in the figure

legend. Infected cells were centrifuged at 290 × *g* for 10 minutes and incubated at

37°C. 1 hour post-infection, cells were treated with 100ng/mL or 50ng/mL of gentamicin

to kill any extracellular *Salmonella* or *Listeria* respectively. Infections proceeded at 37°C

for the indicated length of time for each experiment. Control cells were mock-infected

with PBS for all experiments.

<u>Cell culture of intestinal epithelial cell lines</u>

All cell lines were obtained from American Type Culture Collection (ATCC).

Caco-2 cells (HTB-37; ATCC) were maintained in DMEM supplemented with 10%

(vol/vol) heat-inactivated FBS, 100 IU/mL penicillin, and 100 μg/mL streptomycin. T84

cells (CCL-248; ATCC) were maintained in DMEM F-12 supplemented with 5% (vol/vol)

heat-inactivated FBS, 100 IU/mL penicillin, and 100 μg/mL streptomycin.

One day prior to infection or treatment, cells were dissociated with 0.25%

Trypsin-EDTA (Gibco) diluted 1:1 with 1X PBS. Cells were incubated with trypsin at

37°C for 15 minutes, after which the trypsin was neutralized with serum-containing

media. Cells were replated in media without antibiotics in a 24-well plate at a

concentration of 3×10^5 cells/well. Where indicated, cells were primed with 100ng/mL or

400ng/ml Pam3CSK4 (Invivogen) or 500ng/ml LPS (Sigma-Aldrich) for 3h or 16h prior

to anthrax toxin treatment or bacterial infections.

<u>Culture of intestinal organoids in spheroid culture</u>

Spheroids derived from human duodenum or colon were kindly provided by

Jared Fisher at Oregon Health & Science University. Human and murine organoids

were cultivated in special conditioned medium as described by Myoshi & Stappenbeck

(31). Briefly, spheroids were grown in Corning® Matrigel® Basement Membrane Matrix

(VWR). Spheroids were dissociated with Trypsin-EDTA (1X PBS, 0.25% trypsin, 0.5

mM EDTA) at 37°C for 5 minutes. Subsequently, washing medium (DMEM/F12 with

HEPES (Sigma), 10% FBS (heat-inactivated), 1X P/S) was added to stop trypsinization.

After washing the cells with washing medium at $200 \times g$ for 5 minutes at room

temperature (RT), the supernatant was completely aspirated and the cell pellet

resuspended in Matrigel®. 20 µL matrigel-cell drops were placed into 24-well plates and

polymerized for 5 minutes at 37°C upside down. Afterwards, cells were cultured in

organoid medium consisting of 50% primary culture medium (Advanced DMEM/F12

(Gibco), 20% FBS, 1X P/S, 1X L-Glu) and 50% conditioned medium (L-WRN-cell

supernatant) supplemented with 10 µM of ROCK inhibitor Y27632 (Sigma) and 10 µM

TGF-β inhibitor SB431542 (Millipore) (for murine organoids only).

ELISAs

Supernatants harvest from infected cells were assayed using ELISA kits for

human IL-18 (R&D Systems) and IL-8 (R&D Systems).

Immunoblot analysis

Cell lysates were harvested for immunoblot analysis by adding 1X SDS/PAGE

sample buffer to cells following infection. Cells were incubated and infected in serum-

free media to collect supernatant samples. Supernatant samples were centrifuged at

$200 \times g$ to pellet any cell debris. The supernatant was then treated with trichloroacetic

acid (TCA) (25 µL of TCA for 500 µL of supernatant) overnight at 4°C. The next day, the

samples were centrifuged at maximum speed ($15871 \times g$) for 15 minutes at 4°C.

Precipitated supernatant pellets were washed with ice-cold acetone, centrifuged at

maximum speed ($15871 \times g$) for 10 minutes at 4°C, and resuspended in 1X SDS/PAGE

sample buffer. All protein samples (lysates and supernatants) were boiled for 5 minutes.

Samples were separated by SDS/PAGE on a 12% (vol/vol) acrylamide gel, and

transferred to PVDF Immobilon-P membranes (Millipore). Primary antibodies specific for

human IL-18 (MBL International PM014) and β-actin (4967L; Cell Signaling) and HRP-

conjugated secondary antibody anti-rabbit IgG (7074S; Cell Signaling) were used. ECL

Western Blotting Substrate or SuperSignal West Femto (both from Pierce Thermo

Scientific) were used as the HRP substrate for detection.

<u>Propidium iodide (PI) uptake assay</u>

7.5×10^4 Caco-2 cells per well were plated in a black, flat-bottom 96-well plate (Cellstar) in PI uptake media containing 1X HBSS without phenol red, 20 mM HEPES, and 10% (vol/vol) heat-inactivated FBS. Cells were infected at an MOI=60 and control wells were treated with 1% Triton. After infection, cells were centrifuged at $290 \times g$ for 10 minutes following infection. 5 µM propidium iodide (PI, P3566, Invitrogen) diluted in PI uptake media was added to the cells. The plate was sealed with adhesive optical plate sealing film (Microseal, Bio-Rad) and placed in a Synergy H1 microplate reader (BioTek) pre-heated to 37°C. PI fluorescence was measured every 10 minutes for the indicated number of hours.

<u>Anthrax toxin-mediated delivery of bacterial ligands into organoids</u>

Organoids were seeded in 96-well plates with transparent bottom and opaque walls in 5µl Matrigel domes. After two days in organoid media, organoids were grown for an additional 3 days in differentiation media: DMEM/F12 supplemented with 20% murine R-Spondin1 supernatant, 10% murine Noggin supernatant, 50 ng/mL recombinant murine EGF (Fisher Scientific), 1X P/S, 1X L-Glu, 10 mM HEPES (HiMedia), 1X N2 (Life Technologies), 1X B27 (Life Technologies), and 1 mM N-acetylcysteine (Fisher Scientific). 5 µM DAPT was added for the last 24 hours. Differentiated organoids were then treated with 16µg/ml PA and 8µg/ml LFn-FlaA or 0.1 µg/ml LFn-PrgJ in differentiation media containing 10µg/ml propidium iodide for 4 hours. Total lysis wells were treated with 1%Triton.

<u>Expression of inflammasome genes in human small intestinal organoids</u>

To analyze inflammasome expression under different media conditions, spheroids were cultured in differentiation medium: DMEM/F12 supplemented with 20% supernatant from R-Spondin1 expressing L-cells, 10% supernatant from Noggin expressing cells, 50 ng/mL recombinant murine EGF (Fisher Scientific), 1X P/S, 1X L-Glu, 10 mM HEPES (HiMedia), 1X N2 (Life Technologies), 1X B27 (Life Technologies), 1 mM N-acetylcysteine (Fisher Scientific), and 5 µM DAPT. After 4 days of incubation at 37°C, Matrigel domes were dissolved in PBS-EDTA (5 mM) for 1 hour at 4°C on an orbital shaker. After centrifugation at 300 × g for 5 minutes at 4°C, the cell pellet was resuspended in TRIzol™ to analyze mRNA expression.

<u>Isolation of peripheral blood mononuclear cells (PBMCs)</u>

To compare expression levels of inflammasome components in human intestinal epithelial cells with immune cells, PBMC-derived cDNA was kindly provided by William Messer at Oregon Health & Science University. Briefly, PBMCs were isolated using density gradient centrifugation. After overlay of Lymphoprep™ (Alere Technologies AS) with blood mixed 1:2 with 1X PBS (pH 7.4, Gibco), the sample was centrifuged at 800 × g for 20 minutes at room temperature (RT). Residual erythrocytes were lysed with 1X RBC lysis buffer (10X, BioLegend), followed by three washing steps with fresh PBS for 10 minutes at 250 × g. Subsequently cells were harvested in TRIzol™ Reagent (Thermo Fisher Scientific) for mRNA analysis.

 RNA extraction, cDNA synthesis, and real-time quantitative polymerase chain reaction

(RT-qPCR) of organoid and PBMC samples

After thawing the TRIzol™ samples, chloroform was added and the tubes

centrifuged for 15 minutes at 12,000 × g at 4°C. The aqueous phase was transferred to

a new tube containing linear polyacrylamide (Gene-Elute™ LPA, Sigma). To allow RNA

precipitation, the samples were incubated with isopropanol for 10 minutes and

subsequently centrifuged for 10 minutes at 12,000 × g. After aspiration of the

supernatant, the RNA pellet was washed once with 75% ethanol for 5 minutes at 7,500

× g at 4°C. The supernatant was aspirated and the dried pellet resuspended in

Molecular Biology Grade Water (Corning). After determination of RNA content and

quality (260/280 and 260/230 ratios), 1 µg of RNA was reversely transcribed into cDNA.

Reaction steps were performed in a Biorad T100 Cycler. First, residual DNA was

removed using RQ1 RNAse-free DNase (Promega) in RQ1 DNase 1X Reaction buffer

(Promega) for 30 minutes at 37°C. After stopping the reaction with RQ1 DNase Stop

Solution (Promega) for 10 minutes at 65°C, Oligo dTs (Sigma) and dNTPs (Sigma) were

added for 5 minutes at 65°C. Reverse transcription was performed with SuperScript™

IV Reverse Transcriptase (Invitrogen), 5 mM DTT and SuperScript™ IV Reaction Buffer

(Invitrogen) for 10 minutes at 50-55°C. Subsequently the enzyme was inactivated at

80°C for 10 minutes.

To analyze RNA expression, cDNA was mixed with 10 µM forward and reverse

primers and PowerUp™ SYBR™ Green Master Mix (Applied Biosystems) according to

manufacturer's instructions. The following primers were used:

RPLP0   Forward: 5'- GGCACCATTGAAATCCTGAGTGATGTG -3'

269   Reverse: 5'- TTGCGGACACCCTCCAGGAAG -3'

270 *NAIP*   Forward: 5'- AAGCATCCGCCCAGCTCTTGA -3'

271   Reverse: 5'- TATTGCCCTCCAGATCCACAGACAGTTC -3'

272 *NLRC4*   Forward: 5'- CATAGTCAAGTCTCTGTCAAGTGAACCCTGT -3'

273   Reverse: 5'- GCTGTTCTAGCACGTTCATCCTGTCG -3'

274 *CASP1*   Forward: 5'- GAGGCATTTGCACACCGCCC -3'

275   Reverse: 5'- GGATCTCTTCACTTCCTGCCCACA -3'

Amplification was analyzed in real-time with StepOne Software v2.3. In brief,

samples were incubated for 10 minutes at 95°C, followed by 40 cycles of heating to

95°C for 15 seconds and cooling to 60°C for 1 minute. To monitor specificity of the run,

the melt curves were determined by keeping the samples at 95°C for 15 seconds,

cooling to 60°C for 1 min and then increasing the temperature every 15 seconds by

0.3°C up to 95°C. Expression levels relative to the housekeeping gene (*RPLP0*) were

calculated using the formula x = $2^{-\Delta ct}$.

<u>RNA extraction, cDNA synthesis, and RT-qPCR of Caco-2, T84, and THP-1 samples</u>

RNA was isolated using the RNeasy Plus Mini Kit (Qiagen) following the

manufacturer's instructions. Cells were lysed in 350 µL RLT buffer with β-

mercaptoethanol and centrifuged through a QIAshredder spin column (Qiagen). cDNA

was synthesized from isolated RNA using SuperScript II Reverse Transcriptase

(Invitrogen) following the manufacturer's protocol. Quantitative PCR was conducted with

the CFX96 real-time system from Bio-Rad using the SsoFast EvaGreen Supermix with

Low ROX (Bio-Rad). To calculate relative gene expression, mRNA levels of target

genes were normalized to housekeeping gene *HPRT* and the formula x = $2^{-\Delta CT}$ was

used. The following primers from PrimerBank (PrimerBank identification listed within

parentheses) were used (32–34):

HPRT (164518913c1)   Forward: 5'- CCTGGCGTCGTGATTAGTGAT -3'

296   Reverse: 5' - AGACGTTCAGTCCTGTCCATAA -3'

297 *NAIP* (119393877c3)   Forward: 5' - CCCATTAGACGATCACACCAGA -3'

298   Reverse: 5' - GGAGTCACTTCCGCAGAGG -3'

299 *NLRC4* (312433959c2)   Forward: 5' - TGCATCATTGAAGGGGAATCTG -3'

300   Reverse: 5' - GATTGTGCCAGGTATATCCAGG -3'

301 *NLRP3* (208879435c1)   Forward: 5' - GATCTTCGCTGCGATCAACAG -3'

302   Reverse: 5' - CGTGCATTATCTGAACCCCAC -3'

303 *PYCARD* (313482805c1)   Forward: 5' - TGGATGCTCTGTACGGGAAG -3'

304   Reverse: 5' - CCAGGCTGGTGTGAAACTGAA -3'

305 *CASP1* (380254454c1)   Forward: 5' - TTTCCGCAAGGTTCGATTTTCA -3'

306   Reverse: 5' - GGCATCTGCGCTCTACCATC -3'

<u>Inhibitor experiments</u>

Cells were treated 1 hour prior to infection at the indicated concentrations with

inhibitors: varying concentrations of MCC950 (Sigma Aldrich PZ0280), 20 μM of pan-

caspase inhibitor Z-VAD(OMe)-FMK (SM Biochemicals SMFMK001), 25 μM of

caspase-1 inhibitor Ac-YVAD-cmk (Sigma Aldrich SML0429), 30 μM of disulfiram

(Sigma).

<u>siRNA-mediated knockdown of genes</u>

The following Silencer Select siRNA oligos were purchased from Ambion (Life

Technologies): *CASP4* (s2412), *CASP5* (s2417), two Silencer Select negative control

siRNAs (Silencer Select Negative Control No. 1 siRNA and Silencer Select Negative

Control No. 2 siRNA). Three days before infection, 30 nM of siRNA was transfected into

Caco-2s using Lipofectamine RNAiMAX transfection reagent (Thermo Fisher Scientific)

following the manufacturer's protocol. Cells were primed with 400 ng/ml of Pam3CSK4

(Invivogen) for 3 hours prior to infection.

<u>Statistical analysis</u>

Prism 9.1.1 (GraphPad Software) was used to graph all data and for all statistical

analyses. Statistical significance for experiments was determined using the appropriate

test and are indicated in each figure legend. Differences were considered statistically

significant if the *p* value was <0.05.

Results

<u>*Salmonella* infection induces inflammasome activation in human intestinal epithelial</u>

<u>cells</u>

Once inside the host, *Salmonella* upregulates expression of its SPI-1 T3SS,

which delivers effectors that enable *Salmonella* to invade intestinal epithelial cells (2).

Human IECs infected with *Salmonella* undergo inflammasome activation at late time

points following infection (10hpi) (20, 22). However, early inflammasome responses to

SPI-1-expressing *Salmonella* have not been previously investigated. To test if human

IECs undergo early inflammasome activation in response to *Salmonella* grown under

SPI-1-inducing conditions (28), we infected WT Caco-2 cells, a human colorectal cell

line, with WT *Salmonella* (WT Stm) or *Salmonella* lacking its SPI-1 T3SS (Δ*sipB* Stm)

and assayed for subsequent inflammasome activation by measuring release and

cleavage of the inflammasome-dependent cytokine IL-18 at 6hpi, an early time point

following *Salmonella* infection (Fig. 1A, B). Cells infected with WT Stm released

significantly increased levels of cleaved IL-18 into the supernatant compared to mock-

infected cells (Fig. 1A, B). In contrast, cells infected with Δ*sipB* Stm, which is unable to

invade cells, failed to release cleaved IL-18 (Fig. 1A, B). We measured cell death as

another readout of inflammasome activation by assaying for uptake of the cell-

impermeable dye propidium iodide (PI), which enters cells as they form pores in their

plasma membrane and undergo cell death. In cells infected with WT Stm, PI uptake

began to occur between 4-6hpi, and gradually increased over time, indicating that

infected cells begin to undergo cell death as early as 4-6hpi. As expected, cells infected

with Δ*sipB* Stm, which cannot enter the host cell, did not uptake any PI. We found that

inflammasome activation in response to *Salmonella* also occurs in another human

colorectal cell line, T84 cells (Fig. 1D). T84 cells infected with SPI-1-expressing WT Stm

also released IL-18 into the supernatant at 6hpi (Fig. 1D). As a control, we found that in

both Caco-2 and T84 cells, the inflammasome-independent cytokine, IL-8, is released in

response to WT and Δ*sipB* Stm (Fig. S1A, S1B).

Inflammasome activation leads to cleavage of the pore-forming protein GSDMD

(5, 35). The N-terminal fragment of cleaved GSDMD inserts into the host cell plasma

membrane and oligomerizes to create a pore through which cellular components such

as cleaved IL-1 and IL-18 can be released (36–39). These pores eventually cause

rupture of the cell through osmotic flux, resulting in pyroptosis (3). To determine if pore

formation by GSDMD is required for release of IL-18 and PI uptake in human IECs

infected with *Salmonella*, we pretreated Caco-2 cells with disulfiram, which prevents

cleaved GSDMD from inserting into the plasma membrane and thus abrogates pore

formation (40). Treatment with disulfiram led to loss of IL-18 release and PI uptake

during *Salmonella* infection (Fig. 1E, 1F). Importantly, release of the inflammasome-

independent cytokine IL-8 was not affected by disulfiram treatment (Fig. S1C).

Collectively, these results suggest that as early as 4-6hpi, human IECs infected with WT

Stm grown under SPI-1-inducing conditions undergo inflammasome activation, and

GSDMD-mediated pore formation is required to observe IL-18 release and PI uptake in

response to *Salmonella* in human IECs.

<u>Bacterial T3SS ligands do not activate the inflammasome in human intestinal epithelial</u>

<u>cells</u>

We next sought to determine the bacterial ligands that trigger inflammasome

activation in human IECs. The *Salmonella* SPI-1 T3SS inner rod protein (PrgJ), the SPI-

1 T3SS needle protein (PrgI), and flagellin activate the NAIP/NLRC4 inflammasome in

both murine and human macrophages, (8, 11). Given that *Salmonella* grown under SPI-

1-inducing conditions triggered inflammasome activation in human IECs (Fig. 1), we

hypothesized that this inflammasome activation was due to NAIP-mediated recognition

of *Salmonella* T3SS ligands. First, we asked whether *Salmonella* T3SS ligands are

sufficient to activate the inflammasome in human IECs. We used the Gram-positive

bacterium *Listeria monocytogenes* to deliver *Salmonella* T3SS ligands into IECs, We

have previously used the *Listeria* system to deliver *Salmonella* T3SS ligands into the

cytosol of human macrophages to examine NAIP-dependent inflammasome activation

(17). In this system, ligands of interest are translationally fused to the N-terminus of

truncated ActA, enabling the ligands to be delivered into the host cell cytosol (17, 29).

Human macrophages infected with *Listeria* expressing the *Salmonella* SPI-1 T3SS inner

rod PrgJ or needle PrgI undergo robust inflammasome activation (17). Surprisingly,

IECs infected with *Listeria* expressing either the SPI-1 inner rod (PrgJ) or needle (PrgI)

proteins failed to induce IL-18 release (Fig. 2A, B; Fig. S2A, B). Polarized Caco-2 cells

and C2Bbe1 cells, a Caco-2 subtype, infected with *Listeria* expressing the SPI-1 inner

rod PrgJ also failed to release IL-18 levels above that observed in cells infected with WT

Lm (Fig. S2C, D). Collectively, these data indicate that *Listeria* delivery of bacterial

T3SS ligands is not sufficient to induce inflammasome activation in human IEC lines.

As we did not observe inflammasome activation by bacterial T3SS ligands

delivered using *Listeria*, we tested a second delivery method, the *Bacillus anthracis*

toxin system, to deliver these bacterial ligands into the cytosol of IECs (41). The anthrax

toxin delivery system contains two subunits: a protective antigen (PA) that creates a

pore in the host endosomal membrane and a truncated lethal factor (LFn) that is

delivered through the PA pore into the cytosol. T3SS ligands that are translationally

fused to the N-terminal domain of the *B. anthracis* LFn are delivered into the host cell

cytosol upon treatment with both PA and the LFn fusion (41) (collectively referred to as

Tox). We delivered the *Salmonella* SPI-1 T3SS inner rod protein (PrgJTox) into Caco-2

cells (Fig. S3A), polarized Caco-2 cells (Fig. S3B), C2Bbe1 cells (Fig. S3C), and T84

cells (Fig. S3D) and assayed the release of the inflammasome-dependent cytokine IL-

18. In all cell types, we failed to observe IL-18 secretion in response to PrgJTox that

was above the PA alone control (Fig. S3). Thus, our data indicate that human IEC lines

did not undergo inflammasome activation in response to bacterial T3SS ligands

delivered with the anthrax toxin system.

To determine if this absence of inflammasome responses to T3SS ligands was

limited to immortalized IECs or extended to non-immortalized IECs as well, we delivered

flagellin (FlaTox) or the SPI-1 T3SS inner rod PrgJ (PrgJTox) into human intestinal

organoids and measured levels of cell death (Fig. 2C). Both human small intestinal

organoids and colonoids failed to undergo cell death when treated with FlaTox or

PrgJTox (Fig. 2C). In contrast, murine organoids underwent robust cell death in

response to FlaTox or PrgJTox, as expected, indicating that these preparations of

FlaTox and PrgJTox had the expected biological activity, but were not able to activate

the inflammasome in primary or transformed human intestinal cells (Fig. 2C).

<u>Human intestinal epithelial cells express low levels of *NAIP* and *NLRC4* compared to</u>

<u>human myeloid cells</u>

424   Given the role of NAIP/NLRC4 in detecting and responding to *Salmonella* and

bacterial T3SS ligands in murine IECs as well as murine and human macrophages (7–

12, 14, 16, 17), our findings that human IECs do not respond to T3SS ligands were

surprising. Thus, we next asked if expression of *NAIP* and *NLRC4* in human IECs is

comparable in Caco-2 cells, T84 cells, and human THP-1 macrophages (Fig. 3A, B).

Consistent with their poor responsiveness to cytosolic delivery of NAIP ligands, both

Caco-2 and T84 cells had very low expression of *NAIP* and *NLRC4* mRNA compared to

human macrophages. Moreover, primary small intestinal organoids also expressed very

low levels of *NAIP* and *NLRC4* compared to human PBMCs, indicating that this low

expression was a general feature of human IECs (Fig. 3C, D). Collectively, these data

suggest that human IECs express very low levels of NAIP and NLRC4, and this may

partially explain why human IECs do not mount inflammasome responses to T3SS

ligands.

<u>NAIP is not required for inflammasome activation in response to *Salmonella* infection of</u>

<u>human intestinal cells</u>

440   In murine macrophages and IECS as well as human macrophages, NAIP/NLRC4

contributes to inflammasome responses during *Salmonella* infection (11, 13, 14, 42).

Even low levels of *NAIP* expression are sufficient to induce inflammasome responses in

human macrophages (17). To formally test if the inflammasome activation we observe

in human IECs infected with *Salmonella* (Fig. 1) requires NAIP, we used the Clustered

Regularly Interspersed Palindromic Repeat (CRISPR) system, in conjunction with the

RNA-guided exonuclease Cas9, to disrupt the *NAIP* gene in Caco-2 cells (Fig. S4). We

sequenced two independent single cell clones each of $NAIP^{-/-}$ Caco-2 cells ($NAIP^{-/-}$ #7,

$NAIP^{-/-}$ #8) to confirm appropriate targeting of *NAIP* in each line (Fig. S4). Caco-2 cells

are polyploid, and we therefore found multiple mutant alleles for each CRISPR clone we

sequenced (Fig. S4). For the two $NAIP^{-/-}$ clones, all of the changes resulted in a

premature stop codon (Fig. S4).

To test whether inflammasome activation during *Salmonella* infection of human

IECs requires NAIP, we infected WT or $NAIP^{-/-}$ Caco-2 cells with WT Stm or Δ*sipB* Stm

and assayed for inflammasome activation by measuring IL-18 release and PI uptake

(Fig. 4). As expected, WT Caco-2 cells infected with WT Stm released significantly

increased levels of IL-18 and underwent cell death (Fig. 4A, 4B) and this response was

dependent on the presence of the SPI-1 T3SS, as cells infected with Δ*sipB* Stm failed to

undergo inflammasome activation (Fig. 4A, 4B). In $NAIP^{-/-}$ Caco-2 cells infected with WT

Stm, we did not observe a decrease in inflammasome activation compared to WT Caco-

2 cells (Fig. 4A, 4B). Both WT and $NAIP^{-/-}$ Caco-2 cells secreted similar levels of the

inflammasome-independent cytokine IL-8 (Fig. 4C). Overall, these data indicate that

NAIP is not required for inflammasome responses to SPI-1-expressing *Salmonella* in

human IECs.

<u>NLRP3 and ASC are dispensable for inflammasome activation in response to</u>

<u>*Salmonella* in human intestinal epithelial cells</u>

Since the NAIP/NLRC4 inflammasome did not have a role in responding to

Salmonella grown under SPI-1-inducing conditions in human IECs, we sought to

determine whether another host cytosolic sensor could be responsible for

inflammasome activation. One candidate is the NLRP3 inflammasome, which can be

activated by a variety of stimuli during infection, including potassium efflux (18). In

murine macrophages, the NLRP3 inflammasome is activated late during *Salmonella*

infection (19). *Salmonella* infection also activates the NLRP3 inflammasome in human

macrophages (26, 43). To determine if the inflammasome activation we observed in

human IECs is dependent on the NLRP3 inflammasome, we infected WT Caco-2 cells

that were pre-treated with MCC950, a potent chemical inhibitor of the NLRP3

inflammasome (44) and measured IL-18 release (Fig. 5A). As expected, cells treated

with the DMSO control underwent robust inflammasome activation in response to WT

Stm infection (Fig. 5A). Interestingly, cells treated with varying concentrations of

MCC950 also exhibited similar levels of inflammasome activation as DMSO-control

treated cells (Fig. 5A), suggesting that the NLRP3 inflammasome is not required for

inflammasome responses to *Salmonella* in human IECs. We found that like *NAIP* and

NLRC4, mRNA expression of *NLRP3* is also very low in Caco-2 cells compared to THP-

1 macrophages (Fig. 5B), which may explain why NLRP3 inhibition does not prevent

inflammasome activation in human IECs.

In addition to the NAIP/NLRC4 and NLRP3 inflammasomes, there are many

other inflammasomes that can get activated in response to bacterial infections. The

majority of these inflammasomes, including AIM2, IFI16, NLRP3, NLRP6, NLRP7, and

pyrin, use an adaptor protein called ASC to recruit and activate downstream caspases

(4). To determine if ASC-dependent inflammasomes participate in the response to

Salmonella in human IECs, we tested if ASC is required for inflammasome activation.

Using CRISPR/Cas9, we disrupted *PYCARD*, the gene that encodes for ASC, in Caco-2

cells (Fig. S5). We sequence-validated two independent single cell clones of *PYCARD$^{-/-}$*

Caco-2 cells (*PYCARD$^{-/-}$* #4, *PYCARD$^{-/-}$* #6) (Fig. S5). Most of the mutations resulted in

a premature stop codon or the mutated protein bore no resemblance to the WT protein

sequence (Fig. S5). mRNA expression of *PYCARD* was also abrogated in the KO

clones relative to WT Caco-2 cells (Fig. S6A). We did not detect any ASC protein

expression even in WT Caco-2 cells (Fig. S6B).

499   We next infected WT or *PYCARD$^{-/-}$* Caco-2 cells with WT Stm or Δ*sipB* Stm and

assayed for inflammasome activation (Fig. 5C, D). As expected, WT Caco-2 cells

infected with WT Stm released significantly increased levels of IL-18 and underwent cell

death (Fig. 5C, D). This was dependent on the presence of the SPI-1 T3SS, as cells

infected with Δ*sipB* Stm failed to undergo inflammasome activation (Fig. 5C, D). In

PYCARD$^{-/-}$ Caco-2 cells infected with WT Stm, we did not observe a decrease in

inflammasome activation compared to WT Caco-2 cells, indicating that ASC is

dispensable for IL-18 release and cell death in Caco-2 cells in response to Stm. Both

WT and *PYCARD$^{-/-}$* Caco-2 cells exhibited similar levels of IL-8 release, an

inflammasome-independent cytokine (Fig. 5E). In addition, expression of *PYCARD*

mRNA in Caco-2 cells was very low compared to THP-1 macrophages (Fig. 5F).

Collectively, these data indicate that NLRP3 and ASC-dependent inflammasomes are

not required for inflammasome responses to *Salmonella* infection in human IECs.

<u>Caspase-1 is partially required for inflammasome activation in response to *Salmonella*</u>

<u>in human intestinal epithelial cells</u>

Inflammasomes recruit various caspases which can then cleave and process IL-1

and IL-18 cytokines and mediate pyroptosis (45). For example, the murine NAIP/NLRC4

inflammasome can recruit both caspase-1 and caspase-8 in response to *Salmonella*

infection of murine IECs (14). In human macrophages, NAIP/NLRC4, NLRP3, caspase-

1, and caspase-8 are recruited to the same macromolecular complex during *Salmonella*

infection (26). Notably, although expression of *CASP1* mRNA in Caco-2 cells is lower

than that in THP-1 macrophages (Fig. 6A), *CASP1* expression in Caco-2 cells is still

higher than expression of other inflammasome genes we have assessed so far

(compare axes in Fig. 6A to Fig. 3A, 3B, 5B, 5F). Similarly, expression of *CASP1* in

small intestinal organoids, although lower than that observed in human PBMCs, is

higher than expression of other inflammasome-related genes (compare axes in Fig. 6B

to Fig. 3C, 3D). This suggests that caspase-1 may be an important contributor to

inflammasome responses in IECs.

To test the contribution of caspases to inflammasome responses during

Salmonella infection in human IECs, we pretreated Caco-2 cells with pharmacological

inhibitors targeting caspases of interest (ZVAD: pan-caspase inhibitor or YVAD: inhibitor

for caspase-1). We then infected cells with WT Stm or Δ*sipB* Stm and assayed for

inflammasome activation by measuring IL-18 release (Fig. 6C). DMSO-treated cells

infected with WT Stm released IL-18, whereas cells treated with the inhibitors had a

significant defect in IL-18 release. Treatment with ZVAD, the pan-caspase inhibitor,

resulted in a lower level of IL-18 release compared to treatment with the caspase-1

inhibitor YVAD, suggesting that in addition to caspase-1, other caspases are also

important during *Salmonella* infection of human IECs. As expected, cells infected with

Δ*sipB* Stm demonstrated no inflammasome activation regardless of inhibitor treatment.

These results suggest that caspase-1 contributes to inflammasome responses to

Salmonella in human IECs.

Since pharmacological inhibitors preferentially targeting individual caspases can

have cross-reactivity with other caspases, we used CRISPR/Cas9 to disrupt *CASP1* in

Caco-2 cells (Fig. S7, S8). We sequenced and validated two independent single cell

clones of *CASP1*⁻/⁻ Caco-2 cells (*CASP1*⁻/⁻ #7, *CASP1*⁻/⁻ #12) (Fig. S7, S8). All mutations

resulted in a premature stop codon (Fig. S7, S8). We further confirmed decreased

mRNA expression of *CASP1* in *CASP1*⁻/⁻ #7 Caco-2 cells (Fig. S7B). We were unable to

detect protein expression in both WT Caco-2 cells and *CASP1*⁻/⁻ #7 Caco- 2 cells by

western blot (Fig. S7C), indicating that caspase-1 is poorly expressed in Caco-2 cells.

To test whether inflammasome activation during *Salmonella* infection requires

caspase-1 in human IECs, we infected WT or *CASP1*⁻/⁻ Caco-2 cells with WT or Δ*sipB*

Stm and assayed for subsequent inflammasome activation (Fig. 6D, E). As expected,

WT Caco-2 cells infected with WT Stm released significant levels of IL-18 and

underwent cell death (Fig. 6D, E). This response was dependent on the presence of the

SPI-1 T3SS, as cells infected with Δ*sipB* Stm failed to undergo inflammasome activation

(Fig. 6D). Consistent with YVAD inhibitor treatment, *CASP1*⁻/⁻ Caco-2 cells infected with

WT Stm showed a statistically significant decrease in, but not complete abrogation of,

IL-18 release at 6hpi (Fig. 6D). There was also a slight delay in uptake of PI in $CASP1^{-/-}$

#7 Caco-2 cells relative to WT Caco-2 cells (Fig. 6E). In contrast, both WT and KO

Caco-2 cells exhibited similar levels of IL-8 release, an inflammasome-independent

cytokine (Fig. 6F). Overall, these data suggest that caspase-1 is partially required for

the inflammasome response to *Salmonella* infection in human IECs.

<u>Caspase-4 is required for inflammasome responses to *Salmonella* in human intestinal</u>

<u>epithelial cells</u>

In mice, in addition to caspase-1, a second caspase, caspase-11, responds to

Salmonella infection (19, 21). Caspase-11 and its human orthologs caspases-4/5 detect

cytosolic LPS and form the noncanonical inflammasome (24, 25). Caspases-4/5 are

important for late inflammasome responses to *Salmonella* infection of human intestinal

epithelial cells, specifically when *Salmonella* escapes to the cytoplasm from its

Salmonella-containing vacuole (SCV) (20). To test the contribution of the caspase-4/5

inflammasome earlier during infection with *Salmonella* grown under SPI-1-inducing

conditions, we transfected Caco-2 cells with siRNAs targeting *CASP4, CASP5* or both.

We then infected the cells with WT or Δ*sipB* Stm and assayed for inflammasome

activation by measuring IL-18 release (Fig. 7A). As expected, cells treated with a control

scrambled siRNA exhibited IL-18 secretion upon infection with WT Stm, but failed to

undergo IL-18 secretion when infected with Δ*sipB* Stm (Fig. 7A). However, knockdown

of *CASP4*, either alone, or in conjunction with *CASP5*, resulted in nearly complete

abrogation of IL-18 secretion in cells infected with WT Stm compared to control siRNA-

treated cells, suggesting that caspase-4 is required for inflammasome activation (Fig.

7A). Knockdown of *CASP5* alone resulted in a partial and significant decrease in IL-18

secretion, indicating that while caspase-5 may be playing a role, it is not absolutely

required (Fig. 7A). We observed moderately high (67 – 75%) siRNA-mediated

knockdown efficiencies (Fig. S9A). Release of the inflammasome-independent cytokine

IL-8 was comparable across conditions (Fig. S9B).

To definitively test the requirement of caspase-4 in early inflammasome

responses to SPI-1-expressing *Salmonella*, we disrupted *CASP4* in Caco-2 cells using

CRISPR/Cas9 (Fig. S10, S11). We sequenced and validated two independent single

cell clones of *CASP4*$^{-/-}$ Caco-2 cells (*CASP4*$^{-/-}$ #3, *CASP4*$^{-/-}$ #4) (Fig. S10). All mutations

resulted in a premature stop codon (Fig. S10). Both clones exhibited decreased mRNA

expression of *CASP4* and no protein expression relative to WT Caco- 2 cells (Fig. S11).

We then infected WT or *CASP4*$^{-/-}$ Caco-2 cells with WT or Δ*sipB* Stm and

assayed for subsequent IL-18 secretion and cell death as readouts for inflammasome

activation (Fig. 7B – E). As expected, WT Caco-2 cells infected with WT Stm released

significant levels of cleaved IL-18 and underwent cell death (Fig. 7B – E), whereas cells

infected with Δ*sipB* Stm failed to release IL-18 (Fig. 7B). Consistent with our findings

with siRNA knockdown of *CASP4*, *CASP4*$^{-/-}$ Caco-2 cells infected with WT Stm failed to

release IL-18 and exhibited a delay in PI uptake at 4-6 hpi (Fig. 7B – E). Interestingly,

CASP4$^{-/-}$ Caco-2 cells began to undergo cell death later in infection, suggesting that

while caspase-4 is required for inducing cell death early in infection, it may not be

absolutely required at later timepoints of infection. Like WT cells, *CASP4*$^{-/-}$ cells still

released substantial levels of the inflammasome-independent cytokine IL-8 following

infection (Fig. S9C). Collectively, these data suggest that caspase-4 is required for early

inflammasome responses to SPI-1-expressing *Salmonella* infection in human IECs.

Discussion

<u>Discussion</u>

In this study, we have demonstrated that Caco-2 cells undergo inflammasome

activation in response to early infection with SPI-1-expressing *Salmonella* that requires

the presence of the SPI-1 T3SS, and that GSDMD-mediated pore formation is required

for release of IL-18 and cell death (Fig. 1, S1). Unexpectedly however, individual

bacterial T3SS ligands or flagellin were not sufficient to induce inflammasome activation

in Caco-2 cells or intestinal organoids (Fig. 2, S2, S3). Additionally, we found that

neither the NAIP nor NLRP3 canonical inflammasomes (Fig. 4, S4, 5), are required for

inflammasome responses during *Salmonella* infection, which may be due to low mRNA

expression of *NAIP*, *NLRC4*, and *NLRP3* in human IECs and human intestinal

organoids (Fig. 3, 5). Moreover, we found that ASC, a shared adaptor protein that

several inflammasomes use to recruit caspases, is also not required for inflammasome

activation (Fig. 5, S5, S6), suggesting that inflammasomes that require ASC for their

function are likely not playing a role in human IECs during *Salmonella* infection.

Interestingly, we found that caspase-1 is partially required for inflammasome activation

in human IECs (Fig. 6, S7, S8), leaving the unanswered question of what upstream

sensor is activating caspase-1. Finally, we found that caspase-4 is necessary for early

inflammasome responses to SPI-1-expressing *Salmonella* (Fig. 7, S9 – S11).

NAIP/NLRC4 inflammasome activation in IECs is critical for control of *Salmonella*

infection in mice (13–15). It was therefore surprising that NAIP was dispensable in

human IECs during *Salmonella* infection (Fig. 3, S3 – S6). Perhaps a lack of robust

NAIP/NLRC4 inflammasome responses in human IECs partially underlies why some

intestinal bacterial pathogens with T3SSs, such as *Shigella flexneri,* enterohemorrhagic

Escherichia coli, and enteropathogenic *E. coli,* can cause disease in humans but not

mice. Indeed, mice are normally resistant to *S. flexneri,* but mice lacking the

NAIP/NLRC4 inflammasome can be robustly colonized by *S. flexneri* and develop a

disease resembling human shigellosis (46).

The lack of NAIP activation may be due to low expression of *NAIP* and *NLRC4*

mRNA in human IEC lines as well as human small intestinal organoids (Fig. 3). Single

cell transcriptome analysis of cells intestinal cells from human donors demonstrate that

only a small subset of intestinal cells express *NAIP* at detectable levels, while *NLRC4*

expression is below the limit of detection (47). However, it is possible that a population

of IECs exist that express *NAIP* and *NLRC4* at low levels that may be below the limit of

detection of single-cell transcriptome analysis, but may be sufficient to activate the

inflammasome in response to infection under physiological conditions. Indeed, single

cell transcriptome analysis of murine small intestines reveal that expression of *NLRC4*

is also very low in murine IECs (48), but murine IECs undergo robust NAIP/NLRC4

inflammasome activation despite this relatively low expression (13–15). Despite our

findings that NAIP is dispensable in Caco-2 cells to respond to *Salmonella* infection and

human intestinal organoids do not mount inflammasome responses to T3SS ligands, it

is possible that under physiological conditions, this inflammasome still plays an

important role during bacterial infection or disease pathogenesis. Indeed, human

patients with activating NLRC4 mutations exhibit gastrointestinal symptoms (49–52).

A recent study comparing inflammasome responses of human and murine IECs

during *Salmonella* infection reported that while caspase-1 was required for

inflammasome activation in murine IECs, it was dispensable in late inflammasome

responses to *Salmonella* in human IECs (22). We found caspase-1 to be partially

required for inflammasome activation in IECs infected with *Salmonella* (Fig. 6, S7, S8).

Differences in our experimental conditions likely explain this difference in findings. Our

Salmonella was grown under SPI-1-inducing conditions, and we assayed for

inflammasome activation at a slightly earlier timepoint. Perhaps caspase-1 plays a role

in early inflammasome responses to SPI-1-expressing *Salmonella* in human IECs, and

it is less important at later timepoints when *Salmonella* has shifted to expressing its SPI-

2 T3SS.

The identity of the host sensor that is acting upstream of caspase-1 in human

IECs remains unknown. Humans have 22 NLRs that could serve as potential cytosolic

sensors of bacterial structures and activity (53). Most of these NLRs do not contain a

CARD domain, and they would therefore require an adaptor protein such as ASC to

recruit caspases. Given our finding that ASC is not required for inflammasome

activation in human IECs (Fig. 5, S5, S6), it is likely that the putative host sensor

upstream of caspase-1 contains its own CARD domain. Alternatively, it may interact

with a different CARD domain-containing adaptor protein than ASC. Future studies will

focus on identifying host factors that different caspases interact with during *Salmonella*

infection of human IECs.

In agreement with previous studies (20, 22), we found caspase-4 to be required

for inflammasome activation in response to *Salmonella* in human IECs (Fig 7, S9 –

S11). Importantly, caspase-4 activation restricts bacterial replication, and specifically

restricts replication of a subpopulation of cytosolic *Salmonella* (20, 22). It is unclear

whether *Salmonella*'s access into the cytosol is mediated by the bacteria or host, and

future studies will explore the mechanisms by which cytosolic populations of *Salmonella*

arise and how they influence inflammasome activation in human IECs.

In mice, inflammasome activation mediates control of *Salmonella* by restricting

bacterial replication, extruding infected cells, and preventing systemic dissemination of

Salmonella (13, 15, 20). Human IECs have been shown to undergo extrusion, but

whether this occurs in a caspase-4-dependent manner remains unknown (54). While we

know that human IECs and enteroids restrict bacterial replication in a caspase-4-

dependent manner (20, 22), it remains unknown if inflammasome activation results in

other mechanisms of control of infection. In mice, IL-18 release during *Salmonella*

infection recruits natural killer (NK) cells that are critical for early mucosal inflammatory

responses (55). We observe robust release of the inflammasome-dependent cytokine

IL-18, but the downstream role of this cytokine in the human intestine during *Salmonella*

infection has not been explored. Additionally, while our study focused exclusively on

Salmonella infection, it is worth exploring if enteric pathogens with similar lifestyles to

Salmonella, such as *Shigella*, or ones with different lifestyles, such as the extracellular

pathogen *Yersinia*, elicit similar responses. Future studies that interrogate downstream

consequences of inflammasome activation in human IECs in response to various

enteric pathogens could shed light on human mucosal inflammatory responses to

bacterial pathogens.

Overall, our data indicate that *Salmonella* infection of human IECs triggers

inflammasome pathways that are distinct from that in mice. Pathways that are activated

in mice and important for control of infection, such as the NAIP and NLRP3

inflammasomes, were unexpectedly not required for inflammasome responses in

human IECs under the conditions we investigated. Instead, inflammasome responses in

human IECs required caspases-1 and -4. Our findings provide a foundation for future

studies aimed at uncovering the relative contribution of different caspases, and the

downstream responses that they mediate in human IECs.

<u>Acknowledgements</u>

We thank members of Igor Brodsky's and Sunny Shin's laboratories for scientific

discussion. We thank Meghan Wynosky-Dolfi for technical advice. We thank Russell

Vance, Randilea Nichols, and Jeannette Tenthorey for providing anthrax toxin-based

reagents, and JD Sauer for providing the *Listeria* strains and constructs for generating

ActA fusion proteins. We thank Jared Fisher for providing spheroids of human

duodenum and colon, and William Messer and Zoe Lyski for providing cDNA of human

PBMCs. Work in the Shin laboratory is supported by NIH/NIAID grants AI118861 and

AI123243, and the Linda Pechenik Montague Investigator Award from the University of

Pennsylvania Perelman School of Medicine. Work in the Brodsky laboratory is

supported by NIH/NIAID grants AI128520 and AI139102. S.S. and I.B. are both

recipients of the Burroughs-Wellcome Fund Investigators in the Pathogenesis of

Infectious Disease Award. I.R. is supported by OHSU start-up funding. N.N. is a

recipient of the American Heart Association Predoctoral Fellowship 19PRE34380315.

R.B. was supported by the FWF-Immunity in Cancer & Allergy PhD program of the

University of Salzburg and the Austrian Marshall Plan Foundation Scholarship. J.Z. is a

recipient of the NIH/NIAID Microbial Pathogenesis and Genomics training grant

5T32AI141393-03.

References

1.   Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra

RM, Studies IC on ED "Burden of I. 2010. The global burden of nontyphoidal Salmonella

gastroenteritis. Clinical infectious diseases 50:882–889.

2.   Crowley SM, Knodler LA, Vallance BA. 2016. Salmonella and the inflammasome: battle for

intracellular dominance. Inflammasome signaling and bacterial infections 43–67.

3.   Lamkanfi M, Dixit VM. 2014. Mechanisms and functions of inflammasomes. Cell 157:1013–

22.

4.   Zheng D, Liwinski T, Elinav E. 2020. Inflammasome activation and regulation: toward a

better understanding of complex mechanisms. Cell Discov 6:36.

5.   Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F. 2015.

Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. Nature,

2015/09/16 ed. 526:660–5.

6.   Raupach B, Peuschel S-K, Monack DM, Zychlinsky A. 2006. Caspase-1-Mediated Activation

of Interleukin-1β (IL-1β) and IL-18 Contributes to Innate Immune Defenses against

Salmonella enterica Serovar Typhimurium Infection. IAI 74:4922–4926.

7.   Kofoed EM, Vance RE. 2011. Innate immune recognition of bacterial ligands by NAIPs

determines inflammasome specificity. Nature 477:592–595.

8. Rauch I, Tenthorey JL, Nichols RD, Al Moussawi K, Kang JJ, Kang C, Kazmierczak BI, Vance RE.

2016. NAIP proteins are required for cytosolic detection of specific bacterial ligands in vivo.

Journal of Experimental Medicine 213:657–665.

9. Rayamajhi M, Zak DE, Chavarria-Smith J, Vance RE, Miao EA. 2013. Cutting edge: Mouse

NAIP1 detects the type III secretion system needle protein. The Journal of Immunology

191:3986–3989.

10. Yang J, Zhao Y, Shi J, Shao F. 2013. Human NAIP and mouse NAIP1 recognize bacterial type

III secretion needle protein for inflammasome activation. Proceedings of the National

Academy of Sciences 110:14408–14413.

11. Zhao Y, Shi J, Shi X, Wang Y, Wang F, Shao F. 2016. Genetic functions of the NAIP family of

inflammasome receptors for bacterial ligands in mice. Journal of Experimental Medicine

213:647–656.

12. Zhao Y, Yang J, Shi J, Gong Y-N, Lu Q, Xu H, Liu L, Shao F. 2011. The NLRC4 inflammasome

receptors for bacterial flagellin and type III secretion apparatus. Nature 477:596–600.

13. Sellin ME, Müller AA, Felmy B, Dolowschiak T, Diard M, Tardivel A, Maslowski KM, Hardt W-

D. 2014. Epithelium-intrinsic NAIP/NLRC4 inflammasome drives infected enterocyte

expulsion to restrict Salmonella replication in the intestinal mucosa. Cell host & microbe

16:237–248.

14. Rauch I, Deets KA, Ji DX, von Moltke J, Tenthorey JL, Lee AY, Philip NH, Ayres JS, Brodsky IE,

Gronert K. 2017. NAIP-NLRC4 inflammasomes coordinate intestinal epithelial cell expulsion

with eicosanoid and IL-18 release via activation of caspase-1 and-8. Immunity 46:649–659.

15. Hausmann A, Böck D, Geiser P, Berthold DL, Fattinger SA, Furter M, Bouman JA, Barthel-

Scherrer M, Lang CM, Bakkeren E. 2020. Intestinal epithelial NAIP/NLRC4 restricts systemic

dissemination of the adapted pathogen Salmonella Typhimurium due to site-specific

bacterial PAMP expression. Mucosal immunology 13:530–544.

16. Kortmann J, Brubaker SW, Monack DM. 2015. Cutting edge: inflammasome activation in

primary human macrophages is dependent on flagellin. The Journal of Immunology

195:815–819.

17. Reyes Ruiz VM, Ramirez J, Naseer N, Palacio NM, Siddarthan IJ, Yan BM, Boyer MA,

Pensinger DA, Sauer J-D, Shin S. 2017. Broad detection of bacterial type III secretion system

and flagellin proteins by the human NAIP/NLRC4 inflammasome. Proceedings of the

National Academy of Sciences 114:13242–13247.

18. Muñoz-Planillo R, Kuffa P, Martínez-Colón G, Smith BL, Rajendiran TM, Núñez G. 2013. K^+

efflux is the common trigger of NLRP3 inflammasome activation by bacterial toxins and

particulate matter. Immunity 38:1142–53.

19. Broz P, Newton K, Lamkanfi M, Mariathasan S, Dixit VM, Monack DM. 2010. Redundant

roles for inflammasome receptors NLRP3 and NLRC4 in host defense against Salmonella.

Journal of Experimental Medicine 207:1745–1755.

20. Knodler LA, Crowley SM, Sham HP, Yang H, Wrande M, Ma C, Ernst RK, Steele-Mortimer O,

Celli J, Vallance BA. 2014. Noncanonical inflammasome activation of caspase-4/caspase-11

mediates epithelial defenses against enteric bacterial pathogens. Cell host & microbe

16:249–256.

21. Crowley SM, Han X, Allaire JM, Stahl M, Rauch I, Knodler LA, Vallance BA. 2020. Intestinal

restriction of Salmonella Typhimurium requires caspase-1 and caspase-11 epithelial intrinsic

inflammasomes. PLoS Pathog 16:e1008498.

22. Holly MK, Han X, Zhao EJ, Crowley SM, Allaire JM, Knodler LA, Vallance BA, Smith JG. 2020.

Salmonella enterica Infection of Murine and Human Enteroid-Derived Monolayers Elicits

Differential Activation of Epithelium-Intrinsic Inflammasomes. Infect Immun 88:e00017-20,

/iai/88/7/IAI.00017-20.atom.

23. Broz P, Ruby T, Belhocine K, Bouley DM, Kayagaki N, Dixit VM, Monack DM. 2012. Caspase-

11 increases susceptibility to Salmonella infection in the absence of caspase-1. Nature

490:288–291.

24. Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, Hu L, Shao F. 2014. Inflammatory caspases are

innate immune receptors for intracellular LPS. Nature 514:187–192.

25. Schmid-Burgk JL, Gaidt MM, Schmidt T, Ebert TS, Bartok E, Hornung V. 2015. Caspase-4

mediates non-canonical activation of the NLRP3 inflammasome in human myeloid cells.

European journal of immunology 45:2911–2917.

26. Man SM, Hopkins LJ, Nugent E, Cox S, Glück IM, Tourlomousis P, Wright JA, Cicuta P, Monie

TP, Bryant CE. 2014. Inflammasome activation causes dual recruitment of NLRC4 and NLRP3

to the same macromolecular complex. Proceedings of the National Academy of Sciences

111:7403–7408.

27. Lawley TD, Chan K, Thompson LJ, Kim CC, Govoni GR, Monack DM. 2006. Genome-wide

screen for Salmonella genes required for long-term systemic infection of the mouse. PLoS

Pathog 2:e11.

28. Lee CA, Falkow S. 1990. The ability of Salmonella to enter mammalian cells is affected by

bacterial growth state. Proceedings of the national academy of sciences 87:4304–4308.

29. Sauer J-D, Pereyre S, Archer KA, Burke TP, Hanson B, Lauer P, Portnoy DA. 2011. Listeria

monocytogenes engineered to activate the Nlrc4 inflammasome are severely attenuated

and are poor inducers of protective immunity. Proceedings of the National Academy of

Sciences 108:12419–12424.

30. Lauer P, Chow MYN, Loessner MJ, Portnoy DA, Calendar R. 2002. Construction,

characterization, and use of two Listeria monocytogenes site-specific phage integration

vectors. Journal of bacteriology 184:4177–4186.

31. Miyoshi H, Stappenbeck TS. 2013. In vitro expansion and genetic modification of

gastrointestinal stem cells in spheroid culture. Nat Protoc 8:2471–2482.

32. Spandidos A, Wang X, Wang H, Dragnev S, Thurber T, Seed B. 2008. A comprehensive

collection of experimentally validated primers for Polymerase Chain Reaction quantitation

of murine transcript abundance. BMC Genomics 9:633.

33. Spandidos A, Wang X, Wang H, Seed B. 2010. PrimerBank: a resource of human and mouse

PCR primer pairs for gene expression detection and quantification. Nucleic Acids Research

38:D792–D799.

34. Wang X. 2003. A PCR primer bank for quantitative gene expression analysis. Nucleic Acids

Research 31:154e–1154.

35. Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T, Haley B,

Roose-Girma M, Phung QT, Liu PS, Lill JR, Li H, Wu J, Kummerfeld S, Zhang J, Lee WP, Snipas

SJ, Salvesen GS, Morris LX, Fitzgerald L, Zhang Y, Bertram EM, Goodnow CC, Dixit VM. 2015.

Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. Nature,

2015/09/16 ed. 526:666–71.

36. Ding J, Wang K, Liu W, She Y, Sun Q, Shi J, Sun H, Wang D-C, Shao F. 2016. Pore-forming

activity and structural autoinhibition of the gasdermin family. Nature 535:111–116.

37. Sborgi L, Rühl S, Mulvihill E, Pipercevic J, Heilig R, Stahlberg H, Farady CJ, Müller DJ, Broz P,

Hiller S. 2016. GSDMD membrane pore formation constitutes the mechanism of pyroptotic

cell death. EMBO J 35:1766–1778.

38. Liu X, Zhang Z, Ruan J, Pan Y, Magupalli VG, Wu H, Lieberman J. 2016. Inflammasome-

activated gasdermin D causes pyroptosis by forming membrane pores. Nature 535:153–

158.

39. He W, Wan H, Hu L, Chen P, Wang X, Huang Z, Yang Z-H, Zhong C-Q, Han J. 2015. Gasdermin

D is an executor of pyroptosis and required for interleukin-1β secretion. Cell Res 25:1285–

1298.

40. Hu JJ, Liu X, Xia S, Zhang Z, Zhang Y, Zhao J, Ruan J, Luo X, Lou X, Bai Y, Wang J,

Hollingsworth LR, Magupalli VG, Zhao L, Luo HR, Kim J, Lieberman J, Wu H. 2020. FDA-

approved disulfiram inhibits pyroptosis by blocking gasdermin D pore formation. Nat

Immunol 21:736–745.

41. von Moltke J, Trinidad NJ, Moayeri M, Kintzer AF, Wang SB, van Rooijen N, Brown CR,

Krantz BA, Leppla SH, Gronert K. 2012. Rapid induction of inflammatory lipid mediators by

the inflammasome in vivo. Nature 490:107–111.

42. Miao EA, Mao DP, Yudkovsky N, Bonneau R, Lorang CG, Warren SE, Leaf IA, Aderem A.

2010. Innate immune detection of the type III secretion apparatus through the NLRC4

inflammasome. Proceedings of the National Academy of Sciences 107:3076–3080.

43. Gram AM, Wright JA, Pickering RJ, Lam NL, Booty LM, Webster SJ, Bryant CE. 2021.

Salmonella Flagellin Activates NAIP/NLRC4 and Canonical NLRP3 Inflammasomes in Human

Macrophages. The Journal of Immunology 206:631–640.

44. Coll RC, Robertson AA, Chae JJ, Higgins SC, Muñoz-Planillo R, Inserra MC, Vetter I, Dungan

LS, Monks BG, Stutz A. 2015. A small-molecule inhibitor of the NLRP3 inflammasome for the

treatment of inflammatory diseases. Nature medicine 21:248–255.

45. von Moltke J, Ayres JS, Kofoed EM, Chavarría-Smith J, Vance RE. 2013. Recognition of

Bacteria by Inflammasomes. Annu Rev Immunol 31:73–106.

46. Mitchell PS, Roncaioli JL, Turcotte EA, Goers L, Chavez RA, Lee AY, Lesser CF, Rauch I, Vance

RE. 2020. NAIP–NLRC4-deficient mice are susceptible to shigellosis. eLife 9:e59022.

47. Wang Y, Song W, Wang J, Wang T, Xiong X, Qi Z, Fu W, Yang X, Chen Y-G. 2020. Single-cell

transcriptome analysis reveals differential nutrient absorption functions in human intestine.

Journal of Experimental Medicine 217:e20191130.

48. Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, Burgin G, Delorey TM, Howitt

MR, Katz Y, Tirosh I, Beyaz S, Dionne D, Zhang M, Raychowdhury R, Garrett WS, Rozenblatt-

Rosen O, Shi HN, Yilmaz O, Xavier RJ, Regev A. 2017. A single-cell survey of the small

intestinal epithelium. Nature 551:333–339.

49. Canna SW, de Jesus AA, Gouni S, Brooks SR, Marrero B, Liu Y, DiMattia MA, Zaal KJM,

Sanchez GAM, Kim H, Chapelle D, Plass N, Huang Y, Villarino AV, Biancotto A, Fleisher TA,

Duncan JA, O'Shea JJ, Benseler S, Grom A, Deng Z, Laxer RM, Goldbach-Mansky R. 2014. An

activating NLRC4 inflammasome mutation causes autoinflammation with recurrent

macrophage activation syndrome. Nat Genet 46:1140–1146.

50. Kitamura A, Sasaki Y, Abe T, Kano H, Yasutomo K. 2014. An inherited mutation in NLRC4

causes autoinflammation in human and mice. Journal of Experimental Medicine 211:2385–

2396.

51. Romberg N, Al Moussawi K, Nelson-Williams C, Stiegler AL, Loring E, Choi M, Overton J,

Meffre E, Khokha MK, Huttner AJ, West B, Podoltsev NA, Boggon TJ, Kazmierczak BI, Lifton

RP. 2014. Mutation of NLRC4 causes a syndrome of enterocolitis and autoinflammation. Nat

Genet 46:1135–1139.

52. Steiner A, Reygaerts T, Pontillo A, Ceccherini I, Moecking J, Moghaddas F, Davidson S, Caroli

F, Grossi A, Castro FFM, Kalil J, Gohr FN, Schmidt FI, Bartok E, Zillinger T, Hartmann G, Geyer

M, Gattorno M, Mendonça LO, Masters SL. 2021. Recessive NLRC4-Autoinflammatory

Disease Reveals an Ulcerative Colitis Locus. J Clin Immunol https://doi.org/10.1007/s10875-

021-01175-4.

53. Chen G, Shaw MH, Kim Y-G, Nuñez G. 2009. NOD-Like Receptors: Role in Innate Immunity

and Inflammatory Disease. Annu Rev Pathol Mech Dis 4:365–398.

54. Knodler LA, Vallance BA, Celli J, Winfree S, Hansen B, Montero M, Steele-Mortimer O. 2010.

Dissemination of invasive *Salmonella* via bacterial-induced extrusion of mucosal epithelia.

Proc Natl Acad Sci USA 107:17733–17738.

55. Müller AA, Dolowschiak T, Sellin ME, Felmy B, Verbree C, Gadient S, Westermann AJ, Vogel

J, LeibundGut-Landmann S, Hardt W-D. 2016. An NK Cell Perforin Response Elicited via IL-18

Controls Mucosal Inflammation Kinetics during Salmonella Gut Infection. PLoS Pathog

12:e1005723.

<u>Figures</u>

Figure 1: *Salmonella* infection induces inflammasome activation in human

intestinal epithelial cells. Caco-2 cells (A – C, E, F) or T84 cells (D) were infected with

PBS (Mock), WT *S*. Typhimurium, or Δ*sipB S*. Typhimurium. (A, D) Release of IL-18

into the supernatant was measured by ELISA at 6hpi. (B) Lysates and supernatants

collected 6hpi were immunoblotted for IL-18 and β-actin. (C) Cell death was measured

as percentage uptake of propidium iodide, normalized to cells treated with 1% Triton.

(E, F) Caco-2 cells were treated with 30 μM disulfiram or DMSO as a vehicle control 1

hour prior to infection. Cells were then infected with PBS (Mock), WT *S.* Typhimurium*,*

or Δ*sipB S*. Typhimurium. (E) Release of IL-18 into the supernatant was measured by

ELISA at 6hpi. (F) Cell death as percentage uptake of propidium iodide, normalized to

cells treated with 1% Triton. ns – not significant, **** $p < 0.0001$ by Dunnett's multiple

comparisons test (A), or by unpaired t-test (C, D, F) or by Šídák's multiple comparisons

test (E). Error bars represent the standard deviation or the standard error of the mean

(SEM) (C, F) of triplicate wells from one experiment. Data shown are representative of

at least three independent experiments.

Figure 2: Bacterial T3SS ligands do not activate the inflammasome in human intestinal epithelial cells. (A, B) Caco-2 (A) or T84 cells (B) were primed for 3h with 100 ng/ml of Pam3CSK4 and infected with PBS (Mock), WT *L. monocytogenes* (WT Lm), or *L. monocytogenes* expressing *S.* Typhimurium SPI-1 inner rod (Lm + PrgJ), or SPI-1 needle (Lm + PrgI), at an MOI of 100. Release of IL-18 into the supernatant was measured by ELISA at 16hpi. L.O.D indicates the limit of detection of the assay. (C) Differentiated intestinal organoids were treated with FlaTox (PA + LFn-Fla) or Inner Rod Tox (PA + LFn-Rod) in media containing propidium iodide for 4h. Cell death was

measured as percentage uptake of propidium iodide, normalized to organoids treated

with 1% Triton. ns – not significant, * $p < 0.05$, **** $p < 0.0001$ by Dunnett's multiple

comparisons test. Error bars represent the standard deviation of triplicate wells from

one experiment. Data shown are representative of at least three independent

experiments.

Figure 3: Human intestinal epithelial cells express low levels of NAIP/NLRC4

compared to human myeloid cells. Relative mRNA expression of *NAIP* and *NLRC4*

compared to the housekeeping control *HPRT* or *RPLP0* as measured by qRT-PCR in

(A, B) Caco-2 cells, T84 cells, and THP-1 macrophages, and in (C, D) human peripheral

blood mononuclear cells (PBMCs) and human small intestinal organoids. Error bars

represent the standard deviation of multiple wells. Data shown are representative of at

least three independent experiments.

Figure 4: NAIP is not required for inflammasome responses to *Salmonella* in

human intestinal epithelial cells. WT or two independent single cell clones of *NAIP*$^{-/-}$

Caco-2 cells were infected with PBS (Mock), WT *S*. Typhimurium, or Δ*sipB S*.

Typhimurium for 6hrs. (A, C) Release of IL-18 or IL-8 into the supernatant was

measured by ELISA. (B) Cell death was measured as percentage uptake of propidium

iodide, normalized to cells treated with 1% Triton. (A, C) ns – not significant, * $p < 0.05$,

** $p < 0.01$, *****$p < 0.0001$ by Dunnett's multiple comparisons test. Error bars represent

the standard deviation or the standard error of the mean (SEM) (B) of triplicate wells

from one experiment. Data shown are representative of at least three independent

experiments.

Figure 5: NLRP3 and ASC are dispensable for inflammasome responses to

Salmonella **in human intestinal epithelial cells.** (A) WT Caco-2 cells were primed for

3hrs with 400 ng/ml of Pam3CSK4. One hour prior to infection, cells were treated with

the indicated concentrations of MCC950 or DMSO as a vehicle control. Cells were then

infected with PBS (Mock) or WT *S*. Typhimurium for 6hrs. Release of IL-18 into the

supernatant was measured by ELISA. (B, F) Relative mRNA expression of *NLRP3* and

PYCARD compared to the housekeeping control *HPRT* as measured by qRT-PCR in

Caco-2 cells and THP-1 macrophages. (C, D, E) WT or two independent single cell

clones of *PYCARD*$^{-/-}$ Caco-2 cells were infected with PBS (Mock), WT *S*. Typhimurium,

or Δ*sipB S*. Typhimurium. (C, D) Release of IL-18 or IL-8 into the supernatant were

measured by ELISA at 6hpi. (E) Cell death was measured as percentage uptake of

propidium iodide, normalized to cells treated with 1% Triton. (A, C, E) ns – not

significant, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ by Dunnett's multiple comparisons

test. Error bars represent the standard deviation or the standard error of the mean

(SEM) (D) of triplicate wells from one experiment. Data shown are representative of at

least three independent experiments.

Figure 6: Caspase-1 is partially required for inflammasome responses to

Salmonella **in human intestinal epithelial cells.** (A, B) Relative mRNA expression of

CASP1 compared to the housekeeping control *HPRT* as measured by qRT-PCR in

Caco-2 cells, THP-1 macrophages, human peripheral blood mononuclear cells

(PBMCs), and human small intestinal organoids. (C) WT Caco-2 cells were primed with

400 ng/ml of Pam3CSK4 for 3h. One hour prior to infection, cells were treated with 20

μM of pan-caspase inhibitor Z-VAD(OMe)-FMK, 25 μM of caspase-1 inhibitor Ac-YVAD-

cmk, or DMSO as a vehicle control. Cells were then infected with PBS (Mock), WT *S.*

Typhimurium, or Δ*sipB S.* Typhimurium for 6hrs. Release of IL-18 into the supernatant

was measured by ELISA. (D – F) WT or two independent single cell clones of *CASP1*[-/-]

Caco-2 cells were infected with PBS (Mock), WT *S.* Typhimurium, or Δ*sipB S.*

Typhimurium. (D, F) Release of IL-18 or IL-8 into the supernatant were measured by

ELISA at 6hpi. (E) Cell death was measured as percentage uptake of propidium iodide,

normalized to cells treated with 1% Triton. (C, D, F) ns – not significant, ** $p < 0.01$, ****

$p < 0.0001$ by Dunnett's multiple comparisons test. Error bars represent the standard

deviation or the standard error of the mean (SEM) (E) of triplicate wells from one

experiment. Data shown are representative of at least three independent experiments.

Figure 7: Caspase-4 is required for inflammasome responses to *Salmonella* in human intestinal epithelial cells. (A) WT Caco-2 cells were treated with siRNA targeting *CASP4, CASP5,* or a control scrambled siRNA for 72h. Cells were primed with 400 ng/ml of Pam3CSK4 for 3hrs. Cells were then infected with PBS (Mock), WT *S.* Typhimurium*,* or Δ*sipB S.* Typhimurium for 6hrs. Release of IL-18 was measured by ELISA (B – E) WT or two independent clones of *CASP4⁻ᐟ⁻* or *CASP1⁻ᐟ⁻* (E only) Caco-2 cells were infected with PBS (Mock), WT *S*. Typhimurium, or Δ*sipB S*. Typhimurium for 6hrs. (B) Release of IL-18 into the supernatant was measured by ELISA at 6hpi. (C, D) Cell death was measured as percentage uptake of propidium iodide, normalized to cells treated with 1% Triton. (E) Lysates and supernatants collected 6hpi were immunoblotted for IL-18 and β-actin. (A, B) ns – not significant, **** p < 0.0001 by Tukey's (A) or Dunnett's (B) multiple comparisons test. Error bars represent the standard deviation or the standard error of the mean (SEM) (C, D) of triplicate wells from one experiment. Data shown are representative of at least three independent experiments.

 Figure Legends

Figure 1: *Salmonella* infection induces inflammasome activation in human

intestinal epithelial cells. Caco-2 cells (A – C, E, F) or T84 cells (D) were infected with

PBS (Mock), WT *S*. Typhimurium, or Δ*sipB S*. Typhimurium. (A, D) Release of IL-18

into the supernatant was measured by ELISA at 6hpi. (B) Lysates and supernatants

collected 6hpi were immunoblotted for IL-18 and β-actin. (C) Cell death was measured

as percentage uptake of propidium iodide, normalized to cells treated with 1% Triton.

(E, F) Caco-2 cells were treated with 30 µM disulfiram or DMSO as a vehicle control 1

hour prior to infection. Cells were then infected with PBS (Mock), WT *S*. Typhimurium*,*

or Δ*sipB S*. Typhimurium. (E) Release of IL-18 into the supernatant was measured by

ELISA at 6hpi. (F) Cell death as percentage uptake of propidium iodide, normalized to

cells treated with 1% Triton. ns – not significant, **** $p < 0.0001$ by Dunnett's multiple

comparisons test (A), or by unpaired t-test (C, D, F) or by Šídák's multiple comparisons

test (E). Error bars represent the standard deviation or the standard error of the mean

(SEM) (C, F) of triplicate wells from one experiment. Data shown are representative of

at least three independent experiments.

1006

Figure 2: Bacterial T3SS ligands do not activate the inflammasome in human

intestinal epithelial cells. (A, B) Caco-2 (A) or T84 cells (B) were primed for 3h with

100 ng/ml of Pam3CSK4 and infected with PBS (Mock), WT *L. monocytogenes* (WT

Lm), or *L. monocytogenes* expressing *S*. Typhimurium SPI-1 inner rod (Lm + PrgJ), or

SPI-1 needle (Lm + PrgI), at an MOI of 100. Release of IL-18 into the supernatant was

measured by ELISA at 16hpi. L.O.D indicates the limit of detection of the assay. (C)

Differentiated intestinal organoids were treated with FlaTox (PA + LFn-Fla) or Inner Rod

Tox (PA + LFn-Rod) in media containing propidium iodide for 4h. Cell death was

measured as percentage uptake of propidium iodide, normalized to organoids treated

with 1% Triton. ns – not significant, * $p < 0.05$, **** $p < 0.0001$ by Dunnett's multiple

comparisons test. Error bars represent the standard deviation of triplicate wells from

one experiment. Data shown are representative of at least three independent

experiments.

1020

Figure 3: Human intestinal epithelial cells express low levels of NAIP/NLRC4

compared to human myeloid cells. Relative mRNA expression of *NAIP* and *NLRC4*

compared to the housekeeping control *HPRT* or *RPLP0* as measured by qRT-PCR in

(A, B) Caco-2 cells, T84 cells, and THP-1 macrophages, and in (C, D) human peripheral

blood mononuclear cells (PBMCs) and human small intestinal organoids. Error bars

represent the standard deviation of multiple wells. Data shown are representative of at

least three independent experiments.

1028

Figure 4: NAIP is not required for inflammasome responses to *Salmonella* in

human intestinal epithelial cells. WT or two independent single cell clones of *NAIP*$^{-/-}$

Caco-2 cells were infected with PBS (Mock), WT *S*. Typhimurium, or Δ*sipB S*.

Typhimurium for 6hrs. (A, C) Release of IL-18 or IL-8 into the supernatant was

measured by ELISA. (B) Cell death was measured as percentage uptake of propidium

iodide, normalized to cells treated with 1% Triton. (A, C) ns – not significant, * $p < 0.05$,

** $p < 0.01$, ***** $p < 0.0001$ by Dunnett's multiple comparisons test. Error bars represent

the standard deviation or the standard error of the mean (SEM) (B) of triplicate wells

from one experiment. Data shown are representative of at least three independent

experiments.

1039

Figure 5: NLRP3 and ASC are dispensable for inflammasome responses to

Salmonella **in human intestinal epithelial cells.** (A) WT Caco-2 cells were primed for

3hrs with 400 ng/ml of Pam3CSK4. One hour prior to infection, cells were treated with

the indicated concentrations of MCC950 or DMSO as a vehicle control. Cells were then

infected with PBS (Mock) or WT *S*. Typhimurium for 6hrs. Release of IL-18 into the

supernatant was measured by ELISA. (B, F) Relative mRNA expression of *NLRP3* and

PYCARD compared to the housekeeping control *HPRT* as measured by qRT-PCR in

Caco-2 cells and THP-1 macrophages. (C, D, E) WT or two independent single cell

clones of *PYCARD*$^{-/-}$ Caco-2 cells were infected with PBS (Mock), WT *S*. Typhimurium,

or Δ*sipB S*. Typhimurium. (C, D) Release of IL-18 or IL-8 into the supernatant were

measured by ELISA at 6hpi. (E) Cell death was measured as percentage uptake of

propidium iodide, normalized to cells treated with 1% Triton. (A, C, E) ns – not

significant, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ by Dunnett's multiple comparisons

test. Error bars represent the standard deviation or the standard error of the mean

(SEM) (D) of triplicate wells from one experiment. Data shown are representative of at

least three independent experiments.

1056

Figure 6: Caspase-1 is partially required for inflammasome responses to

Salmonella **in human intestinal epithelial cells.** (A, B) Relative mRNA expression of

CASP1 compared to the housekeeping control *HPRT* as measured by qRT-PCR in

Caco-2 cells, THP-1 macrophages, human peripheral blood mononuclear cells

(PBMCs), and human small intestinal organoids. (C) WT Caco-2 cells were primed with

400 ng/ml of Pam3CSK4 for 3h. One hour prior to infection, cells were treated with 20

μM of pan-caspase inhibitor Z-VAD(OMe)-FMK, 25 μM of caspase-1 inhibitor Ac-YVAD-

cmk, or DMSO as a vehicle control. Cells were then infected with PBS (Mock), WT *S.*

Typhimurium, or Δ*sipB S.* Typhimurium for 6hrs. Release of IL-18 into the supernatant

was measured by ELISA. (D – F) WT or two independent single cell clones of *CASP1*[-/-]

Caco-2 cells were infected with PBS (Mock), WT *S.* Typhimurium, or Δ*sipB S.*

Typhimurium. (D, F) Release of IL-18 or IL-8 into the supernatant were measured by

ELISA at 6hpi. (E) Cell death was measured as percentage uptake of propidium iodide,

normalized to cells treated with 1% Triton. (C, D, F) ns – not significant, ** *p* < 0.01, ****

p < 0.0001 by Dunnett's multiple comparisons test. Error bars represent the standard

deviation or the standard error of the mean (SEM) (E) of triplicate wells from one

experiment. Data shown are representative of at least three independent experiments.

1074

Figure 7: Caspase-4 is required for inflammasome responses to *Salmonella* in

human intestinal epithelial cells. (A) WT Caco-2 cells were treated with siRNA

targeting *CASP4, CASP5,* or a control scrambled siRNA for 72h. Cells were primed with

400 ng/ml of Pam3CSK4 for 3hrs. Cells were then infected with PBS (Mock), WT *S.*

Typhimurium*,* or Δ*sipB S.* Typhimurium for 6hrs. Release of IL-18 was measured by

ELISA (B – E) WT or two independent clones of *CASP4*[-/-] or *CASP1*[-/-] (E only) Caco-2

cells were infected with PBS (Mock), WT *S.* Typhimurium, or Δ*sipB S.* Typhimurium for

6hrs. (B) Release of IL-18 into the supernatant was measured by ELISA at 6hpi. (C, D)

Cell death was measured as percentage uptake of propidium iodide, normalized to cells

treated with 1% Triton. (E) Lysates and supernatants collected 6hpi were

immunoblotted for IL-18 and β-actin. (A, B) ns – not significant, **** p < 0.0001 by

Tukey's (A) or Dunnett's (B) multiple comparisons test. Error bars represent the

standard deviation or the standard error of the mean (SEM) (C, D) of triplicate wells

from one experiment. Data shown are representative of at least three independent

experiments.