Sex-dependent effects of *Setd1a* haploinsufficiency on development and adult behaviour

Matthew L. Bosworth[1], Anthony R. Isles[1], Lawrence S. Wilkinson[1,2,3], & Trevor Humby[1,2,3]*

[1]MRC Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neuroscience, School of Medicine, Cardiff University, Cardiff, UK

[2]School of Psychology, Cardiff University, Cardiff, UK

[3]Neuroscience and Mental Health Research Institute, Cardiff University, Cardiff UK

*Corresponding author: Dr Trevor Humby HumbyT@cardiff.ac.uk Tel. +44(0)2920 876758

ABSTRACT

Loss of function (LoF) mutations affecting the histone methyl transferase *SETD1A* are implicated in the aetiology of a range of neurodevelopmental disorders including schizophrenia. We examined indices of development and adult behaviour in a mouse model of *Setd1a* haploinsufficiency, revealing a complex pattern of sex-related differences spanning the pre- and post-natal period. Specifically, male *Setd1a*$^{+/-}$ mice had smaller placentae at E11.5 and females at E18.5 without any apparent changes in foetal size. In contrast, young male *Setd1a*$^{+/-}$ mice had lower body weight and showed enhanced growth, leading to equivalent weights by adulthood. Embryonic whole brain RNA-seq analysis revealed expression changes that were significantly enriched for mitochondria-related genes in *Setd1a*$^{+/}$ samples. In adulthood, we found enhanced acoustic startle responding in male *Setd1a*$^{+/-}$ mice which was insentitive to the effects of risperidone, but not haloperidol, both commonly used antipsychotic drugs. We also observed reduced pre-pulse inhibition of acoustic startle, a schizophrenia-relevant phenotype, in both male and female *Setd1a*$^{+/-}$ mice which could not be rescued by either drug. In the open field and elevated plus maze tests of anxiety, *Setd1a* haplosufficiency led to more anxiogenic behaviour in both sexes, whereas there were no differences in general motoric ability and memory. Thus, we find evidence for changes in a number of phenotypes which strengthen the support for the use of *Setd1a* haploinsufficient mice as a model for the biological basis of schizophrenia. Furthermore, our data point towards possible underpinning neural and developmental mechanisms that may be subtly different between the sexes.

Keywords: SETD1A; schizophrenia; developmental delay; placenta; startle response; prepulse inhibition; haloperidol; risperidone

Introduction

Epigenetic modifications controlling chromatin structure and organisation, can enhance or suppress gene expression dependent on the mechanism in action. Methylation of lysine residue 4 on histone 3 (H3K4), generally considered an activating histone mark [1], is catalysed by a histone methyltransferase complex, of which *SETD1A* (SET Domain Containing 1A) is an essential subunit [2]. *SETD1A* has been implicated in a range of biological functions, including cell cycle regulation [3-5], maintenance of pluripotency in embryonic stem cells [6-8] and neuronal progenitors [9,10], and DNA repair [4,11,12], highlighting a wide distribution across different cell types, throughout different stages of development and during adulthood [2]. Exome sequencing studies have implicated loss of function (LoF) mutations in the *SETD1A* gene that increase susceptibility for a number of neurodevelopmental disorders (NDD) and conditions, including developmental delay, early-onset epilepsy, intellectual disability and schizophrenia [13-16]. While these genetic variants are rare (occurring in 0.13 % of schizophrenia cases), they are highly penetrant and their effects on gene function (i.e. haploinsufficiency) can be recapitulated in model systems. Consequently, *SETD1A* provides a biologically tractable target for disease modelling.

Homozygous knock-out of *Setd1a* is embryonically lethal[8]. Consequently, non-human models have targeted haploinsufficiency (*Setd1a$^{+/-}$*), with current mouse models employing either frameshift mutations to exon 7 or 15 or disruption to exon 4 via an upstream LacZ/Neo cassette [17-20] to investigate the mechanisms of Setd1a function and how it might influence NDD. Haploinsufficiency also recapitulates the clinical LoF mutations found in human conditions for more direct translational impact. From a neurobiological perspective, changes including axonal branching deficits, impaired synaptic plasticity, regionally-specific gene expression changes linked to morphology and dendritic complexity, synapse formation, and abnormal cortical ensemble activity have been observed [17-21]. *Setd1a* may also have a role in regulating the balance between neuronal progenitor cells (NPC) proliferation and differentiation in neurogenesis [8-10], suggesting that perturbations to this balance may play a role in pathogenic mechanisms of *Setd1a* LoF in neurodevelopmental

disorders. Behaviour in these models has also been affected, with impairments in
working and spatial memory, sensorimotor gating, hyperactivity, and altered juvenile
social behaviour [17-20]. Other work in *Drosophila* has shown that short- and long-
term courtship memory is impaired by conditional knockdown of Set1 in neurons of
the mushroom body [22].

Setd1a may have important roles in early development, as shown in the maternal
high-fat diet model in mice, a manipulation that is associated with schizophrenia-
relevant phenotypes in animal models [23,24], where *Setd1a* expression is reduced
in the placenta [24]. There is further evidence that Setd1a can influence placental
function and development from studies in *Setd1a* haploinsufficient mice via its
actions on embryonic stem cell (ESC) pluripotency and differentiation [6-8], with
some evidence of gestational compromise though no differences in growth to
adulthood [17]. Based on these findings, it might be predicted that *SETD1A* LoF
causes disturbed neurodevelopment with downstream consequences for later brain
function that contribute to the emergence of atypical patterns of behaviour and
psychopathology, possibly via effects on placental function.

Therefore, using a newly created mouse model of Setd1a haploinsufficiency, where
Setd1a is disrupted in exon 4 [25], we investigate how deletion of Setd1a may
influence pre- and post-natal growth and behaviours in adulthood. In this model, we
have previously demonstrated downregulation in genes enriched for mitochondrial
pathways in frontal cortex from E14 to P70, but no differential enrichment in genes
associated with schizophrenia [25], here we replicate these results at a similar time
point (E13.5). We also find evidence for sex-linked developmental changes that
extend into early post-natal growth, with effects on male and female embryo:placenta
ratios that suggest imbalance in demand and delivery capacity at different foetal
ages, and in and male post-weaning catch-up growth. In terms of adult behaviour,
we found evidence for an enhanced acoustic startle response in male Setd1a$^{+/-}$ mice
and increased anxiety-related behaviour in both male and female Setd1a$^{+/-}$ mice.
Prepulse inhibition of the acoustic startle response was attenuated in both sexes.
However, the sensorimotor gating deficits could not be rescued by haloperidol or
risperidone, suggesting that these antipsychotic agents are ineffective for
ameliorating schizophrenia-relevant phenotypes in *Setd1a$^{+/-}$* mice. Taken together

our findings strengthen the support for the use of *Setd1a* haploinsufficient mice as a
model for the biological basis of schizophrenia and point towards possible
underpinning neural mechanisms, that may be sexually dimorphic.
112

Materials and methods

Animals

All procedures were conducted in accordance with the UK Animals (Scientific
Procedures) Act 1986. *Setd1a*$^{+/-}$ mice were produced using a strain created by the
Knockout Mouse Phenotyping Consortium. To generate a model with a constitutive
germline transmissible knockout allele, male C57BL/6NTac-
Setd1a$^{tm1c(EUCOMM)Wtsi}$/WtsiCnrm mice (obtained from MRC Harwell) were paired with
female B6.C-Tg(CMV-cre)1Cgn/J mice (obtained from The Jackson Laboratory) [25].
F1 male progeny were genotyped to identify animals heterozygous for the
Setd1a$^{CMV-cre/+}$ allele. These were crossed with C57BL/6J females (obtained from
Charles River) to enable removal of the X-linked CMV-cre transgene from males in
the F2 generation. Experimental cohorts were generated by pairing male F2
Setd1a$^{+/-}$ mice with C57BL/6J females (wildtype, WT). These mice show
haploinsufficiency as demonstrated by a 48.8% reduction in mRNA at E13.5 in
Setd1a$^{+/-}$ brain compared to WT, with comparable reductions in levels of SETD1A
protein (46.3% reduction) (Supplementary Methods Figure 1). Animals for the
assessment of post-natal growth (N=84) and behavioural studies (N=77 and N=56)
were housed in mixed-genotype cages with littermates of the same sex (2-5 per
cage) in a vivarium maintained on a 12-hour light-dark cycle (lights on from 08:00-
20:00) at a temperature of 21 ($\pm$ 2) °C and 50 ($\pm$ 10) % humidity. Experiments were
conducted during the light phase. Standard laboratory chow and water was available
ad libitum throughout all experiments, and mice were carefully monitored for signs of
ill health throughout the studies.
137

RNA-Seq for schizophrenia common variants

139 RNA was extracted from 16 (8 WT and 8 KO, balanced for sex) E13.5 whole brains
140 using a Direct-zolTM RNA Miniprep kit (Zymo, UK). E13.5 tissue was selected based
141 on the results of the developmental expression work, which showed highest

expression levels at this timepoint, and to supplement the data in Clifton et al (2022) [25]. Detailed methods can be found in Supplementary Methods Figure 2. In brief, following sample purification, concentration and integrity assessment Library preparations were performed according to manufacturer's instructions (KAPA mRNA Hyperprep kit. Roche, Switzerland) with 1 μg total RNA as input. Library concentrations were adjusted prior to pooling to achieve a final DNA molarity of 20 nM and sequencing was performed on a HiSeq 4000 (Illumina, USA) with 75 bp paired-end reads at a read depth of 30 million reads. Reads were trimmed to remove adapters and low-quality bases, and passed FastQC quality control [27], before mapping to the mouse reference genome (GRCm38) using STAR (mean number of reads mapped was 97.2%, SD = 0.5%). Read counts, generated using featureCounts were allocated to genomic features using the mouse Ensembl gene annotation (GRCm.38.95). Differential expression analysis was performed using DEseq2 [28] implemented in R (version 3.5.3). The Benjamini Hochberg correction was used to correct for multiple testing and identify genes that were differentially expressed in *Setd1a*+/- relative to WT ($p$adj < .05), with sex included as a covariate.

GO term enrichment analysis was performed using the Database for Annotation, Visualization and Integrated Discovery [29]. Only protein-coding genes with entrez IDs (219 genes) were included in the differentially expressed gene-set. A custom background gene-set was created using all expressed genes in the dataset (12,295 genes). This was defined as any gene with a FPKM > 1 across 8 samples (i.e., the number of biological replicates in each condition). The differentially expressed gene-set was filtered to include only genes with human entrez IDs (205 genes). As previously, with the hypothesised links between *Setd1a* and risk for schizophrenia*,* a custom background gene-set was assessed for enrichment of schizophrenia common variants using MAGMA [30] with GWAS summary data from 40,675 cases and 65,643 controls [31]. The differentially expressed gene-set was filtered to include only genes with human entrez IDs (205 genes), and gene-set SNPs were filtered to include those with a minor allele frequency of less than 0.01.

Assessment of embryo and placental weights, placental gene expression and postnatal growth.

For the neurodevelopmental investigations, timed-matings were conducted between
C57BL/6J x C57BL/6NTac parents and pregnant dams were culled by cervical
dislocation. Placental and embryo wet weights were recorded at E11.5, E13.5 and
E18.5 and an embryo:placenta weight ratio was calculated by dividing embryo wet
weight by placental wet weight (see Supplementary Methods Table 1 for litter sizes
and N). Whole placentas and embryo tissue biopsies were snap frozen using dry ice
and stored at -80 °C until required. Genotyping was conducted using DNA extracted
from the tissue biopsy to identify $Setd1a^{+/-}$ mice and determine embryonic sex.
Animals for the assessment of post-natal growth (males: WT, N=23, $Setd1a^{+/-}$ mice,
N=18 and females: WT, N=25, $Setd1a^{+/-}$ mice, N=18) were weaned on postnatal day
28 and weighed at maternal separation (P28) and then every seventh day thereafter
until P70. Thus, growth curves were assessed after maternal separation to eliminate
potential effects of early-life stress induced by repeated handling on behaviour in
adulthood.
189
Behavioural testing
Two cohorts of mice were used to assess the effects of $Setd1a$ haplosufficiency on
adult behavior. The first cohort (males: WT, N=21, $Setd1a^{+/-}$ mice, N=18 and
females: WT, N=20, $Setd1a^{+/-}$ mice, N=18), a subsample of the animals used to
evaluate postnatal growth, were tested in adulthood (aged 2-3 months at start of
testing in the following order: elevated-plus maze (EPM), open field test (OFT),
locomotor activity, sensorimotor gating, rotarod test, and novel object recognition.
The second cohort of mice (males: WT, N=16, $Setd1a^{+/-}$ mice, N=10 and females:
WT, N=17, $Setd1a^{+/-}$ mice, N=13), was first used to test whether the effects of
$Setd1a$ haploinsufficiency on ASR and PPI observed in the first cohort could be
replicated (Supplementary Results Figure 3). These mice were then randomly
divided into two approximately equal groups (balanced for sex and genotype, see
Fig. 3 for n) and allocated to either the haloperidol or risperidone condition, to test
the effects of these antipsychotics on sensorimotor gating impairments. All apparatus
was cleaned with 70 % (v/v) ethanol between animals, and testing was conducted
under dimmed lighting (15 lux) except for locomotor activity which was run in the
dark.
207
Sensorimotor gating

The acoustic startle response (ASR) and prepulse inhibition (PPI) were measured
using apparatus from SR-Lab (San Diego Instruments, USA). Animals were placed
in a clear Perspex tube (35 mm internal diameter) mounted on a Perspex plinth in a
sound-attenuating chamber. A 70 dB (A scale) white noise stimulus was
continuously played throughout the session via a loudspeaker positioned 120 mm
above the tube. The whole-body startle response was detected on each trial by a
piezoelectric sensor attached to the plinth, which transduced flexion in the plinth into
a digitised signal. The average startle response (V_{avg}) was recorded in arbitrary
startle units using SR-Lab software over the 65 ms period following stimulus onset.
Startle data were weight-adjusted by dividing V_{avg} by body weight recorded
immediately after the test session. PPI was calculated as the percentage reduction in
startle amplitude between prepulse and pulse-alone trials (excluding the first three
pulse-alone trials). Each session of 96 trials, started with a five-minute habituation
period, and was divided into three blocks. Pulse amplitude was set to 120 dB and
105 dB in block 1 and block 2, respectively and the third block comprised a range of
pulse-alone trials (80 – 120 dB in 10 dB increments), with three of each trial type.
Acoustic stimuli were presented with a mean intertrial interval of 16 seconds
(pseudorandomly varied between 6 and 24 seconds). Each pulse-alone trial
consisted of a 40 ms 120db or 105db startle stimulus. Prepulse trials consisted of a
20 ms prepulse stimulus followed by a 40 ms startle stimulus 80 ms after prepulse
offset. In the first two blocks, 6 consecutive pulse-alone trials were presented
followed by 7 additional pulse-alone trials interspersed with 18 prepulse trials (either
4, 8 or 16 dB (above background) with six trials of each prepulse amplitude). Only
the data from 120 dB pulse alone and the 8 and 16 dB prepulse trials are reported
here for brevity, but all raw data are available in the online repository. For the
pharmacological investigations using the second cohort of mice, with haloperidol and
risperidone, a reduced session protocol was used, whereby all 4 dB prepulse trials
and all trials with 105 dB stimuli were removed.
237

Anxiety-related behaviour
The OFT and EPM were used to assess anxiety-related behaviour under identical
lighting conditions (15 lux) and the position of each mouse in the apparatus was
tracked using EthoVision XT software (Noldus Information Technology, Netherlands)
at a rate of 12 frames/s via a camera mounted above the centre of each piece of

apparatus. The OFT comprised a 750 x 750 mm arena with 45 cm white Perspex

walls. Each animal was placed in the same corner of the arena and allowed to

explore freely for 10 minutes. The arena was divided into two, concentric virtual

zones: the 'inner zone' (central 600 x 600 mm) and the 'outer zone' (surrounding 150

mm). The EPM was constructed of white Perspex and consisted of four arms of

equal size (175 x 78 mm) extending from a central square region (78 x 78 mm) and

positioned 450 mm above the floor. Two of the arms were 'open' (no walls) and two

were enclosed by 150 mm high opaque walls. Arms of the same type were

diametrically opposed. Each animal was placed in the same enclosed arm at the

start of the trial and allowed to freely explore the apparatus for five minutes.

Measures of anxiety were the patterns of exploration between the different arms or

zones of the apparatus, with particular focus on the time in the central zone of the

OFT and the latency to enter and time on the open arms of the EPM.   Other

measures included total distance moved, zone or arm transitions and the numbers of

head dips and stretch attend postures made by the mice on the EPM.

Locomotor activity

Locomotor activity levels were assessed using previous methods [32], with the mice

placed into clear Perspex chambers (210 x 360 x 200 mm) with two transverse

infrared beams positioned 30 mm from either end of the chamber and 10 mm above

the floor of the chamber. The apparatus was linked to a computer using ARACHNID

software (Cambridge Cognition Ltd., UK). Activity levels were recorded as the

number of beam breaks during each session using a custom programme (BBC

BASIC Version 6). Each animal completed three 2 hours sessions over consecutive

days at the same time of day.

Rotarod test

Motor learning and co-ordination were assessed using a mouse Rotarod (Model

47600, Ugo Basile, Italy). The rod (30 mm diameter) was coated with rubber grooves

to provide grip. Each animal completed five trials across two days (three trials on day

one and two trials the next day). During each 5 minutes trial, the speed of rotation

increased from 5 – 50 rpm at a constant rate of 0.15 rpm/s. The latency to fall was

recorded on each trial, when a mouse caused a timer to stop when triggered by a

lever (160 mm below the rod). On trials where the animal stayed on the rod for the
duration of the trial, the latency to fall was recorded as the maximum trial length.
278

Novel object recognition
Recognition memory was assessed using the novel object recognition (NOR)
paradigm [33] with retention intervals of 30 minutes and 24 hours. All mice
completed the experiment at both the 30 minute and 24 hours retention interval, with
order counterbalanced across animals and at least 24 hours between each
condition. For the second test session, new objects that had not been encountered
previously were placed in the quadrants of the arena that were unoccupied in the
previous session. Location of the objects and allocation of objects to retention
interval were counterbalanced across animals. Testing was conducted in a white
Perspex arena (300 x 300 mm) with 300 mm high walls, and objects were common
household items (bottles, cans, containers) of approximately the same size (120, x
50 mm). Prior to testing, mice were habituated to the empty arena for 10 minutes per
day over three consecutive days. Each test session comprised three phases,
conducted over one of two days as appropriate, and between phases mice were
returned to a holding cage in the test room. The 'habituation phase', where the mice
explored the empty arena for 10 minutes, preceded the 'acquisition phase', in which
mice were returned to the arena and allowed to explore two identical objects (in
diagonally opposite quadrants of the arena, 105 mm from the corner) for up to 15
minutes. To control for potential differences in object neophobia that could contribute
to subsequent memory performance, the total amount of object exploration (defined
as when the animal's head was within 20 mm and oriented towards the object) was
timed during the trial by the experimenter. The acquisition phase was ended once 40
seconds of object exploration was achieved or the maximum trial length had
elapsed. Animals were then returned to either a holding cage or their home cage for
a retention interval of 30 minutes or 24 hours. In the 'test phase', mice were placed
back in the arena with one of the objects that they had been exposed to during the
acquisition phase ('familiar') and another object that had not been encountered
previously ('novel'). The objects were presented in the same locations as the
acquisition phase, and the mice allowed to explore freely for five minutes, and the
amount of familiar and novel object exploration was recorded manually by the
experimenter using EthoVision XT software (Noldus Information Technology,

Netherlands). The main measure of novel object recognition was a discrimination

ratio (d1) calculated from the object contact times for each test session, with the

following equation: novel/novel+familiar.

Drug preparation and administration

Haloperidol (Sigma, UK) and risperidone (Sigma, UK) were initially dissolved in 1M

glacial acetic acid and then diluted in 0.9 % saline such that an equal volume (100

µL per gram of bodyweight) was administered for each injection. Drugs were

administered at two doses (0.5 mg/kg and 1.0 mg/kg, i.p.), with a vehicle control

treatment, with a 30 minutes pre-treatment delay before testing. This regimen and

these doses were selected based previous studies [34.35] and pilot dose-response

data showing that a 0.5 mg/kg dose of either haloperidol or risperidone did not have

significant effects on sensorimotor gating in WT C57BL/6J mice (Supplementary

Methods Figure 3). At 1.0 mg/kg, haloperidol caused a significant increase in PPI

and risperidone caused a significant reduction in the ASR. Therefore, 0.5 mg/kg was

selected as a sub-threshold dose to explore drug effects in $Setd1a^{+/-}$ mice in the

absence of non-specific effects of the drugs on sensorimotor gating that occur in WT

mice. Dose order was counterbalanced across animals for vehicle and 0.5 mg/kg

and there was a seven day washout between sessions. All mice received the 1.0

mg/kg dose in the final session to explore drug effects at a dose that was known to

affect sensorimotor gating in WT animals to rule out the possibility that the lack of

effect in $Setd1a^{+/-}$ mice was due to an insufficient dose of drug.

Statistical analysis

Data are available at: https://osf.io/gt7ew/. Data were analysed using IBM SPSS

software (version 25). Placental and embryo weight data were analysed using 3-way

MANOVA with between-subject factors of GENOTYPE (WT and $Setd1a^{+/-}$), SEX

(male and female) and AGE (E11.5, E13.5, E18.5) and a covariate of litter size.

Simple effects were analysed by 1-way ANOVAs with a between-subject factor of

GROUP (male WT, male $Setd1a^{+/-}$, female WT, female $Setd1a^{+/-}$). Analysis of

postnatal growth included the within-subjects factor of AGE (7 levels) and the

between-subject factors of GENOTYPE and SEX. ASR data were analysed using

ANOVA with GENOTYPE and SEX as between-subjects factors, and PPI data were

analysed with PPI-INTENSITY (8 dB and 16 dB) as an additional within-subjects

factor. Effects of haloperidol and risperidone were analysed separately by ANOVA
with GENOTYPE and SEX as between-subjects factors and a within-subjects factor
of DOSE (vehicle, 0.5 mg/kg and 1.0 mg/kg). Data from the EPM and OFT were
analysed by MANOVA with between-subject factors of GENOTYPE and SEX, and
additional within-subjects factors of ZONE (central, outer) or ARM (open, closed)
were included to assess exploration across the OFT or EPM, respectively. Data from
one female WT animal was excluded from analysis of the OFT because it was
identified as a multivariate outlier. Locomotor activity, rotarod and NOR data were
analysed using ANOVAs with between-subject factors of GENOTYPE and SEX and
the additional within-subjects factors of DAY (3 levels), TRIAL (5 levels) or DELAY
(30 minutes and 24 hours), respectively. Data from one male WT subject was
missing from the NOR due to a technical issue during testing. Greenhouse Geisser
corrected results are reported for locomotor activity and the rotarod test because a
significant Mauchly's test result indicated that the assumption of sphericity had been
violated. Bonferroni-corrected *post-hoc* tests were used to probe significant effects
where appropriate. Data are shown as mean±SEM, and criterion level of significance
was set at the 0.05 level.
361
362 **Results**
363
Sex-related effects of *Setd1a* haplosufficiency on placental weight and gene
expression, and post-natal growth.
Analysis of embryonic development through gestation, at time points where *Setd1a*
mRNA expression and protein levels remained relatively constant (Supplementary
Results Figure 1), highlighted that litter size remained relatively consistent at all
gestational ages, with equal numbers of male and female embryos, and Mendelian
ratios of WT:*Setd1a*$^{+/-}$ (Supplementary Results Table 1). Male *Setd1a*$^{+/-}$ embryos
were generally lighter than WT (~6%), and female *Setd1a*$^{+/-}$ embryos slightly heavier
than WT (~4%), however these differences were not significant (Fig. 1a, main effect
of GENOTYPE, $F_{1,105}=0.32$, p=0.57), at any timepoints investigated
(GENOTYPE*AGE, $F_{2,105}=0.61$, p=0.94). Male embryos weighed more than female
(main effect of SEX, $F_{1,105}=6.54$, p=0.01), but genotype did not interact with sex to
influence embryo weight (GENOTYPE*SEX, $F_{1,105}=3.43$, p=0.07). As expected,

embryos increased in weight throughout gestation (main effect of AGE, $F_{2,105}$=5358.26, p<0.001), but growth was not affected by genotype or sex (3-way interaction, $F_{2,105}$=0.40, p=0.67). A significant 3-way interaction demonstrated complex effects of AGE and SEX on the size of placentae from $Setd1a^{+/-}$ and WT embryos (Fig. 1b, $F_{2,105}$=6.21, p=0.003). At E11.5, simple effects analysis showed a significant effect of GENOTYPE ($F_{3,23}$=7.56, p=0.001) with *post-hoc* testing highlighting a significant reduction (~32% smaller) in weight between male $Setd1a^{+/-}$ and male WT placentae (p<0.001). At E18.5, placentae from female $Setd1a^{+/-}$ placentae were significantly reduced in size (~16% smaller) compared with both male and female WT (main effect of GENOTYPE, $F_{3,39}$=7.38, p=0.001 and *post hoc* testing p<0.003). Placentae were generally smaller in $Setd1a^{+/-}$ embryos than WT, across all time points evaluated (main effect of GENOTYPE, $F_{1,105}$=37.11, p<0.001). These placental weight differences were reflected by increased embryo:placenta ratios (EPR) in $Setd1a^{+/-}$ embryos relative to WT (Fig. 1c, main effect of GENOTYPE, $F_{1,105}$=16.98, p<0.001). Further analysis indicated elevated EPR in male $Setd1a^{+/-}$ embryos relative to male WT at E11.5 (main effect of GENOTYPE, $F_{3,23}$=3.98, p=0.02 and *post hoc* testing p=0.05) and female $Setd1a^{+/-}$ embryos at E18.5 (main effect of GENOTYPE, $F_{3,39}$=3.25, p=0.03 and *post hoc* testing p=0.04). As embryo weight increased with embryonic age, the EPR also increased, as placenta size plateaued (main effect of AGE, $F_{2,105}$=2266.42, p<0.001). Litter size, as a covariate, impacted on embryo size ($F_{1,105}$=10.20, p=0.02), but did not affect either placenta size ($F_{1,105}$=0.61, p=0.44) or embryo:placenta ratios ($F_{1,105}$=3.13, p=0.08).

400 *Setd1a* haploinsufficiency, in this model, did not cause embryonic lethality as demonstrated by WT:$Setd1a^{+/-}$ ratios at birth which did not deviate from Mendelian ratios (X^2_{10}=1.39, p>0.05), and median litter size across 13 litters was 7 (Supplementary Results Table 1), which is equivalent to reported sizes for C57BL/6J mice [36]. Proportions of males and females was also as expected (X^2_{10}=1.39, p>0.05). Furthermore, all animals appeared in good general health and low pre-weaning mortality was observed (2.3%). Assessment of body weight (Figs. 1e and 1f) from the point at which pups were separated from dams (P28) revealed that, as expected male mice of both genotypes were heavier than females, with a significant increase over the first few weeks post-weaning (AGE*SEX interaction, $F_{2.26,180.93}$=63.79, *p*<0.001). A significant interaction was found between GENOTYPE

and AGE ($F_{2.26,180.93}$=11.43, p<0.001) and *post-hoc* analysis revealed that male
Setd1a$^{+/-}$ mice were significantly lighter than their WT littermates at maternal
separation (P28; p=0.005), an effect that was sustained one week later (P35,
p<0.001). By P42, male *Setd1a*$^{+/-}$ and WT mice were of equivalent body weight
(*p*=0.06), indicating that catch-up growth had occurred. This was confirmed by
analysing the percentage increase in body weight from P28 (Figs. 1e and 1f, inset
bar charts). Specifically, body weights of *Setd1a*$^{+/-}$ male (but not female) increased
significantly more than their WT littermates from P42 onwards, the age at which
there was no longer a genotype effect on body weight.
420
RNA-Seq highlights mitochondrial differences, but not for schizophrenia
common variants in E13.5 *Setd1a*$^{+/-}$ brain.
267 genes (Benjamini Hochberg p_{adj}<0.05) were differentially expressed (234
downregulated and 33 upregulated; Supplementary Results Table 1) in brains of
Setd1a$^{+/-}$ mice at E13.5 (Fig. 2a). Modest changes in gene expression were
observed, with log2 fold changes ranging from -0.63 to 1.19. Hierarchical clustering
based on normalised read counts was performed using Morpheus
(https://software.broadinstitute.org/morpheus) to produce a dendrogram (Fig. 2b).
This showed that samples of the same genotype clustered together, suggesting
similar patterns of gene expression changes within each group, which were taken
forward for the *Gene ontology* term enrichment analysis.
432
As shown in Table 1, results of the *Gene ontology* term enrichment analysis revealed
14 nominally significant (uncorrected *p*<0 .05) associations, but only the
mitochondrion GO:0005739 (Benjamini Hochberg p_{adj}=0.002) survived correction for
multiple testing. Within the initial analysis, however, were annotations relating to
other mitochondrial terms, cilium, and methylation. Gene-set enrichment analysis
revealed that genes that were differentially expressed in *Setd1a*$^{+/-}$ E13.5 brain were
not significantly enriched for schizophrenia common variant association (*B*=0.004,
SE=0.088, p=0.48).
441
***Setd1a*$^{+/-}$ mice show an elevated acoustic startle response (ASR) and**
diminished prepulse inhibition (PPI)

Both male and female Setd1a$^{+/-}$ mice showed increased levels of startle responding
to the 120 dB pulse-alone stimuli relative to WT littermates (Fig. 3a, main effect of
GENOTYPE, $F_{1,73}=5.35$, $p=0.02$). Although there were no overall effects of SEX
($F_{1,73}=3.25$, $p=0.08$) or a GENOTYPE*SEX interaction ($F_{1,73}=1.98$, $p=0.16$) on ASR,
the difference was much more pronounced in male mice ($t_{37}=2.54$, $p=0.02$ for
comparison between male WT and Setd1a$^{+/-}$ mice), than female mice ($t_{36}=0.67$,
$p=0.51$ for female comparison) suggesting greater impact of Setd1a haplosufficiency
on startle responsivity in male mice. This increase in ASR was relatively consistent
throughout the session (Supplementary Results Figure 2). We were able to replicate
the pattern of increased ASR in male mice, and also demonstrated a similar, but
reduced increase in female mice in a second cohort of mice (Supplementary Results
Figure 3a), further validating increased startle responding as a result of Setd1a
haplosufficiency.

As expected, a weak prepulse stimulus was able to modify the ASR and increased
prepulse intensity led to greater PPI shown by the mice (Fig. 3b, main effect of PPI-
INTENSITY, $F_{1,73}=253.7$, $p<0.001$), however Setd1a$^{+/-}$ mice showed reduced levels
of PPI relative to WT (main effect of GENOTYPE, $F_{1,73}=26.64$, $p<0.001$). This effect
was consisted at both prepulse intensities used (GENOTYPE*PPI-INTENSITY
interaction ($F_{1,73}=2.00$, $p=0.16$) and in both male and female Setd1a$^{+/-}$ mice
(GENOTYPE*PPI-INTENSITY*SEX interaction, $F_{1,73}=0.001$, $p=0.97$). Comparisons
between the groups of mice, demonstrated larger effects between female WT and
Setd1a$^{+/-}$ mice ($t_{36}=3.82$ and 4.75, $p<0.001$, for 8 and 16db prepulse stimuli,
respectively) compared to male mice ($t_{37}=2.04$, $p=0.049$ and $t_{37}=2.88$, $p=0.07$ for 8
and 16db prepulse stimuli, respectively) suggesting greater impact of Setd1a
haplosufficiency on PPI in female mice. This pattern of attenuated PPI in both sexes
of Setd1a$^{+/-}$ mice, was replicated in the second cohort of mice (Supplementary
Results Figure 3b). Thus, these data indicate that Setd1a haplosufficiency led to
robust phenotypes of hyperstartling and reduced PPI, with stronger effects on startle
in male mice and PPI in female mice.

Male Setd1a$^{+/-}$ mice are insentitive to the effects of risperidone on ASR, but
disrupted sensorimotor gating in Setd1a$^{+/-}$ mice cannot be rescued by either
risperidone or haloperidol.

To investigate the effects of *Setd1a* haplosufficiency on ASR and PPI further, and

also to further explore the possible links between *Setd1a* and schizophrenia, we

used the antipsychotic drugs risperidone or haloperidol with a second cohort of mice.

Following a series of pilot studies (Supplementary Methods Figure 3) we used a

dose of each drug (0.5mg/kg) which we had demonstrated to be sub-threshold in WT

mice for both ASR and PPI. We compared the effects at 0.5mg/kg for each drug to

vehicle, and also a higher dose (1mg/kg), where we showed risperidone decreased

ASR and haloperidol increased PPI in WT mice in pilot work.

A significant significant three way interaction between GENOTYPE, DOSE and SEX

($F_{2,46}$=5.51, p=0.01), suggested complex effects of risperidone on the startle

response (Fig. 3c). Investigating this further in males, the main effect of dose

($F_{2,22}$=7.32, p=0.004) was qualified by a significant interaction between dose and

genotype ($F_{2,22}$=6.26, p=0.01). Bonferroni-corrected pairwise comparisons revealed

a dose-dependent reduction of ASR in WT (relative to vehicle, p=0.05 at 0.5 mg/kg

and *p*=0.01 at 1.0 mg/kg) but not *Setd1a*$^{+/-}$ males (relative to vehicle, *p*=0.62 at 0.5

mg/kg and *p*=0.42 at 1.0 mg/kg). Conversely, in females the main effect of DOSE

was significant ($F_{2,24}$=23.09, p<0.001), but there was no GENOTYPE*DOSE

interaction ($F_{2,24}$=1.32, p=0.29). These findings suggest that the startle-inhibiting

effect of risperidone, was reduced in male (but not female) *Setd1a*$^{+/-}$ mice relative to

WT. Risperidone did not have any significant effects on PPI (Fig. 3d, main effect of

DOSE, $F_{2,46}$=2.71, p=0.08), and there were no interactions between the dose of the

drug and *Setd1a* haplosufficiency ($F_{2,46}$=1.38, p=0.26), however the previously

observed reduction of PPI in *Setd1a*$^{+/-}$ mice was maintained (main effect of

GENOTYPE ($F_{1,23}$=6.36, p=0.02). Therefore, these results demonstrate that

risperidone did not rescue the reduction in PPI resulting from *Setd1a*

haplosufficiency in either sex, however, the elevated startle response in *Setd1a*$^{+/-}$

male mice was insensitive to the modifying effects of the drug.

The elevated startle response shown by *Setd1a*$^{+/-}$ mice was unaffected by

haloperidol (main effect of GENOTYPE, $F_{2,50}$=14.01, p=0.001) at all doses used

(GENOTYPE*DOSE interaction: $F_{2,50}$=1.72, p=0.19) and with both sexes of mice

(GENOTYPE*SEX interaction $F_{2,50}$=0.02, p=0.90). Haloperidol did not significantly

alter the startle response from vehicle (Fig. 3e, main effect of DOSE, $F_{2,50}$=3.17,

p=0.05). Haloperidol was also without significant effect on PPI (Fig. 3f, main effect of
DOSE, $F_{2,50}=3.08$, p=0.06), although there was a tendency for the dose-related
increase in PPI we observed in pilot studies. The significant reduction of PPI in
$Setd1a^{+/-}$ mice (main effect of genotype, $F_{1,25}=21.92$, p<0.001) was maintained
across all doses (GENOTYPE*DOSE interaction, $F_{2,50}=0.02$, p=0.98). There was no
effect of SEX on these data ($F_{1,25}=0.03$, p=0.87). Thus, these results show that
neither $Setd1a$ haplosufficiency induced elevated startle responses or attenuations
of PPI were normalised by haloperidol, in either sex.
520

Male and female $Setd1a^{+/-}$ mice show increased anxiety-related behaviour.
In the 10 minutes open field test, all mice, as expected, tended to avoid the inner
zone of the arena (Fig. 4a, main effect of ZONE, $F_{1,72}=5736.72$, p<0.001), however a
significant GENOTYPE*ZONE interaction ($F_{1,72}=4.58$, p=0.04) indicated that
$Setd1a^{+/-}$ mice explored the central zone less (Fig. 4b). Female mice showed
increased avoidance of the central zone (main effect of SEX, $F_{1,72}=5.69$, p=0.02), a
pattern of behaviour that was consistent in both groups of mice (GENOTYPE*SEX
interaction, $F_{1,72}=0.15$, p=0.70), which demonstrates that both sexes of $Setd1a^{+/-}$
mice had a reduced tendency to explore the more anxiogenic central region of the
arena. This was supported by a trend towards reduced transitions into the central
zone (Fig. 4c, main effect of GENOTYPE, $F_{1,72}=3.59$, p=0.06). There were no
differences between $Setd1a^{+/-}$ and WT mice in the total distance moved (locomotion)
during the test (Fig. 4d, main effect of GENOTYPE, $F_{1,72}=2.40$, p=0.13) suggesting
an equivalent propensity to explore the arena, even though $Setd1a$ haplosufficiency
in both males and females led to more anxiogenic behaviour.
536

$Setd1a^{+/-}$ mice also showed an anxiogenic profile on the EPM, with a slower latency
to enter the more anxiety-inducing open arms of the EPM (Fig. 4e, main effect of
GENOTYPE, $F_{1,73}=4.75$, p=0.032) although over the entire 5 min session they
ultimately explored the open arms an equivalent amount to their WT littermates (Fig.
4f, main effect of GENOTYPE, $F_{1,73}=0.09$, p=0.76), and demonstrated equal
amounts of anxiety-induced behaviour such as head dips (Fig. 4g, main effect of
GENOTYPE, $F_{1,73}=1.20$, p=0.28), and stretch attend postures (Fig. 4h, main effect of
GENOTYPE, $F_{1,73}=0.05$, p=0.82). There were no differences between $Setd1a^{+/-}$ and
WT mice in the total distance moved during the test (Fig. 4i, main effect of

GENOTYPE, $F_{1,73}=0.002$, p=0.97) suggesting an equivalent propensity to explore

the maze, and as expected, all of the mice showed an avoidance of the more

anxiety-inducing open arms of the EPM (Fig. 4j, main effect of ARM, $F_{1,73}=7.41$,

p=0.008), and there were no significant effects of SEX or interactions.

Locomotor activity levels, motoric function, and novel object recognition are
not altered in *Setd1a*$^{+/-}$ mice.

Locomotor activity was assessed over three consecutive days, with a 2 hour session

each day. The number of beam breaks made decreased each day demonstrating

habituation (Fig. 5a, main effect of Day, $F_{1.69,123.08}=28.63$, p<0.001) and no significant

GENOTYPE*DAY interaction ($F_{1.69,123.08}=0.18$, p=0.80), showed that the degree of

habituation of locomotor activity levels across days was equivalent in *Setd1a*$^{+/-}$ and

WT mice. *Setd1a* haploinsufficiency did not alter the overall amount of activty over

the three sessions (main effect of GENOTYPE, $F_{1,73}=0.46$, p=0.50). There was no

significant main effect of SEX ($F_{1,73}=0.40$, p=0.84), and no significant interactions

between GENOTYPE, SEX and DAY (all p>0.05). These data would suggest that

both male and female *Setd1a*$^{+/-}$ mice show normal locomotor activity levels and

habituation of activity levels across test sessions.

Assessment of motor co-ordination using the Rotarod revealed that the latency to fall

from the rotating beam over five accelerating trials was not significantly different

between *Setd1a*$^{+/-}$ and WT mice (Fig. 5b, main effect of GENOTYPE, $F_{1,73}=2.70$,

p=0.11). A significant main effect of TRIAL ($F_{2.97,217.0}=16.99$, p<0.001), indicated that

performance increased with training, which was equivalent between *Setd1a*$^{+/-}$ and

WT mice (GENOTYPE*TRIAL interaction $F_{2.97,217.0}=0.50$, p=0.68). There was also no

significant main effect of SEX ($F_{1,73}=0.33$, p=0.57) and no significant

GENOTYPE*SEX interaction ($F_{1,73}=0.001$, p=0.98). The GENOTYPE*SEX*TRIAL

interaction was also not significant ($F_{2.97,217.0}=2.41$, p=0.07). These data indicate

normal motor co-ordination and motor learning in male and female *Setd1a*$^{+/-}$ mice.

Novel object recognition memory was assessed after a 30 minute and 24 hour

retention interval, and as expected, discrimation ratios were higher after 30 minutes

compared to 24 hours (Fig. 5c, main effect of DELAY, $F_{1,72}=19.61$, p<0.001). There

was no significant main effect of GENOTYPE ($F_{1,72}=0.05$, p=0.82) and no

GENOTYPE*DELAY interaction ($F_{1,72}$=1.68, p=0.20), suggesting no differences in

recall by *Setd1a*$^{+/-}$ mice. There were no significant effects of SEX ($F_{1,72}$=0.28,

p=0.61) or significant interactions between these different factors on the discrimation

ratios. Object exploration in these sessions, indexed by duration per contact (Fig.

5d) also showed no differences between WT and *Setd1a*$^{+/-}$ mice (main effect of

GENOTYPE, $F_{1,72}$=0.01, p=0.97), although all mice showed a reduced tendency to

explore the objects in the 24 hours test (main effect of DELAY, $F_{1,72}$=27.13,

p<0.001). There were also no significant effects of SEX ($F_{1,72}$=0.01, p=0.97) or

interactions between the different factors of the ANOVA. Analysis of the acquisition

time data (Supplementary Results Table 2) revealed no significant effect of genotype

on the time taken to achieve 40 seconds of object exploration ($F_{1,72}$=0.09, p=0.76) or

between the two sessions ($F_{1,72}$=0.12, p=0.091), suggesting that all the mice

explored the objects equally in both delay conditions. These findings indicate intact

short- and long-term object recognition memory in both male and female *Setd1a*$^{+/-}$

mice.

Discussion

Here, we examined the developmental and behavioural consequences of

haploinsufficiency of *Setd1a* in a novel mouse model. We provide evidence for a

complex pattern of sex-related differences spanning the pre- and post-natal period:

reduced placental size in males and females at different foetal ages, post-natal

catch-up growth in males, and effects on adult behaviour. Whole brain RNA-Seq

analysis at E13.5 substantiate previous findings on the impact of Setd1a loss of

function (LoF) on markers of mitochondrial function. With a literature demonstrating

Setd1a as a risk candidate for a schizophrenia, we observed a number of

phenotypes of relevance, including increased anxiety-related behaviour, enhanced

acoustic startle response, and decreased pre-pulse inhibition of acoustic startle.

These abnormalities strengthen the support for the use of *Setd1a* haploinsufficient

mice as a model for the biological basis of schizophrenia, although the neural

mechanisms underpinning this may be complex and involve pre-natal effects.

Using the same model of *Setd1a* LoF as we have used here, Clifton et al., 2022 [25]

showed gene expression patterns in cortical samples from E14 through to P70 with

consistent downregulation in genes enriched for mitochondrial function. We have
replicated these findings at a similar developmental time point and demonstrate a
similar relationship with mitochondrial GO terms. Altered mitochondrial gene
expression changes have also been observed in an RNAi SETD1A knockdown
cellular model [4,36] and mitochondrial dysfunction has been repeatedly implicated
in the pathogenesis of schizophrenia [38-40]. A recent study using heterozygous
disruption of SETD1A in hiPSC-derived neurones demonstrated a metabolic
dysfunction that affected neurite growth and differentiation [41]. Common variants for
schizophrenia are dynamically expressed during neurodevelopment, with high
expression in foetal brain [42-44], and the cellular heterogeneity of the E13.5 mouse
brain may have masked subtle gene expression changes, and also may have
reduced the number of differentially expressed genes we were able to observe, in
comparison with previous studies [17,19,25]. However, we were also able to identify
differential expression in several nominally significant cilium-related GO terms, in
addition to the mitochondrial markers. Cilia play an essential role during
neurodevelopment [45] and knockdown of a range of neuropsychiatric risk genes
affects cilia formation *in vitro* [46]. Taken together, these findings provide further
evidence that *Setd1a* haploinsufficiency, early in development, may have a more
generic effect on biological mechanisms, rather than specifically targeting pathways
known to be implicated in schizophrenia.

Our data demonstrate that foetal growth was unaffected by *Setd1a*
haploinsufficiency through gestation, with the expected weight increases up to birth.
However, there were some interesting differences in placental weight with, on
average, smaller placentae in *Setd1a*[+/-] embryos across all time points evaluated,
which led to increased embryo:placenta ratios (EPR) in *Setd1a*[+/-] embryos relative to
WT.   These differences were most notable at E11.5, where male *Setd1a*[+/-]
placentae were ~32% smaller than WT and at E18.5 where female *Setd1a*[+/-]
placentae were ~16% smaller than both male and female WT placentae. Smaller
sizes may reflect impairment in function, and with the placenta acting as the vital
conduit for nutrition, this could imply a paucity of essential substances, such as
neurotransmitters and amino acids, at key developmental stages [47]. This may be
further impacted by the fact that foetal size remains unaltered, which could suggest
an imbalance between foetal demands and the ability of the placentae to match this

need, which could inhibit developmental progression [48]. For example, using a

mouse model of *Igf2* dysfunction, where only placental expression was reduced [49],

we demonstrated that a foetus:placenta imbalance can lead to adult-related changes

in anxiety-related behaviours [48]. In this model, placental size is reduced (~25%

E12 to E19) but the foetus only begins to show intrauterine growth restriction (IUGR)

late on at E19 leading to a low birth weight (31% of WT, a period of catch-up growth

to match adult weight by P100 [48,49]. Other models show similar patterns [47].

Thus, a mismatch between foetal demands and placental function can lead to IUGR,

low birth weight, a period of exaggerated early (catch-up) growth and adult-related

changes in behaviour. We observed similar patterns of pre-natal and post-natal

growth indices in our model of *Setd1a* haploinsufficiency, although we did not

measure birth and early post-natal weights, but male mice demonstrated a period of

catch-up growth adult. We also observed more complex effects suggesting a sex-

related change in placental function, which could affect later behaviour in a sex-

related manner [47,50]. IUGR, as an indicator of placental dysfunction has been

implicated in a number of neuropsychiatric conditions, such as schizophrenia, autism

and other neurodevelopmental disorders, with the association of conceptus sex also

influencing placental (dys)functionality [47,51,26].

In terms of effects on adult behaviour, we demonstrated comparable patterns on

acoustic startle responding (ASR) and the gating of this response by a prepulse

stimulus (prepulse inhibition, PPI), to previous studies in *Setd1a* haploinsufficiency

mice [18]. Deficits in attention and gating, or filtering out intrusive stimuli, are

prominent features shared by both psychiatric and developmental disorders [53-55],

with which haploinsufficiency of *SETD1A* is associated [14,22], therefore the

consistent effect from our and other's work [17,18] in *Setd1a* haploinsufficiency mice

of disrupted PPI may be expected. Altered ASR in male *Setd1a*[+/-] mice only was also

found by Chen et al [18]. No effects on ASR were found in the Naga et al study [17],

but the sex of the mice is not clear from the methods, and results are not split by sex.

We replicated our initial finding in a separate cohort of male and female mice

(Supplementary Methods Figure 2) thus, it is possible to infer a sex-dependent effect

on increased ASR and a sex-independent effect on PPI. In terms of startle

responding, the elevated response we observe in male *Setd1a*[+/-] mice may reflect

initial sensitisation to a fearful stimulus, and reduced habituation to the startle stimuli

through the test session. Increased startle responding may be related to an altered

fear response and/or altered emotional processing via involvement of the amygdala

[56] and may reflect hypervigilance [57] which could link to the subtle anxiogenic

effects we observed. There is some evidence for an interaction between sex and

early life adversity, as demonstrated by the *Igf2*-P0 IUGR model previously

discussed [48], and with male rats demonstrating increased ASR following maternal

separation [58,59] that could also be relevant to the developmental changes we have

observed. Increased startle responding has been observed in neurodevelopmental

conditions such as autism and Fragile X syndrome [53], in addition to PTSD [60] and

schizophrenia [61]. These conditions commonly present more in males than females,

so this phenotype might be expected, however, studies have more commonly

observed increased startle responding in female subjects compared to males

[62,63].

Furthermore, we also demonstrated sex-dependent sensitivity to the typical anti-

psychotic drug risperidone, with male *Setd1a$^{+/-}$* mice showing a blunted effect to the

drug on the ASR, whereas the drug reduced ASR in a dose-dependent way in WTs

and female *Setd1a$^{+/-}$* mice. However, there is little evidence for sex-related effects of

acute risperidone administration, but chronic dosing appears to show a female bias,

with female rats appearing to be more sensitive to chronic dosing at a young age

[64] and female schizophrenia patients responded better to chronic risperidone

treatment than male patients [65]. The mode of action of risperidone, with high

activity via serotonergic 5-HT$_{2A}$ receptors [65] suggests an interesting target to

investigate in the future, especially in male *Setd1a$^{+/-}$* mice. Haloperidol, also a typical

anti-psychotic drug, though mainly dopaminergic in function [66] had no effects on

the ASR in male (or female) *Setd1a$^{+/-}$* mice. Sensorimotor gating deficits (PPI) could

not be rescued by haloperidol or risperidone, although both anti-psychotics have

previously been shown to reverse PPI deficits in several rodent models of

schizophrenia [66]. This suggests that both these anti-psychotic agents are

ineffective for ameliorating schizophrenia-relevant phenotypes in *Setd1a$^{+/-}$* mice, of

both sexes, and point to deficits in neural systems other than the monoamine system

which may be consistent with variable responses to antipsychotic treatment in

individuals with *SETD1A* variant-associated schizophrenia [16]. Of particular

relevance here, is the similarity of the disrupted PPI shown by *Setd1a$^{+/-}$* LoF models

and the effects of NMDAR antagonists, such as PCP, MK-801 and ketamine [68],
which taken together with the possibility that glutamate release is likely reduced
when *Setd1a* is knocked down in postsynaptic pyramidal neurons [17], suggests that
the glutamatergic system may be compromised as a result of Setd1a
haploinsufficiency [21].

In terms of measures of anxiety, then our data from the open field (OFT) and
elevated plus maze (EPM) assays are the first to demonstrate that *Setd1a* LoF can
lead to a subtle anxiogenic phenotype. Exploration patterns in these tests
demonstrated that both male and female *Setd1a*[+/-] mice showed avoidance of the
more anxiety-inducing regions of each piece of apparatus, the central zone of the
OFT and the open arms of the EPM. Previous studies using the OFT [17-19] have
used a procedure more akin to our locomotor activity assessment, with repeated
long sessions and found, as we have, that basic motoric behaviour and investigation
is not affected. Within session measures of activity from the OFT and EPM also
confirm this. It would be interesting if in these studies [17-19] the first few minutes of
the first sessions could be re-examined to gauge if any anxiety phenotype was
present. Taken together, these data illustrate subtle effects of *Setd1a* LoF on
anxiety, which is consistent with emotional issues as co-morbid presentations in
schizophrenia [68]. The other adult behaviour we investigated recapitulated previous
findings in *Setd1a*[+/-] haploinsufficient mice, with no significant effects on novel object
discrimination and recall (NOR), and no profound motoric impairments in either male
or female mice. NOR performance was equivalent to Mukai et a [19] with no
differential recall with a short or long retention delay. However, in the Nagahama et
al [17] study there was an impairment at the short delay, but not the longer timepoint
possibly indicating a subtle phenotype that is subsumed by the difference between
their model, and the ones used here and by Mukai et al [19] which may reflect the
different targeting strategies, and loci, for disrupting *Setd1a* [21].

Taken together these data demonstrate that this new *Setd1a* LoF mouse model
recapitulates some of the core findings of previous models, of relevance to
neurodevelopmental disorders and schizophrenia in a large experimental cohort of
both sexes. We also provide some insight into *Setd1a* haploinsufficiency on
development and how it might interact with sex. The neurodevelopmental hypothesis

of schizophrenia highlights the pre-and peri-natal phase of life as key period in the onset of the disorder, with intrauterine compromise an important factor [70,71]. Sex differences in schizophrenia are well documented, in terms of incidence and symptom presentation [72], thus the altered developmental indices in *Setd1a*^+/- conceptuses and behavioural differences we found add a further dimension to this theory. Taken together our findings would suggest pre-natal effects of *Setd1a* LoF on adult behaviour, mediated by impaired placental functionality and metabolic function in developing neurones. Use of a conditional KO model, where setd1a LoF can be "switched off" either during development or in adulthood would be one way to unpick this conundrum. However, either way, these data add to the growing understanding of the role of *Setd1a* in brain development and behaviour. Our findings, coupled with those from others investigating different *Setd1a* models, demonstrate replicable common phenotypes of relevance that could be used for the development and testing of therapies in rescue studies.

Data Availability

All data are publicly available at the following 'Open Science Framework' link:

https://osf.io/gf3d7/

Funding

This work was supported by a Wellcome Trust Integrative Neuroscience PhD grant (109084/Z/15/Z) and an A. Bruce Naylor Memorial Early Career Research Fellowship from The Waterloo Foundation awarded to MLB; UKRI Medical Research Council (MRC) IMPC: Pump Priming Award (MR/P026176/1) awarded to ARI, LSW and TH; ARI, TH and LSW are members of the MRC Centre for Neuropsychiatric Genetics and Genomics (MR/L010305/1).

Author contribution

MLB, ARI, LSW and TH conceived the project. MLB and TH planned the behavioural experiments, all of which were performed by MLB; Tissue collection, processing and

781 analysis was performed by MLB, with input from ARI. MLB and TH analysed the data
782 and drafted the paper, with editing input from ARI and LSW.
783

784 **Conflict of interest statement**
785 The authors declare no competing financial or non-financial interests.
786

787 **Acknowledgements**

References

Noma K, Allis CD, Grewal SI. (2001) Transitions in distinct histone H3 methylation patterns at the heterochromatin domain boundaries. *Science*, 293(5532):1150-5. doi: 10.1126/science.1064150.

Wang S, Bleeck A, Nadif Kasri N, Kleefstra T, van Rhijn JR, Schubert D. (2021) SETD1A Mediated H3K4 Methylation and Its Role in Neurodevelopmental and Neuropsychiatric Disorders. *Frontiers in Molecular Neuroscience,* 4:772000. doi: 10.3389/fnmol.2021.772000. eCollection 2021.

Tajima, K, Matsuda, S, Yae, T, Drapkin, BJ, Morris, R, Boukhali, M, … Maheswaran, S. (2019) SETD1A protects from senescence through regulation of the mitotic gene expression program. *Nature Communications*, **10**, 2854.

Hoshii, T, Cifani, P, Feng, Z, Huang, C-H, Koche, R, Chen, C-W., … Armstrong, SA. (2018). A Non-catalytic Function of SETD1A Regulates Cyclin K and the DNA Damage Response. *Cell*, *172*(5), 1007-1021.e17. doi: 10.1016/j.cell.2018.01.032

Tajima, K, Yae, T, Javaid, S, Tam, O, Comaills, V, Morris, R, … Maheswaran, S. (2015) SETD1A modulates cell cycle progression through a miRNA network that regulates p53 target genes. *Nature Communications*, **6**, 8257.

Sze, CC, Cao, K, Collings, CK, Marshall, SA, Rendleman, EJ, Ozark, PA, Chen, FX, Morgan, MA, Wang, L and Shilatifard, A. (2017) Histone H3K4 methylation-dependent and -independent functions of Set1A/COMPASS in embryonic stem cell self-renewal and differentiation. *Genes & Development*, **31**, 1732-1737.

Fang, L, Zhang, J, Zhang, H, Yang, X, Jin, X, Zhang, L, Skalnik, DG, Jin, Y, Zhang, Y, Huang, X, Li,, J and Wong, J (2016) H3K4 Methyltransferase Set1a is a key Oct4 coactivator essential for generation of Oct4 positive inner cell mass. *Stem Cells*, **34**, 565-580.

Bledau, AS, Schmidt, K, Neumann, K, Hill, U, Ciotta, G, Gupta, A, Torres, DC, Fu, J, Kranz, A, Stewart, AF and Anastassiadis, K (2014) The H3K4 methyltransferase Setd1a is first required at the epiblast stage, whereas Setd1b becomes essential after gastrulation. *Development*, **141**, 1022-1035.

Qiao, H, Li, Y, Feng, C, Duo, S, Ji, F, & Jiao, J (2018). Nap1l1 Controls Embryonic Neural Progenitor Cell Proliferation and Differentiation in the Developing Brain. *Cell Reports*, *22*(9), 2279–2293. doi: 10.1016/j.celrep.2018.02.019

Li, Y, & Jiao, J. (2017). Histone chaperone HIRA regulates neural progenitor cell proliferation and neurogenesis via β-catenin. *Journal of Cell Biology*, *216*(7), 1975–1992. doi: 10.1083/jcb.201610014

Alsulami, M, Munawar, N, Dillon, E, Oliviero, G, Wynne, K, Alsolami, M, … Cagney G. (2019) SETD1A Methyltransferase Is Physically and Functionally Linked to the DNA Damage Repair Protein RAD18. *Molecular Cell Proteomics*, **18**, 1428-1436.

Higgs, MR, Sato, K, Reynolds, JJ, Begum, S, Bayley, R, Goula, A, … Stewart GS. (2018) Histone Methylation by SETD1A Protects Nascent DNA through the Nucleosome Chaperone Activity of FANCD2. *Molecular Cell*, **71**, 25-41 e26.

Singh, T, Kurki, MI, Curtis, D, Purcell, SM, Crooks, L, McRae, J, … Barrett JC. (2016) Rare loss-of-function variants in SETD1A are associated with schizophrenia and developmental disorders. *Nature Neuroscience*, **19**, 571-577.

Singh T, Poterba T, Curtis D, Akil H, Al Eissa M, Barchas JD, … Daly MJ (2022) *Nature*,

604(7906):509-516. doi: 10.1038/s41586-022-04556-w.

Takata, A, Xu, B, Ionita-Laza, I, Roos, JL, Gogos, JA and Karayiorgou, M. (2014) Loss-of-function variants in schizophrenia risk and SETD1A as a candidate susceptibility gene. *Neuron*, 82, 773-780.

Colijn MA, Carrion P, Poirier-Morency G, Rogic S, Torres I, Menon M … Stowe RM. (2024) SETD1A variant-associated psychosis: A systematic review of the clinical literature and description of two new cases. Prog Neuropsychopharmacol Biol Psychiatry. 129:110888.

Nagahama, K, Sakoori, K, Watanabe, T, Kishi, Y, Kawaji, K, Koebis, M, … Kano M. (2020) Setd1a Insufficiency in Mice Attenuates Excitatory Synaptic Function and Recapitulates Schizophrenia-Related Behavioral Abnormalities. *Cell Reports*, 32, 108126.

Chen R, Liu Y, Djekidel MN, Chen W, Bhattacherjee A, Chen Z, Scolnick E, Zhang Y. (2022) Cell type-specific mechanism of Setd1a heterozygosity in schizophrenia pathogenesis. *Science Advances,* 28(9):eabm1077. doi: 10.1126/sciadv.abm1077.

Mukai J, Cannavò E, Crabtree GW, Sun Z, Diamantopoulou A, Thakur P, Chang CY, Cai Y, Lomvardas S, Takata A, Xu B, Gogos JA. (2019) Recapitulation and Reversal of Schizophrenia-Related Phenotypes in Setd1a-Deficient Mice. *Neuron,* 104(3):471-487.e12. doi: 10.1016/j.neuron.2019.09.014.

Hamm, JP., Shymkiv, Y, Mukai, J, Gogos, JA and Yuste, R. (2020) Aberrant Cortical Ensembles and Schizophrenia-like Sensory Phenotypes in Setd1a(+/-) Mice. *Biological Psychiatry,* 88, 215-223.

Wang S, Rhijn JV, Akkouh I, Kogo N, Maas N, Bleeck A, Ortiz IS, Lewerissa E, Wu KM, Schoenmaker C, Djurovic S, van Bokhoven H, Kleefstra T, Nadif Kasri N, Schubert D. (2022) Loss-of-function variants in the schizophrenia risk gene SETD1A alter neuronal network activity in human neurons through the cAMP/PKA pathway. *Cell Reports,* 39(5):110790. doi: 10.1016/j.celrep.2022.110790.

Kummeling, J, Stremmelaar, DE, Raun, N, Reijnders, MRF, Willemsen, MH, Ruiterkamp-Versteeg, M, … Kleefstra T. (2021) Characterization of SETD1A haploinsufficiency in humans and Drosophila defines a novel neurodevelopmental syndrome. *Molecular Psychiatry*, **26**, 2013-2024. DOI: 10.1038/s41380-020-0725-5

Sullivan, EL, Riper, KM, Lockard, R, & Valleau, JC. (2015). Maternal high-fat diet programming of the neuroendocrine system and behavior. *Hormones and Behavior*, *76*, 153–161. doi: 10.1016/j.yhbeh.2015.04.008

Gabory, A, Ferry, L, Fajardy, I, Jouneau, L, Gothié, J-D., Vigé, A, … Junien, C. (2012). Maternal Diets Trigger Sex-Specific Divergent Trajectories of Gene Expression and Epigenetic Systems in Mouse Placenta. *PLoS ONE, 7*(11), e47986. doi: 10.1371/journal.pone.0047986

Clifton NE, Bosworth ML, Haan N, Rees E, Holmans PA, Wilkinson LS, Isles AR, Collins MO, Hall J. (2022) Developmental disruption to the cortical transcriptome and synaptosome in a model of SETD1A loss-of-function. *Human Molecular Genetics,* 31(18):3095-3106. doi: 10.1093/hmg/ddac105.

Schwenk F, Baron U, Rajewsky K (1995) A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. *Nucleic Acids Research,* 23(24):5080-1. doi: 10.1093/nar/23.24.5080.

Andrews, S. (2010). FastQC A Quality Control tool for High Throughput Sequence Data. Retrieved from https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Love, MI, Huber, W, & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. doi:

10.1186/s13059-014-0550-8

Huang, DW, Sherman, BT, & Lempicki, RA. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, *4*(1), 44–57. doi: 10.1038/nprot.2008.211

de Leeuw, CA, Mooij, JM, Heskes, T, & Posthuma, D. (2015). MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLOS Computational Biology*, *11*(4), e1004219. doi: 10.1371/journal.pcbi.1004219

Pardiñas, AF, Holmans, P, Pocklington, AJ, Escott-Price, V, Ripke, S, Carrera, N, … Walters, JTR. (2018). Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nature Genetics*, *50*(3), 381–389. doi: 10.1038/s41588-018-0059-2

Humby T, Laird FM, Davies W, Wilkinson LS. (1999) Visuospatial attentional functioning in mice: interactions between cholinergic manipulations and genotype. European Journal of Neuroscience, 11(8):2813-23. doi: 10.1046/j.1460-9568.1999.00701.x

Grayson, B, Leger, M, Piercy, C, Adamson, L, Harte, M, & Neill, JC. (2015). Assessment of disease-related cognitive impairments using the novel object recognition (NOR) task in rodents. *Behavioural Brain Research*, *285*, 176–193. doi: 10.1016/j.bbr.2014.10.025

Geyer, MA, Krebs-Thomson, K, Braff, DL, & Swerdlow, NR. (2001). Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: a decade in review. *Psychopharmacology*, *156*(2–3), 117–154. doi: 10.1007/s002130100811

Swerdlow, NR, Weber, M, Qu, Y, Light, GA, & Braff, DL. (2008). Realistic expectations of prepulse inhibition in translational models for schizophrenia research. *Psychopharmacology*, *199*(3), 331–388. doi: 10.1007/s00213-008-1072-4

Silver LM 1995 In Mouse Genetics – Concepts and Applications, ch 4.1. Oxford: Oxford University Press.

Cameron, D, Blake, DJ, Bray, NJ, & Hill, MJ. (2019). Transcriptional Changes following Cellular Knockdown of the Schizophrenia Risk Gene SETD1A Are Enriched for Common Variant Association with the Disorder. *Molecular Neuropsychiatry*, *5*(2), 109–114. doi: 10.1159/000497181

Ben-Shachar, D. (2017). Mitochondrial multifaceted dysfunction in schizophrenia; complex I as a possible pathological target. *Schizophrenia Research*, *187*, 3–10. doi: 10.1016/J.SCHRES.2016.10.022

Flippo, KH, & Strack, S. (2017). An emerging role for mitochondrial dynamics in schizophrenia. *Schizophrenia Research*, *187*, 26–32. doi: 10.1016/J.SCHRES.2017.05.003

Rajasekaran, A, Venkatasubramanian, G, Berk, M, & Debnath, M. (2015). Mitochondrial dysfunction in schizophrenia: Pathways, mechanisms and implications. *Neuroscience & Biobehavioral Reviews*, *48*, 10–21. doi: 10.1016/J.NEUBIOREV.2014.11.005

Chong ZS, Khong ZJ, Tay SH, Ng SY. (2022) Metabolic contributions to neuronal deficits caused by genomic disruption of schizophrenia risk gene SETD1A. *Schizophrenia (Heidelb)*. 29;8(1):115. doi: 10.1038/s41537-022-00326-9.

Birnbaum, R, Jaffe, AE, Chen, Q, Hyde, TM, Kleinman, JE, & Weinberger, DR. (2015). Investigation of the Prenatal Expression Patterns of 108 Schizophrenia-Associated Genetic Loci. *Biological Psychiatry*, *77*(11), e43–e51. doi: 10.1016/j.biopsych.2014.10.008

Clifton, NE, Hannon, E, Harwood, JC, Di Florio, A, Thomas, KL, Holmans, PA, … Hall, J. (2019). Dynamic expression of genes associated with schizophrenia and bipolar disorder

across development. *Translational Psychiatry*, *9*(1), 74. doi: 10.1038/s41398-019-0405-x

Jaffe, AE, Straub, RE, Shin, JH, Tao, R, Gao, Y, Collado-Torres, L, … Weinberger, DR. (2018). Developmental and genetic regulation of the human cortex transcriptome illuminate schizophrenia pathogenesis. *Nature Neuroscience*, *21*(8), 1117–1125. doi: 10.1038/s41593-018-0197-y

Guemez-Gamboa, A, Coufal, NG, & Gleeson, JG. (2014). Primary Cilia in the Developing and Mature Brain. *Neuron*, *82*(3), 511–521. doi: 10.1016/j.neuron.2014.04.024

Marley A, von Zastrow M. (2012) A simple cell-based assay reveals that diverse neuropsychiatric risk genes converge on primary cilia. *PLoS One*, 7(10):e46647. doi: 10.1371/journal.pone.0046647.

Rosenfeld CS. (2021) The placenta-brain-axis. *Journal of Neuroscience Research,* 99(1):271-283. doi: 10.1002/jnr.24603.

Mikaelsson MA, Constância M, Dent CL, Wilkinson LS, Humby T (2013) Placental programming of anxiety in adulthood revealed by Igf2-null models. Nature Communications, 4:2311. doi: 10.1038/ncomms3311.

Constância, M, Hemberger, M, Hughes, J, Dean, W, Ferguson-Smith, A, Fundele, R, … Reik, W. (2002). Placental-specific IGF-II is a major modulator of placental and fetal growth. *Nature*, *417*(6892), 945–948. doi: 10.1038/nature00819

Lodefalk M, Chelslín F, Patriksson Karlsson J, Hansson SR. (2023) Placental Changes and Neuropsychological Development in Children-A Systematic Review. *Cells*, 12(3):435. doi: 10.3390/cells12030435.

Ursini, G, Punzi, G, Chen, Q, Marenco, S, Robinson, JF, Porcelli, A, … Weinberger, DR. (2018). Convergence of placenta biology and genetic risk for schizophrenia. *Nature Medicine*, *24*(6), 792–801. doi: 10.1038/s41591-018-0021-y

Al-Haddad BJS, Oler E, Armistead B, Elsayed NA, Weinberger DR, Bernier R, Burd I, Kapur R, Jacobsson B, Wang C, Mysorekar I, Rajagopal L, Adams Waldorf KM. (2019) The fetal origins of mental illness. American Journal of Obstetrics & Gynecology, 221(6):549-562. doi: 10.1016/j.ajog.2019.06.013.

Sinclair D, Oranje B, Razak KA, Siegel SJ, Schmid S (2017) Sensory processing in autism spectrum disorders and Fragile X syndrome-From the clinic to animal models. Neuroscience & Biobehavior Reviews, 76(Pt B):235-253. doi: 0.1016/j.neubiorev.2016.05.029.

Javitt DC. (2009) Sensory processing in schizophrenia: neither simple nor intact. Schizophr Bull. 35(6):1059-64. doi: 10.1093/schbul/sbp110.

Perry W, Minassian A, Lopez B, Maron L, Lincoln A. (2007) Sensorimotor gating deficits in adults with autism. Biological Psychiatry, 61(4):482-6. doi: 10.1016/j.biopsych.2005.09.025.

Koch, M. (1999). The neurobiology of startle. *Progress in Neurobiology*, *59*(2), 107–128. doi: 10.1016/S0301-0082(98)00098-7

Götz T, Janik VM. (2011) Repeated elicitation of the acoustic startle reflex leads to sensitisation in subsequent avoidance behaviour and induces fear conditioning. *BMC Neuroscience*, 12:30. doi: 10.1186/1471-2202-12-30.

de Jongh R, Geyer MA, Olivier B, Groenink L. (2005) The effects of sex and neonatal maternal separation on fear-potentiated and light-enhanced startle. *Behavioral Brain Research*, 161(2):190-6. doi: 10.1016/j.bbr.2005.02.004.

Granata L, Parakoyi A, Brenhouse HC. (2022) Age- and sex-specific effects of maternal separation on the acoustic startle reflex in rats: early baseline enhancement in females

and blunted response to ambiguous threat. *Frontiers in Behavioral Neuroscience,* 16:1023513. doi: 10.3389/fnbeh.2022.1023513. eCollection 2022.

Glenn DE, Acheson DT, Geyer MA, Nievergelt CM, Baker DG, Risbrough VB; MRS Team. (2016) High and low threshold for startle reactivity associated with PTSD symptoms but not PTSD risk: evidence from a prospective study of active duty marines. *Depression Anxiety*, 33(3):192-202. doi: 10.1002/da.22475.

Meincke U, Light GA, Geyer MA, Braff DL, Gouzoulis-Mayfrank E. (2004) Sensitization and habituation of the acoustic startle reflex in patients with schizophrenia. *Psychiatry Research*, 126(1):51-61. doi: 10.1016/j.psychres.2004.01.003.

de Jongh R, Geyer MA, Olivier B, Groenink (2005) The effects of sex and neonatal maternal separation on fear-potentiated and light-enhanced startle. L. *Behavioral Brain Research*, 161(2):190-6. doi: 10.1016/j.bbr.2005.02.004.

Blanch A, Lucas I, Balada F, Blanco E, Aluja A. (2018) Sex differences and personality in the modulation of the acoustic startle reflex. *Physiology & Behavior,* 195:20-27. doi: 10.1016/j.physbeh.2018.07.020.

Bardgett ME, Downnen T, Crane C, Baltes Thompson EC, Muncie B, Steffen SA, Yates JR, Pauly JR. (2020) Chronic risperidone administration leads to greater amphetamine-induced conditioned place preference. *Neuropharmacology*, 179:108276. doi: 10.1016/j.neuropharm.2020.108276.

Liu H, Liu H, Jiang S, Su L, Lu Y, Chen Z, Li X, Li X, Wang X, Xiu M, Zhang X. (2022) Sex-Specific Association between Antioxidant Defense System and Therapeutic Response to Risperidone in Schizophrenia: A Prospective Longitudinal Study. *Current Neuropharmacology,* 20(9):1793-1803. doi: 10.2174/1570159X19666211111123918.

Cunningham Owens, D, & Johnstone, EC. (2018). The development of antipsychotic drugs. *Brain and Neuroscience Advances*, *2*, 1–6. doi: 10.1177/2398212818817498

Geyer, MA, Krebs-Thomson, K, Braff, DL and Swerdlow, NR. (2001) Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: a decade in review. *Psychopharmacology (Berl)*, 156, 117-154.

Wu, M, Wang, PF, Lee, JS, Martin-Brown, S, Florens, L, Washburn, M, & Shilatifard, A. (2008). Molecular Regulation of H3K4 Trimethylation by Wdr82, a Component of Human Set1/COMPASS. *Molecular and Cellular Biology*, *28*(24), 7337–7344. doi: 10.1128/MCB.00976-08

Braga, RJ, Reynolds, GP, & Siris, SG. (2013). Anxiety comorbidity in schizophrenia. *Psychiatry Research*, *35*(2), 383–402. doi: 10.1016/j.psychres.2013.07.030

Murray, RM. and Lewis, SW. (1987). Is schizophrenia a neurodevelopmental disorder? *British Medical Journal,* 295, 681-682. doi:10.1136/bmj.295.6600.681

Weinberger, DR. (1987). Implications of normal brain development for the pathogenesis of schizophrenia. *Archives of General Psychiatry*, 44, 660-669. doi:10.1001/archpsyc.1987.01800190080012

Freeman HB, Lee J. (2023) Sex Differences in Cognition in Schizophrenia: What We Know and What We Do Not Know. *Current Topics in Behavioral Neuroscience*, 63:463-474. doi: 10.1007/7854_2022_394.

798   END OF DOCUMENT

Figure 1

Figure 2

Table 1. Results of gene ontology (GO) term enrichment analysis.

Category	GO term	Gene count	*P value*	Benjamini Hochberg
Cellular component	Mitochondrion (GO: 0005739)	47	0.00001	0.002
	Cilium (GO: 005929)	10	0.004	0.35
	Ciliary basal body (GO: 0036064)	7	0.005	0.30
	Mitochondrial matrix (GO: 0005759)	8	0.02	0.74
	Integral component of mitochondrial inner membrane (GO: 0031305)	3	0.04	0.80
Biological process	Peptidyl-lysine deacetylation (GO: 0034983)	3	0.006	0.98
	Methylation (GO: 0032259)	8	0.01	0.99
	Cilium morphogenesis (GO: 0060271)	8	0.01	0.95
	Cell projection organization (GO: 0030030)	7	0.02	0.97
	Cilium assembly (GO: 0060271)	6	0.04	1.0
	Nucleotide metabolic process (GO: 0009117)	3	0.04	1.0
Molecular Function	Methyltransferase activity (GO: 0008168)	8	0.009	0.95
	Nucleic acid binding (GO: 0003676)	26	0.01	0.80
	NAD-dependent histone deacetylase activity (H3-K14 specific) (GO: 0032041)	3	0.02	0.82
	Transferase activity (GO: 0016740)	29	0.02	0.84
	Nucleotidyltransferase activity (GO: 0016779)	5	0.03	0.86
	Aspartate-tRNA ligase activity (GO: 0004815)	2	0.03	0.82

Figure 3

Figure 4

Figure 5

a.

b.

c.

d.

Male WT (N=21) Female WT (N=20)
Male Setd1a$^{+/-}$ (N=18) Female Setd1a$^{+/-}$ (N=18)