North and East African mitochondrial genetic variation needs further characterization towards precision medicine

Anke Fähnrich[1,*], Isabel Stephan[1,*], Misa Hirose[2], Mosab Ali Awadelkareem[3,4], Saleh Ibrahim[2,5], Hauke Busch[1,6,+], Inken Wohlers[1,+§]

[1]Medical Systems Biology Division, Lübeck Institute of Experimental Dermatology and Institute for Cardiogenetics, University of Lübeck, Ratzeburger Allee 160, 23562, Lübeck, Germany

[2]Genetics Division, Lübeck Institute of Experimental Dermatology, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

[3]Faculty of Medical Laboratory Sciences, Al-Neelain University, Khartoum, Sudan

[4]Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford OX3 9DU, United Kingdom

[5]Sharjah Institute for Medical Research, University of Sharjah, Sharjah, United Arab Emirates

[6]University Cancer Center Schleswig-Holstein, University Hospital of Schleswig-Holstein, Campus Lübeck, 23538 Lübeck, Germany

[*]Equal contributing first authors

[+] Equal contributing last authors

[§] Correspondence to Inken.Wohlers@uni-luebeck.de

Abstract

Mitochondria are maternally inherited cell organelles with their own genome, and perform various functions in eukaryotic cells such as energy production and cellular homeostasis. Due to their inheritance and manifold biological roles in health and disease, mitochondrial genetics serves a dual purpose of tracing the history as well as disease susceptibility of human populations across the globe. This requires a comprehensive catalogue of commonly observed genetic variations in the mitochondria for all regions throughout the world. So far, however, certain regions, such as North and East Africa have been understudied. Towards this, we have created the most comprehensive quality-controlled North and East African mitochondrial dataset to date by compiling 11 published cohorts with novel data of mitochondrial genomes from 159 Sudanese individuals. We combined these 641 mitochondrial sequences with sequences from the 1000 Genomes (n=2,504) and the Human Genome Diversity Project

(n=828) and used the tool haplocheck for extensive quality control and detection of in-sample

contamination. Using a subset of high-coverage mitochondrial sequences we predict 15

potentially novel haplogroups in North and East African subjects and observe likely

phylogenetic deviations from the established PhyloTree reference for haplogroups L0a1 and

L2a1. This demonstrates common hitherto unexplored variants in mitochondrial genomes of

the North and East African region that lead to novel phylogenetic relationships, calling for

further in-depth population genetic studies in that region.

Author summary

Does currently available human mitochondrial genetic data sufficiently cover North and East

African diversity to investigate precision medicine for mitochondria-related diseases? To

address this question, we have integrated and compared novel with existing sequence data

from that region and world-wide. We find that variant profiles, i.e. haplogroups, carried by

North and East Africans are shared with or related to those observed in Europe, Africa and

Asia alike. Nonetheless, for a few haplogroups, world-wide data is inaccurate and incomplete

for North and East Africans. Thus, we call for additional, future studies to more

comprehensively represent human mitochondrial genetic diversity towards implementing

precision medicine.

Introduction

Mitochondria are energy-producing, double-membrane-bound organelles that contain their

own genome and that are central to many processes in eukaryotic cells (1). The mitochondrial

(MT) genome is circular and typically 16,569 bases long in humans (2). Mitochondrial

dysfunction is often linked to inefficient oxidative phosphorylation (OXPHOS) capacity with

compromised ATP and increased mitochondrial reactive oxygen species (ROS) production that

change cellular function and signaling (3). Dysfunction accompanies various pathologic

conditions, such as metabolic diseases, neurodegenerative disorders (4), and autoimmune

condition as well as cancer (5) and aging (6). Its root cause might stem from variations of the

MT genome, which has thus moved into the center of research as determining factor for

complex diseases and precision medicine.

Since mitochondria are exclusively maternally inherited, the mitochondrial phylogeny

represents a maternal genealogy, which has been summarized in an established and widely

applied reference phylogenetic tree called PhyloTree (7,8). Its current version 17 is based on

more than 24,000 mitochondrial sequences and constitutes more than 5,400 haplogroups,

i.e., recurrent genetic variant profiles common in human mitochondria.

Recent studies assessed mitochondrial haplogroups in world-wide human diversity reference

data sets from the 1000 genomes project (1000G; n=2,504) (9,10), the Human Genome

Diversity Project (HGDP; n=623 in (11); n=102 in (12)) and very recently gnomAD (13). Such

mitochondrial phylogenetics studies have traced modern human settlement of the world (14)

and support the hypothesis of Africa as origin of modern humans. The "Out-of-Africa" theory

proposes the so-called "mitochondrial Eve", which represents the mitochondrial

Reconstructed Sapiens Reference Sequence (RSRS) as the phylogenetic root and modern

human origin (15).

In this work, we address the state of mitochondrial reference data from the regions of North

Africa and East Africa. As of now, there is little genetic data available from this world region,

as it has been forgone by large genome projects such as 1000G (16), gnomAD (17) or Topmed

(18). An exception is Egypt, for which an initial reference dataset has been compiled (19) and

efforts are ongoing to construct a much larger Egyptian genome reference towards

implementing genomics-based precision medicine.

To address this reference data limitation, we sequenced *de novo* mitochondrial genomes of a

Sudanese (n=159) and an Egyptian cohort (n=217) and integrated the results with an Egyptian

cohort having whole genome sequencing data (n=110). Both Egyptian cohorts have previously

been investigated for haplogroup distribution (19,20) and are here combined with

mitochondrial sequencing data of ancient Egyptian mummies (21) as well as mitochondrial

sequences from Genbank with origins of North or East African countries.

In addition to providing additional mitochondrial genomes from this understudied region, we

reanalyzed all available sequencing data using a novel tool, called haplocheck (22), to identify

potential sample contamination and phantom variants (23). Our combined and new quality-

controlled data results in 146 exceptionally high-quality and 641 high quality mitochondrial

genome sequences that we used to phylogenetically re-characterize genetic MT variation in

North and East Africa with respect to world-wide MT diversity. We find that the haplogroups

of the region cover nearly all world-wide clades and identified 15 potential novel

subhaplogroups, among them haplogroups L0a1 and L2a1, for which sequences from North
and East Africa suggest a different phylogenetic relationship as described by the reference
PhyloTree.
99
100
101 **Results & Discussion**
102
North and East African mitochondrial data collection and filtering
To comprehensively cover the phylogenetic tree of the North and East African population we
sequenced *de novo* a cohort of n=159 Sudanese individuals which we combined with
mitochondrial sequencing data of n=217 Egyptians. Additional n=110 MT genomes were
extracted from whole genome sequencing data of n=100 and n=10 Egyptians from Pagani et
al. (20) and Wohlers et al. (19), respectively. We further included mapped mitochondrial
sequencing data of n=90 ancient Egyptian mummies from Schuenemann et al. (21); for three
ancient Egyptian mummies WGS data were available (cf. Supplementary Table 1 for details).
As controls, we used two world-wide whole-genome references, 1000G with n=2,504 samples
(16) and HGDP with n=828 samples (24) from which we extracted the mitochondrial
sequences. Further, we obtained quality-controlled and aligned Genbank sequences from
countries in North and East Africa from the Supplement of McInerny et al. (25), which we
manually traced back to their eight respective publications provided in Suppl. Table 1.
Sampling location or ethnicity as well as number of sequences are provided in Figure 1.
Notably, five of eight studies relating to Genbank entries include only specific haplogroups.
These are two studies on haplogroup U6 (Maca-Meyer et al. (26), n=11; Olivieri et al. (27),
n=28) addressing Berbers in Morocco, two studies on haplogroup L3 in Morocco (Harich et al.
(28), n=8) and Ethiopia (Soares et al. (29), n=70) and one study on both M1 and U6 addressing
Morocco (Pennarun et al. (30), n=56). Notably, we also included n=18 Genbank MT sequences
of ancient Moroccans (Fregel et al. (31)). Genbank-related studies that do not focus on a
specific haplogroup are addressing Egypt (Kujanová et al. (32), n=35) and Tunisia (Costa et al.
(33), n=42).
To exclude study-specific batch effects and sequencing errors, we applied a two-step quality
control (QC) filter on all sequencing-based North and East African data sets. This was
important to correctly assess haplogroup distributions and to identify putatively new

haplogroups. For the filter step, we kept all samples from Genbank as well as those (i) having a haplocheck-reported major haplogroup equal to minor haplogroup, and (ii) having contamination assessed and not detected (i.e. "NO"). An overall of 641 samples from four sequencing-based North and East African data sets and from eight Genbank studies pass this filter (cf. Suppl. Table 11 and in Suppl. Table 17 for aligned sequences). For the second QC filtering step, we excluded Genbank sequences, as contamination checks were unfeasible, and additionally required a minimum sequencing coverage greater than 1000x, which is considerably higher than the minimum 600x coverage required by haplocheck to detect a 1% or greater contamination (22). The number of total and filtered samples in each QC step are provided in Figure 1 for every cohort included. For mitochondrial sequences obtained from Genbank, the overall number is provided, only.

Figure 1: Mitochondrial data compiled from North and East Africa. Blue color denotes cohorts referred to as North African, green color cohorts referred to as Northeast African and yellow color cohorts referred to as East African. Cohorts analyzed from raw sequencing data are denoted by three numbers: The overall number of samples, the number of samples remaining after lenient QC filtering and the number of samples remaining after strict QC filtering. Locations with only one number refer to Genbank MT sequences obtained via McInerny et al.

(25). Whenever we do not explicitly refer to Northeast Africa, it is included in North and/or

East Africa.

<u>Haplogroup-based contamination assessment for diverse sequencing settings</u>

We assessed potential contamination and related sequencing and variant calling metrics for

all six sequencing-based datasets, including the world-wide references 1000G and HGDP (cf.

Supplementary Tables 2-9 and 13-16 for details). Coverage of the two latter is high with a

median coverage greater than 10,000X, while the former four cohorts have a significantly

lower coverage of 840, 670, 163 and 992 for the Egyptian MT sequencing, the Egyptian WGS,

the ancient Egyptian mummies and the Sudanese MT sequencing cohorts, respectively, see

Figure 2 b.

Figure 2: Characterization of the six sequencing-based data sets using numbers reported by

haplocheck. a) Histogram of number of samples with haplocheck contamination status not

available ("NA") (gray), not contaminated ("NO") (blue) and predicted contaminated ("YES")

(red). b) Box plots for sequencing coverage. c) Box plots for haplogroup quality of those

samples not predicted to be contaminated. d) Box plots for distance between major and minor

haplogroup within the phylogenetic tree for those samples predicted to be contaminated. e)

Box plots of the contamination levels for samples predicted as contaminated. Box plots display

median and lower/upper quartiles; whiskers denote the most extreme data point no more

than 1.5 times the interquartile range; outliers are data points extending beyond whiskers.

Contamination is identified using the computational tool haplocheck, which identifies

polymorphic sites within samples. It attempts to decompose the variants into a so-called

major and another, minor haplogroup. If such decomposition is possible and coherent, a

contamination of a sample of the major haplogroup with another sample of the minor

haplogroup is predicted and a respective contamination level reported. Interestingly, we

found a large heterogeneity in contamination calls among the studies. Few of the WGS-based

1000G samples are predicted to be contaminated, yet more than 30% of the WGS-based HGDP

samples (Figure 2a), albeit at a low contamination level of 2% (median 1.2%, Figure 2e).

Comparable, low level high distance contamination is predicted for WGS samples of Wohlers

et al. and Pagani et al. Possibly, these contamination calls are caused by nuclear mitochondrial

DNA sequences (NUMTs) causing mapping artefacts, although a core genome coverage of 30x,

a typically large haplogroup quality (Figure 2c) and a large distance between major and minor

haplogroup (Figure 2d) indicate that this is rather not the case. High contamination levels that

indicate a mix-up during library preparation are restricted to a few samples from the Sudan

cohort. For the ancient Egyptian mummy data set, haplocheck predicts contaminations in

haplogroups related to H2a2a, which is very likely caused by a lack of sequencing coverage,

which causes the revised Reference Sequence (rCRS) bases to be identified that refer to

haplogroup H2a2a1. This is supported by the very low coverage for this cohort.

<u>Distribution of Western Asian and Northeast African mitochondrial haplogroup diversity</u>

The sequential accumulation of polymorphisms in the MT sequence over time across

generations has led to the emergence of geographically isolable variation profiles and MT

haplogroups. As a result, geographic regions have characteristic distributions of mitochondrial

haplogroups. The analysis of n=9,264 world-wide MT sequences (see Methods) yielded

mitochondrial haplogroup distributions for distinct geographical regions as seen in Figure 3a.

a)

b)

Figure 3: World-wide haplogroup distributions by geographic region. a) Macrohaplogroup level b) Macrohaplogroups have been assigned to hypothetical continents or regions of origin. We decided on the following assignment: Africa: L haplogroups; Asia: E, F, G, M, Y, Z; Middle East: N, R, U, I, W; Europe: H, J, K, T, V; America: A, C, D, X; Oceania: B, P, Q.

As described previously, multiple haplogroups are predominant in different continental region. Here, we decided on the following assignment: Africa: L haplogroups; Asia: E, F, G, M, Y, Z; Middle East: N, R, U, I, W; Europe: H, J, K, T, V; Americas: A, C, D, X; Oceania: B, P, Q. Of note, we are aware that haplogroups overlap between continents, e.g., especially in Europe and the Middle East. The grouping above therefore does not entail exclusivity with respect to

the listed haplogroups. Based on our assignment, we determined the relative proportion of haplogroups associated with different continents for each clustered group (see Figure 3b). Regions in Asia, North America, Northeast Africa, and the Middle East are particularly diverse. Historically, this can be attributed to events such as colonization and slave trade in the Americas as well as early human migration in the Middle East (34). In addition to its unique transcontinental location between Africa and Asia, Northeast Africa is adjacent to the center of modern human settlement in the world (14) and has been affected by complex events of prehistoric back migrations (35).

The relative frequency of mitochondrial macrohaplogroups of the 641 QC filtered North and East African samples are depicted in Figure 4a and the individual haplogroup assignment per sample is given in in Supplementary Table 11.

Figure 4: Histograms of macrohaplogroup frequencies. Blue dashed line boxes indicate predominantly North African and yellow dashed line boxes predominantly East African macrohaplogroups within this region. a) For all n=641 North, Northeast and East African

samples passing the lenient QC filter and including Genbank sequences. This includes also

Genbank sequences of studies addressing particular haplogroups (e.g. L3, M1 and U6) and

thus is not a geographically representative distribution. b) Tunisian haplogroup numbers out

of overall n=42. c) Egyptian haplogroup numbers out of overall n=297. d) Sudanese

haplogroup numbers out of overall n=113. e) Haplogroup numbers in ancient Egyptian

mummies. f) Haplogroup numbers in ancient Moroccan individuals.

Country-specific haplogroup frequencies for Tunisia, Egypt, Sudan, ancient Morocco and

ancient Egypt are shown in Figures 4b-f. Geographic neighborhoods are clearly reflected in

the haplogroup distributions and indicate a geographic gradient from North Africa via

Northeast Africa towards East Africa. This can be noted particularly for haplogroups H and U,

that are most prevalent in Tunisia, less abundant in Egypt, and rare in Sudan. Vice versa, L3

and L0 are prevalent in Sudan, and to a lesser extend in Egypt but rarely observed in Tunisia.

Interestingly, L2 is commonly observed in both Tunisia and Sudan, but rarer in Egypt. Overall,

most macrohaplogroups are observed in the Egyptian cohort (compiled from three data sets).

This finding might be explained by the larger sample sizes and compilation of various data sets

available for Egypt, compared to only one data set for Tunisia and Sudan, respectively. Yet,

more macrohaplogroups reach high fractions compared to Tunisia and Sudan, supporting the

hypothesis that Egypt is the region of highest mitochondrial genetic diversity within North-

and East Africa.

<u>Northeast African mitochondrial lineages cover nearly all major world-wide clades</u>

In order to elucidate the relative haplogroup abundance in North and East Africa relative to

the rest of the world, we computed a phylogenetic tree from the 641 sequences in

combination with the 1000G and HGDP reference data. The resulting tree in Figures 5a-b

demonstrates that maternal ancestry is shared with nearly all major, world-wide

mitochondrial clades and is generally not restricted to specific subclades, apart from the

haplogroup M, which is mainly represented by M1. This is a haplogroup known to be restricted

to North and East Africa that is absent from the 1000G and present in HGDP data only once

(see also Supplementary Figures 1 and 2 with added sample names and corresponding

haplogroups).

The world-wide macrohaplogroups not observed in North and East Africa are A, B, D, E, F, G,

P, Y and Z, which are predominantly North Asian, Northeast Asian, East Asian, Southeast Asian,

Native American or Oceanian.

Figure 5: Phylogenetic tree of 641 North and East African mitochondrial sequences together

with sequences from two worldwide reference data sets. The tree is colored according to

major macrohaplogroup assigned by haplocheck (left) and colored according to data source,

with North, Northeast and East African mitochondrial sequences compiled in this study in red

and reference data in gray (right). a) Reference data is 1000G b) Reference data is HGDP.

The predominantly European haplogroup H is more common in Tunisia and Egypt and less

common in Sudan. The same applies to the haplogroups I, J, K, N and U.

Haplogroup I represents a with respect to RSRS not very deep-rooting sister clade of N1a1b,

as described earlier (36). Haplogroup J is also observed in both ancient data sets. North African

sequences belong to J1 as well as J2 and are with respect to RSRS more deep-rooting than

1000G and HGDP sequences.

Haplogroup K represents a subgroup of U8, as expected (37). In the North and East African

data, only K1 is observed, whereas 1000G and HGDP also cover K2.

The African haplogroups L0, L1, L2 and L3 that are commonly found in Africa show a high

abundance in Egypt, Sudan and Tunisia with the exceptions of L0 in Tunisia and L1 in Sudan

being absent and lowly abundant respectively. L3 is the founder of haplogroup N, which has

given rise to all world-wide haplogroups outside of Africa, and as such it has been studied extensively and exclusively. Haplogroup L3 is observed in all data sets, but predominantly in Sudan with a share of 40%. This is in line with previous studies that suggest East Africa as the origin of L3 (29).

Besides African-wide common haplogroups we observe macrohaplogroups L4, L5 and L6. All these macrohaplogroups have previously been observed in East Africa, mainly. Due to low prevalence and due to being restricted to this region, they have not been studied explicitly.

All but one sample relating to North and East African M haplogroups belong to the M1 clade, which is known to be restricted to this region and is the only non-Asian M subclade. The data sets compiled here cover a large diversity of M1. Haplogroup M1 has neither been observed in the comparably small Tunisian nor in the ancient Moroccan data set, but was present in the ancient Egyptian mummy data.

Sequences belonging to the N sub-haplogroups build subclades that are located at different positions in the phylogenetic tree generated here. All of them are in the same phylogenetic branch, the same holds true for the haplogroup R. We only observe sub-haplogroups of R0a (n=17) that has been described and interpreted previously (38), and encounter haplogroup R among the filtered sequences of an ancient Egyptian mummy once.

Nearly all T sub-haplogroups are detected in North or East Africa, and their share is elevated in both Moroccan and Egyptian ancient data sets. The presence of haplogroup T, as well as its sister clade J, has previously been observed and interpreted as Near Eastern genetic impact on the region (21,32).

North and East African sequences of macrohaplogroup U cover all sub-haplogroups (U1 to U8) except the rare U9 subgroup. Haplogroup U6 has been studied in a targeted way (26,27,30) because it is, together with M1, the only haplogroup of a world-wide macrohaplogroup that is autochthonous in Africa, i.e., predominantly occurring there. Corresponding U6 sequences from (26) and (27) that we integrated in our data compilation as well as Egyptian and world-wide reference data support this notion.

Finally, haplogroup X is present in data from all North and East African regions, including data from three ancient individuals. The coalescence of X in the phylogenetic tree with 1000G and HGDP as reference, respectively, is not identical, involving N9b and R11 as sister clades, both of which are haplogroups that are poorly represented in the 1000G and HGDP reference data, resulting in a limited resolution of the phylogenetic relationship of these clades.

Clades occurring predominantly in North and East Africa are often poorly characterized

We next investigated North and East African haplogroup profiles within the geographic

context using selected 1000G and HGDP populations from neighboring regions, see Figure 6.

A corresponding histogram is shown in Supplementary Figure 3.

Figure 6: Haplogroup frequencies of selected 1000G and HGDP populations together with the

Sudanese and Egyptian haplogroups of this work (highlighted in box). The populations are

colored according to geographic region.

This shows that both Egyptian and Sudanese mitochondrial profiles are mediating between

the profiles of populations assessed by these public references, constituting a genetic

continuum between them. For example, 1000G covers African, European, South Asian, East

Asian and Admixed American populations, none of which represents mitochondrial

haplogroup occurrences we observe in Sudan or Egypt according to data generated and

compiled in this work (cf. Figure 6). Middle Eastern and North African populations of the HGDP

reference show a combination of predominantly European, African and Asian haplogroups

comparable to Sudan and Egypt. However, HGDP population sample sizes are low and thus

corresponding haplogroup distributions may not be representative. Further, some

haplogroups from the North and East African region are so far absent from both major global

references 1000G and HGDP. This applies especially to specific haplogroups that exclusively

or predominantly occur in North and/or East Africa, e.g. subclades of L0a1a1, L3, L4, L6, M1,

U6 and X1. Some of those, with a particular interest to world-wide human prehistoric

migration routes, have explicitly been addressed by studies and selectively been sequenced

before, e.g., L3, M1 and U6 and are included in our study. However, local clades possibly not

related to archaeogenetic hypothesis are more poorly characterized, e.g., L0a1a1 and X1.

Finally, there is little information on rare haplogroups such as L6. They differ a lot from all

other, well characterized haplogroups and as such may convey different mitochondrial

function and disease susceptibility. Thus, also and especially such rare haplogroups will need

better characterization in the context of mitochondrial genetics-based precision medicine.

<u>Variation suggesting novel haplogroups</u>

To obtain a dataset of exceptional high-quality we next selected (n=146) samples with high

sequencing coverage and without predicted contamination (provided in Suppl. Table 12). The

phylogenetic tree of these sequences with corresponding haplogroups is depicted in Figure 7.

Figure 7: Phylogenetic tree of the n=146 exceptional high-quality sequences used for detection of novel haplogroups. The circular tree depicts the topological information, while branch lengths are not preserved. Two alternative visualizations preserving branch lengths and a histogram of the macrohaplogroup distribution are shown in Supplementary Figure 4.

In the search for novel and previously undefined subclades, we investigated this set using the phantom mutation module of the online haplogrep2 tool. In this module, sequences are checked for recurrent variants that are not haplogroup-defining and that have a 'Soares score' of less than or equal to 2, i.e., having been detected no more than twice in a large collection of mitochondrial sequences (according to Table S3 of Soares et al. (39)). We checked for such variants occurring in samples of the same haplogroup to ensure that the variation is likely specific to haplogroup carriers and defines a novel, previously undefined subclade. A summary is provided in Table 1.

Haplogroup	Subhaplogroup variant	# HQ	# LQ	Soares score
HV1b+152	8640T;4590G	2	1	0;0
I	567C	3	0	0
I	567C;4772C	2	3	0;2
L0a1a1	8869G	2	1	1
L0a1a1	8017T	3	9	0
L2a1c	5186G;9116C	2	0	2;2
L2a1+143	7897A;398C;9389G	2	0	1;2;2
L2e1	13356C	2	0	0
L3d5a	15457A;14410A	2	1	0;1
L3e4a	470G; 9455G	2	2	0;1
L3e5	711C	2	2	0
L3f1a1	12441C;736T;328G; 15151G;13967T;14554G	3	3	0;0;0;1;2;2
M1a2	16182G;2858G; 15401G;1282A	2	0	1;1;1;1
R0a2c	5981C	2	1	0
V	9126C	2	1	0

Table 1: Haplogroups having potentially novel and previously undescribed sub-haplogroups. The table columns denote, in this order, the haplogroup, the recurrent novel variants, the number of samples after strict filtering (# HQ; out of n=146) and after lenient filtering (#LQ; out of n=641-146=495) and the Soares score (39), which represents the previous occurrence of the variant in world-wide data.

For ten putatively novel haplogroups, we find further support among samples with lower sequencing coverage. We find one, two and 3-6 novel recurrent variants for respectively seven, five and three haplogroups each. Thereby, 14 of 30 potential novel haplogroup variants have a Soares score of 0 (47%), 9 have a score of 1 (30%) and the remaining 7 have a score of 2 (23%). The haplogroups of which potential novel sub-haplogroups have been identified are either haplogroups that are largely restricted to North and/or East Africa, e.g., L2a1, M1, and different L3 sub-haplogrups (d, e and f), or are potential local clades diverging from basic macrohaplogroups, e.g., I and V.

<u>Variation suggesting revision of phylogenetic relationships</u>

For the 146 exceptional high-quality sequences we examined the phylogenetic tree according to the PhyloTree reference for potential new haplogroups using a function from the web-based haplogrep2 tool (see Supplementary Figure 5). For this, we considered all detected and haplogroup-defining expected, but missing, variants in the samples. We find two clades which are better represented, when revising or extending the phylogenetic tree provided by PhyloTree, L0a1a1 and L2a1, see Figure 8

Figure 8: Phylogenetic tree according to the PhyloTree reference for those samples that were assigned to subclades a) L0a1 and b) L2a1. We highlight with magenta, gray and turquoise boxes those variants that indicate that a different phylogenetic tree may better explain the samples from North and East Africa.

There are eight samples assigned to L0a1 sub-haplogroups, see Figure 8a. In most of them, haplogroup-defining variant 95C is not observed, which is already noted within the PhyloTree reference. Further, three samples show a back mutation 185. Such a back mutation is expected for haplogroup L0a2 according to PhyloTree, so the samples we sequenced may link clades L0a1 and L0a2. In one sample, an expected 200 mutation was not observed, showing

the possibility that 95C and 185 together with 310, 8017, 12738 and possibly 16519 are a

hitherto undescribed subclade of L01a1.

The second branch of interest is L2a1. Out of 19 samples to which sub-haplogroups of L2a1

have been assigned, most show extensive, recurrent deviations from the PhyloTree

phylogenetic tree, see Figure 8b. For example, resolving the haplogroup variants 16189 and

16192 within the PhyloTree reference seems not representative for North and East African

samples. Further, variants 3705C, 13752, 7897, 9389, 5186 and 9116, which are recurrent in

the L2a1 clade, are not part of PhyloTree, yet. However, if included, they would implicate

changes in phylogeny.

Conclusion

We investigated whether current global mitochondrial genetic data adequately represents the

genetic mitochondrial variation of the North and East African region. Genetic data from this

region is sparse until now, despite an apparent interest due to autochthonous ancestry

components being identified (19,40) and despite the importance of the region as origin or

transit of modern humans outside of Africa. Beside in-depth analysis of mitochondrial

sequencing data underlying Egyptian haplogroup distributions that we have reported

previously (19), we here generated novel sequencing data of mitochondrial genomes from

Sudan, another region with little genetic data available. We characterized North and East

African mitochondrial variation by combining this data with publicly available sequencing data

as well as with mitochondrial sequences from Genbank relating to 10 further studies. This

results, to the best of our knowledge, in the largest mitochondrial data collection from this

region to date.

With a stringent quality control via the novel tool haplocheck we related North and East

African phylogenetics to the world-wide context by restricting the analysis to mitochondrial

sequences unaffected by potential contamination or other shortcomings such as insufficient

sequencing coverage. An additional filter for a minimal 1000x sequencing coverage results in

exceptionally high-quality mitochondrial sequences, which makes us very confident that novel

mitochondrial variations, novel haplogroups and observed PhyloTree inconsistencies are

indeed caused by a hitherto insufficient genetic representation of this geographic region.

We find that North and East African mitochondrial genomes are phylogenetically diverse and
cover nearly all major clades worldwide. Likewise, specific subclades are prevalent in the
region, supporting that North and East Africa, and particularly Northeast Africa harbors
ancestry components typically attributed to Africa, Europe and Asia (19) , all of which are likely
present in the region since prehistoric times (41). Further, several North or East African clade
representatives highlighted by our work and not studied in-depth so far may still add further
information to worldwide maternal human genetic history, e.g., the haplogroups L4, L6, N, I
and X.
In conclusion, the initial North and East African mitochondrial reference dataset compiled as
part of our work calls for further extension in order to support genetics-based precision
medicine.
437
438
439   **Material and Methods**
440
Haplogroup data compilation for world-wide mitochondrial haplogroup distribution
Based on a comprehensive literature research, n=7,414 publicly available mtDNA sequences
with available haplogroup and geographic information in the form of the country of origin
were collected. Additionally, data from two reference datasets, the 1000 Genomes dataset
(16) (n=2,504) and the Human Genome Diversity Project (24) (n=828) were added, resulting
in a total number of n=10,746 entries. As quality assurance, for all database entries
haplogroups were reassigned using for mitochondrial sequences haplogrep2 (9) and for next
generation sequencing-based cohorts haplocheck (22). Subsequently, entries were filtered
such that only those remained that were representative for the population of a region.
Towards this, studies analyzing only specific haplogroups as well as data from archaeological
studies was removed. Eventually, 9,264 entries were retained for further analysis.
452
Objective geographic grouping for world-wide mitochondrial haplogroup distributions
For determining the relative amount of haplogroups per geographic region, we grouped the
database entries into clusters of minimal geographic distance and having a size of at least 100
entries. At the beginning, only countries with less than 100 entries were considered, i.e., those
that could not form a group on their own. Countries with an initial number less than 100 were

assigned to group 1, all other countries to group 2. In order to classify a country from group
1, an attempt was made to combine it in a first step with other countries or already connected
groups of countries in group 1. A continent-dependent maximum distance was set as a
termination criterion. For Europe, this was set to 1,000 km, for Africa to 1,750 km, for the
Middle East to 1,600 km, for Asia to 1,900 km, for the Americas to 2,600 km and for Oceania
to 1,600 km. This distance was determined as a function of the average area of a country on
the continent and the existing sample density. If after this first step, the sample number of a
country or a group was still less than 100, the countries with an initial sample number greater
than or equal to 100 were included as possible connection partners. In this second step, the
country or group to be classified was in any case merged with the country from group 1 or
group 2 with minimal distance.
469
Mitochondrial DNA sequencing
Genomic DNA samples were processed for library preparation, using the Human mtDNA
Genome   protocol   for   Illumina   Sequencing   Platform
(https://emea.support.illumina.com/content/dam/illumina-
support/documents/documentation/chemistry_documentation/samplepreps_legacy/huma
n-mtdna-genome-guide-15037958-01.pdf) as described previously (19,42) for Egyptian
samples   and   63   Sudanese   samples.   In   brief,   a   primer   set   [MTL-F1
(AAAGCACATACCAAGGCCAC)   and   MTL-R1   (TTGGCTCTCCTTGCAAAGTT);   MTL-F2
(TATCCGCCATCCCATACATT) and MTL-R2 (AATGTTGAGCCGTAGATGCC)] was used to amplify
the mtDNA by long-range PCR (LR-PCR). Library preparation of the LR-PCR products was
performed using a Nextera XT DNA Library Preparation Kit (Illumina Inc., CA, USA). For 96
Sudanese samples, we applied a slightly modified protocol to improve the evenness of the
coverage. More specifically, the LR-PCR was conducted using another primer set [hMT-1_F
(AAATCTTACCCCGCCTGTTT)   and   hMT-1_R   (AATTAGGCTGTGGGTGGTTG);   hMT-2_F
(GCCATACTAGTCTTTGCCGC) and hMT-2_R (GGCAGGTCAATTTCACTGGT)] and the library
preparation of the LR-PCR products of hMT1 and hMT2 primers was conducted using DNA
Prep Tagmentation (Illumina Inc.). This approach allows the fragment size of the tagmented
LR-PCR products to be evenly 300 bp, as the transposome reaction occurs every 300 bp on the
beads, in contrast to the Nextera XT DNA library preparation Kit, by which transposome
reaction occurs randomly, thus, results in various size of the tagmented LR-PCR products (i.e.,

larger fragments were removed by beads purification steps). The libraries prepared by both

methods were further purified, concentration quantified using Qubit Fluorometer (Thermo

Fisher GmbH, Dreieich, Germany) and the library size was determined by 2100 Bioanalyzer

Instrument (Agilent, Santa Clara, CA, USA). The final library was sequenced on the Illumina

MiSeq sequencing platform, using v2 chemistry (2 × 150 bp paired-end reads) (Illumina Inc.).

<u>Sudanese and Egyptian mitochondrial sequencing characteristics and QC</u>

Running FastQC on all Sudanese and Egyptian mitochondrial sequencing FASTQ files resulted

in no systematic, data quality compromising warnings or errors. Sudanese sequencing data

has an average length of 140 bases, Egyptian sequencing comprises samples sequenced with

150 and samples sequenced with 300 bp target read length, resulting in an average length of

205 bases across the cohort. Average bases mapped per sample are about 18 Mb for the

Sudanese cohort and about 30 Mb for the Egyptian cohort, amounting to an approximate

average coverage of more than 1000X. In the Sudanese cohort, five samples have less than

1000 reads and are considered dropouts. The Egyptian cohort does not contain samples with

less than 1000 reads.

<u>Variant calling, haplogroup assignment and contamination detection</u>

Raw reads from FASTQ files of Sudanese and Egyptian sequencing data were mapped to

mitochondrial sequence MT of GRCh38, which resembles the historic revised Cambridge

Reference Sequence (rCRS), a sequence of haplogroup H2a2a1. For samples with available

BAM files, we used these directly. Our analysis is largely based on haplocheck (version 1.1.2)

(22) and its internal accompanying tools haplogrep2 (9) and mutserv (version 1.3.4) as

implemented via the mtDNA-server (43). Haplocheck is a recent tool which allows to detect

sample contaminations from sequencing data by assessing if heteroplasmic variant calls can

be broken down into a mixture of two haplogroups, one denoted as "major", the other as

"minor". Variant caller mutserv was run with default settings, which amounts to reporting

heteroplasmic variants at a level larger than 1% (--level 0.01), calling deletions (--deletions),

base alignment quality turned off (--noBaq), considering a mapping quality of 20 (--mapQ 20)

and a base quality of 20 (--basQ).

<u>Phylogenetic analyses</u>

In order to obtain a FASTA format mitochondrial sequence that relates to the haplogrep2
output reported by haplocheck, we perform variant calling with mutserv using the default
parameters also used within haplocheck. Then we select from the VCF those variants for which
the major base is the non-reference base. Using bcftools consensus, we then convert the VCF
file into FASTA. Note that mutserv does not call insertions, only deletions which are denoted
with a gap symbol (within FASTA, it is "-"), thus ensuring that the sequence length is always
exactly the length of rCRS, i.e. 16,569 bases, and that sequences align correctly without re-
alignment. The same holds for the Genbank files processed by McInerny et al. (25). Thus, we
can simply combine FASTA files of cohorts by pasting them into one combined FASTA file. The
combined FASTA file was then loaded into Jalview (44) where a Neighbour Joining-based tree
was computed and exported into Newick format. This tree was then visualized with ggtree
(45) using various graphical settings.

Ethics statement
This study was approved by the Central Institution Review Board, Al-Neelain University,
Sudan, IRB Serial Number NU-IRB-17-7-7-91. All subjects gave written informed consent in
accordance with the Declaration of Helsinki.

Data availability
The public data compiled as well as the corresponding publications and repository IDs are
provided in Supplementary Table 1. Sudanese raw sequencing data generated as part of this
study is available from the European Genome Phenome Archive (EGA) under study ID
EGAD00001008315. The analysis workflow used to run haplocheck, perform variant calling
and generate and compile mitochondrial sequences is available at
https://github.com/iwohlers/2020_mt_analyses.

Acknowledgements
The authors thank all volunteers who donated blood for the study as well as everyone involved
in the collection of blood samples. We also thank Miriam Freitag for excellent technical
assistance. IW, HB and SI acknowledge funding by the Deutsche Forschungsgemeinschaft

(DFG, German Research Foundation) under Germany's Excellence Strategy—EXC 22167-
390884018. AF, IS, HB and IW acknowledge support through the high-performance computer
cluster (OMICS-Cluster) of the University of Lübeck.

Conflict of interest

560 The authors have declared no conflicting interests.

References

1. Wallace DC, Fan W, Procaccio V. Mitochondrial energetics and therapeutics. Annu Rev
Pathol. 2010;5:297–348.

2. Anderson S, Bankier AT, Barrell BG, de Bruijn MH, Coulson AR, Drouin J, et al. Sequence
and organization of the human mitochondrial genome. Nature. 1981 Apr
9;290(5806):457–65.

3. Chandel NS. Evolution of Mitochondria as Signaling Organelles. Cell Metab. 2015 Aug
4;22(2):204–6.

4. Wang W, Zhao F, Ma X, Perry G, Zhu X. Mitochondria dysfunction in the pathogenesis of
Alzheimer's disease: recent advances. Mol Neurodegener. 2020 May 29;15(1):30.

5. Hsu C-C, Tseng L-M, Lee H-C. Role of mitochondrial dysfunction in cancer progression.
Exp Biol Med (Maywood). 2016 Jun;241(12):1281–95.

6. Wallace DC. A mitochondrial paradigm of metabolic and degenerative diseases, aging,
and cancer: a dawn for evolutionary medicine. Annu Rev Genet. 2005;39:359–407.

7. van Oven M, Kayser M. Updated comprehensive phylogenetic tree of global human
mitochondrial DNA variation. Hum Mutat. 2009 Feb;30(2):E386-394.

8. van Oven M. PhyloTree Build 17: Growing the human mitochondrial DNA tree. Forensic
Science International: Genetics Supplement Series. 2015 Dec 1;5:e392–4.

9. Weissensteiner H, Pacher D, Kloss-Brandstätter A, Forer L, Specht G, Bandelt H-J, et al.
HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput
sequencing. Nucleic Acids Res. 2016 08;44(W1):W58-63.

10. Rishishwar L, Jordan IK. Implications of human evolution and admixture for
mitochondrial replacement therapy. BMC Genomics. 2017 Feb 8;18(1):140.

11. Lippold S, Xu H, Ko A, Li M, Renaud G, Butthof A, et al. Human paternal and maternal
demographic histories: insights from high-resolution Y chromosome and mtDNA
sequences. Investig Genet. 2014;5:13.

12. Rieux A, Eriksson A, Li M, Sobkowiak B, Weinert LA, Warmuth V, et al. Improved
calibration of the human mitochondrial clock using ancient genomes. Mol Biol Evol.
2014 Oct;31(10):2780–92.

13. Laricchia KM, Lake NJ, Watts NA, Shand M, Haessly A, Gauthier L, et al. Mitochondrial
DNA variation across 56,434 individuals in gnomAD. bioRxiv. 2021 Jan
1;2021.07.23.453510.

14. Bergström A, Stringer C, Hajdinjak M, Scerri EML, Skoglund P. Origins of modern human
ancestry. Nature. 2021 Feb;590(7845):229–37.

15. Behar DM, van Oven M, Rosset S, Metspalu M, Loogväli E-L, Silva NM, et al. A
"Copernican" reassessment of the human mitochondrial DNA tree from its root. Am J
Hum Genet. 2012 Apr 6;90(4):675–84.

16. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang
HM, et al. A global reference for human genetic variation. Nature. 2015 Oct
1;526(7571):68–74.

17. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational
constraint spectrum quantified from variation in 141,456 humans. Nature.
2020;581(7809):434–43.

18. Taliun D, Harris DN, Kessler MD, Carlson J, Szpiech ZA, Torres R, et al. Sequencing of
53,831 diverse genomes from the NHLBI TOPMed Program. Nature. 2021
Feb;590(7845):290–9.

19. Wohlers I, Künstner A, Munz M, Olbrich M, Fähnrich A, Calonga-Solís V, et al. An
integrated personal and population-based Egyptian genome reference. Nat Commun.
2020 Sep 18;11(1):4719.

20. Pagani L, Schiffels S, Gurdasani D, Danecek P, Scally A, Chen Y, et al. Tracing the route of
modern humans out of Africa by using 225 human genome sequences from Ethiopians
and Egyptians. Am J Hum Genet. 2015 Jun 4;96(6):986–91.

21. Schuenemann VJ, Peltzer A, Welte B, van Pelt WP, Molak M, Wang C-C, et al. Ancient
Egyptian mummy genomes suggest an increase of Sub-Saharan African ancestry in
post-Roman periods. Nat Commun. 2017 May 30;8:15694.

22. Weissensteiner H, Forer L, Fendt L, Kheirkhah A, Salas A, Kronenberg F, et al.

Contamination detection in sequencing studies using the mitochondrial phylogeny.

Genome Res. 2021 Jan 15;

23. Brandstätter A, Sänger T, Lutz-Bonengel S, Parson W, Béraud-Colomb E, Wen B, et al.

Phantom mutation hotspots in human mitochondrial DNA. Electrophoresis. 2005

Sep;26(18):3414–29.

24. Bergström A, McCarthy SA, Hui R, Almarri MA, Ayub Q, Danecek P, et al. Insights into

human genetic variation and population history from 929 diverse genomes. Science.

2020 20;367(6484).

25. McInerney TW, Fulton-Howard B, Patterson C, Paliwal D, Jermiin LS, Patel HR, et al. A

globally diverse reference alignment and panel for imputation of mitochondrial DNA

variants. BMC Bioinformatics. 2021 Sep 1;22(1):417.

26. Maca-Meyer N, González AM, Pestano J, Flores C, Larruga JM, Cabrera VM.

Mitochondrial DNA transit between West Asia and North Africa inferred from U6

phylogeography. BMC Genet. 2003 Oct 16;4:15.

27. Olivieri A, Achilli A, Pala M, Battaglia V, Fornarino S, Al-Zahery N, et al. The mtDNA legacy

of the Levantine early Upper Palaeolithic in Africa. Science. 2006 Dec

15;314(5806):1767–70.

28. Harich N, Costa MD, Fernandes V, Kandil M, Pereira JB, Silva NM, et al. The trans-

Saharan slave trade - clues from interpolation analyses and high-resolution

characterization of mitochondrial DNA lineages. BMC Evol Biol. 2010 May 10;10:138.

29. Soares P, Alshamali F, Pereira JB, Fernandes V, Silva NM, Afonso C, et al. The Expansion

of mtDNA Haplogroup L3 within and out of Africa. Mol Biol Evol. 2012 Mar;29(3):915–

27.

30. Pennarun E, Kivisild T, Metspalu E, Metspalu M, Reisberg T, Moisan J-P, et al. Divorcing

the Late Upper Palaeolithic demographic histories of mtDNA haplogroups M1 and U6 in

Africa. BMC Evol Biol. 2012 Dec 3;12:234.

31. Fregel R, Méndez FL, Bokbot Y, Martín-Socas D, Camalich-Massieu MD, Santana J, et al.

Ancient genomes from North Africa evidence prehistoric migrations to the Maghreb

from both the Levant and Europe. Proc Natl Acad Sci U S A. 2018 Jun 26;115(26):6774–

9.

32. Kujanová M, Pereira L, Fernandes V, Pereira JB, Cerný V. Near eastern neolithic genetic
input in a small oasis of the Egyptian Western Desert. Am J Phys Anthropol. 2009
Oct;140(2):336–46.

33. Costa MD, Cherni L, Fernandes V, Freitas F, Ammar El Gaaied AB, Pereira L. Data from
complete mtDNA sequencing of Tunisian centenarians: testing haplogroup association
and the "golden mean" to longevity. Mech Ageing Dev. 2009 Apr;130(4):222–6.

34. Rodriguez-Flores JL, Fakhro K, Agosto-Perez F, Ramstetter MD, Arbiza L, Vincent TL, et al.
Indigenous Arabs are descendants of the earliest split from ancient Eurasian
populations. Genome Res. 2016 Feb;26(2):151–62.

35. Cabrera VM, Marrero P, Abu-Amero KK, Larruga JM. Carriers of mitochondrial DNA
macrohaplogroup L3 basal lineages migrated back to Africa from Asia around 70,000
years ago. BMC Evol Biol. 2018 Jun 19;18(1):98.

36. Olivieri A, Pala M, Gandini F, Hooshiar Kashani B, Perego UA, Woodward SR, et al.
Mitogenomes from two uncommon haplogroups mark late glacial/postglacial
expansions from the near east and neolithic dispersals within Europe. PLoS One.
2013;8(7):e70492.

37. González AM, García O, Larruga JM, Cabrera VM. The mitochondrial lineage U8a reveals
a Paleolithic settlement in the Basque country. BMC Genomics. 2006 May 23;7:124.

38. Gandini F, Achilli A, Pala M, Bodner M, Brandini S, Huber G, et al. Mapping human
dispersals into the Horn of Africa from Arabian Ice Age refugia using mitogenomes. Sci
Rep. 2016 May 5;6:25472.

39. Soares P, Ermini L, Thomson N, Mormina M, Rito T, Röhl A, et al. Correcting for purifying
selection: an improved human mitochondrial molecular clock. Am J Hum Genet. 2009
Jun;84(6):740–59.

40. Hodgson JA, Mulligan CJ, Al-Meeri A, Raaum RL. Early back-to-Africa migration into the
Horn of Africa. PLoS Genet. 2014 Jun;10(6):e1004393.

41. Serra-Vidal G, Lucas-Sanchez M, Fadhlaoui-Zid K, Bekada A, Zalloua P, Comas D.
Heterogeneity in Palaeolithic Population Continuity and Neolithic Expansion in North
Africa. Curr Biol. 2019 Nov 18;29(22):3953-3959.e4.

42. Russlies J, Fähnrich A, Witte M, Yin J, Benoit S, Gläser R, et al. Polymorphisms in the
Mitochondrial Genome Are Associated With Bullous Pemphigoid in Germans. Front
Immunol. 2019;10:2200.

43. Weissensteiner H, Forer L, Fuchsberger C, Schöpf B, Kloss-Brandstätter A, Specht G, et al.

mtDNA-Server: next-generation sequencing data analysis of human mitochondrial DNA

in the cloud. Nucleic Acids Res. 2016 Jul 8;44(W1):W64-69.

44. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2--a

multiple sequence alignment editor and analysis workbench. Bioinformatics. 2009 May

1;25(9):1189–91.

45. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc Bioinformatics.

2020 Mar;69(1):e96.