Knockout of liver fluke granulin, *Ov-grn-1*, impedes malignant

transformation during chronic infection with *Opisthorchis viverrini*

Short title: Knockout of liver fluke growth factor impedes bile duct cancer

Sujittra Chaiyadet[1, 2, 3] Sirikachorn Tangkawattana[3,4], Michael J. Smout[3,5,], Wannaporn

Ittiprasert[3,6], Victoria H. Mann[6], Raksawan Deenonpoe[7], Patpicha Arunsan[1,6], Alex

Loukas[5,8], Paul J. Brindley[6, 8], Thewarach Laha[1,8]

10
11 [1] Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002
12 Thailand
13
14 [2] Tropical Medicine Graduate Program, Academic Affairs, Faculty of Medicine, Khon Kaen
15 University, Khon Kaen, 40002 Thailand
16
17 [3]equal contribution
18
19 [4] Faculty of Veterinary Medicine, Khon Kaen University, Khon Kaen, Thailand, and
20 WHO Collaborating Center for Research and Control of Opisthorchiasis, Tropical Disease
21 Research Center, Khon Kaen University, Khon Kaen 40002, Thailand
22
23 [5] Centre for Molecular Therapeutics, Australian Institute of Tropical Health and Medicine, James
24 Cook University, Cairns, Queensland, 4878, Australia
25
26 [6] Department of Microbiology, Immunology and Tropical Medicine, and Research Center for
27 Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington
28 University, Washington, D.C. 20037, USA
29
30 [7]Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002
31 Thailand
32
33 [8]Correspondence: Thewarach Laha, email, thewa_la@kku.ac.th; Alex Loukas, email,
34 alex.loukas@jcu.edu.au; Paul Brindley, email, pbrindley@gwu.edu, Twitter, @PaulJBrindley
35

Abstract

Infection with the food-borne liver fluke *Opisthorchis viverrini* is the principal risk factor for cholangiocarcinoma (CCA) in the Mekong Basin countries of Thailand, Lao PDR, Vietnam, Myanmar and Cambodia. Using a novel model of CCA, involving infection with gene-edited liver flukes in the hamster during concurrent exposure to dietary nitrosamine, we explored the role of the fluke granulin-like growth factor *Ov*-GRN-1 in malignancy. We derived RNA-guided gene knockout flukes (*ΔOv-grn-1*) using CRISPR/Cas9/gRNA materials delivered by electroporation. Genome sequencing confirmed programmed Cas9-catalyzed mutations of the targeted genes, which was accompanied by rapid depletion of transcripts and the proteins they encode. Gene-edited parasites colonized the biliary tract of hamsters and developed into adult flukes, however less hepatobiliary tract disease manifested during chronic infection with *ΔOv-grn-1* worms in comparison to hamsters infected with control gene-edited and non-edited parasites. Specifically, immuno- and colorimetric-histochemical analysis of livers revealed markedly less periductal fibrosis surrounding the flukes and less fibrosis globally within the hepatobiliary tract during infection with *ΔOv-grn-1* genotype worms, minimal biliary epithelial cell proliferation, and significantly fewer mutations of *TP53* in biliary epithelial cells. Moreover, fewer hamsters developed high-grade CCA compared to controls. The clinically relevant, pathophysiological phenotype of the hepatobiliary tract confirmed a role for this secreted growth factor in malignancy and morbidity during opisthorchiasis.

Keywords

Infection-associated cancer; liver fluke; opisthorchiasis; cholangiocarcinoma; CRISPR/Cas;

knockout; granulin

Author summary

Infection with the human liver flukes, *Opisthorchis viverrini*, *O. felineus* and *Clonorchis sinensis*

remains a public health concern in regions where these parasites are endemic. *O. viverrini* is

endemic in the Mekong River drainage countries of including Thailand and the Lao People's

Democratic Republic. Infection follows the consumption of undercooked freshwater fish

harboring the parasite. Liver fluke infection, opisthorchiasis, is associated with diseases of the

liver and bile ducts including cancer of the biliary tract, cholangiocarcinoma, a cancer with a

poor prognosis. This report characterizes for the first time experimental infection with gene-

edited *O. viverrini* liver flukes during concurrent exposure to nitrosamine in a rodent model of

liver fluke infection-associated cancer. Cancer development was slowed in hamsters infected

with the parasite following CRISPR-based knock-out mutation and loss of a parasite gene known

to stimulate growth of cells lining the bile ducts. These findings describe a new model for

investigation of risk factors for infection-associated cholangiocarcinoma and to assess efficacy of

anti-infection/anti-cancer vaccines.

Introduction

Liver fluke infection caused by species of *Opisthorchis* and *Clonorchis* remains a major public health problem in East Asia and Eastern Europe. Infection with *Opisthorchis viverrini* is endemic in Thailand and Laos, where ~10 million people are infected with the parasite. Opisthorchiasis is associated with hepatobiliary diseases including cholangiocarcinoma (CCA), bile duct cancer (1, 2). Northeast Thailand reports the world's highest incidence of CCA, > 80 per 100,000 in some provinces. Indeed, the International Agency for Research on Cancer of the World Health Organization classifies infection with *O. viverrini* as a Group 1 carcinogen, i.e. definitely carcinogenic in humans (1, 3, 4).

Which features or consequences of parasitism by the liver fluke definitely initiate malignant transformation to CCA have yet to be ascertained notwithstanding that opisthorchiasis is the principal risk factor for CCA in regions where this neglected tropical disease remains endemic (1, 4-6). Some factors can be expected to more important than others and the impact of these factors should be quantifiable. Different worm burdens play a role, based on rodent models of liver fluke infection associated CCA (7), as does concurrent exposure to nitrosamines in fermented foods (8, 9) that are culturally important dietary staples in countries of the Lower Mekong River basin (5). Moreover, dose-dependent, synergistic effects of the liver fluke and nitroso-compounds have been documented (7, 10). To survive within the host, parasitic helminths actively release excretory/secretory (ES) proteins and other mediators with diverse effects and roles at the host-parasite interface (11, 12). This interaction is considered to manipulate host cellular homeostasis and, moreover, to underpin malignant transformation during chronic opisthorchiasis, but the molecular mechanisms by which these processes remain

inadequately understood (13). Focusing on the contribution of liver fluke ES to carcinogenesis,

we targeted the *O. viverrini* granulin-like growth factor, *Ov*-GRN-1, a prominent component of

the ES complement that we had determined previously induces phenotypic hallmarks of cancer

(14, 15). *Ov*-GRN-1 and other ES components including extracellular vesicles enter

cholangiocytes, the epithelial cells that line the biliary tract, and drive cellular signaling that

promotes carcinogenesis, including cellular proliferation and migration, angiogenesis and wound

healing (16). We have confirmed the role of *Ov*-GRN-1 in driving proliferation of bile duct

epithelial cells (cholangiocytes) by genetic manipulation of its expression in the liver fluke both

by RNA interference and RNA-guided gene knockout (15, 17, 18). Moreover, we have shown

that infection of hamsters with the gene edited, infectious stage of the live fluke was feasible and

that proliferation of biliary epithelia is markedly suppressed during infection with the *ΔOv-grn-1*

(*Ov-grn-1-/-*) flukes (18).

There is an established and instructive model of the pathogenesis of CCA in experimentally

infected hamsters, *Mesocricetus auratus* that is thought to replicate the epidemiology and

pathogenesis of chronic human opisthorchiasis (1). In this model, malignancy manifests within a

few months following infection with metacercariae of the parasite and concurrent exposure to

otherwise sub-carcinogenic levels of dietary nitrosamine (7, 19, 20). In the hamster, chronic

opisthorchiasis provokes periductal fibrosis, which, coincident with exposure to the nitric oxide

carcinogen facilitates cholangiocyte proliferation, epithelial hyperplasia, DNA damage, and

allied biliary tract lesions (21), which can culminate in CCA (22). Using this model, here we

investigated the outcome of infection of hamsters with gene-edited *O. viverrini* liver flukes in

relation to fluke-induced periductal fibrosis and malignant transformation. For this investigation,

hamsters were infected with juvenile flukes that had been genetically modified using CRISPR

before infection. Specifically, following up on our earlier study (18) which focused on knockout

of the *Ov-grn-1* gene, here we included as comparators juvenile flukes gene-edited for a second

virulence factor, tetraspanin (*Ov-tsp-2*) (23), which also networks at the host-parasite interface

(24, 25) and, as controls, flukes subjected to CRISPR transfection by an irrelevant (non-

targeting) guide RNA (26).

Immuno- and colorimetric-histochemical analysis of thin sections of liver revealed markedly less

fibrosis during infection with *ΔOv-grn-1* worms, reduced proliferation of cholangiocytes,

substantially less expression of mutant forms of the p53 tumor suppressor protein and, overall,

diminished malignancy of the liver. The clinically relevant, pathophysiological phenotype

confirmed a role for *Ov*-GRN-1 in morbidity and malignancy during opisthorchiasis. Moreover,

these findings underscored the utility and tractability of CRISPR-based genome editing for

addressing gene function, essentiality, and pathogenesis in parasitic helminths generally.

Results

Hamster model of malignant transformation during infection with gene-edited

Opisthorchis viverrini

To investigate the effect of programmed gene knockout (KO) in *O. viverrini*, two experiments

were undertaken in which hamsters were infected with newly excysted juveniles (NEJs) of *O.*

viverrini that had been subjected to programmed KO . The CRISPR/Cas systems were delivered

by electroporation of plasmids encoding guide RNAs specific for *Ov-grn-1*, *Ov-tsp-2* and an

irrelevant (control) guide RNA. All groups received plasmids encoding the Cas9 nuclease from

Streptococcus pyogenes. Figure 1A illustrates the experimental approach and timelines, the

findings from which we present below. Initially, juvenile flukes were subjected to transfection

after which reduction in transcription of the targeted genes, *Ov-grn-1* and *Ov-tsp-2*, was verified.

Subsequently, following successful KO of transcription *in vitro*, additional juvenile *O. viverrini*

were transfected before infection of hamsters. The goal of Experiment 1 was to assess impact of

KO on the worm burden. At necropsy14 weeks after infection, the entire liver was examined,

the worms were recovered and counted, and transcriptional changes were investigated in the

worms. With Experiment 2, the primary goal was establishment and assessment of disease

burden including malignant transformation during infection. Given that livers from the hamsters

in Experiment 2 were fixed at necropsy, worm burdens could not be established directly because

recovery of the flukes is a process that damaged the liver and biliary tract and, accordingly, was

incompatible with histological examination of infection-associated disease. Consequently, only a

small number flukes were available from Experiment 2 and these that were available were

collected incidentally during preparation of the liver lobes for fixation. Nonetheless, this sample

of the flukes from each of the three groups was sufficient to assess the performance and level of

programmed KO although total worm burden was unavailable. Nonetheless, the findings from

Experiment 2 provided the first description of gene-edited flukes in chronically-infected

hamsters, i.e., with duration of infection beyond eight weeks and, additionally, the first time

programmed mutation of the liver fluke genome has been combined with exposure to dietary

nitrosamine in the hamster-liver fluke model of human CCA.

173
174
Figure 1. Schematic overview of the experimental design. Hamsters were infected with juvenile
Opisthorchis viverrini worms that had been subjected to gene knockout. Two experiments were
conducted. (A). The goal of experiment 1 was to assess the impact of CRISPR/Cas9 editing on
fluke survival. The goal of experiment 2 was to assess the influence of gene knockout on
pathogenesis and malignant transformation. Groups of hamsters were infected with flukes
transfected with CRISPR/Cas9 plasmids targeting either Ov-grn-1 (red, ΔOv-grn-1), Ov-tsp-2
(green, ΔOv-tsp-2), or irrelevant guide RNA (blue, control) and exposed to dimethyl nitrosamine
(DMN) in the drinking water. At timepoints indicated for each experiment, fecal egg numbers
were measured as eggs per gram of feces (EPG). Experiment 1: From the livers of euthanized
hamsters, all the flukes were recovered and transcript levels assessed. Experiment 2: Liver lobes
were sectioned and stained for histochemical analysis. Flukes incidentally released from bile
ducts were collected and assessed for gene knockout. Before infection, transcript levels of Ov-
grn-1 (B) and Ov-tsp-2 (C) were assessed in juvenile flukes. Transcript levels established by
qPCR were plotted relative to average control transcript levels from 2-4 biological replicates;
average shown with colored bar and with 95% confidence interval error bars. Population
statistics were generated from resampling 1000 times with replacement bootstrap analysis of
untransformed delta-delta Ct values (derived from Figure S1). (Elements of the figure were
created with BioRender.com with copyright and licensing permission.)

193
194
Changes in transcription of the targeted genes induced by programmed mutations were

monitored by RT-qPCR in newly excysted juveniles (NEJs) (Figure S1). Control (irrelevant

gRNA) group flukes was used to normalize expression level from 100% (no change) to 0%

(complete ablation of expression) (Figure 1B, C). For the *ΔOv-grn-1* group, *Ov-tsp-2* was used

as an off-target gene control, and *Ov-grn-1* was used as an off-target control for the *ΔOv-tsp-2*

group. Relative to the control group, *Ov-grn-1* transcript levels were substantially reduced in the

ΔOv-grn-1 flukes (bootstrapped average, 95% confidence interval [CI]: 62.4%, 61.8-63.1%)

whereas the transcription of *Ov-tsp-2* increased marginally (< 20% change) in the *ΔOv-grn-1*

flukes (117.8%, 95% CI: 110.2-127.0%) (Figure 1B). Transcription of *Ov-tsp-2* was substantially

reduced in the *ΔOv-tsp-2* flukes (37.9%, 95% CI: 36.7-39.1%). Levels of *Ov-grn-1* transcripts

(102.1%, 95% CI: 101.8-102.5%) were unchanged (Figure 1C). This outcome revealed gene-

specific, on-target knockout at the *Ov-grn-1* and *Ov-tsp-2* loci. Gene expression analyses were

restricted to the two targeted genes and hence we cannot conclude that off-target mutations did

not occur.

Differential outcomes for adult flukes following knockout of *Ov-grn-1* and *Ov-tsp-2*

In Experiment 1, feces were sampled at both weeks 10 and 12 after infection (Figure 1A).

Significant differences in fecal egg counts (EPG) were not apparent among the three groups

(Figure 2A). Hamsters were euthanized and necropsied at week 14 to investigate the numbers of

adult *O. viverrini*. There were 60.0±3.46 (mean ± SEM), 36.7±3.48, and 21.3±2.96 worms in the

control, *ΔOv-grn-1*, and *ΔOv-tsp*-2 groups, respectively, reflecting reductions of 38.9% for *Ov-*

grn-1 and 64.5% for *Ov-tsp*-2 (Figure 2B). A trend was apparent toward increased egg burdens

in hamsters with higher worm burdens but the correlation not statistically significant (Figure S2).

Of the worms recovered in Experiment 1, gene transcript levels relative to the control flukes

were assessed for 10-13 flukes from each of the three hamsters in each group. Transcript levels

(Figure S3) of both genes expressed by the control parasites clustered around 100% of wild-type

fluke expression levels with bootstrap averages of 114.1% (95% CI: 104.2-126.5%) and 103.2%

(95% CI: 98.4-107.5%) (Figure 2C, D). Transcript levels for *Ov-grn-1* ranged broadly in the

ΔOv-grn-1 flukes from no change (~100%) to complete ablation (~0%) and, overall, were

substantially reduced at 11.8% (95% CI: 4.5-28.8%) of levels seen in wild-type flukes (Figure

2C). This phenotypic range, no change to ablation of transcripts in *Ov-grn-1,* was similar to our

previous findings . In contrast to the juvenile flukes, where there was substantial knockdown of

Ov-tsp-2 (Figure 1C), transcript levels were not markedly reduced in adult *ΔOv-tsp*-2 flukes

(82.5%, 95% CI: 73.6-91.5%) compared to the controls (103.2%, 95% CI: 98.4-103.2%) (Figure

2D). Indeed, most flukes in the *ΔOv-tsp*-2 group were unchanged with only two individual

flukes exhibiting > 50% reduction in *Ov-tsp*-2 expression. We posit that this marked difference,

in comparison to transcript levels for *Ov-tsp*-2 in juvenile flukes, i.e. reduced by 62.1%, in

addition to the 65% reduction of worms recovered at necropsy, *Ov-tsp-2* KO led to a lethal

phenotype and that flukes of this genotype failed to survive *in vivo*. This contrasted with *Ov-grn-*

1 KO where the majority of flukes of the *ΔOv-grn-1* genotype survived even though transcription

of *Ov-grn-1* was not detected in four of 36 flukes, and 10 of 36 exhibited *Ov-grn-1* transcript

levels < 5% of levels expressed by the control worms.

Figure 2. Liver fluke burden and levels of gene transcription. *Fecundity, worm numbers, and*
gene expression levels were determined at 10-14 weeks (Experiment 1) after infection of the
hamsters with 100 gene edited juveniles, and from three hamsters per group. Number of eggs
per gram of feces (EPG) from each hamster at weeks 10 and 12 (A) and worm numbers at week
14 (B) showing mean (horizonal black line) and SEM bars. Each treatment group was
compared to control group with 2-way ANOVA with Holm-Sidak multiple comparison: ns = not
*significant; **, P ≤ 0.01; ***, P ≤ 0.001, and ΔOv-grn-1 against ΔOv-tsp-2: #, P ≤ 0.05. Gene*
transcript levels of Ov-grn-1 (C) and Ov-tsp-2 (D) were determined by qPCR for 10 to 13 flukes
from each animal (30-39 total per group) and plotted with each data point representing the
transcript level of an individual fluke relative to wild-type flukes. Resample with replacement
bootstrap analysis (B=1000) of ddCT scores (Figure S3) used to generate population average,
as denoted by the thick colored line and 95% confidence interval error bars.

Parasitological impact of gene knockout

Quantitative assessment by histopathological and histochemical analysis of hepatobiliary disease

at 24 weeks of infection was a primary goal for Experiment 2. Accordingly, determining the

number of surviving liver flukes was not feasible because the liver and its intrahepatic biliary

tract, occupied by the parasites, was necessarily fixed in formalin at necropsy for downstream

histological processing and analysis. Nonetheless, we collected a sample of the resident *O.*

viverrini flukes before formalin fixation of the liver for molecular screening to assess the

efficacy of programmed gene knockout. As an accepted correlate of the number of worms

parasitizing each hamster (27), fecal egg counts (as eggs per gram of feces, EPG) were

determined at the time of necropsy. EPG values ranged broadly among the groups (Figure 3A),

with feces of the control fluke infected hamsters displaying the highest EPG values, median of

11,062 EPG (95% CI: 4,814-26,536), the *ΔOv-grn-1* group, 5,267 EPG (95% CI: 2,695-8535),

and the *ΔOv-tsp-2 group,* 4,530 EPG (95% CI: 518-10,227) in the same rank order in numbers of

worms recovered from these groups in Experiment 1 (Figure 2B). All the groups showed

substantial variation; the average EPG of *ΔOv-grn-1* hamsters was not significantly different to

controls whereas the *ΔOv-tsp-2* group had a significantly lower median EPG than the control

group ($P \leq 0.05$).

Adult worms recovered from Experiment 2 hamster livers examined at 24 weeks after infection

were evaluated for targeted gene transcripts by qPCR. *Ov-grn-1* transcript levels (Figure S4) of

adult *O. viverrini* from *ΔOv-grn-1* infected hamsters were substantially decreased down to a

bootstrapped average of 10.6% relative to wild-type flukes (95% CI: 3.1-30.9%) compared to the

control group with 91.0% (95% CI: 84.1-97.5%) (Figure 3B). In contrast, numbers of flukes in

the *ΔOv-tsp-2* treatment group were similar, at 101.1% (95% CI: 91.1-111.3%) of wild-type

flukes 98.5% (95% CI: 82.3-114.6%) of the controls (Figure 3C). With Experiments 1 and 2

showing the surviving *Ov-tsp-2* group worms expressing *Ov-tsp-2* at levels comparable to the

control flukes, we posit that substantial *Ov-tsp-2* gene edits were lethal, and worms that survived

to maturity likely had not undergone gene editing and/or few of the cells in the worms had been

edited. Although we retained the *ΔOv-tsp-2* group for comparison of pathogenesis, the genotype

of the flukes from this group was not investigated further.

289
290

Figure 3. ***Cholangiocarcinoma model, fecundity, gene transcript and mutation rates.*** *Eggs per*
gram of feces (EPG) were assessed at week 23 prior to euthanasia at week 24 (Experiment 2).
Panel A, EPG values of the three groups of hamsters. Violin plot denotes each hamster's EPG
with "x" symbols. Solid colored lines indicate the median values and dashed black lines
indicate the quartiles. Kruskal Wallis with Dunn's multiple comparison correction was used to
compare EPG levels against control group: ***,** *P ≤ 0.05; ns, not significant. At necropsy, 12-20*
*flukes were sampled from each group, transcript levels determined for Ov-grn-1 **(B)** and Ov-tsp-*
*2 **(C)**, and plotted with each data point representing the transcript level of individual flukes*
relative to wild-type flukes. Resampling with replacement bootstrap analysis (B=1000) of ddCT
scores (derived from Figure S4) used to generate population average denoted by thick colored
line and 95% confidence interval error bars.

Synopsis of outcomes of CRISPR/Cas9 gene editing of the liver flukes

To characterize mutations from the programmed knockouts, a region of 173 bp flanking the

programmed cleavage site in *Ov-grn-1* was scrutinized by analyzing aligned read-pairs from

targeted amplicon NGS and analysis of the aligned read-pairs with CRISPResso2 (28).

Substitution patterns as determined by the CRISPR-sub tool (29) in the KO groups were not

significantly different from the cognate alleles in the controls (Figure S5). Also, we scanned

insertions and deletions (indels) and, in turn, the potential impact of indels on the open reading

frame. Figure 4A and Table S1 present the indel percentages of juvenile and adult flukes. The

ΔOv-grn-1 pooled NEJs showed 3.26% indel levels (2,723 of 80,571 aligned read-pairs),

significantly more than the control group (18 of 51,402 aligned read-pairs, 0.035%) ($P \leq 0.05$).

The juvenile and adult flukes in the control group showed similar indel % levels, with 0.045% in

the adults (41 of 91,783 aligned read-pairs). Individual *ΔOv-grn-1* adult *O. viverrini* flukes

displayed a broad range of editing efficiency in terms of indel profiles. These ranged from an

apparent absence of programmed mutation (no indels) to near complete KO (91% indels), with a

median of 3.1% indels (M*ΔOv-grn-1*), which was significantly higher than in the control group

flukes ($P \leq 0.01$). As noted for levels of transcription, however, there were apparently distinct

groupings consisting of six low mutation status flukes (termed L*ΔOv-grn-1*) and six highly

mutation flukes (H*ΔOv-grn-1*) observed. From a total of 711 megabase pairs sequenced with 1.7

million aligned read pairs, programmed deletions (0.5 million) were overwhelmingly more

common than insertions, with only seven insertions identified (Table S1). There was an inverse

correlation between efficiency of KO of *Ov-grn-1* (indel percentage) and with a two-tailed non-

parametric Spearman correlation co-efficient r_s = -0.74 (Figure 4B; $P \leq 0.01$). Only minimal

expression of *Ov-grn-1* was detected in the highly mutated, H*ΔOv-grn-1* flukes, < 11% level of

transcription of control liver flukes. However, the highly edited genotype/highly reduced

transcription phenotype contrasted with the wide range of transcription in the flukes with low or

moderate levels of editing, $_L\Delta Ov$-*grn-1* and $_M\Delta Ov$-*grn-1,* with a wide range of transcription from

6% to 94% of the levels of the control group worms.

Evaluation of indel positions

With respect to indel length and position, mutations that were detected in the amplicon NGS

reads were observed at positions ranging from 29 bp upstream to 54 bp downstream of the

double stranded break (DSB+ (ORF nucleotide [nt] position -10 to +74)). Most indels were

deletions of a single nucleotide, others were several nucleotides in length, and one of 62 nt was

observed. Deletions were noted along the length of the amplicon, with several higher frequency

sites indicated with bubbles of greater diameter in Figure 4C. The location of indels in the

genomic DNAs pooled from juvenile flukes generally conformed with the location of

programmed mutations detected in the genome of the adult stage liver flukes (Figure 4C, large

bubbles). Insertions were seen only infrequently but these associated around these high

frequency indel locations in juvenile and adult flukes. A cluster of mutations at nucleotide

position -1 to -10 bp, within the 5' untranslated region (UTR) of the ORF, was notable given that

48 of the 216 indels (22%) observed in adult flukes occurred in this region, and five specific sites

included indels detected in the genome of at least seven individual adult worms.

Figure 4. ***Gene mutation rates among liver flukes***. ***A***, *Programmed gene knock-out was highly efficient although not in all the flukes. The percentage of total indels (insertions/deletions) determined by next-generation sequencing was plotted on the vertical axis of juvenile and 24 wk adult flukes from Experiment 2. Control group juvenile (NEJ) and adult flukes were each from a single pooled sample while juvenile ΔOv-grn-1 were from pools of two biological replicates s, and ΔOv-grn-1 adults from 13 individual flukes. The highly edited flukes were denoted HΔOv-grn-1, flukes with low editing denoted LΔOv-grn-1, and the single fluke with median level editing denoted MΔOv-grn-1. One sample t-test for either juvenile or adult worms comparing ΔOv-grn-1 and control:* *******, *P ≤ 0.05;* ********, *P ≤ 0.01. The thick solid line is the median and black dashed lines represent the inter-quartiles. A broken Y-axis with a magnified lower portion highlights the near zero values.* ***B***. *Adult fluke indel mutation rate was inversely correlated with transcript level. The indel and transcript levels were plotted for each individual ΔOv-grn-1 fluke (red circles, combining data from Figures 3C and 4A). Two-tailed non-parametric correlation determined by Spearman co-efficient:* ********, P ≤ 0.01. For context, the control indel percentages were plotted against the transcript median (blue triangle) with interquartile range error bars.* ***C***. *ΔOv-grn-1 indel location and size. The NGS reads revealed distinct indel patterns in 12/13 adult flukes. Shown as a multivariate bubble plot, the amplicon base pair open reading frame (ORF position) was plotted against the average indel length. The diameter of the bubbles (1-11) reflected how many of 13 adult flukes recorded a matching indel. The programmed double stranded break between residues 19 and 20 was indicated on the X-axis by the term "cut". For clarity, deletions in adult worm genomes (blue) and insertions (red) have been nudged up/down on the y-axis ±0.1. The deletions in juvenile worms (yellow) are shown from one pooled sample. Insertions were not seen. Position -9 was highlighted with a vertical dotted line and the black horizontal square bracket (⌐) highlighted a cluster of mutations. The sequence around this cluster was shown below the x-axis and the initiator ATG codon indicated in red.* ***D***. *Mutation rate (indels) at each location: the graph plotted the nucleotide position against the percentage frequency of indels in individual flukes. A vertical dotted line highlighted a mutational hotspot at nucleotide -9 and the black horizontal bracket (⌐) marked a mutational cluster. Other indels of note were labeled with the nucleotide and position.*

349 Figure 4D shows the frequency within each sample at each base pair position. There were

350 common locations in the ORF at which both control and *ΔOv-grn-1* juvenile and adults all

351 exhibited a mutation, albeit at low frequency (<0.02%). The location of these indels was denoted

352 with a number/letter to signify the nucleotide position and mutated residue. The cluster of

353 mutations was situated within nucleotide positions -1 to -10 of the 5'UTR of *Ov-grn-1* in adult

354 and juvenile flukes. Similar alleles were not present in the control worms. Although the

355 mutation rate in the 5'-UTR was low overall (<0.09%), numerous mutant alleles were seen at -

9T and this single position comprised ~99% of the mutations for the $_{M/H}\Delta Ov$-grn-1 adult and

ΔOv-grn-1 juvenile flukes.

Programmed knockout of *Ov-grn-1* impeded malignant transformation

Malignant transformation was induced in hamsters by the end point of Experiment 2, at 24 weeks

after infection. In the version of the model (7) that we adapted, Syrian hamsters were infected

with gene-edited juvenile *O. viverrini* flukes during concurrent exposure to exogenous

nitrosamine (Figure 1A). At necropsy, prominent malignant and premalignant lesions of several

type were diagnosed frequently in all three treatment groups of hamsters. Figure 5 panels A and

B present gross anatomical appearance of livers from representative hamsters, where the CCA

was visible to the unaided eye, highlighting the severity of disease that manifested using this

model. Specifically, multiple CCA nodules were obvious on both diaphragmatic (Figure 5A) and

visceral surfaces (Figure 5B). When micrographs of thin sections of liver were examined,

precancerous lesions were evident including biliary dysplasia (a precancerous precursor of CCA

(30) in many of the bile ducts and was frequently accompanied by periductal fibrosis (Figure

5C). Figure 5 D-F presents representative photomicrographs from each of the treatment groups

that highlighted the high-grade, malignant transformation. Figure 5G and Table S2 summarize

the findings from the treatment groups. Ten of 12 (83.3%) hamsters in the control group were

diagnosed with CCA; high-grade CCA in eight of them and low-grade CCA in other two.

Dysplasia also was apparent in the remaining two of the 12 (one mild, one moderate) hamsters in

the control group. CCA emerged in seven of 13 hamsters (six with high-grade CCA) in the ΔOv-

tsp-2 group. Of the remaining hamsters in this group, two showed mild biliary tract dysplasia,

one showed biliary tract proliferation, two exhibited hepatobiliary inflammation, and a single

hamster was free of apparent lesions. CAA was diagnosed in nine of 13 in the *ΔOv-grn-1* group,

four of which showed high-grade CCA. Of the remaining hamsters, one showed dysplasia, two

showed proliferation, and one was free of apparent lesions. Several hamsters infected with *ΔOv-*

tsp-2 (4/13 hamsters) and *ΔOv-grn-1* flukes (3/13 hamsters) exhibited lesions less severe than

dysplasia, i.e. inflammation or proliferation, or were free of lesions.

Figure 5. Burden of disease in liver fluke infection associated cholangiocarcinoma. *After resection of the livers at necropsy, a piece of each lobe either fixed in formalin for downstream thin sectioning or was manually disrupted to release flukes, which in turn were examined for gene editing events (Figure 1, Experiment 2). Gross anatomical appearance and histopathological results during induction of cholangiocarcinoma (31) Multiple CCA nodules in the hamster liver presented on both diaphragmatic **(A)** and visceral surfaces **(B)**. Micrographs of H&E-stained thin sections of liver highlighting foci of moderate dysplasia **(C)**. This image shows bile ducts (blue #) encircled by dysplastic biliary epithelium (yellow arrow) surrounded by fibrosis (fb) with hepatocytes (h) to the left. H&E stained images of CCA from each of the groups of hamsters group: control **(D)**, ΔOv-grn-1 **(E)**, and ΔOv-tsp-2 **(F)**. Inflammation marked with green asterisk (*), cholangiocarcinoma labeled as CCA; other labels as in panel C. **G**. Assessment and scoring of lesions was undertaken independently by*

*two co-authors (both veterinary pathologists) using anonymously labeled (blinded) micrographs. The severity of lesions increased from normal tissue (grey) to high grade CCA spanning multiple liver lobes (red). **H**. EPG from individual hamsters plotted against disease burden on a scale of zero (0, no lesion) to 6 (high CCA) scale. Data plots were slightly reformatted (nudged ± 0.1 on Y-axis) to enable display of overlapping points. Linear regression lines (which were not statistically significant) are shown in shaded color with 95% confidence intervals.*

Substantial differences were not evident among the treatment groups in the location or subtype of

CCA tumors (Table S2). Cholangiocarcinoma mass ranged from microscopic neoplasms with

multifocal distributions (12/26) to tumor masses apparent to the unaided eye (14/26) with

representative images in Figure 5A, B. With respect to histological classification, 21/26 CCAs

were the tubular type, and were seen in the three treatment groups. The papillary/cystic type was

seen in a single instance, in an *ΔOv-grn-1* group hamster. Additionally, four mucinous type

tumors were observed, one in the control and three in the *ΔOv-tsp-2* group hamsters. The right

lobe was the common tumor location (20/26 livers), with nine in the left lobe, and three in the

middle lobe. In some cases, tumors had developed in more than one lobe (5/26). Last, we

assessed pathology in relation to EPG levels (Figure 5H) and noted there was less disease in

ΔOv-tsp-2 hamsters where EPG was lower, especially in hamsters with EPG < 1,000. However,

significant correlation between pathogenesis and EPG was not apparent among the groups.

Reduced fibrosis during infection with knockout parasites

Hepatic fibrosis was detected in Picro-Sirius Red (PSR)-stained thin tissue sections and enabled

investigation and quantification of development of peribiliary fibrosis (Figure 6A). Fibrosis was

evaluated, firstly by Ishak staging, a semi-quantitative classification of the degree of fibrosis

spread across the liver parenchyma (32) and, secondly, on the fibrotic deposition as

demonstrated by staining with PSR localized around the liver flukes, i.e. the amount of collagen

deposition surrounding bile ducts occupied where individual liver flukes were situated at the

time of necropsy.

The Ishak scores corresponded to degrees of injurious fibrosis and the levels reflect expansion of

fibrosis into periportal regions and the degree of bridging between portal regions, culminating

full cirrhosis which is scored as Ishak level 6. The control group was severely affected, with the

majority (13/16) of the liver lobes assessed with an Ishak score of 4 (median 4, range 3-4)

(Figure 6B). Level 4 indicated that fibrosis had progressed extensively with marked portal-portal

and portal-central bridging. Hamsters in the *ΔOv-grn-1* group showed only a single lobe graded

at 4 and a majority of lobes (11/17) Ishak grade of 2 (median 2, range 2-4). Level 2 indicated

less pathology, with fibrosis in most portal areas with/without short fibrous septa that had not

bridged to other portal regions. Hamsters infected with the *ΔOv-tsp-2* liver flukes exhibited

similar levels of pathology among the group with Ishak scores ranging from 2 to 4 (median 3).

Level 3 was defined as fibrous expansion of most portal areas with occasional bridging among

them (32). Periductal fibrosis of the hepatobiliary tract in the *ΔOv-tsp-2* group hamsters was not

significantly different in Ishak score from the control group whereas the *ΔOv-grn-1* group

displayed significantly less fibrosis than the control ($P \leq 0.001$) or *ΔOv-tsp-2* ($P \leq 0.05$) groups

(Figure 6B).

Figure 6. ***Attenuated liver fibrosis during infection with ΔOv-grn-1 knockout parasites. A.***
*Representative images of hepatic fibrosis stained by Picro-Sirius Red with CRISPR-Cas9
edited O. viverrini. Fibrosis was denoted as pink/red thick bands around the bile ducts
(periductal fibrosis, fb) and expanded from each portal triad with fibrous septa. OV =
Opisthorchis viverrini, H = hepatocyte, BD = bile duct, BE = biliary epithelium. **B**. Global
liver fibrosis plotted as a violin plot (n = 14-17 liver lobes per group). Livers were scored for
fibrosis with an Ishak Stage Grading scale and plotted on a scale spanning from zero (no
fibrosis) to six (cirrhosis). **C.** Fibrosis proximal to flukes plotted as a violin plot (n = 46-54
per group). Automated ImageJ fibrosis evaluation of the percentage of collagen deposition in
images surrounding fluke-containing bile ducts. Panels B+C: median shown as thick colored
line and dashed black lines mark the inter-quartile ranges. Comparing groups with Kruskal-
Wallis test with Dunn's multiple comparisons against control: ns =not significant; *, P ≤
0.05; ***, P ≤ 0.001, and against ΔOv-tsp-2 group: #, P ≤ 0.05; ##, P ≤ 0.01.*

While the Ishak scores showed a significant difference between *ΔOv-grn-1* hamster livers and

both other groups, Experiment 1 data showed a higher worm burden in control animals than the

gene edited groups, and these extra worms may have contributed to greater fibrosis.

Surprisingly, there was no significant correlation between liver fibrosis and EPG for the control

or *ΔOv-grn-1* fluke-infected groups, but a significant inverse correlation ($P \leq 0.05$) was detected

for the *ΔOv-tsp-2* fluke-infected hamsters (Figure S6). To minimize the impact on fibrosis of

different worm burdens, we used automated image analysis to assess periductal fibrosis

immediately proximal to liver flukes (which live in the lumen of the bile ducts). An ImageJ

driven fibrosis quantification tool for scoring PSR-stained collagen was deployed for automated

analysis of the collagen deposition in periductal regions. Of the periductal regions proximal to

flukes in the control and the *ΔOv-tsp-2* fluke-infected hamsters, median values of 14.55% (95%

CI: 13.6-16.0) and 14.56% (95% CI: 13.5-16.0%) of fibrotic tissue were detected, whereas

significantly less (12.66%; 95% CI: 12.1-13.5%) bile duct tissue surrounding *ΔOv-grn-1* flukes

was fibrotic (Figure 6C, $P \leq 0.05$). While concerns over worm burden influencing pathology

remain, both the control and *ΔOv-tsp-2* group showed very similar localized collagen deposition

factors despite very different worm burdens, and suggests that focusing on pathology

immediately adjacent to detectable flukes is an accurate and meaningful way to compare

pathogenesis on a per worm basis.

***Ov-grn-1* KO flukes provoked less cell proliferation**

Here we explored the *in vivo* effects of programmed knockout of *Ov-grn-1* expression, as

opposed to our earlier reports which centered on proliferation of the biliary epithelium and/or

cultured cholangiocytes in response to *in vitro* exposure to recombinant *Ov*-GRN-1 (14-17, 33).

Proliferation of hamster biliary cells *in situ* was investigated using incorporation of the thymine

analogue into cellular DNA. Visualization was performed using immunohistochemistry and

evaluated quantitatively (Figure 7A). Concerning worm survival, and its corollary, the fitness

cost of the programmed mutation, we examined proliferation but only in the bile ducts where

flukes were situated. Median proliferation in bile duct tissue surrounding liver flukes in the

control (15.0%, 95% CI: 12.4-22.8%) and *ΔOv-tsp-2* (11.1%, 95% CI: 4.1-20.0%) groups

showed wide ranging values but not significantly different from each other (Figure 7B).

Whereas the biliary epithelia surrounding *ΔOv-grn-1* flukes showed substantial variation from 0-

30%, the majority of the readings were below the other groups with a median of only 3.1% (95%

CI: 1.6-7.6%) of cholangiocytes incorporating BrdU ($P \le 0.001$ vs control; $P \le 0.05$ vs *ΔOv-tsp-*

2, Figure 7B). By contrast, the *ΔOv-grn-1* group (3.1%) showed significantly less proliferation

than both the control group (4.8-fold reduction, $P \le 0.001$) and the *ΔOv-tsp-2* group (3.6-fold

reduction, $P \le 0.05$).

Mutant *TP53* less frequent during infection with *Ov-grn-1* KO flukes

Tumor protein p53 plays a well-recognized role in cholangiocarcinogenesis and is highly

expressed in fluke infection associated malignancy (1, 34). A brown nuclear staining pattern

presents only in neoplastic biliary cells (Figure 7C). Wide angle images of flukes in the biliary

tract showed mutant p53-positive and -negative cells in the epithelium. The profile of p53

positive cells differed markedly among the groups: control and *ΔOv-tsp-2* fluke-infected

hamsters showed similar levels, 61.1% (95% CI: 43.6-68.3%) and 67.7% (95% CI: 41.5-73.5) of

cholangiocytes stained for mutant p53, respectively (Figure 7D). By contrast, of the *ΔOv-grn-1*

fluke-infected hamster bile ducts, only 7.5% (95% CI: 2.3-27.5%) exhibited mutant p53-positive

cells, ~13% of the levels that manifested in the other groups ($P \leq 0.01$) (Figure 7D).

***Figure 7. Reduced proliferation and minimal mutant p53 expression in cholangiocytes in
hamsters infected with ΔOv-grn-1 genotype liver flukes.*** *Representative images of biliary
cells that incorporated BrdU from regions proximal to flukes in control, ΔOv-grn-1 and ΔOv-
tsp-2 groups **(A)**. The boxed region in the upper image is magnified in the lower panel. The
brown arrow highlights the positive BrdU-stained nuclei and the blue arrow highlights a bile
duct cell that did not incorporate BrdU. Violin plots (n = 27-42 per group) of BrdU index
measured from cholangiocytes adjacent to where a fluke was located **(B)**. Representative
micrograph of p53 immunohistochemical staining of biliary epithelium during infection with
gene edited flukes **(C)**. Anti-mutant p53 antibody stained the nuclei brown (brown arrows);
blue arrows indicate negative cells. Black dashed box in upper wide-angle image magnified in*

*the lower image to aid visualization. Violin plot (n = 29-39 per group) of mutant p53 positive cholangiocytes as a percentage **(D)**. When available, 500 to 800 cells were scored from sections of each of the left, middle, and right lobes of the liver marked by "X". Fewer cholangiocytes (300-500) were available for assessment in several samples, denoted byⓍ Panels A and C: OV = Opisthorchis viverrini, H = hepatocytes, BD = bile duct, BE = biliary epithelium. Panels B, D: non-parametric Kruskal-Wallis test with Dunn's multiple comparison correction compared against control: ns = not significant; ***, P ≤ 0.001, or against ΔOv-tsp-2: #, P ≤ 0.05; ###, P ≤ 0.001. Thick colored lines signify the median and the dashed black lines mark the inter-quartile range.*

Discussion

CCA accounts for ~15% of all primary liver cancers globally and its incidence is increasing (35).

Infection with *O. viverrini* is the principal risk factor for CCA in the Lower Mekong River Basin

countries including Thailand and Laos PDR, where CCA is the dominant form of liver cancer (1,

22, 36). In an earlier report, we exploited this link to explore the role of *Ov*-GRN-1 secreted by

the parasite in tumorigenesis using programmed gene knockout, and reported that the infection

was less severe even though gene-edited parasites colonized the biliary tract of hamsters and

developed into adult flukes (18). In this follow-up investigation, we report findings during

concurrent exposure to dietary nitrosamine and infection with the gene edited parasites, and that

KO of the granulin gene retards malignant transformation to CCA, including the emergence of

mutant p53, in a rodent model of human opisthorchiasis-associated CCA. These novel results

build upon and advance the findings from our original report (18) and, notably, confirmed the

role of liver fluke granulin in malignant transformation during chronic opisthorchiasis (15).

We utilized an established model of opisthorchiasis-associated CCA in hamsters that were

infected with the parasite during concurrent exposure to exogenous nitrosamine. CCA manifests

under these conditions, and this rodent model reflects the human situation where chronic

opisthorchiasis in the context of a diet that is rich in fermented fish (in turn, rich in nitrosamines)

culminates in a high incidence of CCA (20, 22, 37, 38). In hamsters, opisthorchiasis leads to

periductal fibrosis. Chronic periductal fibrosis combined with a nitric oxide carcinogen, such as

DMN, results in epithelial cholangiocyte proliferation, hyperplasia, dysplasia, and DNA damage,

eventually and reliably manifesting as malignant neoplasia of the biliary tract (20, 21, 39). By

contrast, conspicuously less proliferation of the biliary epithelium, reduced mutant p53

expression by cholangiocytes, and less periductal fibrosis accompanied infection here with *ΔOv-*

grn-1 genotype worms compared to controls. As noted, our approaches and findings represent a

functional genomics (forward genetics)-focused extension of the model pioneered by Thai

investigators more than 30 years ago (7).

Fitness cost of gene knockout can be assessed from programmed gene editing, an approach that

is employed for the unbiased identification of essential genes in other organisms and disease

settings (40, 41). The present findings confirmed the power of RNA-guided targeted mutation to

define essentiality and relevance of two parasite proteins in infection-associated morbidity and

malignancy. The *Ov-grn-1* gene does not appear to be essential for *in vivo* development and

survival, which has enabled investigation here on the role of this protein in driving cell

proliferation, pathology and ultimately contributing to CCA. Nonetheless, the reduced fecundity

of *ΔOv-grn-1* liver flukes likely reflected a fitness deficit as the result of the targeted KO. By

contrast, *Ov-tsp-2* appears to be essential to parasitism. The *ΔOv-tsp-2* genotype did not survive

in vivo, and sequencing of the indels across the relevant region of the genome confirmed that

most of the surviving flukes from hamsters had not undergone editing at the *Ov-tsp-2* gene locus.

These findings build upon earlier RNA interference-mediated silencing of *Ov-tsp-2* gene

expression and the resultant malformation of the tegument observed *in vitro* (42). Although

infection of hamsters with *Ov-tsp-2* dsRNA-treated parasites was not investigated in this earlier

report, the damage to the tegument following exposure to *Ov-tsp-2* dsRNA in worms cultured for

several days appeared to be so extensive and debilitating that worms damaged to that extent by

either CRISPR-based genome knockout or dsRNA likely did not establish or survive for long

periods *in vivo*.

In *O. viverrini*, *Ov*-GRN-1 and *Ov*-TSP-2 share key, though dissimilar functions at the parasite

interface with the mammalian host. *Ov*-GRN-1 induces proliferation of cholangiocytes whereas

Ov-TSP-2 is a key structural protein of the tegument of liver flukes (and indeed in schistosomes

(43, 44)) and of extracellular vesicles that are taken up by host cholangiocytes, among other roles

(15, 16, 24). Accordingly, *Ov-tsp-2* was included here as a comparator gene for *Ov-grn-1* KO. A

non-targeting guide RNA encoded here by pCas-*Ov*-scramble, also was included, to provide a

negative control for off-targeting by the Cas9 nuclease (26). Ultimate lethality of *Ov-tsp-2* KO

was borne out in our observation that *ΔOv-tsp-2* genotype flukes failed to survive to the adult

stage in the hamsters. Whereas this diminished the value of *ΔOv-tsp-2* worms as controls, the

findings highlighted the apparent essentiality of this tetraspanin to the intra-mammalian stages of

the liver fluke. This essentiality of tetraspanin contrasted with *Ov*-GRN-1, the absence of which

was not lethal to the parasite but which, fortunately, enabled inferences on its contribution to

malignant transformation and which reinforced earlier hypotheses on its role as a carcinogen

(15). These approaches and findings are novel in the field of functional genomics for helminth

parasites and hence, even if targeting *Ov-tsp-2*, in hindsight, was a misstep, the outcome

provided a programmed mutation-based demonstration of the essentiality and lethality of

mutations of these liver fluke genes, which represents vanguard progress in forward genetics for

helminth parasites. Indeed, for context concerning the pioneering significance of this advance,

establishing the essentiality of human genes is an active and fertile field in functional genomics

and gene therapy (45, 46).

Notwithstanding that the primary goal of this investigation was to characterize pathogenesis and

carcinogenesis associated with infection with *ΔOv-grn-1* flukes in this singular infection/nitroso-

compound hamster model, we investigated the mutation profile induced by the CRISPR-based

targeted genome editing. Intriguingly, numerous mutations at the targeted *Ov-grn-1* locus were

situated within the 5'UTR, rather than in the targeted exon, and most were detected at or

proximal to a single residue, referred to here as -9T. This was situated 28 bp 5' to the

programmed CRISPR/Cas9 double strand break at nucleotide positions 19 and 20 of the open

reading frame. This outcome was unforeseen given that mutations are usually expected at and

adjacent to the programmed double stranded break, although guide RNAs have individual, cell-

line dependent biases toward particular outcomes (47). The sequence of the 5'UTR of *Ov-grn-1*

does not exhibit identity to regulatory elements (UTR database, http://utrdb.ba.itb.cnr.it/), which

was unsurprising given that few helminth parasite UTR regulatory elements have been

characterized (48). Why this position was preferentially mutated is unclear although the marked

reduction of transcription of *Ov-grn-1* that accompanied this mutation profile may signal the

presence of a regulatory control element within the UTR.

Chronic inflammation and fibrosis are risk factors for liver cancer (49). The traditional lifestyle

of people living in *O. viverrini*-endemic areas, notably a diet enriched in nitrosamines as well as

routine alcohol consumption, in tandem with the assault on the biliary epithelium by the

attachment, feeding, movement, and secretions of the liver flukes that result in repeated cycles of

injury and repair, establishes a compelling and conducive setting for malignant transformation

(5, 50, 51). The secretion of liver fluke granulin into the bile duct and the ability of this growth

factor to drive relentless cell proliferation during infection and to (re)heal wounds inflicted by

the helminth plays a central role in this process (15). Whereas knockout mutation of *Ov-grn-1*

did not prevent development and survival of the liver fluke *in vivo*, infection with these *ΔOv-grn-*

1 flukes failed to lead to marked cell proliferation and fibrosis in the immediate vicinity of the

parasites, and consequently fewer hamsters developed high-grade CCA compared with hamsters

infected with control and *ΔOv-tsp-2* parasites. Indeed, more hamsters infected with *ΔOv-grn-1*

flukes were diagnosed in the categories termed either low-CCA and/or proliferation than in the

other two groups. Knockout mutation of *Ov-grn-1* clearly impeded malignant transformation

during chronic opisthorchiasis.

Infected hamsters exhibited elevated rates of *TP53* mutation, however the level was markedly

less during infection with *ΔOv-grn-1* flukes. The mutational signatures and related molecular

pathways characteristic of human CCAs have been reviewed in depth, and the signature profiles

differ between fluke-associated and non-fluke-associated CCAs (1, 34). Fluke-associated CCAs

exhibit substantially more somatic mutations than non-fluke related CCAs (34), likely the

consequence of opisthorchiasis-associated chronic inflammation. In conformity with the human

situation, reduced inflammation and fibrosis were seen during infection with the *ΔOv-grn-1*

flukes, further emphasizing the virulence of this growth factor in chronic opisthorchiasis.

Inactivating mutations of *TP53* are more prevalent in CCA with a fluke infection etiology, as are

mutations of *ARID1A, ARID2*, *BRCA1* and *BRCA2*, than in non-fluke related CCAs (34, 52-55).

In addition, hypermethylation has been noted for the promoter CpG islands of several other

aberrantly expressed genes (34).

Mosaicism of gene knock-out is a limitation of our somatic gene-editing approach. Obviating

mosaicism by access to transgenic worms following germline transgenesis would clearly be

preferable. However, access to transgenic lines of *O. viverrini,* while desirable seems unlikely in

the near future, especially considering the genetic complexity of this hermaphroditic

platyhelminth obligate parasite with a diploid genome and a multiple host developmental cycle

which cannot reliably be established in the laboratory (56). Nonetheless, somatic genome editing

is increasingly expedient in the clinic including for the treatment of hemoglobinopathies (57) and

the identification of targets for disease intervention in translational medicine (58). Given the role

of liver fluke granulin as a virulence factor, the cogent link between CCA and liver fluke

infection, and the dismal prospects following a diagnosis of CCA in resource poor settings (1,

59, 60), interventions that target this growth factor should be beneficial. Indeed, antibodies raised

against liver fluke granulin block its ability to drive proliferation of CCA cell lines (14) and,

hence, bootstrap support for a vaccination strategy targeting *Ov*-GRN-1 in the gastro-intestinal

tract, for example through induction of mucosal IgA and IgG responses (23, 61). Such an

intervention might contribute a productive component to a multivalent, orally administered, anti-

infection and anti-cancer vaccine (1, 23), which in turn would augment current tools for the

public health intervention and control of this neglected tropical disease and its cancer burden (62,

63).

Materials and methods

Ethics

The protocol for this research was approved by the Animal Ethics Committee of Khon Kaen

University, approval number ACUC-KKU-61/60, which adhered to the guidelines prescribed by

the National Research Council of Thailand for the Ethics of Animal Experimentation. All the

hamsters were maintained at the animal husbandry facility of the Faculty of Medicine, Khon

Kaen University, Khon Kaen.

Metacercariae, newly excysted juvenile and adult developmental stages of *O. viverrini*

Metacercariae (MC) of *O. viverrini* were obtained from naturally infected cyprinid fish

purchased from local food markets in the northeastern provinces of Thailand (64). MC were

isolated from fishes by using pepsin digestion as described previously (65). Briefly, whole fishes

were minced by electric blender and digested with 0.25% pepsin with 1.5% HCl in 0.85% NaCl

at 37 ºC for 120 min. The digested fishes were filtered sequentially through sieves of 1100, 350,

250, and 140 μm mesh apertures. The filtered, digested tissues were subjected to gravity

sedimentation through several changes of 0.85% NaCl until the supernatant was clear.

Sedimented MC were identified under a dissecting microscope as *O. viverrini*, and active (i.e.,

exhibiting larval movement within the cyst) MC were stored in 0.85% NaCl at 4ºC until used.

Newly excysted juveniles (NEJ) of *O. viverrini* were induced to escape from the metacercarial

cyst by incubation in 0.25% trypsin in PBS supplemented with 2× 200 U/ml penicillin, 200

µg/ml streptomycin (2× Pen/Strep) for 5 min at 37°C in 5% CO_2 in air. The juvenile flukes were

isolated free of discarded cyst walls by mechanical passage through a 22 G needle (18). We also

use the term NEJ for the juvenile flukes because NEJ is also widely used for juveniles of related

liver flukes (66, 67).

Plasmid constructs and transfection of *O. viverrini*

The CRISPR plasmid encoding a guide RNA (gRNA) complimentary to *Ov-grn-1* exon 1 termed

pCas-*Ov-grn-1* was constructed using the GeneArt CRISPR Nuclease Vector kit (Thermo Fisher

Scientific), as described . The programmed cleavage site at nucleotide position 1589–1608, 5'-

GATTCATCTACAAGTGTTGA with a CGG proto-spacer adjacent motif (PAM) which

determined the cleavage site located at three nucleotides upstream on *Ov-grn-1* was designed

using the online tools, http://crispr.mit.edu/ (68) and CHOPCHOP,

 http://chopchop.cbu.uib.no/ (69, 70) using the *Ov-grn-1* gene (6,287 bp, GenBank FJ436341.1)

as the reference. A second plasmid, termed pCas-*Ov-tsp-2* was constructed using the same

approach; pCas-*Ov-tsp-2* encodes a gRNA targeting exon 5 of the *Ov-tsp-2* gene (10,424 bp,

GenBank JQ678707.1) (71, 72). The guide RNAs encoded by *Ov-grn-1* and *Ov-tsp-2* exhibited

theoretically high, on-target efficiency and little or no off-target matches to the *O. viverrini*

genome. A third construct, termed pCas-*Ov*-scramble was also prepared and included as a

control to normalize analysis of gene expression and programmed gene knockout. The pCas-*Ov-*

scramble construct included as the gRNA, a transcript of 20 nt, 5'-

GCACTACCAGAGCTAACTCA which exhibits only minimal identity to the *O. viverrini*

genome and which lacks a PAM (73). A non-targeting guide RNA (26), encoded here by pCas-

Ov-scramble, provided a negative control for off-targeting by the Cas9 nuclease. A mammalian

U6 promoter drives transcription of the gRNAs in all three plasmids and the CMV promoter

drives expression of the *Streptococcus pyogenes* Cas9 nuclease, modified to include the

eukaryotic nuclear localization signals 1 and 2. To confirm the orientation and sequences of

gRNA in the plasmid vector, *Escherichia coli* competent cells (TOP10) were transformed with

the plasmids, plasmid DNAs were recovered from ampicillin resistant colonies using a kit

(NucleoBond Xtra Midi, Macherey-Nagel GmbH, Düren, Germany), and the nucleotide

sequences of each construct confirmed as correct by Sanger direct cycle sequencing using a U6-

specific sequencing primer.

Two hundred juvenile *O. viverrini* were dispensed into an electroporation cuvette, 4 mm gap

(Bio-Rad, Hercules, CA) containing 20 µg pCas-*Ov-grn-1,* pCas-*Ov-tsp-2* or pCas-*Ov*-scramble

in a total volume of 500 µl RPMI, and subjected to a single square wave pulse at 125 V for 20

ms (Gene Pulser Xcell, Bio-Rad). These juvenile flukes were maintained in culture in RPMI

supplemented with 1% glucose for 60 min after which they were used for infection of hamsters

by stomach gavage (below).

Infection of hamsters with gene edited *O. viverrini* juveniles

Wild-type (WT) flukes were collected and prepared for qPCR from infected hamsters 8 weeks

after infection as previously described . Figure 1 provides a timeline of the CCA model,

employed in Experiments 1 and 2. In Experiment 1, nine male hamsters (Syrian golden hamster,

Mesocricetus auratus) aged between 6-8 weeks were randomly divided into three experimental

groups (Figure 1). Each hamster was infected with 100 *O. viverrini* NEJs by gastric gavage.

These juvenile flukes had been transfected with pCas-*Ov-grn-1* plasmid, pCas-*Ov-tsp-2*, or the

control pCas-*Ov*-scramble, and assigned the following identifiers: delta(Δ)-*gene* name, *ΔOv-grn-*

1, ΔOv-tsp-2, or control, respectively. The infected hamsters were maintained under a standard

light cycle (12 hours dark/light) with access to water and food *ad libitum*. Two weeks following

infection, the drinking water provided to hamsters was supplemented with dimethylnitrosamine

(DMN) (synonym, *N*-nitrosodimethylamine) (Sigma-Aldrich, Inc., St. Louis, MO) at 12.5 ppm

until 10 weeks following infection (7, 19, 20). Feces from each hamster were collected for fecal

egg counts at weeks 10 and 12 after infection. The hamsters were euthanized at week 14 by

inhalation overdose of isoflurane, followed by removal of the liver. Liver flukes were recovered

from the liver, counted, and prepared for qPCR analysis of targeted genes.

In Experiment 2, 45 male hamsters, 6–8 weeks of age, were randomly divided into three groups

each with 15 hamsters and infected with gene edited NEJs and DMN added to drinking water (as

in Experiment 1). Hamster feces were collected at week 23 after infection for fecal egg counts.

At week 24, 40 mg/kg thymine analogue 5-bromo-2'-deoxyuridine (BrdU, Abcam, College

Science Park, UK) was introduced into the peritoneum at 30 min before euthanasia, for

incorporation into the nuclei for investigation of proliferation of the biliary epithelia (74).

Hamsters were euthanized, the liver resected from each hamster, the liver lobes separated, and

the left, middle and right lobes fixed in 10% formalin. We used a shorthand to label the lobes:

left (left dorsocaudal), middle (combined ventral and dorsal median lobes), and right (right

dorsocaudal). Liver flukes that were incidentally released from bile ducts during liver

preparation and o fixation for thin sectioning were retained and stored in RNAlater (Thermo

Fisher). Levels of gene expression, mutation efficiency, and mutation profile were assessed in

this sample of the liver flukes.

Histopathological investigation

From each hamster, the entire liver was dissected and immersed in 10% buffered formalin. After

overnight fixation, the liver lobes were processed for embedding in paraffin by dehydration

through series of a 70%, 90%, and 100% ethanol, cleared in xylene, subjected to infiltration by

paraffin at 56°C, and last, embedding in paraffin. Four μm sections were sliced by microtome

from the paraffin-embedded liver, the sections stained with hematoxylin and eosin (H&E).

Histopathological grading was undertaken by examination of the stained sections of liver and

bile duct for inflammation, bile duct changes, dysplasia (including dysplasia in

cholangiofibrosis), and stage of CCA as described (21, 75-77), with modifications (Table 1).

717
718

719 **Table 1**. Criteria for histopathological and histochemical assessment and grading.
720
721

Histopathological lesion	Grade description	Reference
Inflammation	0 = None (no/minimal liver tissue or portal inflammation) 1 = Mild (1-2 foci per 4× objective at hepatocyte & periportal area) 2 = Moderate (3-5 foci per 4× objective at hepatocyte & periportal area) 3 = Severe (> 5 foci per 4× objective at hepatocyte & periportal area)	(21, 77, 78)
Bile duct changes	0 = None (absence of proliferation and cholangiofibrosis) 1 = Mild (bile duct proliferation without cholangiofibrosis or periductal fibrosis) 2 = Moderate (bile duct proliferation with cholangiofibrosis) 3 = Severe (bile duct proliferation with cholangiofibrosis and periductal fibrosis)	(21, 77, 78)
Dysplasia	0 = None (No cellular atypia, no nuclear polarity, no nuclear protrusions, no nuclear pseudostratification) 1 = Mild (Cellular atypia+, no nuclear polarity, no nuclear protrusions, nuclear pseudostratification+, nuclei within the lower two-thirds) 2 = Moderate (Cellular atypia+, nuclear polarity+, protruding of nuclei+, nuclear pseudostratification+) 3 = High (Cellular atypia++, nuclear polarity++, protruding of nuclei+, nuclear pseudostratification+)	(76)
Cholangiocarcinoma	0 = None (no evidence of CCA) Low CCA: 1 = Mild (CCA area 1-2 foci per 4× objective) High CCA: combined 2+3: 2 = Moderate (CCA area 3-5 foci per 4× objective) and 3 = Severe (CCA area> 5 foci per 4× objective)	(21, 77, 78)
Fibrosis (PSR stain):	0 = No fibrosis	(32, 79)

| Ishak score | 1 = Fibrous expansion of some portal areas, with or without short fibrous septa
2 = Fibrous expansion of most portal areas, with or without short fibrous septa
3 = Fibrous expansion of most portal areas with occasional portal to portal bridging
4 = Fibrous expansion of portal areas with marked bridging; portal to portal as well as portal to central
5 = Marked bridging (portal–portal and/or portal–central) with occasional nodules (incomplete cirrhosis)
6 = Cirrhosis, probable or definite | |
| **Assessment of collagen proximal to liver flukes** | Quantitative automated evaluation of collagen deposition percentage surrounding bile ducts | ImageJ MRI Fibrosis Tool BioCampus, Montpellier, France (Volker Bäcker 2015), www.mri.cnrs.fr https://dev.mri.cnrs.fr/projects/imagej-macros/wiki/Fibrosis_Tool |

722
723

Fecal egg counts and worm counts

Feces from each hamster were individually collected, weighed and *O. viverrini* eggs per gram of

feces (EPG) calculated using a modified formalin-ethyl acetate technique (80). In brief, hamster

feces were collected and fixed in 10 ml of 10% formalin. Thereafter, the slurry of formalin-fixed

feces was filtered through two layers of gauze, and clarified by centrifugation at 500 *g* for 2 min.

The pellet was resuspended with 7 ml of 10% formalin, mixed with 3 ml ethyl-acetate and

pelleted at 500 *g* for 5 min. The pellet was resuspended in 10% formalin solution and examined

at 400× by light microscopy. EPG was calculated as follows: (average number eggs × total drops

of fecal solution)/ gram of feces. To recover the adult liver flukes, food was withdrawn from the

hamsters 16 hours before euthanasia. Intact mature *O. viverrini* from the hepatobiliary tract were

recovered during observation of the livers using a stereo dissecting microscope and stored for

downstream gene-editing investigation.

Extraction of nucleic acids

Pooled NEJ or single mature worms from either experimental or control groups were

homogenized in RNAzol RT (Molecular Research Center, Inc., Cincinnati, OH) before dual

RNA and DNA extraction as described . Briefly, the parasite(s) were homogenized in RNAzol

RT using a motorized pestle, after which the DNA and protein were precipitated in nuclease free

water. The aqueous upper phase was transferred into a new tube for total RNA precipitation by

isopropanol (50% v/v). The DNA/protein pellet was resuspended in DNAzol and genomic DNA

extracted according to the manufacturer's instructions (Molecular Research Center).

Concentration and integrity of genomic DNA and total RNA were independently quantified by

spectrophotometry (NanoDrop 1000, Thermo Fisher, Waltham, MA). Transcription and

expression were investigated in pools of NEJs and in individual adult flukes after normalization

against the controls.

Quantitative real-time PCR

cDNA was synthesized from DNase I-treated-total RNA (10 ng) using Maxima First Strand

cDNA synthesis with a DNase kit (Thermo Scientific) prior to performing quantitative real-time

PCR (qPCR). Each cDNA sample was prepared for qPCR in triplicate using SSoAdvanced

Universal SYBR Green Supermix (Bio-Rad). Each qPCR reaction consisted of 5 μl of SYBR

Green Supermix, 0.2 μl (10 μM) each of specific forward and reverse primers for *Ov-grn-1*

(forward primer, *Ov*-GRN-1-RT-F: 5'-GACTTGTTGTCGCGGCTTAC-3' and reverse

primer, *Ov*-GRN1-RT-R: 5'-CGCGAAAGTAGCTTGTGGTC-3'), amplifying 147 base pairs

(bp) of 444 nt of *Ov-grn-1* mRNA, complete cds GenBank FJ436341.1) or primers for *Ov-tsp-2*

(forward primer, *Ov-TSP-2*-F 5'- ACAAGTCGTATGTGGAATCA- 3' and reverse primer *Ov*-

TSP-2-R 5'- CCGTCTCGCCTTCTCCTTT- 3', product size 377 bp of 672 nt of *Ov-tsp-2*A

mRNA, complete cds (GenBank JQ678707.1), 2 µl of cDNA and distilled water to a final

volume of 10 µl were used in the reaction. The thermal cycle was a single initiation cycle at

95°C for 10 min followed by 40 cycles of denaturation at 95°C for 15s, annealing at 55°C for 30s

using CFX Connect Real-Time PCR system (Bio-Rad). The endogenous actin gene (1301 nt of

Ov-actin mRNA, GenBank EL620339.1) was used as a reference (17, 81, 82) (forward

primer, *Ov*-actin-F: 5'-AGCCAACCGAGAGAAGATGA and reverse primer, *Ov*-actin-R: 5'-

ACCTGACCATCAGGCAGTTC. The fold change in *Ov-grn-1* and *Ov-tsp*-2 transcripts was

calculated using the $2^{(-\Delta\Delta Ct)}$ method using the *Ov-actin* gene as a reference for normalization (17,

81, 82). Transcript ddCT qPCR data were resampled with replacement bootstrap analysis in

Microsoft Excel with 1000 bootstrap resamples (B = 1000, n = original sample number) to

generate mean values and 95% confidence intervals (83).

Illumina based targeted next generation sequencing of targeted amplicons

The Amplicon-EZ next generation sequencing service (GENEWIZ, South Plainfield, NJ) was

used to obtain deeper coverage of *Ov-grn-1* exon 1 from individual mature worms or pooled

NEJ, providing > 50,000 complete amplicon aligned read-pairs per sample. A 173-nucleotide

region flanking the programmed DSB was amplified with forward primer 5'-

TTCGAGATTCGGTCAGCCG-3' and reverse primer 5'-GCACCAACTCGCAACTTACA-3',

and was sequenced directly using Illumina chemistry. The CRISPR RGEN Tool web platform

(http://www.rgenome.net/about/) analysis, comparison range 60 nt, was used to screen for Cas9-

catalyzed substitutions in the aligned read-pairs with comparisons among the treatment groups

(84, 85). In addition, CRISPResso2 with a quantification window set at 30 nt (28) was employed

for indel estimation, as described (18). The NGS reads are available at GenBank Bioproject

PRJNA385864, BioSample SAMN07287348, SRA study PRJNA385864, accessions SRR

15906234-15906251.

BrdU-staining for proliferation of the biliary epithelium

Proliferation of biliary epithelial cells was investigated by using incorporation of BrdU. In brief,

the liver sections of a paraffin-embedded sample were soaked in xylene, rehydrated in graded

alcohol solution (100%, 90%, and 70% ethanol for 5 min each), and antigen was retrieved in

citrate buffer (pH 6) for 5 min in a high-pressure cooker. The tissue sections were blocked with

3% H_2O_2 in methanol for 30 min and subsequently incubated with 5% fetal bovine serum in

phosphate buffered saline for 30 min at room temperature (RT). The sections were incubated

with monoclonal mouse anti-BrdU (Abcam, catalogue no. ab8955) diluted 1:200 in PBS at 4°C

overnight, and then probed with goat anti-mouse IgG-HRP (Invitrogen, Thermo Fisher) diluted

1:1,000 in PBS for 60 min at RT. The peroxidase reaction was developed with 3, 3'-

diaminobenzidine (DAB). Sections were counterstained with Mayer's hematoxylin for 5 min

before dehydrating and mounting. A positive signal was indicated by a brown color under light

microscopy. The image was captured by a Zeiss Axiocam microscope camera ICc5 and NIS-

Element software (Nikon, Minato, Tokyo, Japan). To quantify BrdU-positive nuclei,

cholangiocytes were counted in 10 non-overlapping fields of 400x magnification, with a total of

1,000 biliary cholangiocytes counted using the counter plug-in tool at ImageJ 1.52P. The cell

proliferation index was calculated as a percentage using the formula: positive biliary nuclei/total

biliary cells x100%.

Staining for mutant forms of p53

To investigate levels of p53 mutation (86) in cholangiocytes, paraffin-embedded tissue sections

were deparaffinized and rehydrated by standard methods. Thereafter, sections were incubated

with monoclonal mouse anti-p53 (mutant, clone Ab-3 PAb240 catalogue no. OP29-200UG)

(Merck, Darmstadt, Germany) diluted 1:100 in PBS at 4°C overnight, and after thorough

washing, probed with goat anti-mouse IgG-HRP (Invitrogen, Carlsbad, CA) diluted 1:1,000 in

PBS for 60 min at 25°C. The peroxidase reaction was developed with 3,3'- DAB and sections

counterstained with Mayer's hematoxylin for 5 min. A human CCA cell line served as the

positive control for p53 positivity (87, 88). Images of high-power fields (400x magnification) of

the biliary epithelium were taken in five non-overlapping fields of each of the right, middle, and

left liver lobes using a Zeiss Axiocam fitted with a ICc5 camera and NIS-Element software

(Nikon). The percentage of mutant p53-positive cholangiocytes was determined by calculating

positive cells from 500 to 800 cholangiocytes from the right, middle, and left lobes of the liver

using imageJ.

PSR staining to evaluate fibrosis

Liver tissue sections where *O. viverrini* reside were selected for fibrosis measurements with PSR

(Abcam, catalogue ab150681). The thin sections were deparaffinized in xylene and rehydrated

through an ethanol gradient. PSR solution was applied to the sections and incubated at 25°C for

60 min. Excess dye was removed by washing twice in dilute acetic acid (0.5%) after which

sections were dehydrated through graded series of ethanol and xylene, the slides cleared with

100% xylene, mounted in Per-mount, and air dried overnight. Fibrosis surrounding the bile duct

(periductal fibrosis, PF) proximal to the liver flukes was evaluated by two approaches. First, by

scoring according to accepted criteria (32): samples were blinded and fibrosis scores (0-6) were

graded semi-quantitatively by Ishak stage (Table 1) by two experienced pathologists. Second,

localized fibrosis was evaluated by capturing images for quantification of collagen deposition.

Specifically, the PSR-stained fibrotic lesions was measured using the plug-in fibrosis tool

developed by Volker Bäcker (New FUJI toolsets for bioimage analysis, available at

https://github.com/MontpellierRessourcesImagerie/imagej_macros_and_scripts/wiki/MRI_Fibro

sis_Tool. We (18) and others (89, 90) have previously used this tool with PSR-stained tissues.

Statistical analysis

One-way ANOVA with Tukey multiple comparisons was used for comparisons with two to four

replicates (worm burden and EPG at week 14). The Krustal-Wallis non parametric test with

Dunn's multiple comparisons was used for datasets that were not normally distributed including

the 24-week EPGs, Ishak and periductal fibrosis, BrdU scores, and mutant p53 signals.

Replicates and errors bars are as listed in each figure legend. Statistical analysis and graphic

presentation of the results were undertaken using GraphPad Prism version 9 (GraphPad Software

Inc, San Diego, CA). To compare mutation rate, a one sample *t*-test compared the replicate

%indel values for the *ΔOv-grn-1* against the single control group % indel for either juvenile or

adult *O. viverrini* worms. Assessment for correlation between % indel and transcription was

carried out using a two-tailed non-parametric Spearman correlation co-efficient (r_s). Values of *P*

≤ 0.05 were considered to be statistically significant: an asterisk (*) corresponds to the control vs

ΔOv-grn-1 group comparison; *, $P \leq 0.05$, **, $P \leq 0.01$, ***, $P \leq 0.001$, hashtag (#) for *ΔOv-grn-*

1 vs *ΔOv-tsp-2* group comparison; #, $P \leq 0.05$; ##, $P \leq 0.01$; ###, $P \leq 0.001$.

Acknowledgements

We thank Suwit Balthaisong for technical assistance. This study was supported by the Research

Program, Research and Graduate Studies, Khon Kaen University to SC, ST and TL, the National

Cancer Institute, National Institutes of Health (NIH) USA award R01CA164719 (AL, TL, PJB,)

and the Australian National Health and Medical Research Council, NHMRC award APP1085309

to AL, JS and TL, and senior principal research fellowship APP1117504 to AL. This research

was funded in part, by the Wellcome Trust, grant number 107475/Z/15/Z to PJB. For the purpose

of Open Access, the authors have applied a CC BY public copyright license to any Author

Accepted Manuscript version arising from this submission. The funders had no role in the study

design, data collection and analysis, decision to publish, and/or preparation of the manuscript.

The content is solely the responsibility of the authors and does not necessarily represent the

official views of the NCI, NIH, NHMRC, or WT.

Author contributions

T. Laha, A. Loukas, P. Brindley, S. Tangkawattana, S. Chaiyadet, W. Ittiprasert, P. Arunsan, and

M. Smout conceived and designed the research; S. Chaiyadet, S. Tangkawattana, R. Deenonpoe,

V. Mann, M. Smout, and W. Ittiprasert performed the research; S. Chaiyadet, S. Tangkawattana,

R. Deenonpoe, and M. Smout recorded the findings ; T. Laha, P.J. Brindley, A. Loukas, M.

Smout and W. Ittiprasert contributed reagents and analytical tools; T. Laha, S. Chaiyadet, P.

Arunsan, S. Tangkawattana, V. Mann, M. Smout, and W. Ittiprasert completed the experiments;

S. Chaiyadet, S. Tangkawattana, W. Ittiprasert, and M. Smout analyzed data; S. Chaiyadet, S.

Tangkawattana, T. Laha, P. Brindley, W. Ittiprasert, M. Smout, and A. Loukas prepared figures

and wrote the paper.

Author ORCiD

Wannaporn Ittiprasert, http://orcid.org/0000-0001-9411-8883

Michael J. Smout, https://orcid.org/0000-0001-6937-0112

Paul J. Brindley, https://orcid.org/0000-0003-1765-0002

Alex Loukas, https://orcid.org/0000-0002-0896-8441

Conflict of interest. The authors declare that this study was performed without any commercial

or financial relationships construable as potential conflicts of interest.

891
892
893 **References**
894

1.   Brindley PJ, Bachini M, Ilyas SI, Khan SA, Loukas A, Sirica AE, et al.
Cholangiocarcinoma. Nat Rev Dis Primers. 2021;7(1):65.

2.   Fedorova OS, Kovshirina AE, Kovshirina YV, Hattendorf J, Onishchenko SV,
Katanakhova LL, et al. Opisthorchis felineus infection is a risk factor for cholangiocarcinoma in
Western Siberia: a hospital-based case-control study. Clin Infect Dis. 2022.
3.   IARC. Biological agents. Volume 100 B. A review of human carcinogens. IARC
monographs on the evaluation of carcinogenic risks to humans. 2012;100(Pt B):1-441.
4.   Brindley PJ, da Costa JM, Sripa B. Why does infection with some helminths cause
cancer? Trends Cancer. 2015;1(3):174-82.
5.   Sripa B, Brindley PJ, Mulvenna J, Laha T, Smout MJ, Mairiang E, et al. The tumorigenic
liver fluke Opisthorchis viverrini--multiple pathways to cancer. Trends Parasitol.
2012;28(10):395-407.
6.   Shin HR, Oh JK, Masuyer E, Curado MP, Bouvard V, Fang YY, et al. Epidemiology of
cholangiocarcinoma: an update focusing on risk factors. Cancer Sci. 2010;101(3):579-85.
7.   Thamavit W, Kongkanuntn R, Tiwawech D, Moore MA. Level of Opisthorchis
infestation and carcinogen dose-dependence of cholangiocarcinoma induction in Syrian golden
hamsters. Virchows Arch B Cell Pathol Incl Mol Pathol. 1987;54(1):52-8.
8.   Srivatanakul P, Ohshima H, Khlat M, Parkin M, Sukaryodhin S, Brouet I, et al.
Opisthorchis viverrini infestation and endogenous nitrosamines as risk factors for
cholangiocarcinoma in Thailand. Int J Cancer. 1991;48(6):821-5.
9.   Sriraj P, Boonmars T, Aukkanimart R, Songsri J, Sripan P, Ratanasuwan P, et al. A
combination of liver fluke infection and traditional northeastern Thai foods associated with
cholangiocarcinoma development. Parasitol Res. 2016;115(10):3843-52.
10.   Thamavit W, Ngamying M, Boonpucknavig V, Boonpucknavig S, Moore MA.
Enhancement of DEN-induced hepatocellular nodule development by Opisthorchis viverrini
infection in Syrian golden hamsters. Carcinogenesis. 1987;8(9):1351-3.
11.   Maizels RM, Smits HH, McSorley HJ. Modulation of Host Immunity by Helminths: The
Expanding Repertoire of Parasite Effector Molecules. Immunity. 2018;49(5):801-18.
12.   Drurey C, Maizels RM. Helminth extracellular vesicles: Interactions with the host
immune system. Mol Immunol. 2021;137:124-33.
13.   Brindley PJ, Loukas A. Helminth infection-induced malignancy. PLoS Pathog.
2017;13(7):e1006393.
14.   Smout MJ, Laha T, Mulvenna J, Sripa B, Suttiprapa S, Jones A, et al. A granulin-like
growth factor secreted by the carcinogenic liver fluke, Opisthorchis viverrini, promotes
proliferation of host cells. PLoS Pathog. 2009;5(10):e1000611.
15.   Smout MJ, Sotillo J, Laha T, Papatpremsiri A, Rinaldi G, Pimenta RN, et al.
Carcinogenic Parasite Secretes Growth Factor That Accelerates Wound Healing and Potentially
Promotes Neoplasia. PLoS Pathog. 2015;11(10):e1005209.
16.   Haugen B, Karinshak SE, Mann VH, Popratiloff A, Loukas A, Brindley PJ, et al.
Granulin Secreted by the Food-Borne Liver Fluke Opisthorchis viverrini Promotes Angiogenesis
in Human Endothelial Cells. Front Med (Lausanne). 2018;5:30.
17.   Papatpremsiri A, Smout MJ, Loukas A, Brindley PJ, Sripa B, Laha T. Suppression of Ov-
grn-1 encoding granulin of Opisthorchis viverrini inhibits proliferation of biliary epithelial cells.
Exp Parasitol. 2015;148:17-23.
18.   Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, et al.
Programmed knockout mutation of liver fluke granulin attenuates virulence of infection-induced
hepatobiliary morbidity. Elife. 2019;8:e41463.

19.   Loilome W, Yongvanit P, Wongkham C, Tepsiri N, Sripa B, Sithithaworn P, et al.
Altered gene expression in Opisthorchis viverrini-associated cholangiocarcinoma in hamster
model. Molecular carcinogenesis. 2006;45(5):279-87.
20.   Thamavit W, Pairojkul C, Tiwawech D, Shirai T, Ito N. Strong promoting effect of
Opisthorchis viverrini infection on dimethylnitrosamine-initiated hamster liver. Cancer Lett.
1994;78(1-3):121-5.
21.   Lvova MN, Tangkawattana S, Balthaisong S, Katokhin AV, Mordvinov VA, Sripa B.
Comparative histopathology of Opisthorchis felineus and Opisthorchis viverrini in a hamster
model: an implication of high pathogenicity of the European liver fluke. Parasitol Int.
2012;61(1):167-72.
22.   Sripa B, Kaewkes S, Sithithaworn P, Mairiang E, Laha T, Smout M, et al. Liver fluke
induces cholangiocarcinoma. PLoS Med. 2007;4(7):e201.
23.   Phumrattanaprapin W, Chaiyadet S, Brindley PJ, Pearson M, Smout MJ, Loukas A, et al.
Orally Administered Bacillus Spores Expressing an Extracellular Vesicle-Derived Tetraspanin
Protect Hamsters Against Challenge Infection With Carcinogenic Human Liver Fluke. J Infect
Dis. 2021;223(8):1445-55.
24.   Suttiprapa S, Sotillo J, Smout M, Suyapoh W, Chaiyadet S, Tripathi T, et al. Opisthorchis
viverrini Proteome and Host-Parasite Interactions. Adv Parasitol. 2018;102:45-72.
25.   Chaiyadet S, Sotillo J, Krueajampa W, Thongsen S, Smout M, Brindley PJ, et al.
Silencing of Opisthorchis viverrini Tetraspanin Gene Expression Results in Reduced Secretion
of Extracellular Vesicles. Front Cell Infect Microbiol. 2022;12:827521.
26.   Morgens DW, Wainberg M, Boyle EA, Ursu O, Araya CL, Tsui CK, et al. Genome-scale
measurement of off-target activity using Cas9 toxicity in high-throughput screens. Nat Commun.
2017;8:15178.
27.   Sithithaworn P, Tesana S, Pipitgool V, Kaewkes S, Pairojkul C, Sripa B, et al.
Relationship between faecal egg count and worm burden of Opisthorchis viverrini in human
autopsy cases. Parasitology. 1991;102 Pt 2:277-81.
28.   Clement K, Rees H, Canver MC, Gehrke JM, Farouni R, Hsu JY, et al. CRISPResso2
provides accurate and rapid genome editing sequence analysis. Nat Biotechnol. 2019;37(3):224-
6.
29.   Hwang G-H, Yu J, Yang S, Son WJ, Lim K, Kim HS, et al. CRISPR-sub: Analysis of
DNA substitution mutations caused by CRISPR-Cas9 in human cells. Computational and
Structural Biotechnology Journal. 2020;18:1686-94.
30.   Shimonishi T, Sasaki M, Nakanuma Y. Precancerous lesions of intrahepatic
cholangiocarcinoma. J Hepatobiliary Pancreat Surg. 2000;7(6):542-50.
31.   Kitadokoro K, Bordo D, Galli G, Petracca R, Falugi F, Abrignani S, et al. CD81
extracellular domain 3D structure: insight into the tetraspanin superfamily structural motifs.
EMBO J. 2001;20(1-2):12-8.
32.   Goodman ZD. Grading and staging systems for inflammation and fibrosis in chronic liver
diseases. J Hepatol. 2007;47(4):598-607.
33.   Bansal PS, Smout MJ, Wilson D, Cobos Caceres C, Dastpeyman M, Sotillo J, et al.
Development of a Potent Wound Healing Agent Based on the Liver Fluke Granulin Structural
Fold. J Med Chem. 2017;60(10):4258-66.
34.   Jusakul A, Cutcutache I, Yong CH, Lim JQ, Huang MN, Padmanabhan N, et al. Whole-
Genome and Epigenomic Landscapes of Etiologically Distinct Subtypes of Cholangiocarcinoma.
Cancer Discov. 2017;7(10):1116-35.

35.   Banales JM, Marin JJG, Lamarca A, Rodrigues PM, Khan SA, Roberts LR, et al.
Cholangiocarcinoma 2020: the next horizon in mechanisms and management. Nature Reviews
Gastroenterology & Hepatology. 2020;17(9):557-88.
36.   Petney TN, Andrews RH, Saijuntha W, Wenz-Mucke A, Sithithaworn P. The zoonotic,
fish-borne liver flukes Clonorchis sinensis, Opisthorchis felineus and Opisthorchis viverrini. Int J
Parasitol. 2013;43(12-13):1031-46.
37.   Mitacek EJ, Brunnemann KD, Suttajit M, Martin N, Limsila T, Ohshima H, et al.
Exposure to N-nitroso compounds in a population of high liver cancer regions in Thailand:
volatile nitrosamine (VNA) levels in Thai food. Food Chem Toxicol. 1999;37(4):297-305.
38.   Sripa B, Tangkawattana S, Laha T, Kaewkes S, Mallory FF, Smith JF, et al. Toward
integrated opisthorchiasis control in northeast Thailand: the Lawa project. Acta Trop.
2015;141(Pt B):361-7.
39.   Gouveia MJ, Pakharukova MY, Laha T, Sripa B, Maksimova GA, Rinaldi G, et al.
Infection with Opisthorchis felineus induces intraepithelial neoplasia of the biliary tract in a
rodent model. Carcinogenesis. 2017;38(9):929-37.
40.   Kim E, Hart T. Improved analysis of CRISPR fitness screens and reduced off-target
effects with the BAGEL2 gene essentiality classifier. Genome Med. 2021;13(1):2.
41.   Chai AWY, Yee PS, Price S, Yee SM, Lee HM, Tiong VK, et al. Genome-wide CRISPR
screens of oral squamous cell carcinoma reveal fitness genes in the Hippo pathway. Elife.
2020;9.
42.   Chaiyadet S, Krueajampa W, Hipkaeo W, Plosan Y, Piratae S, Sotillo J, et al.
Suppression of mRNAs encoding CD63 family tetraspanins from the carcinogenic liver fluke
Opisthorchis viverrini results in distinct tegument phenotypes. Sci Rep. 2017;7(1):14342.
43.   Tran MH, Pearson MS, Bethony JM, Smyth DJ, Jones MK, Duke M, et al. Tetraspanins
on the surface of Schistosoma mansoni are protective antigens against schistosomiasis. Nat Med.
2006;12(7):835-40.
44.   Wendt GR, Collins JN, Pei J, Pearson MS, Bennett HM, Loukas A, et al. Flatworm-
specific transcriptional regulators promote the specification of tegumental progenitors in
Schistosoma mansoni. Elife. 2018;7.
45.   Luo H, Lin Y, Liu T, Lai F-L, Zhang C-T, Gao F, et al. DEG 15, an update of the
Database of Essential Genes that includes built-in analysis tools. Nucleic Acids Research.
2020;49(D1):D677-D86.
46.   Pavani G, Amendola M. Targeted Gene Delivery: Where to Land. Front Genome Ed.
2020;2:609650.
47.   Allen F, Crepaldi L, Alsinet C, Strong AJ, Kleshchevnikov V, De Angeli P, et al.
Predicting the mutations generated by repair of Cas9-induced double-strand breaks. Nat
Biotechnol. 2018.
48.   International Helminth Genomes C. Comparative genomics of the major parasitic worms.
Nat Genet. 2019;51(1):163-74.
49.   Yang YM, Kim SY, Seki E. Inflammation and Liver Cancer: Molecular Mechanisms and
Therapeutic Targets. Semin Liver Dis. 2019;39(1):26-42.
50.   Sripa B, Tangkawattana S, Sangnikul T. The Lawa model: A sustainable, integrated
opisthorchiasis control program using the EcoHealth approach in the Lawa Lake region of
Thailand. Parasitol Int. 2017;66(4):346-54.

51.   Fedorova OS, Fedotova MM, Zvonareva OI, Mazeina SV, Kovshirina YV, Sokolova TS,
et al. Opisthorchis felineus infection, risks, and morbidity in rural Western Siberia, Russian
Federation. PLoS Negl Trop Dis. 2020;14(6):e0008421.
52.   Chan-On W, Nairismagi ML, Ong CK, Lim WK, Dima S, Pairojkul C, et al. Exome
sequencing identifies distinct mutational patterns in liver fluke-related and non-infection-related
bile duct cancers. Nat Genet. 2013;45(12):1474-8.
53.   Ong CK, Subimerb C, Pairojkul C, Wongkham S, Cutcutache I, Yu W, et al. Exome
sequencing of liver fluke-associated cholangiocarcinoma. Nat Genet. 2012;44(6):690-3.
54.   Chaisaingmongkol J, Budhu A, Dang H, Rabibhadana S, Pupacdi B, Kwon SM, et al.
Common Molecular Subtypes Among Asian Hepatocellular Carcinoma and
Cholangiocarcinoma. Cancer Cell. 2017;32(1):57-70 e3.
55.   Montal R, Sia D, Montironi C, Leow WQ, Esteban-Fabro R, Pinyol R, et al. Molecular
classification and therapeutic targets in extrahepatic cholangiocarcinoma. J Hepatol.
2020;73(2):315-27.
56.   Young ND, Gasser RB. Opisthorchis viverrini Draft Genome - Biomedical Implications
and Future Avenues. Adv Parasitol. 2018;101:125-48.
57.   Frangoul H, Altshuler D, Cappellini MD, Chen YS, Domm J, Eustace BK, et al.
CRISPR-Cas9 Gene Editing for Sickle Cell Disease and β-Thalassemia. N Engl J Med.
2021;384(3):252-60.
58.   Fellmann C, Gowen BG, Lin PC, Doudna JA, Corn JE. Cornerstones of CRISPR-Cas in
drug discovery and therapy. Nat Rev Drug Discov. 2017;16(2):89-100.
59.   Rizvi S, Khan SA, Hallemeier CL, Kelley RK, Gores GJ. Cholangiocarcinoma - evolving
concepts and therapeutic strategies. Nat Rev Clin Oncol. 2018;15(2):95-111.
60.   Chansitthichok S, Chamnan P, Sarkhampee P, Lertsawatvicha N, Voravisutthikul P,
Wattanarath P. Survival of Patients with Cholangiocarcinoma Receiving Surgical Treatment in
an O. viverrini Endemic Area in Thailand: A Retrospective Cohort Study. Asian Pac J Cancer
Prev. 2020;21(4):903-9.
61.   Chaiyarit P, Sithithaworn P, Thuwajit C, Yongvanit P. Detection of salivary antibodies to
crude antigens of Opisthorchis viverrini in opisthorchiasis and cholangiocarcinoma patients. Clin
Oral Investig. 2011;15(4):477-83.
62.   Mairiang E, Laha T, Kaewkes S, Loukas A, Bethony J, Brindley PJ, et al. Hepatobiliary
morbidities detected by ultrasonography in Opisthorchis viverrini-infected patients before and
after praziquantel treatment: a five-year follow up study. Acta Trop. 2021;217:105853.
63.   Sripa B, Suwannatrai AT, Sayasone S, Do DT, Khieu V, Yang Y. Current status of
human liver fluke infections in the Greater Mekong Subregion. Acta Trop. 2021;224:106133.
64.   Charoensuk L, Ribas A, Chedtabud K, Prakobwong S. Infection rate of
Opisthorchis viverrini metacercariae in cyprinoid fish from the markets and its association to
human opisthorchiasis in the local community in the Northeast Thailand. Acta Trop.
2022;225:106216.
65.   Pinlaor S, Onsurathum S, Boonmars T, Pinlaor P, Hongsrichan N, Chaidee A, et al.
Distribution and abundance of Opisthorchis viverrini metacercariae in cyprinid fish in
Northeastern Thailand. The Korean journal of parasitology. 2013;51(6):703-10.
66.   Piedrafita D, Spithill TW, Dalton JP, Brindley PJ, Sandeman MR, Wood PR, et al.
Juvenile Fasciola hepatica are resistant to killing in vitro by free radicals compared with larvae
of Schistosoma mansoni. Parasite Immunol. 2000;22(6):287-95.

67.   De Marco Verissimo C, Jewhurst HL, Dobó J, Gál P, Dalton JP, Cwiklinski K. Fasciola
hepatica is refractory to complement killing by preventing attachment of mannose binding lectin
(MBL) and inhibiting MBL-associated serine proteases (MASPs) with serpins. PLoS Pathog.
2022;18(1):e1010226.
68.   Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using
the CRISPR-Cas9 system. Nature Protocols. 2013;8(11):2281-308.
69.   Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing. Nucleic Acids Research.
2014;42(W1):W401-W7.
70.   Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a web tool
for the next generation of CRISPR genome engineering. Nucleic Acids Research.
2016;44(W1):W272-W6.
71.   Young ND, Nagarajan N, Lin SJ, Korhonen PK, Jex AR, Hall RS, et al. The Opisthorchis
viverrini genome provides insights into life in the bile duct. Nat Commun. 2014;5:4378.
72.   Laha T, Pinlaor P, Mulvenna J, Sripa B, Sripa M, Smout MJ, et al. Gene discovery for the
carcinogenic human liver fluke, Opisthorchis viverrini. BMC Genomics. 2007;8:189.
73.   Hulme BJ, Geyer KK, Forde-Thomas JE, Padalino G, Phillips DW, Ittiprasert W, et al.
Schistosoma mansoni α-N-acetylgalactosaminidase (SmNAGAL) regulates coordinated parasite
movement and egg production. PLoS Pathog. 2022;18(1):e1009828.
74.   Suyapoh W, Tangkawattana S, Suttiprapa S, Punyapornwithaya V, Tangkawattana P,
Sripa B. Synergistic effects of cagA+ Helicobacter pylori co-infected with Opisthorchis viverrini
on hepatobiliary pathology in hamsters. Acta Tropica. 2021;213:105740.
75.   Zen Y, Adsay NV, Bardadin K, Colombari R, Ferrell L, Haga H, et al. Biliary
intraepithelial neoplasia: an international interobserver agreement study and proposal for
diagnostic criteria. Modern pathology : an official journal of the United States and Canadian
Academy of Pathology, Inc. 2007;20(6):701-9.
76.   Sato Y, Sasaki M, Harada K, Aishima S, Fukusato T, Ojima H, et al. Pathological
diagnosis of flat epithelial lesions of the biliary tract with emphasis on biliary intraepithelial
neoplasia. J Gastroenterol. 2014;49(1):64-72.
77.   Sudsarn P, Boonmars T, Ruangjirachuporn W, Namwat N, Loilome W, Sriraj P, et al.
Combination of Praziquantel and Aspirin Minimizes Liver Pathology of Hamster Opisthorchis
viverrini Infection Associated Cholangiocarcinoma. Pathology oncology research : POR.
2016;22(1):57-65.
78.   Sudsarn P, Wongchalee N, Boonmars T, Laummaunwai P, Chamgramol Y, Pairojkul C,
et al. Sex differences in opisthorchiosis and the development of cholangiocarcinoma in Syrian
hamster model. Parasitol Res. 2014;113(3):829-35.
79.   Ishak K, Baptista A, Bianchi L, Callea F, De Groote J, Gudat F, et al. Histological
grading and staging of chronic hepatitis. J Hepatol. 1995;22(6):696-9.
80.   Elkins DB, Haswell-Elkins M, Anderson RM. The epidemiology and control of intestinal
helminths in the Pulicat Lake region of Southern India. I. Study design and pre- and post-
treatment observations on Ascaris lumbricoides infection. Trans R Soc Trop Med Hyg.
1986;80(5):774-92.
81.   Piratae S, Tesana S, Jones MK, Brindley PJ, Loukas A, Lovas E, et al. Molecular
characterization of a tetraspanin from the human liver fluke, Opisthorchis viverrini. PLoS Negl
Trop Dis. 2012;6(12):e1939.

82.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method.
Nature Protocols. 2008;3(6):1101-8.
83.   Theodorsson E. Resampling methods in Microsoft Excel for estimating reference
intervals. Biochem Med (Zagreb). 2015;25(3):311-9.
84.   Hwang GH, Yu J, Yang S, Son WJ, Lim K, Kim HS, et al. CRISPR-sub: Analysis of
DNA substitution mutations caused by CRISPR-Cas9 in human cells. Comput Struct Biotechnol
J. 2020;18:1686-94.
85.   Park J, Lim K, Kim JS, Bae S. Cas-analyzer: an online tool for assessing genome editing
results using NGS data. Bioinformatics. 2017;33(2):286-8.
86.   Surget S, Khoury MP, Bourdon JC. Uncovering the role of p53 splice variants in human
malignancy: a clinical perspective. Onco Targets Ther. 2013;7:57-68.
87.   Nakayama M, Oshima M. Mutant p53 in colon cancer. Journal of Molecular Cell
Biology. 2018;11(4):267-76.
88.   Batheja N, Suriawinata A, Saxena R, Ionescu G, Schwartz M, Thung SN. Expression of
p53 and PCNA in Cholangiocarcinoma and Primary Sclerosing Cholangitis. Modern Pathology.
2000;13(12):1265-8.
89.   Rabacal W, Schweitzer F, Rayens E, Tarantelli R, Whang P, Jimenez VC, et al. Statin
treatment prevents the development of pulmonary arterial hypertension in a nonhuman primate
model of HIV-associated PAH. Scientific Reports. 2019;9(1):19832.
90.   Hollenbach M, Thonig A, Pohl S, Ripoll C, Michel M, Zipprich A. Expression of
glyoxalase-I is reduced in cirrhotic livers: A possible mechanism in the development of cirrhosis.
PLoS One. 2017;12(2):e0171260.

1144

1145

1167 **Supporting information: Figures S1-S6 and Tables S1 and S2 (eight items).**
1168
1169
1170

1171
1172
Figure S1. Transcript levels of gene edited NEJ flukes with bootstrapped population values.
Each group of flukes was subjected to gene editing targeting Ov-grn-1 (ΔOv-grn-1 flukes), Ov-
tsp-2 (ΔOv-tsp-2 flukes), or an irrelevant guide RNA as a control (Control). Relative transcript
levels were plotted for both Ov-grn-1 (A) and Ov-tsp-2 genes (B) for all three groups: control
flukes; ΔOv-grn-1 flukes, and ΔOv-tsp-2 flukes. Each panel shows ddCt (delta-delta cycle
threshold) biological replicate values plotted relative to newly excysted juvenile (NEJ) control
average. Resampling with replacement bootstrap analysis (B=1000) of ddCT scores used to
generate population average denoted by thick colored line and 95% confidence interval error
bars.

Figure S2. Experiment 1: Relationship of fecal egg count to worm burden. *Assessment of eggs per gram of feces (EPG) at 10 (A) and 12 (B) weeks after infection compared to worm burden at necropsy (week 14). The worm burden for each hamster was plotted against the EPG at weeks 10 and 12. Comparing these timepoints (C) with linear regression did not reveal variation from a line with zero slope (horizontal line) at both intervals. Each hamster was designated by letter, C= control, G=ΔOv-grn-1, T=ΔOv-tsp-2, with the number of the hamster, as 1, 2, and 3.*

1216
1217
1218
Figure S3. Experiment 1: Transcript levels of gene edited adult flukes with bootstrapped
population values*. Each group of the flukes was subjected to gene editing targeting Ov-grn-1*
(ΔOv-grn-1 flukes), Ov-tsp-2 (ΔOv-tsp-2 flukes), or an irrelevant guide RNA as a control
(Control). Each panel shows ddCt (delta-delta cycle threshold) of individual flukes plotted
*relative to transcript levels of wild-type flukes for Ov-grn-1 (**A**) and Ov-tsp-2 (**B**). The dashed*
line purple box inset is an enlarged section of panel B, included for clarity. Resampling with
replacement bootstrap analysis (B=1000) of ddCT scores used to generate population average
denoted by thick colored line and 95% confidence interval error bars.
1227
1228

1229

1230

1231

1232

1233

1234

1235

Figure S4. Transcript levels of adult flukes from Experiment 2 with bootstrapped population
values. Each group of the 24 wk old flukes was subjected to gene editing targeting Ov-grn-1
(ΔOv-grn-1 flukes), Ov-tsp-2 (ΔOv-tsp-2 flukes), or an irrelevant guide RNA as a control
(Control). Each panel shows ddCt (delta-delta cycle threshold) for individual flukes plotted
relative to wild-type (WT) fluke transcript levels for both Ov-grn-1 (A) and Ov-tsp-2 genes (B).
The dashed line purple box inset is an enlarged section of panel B for clarity. Resampling with
replacement bootstrap analysis (B=1000) of ddCT scores used to generate population average
denoted by thick colored line and 95% confidence interval error bars.

1267

1268

1269

1270

Figure S5. Profiles of nucleotide substitutions. *Nucleotide substitution profiles detected in the*
173 bp amplicon spanning the programmed cleavage site of Ov-grn-1 from both juvenile (NEJ)
and single adult O. viverrini flukes of the ΔOv-grn-1 treatment group compared with the
irrelevant guide RNA treated control group. RGEN's CRISPR-sub analysis tool,
http://www.rgenome.net/crispr-sub/#!, aligns read-pairs to plot the substitution patterns among
Illumina sequence reads from amplicon libraries derived from CRISPR/Cas9 editing-focused
datasets. Experimental group (red, upper axis) versus control group (blue, lower axis); the X-
axis shows the targeted gene including programmed cleavage site (position 0) between
nucleotides 19 and 20 of ORF 1 of Ov-grn-1. Juvenile flukes are shown in the top left and each
adult fluke is shown separately and designated with the number of its host hamster number (x) 1-
15 and worm number (y) 1-3: (h "x" worm "y"). Substantially differences in patterns of
substitutions detected between the experimental and control groups were not apparent.

1283

1291
1292
1293 ***Figure S6. Correlation between fecal EPG and fibrosis****. For Experiment 2, each liver lobe was*
1294 *plotted as eggs per gram against the Ishak fibrosis (data combined from figure 3A and 6B).*
1295 *Data points have been nudged ± 0.1 on the vertical axis for clarity among overlapping points.*
1296 *The linear regression line for each group is shown; ns = not statistically significant; *, P ≤ 0.05.*
1297
1298

1299

Sample	Bases sequenced (Mbp)	Read depth	Read-pairs sequences	Read-pairs aligned (RPA)	RPA Unmodified reads	RPA Modified reads	RPA Insertions	RPA Deletions	Frameshift mutation (number,%)	Indel (%)
NEJs										
Control NEJs	26.3	152,023	113,442	51,402	51,384	18	0	18	0	**0.035%**
ΔOv-grn-1 NEJs	40.8	235,838	159,583	80,571	77,848	2,723	0	2,723	2 (0.02%)	**3.380%**
ADULT FLUKES										
Hamster 1-15: Control flukes	37.9	219,075	129,859	91,783	91,742	41	0	41	0	**0.045%**
$_L$ΔOv-grn-1 flukes (0-0.04% indels)										
Hamster 6: fluke 1	47.8	276,301	185,269	105,363	105,363	0	0	0	0	**0.000%**
Hamster 8: fluke 1	46.6	269,364	171,164	110,079	110,078	1	0	1	0	**0.001%**
Hamster 3: fluke 1	33.5	193,642	150,356	60,954	60,935	19	0	19	0	**0.031%**
Hamster 8: fluke 3	42.5	245,665	154,598	101,288	101,252	36	2	34	0	**0.036%**
Hamster 10: fluke 1	44.7	258,382	143,186	118,253	118,211	41	0	41	19 (48.8%)	**0.035%**
Hamster 9: fluke 1	62.7	362,428	193,741	168,806	168,741	65	0	65	0	**0.039%**
$_M$ΔOv-grn-1 flukes (3.1% indels)										
Hamster 8: fluke 2	48.1	278,035	169,500	116,021	112,458	3,563	1	3,563	8 (0.20%)	**3.071%**
$_H$ΔOv-grn-1 flukes (51-91% indels)										
Hamster 10: fluke 2	49.4	285,549	169,733	128,391	63,378	65,013	0	65,013	20 (0.03%)	**50.637%**
Hamster 15: fluke 2	53.1	306,936	166,531	143,206	64,899	78,307	1	78,307	0	**54.681%**
Hamster 7: fluke 1	46.7	269,942	182,260	102,168	27,617	74,551	0	74,551	0	**72.969%**
Hamster 10: fluke 3	47.5	274,566	161,903	120,064	26,447	93,617	0	93,617	22 (0.02%)	**77.973%**
Hamster 9: fluke 2	38.2	220,809	125,032	99,750	19,707	80,043	2	80,042	17 (0.02%)	**80.244%**
Hamster 15: fluke 1	45.3	261,850	155,792	115,881	10,413	105,468	1	105,468	0	**91.014%**

1300
1301
Table S1. NGS sequencing data summary. *Frequency of successful CRISPR/Cas9 gene knock*
out editing as determined by frameshift mutations. The CRISPResso2 analysis used a window
size (-3 option) that include the whole 173 bp amplicon, except 25 bp at each end in order to the
exclude the primer regions. Combined insertions and deletions are used to generate the Indel%,
the proportion of read-pairs aligned (RPA). To match the color scheme in the figures the control
groups highlighted in blue and low, medium and highly edited adults are highlighted in red,
purple and green. Note that the vast majority of indels are located in the 5' UTR and are not
frameshift mutations.

Parameters Groups	Control	ΔOv-grn-1	ΔOv-tsp-2
CCA development			
CCA positive hamsters/total hamsters (% positive)	10/12 (83.33%)	9/13 (69.23%)	7/13 (53.85%)
Histopathological diagnosis			
Dysplasia (precancerous lesion)	2	1	2
Low grade CCA	2	5	1
High grade CCA	8	4	6
CCA histopathological type			
Tubular	9	8	4
Papillary/Cystic	0	1	0
Mucinous	1	0	3
CCA location (lobe)			
Right	6	5	4
Left	1	2	2
Middle	0	0	1
Right and Left	2	1	0
Right and Middle	1	0	0
Right, Middle and Left	0	1	0
Tumor progression			
Focal	6	5	1
Fully developed	4	4	6

1315
1316
1317
Table S2. Pathology assessment on development of pre-malignant and malignant lesions. *The*
table shows the pathogenesis outcomes from the 3 groups of hamsters and the histopathological
diagnosis data was summarized as graphs in Figure 5. Beyond Figure 5, the table describes the
CCA histopathological type, location, and tumor progression. Below the table histological
*images show the three major histopathological types: **A** = tubular; **B** = papillary; and **C** =*
mucinous CCA.

1324

1325

1326

1327

1328

1329

1330