Mutations in RBD of SARS-CoV-2 Omicron variant result stronger binding to human ACE2 protein

Cecylia S. Lupala[a], Yongjin Ye[a], Hong Chen[b], Xiao-dong Su[b,]*

Haiguang Liu[a,c,]*,

[a]Complex Systems Division, Beijing Computational Science Research Center, Haidian, Beijing 100193, People's Republic of China

[b]School of Life Sciences, State Key Laboratory of Protein and Plant Gene Research and Biomedical Pioneering Innovation Center (BIOPIC), Peking University, Beijing 100871, People's Republic of China

[c]Physics Department, Beijing Normal University, Haidian, Beijing 100875, People's Republic of China

*Corresponding authors: Xiao-dong Su, xdsu@pku.edu.cn Or Haiguang Liu, hgliu@csrc.ac.cn

19
Abstract

The spreading of SARS-CoV-2 virus resulted the COVID-19 pandemic, which has caused

more than 5 millions of death globally. Several major variants of SARS-CoV-2 have

emerged and placed challenges in controlling the infections. The recently emerged

Omicron variant raised serious concerns about reducing efficacy of antibodies or vaccines,

due to its vast mutations. We modelled the complex structure of human ACE2 protein and

the receptor binding domain of Omicron variant, then conducted atomistic molecular

dynamics simulations to study the binding interactions. The analysis shows that the

Omicron variant RBD binds more strongly to the human ACE2 protein than the original

strain. The mutation at the ACE2-RBD interface enhanced the tight binding by increasing

hydrogen bonding interaction and enlarging buried solvent accessible surface area.

Keywords: SARS-CoV-2, Omicron, ACE2, mutation, molecular dynamics simulation

Introduction

The COVID-19 pandemic caused by the SARS-CoV-2 is still affecting global health and economy seriously[1]. According to JHU CSSE COVID-19 Data[2], there are 266 million infections and over 5.26 million fatalities as of December 6, 2021. Several vaccines have been developed and applied to prevent the spreading of SARS-CoV-2 viruses[3], however, these efforts are challenged by emerged virus variants due to mutations [4–7]. Among major variants, several strains were called out to be 'variant of concerns (VOC)' by the world health organization (WHO). On November 26, 2021, the WHO named a new variant (B.1.1.529) to be Omicron, designated to be a VOC [8]. The Omicron variant has accumulated a vast number of mutations, particularly in spike proteins that are responsible for the initiation of infection, i.e., the cell entry. There are 15 mutations on the receptor binding domain (RBD) of the spike protein, which has over 30 mutations in total (see Figure 1) [8,9]. Such a large number of accumulated mutations is unprecedent. Because the spike protein is the target of many antibodies or drugs, it is crucial to investigate the impacts to the efficacy of neutralizing antibodies, under the concerns of immune escapes. Furthermore, about 10 mutations are at the RBD binding interface to the receptor protein, the Angiotensin converting enzyme 2 (ACE2)[10,11]. This level of mutation also raised a question whether the RBD of Omicron variant binds tightly to the ACE2, and whether there is an alternative receptor that facilitate the infection of healthy cells.

Computational modeling and dynamics simulations have been applied to investigate the interactions between the SARS-CoV-2 RBD and the ACE2 proteins [12,13]. Before the structures of RBD-ACE2 complex were resolved experimentally, homology modeling and simulations have successfully predicted the model and quantified the interactions [13,14]. Computer simulations were also used to study the interactions between RBD and ACE2 proteins from other mammals, and the results provide hints on molecular mechanism for

SARS-CoV-2 infection to other animals [15,16]. Here, we followed the same approach,

constructed the structure of human ACE2 and the RBD of Omicron variant (denoted as

ACE2-RBD^O, where the superscript indicates Omicron). Then the structure is subjected to

atomistic molecular dynamics simulations to refine the model and to probe the dynamics.

After comparing to the wild type ACE2-RBD complex system, we found that the RBD^O

exhibits stronger binding to human ACE2 protein, suggesting that the Omicron variant

infects cells via the same mechanism and the infectivity might be enhanced due to the

stronger binding interactions.

Figure 1. Mutations and the diversity of SARS-CoV-2. (**a**) The phylogenetic tree of

SARS-CoV-2. Major variants are labelled on the graph, and the color of clans is according

to the number of spike protein mutations. The tree is generated at https://nextstrain.org. (**b**)

Mutation sites of the receptor binding domain. The residues below the red line are at or

near the ACE2 binding interface.

Results

The structures of RBD and ACE2-RBD complex are stable. The averaged backbone

root-mean-square-deviation (RMSD) of the RBD is less than 1.4 Å compared to the

starting model (Figure 2a). For the wild type RBD, the structure ensembles from two

independent simulations (each 500 ns) deviated by 1.2 Å on average; interestingly, the

RBD^O has averaged RMSD values of 1.4 Å, indicating that mutations only slightly alters

the structure of RBD^O from the wild type RBD. Similarly, the ACE2-RBD complexes are

stable from simulations, reflected on the RMSD with respect to starting complex structures

(Figure 2b). The RMSD for the wild type ACE2-RBD complex are averaged to 3.0 Å and

2.5 Å in the structures sampled from the two trajectories; while the values are 2.2 Å and

2.6 Å for the two simulation trajectories of ACE2-RBD^O. Therefore, we predict that the

mutations in Omicron variant do not significantly reduce the RBD stability, instead, the

ACE2-RBD^O complex is even slightly more stable than the wild type complex according to

the RMSD analysis. The residue fluctuations were analyzed by calculating the root-mean-

square-fluctuations (RMSF) of the RBD (Figure 2c). According to the average values of

RMSF, the RBD^O is more rigid than its wild type (1.5 Å vs. 2.1 Å). The reduction of the

RMSF is more pronounced at the interfacing residues of RBD, also known as the receptor

binding motif (RBD, residues 434-508) [17]. We also closely examined the flucutations of

residues that were mutated (Figure 2c, right panel). We found that the 15 mutated residues

in the Omicron variant consistently exhibit smaller fluctuations, compared to their wild

type counterparts. It is plausible that the binding of ACE2 stabilize these residues, which in

turn enhanced the stability of the ACE2-RBD^O complex. Detailed quantification on

interactions between ACE2 and RBD are elaborated in the following sections.

Figure 2. Stability of the RBD and ACE2-RBD complex structures. (**a**) The RMSD of

RBD with respect to the starting structure. The histogram of each RMSD time trace is

drawn on the right. (**b**) The RMSD of the whole complex with respect to the starting

complex structure, with the histograms shown on the right. (**c**) The RBD residue

fluctuations measured with RMSF. The residue fluctuations at the mutation sites are shown

on the right.

The interactions between ACE2 and RBD is enhanced in Omicron variant. We

extracted the hydrogen bonds formed directly between ACE2 and RBD^{O}, and further

compared to the wild type system (Figure 3). On average, there are 6.5 ± 2.2 hydrogen

bonds formed between ACE2 and RBD^O, about 10% more than 5.9 ± 2.4 hydrogen bonds

observed in the wild type system. A closer examination on the specific hydrogen bonds

reveals that the Q493K and N501Y play important roles in forming new hydrogen bonds

(Table 1). It is worthwhile to note that the hydrogen bonds are very dynamical, and the

total number of hydrogen bonds at any instant time fluctuate significantly. Therefore, in

the table we only listed seven hydrogen bonds that are frequently observed during

simulations, with the occupancy near or above 20%. As shown in Table 1, only the

hydrogen bond between ACE2 S19 and RBD A475 has an occupancy below 20%

(18.73%). In the case of ACE2-RBDO, the next frequently observed hydrogen bond is

between K31 of ACE2 and W456 of RBD^O with an occupancy of 16.25%. As shown in

Table 1, there are five common stable hydrogen bonds observed in both the wild type and

Omicron variant systems. The mutations resulted the loss of two hydrogen bonds: (1) the

K417N mutation caused the loss of hydrogen bonding with ACE2 residue D30, and (2) the

Y505H mutation significantly reduced its bonding to E37 of ACE2. The Q493K mutation

maintains the hydrogen bond between Q493 and E35 of ACE2 in the wild type complex,

adding the possibility of forming a new hydrogen bond between K493 and the D38 of

ACE2. The hydrogen bond between Y501 of RBD^O and the Y41 of the ACE2 is also a new

hydrogen bond frequently observed in simulations. The hydrogen bond between the S19 of

ACE2 and the A475 of RBD^O is stronger than that in the wild type system, although

neither residues were mutated in the Omicron variant. It is possibly influenced by the local

changed due to the S477N and T478K mutations. By comparing the occupancies, we

conclude that the hydrogen bonds between ACE2 and RBD^O is more stable through the

simulations, and therefore resulting more hydrogen bonds on average.

Figure 3. Quantitative analysis of the ACE2-RBD interactions. (**a**) Hydrogen bonds

between ACE2 and RBD/RBD^O. The time traces of hydrogen bond numbers observed

during the simulations. The histograms are shown on the right to compare the statistics

between the wild type system and the Omicron variant system. (**b**) The number of residue

contacts between ACE2 and RBD. (**c**) The buried surface area at the binding interface.

Similar to (a), the histograms are shown to facilitate the comparison in (b,c).

Table 1. Hydrogen bonds between the RBD and the ACE2.

Wild type ACE2-RBD			Omicron ACE2-RBD		
ACE2	RBD	Occupancy	ACE2	RBD^O	Occupancy
K353-Main	G502-Main	57.97%	E35-Side	K493-Side *	75.67%
Y83-Side	N487-Side	50.60%	D355-Side	T500-Side	60.52%
E35-Side	Q493-Side	38.84%	K353-Main	G502-Main	57.03%
D30-Side	K417-Side	34.36%	D38-Side	K493-Side *	52.34%
D355-Side	T500-Side	25.60%	Y83-Side	N487-Side	52.04%
E37-Side	Y505-Side	21.71%	S19-Side	A475-Main	36.59%
S19-Side	A475-Main	18.73%	Y41-Side	Y501-Side *	35.00%

* The residue was mutated from the wild type RBD
The entries shaded in blue color are either not lost or with low occupancy in the Omicron variant
system; the entries with yellow shading are the new hydrogen bonds; the other entries are the
common hydrogen bonds.

We computed the number of van der Waals contacts between the ACE2 and RBD, as well

as the buried surface area, to further assess the binding affinity between ACE2 and RBD.

For the wild type system, the two simulations yields 137 ± 12 contacts on average, while

the ACE2-RBD^O has 148 ± 9 contacts on average (Figure 3b). The statistics on the buries

surface areas are consistent with the level of contacts. The Omicron variant resulted an

increase of buried surface area from 18.5 nm^2 to 19.1 nm^2 (Figure 3c).

The representative structures are highly similar. The representative structures were

selected from the most populated clusters for the ACE2-RBD complexes, in the wild type

and Omicron variant. The RBD structures are similar in the representative models, both

within 1.4 Å RMSD from the crystal structure (see Figure 4). In particular, the RBM

region are highly structurally consistent, in accordance with the tight binding to ACE2. We

computed the electrostatic potentials by solving the Poisson-Boltzmann equation for the

RBM region in three structures: the crystal structure and representative structure of the

wild type RBD, as well as the representative structure of the RBD^O. For the wild type

RBD, positive and negative potential patches are dispersively located at the binding

interface. Strikingly, the same interface has larger patches with positive potentials in the

RDB[O]. For instance, the region centered around G446S-Q493K-G496S-Q498R-N501Y-

Y505H mutation sites exhibits stronger positive electrostatic potentials, improving its

complementary to the charge surface of ACE2 protein. In the corresponding region, the

key residues from ACE2 is composed of D38-Y41-Q42-D355-S446, forming a negatively

charged patch (Figure 4c). We computed the binding energies for the representative

models. In this case, we obtained a representative structure from each simulation trajectory

using the same clustering algorithm, e.g., two for the wild type ACE-RBD, and two for the

Omicron variant system. The binding energies for the two wild type ACE2-RBD structures

are -104.17 kcal/mol and -97.73 kcal/mol. The binding energies for ACE2-RBD[O]

structures have lower values (-112.25 kcal/mol and -107.04 kcal/mol), indicating stronger

binding between ACE2 and RBD[O].

Crystal structure (Wild type)
Representative (Wild type)
Representative (Omicron Variant)

crystal structure
(Wild type 6LZG)

representative
(Wild type 38.4%)

representative
Omicron variant 19.8%

Figure 4. Representative structures and the electrostatic potential surfaces. (a) The

representative structures of wild type RBD (blue) and RBD[O] (red) are superposed to the

crystal structure (green). The bottom panel shows the RBD binding interface to ACE2. (**b**)

The electrostatic potentials on the RBD/RBD[O] surface. (**c**) The RBD binding interface of

ACE2 and its electrostatic potentials (crystal structure of ACE2). The black arrows point to

the positive and negative potential patches.

Detailed structure features at the interface

The interactions at the interface of ACE2-RBD complex for the wildtype have been

previous reported for both the crystal structures [18,19] and the dynamic conformations

from computational simulations [13]. Generally, ACE2 residues 19-42 of the N-terminal

helix, 82-83 near the η1, N330 at helix-13 and 352-357 at the β-hairpin-4,5 are known to

be interacting with RBD. For the RBD, crystal structures show residues K417, G446,

Y449, Y453, L455, F456, A475, F486, N487, Y489, Q493, Y495, G496, Q498, T500,

N501, G502 and Y505 interacting with hACE2, while simulations have reported additional

residues Q474, G476, S477, T478 , E484 and G485 all at the loop (L67) of RBD to

enhance the interactions. Out of the 15 RBD mutations found in the omicron variant, 10

residues (K417N, N440K, G446S, S477N, T478K, E484A, Q493K, G496S, Q498R,

N501Y and Y505H) are located at interface, consequently changing the electrostatics

surface charges at the interface and may have additional effects on the binding of

antibodies and drugs targeting the interface due to the bulkier size of the mutant sidechains

such as in T478K. This also applies for the mutant residue N400K at a loop near the

binding interface with ACE2.

As a result of these mutations, wildtype RBD-ACE2 interactions (Figure 5) such as salt

bridge E484-K31 are lost , K417-D30 are weakened in the Omicron variant due to

shortened side chain, while hydrogen bonds Q493-E35, Q498-K353, Y505-E37 are

enhanced by substitutions, repositioning and formation of new interactions, K493-D38,

R498-Y41, R498-Q42 H505-K353 and N377-Q24 respectively and by supplementing

with additional π-π stacking interaction Y501-Y41. The rest of interactions observed in

wildtype ACE2-RBD (e.g., hydrogen bonds and contacts between Y449-D38, Y453-H34

A475-S19, N487-Y83, T500-N330 and T500-D355) are also maintained in the omicron

variant (Figure 5).

209
210
211
Figure 5. The detailed structure at the ACE2-RBD binding interface. (**a**) the interface of
the wild type ACE2-RBD complex, the amino acids at mutation sites are shown with stick
representations.. The upper panels shows the side chain positions of RBD on the surface of
ACE2, where the surface is colored according to electrostatic potentials (blue for positive and
red for negative potentials); lower panel shows the side chains of both ACE2 and RBD. (**b**)
the interactions between ACE2-RBDO of the Omicron variant. The figure labeling and
coloring scheme are the same as in (a), except residues are the mutated.

220
Discussions and Conclusion

The large number of mutations observed in the spike protein of SARS-CoV-2 raised

serious concerns about the new variant Omicron. Using computational modeling and

simulations, we carried out quantitative analysis on the stability of ACE2-RBD complex

for the Omicron variant, and compared to that of the wild type system. The interactions are

assessed using several quantities, including hydrogen bonding, van der Waals contacts,

buried surface areas, and the binding free energies. The dynamics simulation results and

the quantitative comparison show that the binding interactions between ACE2 and RBD

are slightly stronger for the Omicron variant than for the wild type. This information

provides molecular basis for enhanced infectivity of the Omicron variant. For antibodies

targeting the RBD at the ACE2 binding interface, the efficacy might not be influenced by

the mutation. The study also confirms that the ACE2 can still serve as the receptor for

Omicron variant, despite the large number of mutations.

Methods

Molecular Dynamics Simulation and Analysis

The mutation information of Omicron is retrieved from the US CDC website [9]. We

included 15 mutations occurred in the RDB (see Figure 1). The mutations were

implemented based on the wild type ACE2-RBD complex structure using the Charmm-

GUI webserver [20]. The protonation state was determined under PH 7.0 solvent

environment.

The wild type ACE2-RBD and its Omicron variant were prepared using the CHARMM36

force fields, following the procedure of the CHARMM-GUI webserver. Each system was

solvated in 150 mM sodium chloride solvent with TIP3P water models. Steepest descent

algorithm was applied to minimize the system energy, then each system was equilibrated to

310.15 K (37 °C) within 125 ps. The temperature was maintained by Nose-Hoover scheme

with 1.0 ps coupling constant in the NVT ensemble (constant volume and temperature).

During the equilibration stage, harmonic restraint forces were applied to the molecules (400

kJ mol^{-1} nm^{-2} on backbone and 40 kJ mol^{-1} nm^{-2} on the side chain atoms) [21,22].

Subsequently, the harmonic restraints were removed and the NPT ensembles (constant

pressure and temperature) were simulated at one atmosphere pressure (10^5 Pa) and 310.15 K.

The pressure was maintained by isotropic Parrinello-Rahman barostat [23], with a

compressibility of 4.5×10^{-5} bar^{-1} and a coupling time constant of 5.0 ps. The wild type and

Omicron variant ACE2-RBD systems were both simulated for 2 x 500 ns using the

GROMACS 5.1.2 package [24]. In all simulations, a time step of 2.0 fs was used and the

PME (particle mesh Ewald) [25] was applied for electrostatic interactions beyond 12.0 Å.

The van der Waals interaction cutoff was set to 12.0 Å. Hydrogen atoms were constrained

using the LINCS algorithm [26].

Analyses were carried out with tools in GROMACS (rmsd, rmsf, mindist, sasa) to examine

the system stability. The buried surface area is computed as

$$\Delta A = A_{ACE2} + A_{RBD} - A_{ACE2-RBD} \qquad (1)$$

Where A_{ACE2}, A_{RBD}, and $A_{ACE2-RBD}$ are the solvent accessible surface area computed using

gmx sasa function. The mindist command was used to compute the residue distances, the

residue pairs with distance below 4.0 Å were considered as contacting residues.

VMD was used to analyze hydrogen bonding interactions [27], with the following criteria: D-

A distance cutoff=3.9 Å and D-H-A angle cutoff=20 degrees, where D,A,H are Donor atom,

Acceptor atom, and the Hydrogen atom linked to the Donor atom. Pymol was used for

molecular binding interface, water distributions, visualization, and rending model images

[27]. The adaptive Poisson-Boltzmann equation solver (APBS) was used to compute the

electrostatic potentials [28].

The binding energy was calculated using Prime 3.0 MM-GBSA module of the Schrodinger

24 package . In each ACE2-RBD complex, the ACE2 was treated as the receptor and RBD

was considered as the ligand. Prime MM-GBSA uses OPLS-AA force field and VSGB 2.0

implicit solvation model to estimate the binding energy of the receptor-ligand complex. The

binding energy is calculated as:

$$\Delta G \text{ (bind)} = E_{ACE2-RBD} - (E_{ACE2} + E_{RBD}) \qquad (2)$$

Acknowledgement

Funding from National Natural Science Foundation of China (grant numbers: U1930402,

31971136). The work is supported by a Tianhe-2JK computing time award at Beijing

Computational Science Research Center (CSRC).

283   Competing interests

284   The authors declare no competing interests.

Reference:

[1] I. Ishigami, N.A. Zatsepin, M. Hikita, C.E. Conrad, G. Nelson, J.D. Coe, S. Basu, T.D. Grant, M.H. Seaberg, R.G. Sierra, M.S. Hunter, P. Fromme, R. Fromme, S.R. Yeh, D.L. Rousseau, Crystal structure of CO-bound cytochrome c oxidase determined by serial femtosecond X-ray crystallography at room temperature, Proc. Natl. Acad. Sci. U. S. A. 114 (2017) 8011–8016. https://doi.org/10.1073/pnas.1705628114.

[2] Johns Hopkins Coronavirus Resource Center, (n.d.). https://coronavirus.jhu.edu/vaccines/vaccines-faq (accessed December 6, 2021).

[3] E. Mathieu, H. Ritchie, E. Ortiz-Ospina, M. Roser, J. Hasell, C. Appel, C. Giattino, L. Rodés-Guirao, A global database of COVID-19 vaccinations, Nat. Hum. Behav. 5 (2021) 947–953. https://doi.org/10.1038/s41562-021-01122-8.

[4] J.S. Tregoning, K.E. Flight, S.L. Higham, Z. Wang, B.F. Pierce, Progress of the COVID-19 vaccine effort: viruses, vaccines and variants versus efficacy, effectiveness and escape, Nat. Rev. Immunol. 2021 2110. 21 (2021) 626–636. https://doi.org/10.1038/s41577-021-00592-1.

[5] P.R. Krause, T.R. Fleming, R. Peto, I.M. Longini, J.P. Figueroa, J.A.C. Sterne, A. Cravioto, H. Rees, J.P.T. Higgins, I. Boutron, H. Pan, M.F. Gruber, N. Arora, F. Kazi, R. Gaspar, S. Swaminathan, M.J. Ryan, A.M. Henao-Restrepo, Considerations in boosting COVID-19 vaccine immune responses, Lancet. 398 (2021) 1377–1380. https://doi.org/10.1016/S0140-6736(21)02046-8/ATTACHMENT/99B4689E-FA37-476A-B30E-8D1E3B4E33BC/MMC1.PDF.

[6] A.P. Vashi, O.C. Coiado, The future of COVID-19: A vaccine review, J. Infect. Public Health. 14 (2021) 1461–1465. https://doi.org/10.1016/J.JIPH.2021.08.011.

[7] L. Wang, T. Zhou, Y. Zhang, E.S. Yang, C.A. Schramm, W. Shi, A. Pegu, O.K. Oloniniyi, A.R. Henry, S. Darko, S.R. Narpala, C. Hatcher, D.R. Martinez, Y. Tsybovsky, E. Phung, O.M. Abiona, A. Antia, E.M. Cale, L.A. Chang, M. Choe, K.S. Corbett, R.L. Davis, A.T. DiPiazza, I.J. Gordon, S.H. Hait, T. Hermanus, P. Kgagudi, F. Laboune, K. Leung, T. Liu, R.D. Mason, A.F. Nazzari, L. Novik, S. O'Connell, S. O'Dell, A.S. Olia, S.D. Schmidt, T. Stephens, C.D. Stringham, C.A. Talana, I.T. Teng, D.A. Wagner, A.T. Widge, B. Zhang, M. Roederer, J.E. Ledgerwood, T.J. Ruckwardt, M.R. Gaudinski, P.L. Moore, N.A. Doria-Rose, R.S. Baric, B.S. Graham, A.B. McDermott, D.C. Douek, P.D. Kwong, J.R. Mascola, N.J. Sullivan, J. Misasi, Ultrapotent antibodies against diverse and highly transmissible SARS-CoV-2 variants, Science. 373 (2021). https://doi.org/10.1126/science.abh1766.

[8] Classification of Omicron (B.1.1.529): SARS-CoV-2 Variant of Concern, (n.d.). https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern (accessed December 9, 2021).

[9] Science Brief: Omicron (B.1.1.529) Variant | CDC, (n.d.). https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-omicron-variant.html (accessed December 9, 2021).

[10] J. Shang, G. Ye, K. Shi, Y. Wan, C. Luo, H. Aihara, Q. Geng, A. Auerbach, F. Li, Structural basis of receptor recognition by SARS-CoV-2, Nature. 581 (2020) 221–224. https://doi.org/10.1038/s41586-020-2179-y.

[11] R. Yan, Y. Zhang, Y. Li, L. Xia, Y. Guo, Q. Zhou, Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2, Science. 367 (2020) 1444–1448. https://doi.org/10.1126/science.abb2762.

[12] Y. Wang, M. Liu, J. Gao, Enhanced receptor binding of SARS-CoV-2 through networks of hydrogen-bonding and hydrophobic interactions, Proc. Natl. Acad. Sci. 117 (2020) 13967–13974. https://doi.org/10.1073/pnas.2008209117.

[13] C.S. Lupala, X. Li, J. Lei, H. Chen, J. Qi, H. Liu, X.-D. Su, Computational simulations

reveal the binding dynamics between human ACE2 and the receptor binding domain
of SARS-CoV-2 spike protein, Quant. Biol. 9 (2021) 61. https://doi.org/10.15302/J-
QB-020-0231.

[14] X. Xu, P. Chen, J. Wang, J. Feng, H. Zhou, X. Li, W. Zhong, P. Hao, Evolution of the
novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein
for risk of human transmission, Sci. China Life Sci. 63 (2020) 457–460.
https://doi.org/10.1007/s11427-020-1637-5.

[15] J. Damas, G.M. Hughes, K.C. Keough, C.A. Painter, N.S. Persky, M. Corbo, M.
Hiller, K.P. Koepfli, A.R. Pfenning, H. Zhao, D.P. Genereux, R. Swofford, K.S.
Pollard, O.A. Ryder, M.T. Nweeia, K. Lindblad-Toh, E.C. Teeling, E.K. Karlsson,
H.A. Lewin, Broad host range of SARS-CoV-2 predicted by comparative and
structural analysis of ACE2 in vertebrates, Proc. Natl. Acad. Sci. U. S. A. 117 (2020)
22311–22322. https://doi.org/10.1073/pnas.2010146117.

[16] C.S. Lupala, V. Kumar, X. Su, C. Wu, H. Liu, Computational insights into differential
interaction of mammalian angiotensin-converting enzyme 2 with the SARS-CoV-2
spike receptor binding domain, Comput. Biol. Med. (2021) 105017.
https://doi.org/10.1016/j.compbiomed.2021.105017.

[17] C. Yi, X. Sun, J. Ye, L. Ding, M. Liu, Z. Yang, X. Lu, Y. Zhang, L. Ma, W. Gu, A.
Qu, J. Xu, Z. Shi, Z. Ling, B. Sun, Key residues of the receptor binding motif in the
spike protein of SARS-CoV-2 that interact with ACE2 and neutralizing antibodies,
Cell. Mol. Immunol. 17 (2020) 621–630. https://doi.org/10.1038/s41423-020-0458-z.

[18] Q. Wang, Y. Zhang, L. Wu, S. Niu, C. Song, Z. Zhang, G. Lu, C. Qiao, Y. Hu, K.Y.
Yuen, Q. Wang, H. Zhou, J. Yan, J. Qi, Structural and Functional Basis of SARS-
CoV-2 Entry by Using Human ACE2, Cell. 181 (2020) 894-904.e9.
https://doi.org/10.1016/j.cell.2020.03.045.

[19] J. Lan, J. Ge, J. Yu, S. Shan, H. Zhou, S. Fan, Q. Zhang, X. Shi, Q. Wang, L. Zhang,
X. Wang, Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
ACE2 receptor., Nature. (2020). https://doi.org/10.1038/s41586-020-2180-5.

[20] J. Lee, X. Cheng, J.M. Swails, M.S. Yeom, P.K. Eastman, J.A. Lemkul, S. Wei, J.
Buckner, J.C. Jeong, Y. Qi, S. Jo, V.S. Pande, D.A. Case, C.L. Brooks, A.D.
MacKerell, J.B. Klauda, W. Im, CHARMM-GUI Input Generator for NAMD,
GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the
CHARMM36 Additive Force Field, J. Chem. Theory Comput. 12 (2016) 405–413.
https://doi.org/10.1021/acs.jctc.5b00935.

[21] S. Nosé, A unified formulation of the constant temperature molecular dynamics
methods, J. Chem. Phys. 81 (1984) 511–519. https://doi.org/10.1063/1.447334.

[22] W.G. Hoover, Canonical dynamics: Equilibrium phase-space distributions, Phys. Rev.
A. 31 (1985) 1695–1697. https://doi.org/10.1103/PhysRevA.31.1695.

[23] M. Parrinello, A. Rahman, Polymorphic transitions in single crystals: A new molecular
dynamics method, J. Appl. Phys. 52 (1981) 7182–7190.
https://doi.org/10.1063/1.328693.

[24] M.J. Abraham, T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, E. Lindah,
Gromacs: High performance molecular simulations through multi-level parallelism
from laptops to supercomputers, SoftwareX. 1–2 (2015) 19–25.
https://doi.org/10.1016/j.softx.2015.06.001.

[25] T. Darden, D. York, L. Pedersen, Particle mesh Ewald: An N·log(N) method for
Ewald sums in large systems, J. Chem. Phys. 98 (1993) 10089–10092.
https://doi.org/10.1063/1.464397.

[26] B. Hess, H. Bekker, H.J.C. Berendsen, J.G.E.M. Fraaije, LINCS: A Linear Constraint
Solver for molecular simulations, J. Comput. Chem. 18 (1997) 1463–1472.

385   https://doi.org/10.1002/(SICI)1096-987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-
386   H.

387 [27] W. Humphrey, A. Dalke, K. Schulten, VMD: Visual molecular dynamics, J. Mol.
388   Graph. 14 (1996) 33–38. https://doi.org/10.1016/0263-7855(96)00018-5.

389 [28] E. Jurrus, D. Engel, K. Star, K. Monson, J. Brandi, L.E. Felberg, D.H. Brookes, L.
390   Wilson, J. Chen, K. Liles, M. Chun, P. Li, D.W. Gohara, T. Dolinsky, R. Konecny,
391   D.R. Koes, J.E. Nielsen, T. Head-Gordon, W. Geng, R. Krasny, G.W. Wei, M.J. Holst,
392   J.A. McCammon, N.A. Baker, Improvements to the APBS biomolecular solvation
393   software suite, Protein Sci. 27 (2018) 112–128. https://doi.org/10.1002/pro.3280.

394
395
396
397 **Supporting Information**
398