Molecular architecture of 40S initiation complexes on the Hepatitis C virus

IRES: from ribosomal attachment to eIF5B-mediated reorientation of initiator

tRNA

Author List and Affiliations

Zuben P. Brown[1], Irina S. Abaeva[2], Swastik De[1], Christopher U.T. Hellen[2], Tatyana V. Pestova*[2],

and Joachim Frank*[1,3].

[1]Department of Biochemistry and Molecular Biophysics, Columbia University, New York, NY,

USA

[2]Department of Cell Biology, SUNY Downstate Health Sciences University, Brooklyn, NY, USA

[3]Department of Biological Sciences, Columbia University, New York, NY, USA

*Authors for correspondence (Tatyana.Pestova@downstate.edu; jf2192@cumc.columbia.edu)

Lead Contact: Joachim Frank*[1,3]

SUMMARY

Hepatitis C virus mRNA contains an internal ribosome entry site (IRES) that mediates end-independent

translation initiation, requiring a subset of eukaryotic initiation factors (eIFs). Direct binding of the IRES

to the 40S subunit places the initiation codon into the P site, where it base-pairs with eIF2-bound Met-

tRNA$_i^{Met}$ forming a 48S initiation complex. Then, eIF5 and eIF5B mediate subunit joining. Initiation can

also proceed without eIF2, in which case Met-tRNA$_i^{Met}$ is recruited directly by eIF5B. Here, we present

cryo-EM structures of IRES initiation complexes at resolutions up to 3.5 Å that cover all major stages from

initial ribosomal association, through eIF2-containing 48S initiation complexes, to eIF5B-containing

complexes immediately prior to subunit joining. These structures provide insights into the dynamic network

of 40S/IRES contacts, highlight the role for IRES domain II, and reveal conformational changes that occur

during the transition from eIF2- to eIF5B-containing 48S complexes that prepare them for subunit joining.

Keywords

hepatitis C virus IRES, ribosome, eIF2, eIF5B, translation initiation, cryo-EM

INTRODUCTION

The canonical initiation process begins with formation of the 43S preinitiation complex (PIC) comprising the 40S ribosomal subunit, the eIF2•GTP/Met-tRNA$_i^{Met}$ ternary complex (eIF2-TC), eIF1, eIF1A and eIF3 (Jackson et al., 2010). The 43S PIC attaches to the capped 5′-terminal region of mRNA and then scans to the initiation codon in a favorable nucleotide context (containing A/G and G at the -3 and +4 positions relative to the AUG, respectively) where it stops and forms the 48S initiation complex (IC) with established codon-anticodon base-pairing. Attachment is mediated by eIFs 4A, 4B and eIF4F, which cooperatively unwind the cap-proximal region allowing attachment and also assist 43S PIC scanning. eIF1, in cooperation with eIF1A, induces an 'open' scanning-competent conformation of the 43S PIC and monitors the fidelity of initiation codon selection (Pestova et al., 1998a; Pestova and Kolupaeva, 2002; Passmore et al., 2007; Hussain et al., 2014). Establishment of codon-anticodon base-pairing in the 48S IC leads to dissociation of eIF1 and eIF5-induced hydrolysis of eIF2-bound GTP, and thereby switches the 40S subunit to the 'closed' conformation (Unbehaun et al., 2004; Maag et al., 2005). After that, eIF5B, in its GTP-bound form, displaces residual eIF2•GDP (Pisarev et al., 2006) and promotes joining of the 60S subunit (Pestova et al., 2000). Interaction of eIF5B with eIF1A enhances eIF5B's subunit joining activity and the hydrolysis of eIF5B-bound GTP, leading to coupled release of eIF5B•GDP and eIF1A from the assembled 80S ribosome (Marintchev et al., 2003; Acker et al., 2006; Nag et al., 2016).

A number of viral mRNAs contain internal ribosomal entry sites (IRESs), structured RNA regions that mediate cap-independent initiation of translation using a subset of the eIFs that are required by canonical initiation. All IRES-mediated initiation mechanisms are based on non-canonical interactions of IRESs with canonical components of the translation apparatus (Jackson et al, 2010). The ~300nt-long hepatitis C virus (HCV) IRES is located in the 5′-terminal region of the viral genome and epitomizes a class of related RNA elements. HCV-like IRESs occur in the genomes of pestiviruses (e.g., classical swine fever virus (CSFV)), some pegiviruses and numerous members of *Picornaviridae* (Arhab et al., 2020). The HCV IRES comprises three domains (II–IV), with domain III divided into several subdomains (Figure 1A). Ribosomal recruitment of HCV and HCV-like IRESs occurs by direct binding of the IRES to the 40S subunit and does not involve scanning, group 4 eIFs or eIF1 (Pestova et al., 1998b). Domain III binds at the back of the 40S subunit, whereas the long, bent domain II loops out and reaches into the E site. The sites of

interaction with the 40S subunit include domains IIIa and IIIc (which bind to eS1, eS26 and eS27),

the apex of domain IIId (which base-pairs to expansion segment (ES) 7 of 18S rRNA), domain

IIIe (which interacts with helix (h) 26 of 18S rRNA), and the apex of domain II, which interacts

with uS7 and eS25 in the head and uS11 on the platform of the 40S subunit, intercalating into the

mRNA binding channel and causing tilting of the head and forcing the 40S subunit to adopt the

open conformation (Kolupaeva et al., 2000; Kieft et al., 2001; Malygin et al., 2013a; 2013b;

Hashem et al., 2013; Angulo et al., 2016; Matsuda and Mauro, 2014; Quade et al., 2015;

Yamamoto et al., 2015; Yokoyama et al., 2019).

Although domain II does not contribute to the affinity of the HCV IRES to the 40S subunit

(e.g. Kieft et al., 2001; Spahn et al., 2001), the open conformation of the 40S subunit promoted by

domain II facilitates loading of the region containing the initiation codon into the mRNA binding

channel, accounting for the stimulatory role of domain II during initiation on HCV-like IRESs

(Honda et al., 1996; Reynolds et al., 1996; Filbin and Kieft, 2011; Odreman-Macchioli et al.,

2001). Upon binding to the 40S subunit, domain IV of the HCV IRES unfolds, and the initiation

codon is placed in the immediate vicinity of the P site, where it base-pairs with the anticodon of

Met-tRNA$_i$^{Met} as a part of the eIF2-TC, leading to formation of the 48S IC (Pestova et al., 1998b).

After that, eIF5 and eIF5B mediate the subunit joining step to complete formation of the

elongation-competent 80S ribosome (Locker et al., 2007; Pestova et al., 2008; Terenin et al., 2008).

Notably, in 80S complexes assembled on the HCV IRES, the P-site Met-tRNA$_i$^{Met} and eIF5B•GTP,

which correspond to the last stage in the initiation process prior to formation of an elongation-

competent ribosome, the tilt of the 40S subunit is reversed, and the apex of domain II is released

from its position on the head of the 40S subunit (Yamamoto et al., 2014). Remarkably, when levels

of active eIF2 are reduced due to stress-induced phosphorylation, Met-tRNA$_i$^{Met} can be recruited

by eIF5B instead to the IRES/40S complexes (Pestova et al., 2008; Terenin et al., 2008). In both

eIF2- and eIF5B-mediated pathways, eIF1A enhances 48S complex formation (de Breyne et al.,

2008; Jaafar et al., 2016), whereas eIF1 inhibits the process and even induces dissociation of pre-

assembled 48S ICs, but this inhibition can be alleviated by deletion of domain II (Pestova et al.,

2008).

In addition to 40S subunits, HCV and related IRESs also bind to eIF3 via their apical IIIa

and IIIb domains (Pestova et al., 1998b; Sizova et al., 1998; Ji et al., 2004; Hashem et al., 2013).

Strikingly, in 40S/IRES/eIF3 complexes, HCV-like IRESs displace eIF3 from its ribosomal

position (Hashem et al., 2013), usurping eIF3's key ribosomal contacts involving eS1, eS26 and

eS27 (des Georges et al., 2015). Moreover, the ribosome-binding surface of eIF3 is now involved

in interaction with the IRES (Hashem et al., 2013). In *in vitro* reconstituted initiation reactions,

eIF3 only modestly enhances 48S complex formation on HCV-like IRESs (Pestova et al., 1998b;

Hashem et al., 2013), which led to the suggestion that *in vivo*, the role of the eIF3/IRES interaction

is likely to relieve the competition between the IRES and eIF3 for a common ribosomal binding

site, and to reduce formation of 43S PICs, thereby favoring translation of viral mRNAs (Hashem

et al., 2013).

Cryo-EM studies have been indispensable in providing insights into the architecture and

interactions of HCV and HCV-like IRES ribosomal complexes, as well as the mechanism of the

IRES function, initially through low-resolution 40S/HCV IRES and 80S/HCV IRES structures

(Spahn et al. 2001; Boehringer et al, 2005) and continuing with sub-nanometer resolution

structures of 40S/eIF3/CSFV IRES and 80S/Met-tRNA$_i^{Met}$/eIF5B•GMPPNP/HCV IRES

functional complexes (Hashem et al., 2013; Yamamoto et al., 2014), and the more recent near-

atomic resolution reconstructions of 80S•HCV IRES complexes (Yamamoto et al., 2015; Quade

et al., 2015; Yokoyama et al., 2019). However, despite these advances, the structures of 48S ICs

assembled on the HCV IRES, as well as the transitions between different states in the initiation

pathways and accompanying conformational changes have remained unknown. To fill these gaps,

we present cryo-EM structures of HCV IRES ribosomal complexes up to 3.5 Å resolution that

cover all major stages of IRES-mediated initiation pathways from IRES binding to the 40S subunit

through eIF2-containing 48S ICs to the final eIF5B-containing 48S ICs immediately prior to the

joining of the 60S subunit. Individually, these structures also provide detailed insights into the

dynamic network of contacts between the IRES and the 40S subunit, highlight the role for IRES

domain II, and importantly, include the first structure of eIF5B bound to the 40S subunit, prior to

subunit joining.

RESULTS AND DISCUSSION

Overview of cryo-EM analysis of initiation complexes assembled on the wt and the ΔdII HCV
IRES with eIF2 or eIF5B

To capture discrete states within either the eIF2- or eIF5B-containing IRES-mediated
initiation pathways and to visualize the role of IRES domain II in these processes, initiation
complexes were assembled *in vitro* by incubating the *wt* or the ΔdII mutant IRES (Figure 1A) with
individual purified translation components. To follow the eIF2-mediated pathway, reaction
mixtures were prepared containing the *wt* or the ΔdII IRES, 40S ribosomal subunits, eIF2, eIF3,
eIF1A and Met-tRNA$_i^{Met}$, and to follow the eIF5B-mediated pathway, eIF2 was replaced by eIF5B,
thus yielding four discrete sample types (Table S1). Cryo-EM grids of each complex were imaged
at 300 kV producing high-contrast micrographs with easily identifiable 40S ribosomal particles
(Figure S1A-E; Table S2). The images were processed using maximum-likelihood classification
techniques implemented in Relion 3.1 (Scheres, 2012; 2016; Zivanov et al., 2018; 2019) yielding
18 structures containing different sets of components at resolutions as high as 3.5 Å (Figure S2;
Table S3). Although some flexible regions had a poor local resolution (e.g., eS12 in the 40S head
or IRES domain IIIb), most of the ribosome, all IRES-ribosome contacts, and all initiation factors
present had resolutions, between 3-7 Å (Figures S2 and S3), that allowed modeling of all these
components. None of the structures obtained contained eIF3. During initiation on HCV-like
IRESs, eIF3 interacts with the apical region of IRES domain III rather than with the 40S subunit
(Hashem et al., 2013). This interaction is sensitive to the process of grid preparation and is more
stable when grids have thicker ice so that imaging complexes that contain eIF3 requires the
intentional selection of regions with sufficiently thick ice (e.g., Hashem at al., 2013; Neupane et
al., 2020); however, our study aimed to determine the details of ribosomal interactions with the
IRES, initiation factors and Met-tRNA$_i^{Met}$ at high resolution, which relies on imaging in regions
with thinner ice. Importantly, however, the absence of eIF3 does not affect data interpretation
because all studied complexes can be assembled efficiently without eIF3 (Pestova et al., 1998b;
2008).

The structures obtained comprise the 40S/IRES$_{ΔdII}$ binary complex in various
conformational states (structures 1$_{ΔdII}$-9$_{ΔdII}$); the 40S/IRES$_{wt}$ binary complex in a single
conformational state (structure 10$_{wt}$); the 40S/eIF1A/IRES$_{wt}$ ternary complex (structure 11$_{wt}$); 48S

ICs containing eIF2, Met-tRNA$_i^{Met}$, eIF1A and the *wt* or the ΔdII IRES (structures 12_{wt} and $12_{ΔdII}$);

48S complexes containing the *wt* or the ΔdII IRES base-paired with the P-site Met-tRNA$_i^{Met}$ but

lacking eIF2 and thus mimicking the stage after eIF2 dissociation following hydrolysis of GTP

(structures 13_{wt} and $13_{ΔdII}$); the pre-48S IC containing eIF5B, eIF1A, the *wt* IRES and P-site Met-

tRNA$_i^{Met}$ that is not base-paired with the initiation codon (structure 14_{wt}); and 48S ICs containing

eIF5B, eIF1A, Met-tRNA$_i^{Met}$ and the *wt* or the ΔdII IRES (structures 15_{wt} and $15_{ΔdII}$) (Figure 1B).

Thus, the structures obtained cover the entire initiation pathway, starting with initial

binding of the IRES to the 40S subunit and finishing with the eIF5B-containing 48S complex prior

to subunit joining, and also provide details of the dynamic interactions between the IRES and the

40S subunit (Figures 1C and S4; Table S4).

Stepwise binding of the IRES to the 40S subunit

Binding of the IRES to the 40S subunit involves multiple contacts formed by several IRES

domains (IIIa, IIIc, IIId, IIIe, S2) and ribosomal proteins eS1, eS27, eS28 as well as h26 in ES7 of

18S rRNA (e.g., Quade et al., 2015; Yamamoto et al., 2015). Deletion or mutation of these domains

impairs binding of the IRES to the 40S subunit to different extents, reflecting their cumulative

importance for IRES function (Kieft et al., 2001). Strikingly, classification of ribosomal complexes

formed on the ΔdII IRES identified a small proportion of 40S/IRES binary complexes that showed

conformational differences in the individual positions of IRES domains IIIa/IIIc/IIId/IIIe/IIIf

(structures $1_{ΔdII}$-$6_{ΔdII}$) compared to other complexes, in all of which these domains had similar

positions (structures $7_{ΔdII}$-$15_{wt/ΔdII}$) (Figure 1B). The local resolution of the IRES contacts with the

40S subunit in these maps was sufficient for model building of all components (Figures S2 and

S3) and allowed detailed examination of the relative motion of the ΔdII IRES as it transitioned

from a minimally associated state (structure $1_{ΔdII}$) to the canonically bound conformation in which

domains IIIa, IIIc, IIId, and IIIe contact eS1, eS27 and h26 (structure $6_{ΔdII}$ and structures $7_{ΔdII}$-

$15_{wt/ΔdII}$).

By ordering these structures based on the number of IRES-40S subunit contacts and

displacement from the canonically-bound IRES position we produced a putative sequence of

binding events between the ΔdII IRES and the 40S subunit (Figure 2A; Table S4). Comparison

between the least- and the most- bound states (structures $1_{ΔdII}$ and $6_{ΔdII}$) shows that the IRES

domains undergo displacement of varying extents during IRES binding (Figure 2B). Across all

structures, the most uniform regions of the IRES are the linked domains IIIa and IIIc (Figures 2A-
B), which contact eS27 via nt. 163 and 233 as reported (Quade et al., 2015; Yamamoto et al.,
2015). Given that these domains undergo minimal structural change regardless of the conformation
of the rest of the IRES, the observed impairment of IRES activity by nucleotide substitutions (e.g.,
Tang et al., 1999) suggests that these interactions are critical for correct IRES function. Other IRES
domains, however, are more dynamic. Thus, to transition from the least- to the most-bound states
requires translation of domain IIId by 10.6 Å toward the intersubunit face and domains
hIII$_1$/IIIe/IIIf by 23.3 Å toward the mRNA exit channel, whereas domains IIIa/IIIc only move by
5.9 Å (Figure 2B).

The fully bound IRES forms Watson-Crick base pairs between ES7 nt U$_{1114\text{-}1118}$ and IRES
domains IIId (GGG$_{266\text{-}268}$), IIIe (A$_{296}$), and hIII$_1$ (A$_{136}$) as well as a stacking interaction between
domain IIIe (G$_{295}$) and U$_{1115}$ of ES7 (Quade et al., 2015; Yamamoto et al., 2015). This network of
interactions also occurred in the later-stage complexes (structures 6$_{\Delta dII}$ and 7$_{\Delta dII}$-15$_{wt/\Delta dI}$) but they
were not present during early-stage association of the IRES with the 40S subunit (structure 1$_{\Delta dII}$)
(Figure 2C; summarized in Figure S4 and Table S4). Initially, the only contact between the IRES
and ES7 is a transient and previously undescribed stacking pair between G$_{296}$ in domain IIIe and
U$_{1115}$ of 18S rRNA. Structure 1$_{\Delta dII}$ also shows another transient, previously undescribed hydrogen
bond between U$_{265}$ in domain IIId and Lys199 in eS1, a residue that instead interacts with IIIe in
the fully bound IRES (Quade et al., 2015; Yamamoto et al., 2015). These two contacts along with
the domain IIIa/IIIc interactions with eS27 are the only bonds between the IRES and the 40S
subunit in this complex (Table S4). In structure 2$_{\Delta dII}$, domain IIId begins to be repositioned, moving
by 4.4 Å relative to structure 1$_{\Delta dII}$, and the transient domain IIIe-U$_{1115}$ contacts are lost. Although
the full complement of interactions with ES7 is not present, the Watson-Crick base-pairs between
domain IIId (GG$_{266\text{-}267}$) and CC$_{1117\text{-}1118}$ and stacking interactions of G$_{295}$ and A$_{296}$ in domain IIIe
with ES7 have formed. These Watson-Crick base pairs and the G$_{295}$ stacking interaction are
maintained in all subsequent (structures 2$_{\Delta dII}$-6$_{\Delta dII}$) and fully bound complexes (structures 7$_{\Delta dII}$-
15$_{wt/\Delta dI}$), but the A$_{296}$ stacking interaction with U$_{1115}$ is only present for this and the following
complex (structures 2$_{\Delta dII}$ and 3$_{\Delta dII}$). Another interaction that first appears in structure 2$_{\Delta dII}$ is the eS27
Glu75 hydrogen bond with domain IIId nts. 266-267. It is maintained in all subsequent structures
except those that have domain II inserted into the E site of the 40S subunit (structures 10$_{wt}$, 11$_{wt}$
and 14$_{wt}$; Table S4). In the following complex (structure 3$_{\Delta dII}$), domain IIId is located closer to the

platform of the 40S subunit, moving by 2.8 Å; the $S1/S2/hIII_1/IIIe/IIIf$ region moves by 5.0 Å,

whereas the position of the stable domain IIIa/IIIc changes by only 1.5 Å. This repositioning breaks

none of the contacts formed in structure $2_{\Delta dII}$ and allows the formation of a new base-pair between

G_{268} in domain IIId and C_{1116} in ES7. The IRES domains continue to move closer to their canonical

bound positions in structure $4_{\Delta dII}$, which lacks the transient A_{296}/U_{1115} interaction but maintains all

other ribosomal contacts (Table S4). In structures $5_{\Delta dII}$ and $6_{\Delta dII}$, the final canonical interactions of

$hIII_1$ (A_{136}) and domain IIIe (A_{296}) with ES7 are present. The contact between A_{136} and U_{1115} is

enabled because the base pairing between domain IIIe U_{297} and A_{288} in $hIII_1$ that induces the

flipping-out of A_{136} exists in all structures ($1_{\Delta dII}$-$15_{wt/\Delta dII}$) (Easton et al., 2009). Structures $5_{\Delta dII}$ and

$6_{\Delta dII}$ are also the first complexes in which the IRES is in a position to form a hydrogen bond between

Asn147 in eS1 and the phosphate backbone of $GG_{300\text{-}301}$, an interaction that is maintained

throughout all subsequent complexes (Table S4).

Taken together, this series of structures ($1_{\Delta dII}$-$6_{\Delta dII}$) indicates a likely sequence of binding

events between the 40S subunit and the IRES from initial encounter to the canonically-bound

conformation in which domain IIIa/IIIc, the first element of the IRES to bind to the 40S subunit,

acts as a pivot to dock domain IIId onto ES7. These structures may represent transient states in

binding of both the ΔdII and *wt* IRES, which we were able to capture in the former case because

the altered kinetic landscape of the initiation pathway in the absence of domain II allowed them to

accumulate and be observed.

An important corollary of IRES binding is the conformational changes that occur in the

40S subunit. Whereas complexes with an incompletely accommodated IRES (structures $1_{\Delta dII}$-$5_{\Delta dII}$)

contain 40S subunits in the analogous closed conformation, the ribosomal structures with the full

complement of IRES/40S contacts (structure $6_{\Delta dII}$-$9_{\Delta dII}$) show a striking difference between the

position of the head, from the closed conformation in structure $6_{\Delta dII}$ (matching the head position in

structures $1_{\Delta dII}$-$5_{\Delta dII}$) to the fully open state in structure $9_{\Delta dII}$ (Figure 2D-E). Although structures $7_{\Delta dII}$-

$9_{\Delta dII}$ have a canonically bound IRES with domain III contacting eS1, eS27 and ES7 as in structure

$6_{\Delta dII}$, they show large-scale conformational changes to the head as the 40S subunit transitions from

semi-closed (structure $7_{\Delta dII}$) to fully open (structure $9_{\Delta dII}$) states. Thus, structure $7_{\Delta dII}$ opens by 3.9°,

structure $8_{\Delta dII}$ by 8.1°, and structure $9_{\Delta dII}$ by 10.0° compared to the conformation of the 18S rRNA

in structures $1_{\Delta dII}$ -$6_{\Delta dII}$. These global changes to the position of the 40S head are reflected in changes

in the P site as the distance between U_{1248} and C_{1701} increases from 7.3 Å, to 9.9 Å and finally to

11 Å in structures $7_{\Delta dII}$, $8_{\Delta dII}$ and $9_{\Delta dII}$, respectively. Thus, this series of structures shows that even

in the absence of domain II, establishment of the full complement of IRES/40S contacts results in

the transition of the 40S subunit from the closed to the open state, which is required for

accommodation of the initiation codon and surrounding regions in the mRNA-binding channel.

Accommodation of the IRES in the mRNA-binding channel

In contrast to 40S/IRES binary complexes assembled on the ΔdII IRES, which showed

remarkable differences in the position of the 40S subunit head, binary complexes assembled on

the *wt* IRES yielded a uniform structure that was refined to 3.8 Å resolution from 119,320 particles

(structure 10_{wt}) (Figure 1B). Both the conformation of the 40S subunit and the structure of IRES

domains IIIa-f in structure 10_{wt} are identical to those in the open state of the 40S/IRES$_{\Delta dII}$ binary

complex (structure $9_{\Delta dII}$) (Figures 3A-B). There is, however, additional density that corresponds

to IRES domain II inserted into the 40S subunit E site, in a conformation that was observed in

purified IRES/80S complexes (Quade et al., 2015; Yamamoto et al., 2015). Superposition of

structure 10_{wt} and a closed-state 40S subunit shows that in the latter, steric clashes between uS7

and domain II would prevent insertion of this domain into the E site. Thus, domain II locks the

40S subunit into an open state, which is not similarly imposed in the case of the ΔdII IRES.

For the *wt* IRES, we also obtained the structure of the 40S•IRES•eIF1A ternary complex

that was refined to 3.8 Å resolution from 204,320 particles (structure 11_{wt}, Figure 1B). The

conformation of the 40S subunit and the position of the IRES in it were identical to those in the

40S/IRES$_{wt}$ binary complex (structure 10_{wt}) (Figures 3A-B). The complex clearly showed density

for eIF1A located between 18S rRNA helix (h) 44 and the ribosomal proteins eS30 and uS12,

allowing us to model the OB domain and the C-terminal subdomain of eIF1A (residues 22-122).

Although structure 11_{wt} lacks tRNA and the 40S subunit is in the open state, the position of eIF1A

on the ribosome as well as its overall conformation are identical to those in the structures of eIF2-

containing 48S complexes (Brito Querido et al.*,* 2020; Simonetti et al.*,* 2020).

As expected, binary complexes assembled on the ΔdII IRES and containing the 40S subunit

in the closed conformation ($1_{\Delta dII}$-$6_{\Delta dII}$) do not have mRNA in the mRNA-binding channel.

However, although structures $9_{\Delta dII}$, 10_{wt} and 11_{wt} all have 40S subunits in the identical open

conformation that is required for loading of mRNA into the channel, they differ strongly in the

degree of ribosomal accommodation of the initiation codon and surrounding regions (Figure 3C).

In the 40S/IRES$_{\Delta dII}$ binary complex (structure $9_{\Delta dII}$), clear mRNA density was seen only in the exit portion of the channel up to the -8 position of mRNA (HCV nt 334), after which the mRNA became disordered (Figure 3D). In contrast, in the *wt* 40S/IRES binary complex (structure 10_{wt}), mRNA nucleotides could be identified at the exit channel through the E site where it is stabilized by domain II, to AU$_{342-3}$ located in the P site (Figure 3E). The (+3) nucleotide linked to U$_{343}$ could not be identified due to disorder in the map. However, there is additional mRNA density in the mRNA channel ~8 Å from the P-site G$_{344}$, but the identity of these nucleotides does not correspond to those that immediately follow the start codon (A$_{345}$ onwards) as this density extends 20Å out from the mRNA channel beyond the binding site of eIF1A, suggesting that after the P site, the mRNA is looped out (Figure 3E). Thus, the presence of domain II results in insertion of mRNA into the entire channel, but only the additional 9 nucleotides from the exit to the P site (i.e., from G$_{335}$ to U$_{343}$) can be reliably identified. Density corresponding to mRNA is present from the P site and entry channel but is likely a mixture of different registers of mRNA and so the sequence could not be determined (Figure 3E). Strikingly, the presence of eIF1A in ribosomal complexes resulted in accommodation of sequential mRNA along the entire mRNA binding cleft (Figure 3F). Examination of critical P-site nucleotides for all complexes shows that in each case C$_{1701}$ of 18S rRNA is in a single conformation, except for the binary complex prepared with the *wt* IRES (structure 10_{wt}) where it is present in two states as determined by examination of the Coulomb potential around that nucleotide (Figure 3G). In the conformation of the second state, C$_{1701}$ contacts the upstream mRNA base U$_{343}$, possibly contributing to stabilizing the mRNA when it has not undergone complete accommodation in the mRNA channel at the P site. The highly conserved C$_{1698}$ (Prince et al., 1982) contacts downstream mRNA and may act as a sensor to stimulate C$_{1701}$ adopting the second conformation in which it can stabilize the incompletely loaded mRNA (Figure 3G).

The position of the mRNA in structures 10_{wt} and 11_{wt}, however, does not match its position in 48S complexes with an established codon-anticodon interaction (Brito Querido et al., 2020; Simonetti et al., 2020). Beginning from the (-1) position, the following (+1), and (+2) nucleotides are ~4 Å above their position when tRNA is inserted (Figures 3H-I). However, the raised position of these nucleotides in the complex containing eIF1A nevertheless allows the (+4) adenine base to form the stacking triple with eIF1A Trp70 and 18S rRNA A$_{1825}$ (Figure 3I), which is a key function

of eIF1A (Battiste et al., 2000). These contacts are maintained up to formation of 48S complexes

(Simonetti et al., 2020).

Taken together, the structures of binary complexes assembled on ΔdII and *wt* IRESs and

the eIF1A-containing complex assembled on the *wt* IRES provide structural rationalization for the

roles of domain II and eIF1A in sequentially loading the mRNA channel. Even without domain II,

binding of the IRES induces conformational changes in the 40S subunit that are required for

accommodation of mRNA in the binding channel. However, accommodation in this case is only

partial, and the 40S subunit is not stably present in the open conformation, which is consistent, on

one hand, with the ability of ΔdII IRES to function in initiation, but on the other hand, with lower

initiation activity than the *wt* IRES (e.g., Reynolds et al., 1996). The presence of domain II results

in accommodation of the initiation codon and the upstream region in the mRNA-binding channel,

thereby enhancing the efficiency of initiation (Reynolds et al., 1996), whereas addition of eIF1A

results in further accommodation of the mRNA in the entire mRNA-binding channel, accounting

for its enhancement of initiation on the IRES (Jaafar et al., 2016).

The structure of eIF2-containing 48S complexes assembled on the HCV IRES

The structures of eIF2-containing 48S ICs assembled on the *wt* and ΔdII IRES were refined

from 46,904 and 103,813 particles to resolutions of 3.6 and 3.5 Å, respectively (structures 12_{wt}

and $12_{\Delta dII}$, Figures 1B and S1). Both *wt* and ΔdII IRES 48S ICs form identical complexes with

respect to the 40S subunit's closed conformation, the positions of Met-tRNA$_i^{Met}$, eIF2 and eIF1A,

and the established P-site codon-anticodon base pairing (Figures 4A, and S5A-B). The IRES-

containing 48S IC is also structurally identical to the canonical 48S complex formed by cap-

dependent initiation (e.g., Simonetti et al., 2020) with respect to the global conformation of the

40S subunit as well as the placement of Met-tRNA$_i^{Met}$, eIF2α, and eIF1A. Thus, the E site-

associated eIF2α contacts the highly conserved uS7/Asp194 directly and interacts with Met-

tRNA$_i^{Met}$ via Thr103, Arg67 and His114 (Figure 4B). As with yeast 48S ICs (Hussain et al., 2014),

eIF2α forms hydrogen bonds with the (-3) adenosine via Arg55, a contact that enhances codon

selection in the scanning mode of initiation, presumably by stabilizing the arrested 48S IC (Pisarev

et al., 2006; Thakur et al., 2020). Contacts between eIF2α Arg57 and the tRNA acceptor stem loop

(ASL) and the 18S rRNA that occur in canonical 48S ICs are also present (Simonetti et al., 2020)

(Figure 4B). The position of eIF2α is identical in *wt* and ΔdII IRES 48S ICs (Figure S5E). The

unsharpened map also contained density corresponding to eIF2γ (Figure S5D) identical to that seen

in cap-dependent 48S initiation complexes (Simonetti et al., 2020), but it had a low local resolution

at the acceptor end of Met-tRNA$_i^{Met}$ and was not modelled after map sharpening.

The position of domain II differs substantially between the 48S IC assembled on the *wt*

IRES (structure 12_{wt}) and the corresponding 40S/IRES binary and 40S/eIF1A/IRES ternary

complexes (structures 10_{wt} and 11_{wt}). In both the binary and ternary complexes, it is inserted into

the E site, a position that is incompatible with the binding of eIF2. Thus, in the 48S IC, domain II

is oriented away from the subunit interface, towards the solvent side of the 40S subunit, and shows

an attenuation of density so that a model of domain II is not fully enclosed by the map. Focused

classification of this region revealed that domain II is flexible and occupies multiple conformations

oriented away from the E site (Figure 4C; Figure S5F; Table S5).

Compared with the binary or ternary complexes (structures 10_{wt} and 11_{wt}), the head of 48S

ICs formed on both *wt* and ΔdII IRESs is in a closed position, having moved by 11.2° relative to

the open states (Figure 4D), and is in an even more closed conformation than in closed binary

complexes (structures $1_{\Delta dII}$-$6_{\Delta dII}$), in which the position of the head differed from that in structures

10_{wt} and 11_{wt} by ~9.0° (Figure 3A).

Comparison between the ternary complex (structure 11_{wt}) and the 48S IC (structure

$12_{wt/\Delta dII}$) shows the effect that incorporation of tRNA into the P site and rearrangement of 18S

rRNA into the closed conformation has on the position of mRNA in the mRNA-binding channel.

The raised position of P-site mRNA in the ternary complex (Figure 4E) cannot be maintained as

this would cause a clash with the P-site tRNA in the 48S IC (Figure 4F). Thus, upon 40S subunit

closure and incorporation of P-site tRNA, the mRNA repositions ~4 Å deeper into the mRNA

channel and the A(+4) base flips out to maintain the stacking triple with eIF1A Trp70 and 18S

rRNA A$_{1825}$ that is seen in the ternary complex (Figures 4E-F). The mRNA is also shifted upstream

due to the movement of eS28 in the ribosome head (Figure 4G). A network of hydrogen bonds in

the ternary complex (structure 11_{wt}) between multiple arginines in eS28 and the mRNA is reformed

by the displacement of eS28 in the closed complex (structure $12_{wt/\Delta dII}$). Arg69 contacts the guanine

at (-7) position in the ternary complex, but the shift of the 18S rRNA to the closed position causes

this leading contact to be broken and reformed with G(-5). Other contacts are similarly

reorganized, and the Arg31 contact with the stacked G$_{331}$-G$_{307}$ pair is lost completely (Figure 4G-

H). This reorganization of contacts causes the mRNA to shift by one base pair at the P site and

allows contact with the tRNA anticodon arm once tRNA is inserted into the P site. Interestingly,

eIF2α Arg55 contacting the mRNA at A(-3) is in the same position as C_{83} in IRES domain II in

the 40S/eIF1A/IRES ternary complex, which could indicate that this location in the E site is

important for stabilizing the mRNA regardless of the conformation of the 40S subunit and

differences in position of mRNA at other locations along the mRNA-binding channel (Figure 4I).

378   Interestingly, we also obtained 40S ribosomal complexes containing platform-bound *wt* or

ΔdII IRES and P-site Met-tRNA$_i^{Met}$ but lacking all initiation factors (structures 13_{wt} and $13_{ΔdII}$)

(Figure 1C). These closed complexes, refined from 15,906 and 15,598 particles to 4.6 and 4.4 Å,

respectively, clearly showed density for codon-anticodon base pairing (Figure S3G; Table S3). It

is unclear whether initiation factors dissociated due to slow eIF5-independent hydrolysis of eIF2-

bound GTP (Unbehaun et al., 2004), or through denaturation at the air-water interface and/or due

to the shear forces associated with blotting (Glaeser, 2021; d'Imprima et al., 2019). In any case,

this complex likely mimics an intermediate state immediately after eIF2 dissociation and prior to

the binding of eIF5B.

The structure of eIF5B-containing 48S complexes assembled on the IRES

389   Classification of eIF5B-containing ribosomal complexes yielded two structures formed on

the *wt* IRES (structures 14_{wt} and 15_{wt}) and one structure ($15_{ΔdII}$) assembled on the ΔdII IRES

(Figures 1B and S1). Structures 14_{wt}, 15_{wt} and $15_{ΔdII}$ were refined to 3.8 Å, 3.7 Å and 3.7 Å

resolution from 60,578, 133,782 and 61,648 particles, respectively (Figures 5A-B and S6A). All

structures contain eIF5B, eIF1A and the P-site Met-tRNA$_i^{Met}$. However, whereas structures 15_{wt}

and $15_{ΔdII}$ showed density for the base-paired codon-anticodon and were accordingly classified as

48S ICs, the P-site Met-tRNA$_i^{Met}$ was not base-paired with the initiation codon in structure 14_{wt},

which was therefore designated as a pre-48S IC (Figure S3G). Similar to eIF2-containing 48S ICs

assembled on the *wt* IRES, in eIF5B-containing ICs assembled on the *wt* IRES (structure 15_{wt})

domain II is oriented away from the subunit interface, toward the solvent side of the 40S subunit,

and shows an attenuation of density that can be resolved by focused classification into multiple

states (Figure 1B and S6B; Table S5). In contrast to structures $15_{wt/ΔdII}$ that are in an identical

closed conformation, the 40S subunit in structure 14_w is in an open conformation with domain II

inserted into the E site (Figure S6C). In all structures, eIF1A occupies its usual position over h44

and eS30 and uS12, whereas eIF5B resides on the intersubunit face of the 40S subunit as in 80S

ribosomal complexes (Yamamoto et al., 2014; Huang and Fernandez, 2020; Wang et al., 2020).

The structures show clear density for eIF5B residues 592-1218, corresponding to all major

domains. The G domain, and domains II and III form the central domains of eIF5B that connect

via domain III and helix 12 to the tRNA acceptor stem binding domain IV. (Figure 5C). This is

the first high-resolution structure of mammalian eIF5B and also the first structure of the 40S/eIF5B

subunit complex prior to subunit joining. Although we used full-length native eIF5B, extensive

3D classification and processing did not reveal any structure corresponding to the N-terminal

region, suggesting that the structure in this region is highly disordered, an observation that is

supported by the structure of full-length eIF5B from a range of species predicted using AlphaFold

(Jumper et al., 2021). Consistent with the requirement of the 60S subunit for induction of eIF5B's

GTPase activity (Pestova et al., 2000), all complexes contained eIF5B-bound GTP (Figure S6E).

Previous structures of mammalian eIF5B determined at ~9 Å resolution used GMPPNP and

showed the GDP-bound conformation of switch 1 (Yamamoto et al., 2014) (Figure S7A). In

contrast, all our complexes showed switch 1, switch 2, and the $\beta 9$-$\beta 10$ loop in domain II in the

GTP-bound conformation (Kuhle and Ficner, 2014). The conformation of this conserved GTP-

binding region of mammalian eIF5B is identical to that in fungal eIF5B (Kuhle and Ficner, 2014;

Wang et al., 2020) (Figure S7B-D).

We identified multiple contacts between eIF5B domains II and III and 18S rRNA h5, h14,

and h15, as well as an interaction between eIF5B domain II and uS12 (Table S6; Figure S6D), and

noted that the $\beta 13$-$\beta 14$ loop in our complexes could interfere with the transition of switch 1 from

GTP-bound to GDP-bound conformations (Kuhle and Ficner, 2014) (Figure S6F). On the yeast

80S ribosome, the $\beta 13$-$\beta 14$ loop of eIF5B contacts A_{415} in 18S rRNA h14 and is positioned away

from the path that switch 1 might take as it changes to the GDP-bound conformation (Wang et al.,

2020). Although there is no contact between the equivalent nucleotide (A_{464}) and $\beta 13$-$\beta 14$ in our

complexes, h14 is accessible. These observations suggest that the 60S subunit might stimulate the

interaction between A_{415} and the $\beta 13$-$\beta 14$ loop to reposition this loop away from switch 1 so that

a transition from GTP- to GDP-bound conformation. In both pre-48S ICs and 48S ICs, the position

of domain IV also allows it to contact eIF1A via interactions between His112 in the helical

subdomain of eIF1A (Battiste et al., 2000) and the extreme C-terminal region of eIF5B, as well as

between the eIF1A L23 beta-turn (near Gly54) and Arg1186 in domain IV of eIF5B (Figure 5D).

These interactions are distinct from the previously reported binding of eIF1A's extreme C-terminal

DDIDI sequence to the h12/h13 loop in eIF5B domain IV (Marintchev et al., 2003; Zheng et al.,

2014) and of the eIF1A L45 loop to eIF5B domain III (Nag et al., 2016).

Comparison of the position of the P-site Met-tRNA$_i^{Met}$ in the eIF2-containing 48S IC

(Simonetti et al., 2020) and in the 80S ribosome (Yamamoto et al., 2014; Wang et al., 2020) shows

when the large subunit is present, tRNA rotates 14º towards the 40S subunit body and the T-loop

moves by ~ 15 Å to allow placement of the acceptor stem into the P site of the 60S subunit. We

therefore examined our maps (structures 12$_{wt/\Delta dII}$-15$_{wt/\Delta dII}$) to determine how the orientation of the

P-site tRNA differs in 40S ribosomal complexes depending on the presence of eIF2 or eIF5B.

Compared to eIF2-containing 48S ICs, in eIF5B-containing complexes, Met-tRNA$_i^{Met}$ rotates by

~14° and moves by 15 Å from the head of the 40S subunit to a position that matches the orientation

seen in 80S structures (Wang et al., 2020) (Figure 5C). This repositioning of tRNA was observed

for all structures that contained eIF5B, regardless of whether the 40S subunit was in the open (pre-

48S IC) or closed (48S IC) conformation (Figure 5C), indicating that eIF5B re-orients Met-

tRNA$_i^{Met}$ on the 40S subunit at the 48S PIC stage prior to subunit joining. Interestingly, although

the overall position of eIF5B-Met-tRNA$_i^{Met}$ in eIF5B-containing 48S ICs and in 80S ribosomes is

similar, upon binding of a 60S subunit, domain IV of eIF5B undergoes a 33° rotation towards the

platform side of the 40S subunit as well as a translation by 6.4 Å parallel to the mRNA channel

towards the platform of the 40S subunit, which results in repositioning of the tRNA acceptor stem

by 6.4 Å toward the ribosome head (i.e., toward ribosomal protein eS25) without changing the

position of the ASL (Figure 5E-F). The repositioning of eIF5B domain IV occurs despite an only

minor movement of helix 12 (Figure 5F; Figure S6G). Such repositioning of domain IV and the

tRNA acceptor stem upon binding of a 60S subunit would avert steric clashes of domain IV with

H84 of 28S rRNA and between uL16 and the tRNA acceptor stem (~A74) (Figure 5F). Similar

repositioning of IF2 domain C2 (equivalent to eIF5B domain IV) to avoid analogous steric clashes

was observed in bacteria upon joining of 50S subunits to 30S ICs (Hussain et al., 2016).

Adjustment of the orientation of initiator tRNA prior to subunit joining is a critical step in

initiation, and the mechanism by which it is mediated, by rotation and translation of domain IV of

the universally conserved initiation factor IF2/eIF5B, likely appeared early in evolution.

Despite the differences in the orientation of the acceptor arm of the P-site tRNA in

ribosomal complexes containing eIF2 and eIF5B, as well as differences in the conformation of the

40S subunit in eIF2/eIF5B-containing 48S ICs and eIF5B-containing pre-48S ICs, the anticodon

loop in all these complexes is identically positioned (Figure 5D). eIF5B- and eIF2-containing 48S

ICs assembled on the IRES are also in the same closed conformation (Figures S6C), identical to

that in 48S ICs assembled on canonical cap-dependent mRNA (Simonetti et al., 2020). Thus, in

eIF5B- and eIF2-containing 48S ICs, 18S rRNA nucleotides C_{1701} and U_{1248} are separated by ~3.5

Å, and the contacts of the P-gate nucleotides G_{1639} and A_{1640} in the 18S rRNA and Arg146 in uS9

with the tRNA anticodon arm on the opposite side to the anticodon are also present in both 48S

ICs.

In contrast, the 40S subunit in the eIF5B-containing pre-48S IC is in the open

conformation, in which the separation between C_{1701} and U_{1248} is increased to ~11 Å, and there are

no contacts between the tRNA and the P gate or uS9 (Figure 5H). As with the 40S/eIF1A/IRES

ternary complex (structure 11_{wt}), the open configuration of the ribosome head present in the pre-

48S IC creates a network of hydrogen bonds between the mRNA and eS28 (Figure S7E) that causes

the mRNA to be shifted by one nucleotide relative to the closed configuration (Figure S3G). This

results in the final nucleotide of the start codon (G_{344}) being out of place in the P site and not

participating in any contacts with the tRNA anticodon. Instead, G_{344} is available to form a stacking

triple with eIF1A Trp70 and A_{1825} of 18S rRNA (Figure 5H). In the closed conformation of 48S

ICs, the hydrogen bonds with eS28 have been partially broken (Figure S7F) and so all three

nucleotides of the start codon are accommodated in the P site, and the A(+4) nucleotide is available

to form a stacking triple with eIF1A Trp70 and A_{1825} of 18S rRNA (Figure 5F and 5I).

Comparison of eIF5B-containing pre-48S initiation complexes with eIF2-containing scanning

43S complexes

In contrast to initiation on the HCV IRES, eIF5B cannot substitute for eIF2 in recruiting

Met-tRNA$_i^{Met}$ to the 40S subunit in the canonical scanning mechanism. Canonical initiation also

requires eIF1, which binds to the 40S subunit below the mRNA channel at the P site between h24

and the region connecting h44 to h45 and, in cooperation with eIF1A, induces the open

conformation of the 40S subunit (Passmore et al., 2007; Llácer et al., 2015), thereby promoting

ribosomal attachment to mRNA, scanning and initiation codon selection (e.g., Pestova and

Kolupaeva, 2002). We therefore compared the conformation of the 40S subunit and the positions

of mRNA and Met-tRNA$_i^{Met}$ in eIF5B-containing pre-48S complexes formed on the *wt* IRES

(structure 14_{wt}) and in canonical scanning eIF2-containing 43S complexes that also contained eIF1

(Brito Querido et al., 2020).

Even though the individual mRNA bases could not be identified in the scanning, eIF2-

containing 43S complex because the P site nucleotides were heterogenous (Brito Querido et al.,

2020), the positions of mRNA in this complex and in the eIF5B-containing pre-48S complex are

very similar (Figure 6A). Although codon-anticodon base-pairing is not established in either

complex, the anticodon loops of Met-tRNA$_i^{Met}$ in both are poised to contact mRNA and base-pair

with an initiation codon, which is in contrast to the P_{OUT} position of tRNA in fully open mRNA-

free 43S complexes (Kratzat et al., 2021), in which the anticodon loop is positioned too far away

to be able to interact with mRNA if it had bound in the P site (Figure S7G). However, whereas the

conformations of the anticodon in the P site in both complexes are similar, in the eIF5B-containing

complex, the T-arm and the acceptor stem are shifted toward the body of the 40S subunit by

bending of the anticodon loop region (Figure 6B). Consequently, the contacts between the ASL of

Met-tRNA$_i^{Met}$ and 18S rRNA nucleotides $GA_{1639-40}$ and the N-terminal region of uS9 in the head

of the 40S subunit, which are present in all eIF2-containing complexes, including mRNA-free and

scanning 43S complexes and 48S ICs (e.g., Hussain et al., 2014; Brito Querido et al., 2020;

Simonetti et al., 2020), do not exist in the eIF5B-containing pre-48S ICs and only form after codon-

anticodon base-pairing (Figure 6B). Thus, interaction with eIF2 allows tRNA to maintain the

contacts with the head of the 40S subunit in all conformations, from fully open to fully closed, and

the position of the anticodon loop that allows it to inspect mRNA in the scanning 43S complexes

is determined by the rotation of the head that is in the intermediate conformation compared to fully

open and fully closed states. In contrast, when interacting with eIF5B, tRNA establishes these

contacts only in the closed position of the head, whereas in the open pre-48S ICs tRNA is stabilized

by contacts between eIF5B domain IV and the acceptor stem, tRNA anticodon U_{35} and mRNA

A(+1), and a single 18S rRNA nucleotide (C_{1701}) that partially stacks with C_{33} (instead of the

stacking of both U_{1248}/C_{1701} as in the closed case) (Figure 6B). Thus, whereas some aspects of

eIF5B-containing pre-48S complexes are analogous to those of eIF2-containing scanning 43S

complexes, the overall orientation, and the specific interactions of tRNA in them differ.

In eIF5B pre-48S complexes, the open conformation of the 40S subunit is supported by the

insertion of IRES domain II into the E site, whereas in eIF2-containing complexes, the

conformation of the 40S subunit is determined by the binding of eIF1. We therefore analyzed

whether binding of eIF1 would be compatible with the structure of eIF5B-containing pre-48S ICs.

The position of eIF1 placed into such complexes suggests that it would clash with tRNA. Thus,

the repositioning of tRNA in eIF5B-containing complexes causes the AAC_{38-40} nucleotides of Met-

$tRNA_i^{Met}$ to move toward the 40S subunit body by ~3.0 Å, so that the binding of eIF1 as in the

43S scanning complex would create a clash between A_{36} of Met-$tRNA_i^{Met}$ and the β1-β2 loop

(Figure 6C). Accommodation of eIF1 would require either reorganization of this loop or

displacement of the P-site tRNA. Examination of human (Fletcher et al., 1999) and yeast (Reibarkh

et al., 2008) solution NMR structures of eIF1 did not identify any conformations of the β1-β2 loop

that would allow a clash with the anticodon stem of tRNA to be avoided in the eIF5B-containing

pre-48S and 48S ICs. Moreover, the eIF1 β3-β4 loop would also clash with tRNA nucleotides

GU_{23-24}, and a clash between Phe113 and tRNA nucleotide G_{25} is also possible (Figure 6C). These

observations suggest that even if eIF5B were able to bind Met-$tRNA_i^{Met}$ with high affinity and

recruit it to the 40S subunit efficiently, the structure of the resulting complexes would not be

compatible with binding of eIF1 and hence, with the scanning mechanism of initiation. On the

other hand, the stabilizing interaction of the acceptor arm of Met-$tRNA_i^{Met}$ with domain IV of

eIF5B in the closed 48S complexes following dissociation of eIF2•GDP and eIF1 would lock the

complex preventing leaky scanning from occurring.

The incompatibility of Met-$tRNA_i^{Met}$ and eIF1 on eIF5B-containing pre-48S complexes

likely explains why eIF1 disrupts 48S complexes prepared using *wt* but not the ΔdII variant of the

HCV-like CSFV IRES (Pestova et al., 2008). HCV domain II has the propensity to insert into the

E site (Quade et al., 2015; Yamamoto et al., 2015), locking the ribosome into the open

conformation, and the structurally related CSFV domain II likely behaves similarly. As outlined

above, the tRNA consequently loses contacts with uS9, with $GA_{1639-40}$ as well as with U_{1248}. If

eIF1 binds to this complex, then the insertion of its β1-β2 loop into the mRNA channel creates

steric hindrance between mRNA and tRNA in the P site, dislodging the tRNA.

CONCLUDING REMARKS

Here we present the most comprehensive structural overview of the HCV IRES-mediated

initiation pathway to date (Figure 7). The IRES initially binds to the 40S subunit through domains

IIIa/IIIc and then pivots onto its platform side where it establishes the complete set of contacts

(structures $1_{ΔdII}$-$6_{ΔdII}$). Once the canonical set of contacts are made, this induces the head of the

ribosome to open (structure $8_{ΔdII}$-$9_{ΔdII}$). Although head opening can occur in the absence of IRES

domain II, such complexes are nevertheless characterized by remarkable heterogeneity in the position of the 40S subunit head. In contrast, 40S/IRES binary complexes assembled on the *wt* IRES yield a uniform structure, in which the 40S subunit is in the open conformation, and domain II is inserted into the E site (structure 10_{wt}). Importantly, in the absence of domain II, mRNA density was clearly seen only in the exit portion of the channel up to the -8 position of mRNA (structure $9_{\Delta dII}$), whereas in the *wt* 40S/IRES binary complex, mRNA nucleotides could be identified at the exit channel through the E site where it is stabilized by domain II, to AU_{342-3} located in the P site (structure 10_{wt}), and eIF1A induces further accommodation of the mRNA in the entire mRNA-binding channel (structure 11_{wt}). Thus, these complexes provide structural insights into the functions of multiple IRES domains, including IIIa/IIIc in establishing the initial ribosome contacts, IIId in fixing the IRES to the 40S subunit and inducing ribosomal head opening, and II in imposing the open conformation and promoting fixation of mRNA in and upstream of the P site. Our analysis also revealed the role of eIF1A in completing mRNA accommodation.

Once the mRNA is loaded, initiation can proceed either along the canonical initiation pathway, in which eIF2 promotes attachment of Met-tRNA$_i^{Met}$ to form the 48S IC (structure $12_{wt/\Delta dII}$) and then dissociates after GTP hydrolysis (likely structure $13_{wt/\Delta dII}$) followed by binding of eIF5B (structure $15_{wt/\Delta dII}$), or by a shortcut route, in which Met-tRNA$_i^{Met}$ is loaded directly by eIF5B, first forming open pre-48S ICs without established codon-anticodon interaction (structure 14_{wt}) and then locking upon codon-anticodon base-pairing (structure 15_{wt}). The eIF2-containing 48S ICs assembled on the IRES are structurally identical to canonical 48S ICs with respect to the conformation of the 40S subunit and the positions of Met-tRNA$_i^{Met}$, eIF2 and eIF1A. The position of domain II of the IRES in 40S/IRES$_{wt}$ binary complexes is incompatible with binding of eIF2 and the closed conformation of the 40S subunit, and consequently, in 48S ICs, domain II is oriented away from the subunit interface, toward the solvent side, occupying multiple conformations oriented away from the E site. In eIF5B-containing pre-48S ICs, the 40S subunit is in the open conformation and domains II is inserted into the E site. Upon codon-anticodon base-pairing in eIF5B-containing 48S ICs, the 40S subunit adopts the closed conformation, and domain II becomes displaced from the E site and is oriented away from the subunit interface like in eIF2-containing 48S ICs. Importantly, compared to eIF2-containing 48S ICs, in eIF5B-containing pre-48S ICs and 48S ICs, Met-tRNA$_i^{Met}$ rotates by ~14° and moves 15 Å from the head of the 40S subunit to a position that matches the orientation in 80S ribosomes. Thus, our data show how

589   eIF5B repositions tRNA already on the 48S complex, thus preparing it for joining with the 60S

590   subunit.

Acknowledgments

This work was supported by NIH R01 GM55440 (to J.F.), R35 GM139453 (to J.F.), R35 GM122602 (to T.V.P.), and R01 AI123406 (to C.U.T.H.) Cryo-EM data were collected at the Columbia University Cryo-Electron Microscopy Center.

Author Contributions

Conceptualization, T.V.P., C.U.T.H., and J.F.; Investigation, Z.P.B., and I.A.; Validation Z.P.B., I.A., and S.D.; Writing – Original Draft, T.V.P. and Z.P.B.; Writing – Review and Editing, Z.P.B., I.A., S.D., C.U.T.H., T.V.P., and J.F.; Funding Acquisition, C.U.T.H., T.V.P., and J.F.; Supervision, C.U.T.H., T.V.P., and J.F.

Declaration of Interests

Authors declare no competing interests.

FIGURE LEGENDS

Figure 1. Overview of HCV IRES-mediated initiation complexes.

(**A**) Secondary structure of the HCV IRES.

(**B**) Segmented maps of labeled IRES ICs, showing the 40S subunit (yellow), IRES (red), eIF1A (blue), Met-tRNA$_i^{Met}$ (magenta), and initiation factors eIF2 or eIF5B (green). Complexes were assembled on the *wt* or ΔdII IRES. Complexes that share identical 40S subunit conformation and factor composition are enclosed by dashed lines.

(**C**) Contacts between the IRES and 40S subunit. Ribosomal proteins (red), 18S rRNA (yellow), and the AUG codon (blue) are marked for the labeled complexes. Complete pathway in Figure S4 and Table S4.

See also Figures S1-S4 and Tables S1-S4.

Figure 2. Initial events during binding of the HCV IRES to the 40S subunit.

(**A**) IRES models from structures $1_{\Delta dII}$-$6_{\Delta dII}$ aligned on the 40S subunit. IRES domains and ribosomal proteins eS1 (blue), eS27 (magenta) and 18S rRNA ES7 (yellow) are indicated (inset).

(**B**) Comparison between minimally bound (structure $1_{\Delta dII}$) and fully bound (structure $6_{\Delta dII}$) IRES complexes showing the displacement of labeled IRES domains.

(**C**) Formation of critical contacts between ES7 and the IRES requires reorganization of domains hIII$_1$, IIIe, and IIId. The arrows indicate displacement of IRES nucleotides A$_{136}$ (hIII$_1$), G$_{295}$/A$_{296}$ (IIIe) and GGG$_{266-268}$ (IIId) interacting with ES7 from the minimally bound (structure $1_{\Delta dII}$) to the fully bound state (structure $6_{\Delta dII}$) indicated with a prime (′).

(**D**) RMSD (Å) of 18S rRNA for complexes labeled as in (E) compared to the minimally bound state (structure $1_{\Delta dII}$), color-coded as in the inset key. The angle formed between helix 33 (h33′) and helix 28 (h28′) in structure $1_{\Delta dII}$ and helix 33 (h33″) in the labeled complex is also marked.

(**E**) Segmented maps for labeled complexes showing 40S subunit (yellow) and IRES (red) organized in a putative sequence showing the minimally bound state (structure $1_{\Delta dII}$), canonical IRES binding (structure $6_{\Delta dII}$) and the induction of opening of the head of the 40S subunit (structure $7_{\Delta dII}$-$9_{\Delta dII}$).

See also Figure S4 and Table S4.

Figure 3. IRES domain II is required for loading mRNA into the mRNA channel

(A) RMSD (Å) of 18S rRNA for complexes labeled as in (B) compared to the minimally bound

state (structure $1_{\Delta dII}$), color-coded as in the inset key. The angle formed between helix 33 (h33′)

and helix 28 (h28′) in structure $1_{\Delta dII}$ and helix 33 (h33″) in the labeled complex is also marked.

(B) Segmented maps for labeled complexes showing 40S subunit (yellow), IRES (red), and eIF1A

(blue) organized in a putative sequence showing the minimally bound state (structure $1_{\Delta dII}$), fully

opened head of the 40S subunit (structure $9_{\Delta dII}$), binary complex (structure 10_{wt}), and the eIF1A-

containing ternary complex (structure 11_{wt}).

(C) Diagram of structures $1_{\Delta dII}$, $9_{\Delta dII}$, 10_{wt} and 11_{wt} showing the sequential loading of mRNA into

the mRNA channel.

(D-F) The mRNA channel spanning the entry and exit channels, A, P, and E sites viewed through

the ribosome head towards the body for the labeled complexes. The IRES (red), Coulomb potential

(blue mesh), and eIF1A (blue) are shown.

(G) Position of key nucleotides in the P site in structure 10_{wt} showing interactions between

incompletely loaded mRNA and C_{1698} resulting in C_{1701} sampling dual conformations near the (+2)

position.

(H-I) P site showing mRNA interactions for the (H) binary complex (structure 10_{wt}) and the (I)

eIF1A-containing ternary complex (structure 11_{wt}). 18S rRNA (yellow), eIF1A (blue), IRES

mRNA (red) with start codon (cyan) are all marked. The position of mRNA modelled in the 48S

cap-dependent IC (PDB: 6ZMW) is shown in grey.

Figure 4. The HCV IRES•eIF2-containing 48S initiation complex.

(A) Overview of the *wt* IRES eIF2-containing 48S IC (structure 12_{wt}).

(B) Contacts between eIF2α and HCV IRES, Met-tRNA$_i^{Met}$, and 40S subunit.

(C) IRES domain II occupies multiple positions in the eIF2-containing 48S IC.

(D) Position of 18S rRNA in the open ternary complex (structure 11_{wt}) and eIF2-containing 48S

IC (structure 12_{wt}).

(E-F) The P site in the (E) ternary complex (structure 11_{wt}) and the (F) eIF2-containing 48S IC

(structure 12_{wt}) showing 18S rRNA (yellow), Met-tRNA$_i^{Met}$ (magenta), eIF1A (blue), and IRES

mRNA (red) with the start codon (cyan) marked. **(E)** The position of mRNA in the cap-dependent

48S IC is marked (grey).

(G-H) Contacts between eS28 (salmon) and the IRES (red) in (G) the ternary complex (structure

11_{wt}) and (H) the eIF2-containing 48S IC (structure 12_{wt}).

(I) Global conformation of mRNA in the ternary complex (structure 11_{wt}) and the eIF2-containing

48S IC (structure 12_{wt}). The arrow indicates the extent of movement of eS28 between the binary

complex (structures 11_{wt}) and the eIF2-containing 48S IC complex (structure 12_{wt}).

See also Figure S5 and Table S5.

Figure 5. The HCV IRES•eIF5B-containing 48S initiation complex

(A) Overview of the *wt* IRES eIF5B-containing pre-48S IC (structure 14_{wt}).

(B) Overview of the *wt* IRES eIF5B-containing 48S IC (structure 15_{wt}).

(C) Changes to the position of P site tRNA depending on the presence of either eIF2 or eIF5B.

Upon binding of eIF5B, the T-arm, D-arm and acceptor stem of Met-tRNA$_i^{Met}$ move by 15.6 Å

and 14^o relative to their positions in the eIF2-containing 48S complex.

(D) Inset showing the conformation of tRNA acceptor stem loop, and contacts between eIF5B

domain IV and eIF1A. The conformation of the ASL is unchanged between the pre-48S IC

(eIF5B), and the 48S IC (eIF2 or eIF5B).

(E) Position of eIF5B and tRNA in the 48S and pre-elongation 80S ribosome complexes. eIF5B

undergoes relatively little movement between the 48S (green) and 80S stages (red) in the II, G,

and III domains, but domain IV translates and rotates (see F) causing movement in the tRNA

between the two complexes (magenta and orange respectively). Arrows show displacement for

labeled domains or components between the eIF5B-containing 48S IC and pre-elongation 80S

ribosome.

(F) The position of eIF5B domain IV in the 48S IC (green) would clash with 28S rRNA H89 and

uL16 in the 60S subunit. Upon binding of the 60S subunit, eIF5B domain IV (red) translates by

6.4Å towards to the platform side of the 40S subunit and rotates by 30.9° causing the tRNA to

rotate by 12.4° and translate by 6.4Å towards the head.

(G-I) P site for (G) eIF2-containing 48S IC (structure 12_{wt}), (H) eIF5B-containing pre-48S IC

(structure 14_{wt}), and (I) eIF5B-containing 48S IC (structure 15_{wt}) showing 18S rRNA (yellow),

Met-tRNA$_i^{Met}$ (magenta), eIF1A (blue), and IRES mRNA (red) with the start codon (cyan) marked.

See also Figures S6-S7 and Tables S5-S6.

Figure 6. Position of mRNA in the ribosomal P-site and codon-anticodon recognition.

(A) Position of tRNA in the P site for the P_{OUT} complex (PDB: 7A09) (blue), 43S scanning complex (PDB: 6ZMW) (yellow), and the pre-48S IC (structure 14_{wt}) (magenta). mRNA from the 43S scanning complex (green) and pre-48S IC (red) is shown.

(B) Contacts between tRNA, the initiation codon and the P site for the labeled complexes.

(C) The β1-β2 and β3-β4 loops of eIF1 bound as in the 43S scanning complex (PDB: 6ZMW) would clash with the pre-48S IC position of Met-tRNA$_i^{Met}$.

See also Figure S7.

Figure 7. The HCV IRES-mediated initiation pathway.

Ribosomal complexes organized in a putative IRES-mediated initiation pathway. Maps are segmented showing the 40S subunit (yellow), IRES (red), eIF1A (blue), Met-tRNA$_i^{Met}$ (magenta), and eIF2 or eIF5B (green).

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Classification of HCV IRES containing ribosomal complexes, related to Figure 1.

(A-D) Example micrographs for (A) the *wt* eIF2-containing sample, (B) the *wt* IRES eIF5B-

containing sample, (C) the ΔdII IRES eIF2-containing sample, and (D) the ΔdII IRES eIF5B-

containing sample. Further details in Table S2. Scale bar 2000 Å.

(E) Example 2D classification results for the *wt* IRES eIF2-containing sample.

(F-G) Classification scheme for (F) *wt* IRES and (G) ΔdII IRES containing samples. For full details

regarding classification see methods. Final maps are colored showing the 40S subunit (yellow),

IRES (red), eIF1A (blue), Met-tRNA$_i^{Met}$ (magenta), and initiation factors eIF2 or eIF5B (green).

Full details for complexes are available in Table S3 (structures $1_{\Delta dII}$-$15_{wt/\Delta dII}$), Table S5 (16_{wt}-28_{wt}),

and Table S7 (structures 29_{wt}-$46_{\Delta dII}$).

Figure S2. Local resolution, related to Figure 1.

(A) Local resolution of labeled HCV IRES complexes.

(B) Central slice of each complex.

Local resolution values and filtered maps can be found under EMDB ascension codes for each

structure (see Table S3).

Figure S3. Example densities, related to Figure 1.

(**A**) Domain IIId (red) and ES7 (yellow) for the pre-48S IC (structure 14_{wt}).

(**B**) Domain IIId (red) and ES7 (yellow) for the minimally bound complex (structure $1_{\Delta dII}$)

(**C**) Contact between domain IIIa/IIIc (red) and riboprotein eS27 (magenta) in the ternary complex

(structure 12_{wt}).

(**D**) eIF2 alpha helix 1 in the *wt* IRES 48S IC (structure 12_{wt}), top, and the ΔdII IRES 48S IC

(structure $12_{\Delta dII}$), bottom.

(**E**) eIF5B alpha helix 12 in the eIF5B-containing pre-48S IC (structure 14_{wt}), top, and the 48S IC

(structure 15_{wt}), bottom.

(**F**) eIF1A example densities from the eIF5B-containing pre-48S IC (structure 14_{wt}), left, and the

eIF2-containing 48S IC (structure 12_{wt}), right.

(G) Initiation codon (red) and tRNA anticodon (magenta) for all complexes where mRNA is

present in the P site. Diagram shows identity of nucleotides and hydrogen bonds for tRNA and

initiation codon bases in each accompanying figure.

Figure S4. Interactions between HCV IRES and 40S ribosomal subunit for all complexes,

related to Figures 1 and 2.

Contacts between ribosomal proteins (red), 18S rRNA (yellow), and the AUG codon (blue) are

marked. Labels refer to ribosomal proteins and elements of 18S rRNA, and to domains of the

IRES. Complexes assembled using the ΔdII IRES show the deleted domain marked in red. Also

see Table S4 for more details.

Figure S5. eIF2 related supplement, related to Figure 4.

(A) Overview of the ΔdII IRES eIF2-containing 48S initiation complex (structure $12_{\Delta dII}$).

(B-C) Presence of Domain II in the E site causes the head of the 40S subunit to open. (B) Position

of 18S rRNA in the open ternary complex (structure 11_{wt}) (orange), *wt* IRES eIF2-containing 48S

IC (structure 12_{wt}) (yellow), and ΔdII IRES eIF2-containing 48S IC (structure $12_{\Delta dII}$) (magenta).

(C) Position of 18S rRNA in the open ternary complex (structure 11_{wt}) (orange), *wt* IRES w/o eIF2

48S IC (structure 13_{wt}) (yellow), and ΔdII IRES w/o eIF2 48S IC (structure $13_{\Delta dII}$) (magenta).

Measured angle between 18S rRNA h33 and h28 in the ternary complex (denoted with ′) to h33 in

the *wt* IRES complex (denoted with ″) shown for (C-D).

(D) Unsharpened maps for the *wt* IRES eIF2-containing 48S IC (structure 12_{wt}), left, and the ΔdII

IRES eIF2-containing 48S IC (structure $12_{\Delta dII}$), right. Comparison to canonical cap-dependent

initiation complexes (PDB: 6YAL) identifies the presence of eIF2γ (cyan).

(E) The position and conformation of mRNA (magenta) and the eIF2 α-subunit for the *wt* IRES

eIF2-containing 48S IC (structure 12_{wt}, green) and the ΔdII IRES eIF2-containing 48S IC

(structure $12_{\Delta dII}$ (orange) is identical.

(F) Maps showing that domain II occupies multiple positions in the *wt* IRES eIF2-containing 48S

IC.

(G-J) Diagram of P site for the (G) IRES binary complex (structure $6_{\Delta dII}$), **(H)** P site for pre-48S

IC (structure 14_{wt}), **(I)** P site for eIF2-containing 48S IC (structure 12_{wt}), and **(J)** P site for eIF5B-

containing 48S IC (structure 15$_{wt}$). Ribosomal proteins and 18S rRNA (yellow), mRNA (red),

tRNA (white), and eIF2 (green) are shown.

Figure S6, related to Figure 5.

(A) Overview of the ΔdII IRES eIF5B-containing 48S initiation complex (structure 15$_{ΔdII}$).

(B) Maps showing that domain II occupies multiple positions in the *wt* IRES eIF5B-containing

48S IC.

(C) Position of 18S rRNA in the *wt* IRES pre-48S IC (structure 14$_{wt}$) (orange), *wt* IRES eIF5B-

containing 48S IC (structure 15$_{wt}$) (magenta), and ΔdII IRES eIF5B-containing 48S IC (structure

15$_{ΔdII}$) (yellow). Measured angle between 18S rRNA h33 and h28 in the eIF5B-containing 48S IC

(denoted with ′) to h33 in the pre-48S IC (denoted with ″).

(D) Global position of eIF5B bound to the intersubunit face of the 40S ribosomal subunit.

Interactions between 18S rRNA (yellow), uS12 (salmon), and eIF1A (blue) are shown.

(E) Coulomb potential for the GTP nucleotide in eIF5B G domain for pre-48S IC (structure 14$_{wt}$),

left, and 48S IC (structure 15$_{wt}$), right.

(F) Comparison of conformation of switch 1 in GTP (orange) and GDP (blue), and switch 2 in

GTP (magenta) and GDP (yellow) conformations from the *C. thermophilum* G domain (grey)

(PDB: 4NCN). Conformational changes in domain II seen in the *O. cuniculus* 48S IC state (green)

and the *S. cerevisiae* pre-elongation 80S state (red) (PDB: 6WOO).

(G) Domain alignment between *O. cuniculus* pre-48S IC (structure 14$_{wt}$)/48S IC (structure 15$_{wt}$,

green) and *S. cerevisiae* eIF5B from the pre-elongation 80S (red) (PDB: 6WOO). Position of G

domain switch 1 (magenta) and switch 2 (orange) are marked

Figure S7, related to Figures 5 and 6.

(A) Structure of the *O. cuniculus* eIF5B G domain (PDB: 4UJD) bound to the pre-elongation 80S

ribosome (yellow) compared to the *O. cuniculus* pre-48S IC (structure 14$_{wt}$, green).

(B) Structure of the *O. cuniculus* eIF5B G domain in the pre-48S IC (structure 14$_{wt}$) (green).

(C) Structure of the *C. thermophilum* eIF5B G domain (PDB: 4NCN) (blue) compared to the *O.*

cuniculus eIF5B G domain bound to the pre-48S IC (structure 14$_{wt}$) (green).

(D) Structure of the *S. cerevisiae* eIF5B G domain (PDB: 6WOO) bound to the pre-elongation 80S

ribosome (red) compared to the *O. cuniculus* pre-48S IC (structure 14$_{wt}$) (green).

(E-F) Contacts between eS28 (salmon) and the HCV IRES (red) in the (E) pre-48S IC (structure

14_{wt}) and (F) 48S IC (structure 15_{wt}).

(G) Distance between P_{OUT} tRNA (PDB: 7A09) and the initiation codon in ternary complex

(structure 11_{wt})

(H) Contacts between tRNA and the P site for the eIF2-containing 48S IC.

Table S1. **Sample composition, related to Figure 1.**

Table S2. Data collection statistics, related to Figure 1.

Table S3. Refinement and validation statistics, related to Figure 1.

Table S4. Contacts between the HCV IRES and 40S subunit, related to Figure 2. Distances

below 3.5 Å are marked as 'yes', between 3.5-4.5 Å are marked as 'weak', and above 4.5 Å

marked as 'no'.

Table S5. Maps showing HCV IRES domain II movement, related to Figures 4 and 5.

Table S6. Contacts between eIF5B and the 40S subunit, related to Figure 5.

Table S7. Additional maps produced during data processing, related to Figure 1 and S1.

STAR Methods

Resource availability

Materials availability

Requests for materials and additional information can be directed to Dr. Tatyana Pestova

(Tatyana.Pestova@downstate.edu;) or Dr. Joachim Frank (jf2192@cumc.columbia.edu).

Data availability

Primary models and maps (Table S3) reported in this study were deposited in the Protein

Data Bank (PDB) and Electron Microscopy Data Bank (EMDB) under the following accession

codes: structure $1_{\Delta dII}$ (PDB: 7SYG; EMD-25527), structure $2_{\Delta dII}$ (PDB: 7SYH; EMD-25528),

structure $3_{\Delta dII}$ (PDB: 7SYI; EMD-25529), structure $4_{\Delta dII}$ (PDB: 7SYJ; EMD-25530), structure $5_{\Delta dII}$

(PDB: 7SYK; EMD-25531), structure $6_{\Delta dII}$ (PDB: 7SYL; EMD-25532), structure $7_{\Delta dII}$ (PDB:

7SYM; EMD-25533), structure $8_{\Delta dII}$ (PDB: 7SYN; EMD-25534), structure $9_{\Delta dII}$ (PDB: 7SYO;

EMD-25535), structure 10_{wt} (PDB: 7SYP; EMD-25536), structure 11_{wt} (PDB: 7SYQ; EMD-

25537), structure 12_{wt} (PDB: 7SYR; EMD-25538), structure $12_{\Delta dII}$ (PDB: 7SYS; EMD-25539),

structure 13_{wt} (PDB: 7SYT; EMD-25540), structure $13_{\Delta dII}$ (PDB: 7SYU; EMD-25541), structure

14_{wt} (PDB: 7SYV; EMD-25542), structure 15_{wt} (PDB: 7SYW; EMD-25543), and structure $15_{\Delta dII}$

(PDB: 7SYX; EMD-25544). For each entry half maps, the mask used for post processing, Fourier

correlation curve, and local resolution maps have been deposited as additional files.

Additional maps (table S5) showing the movement of HCV IRES domain II were deposited

in the EMDB under the following accession codes: structure 16_{wt} (EMD-25545), structure 17_{wt}

(EMD-25546), structure 18_{wt} (EMD-25547), structure 19_{wt} (EMD-25548), structure 20_{wt} (EMD-

25549), and structure 21_{wt} (EMD-25550), structure 22_{wt} (EMD-25551), structure 23_{wt} (EMD-

25552), structure 24_{wt} (EMD-25553), structure 25_{wt} (EMD-25554), structure 26_{wt} (EMD-25555),

structure 27_{wt} (EMD-25556), and structure 28_{wt} (EMD-25557). For each entry half maps, the mask

used for post processing, and Fourier correlation curve have been deposited as additional files.

Maps obtained during data processing (Table S7) from particle stacks that were

compositionally and conformationally identical that were later combined were deposited in the

EMDB under the following accession codes: structure 29_{wt} (EMD-25588), structure 30_{wt} (EMD-

25589), structure $31_{\Delta dII}$ (EMD-25590), and structure $32_{\Delta dII}$ (EMD-25591). For each entry half

maps, the mask used for post processing, and Fourier correlation curve have been deposited as
additional files.

Consensus maps (Table S7) that were used for focused classification were deposited in the
EMDB under the following accession codes: structure $33_{\Delta dII}$ (EMD-25592), structure $34_{\Delta dII}$ (EMD-
25593), structure 35_{wt} (EMD-25594), structure 36_{wt} (EMD-25595), structure $37_{\Delta dII}$ (EMD-25596),
structure 38_{wt} (EMD-25597), structure 39_{wt} (EMD-25598), and structure $40_{\Delta dII}$ (EMD-25599).
Other high-resolution maps obtained during classification were deposited in the EMDB under the
following accession codes: structure 41_{wt} (EMD-25600), structure 42_{wt} (EMD-25601), structure
43_{wt} (EMD-25602), structure 44_{wt} (EMD-25603), structure 45_{wt} (EMD-25604), and structure $46_{\Delta dII}$
(EMD-25620). For each entry half maps, the mask used for post processing, mask used for focused
classification, and the Fourier correlation curve have been deposited as additional files.

Method details
Plasmids

Vectors for expression of His$_6$-tagged eIF1A (Pestova et al., 1998a) and *Escherichia coli*
methionyl tRNA synthetase (Lomakin et al., 2006) have been described. The plasmid HCV-
MSTN-Stop (Hashem et al., 2013) containing HCV Type 1 nt. 40–375 was used for transcription
of mRNA containing the *wt* HCV IRES. A derivative for transcription of HCV IRES lacking
domain II (containing HCV nt. 125-375) was made by GenScript Corp. (Piscataway, NJ). The
HCV plasmids were linearized by BamHI, and mRNAs were transcribed using T7 RNA
polymerase (Thermo Scientific).

Purification of factors, ribosomal subunits and aminoacylation of tRNA
Native mammalian 40S subunits, eIF2, eIF3 and eIF5B were purified from rabbit
reticulocyte lysate (RRL) (Green Hectares), as described (Pisarev et al., 2007). Recombinant
eIF1A and *Escherichia coli* methionyl tRNA synthetase were expressed and purified from *E.*
coli as described (Pisarev et al., 2007).
For purification of native rabbit total tRNA, 200 ml RRL were centrifuged at 45,000 rpm
for 4.5 h in a Beckman 50.2 Ti rotor at 4°C in order to pellet polysomes. The supernatant was
dialyzed overnight against buffer A (20 mM Tris [pH 7.5], 4 mM MgCl$_2$, 250 mM KCl, 2 mM
DTT) and applied to a DE52 (Whatman) column equilibrated with buffer A. The tRNA was eluted

with buffer B (20 mM Tris [pH 7.5], 3 mM $MgCl_2$, 700 mM NaCl, 2 mM DTT) and precipitated

overnight with 2.5 volumes of ethanol at -80°C. The precipitate was centrifuged at 13,000 rpm for

15 minutes and resuspended in 5 ml buffer C (100 mM Tris [pH 7.5], 5 mM $MgCl_2$), phenol-

chloroform (pH 4.7)) extracted and precipitated again with 0.3 M NaOAc and 2.5 volumes of

ethanol. To isolate tRNA, the pellet was dissolved and subjected to gel filtration on a Superdex 75

column (Pestova and Hellen, 2001). Purified total tRNA was aminoacylated using *E. coli*

methionyl tRNA synthetase (to obtain Met-tRNA$_i^{Met}$) as described (Pisarev et al., 2007).

Assembly of ribosomal complexes

To form 48S initiation complexes, 7 pmol HCV IRES mRNA (*wt* or Δdomain II mutant)

were incubated with 3.5 pmol 40S subunits, 10 pmol eIF1A, 4.5 pmol eIF3, total native rabbit

tRNA containing 3.5 pmol Met-tRNA$_i^{Met}$, and 6 pmol eIF2 or 10 pmol eIF5B in 40 μl buffer D

(20 mM Tris [pH 7.5], 100 mM KAc, 2.5 mM $MgCl_2$, 2 mM DTT, 0.25 mM spermidine, 1 mM

ATP and 0.2 mM GTP) for 10 minutes at 37°C. The obtained complexes (containing 87.5 nM 40S

subunits) were applied directly onto grids without dilution.

Grid preparation and electron microscopy

Gold foil grids were prepared from Quantifoil gold mesh grids (Passmore and Russo,

2016). Initially, Quantifoil R0.6/1.0 300 mesh gold grids (Quantifoil Micro Tools GmbH) were

visually inspected to check for uniformity and intactness of the Quantifoil layer and then placed

into an Auto 306 Turbo Vacuum Coater (Edwards Vacuum) at a pressure of 10^3 Pa and then gold

wire (Ted Pella, Inc) was evaporated for approximately 8 minutes to create a 500 Å layer.

Deposition thickness was determined using the inbuilt film thickness monitor. To remove the

underlying Quantifoil carbon foil layer the grids were then treated with plasma using a Gatan

Solarus 950 (Gatan Inc) operated at 25 W for 4 minutes with an argon/oxygen gas mixture.

To prepare hydrophilic grids, 30 minutes prior to sample application, grids were treated

with plasma using a Gatan Solarus 950 (Gatan Inc) operated at 25W for 25 seconds with a

hydrogen/oxygen gas mixture. These grids were then transferred to the environmental chamber of

a Vitrobot Mark IV (Thermo Fisher Scientific) maintained at 4°C and 100% humidity. Here 3 μL

of sample were applied and then blotted for 4 seconds with blot force 3 before immediate plunging

into an a cooled (77K) ethane-propane mixture (Tivol et al., 2008) and then transferred to liquid

nitrogen. Selected grids were screened to confirm sample composition and ice thickness using a

Tecnai F20 electron microscope (Thermo Fisher Scientific) equipped with a field emission gun

(FEG) operating at 200kV and a K2 summit direct electron detector (Gatan, Inc).

After screening grids from each sample, data collection was performed on a Tecnai Polara

F30 (Thermo Fisher Scientific) equipped with an FEG operating at 300 kV and a K3 direct electron

detector (Gatan, Inc). Movies were collected at a nominal magnification of 52,000× and defocus

range of -0.5 to -2.5 μm in counting mode with a pixel size of 0.95 Å/pixel using the automated

collection software Leginon (Potter et al., 1999; Carragher et al., 2000; Suloway et al., 2005). Each

movie consisted of 40 frames recorded over 4 seconds with a total dose of 70.9 e$^-$/Å^2. Due to

sample conditions the 40S ribosome may enter a preferred orientation and so potions of the data

were collected with a 30° stage tilt (Table S2). For the *wt* IRES eIF2-containing sample 14,815

micrographs (14,815 at 30° stage tilt) were collected over 2 sessions, for the *wt* IRES eIF5B-

containing sample 27,263 micrographs (20,509 at 30° stage tilt) were collected over 4 sessions,

for the ΔdII IRES eIF2-containing sample 22,735 micrographs (17,695 at 30° stage tilt) were

collected over 3 sessions, and for the ΔdII IRES eIF5B-containing sample 13,809 micrographs

(13,809 at 30° stage tilt) were collected over 2 sessions (Table S2). Optical groups of micrographs

with similar beam tilt values were identified using k-means clustering on the image shift values

recorded by the microscope during data collection.

1009

Image processing

Gain references for each session were produced by visually screening ~1000 micrographs

to remove images that contained gold foil and then summing them using cisTEM *sum_all_tifs*

(Grant et al., 2018). Movies were then aligned using MotionCor2 (Zheng et al., 2017) with dose

weighting of 1.77 e$^-$/Å^2/frame and local patch correction with 8×5 patches. Initial CTF parameters

were estimated using CTFFIND4 (Rohou and Grigorieff, 2015). Particle locations were identified

using Topaz version 0.2.3 (Bepler et al. 2019) by initially downscaling all micrographs by 8× and

then using the Topaz general model to identify particles. Particles with a confidence score below

0 were removed and the remaining positions rescaled for subsequent processing in Relion 3.1

(Scheres, 2012; 2016; Zivanov et al., 2018; 2019).

Initially, we identified 2,183,185 particles for the *wt* IRES eIF2-containing sample

(147 particles per micrograph), 1,133,335 particles for the *wt* IRES eIF5B-containing sample

(42 particles per micrograph), 2,213,826 particles for the ΔdII IRES eIF2-containing sample

(97 particles per micrograph), and 1,459,506 particles for the ΔdII IRES eIF5B-containing sample

(106 particles per micrograph). Particle locations were extracted from micrographs into

downsampled boxes of 100×100 pixels (at 3.8 Å/pixel) to speed initial classification. This

corresponds to 400×400-pixel boxes (at 0.95 Å/pixel) without downsampling. 25 iterations of 2D

classification were performed to identify incorrectly picked particles, contamination, and other

particles that were unable to be correctly aligned (e.g., due to poor SNR). Particles that were

selected for removal were subjected to additional 2D classification to confirm that they did not

contain clear 40S ribosome particles.

After initial screening of the 2D classification data the remaining particles for each sample

were 1,201,923 particles for the *wt* IRES eIF2-containing sample (81 particles per micrograph),

736,700 particles for the *wt* IRES eIF5B-containing sample (36 particles per micrograph),

1,484,658 particles for the ΔdII IRES eIF2-containing sample (84 particles per micrograph), and

1,119,610 particles for the ΔdII IRES eIF5B-containing sample (81 particles per micrograph). For

each sample, all particles were refined into a single model which was used to estimate the defocus

values on a per-particle basis, followed by an additional refinement step, and then 3D classification

without alignment into 10 classes for 25 iterations. This initial 3D classification was used to

identify the major conformational states present in each sample, as well as further removal of poor-

quality particles. For the *wt* IRES eIF2-containing sample we identified 580,938 particles in the

closed state, 176,793 particles in the open sate, and removed 444,192 particles. For the *wt* IRES

eIF5B-containing sample we identified 378,325 open state, 360,338 closed state, and removed

58,037 particles. For the ΔdII IRES eIF5B-containing sample we identified 883,893 particles in

the closed state, and removed 62,455 particles. For the ΔdII IRES eIF2-containing sample we

identified 615,125 particles in the closed state, 198,920 particles in the intermediate-open state,

and removed 670,433 particles. All particles that were selected for removal were subjected to

additional 2D classification and 3D refinement steps to confirm that they did not contain 40S

ribosome complexes.

1049

General processing pathway

All particles were re-extracted at full-size (400×400-pixel box, 0.95 Å/pixel) and

underwent iterations of 3D refinement, following by anisotropic magnification correction, defocus

refinement, and beam tilt estimation. Multiple rounds of 3D classification (25 iterations, without
alignment) were used to progressively remove poor quality particles. After each round of CTF
refinement each particle stack underwent 3D refinement and was checked for increase in resolution
and visual improvement of map density. Once CTF refinement no longer improved map quality,
particle polishing using all frames was performed and then iterations of CTF refinement as outline
above were completed. Focused classification on consensus maps was performed to isolate desired
conformational or compositional states (see below).
1060

_wt_ IRES eIF2-containing sample

After 3D classification the consensus map of closed 40S ribosome particles (Table S7)
from the _wt_ IRES eIF2-containing sample contained 530,720 particles at 3.1 Å resolution
(structure 36_{wt}; EMD-25595) underwent focused classification by enclosing Met-tRNA$_i^{Met}$ and the
eIF2α subunit in a mask. Iterations of focused classification were able to produce two high quality
classes of either the eIF2-containing 48S IC with 46,904 particles at 3.6 Å resolution (structure 12_{wt};
PDB: 7SYR; EMD-25538) and the 48S IC w/o eIF2 with 15,598 particles at 4.4 Å resolution
(structure 13_{wt}; PDB: 7SYT; EMD-25540). An additional class of the eIF2-containing 48S IC
lacking eIF1A with 45,571 particles at 3.6 Å resolution (structure 45_{wt}; EMD-25604) was
obtained. Extensive classification of the remaining 422,647 particles produced maps that showed
the 40S ribosome bound to eIF2 and/or Met-tRNA$_i^{Met}$ but with a low resolution, or the ribosome
lacking both factors.

To determine the positions that IRES domain II occupies the region around domain II was
masked and classified into 10 classes over 25 iterations without alignment. Six classes were able
to be clearly resolved showing IRES domain II sampling different positions around the ribosome
head. Class 1 containing 41,393 particles at 3.8 Å resolution (structure 16_{wt}; EMD-25545), class
2 containing 39,148 particles at 3.8 Å resolution (structure 17_{wt}; EMD-25546), class 3 containing
49,034 particles at 3.7 Å resolution (structure 18_{wt}; EMD-25547), class 4 containing 66,090
particles at 3.6 Å resolution (structure 19_{wt}; EMD-25548), class 5 containing 24,190 particles at
3.9 Å resolution (structure 20_{wt}; EMD-25549), and class 6 containing 36,155 particles at 3.8 Å
resolution (structure 21_{wt}; EMD-25550). Extensive classification of the remaining 274,710
particles could not further resolved additional positions of domain II.

After extracting at full size the consensus map of the open 40S ribosome particles (Table

S7) from the *wt* IRES eIF2-containing sample contained 176,793 particles at 3.8 Å resolution

(structure 29_{wt}; EMD-25588).

1086

***wt* IRES eIF5-containing sample**

The consensus map of closed 40S ribosome particles (Table S7) from the *wt* IRES eIF5B-

containing sample contained 360,338 particles at 3.4 Å resolution (structure 39_{wt}; EMD-25598)

underwent focused classification by enclosing eIF5B in a mask. This could be classified into three

high-resolution classes of the eIF5B-containing 48S IC eIF5B: one containing 133,782 particles

at 3.7 Å resolution (structure 15_{wt}; PDB: 7SYW; EMD-25543), one containing 109,025 particles

at 3.7 Å resolution (structure 41_{wt}; EMD-25600), and one containing 55,367 at 3.8 Å resolution

(structure 42_{wt}; EMD-25601). The remaining 62,164 particles showed the 48S IC at 3.9 Å

resolution but the occupancy of eIF5B was lower (structure 43_{wt}; EMD-25602).

To determine the positions that IRES domain II occupies the region around domain II was

masked and classified into 10 classes over 25 iterations without alignment. Seven classes were

able to be resolved showing IRES domain II sampling different positions around the ribosome

head. Class 1 containing 20,138 particles at 5.0 Å resolution (structure 22_{wt}; EMD-25551), class

2 containing 11,082 particles at 5.4 Å resolution (structure 23_{wt}; EMD-25552), class 3 containing

16,584 particles at 5.2 Å resolution (structure 24_{wt}; EMD-25553), class 4 containing 16,217

particles at 4.9 Å resolution (structure 25_{wt}; EMD-25554), class 5 containing 8,102 particles at 6.0

Å resolution (structure 26_{wt}; EMD-25555), class 6 containing 13,152 particles at 5.3 Å resolution

(structure 27_{wt}; EMD-25556), and class 7 containing 7,029 particles at 6.2 Å resolution (structure

28_{wt}; EMD-25557). Extensive classification of the remaining 268,034 particles could not further

resolved the position of domain II.

The consensus map of open 40S ribosome particles (Table S7) from the *wt* IRES eIF5B-

containing sample was classified into two classes: a consensus eIF5B-containing pre-48S IC with

199,047 particles at 3.6 Å (structure 38_{wt}; EMD-25597) and the open binary complex with 147,309

particles at 4.1 Å resolution (structure 30_{wt}; EMD-25589). The 31,969 remaining particles were

subjected to 3D classification but did now contain high-quality classes. The consensus map of the

eIF5B containing pre-48S IC (structure 38_{wt}) was masked around eIF5B and focused classification

produced two classes, one with 60,578 particles at 3.8 Å (structure 14_{wt}; PDB: 7SYV; EMD-

25542), and a lower-resolution map from 29,072 particles at 4.2 Å (structure 44_{wt}; EMD-25603).

1115

ΔdII IRES eIF2-containing sample

The consensus map of closed 40S ribosome particles (Table S7) from the ΔdII IRES eIF2-

containing sample contained 615,195 particles at 3.3 Å resolution (structure $37_{ΔdII}$; EMD-25596)

underwent focused classification by enclosing Met-tRNA$_i^{Met}$ and the eIF2α subunit in a mask.

Iterations of focused classification were able to produce two high quality classes of either the eIF2-

containg 48S IC (structure $12_{ΔdII}$; PDB:7SYS; EMD-25539), and the 48S IC w/o eIF2 (structure

$13_{ΔdII}$; PDB:7SYU; EMD-25541). An additional class of 147,713 particles at 3.6 Å (structure

$46_{ΔdII}$; EMD-25620) showed the 48S IC but with very poor density for eIF2. Extensive

classification of the remaining 348,071 particles did not produced any high-quality maps. The

consensus map of the intermediate conformation 40S ribosome particles (Table S7) from the ΔdII

IRES eIF2-containing sample had 198,920 particles at 3.5 Å resolution (structure $31_{ΔdII}$; EMD-

25590).

1128

ΔdII IRES eIF5B-containing sample

3D classification of the closed state eIF5B-containing 40S ribosome produced three classes

that were processed further (Figure S1G; Table S7). The consensus map of closed 40S ribosome

particles (Table S7) from the ΔdII IRES eIF5B-containing sample contained 148,763 particles at

3.4 Å resolution (structure $40_{ΔdII}$; EMD-25599) and underwent focused classification by enclosing

eIF5B in a mask. Iterations of focused classification were able to produce one high quality class

of the eIF5B-containg 48S IC (structure $15_{ΔdII}$; PDB: 7SYX; EMD-25544). Classification of the

remaining 99,239 particles produced maps at either low resolution or lacking eIF5B. The

consensus map showing the closed 40S ribosome with multiple HCV IRES states (Table S7) had

346,516 particles at 3.6 Å resolution (structure $33_{ΔdII}$; EMD-25592). A mask was prepared around

IRES domains S1/S2/IIIe/IIIf and focused classification resolved six classes of the IRES in various

association/dissociation states: (1) the early stage association between the IRES and 40S/IRES$_{ΔdII}$

binary complex with 42,271 particles at 4.3 Å resolution (structure $1_{ΔdII}$; PDB: 7SYG; EMD-

25527), (2) the early stage association between the IRES and 40S/IRES$_{ΔdII}$ binary complex with

28,684 particles at 4.6 Å resolution (structure $2_{ΔdII}$; PDB: 7SYH; EMD-25528), (3) the early stage

association between the IRES and 40S/IRES$_{\Delta dII}$ binary complex with 24,545 particles at 4.5 Å

resolution (structure $3_{\Delta dII}$; PDB: 7SYI; EMD-25529), (4) the early stage association between the

IRES and 40S/IRES$_{\Delta dII}$ binary complex with 27,043 particles at 4.8 Å resolution (structure $4_{\Delta dII}$;

PDB: 7SYJ; EMD-25530), (5) the early stage association between the IRES and 40S/IRES$_{\Delta dII}$

binary complex with 48,757 particles at 4.2 Å resolution (structure $5_{\Delta dII}$; PDB: 7SYK; EMD-

25531), and (6) the canonically bound IRES 40S/IRES$_{\Delta dII}$ binary complex with 29,657 particles at

4.5 Å resolution (structure $6_{\Delta dII}$; PDB: 7SYL; EMD-25532). The remaining 145,559 particles were

classified into either low-resolution maps or with the 40S ribosome lacking the HCV IRES. The

consensus map of the intermediate 40S ribosome particles (Table S7) from the ΔdII IRES eIF5B-

containing sample had 287,087 particles at 4.5 Å resolution (structure $32_{\Delta dII}$; EMD-25591).

1154

Combining ΔdII IRES intermediate 40S ribosome particles and focused classification

Both eIF2-, and eIF5B-containing samples prepared on the ΔdII IRES yielded

classes that showed the HCV IRES bound to the 40S ribosome where the head of the ribosome

was in multiple states (structure $31_{\Delta dII}$-$32_{\Delta dII}$). Both maps were visually inspected and showed high

similarity (unsharpened maps a correlation of 0.9834, and after B-factor sharpening 0.9295). We

combined these particles into a consensus map of 456,311 particles at 3.5 Å resolution (structure

$34_{\Delta dII}$; EMD-25593) that underwent focused classification of the entire head region of the ribosome

that produced three classes: the closed 40S/IRES$_{\Delta dII}$ binary complex with 59,660 particles at 4.8

Å resolution (structure $7_{\Delta dII}$; PDB: 7SYM; EMD-25533), the intermediate-open 40S/IRES$_{\Delta dII}$

binary complex with 144,252 particles at 4.0 Å resolution (structure $8_{\Delta dII}$; PDB: 7SYN; EMD-

25534), and the open 40S/IRES$_{\Delta dII}$ binary complex with 46,095 particles at 4.6 Å resolution

(structure $9_{\Delta dII}$; PDB: 7SYO; EMD-25535). The remaining 206,304 particles were classified into

low resolution maps.

1168

Combining *wt* IRES open 40S ribosome particles and focused classification

Both eIF2-, and eIF5B-containing samples prepared on the *wt* IRES yielded classes that

showed the HCV IRES bound to the 40S ribosome with IRES domain II inserted into the E site

(structure 29_{wt}-30_{wt}). Both maps were visually inspected and showed high similarity (unsharpened

maps correlation of 0.9945, and after B-factor sharpening 0.9464). We combined these particles

into a consensus map of 324,102 particles at 3.3 Å resolution (structure 35_{wt}; EMD- 25594) that

underwent focused classification of eIF1A that produced two classes: the binary complex with

119,320 particles at 3.8 Å resolution (structure 10_{wt}; PDB:7SYP; EMD-25536), and the ternary

complex with 204,782 particles at 3.8 Å resolution (structure 11_{wt}; PDB:7SYQ; EMD-25537).

1178

Model building, and refinement

For all data, where applicable, we were able to unambiguously fit the head and body of the

40S (PDB: 6D9J; Pisareva et al. 2018), HCV IRES (PDB: 5FLX; Yamamoto et al. 2015), eIF1A

(PDB: 4KZZ; Lomakin and Steitz 2013), tRNA (PDB: 5K0Y; Simonetti et al. 2016) eIF2α subunit

(PDB: 6O85; Kenner et al. 2019), and eIF5B (PDB: 4UJD; Yamamoto et al. 2014). Initial model

fitting was performed using UCSF Chimera v1.14 (Pettersen et al., 2004) with additional

modelling in Coot (Emsley and Cowtan, 2004). For regions of eIF5B that did not have available

models (e.g., Switch 1) model building was performed independently and then cross-checked for

consistency. All models underwent one round of Phenix geometry minimization and multiple

rounds of PHENIX real-space refinement (Adams et al., 2010; Afonine et al., 2018).

1189

Figures

1191   All figures were prepared using UCSF Chimera v1.14 (Pettersen et al., 2004).

1192

Quantification and statistical analysis

Global resolution estimates were calculated using the 0.143 FSC criterion (Rosenthal and

Henderson, 2003). Local resolution maps were calculated using Relion 3.1 (Scheres, 2012; 2016;

Zivanov et al., 2018; 2019) using the B-factor determined during post processing and the

Modulation transfer function (MTF) curve for the K3 camera at 300 kV provided by the

manufacturer. (https://www.gatan.com/techniques/cryo-em#MTF). RMSD calculations for 18S

rRNA chains were performed using Pymol (Schrödinger, 2015). Model validation for all models

were calculated using PHENIX (Adams et al., 2010; Afonine et al., 2018) installed as part of the

SBGrid package (Morin et al., 2013).

1202

1203

REFERENCES

Acker, M.G., Shin, B.S., Dever, T.E., and Lorsch, J.R. (2006). Interaction between eukaryotic initiation factors 1A and 5B is required for efficient ribosomal subunit joining. J. Biol. Chem. 281, 8469-8475. 10.1074/jbc.M600210200.

Adams, P.D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., McCoy, A.J., Moriarty, N.W., Oeffner, R., Read, R.J., Richardson, D.C., Richardson, J.S., Terwilliger, T.C., Zwart, P.H. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr. D Biol. Crystallogr. 66, 213–221. 10.1107/S0907444909052925.

Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A., and Adams, P.D. (2018). Real-space refinement in PHENIX for cryo-EM and crystallography. Acta Crystallogr. D Struct. Biol. 74, 531-544. 10.1107/S2059798318006551.

Angulo, J., Ulryck, N., Deforges, J., Chamond, N., Lopez-Lastra, M., Masquida, B., and Sargueil, B. (2016). LOOP IIId of the HCV IRES is essential for the structural rearrangement of the 40S-HCV IRES complex. Nucleic Acids Res. 44, 1309-25. 10.1093/nar/gkv1325.

Arhab, Y., Bulakhov, A.G., Pestova, T.V., and Hellen, C.U.T. (2020). Dissemination of internal ribosomal entry sites (IRES) between viruses by horizontal gene transfer. Viruses 12, 612. 10.3390/v12060612.

Battiste, J.L., Pestova, T.V., Hellen, C.U., and Wagner, G. (2000). The eIF1A solution structure reveals a large RNA-binding surface important for scanning function. Mol. Cell 5, 109-119. 10.1016/s1097-2765(00)80407-4

Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A.J., and Berger, B. (2019). Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. Nat. Methods 16, 1153-1160. 10.1038/s41592-019-0575-8.

Boehringer, D., Thermann, R., Ostareck-Lederer, A., Lewis, J.D., and Stark, H. (2005). Structure of the hepatitis C virus IRES bound to the human 80S ribosome: remodeling of the HCV IRES. Structure 13, 1695-706. 10.1016/j.str.2005.08.008.

Brito Querido, J., Sokabe, M., Kraatz, S., Gordiyenko, Y., Skehel, J.M., Fraser, C.S., and Ramakrishnan, V. (2020). Structure of a human 48*S* translational initiation complex. Science 369, 1220-1227. 10.1126/science.aba4904.

Carragher, B., Kisseberth, N., Kriegman, D., Milligan, R.A., Potter, C.S., Pulokas, J., and Reilein, A. (2000). Leginon:
an automated system for acquisition of images from vitreous ice specimens. J. Struct. Biol. 132, 33-45.
10.1006/jsbi.2000.4314.

de Breyne, S., Yu, Y., Pestova, T.V., and Hellen, C.U. (2008). Factor requirements for translation initiation on the
Simian picornavirus internal ribosomal entry site. RNA 14, 367-380. 10.1261/rna.696508.

des Georges, A., Dhote, V., Kuhn, L., Hellen, C.U., Pestova, T.V., Frank, J., and Hashem, Y. (2015) Structure of
mammalian eIF3 in the context of the 43S preinitiation complex. Nature 525, 491-495. 10.1038/nature14891.

d'Imprima, E., Floris, D., Joppe, M., Sánchez, R., Grininger, M., and Kühlbrandt, W. (2019). Protein denaturation at
the air-water interface and how to prevent it. Elife 8, e42747. 10.7554/eLife.42747.

Easton, L.E., Locker, N., and Lukavsky, P.J. (2009). Conserved functional domains and a novel tertiary interaction
near the pseudoknot drive translational activity of hepatitis C virus and hepatitis C virus-like internal ribosomal entry
sites. Nucleic Acids Res. 37, 5537-5549. 10.1093/nar/gkp588.

Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. Acta Crystallogr. D Biol.
Crystallogr. 60, 2126-2132. 10.1107/S0907444904019158.

Filbin, M.E., and Kieft, J.S. (2011). HCV IRES domain IIb affects the configuration of coding RNA in the 40S
subunit's decoding groove. RNA 17, 1258-1273. 10.1261/rna.2594011.

Fletcher, C.M., Pestova, T.V., Hellen, C.U., and Wagner, G. (1999). Structure and interactions of the translation
initiation factor eIF1. EMBO J. 18, 2631-2637. 10.1093/emboj/18.9.2631.

Glaeser, R.M. (2021). Preparing better samples for cryo-electron microscopy: Biochemical challenges do not end with
isolation and purification. Annu. Rev. Biochem. 90, 451-474. 10.1146/annurev-biochem-072020-020231

Grant, T., Rohou, A., and Grigorieff, N. (2018). *cis*TEM, user-friendly software for single-particle image
processing. Elife 7, e35383. 10.7554/eLife.35383.

Hashem, Y., des Georges, A., Dhote, V., Langlois, R., Liao, H.Y., Grassucci, R.A., Pestova, T.V., Hellen, C.U., and
Frank, J. (2013). Hepatitis-C-virus-like internal ribosome entry sites displace eIF3 to gain access to the 40S subunit.
Nature 503, 539-543. 10.1038/nature12658.

1274

1275   Honda, M., Ping, L.H., Rijnbrand, R.C., Amphlett, E., Clarke, B., Rowlands, D., and Lemon, S.M. (1996). Structural

1276   requirements for initiation of translation by internal ribosome entry within genome-length hepatitis C virus RNA.

1277   Virology 222, 31-42. 10.1006/viro.1996.0395.

1278

1279   Huang, B.Y., and Fernández, I.S. (2020). Long-range interdomain communications in eIF5B regulate GTP hydrolysis

1280   and translation initiation. Proc. Natl. Acad. Sci. U.S.A. 117, 1429-1437. 10.1073/pnas.1916436117.

1281

1282   Hussain, T., Llácer, J.L., Fernández, I.S., Munoz, A., Martin-Marcos, P., Savva, C.G., Lorsch, J.R., Hinnebusch, A.G.,

1283   and Ramakrishnan, V. (2014). Structural changes enable start codon recognition by the eukaryotic translation initiation

1284   complex. Cell 159, 597-607. 10.1016/j.cell.2014.10.001.

1285

1286   Hussain, T., Llácer, J.L., Wimberly, B.T., Kieft, J.S., and Ramakrishnan, V. (2016). Large-Scale Movements of IF3

1287   and tRNA during Bacterial Translation Initiation. Cell 167, 133-144.e13. 10.1016/j.cell.2016.08.074.

1288

1289   Jaafar, Z.A., Oguro, A., Nakamura, Y., and Kieft, J.S. (2016). Translation initiation by the hepatitis C virus IRES

1290   requires eIF1A and ribosomal complex remodeling. Elife 5, e21198. 10.7554/eLife.21198.

1291

1292   Jackson, R.J., Hellen, C.U., and Pestova, T.V. (2010). The mechanism of eukaryotic translation initiation and

1293   principles of its regulation. Nat. Rev. Mol. Cell. Biol. 11, 113-127. 10.1038/nrm2838.

1294

1295   Ji, H., Fraser, C.S., Yu, Y., Leary, J., and Doudna, J.A. (2004). Coordinated assembly of human translation initiation

1296   complexes by the hepatitis C virus internal ribosome entry site RNA. Proc. Natl. Acad. Sci. U.S.A. 101, 16990-16995.

1297   10.1073/pnas.0407402101.

1298

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek,

A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov,

S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M.,

Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis, D.

(2021). Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589. 10.1038/s41586-021-

03819-2.

1305

1306   Kenner, L.R., Anand, A.A., Nguyen, H.C., Myasnikov, A.G., Klose, C.J., McGeever, L.A., Tsai, J.C., Miller-Vedam,

1307   L.E., Walter, P., and Frost, A. (2019). eIF2B-catalyzed nucleotide exchange and phosphoregulation by the integrated

1308   stress response. Science 364, 491-495. 10.1126/science.aaw2922.

1309

1310   Kieft, J.S., Zhou, K., Jubin, R., and Doudna, J.A. (2001). Mechanism of ribosome recruitment by hepatitis C IRES

1311   RNA. RNA 7, 194-206. 10.1017/s1355838201001790.

1312

1313 Kolupaeva, V.G., Pestova, T.V., and Hellen, C.U. (2000). An enzymatic footprinting analysis of the interaction of

1314 40S ribosomal subunits with the internal ribosomal entry site of hepatitis C virus. J Virol. 2000 Jul;74(14):6242-50.

1315 10.1128/jvi.74.14.6242-6250.2000.

1316

1317 Kratzat, H., Mackens-Kiani, T., Ameismeier, M., Potocnjak, M., Cheng, J., Dacheux, E., Namane, A., Berninghausen,

1318 O., Herzog, F., Fromont-Racine, M., Becker, T., and Beckmann, R. (2021). A structural inventory of native ribosomal

1319 ABCE1-43S pre-initiation complexes. EMBO J. 40, e105179. 10.15252/embj.2020105179.

1320

1321 Kuhle, B., and Ficner, R. (2014). eIF5B employs a novel domain release mechanism to catalyze ribosomal subunit

1322 joining. EMBO J. 33, 1177-1191. 10.1002/embj.201387344.

1323

1324 Llácer, J.L., Hussain, T., Marler, L., Aitken, C.E., Thakur, A., Lorsch, J.R., Hinnebusch, A.G., and Ramakrishnan, V.

1325 (2015). Conformational Differences between Open and Closed States of the Eukaryotic Translation Initiation

1326 Complex. Mol. Cell 59, 399-412. 10.1016/j.molcel.2015.06.033.

1327

1328 Locker, N., Easton, L.E., and Lukavsky, P.J. (2007). HCV and CSFV IRES domain II mediate eIF2 release during

1329 80S ribosome assembly. EMBO J. 26, 795-805. 10.1038/sj.emboj.7601549.

1330

1331 Lomakin, I.B., and Steitz, T.A. (2013). The initiation of mammalian protein synthesis and mRNA scanning

1332 mechanism. Nature 500, 307-311. 10.1038/nature12355.

1333

1334 Lomakin, I.B., Shirokikh, N.E., Yusupov, M.M., Hellen, C.U., and Pestova, T.V. (2006). The fidelity of translation

1335 initiation: reciprocal activities of eIF1, IF3 and YciH. EMBO J. 25, 196-210. 10.1038/sj.emboj.7600904.

1336

1337 Maag, D., Fekete, C.A., Gryczynski, Z., and Lorsch, J.R. (2005). A conformational change in the eukaryotic

1338 translation preinitiation complex and release of eIF1 signal recognition of the start codon. Mol. Cell 17, 265-275

1339 10.1016/j.molcel.2004.11.

1340

1341 Malygin, A.A., Kossinova, O.A., Shatsky, I.N., and Karpova, G.G. (2013a). HCV IRES interacts with the 18S rRNA

1342 to activate the 40S ribosome for subsequent steps of translation initiation. Nucleic Acids Res. 41, 8706-8714.

1343 10.1093/nar/gkt632.

1344

1345 Malygin, A.A., Shatsky, I.N., and Karpova, G.G. (2013b). Proteins of the human 40S ribosomal subunit involved in

1346 hepatitis C IRES binding as revealed from fluorescent labeling. Biochemistry (Mosc). 78, 53-59.

1347 10.1134/S0006297913010069.

1348

Marintchev, A., Kolupaeva, V.G., Pestova, T.V., and Wagner, G. (2003). Mapping the binding interface between
human eukaryotic initiation factors 1A and 5B: a new interaction between old partners. Proc. Natl. Acad. Sci. U.S.A.
100, 1535-1440. 10.1073/pnas.0437845100.

Matsuda, D., and Mauro, V.P. (2014). Base pairing between hepatitis C virus RNA and 18S rRNA is required for
IRES-dependent translation initiation in vivo. Proc. Natl. Acad. Sci. U.S.A. 111, 15385-15389.
10.1073/pnas.1413472111.

Morin, A., Eisenbraun, B., Key, J., Sanschagrin, P.C., Timony, M.A., Ottaviano, M. and Sliz, P., (2013). Cutting edge:
Collaboration gets the most out of software. Elife. 2, e01456. 10.7554/eLife.01456.

Nag, N., Lin, K.Y., Edmonds, K.A., Yu, J., Nadkarni, D., Marintcheva, B., and Marintchev, A. (2016). eIF1A/eIF5B
interaction network and its functions in translation initiation complex assembly and remodeling. Nucleic Acids Res.
44, 7441-7456. 10.1093/nar/gkw552.

Neupane R, Pisareva VP, Rodriguez CF, Pisarev AV, Fernández IS. (2020) A complex IRES at the 5'-UTR of a viral
mRNA assembles a functional 48S complex via an uAUG intermediate. Elife. 9, e54575. 10.7554/eLife.54575.

Odreman-Macchioli, F., Baralle, F.E., and Buratti, E. (2001). Mutational analysis of the different bulge regions of
hepatitis C virus domain II and their influence on internal ribosome entry site translational ability. J Biol Chem. 276,
41648-41655. 10.1074/jbc.M104128200.

Passmore, L.A., and Russo, C.J. (2016). Specimen Preparation for High-Resolution Cryo-EM. Methods Enzymol.
579, 51-86. 10.1016/bs.mie.2016.04.011.

Passmore, L.A., Schmeing, T.M., Maag, D., Applefield, D.J., Acker, M.G., Algire, M.A., Lorsch, J.R., and
Ramakrishnan, V. (2007). The eukaryotic translation initiation factors eIF1 and eIF1A induce an open conformation
of the 40S ribosome. Mol. Cell 26, 41-50. 10.1016/j.molcel.2007.03.018.

Pestova, T.V., and Hellen, C.U. (2001). Preparation and activity of synthetic unmodified mammalian tRNAi(Met) in
initiation of translation in vitro. RNA 7, 1496-1505. 10.1017/s135583820101038x.

Pestova, T.V., and Kolupaeva, V.G. (2002). The roles of individual eukaryotic translation initiation factors in
ribosomal scanning and initiation codon selection. Genes Dev. 16, 2906-2922. 10.1101/gad.1020902.

Pestova, T.V., Borukhov, S.I., and Hellen, C.U. (1998a). Eukaryotic ribosomes require initiation factors 1 and 1A to
locate initiation codons. Nature 394, 854-859. 10.1038/29703.

1386

1387 Pestova, T.V., Shatsky, I.N., Fletcher, S.P., Jackson, R.J., and Hellen, C.U. (1998b). A prokaryotic-like mode of
1388 cytoplasmic eukaryotic ribosome binding to the initiation codon during internal translation initiation of hepatitis C
1389 and classical swine fever virus RNAs. Genes Dev. 12, 67-83. 10.1101/gad.12.1.67.

1390

1391 Pestova, T.V., Lomakin, I.B., Lee, J.H., Choi, S.K., Dever, T.E., and Hellen, C.U. (2000). The joining of ribosomal
1392 subunits in eukaryotes requires eIF5B. Nature 403, 332-335. 10.1038/35002118.

1393

1394 Pestova, T.V., de Breyne, S., Pisarev, A.V., Abaeva, I.S., and Hellen, C.U. (2008). eIF2-dependent and eIF2-
1395 independent modes of initiation on the CSFV IRES: a common role of domain II. EMBO J. 27, 1060-1072.
1396 10.1038/emboj.2008.49.

1397

1398 Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E. (2004).
1399 UCSF Chimera--a visualization system for exploratory research and analysis. J. Comput. Chem. 25, 1605-12.
1400 10.1002/jcc.20084.

1401

1402 Pisarev, A.V., Kolupaeva, V.G., Pisareva, V.P., Merrick, W.C., Hellen, C.U., and Pestova, T.V. (2006). Specific
1403 functional interactions of nucleotides at key -3 and +4 positions flanking the initiation codon with components of the
1404 mammalian 48S translation initiation complex. Genes Dev. 20, 624-36. 10.1101/gad.1397906

1405

1406 Pisarev, A.V., Unbehaun, A., Hellen, C.U., and Pestova, T.V. (2007). Assembly and analysis of eukaryotic translation
1407 initiation complexes. Methods Enzymol. 430, 147-77. 10.1016/S0076-6879(07)30007-4.

1408

1409 Pisareva, V.P., Pisarev, A.V., and Fernández, I.S. (2018). Dual tRNA mimicry in the cricket paralysis virus IRES
1410 uncovers an unexpected similarity with the hepatitis C virus IRES. Elife 7, e34062. 10.7554/eLife.34062.

1411

1412 Potter, C.S., Chu, H., Frey, B., Green, C., Kisseberth, N., Madden, T.J., Miller, K.L., Nahrstedt, K., Pulokas, J.,
1413 Reilein, A., Tcheng, D., Weber, D., and Carragher, B. (1999). Leginon: a system for fully automated acquisition of
1414 1000 electron micrographs a day. Ultramicroscopy 77, 153-161. 10.1016/s0304-3991(99)00043-1.

1415

1416 Prince, J.B., Taylor, B.H., Rhurlow, D.L., Ofengand, J., and Zimmermann, R.A. (1982). Covalent crosslinking of
1417 tRNA$_1^{Val}$ to 16S RNA at the ribosomal P site: identification of crosslinked residues. Proc. Natl. Acad. Sci. U.S.A. 79,
1418 5450-5454. 10.1073/pnas.79.18.5450.

1419

1420 Quade, N., Boehringer, D., Leibundgut, M., van den Heuvel, J., and Ban, N. (2015). Cryo-EM structure of Hepatitis
1421 C virus IRES bound to the human ribosome at 3.9-Å resolution. Nat. Commun. 6, 7646. 10.1038/ncomms8646.

1422

1423 Reibarkh, M., Yamamoto, Y., Singh, C.R., del Rio, F., Fahmy, A., Lee, B., Luna, R.E., Ii, M., Wagner, G., and Asano,
1424 K. (2008). Eukaryotic initiation factor (eIF) 1 carries two distinct eIF5-binding faces important for multifactor
1425 assembly and AUG selection. J. Biol. Chem. 283, 1094-1103. 10.1074/jbc.M708155200.
1426
1427 Reynolds, J.E., Kaminski, A., Carroll, A.R., Clarke, B.E., Rowlands, D.J., and Jackson, R.J. (1996). Internal initiation
1428 of translation of hepatitis C virus RNA: the ribosome entry site is at the authentic initiation codon. RNA 2, 867-878.
1429
1430 Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus estimation from electron micrographs.
1431 J. Struct. Biol. 192, 216-221. 10.1016/j.jsb.2015.08.008.
1432
1433 Rosenthal, P.B., and Henderson, R., (2003). Optimal determination of particle orientation, absolute hand, and contrast
1434 loss in single-particle electron cryomicroscopy. J. Mol. Biol. 333, 721-745. 10.1016/j.jmb.2003.07.013.
1435
1436 Scheres, S.H. (2012). RELION: implementation of a Bayesian approach to cryo-EM structure determination. J. Struct.
1437 Biol. 180, 519-530. 10.1016/j.jsb.2012.09.006.
1438
1439 Scheres, S.H. (2016). Processing of structurally heterogeneous cryo-EM data in RELION. Methods. Enzymol. 579,
1440 125-157. 10.1016/bs.mie.2016.04.012
1441
1442 Schrödinger, L.L.C. (2015). The PyMOL Molecular Graphics System, Version 2.4.1 (Shrodinger LLC).
1443 https://pymol.org/
1444
1445 Simonetti, A., Brito Querido, J., Myasnikov, A.G., Mancera-Martinez, E., Renaud, A., Kuhn, L., and Hashem Y.
1446 (2016). eIF3 Peripheral Subunits Rearrangement after mRNA Binding and Start-Codon Recognition. Mol. Cell 63,
1447 206-217. 10.1016/j.molcel.2016.05.033.
1448
1449 Simonetti, A., Guca, E., Bochler, A., Kuhn, L., and Hashem, Y. (2020). Structural insights into the mammalian late-
1450 stage initiation complexes. Cell Rep. 31, 107497. 10.1016/j.celrep.2020.03.061
1451
1452 Sizova, D.V., Kolupaeva, V.G., Pestova, T.V., Shatsky, I.N., and Hellen, C.U. (1998). Specific interaction of
1453 eukaryotic translation initiation factor 3 with the 5' nontranslated regions of hepatitis C virus and classical swine fever
1454 virus RNAs. J. Virol. 72, 4775-4782. 10.1128/JVI.72.6.4775-4782.1998.
1455
1456 Spahn, C.M., Kieft, J.S., Grassucci, R.A., Penczek, P.A., Zhou, K., Doudna, J.A., and Frank, J. (2001). Hepatitis C
1457 virus IRES RNA-induced changes in the conformation of the 40s ribosomal subunit. Science 291, 1959-
1458 1962. 10.1126/science.1058409.
1459

1460 Suloway, C., Pulokas, J., Fellmann, D., Cheng, A., Guerra, F., Quispe, J., Stagg, S., Potter, C.S., and Carragher, B.
1461 (2005). Automated molecular microscopy: the new Leginon system. J. Struct. Biol. 151, 41-60.
1462 10.1016/j.jsb.2005.03.010.
1463
1464 Tang, S., Collier, A.J., and Elliott, R.M. (1999). Alterations to both primary and predicted secondary structure of stem-
1465 loop IIIc of the hepatitis C virus 1b 5' untranslated region 5'UTR) lead to mutants severely defective in translation
1466 which cannot be complemented in trans by the wild-type 5'UTR sequence. J. Virol. 73, 2359-2364.
1467 10.1128/JVI.73.3.2359-2364.1999.
1468
1469 Terenin, I.M., Dmitriev, S.E., Andreev, D.E., and Shatsky, I.N. (2008). Eukaryotic translation initiation machinery
1470 can operate in a bacterial-like mode without eIF2. Nat. Struct. Mol. Biol. 15, 836-841. 10.1038/nsmb.1445.
1471
1472 Thakur, A., Gaikwad, S., Vijjamarri, A.K., and Hinnebusch, A.G. (2020). eIF2α interactions with mRNA control
1473 accurate start codon selection by the translation preinitiation complex. Nucleic Acids Res. 48, 810280-10296.
1474 10.1093/nar/gkaa761.
1475
1476 Tivol, W.F., Briegel, A., and Jensen, G.J. (2008). An improved cryogen for plunge freezing. Microsc. Microanal. 14,
1477 375-379. 10.1017/S1431927608080781.
1478
1479 Unbehaun, A., Borukhov, S.I., Hellen, C.U., and Pestova, T.V. (2004). Release of initiation factors from 48S
1480 complexes during ribosomal subunit joining and the link between establishment of codon-anticodon base-pairing and
1481 hydrolysis of eIF2-bound GTP. Genes Dev. 18, 3078-3093. 10.1101/gad.1255704.
1482
1483 Wang, J., Wang, J., Shin, B.S., Kim, J.R., Dever, T.E., Puglisi, J.D., and Fernández, I.S. (2020). Structural basis for
1484 the transition from translation initiation to elongation by an 80S-eIF5B complex. Nat. Commun. 11, 5003.
1485 10.1038/s41467-020-18829-3.
1486
1487 Yamamoto, H., Unbehaun, A., Loerke, J., Behrmann, E., Collier, M., Bürger, J., Mielke, T., and Spahn, C.M. (2014).
1488 Structure of the mammalian 80S initiation complex with initiation factor 5B on HCV-IRES RNA. Nat. Struct. Mol.
1489 Biol. 21, 721-727. 10.1038/nsmb.2859.
1490
1491 Yamamoto, H., Collier, M., Loerke, J., Ismer, J., Schmidt, A., Hilal, T., Sprink, T., Yamamoto, K., Mielke, T., Bürger,
1492 J., Shaikh, T.R., Dabrowski, M., Hildebrand, P.W., Scheerer, P., and Spahn, C.M. (2015). Molecular architecture of
1493 the ribosome-bound Hepatitis C Virus internal ribosomal entry site RNA. EMBO J. 34, 3042-3058.
1494 10.15252/embj.201592469.
1495

1496 Yokoyama, T., Machida, K., Iwasaki, W., Shigeta, T., Nishimoto, M., Takahashi, M., Sakamoto, A., Yonemochi, M.,
1497 Harada, Y., Shigematsu, H., Shirouzu, M., Tadakuma, H., Imataka, H., and Ito, T. (2019). HCV IRES captures an
1498 actively translating 80S ribosome. Mol. Cell 74, 1205-1214.e8. 10.1016/j.molcel.2019.04.022.

1499

1500 Zheng, A., Yu, J., Yamamoto, R., Ose, T., Tanaka, I., and Yao, M. (2014). X-ray structures of eIF5B and the eIF5B-
1501 eIF1A complex: the conformational flexibility of eIF5B is restricted on the ribosome by interaction with eIF1A. Acta
1502 Crystallogr. D Biol. Crystallogr. 70, 3090-3098. 10.1107/S1399004714021476.

1503

1504 Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017). MotionCor2: anisotropic
1505 correction of beam-induced motion for improved cryo-electron microscopy. Nat. Methods 14, 331-332.
1506 10.1038/nmeth.4193.

1507

1508 Zivanov, J., Nakane, T., Forsbery, B.O., Kimanius, D., Hagen, W.J., Lindahl, E. and Scheres, S.H.W. (2018). New
1509 tools for automated high-resolution cryo-EM structure determination in RELION-3. Elife 7, e42166.
1510 10.7554/eLife.42166

1511

1512 Zivanov, J., Nakane, T., and Scheres, S.H.W. (2019). A Bayesian approach to beam-induced motion correction in
1513 cryo-EM single-particle analysis. IUCr. J. 6, 5-17. 10.1107/S205225251801463X

1514

Brown et al., Figure 1

A Pre-48S IC (eIF5B) Structure 14wt

Head — Beak — Met-tRNAiMet — eIF1A — eIF5B — Body — Intersubunit side — Domain II — IRES — Domain IIId — Platform side

B 48S IC (eIF5B) Structure 15wt

Head — Beak — Met-tRNAiMet — eIF1A — eIF5B — Body — Intersubunit side — IRES — Platform side

C eIF5B repositions tRNA

14° 15.6Å — tRNA (eIF5B) — tRNA (eIF2) — eIF2α — 48S IC (eIF5B) — eIF1A

D mRNA — U(35) — A(34) — C(33) — E1216 — R1186 — H112 — I1218 — OB domain — E1216 — R1186 — H112 — I1218 — 90° rotation

E tRNA (48S IC) — 6.4Å Domain IV — 3.1Å G domain — 6.4Å tRNA — tRNA (80S) — eIF1A — 48S IC (eIF5B) — 80S ribosome

F 48S IC (eIF5B) — uL16 — tRNA — Clash — 28S — Domain IV — 80S ribosome — uL16 — tRNA — 28S — Domain IV — Domain movement — 6.4Å tRNA movement — C(56) — C(74) — 30.9° — 12.4° — His1143 — His924 — Phe1082 — Phe858 — 6.4Å Domain IV

G 48S IC (eIF2) Structure 12wt

U(35) — A(34) — C(33) — Met-tRNAiMet — U1248 — eIF1A — W70 — A(+4) — A1825 — G(+3) — U(+2) — A(+1) — 18S

H Pre-48S IC (eIF5B) Structure 14wt

U1248 — U(35) — Met-tRNAiMet — A(34) — C(33) — eIF1A — W70 — G(+3) — C(-1) — A(+1) — U(+2) — A1825 — 18S

I 48S IC (eIF5B) Structure 15wt

U(35) — A(34) — C(33) — Met-tRNAiMet — U1248 — eIF1A — W70 — A(+4) — A1825 — G(+3) — U(+2) — A(+1) — 18S

A

43S scanning

Pre-48S IC

eIF1

P_OUT

40S head

mRNA
Pre-48S IC
43S scanning

C

β3-β4 loop

G(23)

U(24)

β1-β2 loop

A(36)

eIF1

B

43S scanning

G1639
A1640

Arg146

U1248

(+1)
(+2)
(+3)

Pre-48S IC (eIF5B)
Structure 14_wt

C1701

(+1)
(+2)
(+3)

48S IC (eIF5B)
Structure 15_wt

G1639
A1640

Arg146

U1248

C1701

(+1)
(+2)
(+3)