Sensory convergence in the world's largest cavefish diversification: patterns of

neuromast evolution, distribution and associated behaviour in *Sinocyclocheilus*

Bing Chen[1], Tingru Mao[1], Yewei Liu[1], Wenzhang Dai[1], Xianglin Li[1], Amrapali P. Rajput[1], Jian Yang[2], Joshua B. Gross[3], Madhava Meegaskumbura[1*]

[1] Guangxi Key Laboratory for Forest Ecology and Conservation, College of Forestry, Guangxi University, Nanning, Guangxi, China

[2] Key Laboratory of Environment Change and Resource Use, Beibu Gulf, Nanning Normal University, Nanning, Guangxi, China

[3] Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45221, U.S.A.

Corresponding author *E–mail* : madhava_m@mac.com

Abstract

The genus *Sinocyclocheilus* represents the largest freshwater cavefish genus in the world. This emerging model system is endemic to the southern Chinese karstic landscape, and demonstrates multiple adaptations for caves (troglomorphism), with eye-degeneration being the most pronounced. The less-apparent lateral line system, which is often expanded in cave-dwellers, has been studied in other cavefish systems, but never in the context of this diversification. Here we investigated the distribution and evolution of cephalic neuromasts in 26 *Sinocyclocheilus* species. We used live-staining and behavioural assays, and interpreted results in a phylogenetic context. We show that asymmetry in neuromast features and the rate of evolution is greater in cave-adapted species. Ancestral state reconstructions show that most *Sinocyclocheilus* are right-biased with some scatter, and show convergence of neuromast phenotypes. There is substantial variation in cephalic neuromast distribution patterns between and (to a lesser extent) within species. Behavioural assays show blind species have a distinctive wall-following behaviour. We explain these patterns in the context of the deep evolutionary history associated with this karstic region, other organismal traits, and habitat occupation of these remarkable diversifications of fishes. Interestingly, some of these neuromast patterns and behaviour show convergence with other phylogenetically distant cavefish systems.

Key words: Asymmetry, convergence, ancestral state reconstructions, wall-following behaviour, live staining, troglomorphisim

Background

Cave-dwelling (hypogean) fish provide a valuable system to study evolutionary novelty [1-4], owing to the extreme conditions associated with these habitats, such as limited food and perpetual darkness. Strong selective pressures arising from this extreme environment are associated with a suite of regressive morphological changes, such as loss of eyes and pigment [5, 6]. However, numerous constructive features also evolve in cave-dwellers. Among the most notable are expansions of the mechanosensory lateral line system [7]. Here, we examine the evolution of the lateral line system in the species-rich genus *Sinocyclocheilus*, an emerging model system and the largest diversification of freshwater cavefish in the world [8, 9].

Neuromast organs are composed of hair cells with cilia embedded in a gelatinous cupula, surrounded by a complex set of support cells [10, 11]. There are two general neuromast types: canal neuromasts (CN), which are larger, and embedded in a bony canal, and superficial neuromasts (SN), which project directly from the skin epithelium [12-14]. In the Mexican tetra (*Astyanax*), cave morphs harbor variation in the number and size of both types of neuromasts in the lateral line, including several-fold more SNs compared to surface fish. Moreover, *Astyanax* cave morphs have a highly asymmetric distribution of neuromasts across the left/right axis compared to surface-dwelling morphs [15]. At present, it is unclear the relevance of this asymmetry, however, some studies suggest it facilitates lateral swimming preference, rheotaxis (orientation towards flow), foraging, shoaling, predator avoidance, and mate-finding [7, 16-20].

Sinocyclocheilus are distributed across a vast 62,000 km^2 karstic area in Yunnan

and Guizhou Provinces and Guangxi Zhuang Autonomous Region [21, 22]. Based on their habitat preference, these fish can be divided into three principal groups: troglobitic (cave-restricted); troglophilic (cave-associated); and surface-dwelling [9]. However, one study characterized neuromast size differences between two *Sinocyclocheilus* species [20]. *Sinocyclocheilus* species span the macroevolutionary transition from surface to cave-restricted species and thus offer an exceptional opportunity to examine the evolution of traits associated with this transition.

Here we examined numerical variation, lateral distribution and behavioural differences associated with neuromasts in *Sinocyclocheilus*. Neuromasts may provide functional compensation for eye loss, therefore we hypothesized that more severe eye/vision loss may be associated with compensatory expansions of the lateral line neuromasts. Specifically, troglomorphy was predicted to be associated with more, and larger, neuromasts. Additionally, we anticipated a higher prevalence of distributional asymmetry of neuromasts in troglomorphic species compared to surface species, owing to relaxation of symmetry.

We found that neuromast distributions were asymmetric among all groups, however the degree of asymmetry was most apparent for Regress-Eyed (troglomorphic; collectively both Blind and Micro-Eyed) species. Additionally, in phylogenetic contexts the pace of neuromast evolution is faster among species with smaller eyes, compared to normal eyed species. Taken together, this work provides the first diversification-wide description of lateral line evolution, and clarifies the shared evolutionary pressure on constructive trait evolution among distantly-related species that colonize caves.

2. Methods

Sinocyclocheilus adult fishes used in this study were examined using live, biosafe

staining techniques and behavioural assays. The project was approved by Guangxi

Autonomous Region Government and Guangxi University Ethical Clearance

Committee (# GXU-2021-125).

(a) Fish maintenance in captivity

Adult fish used in this study were collected from the field from 2017-2020 across

Yunnan and Guizhou Provinces and Guangxi Zhuang Autonomous Region of China

(figure 1, see geographic coordinates in electronic supplementary material: table S1,

figure S5). All fish were maintained in a centralized fish aquarium system, and

maintained at (pH: 7.0 - 8.0, temperature 19℃ ± 1℃, dissolved oxygen 8.5mg/L).

Specimens larger than 5 cm were maintained in large glass aquaria ($90 \times 50 \times 50$ cm,

$150 \times 80 \times 80$ cm), with individual mechanical and bio-filters. For this study, 76

individuals from 26 *Sinocyclocheilus* species were used (table 1). *Carassius auratus* (n

= 3) and *Cyprinus carpio* (n = 3), two species from a closely related clade to

Sinocyclocheilus for comparison.

Table 1. Summary of 26 *Sinocyclocheilus* species used in the current analysis grouped

according to eyes morphology and habitat.

	Troglobitic (TB)	Troglophilic (TP)	Surface (S)
Blind (B)	*S. furcodorsalis* *S. tianlinensis* *S. tianeensis* *S. xunlensis*		
Micro-eyed (ME)	*S. mashanensis* *S. altishoulderus* *S. microphthalmus* *S. bicornutus* *S. rhinocerous* *S. cyphotergous*	*S. multipunctatus*	
Normal-eyed (NE)	*S. guilinensis* *S. huanjiangensis* *S. macrophthalmus* *S. guanyangensis* *S. brevis* *S. lingyunensis* *S. zhenfengensis* *S. jii* *S. yishanensis*	*S. longibarbatus*	*S. angustiporus* *S. oxycephalus* *S. malacopterus* *S. purpureus* *S. maitianheensis*

Figure 1. Map showing the distribution of sampling sites of the 26 species of
Sinocyclocheilus **(n = 76) used in this study.** These fishes are mainly distributed across
South China's Karstic habitats on three provinces: Guangxi Autonomous region (GX),
Guizhou (GZ) and Yunnan (YN). The four main clades (A – D, represented by a color
code), the species used from each clade, and a representative photo of species from
these clades are shown. Early emerging Clade A is mainly distributed in the eastern
range of the diversification and Clade D to the hills to the West; Clade B and C contain
the most troglomorhpic species and are distributed mainly across GX and GZ. The eye-
condition (Blind, Micro-eyed and Normal-eyed) are depicted using a color code.

(b) Vital staining of neuromasts

To visualize bones and neuromasts, live fish were immersed in 1mg/L Calcein-AM

(C0875-5G; Sigma-Aldrich) and 20μg/ml DASPEI (D0815; Sigma-Aldrich) [15, 23,

24]. Fish were anesthetized using MS222 (E808894-5g; Macklin) 0.01-0.02g/L for 3-6

min followed by brief immersion in ice water for ~20s, depending on size and age (e.g.,

smaller specimens were exposed to ice water for shorter periods). All individuals were

continuously monitored to ensure the health and safety of experimental individuals.

(c) Digital Analysis of Neuromast Position and between Distances

This study focused on the cephalic region anterior to the opercular and interopercular

bones, near the arc on the lateral aspect [20, 22]. Images were collected using a Leica

M165FC fluorescent stereomicroscope outfitted with Leica Application Imaging

software (LAS X v3.4.2.18368). The dorsal, left and right aspects of each individual

were imaged under identical conditions. High-definition "montage" image were

consolidated following automated alignment and 'flattening' of 30 images collected in

the z-plane (× 9/ 13/ 16 magnification) using LASX or Helicon Focus (Pro v7.6.1), to

perform the two-dimensional shape images. The use of montage imaging minimized

potentially confounding variables arising from the z-plane for each specimen.

The outline of fish and neuromast positions were obtained using the "Pencil" and

"Point" tool in GIMP (v2.10.24). Neuromast numbers were quantified using an

automated method in ImageJ (v1.8.0), following the method of Gross et al. (2016) [15].

For ambiguous images, we performed direct, manual counting (figure 3*a*). We

calculated the size (2-D area) of individual neuromasts, lens, and eye orbit diameter by

determining their size in pixels using the "measure" function in ImageJ and converting

to mm^2. We also used a vernier caliper to measure standard length (SL). To determine

the density of neuromast within particular unit areas, we used "Delaunay Voronoi"

triangulation in ImageJ to define a proxy for the mean distance between neuromasts

[25-27] (figure 3*a*).

(d) Quantifying symmetry

To examine positional symmetry of neuromasts across the left-right axis, we

manually superimposed the fluorescent images of neuromasts (figure 3*b*). We used

excitation filters for three colors (Blue 470/40 nm, Texas Red 560/40 nm and Green

546/10 nm). In GIMP, we then reflected the left images to align with the right images,

creating a single image. We measured the scoring of neuromasts in the reverse sequence

(right-side first) relative to the initial scoring of neuromasts (left-side first) to avoid bias

in our calculations [28].

The "colocalization" [29, 30], and "JACop" [31, 32] plugins in FIJI were used to

calculate an asymmetry coefficient [15, 33, 34]. We performed a Pearson's correlation

to compare the intensity distribution between channels [35]. Next, we calculated an

Overlap Coefficient (OC), to identify positional overlap of signals from the left and

right sides [36]. This enabled us to quantify the extent to which positions of neuromasts

on the left and right sides of the cranium were symmetric (figure 3*a*).

We divided 26 species into Regressed-Eyed groups (containing Blind: n = 12 and

Micro-Eyed: n = 21) and Normal-Eyed groups (n = 45), following the categorization

and phylogeny of Mao et al. (2021). Importantly, a few *Sinocyclocheilus* species do not

have uniform eye sizes [22]. For instance, *S. guanyangensis* have highly variable eye

sizes, so for this study we selected individuals with the most common Normal-Eyed

morphology. All parametric neuromast measures were subjected to one-way ANOVA

with a post-hoc Tukey test analysis. Non-parametric distributions were subjected to

Wilcoxon Signed Rank Test (2-tailed, Holm correction) or a Kruskal-Wallis test with a

post-hoc analysis Dunnett test (2-tailed, Bonferroni correction). Statistical significance

was set at $p < 0.05$. We used the packages "wmc" and "FSA" in R (v3.6.3). Principal

Component Analysis (PCA) was used to evaluate the following variables: number, area,

asymmetry coefficients and mean distance. To analyse the relationships between

Regressed-Eyed and Normal-Eyed groups, we performed Spearman's rank correlation

coefficients using SL, eye traits, area, number, mean distance and OC of neuromasts

patterns.

(e) Phylogenetic inference

To study the patterns of neuromast evolution in *Sinocyclocheilus,* we inferred a

phylogenetic tree for the 26 species, using two available gene fragments from Genbank

(NADH-ubiquinone oxidoreductase chain 4 - ND4 and cytochrome b gene – Cyt b),

together with an outgroup species (*Cyprinus carpio*) (table S1). We used two different

methods for phylogenetic inference [37] – Bayesian and Maximum Likelihood for tree

construction (alignment, model selection and phylogenetic inference method details are

available in the electronic supplementary material, Supplementary Methods).

(f) Evolution rate analysis

To test the allometric evolution rate between neuromasts in different morphs [9],

we analysed neuromast numbers, left/right-side numbers, mean distance coefficient,

area and density (i.e. the proportion of number and area anterior to the gill). We used

100 potential trait histories from stochastic character mapping, and fit two alternative

models (single or multiple rate model, calculated by AIC) of evolution for each studied

trait. In the case of small samples, we assumed the AIC and AICc to assess, and then

weighted from Bayesian analyses on the trees using the brownie.lite function. We used

the package "rgl", "ape" and "phytools" in R to calculate the model-averaged estimates

[38].

(g) Patterns of neuromasts evolution using ancestral state reconstructions

To understand the broad patterns of neuromast evolution in *Sinocyclocheilus*, we

classified 26 species into the following three morphological groups: (1) Left-right axis

asymmetry: according to different degree of overlapping coefficient, divided as

Absolute-asymmetry (OC < 0.1) and Slight-asymmetry (OC ≥ 0.1) based on Gross et

al. (2016) [15]. (2) Left/right-bias handedness: we used the normalized SN number by

signed (directional) asymmetry rate: $\text{SAR} = \frac{R-L}{R+L} \times 100\%$ following Planidin et al. (2021)

[39], which suggests the presence of two morphological categories: Right-biased (SAR

> 0, neuromasts on right-side mainly) and Left-biased (SAR < 0, figure S1). (3)

Distance expansion: depending on the results of neuromasts mean distance coefficient,

we divided species into Scattered (DEL > 0.2) and Serried (DEL ≤ 0.2), following

Gross et al. (2016) [15] (table S1).

To understand the evolution of neuromast patterns, we performed ancestral state

reconstructions using these three morphological categories. One hundred stochastic

reconstructions were simulated through the stochastic mapping approach that was

conducted using function "make.simmap" (model = "ER") in the R packages "ape",

"phytools" and "viridisLite".

(h) Cave-dwelling behaviour associated with neuromast variation

To understand how neuromast patterns may facilitate certain behaviour, we carried out

a series of behavioural assays. All assays were performed using the following 14 species

(n = 3 individuals for each): Blind - *S. furcodorsalis, S. tianlinensis, S. tianeensis*;

Micro-Eyed - *S. mashanensis, S. microphthalmus, S. bicornutus, S. multipunctatus*;

Normal-Eyed - *S. guilinensis, S. longibarbatus, S. macrophthalmus, S. oxycephalus, S.*

zhenfengensis, S. purpureus, S. maitianheensis. Each experimental fish was

acclimatized for 30 min in a rectangle assay arena (45 × 28 × 28 cm) in system water.

An infrared camera (Cannon XF400/405) was used to capture video under infrared light

in a quiet, dimly lit room (frame rate: 4Mbps (VBR, YCC 4:2:0, 25p, 1280 × 720).

Wall-following behaviour has been observed in *A. mexicanus* cavefishes; therefore,

we examined a range of wall-following within the arena. This included fish swimming

a minimum distance of its SL along the wall, or a distance of ≤ 0.5 SLs away from the

wall (figure 5*a-b*) [40, 41]. We recorded normal tracking without stimulation for 10-

min to determine if wall-following behaviour was present using EthoVision XT (v15.0 ,

Noldus) alongside direct visual monitoring. (2) We next used an aeration pump (45~50

Hz vibration) for 5-min to record the approaching of a novel object in modified

vibration attraction behaviour test [42-44]. We selected the area centered on the

stimulation model based on tank shape (10 × 16 cm). We monitored the frequency of

time spent in the stimulation range from left/right-side. Data analysis were performed

in R's basic functions as mentioned.

Results and Discussion

Neuromast distribution is asymmetric in all cave fish studied to date [15, 45-48]. Accordingly, we found the two outgroup species of Cyprinidae showed several neuromasts on observation region, which did not show a comparable pattern of neuromast asymmetry as was observed in *Sinocyclocheilus*. Fish living in perpetual darkness frequently lose vision while enhancing non-visual sensation, such as the lateral line [42, 49, 50]. Here, for the first time, we show that *Sinocyclocheilus* similarly conform to this pattern (figure 2). Additionally, widespread neuromast distributional asymmetry in *Sinocyclocheilus* is convergent with other distantly related cavefish species inhabiting similar cave microenvironments. Interestingly, our results showed that this asymmetric neuromast pattern is variable across the 26 *Sinocyclocheilus* species tested, with most species demonstrating a right-sided bias (figure 4). Evidence for neuromast asymmetry patterns in other lineages comes from two lineages from Central and North America: *Astyanax mexicanus,* with a left-side bias and the *Gasterosteus aculeatus* (stickleback fish), with a right-side bias [39, 51]. These convergent results reveal variation in "sidedness" across taxa, despite convergence in the asymmetry of neuromasts traits.

(a) Patterns of neuromast symmetry and asymmetry

The neuromasts distribution pattern showed asymmetry across the left-right axis, with variation both within and across species (figure 2*a*, table S2). Interestingly, mean overlap coefficients indicated that all species showed a degree of neuromast asymmetry.

Normal-Eyed species showed the least asymmetry, while Blind species showed the

highest and Micro-eyed species had intermediate values (mean ± s.d.: Normal-Eyed =

0.098 ± 0.05; Blind = 0.091 ± 0.031; Micro-eyed = 0.096 ± 0.027; $H_2 = 0.64$, p > 0.05).

This finding is consistent with prior work suggesting that asymmetry may facilitate

navigation in darkness [52, 53], foraging [54] and the ability to maximize sensory

information using fewer receptors [39, 51]. In comparing asymmetry measures across

clades, Clade D (all Normal-Eyed, surface-dwelling species, showed the least

asymmetry (mean ± s.d. = 0.119 ± 0.058), while Clade C (mean ± s.d. = 0.079 ± 0.031)

showed the highest asymmetry; with Clade A and B demonstrating intermediate values

(Clade A – 0.086 ± 0.029 and Clade B – 0.104 ± 0.038).

Figure 2. Comparison of neuromast-related measurements for 26 species of *Sinocyclocheilus* (n = 76).

(a) The average scores of Overlapping Coefficient (OC) measured as the pattern of

neuromasts on left/right-side of the fish. The standard of asymmetry results is defined

as: asymmetry (OC < 0.1), symmetry (OC ≥ 0.1). Clade A-D represent the clade level

relationships of these 26 species in the phylogenetic tree. All statistical results are

available in table S1, S2.

(b) The bars represent the individual's canal neuromasts (CN) total counts and the lines

represent the left and right sides of cephalic CN in different species. The CN of Micro-

Eyed species was significantly less than Normal-Eyed species (H = -2.64, p = 0.025).

(c) The mean number of superficial neuromasts (SN) on the cephalic area (dorsal, left

and right sides) in different *Sinocyclocheilus* species.

(d) The mean surface area covered by SN on the dorsal/left/right-side of different

Sinocyclocheilus species. The surface species covered by the dorsal SN are less than

that of the surface are covered by the right and left sides of fish.

(e-g). The comparisons about: (e) the number of SN; (f) neuromasts mean distance

coefficient; (g) area of SN on Dorsal/Left/Right-side on cephalic area. Group divided

by the Blind (B in orange), Micro-Eyed (M in blue) and Normal-Eyed (N in yellow).

The Wilcoxon signed rank (2-tailed, Holm correction) test suggested that the surface

covered by dorsal neuromasts are significantly less than that covered by the right and

left sides. *: P < 0.05, **: P < 0.01, and ***: P < 0.001. All statistical results are

available in table S2, S5.

(h-j). The comparisons of the non-parametric statistical tests performed for (h), the

mean number of SN; (i) neuromasts mean distance coefficient; (j) mean area of SN on

the cephalic area of the Blind, Micro-Eyed and Normal-Eyed morphs. The comparisons

were sub-divided into dorsal-side (D in orange), left-side (L in blue) and right-side (R

in yellow) for visualization purposes.

(b) Patterns of neuromast distribution

Generally, we found more SNs on the right compared to the left-side, however we

observed substantial variation between and across species. In the dorsal aspect, Blind

fish have the fewest, Normal-Eyed fish have the most, and the Micro-Eyed group was

intermediate (figure 2*e*). Unexpectedly, there were significantly fewer SNs in Blind

species (mean ± s.d.: Blind = 1543 ± 248; Micro-Eyed = 2252 ± 802; Normal-Eyed =

1859 ± 504; H_2 = 10, p = 0.0065, table S5). In terms of lateral distribution, Blind and

Normal eyed fish had higher numbers on the right-side, while Micro-Eyed species had

more neuromasts on the left-side (median: left = 767, right = 758; W = 227, p = 0.890;

figure 2*h*; table S5).

We found that dorsal neuromast numbers are significantly lower than the lateral

sides. An interesting exception was *S. lingyunensis* (N-TB, Clade C), dorsally

distributed neuromasts outnumbered laterally-situated neuromasts (mean dorsal = 557,

left = 419, right = 453; figure 2*c*, table S5). In the dorsal aspect, Normal-Eyed species

had the most neuromasts, while the Blind group had the fewest. One potential

explanation for this difference could be that obligate subterranean fishes experience

relaxed selective pressure (e.g. aerial predation from birds) [21, 55], but perhaps dorsal

neuromasts are necessary for navigation within caves [56].

With respect to neuromast expansion (scatter), we found that the most scattered

SNs are dorsal, especially in Blind species (mean ± s.d.: Blind = 0.31 ± 0.06; Micro-Eyed = 0.28 ± 0.10; Normal-Eyed = 0.24 ± 0.04; H_2 = 12, p = 0.002; figure 2*f*, table S5). Additionally, neuromasts tend to be more scattered on right- compared to the left-side across all three groups (median right/left Blind = 0.26, 0.25; Micro-Eyed = 0.22, 0.21; Normal-Eyed = 0.24, 0.23; figure 2*d*). Interestingly, the Blind cavefish group demonstrated the most scatter, while the Micro-Eyed group had the least scatter (figure 2*i*, table S5). The Mexican cavefish (*Astyanax mexicanus*) demonstrates more scatter in the surface forms compared to *Sinocyclocheilus* [15, 57], however this is likely due to the fact that *Astyanax* Blind forms have more neuromasts, and hence less scatter.

In the comparison of surface area covered by neuromasts, we found a significantly lower surface area populated by neuromast on the cephalic region dorsally, with the Blind group having the smallest area (mean ± s.d. Blind = 2.14 ± 0.59; Micro-Eyed = 3.13 ± 1.94; Normal-Eyed = 3.88 ± 1.33; H_2 = 15.9, p = 0.0003; figure 2*j*). When comparing lateral sides, we found the total distributional area of neuromasts to be highly comparable within groups (median right/left-side Blind = 4.44, 4.6; Micro-Eyed = 6.19, 6.21; Normal-Eyed = 5.31, 5.14; table S5). Normal-Eyed species have the largest area of neuromast distribution, Micro-Eyed fish have an intermediate area, and the Blind group had the smallest area covered by (mean ± s.d. = 10.56 ± 2.95; Micro-Eyed = 16.44 ± 8.80; Normal-Eyed = 14.49 ± 4.59; F = 3.83, p=0.026; figure 2*g*). Although it has been shown that the neuromast complexity of the Blind species is greater than Normal-Eyed species [20], this however, has not been established at a diversification-wide scale.

Blind *Sinocyclocheilus* have fewer CNs compared to Normal-Eyed

Sinocyclocheilus (median Blind = 5, Micro-Eyed = 13; Normal-Eyed = 17; table S5),

reflecting the same pattern as SNs. For *Astyanax*, surface fish are invariant in terms of

numbers, but cavefish are highly variable. Surprisingly, two species, *S. rhinocerous*

and *S. cyphotergou* appeared to possess no CNs (both M-TB, figure 2*b*), suggesting

the canal lateral line system may have regressed in this species. Interestingly, a similar

phenomenon has been observed in Amblyopsid cavefish (Teleostei: Percopsiformes)

[47] and in Threespine Sticklebacks (*Gasterosteus aculeatus*) [58].

In sum, neuromasts are generally reduced in number and area in Blind species, but

not in distribution (expansion). Somewhat surprisingly, Micro-Eyed species had the

most neuromasts, greater area, and least dense distribution of neuromasts. We propose

that Blind species may have optimized adaptation to the subterranean biome by using

fewer, but more complex, neuromasts. Blind *S. tianlinensis*, for instance, have SNs with

greater diameters and more hair cells compared to Normal-eyed *S. macrophthalmus*

[20].

(c) Correlations between asymmetry and eye condition

We performed a Spearman Rank's correlation of the highly loading variables for

the Regressed-Eyed and Normal-Eyed. In the Regressed-Eyed group, eye measures

demonstrated a significantly positive correlation with neuromasts number (ρ= 0.7, p <

0.001) and area (ρ= 0.7, p < 0.001) , but a slightly negative correlation with mean

distance between neuromasts (ρ= -0.3, p < 0.05). In the Normal-Eyed group, eye

measures had a significantly positive correlation with neuromasts area (ρ= 0.4, p <

0.005) and mean distance (ρ= 0.4, p < 0.05) , but significantly negative correlation with

the results of asymmetry (ρ= -0.4, p < 0.001; figure 3*d,e*, table S4). Overall, the

Regressed-Eyed group is more asymmetric compared to Normal-Eyed group.

Figure 3. Summary of fluorescent staining results for different cavefish morphs.

(a) the fluorescent staining results of *S. tianlinensis* (Blind species-Troglobitic, B-TB).

(table 1 & table S1).

(a-c) photos taken under normal lights. (d-f) dorsal, left and right sides of the

neuromasts under fluorescent light. (small green dots are SN, bigger green dots are CN).

(g-i) The diagrammatic representations of the dorsal, right and left side neuromasts. SNs denoted by red dots and the area outlined in black represent cephalic region and the olfactory area. (j-l) The neuromasts mean distance measures as the "Delaunay Mean Distance" are indicated by yellow lines. (m) The asymmetry values calculated by overlap coefficient. Red and green dots represent the left and right sides of the fish. Note that the scale of the images is the same at the length of 1.0 mm.

(b) The results of the fluorescent staining of three cavefishes, (i-iii) *S. furcodorsalis,* (iv-vi) *S. microphthalmus* and (vii-ix) *S. oxycephalus,* used as representatives of each eye-related morphs. These three species are classified as (B-TB), Absolute-asymmetry/Scattered/Left-bias; (M-TB), Slight-asymmetry/ Scattered/Right-bias and (N-S), Slight-asymmetry/Serried/Right-bias respectively. Scales of the images are the same at the length of 1.0 mm.

(c) Principal Component Analysis (PCA) bi-plot for the distribution traits of neuromasts in *Sinocyclocheilus*.

The dorsal/left/right-side and mean counts number (Num-D, Num-L, Num-R, NUM); dorsal/left/right-side and mean distance coefficient (Del-D, Del-L, Del-R, DEL); asymmetry coefficients of neuromasts (PearsonsC, OverlapC); eyes traits (area of lens/orbit (Area-Lens, Area-Orbit) and length of lens/orbit (Length-Lens, Length-Orbit)) with standard length [5] were included as variables in the PCA. All results are available in Supplemental table S2, S3.

(d) The Correlations between SN number (NUM), area (AREA), mean distances

coefficient (DEL) and asymmetry coefficient (OC) of neuromasts; standard length [5]

and the length of lens (LENGTHEYE) and area of lens (AREAEYE). The   scores

indicate the results of the spearman's rank correlation coefficient $|\rho|$; < 0.3 = no

correlation, 0.3-0.8 = low correlation, > 0.8 = high correlation. Positive correlations (ρ >

0) and negative correlations ($\rho \leq 0$) are shown in blue and red colors respectively. *: P

< 0.05, **: P < 0.01, and ***: P < 0.001 indicate the level of statistical significance. All

results are available in table S4.

(d) Dimensionality of the neuromast variables

We performed a principal component analysis (PCA) based on 26 species of

Sinocyclocheilus and their neuromast-related variables, to identify key features

underlying neuromast variation. Variation on PC1 was explained mostly by the number

of neuromasts (mean number: NUM, counts on dorsal side: Num-D) and the total

neuromast distributional area (AREA). PC2 was explained mostly by the distances

between neuromasts (mean distance coefficient on left side: Del-L, on right side: Del-

R; table S3). The 95% confidence intervals for each group shows that intraspecific

variation in the Blind group was the narrowest, while the Micro-Eyed group showed

the highest variation (figure 3*c*). This may be explained by the fact that Micro-Eyed

forms are subjected to selective pressures of both surface habitats, subterranean habitats

and the transitional habitats between these [22, 59, 60]; i.e. habitat heterogeneity.

(e) Patterns of neuromasts evolution

We next performed an ancestral reconstruction for three character states: asymmetry,

handedness and neuromasts expansion. We found the deeper nodes in the phylogeny

showed ambiguity for asymmetry and handedness (see maximum credibility tree;

figure 4). However, the deeper nodes for neuromast distribution suggested a scattered

distribution of neuromasts is the ancestral condition. In *Sinocyclocheilus,* ~ 70% of

the species examined demonstrate right-handedness. Our ancestral state

reconstruction shows an entire clade of 7 species is right-handed. However, both left-

and right-handed mixed clades and sister taxa are present within the diversification,

suggesting this is a variable trait. In contrast, ancestral state reconstructions show

nearly 80% of the *Sinocyclocheilus* have scattered neuromast distributions.

Additionally, all Blind species showed a scattered neuromast pattern of distribution.

(figure 4*c*; black arrow).

Figure 4. Patterns of cephalic neuromast trait evolution of *Sinocyclocheilus* based

on ancestral state reconstruction on the maximum-likelihood tree. Ancestral state

reconstructions based on the Bayesian inference method is shown in figure S2-S4.

(a) Ancestral state reconstruction using stochastic character mapping for symmetry in

neuromasts pattern (Absolute/Slight-asymmetry). The tip colors represent state of the

extant species and each node indicates the ancestral state as a proportion of the tip state.

A, B, C and D showed on the extreme right of the figure indicate four major clades.

The bootstrap values > 95% are indicated as * on nodes.

(b) Ancestral state reconstruction using stochastic character mapping for handedness

bias (Left/Right-bias) on the phylogeny.

(c) Ancestral state reconstruction using stochastic character mapping for neuromast

expansion (Scattered/Serried) on the phylogeny.

(f) Evolution rates analysis

Among cavefish, the rate of neuromast evolution has only been determined for

Sinocyclocheilus. The rate of scatter for Blind species is higher compared to others

(mean distance and density - 3.5 and 6.8 times increased than Normal-Eyed species),

and the rate of numerical expansion of neuromasts is 1.2 times greater in Micro-Eyed

species (table 3). Normal-Eyed *Sinocyclocheilus* species demonstrated lower

evolutionary rates for every character state. A multiple-rate model of evolution

provided the best fit for mean distance between neuromasts, relative neuromast

density and neuromast distribution area. However, a single-rate model of evolution

provided the best fit for traits associated with neuromast numbers (table S6). Thus, the

evolutionary rate of cave-related neuromast patterns were highest in cave adapted

forms. Further, rates of evolution for surface area covered by SN neuromasts, and

right-sided neuromast numbers, are moderately increased in Blind and Micro-Eyed

species.

Table 2. Model-averaged rate parameters for the measured traits of neuromasts in

eyes-related morphs of *Sinocyclocheilus* species. **Bold font** = highest rate

Trait	Average rate		
	Blind	Micro-Eyed	Normal-Eyed
Number	155.1042	**179.4815**	150.6202
Left Number	**0.6395**	0.4106	0.4249
Right Number	**0.7095**	0.5056	0.5329
Area	102.8836	**106.2819**	101.9520
Mean distance	**34.4129**	11.7357	7.9001
Density	**362.2959**	64.3977	42.2944

(g) Behavioural correlates of neuromasts

We performed a series of behavioural assays, and found that Blind species

navigate markedly differently from eyed-species. Blind species have a well-

established wall-following behaviour, while sighted species utilize an entire arena

space (figure 5). We found that Blind species prefer to use the side with fewer

neuromasts for wall-following, but the side with more neuromasts for exploring (i.e.,

stimulation during VAB test). Micro-Eyed species having a left-bias in neuromast

distribution, and they preferred approaching the stimulation from the left-side (median

= 4) more frequently than right-side (median = 3; W = 612, p > 0.05). These species,

however, followed the wall on their right-side (median = 154) for a significantly

longer time compared to the left-side (median = 142; W = 967, p < 0.001; figure 5*c,g*;

table S5). Behavioural heatmaps depicting wall-following behaviour showed that

Blind species prefer swimming in a fixed route, but eyed-species swam in irregular

patterns (figure 5 *d-f'*). Wall following behaviour has also been characterized in

Astyanax mexicanus. In this species, blind morphs similarly use the side with fewer

neuromasts towards the wall, perhaps to use the contralateral side for detecting stimuli

important for feeding, communication and spatial learning [17, 42].

Figure 5. The experimental setup and results of the behavioural assays.

(a) Schematic diagram showing the assay system used to for the wall-following and

vibration attraction behaviour (VAB).

(b) The vertical view of the arena's schematic diagram. The pale yellow area within the

dotted line depict the near-wall belt ("Near wall"). The orange dotted line showed the

stimulating area. The black rectangle in center shows the Aeration pump.

(c) The boxplot showing a comparison of the wall following time of Blind (3 species;

n = 9), Micro-Eyed (4 species; n = 12) and Normal-Eyed (7 species; n = 21) morphs.

Their handedness was also considered and the Kruskal-wallis Rank Sum Tests showed

that the groups spent significantly different (U = 79.49, p < 0.001) times following the

wall. Levels of significance indicated by ***.

(d-f'). Representative result of a 10-min wall-following behaviour assay for three

species. Vertical view of the swimming path of Blind fish (d,d'; *S. tianeensis*, B-TB),

Micro-Eyed fish (e,e'; *S. microphthalmus*, M-TB) and Normal-Eyed fish (f,f'; *S.*

macrophthalmus, N-TB).

(d, e, f) Visualization of the traces of the fish swimming paths as depicted by EthoVision.

(d', e', f') Heatmaps generated from the trial results of the wall-following behaviour.

The color bar represents the total time (min) the fish stayed in one place. Warmer colors

denote areas with a longer time spent, whereas cooler colors denote areas of shorter

time spent.

(g) The results of the frequency of approach in stimulation area of VAB test.

Sinocyclocheilus is an ancient species complex, with an estimated age for the

clade of nearly 10-11 Mya [9, 62].They are distributed across an enormous karstic

area spanning three provinces in China [21]. One of the main forms of speciation

seems to be isolation of populations over long periods, and therefore geographic

speciation appears to have dominated this diversification [61]. This divergence

occurred during the geological uplift of the Yunnan-Kweichow Plateau together with

the aridification of China, which occurred during the Pliocene and the Pleistocene

[61]. The neuromast variability seen in these fishes is most likely attributable to the

collective influences of both selection- and drift-related evolutionary mechanisms that

have played on these fishes over long periods. However, the exact dynamics of the

evolution of neuromasts is still an intriguing question to be explored.

Conclusion

In *Sinocyclocheilus,* alongside some basic patterns, there is widespread variation in

cephalic neuromast patterns between species and to some extent within species. We

showed neuromast asymmetry with right-side enhanced for most species. For almost

all species, the dorsal neuromast numbers were lesser than either of the sides.

Furthermore, Regressed-Eyed (Blind and Micro-Eyed) species are more asymmetric

than the Normal-Eyed forms. Interestingly, we found the greatest degree of neuromast

variation and rate of evolution in Micro-Eyed species (living outside and inside caves

- troglophilic), this is possibly an adaptation for life in two markedly different habitats

types. Assays of swimming behaviour suggest a functional role of neuromasts in

habitat exploration. These patterns of neuromast distribution and swimming

behaviours are convergent with other cavefish lineages. The diversity of patterns and

variation can be explained by the deep evolutionary history associated with the karstic

region and the associated traits of this remarkable diversifications of fishes.

Abbreviations Section:

TB: Troglobitic; TP: Troglophilic; S: Surface; B: Blind; M: Micro-Eyed; N: Normal-

Eyed; CN: Canal neuromast; SN: Superficial neuromast; GXU: Guangxi University;

SL: Standard length; DEL: Delaunay Mean Distance (neuromast mean distance

coefficient); L: Left-Side; R: Right-Side; D: Dorsal-Side; PCC: asymmetry coefficient

1 = Pearson's Correlation Coefficient; OC: asymmetry coefficient 2 = Overlap-

Coefficient; PCA: Principal component analysis; AIC: Akaike information criterion;

mtDNA: Mitochondrial DNA; ND4: NADH-ubiquinone oxidoreductase chain 4; Cyt

b: cytochrome b gene.

Declarations

Acknowledgements: We thank the following individuals: members of EDD lab for

their cooperation and support; Chenghai Fu for assistance in the field; Gajaba Ellepola

for suggestions with the data analyses.

Funding

This work was supported by the (1) Startup funding for MM though Guangxi

University for fieldwork, lab work and student support. (2) National Natural Science

Foundation of China (#31860600) to JY for lab and fieldwork. (3) BC, TM and YL

537   were supported also by Innovation Project of Guangxi Graduate Education

538   YCBZ2021008. These funding bodies played no role in the design of the study and

539   collection, analysis, and interpretation of data or in the writing of the manuscript.

Conflicts of interest/Competing interests

542   We declare no conflicts of interest.

Ethics approval

545   The project was approved by Guangxi Autonomous Region Government and Guangxi

546   University Ethical Clearance Committee (protocol number: GXU-2021-125).

Consent to participate

549   Not applicable.

Availability of data and material

552   All the data used in the study are provided in the electronic supplementary material.

Authors' contributions

Conceptualization – MM, BC, JBG; Fieldwork – YL, BC, TM, MM, JY;

Experimentation and Lab Work – BC, XL, APR; Data Analyses – BC, TM, WD, XL;

Interpretation – All; Figures – BC, WD, TM; Writing original draft – BC, MM, JBG,

APR; Subsequent Drafts – All; Funding acquisition – MM, JY, BC, TM, YL;

Supervision – MM, JBG, JY.

References

1 Jeffery WR. 2001 Cavefish as a model system in evolutionary developmental biology. *Dev. Biol.* **231**,
1-12. (doi:10.1006/dbio.2000.0121)

2 Berti R, Durand JP, Becchi S, Brizzi R, Keller N, Ruffat G. 2001 Eye degeneration in the blind cave-
dwelling fish *Phreatichthys andruzzii*. *Can. J. Zool.* **79**, 1278-1285. (doi:10.1139/z01-084)

3 Jeffery WR. 2009 Regressive evolution in *Astyanax* cavefish. *Annu. Rev. Genet.* **43**, 25-47.
(doi:10.1146/annurev-genet-102108-134216)

4 Niemiller ML, Fitzpatrick BM, Miller BT. 2008 Recent divergence with gene flow in Tennessee cave
salamanders (Plethodontidae: *Gyrinophilus*) inferred from gene genealogies. *Mol. Ecol.* **17**, 2258-2275.
(doi:10.1111/j.1365-294X.2008.03750.x)

5 Jeffery WR, Strickler AG, Guiney S, Heyser DG, Tomarev SI. 2000 *Prox 1* in eye degeneration and
sensory organ compensation during development and evolution of the cavefish *Astyanax*. *Dev. Genes
Evol.* **210**, 223-230. (doi:10.1007/s004270050308)

6 Wilkens H, Strecker U. 2003 Convergent evolution of the cavefish *Astyanax* (Characidae, Teleostei):
genetic evidence from reduced eye-size and pigmentation. *Biol. J. Linn. Soc.* **80**, 545-554.
(doi:10.1111/j.1095-8312.2003.00230.x)

7 Yoshizawa M, Jeffery WR. 2011 Evolutionary tuning of an adaptive behavior requires enhancement
of the neuromast sensory system. *Commun. Integr. Biol.* **4**, 89-91. (doi:10.4161/cib.14118)

8 Yang JXet al.. 2016 The *Sinocyclocheilus* cavefish genome provides insights into cave adaptation.
BMC Biol. **14**, 1. (doi:10.1186/s12915-015-0223-4)

9 Mao TR, Liu YW, Meegaskumbura M, Yang J, Ellepola G, Senevirathne G, Fu C, Gross JB, Pie MR.
2021 Evolution in *Sinocyclocheilus* cavefish is marked by rate shifts, reversals, and origin of novel traits.
BMC Ecol. Evol. **21**, 45. (doi:10.1186/s12862-021-01776-y)

10 Cahn PH. 1989 The mechanosensory lateral line: neurobiology and evolution. *BioScience.* **40**, 215-
216. (doi:10.2307/1311373)

11 Webb JF, Shirey JE. 2003 Postembryonic development of the cranial lateral line canals and
neuromasts in zebrafish. *Dev. Dynam.* **228**, 370-385. (doi:10.1002/dvdy.10385)

12 Fritzsch B, Wahnschaffe U. 1983 The electroreceptive ampullary organs of urodeles. *Cell Tissue Res.*
229, 483-503. (doi:10.1007/BF00207693)

13 Smith S. 1996 Pattern formation in the urodele mechanoreceptive lateral line: What features can be
exploited for the study of development and evolution? *Int. J. Dev. Biol.* **40**, 727-733. (doi:10.1016/0141-
8130(96)01109-9)

14 Bird NC, Webb JF. 2014 Heterochrony, modularity, and the functional evolution of the
mechanosensory lateral line canal system of fishes. *EvoDevo.* **5**, 21. (doi:10.1186/2041-9139-5-21)

15 Gross JB, Gangidine A, Powers AK. 2016 Asymmetric facial bone fragmentation mirrors asymmetric
distribution of cranial neuromasts in blind Mexican cavefish. *Symmetry.* **8**, 118.
(doi:10.3390/sym8110118)

16 Elipot Y, Hinaux H, Callebert J, Rétaux S. 2013 Evolutionary shift from fighting to foraging in blind
cavefish through changes in the serotonin network. *Curr. Biol.* **23**, 1-10. (doi:10.1016/j.cub.2012.10.044)

17 Yoshizawa M. 2015 Behaviors of cavefish offer insight into developmental evolution. *Mol. Reprod.*

Dev. **82**, 268-280. (doi:10.1002/mrd.22471)

Jiang Y, Peichel CL, Torrance L, Rizvi Z, Thompson S, Palivela VV, Pham H, Ling F, Bolnick DI. 2017 Sensory trait variation contributes to biased dispersal of threespine stickleback in flowing water. *J. Evol. Biol.* **30**, 681-695. (doi:10.1111/jeb.13035)

Simon V, Hyacinthe C, Rétaux S. 2019 Breeding behavior in the blind Mexican cavefish and its river-dwelling conspecific. *PLoS ONE.* **14**, e0212591. (doi:10.1371/journal.pone.0212591)

Ma ZQ, Herzog H, Jiang YG, Zhao YH, Zhang DY. 2020 Exquisite structure of the lateral line system in eyeless cavefish *Sinocyclocheilus tianlinensis* contrast to eyed *Sinocyclocheilus macrophthalmus* (Cypriniformes: Cyprinidae). *Integr. Zool.* **15**, 314-328. (doi:10.1111/1749-4877.12430)

Romero A, Zhao YH, Chen XY. 2009 The hypogean fishes of China. *Environ. Biol. Fishes.* **86**, 211-278. (doi:10.1007/s10641-009-9441-3)

Zhao YH, Zhang CG. 2009 *Endemic fishes of Sinocyclocheilus (Cypriniformes: Cyprinidae) in China -- Species diversity, cave adaptation, systematics and zoogeography.* Beijing: Science Press.

Webb JF, Northcutt RG. 1997 Morphology and distribution of pit organs and canal neuromasts in non-teleost bony fishes. *Brain Behav. Evol.* **50**, 139-151. (doi:10.1159/000113328)

Powers AK, Berning DJ, Gross JB. 2020 Parallel evolution of regressive and constructive craniofacial traits across distinct populations of *Astyanax mexicanus* cavefish. *J. Exp. Zool. B Mol. Dev. Evol.* **334**, 450-462. (doi:10.1002/jez.b.22932)

George PL, Borouchaki H. 1998 *Delaunay triangulation and meshing: application to finite elements.* Paris: Hermès.

Aurenhammer F, Klein R, Lee D-T. 2012 *Voronoi diagrams and delaunay triangulations.* World Scientific.

Lee DT, Schachter BJ. 1980 Two algorithms for constructing a delaunay triangulation. *Int. J. Parallel Program.* **9**, 219-242. (doi:10.1007/BF00977785)

Palmer AR, Strobeck C, Chippindale AK. 1993 Bilateral variation and the evolutionary origin of macroscopic asymmetries. *Genetica.* **89**, 201-218. (doi:10.1007/BF02424514)

Gorlewicz A, Krawczyk K, Szczepankiewicz AA, Trzaskoma P, Mulle C, Wilczynski GM. 2020 Colocalization colormap – an ImageJ Plugin for the quantification and visualization of colocalized signals. *Neuroinformatics.* **18**, 661-664. (doi:10.1007/s12021-020-09465-9)

Zeitvogel F, Schmid G, Hao L, Ingino P, Obst M. 2016 ScatterJ: An ImageJ plugin for the evaluation of analytical microscopy datasets. *J. Microsc.* **261**, 148-156. (doi:10.1111/jmi.12187)

Cordelières F, Bolte S. 2008 JACoP v2.0: improving the user experience with co-localization studies. *ImageJ User & Developer Conference.* 174-181.

Wang YS, Wu QH, Hu MQ, Liu B, Chai ZY, Huang R, Wang Y, Xu HD, Zhou L, Zheng LH, *et al.* 2017 Ligand- and voltage-gated Ca2+ channels differentially regulate the mode of vesicular neuropeptide release in mammalian sensory neurons. *Sci. Signal.* **10**, eaal1683. (doi:10.1126/scisignal.aal1683)

Bolte S, Cordelières FP. 2007 A guided tour into subcellular colocalization in light microscopy. *J. Microsc.* **224**, 213-232. (doi:10.1111/j.1365-2818.2006.01706.x)

Dunn KW, Kamocka MM, McDonald JH. 2011 A practical guide to evaluating colocalization in biological microscopy. *Am. J. Physiol. Cell Physiol.* **300**, C723-C742. (doi:10.1152/ajpcell.00462.2010)

Manders EM, Stap J, Brakenhoff GJ, van Driel R, Aten JA. 1992 Dynamics of three-dimensional replication patterns during the S-phase, analysed by double labelling of DNA and confocal microscopy. *J. Cell Sci.* **103**, 857-862. (doi:10.1242/jcs.103.3.857)

Zinchuk V, Grossenbacher Zinchuk O. 2009 Recent advances in quantitative colocalization analysis:

focus on neuroscience. *Prog. Histochem. Cytochem*. **44**, 125-172. (doi:10.1016/j.proghi.2009.03.001)

Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP. 2001 Bayesian inference of phylogeny and its impact on evolutionary biology. *Science*. **294**, 2310-2314. (doi:10.1126/science.1065889)

Revell LJ. 2012 Phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol*. **3**, 217-223. (doi:10.1111/j.2041-210X.2011.00169.x)

Planidin NP, Reimchen TE. 2021 Ecological predictors of lateral line asymmetry in stickleback (*Gasterosteus aculeatus*). *Evol. Ecol*. **35**, 609-629. (doi:10.1007/s10682-021-10117-w)

Sharma S, Coombs S, Patton P, de Perera TB. 2009 The function of wall-following behaviors in the Mexican blind cavefish and a sighted relative, the Mexican tetra (*Astyanax*). *J. Comp. Physiol. A*. **195**, 225-240. (doi:10.1007/s00359-008-0400-9)

Patton P, Windsor S, Coombs S. 2010 Active wall following by Mexican blind cavefish (*Astyanax mexicanus*). *J. Comp. Physiol. A*. **196**, 853-867. (doi:10.1007/s00359-010-0567-8)

Burt de Perera T, Braithwaite VA. 2005 Laterality in a non-visual sensory modality — the lateral line of fish. *Curr. Biol*. **15**, 241-242. (doi:10.1016/j.cub.2005.03.035)

Jiang YG, Fu JC, Zhang DY, Zhao YH. 2016 Investigation on the lateral line systems of two cavefish: *Sinocyclocheilus Macrophthalmus* and *S. Microphthalmus* (Cypriniformes: Cyprinidae). *J. Bionic Eng*. **13**, 108-114. (doi:10.1016/S1672-6529(14)60164-5)

McGaugh SE, Weaver S, Gilbertson EN, Garrett B, Rudeen ML, Grieb S, Roberts J, Donny A, Marchetto P, Gluesenkamp AG. 2020 Evidence for rapid phenotypic and behavioural shifts in a recently established cavefish population. *Biol. J. Linn. Soc*. **129**, 143-161. (doi:10.1093/biolinnean/blz162)

Dezfuli BS, Magosso S, Simoni E, Hills K, Berti R. 2009 Ultrastructure and distribution of superficial neuromasts of blind cavefish, *Phreatichthys andruzzii*, juveniles. *Microsc. Res. Tech*. **72**, 665-671. (doi:10.1002/jemt.20714)

Powers AK, Kaplan SA, Boggs TE, Gross JB. 2018 Facial bone fragmentation in blind cavefish arises through two unusual ossification processes. *Sci. Rep*. **8**, 7015. (doi:10.1038/s41598-018-25107-2)

Soares D, Niemiller ML. 2020 Variation in cephalic neuromasts surface and cave-dwelling fishes of the family amblyopsidae (teleostei: percopsiformes). *J. Cave Karst Stud*. **82**, 198-209. (doi:10.4311/2019LSC0115)

Trokovic N, Herczeg G, Ghani NIA, Shikano T, Merilä J. 2012 High levels of fluctuating asymmetry in isolated stickleback populations. *BMC Evol. Biol*. **12**, 115. (doi:10.1186/1471-2148-12-115)

Soares D, Niemiller ML. 2020 Extreme Adaptation in Caves. *Anat. Rec*. **303**, 15-23. (doi:10.1002/ar.24044)

Krishnan J, Rohner N. 2017 Cavefish and the basis for eye loss. *Philos. Trans. R. Soc. Lond., B. Biol. Sci*. **372**, 20150487. (doi:10.1098/rstb.2015.0487)

Fernandes VFL, Macaspac C, Lu L, Yoshizawa M. 2018 Evolution of the developmental plasticity and a coupling between left mechanosensory neuromasts and an adaptive foraging behavior. *Dev. Biol*. **441**, 262-271. (doi:10.1016/j.ydbio.2018.05.012)

Coombs S, Janssen J, Webb JF. Year Diversity of lateral line systems: evolutionary and functional considerations. In: J. Atema, R. R. Fay, A. N. Popper, W. N. Tavolga, editors. Sensory Biology of Aquatic Animals; 1988; New York, NY: Springer New York; 1988. 553-593.

Montgomery JC, Coombs S, Baker CF. 2001 The mechanosensory lateral line system of the hypogean form of *Astyanax Fasciatus*. *Environ. Biol. Fishes*. **62**, 87-96. (doi:10.1023/A:1011873111454)

Yamamoto Y, Byerly MS, Jackman WR, Jeffery WR. 2009 Pleiotropic functions of embryonic *sonic hedgehog* expression link jaw and taste bud amplification with eye loss during cavefish evolution. *Dev.*

Biol. **330**, 200-211. (doi:10.1016/j.ydbio.2009.03.003)

55 Zhao YH, Zhang CG. 2006 Past research and future development on endemic Chinese cavefish of

the genus *Sinocyclocheilus* (Cypriniformes, Cyprinidae). *Acta Zootaxonomica Sinica*. **31**, 769-777.

56 He Y, Chen XY, Xiao TQ, Yang JX. 2013 Three-dimensional morphology of the *Sinocyclocheilus*

hyalinus (Cypriniformes : Cyprinidae) horn based on synchrotron X-ray microtomography. *Zool. Res*.

34, 128-134. (doi:10.11813/j.issn.0254-5853.2013.E4-5.E128)

57 Teyke T. 1990 Morphological differences in neuromasts of the blind cave fish *Astyanax hubbsi* and

the sighted river fish *Astyanax mexicanus*. *Brain Behav. Evol*. **35**, 23-30. (doi:10.1159/000115853)

58 Wark AR, Peichel CL. 2010 Lateral line diversity among ecologically divergent threespine

stickleback populations. *J. Exp. Biol*. **213**, 108-117. (doi:10.1242/jeb.031625)

59 Zhao YH, Watanabe K, Zhang CG. 2006 *Sinocyclocheilus donglanensis*, a new cavefish (Teleostei:

Cypriniformes) from Guangxi, China. *Ichthyol. Res*. **53**, 121-128. (doi:10.1007/s10228-005-0317-z)

60 Culver D, Pipan T. 2019 *The biology of caves and other subterranean habitats*.

61 Xiao H, Chen SY, Liu ZM, Zhang RD, Li WX, Zan RG, Zhang YP. 2005 Molecular phylogeny of

Sinocyclocheilus (Cypriniformes: Cyprinidae) inferred from mitochondrial DNA sequences. *Mol.*

Phylogenet. Evol. **36**, 67-77. (doi:10.1016/j.ympev.2004.12.007)

62 Li ZQ, Guo BC, Li JB, He SP, Chen YY. 2008 Bayesian mixed models and divergence time estimation

of Chinese cavefishes (Cyprinidae: *Sinocyclocheilus*). *Chinese Science Bulletin*. **53**, 2342-2352.

(doi:10.1007/s11434-008-0297-2)