# Predicting proprioceptive cortical anatomy and neural coding with topographic autoencoders

Max Grogan[1,¶], Kyle P. Blum[2,¶], Yufei Wu[1,¶], J. Alex Harston[1], Lee E. Miller[2] & A. Aldo Faisal[1,3,*]

[1] Department of Bioengineering, Imperial College London, London, UK

[2] Department of Physiology, Northwestern University, Illinois, USA

[3] Institute of Artificial & Human Intelligence, University of Bayreuth, Bayreuth, Germany

* Corresponding author

E-mail: A.faisal@imperial.ac.uk

¶ These authors contributed equally to the work

## Abstract

Proprioception is one of the least understood senses, yet fundamental for the control of movement.

Even basic questions of how limb pose is represented in the somatosensory cortex are unclear. We

developed a topographic variational autoencoder with lateral connectivity (topo-VAE) to compute

a putative cortical map from a large set of natural movement data. Although not fitted to neural

data, our model reproduces two sets of observations from monkey centre-out reaching: 1. The

shape and velocity dependence of proprioceptive receptive fields in hand-centered coordinates

despite the model having no knowledge of arm kinematics or hand coordinate systems. 2. The

distribution of neuronal preferred directions (PDs) recorded from multi-electrode arrays. The

model makes several testable predictions: 1. Encoding across the cortex has a blob-and-pinwheel-

type geometry of PDs. 2. Few neurons will encode just a single joint. Our model provides a

principled basis for understanding of sensorimotor representations, and the theoretical basis of

neural manifolds, with applications to the restoration of sensory feedback in brain-computer

interfaces and the control of humanoid robots.

## Author Summary

It is well established that proprioception is essential for effective for motor control, yet our

understanding of proprioceptive coding in somatosensory cortex is far behind that of more

established sensory modalities such as vision and touch. Here, we use unsupervised learning of a

deep neural network imposed with biological constraints to reproduce coding properties of

proprioceptive neurons in area 2 of primary somatosensory cortex. With this model, we

demonstrate that the tendency for area 2 neurons in close physical proximity to share similar

directional tuning can be explained by local topographic organisation driven by Mexican hat

lateral effects, a phenomenon that is otherwise unobservable with the spatial resolution of

available electrode microarray recordings. This provides, to the best of our knowledge, the first

evidence of local topographic organisation in area 2 proprioceptive neurons. We also predict that

the structure of proprioceptive receptive fields are typically multi-joint rather than single-joint in

nature and provide, to the best of our knowledge, and find that our model best reproduces the

tuning properties of neural data when training on natural kinematic recordings, rather than those

from a stereotyped task, underlining the importance of training on data that reflect the natural

distribution of sensory stimuli.

# Introduction

Somatosensation includes the familiar sense of touch, provided by receptors in the skin, and proprioception, the much less consciously perceived sense that informs us about the pose, motion and associated forces acting on our limbs. While the former has received much scientific attention, proprioception is often overlooked, yet this modality of sensory feedback is essential for our ability to plan, control and adapt movements. In engineering, the control of robotic movement would be impossible if the controller did not know the location of its actuators; correspondingly in human motor control, optimal feedback control theory is the preeminent explanation for the computations underlying limb control [1,2]. Moreover, individuals with proprioceptive neurological deficits, such as patient IW, have profound motor deficits even in the presence of vision and an intact motor system [3,4]. Similarly, recent major developments in neuroprosthetics are centred on restoring sensory feedback as well as limb motion through bi-directional interfaces [5] and will likely require not only touch but also proprioceptive feedback to restore functional capability [6]. Thus, understanding proprioceptive encoding is essential for restoration of both motor action and sensory function in clinical rehabilitation [7].

Unlike hearing or vision, proprioceptive afferent pathways originate not from a single organ but from diverse families of mechanoreceptors within muscles, tendons, joints, and the skin itself [8]. Proprioceptive information ascends within the dorsal column pathway through the dorsal root ganglia, the dorsal column nuclei, and thalamus before arriving in the primary somatosensory cortex (S1). Crucially, neurons are somatotopically organised throughout this pathway, meaning that neighbouring neurons encode stimuli from closely related parts of the body. This gives rise to

the sensory homunculus, which results from the ordered mapping of tactile representations of the

body's surface across the cortical surface [9].

Although proprioception is generally acknowledged to be critical to motor behaviour, the

corresponding proprioceptive maps - particularly that of primate area 3a, but also the mixed

modality area 2 - are much less distinct and well understood than those of the touch (areas 1 and

3b). We know that the proprioceptive and tactile systems often encode overlapping information,

e.g. mechanoreceptors in our skin and interosseous membranes respond to deformation and

vibration, contributing to the sense of body position and movement [3]. Indeed, the firing rates of

somatosensory neurons with cutaneous receptive fields in area 2 can be used to decode limb

movement as accurately as those with muscle fields [10]. While proprioceptive cortical areas are

critical for our ability to generate goal-directed complex behaviour it is unclear what properties of

proprioceptive cortical coding facilitate this capability. Crucially, we lack an accepted hypothesis

about the computational principles that drive the mapping of proprioceptive arm representations

onto the cortex.

The limited nature of our understanding of proprioceptive neural representations in the brain has

two major causes. First is the difficulty recording with many electrodes from sulcal proprioceptive

areas in primates [11],  and second is the difficulty delivering independent proprioceptive stimuli

in comparison to other senses, such as vision [12] and touch [13]. To overcome these limitations,

we combined computational modelling and natural movement kinematics data to test hypotheses

of organisational mechanisms that are currently beyond the capability of experimental recording

techniques.  This combination of experiment and theory has been useful for explaining population-

level coding in other sensory modalities, such as vision [14], olfaction, and touch [15,16], but has

only recently begun to be applied to proprioception [17]. Much of the computational theoretical

work on other senses has either focused on predicting specific neural coding features from natural

sensory statistics [15,18],  or has ignored the details of neural coding and focused on the spatial

distribution of stimulus representation across the cortex [19].

Our chosen system was the proprioception of the right arm (Fig. 1.A) for which we had collected

natural behavioural data from daily life in humans (food preparation, eating, etc; Fig. 1.B and

Supplemental Fig. S1) as well as constrained, planar centre-out reaching in both human and

monkeys (Fig. 1.C). This data describes the joint angle kinematics of the of the body over time

and thus stands in for the proprioceptive sensory state. We aimed to relate kinematics to

proprioceptive encoding with modelling and comparisons to existing single-unit recordings from

monkeys. We formulated a novel computational model that predicts both neural coding of single

neurons and spatial organisation of these neurons across the cortex (Fig. 1.D-G). We chose a small

set of general computational elements that reflect principles and mechanisms found in sensory

systems but have not been previously unified, which are as follows: Models should 1) use the

information maximisation principle, which postulates that efficient sensory representations in the

brain reflect natural sensory statistics, and implies that they are essential in shaping neural

representations, 2) be stochastic, generative, and decodable, to reflect the natural variability in the

data, produce neural activity to mimic that of the biological system, and offer the means to

reconstruct the relevant original sensory input from its output, 3) implement (in Marr's sense [20])

neural computations that are performed by locally interacting neurons through synaptic

interactions, rather than by an abstract computational machine.

In the context of proprioception, these principles require that our model should learn how to translate proprioceptive inputs from movements of the body into a latent representation of proprioception (which is read out as neural firing). In addition to performing efficient feature learning of a spike-based code, the model should have minimal human induction bias, except for specific characteristics relating to the principles laid out above. By training the model on natural movement stimuli, it should learn representations that can maximise information about the natural environment with the limited bandwidth of neurons that is available at any given encoding step, thereby implementing efficient coding [21–23]. Given these requirements, we chose to use an unsupervised deep neural network model, a variational autoencoder (VAE) [24] (Fig. 1.D) to perform efficient feature learning of proprioceptive stimuli. The primary training objective of an autoencoder is to reconstruct the input stimuli, enabling the model to learn in an unsupervised manner. Furthermore, the bottleneck of the VAE model parameterises a latent Bernoulli distribution to imitate the activity of spiking neurons.

**A**

**B** Motion Sensors

Natural Movement
(e.g. Kitchen Scenario)

**C**

**D**

x → Encoder → q(z) → Sampling → z ~ q(z) → Decoder → x̂

Input          Distribution of Latent Variable          Sampled Latent Variable          Reconstruction

**E**

x → MLP Encoder → [M × M grid] → Sample + Flatten → Spike Count → Linear Decoder → x̂

Stimuli          M × M Cortical Map (Bernoulli Neurons)          Spike Count          Reconstruction

**F**

Ψ(P,:)          Ψ(P,Q)

1 unit = 1 cell

P

M × M Cortical Map

Lateral inhibition
Lateral excitation
No effect

**G**

Ψ

d

d = 2σ

**H**

Natural Movement (Human) ──Train──┐
                                   ↓
Planar Movement (Human) → Model → Predict → Representation of Planar Motion → Compare → S1 Recording (Macaque)

***Figure 1: Task Context and model architecture. (A)*** *9-dimensional arm movement data acquired*

*from motion capture suit is represented by Euler angles between body segments, following the ISB*

*Euler angle extractions* [25] *in a ZXY coordinate. **(B)** Illustrative example of movements carried*

*out in the natural scenario (cf. Fig. S1 for further examples). **(C)** Illustrative example of planar*

*centre-out reaching movements. **(D)** General architecture of a VAE: Inputs are encoded as a latent*

*distribution, from which samples are decoded to reconstruct the original input. **(E)** Topo-VAE*

*architecture. Our cortical neural representation is modelled by an 80x80 cortical grid of artificial*

*neurons in the latent layer q(z). In contrast to conventional VAEs which model these latent neurons*

*as multi-dimensional Gaussian random variables, we used Poisson random variables. Input*

*stimuli drive these model neurons via a multi-layer perception (MLP) encoder network (2 layers*

*of 500 neurons each). A linear decoder is applied to the spike counts emitted from the cortical map*

*within a time interval Δt, to reconstruct the sensory input stimuli. **(F)** To embed the model neurons*

*in a cortex-like topographic context we use a Mexican-hat lateral effect between the latent layer*

*neurons. **(G)** This interaction $\Psi(P,Q)$ is a function of the Euclidean distance $d = ||P\text{-}Q||$ between*

*a pair of neurons P,Q and is characterised by a length scale σ. Nearby neurons are excited,*

*intermediate-range neurons are inhibited and there is no effect on distant neurons. **(H)** Flow of*

*natural and planar movement data (joint angle velocities) in this work.*

In following with our third outlined principle, and the observation that spatial organization is

relevant to cortical coding, we introduce a 2-dimensional structure to the latent ("bottleneck") layer

of the VAE, representing a simplified proprioceptive cortex (Fig. 1.E). A conventional "vanilla"

VAE model will capture the properties of encoded sensory features but would be devoid of any of

the spatial properties of cortical neurons that are critical for understanding how sensory stimuli are

represented across the cortical surface. Incorporating spatial relationships in neural coding models

has not been well investigated, yet anatomical structure and function in neural computation are

fundamentally linked by biophysical constraints [26]. We therefore also implemented a

topographical mechanism, using a lateral interaction term between the units in the VAE's latent

layer (see Methods for detailed motivation and mathematical description; Fig. 1.F). This term

corresponds biologically to a spatial distribution of short-range excitatory and longer-range

inhibitory synaptic interactions between neurons (Fig. 1.G).

We call our model the "topo-VAE" and compared it with existing recordings from area 2 of

somatosensory cortex. It could equally well be applied to other proprioceptive areas such as 3a or

even 5. Models with its basic form could be applied to other sensory modalities, including touch,

vision, and hearing. Here, the approach has enabled us to uncover novel insights into

proprioceptive coding by linking anatomical structure and function so that we can test our model's

predictions using the sparse set of obtainable data.

# **Results**

We developed a novel model of cortical sensory representation (see Methods for details) that

predicts both function and structure – the topo-VAE model. We trained, tested and validated our

model as follows (Fig. 1.H): We trained our topo-VAE model on natural daily human arm

movement kinematics and explored the emergent proprioceptive representations in its cortical

layer (i.e. the latent layer of the topo-VAE). After training, we used the model to generate neural

responses by providing it kinematic data from a centre-out reaching task performed by human

subjects. We then compared the properties of these simulated neural activities (i.e. the generated

spike trains) to those of S1 neurons which were previously recorded from monkeys performing a

planar centre-out reaching task. The human and monkey reaching data were rescaled so that the

biomechanical differences between human and monkey arm movements are kept small. We

characterised the proprioceptive neural representations at two levels of description: first, we

considered the spatial tuning curves of individual neurons and second, and second, we considered

the organisation of tuning preference maps across our model's cortical surface.

We used three sets of measurements to summarise the neural coding properties of S1 neurons for

this centre-out task: 1) the preferred direction (PD) of single-neuron firing rates, 2) features of the

tuning surface defined by firing rate modulation as a function of both direction and speed of

movement, and 3) the overall distribution of firing rates throughout the behaviour. First, we

analysed PDs of neurons in the recordings and in our modelling (Fig. 2). The PD refers to the

movement direction of the hand (when performing the centre-out reaching task) in which a neuron

was most active. We found that generally, modelled neurons exhibited PD tuning surfaces that

were similar in appearance to those of recorded neurons (Fig. 2.A,B respectively). At the

population level, while there was a statistically significant difference ($p<0.005$; Kolmogorov-

Smirnov test) between the distribution of PDs of modelled and recorded neurons (Fig. 2.C), the

distributions were similar in that they were both bimodal (Supplemental Fig. 2), with a bias

towards the 120º/300º reach axis (Fig. 2.D). Previous experimental studies have suggested such

coding biases are a natural consequence of musculoskeletal constraints in the arm [27,28], which

cause the natural distribution of right arm reach movements to favour the 120º/300º reach axis.

Under the efficient coding hypothesis, this becomes reflected in the coding properties of the

sensory neurons. To test this theory, we trained a second model on stereotyped movement stimuli

from the centre-out reach task only. We found that this new model was able to reconstruct natural

movement data almost as well as the natural-trained model (0.80 ±0.01 vs 0.96 ±0.008, planar-

trained vs natural-trained; $R^2$ scores; p<0.005; Wilcoxon) and produced a PD distribution that was

also comparable to the recorded neuron distribution (0.241 vs 0.248, planar-trained vs natural-

trained; Jensen-Shannon divergence). However, unlike the natural-trained model, we observed a

reach axis bias that was perpendicular to that of recorded neurons (Fig. 2.D,E). This suggests that

certain features of proprioceptive coding are only emergent when the coding is conditioned on the

natural distribution of stimuli.

While preferred direction is an informative neural correlate, we looked to better compare tuning

properties by extracting geometric features from the tuning surfaces of modelled and recorded

neurons and evaluating how different model properties affect similarity. Two such features were

compared (Fig. 3.A): Half-peak widths measure how sharp the tuning of a neuron is, where smaller

half-peak widths indicate sharper tuning, and velocity gradients measure how sharply the firing

rate of a neuron changes as a function of the endpoint velocity in the direction of tuning (note that

the velocity gradient measurements were adjusted to control for differences in mean firing rate

between modelled and recorded neurons). While there was a statistically significant difference

between the feature distributions of modelled and recorded neurons (p<0.005 for both features and

all model variations; Kolmogorov-Smirnov test), Jensen-Shannon divergence between

distributions was minimised (Fig. 3.B,C) when the model included lateral effects and was trained

on natural movement data. This would suggest that both model constraints are important for

reproducing the tuning properties of recorded neurons. One concern was that the distributions of

tuning curve geometry between modelled and recorded neuron populations were similar only

because of the Poisson GLM fitting process -- perhaps fitting a Poisson GLM to any random data

would produce distributions like these. To verify that this was not the case, we shuffled the firing

rate data from modelled neurons to remove any correlation between endpoint velocities and firing

rates and trained an additional Poisson GLM on this data. The shuffling produced a very different

pair of distributions for the geometric features, with flat, poorly tuned curves - suggesting that any

similarities we saw between the original unshuffled curves were in fact due to similarities in the

underlying structure of the modelled and recorded neuron data. This provides an empirical lower

bound of the similarity between feature distributions, and would suggest that similarity between

the distributions of modelled and recorded neurons is not trivially the result of using a GLM (Fig.

3.B,C).

Finally, we examined endpoint-velocity modulation of firing rate distributions, which were well

fit by Gamma curves for both modelled (Fig. 3.D) and recorded neurons (Fig. 3.E; $R^2 > 0.95$ for

equal numbers of neurons). While the topo-VAE generates spike count distributions over an

arbitrary time window (cf. Fig. 3.D, x-axis), the functional form and relative change of the

distributions with altered velocity are not arbitrary. Therefore, we compared the distributions at

different endpoint velocities and found that they differed significantly in both modelled and

recorded neurons ($p<0.005$; two-sided Wilcoxon signed-rank test), indicating velocity dependence

in both the modelled and recorded neurons.

In summary, modelled neurons in the topo-VAE exhibit various tuning properties such as preferred

direction and endpoint-velocity modulation of firing rates which are structurally similar to those

of recorded neurons. While the population statistics of these tuning properties differ significantly

between modelled recorded neurons, the combined lateral effects that constrain our model and the

training data that reflect the natural distribution of arm movement, increase the similarity, all

without fitting our model to any neural data.

Figure 2. Comparing tuning of modelled and recorded neurons. *(A)* *Tuning maps of example*

*area 2 neurons during planar movements.* *(B)* *Tuning maps from example neurons observed in the*

*latent layer of the topo-VAE under planar movements, matched to example neurons in A.* *(C)*

*Circular density histograms of preferred directions in our topo-VAE model trained on natural data*

*(n=6400) (blue) and neurons recorded from area 2 of 3 monkeys (n=383) (orange). Note, that the*

*neural models were not fitted to monkey neural data, but directly predicted from the statistics of*

 *natural human body kinematics. **(D)** Frequency histogram of mode bias in PD distributions for*

*topo-VAE models trained on natural data (blue) and stereotyped centre-out reach data (n=10 for*

*each condition) vs mode bias of PD distribution in recorded neurons (Orange line). **(E)** Circular*

*density histograms of preferred directions in our topo-VAE trained on stereotyped centre-out*

*reach data (n=6400) (green) and neurons recorded from area 2 of 3 monkeys (n=383) (orange).*

***Figure 3. Comparing tuning surface features and firing rate distributions between modelled***

***and recorded neurons.*** *(**A)** Tuning surface from a single modelled neuron, illustrating two*

*measures we computed to summarise neuronal modulation with velocity: half-peak width of spatial*

*tuning (at maximum end-point speed) and velocity gradient. **(B)** Jensen-Shannon divergence*

*between the distributions of half-peak widths in modelled neurons trained under different*

*conditions (with (+) and without (-) lateral effects, trained on natural data, trained on planar data)*

*(6400 neurons, 10 models per condition) and recorded neurons (383 neurons). The shuffled*

*control is the same as the natural trained model with lateral effects, except the neural activity is*

*shuffled prior to training the Poisson-GLM in the tuning surface calculation step. **(C)** Equivalent*

*plot to (B) for velocity gradient distributions. **(D)** Firing rate modulation (peak rate minus mean*

*rate) by endpoint-velocity across all reach directions in modelled (6400 neurons) and **(E)** in*

*recorded neurons (383 neurons). Histograms represent actual distributions, whereas lines*

*represent fitted gamma distributions ($R^2$>0.95 for both; Kolmogorov-Smirnov test).*

Having demonstrated similarity in the tuning statistics between modelled and recorded neurons,

we next compared the spatial relationship of PDs in our model neurons to those in the monkey

brain within the constraints of the 400 μm spacing of the recording electrodes (Fig. 4). While

neurons recorded at two adjacent electrodes corresponded to a distance that was substantially

beyond the effective range of the Mexican hat distance function, many electrodes recorded data

from more than one distinguishable neuron. The distance between these neurons was within the

local neighbourhood of neurons in our model. We could thus compare the similarity of neural

coding properties (specifically, the PDs) between neurons recorded on the same electrode to the

similarity between neurons on different electrodes (cf Fig. 4.A). In modelled neurons, same-

electrode PD differences followed an exponential distribution, such that the probability that two

neurons had similar (within 30 degrees) PDs was much higher than chance (p=0.77; two-sided

Wilcoxon signed-rank test; Fig. 4.C., blue histogram). Conversely, the probability that two neurons

on the same electrode had nearly opposite PDs (between 150º and 180º) was very low (p=0.03;

two-sided Wilcoxon signed-rank test; Fig. 4.C orange histogram). We then ran the same analysis

on recorded neurons and found that the PD difference distributions appeared similar to those

predicted by the model (Fig. 4.C). Given this similarity, we use the topo-VAE model to directly

predict the spatial organization of PD tuning across the proprioceptive cortex, which the 400um

electrode spacing lacks the adequate spatial resolution to observe. The model predicts that PDs are

clustered in blob-like structures of similar PDs with boundary regions where the PDs rotate

smoothly toward neighbouring directions (Fig. 4.D). This arrangement is interrupted by small,

sparsely spaced "pinwheel" regions, representing all PDs in a small neighbourhood, which is

consistent with same-electrode similarity of Fig. 4.B,C.

In the case of the model, different choices of the neighbourhood hyperparameter, $\sigma$, led to

differences in the exponential distributions of PD difference (Supplemental Fig. S3.B,C). Small

neighbourhood values ($\sigma$=1, Supplemental Fig. S3.B,C, left) led to more uniform distributions,

whereas larger neighbourhood values ($\sigma$={2,3}, Supplemental Fig. 3.B,C, middle and right) led to

exponential relationships like those we found in the recorded data, with the closest match to the

model being $\sigma = 2$. Note, that for our simple cortical grid model we only considered integer values

of $\sigma$.

*Figure 4. Predicting the topography of proprioceptive cortex. (A) Illustration of pairwise*

*comparisons of preferred directions recorded on same (blue) and different (orange) electrodes,*

*performed in recorded and modelled neurons. Neuron A and B are recorded from the same*

*electrode, Neuron C is recorded from a distant electrode. (B) Distribution of PD difference for*

*same electrode (blue histograms) and different electrode (orange histograms) comparisons in*

*modelled neurons and (C) recorded neurons (length scale σ = 2); error bars are* standard

deviation. *(D) The PD map in a topo-VAE with σ = 2, the hyperparameter value which well*

*approximates the recorded neuron topography (cf. Supplemental Fig. 3 for other values of length*

*scale σ).*

Next, we examined which specific elements of the lateral interaction were important for

reproducing the recorded data. We verified through model ablation the importance of the shape of

the Mexican hat function (blue curve in Fig. 5.A), which determines the strength of excitatory and

inhibitory connections as a function of distance between two neurons. We found no lateral effect

function with a significant effect on input data reconstruction accuracy when compared to a model

with no lateral effects (P<0.005; ANOVA). Flipping the function upside down (i.e., short range

inhibition and intermediate range excitation, yellow curve in Fig. 5.A) eliminated the obvious

topographic structure (Fig. 5.B, yellow box). Similarly, removing either the excitatory (purple

curve in Fig. 5.A) or inhibitory component (red curve in Fig 5.A) produced results unlike the

localised topographic structure of our primary model against which the recorded data was validated

(Fig. 5.B, purple and red boxes, respectively), suggesting that the specific Mexican hat shape was

important for reproducing biological results.

***Figure 5. Mexican hat lateral effects are essential for reproducing spatial organisation. (A)***

*Alternative lateral effect functions tested in control experiments, compared to the Mexican hat*

*(blue line). **(B)** resulting PD maps for each function (colour matched to 5.D). The PD map for the*

*inverted Mexican hat model is not shown due to space constraints, but produces no topographic*

*structure, much like models with no lateral effect or inhibition only.*

In addition to predicting the spatial organisation of proprioceptive coding, our model enables

testable predictions about the representation of movements involving multiple joints (wrist, elbow,

shoulder) across the neural population. To obtain a measure of how the 3 joints (Fig 6.A) are

encoded across the cortical surface relative to each other, we found the average correlation with

firing rate for all the degrees of freedom of each joint. All joints were evenly represented across

the whole population (Fig 6.B), but no neurons correlated with only one joint. Many modelled

neurons were correlated even with non-adjacent joints, e.g. elbow-wrist or even all three joints,

possibly a result of the natural covariance patterns that exist between joints. For pairwise joint

sensitivity comparisons see scatter plots (Fig. S3).

Fig. 6.C shows in more detail, the response strength of our cortical model neurons for individual

degrees of freedom. The maps had a spatial pattern of ripple-like transitions between positive

(white) and negative (black) correlations (Fig. 6.C), producing topographic clusters of neurons that

are sensitive to particular joints (Fig. 6.D). The scale of these clusters approximately matched the

that of the Mexican-hat lateral connection range. Regions of pure red, green, or blue in Fig. 6D

would correspond to neurons responsive to only shoulder, elbow or wrist respectively. Instead,

colours that are the result of blending two or more joints predominate.

**Figure 6. Sensitivity of latent layer neurons in topo-VAE to different input dimensions. (A)** *Joint angle axes, 3 per joint.* **(B)** *3D plot showing the relative sensitivity of each neuron to joint inputs. Sensitivity is defined as the sum of correlations across X/Z/Y axes of all joints for a given neuron (colour normalised by maximum values per joint) (n=6400).* **(C)** *Correlations between each dimension of input joint motion and neural response.* **(D)** *Joint sensitivity map where each pixel represents a neuron and the pixels RGB values (colour wheel inset) are reflecting the correlation with shoulder ("S"), elbow ("E"), and wrist ("W") angular velocity, respectively (colours normalised by maximum value across joints).*

# Discussion

Here, we combine natural proprioceptive stimuli with a novel model of neural coding and

topographical organization of neurons in somatosensory cortex to help develop our

understanding of proprioceptive coding and allowing us to paint a picture beyond that we can

see through the restricted lens of neural recordings. Core to our approach was allowing the

model to learn to represent the statistics of natural human movements rather than stereotyped

movements, which under the efficient coding hypothesis is key to understanding sensory

representations in the nervous system [23,29–31].

Our topo-VAE is designed to reflect information processing in cortex: we hypothesise that the

cortex learns, through experience, an efficient representation of the somatosensory world

which we replicate here as a deep autoencoder network. The network learns a nonlinear

mapping from kinematic inputs into a "cortical" latent space of spiking neurons, which

embodies a stochastic, generative model of movement-related neural activity. While VAEs

have been used before to learn both single neuron and population level features from spike

train data [32,33] a vanilla VAE (with a Euclidean latent space) cannot induce a topographic

mapping on the data. Consequently, they cause nearby input variables (stimuli) to be

represented in arbitrary locations across the latent space. By contrast, the latent space of a

topo-VAE is arranged as a two-dimensional cortical surface, thereby creating a population of

neurons on a square grid. This enables a lateral interaction term between neighbouring neurons

to be added, that shapes cortical activity into local neighbourhoods, allowing it to learn

cortical-like spatial structure as well as temporal single-neuron activity patterns, both of which

we compare to recorded neural data.

Previous self-organising maps [19,34,35] operate on less biologically plausible grounds. For example, Kohonen-type maps operate on winner-takes-all, non-spiking activation (and consequently, their training updates synaptic connections in a winner-takes-all form as well) [35]. Therefore, only one neuron can ever be active for given a sensory stimulus in a Kohonen map, a property very unlike actual proprioceptive cortex. In contrast, any number of neurons in our topo-VAE can be active simultaneously (and consequently, all synaptic connections are updated as sensory information is processed). Likewise, Poisson GLMs consider only the input statistics when producing simulated neural responses. While it is possible to reproduce the coding properties of individual neurons from convergence of peripheral inputs to a GLM, those models can tell us little about the role of neighbouring cells in shaping receptive fields or neuroanatomy-neural function relationships in general.

Emergent from our model, were neurons with tuning surfaces comparable to those of recorded neurons as monkeys performed the same centre-out task (Fig. 2.A,B). Furthermore, higher-level features of the modelled neurons, such as PD distributions and the distributions of tuning curve features, were comparable to recorded neurons, particularly when the model was constrained by lateral effects and trained on stimuli that reflected the natural distributions of arm movement (Fig. 2&3). However, they were not captured perfectly (e.g. the PD and tuning geometry distributions of modelled neurons were still significantly different from those of recorded neurons, $p<0.005$; Kolmogorov-Smirnov test). Despite this, the current model is able provide multiple predictions about the nature of proprioceptive coding, one of which we

partially validate with our limited neural data. Future work includes identifying additional

constraints that can allow the model to better reproduce neural data from proprioceptive cortex.

The well-known somatosensory homunculus represents only the tactile component of

somatosensation and its well-ordered map of the skin receptors. Since the human tactile

homunculus has driven much of neuroscience's intuition about somatosensory representations,

it is tempting to hypothesise that proprioceptive representations might also be similarly

structured. However, proprioception is driven not by receptors embedded in a "simple" two-

dimensional sheet, but rather by different types of muscle receptors with varying dynamics,

that span one, two, and even three joints. Thus, the expectation that proprioception and touch

might share a similar homunculus may not be reasonable. Indeed, instead of a point-to-point

mapping to the limb, in our model, we find structure akin to the pinwheels of the visual cortex,

here representing hand movement direction instead of orientation selectivity. While the

relatively large spacing of electrodes used to record neural data from area 2 does not allow us

to confirm the pinwheel anatomical structure directly, there are signatures of it in the

recordings, such as the fact that neurons recorded on a given electrode tend to have more

similar PDs than those recorded on separate electrodes – a necessary, but not sufficient

property for proprioceptive pinwheels.

Our model predicts not only the spatial organisation of tuning, but also the receptive fields of

single neurons – although this prediction remains difficult to test with our currently available

data. We find that modelled neurons encode combinations of joints, both adjacent (shoulder-

elbow) and distant (shoulder-wrist). Such convergence in monkey somatosensory cortex has

been observed for the hand [36,37] and must be present to some extent in the proximal arm,

given its multi-articular muscles. Our model used only joint-based inputs (i.e., the kinematic

state or pose of the arm) but knows nothing about the musculoskeletal mechanics of the limb

(e.g., the fact that bi-articulate muscles span multiple joints). Nonetheless, multi-joint coding

emerged for most neurons in our model. The correlations between motion of different joints

that we observe in the natural data are substantial (4 principal components explain 80% of the

variance of the seven degrees of freedom of the arm) and are the result of three main factors:

1) the biomechanics of the body, including the way many muscles span multiple joints, 2) the

way the brain controls movements and 3) the tasks performed. Arguably, task requirements

drive a substantial amount of the correlations. By training the topo-VAE on a highly varied

dataset of natural movement, we enable it to generalise from these natural tasks to planar

centre-out movements. Using this same dataset, Ejaz et al showed that the pairwise similarity

of finger-specific fMRI activity patterns in human sensorimotor cortex was better explained

by the correlation structure of hand movements than muscle activity [29]. Therefore, the

emergence of multi-joint proprioceptive receptive fields may be a further example of the

shaping of neural activity by higher-order features of movements [38], analogous to higher-

order features of visual receptive fields, such as edges in V1 [18].

It is important to acknowledge that while the comparisons between modelled and recorded

neurons are promising, it is largely a qualitative similarity, particularly at the level of

population statistics. Understanding what further constraints might enable our model to better

reproduce these features will be an important step in further elucidating the principles that

shape proprioceptive representations in the brain. We also make the key assumption that the

proprioceptive organizational principles captured by our topo-VAE model based on the

kinematics of human movement would extend to monkeys. While this assumption seems to

have largely held, there may still exist differences that contribute to the inconsistencies

between our model and recorded neurons.

Another key limitation of the current model is the use of only joint kinematics. Biological

inputs to proprioceptive cortex are received from muscle spindles and Golgi tendon organs,

which convey muscle length, velocity, and force information. An important future direction

will be to test the topo-VAE on such inputs, whether they be recorded experimentally or

inferred from existing kinematic data using musculoskeletal simulations. Furthermore,

because of its accessibility with Utah multielectrode arrays, we compared our modelled

neurons to those recorded in area 2 of the somatosensory cortex, an area that combines

cutaneous and muscle information in single neurons. Area 3a, on the other hand, has only

muscle-receptor inputs, and shares many features with the adjacent motor cortex. One might

expect an even closer correspondence between our modelled cortex and area 3.

Lastly, we would like to highlight the observation that, when trained on the stereotyped centre-

out-data only, the topo-VAE poorly predicted recorded neuron data more poorly than when it

was trained on the richer natural movement data. We interpret this to reflect the efficient

coding hypothesis [22], whereby a population of sensory neurons are optimised to code stimuli

representative of those found in their natural environment. Such codes are well documented in

the visual [18] and auditory systems [39] – here we provide evidence that this may also be the

case for proprioceptive neurons. This points towards the importance of working with training

data that matches the natural distribution of stimuli when modelling sensory neurons, as it

suggests important features of the neural code (such as the complex receptive fields predicted

by our model) will only be elucidated under these conditions. This observation may be of great

importance for the many proprioceptive and motor neurophysiology experiments that have

been conducted in highly constrained lab settings, settings that may not contain adequate

ethologically relevant kinematic statistics to uncover the true coding of cortical neurons.

Undertaking electrophysiological experiments with a broader repertoire of movements may

affect proprioceptive neuroscience as much as the adoption of natural images did for

understanding vision.

# Methods

## *Human Behaviour Scenarios and Natural Movement Data*

We recorded full-body movements from 18 healthy right-handed participants in two experimental scenarios. In the natural behaviour scenario (see Fig. 1.B & Supplemental Fig. 1), subjects performed unconstrained daily tasks in a working kitchen environment. As food preparation and feeding are universal behaviours, the only direction given to subjects was to prepare and eat an omelette. For this modelling work we used only the arm movement data (including wrist but not digits). The average recording time across subjects was 22 minutes. In the second movement scenario, a subject performed planar centre-out reaches in a 20x20 cm horizontal task space (see Fig. 1.C) to mimic the movement data for the monkey task. The horizontal task space was aligned 20 cm below the subject's shoulder and centred on the mid-line at 30 cm forward of the chest.

Arm movements in both scenarios were recorded at 60 Hz by an XSENS 3D motion tracking suit, a full-body sensor network based on inertial sensors. We used biomechanical models and fusion algorithms (including calibration and validation) to estimate joint angles. Fig. 1.A shows the biomechanical structure and coordinate system used in this paper. Arm movement datasets were formatted as time series of angles between segments following the International Society of Biomechanics (ISB) Euler angle extractions [25] in a ZXY coordinate. For the elbow and the wrist, angular rotations of Z, X and Y represent flexion/extension, abduction/adduction, and internal/external rotation, respectively, since the biomechanics of the human body cannot be fully described by a rotation around a single axis and must instead be described with respect to 3

509   rotational axes. During planar movements, we used optical tracking as well as the motion tracking

510   suit for capturing the endpoint (hand) position on the task square. Data from the inertial and optical

511   motion tracking systems were synchronised manually via cue-based movements before, during

512   and after the recording period.

## *Variational Autoencoder with Topographic Latent Space*

In the following we lay out the rationale for building our model and the model itself, the various

forms of data we collected for model training and validation, and the validation methodology.

The topo-VAE model (cf. Fig. 1.D-G) uses the autoencoder framework to model sensory

representations. Autoencoders are a type of artificial neural network used to learn efficient

encodings of unlabelled data [40]. The neuroscientific mechanism is referred to as the Infomax

principle, i.e. unsupervised learning to discover structure in the sensory data by maximising the

match between the inputs (in our case, the somatosensory world) and their neural representation

[21,22]. We choose specifically a variational autoencoder [24] because we want the latent cortical

layer to be able to capture the stochasticity and variability inherent to neural representations [41].

We modelled the latent neurons as Bernoulli processes using the Gumbel-max trick [42] where

neuron firing is sampled from a categorial distribution of $\{0,1\}$. This forces the model to learn an

encoding based on spike counts, which for an unchanging input will approximate the Poisson

distributed statistics of biological neurons. This allowed us to use the same data analysis pipeline

as for our recorded neural data. However, this simple VAE model would be devoid of any spatial

relationships between the neurons. Therefore, we added a simple organisational mechanism that

would link learning between neighbouring latent neurons, effectively implementing cortical lateral

connectivity with short range excitation and longer-range inhibition. Thus, our enhanced VAE

learns both receptive field tuning properties (in an unsupervised manner through deep feature

learning) and establishes a topographic relationship between neurons.

Let $[X] = \{\vec{x_n}\}_{n=1,...,N}$, $x_n \in \mathbb{R}^D$ be the sensory stimuli, following the natural behaviour

distribution $p(\vec{x})$. A group of cortical neurons, arranged with a topographic structure, are

activated by the sensory stimuli $[X]$ and generate firing patterns $[Z] = \{\vec{z_n}\}_{n=1,...,N}$, $z_n \in$

$\mathbb{R}^{M \times M}$, $(M^2 \gg D)$. We aim to find a decoder $p_\Phi([X] \mid [Z])$ and corresponding neural responses

$[Z]$ to optimally represent the sensory stimuli $[X]$, namely maximise the marginal likelihood of

$p([X])$. The variational lower bound of the log likelihood $\log p(X)$ is derived as:

$$\log p([X]) = log \int p([X]|[Z])p([Z]) \, d \, [Z]$$
$$\geq \mathbb{E}_{Z \sim q([Z])}[log p_\phi([X]|[Z]) - KL(q([Z])||p([Z]))]$$

(1)

where $q(Z)$ is the variational parameter approximating the intractable true posterior $p(Z|X)$.

Our topo-VAE encoder can be considered as a multi-layer neuronal structure delivering sensory

stimuli to the cortex from sensory afferents, through brainstem nuclei, to the thalamus and cortex.

This implies that at the encoder level we do not attribute or consider specific representations at

these intermediate stages of proprioceptive processing (including how different sensory systems

are integrated. From the perspective of computational modelling, it is also the amortised inference

for inferring the optimal representation $q([Z])$, helping to avoid smoothness problems in over-

complete representations (`back-constraint'). Fig. 1b illustrates the detailed structure of our topo-

VAE, with a multi-layer perceptron (MLP) encoder and a linear decoder. The latent layer contains

spiking neurons whose spike counts $z_n$ within a given time interval $\Delta t$ (i.e. over multiple input

steps) follow a Poisson distribution. The encoder infers the distribution $q([Z])$ of responses for a

group of cortical neurons, and $z_n$ is sampled from this distribution using the Gumbel-max trick:

$$q(\vec{z}_n) = \underset{i}{\mathrm{argmax}}[g_i + \log \pi_i]$$
(2)

Where $\pi_i = [p \quad 1-p]$ is a vector defining the probability of a neuron spiking vs. not spiking,

and $p$ is set by the output of the encoder network for each neuron.

The decoder is a linear mapping from neural activities $[Z]$ to the reconstructions $[\hat{Z}]$ of the

sensory stimuli. To summarise, the standard VAE model (which typically uses normally

distributed random variables in the latent layer) is replaced by a latent layer of Bernoulli-type

spike count distributions that are immediately applicable to neural signal analysis.

In addition to the log-likelihood function (Eq. 1) of the standard variational encoder, we include

lateral effects in the latent layer. This was done by defining a distance-dependent function acting

on the neurons, which are arranged in an $M \times M$ topographic map. A natural choice for cortical

neurons is the Mexican-hat function, which transitions with distance from excitation to inhibition

before vanishing [43] (Fig. 1.F,G). As we are modelling an entire population, we can represent the

interaction between neurons as a matrix $[\Psi]$. Each element $[\Psi_{p,q}]$ represents the lateral effect

between neurons p and q, calculated as:

$$\Psi_{p,q} = (1 - \frac{d_{p,q}^2}{2\sigma^2})e^{-\frac{d_{p,q}^2}{2\sigma^2}} \tag{3}$$

576 where $d_{i,j}$ represents the Euclidean distance and $\sigma$ is a hyperparameter defining the common

577 length scale of local excitation and intermediate-range inhibition. As shown in Fig. 1.G, the

578 transition from maximum excitation to maximum inhibition spans a distance of $2\sigma$ and the lateral

579 effect vanishes at about $4\sigma$.

581 The total loss function governing our topo-VAE model is given by:

$$L(\Phi, \Theta) = -\mathbb{E}_{[Z] \sim q_\Phi([Z])}[log\, p_\Theta([X]|[Z])] + KL(q([Z]) \,||\, p([Z]))$$
$$- \gamma \mathbb{E}_{Z \sim q_\Phi([Z])}[Tr([Z']^T[\Psi][Z'])] \tag{4}$$

where $p([Z])$ is the prior distribution, set to be an independent Poisson distribution with rate $r_p$,

$Z' = Z - z_b$. $z_b$ is the base firing rate and $\gamma$ is a constant controlling the impact of the lateral

effects. $\Phi$ and $\Theta$ represent the trainable parameters in the encoder and the decoder, respectively.

The KL divergence term in the loss function performs as a constraint on temporal sparsity,

penalizing firing rates far from the expected rate $r_p$ in the prior distribution $p([Z])$. $r_p$ is usually

set to be small, due to constraints such as metabolic cost [44], and allows us to naturally control

the temporal sparsity of neural activity. In addition to the temporal sparsity, our topo-VAE also

involves structured spatial sparsity. The lateral effect, as represented by the topographic term

$\gamma \mathbb{E}_{[Z] \sim q_\Phi([Z])}[Tr([Z']^T[\Psi][Z'])]$ in Eq. 4 introduces topographic structure on the latent space and

specifies the firing dependencies between neurons. Lateral excitation dominates the formation of

pattern patterns in a structured space while lateral inhibition encourages spatial sparsity by

penalising co-activation of non-nearby neurons. From the perspective of probabilistic inference,

this regularisation item is equivalent to amending the prior distribution $p([Z])$ and modifies the

target function as:

$$L(\Phi, \Theta) = -\mathbb{E}_{[Z] \sim q_\Phi([Z])}[log\, p_\Theta([Z]|[Z])] + KL\big(q([Z]) \,\|\, p^*([Z])\big) + const,$$

$$p^*([Z]) = \frac{p([Z])e^{\gamma Tr([z']^T[\Psi][z'])}}{B},$$

(5)

where the normalisation factor $B = \int p([Z])e^{Tr\big(\gamma[z']^T[\Psi][z']\big)}\, dZ$ is a constant. In this form, both

the firing sparsity and the lateral effect are expressed within the amended prior distribution

$q^*([Z])$ and the target function is maintained in a standard variational autoencoder framework.

We demonstrate through ablation and parameter variation experiments the need for the specific

design elements of our model to explain the biological data.

The encoder component of the model contains two fully connected feed-forward layers of size 50

and 100 neurons, respectively, and a final linear readout layer. Tanh activation functions were

used for all neurons in the first two layers. The reconstruction error of the model is measured

using mean squared error.

To compare our modelling and recording results, we chose hyperparameters of the topographic

map with consideration of 1) spatial densities of S1 neurons and 2) characteristics of the neural

recording devices. The density of neurons in S1 is about 8M – 17M per cm$^3$ [45,46] of which 70%

to 90% are pyramidal cells [47]. The thickness of cortex varies from 1 mm to 4.5 mm [48,49]. The

Utah electrode array has 100 microelectrodes arranged in a 10 x 10 configuration with 400 μm

separation along each axis, thus spanning 3.6 mm x 3.6 mm of the cortex. We model neuronal

anatomy as voxels or cubes, where the number of pyramidal cells contained within 1mm x 1mm x

1mm of cortex varies from 12,000 to 1,000,0000, which means every 1 mm along the cortical

surface crosses about 20 - 50 pyramidal cells (see Fig. 1c). A surface-parallel slice captures a grid

of 80 x 80 neurons. This conceptually simplified arrangement allows us to formulate a

computationally tractable design of the latent layer neurons in topo-VAE. We can give a bit of

intuition of our topographic model's parameters to neuroanatomy: the range of the lateral effect

parameter $\sigma$ translates to about 1-2 neuron spacing on our cortical model grid (assuming a radius

of dendritic input to these cortical neurons of around 200 μm [50]). Its precise value was

determined pot-hoc using model selection by numerically sweeping for a range of $\sigma$ and selecting

the best fitting value.

To observe the effect of the neighbourhood range parameter σ on topography in the latent

representation, we test values σ={1,2,3}. For the expected firing rate *p(x)*, we find stable

topography around *p(x)*=0.01. The training loss function contains multiple components which can

be given individual weightings. Here we use weightings of 10, 0.4, and 0.005 for the

reconstruction, lateral effect, and firing sparsity components, respectively. In addition, we include

an L2 norm on the model weights β=0.1. When sampling from the latent space during decoding,

we parameterise the scaling of the rate parameter, to simulate sampling across different time

windows. Varying this parameter in the range [0,1000] yielded optimal reconstruction at a scaling

factor of 40.

*Model Training & Validation overview*

The topo-VAE model was implemented in Python using PyTorch [51] and run on a GPU

workstation. All models were trained for 4000 epochs using the Adam optimiser with a learning

rate of $10^{-5}$ and a batch size of 400.

To train our topo-VAE we needed to pre-process the input data. Empirical distribution of joint

angular velocities during movements in our tasks is symmetric, unimodal, with sharp peaks at zero

and heavy tails towards large speeds. To ensure equal contributions to model optimisation, these

features are standardised without mean centring (since they are already approximately zero-

centred, and to preserve the correspondence of a zero-valued input to zero joint angle velocity).

This pre-processing also improves the training efficiency and convergence of our model without

breaking the spatial structure of natural movements. In addition, since the original kinematic data

were sampled at a frequency of 60Hz, we subsampled the training data at a depth of 1%, to remove

redundancy and improve training times, without significant effects on the outcome of the model.

Optimisation of VAEs by backpropagation requires estimation of parameter gradients with respect

to loss. Since gradients cannot be estimated for the Gumbel-max function, we implemented the

Gumbel-softmax reparameterisation trick [52] during training, where *z* is sampled from *q(z)* using

the Gumbel-softmax function:

$$q(\vec{z}_n) = \frac{\exp(\frac{\log(\pi_i) + g_i}{\tau})}{\sum_{j=1}^{2} \exp(\frac{\log(\pi_j) + g_j}{\tau})} \tag{2}$$

658    And $g_i$ is a random sample from the Gumbel distribution $G(0,1)$:

$$P(x) = e^{-(x+e^{-x})} \tag{3}$$

We wanted to perform feature learning of proprioceptive representations with our topo-VAE and

this required large amounts of natural movement data. Therefore, we performed human

experiments (see above) and measured the full-body kinematics of human subjects. We use human

full-body behaviour as a functional proxy for non-human primate movements of the arm [53], as

human data is much easier and more precisely obtainable. We used the human data to drive the

training of our topo-VAE, then froze our model parameters to evaluate it. Our topo-VAE model is

generative, so by playing back any limb movement data (time series of body poses) we obtain

spike trains for each neuron in our cortical grid.

To compare our model's predictions to those of recorded neurons, we use the kinematic data from

humans performing the same centre-out task as the monkeys (see above) to drive the frozen topo-

VAE model and compare its output to the actual recorded neural data.

*Model Robustness and Parameter Variation*

The topographic property of the VAE arises from the lateral effect component of the loss function

(Eq. 4). As a control, we tested our model with no lateral effects (Fig. 5.A-D), and with several

different distance functions (Mexican hat, inverted Mexican hat, excitation only, inhibition only;

Fig. 5.F,G). The inverted Mexican hat lateral effect is the additive inverse of the Mexican hat

function (Eq. 3). Excitation-only lateral effect is defined as:

$$\Psi_{p,q} = e^{-\frac{d_{p,q}^2}{2\sigma^2}} \tag{6}$$

681    and the inhibition-only lateral effect is the additive inverse of the excitation-only lateral effect (Eq.

682    6).

## *Model Analysis*

To quantify the sensitivity of neurons to specific joints (Fig. 6), we individually perturb each input feature of the model and measure the Pearson correlation between individual neurons in the latent space and that feature. Since each joint is represented by three input features (angular velocity in the Z, X, and Y axes), we use the mean of the absolute correlation across all three axes to quantify the sensitivity of a given neuron to a particular joint.

We computed angular velocity profiles from the recorded data and analysed the natural movement dataset $X_{nat}$ and the planar movement dataset $X_{pl}$. The first two principal components of the planar movements explained over 95% of the total variance, but only half of the variance of the natural movements. This reveals that joint velocities of planar movements are highly constrained, as expected, restricted largely to a 2-dimensional subspace. We applied the manipulative complexity metric [54] to quantify the complexity of the movements, which is defined as:

$$C = 1 - \frac{2}{D-1}\sum_{j=1}^{D}\sum_{i=1}^{j}\left(VAF_i - \frac{1}{D}\right)$$

Where $VAF_i$ is the variance captured by the i[th] PC. Larger values of C indicate higher complexity and a value of $C = 1$ means that all PCs contribute equally to the total variance. The complexity of planar movements was 0.06, much lower than the natural movement complexity of 0.5. We defined the direction θ and speed $v$ of planar movements in world coordinates: 90°/270° are respectively away from and towards the chest, 180°/0° are to the left and right.

Fig. 1.H illustrates our use of the two human datasets with our computational model. The natural

movement dataset $X_{nat}$ can be viewed as a group of samples generated from the natural movement

statistic. Following the idea of natural sensory coding [18], we used this natural movement dataset

as training data to learn an optimal neural coding scheme. After training, we tested the converged

model with data from the planar reaching task. To compare with hand-based coding properties of

area 2 neurons [55,56], we found a linear mapping between joint angular velocities and planar

hand velocity. This allowed us to assess the relationship between hand movement direction and

topo-VAE firing rates.

*Nonhuman Primate Behaviour and Data Collection*

We used a combination of previously recorded data in which three rhesus macaques performed a

planar, centre-out reaching task while seated, using a two-link planar manipulandum. A cursor

displayed on a monitor tracked the position of the manipulandum and provided visual feedback

for the monkey as he reached for a target on-screen. The monkey moved the cursor to a central

target in the workspace. After a random delay period, 1 of 8 targets spaced evenly in a circle around

the central target appeared on the screen and the monkey moved the cursor toward it upon an

audible 'go' cue. After placing the cursor in the target for a random hold time of 0-500 ms, the

monkey received a liquid reward and returned the cursor to the central target. We used 6

experimental sessions across three monkeys; two contain data that has been previously published

[56], and the rest are unpublished. All procedures were in accordance with the Guide for the Care

and Use of Laboratory Animals and were approved by the institutional animal care and use

committee of Northwestern University under protocol #IS00000367.

Once a monkey was trained on the experimental apparatus, a 96-channel microelectrode array with

1 mm iridium-oxide coated electrodes (Blackrock Microsystems, Inc.) was pneumatically inserted

in Brodmann's area 2, near the intraparietal sulcus [56]. The implantation site was chosen to avoid

cerebral vasculature and maximise proximal arm representation. All surgery was performed under

isoflurane gas anaesthesia (1-2 percent) except during intraoperative recording to identify the arm

representations, when the monkey was transitioned to a mixture of <0.5% isoflurane and

remifentanil (0.4 ug/kg/min).

The data were recorded from the microelectrode array using the Cerebus multichannel data

recording system (Blackrock Microsystems, Inc.). Thresholded waveforms and timing of

behavioural task events were synchronised and recorded for offline analyses. The position of the

handle was recorded at 1kHz. We discriminated single neurons using Offline Sorter (Plexon, Inc.,

Dallas TX). 383 recorded neurons were found to be suitable for subsequent analysis.

To calculate the preferred direction of a neuron, we used a simple bootstrapping procedure. For

each iteration, we drew random points from the dataset conforming to a uniform distribution of

movement directions. We then fit Poisson generalised linear models (GLM) with angular velocity

inputs to the firing rate for the sampled timepoints. The GLM models are defined by:

$$[\hat{f}] = Poisson(\lambda), \qquad \lambda = e^{[X][\beta]}$$

where $[\hat{f}]$ is a $T$ (number of time points) by $N$ (number of neurons) matrix of firing rate estimates

of the recorded rates $[f]$, $X$ is a $T$ by 2 (number of velocity inputs) matrix, and $\beta$ is a $P$ by $N$ matrix

of encoding parameters. $\beta$ was found using maximum likelihood estimation. The preferred

direction was then calculated from the encoding vector of the GLM as:

754
$$PD_i = \tan^{-1}(\beta_y, \beta_x), \qquad r_i = \sqrt{\beta_y^2, \beta_x^2}$$

In these equations, a bootstrap PD estimate of neuron $i$ was defined by $\beta_y$ and $\beta_x$, the bootstrap

encoding parameters for hand velocity in the $y$ and $x$ directions, respectively. We took the circular

mean of the PD estimates over all bootstrap iterations to find the PD for each neuron. This method

of PD calculation was used for both recorded and modelled neurons.

**Data Availability**

All code is available at https://doi.org/10.6084/m9.figshare.17393591.v3.

All data is available at https://doi.org/10.6084/m9.figshare.17393591.v3.

**Author contributions**

AAF conceived the study. LEM & AAF supervised the study. YW, AAF developed the theory and

model. JAH, AAF collected the behavioural data. KPB collected the unpublished neural data.

YW, MG implemented the model code. YW, MG implemented the behavioural data analysis code.

KPB implemented the neural data analysis code. KPB, MG, YW, AAF, LEM analysed the data.

KPB, LEM, AAF drafted the manuscript. All authors discussed the results and reviewed the

manuscript.

## Funders

We acknowledge: KPB was supported by NIH BRAIN NRSA Postdoctoral Fellowship

F32MH120893 and [S1R01]. YW was supported by an NIHR Imperial College BRC Deep

Phenotyping Grant and Project eNHANCE (http://www.enhance-motion.eu) under the European

Union's Horizon2020 research and innovation programme (Grant No. 644000). MG was supported

by the Wellcome Trust PhD Program "Bioinformatics & Theoretical Systems Biology"

(222888/Z/21/Z). JAH was supported by an EPSRC Doctoral Training Award. LEM was

supported by NINDS grant #R01NS095251. AAF acknowledges his UKRI Turing AI Fellowship

(EP/V025449/1).

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

## Financial Disclosures

**REFERENCES**

1.  Todorov E, Jordan MI. Optimal feedback control as a theory of motor coordination. Nat Neurosci. 2002. doi:10.1038/nn963

2.  Scott SH. Optimal feedback control and the neural basis of volitional motor control. Nature Reviews Neuroscience. 2004. doi:10.1038/nrn1427

3.  Tuthill JC, Azim E. Proprioception. Current Biology. 2018. doi:10.1016/j.cub.2018.01.064

4.  Sainburg RL, Poizner H, Ghez C. Loss of proprioception produces deficits in interjoint coordination. J Neurophysiol. 1993. doi:10.1152/jn.1993.70.5.2136

5.  Flesher SN, Downey JE, Weiss JM, Hughes CL, Herrera AJ, Tyler-Kabara EC, et al. A brain-computer interface that evokes tactile sensations improves robotic arm control. Science (80- ). 2021. doi:10.1126/science.abd0380

6.  Faisal AA. Putting touch into action. Science (80- ). 2021. doi:10.1126/science.abi7262

7.  Formento E, Minassian K, Wagner F, Mignardot JB, Le Goff-Mignardot CG, Rowald A, et al. Electrical spinal cord stimulation must preserve proprioception to enable locomotion in humans with spinal cord injury. Nat Neurosci. 2018. doi:10.1038/s41593-018-0262-6

8.  Blum KP, Versteeg C, Sombeck J, Chowdhury RH, Miller LE. Proprioception: a sense to facilitate action. Somatosensory Feedback for Neuroprosthetics. 2021. doi:10.1016/b978-0-12-822828-9.00017-4

9.  Penfield W, Boldrey E. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. Brain. 1937. doi:10.1093/brain/60.4.389

10. Weber DJ, London BM, Hokanson JA, Ayers CA, Gaunt RA, Torres RR, et al. Limb-state information encoded by peripheral and central somatosensory neurons: Implications for an

afferent interface. IEEE Trans Neural Syst Rehabil Eng. 2011.

doi:10.1109/TNSRE.2011.2163145

11.  Huffman KJ, Krubitzer L. Area 3a: Topographic organization and cortical connections in

marmoset monkeys. Cereb Cortex. 2001. doi:10.1093/cercor/11.9.849

12.  Rossi LF, Harris KD, Carandini M. Spatial connectivity matches direction selectivity in

visual cortex. Nature. 2020. doi:10.1038/s41586-020-2894-4

13.  Killebrew JH, Bensmaïa SJ, Dammann JF, Denchev P, Hsiao SS, Craig JC, et al. A dense

array stimulator to generate arbitrary spatio-temporal tactile stimuli. J Neurosci Methods.

2007. doi:10.1016/j.jneumeth.2006.10.012

14.  Mohsenzadeh Y, Mullin C, Lahner B, Oliva A. Emergence of Visual Center-Periphery

Spatial Organization in Deep Convolutional Neural Networks. Sci Rep. 2020.

doi:10.1038/s41598-020-61409-0

15.  Stringer C, Pachitariu M, Steinmetz N, Carandini M, Harris KD. High-dimensional

geometry of population responses in visual cortex. Nature. 2019. doi:10.1038/s41586-019-

1346-5

16.  Pehlevan C, Genkin A, Chklovskii DB. A clustering neural network model of insect

olfaction. Conference Record of 51st Asilomar Conference on Signals, Systems and

Computers, ACSSC 2017. 2018. doi:10.1109/ACSSC.2017.8335410

17.  Sandbrink KJ, Mamidanna P, Michaelis C, Mathis MW, Bethge M, Mathis A. Task-driven

hierarchical deep neural network models of the proprioceptive pathway. bioRxiv. 2020.

18.  Olshausen BA, Field DJ. Sparse coding with an overcomplete basis set: A strategy

employed by V1? Vision Res. 1997. doi:10.1016/S0042-6989(97)00169-7

19.  Obermayer K, Blasdel GG. Geometry of orientation and ocular dominance columns in

monkey striate cortex. J Neurosci. 1993. doi:10.1523/jneurosci.13-10-04114.1993

20.  Marr D. Vision: a computational investigation into the human representation and

processing of visual information. Vis a Comput Investig into Hum Represent Process Vis

information. 1982. doi:10.1016/0022-2496(83)90030-5

21.  Linsker R. Self-Organization in a Perceptual Network. Computer (Long Beach Calif).

1988. doi:10.1109/2.36

22.  Barlow HB. Possible Principles Underlying the Transformations of Sensory Messages.

Sensory Communication. 2013. doi:10.7551/mitpress/9780262518420.003.0013

23.  Laughlin S. A simple coding procedure enhances a neuron's information capacity.

Zeitschrift fur Naturforschung - Section C Journal of Biosciences. 1981. doi:10.1515/znc-

1981-9-1040

24.  Kingma DP, Welling M. Auto-encoding variational bayes. 2nd International Conference

on Learning Representations, ICLR 2014 - Conference Track Proceedings. 2014.

25.  Wu G, Van Der Helm FCT, Veeger HEJ, Makhsous M, Van Roy P, Anglin C, et al. ISB

recommendation on definitions of joint coordinate systems of various joints for the

reporting of human joint motion - Part II: Shoulder, elbow, wrist and hand. J Biomech.

2005. doi:10.1016/j.jbiomech.2004.05.042

26.  Sterling P, Laughlin S. Principles of neural design. Principles of Neural Design. 2015.

doi:10.7551/mitpress/9780262028707.001.0001

27.  Kurtzer I, Herter TM, Scott SH. Nonuniform distribution of reach-related and torque-

related activity in upper arm muscles and neurons of primary motor cortex. J

Neurophysiol. 2006. doi:10.1152/jn.00110.2006

28.  Herter TM, Kurtzer I, Cabel DW, Haunts KA, Scott SH. Characterization of torque-

related activity in primary motor cortex during a multijoint postural task. J Neurophysiol.

2007. doi:10.1152/jn.00757.2006

29.  Ejaz N, Hamada M, Diedrichsen J. Hand use predicts the structure of representations in

sensorimotor cortex. Nat Neurosci. 2015. doi:10.1038/nn.4038

30.  Simoncelli EP, Olshausen BA. Natural image statistics and neural representation. Annual

Review of Neuroscience. 2001. doi:10.1146/annurev.neuro.24.1.1193

31.  Ganguli D, Simoncelli EP. Efficient sensory encoding and Bayesian inference with

heterogeneous neural populations. Neural Comput. 2014. doi:10.1162/NECO_a_00638

32.  Wu A, Roy NA, Keeley S, Pillow JW. Gaussian process based nonlinear latent structure

discovery in multivariate spike train data. Advances in Neural Information Processing

Systems. 2017.

33.  Higgins I, Chang L, Langston V, Hassabis D, Summerfield C, Tsao D, et al. Unsupervised

deep learning identifies semantic disentanglement in single inferotemporal face patch

neurons. Nat Commun. 2021. doi:10.1038/s41467-021-26751-5

34.  Aflalo TN, Graziano MSA. Possible origins of the complex topographic organization of

motor cortex: reduction of a multidimensional space onto a two-dimensional array. J

Neurosci. 2006. doi:10.1523/JNEUROSCI.0768-06.2006

35.  Kohonen T. Self-organized formation of topologically correct feature maps. Biol Cybern.

1982. doi:10.1007/BF00337288

36.  Costanzo RM, Gardner EP. Multiple-joint neurons in somatosensory cortex of awake

monkeys. Brain Res. 1981. doi:10.1016/0006-8993(81)91197-5

37.  Warren JP, Santello M, Tillery SI. Effects of fusion between tactile and proprioceptive

inputs on tactile perception. PLoS One. 2011. doi:10.1371/journal.pone.0018073

38.    Thomik AAC, Fenske S, Faisal AA. Towards sparse coding of natural movements for

883           neuroprosthetics and brain-machine interfaces. International IEEE/EMBS Conference on

884           Neural Engineering, NER. 2015. doi:10.1109/NER.2015.7146780

885    39.    Lewicki MS. Efficient coding of natural sounds. Nat Neurosci. 2002. doi:10.1038/nn831

886    40.    Hinton GE, Salakhutdinov RR. Reducing the dimensionality of data with neural networks.

887           Science (80- ). 2006. doi:10.1126/science.1127647

888    41.    Orbán G, Berkes P, Fiser J, Lengyel M. Neural Variability and Sampling-Based

889           Probabilistic Representations in the Visual Cortex. Neuron. 2016.

890           doi:10.1016/j.neuron.2016.09.038

891    42.    Gumbel EJ. Statistical Theory of Extreme Values and Some Practical Applications: A

892           series of lectures. US Government Printing Office. 1954.

893    43.    Amari S ichi. Dynamics of pattern formation in lateral-inhibition type neural fields. Biol

894           Cybern. 1977. doi:10.1007/BF00337259

895    44.    Attwell D, Laughlin SB. An energy budget for signaling in the grey matter of the brain.

896           Journal of Cerebral Blood Flow and Metabolism. 2001. doi:10.1097/00004647-

897           200110000-00001

898    45.    Turner EC, Young NA, Reed JL, Collins CE, Flaherty DK, Gabi M, et al. Distributions of

899           Cells and Neurons across the Cortical Sheet in Old World Macaques. Brain Behav Evol.

900           2016. doi:10.1159/000446762

901    46.    Collins CE, Turner EC, Sawyer EK, Reed JL, Young NA, Flaherty DK, et al. Cortical cell

902           and neuron density estimates in one chimpanzee hemisphere. Proc Natl Acad Sci U S A.

903           2016. doi:10.1073/pnas.1524208113

47.    Kaas JH. Evolution of the neocortex. Current Biology. 2006.

doi:10.1016/j.cub.2006.09.057

48.    Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic

resonance images. Proc Natl Acad Sci U S A. 2000. doi:10.1073/pnas.200033797

49.    Wagstyl K, Ronan L, Goodyer IM, Fletcher PC. Cortical thickness gradients in structural

hierarchies. Neuroimage. 2015. doi:10.1016/j.neuroimage.2015.02.036

50.    Braitenberg V, Schüz A. Cortex: Statistics and Geometry of Neuronal Connectivity.

Cortex: Statistics and Geometry of Neuronal Connectivity. 1998. doi:10.1007/978-3-662-

03733-1

51.    Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, et al. PyTorch: An

imperative style, high-performance deep learning library. Advances in Neural Information

Processing Systems. 2019.

52.    Jang E, Gu S, Poole B. Categorical reparameterization with gumbel-softmax. 5th

International Conference on Learning Representations, ICLR 2017 - Conference Track

Proceedings. 2017.

53.    Young NM, Wagner GP, Hallgrímsson B. Development and the evolvability of human

limbs. Proc Natl Acad Sci U S A. 2010. doi:10.1073/pnas.0911856107

54.    Belić JJ, Faisal AA. Decoding of human hand actions to handle missing limbs in

neuroprosthetics. Front Comput Neurosci. 2015. doi:10.3389/fncom.2015.00027

55.    Prud'homme MJL, Kalaska JF. Proprioceptive activity in primate primary somatosensory

cortex during active arm reaching movements. J Neurophysiol. 1994.

doi:10.1152/jn.1994.72.5.2280

56.    Chowdhury RH, Glaser JI, Miller LE. Area 2 of primary somatosensory cortex encodes

kinematics of the whole arm. Elife. 2020. doi:10.7554/eLife.48198