A novel camera trap design for studying wildlife in mountain glacier ecosystems yields

new insight for glacier biodiversity in the Pacific Northwest, USA

Scott Hotaling[1], Jordan Boersma[2], Neil A. Paprocki[3], Alissa Anderson[4], Logan Whiles[4], Lucy

Ogburn[5], Sophia Kasper[5], Catharine White[6], Daniel H. Thornton[4], and Peter Wimberger[5,7]

Affiliations:

[1] Department of Watershed Sciences, Utah State University, Logan, UT, USA; ORCID #0000-

0002-5965-0986

[2] Cornell Lab of Ornithology, Ithaca, NY, USA

[3] Idaho Cooperative Fish and Wildlife Research Unit, College of Natural Resources, University

of Idaho, Moscow, ID, USA

[4] School of the Environment, Washington State University, Pullman, WA, USA

[5] Biology Department, University of Puget Sound, Tacoma, WA, USA

[6] Coast Mountain College, Terrace, BC, Canada

[7] Slater Museum of Natural History, Tacoma, WA, USA

Correspondence: Scott Hotaling, Department of Watershed Sciences, Utah State University,

Logan, UT, USA; Email: scott.hotaling1@gmail.com; Phone: (828) 507-9950

Running head: Human and wildlife usage of a glacier

Abstract

Context

The global recession of glaciers and perennial snowfields is reshaping mountain ecosystems.

Beyond physical changes to the landscape and altered downstream hydrology, the implications

of glacier decline for biodiversity are poorly known. Before predictions can be made about how

climate change will affect wildlife in glacier-associated ecosystems, a more thorough accounting

of the role that glaciers play in species' life histories is needed. However, typical approaches for

documenting wildlife presence and behavior—remote camera traps—are difficult to use in

glaciated terrain due to limited options for securing them (e.g., no trees) and dramatic seasonal

changes in snowpack.

Aims

In this study, we sought to test a novel camera trap designed for glaciated mountain

ecosystems. We also aimed to use this approach to gain insight into wildlife and human usage

of a mountain glacier in western North America.

Methods

We deployed an elevational transect of uniquely designed camera traps along the western

margin of the Paradise Glacier, a rapidly receding mountain glacier on the south side of Mount

Rainier, WA, USA. Our simple camera trap design consisted of a wildlife camera attached to a

camouflaged cylindrical cooler filled with snow and rocks.

Key results

Our camera design proved ideal for a mountain glacier ecosystem and from June to September

2021, we detected at least 16 vertebrate species (seven birds, nine mammals) over 770 trap

nights using glacier-associated habitats. Humans, primarily skiers, were the most common

species detected, but we also recorded 99 observations of wildlife (birds and mammals). These

included three species of conservation concern in Washington: wolverine (*Gulo gulo*), Cascade

red fox (*Vulpes vulpes cascadensis*), and white-tailed ptarmigan (*Lagopus leucura*).

Conclusions

Collectively, our results provide proof-of-concept for a novel camera trap design that is ideal for

treeless, perennially snow-covered landscapes and revealed a rich diversity of wildlife using

mountain glacier habitat in the Pacific Northwest. We highlight the global need for similar

studies to better understand the true scale of biodiversity that will be impacted by glacier

recession in mountain ecosystems.

Keywords: mountain biodiversity, wolverine, Pacific Northwest, Mount Rainier, camera

trapping, climate change, glacier recession, white-tailed ptarmigan

Introduction

As climate change proceeds and mountain glaciers are lost, there is a pressing need to

understand how the loss of glacier ice will impact habitats and ecosystems (Hotaling *et al.* 2017;

Milner *et al.* 2017). Well-known impacts of glacier recession include the loss of key meltwater

resources which transport nutrients to downstream habitats (Ren et al. 2019), underpin

hydropower infrastructure (Finger *et al.* 2012), and provide key habitats for specialized aquatic

communities (Giersch *et al.* 2017; Hotaling *et al.* 2019; Muhlfeld *et al.* 2020). Often overlooked

in this discussion, however, is the impact of glacier loss on the biodiversity living on and

adjacent to glaciers (Hardy *et al.* 2018; Schano *et al.* 2021; Stibal *et al.* 2020).

The biodiversity of glacier ecosystems has gained appreciation in recent decades. Long

considered devoid of life, it is now established that glaciers host a thriving, multi-kingdom

community (Anesio *et al.* 2017; Shain *et al.* 2021), particularly in the uppermost layer of snow

and ice where intense solar radiation, atmospheric deposition, and available water during melt

yield conditions that are more hospitable to life (Anesio and Laybourn-Parry 2012; Hotaling *et al.*

2021). Still, however, most studies have focused on the most dominant life form on glaciers—

microbes—leaving larger-bodied residents of glaciers and surrounding habitats overlooked.

Globally, 36 species of birds and mammals have been observed using glacier and perennial

snow habitat for some portion of their life history (Rosvold 2016). Mammals typically use these

habitats for relief from lower elevation conditions during warmer parts of the year, whereas birds

commonly forage on arthropods and seeds that have been atmospherically deposited on the ice

(Antor 1995; Hotaling *et al.* 2021; Rosvold 2016). However, closer links between wildlife and

glacier habitats have also been described. In western North America, the high abundance of

glacier ice worms (*Mesenchytraeus solifugus*)—which are commonly present at densities >200

per m^2 on glaciers during summer—may be a key resource subsidy for nesting birds (e.g., gray-

crowned rosy finches, Hotaling *et al.* 2020). On glaciers of the high Andes in South America, the

white-winged diuca finch nests directly within crevasses (Hardy *et al.* 2018). Throughout higher

latitudes of the Northern Hemisphere, wolverines cache food in snow and ice, and the

persistence of spring snow cover and cache longevity has been hypothesized as a primary

reason underpinning their cold, snowy distributions (i.e., the "refrigeration-zone" hypothesis,

Inman *et al.* 2012).

While links between larger-bodied vertebrates and icy habitats have been made for a range of

species, these observations typically stem from focused species-specific studies and likely

underestimate the true scale of wildlife use in the cryosphere (the global collection of Earth's

frozen waters). Beyond researcher focus on specific species, this underestimation likely also

stems from two other, intertwined challenges. Mountain ecosystems, and particularly glaciers,

are poorly studied relative to more convenient habitats (e.g., those in close proximity to

universities and museums, Freitag *et al.* 1998). Moreover, mountain ecosystems have additional

challenges associated with their study. Indeed, for mountain research, typical field methods are

often difficult to implement due to, for example, the remoteness, rugged terrain, and harsh

weather conditions (and high rates of snowfall) associated with these habitats. For instance,

most camera trapping studies affix cameras to trees or T-Post stakes driven into soil (Rich *et al.*

2019; Rovero *et al.* 2010). In mountain glacier ecosystems, trees are typically absent, soil is

usually covered by deep snowpack or non-existent due to glacier activity scouring the

landscape down to bedrock, and affixing cameras to boulders limits locations for deployment

and timing (e.g., early season deployments might not be possible due to boulders still being

covered by seasonal snow). Thus, an easy to transport, flexible, but robust camera trap design

to ameliorate the many challenges of high mountain ecosystems is needed.

In this study, we sought to gain a more holistic understanding of human and wildlife usage of

alpine glacier habitat while also testing a simple camera trap designed to alleviate the difficulties

associated with high-mountain ecosystems. We deployed our camera-trapping array along an

elevational transect of a mountain glacier on Mount Rainier, Washington, USA. While we

expected our camera trap array to detect many species known to occur in these ecosystems

(e.g., marmot), we also expected to observe species not previously associated with glacier or

snow ecosystems. If detected, these results would underscore the likely underestimation of

glacier-linked vertebrate biodiversity globally and the pressing need to better understand how

climate change may impact enigmatic mountain species.

Materials and methods

Study area

Mount Rainier is a large stratovolcano in south-central Washington, USA (Figure 1a), and the

centerpiece of Mount Rainier National Park (MORA). At 4,392 m (14,411 feet), Mount Rainier is

the highest peak in Washington state and is flanked by 29 named glaciers covering ~78 km^2

(Beason 2017). The Paradise Glacier lies on the southern side of the peak—with lower and

upper extents at 2,100 and 2,585 m, respectively—and is one of the smaller glaciers in MORA

at ~0.6 km^2. Like most glaciers around the world, Mount Rainier's glaciers are in rapid retreat.

From 1896 to 2015, glaciated area on Mount Rainier has decreased by ~40%, a loss of 50.6

km^2 (Beason 2017). The Paradise Glacier, which was part of a larger Paradise-Stevens Glaciers

complex in 1896, has retreated by 87% over the same period (Beason 2017). Still, Mount

Rainier and its glaciers remain some of the snowiest places on Earth. Annual snowfall at 1,646

m—as measured near the park visitor center at Paradise (Figure 1)—has averaged 1,620 cm

(639 inches) from 1920 - 2020 (National Park Service 2021a). On Mount Rainier, treeline occurs

from roughly 1,829 - 1,981 m (6,000 - 6,500 feet).

Experimental design

From 17 June 2021 to 31 October 2021, we deployed wildlife cameras (Browning Strike Force

850 Extreme, Browning, USA) along the western margin of the Paradise Glacier (Figures 1b,c).

However, we only analyzed data through 25 September 2021 because variable but often

persistent snowpack either buried or obscured cameras during the final five weeks of the field

season (25 September - 31 October 2021). We placed six cameras along an elevational

transect from 2,164 to 2,469 m (7,100 - 8,100 feet) and re-visited the cameras every 3-5 weeks

to adjust positioning, exchange memory cards, and perform similar routine activities. We

focused our cameras to target the margins of the glacier where talus and boulders give way to

ice. Because the Paradise Glacier is such a dynamic environment during summer, with a

dramatically shifting physical surface (both in area and elevation), we moved the cameras as

needed to maintain close proximity (generally 0-10 m) to the ice surface. When triggered,

cameras took bursts of five photos in quick succession with a 30-second rest period between

triggers. While our camera traps were largely reliable and stayed in working order throughout

the study, we did experience a memory card failure at C5 from 20 August - 25 September. Due

to the variable terrain and study goals, we made no attempts to control for the amount of area

each camera could "see" which likely affected the comparability of our results across cameras.

To account for the challenges of the alpine glacier ecosystem, including the need to move the

cameras throughout the melt season, we employed a unique camera trap design (Figures 2a,b).

Cameras were mounted to 5-gallon (18.9 liter) cylindrical Igloo$^{©}$ coolers that were covered in

camouflage duct tape with small, camouflage ratchet straps. The coolers were filled with snow

and loose rocks from the trapping area and sealed with ⅛" coated steel cable and a small U-

clamp. Because we expected visitors to notice our cameras despite efforts to blend them into

the landscape, we added a small label to the top of each trap noting its purpose with contact

information. The low-profile but robust footprint of our camera traps when the coolers were filled

with snow/rock provided a stable platform that remained in place despite high winds and an

unprecedented heat wave that occurred during the course of our study and rapidly melted a

large amount of snowpack (Pelto *et al.* 2022).

Animal identification and data analysis

Images were visually inspected for any evidence of vertebrate detection (see example images in

Figure 3). When humans or wildlife (birds and mammals) were detected, they were identified to

the lowest taxonomic level possible by consulting guides for Mount Rainier National Park and

the Pacific Northwest as well as our own research team's expertise in mammals (A.A. and L.W.,

specifically) and birds (J.B., N.A.P., and P.W., specifically).Given the proximity of our cameras

to glacier ice or perennial snow, we considered any species recorded in this study to have a link

to glacial habitats. This definition is slightly expanded relative to Rosvold (2016), the most

similar study to ours, which focused exclusively on birds and mammals that were directly on

snow and ice.

When multiple images from a camera burst included the same organism(s), it was only counted

once. The same is true for multiple bursts within minutes of each other. However, because our

camera array was arranged along an elevational transect, it is possible, and perhaps likely, that

some humans or wildlife were detected multiple times across different cameras. We did not

attempt to control for multiple identifications in space nor time beyond the methods described

above. Photos of the research team were also excluded from all analyses. Observations were

totaled and analyzed for each of the 14 weeks of the study. We note that the final "week" (week

14) contained two extra days due to where the study endpoint fell (25 September) relative to the

end of that particular week (23 September).

To test for a relationship between human and wildlife occurrences in our data set, we U a series

of statistical analyses in R v3.6.3 (R Core Team 2021). First, we grouped all human and wildlife

observations separately for each camera on a week-by-week basis. Then, we tested for

normality of the human and wildlife data sets using a Shapiro-Wilk normality test ("shapiro.test").

Because both data sets were significantly different from normal (*P,* Shapiro-Wilk < 0.001), we

used correlation analyses with a Spearman's rank correlation. We tested for a correlation

between the number of human versus wildlife detections across all cameras and weeks in the

study.

Results

Across 770 trap nights we had 307 human and wildlife detections in the Mount Rainier alpine.

We identified at least 16 vertebrate species (seven birds, nine mammals) using glacier-

associated habitats (Table 1). Aside from a single camera failure that was unrelated to the trap

itself, our low-cost, alpine camera trap design performed well. Across our data set, humans

were the most frequently observed taxon ($N = 208$), followed by American pipits ($N = 18$),

mountain goats ($N = 13$), hoary marmot ($N = 13$), and wolverines ($N = 8$; Table 1, Figure 4). Just

under one-fifth of birds could not be identified to species but for those that were identified,

American pipits and white-tailed ptarmigan accounted for more than half of the observations.

However, a number of nondescript brown and gray "blurs" were also recorded, of which many

were likely small birds (i.e. gray-crowned rosy finch, horned lark), and thus our results for birds

are likely underestimations.

We observed the most humans at the third-highest camera, C3 ($N = 59$), and the most wildlife at

the lowest camera, C6 ($N = 36$; Figure 4). Interestingly, C6 also recorded the fewest humans (N

$= 8$) and the most taxonomically diverse wildlife: five mammals (golden-mantled ground squirrel,

hoary marmot, human, mountain goat, wolverine), three identifiable birds (American pipit, gray-

crowned rosy finch, white-tailed ptarmigan), and several unidentifiable passerines and other

birds. Because we did not control for the amount of area each camera could "see," comparisons

across cameras (and detection periods since cameras were adjusted throughout the study),

should be interpreted with caution.

We observed the highest number of detections during the first week of the study (18-24 June

2021) but this pattern was overwhelmingly driven by humans (Figure 4). Indeed, ~54% of all

human observations were within this period. For wildlife (mammals and birds), no clear temporal

pattern was present. At cameras where humans and wildlife were regularly observed (e.g., C4,

C5), we observed a temporal separation with humans being seen early in the season from mid-

June to mid-July and other taxa appearing later (after mid-July; Figure 4). However, we did not

observe a correlation—positive or negative—between the presence of humans and wildlife

across all cameras and weeks of the study (P, Spearman's rho = 0.106).

During the study, we identified three taxa of conservation concern in Washington state:

wolverines (including two kits playing on the glacier), a Cascade red fox, and white-tailed

ptarmigan (Figures 3e-g). We observed wolverines at the three lowest cameras (C4-C6) in late

July/early August, a Cascade red fox at our C4 camera in mid-August, and white-tailed

ptarmigans at the two highest cameras (C1-C2) and the lowest camera (C6), including a pair of

ptarmigans together in mid-July at C2.

Discussion

Discussion

Mountain glaciers are in rapid decline around the world (Hugonnet *et al.* 2021). Concurrent with

this recession of glacier ice is the loss of critical habitat for glacier-associated biodiversity (Stibal

et al. 2020). To date, most of the research emphasis in glacier biology has centered on

understanding the distribution and functional roles of microbial life (Hotaling *et al.* 2021). This is

not surprising since microbes are the dominant life form in glacier environments and play critical

roles in the cryosphere, even altering the albedo of ice and impacting its rate of melt (Ganey *et*

al. 2017; Hotaling *et al.* 2021). However, overlooked in this emphasis have been larger

vertebrates that use glaciers and adjacent habitats for travel, nesting, foraging, and other key

aspects of their life histories (Rosvold 2016). This lack of research emphasis stems, in part,

from the difficulties associated with studying wildlife in rugged, glaciated ecosystems, including

the limitations of traditional methods when working in harsh, snow-covered habitat.

In this study, we deployed an array of novel camera traps along the margin of a rapidly receding

mountain glacier to test our camera design and gain a better understanding of human and

wildlife use of these habitats. We found considerable success with our approach and identified

an array of birds and mammals, from songbirds to mesocarnivores, highlighting the ecological

complexity and potential importance of glaciers and alpine ecosystems to mountain biodiversity.

Indeed, many species observed in this study (e.g., American kestrel) have not been previously

linked to glacier habitats, raising questions about the degree to which components of their life

history have been overlooked. We also found limited, albeit non-significant, evidence for

interactions between human and wildlife usage of glacier ecosystems with little to no overlap

between the two groups in space and time.

From a conservation and management perspective, we observed three wildlife species that are

particularly notable: wolverine (*Gulo gulo*), Cascade red fox (*Vulpes vulpes cascadensis*), and

white-tailed ptarmigan (*Lagopus leucura*). All three are listed as "Species of Greatest

Conservation Need" under the Washington State Wildlife Action Plan (Washington Department

of Fish and Wildlife 2021), with a need for monitoring and distribution information specifically

highlighted for wolverine and Cascade red fox. The successful identification of all three species

during the course of this study shows the value of wildlife camera trapping above treeline.

Furthermore, wolverines are exceptionally rare in the South Cascades with the family (adult

female, two kits) observed in this study likely representing just the third family in the range in a

century (National Park Service 2021). Given their rarity in MORA and the South Cascades, we

assume that all of our observations were of the same wolverines, which were seen on three

separate cameras during the week of 30 July 2021. This burst of activity might indicate

successful foraging in the area as wolverines are known to use rendezvous sites near food

caches, particularly when kits are young (Inman *et al.* 2012). Notably, we only observed

mesocarnivores (e.g., wolverines) where we also observed common prey species (e.g.,

marmots; Table 1).

While our camera trap design and array in the Mount Rainier alpine was successful, it was not

without difficulties. Indeed, the challenging nature of camera trapping in a high-alpine landscape

should not be overlooked. In addition to highly variable, and often difficult, weather conditions,

the landscape does not lend itself well to traditional camera trapping approaches for three

reasons. (1) The physical landscape dramatically changes during summer due to snowmelt.

During 2021, seasonal snow depth at a substantially lower elevation than our study area peaked

in late winter at 5.36 m (213 inches; Paradise SNOTEL #679). This seasonal snow completely

melted by mid-July, effectively lowering the habitat surface by several meters during the first

month of our study (mid-June to mid-July). Thus, camera trap locations were limited either to

areas that were snow-free early in the melt season or required revisiting as the snow surface

changed. In our case, portability was important as we moved cameras small distances

throughout the season to track shifting ice margins. Changing snow depth also meant that

framing the view, and controlling for the field of vision across cameras, was nearly impossible.

For instance, one view that was completely snow-covered in early July was entirely rock and

talus within a matter of days. The end of the season offers another complication as seasonal

snow can accumulate rapidly, particularly in "sheltered" areas that are ideal for obscuring

camera traps. Thus, we recommend that similar future studies in the PNW or comparable areas

conclude by the end of September or that researchers devise a strategy for finding snow-

covered cameras and/or keeping them free of snow. (2) Hiking and/or game trails are minimal or

non-existent above treeline. Thus, an exceptionally large and consistent survey area exists with

little to no mechanism for targeting areas where wildlife may be more abundant. To this end, a

denser camera array would have certainly yielded better quantifications of vertebrate presence

and interactions (e.g., links between carnivores and prey or humans and wildlife). Finally, (3) we

did not observe many songbirds on the glacier and in surrounding habitats despite well-known

connections between the two (e.g., Hotaling *et al.* 2020). This was likely due to a combination of

our cameras not being sensitive enough to detect small birds and images with potential birds in-

flight being too blurry for identification. Indeed, a number of photographs included unidentifiable

brown or gray blurs that we could not confidently categorize as a bird or something else (e.g., a

flying insect). One way to overcome this, or at least improve the detection rate of small birds,

would be to pair cameras with acoustic recorders at each site so song could be used in

combination with (or in lieu of) imagery.

Conclusion

In this study, we showed that a simple, cost-effective camera trap design could reveal a rich

community of birds and mammals, including species of conservation concern, are using difficult-

to-study glacier habitat in western North America. Given the relatively small number of camera

traps placed in the vast alpine habitat of Mount Rainier with no game-trails and few natural

chokepoints to increase success, it is likely that our results are a significant underestimation of

the true scale of biodiversity in these habitats. For instance, in a global meta-analysis, Rosvold

(2016) identified 36 species of birds and mammals that are known to use snow and ice habitats

for some portion of their life history. Here, we observed 16 species on a small glacier during a

single summer with a limited array. Thus, our current understanding clearly underestimates the

true scale of wildlife associations with mountain glaciers. As glaciers continue to rapidly recede

worldwide, there is a growing need to document and understand these wildlife-ice connections

to properly account for the biodiversity impacts of cryosphere decline.

Acknowledgements

This research was conducted under Mount Rainier National Park permit #MORA-2021-SCI-

0016. We thank Ben Lee for assistance in the field and park staff for their support during the

permitting process. We also thank David Sousa and the co-owners of Tatoosh Meadows for

allowing us to camp on their property during this research. We gratefully acknowledge that our

research was conducted on the traditional lands of the Cowlitz, Muckleshoot, Nisqually,

Puyallup, Squaxin Island, and Yakama tribes.

Data availability

The data and scripts used in this study are available in this study's GitHub accession:

https://github.com/scotthotaling/glacier_cams.

Declaration of funding

This project was funded by grants from the Northwest Ecological Research Institute and

Western North American Naturalist. S.H. was also supported by NSF award #OPP-1906015.

References

Anesio AM and Laybourn-Parry J (2012) Glaciers and ice sheets as a biome. *Trends Ecol Evol* **27**, 219-225.

Anesio AM, Lutz S, Chrismas NA, and Benning LG (2017) The microbiome of glaciers and ice sheets. *NPJ biofilms and microbiomes* **3**, 10.

Antor RJ (1995) The importance of arthropod fallout on snow patches for the foraging of high-alpine birds. *Journal of Avian Biology*, 81-85.

Beason S (2017) Change in glacial extent at mount rainier national park from 1896–2015. *Natural Resource Report NPS/MORA/NRR-2017/1472, National Park Service.*

Finger D, Heinrich G, Gobiet A, and Bauder A (2012) Projections of future water resources and their uncertainty in a glacierized catchment in the swiss alps and the subsequent effects on hydropower production during the 21st century. *Water Resources Research* **48**.

Freitag S, Hobson C, Biggs H, and Van Jaarsveld A Testing for potential survey bias: The effect of roads, urban areas and nature reserves on a southern african mammal data set. 1998 pp. 119-127. (Cambridge University Press)

Ganey GQ, Loso MG, Burgess AB, and Dial RJ (2017) The role of microbes in snowmelt and radiative forcing on an alaskan icefield. *Nature Geoscience* **10**, 754.

Giersch JJ, Hotaling S, Kovach RP, Jones LA, and Muhlfeld CC (2017) Climate-induced glacier and snow loss imperils alpine stream insects. *Global Change Biology* **23**, 2577-2589. doi: 10.1111/gcb.13565.

Hardy SP, Hardy DR, and Gil KC (2018) Avian nesting and roosting on glaciers at high elevation, cordillera vilcanota, peru. *The Wilson Journal of Ornithology* **130**, 940-957.

Hotaling S, Finn DS, Giersch JJ, Weisrock DW, and Jacobsen D (2017) Climate change and alpine stream biology: Progress, challenges, and opportunities for the future. *Biological Reviews* **92**, 2024-2045. doi: 10.1111/brv.12319.

Hotaling S, Foley ME, Zeglin LH, Finn DS, Tronstad LM, Giersch JJ, Muhlfeld CC, and Weisrock DW (2019) Microbial assemblages reflect environmental heterogeneity in alpine streams. *Global Change Biology* **25**, 2576-2590. doi: 10.1111/gcb.14683.

Hotaling S, Lutz S, Dial RJ, Anesio AM, Benning LG, Fountain AG, Kelley JL, McCutcheon J, Skiles SM, and Takeuchi N (2021) Biological albedo reduction on ice sheets, glaciers, and snowfields. *Earth-science reviews* **220**, 103728.

Hotaling S, Wimberger PH, Kelley JL, and Watts HE (2020) Macroinvertebrates on glaciers: A key resource for terrestrial food webs? *Ecology* **101**, e02947. doi: 10.1002/ecy.2947.

Hugonnet R, McNabb R, Berthier E, Menounos B, Nuth C, Girod L, Farinotti D, Huss M, Dussaillant I, and Brun F (2021) Accelerated global glacier mass loss in the early twenty-first century. *Nature* **592**, 726-731.

Inman RM, Magoun AJ, Persson J, and Mattisson J (2012) The wolverine's niche: Linking
reproductive chronology, caching, competition, and climate. *Journal of Mammalogy* **93**, 634-
644. doi: 10.1644/11-MAMM-A-319.1.

Milner AM, Khamis K, Battin TJ, Brittain JE, Barrand NE, Füreder L, Cauvy-Fraunié S, Gíslason
GM, Jacobsen D, and Hannah DM (2017) Glacier shrinkage driving global changes in
downstream systems. *Proceedings of the National Academy of Sciences* **114**, 9770-9778.

Muhlfeld CC, Cline TJ, Giersch JJ, Peitzsch E, Florentine C, Jacobsen D, and Hotaling S (2020)
Specialized meltwater biodiversity persists desite widespread deglaciation. *Proceedings of the*
National Academy of Sciences **117**, 12208-12214. doi: 10.1073/pnas.2001697117.

National Park Service (2021) Second family of wolverines documented at mount rainier national
park [online].

Pelto MS, Dryak M, Pelto J, Matthews T, and Perry LB (2022) Contribution of glacier runoff
during heat waves in the nooksack river basin USA. *Water* **14**, 1145.

R Core Team (2021) R: A language and environment for statistical computing.

Rich LN, Beissinger SR, Brashares JS, and Furnas BJ (2019) Artificial water catchments
influence wildlife distribution in the mojave desert. *The Journal of Wildlife Management* **83**, 855-
865.

Rosvold J (2016) Perennial ice and snow-covered land as important ecosystems for birds and
mammals. *Journal of Biogeography* **43**, 3-12.

Rovero F, Tobler M, and Sanderson J (2010) Camera trapping for inventorying terrestrial
vertebrates. *Manual on field recording techniques and protocols for all taxa biodiversity*
inventories and monitoring. The Belgian National Focal Point to the Global Taxonomy Initiative
8, 100-128.

Schano C, Niffenegger C, Jonas T, and Korner-Nievergelt F (2021) Hatching phenology is
lagging behind an advancing snowmelt pattern in a high-alpine bird. *Sci Rep* **11**, 1-11.

Shain DH, Novis PM, Cridge AG, Zawierucha K, Geneva AJ, and Dearden PK (2021) Five
animal phyla in glacier ice reveal unprecedented biodiversity in new zealand's southern alps.
Sci Rep **11**, 1-7.

Stibal M, Bradley JA, Edwards A, Hotaling S, Zawierucha K, Rosvold J, Lutz S, Cameron KA,
Mikucki JA, and Kohler TJ (2020) Glacial ecosystems are essential to understanding
biodiversity responses to glacier retreat. *Nature Ecology & Evolution* **4**, 686-687. doi:
10.1038/s41559-020-1163-0.

Washington Department of Fish and Wildlife (2021) Species in washington [online].
Anesio AM and Laybourn-Parry J (2012) Glaciers and ice sheets as a biome. *Trends Ecol Evol*
27, 219-225.

Anesio AM, Lutz S, Chrismas NA, and Benning LG (2017) The microbiome of glaciers and ice
sheets. *NPJ biofilms and microbiomes* **3**, 10.

Antor RJ (1995) The importance of arthropod fallout on snow patches for the foraging of high-
alpine birds. *Journal of Avian Biology*, 81-85.

Beason S (2017) Change in glacial extent at mount rainier national park from 1896–2015.
Natural Resource Report NPS/MORA/NRR-2017/1472, National Park Service.

Finger D, Heinrich G, Gobiet A, and Bauder A (2012) Projections of future water resources and
their uncertainty in a glacierized catchment in the swiss alps and the subsequent effects on
hydropower production during the 21st century. *Water Resources Research* **48**.

Freitag S, Hobson C, Biggs H, and Van Jaarsveld A Testing for potential survey bias: The effect
of roads, urban areas and nature reserves on a southern african mammal data set. 1998 pp.
119-127. (Cambridge University Press)

Ganey GQ, Loso MG, Burgess AB, and Dial RJ (2017) The role of microbes in snowmelt and
radiative forcing on an alaskan icefield. *Nature Geoscience* **10**, 754.

Giersch JJ, Hotaling S, Kovach RP, Jones LA, and Muhlfeld CC (2017) Climate-induced glacier
and snow loss imperils alpine stream insects. *Global Change Biology* **23**, 2577-2589. doi:
10.1111/gcb.13565.

Hardy SP, Hardy DR, and Gil KC (2018) Avian nesting and roosting on glaciers at high
elevation, cordillera vilcanota, peru. *The Wilson Journal of Ornithology* **130**, 940-957.

Hotaling S, Finn DS, Giersch JJ, Weisrock DW, and Jacobsen D (2017) Climate change and
alpine stream biology: Progress, challenges, and opportunities for the future. *Biological Reviews*
92, 2024-2045. doi: 10.1111/brv.12319.

Hotaling S, Foley ME, Zeglin LH, Finn DS, Tronstad LM, Giersch JJ, Muhlfeld CC, and Weisrock
DW (2019) Microbial assemblages reflect environmental heterogeneity in alpine streams. *Global
Change Biology* **25**, 2576-2590. doi: 10.1111/gcb.14683.

Hotaling S, Lutz S, Dial RJ, Anesio AM, Benning LG, Fountain AG, Kelley JL, McCutcheon J,
Skiles SM, and Takeuchi N (2021) Biological albedo reduction on ice sheets, glaciers, and
snowfields. *Earth-science reviews* **220**, 103728.

Hotaling S, Wimberger PH, Kelley JL, and Watts HE (2020) Macroinvertebrates on glaciers: A
key resource for terrestrial food webs? *Ecology* **101**, e02947. doi: 10.1002/ecy.2947.

Hugonnet R, McNabb R, Berthier E, Menounos B, Nuth C, Girod L, Farinotti D, Huss M,
Dussaillant I, and Brun F (2021) Accelerated global glacier mass loss in the early twenty-first
century. *Nature* **592**, 726-731.

Inman RM, Magoun AJ, Persson J, and Mattisson J (2012) The wolverine's niche: Linking
reproductive chronology, caching, competition, and climate. *Journal of Mammalogy* **93**, 634-
644. doi: 10.1644/11-MAMM-A-319.1.

Milner AM, Khamis K, Battin TJ, Brittain JE, Barrand NE, Füreder L, Cauvy-Fraunié S, Gíslason
GM, Jacobsen D, and Hannah DM (2017) Glacier shrinkage driving global changes in
downstream systems. *Proceedings of the National Academy of Sciences* **114**, 9770-9778.

492
493 Muhlfeld CC, Cline TJ, Giersch JJ, Peitzsch E, Florentine C, Jacobsen D, and Hotaling S (2020)
494 Specialized meltwater biodiversity persists desite widespread deglaciation. *Proceedings of the*
495 *National Academy of Sciences* **117**, 12208-12214. doi: 10.1073/pnas.2001697117.
496
497 National Park Service (2021) Second family of wolverines documented at mount rainier national
498 park [online].
499
500 Pelto MS, Dryak M, Pelto J, Matthews T, and Perry LB (2022) Contribution of glacier runoff
501 during heat waves in the nooksack river basin USA. *Water* **14**, 1145.
502
503 R Core Team (2021) R: A language and environment for statistical computing.
504
505 Rich LN, Beissinger SR, Brashares JS, and Furnas BJ (2019) Artificial water catchments
506 influence wildlife distribution in the mojave desert. *The Journal of Wildlife Management* **83**, 855-
507 865.
508
509 Rosvold J (2016) Perennial ice and snow-covered land as important ecosystems for birds and
510 mammals. *Journal of Biogeography* **43**, 3-12.
511
512 Rovero F, Tobler M, and Sanderson J (2010) Camera trapping for inventorying terrestrial
513 vertebrates. *Manual on field recording techniques and protocols for all taxa biodiversity*
514 *inventories and monitoring. The Belgian National Focal Point to the Global Taxonomy Initiative*
515 **8**, 100-128.
516
517 Schano C, Niffenegger C, Jonas T, and Korner-Nievergelt F (2021) Hatching phenology is
518 lagging behind an advancing snowmelt pattern in a high-alpine bird. *Sci Rep* **11**, 1-11.
519
520 Shain DH, Novis PM, Cridge AG, Zawierucha K, Geneva AJ, and Dearden PK (2021) Five
521 animal phyla in glacier ice reveal unprecedented biodiversity in new zealand's southern alps.
522 *Sci Rep* **11**, 1-7.
523
524 Stibal M, Bradley JA, Edwards A, Hotaling S, Zawierucha K, Rosvold J, Lutz S, Cameron KA,
525 Mikucki JA, and Kohler TJ (2020) Glacial ecosystems are essential to understanding
526 biodiversity responses to glacier retreat. *Nature Ecology & Evolution* **4**, 686-687. doi:
527 10.1038/s41559-020-1163-0.
528
529 Washington Department of Fish and Wildlife (2021) Species in washington [online].
530
531

532 **Tables**

533 Table 1. Taxa detected on the Paradise Glacier and directly adjacent habitat of Mount Rainier, WA,

534 USA. N_{DET}: number of detections. Taxa were classified to the lowest taxonomic possible.

Species	Order, Family	Camera	N_{DET}
Birds			
American kestrel (*Falco sparverius*)	Falconiformes, Falconidae	C5	1
American pipit (*Anthus rubescens*)	Passeriformes, Motacillidae	C1-4, C6	18
Common raven (*Corvus corvax*)	Passeriformes, Corvidae	C2	1
Gray-crowned rosy finch (*Leucosticte tephrocotis*)	Passeriformes, Fringillidae	C3-4, C6	5
Horned lark (*Eremophila alpestris*)	Passeriformes, Alaudidae	C1, C3	5
Mountain bluebird (*Sialia corrucoides*)	Passeriformes, Turdidae	C4	1
White-tailed ptarmigan (*Lagopus leucura*)	Galliformes, Phasianidae	C1-2, C6	6
Unidentified passerines	Passeriformes	C6	4
Unidentified birds	n/a	C2, C4-6	4
Mammals			
Bushy-tailed wood rat (*Neotoma cinereal*)	Rodentia, Cricetidae	C1-2	2
Cascade red fox (*Vulpes vulpes cascadensis*)	Carnivora, Canidae	C4	1
Cascade golden-mantled ground squirrel (*Callospermophilus saturatus*)	Rodentia, Sciuridae	C1-2, C4, C6	2
Hoary marmot (*Marmota caligata*)	Rodentia, Sciuridae	C1, C4, C6	13
Human (*Homo sapiens*)	Primates, Hominidae	C1-6	208
Mountain goat (*Oreamnos americanus*)	Artiodactyla, Bovidae	C4-6	13
Pacific jumping mouse (*Zapus trinotatus*)	Rodentia, Cricetidae	C1	4
Wolverine (*Gulo gulo*)	Carnivora, Mustelidae	C4-6	8
Unknown *Mustela*	Carnivora, Mustelidae	C4	1

536   **Figure captions**

538
539   Figure 1. (a) Mount Rainier, Washington, USA, viewed from the south with our camera locations
540   labeled. Satellite imagery from Google Earth. (b, c) Sentinel-2 satellite imagery of our camera
541   locations when they were first deployed (b) and once the bulk of seasonal snow had melted (c).
542   The outline of the Paradise Glacier (thin green line) in (b) and (c) was approximated from
543   Beason (2017).

545
546   Figure 2. (a) A close-up view of an example camera trap on the margin of the Paradise Glacier
547   on Mount Rainier, Washington, USA. (b) A more distant view of the camera (within the red
548   square) showing how it blends into the surrounding talus. Photographs: © Scott Hotaling.

Figure 3. Examples of taxa detected with camera traps along the western margin of the
Paradise Glacier, Mount Rainier, WA, USA. (a) Common raven (*Corvus corvax*), (b) humans
(*Homo sapiens*), (c) mountain bluebird (*Sialia corrucoides*), (d) mountain goat (*Oreamnos
americanus*), (e) wolverine (*Gulo gulo*), (f) Cascade red fox (*Vulpes vulpes cascadensis*), (g)
white-tailed ptarmigan (*Lagopus leucura*), (h) hoary marmot (*Marmota caligata*).

Figure 4. Humans and wildlife observed in this study by elevation (top to bottom) and seasonal

timing (grouped on a per week basis; left to right for each taxonomic group). Observations refer

to unique sightings and the total observations for a given taxonomic group are given at the top

of the plot (N_{ALL}). The viewer should note the substantial scale difference of the y-axis for human

observations relative to wildlife. A short-term camera failure at C5 is denoted by the labeled light

pink square.