1 **Original Article**
2
3
4 **Manipulating Histone Acetylation Leads to Adverse Effects in Hemangiosarcoma Cells**
5
6

7 Tamami Suzuki [a], Keisuke Aoshima [a, *], Jumpei Yamazaki [b], Atsushi Kobayashi [a], Takashi
8 Kimura [a]
9
10 [a] *Laboratory of Comparative Pathology, Department of Clinical Sciences, Faculty of*
11 *Veterinary Medicine, Hokkaido University, Sapporo, Hokkaido, 060-0818, Japan*
12 [b] *Translational Research Unit, Veterinary Teaching Hospital, Faculty of Veterinary Medicine,*
13 *Hokkaido University, Sapporo, Hokkaido, 060-0818, Japan*
14
15
16
17
* Corresponding author. Tel.: +81 11 7065193.
E-mail address: k-aoshima@vetmed.hokudai.ac.jp (Keisuke Aoshima).

Abstract

Canine hemangiosarcoma (HSA) is a malignant tumour derived from endothelial cells.

No effective treatment has yet been developed because of the lack of understanding of its

pathogenesis. Histone acetylation, an epigenetic modification, is highly associated with

cancer pathogenesis. Manipulating histone acetylation by histone deacetylase inhibitors

(HDACi) or bromodomain and extraterminal domain inhibitors (BETi) is one approach to

treat various cancers. However, the role of histone acetylation in HSA remains unknown. This

study aimed to investigate how histone acetylation functions in HSA pathogenesis using two

HDACi, suberanilohydroxamic acid (SAHA) and valproic acid (VPA), and one BETi, JQ1, *in*

vitro and *in vivo*. Histone acetylation levels were high in cell lines and heterogeneous in

clinical cases. SAHA and JQ1 induced apoptosis in HSA cell lines. SAHA and VPA

treatment in HSA cell lines upregulated inflammatory-related genes, thereby attracting

macrophages. This implies that SAHA and VPA can induce anti-tumour immunity. JQ1

stimulated autophagy and inhibited the cell cycle. Finally, JQ1 suppressed HSA tumour cell

proliferation *in vivo*. These results suggest that HDACi and BETi can be alternative drugs for

HSA treatment. Although further research is required, this study provides useful insights for

developing new treatments for HSA.

Keywords: BETi; HDACi; hemangiosarcoma; histone acetylation

Introduction

Canine hemangiosarcoma (HSA) is an aggressive malignant tumour derived from

endothelial cells. Surgery and chemotherapy are conventional therapies for HSA, but a review

article summarised that the efficacy of these treatments was limited (Clifford et al., 2000). No

other treatments are available because little is known about the detailed molecular

pathogenesis of HSA. Thus, more basic research on HSA is warranted.

Epigenetics such as DNA methylation and histone modifications regulates

transcription without altering DNA sequences, and these mechanisms are involved in various

biological processes such as embryonic development, metabolism, and diseases, including

cancer (Kaelin and McKnight, 2013; Virani et al., 2012; Xu et al., 2021). Acetylated histone

can promote transcription by binding to bromodomain and extraterminal (BET) proteins and

loosening the chromatin structure by neutralizing the electric charges between histone and

DNA (Josling et al., 2012; Clayton et al., 2006).

In recent years, a growing number of reports have highlighted that histone

acetylation is involved in cancer pathogenesis (Calcagno et al., 2019; Liu et al., 2018).

Histone deacetylase inhibitors (HDACi) and BET inhibitors (BETi) have been used to

investigate the function of histone acetylation and their treatment efficacy. HDACi inhibits

histone deacetylase function, thereby increasing histone acetylation levels (West and

Johnstone, 2014). BETi binds to the functional domain of BET proteins through which BET

proteins cannot recognise acetylated histones (Bechter and Schöffski, 2020).

Suberanilohydroxamic acid (SAHA) and valproic acid (VPA) are HDACis that have been

reported to induce apoptosis and cell death *in vitro* and suppress tumour proliferation *in vivo*

in various human cancers (Claerhout et al., 2011; Lee et al., 2012; Uehara et al., 2012; Yagi et

al., 2010). HDACi can also improve tumour immunity, and combination therapies with

immune checkpoint inhibitors or chemotherapeutics enhance their efficacy in human cancers

(Huang et al., 2017; Kim et al., 2020; Xue et al., 2016). BETi has also been broadly

investigated as cancer drugs, including JQ1, which can cause apoptosis, cell cycle arrest, and

autophagy *in vitro*, and impede tumour growth *in vivo* (Jostes et al., 2017; Li et al., 2019;

Wen et al., 2019).

Regarding epigenetics in HSA, KDM2B—a histone demethylase—has recently been

found to promote tumour viability by enhancing the DNA damage response (Gulay et al.,

2021). However, the role of histone acetylation in HSA remains unknown. In this study, we

aimed to investigate the role of histone acetylation and the effects of HDACi and BETi on

HSA.

Materials and Methods

Histopathology and immunohistochemistry (IHC)

Tumour samples were obtained from patients presenting to Hokkaido University

Veterinary Teaching Hospital with written informed consent (Table S1). Haematoxylin and

eosin staining was performed as previously described (Maharani et al., 2018). IHC analysis

was performed as previously described (Gulay et al., 2021) with slight modifications

described below. Heat-induced antigen retrieval was performed twice in Tris-EDTA buffer

(pH 9.0) in a microwave for 5 min. Tissues were stained with anti-acetylated histone H3

antibody (1:250; #39140, Active Motif, CA, USA) overnight at 4°C. Quantification of IHC

results using QuPath was performed based on Nucleus DAB OD mean, and the values of

acetylated histone H3 in tumour cells were normalised against those in normal endothelial

cells on the same slides for further analysis (Bankhead et al., 2017).

Cell culture

Normal canine aortic endothelial cells (CnAOEC: #Cn304-05, Cell Applications, CA,

USA), seven canine HSA cell lines (JuA1, JuB2, JuB4, Re12, Re21, Ud2, and Ud6. (Murai et

al., 2012), ISOS-1 obtained from the Cell Resource Center for Biomedical Research Cell

Bank (Tohoku University) (Masuzawa et al., 1998), and RAW264 obtained from RIKEN

Bioresource Center were cultured with Dulbecco's Modified Eagle Medium High Glucose

(DMEM; #044-29765, Fujifilm Wako Pure Chemical Industries, Osaka, Japan) supplemented

with 10% foetal bovine serum (FBS; #S1580-500, Biowest, UT, USA) and penicillin–

streptomycin solution (#168-23191, Fujifilm Wako) at 37□ with 5% CO_2.

Drug preparation

SAHA (#10009929, Cayman Chemical, MI, USA) was initially dissolved in

dimethyl sulfoxide (DMSO) to a stock solution of 150 mg/ml. For *in vivo* studies, the stock

solution of SAHA was diluted with 30% polyethylene glycol 400 and 5% Tween 80 in double

distilled water. VPA (#227-01071, Fujifilm Wako) was dissolved in PBS. JQ1 (#T117580,

Toronto Research Chemicals, Toronto, Canada) was initially dissolved in DMSO to a stock

solution of 50 mg/ml. For *in vivo* studies, the stock solution of JQ1 was diluted with 5%

Tween 80 in PBS.

Cell viability assay

Two thousand cells were seeded in 96-well cell culture plates and cultured in 100 µL

culture medium. The next day, they were treated with DMSO, SAHA, or JQ1. Survival rates

and growth inhibition rates were analysed as previously described (Aoshima et al., 2018;

Morita et al., 2019)[1].

Apoptosis assay

---

[1] See: NCI 60 Cell Five-Dose Screen.
    https://dtp.cancer.gov/discovery_development/nci-60/methodology.htm (Accessed 21 October, 2021).

Apoptosis assay was performed using the FITC Annexin V Apoptosis Detection Kit

□ (#556547, BD Biosciences, NJ, USA) according to the manufacturer's instructions. Briefly,

JuB4 and Re12 cells were treated with 10 μM SAHA or JQ1 for 24 h, and detached from

dishes using TrypLE Express Enzyme (#12604013; Thermo Fisher Scientific). One million

cells were suspended in the binding buffer and divided into stained with annexin V. Cells

were analysed with a BD FACSVerse flow cytometer (BD Biosciences), and the results were

analysed with FCS Express 4 software (De Novo Software, CA, USA).

RNA-Sequencing

ISOS-1 cells were treated with 1 μM SAHA for 24 h, 1 μM JQ1 for 36 h, or 2 mM

VPA for 36 h in triplicate. Total RNA was extracted with a NucleoSpin RNA isolation kit

(#740955.50, Macherey-Nagel GmbH & Co. Düren, Germany) according to the

manufacturer's instructions. RNA samples were submitted to Kazusa DNA Research

Institution (Chiba, Japan) for further analyses. RNA-seq libraries were constructed using a

QuantSeq 3'mRNA-Seq Library Prep Kit (LEXOGEN, Vienna, Austria) and sequenced with

the NextSeq500 (Illumina, CA, USA) to generate a minimum of two million single-end 75-bp

reads. Sequencing reads were mapped to the mm39 mouse reference genome using STAR,

and expression levels were estimated using RSEM (Dobin and Gingeras, 2015; Li and Dewey,

2011). Differential expression and gene expression profiles were analysed by edgeR and

GSEA v4.1.0 (Robinson et al., 2009; Mootha et al., 2003; Subramanian et al., 2005),

respectively.

SA-β-gal staining

SA-β-gal staining was performed using a Senescence β-Galactosidase Staining Kit

(#9860, Cell Signaling Technology, MA, USA) according to the manufacturer's instructions.

Briefly, ISOS-1 was treated with DMSO or 10 µM JQ1 for 24 h and fixed with 4%

paraformaldehyde (PFA) for 10 min at room temperature (RT). Cells were stained with β-

Galactosidase Staining Solution containing 5-bromo-4-chloro-3-indolyl β-D-

galactopyranoside (X-Gal: #16495, Cayman Chemical) for 13 h at 37□. Cells were covered

with 70% glycerol. Positive cells were counted manually using an inverted microscope

(BIOREVO BZ-9000; KEYENCE, Tokyo, Japan).

Cell cycle analysis

ISOS-1, JuB2, JuB4, and Re12 cells were treated with DMSO or JQ1 for 24 h. ISOS-

1 was treated with 10 µM JQ1, and other cell lines were treated with 5 µM JQ1. Cell cycle

analysis was performed as previously described (Gulay et al., 2021).

Migration assay

JuB4 and Re12 were treated with DMSO, 2 µM SAHA, or 2 mM VPA for 48 h in 6-

well plates. Then, the medium was replaced to the one without inhibitors, and the cells were

co-cultured with RAW264 seeded on ThinCert Cell Culture Inserts (#657630, Greiner Bio-

One) for 24 h. RAW264 cells on the culture surface of the inserts were removed with a cotton

swab. Migrated RAW264 cells on the bottom side of the inserts were fixed with 4% PFA for

30 min at RT and then stained with 0.01% crystal violet for 30 min at RT. The number of

cells was counted manually in 10 fields at 200× under a light microscope (BH-2; Olympus,

Tokyo, Japan).

Protein extraction and western blotting

Protein extraction and western blotting were performed as previously described

(Gulay et al., 2021). Antibodies used in this study are listed in Table S2.

RNA extraction and reverse transcription quantitative polymerase chain reaction (RT-qPCR)

Primers and samples were prepared as described previously (Gulay et al., 2021).
qPCR was performed using the primers listed in Table S3. Results were normalised according
to the geometric mean of reference genes (*RPL32* and *HPRT* for canine genes and *Hprt* and
Tbp for murine genes), which were selected from potential internal controls by geNorm
analysis (Fig. S1) (Vandesompele et al., 2002). Relative expression levels were calculated by
setting the DMSO-treated cells as the control.

Animal studies

All mouse experiments were performed under the guidelines of Hokkaido University
(protocol number: 20-0083), which follows the ARRIVE guidelines. Five-week-old female
Balb/c and KSN/Slc mice (Japan SLC, Inc. Shizuoka, Japan) were used. A day before tumour
cell inoculation, KSN/Slc mice were treated with 100 μL 2.5 mg/ml anti-asialo GM1 (014-
09801, Fujifilm Wako) to increase the success rate of JuB2 transplantation (Yoshino et al.,
2000). Five million cells were inoculated subcutaneously in both flanks of Balb/c and
KSN/Slc mice, respectively, which were anesthetised with 0.3 mg/kg medetomidine (Domitor,
ZENOAQ, Tokyo, Japan), 4 mg/kg midazolam (Dormicum, Maruishi Pharmaceutical Co.,
Ltd. Osaka, Japan) and 5 mg/kg butorphanol (Vetorphale, Meiji Seika Pharma Co., Ltd.
Tokyo, Japan). After tumour cell inoculation, mice were awoken by intraperitoneal injection
of 3 mg/kg atipamezole (Atipame, Kyoritsu Seiyaku Corporation, Tokyo, Japan). Tumour
volumes were calculated using the formula: Volume = (Length × Width^2)/2. When the tumour
volume reached 100 mm^3, mice were treated with the vehicle control or inhibitors. Mice were
intraperitoneally injected with SAHA (150 mg/kg) daily, VPA (200 mg/kg) five times weekly,
or JQ1 (50 mg/kg) daily. Animals in all experimental groups were examined at least twice

weekly to check their health status and to measure tumour sizes. Balb/c mice were euthanized

with CO_2 when tumours reached 1000 mm^3 in volume. KSN/Slc mice were euthanized with

CO_2 11 or 12 days after treatment initiation.

Statistical analysis

Statistical analyses were performed with R (version 4.1.0). Fisher's exact test was

used to analyse the ratio of acetylated H3 staining levels. Student's *t*-test or Mann-whiteny test

were used to analyse the differences between two groups, whereas Tukey's test was used to

analyse differences among multiple groups. The significance of the differences in the survival

curves of ISOS-1 tumours was analysed using the log-rank test. Overall JuB2 tumour growth

was compared using two-way ANOVA.

Results

Histone acetylation levels in HSA cells are high in vitro and heterogeneous in vivo

We first analysed histone acetylation levels in seven HSA cell lines and in CnAOEC

normal canine endothelial cells. We found that global acetylation of H2B, H3, and H4 was

significantly enriched in all HSA cell lines compared with CnAOEC (Fig. 1A). Then, we

examined global acetylated histone H3 levels in 10 clinical HSA cases. These cases were

classified into three types according to their histological pattern: solid, capillary, or cavernous

(Kim et al., 2015). Five cases had a single histological pattern, while the other five possessed

multiple proliferation patterns (Fig, 1B). We also classified histone H3 acetylation levels into

three groups according to the normalized values: negative (< 0.5), weak (0.5–2.0), and strong

(> 2.0) (Figs. 1C and 1D). The results showed that the histone H3 acetylation levels of tumour

cells were heterogeneous, and the ratio was different depending on their histological pattern

even in the same tumour tissue (Figs. 1E and 1F). Furthermore, the ratio of tumour cells

classified as negative was the highest in the solid pattern followed by the capillary pattern,

and the lowest in the cavernous pattern (Fig. 1F). These results suggest that histone H3

acetylation levels are heterogeneous in tumour cells and are associated with their

histopathological patterns.

SAHA and JQ1 induce apoptosis in HSA cell lines

Next, to examine the roles of histone acetylation in HSA pathogenesis, we modified

histone acetylation levels using HDACi (SAHA and VPA) and BETi (JQ1). First, we

evaluated whether global histone acetylation levels were altered by these inhibitors. SAHA

and VPA treatments increased global histone acetylation levels in both canine and murine

HSA cell lines, whereas JQ1 decreased them in canine HSA cell lines but not in ISOS-1 (Fig.

2A). Then, we examined the effects of HDACi and BETi on HSA cell proliferation. SAHA

and JQ1 treatment significantly decreased the number of cells, while VPA treatment did not

affect their proliferation (Fig. 2B). IC_{50} values of the inhibitors in HSA cell lines were almost

or less than one-tenth of those in CnAOEC, which indicated that HSA cell lines were more

susceptible to these inhibitors than CnAOEC (Fig. 2C). Then, to investigate whether the cells

actively died by SAHA or JQ1, we examined apoptosis markers: cleaved caspase 3 and

annexin V. The results demonstrated that SAHA and JQ1 increased cleaved caspase 3

expression levels and annexin V positivity in HSA cell lines except for JQ1-treated ISOS-1

cells (Figs. 2D–F). Finally, we evaluated total protein expression and phosphorylation levels

of cell survival proteins AKT and ERK. Although the results were different among the cell

lines, either total or phosphorylated AKT or ERK levels were downregulated by SAHA and

JQ1 but not by VPA (Fig. S2B). These results suggested that SAHA and JQ1 induced

apoptosis in HSA cell lines except for ISOS-1 treated with JQ1, and that VPA did not induce

cell death in all HSA cell lines.

SAHA and VPA activate inflammatory responses in HSA cell lines

Next, we further investigated the mechanisms of their effects in HSA cells. GSEA

results of ISOS-1 cells indicated that inflammatory-related gene expression was positively

correlated in SAHA and VPA treatment (Fig. 3A). Consistent with this, *Il6*, *Cxcl1*, *Ccl2*, *Ccl7*,

and *Oas1a* were upregulated more than two-fold following treatment with either SAHA or

VPA or both (Fig. 3B). We wanted to extrapolate these findings to canine HSA cell lines, but

CXCL1 and *CCL2* do not exist in the canine genome and *CCL7* expression was not detected.

However, *OAS1*, the orthologue of murine *Oas1a*, and other inflammatory-related genes were

upregulated more than two-fold in canine HSA cell lines treated with SAHA and/or VPA (Fig.

3C). Inflammatory responses in tumour cells can induce immune cell migration, leading to

host immune responses to tumour tissues. Therefore, to corroborate whether SAHA and VPA

treatment in HSA cell lines can attract immune cells, a macrophage migration assay was

performed using RAW264 cells (Fig. 3D). RAW264 cells migrated when they were co-

cultured with SAHA-treated JuB4 or VPA-treated Re12 (Fig. 3E).

These results suggest that SAHA and VPA treatment can induce inflammatory

responses in HSA cell lines. This could attract macrophages, although it depends on the

combination of HSA cell lines and HDACi.

JQ1 treatment induces autophagy and impedes the cell cycle in HSA cell lines

We also investigated the genome-wide gene expression changes in JQ1-treated ISOS-

1 cells. GSEA results showed that autophagy-related gene expression and cell cycle related-

gene expression were positively and negatively correlated, respectively (Figs 4A and 4B).

These expression changes were validated by RT-qPCR (Fig. 4C). LC3-II, an active autophagy

marker, was expressed at significantly higher levels in all JQ1-treated HSA cell lines except

for Re12 compared with DMSO controls (Fig. 4D).

Then, we generated growth inhibition curves through which we could evaluate

whether the cells actively died or slowed their proliferation by JQ1 treatment. The results

showed that growth inhibition rates were positive even when the cells were treated with 10

µM JQ1, which indicated that the cell cycle was impeded, and active cell death did not occur

(Fig. 5A). BrdU and PI staining revealed that proliferating cells (BrdU+ cells) were

significantly decreased in both murine and canine HSA cell lines by JQ1 treatment (Fig. 5B).

propidium iodide (PI) signals showed that the percentage of dead cells was slightly increased

by JQ1 in ISOS-1, but its extent was significantly lower than in canine HSA cell lines (Fig.

5C). We also examined cell senescence by performing SA-β-gal staining in ISOS-1 treated

with DMSO or JQ1. The results indicated that JQ1-treated ISOS-1 showed a significantly

higher number of positive cells than DMSO-treated cells (Fig. 5D).

These results suggest that autophagy activation and cell cycle downregulation are

induced by JQ1 treatment, and that cell senescence but not apoptosis is caused by JQ1

treatment in ISOS-1.

JQ1 suppresses HSA tumour growth in vivo

Finally, to evaluate the treatment effects of SAHA, VPA, and JQ1 *in vivo*, we treated

ISOS-1 tumour-bearing Balb/c mice with SAHA, VPA, or JQ1. The results showed that only

JQ1 slowed tumour growth and extended the survival rates (Fig. 6A and B). We also treated

JuB2 tumour-bearing KSN/Slc mice with JQ1 and showed that JQ1 also suppressed JuB2

tumour growth (Fig. 6C). These treatments did not cause body weight loss in mice (Fig. S3).

These results suggest that JQ1 has suppressive effects on HSA tumour growth i*n vivo*.

Discussion

HSA tumour cells in clinical cases showed various H3 acetylation levels even in the

same tumour tissue. Moreover, this heterogeneity differed depending on the histological

pattern. This can be explained by the differentiation status of the tumour cells. Global histone

acetylation levels have been reported to be increased as cells differentiate (Ashktorab et al.,

2009; Golob et al., 2008). Furthermore, HSA tumour cells forming the solid pattern are

considered to be undifferentiated (Göritz et al., 2013). Given that the percentage of tumour

cells classified as H3 acetylation-negative was high in the solid pattern, they could be more

undifferentiated than other tumour cells. Further research is required to clarify whether global

histone acetylation levels reflect the differentiation status of HSA tumour cells.

We demonstrated that SAHA and VPA upregulated inflammatory-related genes.

Moreover, SAHA-treated JuB4 and VPA-treated Re12 attracted RAW264 cells. Several

reports demonstrated that HDACi activated immune responses and enhanced immune cell

functions (Cox et al., 2020; Laengle et al., 2020; Sun et al., 2019); thus, SAHA and VPA

could also be used as alternative therapeutics for HSA. In our study, however, SAHA and

VPA treatment of ISOS-1 in an immunocompetent mouse model did not suppress tumour

growth. This implies that the condition we tested was not able to sufficiently induce the

immune response to attack tumour cells. Considering this, co-treatment with other drugs such

as doxorubicin or immune checkpoint inhibitors would be beneficial for HSA treatment by

further stimulating immune responses (Le et al., 2017; Tubbs and Nussenzweig, 2017；Igase

et al., 2020; Maekawa et al., 2017). In the present study, we only showed the results from the

RAW264 migration assay, and further studies are required to investigate if canine-derived

macrophages exhibit the same results and if the migrated macrophages possess anti-tumour

functions.

We revealed that JQ1 suppressed tumour growth in HSA cell lines. Several reports

demonstrated that JQ1 or other BETi had anti-tumour effects, and clinical trials are ongoing

(Shorstova et al., 2021; Wen et al., 2019). Given that our results could not demonstrate

tumour regression upon JQ1 treatment, further research is required to enhance its efficacy and

to identify another BETi that can regress HSA tumour growth *in vivo*.

Conclusions

We demonstrated that HDACi and BETi can be used for HSA treatment. Although

further research is required, this work provides useful insights for developing new HSA

treatments.

Conflict of interest statement

The authors declare that there is no conflict of interest in this study.

Acknowledgements

We would like to extend our sincerest gratitude to Dr. Hiroki Sakai, Gifu University,

for providing canine hemangiosarcoma cell lines. We also acknowledge the efforts of Drs.

Mitsuyoshi Takiguchi, Shintaro Kobayashi, and Hironobu Yasui, Faculty of Veterinary

Medicine, Hokkaido University for giving useful pieces of advice and constructive discussion.

We are grateful to all the members of the Laboratory of Comparative Pathology, Faculty of

Veterinary Medicine, Hokkaido University for their helpful discussions, encouragement, and

support. This research was supported by KAKENHI Grant-in-Aid for Young Scientists (KA,

Number 20K15654) provided by Japan Society for the Promotion of Science. We thank H.

Nikki March, PhD, from Edanz (https://jp.edanz.com/ac) for editing a draft of this manuscript.

References
Aoshima, K., Fukui, Y., Gulay, K.C.M., Erdemsurakh, O., Morita, A., Kobayashi, A., Kimura,
T., 2018. Notch2 signal is required for the maintenance of canine hemangiosarcoma
cancer stem cell-like cells. BMC Vet. Res. 14.

340
341 Ashktorab, H., Belgrave, K., Hosseinkhah, F., Brim, H., Nouraie, M., Takkikto, M., Hewitt,
342   S., Lee, E.L., Dashwood, R.H., Smoot, D., 2009. Global histone H4 acetylation and
343   HDAC2 expression in colon adenoma and carcinoma. Dig. Dis. Sci. 54, 2109–2117.
344
345 Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G., Dunne, P.D.,
346   McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al., 2017. QuPath: Open
347   source software for digital pathology image analysis. Sci. Rep. 7.
348
349 Bechter, O., Schöffski, P., 2020. Make your best BET: The emerging role of BET inhibitor
350   treatment in malignant tumors. Pharmacol. Ther.
351
352 Bellarosa, D., Bressan, A., Bigioni, M., Parlani, M., Maggi, C.A., Binaschi, M., 2012.
353   SAHA/Vorinostat induces the expression of the CD137 receptor/ligand system and
354   enhances apoptosis mediated by soluble CD137 receptor in a human breast cancer cell
355   line. Int. J. Oncol. 41, 1486–1494.
356
357 Benton, C.B., Fiskus, W., Bhalla, K.N., 2017. Targeting Histone Acetylation: Readers and
358   Writers in Leukemia and Cancer. Cancer J. (United States).
359
360 Calcagno, D.Q., Wisnieski, F., Da Silva Mota, E.R., Maia De Sousa, S.B., Costa Da Silva,
361   J.M., Leal, M.F., Gigek, C.O., Santos, L.C., Rasmussen, L.T., Assumpção, P.P., et al.,
362   2019. Role of histone acetylation in gastric cancer: Implications of dietetic compounds
363   and clinical perspectives. Epigenomics.
364
365 Claerhout, S., Lim, J.Y., Choi, W., Park, Y.Y., Kim, K.H., Kim, S.B., Lee, J.S., Mills, G.B.,
366   Cho, J.Y., 2011. Gene expression signature analysis identifies vorinostat as a candidate
367   therapy for gastric cancer. PLoS One 6.
368
369 Clayton, A.L., Hazzalin, C.A., Mahadevan, L.C., 2006. Enhanced Histone Acetylation and
370   Transcription: A Dynamic Perspective. Mol. Cell.
371
372 Clifford, C.A., Mackin, A.J., Henry, C.J., 2000. Treatment of canine hemangiosarcoma: 2000
373   and beyond. J. Vet. Intern. Med.
374
375 Cox, D.J., Coleman, A.M., Gogan, K.M., Phelan, J.J., Ó Maoldomhnaigh, C., Dunne, P.J.,
376   Basdeo, S.A., Keane, J., 2020. Inhibiting Histone Deacetylases in Human Macrophages
377   Promotes Glycolysis, IL-1β, and T Helper Cell Responses to Mycobacterium
378   tuberculosis. Front. Immunol. 11.
379
380 Dobin, A., Gingeras, T.R., 2015. Mapping RNA-seq Reads with STAR. Curr. Protoc.
381   Bioinforma. 51, 11.14.1-11.14.19.
382
383 Golob, J.L., Paige, S.L., Muskheli, V., Pabon, L., Murry, C.E., 2008. Chromatin remodeling
384   during mouse and human embryonic stem cell differentiation. Dev. Dyn. 237, 1389–
385   1398.
386
387 Göritz, M., Müller, K., Krastel, D., Staudacher, G., Schmidt, P., Kühn, M., Nickel, R., Schoon,
388   H.A., 2013. Canine splenic haemangiosarcoma: Influence of metastases, chemotherapy
389   and growth pattern on post-splenectomy survival and expression of angiogenic factors. J.

390   Comp. Pathol. 149, 30–39.
391

392   Gulay, K.C.M., Aoshima, K., Shibata, Y., Yasui, H., Yan, Q., Kobayashi, A., Kimura, T.,
393   2021. KDM2B promotes cell viability by enhancing DNA damage response in canine
394   hemangiosarcoma. J. Genet. Genomics 48, 618–630.
395

396   Guo, P., Chen, W., Li, H., Li, M., Li, L., 2018. The Histone Acetylation Modifications of
397   Breast Cancer and their Therapeutic Implications. Pathol. Oncol. Res.
398

399   Han, W., Sun, J., Feng, L., Wang, K.F., Li, D., Pan, Q., Chen, Y., Jin, W., Wang, X., Pan, H.,
400   et al., 2011. Autophagy inhibition enhances daunorubicin-induced apoptosis in K562
401   cells. PLoS One 6.
402

403   Hou, Y.J., Dong, L.W., Tan, Y.X., Yang, G.Z., Pan, Y.F., Li, Z., Tang, L., Wang, M., Wang,
404   Q., Wang, H.Y., 2011. Inhibition of active autophagy induces apoptosis and increases
405   chemosensitivity in cholangiocarcinoma. Lab. Investig. 91, 1146–1157.
406

407   Huang, P., Almeciga-Pinto, I., Jarpe, M., van Duzer, J.H., Mazitschek, R., Yang, M., Jones,
408   S.S., Quayle, S.N., 2017. Selective HDAC inhibition by ACY-241 enhances the activity
409   of paclitaxel in solid tumor models. Oncotarget 8, 2694–2707.
410

411   Igase, M., Nemoto, Y., Itamoto, K., Tani, K., Nakaichi, M., Sakurai, M., Sakai, Y., Noguchi,
412   S., Kato, M., Tsukui, T., et al., 2020. A pilot clinical study of the therapeutic antibody
413   against canine PD-1 for advanced spontaneous cancers in dogs. Sci. Rep. 10.
414

415   Jones, P.A., Baylin, S.B., 2007. The Epigenomics of Cancer. Cell.
416

417   Josling, G.A., Selvarajah, S.A., Petter, M., Duffy, M.F., 2012. The role of bromodomain
418   proteins in regulating gene expression. Genes (Basel).
419

420   Jostes, S., Nettersheim, D., Fellermeyer, M., Schneider, S., Hafezi, F., Honecker, F.,
421   Schumacher, V., Geyer, M., Kristiansen, G., Schorle, H., 2017. The bromodomain
422   inhibitor JQ1 triggers growth arrest and apoptosis in testicular germ cell tumours in vitro
423   and in vivo. J. Cell. Mol. Med. 21, 1300–1314.
424

425   Kabeya, Y., Mizushima, N., Ueno, T., Yamamoto, A., Kirisako, T., Noda, T., Kominami, E.,
426   Ohsumi, Y., Yoshimori, T., 2000. LC3, a mammalian homologue of yeast Apg8p, is
427   localized in autophagosome membranes after processing. EMBO J. 19, 5720–5728.
428

429   Kaelin, W.G., McKnight, S.L., 2013. Influence of metabolism on epigenetics and disease.
430   Cell.
431

432   Kim, J.H., Graef, A.J., Dickerson, E.B., Modiano, J.F., 2015. Pathobiology of
433   hemangiosarcoma in dogs: Research advances and future perspectives. Vet. Sci.
434

435   Kim, Y.D., Park, S.M., Ha, H.C., Lee, A.R., Won, H., Cha, H., Cho, S., Cho, J.M., 2020.
436   HDAC Inhibitor, CG-745, enhances the anti-cancer effect of anti-PD-1 immune
437   checkpoint inhibitor by modulation of the immune microenvironment. J. Cancer 11,
438   4059–4072.
439

440 Laengle, J., Kabiljo, J., Hunter, L., Homola, J., Prodinger, S., Egger, G., Bergmann, M., 2020.
441   Histone deacetylase inhibitors valproic acid and vorinostat enhance trastuzumab-
442   mediated antibody-dependent cell-mediated phagocytosis. J. Immunother. Cancer 8.
443

444 Le, D.T., Durham, J.N., Smith, K.N., Wang, H., Bartlett, B.R., Aulakh, L.K., Lu, S.,
445   Kemberling, H., Wilt, C., Luber, B.S., et al., 2017. Mismatch repair deficiency predicts
446   response of solid tumors to PD-1 blockade. Science (80-.). 357, 409–413.
447

448 Lee, Y.J., Won, A.J., Lee, J., Jung, J.H., Yoon, S., Lee, B.M., Kim, H.S., 2012. Molecular
449   mechanism of SAHA on regulation of autophagic cell death in tamoxifen-resistant MCF-
450   7 breast cancer cells. Int. J. Med. Sci. 9, 881–893.
451

452 Li, B., Dewey, C.N., 2011. RSEM: Accurate transcript quantification from RNA-Seq data
453   with or without a reference genome. BMC Bioinformatics 12.
454

455 Li, F., Yang, C., Zhang, H.B., Ma, J., Jia, J., Tang, X., Zeng, J., Chong, T., Wang, X., He, D.,
456   et al., 2019. BET inhibitor JQ1 suppresses cell proliferation via inducing autophagy and
457   activating LKB1/AMPK in bladder cancer cells. Cancer Med. 8, 4792–4805.
458

459 Liu, K.Y., Wang, L.T., Hsu, S.H., 2018. Modification of epigenetic histone acetylation in
460   hepatocellular carcinoma. Cancers (Basel).
461

462 Maekawa, N., Konnai, S., Takagi, S., Kagawa, Y., Okagawa, T., Nishimori, A., Ikebuchi, R.,
463   Izumi, Y., Deguchi, T., Nakajima, C., et al., 2017. A canine chimeric monoclonal
464   antibody targeting PD-L1 and its clinical efficacy in canine oral malignant melanoma or
465   undifferentiated sarcoma. Sci. Rep. 7.
466

467 Maharani, A., Aoshima, K., Onishi, S., Gulay, K.C.M., Kobayashi, A., Kimura, T., 2018.
468   Cellular atypia is negatively correlated with immunohistochemical reactivity of CD31
469   and vWF expression levels in canine hemangiosarcoma. J. Vet. Med. Sci. 80, 213–218.
470

471 Masuzawa, M., Fujimura, T., Tsubokawa, M., Nishiyama, S., Katsuoka, K., Terada, E.,
472   Kunita, S., Sakurai, Y., Kato, H., 1998. Establishment of a new murine-phenotypic
473   angiosarcoma cell line (ISOS- 1). J. Dermatol. Sci. 16, 91–98.
474

475 Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J.,
476   Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al., 2003. PGC-1α-
477   responsive genes involved in oxidative phosphorylation are coordinately downregulated
478   in human diabetes. Nat. Genet. 34, 267–273.
479

480 Morita, A., Aoshima, K., Gulay, K.C.M., Onishi, S., Shibata, Y., Yasui, H., Kobayashi, A.,
481   Kimura, T., 2019. High drug efflux pump capacity and low DNA damage response
482   induce doxorubicin resistance in canine hemangiosarcoma cell lines. Res. Vet. Sci. 127,
483   1–10.
484

485 Murai, A., Asa, S.A., Kodama, A., Hirata, A., Yanai, T., Sakai, H., 2012. Constitutive
486   phosphorylation of the mTORC2/Akt/4E-BP1 pathway in newly derived canine
487   hemangiosarcoma cell lines. BMC Vet. Res. 8.
488

Pagiatakis, C., Musolino, E., Gornati, R., Bernardini, G., Papait, R., 2021. Epigenetics of

aging and disease: a brief overview. Aging Clin. Exp. Res.

Pratap, J., Akech, J., Wixted, J.J., Szabo, G., Hussain, S., McGee-Lawrence, M.E., Li, X.,
Bedard, K., Dhillon, R.J., Van Wijnen, A.J., et al., 2010. The histone deacetylase
inhibitor, vorinostat, reduces tumor growth at the metastatic bone site and associated
osteolysis, but promotes normal bone loss. Mol. Cancer Ther. 9, 3210–3220.

Proske, J., Walter, L., Bumes, E., Hutterer, M., Vollmann-Zwerenz, A., Eyüpoglu, I.Y.,
Savaskan, N.E., Seliger, C., Hau, P., Uhl, M., 2016. Adaptive immune response to and
survival effect of temozolomide-and valproic acid-induced autophagy in glioblastoma.
Anticancer Res. 36, 899–906.

Qiu, H., Jackson, A.L., Kilgore, J.E., Zhong, Y., Chan, L.L.Y., Gehrig, P.A., Zhou, C., Bae-
Jump, V.L., 2015. JQ1 suppresses tumor growth through downregulating LDHA in
ovarian cancer. Oncotarget 6, 6915–6930.

Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2009. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26, 139–
140.

Shorstova, T., Foulkes, W.D., Witcher, M., 2021. Achieving clinical success with BET
inhibitors as anti-cancer agents. Br. J. Cancer.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al., 2005. Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. Proc. Natl. Acad. Sci. U. S. A. 102, 15545–15550.

Sun, T., Li, Y., Yang, W., Wu, H., Li, X., Huang, Y., Zhou, Y., Du, Z., 2019. Histone
deacetylase inhibition up-regulates MHC class I to facilitate cytotoxic T lymphocyte-
mediated tumor cell killing in glioma cells. J. Cancer 10, 5638–5645.

Tubbs, A., Nussenzweig, A., 2017. Endogenous DNA Damage as a Source of Genomic
Instability in Cancer. Cell.

Uehara, N., Kanematsu, S., Miki, H., Yoshizawa, K., Tsubura, A., 2012. Requirement of p38
MAPK for a cell-death pathway triggered by vorinostat in MDA-MB-231 human breast
cancer cells. Cancer Lett. 315, 112–121.

Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., Speleman,
F., 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. Genome Biol. 3.

Virani, Shama, Virani, Shami, Colacino, J.A., Kim, J.H., Rozek, L.S., 2012. Cancer
epigenetics: a brief review. ILAR J.

Wen, N., Guo, B., Zheng, H., Xu, L., Liang, H., Wang, Q., Wang, D., Chen, X., Zhang, S., Li,
Y., et al., 2019. Bromodomain inhibitor jq1 induces cell cycle arrest and apoptosis of
glioma stem cells through the VEGF/PI3K/AKT signaling pathway. Int. J. Oncol. 55,
879–895.

540
541 West, A.C., Johnstone, R.W., 2014. New and emerging HDAC inhibitors for cancer treatment.
542   J. Clin. Invest.
543
544 Xia, Q., Sung, J., Chowdhury, W., Chen, C.L., Höti, N., Shabbeer, S., Carducci, M.,
545   Rodriguez, R., 2006. Chronic administration of valproic acid inhibits prostate cancer cell
546   growth in vitro and in vivo. Cancer Res. 66, 7237–7244.
547
548 Xu, R., Li, C., Liu, X., Gao, S., 2021. Insights into epigenetic patterns in mammalian early
549   embryos. Protein Cell.
550
551 Xue, K., Gu, J.J., Zhang, Q., Mavis, C., Hernandez-Ilizaliturri, F.J., Czuczman, M.S., Guo, Y.,
552   2016. Vorinostat, a histone deacetylase (HDAC) inhibitor, promotes cell cycle arrest and
553   re-sensitizes rituximab- and chemo-resistant lymphoma cells to chemotherapy agents. J.
554   Cancer Res. Clin. Oncol. 142, 379–387.
555
556 Yagi, Y., Fushida, S., Harada, S., Kinoshita, J., Makino, I., Oyama, K., Tajima, H., Fujita, H.,
557   Takamura, H., Ninomiya, I., et al., 2010. Effects of valproic acid on the cell cycle and
558   apoptosis through acetylation of histone and tubulin in a scirrhous gastric cancer cell line.
559   J. Exp. Clin. Cancer Res. 29.
560
561 Yang, W., Hosford, S.R., Traphagen, N.A., Shee, K., Demidenko, E., Liu, S., Miller, T.W.,
562   2018. Autophagy promotes escape from phosphatidylinositol 3-kinase inhibition in
563   estrogen receptor-positive breast cancer. FASEB J. 32, 1222–1235.
564
565 Yoshino, H., Ueda, T., Kawahata, M., Kobayashi, K., Ebihara, Y., Manabe, A., Tanaka, R.,
566   Ito, M., Asano, S., Nakahata, T., et al., 2000. Natural killer cell depletion by anti-asialo
567   GM1 antiserum treatment enhances human hematopoietic stem cell engraftment in
568   NOD/Shi-scid mice. Bone Marrow Transplant. 26, 1211–1216.
569
570 Zheng, H., Zhao, W., Yan, C., Watson, C.C., Massengill, M., Xie, M., Massengill, C., Noyes,
571   D.R., Martinez, G. V., Afzal, R., et al., 2016. HDAC inhibitors enhance T-cell
572   chemokine expression and augment response to PD-1 immunotherapy in lung
573   adenocarcinoma. Clin. Cancer Res. 22, 4119–4132.
574
575

Figure legends

Figure 1. Histone acetylation levels in HSA cells are high *in vitro* and heterogenous *in*

vivo.

(A) Western blot analysis of acetylated histone H2B, H3, and H4. (B) Representative images

of HE and IHC staining in clinical HSA cases. Scale bars = 50 μm. (C) Box plots for

normalized nucleus DAB OD mean values in 10 clinical HSA cases. (D) Representative IHC

images of the classifications according to normalised acetylated H3 intensities in each

histological pattern. Arrowheads indicate the nuclei of tumour cells. (E) Stacked bar graphs

indicating the ratio of each staining level in 10 clinical cases. (F) Stacked bar graphs

comparing the ratio of each staining level according to the histological patterns in all cases or

within each case. *$P < 0.05$, **$P < 0.01$. Fisher's exact test.

Figure 2. SAHA and JQ1 induce apoptosis in HSA cell lines.

(A) Western blot analysis of acetylated H2B, H3, and H4 in canine HSA cell lines (JuB2,

JuB4, Re12) and a murine HSA cell line (ISOS-1) treated with DMSO, SAHA, JQ1, or VPA.

(B) Phase contrast images of canine and murine HSA cell lines 48 h after treatment with

DMSO, SAHA, VPA, or JQ1. Scale bars = 100 μm. (C) Survival curves (top) and IC_{50} values

(bottom) of canine and murine HSA cell lines treated with SAHA or JQ1 for 60 h. All

samples were analysed in triplicate. Data are plotted as average percentages ± SD. (D)

Western blot analysis of cleaved caspase 3 in canine and murine HSA cells treated with

DMSO, SAHA, or JQ1. Uncropped images of JuB4 and Re12 are shown in Figure S2A. (E)

Annexin V staining of canine and murine HSA cells treated with DMSO, SAHA, or JQ1. (F)

Percentages of apoptotic cells (annexin V-positive) in canine and murine HSA cells treated

with DMSO, SAHA, or JQ1. All samples were analysed in triplicate. Data are plotted as

average percentages ± SD. *$P < 0.05$, **$P < 0.01$. Tukey's *t* test.

Figure 3. SAHA and VPA activate inflammatory responses in HSA cell lines.

(A) (Left) GSEA results in ISOS-1 treated with SAHA (top) or VPA (bottom). Inflammatory-related gene sets are highlighted in red. The number of genes in each gene set is shown next to each bar. (Right) Representative GSEA plots of enriched gene sets in ISOS-1 treated with SAHA (top) or VPA (bottom). (B) RT-qPCR results of genes related to inflammatory responses in ISOS-1 treated with SAHA or VPA. (C) RT-qPCR results of inflammatory-related genes in JuB4 and Re12 treated with SAHA or VPA. (D) Graphical images of migration assay. (E) Quantitative analysis of migrated RAW264 cells co-cultured with SAHA or VPA-treated Re12. *$P < 0.05$, **$P < 0.01$. Tukey's t test.

Figure 4. JQ1 treatment induces autophagy in canine and murine HSA cell lines.

(A) Positively (top) or negatively (bottom) enriched gene sets by GSEA in ISOS-1 treated with JQ1 compared with DMSO-treated cells. The number of genes in each gene set is shown next to each bar. (B) Representative GSEA plots of positively (top) and negatively (bottom) enriched gene sets. (C) RT-qPCR of genes related to autophagy (top) and cell cycle (bottom) in ISOS-1 treated with DMSO or JQ1. (D) Western blot analysis of LC3 protein in canine and murine HSA cell lines treated with DMSO or JQ1.

Fig. 5. JQ1 treatment impedes the cell cycle in canine and murine HSA cell lines.

(A) Growth inhibition curve of ISOS-1 treated with JQ1. All samples were analysed in triplicate. Data are plotted as average percentages ± SD. (B) Density plots (left) and quantitative analysis (right) of BrdU-positive cells in canine and murine HSA cell lines treated with DMSO or JQ1. All samples were analysed in triplicate. (C) Histograms of PI-positive cells (left) and quantitative analysis of dead cells (right) in canine and murine HSA

626 cell lines treated with DMSO and JQ1. All samples were analysed in triplicate. (D) (Left)

627 Representative images of SA-β-gal staining in ISOS-1 treated with DMSO or JQ1.

628 Arrowheads indicate positive cells. (Right) Quantitative analysis of SA-β-gal-positive cells.

629 Positive cells were counted in 10 fields under 200× magnification with a light microscope.

630 The average numbers were plotted with ± SD. Scale bars = 100 μm. $*P < 0.05$, $**P < 0.01$.

631 Student's t-test.

633 **Figure 6. JQ1 suppresses HSA tumour growth *in vivo*.**

634 (A) Kaplan–Meier survival curves of ISOS-1-inoculated Balb/c mice treated with DMSO

635 (n=5), SAHA (n=5), VPA (n=4), or JQ1 (n=5). Days were counted after initiating drug

636 treatment. P-values were calculated using the log-rank test. (B) Individual growth curves of

637 ISOS-1 tumours in Balb/c mice treated with DMSO, SAHA, VPA, or JQ1 after treatment

638 initiation. (C) (Left) Individual tumour growth curves of JuB2 tumours in KSN/Slc mice

639 treated with DMSO or JQ1. The day when the treatment began was defined as day 1. $***P <$

640 0.001. Two-way ANOVA. (Right) Box plots of relative tumour volumes on the last day in

641 KSN/Slc mice treated with DMSO or JQ1. Relative tumour volumes were calculated by

642 dividing the tumour volume on the last day by that at day 1. $*P < 0.05$, Mann-whitney U test.

Fig.1

Fig.2

Fig.3

Fig.5

Fig.6