A genetic model of congenital intestinal atresia in medaka (*Oryzias latipes*) implicates

Mypt1 in epithelial organisation

Daisuke Kobayashi[1],*, Akihiro Urasaki[1],[2], Tetsuaki Kimura[3], Satoshi Ansai[4], Kazuhiko

Matsuo[1], Hayato Yokoi[5], Shigeo Takashima[6], Tadao Kitagawa[7], Takahiro Kage[8], Takanori

Narita[9], Tomoko Jindo[8], Masato Kinoshita[10], Kiyoshi Naruse[11], Yoshiro Nakajima[1], Masaki

Shigeta[1], Shinichiro Sakaki[1], Satoshi Inoue[1], Rie Saba[12], Kei Yamada[12], Takahiko

Yokoyama[1], Yuji Ishikawa[13], Kazuo Araki[14], Yumiko Saga[8],[15],[16], Hiroyuki Takeda[8],[17],*,

Kenta Yashiro[1],*

Affiliations:

[1] Department of Anatomy and Developmental Biology, Kyoto Prefectural University of

Medicine, 465 Kajii-cho, Kamigyo-ku, Kyoto 602-8566, Japan

[3] Human Genetics Laboratory, National Institute of Genetics, 1111 Yata, Mishima, Shizuoka,

Japan

[4] Graduate School of Life Sciences, Tohoku University, Sendai 980-8577, Japan

[5] Graduate School of Agricultural Science, Tohoku University, Sendai 980-8572, Japan

[6] Institute for Glyco-core Research (iGCORE)/Life Science Research Centre, Gifu

University, 1-1 Yanagido, Gifu 501-1193, Japan

[7] Program in Environmental Management, Graduate School of Agriculture, Kindai

University, 3327-204 Nakamachi, Nara, Nara 631-8505, Japan

[8] Department of Biological Sciences, Graduate School of Science, The University of Tokyo,

Bunkyo-ku, Hongo, Tokyo, 113-0033, Japan

[9] Laboratory of Molecular Biology, Department of Veterinary Medicine, College of

Bioresource Sciences, Nihon University, 1866 Kameino, Fujisawa, Kanagawa 252-0880,

Japan

[10] Department of Applied Biosciences, Graduate School of Agriculture, Kyoto University,

Kyoto 606-8502, Japan

[11] Laboratory of Bioresources, National Institute for Basic Biology, 38 Nishigonaka,

Myodaiji, Okazaki 444-8585, Aichi, Japan

[12] Department of Radiology, Kyoto Prefectural University of Medicine, 465 Kajii-cho,

Kamigyo-ku, Kyoto 602-8566, Japan

[13] Research Centre for Radiation Protection, National Institute of Radiological Sciences, 4-9-

1 Anagawa, Inage-ku, Chiba 263-8555, Japan

[14] Research Center for Aquatic Breeding, National Research Institute of Aquaculture,

Fisheries Research Agency, 224 Hiruda, Tamaki-cho, Watarai, Mie 519-0423, Japan

[15] Department of Gene Function and Phenomics, Mammalian Development Laboratory,

National Institute of Genetics, Mishima 411-8540, Japan

[16] Department of Genetics, The Graduate University for Advanced Studies, SOKENDAI,

Mishima 411-8540, Japan

*Address correspondence to: Daisuke Kobayashi (kdaisuke@koto.kpu-m.ac.jp), Hiroyuki

Takeda (htakeda@bs.s.u-tokyo.ac.jp) and Kenta Yashiro (kyashiro@koto.kpu-m.ac.jp)

Present address:

[2] Graduate School of Science, Technology and Innovation, Kobe University

[17] Faculty of Life Sciences, Kyoto Sangyo University, Kyoto 603-8555, Japan

Keywords: intestinal atresia, Mupt1, disease model, actomyosin regulation, intestinal

development

Conflict of interest disclosure: The authors declare no competing interests.

Funding statement: This work was supported by JSPS (Japan Society for the Promotion of

Science) KAKENHI Grant # 15K08139 and 21K08601 and NIBB Individual Collaborative

Research Program (14-376) to DK.

Abstract

Congenital intestinal atresia (IA) is a birth defect characterised by the absence or closure of

part of the intestine. Although genetic factors are implicated, mechanistic understanding has

been hindered by the lack of suitable animal models. Here, we describe a medaka (*Oryzias*

latipes) mutant, generated by N-ethyl-N-nitrosourea (ENU) mutagenesis, that develops IA

during embryogenesis. Positional cloning identified a nonsense mutation in *mypt1,* encoding

myosin phosphatase target subunit 1. Mutant embryos exhibited ectopic accumulation of F-

actin and phosphorylated myosin regulatory light chain (Mrlc) in the intestinal epithelium,

consistent with disrupted actomyosin regulation. These cytoskeletal abnormalities were

accompanied by epithelial disorganisation without notable alterations in cell proliferation,

motility, or apoptosis. Inhibition of *myh11a*, encoding smooth muscle (SM) myosin heavy

chain, ameliorated the IA phenotype but Blebbistatin treatment completely rescued the

defect, suggesting a non-contractile role prior to SM maturation. Together, these findings

demonstrate that *mypt1* loss disrupts intestinal morphogenesis through actomyosin

dysregulation. Given the recent clinical identification of IA associated with MYPT1

mutations, this medaka model offers a valuable platform to investigate the developmental and

molecular basis of *MYPT1*-associated IA in human.

Introduction

Intestinal atresia (IA) is a congenital defect characterised by complete occlusion of the

intestinal lumen, with an estimated incidence of 1.3 to 2.9 per 10,000 live births (Frischer and

Azizkhan, 2012). Affected neonates require immediate surgical intervention to restore

gastrointestinal continuity. Historically, IA has been attributed to in utero vascular accidents

that impair blood supply to the developing gut (Louw, 1959; Louw and Barnard, 1955) .

However, familial clustering and increased prevalence in individuals with trisomy 21 strongly

suggest a genetic contribution to disease aetiology (Celli, 2014; Gupta et al., 2013; Louw,

1959; Shorter et al., 2006).

Mouse models support for this genetic hypothesis. Targeted deletion of *Fgfr2IIIb* or its

ligand *Fgf10* results in IA in the absence of mesenteric vascular occlusion (Fairbanks et al.,

2004a; Fairbanks et al., 2004b; Fairbanks et al., 2005; Kanard et al., 2005) . More recently,

mutations in *PPP1R12A*, which encodes myosin phosphatase target subunit 1 (MYPT1), have

been identified in human patients with IA and other developmental anomalies, including

holoprosencephaly, urogenital malformations and persistent Müllerian duct syndrome

(PMDS).(Hughes et al., 2020) (Picard et al., 2022). Despite these insights, mechanistic

studies of MYPT1-related IA remain limited, as conventional *Mypt1* knockout mice exhibit

early embryonic lethality (Okamoto et al., 2005). Actomyosin dynamics play a vital role in

morphogenesis (Munjal and Lecuit, 2014). Non-muscle myosin II (NMII)-mediated

contractility regulates cell shape, adhesion, migration, and tissue organisation (Agarwal and

Zaidel-Bar, 2018; Munjal and Lecuit, 2014; Munjal et al., 2015; Vasquez et al., 2014).

Phosphorylation of myosin regulatory light chain Mrlc promotes contraction (Vicente-

Manzanares et al., 2009), while myosin light chain phosphatase (MLCP) protein complex

dephosphorylates Mrlc to induce relaxation. MLCP comprises a catalytic subunit Pp1cδ,

Mypt1, and a small regulatory subunit, M20 (Ito et al., 2004). Mypt1 enhances MLCP

activity and substrate specificity by modulating the catalytic domain conformation (Grassie et

al., 2011; Hartshorne et al., 2004). Mypt1 is essential for various developmental processes,

including gastrulation, axon guidance, vascular remodelling, and the morphogenesis of the

liver, pancreas, and central nervous system (Angulo-Urarte et al., 2018; Barresi et al., 2010;

Bremer and Granato, 2016; Dong et al., 2019; Gutzman and Sive, 2010; Huang et al., 2008;

Hughes et al., 2020; Kim et al., 2011; Okamoto et al., 2005; Weiser et al., 2009).

In medaka (*Oryzias latipes*), intestinal morphogenesis begins at stage (st.) 21 with medial

migration of the endodermal epithelial monolayer to the ventral midline,   (Kobayashi et al.,

2006). This monolayer then forms a bilayer of mediolaterally elongated cells. In the anterior

region, endodermal cells stack and converge at the dorsal midline to form a radial

endodermal rod—the intestinal anlage. In contrast, intermediate and posterior gut regions

undergo dorsoventral elongation and subsequent cavitation to form a tubular gut. Tube

formation proceeds in an anterior-to-posterior sequence, with the anterior and intermediate

regions becoming lumenized by st. 25, and the posterior region completing this process by st.

26. During this period, buds of accessory organs such as the liver and swim bladder also

emerge (Kobayashi et al., 2006).

In this study, we report a novel medaka mutant identified through N-ethyl-N-nitrosourea

(ENU) mutagenesis that develops IA during embryogenesis. Positional cloning identified a

nonsense mutation in *mypt1*. Despite normal endodermal migration and gut anlage formation,

mutant embryos failed to maintain epithelial continuity in the intermediate intestinal region

immediately following lumen formation. This disruption coincided with localized

accumulation of phosphorylated Mrlc (pMrlc) and F-actin, consistent with enhanced

actomyosin contractility due to loss of Mypt1 function. Our findings establish a genetically

defined vertebrate model of Mypt1-associated IA and provide new mechanistic insights into

how dysregulated cytoskeletal dynamics compromise epithelial integrity during gut

development.

Results

The *g1-4* medaka mutant develops IA

The *g1-4* medaka mutant was isolated as a recessive lethal line through our ENU

mutagenesis screening for mutations affecting embryonic development and

organogenesis.(Yokoi et al., 2007) Examination of homozygous mutant embryos revealed the

presence of IA, which developed at st. 32–33, although the penetrance rate of the IA

phenotype varied and did not reach 100% (Table 1). In adult medaka, small and large

intestines distinguishable by both morphological characteristics and distinct gene expression

profiles.(Aghaallaei et al., 2016) However, we could not distinguish small from large

intestine in the embryos a either morphologically or by the expression of known molecular

markers. Therefore, hereafter, we refer to the region between the liver bud and the cloaca

simply as the intestine. Despite of the existence of IA, the remaining intestinal tissue

appeared to be properly developed at st. 40, the hatching sage (Fig. 1A, B).

The *g1-4* allele encodes *mypt1*

We hypothesized that the inheritance mode of the *g1-4* mutant was recessive, and to identify

the affected locus, we performed positional cloning. Initial mapping using M-marker analysis

placed the mutation on linkage group 23 (LG23).(Kimura et al., 2004) We then conducted

high-resolution linkage analysis using an F2 mapping panel comprising 766 embryos. This

analysis narrowed the *g1-4* locus to a 0.13 cM interval between the markers LG23-4.4 and

LG23-4.6, corresponding to a genomic region of 24.227 kb (Fig. 1C). Within this region,

only one gene, *mypt1*, also known as *ppp1r12a* (*protein phosphatase 1 regulatory subunit*

12a) was identified in the medaka reference genome.

The open reading frame (ORF) of medaka *mypt1* (3,234 bp: LC662536) consists of 25

exons and encodes a predicted protein of 1,078 amino acids that is closely related to human

MYPT1 (NP_001137357.1). Medaka Mypt1 has three highly conserved domains, an RVxF

motif, an ankyrin repeat, and a protein kinase cGMP-dependent (Prkg) interacting domain

(Fig. 1D). Sequencing the *g1-4* allele revealed a C-to-A transversion in exon 22 (C2952A),

resulting in a premature stop codon (Fig. 1C, D). The predicted premature termination occurs

within the Prkg-interacting domain and eliminates the C-terminal leucine-zipper (LZ) motif,

which is essential for interaction with Prkg1α.(Grassie et al., 2011; Surks et al., 1999) As

interaction with Prkg1α is essential for actomyosin activation, the C2952A mutation is

expected to impair the function of Mypt1.

To confirm whether the *mypt1* mutation causes the IA phenotype observed in the *g1-4*

mutants, we tried to rescue the defect using *wild-type* (*WT*) *mypt1* mRNA. However, for

unknown reasons, we could not synthesize the full-length *mypt1* mRNA *in vitro*.

Alternatively, we generated *mypt1* knockout medaka using the CRISPR/Cas9 system.(Ansai

and Kinoshita, 2014) We isolated a mutant line possessing an 11 nucleotide deletion with a 7

nucleotide insertion (*mypt1*[#4-6], c.127_137delinsTCTGTAT, Fig. 1D, Fig. 2A, B). This

mutation creates a transcriptional frame shift that alters 45 codons before introducing a

premature stop codon at the 88th codon. Crosses between F1 *mypt1*[#4-6] heterozygotes yielded

embryos displaying IA at a frequency of 23.7% (n = 59), and we successfully established a

stable mutant line. In the F2 generation, the penetrance of the IA phenotype varied among

sibling, similar to the original *g1-4* mutant line (Table 2), with approximately 15–30% of

total siblings exhibiting IA, - higher penetrance than in the original mutant. For further

phenotypic analysis, we used this newly generated mutant line, *mypt1*[#4-6]. We could also see

dilation of the intestine upstream of the atretic region at the hutching stage, a hallmark

commonly associated with IA in *mypt1*[#4-6] embryos (Fig. 3A–D). Taken together, we

concluded that *mypt1* mutation is responsible for the IA phenotype.

Intestinal Development Is Largely Preserved in *mypt1#4-6* Mutants, Except at the

Atresia Lesion

To investigate the phenotype of the $mypt1^{\#4-6}$ mutant, we performed histological analyses

on st. 40 larvae, immediately after hatching. Serial cross-sections were prepared from *WT*

($n = 3$) and $mypt1^{\#4-6}$ mutant larvae ($n = 3$) and compared. The mutants exhibited intestinal

atresia, whereas other organs appeared to develop normally (Fig. 3E–P). To further assess the

epithelial architecture at earlier stages, we performed conventional histological analysis using

plastic sections at st. 32. These sections showed that the embryonic intestine appeared

morphologically intact in $mypt1^{\#4-6}$ mutants, except at the atretic lesion. The mesenchymal

cells, which give rise to SM and connective tissue, surrounded the endodermal epithelium

similarly to *WT*, and this organization was also preserved in unaffected regions of the mutant

intestine (Fig. 3Q–X) (Wallace et al., 2005a).   Homozygous mutants were generally lethal,

with only few individuals surviving to adulthood. Most homozygous mutants die after

hatching. Thus, intestinal atresia is very likely one of the major causes for lethality, as

affected larvae are unable to feed after hatching. However, we cannot rule out the possibility

that other factors contribute to lethality. Importantly, while most survivors exhibited poor

health, some were indistinguishable from *WT* fish (data not shown).

Cytokeratin, a marker of epithelial cells and intermediate filament protein, predominantly

localises to the apical plasma membrane in polarized epithelia (Fig. 4A) (Yang et al., 2020).

Consistent with this, the intestinal epithelium of st. 31 *WT* embryos showed strong apical

localization of cytokeratin (Fig. 4Bb). In $mypt1^{\#4-6}$ mutants, the intestinal epithelium—

excluding the atretic lesion—displayed cytokeratin distribution comparable to *WT* (Fig. 4Cb).

SM cells, labelled by SM-specific myosin (SM-myosin), properly surrounded the

intestinal epithelium in both *WT* and $mypt1^{\#4-6}$ embryos at st. 31 (Fig. 4Ba–Cb). Notably, the

blind end of the atresia lesion in mutants was also lined by SM. Furthermore, the basement

membrane, labelled by laminin, was present beneath the epithelium even at the blind-end of

the atresia in the mutant (Fig. 4Da–Eb). These findings indicate that the fundamental

structure of the gut—including epithelial apico-basolateral polarity, SM layer, and basement

membrane—is preserved normally in the region outside the atresia. This is consistent with

observations in human IA cases.

To elucidate the sequence of events that lead to atresia formation in $mypt1^{\#4-6}$ embryos,

we carefully monitored live embryos but could not detect morphological signs of IA until st.

28. We first examined endodermal development, as the endoderm gives rise to the intestine,

to determine whether endodermal defects were present before or during intestinal tube

formation. Expression of *foxa2*, a marker of the endoderm and its derivatives including the

intestine (Kobayashi et al., 2006), was indistinguishable between *WT* and $mypt1^{\#4-6}$ embryos

up to st. 25 (Fig. 5), suggesting that early endoderm formation and migration are not

disrupted in the mutant.

Importantly, at st. 26—when intestinal tube formation is completed (Kobayashi et al.,

2006)—fragmentation of the basement membrane was observed in 8 out of 9 *mypt1#4-6*

mutants (Fig. 4G), whereas none of the 11 *WT* embryos showed such fragmentation (Fig. 4F).

These results indicate that molecular events leading to IA had already begun by st. 26, even

though no obvious morphological anomalies were yet visible.

To investigate whether epithelial disintegration due to apoptosis or altered proliferation

contributed to basement membrane fragmentation, we performed acridine orange staining at

st. 27. This revealed no detectable apoptotic cells in either *WT* or $mypt1^{\#4-6}$ intestines (Fig.

6A, B). TUNEL assays also showed no significant difference between *WT* and mutant

embryos (data not shown). Furthermore, proliferation, assessed by anti-phospho-histone H3

staining, showed no significant difference in the number of mitotic cells between *WT* and

mutant intestines (Fig. 6E–I). These data suggest that neither apoptosis nor altered

proliferation underlies the basement membrane fragmentation and IA in the mutant.

Although apico-basolateral polarity was preserved in the gut epithelium, localised

epithelial-mesenchymal transition (EMT) could hypothetically result in loss of polarity and

basement membrane fragmentation, as occurs during gastrulation (Savagner, 2015). To

explore this possibility, we examined the expression of EMT-inducing transcription factors,

snai1a, *snai1b*, and *snai2* (Scheibner et al., 2021), via whole-mount in situ hybridization

(WISH). None of these genes were expressed in the intestine of either *WT* or mutant embryos

at st. 26 (Fig. 7A–F). Consistently, the subcellular localization of Pkcζ, an apical epithelial

marker, was not disrupted in *mypt1*#4-6 intestines (Fig. 4H–I). Taken together, these

observations indicate that basement membrane fragmentation and IA formation in *mypt1*[#4-6]

embryos do not result from EMT or loss of epithelial polarity.

Mypt Family Members Are Unlikely to Compensate for the Loss of MYPT1 in a

Tissue-Specific Manner

In zebrafish, *mypt1* mutations cause a spectrum of morphogenetic abnormalities,

including defects in liver and pancreas development, dilation of brain ventricles, and

mispositioning of motoneurons (Bremer and Granato, 2016; Dong et al., 2019; Gutzman and

Sive, 2010; Huang et al., 2008). In contrast, these phenotypes were not observed in *mypt1*

mutant medaka, suggesting possible species-specific differences in gene function or

redundancy within *mypt* gene family. To explore whether this phenotypic divergence could

be explained by differential expression or functional compensation by other *mypt* family

members, we identified *mypt* homologs in the medaka genome and analysed their spatial

expression patterns by WISH.

In humans, the myosin phosphatase regulatory subunit family consists of *MYPT1*

(PPP1R12A), *MYPT2* (PPP1R12B), and *MBS85* (PPP1R12C) (Gutzman and Sive, 2010;

Huang et al., 2008). To determine whether other *mypt* family genes might compensate for the

loss of *Mypt1* function in tissues other than the intestine in medaka mutants, we searched the

medaka genome and identified three members of the *mypt* family: *mypt1*, *mbs85*, and a novel

gene that diverges from the ancestral lineage shared by *mypt1* and *mypt2*, which we

designated *mypt1/2-related* (*mypt1/2-r*) (Table 3; Fig. 8). Notably, no clear *mypt2* ortholog

was found in the medaka genome. Although medaka possesses a gene annotated as *mypt3*,

this gene belongs to the *Ppp1r16* family rather than the *Ppp1r12* family and therefore was

excluded from further analysis.

We examined the expression patterns of *mypt1*, *mypt1/2-r*, and *mbs85* by WISH. Both

mypt1 and *mypt1/2-r* were ubiquitously expressed at low levels, with slightly elevated

expression in the head region. *mbs85* exhibited lower overall expression than the other two

genes but displayed a similar spatial pattern. No significant differential expression was

detected in the developing intestine (Fig. 9A–F). RT-PCR analysis of dissected head and

intestinal tissues confirmed the expression of all three genes, supporting their broad, low-

level distribution. These findings suggest that the intestinal specificity of the atresia

phenotype in *mypt1* mutants is not due to tissue-specific expression of *mypt* family genes.

Furthermore, given the ubiquitous—albeit low—expression of these genes, it is possible that

the phenotypes observed in the zebrafish *mypt1* mutant were not reproduced in the medaka

mutant due to functional compensation by related genes.

Mypt1 Is Maternally Expressed

Previous studies in zebrafish have shown that maternal knockdown of *mypt1* using

translation-blocking morpholinos leads to convergent extension (CE) defects during

gastrulation (Weiser et al., 2009). In contrast, such defects were not observed in the *mypt1[#4-6]*

mutant medaka line. To investigate whether maternal *mypt1* transcripts contribute to early

embryogenesis and mask early developmental phenotypes, we performed RT-PCR on one-

cell stage embryos. The results confirmed the presence of *mypt1* transcripts at this stage,

indicating that maternal *mypt1* mRNA is supplied during oogenesis. This maternal expression

is likely to compensate for the absence of zygotic *mypt1*, thereby preventing CE defects in

mypt1[#4-6] mutants (Fig. 9G).

Actomyosin Regulation Is Perturbed in *mypt1* Mutant Intestine

Phosphorylation of myosin regulatory light chain (Mrlc) governs the contractility of

non-muscle myosin II (NMII). (Ito et al., 2004) Dephosphorylated Mrlc reduces NMII

activity and cytoskeletal contractility, while pMrlc activates NMII. Thus, NMII-driven

contractility is determined by the balance between phosphorylation by myosin light chain

kinase (Mlck) and dephosphorylation by myosin light chain phosphatase (MLCP). *Mypt1*

encodes the regulatory subunit of MLCP that is essential for its enzymatic function.

Therefore, in *mypt1* mutants, elevated levels of pMrlc are expected.

In *WT* embryos, pMrlc exhibited weak and diffuse localisation specifically on both the

apical and basal surfaces of intestinal epithelial cells (Fig. 10A, st. 25; 10C, st. 27). As

predicted, *mypt1* mutants displayed increased pMrlc levels at st. 25 compared with *WT*

embryos (Fig. 10B). This difference was not apparent at st. 24 (data not shown). Notably,

pMrlc was not uniformly increased along the intestinal axis but appeared enriched in regions

corresponding to presumptive intestinal atresia (IA) lesions (Fig. 10B). Higher magnification

revealed that this increased pMrlc signal localized specifically to the intestinal epithelium, as

marked by *foxa2:memGFP* expression (Fig. 10Bd). By st. 27, this abnormal expression

pattern became more pronounced (Fig. 10D), and in some embryos, pMrlc was strongly

expressed along the entire intestinal tract (Fig. 10E; 2 out of 12 samples). These findings

indicate that pMrlc regulation is markedly altered in *mypt1* mutants from st. 25 onward,

particularly in regions prone to IA formation (Fig. 10F).

Because pMrlc binds NMII and promotes actin-myosin interaction, increased and mis-

localised pMrlc is likely to enhance actomyosin contractility and redistribute mechanical

forces (Vicente-Manzanares et al., 2009). In *WT* intestinal epithelial cells, F-actin was

localised to the cortical region beneath the plasma membrane and was slightly enriched at the

apical surface (Fig. 10G and I). In contrast, although intestinal lumen formation appeared

morphologically similar between *WT* and *mypt1* mutants, F-actin accumulation was elevated

in both the apical and basal cortical regions of *mypt1* mutant cells (Fig. 10H, J, and K). This

phenotype is consistent with previous reports showing that increased pMrlc enhances actin

filament assembly (Gutzman and Sive, 2010), whereas reduced pMrlc leads to decreased F-

actin levels.(Bavaria et al., 2011; Ray et al., 2007) Together, these results indicate that loss of

mypt1 function leads to upregulation of pMrlc, suggesting hyperactive actomyosin

contractility of intestinal epithelium. This aberrant mechanical activity may compromise

epithelial integrity and contribute to IA development in *mypt1* mutant embryos.

Smooth Muscle Myosin Function Partially Contributes to the Development of

Intestinal Atresia (IA)

Given that Mypt1 regulates SM myosin activity, abnormalities in the SM layer

surrounding the intestinal epithelium may play a role in the pathogenesis of intestinal atresia

(IA). In the medaka genome, we identified two homologs of *MYH11*: *myh11a* on linkage

group (LG) 1 and *myh11b* on LG 8. However, RT-PCR analysis at 2 days post-fertilization

(dpf) revealed expression of *myh11a* only (data not shown), prompting us to focus our

investigation on *myh11a*.

To disrupt *myh11a* function, we designed three overlapping single-guide RNAs

(sgRNAs) targeting the ATP-binding domain of *myh11a* (Fig. 11A). These sgRNAs were

injected into medaka embryos at the 1–2 cell stage, either alone or in combination with Cas9

nuclease. To confirm the effect of CRISPR-mediated knock-out, we performed

immunohistochemical analysis of SM myosin expression at a developmental stage when the

SM layer is well established. In the embryos injected without Cas9 at 6 dpf, SM myosin was

robustly detected in the smooth muscle layer surrounding the intestinal epithelium (Fig. 11B;

6/6 embryos). In contrast, embryos co-injected with Cas9 exhibited either a complete loss of

SM myosin expression (Fig. 11C; 2/6 embryos) or an aberrant, patchy expression pattern

(Fig. 11D; 4/6 embryos), indicating effective disruption of functional SM myosin. We then

assessed whether IA still occurred under these SM myosin-deficient conditions in $mypt1^{\#4-6}$

homozygous mutant embryos. Notably, IA was absent in 4 out of 10 $mypt1^{\#4-6}$ embryos co-

injected with *myh11a* sgRNAs and Cas9, whereas all 9 embryos injected with *myh11a*

sgRNAs alone (without Cas9) developed IA (Table 4). To further validate the role of

myh11a, we performed a morpholino (Mo)-mediated knockdown using a splice-blocking

morpholino targeting the exon 2–intron 2 junction (Mo-*myh11a*-E2I2; Fig. 11E). RT-PCR

confirmed efficient disruption of normal splicing (Fig. 11F), and partial rescue of IA was

observed in morpholino-treated embryos (Table 5). These findings suggest that *myh11a*-

dependent SM function contributes to IA pathogenesis. However, as immunohistochemical

signal of Myh11a is not detected around the region where IA occur at st. 28 when IA

becomes morphologically detectable, the SM contraction is not likely involved in IA

formation (data not shown). Non-smooth muscle expression of *myh11a* (e.g., in intestinal

epithelial cells), or alternatively, cells expressing *myh11a* located distant from the intestinal

epithelium, may contribute non-cell-autonomously to epithelial integrity at early stages in

medaka, but likely play only a partial, not primary, role (see Discussion).

Augmented Actomyosin Contraction Is Responsible for IA in *mypt1* Mutants.

To further investigate whether enhanced cytoskeletal contractility contributes to the

development of intestinal atresia (IA) in *mypt1* mutants, we tested whether blebbistatin—a

non-muscle myosin II (NMII) inhibitor—could rescue the IA phenotype in *mypt1#4-6* mutant

embryos. Embryos were treated with 50 µM blebbistatin at st. 25 for two hours (Fig. 12A).

Under these conditions, no major morphological abnormalities were observed other than a

pronounced curly tail phenotype (Fig. 12B, C). Remarkably, IA was not observed in all

treated *mypt1* mutant embryos (Fig. 12D), strongly suggesting that excessive actomyosin

contractility is a key driver of IA formation.

Discussion

Here, we report a novel animal model of IA. Through ENU mutagenesis of medaka, we

isolated the *g1-4* mutant, in which early intestinal development proceeds normally, but loss of

epithelial integrity leads to IA after the intestinal lumen has formed. Positional cloning

identified a loss-of-function mutation in the *mypt1* gene. A genome-edited *mypt1* mutant

allele (*mypt1*[#4-6]) recapitulated the *g1-4* phenotype, confirming that loss of Mypt1 function is

responsible for IA. In *mypt1* mutants, we observed elevated levels of pMrlc and abnormal

accumulation of F-actin in the developing intestine (Fig. 10). Treatment with blebbistatin, a

non-muscle myosin II (NMII) inhibitor, significantly suppressed IA in *mypt1* mutants (Fig.

12), suggesting that hyperactive cytoskeletal contraction is a likely cause of epithelial

disruption and IA formation. Given the limited understanding of the molecular

pathophysiology of human IA and the scarcity of appropriate animal models, the medaka

mypt1 mutant provides a valuable platform for elucidating the mechanisms underlying this

congenital condition.

A long-standing hypothesis proposes that IA arises from an accidental vascular injury

during development (Louw, 1959; Louw and Barnard, 1955). However, in our medaka

model, no overt abnormalities in blood circulation were observed in *mypt1*[#4-6] embryos.

Moreover, prior studies have shown that medaka embryos establish steady and robust

digestive tract blood flow by st. 30 (Fujita et al., 2006), a time point after IA formation

begins. These observations suggest that IA in *mypt1* mutants does not result from vascular

disruption, but instead arises from non-vascular mechanisms, such as dysregulated

actomyosin contractility and impaired epithelial integrity during intestinal morphogenesis.

The developmental role of Mypt1 has been documented in some species. In *Drosophila*,

mypt1 mutants have various developmental defects, including failure of dorsal

closure.(Mizuno et al., 2002; Tan et al., 2003) In these mutants, ectodermal cell sheet

movement is impaired and the subcellular localization of actomyosin regulatory proteins,

such as pMrlc and actin, is perturbed.(Mizuno et al., 2002) In zebrafish, *mypt1* mutants

display impaired migration of bone morphogenetic protein 2a (*bmp2a*)-expressing lateral

plate mesoderm (LPM) cells, causing agenesis/hypoplasia of the liver and exocrine

pancreas.(Dong et al., 2019; Huang et al., 2008) Brain ventricle formation is also affected in

the zebrafish mutant because of ectopically induced tension in the neuroepithelium

layer.(Gutzman and Sive, 2010) In these mutants, increased pMrlc levels and aberrant

localisation of NMII and F-actin are observed in the neuroepithelium. Importantly, the

upregulation and mislocalisation of actomyosin regulatory components, such as pMrlc, NMII

and F-actin, are likely to be common features in *mypt1* mutants across species, including

medaka. Furthermore, in the context of brain ventricle development in zebrafish, blebbistatin

treatment to inhibit MNII successfully rescued the *mypt1* mutant phenotype. This supports

the notion that "epithelial relaxation" regulated by Mypt1 is probably critical for proper tube

inflation, as occurs in brain ventricle formation. Our findings are consistent with the view that

loss of *mypt1* function affects actomyosin dynamics during embryogenesis.

Despite these commonalities, notable differences exist between *mypt1* mutants in

medaka and previously reported *mypt1* mutants in zebrafish. IA was not observed in zebrafish

mutants, whereas neither liver nor brain ventricle abnormalities were found in medaka

mutants. Although both species show F-actin accumulation in endodermal cells, the intestine

of zebrafish mutants exhibits only a mildly reduced size without significant morphological

changes (Huang et al., 2008). Interestingly, treatment with a BMP inhibitor phenocopies the

liver-loss phenotype but does not affect intestinal size in zebrafish, suggesting that mis-

localisation of *bmp2a*-expressing lateral plate mesoderm (LPM) cells is primarily responsible

for the liver phenotype. In medaka, the penetrance of the IA phenotype was incomplete

(Table 2). Our data indicate that differential expression of *mypt* family genes does not explain

the intestinal specificity, implicating genetic background as a modifier of IA manifestation.

These findings raise the possibility that genetic modifiers contribute to the variability and

expressivity of the IA phenotype in medaka and may underlie interspecies differences.

Supporting this, individuals with MYPT1 mutations show highly variable clinical

presentations (Contreras-Capetillo et al., 2024; Hughes et al., 2020; Picard et al., 2022), and

IA has been reported in only a subset of cases. Notably, Picard et al. (2022) described a

patient with IA at birth and persistent Müllerian duct syndrome (PMDS), who remained

otherwise healthy until age 29. This observation is consistent with our finding that a small

number of medaka *mypt1* homozygous mutants can survive to adulthood, although this is

rare. During the establishment of our mutant lines, we selectively bred individuals in which

IA was frequently observed, and other developmental defects were relatively rare. This

selective breeding likely enriched a genetic background predisposed to IA but less susceptible

to other phenotypes. Given the more than 80 established laboratory medaka strains with

diverse geographic and genetic origins (Katsumura et al., 2019), it would be interesting to

cross our *mypt1* mutant with other genetic backgrounds to explore the influence of genetic

modifiers on IA penetrance and to reveal additional phenotypes associated with *mypt1*

disruption..

Mypt1 acts as a scaffold to provide a platform for assembly of the regulatory and

catalytic subunits of the MLCP complex and for the recruitment of phosphatase substrates

such as NMII. Conserved domains of Mypt1 that are vital for such molecular interaction are

also found in medaka Mypt1. Our original *g1-4* mutant provides insight into one such

domain: C2952A mutation introduces a premature stop codon after amino acids (aa) 983,

thereby deleting a large part of the Prkg interacting domain of the *WT* 1078-aa protein (Fig

1D). The LZ motif of this domain specifically mediates the interaction with Prkg1α.(Surks et

al., 1999) In vascular SM cells, activation of Mlcp mediated by Prkg dephosphorylates

pMrlc, which in turn induces vascular SM cell relaxation.(Surks et al., 1999) Thus, loss of the

LZ motif could explain increased contraction in the *g1-4* mutant. However, it remains to be

clarified whether Prkg1α or another PRKG family members regulate intestinal epithelium

integrity in developing intestine.

Our experiments using CRISPR-mediated knockout and morpholino knockdown (Mo-KD)

of *myh11a* suggest potential involvement of SM-myosin in IA pathogenesis. Yet, SM-myosin

expression was not detected in the intestinal region where IA typically forms, suggesting that

the SM layer is either absent or immature at the time of IA onset. MYH11 encodes the SM-

specific myosin heavy chain, a key component of the contractile machinery required for force

generation and structural stability in SM cells. However, the lack of fully developed SM cells

at the timing of IA formation argues against a direct contribution of MYH11-mediated

contraction of SM cells to the initial pathogenesis. In zebrafish, the *meltdown* (*mlt*) mutant—

harbouring a constitutively active form of Myh11—develops cystic expansion of the

posterior intestine. (Wallace et al., 2005b) This phenotype is thought to arise from Myh11-

dependent stromal–epithelial signalling that disrupts epithelial architecture. A similar

mechanism may operate in medaka *mypt1* mutants, where loss of Mypt1 could enhance

Myh11a activity. However, several distinctions suggest divergent downstream pathways:

snail family genes are upregulated in zebrafish *mlt* mutants but not in medaka; and whereas

the *mlt* phenotype arises at 74 dpf —after SM encasement—the IA phenotype in medaka

emerges by st. 28, before smooth muscle cslls are identified surrounding the region where IA

develop. (Gays et al., 2017; Wallace et al., 2005a) Thus, although Myh11a dysregulation and

stromal–epithelial signalling may be involved in both models, their timing and molecular

mediators differ.

Interestingly, MYH11 also functions in contractile non-SM cells such as myofibroblasts,

which can exert mechanical forces during development and repair (Pakshir et al., 2020). It is

possible that *myh11a* functions in mesenchymal cells other than SM, contributing to

mechanical stress in the developing gut and influencing epithelial morphogenesis. Although

speculative, this possibility warrants further investigation. In addition, SM-specific *Mypt1*

knockout (Mypt1[SMKO]) mice showed altered contractility but no apparent intestinal

malformations or motility defects (He et al., 2013), further supporting the notion that

canonical SM contraction is not directly responsible for IA pathogenesis. Together, these

findings suggest that *myh11a* may promote IA through a SM-independent mechanism,

potentially involving mechanically active mesenchymal populations such as myofibroblasts.

In addition to MYPT1 mutations, ACTB mutations have also been reported to cause IA

(Fakhro et al., 2019; Saskin et al., 2017; Sibbin et al., 2021), highlighting the importance of

tightly regulated actomyosin dynamics during intestinal development. Notably, mutations in

ACTB and ACTG1 are known to cause Baraitser-Winter Syndrome 1 (BRWS1, OMIM

#243310) in human, in which IA is an uncommon manifestation. This suggests that while

disruption of actomyosin dynamics contributes to IA pathogenesis, it is not solely sufficient

to induce the condition. Supporting this, pharmacological inhibition of NM II using

blebbistatin completely rescued the IA phenotype in *mypt1* mutants, whereas treatment with

calyculin A treatment—used to inhibit PP1 and thus supress dephosphorylation of NMII—

failed to phenocopy the IA phenotype. This discrepancy implies that Mypt1 may exert

additional, non-canonical functions beyond its regulatory role in MLCP (Kiss et al., 2019), or

that inhibition of PP1 alone is is insufficient to increase the amount of phosphorylated posho-

MII and contractility ofin intestinal epithelial cells. Identification of genetic modifiers that

influence IA development in the *mypt1* mutant background may provide further mechanistic

insight.

Our data show that ectopic accumulation of pMrlc and F-actin, likely indicative of

actomyosin hypercontractility, occurs in a non-uniform pattern along the intestine in *mypt1*

mutants. The cause of this spatial variability remains unknown. Moreover, we have not yet

validated the molecular mechanism that underlies how hypercontractility in the intestinal

epithelium disintegrates the epithelial layer structure. Further study is required to assess

completely the mechanics underpinning embryonic epithelial integrity.

Experimental Procedures

Ethics statement and Fish Strains

All experiments were approved by the animal experimentation committee of Kyoto

Prefectural University of Medicine (KPUM), licence numbers 27-123 and 2019-154. Fish

were kept at 26 °C on a 14 hours light / 10 hours dark cycle in a constant re-circulating

system of the in-house KPUM facility. The d-rR medaka strain original *mypt1* mutant *g1-4*

and hatching enzyme were obtained from NBRP (National Institute for Basic Biology,

Okazaki, Japan). Embryos were staged according to Iwamatsu's staging system.(Iwamatsu,

2004)

Knock-in transgenic (KIT) medaka

To visualise the intestinal epithelium, we generated a KIT medaka line that expresses

membrane-bound GFP (memGFP) under the control of the endogenous *foxa2* promotor (Fig.

3E), *Tg*[*foxa2:memGFP*]. *foxa2* is a well-known marker for the endoderm and its derived

organs.(Kobayashi et al., 2006) The targeting vector was constructed as described.(Ansai and

Kinoshita, 2017; Okuyama et al., 2013) This construct was integrated into the genome as

previously described.(Murakami et al., 2017) The sgRNA for the genomic target site was

designed as previously described (Kimura et al., 2014).

Detection of Apoptosis by Acridine Orange and TUNEL Assay

To detect apoptotic cells, acridine orange (AO) staining and TUNEL assays were

performed. For AO staining, a stock solution of AO (0.05 mg/mL; Tokyo Chemical

Industry, Tokyo, Japan) was prepared in distilled water and stored at 4°C in the dark.

Prior to use, the stock solution was diluted to 17 μg/mL in hatching buffer (HB).

Embryos at st. 27 were dechorionated and incubated in the AO solution for 30 minutes,

then washed three times in fresh HB for 5 minutes each to remove excess dye.

Fluorescent images were acquired using a DP70 digital camera mounted on an SZX16

stereomicroscope (Olympus, Tokyo, Japan). TUNEL assays were performed using the

In Situ Apoptosis Detection Kit (Takara) according to the manufacturer's instructions.

To visualise the intestinal epithelial membrane, embryos carrying the

Tg[foxa2:memGFP] transgene were used. GFP signals were detected using a rabbit anti-

GFP antibody (Invitrogen, A6455), followed by a goat anti-rabbit IgG antibody

conjugated with Alexa Fluor 633 (Thermo Fisher Scientific).

Whole mount *in situ* hybridization

A *foxa2* probe was designed and whole mount *in situ* hybridisation was performed as

described previously.(Kobayashi et al., 2006) (Takashima et al., 2007) Template DNAs for

snai1a (EST clone number, olea55k06), *snai1b* (olea23g07) and *snai2* (olsp48a08) were

obtained from the National BioResource Project (NBRP) medaka

(https://shigen.nig.ac.jp/medaka/).

Histology

Embryos fixed with 4% paraformaldehyde (PFA) were embedded in paraffin or resin

(Technovit 7100; Heraeus, Werheim Germany) and 6 µm tissue sections were prepared.

Paraffin sections were stained with haematoxylin and eosin, while plastic sections were

stained with haematoxylin only, followed by analysis using a BX51 microscope (Olympus,

Tokyo, Japan).

Immunohistology

For staining SM myosin, laminin and cytokeratin, 80% methanol/20% DMSO (Dent's

solution) was used for fixation. In other cases, embryos were fixed in 4% PFA in 1.5× PBS

containing 0.1% Tween 20. If Dent's fixation was used, embryos were de-chorionated with

medaka hatching enzyme before fixation. Then, non-specific antibody binding sites were

blocked with 1% dimethyl sulfoxide, 2% bovine serum albumin, 10% normal goat serum,

0.1% Triton X-100 in 1× PBS. Embryos fixed in PFA were permeabilised by incubation in

1× PBS containing 2% Triton X-100 for at least 2 hours before blocking of non-specific

binding sites. The following commercially available antibodies were used: GFP (1/2,000;

ab13970; Abcam), cytokeratin (1/100; clone AE1/AE3, ab27988; Abcam), SM myosin

(MYH11, 1/50; BT562; Biomedical Technologies), laminin (1/100; L9393; Sigma), phospho-

myosin light chain 2 (Ser19) (1:20; #3671; Cell Signaling Technology), phospho-Histone H3

(Ser10) (1/500; #06-570, Millipore), and PKCζ (1/800; ab5964; Abcam,). Alexa Fluor-633-

Phalloidin (1/20; A22284; Thermo Fisher Scientific) was used to mark F-actin. Detection of

primary antibodies was performed using Alexa Fluor-488, -555 goat anti-rabbit, chicken or

mouse IgG (1/400; Invitrogen). Images were acquired with an Olympus FV1200 confocal

microscope or Zeiss LSM900 and yz-planes were reconstructed using FIJI.(Schindelin et al.,

2012) Abnormal accumulation of pMrlc or F-actin was judged based on: (a) obviously higher

immunofluorescence signal in the middle part of the intestine compared with other areas, and

(b) obviously higher immunofluorescence signal at apical and also basal areas of the

intestinal epithelium where strong signal was never observed in the *WT*. Representative

single-plane confocal images (not z-projections) are shown in the figures.

CRISPR/Cas9-Mediated Mutagenesis of *mypt1*

The sgRNA targeting exon 1 of *mypt1* was designed using the web tool "Search for

CRISPR target site with micro-homology sequences" (http://viewer.shigen.info/cgi-

bin/crispr/crispr.cgi), with the parameter "micro-homology sequences" set to 4 bases. sgRNA

synthesis was performed as previously described.(Ansai and Kinoshita, 2014) The target

sequence is shown in Fig. 2A. A mixture containing 150 ng/µL sgRNA and 200 ng/µL Cas9

mRNA was microinjected into 1–2 cell stage embryos. Injected embryos were raised to

adulthood, and eight germline-transmitting F0 founders were identified by genotyping pooled

F1 offspring. One of these founders, #4-6, was outcrossed to *WT* fish, and the resulting F1

progeny were genotyped. F1 fish harbouring the mutation (Fig. 1, 2) were used for

subsequent analyses.

CRISPR/Cas9 knockout of *myh11a*.

To knock out *myh11a*, three overlapping sgRNAs targeting exon 5—which encodes part of

the ATP-binding domain—were designed using the CRISPR-Cas9 guide RNA design

checker (Integrated DNA Technologies, IDT). Off-target effects were evaluated via BLAST

search against the medaka genome using the Ensembl genome browser. The overlapping

design of sgRNAs was intended to increase mutagenesis efficiency in the targeted region

(Fig. 11A). Each sgRNA was prepared following the manufacturer's protocol (IDT). A

mixture containing 1.0 µg/µL Alt-R™ S.p. Cas9 Nuclease V3 (IDT) and 5.0 µM of each

sgRNA, diluted in RNase-free water was injected into 1–2 cell stage medaka embryos.

Morpholino knockdown of myh11a

Antisense Morpholino (Gene Tools) targeting the splice donor site of exon 2of medaka

myh11a was used for knockdown (Fig. 11E). The Standard Control Oligo supplied by Gene

Tools was used as a negative control.

Genotyping

WT and mutant *mypt1* alleles were distinguished using a polymerase chain reaction-

restriction fragment length polymorphism (PCR-RFLP) method. Genomic DNA from lysed

tissue was subjected to PCR using Quick Taq® HS DyeMix Taq polymerase (TOYOBO),

with an initial denaturation at 94°C for 2 min, followed by 45 cycles of 94°C denaturation for

30 s, annealing at 55°C for 30 s, and extension at 68°C for 50 s,. The primers used are shown

in Table 6. PCR products were digested with the restriction enzyme *Hae*III (TOYOBO), and

analysed by electrophoresis in 3% agarose gels (02468-95, Nacalai) in 1× TAE buffer.

Chemical Inhibition of Actomyosin Contractility

To examine the effect of NMII inhibition on IA development, st. 25 embryos were

treated with either 0.1% DMSO (control) or 50 µM (-)-blebbistatin (021-17041, FUJIFILM),

dissolved in 0.1% DMSO. After a 2-hour incubation, embryos were thoroughly washed and

raised until 4 dpf for phenotypic analysis.. Furthermore, embryos were raised to 9 dpf for

extended observation, imaging and genotyping. To assess the effect of increased myosin

phosphatase inhibition, *WT* embryos were treated from st. 23 to st. 32 with either 0.1%

DMSO (control) or 0.1 µM Calyculin A (101932-71-2, FUJIFILM), also dissolved in 0.1%

DMSO. Phenotypic evaluation was performed after the treatment, and embryos were further

raised to 9 dpf for evaluation.

Phylogenetic analysis

Multiple sequence alignments were performed using CLUSTALW with default

parameters, followed by phylogenetic tree construction using the PhyML-bootstrap method

(Kyoto University Bioinformatics Center, https://www.genome.jp/tools-bin/clustalw). The

protein sequences used for the alignment are listed in Table 3.

RT-PCR

Total RNA was extracted using TRIzol reagent (Invitrogen) according to the

manufacturer's protocol. For small tissue samples, RNA was purified using the Direct-zol

RNA Kit (Zymo Research). Reverse transcription and PCR were carried out using superscript

III (Invtrogen) and Quick Taq® HS DyeMix (TOYOBO) , respectively, following the

manufacturer's instructions.

Statistical analysis

Statistical analyses were performed using EZR(Kanda, 2013)

Acknowledgements

The medaka National Bioresource Project (NBRP: https://shigen.nig.ac.jp/medaka/),

governed by the Ministry of Education, Culture, Sports, Science and Technology, Japan,

provided the hatching enzyme, EST clones, d-rR and original mutant *g1-4* medaka strains

used in this study.

Author Contributions

DK designed the research; DK, AK, TK, KM, SA, MK, and KN examined the mutant

phenotype; DK, TK, HY, ST, TK, TK, TN and TJ performed mutant screening; YI, KA, YS

and HT organized mutant screening; DK, KM, YN, MS, SS, SI, TY, HT and KY analysed

data; and DK and KY wrote the paper.

References

Agarwal, P. and Zaidel-Bar, R. (2018). Principles of Actomyosin Regulation In Vivo.
Trends Cell Biol.
Aghaallaei, N., Gruhl, F., Schaefer, C. Q., Wernet, T., Weinhardt, V., Centanin, L.,
Loosli, F., Baumbach, T. and Wittbrodt, J. (2016). Identification, visualization and clonal
analysis of intestinal stem cells in fish. *Development* **143**, 3470-3480.
Angulo-Urarte, A., Casado, P., Castillo, S. D., Kobialka, P., Kotini, M. P., Figueiredo, A.
M., Castel, P., Rajeeve, V., Mila-Guasch, M., Millan, J. et al. (2018). Endothelial cell
rearrangements during vascular patterning require PI3-kinase-mediated inhibition of
actomyosin contractility. *Nat Commun* **9**, 4826.
Ansai, S. and Kinoshita, M. (2014). Targeted mutagenesis using CRISPR/Cas system in
medaka. *Biol Open.*
Ansai, S. and Kinoshita, M. (2017). Genome Editing of Medaka. *Methods Mol Biol* **1630**,
175-188.
Barresi, M. J., Burton, S., Dipietrantonio, K., Amsterdam, A., Hopkins, N. and
Karlstrom, R. O. (2010). Essential genes for astroglial development and axon pathfinding during
zebrafish embryogenesis. *Dev Dyn* **239**, 2603-18.

Bavaria, M. N., Ray, R. M. and Johnson, L. R. (2011). The phosphorylation state of
MRLC is polyamine dependent in intestinal epithelial cells. *Am J Physiol Cell Physiol* **300**, C164-
75.
Bremer, J. and Granato, M. (2016). Myosin phosphatase Fine-tunes Zebrafish
Motoneuron Position during Axonogenesis. *PLoS Genet* **12**, e1006440.
Celli, J. (2014). Genetics of gastrointestinal atresias. *Eur J Med Genet* **57**, 424-439.
Contreras-Capetillo, S. N., Abreu-González, M., Centeno-Navarrete, Y., Ferro-Muñoz, S.
R., Ceballos-Zapata, J. and García-Martínez, C. (2024). Loss-of-Function Variant in PPP1R12A-
Related Urogenital and/or Brain Malformation Syndrome: Expanded Phenotype of Sex Reversal.
Am J Med Genet A, e63876.
Dong, G., Huang, Y., Ding, H., Luo, L., Zhang, Y., Huang, H. and Ruan, H. (2019).
Mypt1 regulates Bmp signaling to promote embryonic exocrine pancreas growth in zebrafish.
genesis, e23345.
Fairbanks, T. J., Kanard, R., Del Moral, P. M., Sala, F. G., De Langhe, S., Warburton,
D., Anderson, K. D., Bellusci, S. and Burns, R. C. (2004a). Fibroblast growth factor receptor 2
IIIb invalidation--a potential cause of familial duodenal atresia. *J Pediatr Surg* **39**, 872-4.
Fairbanks, T. J., Kanard, R. C., De Langhe, S. P., Sala, F. G., Del Moral, P. M.,
Warburton, D., Anderson, K. D., Bellusci, S. and Burns, R. C. (2004b). A genetic mechanism for
cecal atresia: the role of the Fgf10 signaling pathway. *J Surg Res* **120**, 201-9.
Fairbanks, T. J., Kanard, R. C., Del Moral, P. M., Sala, F. G., De Langhe, S. P., Lopez,
C. A., Veltmaat, J. M., Warburton, D., Anderson, K. D., Bellusci, S. et al. (2005). Colonic atresia
without mesenteric vascular occlusion. The role of the fibroblast growth factor 10 signaling
pathway. *J Pediatr Surg* **40**, 390-6.
Fakhro, K. A., Robay, A., Rodrigues-Flores, J. L., Mezey, J. G., Al-Shakaki, A. A.,
Chidiac, O., Stadler, D., Malek, J. A., Imam, A. B., Sheikh, A. et al. (2019). Point of Care Exome
Sequencing Reveals Allelic and Phenotypic Heterogeneity Underlying Mendelian disease in
Qatar. *Hum Mol Genet* **28**, 3970-3981.
Frischer, J. S. and Azizkhan, R. G. (2012). Chapter 82 - Jejunoileal Atresia and
Stenosis. In *Pediatric Surgery (Seventh Edition)*,   (ed. A. G. Coran), pp. 1059-1071.
Philadelphia: Mosby.
Fujita, M., Isogai, S. and Kudo, A. (2006). Vascular anatomy of the developing medaka,
Oryzias latipes: a complementary fish model for cardiovascular research on vertebrates. *Dev Dyn*
235, 734-46.
Gays, D., Hess, C., Camporeale, A., Ala, U., Provero, P., Mosimann, C. and Santoro, M.
M. (2017). An exclusive cellular and molecular network governs intestinal smooth muscle cell
differentiation in vertebrates. *Development* **144**, 464-478.

Grassie, M. E., Moffat, L. D., Walsh, M. P. and MacDonald, J. A. (2011). The myosin
phosphatase targeting protein (MYPT) family: a regulated mechanism for achieving substrate
specificity of the catalytic subunit of protein phosphatase type 1delta. *Arch Biochem Biophys* **510**,
147-59.
Gupta, T., Yang, W., Iovannisci, D. M., Carmichael, S. L., Stevenson, D. K., Shaw, G. M.
and Lammer, E. J. (2013). Considering the vascular hypothesis for the pathogenesis of small
intestinal atresia: a case control study of genetic factors. *Am J Med Genet A* **161A**, 702-10.
Gutzman, J. H. and Sive, H. (2010). Epithelial relaxation mediated by the myosin
phosphatase regulator Mypt1 is required for brain ventricle lumen expansion and hindbrain
morphogenesis. *Development* **137**, 795-804.
Hartshorne, D. J., Ito, M. and Erdodi, F. (2004). Role of protein phosphatase type 1 in
contractile functions: myosin phosphatase. *The Journal of biological chemistry* **279**, 37211-4.
He, W. Q., Qiao, Y. N., Peng, Y. J., Zha, J. M., Zhang, C. H., Chen, C., Chen, C. P.,
Wang, P., Yang, X., Li, C. J. et al. (2013). Altered contractile phenotypes of intestinal smooth
muscle in mice deficient in myosin phosphatase target subunit 1. *Gastroenterology* **144**, 1456-65,
1465 e1-5.
Huang, H., Ruan, H., Aw, M. Y., Hussain, A., Guo, L., Gao, C., Qian, F., Leung, T.,
Song, H., Kimelman, D. et al. (2008). Mypt1-mediated spatial positioning of Bmp2-producing
cells is essential for liver organogenesis. *Development* **135**, 3209-18.
Hughes, J. J., Alkhunaizi, E., Kruszka, P., Pyle, L. C., Grange, D. K., Berger, S. I.,
Payne, K. K., Masser-Frye, D., Hu, T., Christie, M. R. et al. (2020). Loss-of-Function Variants in
PPP1R12A: From Isolated Sex Reversal to Holoprosencephaly Spectrum and Urogenital
Malformations. *Am J Hum Genet* **106**, 121-128.
Ito, M., Nakano, T., Erdodi, F. and Hartshorne, D. J. (2004). Myosin phosphatase:
structure, regulation and function. *Mol Cell Biochem* **259**, 197-209.
Iwamatsu, T. (2004). Stages of normal development in the medaka Oryzias latipes.
Mech Dev **121**, 605-18.
Kanard, R. C., Fairbanks, T. J., De Langhe, S. P., Sala, F. G., Del Moral, P. M., Lopez,
C. A., Warburton, D., Anderson, K. D., Bellusci, S. and Burns, R. C. (2005). Fibroblast growth
factor-10 serves a regulatory role in duodenal development. *J Pediatr Surg* **40**, 313-6.
Kanda, Y. (2013). Investigation of the freely available easy-to-use software 'EZR' for
medical statistics. *Bone Marrow Transplant* **48**, 452-8.
Katsumura, T., Oda, S., Mitani, H. and Oota, H. (2019). Medaka Population Genome
Structure and Demographic History Described via Genotyping-by-Sequencing. *G3 (Bethesda)* **9**,
217-228.

1   **Kim, J. H., Lee, S. R., Li, L. H., Park, H. J., Park, J. H., Lee, K. Y., Kim, M. K., Shin, B.**
2 **A. and Choi, S. Y.** (2011). High cleavage efficiency of a 2A peptide derived from porcine
3 teschovirus-1 in human cell lines, zebrafish and mice. *PLoS ONE* **6**, e18556.
4   **Kimura, T., Jindo, T., Narita, T., Naruse, K., Kobayashi, D., Shin, I. T., Kitagawa, T.,**
5 **Sakaguchi, T., Mitani, H., Shima, A. et al.** (2004). Large-scale isolation of ESTs from medaka
6 embryos and its application to medaka developmental genetics. *Mech Dev* **121**, 915-32.
7   **Kimura, Y., Hisano, Y., Kawahara, A. and Higashijima, S.** (2014). Efficient generation
8 of knock-in transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-mediated
9 genome engineering. *Sci Rep* **4**, 6545.
10   **Kiss, A., Erdodi, F. and Lontay, B.** (2019). Myosin phosphatase: Unexpected functions of
11 a long-known enzyme. *Biochim Biophys Acta Mol Cell Res* **1866**, 2-15.
12   **Kobayashi, D., Jindo, T., Naruse, K. and Takeda, H.** (2006). Development of the
13 endoderm and gut in medaka, Oryzias latipes. *Dev Growth Differ* **48**, 283-95.
14   **Louw, J. H.** (1959). Congenital intestinal atresia and stenosis in the newborn.
15 Observations on its pathogenesis and treatment. *Ann R Coll Surg Engl* **25**, 209-34.
16   **Louw, J. H. and Barnard, C. N.** (1955). Congenital intestinal atresia; observations on its
17 origin. *Lancet* **269**, 1065-7.
18   **Mizuno, T., Tsutsui, K. and Nishida, Y.** (2002). Drosophila myosin phosphatase and its
19 role in dorsal closure. *Development* **129**, 1215-23.
20   **Munjal, A. and Lecuit, T.** (2014). Actomyosin networks and tissue morphogenesis.
21 *Development* **141**, 1789-93.
22   **Munjal, A., Philippe, J. M., Munro, E. and Lecuit, T.** (2015). A self-organized
23 biomechanical network drives shape changes during tissue morphogenesis. *Nature* **524**, 351-5.
24   **Murakami, Y., Ansai, S., Yonemura, A. and Kinoshita, M.** (2017). An efficient system for
25 homology-dependent targeted gene integration in medaka (Oryzias latipes). *Zoological Lett* **3**, 10.
26   **Okamoto, R., Ito, M., Suzuki, N., Kongo, M., Moriki, N., Saito, H., Tsumura, H.,**
27 **Imanaka-Yoshida, K., Kimura, K., Mizoguchi, A. et al.** (2005). The targeted disruption of the
28 MYPT1 gene results in embryonic lethality. *Transgenic Res* **14**, 337-40.
29   **Okuyama, T., Isoe, Y., Hoki, M., Suehiro, Y., Yamagishi, G., Naruse, K., Kinoshita, M.,**
30 **Kamei, Y., Shimizu, A., Kubo, T. et al.** (2013). Controlled Cre/loxP site-specific recombination in
31 the developing brain in medaka fish, Oryzias latipes. *PLoS ONE* **8**, e66597.
32   **Pakshir, P., Noskovicova, N., Lodyga, M., Son, D. O., Schuster, R., Goodwin, A.,**
33 **Karvonen, H. and Hinz, B.** (2020). The myofibroblast at a glance. *J Cell Sci* **133**.
34   **Parkin, C. A., Allen, C. E. and Ingham, P. W.** (2009). Hedgehog signalling is required for
35 cloacal development in the zebrafish embryo. *Int J Dev Biol* **53**, 45-57.
36   **Picard, J. Y., Morin, G., Devouassoux-Shisheboran, M., Van der Smagt, J., Klosowski,**
37 **S., Pienkowski, C., Pierre-Renoult, P., Masson, C., Bole, C. and Josso, N.** (2022). Persistent

Müllerian duct syndrome associated with genetic defects in the regulatory subunit of myosin phosphatase. *Hum Reprod* **37**, 2952-2959.

Ray, R. M., Guo, H., Patel, M., Jin, S., Bhattacharya, S. and Johnson, L. R. (2007). Role of myosin regulatory light chain and Rac1 in the migration of polyamine-depleted intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* **292**, G983-95.

Saskin, A., Tischkowitz, M. and Study, D. D. D. (2017). Jejunal atresia, periodic fevers and psoriatic arthropathy in Baraitser-Winter malformation syndrome. *Clin Dysmorphol* **26**, 235-237.

Savagner, P. (2015). Epithelial-mesenchymal transitions: from cell plasticity to concept elasticity. *Curr Top Dev Biol* **112**, 273-300.

Scheibner, K., Schirge, S., Burtscher, I., Büttner, M., Sterr, M., Yang, D., Böttcher, A., Ansarullah, Irmler, M., Beckers, J. et al. (2021). Epithelial cell plasticity drives endoderm formation during gastrulation. *Nature Cell Biology* **23**, 692-703.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B. et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-82.

Shorter, N. A., Georges, A., Perenyi, A. and Garrow, E. (2006). A proposed classification system for familial intestinal atresia and its relevance to the understanding of the etiology of jejunoileal atresia. *J Pediatr Surg* **41**, 1822-1825.

Sibbin, K., Yap, P., Nyaga, D., Heller, R., Evans, S., Strachan, K., Alburaiky, S., Nguyen, H. M. A., Hermann-Le Denmat, S., Ganley, A. R. D. et al. (2021). A de novo ACTB gene pathogenic variant in identical twins with phenotypic variation for hydrops and jejunal atresia. *Am J Med Genet A*.

Surks, H. K., Mochizuki, N., Kasai, Y., Georgescu, S. P., Tang, K. M., Ito, M., Lincoln, T. M. and Mendelsohn, M. E. (1999). Regulation of myosin phosphatase by a specific interaction with cGMP- dependent protein kinase Ialpha. *Science* **286**, 1583-7.

Takashima, S., Shimada, A., Kobayashi, D., Yokoi, H., Narita, T., Jindo, T., Kage, T., Kitagawa, T., Kimura, T., Sekimizu, K. et al. (2007). Phenotypic analysis of a novel chordin mutant in medaka. *Dev Dyn* **236**, 2298-310.

Tan, C., Stronach, B. and Perrimon, N. (2003). Roles of myosin phosphatase during Drosophila development. *Development* **130**, 671-81.

Vasquez, C. G., Tworoger, M. and Martin, A. C. (2014). Dynamic myosin phosphorylation regulates contractile pulses and tissue integrity during epithelial morphogenesis. *J Cell Biol* **206**, 435-50.

Vicente-Manzanares, M., Ma, X., Adelstein, R. S. and Horwitz, A. R. (2009). Non-muscle myosin II takes centre stage in cell adhesion and migration. *Nat Rev Mol Cell Biol* **10**, 778-90.

Wallace, K. N., Akhter, S., Smith, E. M., Lorent, K. and Pack, M. (2005a). Intestinal

growth and differentiation in zebrafish. *Mech Dev* **122**, 157-73.

Wallace, K. N., Dolan, A. C., Seiler, C., Smith, E. M., Yusuff, S., Chaille-Arnold, L.,

Judson, B., Sierk, R., Yengo, C., Sweeney, H. L. et al. (2005b). Mutation of smooth muscle myosin

causes epithelial invasion and cystic expansion of the zebrafish intestine. *Dev Cell* **8**, 717-26.

Weiser, D. C., Row, R. H. and Kimelman, D. (2009). Rho-regulated myosin phosphatase

establishes the level of protrusive activity required for cell movements during zebrafish

gastrulation. *Development* **136**, 2375-84.

Yang, J., Antin, P., Berx, G., Blanpain, C., Brabletz, T., Bronner, M., Campbell, K.,

Cano, A., Casanova, J., Christofori, G. et al. (2020). Guidelines and definitions for research on

epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol* **21**, 341-352.

Yokoi, H., Shimada, A., Carl, M., Takashima, S., Kobayashi, D., Narita, T., Jindo, T.,

Kimura, T., Kitagawa, T., Kage, T. et al. (2007). Mutant analyses reveal different functions of

fgfr1 in medaka and zebrafish despite conserved ligand-receptor relationships. *Dev Biol* **304**, 326-

37.

Figure Legends

Fig. 1. Characterisation of medaka IA mutant, *g1-4*. (A, B) Lateral views of *WT* (A) and

mypt1 mutant (*g1-4*) (B) embryos at st. 40. Note the development of IA in the mutant,

represented by the blind ends of the intestine (arrowheads). Anterior is to the left. Arrow,

cloaca; GB, gallbladder; Dotted brackets, intestine. Scale bar: 200 μm. **(C)** Schematic

presentation of part of LG23. The *g1-4* locus is confined to a 0.13 cM interval between the

markers LG23-4.4 and LG23-4.6. Inset: chromatogram of a point mutation in *g1-4*, which

gives rise to a premature stop codon. (D) Schematics outlining Mypt1 domains of *WT* (top),

g1-4 (middle), and *mypt1*[#4-6] (bottom) proteins. Asterisks indicate premature translation

termination. In *mypt1*[#4-6], an 11-nucleotide deletion with a 7-nucleotide insertion cause a

frame-shift (green), leading to truncation.

Fig. 2. Comparison of *WT* and *mypt1#4-6* allele sequences. (A) The 11-nucleotide deletion

and 7 nucleotide insertion are shown in red and blue, respectively. The CRISPR target site is

underlined. The predicted amino acid sequence resulting from the frame shift is shown in

green. Asterisk indicates premature stop codon. *Hae*III sites (GGCC) that are used for

genotyping by restriction fragment length polymorphism (RFLP) assays are indicated. (B)

Schematic outline of PCR products amplified from *WT* and *mypt1#4-6* alleles. Primer positions

(*ppp1r12a*-F15 and *ppp1r12a*-R16), the CRISPR target site and *Hae*III sites are indicated.

The *Hae*III site lost in *mypt1#4-6* is shown in grey. The introduced stop codon in *mypt1#4-6* is

marked by an asterisk.

Fig. 3. Organogenesis proceeds normally in *mypt1* mutants except for the presence of

IA. (A–D) Dilation of the intestine in *mypt1* mutants. Ventral views of *WT* (A, C) and

mypt1#4-6 (B, D) embryos at st. 40 in the Tg[*foxa2:memGFP*] medaka line. GFP

fluorescence and brightfield images are shown in (A, B) and (C, D), respectively. White

dotted lines outline the intestine. Note the significant dilation anterior to the IA lesion in the

mutant (asterisk). LV, liver. Scale bar: 200 µm. (E–P) Histological sections of st. 40

embryos. *WT*: (E–J); *mypt1* mutant: (K–P). Panels show comparable anatomical levels: (E

and K), (F and L), (G and M), (H and N), (I and O), and (J and P). Abbreviations: EnP,

endocrine pancreas; Es, esophagus; ExP, exocrine pancreas; GB, gall bladder; Gd, gonad; Gl,

glomerulus; Ht, heart; IE, inner ear; In, intestine; Lv, liver; NT, nephric tubule; Sp, spleen;

AB, swim bladder; Ty, thyroid; R, right; L, left. Scale bar: 50 µm. (Q–X) Histological

micrographs of the intestine in *WT* (Q–T) and *mypt1#4-6* (U–X) embryos at st. 32, stained with

haematoxylin. (U) Region anterior to IA; (V) IA lesion; (W) anterior blind-end of posterior

intestine; (X) region posterior to the IA. *WT* sections (Q–T) correspond anatomically to (U–

X). Note normal morphology in (U) and (X). MS, mesentery; NT, nephric tubule. Scale bar:

20 μm.

Fig. 4. Apico-Basolateral polarity is maintained in *mypt1* mutants. (A) Schematic diagram

of the distribution of cytokeratin, laminin and SM myosin in the developing intestine. (B–I)

Whole mount immunofluorescence micrographs of cytokeratin and SM myosin (B, C),

laminin (D–G) and Pkcζ (H, I) in *WT* (B, D, F, H) and *mypt1*[#4-] (C, E, G and I) as left lateral

view. White dotted lines mark the outline of intestine in a differential interference contrast

(DIC) micrograph corresponding to a fluorescence image. Note that basement membrane was

fragmented in the mutant embryos at st. 26 before IA onset (F, n = 11; G, n=9). (H, I)

Immunohistochemistry for Pkcζ in long-axis optical sections of st. 26 embryo intestines. The

epithelium of the intestine was visualized with *Tg[foxa2:memGFP]* (H, n = 9; I, n = 10). Note

that the apical localization of Pkcζ was not disturbed in *mypt1* mutants (I). Arrowheads,

blind-ends of IA. Scale bars: 50 μm.

Fig. 5. Endodermal migration to form the intestinal anlage is intact in *mypt1* mutants.

WT (A, C, E, n = 10) and *mypt1* mutant (B, D, F, n = 10) embryos were subjected to whole

mount *in situ* hybridization for *foxa2*. Dorsal (A–D) and dorsolateral views (E–F) are shown.

Embryonic stages are indicated in the top right. Brackets mark endoderm and intestinal

anlage. Scale bar: 100 μm.

Fig. 6. Cell death and cell proliferation are not involved in IA development. (A–H) Cell

death and cell proliferation in the intestine are not significantly altered in *mypt1* mutants. (A,

B) Acridine orange staining of *WT* embryos (Aa, *n* = 12) and *mypt1* mutants (Ba, *n* = 5) at st.

27. (Ab, Bb) Brightfield images corresponding to (Aa) and (Ba), respectively. Dotted ellipses

mark apoptotic cells surrounding the cloacal opening, where apoptosis is normally observed

(Parkin et al., 2009). (C–H) *WT* (C, E, G) and *mypt1* mutant (D, F, H) embryos at st. 26 were

stained with anti-phosphorylated histone H3 (PH3) antibody and/or propidium iodide (PI).

(I) Quantification of PH3-positive cell ratio to total cell number in the intestinal region was

not significantly different between *WT* (n = 16) and *mypt1* mutants (n = 11; chi-square test, P

= 1). Scale bar: 50 μm.

Fig. 7. Epithelial–mesenchymal transition (EMT) is not involved in IA formation in

***mypt1* mutants.** Lateral views of st. 26 embryos subjected to whole mount *in situ*

hybridization to detect EMT markers: WT (A, C, E); mutants (B, D, F). (A, B) for *snai1a*, (C,

D) for *snai1b*, and (E, F) for *snai2*. To enhance visibility, the intestine was detached from the

dorsal mesentery. Dotted boxes in (Aa-Fa) are enlarged in (Ab-Fb), respectively. The number

of samples examined is indicated in the bottom left corner. Scale bar: 200 μm.

Fig. 8. Phylogenetic analysis of the *Mypt* gene family. Phylogenetic tree conducted from

protein sequences of Mypt family genes listed in Table 3.

Fig. 9. Expression patterns of *Mypt1* family genes. (A–F) Whole-mount *in situ*

hybridization of *mypt1* (A, D), *mypt1/2-r* (B, E), and *mbs85* (C, F). (A–C) st. 25; (D–F) of st.

27 in *WT*. Dotted boxes in (Aa-Fa) are enlarged in (Ab-Fb), respectively. The number of

samples examined is indicated in the bottom left corner. Scale bar: 200 μm. MW: Molecular

weight marker (DM2100, SMBIO, Taiwan)

(G) RT-PCR analysis. PCRs using first-strand cDNA from: (a) negative control, (b) 1-cell

stage, (c) blastula stage, (d) head region at stage 26, (e) intestine region at stage 25, and (f)

adult whole body.

Fig. 10. Actomyosin is abnormally activated in the developing intestine of *mypt1*

mutants. (A–E) Fluorescence micrographs of pMrlc for *WT* (A, C) and *mypt1*[#4-6] (B, D, E).

st. 25 embryos are shown in A and B, whereas st. 27 embryos are shown in C, D and E. Note

significant accumulation of pMrlc in mutant intestinal epithelium (Bb, Db, Eb). Dotted boxes

in Bc is enlarged in Bd. Scale bar, 50 μm. (F) The proportion of embryos possessing an

abnormal increase of pMrlc in embryonic intestine (Fisher's exact test). (G-J) Fluorescence

micrographs of F-actin stained with phalloidin. (a–e) represent short axis sections of intestine.

WT (G, I) and *mypt1*[#4-6] (H, J) st.25 (G, H) and st. 27 (I, J) embryos are shown. (H, J) Note

abnormal accumulation of F-actin in the *mypt1* mutant intestine. Scale bar: 50 μm. *, an

intestinal lumen. (K) The proportion of embryos possessing an abnormal increase of F-actin

in embryonic intestine (Fisher's exact test).

Fig. 11. Targeting of SM-myosin *myh11a*. (A) CRISPR target sites within the *myh11a* gene.

(B–D) Expression of SM-myosin in 6 dpf embryos, into which *myh11a* sgRNA was injected

at 1-2 cell stage,: without Cas9 nuclease (B); with Cas9 (C and D). Scale bar: 100 μm.

(E) Morpholino target site within *myh11a*. (F) RT-PCR analysis of *myh11a* morpholino-

injected embryos. Lane 1: control; lanes 2 and 3: morpholino-injected; lane 4: *WT* cDNA;

lane 5: *WT* genomic DNA

Fig. 12 Attenuation of actomyosin activation rescues IA phenotype in *mypt1* mutants.

(A) Experimental design for blebbistatin treatment. (B, C) Images at 9 dpf: control (B) and

blebbistatin-treated. Arrowheads indicate IA. Scale bars: 200 μm. (Ba, Ca) Dotted boxes in

(B) and (C) are enlarged in (Bb) and (Cb), respectively. (Bc, Cc) Corresponding fluorescent

images of (Bb) and (Cb), respectively. (D) Quantification of rescue of the IA phenotype in

mypt1[#4-6] embryos following blebbistatin treatment (Fisher's exact test, $P = 1.9 \times 10^{-5}$). (E)

Diagram of NMII regulation. *Mypt1* is a critical subunit of MLCP and is essential for its

function. Loss of *Mypt1* leads to reduced MLCP activity, resulting in ectopic activation of

actomyosin contraction. (F) Hypothetical molecular events following the loss of *mypt1*

function. (G) Schematic representation of the molecular state under blebbistatin treatment.

Table 1. Variation in penetrance among *g1-4* pairs

Fish pair		IA	TOTAL	%
Female 1	Male 1	1	37	2.7
Female 2	Male 2	8	45	17.8
Female 2	Male 3	8	36	22.2
Female 3	Male 2	5	30	16.7
Female 4	Male 3	2	20	10.0
Female 5	Male 1	4	45	8.9

IA: The number of embryos that develop intestinal atresia.

Total: The number of embryos after deduction of those for which intestinal phenotype could not be determined (dead or abnormal development) .

Table 2. Variation in penetrance among *mypt1* [#4-6] pairs

Generation	Fish pair		IA	TOTAL	%
	Female	Male			
F1	16	17	14	59	23.7
F2	33	34	19	97	19.6
	35	36	9	29	31.0
	37	38	8	30	26.7
	39	40	13	71	18.3
	41	43	13	84	15.5
	44	45	9	44	20.5
	46	47	17	62	27.4
	48	49	13	90	14.4
	50	51	12	72	16.7
	54	55	5	39	12.8
	56	57	15	83	18.1
F2 average			133	701	19.0

IA: The number of embryos that develop intestinal atresia.

Total: The number of embryos after deduction of those for which intestinal phenotype could not be determined (dead or abnormal development) .

Table 3. List of protein sequences used for ClustalW analysis

Species	Protein Name	Accession Number
Homo sapiens	MYPT1	NP_002471.1
Gallus gallus	Mypt1	NP_990454.1
Danio rerio	Mypt1	NP_001003870.2
Oryzias latipes	Mypt1	XP_023807904.1
Homo sapiens	MYPT2	NP_002472.2
Gallus gallus	Mypt2	XP_040508700.1
Homo sapiens	MBS85	NP_060077.1
Gallus gallus	Mbs85	XP_025002065.1
Danio rerio	Mbs85	NP_001071047.1
Oryzias latipes	Mbs85	XP_023810293.1
Danio rerio	Mypt1/2-r	XP_002666820.2
Oryzias latipes	Mypt1/2-r	XP_020559658.2
Homo sapiens	MYPT3	NP_001316373.1
Danio rerio	Mypt3	XP_001334473.1
Oryzias latipes	Mypt3	XP_011487151.1
Homo sapiens	TIMAP	NP_056383.1
Gallus gallus	Timap	NP_001026022.2
Danio rerio	Timap	XP_001340092.6
Oryzias latipes	Timap	XP_004068819.1
D. melanogaster	mbs	NP_001261919.1
C. elegans	MEL-11	AAB47273.1

Table 4. Intestinal atresia phenotype in homozygous *myh11a* mutant embryos injected with sgRNAs with or without Cas9 nuclease

Cas9(+)			Cas9(-)		
IA(+)	IA(-)	Total	IA(+)	IA(-)	Total
6	4	10	9	0	9

Embryos were injected at the one-cell stage with a mixture of three sgRNAs targeting *myh11a*, either alone or in combination with Cas9 protein. The presence or absence of intestinal atresia was scored in mutant homozygous embryo. Injection of sgRNAs with Cas9 partially rescued the atresia phenotype, whereas sgRNsA injection alone had no significant effect.

Table 5 Intestinal atresia phenotype in homozygous *myh11a* mutant embryos injected with Mo-*myh11a*-E2I2 morhplino

Morpholino	Conc. (µM)	IA (+)	IA(-)	JNP	Total
Control	200	22	1	1	23
myh11a E1I1	150	14	6	0	20
	200	21	9	11	41

JNP: Judgement Not Possible.
Embryos were injected at the one-cell stage with Mo-*myh11a*-E2I2 or control morpholinos .The presence or absence of intestinal atresia was scored in mutant homozygous embryo. Injection of Mo-*myh11a*-E2I2 partially rescued the atresia phenotype, whereascontrol morpholino injection had no significant effect.

Table 6 Primers

gene	Forward	Reverse
mypt1	TCCTCCACTTTCCTCATATCTGCTGATGG	TGGCCAAAAATCGTCAGGTGTGTTCG
myh11a	ATGTCTAAGAAGGCCCCGAGTG	GGAACTTCATGCCGTTTCTTCC
dagph	GTATCAATGGTTTTGGCCGTATC	AGTGATGGCATGAACTGTGCTC
actb	GATGCCCCTCGTGCTGTCTTTCC	GAGCAGGACAGGGTGCTCCTCA
mypt1/2-r	GACGGCAGGATCAGTTGCAG	CTCCTCTTTTCGGGCCTTGTC
mbs85	CGCCACCGTATGTCTTGTTT	CACCGTGATCCAAGAGGAAC
snail1a	CACTAAAGCTATCACTCCGTTGCT	GGTCCCAGCCTTCTTATGGTTC
snail1b	CGCGTCTTCCAACATTTACGCA	ACCATACACAGACAGCTGTCTCA
snail2	GGAGTTTGTTCTTTACGCAGCGAA	AGCTGGCAAACTGTTTCTACAGG

Fig. 1. Characterization of medaka IA mutant, *g1-4.*

A

	10	20	30	40	50	60	

WT   ATGAAGATGGCGGACGCCAAGCAGAAAAGGAATGAACAGCTAAAGCGATGGATGGGCTCG
      M K M A D A K Q K R N E Q L K R W M G S

| | 70 | 80 | 90 | 100 | 110 | 120 |

mypt1#4-6   GAAACGGACCAGGAGCCTCCTGTTTTCAAAAAGAAGAAGACGAAGGTGAAGTTCGATGAT
            E T D Q E P P V F K K K K T K V K F D D

HaeIII
| | 130 | 140 | 150 | 160 | 170 | 180 |

WT   GGCGCCGTGTTTCTGGCCGCCTGCTCAAGCGGCGATACAGAGGAGGTGCTCCGTATGCTG
      G A V F L A A C S S G D T E E V L R M L

mypt1#4-6   GGCGCCTCTGTAT----CGCCTGCTCAAGCGGCGATACAGAGGAGGTGCTCCGTATGCTG
            G A S V S P A Q A A I Q R R C S V C W
            in/del

HaeIII
| | 190 | 200 | 210 | 220 | 230 | 240 |

WT   GACCGGGGTGCTGACATCAACTATGCCAATGTGGACGGCCTCACAGCGCTCCACCAGGCA
      D R G A D I N Y A N V D G L T A L H Q A

mypt1#4-6   GACCGGGGTGCTGACATCAACTATGCCAATGTGGACGGCCTCACAGCGCTCCACCAGGCA
            T G V L T S T M P M W T A S Q R S T R H

| | 250 | 260 |

WT   TGCATCGACGACAACGTGGACATGGTGA
      C I D D N V D M V

mypt1#4-6   TGCATCGACGACAACGTGGACATGGTGA
            A S T T T W T W *

B

Fig. 2. Comparison of *WT* and *mypt1#4-6* mutant allele sequences.

Fig. 3 Organogenesis appears normal except for intestinal atresia

Fig. 4. Apical-basolateral polarity is not perturbed in *mypt1* mutant

Fig. 5 Endodermal migration to form intestinal anlage is intact in mypt1 mutants.

Fig. 6. Cell death, cell proliferation are not involved in IA development.

Fig. 7 EMT is not involved in IA formation in *mypt1* mutants

Fig. 8 mypt1_homolog phylogenic tree

Fig. 9 Expression patterns of *mypt* family genes

Fig. 10. Actomyosin is activated in the developing intestine of *mypt1* mutants

Fig. 11. Suppression of SM-myosin expression

Fig. 12 Attenuation of actomyosin activation rescues IA phenotype in *mypt1* mutants.