The human thalamus orchestrates neocortical oscillations during NREM sleep.

Thomas Schreiner[1], Elisabeth Kaufmann[2], Soheyl Noachtar[2], Jan-Hinnerk Mehrkens[3] & Tobias Staudigl[1]

1 Department of Psychology, Ludwig-Maximilians-Universität München, Munich, Germany

2 Epilepsy Center, Department of Neurology, Ludwig-Maximilians-Universität München, Munich, Germany

3 Department of Neurosurgery, Ludwig-Maximilians-Universität München, Munich, Germany

Correspondence: Tobias Staudigl (tobias.staudigl@lmu.de)

ABSTRACT

A hallmark of non-rapid eye movement sleep is the coordinated interplay of slow oscillations (SOs) and sleep spindles. Traditionally, a cortico-thalamo-cortical loop is suggested to coordinate these rhythms: neocortically-generated SOs trigger spindles in the thalamus that are projected back to neocortex. Here, we used intrathalamic recordings from human epilepsy patients to test this canonical interplay. We show that SOs in the anterior thalamus precede neocortical SOs (peak -50 ms), whereas concurrently-recorded SOs in the mediodorsal thalamus are led by neocortical SOs (peak +50 ms). Sleep spindles, detected in both thalamic nuclei, preceded their neocortical counterparts (peak -100 ms) and were initiated during early phases of thalamic SOs. Our findings indicate an active role of the anterior thalamus in organizing sleep rhythms in the neocortex and highlight the functional diversity of thalamic nuclei in humans. The thalamic coordination of sleep oscillations could have broad implications for the mechanisms underlying memory consolidation.

INTRODUCTION

The presence and coordinated interplay of slow oscillations (SOs) and sleep spindles hallmarks non-rapid eye movement (NREM) sleep [1]. SOs (~1 Hz) reflect neuronal network alterations of the membrane potential between periods of neuronal silence ('hyperpolarization', i.e., down state) and neuronal excitation (depolarization, i.e., up state) [2,3]. The origin of SOs has been traditionally located to neocortical circuits [3,4], triggering time windows of excitability and inhibition not only in neocortex but also in the thalamus and the hippocampus [3]. However, evidence from animal models casts doubt on the view that SOs are exclusively initiated and orchestrated by neocortical activity and suggest that the thalamus plays a critical role in synchronizing and coordinating SO activity [5–10].

The depolarizing phase of SOs is assumed to initiate the generation of sleep spindles within the thalamic circuitry [11]: reciprocal interactions between the thalamic reticular nucleus and thalamocortical neurons result in waxing and waning oscillations in the range of 11-16 Hz. Sleep spindles are usually nested towards the excitable up states of neocortical SOs [12–14], and are also found in the hippocampus, where they are thought to synchronize hippocampal ripples [15,16].

Importantly, this triple-coupling of sleep related oscillations has been suggested to facilitate memory consolidation, by synchronizing neuronal activity across brain regions and relaying memory representations between the hippocampus and neocortical long-term stores [17]. While a plethora of studies bolster the critical role of sleep oscillations and their coordinated interplay for the memory function of sleep [1,18], the exact neural circuits enabling their complex orchestration across brain areas are less clear, ultimately impeding our understanding of the prime mechanistic vehicle of memory consolidation.

This is particularly striking with respect to the neural circuits facilitating memory consolidation in humans, since direct neural recordings from one of the key players, the thalamus, are scarce. The anterior thalamic nuclei (ANT) and the mediodorsal thalamus (MD) have recently taken center-stage as thalamic key areas for different aspects of human memory functions [19–21], and might therefore also play a key role in coordinating sleep oscillations relevant for memory consolidation. Leveraging the rare opportunity to record intracranial electroencephalography (iEEG) from the human ANT and MD, together with simultaneous scalp electroencephalography (EEG), we investigated the interplay of NREM sleep oscillations within a thalamocortical network that is known to serve memory functions during wake [22].

We show that SOs in the ANT, but not the MD, lead neocortical SOs. These results

undermine the notion of an exclusive generation of SOs in the human neocortex and

highlight the functional diversity of specific thalamic nuclei. Sleep spindles in both the ANT

and MD preceded neocortical spindles, in line with their thalamic origin [23]. Furthermore,

we show that the nesting of sleep oscillations in the thalamus differs from the nesting in

neocortex. Thalamic spindles locked to earlier thalamic SO-phases than their neocortical

counterparts.

RESULTS

To examine thalamo-cortical interactions during human NREM sleep, we analyzed

simultaneously-recorded intracranial thalamic and scalp EEG collected across 11 full

nights of sleep from 8 patients with pharmaco-resistant epilepsy. On average, participants

slept for 8.82 ± 1.41 hours, with 63.51 ± 0.07 % spent in NREM sleep (stages N2 and N3;

see Supplementary Table 1 for proportions of times spent in each stage, Fig. 1c).

Thalamic activity was recorded from the ANT and the MD (see Fig. 1a and methods for

details on electrode localization). We isolated sleep oscillations from 27 bipolar intracranial

thalamic channels (12 ANT; 15 MD) and scalp EEG electrodes (see Supplementary Table

2 for details).To determine thalamo-cortical coordination during NREM sleep and their

specificity, we detected SOs, spindles and SO-spindle events independently in thalamic

iEEG (ANT & MD) and scalp EEG recordings, assessed their coupling and delineated the

temporal relationship of thalamo-cortical interactions (for descriptive values and

characteristics see Supplementary Tables 3-6; Fig. 1b).

Fig 1. | **Thalamic electrode placement and sleep architecture.** (a) Electrodes were implanted in the left and right ANT (blue) and MD (red). Localization of an exemplar patient's DBS leads [24] is shown on a T1-weighted template MRI with superimposed thalamus atlas [25]. (b) Example of NREM sleep segment (15 sec), comprising SOs and sleep spindles (top row: scalp recording; bottom row: ANT). (c) Hypnogram of a sample participant, showing time spent in different sleep stages across one recording night. The grey shading indicates NREM sleep stages N2 and SWS.

94
95 [24]
96
97 [25]
98
99
100

ANT SOs precede neocortical SOs

ANT SOs significantly preceded neocortical SOs, as revealed by occurrence probabilities

of ANT SOs relative to neocortical SOs (Fig. 2a, positive cluster from -0.15 to 0.05 s, p <

0.005; tested against event-free occurrence probabilities; corrected for multiple

comparisons across time; neocortical SOs were derived from frontal scalp EEG

electrodes, see Supplementary table 2 for details). ANT SO occurrence preferentially

peaked, on average, briefly before the emergence of neocortical SOs (time of peak: -0.05

sec). To further validate this outcome, we specifically determined the phase of ANT SOs

for all paired SO-events (i.e., all ANT SOs within ± 750 ms of neocortical SOs) at the time

of neocortical SO down state (i.e., thalamo-cortical SO phase-phase coupling). We found

significant nonuniform distributions in each of the ANT contacts (12/12) (p < 0.05,

corrected for multiple comparisons using the False Discovery Rate (FDR) [26]; Rayleigh test,

mean vector length: 0.46 ± 0.05). Moreover, we found a significant nonuniform

distribution across contacts (Rayleigh z = 10.95, p < 0.0001), with the phase of ANT SOs

being ahead of their neocortical counterparts (the phase of neocortical SO down states

corresponds to ± π; mean coupling direction: −176.39 ± 4.94°; see Fig. 2b).

To showcase the generic features of ANT and neocortical detected SOs, time-frequency

representations (TFRs) of SO-locked neocortical (Fig. 2c) and ANT (Fig. 2e) activity were

contrasted against event-free events. TFRs time locked to scalp-derived SO down state

peaks (see Supplementary table 2 for details) exhibited the prototypical modulation of low

frequency (< 5 Hz) and spindle power (11–16 Hz) during SO events (Fig. 2c). In particular,

low frequency power peaked before the SO down state. In contrast, spindle power was

diminished during the down state, increased during the positive deflections of the SO and

peaked during the SO up state (significant differences: p < 0.05, corrected for multiple

comparisons across time and frequency). Remarkably, while low frequency power for ANT

derived SOs showed similar pre-down state increases, power in the spindle band (11-16

Hz) and beyond (> 20 Hz) showed an early power increase tightly locked to the ANT SO

down state (Fig. 2e; p < 0.05, corrected for multiple comparisons across time and

frequency). This pattern deviates from generic modulations of spindles by SOs as

described in our and previous scalp EEG recordings, where spindles nest towards SO up

states [14]. Figure 2d illustrates the scalp EEG grand average, locked to the minimum of

neocortical SO down states. Figure 2f illustrates the ANT grand average locked to the

minimum of ANT SO down states.

Fig 2. | **ANT SOs precede neocortical SOs.** (a) Occurrence probabilities of ANT SO down state peaks relative to neocortical SO down state peaks (time = 0; dashed line; bin size = 50 ms), indicating that ANT SOs precede neocortical SOs. The solid black bar indicates significant differences, resulting from comparison with SO-free control events (positive cluster from -0.15 to 0.05 sec, p < 0.005; time of peak: -0.05 sec). (b) Phase of ANT SOs at the time of neocortical SO downstate for paired SO-SO events, illustrating that ANT SO phases preceded their neocortical counterparts (phase of neocortical SO down states corresponds to ± π; mean coupling direction: −176.39 ± 4.94°; Rayleigh test: p < 0.0001; z = 10.95). (c) Time–frequency representation of neocortical SOs (locked to neocortical SO down states), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Grand average EEG trace of neocortical SOs (mean ± SEM, negative peak, time 0, N = 1513.5 ± 130.1). (e) Time–frequency representation of all ANT SOs (locked to ANT SO down states), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (f) Grand average iEEG trace of ANT SOs (mean ± SEM, negative peak, time 0, N = 1573.8 ± 129.9).

Neocortical SOs precede SOs in the MD

Next, we asked, whether SOs leading neocortical SOs is specific to the ANT or can also be found in other thalamic nuclei. To address this question, we investigated the interplay between neocortical SOs and SOs in the mediodorsal thalamus (MD).

In stark contrast to ANT SOs, SOs in the MD did not precede, but were on average led by neocortical SOs, as evidenced by occurrence probabilities (Fig. 3a; positive cluster from -0.05 to 0.1 s, p = 0.021; corrected for multiple comparisons across time; time of peak: 0.05 sec; neocortical SOs were derived from frontal scalp EEG electrodes, see Supplementary table 2 for details). Again, we determined the phase of thalamic SOs for all paired SO-events at the time of neocortical SO down state (± 750 ms). We found significant nonuniform distributions in all MD contacts (15/15, p < 0.05, corrected for

multiple comparisons using FDR; Rayleigh test, mean vector length: 0.41 ± 0.06) and a

significant nonuniform distribution across contacts (Rayleigh z = 7.94, p = 0.0001), with

the phase of MD SOs following their neocortical counterparts (mean coupling direction:

133.51 ± 10.91°; see Fig. 3b). TFRs time-locked to scalp-derived SO down states

exhibited the typical modulation of low frequency (< 5 Hz) and spindle power (11–16 Hz)

during SO events (Fig. 3c; p < 0.05, corrected for multiple comparisons across time and

frequency). Similar to the ANT, TFRs locked to the down states of MD SOs exhibited

significant low frequency power increases before the down state-peak (Fig. 3e, time = 0),

and power increases in the sleep spindle band and beyond locked to the down state

(significant difference: p < 0.05, corrected for multiple comparisons across time and

frequency). Figure 3d illustrates the scalp EEG grand average, locked to the minimum of

neocortical SO down states. Figure 3f illustrates the MD grand average locked to the

minimum of MD SO down states.

Next, we compared ANT-neocortical and MD-neocortical interactions with regards to SOs

and tested whether ANT SOs would systematically emerge earlier (in relation to

neocortical SOs) than MD SOs. The occurrence probabilities of ANT SOs relative to

neocortical SOs differed significantly from those obtained from MD SOs relative to

neocortical SOs, with ANT SOs preferentially emerging before MD SOs (Fig 3g; positive

cluster from -0.25 to -0.05 s, p = 0.018; corrected for multiple comparisons across time).

Also, the phase distribution for the preferential coupling between thalamus and neocortex

differed significantly when comparing ANT and MD related SO coupling (Fig. 3h; Watson-

Williams test: F = 12.7, p = 0.0015), supporting the leading role of ANT SOs as compared

to the MD. Finally, we directly assessed ANT – MD related SO interactions and tested

whether ANT SOs would precede MD SOs (here occurrence probabilities were obtained

for all possible ANT – MD combinations per participant, leading to 13 distributions). In line

with the previous findings, results of this direct comparison revealed that ANT SOs

preferentially emerge before MD SOs (positive cluster from -0.15 to 0 sec, p = 0.015; time

of peak: -0.05 sec; Fig. 3i). Again, we obtained the phase of ANT SOs for all paired SO-

events at the time of MD SO down state (± 750 ms). We found a significant nonuniform

distribution across contacts (Rayleigh z = 7.44, p = 0.0001), with the phase of ANT SOs

preceding their MD counterparts (mean coupling direction: -138.34 ± 11.08°; see Fig. 3j).

Fig 3. | **Neocortical SOs precede SOs in the MD.** (a) Occurrence probabilities of MD SO down state peaks relative to neocortical SO down state peaks (time = 0, dashed line; bin size = 50 ms), illustrating that, on average, neocortical SOs lead MD SOs. The solid black line indicates significant differences, resulting from comparison with SO-free control events (positive cluster from -0.05 to 0.1 sec, p = 0.021; time of peak: 0.05 sec). (b) Phase of MD SOs at the time of neocortical SO downstate for paired SO-SO events, depicting that MD SO phases followed their neocortical counterparts (phase of neocortical SO down states corresponds to ± π; mean coupling direction: 133.51 ± 10.91°; Rayleigh test: p < 0.0001; z = 7.94). (c) Time–frequency representation of neocortical SOs (locked to neocortical SO down states), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Grand average EEG trace of neocortical SOs (mean ± SEM, negative peak, time 0, N = 1476.9 ± 129.1). (e) Time–frequency representation of all MD SOs (locked to MD SO down states), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (f) Grand average iEEG trace of MD SOs (mean ± SEM, negative peak, time 0, N = 1408.5 ± 126.2). (g) Occurrence probabilities for the comparison of ANT-neocortical (blue) and MD-neocortical SO interactions (red), showing that ANT SOs preferentially emerge before MD SOs with regards to their neocortical counterparts (positive cluster from -0.25 to -0.05 sec, p = 0.018; corrected for multiple comparisons across time;). (h) The phase distribution for the coupling between thalamus and neocortex differed significantly when comparing ANT and MD related SO coupling (Watson-Williams test: F = 12.7, p = 0.0015). (i) Occurrence probabilities of ANT SO down state peaks relative to MD SO down state peaks, showing that in a direct comparison, ANT SOs preferentially emerged before MD SOs (positive cluster from -0.15 to 0 sec, p = 0.015; time of peak: -0.05 sec). (j) Phase of ANT SOs at the time of MD SO downstate for paired SO-SO events, depicting that ANT SO phases preceded their MD counterparts (phase of MD SO down states corresponds to ± π; mean coupling direction: -138.34 ± 11.08°; Rayleigh test: p < 0.0001; z = 7.44).

ANT spindles precede neocortical spindles

Animal models established that sleep spindles are generated in the thalamus and spread to the neocortex along thalamocortical fibers [23]. However, it is unclear which of the many thalamic nuclei contribute to the generation of spindles and whether this translates to humans. Here, we show that the human ANT is putatively part of the spindle generating thalamic circuit, as ANT spindle occurrence peaked on average before the emergence of neocortical spindles (Fig. 4a; positive clusters: -0.25 to 0.05 s, $p < 0.005$ & 0.15 to 0.2 s, $p = 0.019$; corrected for multiple comparisons across time; time of peak: -0.1 sec; neocortical spindles were derived from frontal, central or parietal scalp EEG electrodes, see Supplementary table 2 for details).

We further assessed the interplay of ANT and neocortical sleep spindles using a complementary analytical approach. Specifically, we tested to what extend spindle-related oscillatory power-modulations would precede those in the neocortex, by computing power-power correlations in the sleep spindle band (see methods for details). We found a significant, off-diagonal cluster of temporal cross-correlations with regards to spindle power between ANT and neocortical sites ($p = 0.004$, cluster corrected across time; Fig. 4b), supporting our initial finding that ANT spindles lead their neocortical counterparts. Figure 4c and e depict the spindle-locked (time 0 = maximally negative amplitude) neocortical (Fig. 4c) and ANT (Fig. 4e) related TFRs. While the power of scalp derived spindles was confined to the classical sleep spindle range (~11-16 Hz; Fig. 4c), spindle – related power increases in case of ANT recordings were also evident in higher frequencies (> 20 Hz; Fig. 4e; $p < 0.05$, corrected for multiple comparisons across time and frequency). Figure 4d and f illustrate the grand average electrophysiological traces across scalp electrodes / ANT contacts (mean ± SEM, respectively), locked to the neocortical (Fig. 4d) and ANT (Fig. 4f) sleep spindle peak (maximally negative amplitude, time 0). Figure 4d illustrates the scalp EEG grand average, locked to the neocortical sleep spindle peak. Figure 4f illustrates the ANT grand average, locked to the ANT sleep spindle peak. See Supplementary Fig. 1 for results concerning spindles detected simultaneously in ANT and MD and spindles detected in ANT only.

222
223

Fig 4. | ANT spindles lead neocortical spindles. (a) Occurrence probabilities of ANT spindle peaks relative to neocortical spindle peaks (maximal negative amplitude, time = 0; dashed line; bin size = 50 ms), indicating that ANT spindles precede neocortical spindles. The solid black line indicates significant differences, resulting from comparison with spindle-free control events (positive clusters from -0.25 to -0.05 sec, p < 0.005 & 0.15 to 0.2 sec, p = 0.019; corrected for multiple comparisons across time; time of peak: -0.1 sec). (b) Power-power correlations in the sleep spindle band for paired spindle events (locked to neocortical spindle peak), contrasted against event-free correlation maps. The contour lines indicate the cluster (p = 0.004, cluster corrected across time), depicting that ANT spindle power precedes neocortical spindle power. (c) Time–frequency representation of neocortical spindles (locked to neocortical spindle peak), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Grand average EEG trace of neocortical spindles (mean ± SEM, maximally negative amplitude, time 0, N = 2203.8 ± 213.5). (e) Time–frequency representation of all ANT spindles (locked to ANT spindle peak), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (f) Grand average iEEG trace of ANT spindles (mean ± SEM, maximally negative amplitude, time 0, N = 2203.8 ± 213.5).

MD spindles lead neocortical spindles

Next, we tested the temporal relationship between MD and neocortical spindles. MD spindles occurred on average before neocortical spindles (Fig. 5a; positive clusters: -0.3 to 0.05 s, $p < 0.005$; corrected for multiple comparisons across time; time of peak: -0.05 sec neocortical spindles were derived from central or parietal scalp EEG electrodes, see Supplementary table 2 for details). We also assessed power-power correlations in the sleep spindle band for neocortical- MD paired spindle events (locked to neocortical spindle peaks). We found a significant, off-diagonal cluster of temporal cross-correlation with regards to spindle power between MD and neocortical sites (Fig. 5b; tested against event-free segments; $p = 0.002$, cluster corrected across time), indicating that MD spindle-related power modulations preceded neocortical spindle-related power.

Fig. 5c and e depict the spindle-locked (maximally negative amplitude) neocortical and MD related TFRs. Power of scalp derived spindles was, as expected, confined to the classical sleep spindle range (~11-16 Hz; Fig. 5c), while TFRs related to MD spindles exhibited power increases in the spindle band and beyond (> 20 Hz). Overall, the relationship between MD spindles and neocortical spindles was highly similar to the relationship between ANT spindles and neocortical spindles (see Figs. 4a + b). Figure 5d illustrates the scalp EEG grand average, locked to the neocortical sleep spindle peak. Figure 5f illustrates the MD grand average, locked to the MD sleep spindle peak. Figure 5g shows a comparison of ANT-neocortical and MD-neocortical interactions with regards to spindles. The timing of ANT and MD-related spindles in relation to neocortical spindles was highly similar. No significant differences emerged (cluster with lowest $p = 0.52$). See Supplementary Fig. 2 for results concerning spindles detected simultaneously in MD and ANT and spindles detected in MD only.

Fig 5. | **MD spindles lead neocortical spindles.** (a) Occurrence probabilities of MD spindle peaks relative to neocortical spindle peaks (maximal negative amplitude, time = 0; dashed line; bin size = 50 ms), indicating that MD spindles precede neocortical spindles. The solid black line indicates significant differences, resulting from comparison with spindle-free control events (positive cluster from -0.3 to 0.05 sec, p < 0.005; corrected for multiple comparisons across time; time of peak: -0.05 sec). (b) Power-power correlations in the sleep spindle band for paired spindle events (locked to neocortical spindle peak), contrasted against event-free correlation maps. The contour lines indicate the cluster (p = 0.002, cluster corrected across time), showing that MD related spindle power precedes neocortical spindle power. (c) Time–frequency representation of neocortical spindles (locked to neocortical spindle peak), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Grand average EEG trace of neocortical spindles (mean ± SEM, maximally negative amplitude, time 0, N = 1796.5 ± 229.8). (e) Time–frequency representation of all MD spindles (locked to MD spindle peak), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (f) Grand average iEEG trace of MD spindles (mean ± SEM, maximally negative amplitude, time 0, N = 2172.8 ± 238.2). (g) Direct comparison of ANT-neocortical (blue) and MD-neocortical interactions (red) with regards to spindles yielded that the timing of ANT and MD-related spindles in relation to neocortical spindles was highly similar (cluster with lowest p = 0. 52).

ANT spindles lock to early SO phases

So far, we established that both, neocortical SOs and spindles are preceded by

corresponding activity in the ANT. These results also imply that the properties of local SO-

spindle interactions, which are thought to shape memory consolidation [12], may differ

between thalamic and neocortical sites, to allow for a concomitant thalamo-cortical

coordination of SOs and spindles. To address this, we isolated SO-spindles complexes in

ANT as well as scalp recordings and extracted the onset latencies of spindles relative to

their corresponding SO down states within both regions, respectively.

In line with previous findings [16,27], occurrence probabilities of neocortical spindles with

respect to neocortical SOs (Fig. 6a), indicated that spindles started specifically at the SO

down-to-up transition (0.05 to 0.15 s; $p < 0.0001$; peak: 0.1 sec). In contrast, ANT

spindles preferential emerged around ANT SO down states (-0.15 to 0.25 s; $p = 0.0012$;

peak: 0 s; Fig. 6b). These diverging dynamics also became apparent when depicting the

SO-locked (down state) neocortical and ANT derived SO-spindle TFRs (Fig. 6c: locked to

neocortical SO down state; Fig. 6d: locked to ANT SO down state). TFRs for both ANT

and scalp derived SO-spindle complexes exhibited characteristic power-increases in the

SO and sleep spindle band ($p < 0.05$, corrected for multiple comparisons across time and

frequency). However, while neocortical spindles showed their typical power peaks well

after the corresponding down states [14], increases in the ANT sleep spindle band were

initiated around the ANT SO down states.

Finally, directly comparing the occurrence probabilities of thalamic and neocortical SO-

spindle complexes affirmed that ANT spindles locked significantly earlier to SOs than

neocortical spindles (-0.3 to -0.1, $p = 0.0039$; Fig. 6e). In a complementary analytical

approach, we determined the preferred phase of SO-spindle modulation (i.e., SO-spindle

coupling), for both ANT and neocortical events, respectively, by assessing the SO phases

corresponding to the spindle onset latencies in each contact / scalp electrode. In 12/12

ANT contacts we found significant nonuniform distributions ($p < 0.05$, corrected for

multiple comparisons using FDR; Rayleigh test, mean vector length: 0.41 ± 0.03) and a

significant nonuniform distribution across contacts (Rayleigh $z = 9.33$, $p < 0.0001$), with

spindles starting near the SO down state (corresponding to $\pm 180°$; mean coupling

direction: $-175.22° \pm 6.92°$; see Fig. 6f). Similarly, we found in 7/8 scalp contacts a

significant nonuniform distribution ($p < 0.05$, corrected for multiple comparisons using

FDR; Rayleigh test, mean vector length: 0.27 ± 0.02). Again, a significant nonuniform

distribution across participants was present (Rayleigh $z = 6.91$, $p < 0.0001$), with spindles

preferentially starting at the SO down-to-up transition (mean coupling direction: -137.14

$\pm 7.58°$; see Fig. 6f). Next, we tested whether the preferred coupling phases would vary

systematically between ANT and neocortical areas, using the circular Watson-Williams

test. Indeed, we found a significant difference in relation to the preferred phase of ANT

and neocortical SO-spindle modulation ($F = 9.31$; $p = 0.0069$), corroborating the finding

that spindles in the thalamus lock to earlier phases of the SO. Finally, we quantified the

directional influence of SOs on spindles in both ANT and neocortical data, using the PSI [28]. We found that SOs predicted sleep spindle activity within ANT and neocortex, respectively, as evidenced by a positive PSI (ANT: 0.0040 ± 0.0017; t-test against zero: t = 2.37, p = 0.037; neocortex: 0.0048 ± 0.002; t-test against zero: t = 2.44, p = 0.044; Fig. 6g).

Fig 6. | ANT spindles lock to early SO phases. (a) Occurrence probabilities of neocortical spindle onsets with respect to neocortical SO down states (bin size = 50 ms), illustrating that spindles started specifically at the SO down-to-up transition (0.05 to 0.15 sec; p < 0.0001; peak: 0.1 sec). (b) ANT spindles preferential emerged around ANT SO down states (-0.15 to 0.25 sec; p = 0.0012; peak: 0 sec). (c) Time–frequency representation of neocortical SO-spindle events (locked to neocortical SO down state peaks), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Time–frequency representation of all ANT SO-spindle events (locked to ANT SO down state peaks), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (e) Direct comparison of the occurrence probabilities of ANT (blue) and neocortical SO-spindle complexes (gray) yielded that ANT spindles locked significantly earlier to SOs as compared to neocortical spindles (-0.3 to -0.1, p = 0.0039). (f) Phases of the SO-spindle modulation derived from neocortical (gray circles) and ANT (blue circles) events. Neocortical spindle onsets preferentially emerged at the neocortical SO down-to-up transition (phase of neocortical SO down states corresponds to ± π; mean coupling direction: −137.14 ± 7.58°; Rayleigh z = 6.91, p < 0.0001), while ANT spindles started specifically around the ANT SO down states (mean coupling direction: −175.22° ± 6.92°; Rayleigh z = 9.33, p < 0.0001). The preferred phases of SO-spindle modulation differed significantly between the ANT and neocortical sites (Watson-Williams test: F = 9.31; p = 0.0069). (g) Directional SO-spindle coupling as obtained by the phase-slope index (PSI; mean ± SEM) for ANT (blue circles) and neocortical (gray circles) SO-spindle complexes. SO-phases significantly predicted spindle amplitudes both in case of ANT (t-test against zero: p = 0.037) and neocortical SO-spindle events (t-test against zero: p = 0.044).

MD spindles lock to early SO phases

We have shown that neocortical SOs precede MD SOs, while MD spindles lead

neocortical spindles. Accordingly, the coupling within MD related SO-spindle complexes

is expected to be shifted compared to neocortical SO-spindle complexes. Occurrence

probabilities indicated that neocortical spindles started specifically at the down-to-up

transition of neocortical SOs (0.1 to 0.3 s; p = 0.003; peak: 0.1 s; Fig 7a). MD spindles,

however, started preferentially briefly before the MD SO down states (-0.35 to 0 s; p <

0.0001; peak: -0.05 s; Fig 7b). Figures 7c and d show the neocortical and MD SO-spindle

TFRs (Fig. 7c: locked to neocortical SO down states; Fig. 7d: locked to MD SO down

state). TFRs for both scalp and MD derived SO-spindle complexes featured power-

increases in the SO and sleep spindle band (p < 0.05, corrected for multiple comparisons

across time and frequency). Neocortical spindles exhibited their typical power peaks after

the corresponding down states, while power peaks of MD sleep spindles started around

MD SO down states. Accordingly, the direct comparison of occurrence probabilities

revealed that MD spindles locked significantly earlier to accompanying MD SOs as did

neocortical spindles to neocortical SOs (-0.35 to 0.05, p < 0.0001; Fig. 7e).

Next, we determined the preferred phase of SO-spindle modulation for MD and

neocortical events, respectively, by assessing the SO phases corresponding to the

spindle onset latencies in each contact / scalp electrode. In 15/15 thalamic contacts, we

found significant nonuniform distributions (p < 0.05, corrected for multiple comparisons

using FDR; Rayleigh test, mean vector length: 0.42 ± 0.03). Also, across contacts the

distribution was significantly nonuniform (Rayleigh z = 12.10, p < 0.0001), with spindles

starting near the SO down state (corresponding to ± 180°; mean coupling direction:

172.26° ± 6.67°; see Fig. 7f).

For scalp derived data, we found in 9/11 contacts a significant nonuniform distribution (p

< 0.05, corrected for multiple comparisons using FDR; Rayleigh test, mean vector length:

0.21 ± 0.02). Again, a significant nonuniform distribution across participants was present

(Rayleigh z = 8.18, p < 0.0001), with spindles preferably starting at the down-to-up

transition of neocortical SOs (mean coupling direction: −138.43 ± 9.05°; see Fig. 7f).

When directly comparing the preferred coupling phases between MD and neocortical

areas, we found a significant difference (Watson-Williams test: F = 17.33; p = 0.0003),

supporting the outcome that spindles in the MD start at earlier phases of the SO then

neocortical spindles. We also quantified the directional influence of SOs on spindles in

both MD and neocortical data, using the phase-slope index [28]. Again, we show that SOs

predicted sleep spindle activity within MD and neocortex, respectively, as evidenced by a

positive PSI (MD: 0.0016 ± 0.0006; t-test against zero: t = 2.36, p = 0.032; neocortex:

0.0028 ± 0.002; t-test against zero: t = 2.23, p = 0.049; Fig. 7g). Finally, we compared

the features of ANT- and MD-derived SO-spindle complexes. While MD related sleep

spindles had a tendency to lock to even earlier phases of MD related SOs (Fig. 7h), this

difference was not significant (occurrence probabilities: p = 0.11; Fig. 7i: phase

distribution: F = 1.29; p = 0.25). There was no difference in the SO-spindle coupling rate

between ANT and MD (t = 0.6057, p = 0.55).

Fig7. | MD spindles lock to early SO phases. (a) Occurrence probabilities of neocortical spindle onsets with respect to neocortical SO down states (bin size = 50 ms), indicating that spindles started specifically at the SO down-to-up transition (0.05 to 0.15 sec; p < 0.0001; peak: 0.1 sec). (b) MD spindles started around MD SO down states (-0.35 to 0 sec; p < 0.0001; peak: -0.05 sec). (c) Time–frequency representation of neocortical SO-spindle events (locked to neocortical SO down state peaks), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (d) Time–frequency representation of all MD SO-spindle events (locked to MD SO down state peaks), contrasted against event-free segments. The contour lines indicate clusters (p < 0.05, corrected for multiple comparisons across time and frequency). (e) Direct comparison of the occurrence probabilities of MD (red) and neocortical SO-spindle complexes (gray) revealed that MD spindles start at significantly earlier SO phases as compared to neocortical spindles (-0.35 to 0.05, p < 0.0001). (f) Phases of the SO-spindle modulation derived from neocortical (gray circles) and MD (red circles) events. Neocortical spindle onsets preferentially emerged at the neocortical SO down-to-up transition (phase of neocortical SO down states corresponds to ± π; mean coupling direction: −138.43 ± 9.05°; Rayleigh z = 8.18, p < 0.0001), while MD spindles started specifically around the MD SO down states (mean coupling direction: 172.26 ± 6.67°; Rayleigh z = 12.10, p < 0.0001). The preferred phases of SO-spindle modulation differed significantly between the MD and neocortical sites (Watson-Williams test: F = 17.33; p = 0.0003). (g) Directional SO-spindle coupling obtained by the phase-slope index (PSI; mean ± SEM) for MD (red circles) and neocortical (gray circles) SO-spindle complexes. SO-phases significantly predicted spindle amplitudes both in case of MD (t-test against zero: p = 0.032) and neocortical SO-spindle events (t-test against zero: p = 0.049). (h + i) Directly comparing ANT- (blue) and MD-derived SO-spindle complexes (red) did not yield a significant difference (h) occurrence probabilities: p = 0.11; (i) phase distribution: F = 1.29; p = 0.25).

Fig. 8 | Schematic overview of the interplay between sleep oscillations recorded in the ANT and on the scalp. (a) ANT SOs (bottom, blue) preferentially peak on average ~50 ms before the emergence of neocortical SOs (top, grey). (b) ANT spindles (bottom, blue) peak on average 100 ms before neocortical spindles (top, grey). (c) ANT spindles couple to earlier ANT SO phases (bottom, blue) than neocortical spindles couple to neocortical SOs (top, grey). While neocortical oscillations show the prototypical SO-spindle modulation (i.e., nesting of spindles towards the SO upstate), spindles in the ANT start closer to the ANT SO down-state. The insets illustrate the difference in the phase locking of spindle start to their respective SO (the phase of SO down states corresponds to ± π).

DISCUSSION

Our results suggest that anterior thalamic activity might play a fundamental role in the

orchestration of cardinal sleep rhythms in humans. In particular, we found that SOs

recorded in the human ANT preceded their neocortical counterparts (average peak: -50

ms), in contrast to SOs recorded in the MD, which were led by neocortical SOs (average

peak: +50 ms). Spindles recorded in ANT as well as MD preceded neocortical spindles

(average peak: -100 ms). Finally, both in the ANT and the MD, thalamic spindles locked

to early SO-phases, facilitating the prototypical synchrony between SOs and spindles in

the neocortex.

Sleep oscillations have gained considerable interest over the last decade, mainly due their

presumed role in facilitating the memory function of sleep [18]. SOs are thought to represent

the time-giving pacemakers of memory consolidation and have been studied extensively

in both rodent models and during human sleep [14,29]. Traditionally, they have been assumed

to be solely generated within neocortical circuits [3,4]. However, recent animal studies cast

doubt on this view by identifying essential thalamic contributions for SO generation and

coordination [5-10]. Thalamic neurons have been shown to exhibit intrinsic, rhythmic up and

down states in isolated conditions (i.e., without neocortical inputs [5,30]), while phasic burst-

firing of thalamocortical neurons at the onset of up states often precedes the firing of

neocortical neurons by 20 - 50 ms [7,31]. Optogenetic stimulation of thalamocortical neurons

efficiently initiates neocortical up states [6,32]. In addition, thalamic activity terminates SO up

states synchronously across neocortical areas, thus mediating a well-controlled down

state transition [33]. In sum, these findings suggest that thalamic activity exerts a controlling

influence on neocortical SOs, by coordinating up and down state dynamics in neocortical

circuits. However, the generalization of thalamic contributions, and specifically of distinct

thalamic nuclei, to sleep oscillations has been shown to be challenging across species

(e.g., [34,35]), prompting their direct assessment during human sleep.

To investigate potential contributions of the ANT to neocortical SO coordination, we

detected SOs during NREM sleep in intracranial contacts located in the ANT as well as

scalp EEG. We show that the ANT not only exhibits a slow oscillatory rhythm (note that

the ANT signal is likely to be of local origin due to a bipolar montage), but, importantly,

that ANT SOs precede their neocortical counterparts. This might indicate that the ANT

guides SO activity in the neocortex. Focusing on paired events only (i.e., time-windows

where both neocortical and ANT SOs were present), we found that ANT SO phases

significantly lead neocortical SO phases. Notably, this pattern of results was specific to

ANT recordings. SOs in the MD did not precede but rather followed neocortical SOs,

which is in line with a previous report of neocortical SOs leading SOs in the human pulvinar

[13]. Our findings thus highlight both a specific role of the ANT for SO dynamics and, more

generally, the functional diversity of specific thalamic nuclei in humans [36], also with regards

to sleep oscillations potentially originating there. Since scalp EEG recordings provide an

indirect measure of the underlying cortical sources, one might ask whether a delay in

cortical SOs relative to ANT SOs might be driven by effects of interjacent tissue (e.g., the

skull) delaying the timing of the signal picked up at frontal scalp EEG channels. To mitigate

these concerns, we show, in an independent data set, that intracranially recorded SOs in

the frontal lobe exhibit no temporal delay with regards to SOs simultaneously recorded by

frontal scalp EEG (see Supplementary Fig. 3). Another concern might arise considering

volume conduction contaminating the thalamic signal. We used a bipolar montage for

thalamic contacts to ensure a local origin of the signal [37–39]. However, there might still be

a possibility that the signal recorded from these contacts is contaminated by volume

conducted activity from adjacent brain tissue outside the thalamus. We address this issue

in a dedicated control analysis (see Supplementary Figure 4) and conclude that none of

these factors is likely to systematically bias our findings. The dissociation of SO timing

between ANT and MD SOs, which are anatomical neighbors, also speaks against a global,

volume conducted source driving the effects. In addition, our data were recorded in

epilepsy patients in whom anti-epileptic medication and epileptiform events may affect
sleep architecture and sleep related oscillations [40–43], warranting caution when
generalizing to healthy population. However, our patient population was heterogeneous
in terms of medication, age, epilepsy form and affected cortical area. Furthermore, sleep
architecture and SO- and spindle quantity and densities were within normal ranges of
healthy participants (see Supplementary tables 3 and 4, [44,45]), making it unlikely that a
single aspect had a systematic impact on our results. In addition, while only seizure-free
nights were included in the analysis, our strict artifact rejection procedure minimized the
impact of epileptiform activity on our findings.
425
What makes the ANT a favorable candidate for sculpting neocortical SOs? The ANT is a
key node in the limbic circuit [46] and functionally relevant for human memory [20,22]. It has
extensive connections with the anterior cingulate, orbitofrontal cortex and the
hippocampus [47,48]. Notably, medial prefrontal regions, such as the orbitofrontal and the
cingulate cortex have been identified as neocortical hot spots for the emergence of SO
during human NREM sleep [49,50]. In addition, work in rodent models has shown that midline
and anterior thalamic nuclei are likely candidates to influence global neocortical activity
during sleep due to their widespread projections [7,33]. Hence, the dense connectivity of the
ANT with medial prefrontal regions and within the limbic circuit sets the stage for ANT
activity to shape neocortical SOs and coordinate sleep oscillations between regions
implicated in memory consolidation.
We here show ANT SOs leading neocortical SOs on the basis of approximately 30% of
neocortical SOs that were coupled to ANT SOs. Neocortical SOs that were not coupled
to the ANT may as well have been initiated in the neocortex, in line with traditional
accounts of the origin of SOs [3,4]. Future research needs to investigate whether ANT
coordinated SOs are functionally distinct from neocortically coordinated SOs, for example
with respect to memory reprocessing during sleep. The precise coupling between SOs
and thalamic spindles might be of relevance here, given that the precision of the coupling
between SOs and spindles has been shown to be instrumental for memory consolidation
to unfold (e.g., [12,27]). Due to its prominent location within the limbic circuit and direct
projections to the hippocampal formation [21], the ANT could also be important for
integrating hippocampal ripples into SO-spindle complexes, facilitating the full triple-
coupling of sleep oscillations that is assumed to mediate the memory functions of sleep
[18]. Hence, our work represents a starting point for elucidating the distinct role of the ANT
in coordinating sleep oscillations and memory consolidation.

We also found clear evidence for spindles in ANT and MD. Spindles emerge in recurrent

loops between the thalamic nucleus reticularis (TRN) and thalamocortical neurons, which

in turn forward them to the neocortex and hippocampus [23]. There have been conflicting

findings across animal species about whether the TRN innervates the ANT and, hence,

whether or not spindles are present in the ANT [19,34,35,51–53]. Our results provide a clear

picture in humans: spindles are not only present both in the ANT and MD but also precede

neocortical spindles, indicating an involvement of both nuclei in spindle generation and

their projection to neocortical areas. In line with our interpretation, optogenetic stimulation

of the rodent anterodorsal TRN, which innervates the ANT [35,54], has been shown to initiate

spindles in the neocortex and hippocampus [55], indicating that the ANT might synchronize

key areas of memory consolidation [56]. With regards to the MD it has been found that its

neurons exhibit increased firing during spindle events [57] in rodents. In humans, reduced

MD volume in persons with schizophrenia is associated with dampened neocortical

spindle activity [58], further pointing towards a role of the MD in the thalamo-cortical spindle

transfer. In line with previous work on spindles in posterior thalamic nuclei [59], we identify

local and non-local spindles in ANT and MD. With respect to the neural circuitry

generating spindles in ANT and MD, one could assume recurrent loops between TRN and

thalamocortical neurons, similar to what has been shown in animals [11,23]. Extrapolating

these findings from animals to humans needs to be done with caution, however, since

result from animal models concerning TRN innervation of thalamic nuclei is not always

univocal [34,35]. In general, it is assumed that TRN is organized in sectors that

topographically project to thalamic nuclei [35], which can explain why local spindles emerge

in both MD and ANT. However, the mapping of TRN sections to thalamic nuclei is not one-

to-one [60]: for example, some TRN sectors projecting to the MD also project to ANT [61–63].

Such overlapping projections seem plausible when considering that MD and ANT

thalamocortical neurons share common targets, for example in the prefrontal cortex.

In sum, our results extend previous research identifying sleep spindles in other human

thalamic nuclei [13,59], by showing that sleep spindles in the ANT and MD precede

neocortical spindles during human sleep, indicating a role of both nuclei in thalamo-

cortical spindle projections.

Finally, we set out to characterize the properties of SO-spindle interactions within

the ANT and the MD. Spindles tend to nest within SO up states in scalp EEG recordings

[12,14,27]. Importantly, the precise timing of neocortical spindle peaks with respect to the SO

upstate has been shown to be tightly linked to the behavioral expressions of memory

consolidation [12,27] putatively by igniting structural changes in neocortical sites [64]. Hence,

the characteristics of SO-spindle interactions in the ANT might highlight how the thalamus

simultaneously groups SOs and spindles in the neocortex, leading to the well-known

oscillatory nesting (i.e., spindles nesting towards SO up-states). We found that spindles

in the ANT were initiated around the thalamic SO down state, in contrast to neocortical

spindles that preferentially started later, at the neocortical SO down to up state transition

(leading to the well-known nesting of neocortical spindles towards the up states) [1,14].

Thus, ANT spindles started at significantly earlier ANT SO phases as compared to

neocortical SO-spindle interactions. Notably, this divergent property of SO-spindle

coordination in the ANT is a consequence of the delays between ANT and cortical sleep

oscillations we found. It allows spindles, putatively generated in the ANT, to arrive during

SO up-states on neocortical sites, generating the prototypical SO-spindle coupling found

in scalp EEG. Specifically, while we show that both ANT SOs and spindles preceded their

neocortical counterparts, the time lag between ANT and neocortical spindles exceeded

the time lag between ANT and neocortical SOs (100 ms vs. 50 ms). Hence, to enable the

prototypical SO-spindle modulation in neocortical circuits (i.e., nesting of spindles towards

the SO upstate) under the assumption that both graph-elements in the neocortex are

governed by the ANT, it is inevitable that spindles start at relatively earlier phases of the

SO in the ANT as compared to spindles of neocortical SO-spindle complexes. Without

such a shifted coordination on the thalamic level, cortical SO-spindle coupling would be

systematically misaligned, with spindles regularly peaking after SO up-states (see Fig. 8

for a schematic illustration).

We found a similar pattern of SO-spindle interactions within the MD, with spindles starting

at early phases of the SO. Interestingly, even though not significant, spindles in the MD

seem to start even earlier with respect to SOs as it is the case in the ANT. Again, in order

to project spindles to the neocortex that nest in neocortical up states in a timely manner,

a shifted coordination of SOs and spindles in the MD is necessary (i.e., spindles starting

at early SO phases). This seems to be even more relevant in the case of the MD, where

SOs were preceded (and potentially driven) by neocortical SOs, while spindles preceded

neocortical spindles. Hence, for MD spindles to arrive at the neocortex during the

presence of an SO-upstate, it is inevitable to start at an early MD SO-phase. The tight

coupling of thalamic spindles to thalamic down-states is in line with previous work showing

a very similar temporal relationship in the human pulvinar [13]. These results indicate that

the hyperpolarization during thalamic down-states is required to trigger the neuronal

cascade underlying spindle generation.

Taken together, our findings provide first evidence for the ANT as a major hub for
coordinating the cardinal NREM-sleep related oscillations. In the neocortex, both spindles
and SOs were led by ANT-related activity, while the interplay of local ANT SO-spindles
was tuned in a way that allows for neocortical spindles to nest within SO up states (ANT
spindles starting at early ANT SO phases, see Fig. 8). Importantly, this exact coordination
is thought to be instrumental for the consolidation of memories [1]. Hence, the ANT might
represent a key-player in governing memory re-processing during human NREM sleep.
Due to its projections to the medial prefrontal cortex and the hippocampus, the ANT has
already been implicated in facilitating memory functioning and spatial navigation during
wake, putatively by facilitating hippocampal-neocortical interactions [22,56,65–67]. Notably, the
same hippocampal-cortical dialogue is assumed to underly sleep-related memory
consolidation, facilitated by a precise coupling of hippocampal ripples, neocortical SOs
and thalamic spindles [1]. The standard model suggests that neocortical SOs are imposed
on a rather passive thalamus, where they initiate spindles. Spindles in turn are thought to
synchronize hippocampal ripples and reactivated memory information [16,68]. Through this
coordination the reactivated memory information is suggested to arrive at the neocortex
during periods of high plasticity (i.e., during the presence of SO up states and spindles)
[64].
Our data critically extend these theoretical considerations, by spotlighting the human
thalamus, and in specific the ANT, as a putative active agent in interfacing sleep-related
oscillations between brain regions and thereby, potentially facilitating memory
consolidation. While our current data remain agnostic with regards to hippocampal
activity, it has been shown in a single patient study, that hippocampal spindles are aligned
with SOs in the ANT [69]. Hence, ANT related oscillations might not only migrate to the
neocortex but likewise to the hippocampus, where spindles are known to govern ripples
and associated memory reactivation [70].Taken together, ANT related activity might
critically contribute to the triple-coupling of sleep oscillations and thus, the memory
function of sleep. Future work relating the association of ANT-neocortical interactions with
the behavioral expressions of memory consolidation will elucidate the memory function of
the ANT and further characterize the functionally diverse human thalamus.
551

METHODS

Participants.

Intracranial and scalp EEG were simultaneously recorded from patients with bilateral

depth electrodes implanted in the thalamus for deep brain stimulation (DBS) therapy of

pharmaco-resistant epilepsy. Data was recorded at the Epilepsy Center, Department of

Neurology, Ludwig-Maximilian Universität (Munich, Germany), while DBS leads were

externalized post-surgery, prior to their connection to the pulse generator located in a

subcutaneous infraclavicular or abdominal pouch [71]. The availability of direct iEEG

recordings from the human thalamus is highly limited due to (i) the rarity of patients being

treated with thalamic DBS, (ii) access to thalamic iEEG in these patients (post-surgery

externalization of DBS leads is performed only in a fraction of these patients), and (iii) the

limited amount of time for externalization (after surgery and before connecting the leads

to the pulse generator).

A total of 13 patients participated. The data of 5 patients were discarded due to excessive

epileptic activity. Out of the remaining 8 patients (mean age: 38.65 ± 3.59; 5 female), 6

patients contributed single full-night recordings, one patient contributed two consecutive

full-night recordings and one patient contributed three consecutive full-night recordings.

In sum, 11 full-night recordings entered the analyses of the present study. All patients

received anticonvulsive medication (see Supplementary Table 7 and 8 for each

participant's drug regimen at the time of recordings and their influence on sleep

architecture). Informed consent was obtained from all patients. The study was approved

by the ethics committee of the Medical Faculty of the Ludwig-Maximilian Universität.

Intracranial data acquisition and electrode localization

iEEG data was recorded from thalamic depth electrodes that each had four intracranial

electrode contacts (platinum–iridium contacts, 1.5 mm wide with 1.5 mm edge-to-edge

distance; Medtronic 3387). Data were recorded using XLTEK Neuroworks software

(Natus Medical, San Carlos, California, US) and an XLTEK EMU128FS amplifier, with

voltages referenced online to a CPz (250 Hz sampling rate in 7 patients; 200 Hz sampling

rate in 1 patient).

The ANT in both hemispheres are the clinically-relevant implantation target for DBS

therapy of epilepsy [72]. Due their small and high inter- and intraindividual variable size and

the transventricular implantation trajectory, a subset of the electrode contacts ends up

outside the ANT, frequently in the mediodorsal thalamus (see Fig. 1a).

The locations of the electrode contacts were estimated using the Lead-DBS toolbox [24].
First, the post-operative CT scan was co-registered to the pre-operative T1-weighted
MRI, as implemented in the Advanced Normalisation Tools [73]. The scans were then
spatially normalized to MNI space, based on the pre-operative T1 image using the unified
segmentation approach as implemented in SPM12 [74]. Next, the trajectories of the DBS
electrodes and positions of the electrode contacts were reconstructed based on the post-
operative CT scan. The positions of the electrode contacts where then visually confirmed
using the DISTAL atlas [25]. To mitigate effects of volume conduction on our analyses [75], the
iEEG recordings were then re-referenced to their immediate neighboring contact (bipolar
montage; see Supplementary Fig. 4 for an estimation of the potential influence of
remaining volume conducted SOs on the main results). A bipolar pair of contacts was
considered to be in the ANT if both or at least one contact was localized to the ANT, but
no contact was localized in the MD (see Supplementary Fig. 5 for a comparison of full
ANT contacts and single ANT contacts with regards to occurrence probabilities of SOs,
spindles and SO-spindles). A bipolar pair of contacts was considered to be in the MD if
both or at least one contact was localized to the MD, but no contact was localized in the
ANT. This procedure yielded 20 thalamic contacts (7 patients) in the ANT, and 24
contacts (8 patients) in the MD (see Supplementary Fig. 6 for a comparison of left and
right ANT / MD contacts with regards to occurrence probabilities of SOs and spindles).
To investigate whether the signal picked-up at the scalp level might have been delayed
by interjacent tissue (e.g., the skull), we assessed SO coupling between frontal scalp
electrodes (electrode FP1) and frontal intracranial contacts in an independent data set
consisting of three pre-surgical epilepsy patients (for details see Supplementary Fig. 3).
610

Scalp EEG acquisition
Scalp EEG was simultaneously recorded with iEEG via a common amplifier. Scalp EEG
electrodes were placed on the accessible scalp according to the international 10-20
system for electrode placement. The sampling rate of the scalp EEG was identical to the
sampling rate of the iEEG. Scalp EEG was recorded from 20 (2 patients), 22 (3 patients),
24 (1 patient) or 36 electrodes (2 patients). Scalp EEG was online referenced to CPz.
Scalp EEG data was re-referenced offline against the mean of all available artifact-free
scalp EEG electrodes.
619

621   Sleep staging

622 All available scalp electrodes were low-pass filtered at 40 Hz, down-sampled to 200 Hz (if
623 necessary). Sleep staging was carried out according to standard criteria [76]. For
624 subsequent analyses, sleep stages N2 and N3 were collapsed and referred to as non-
625 rapid eye movement (NREM) sleep.
626
Data pre-processing
iEEG and scalp EEG were manually inspected to discard noisy or artefact-contaminated
(e.g., excessive movements during NREM) channels (see below for detection of epileptic
activity). All data were down-sampled to 200 Hz, demeaned and de-trended. The
recordings were then filtered using a 150 Hz Butterworth low-pass filter and a two
Butterworth band-stop filters (to attenuate line noise; 48-52 Hz).
633
Interictal epileptic discharge (IED) detection: In general, only seizure-free nights were
included in the analysis. Epileptic activity in scalp and thalamic recordings was semi-
automatically detected. First, interictal epileptic discharges were detected automatically
by high- and low pass filtering the data below 20 and above 80 Hz and z-scoring the data.
Subsequently, data segments with amplitudes exceeding the mean signal by 6 SDs for
less than 100 ms were identified. Second, iEEG and scalp EEG data was visually
inspected for epileptic activity by two investigators, independently. Data segments
comprising epileptic events at any given channel were discarded (13.91 ± 2.41 % of all
sleep data). Four thalamic contacts (2 ANT contacts, 2 MD contacts) and on average 7.5
± 0.92 scalp electrodes were discarded.
644
Event detection
SOs and sleep spindles were identified for each patient, based on established detection
algorithms [16,27].
SO detection: Data were filtered between 0.3–2 Hz (two-pass FIR bandpass filter, order
= three cycles of the low frequency cut-off). Only movement-free data (as determined
during sleep scoring) from NREM sleep stages 2 and 3 were considered. All zero-
crossings were determined in the filtered signal of each channel (iEEG and scalp EEG,
respectively), and event duration was determined for SO candidates (that is, down states
followed by up states) as time between two successive positive- to-negative zero-
crossings. Events that met the SO duration criteria (minimum of 0.8 and maximum of 2
sec) and exceeded the mean amplitude of all detected events by 1.25 SD entered the
analysis. For subsequent time-locked analyses (ERPs, time-frequency representations,

etc.), 5-sec-long segments centered on the down state (± 2.5 sec) were extracted from

the unfiltered raw signal.

Spindle detection: Data was filtered between 11–16 Hz (two-pass FIR bandpass filter,

order = three cycles of the low frequency cut-off). Again, only artifact-free data from NREM

sleep stages 2 and 3 were used for event detection. The root mean square (RMS) signal

was calculated for the filtered signal at each channel using a moving average of 200 ms,

and spindles that exceeded the mean amplitude of all detected events by 1.5 SD entered

the analysis. Whenever the signal exceeded this threshold for more than 0.5 sec but less

than 3 sec (duration criteria), a spindle event was detected. Spindle onsets were defined

as the points in time when the signal initially crossed the threshold. Epochs time-locked

to the maximally negative spindle trough (−2.5 to +2.5 sec) were extracted from the

unfiltered raw signal for subsequent time-locked analyses.

SO-spindle complexes: To isolate SO-spindle complexes in scalp and thalamic

recordings, we determined for all SOs whether a spindle was detected following the SO

(SO down state + 750 ms). Again, SO-spindle events were extracted (−2.5 to +2.5 sec

with regards to the SO down state) from the raw signal for subsequent time-locked

analyses.

Paired events: To identify paired SO events (scalp & thalamus), we determined for all

scalp detected SOs whether corresponding SOs were detectable in thalamic recordings

within a time-window of ± 750 ms. Data segments locked to the scalp detected down

states (± 2.5 sec) were extracted. Likewise, paired sleep spindles were isolated by

determining for all scalp detected sleep spindles whether related spindles emerged (±

750 ms) in thalamic recordings. Data segments locked to the scalp detected maximally

negative spindle peak (± 2.5 sec) were extracted.

Electrode selection [scalp EEG]: As outlined above, all electrodes exhibiting significant

epileptic activity were discarded to ensure that our results were not influenced by

epileptiform activity. This conservative procedure together with restricted access to the

scalp due to post-surgical wound care impeded us from using identical scalp EEG

electrodes across patients to delineate thalamo-cortical interactions during sleep (e.g.,

electrode Fz for SO related and electrode Cz for sleep spindle related analyses [16]. Instead,

we pursued a data-driven and individualized approach. For SO-related analyses, we

selected for each participant the frontal scalp electrode that exhibited the highest number

of associated thalamic SOs within a time of ± 750 ms. This procedure was accomplished

independently for data stemming from the ANT and the MD. The respective scalp

electrodes (one related to the ANT, one related to the MD) were used for all subsequent

SO-related analyses (see Supplementary Table 1 for on overview of the selected scalp

electrode; see Supplementary Fig. 7 for ANT SO occurrence probabilities relative to scalp

electrode Fz; see Supplementary Fig. 8 for ANT SO occurrence probabilities relative to

occipital scalp electrodes). Similarly, for sleep spindle related analyses, we selected the

frontal, central or parietal scalp electrode that exhibited the highest number of associated

thalamic spindles (ANT or MD, respectively; see Supplementary Table 2 for on overview

of the selected scalp electrodes; see Supplementary Fig. 9 for spindle-spindle occurrence

probabilities, were scalp spindles were exclusively derived from frontal or parietal

electrodes; for slow and fast spindle related results see Supplementary Fig. 10, 11 and

Supplementary table 9). The respective scalp electrodes (one related to the ANT, one

related to the MD) were subsequently used for all sleep spindle-related analyses. Data

from scalp electrodes was used as a proxy of neocortical activity.

Event-free segments: For statistical comparisons, we extracted 5-sec-long intervals

during NREM sleep, which did not exhibit any SO, spindle or SO-spindle event,

respectively. In each participant and electrode, the number of control events

corresponded to the number of prior detected events of interest. Event-free events were

only drawn from time window starting 5 mins before and ending 5 mins after the

corresponding oscillatory event.

Analyses

Temporal relation between sleep oscillations: To assess the temporal relationship

between scalp- and thalamus-derived SO events, we created peri-event time histograms

(bin size = 50 ms) where scalp detected SO down states served as seed (time = 0), while

the targets (thalamic SO down states) are depicted relative to the seed. The resulting

histogram was normalized by the total number of detected scalp SOs (multiplied by 100)

per participant and electrode site. Similarly, for sleep spindles, peri-event histograms (bin

size = 50 ms) of thalamic sleep spindle peaks (maximally negative amplitude) in relation

to scalp sleep spindles peaks (maximally negative amplitude: ± 1.5 sec) were created.

The resulting histogram was normalized by the total number of detected scalp spindles

(multiplied by 100). To determine the interplay of SOs and sleep spindles (separately

within scalp and thalamic recording), a peri-event histogram of sleep spindles (onsets)

around SO down states (± 1.5 sec) was created. The resulting histogram was normalized

by the total number of detected spindles (multiplied by 100). The same procedure was

applied to spindle-free control segments.

Phase relationship between sleep oscillations: To estimate the phase-relationship

between scalp and thalamic SOs, data epochs including paired events were filtered in the

SO range (0.3–2 Hz, two-pass Butterworth bandpass filter). As described above, these

segments included both scalp and thalamic SOs, and were time-locked to the down states

of scalp detected SOs. A Hilbert transform was applied to the data and the instantaneous

phase angle of the thalamic recording at the time of the scalp SO down state was

extracted. Each participant's preferred phase of SO-SO coupling was then obtained by

taking the circular mean of all individual events' preferred phases.

For the analysis of SO-spindle coupling, we filtered the SO-spindle data (locked to spindle

onsets) in the SO range (0.3–2 Hz, two-pass Butterworth bandpass filter), applied a

Hilbert transform and extracted the instantaneous phase angle. Next, we isolated the SO

phase angle at the time of spindle onsets. Each participant's t preferred SO phase at

spindle onsets was obtained by taking the circular mean of all individual events' preferred

phases.

Spindle related across-channel power correlations: We further estimated the temporal

relationship of thalamic and scalp detected spindles using channel to channel correlations

of oscillatory power in the sleep spindle band. To this end, a Hanning window was applied

to paired sleep spindles segments (locked to scalp detected maximally negative

amplitude) and power values between 11 and 16 Hz were extracted for each time-bin.

Power values were z-scored per channel and averaged. Subsequently, power values

derived from thalamic channels were correlated with scalp related power values at each

time-point (Spearman correlation), resulting in a 2d correlation map per contact pair.

Finally, the correlation maps were normalized using the Fisher z-transform. The same

procedure was applied to spindle-free control segments. Off-diagonal correlations

indicate a time-lag between spindle power in scalp and thalamic channels.

Phase Slope Index: We assessed whether SOs drive activity in the sleep spindle range or

vice versa in NREM sleep data segments comprising SO-spindle complexes, separately

for thalamic and scalp recordings. The cross-frequency phase-slope index [34,] was

calculated for each contact between the signal and the signal filtered in the sleep spindle

range (11 – 16 Hz). After applying a Hanning window and extracting the complex Fourier

coefficients, all SO frequencies < 2 Hz were considered. In this context, positive values
indicate SOs driving sleep spindle activity, while negative values indicate sleep spindles
driving SOs. The obtained data distributions were tested against zero, using paired
samples t-tests.

Time frequency representations: Time–frequency analyses of SO, sleep spindle and SO-
spindle segments were performed using FieldTrip [78]. Frequency decomposition of the data
was achieved using Fourier analysis based on sliding time windows (moving forward in 50
ms increments). The window length was set to five cycles of a given frequency (frequency
range: 1–30 Hz in 1 Hz steps). The windowed data segments were multiplied with a
Hanning taper before Fourier analysis [−2.5 to 2.5 sec]. The longer time segments were
chosen to allow for resolving low frequency activity within the time windows of interest
[−1.5 to 1.5 sec] and avoid edge artifacts. Resulting power values were z-scored across
time. The same procedure was applied to event-free control segments.

Statistics: Statistical analyses were performed at the individual electrodes / contacts level
(fixed-effects analysis), considering all electrodes / contacts that were eligible based on
our criteria (e.g., artefact free, etc.). Unless stated otherwise, we used cluster-based
permutation tests to correct for multiple comparisons as implemented in fieldtrip. A
dependent-samples t-test was used at the sample level and values were thresholded at p
= 0.05 (1000 randomizations). The sum of all t-values in cluster served as cluster statistic
and Monte Carlo simulations were used to calculate the cluster p-value (alpha = 0.05,
two-tailed) under the permutation distribution. The input data were either occurrence
probabilities across time (e.g., Fig. 2a), time x frequency values (e.g., Fig. 2c) or time x
time correlation maps (e.g., Fig. 4b) which were tested against corresponding data
stemming from event-free events. In case of between area comparisons (e.g., Fig. 3g)
independent samples t-tests (cluster corrected) were employed to conform with the
varying number of observation units (i.e., contacts).
For circular statistics the phase distributions (within or across participants) were tested
against uniformity using the Rayleigh test (CircStat toolbox [79]). When directly comparing
phase distributions across recording sites, the circular Watson-Williams test was used [79].

References:

1.   Klinzing, J. G., Niethard, N. & Born, J. Mechanisms of systems memory
consolidation during sleep. *Nature Neuroscience* (2019). doi:10.1038/s41593-
019-0467-3
2.   Amzica, F. & Steriade, M. The functional significance of K-complexes. *Sleep Med.*
Rev. **6**, 139–149 (2002).
3.   Steriade, M., McCormick, D. A. & Sejnowski, T. J. Thalamocortical oscillations in
the sleeping and aroused brain. *Science (80-.).* (1993).
doi:10.1126/science.8235588
4.   Timofeev, I., Grenier, F., Bazhenov, M., Sejnowski, T. J. & Steriade, M. Origin of
slow cortical oscillations in deafferented cortical slabs. *Cereb. Cortex* (2000).
doi:10.1093/cercor/10.12.1185
5.   Hughes, S. W., Cope, D. W., Blethyn, K. L. & Crunelli, V. Cellular Mechanisms of
the Slow (. *Neuron* (2002).
6.   David, F. *et al.* Essential thalamic contribution to slow waves of natural sleep. *J.*
Neurosci. (2013). doi:10.1523/JNEUROSCI.3169-13.2013
7.   Gent, T. C., Bandarabadi, M., Herrera, C. G. & Adamantidis, A. R. Thalamic dual
control of sleep and wakefulness. *Nat. Neurosci.* (2018). doi:10.1038/s41593-
018-0164-7
8.   Crunelli, V. & Hughes, S. W. The slow (<1 Hz) rhythm of non-REM sleep: a
dialogue between three cardinal oscillators. *Nat. Neurosci.* **13**, 9 (2010).
9.   Crunelli, V., David, F., Lorincz, M. L. & Hughes, S. W. The thalamocortical
network as a single slow wave-generating unit. *Curr. Opin. Neurobiol.* **31**, 72–80
(2015).
10. Neske, G. T. The Slow Oscillation in Cortical and Thalamic Networks:
Mechanisms and Functions. *Front. Neural Circuits* **9**, (2016).
11. Steriade, M. Grouping of brain rhythms in corticothalamic systems. *Neuroscience*
137, 1087–1106 (2006).
12. Helfrich, R. F., Mander, B. A., Jagust, W. J., Knight, R. T. & Walker, M. P. Old
Brains Come Uncoupled in Sleep: Slow Wave-Spindle Synchrony, Brain Atrophy,
and Forgetting. *Neuron* **97**, 221-230.e4 (2018).
13. Mak-McCully, R. A. *et al.* Coordination of cortical and thalamic activity during
non-REM sleep in humans. *Nat. Commun.* **8**, 15499 (2017).
14. Mölle, M., Marshall, L., Gais, S. & Born, J. Grouping of spindle activity during
slow oscillations in human non-rapid eye movement sleep. *J. Neurosci.* **22**,
10941–7 (2002).
15. Sirota, A., Csicsvari, J., Buhl, D. & Buzsáki, G. Communication between
neocortex and hippocampus during sleep in rodents. *Proc. Natl. Acad. Sci. U. S.*
A. **100**, 2065–2069 (2003).
16. Staresina, B. P. *et al.* Hierarchical nesting of slow oscillations, spindles and
ripples in the human hippocampus during sleep. *Nat. Neurosci.* **18**, 1679–1686
(2015).
17. Schreiner, T. & Staudigl, T. Electrophysiological signatures of memory
reactivation in humans. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **375**, (2020).
18. Girardeau, G. & Lopes-dos-Santos, V. Brain neural patterns and the memory
function of sleep. *Science (80-.).* **374**, 560–564 (2021).
19. Pergola, G. *et al.* The Regulatory Role of the Human Mediodorsal Thalamus.
Trends Cogn. Sci. **22**, 1011–1025 (2018).
20. Aggleton, J. P., Pralus, A., Nelson, A. J. D. & Hornberger, M. Thalamic pathology
and memory loss in early Alzheimer's disease: moving the focus from the medial
temporal lobe to Papez circuit. *Brain* **139**, 1877–1890 (2016).

21.   Aggleton, J. P. & O'Mara, S. M. The anterior thalamic nuclei: core components of
a tripartite episodic memory system. *Nat. Rev. Neurosci.* (2022).
doi:10.1038/S41583-022-00591-8
22.   Aggleton, J. P. & Brown, M. W. Episodic memory, amnesia, and the
hippocampal–anterior thalamic axis. *Behav. Brain Sci.* **22**, 425–444 (1999).
23.   Fernandez, L. M. J. & Lüthi, A. Sleep Spindles: Mechanisms and Functions.
Physiological reviews **100**, 805–868 (2020).
24.   Horn, A. & Kühn, A. A. Lead-DBS: a toolbox for deep brain stimulation electrode
localizations and visualizations. *Neuroimage* **107**, 127–135 (2015).
25.   Ewert, S. *et al.* Toward defining deep brain stimulation targets in MNI space: A
subcortical atlas based on multimodal MRI, histology and structural connectivity.
Neuroimage **170**, 271–282 (2018).
26.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. *Source J. R. Stat. Soc. Ser. B* **57**,
289–300 (1995).
27.   Schreiner, T., Petzka, M., Staudigl, T. & Staresina, B. P. Endogenous memory
reactivation during sleep in humans is clocked by slow oscillation-spindle
complexes. *Nat. Commun.* (2021). doi:10.1038/s41467-021-23520-2
28.   Jiang, H., Bahramisharif, A., van Gerven, M. A. J. & Jensen, O. Measuring
directionality between neuronal oscillations of different frequencies. *Neuroimage*
118, 359–367 (2015).
29.   Adamantidis, A. R., Gutierrez Herrera, C. & Gent, T. C. Oscillating circuitries in
the sleeping brain. *Nat. Rev. Neurosci.* **20**, 746–762 (2019).
30.   Blethyn, K. L., Hughes, S. W., Tóth, T. I., Cope, D. W. & Crunelli, V. Neuronal
Basis of the Slow (<1 Hz) Oscillation in Neurons of the Nucleus Reticularis
Thalami In Vitro. *J. Neurosci.* **26**, 2474–2486 (2006).
31.   Slézia, A., Hangya, B., Ulbert, I. & Acsády, L. Phase Advancement and Nucleus-
Specific Timing of Thalamocortical Activity during Slow Cortical Oscillation. *J.
Neurosci.* **31**, 607–617 (2011).
32.   Poulet, J. F. A., Fernandez, L. M. J., Crochet, S. & Petersen, C. C. H. Thalamic
control of cortical states. *Nat. Neurosci. 2012 153* **15**, 370–372 (2012).
33.   Hay, Y. A. *et al.* Thalamus mediates neocortical Down state transition via GABA
B-receptor-targeting interneurons. *Neuron* **109**, 2682-2690.e5 (2021).
34.   Pare, D., Steriade, M., Deschenes, M. & Oakson, G. Physiological characteristics
of anterior thalamic nuclei, a group devoid of inputs from reticular thalamic
nucleus. *https://doi.org/10.1152/jn.1987.57.6.1669* **57**, 1669–1685 (1987).
35.   Gonzalo-Ruiz, A. & Lieberman, A. R. Topographic organization of projections
from the thalamic reticular nucleus to the anterior thalamic nuclei in the rat. *Brain
Res. Bull.* **37**, 17–35 (1995).
36.   Halassa, M. M. & Sherman, S. M. Thalamocortical Circuit Motifs: A General
Framework. *Neuron* **103**, 762–770 (2019).
37.   Lachaux, J. P., Rudrauf, D. & Kahane, P. Intracranial EEG and human brain
mapping. in *Journal of Physiology Paris* (2003).
doi:10.1016/j.jphysparis.2004.01.018
38.   Marmor, O. *et al.* Local vs. Volume conductance activity of field potentials in the
human subthalamic nucleus. *J. Neurophysiol.* (2017).
doi:10.1152/jn.00756.2016
39.   Wennberg, R. & Lozano, A. M. Restating the importance of bipolar recording in
subcortical nuclei [1]. *Clinical Neurophysiology* (2006).
doi:10.1016/j.clinph.2005.09.020
40.   Liguori, C., Toledo, M. & Kothare, S. Effects of anti-seizure medications on sleep
architecture and daytime sleepiness in patients with epilepsy: A literature review.

Sleep Medicine Reviews (2021). doi:10.1016/j.smrv.2021.101559

41. Frauscher, B., von Ellenrieder, N., Dubeau, F. & Gotman, J. Scalp spindles are
associated with widespread intracranial activity with unexpectedly low synchrony.
Neuroimage (2015). doi:10.1016/j.neuroimage.2014.10.048

42. Frauscher, B. *et al.* Facilitation of epileptic activity during sleep is mediated by
high amplitude slow waves. *Brain* (2015). doi:10.1093/brain/awv073

43. Steriade, M. Sleep, epilepsy and thalamic reticular inhibitory neurons. *Trends in*
Neurosciences (2005). doi:10.1016/j.tins.2005.03.007

44. Simon, K. C. *et al.* Slow oscillation density and amplitude decrease across
development in pediatric Duchenne and Becker muscular dystrophy. *Sleep*
(2021). doi:10.1093/sleep/zsaa240

45. Gais, S., Mölle, M., Helms, K. & Born, J. Learning-dependent increases in sleep
spindle density. *J. Neurosci.* **22**, 6830–4 (2002).

46. Jankowski, M. M. *et al.* The anterior thalamus provides a subcortical circuit
supporting memory and spatial navigation. *Front. Syst. Neurosci.* **7**, (2013).

47. Grodd, W., Kumar, V. J., Schüz, A., Lindig, T. & Scheffler, K. The anterior and
medial thalamic nuclei and the human limbic system: tracing the structural
connectivity using diffusion-weighted imaging. *Sci. Reports 2020 101* **10**, 1–25
(2020).

48. Van Groen, T. & Wyss, J. M. Projections from the anterodorsal and anteroveniral
nucleus of the thalamus to the limbic cortex in the rat. *J. Comp. Neurol.* **358**,
584–604 (1995).

49. Massimini, M., Huber, R., Ferrarelli, F., Hill, S. & Tononi, G. The sleep slow
oscillation as a traveling wave. *J. Neurosci.* **24**, 6862–70 (2004).

50. Riedner, B. A., Hulse, B. K., Murphy, M. J., Ferrarelli, F. & Tononi, G. Temporal
dynamics of cortical sources underlying spontaneous and peripherally evoked
slow waves. *Prog. Brain Res.* **193**, 201–218 (2011).

51. Steriade, M. & Deschenes, M. The thalamus as a neuronal oscillator. *Brain Res.*
320, 1–63 (1984).

52. Gentet, L. J. & Ulrich, D. Strong, reliable and precise synaptic connections
between thalamic relay cells and neurones of the nucleus reticularis in juvenile
rats. *J. Physiol.* **546**, 801–811 (2003).

53. Velayos, J. L., Oliva, M. & Alfageme, F. Afferent projections to the mediodorsal
and anterior thalamic nuclei in the cat. Anatomical-clinical correlations. *Brain*
Pathol. **8**, 549–552 (1998).

54. Lozsádi, D. A. Organization of connections between the thalamic reticular and
the anterior thalamic nuclei in the rat. *J. Comp. Neurol.* **358**, 233–246 (1995).

55. Latchoumane, C.-F. V., Ngo, H.-V. V., Born, J. & Shin, H.-S. Thalamic Spindles
Promote Memory Formation during Sleep through Triple Phase-Locking of
Cortical, Thalamic, and Hippocampal Rhythms. *Neuron* 1–12 (2017).
doi:10.1016/j.neuron.2017.06.025

56. Viejo, G. & Peyrache, A. Precise coupling of the thalamic head-direction system
to hippocampal ripples. *Nat. Commun. 2020 111* **11**, 1–14 (2020).

57. Yang, M., Logothetis, N. K. & Eschenko, O. Occurrence of Hippocampal Ripples
is Associated with Activity Suppression in the Mediodorsal Thalamic Nucleus. *J.*
Neurosci. (2019). doi:10.1523/jneurosci.2107-18.2018

58. Ferrarelli, F. & Tononi, G. Reduced sleep spindle activity point to a TRN-MD
thalamus-PFC circuit dysfunction in schizophrenia. *Schizophr. Res.* **180**, 36–43
(2017).

59. Bastuji, H., Lamouroux, P., Villalba, M., Magnin, M. & Garcia-Larrea, L. Local
sleep spindles in the human thalamus. *J. Physiol.* **598**, 2109–2124 (2020).

60. Guillery, R. W., Feig, S. L. & Lozsádi, D. A. Paying attention to the thalamic

reticular nucleus. *Trends Neurosci.* (1998). doi:10.1016/S0166-2236(97)01157-
0

61. Kultas-Ilinsky, K., Yi, H. & Ilinsky, I. A. Nucleus reticularis thalami input to the
anterior thalamic nuclei in the monkey: a light and electron microscopic study.
Neurosci. Lett. (1995). doi:10.1016/0304-3940(95)11273-Y

62. Ilinsky, I. A., Ambardekar, A. V. & Kultas-Ilinsky, K. Organization of projections
from the anterior pole of the nucleus reticularis thalami (NRT) to subdivisions of
the motor thalamus: Light and electron microscopic studies in the rhesus
monkey. *J. Comp. Neurol.* (1999). doi:10.1002/(SICI)1096-
9861(19990705)409:3<369::AID-CNE3>3.0.CO;2-H

63. Tai, Y., Yi, H., Ilinsky, I. A. & Kultas-Ilinsky, K. Nucleus reticularis thalami
connections with the mediodorsal thalamic nucleus: A light and electron
microscopic study in the monkey. *Brain Res. Bull.* (1995). doi:10.1016/0361-
9230(95)02018-M

64. Niethard, N., Ngo, H. V. V., Ehrlich, I. & Born, J. Cortical circuit activity underlying
sleep slow oscillations and spindles. *Proc. Natl. Acad. Sci. U. S. A.* (2018).
doi:10.1073/pnas.1805517115

65. Taube, J. S. Head direction cells recorded in the anterior thalamic nuclei of freely
moving rats. *J. Neurosci.* **15**, 70–86 (1995).

66. Peyrache, A., Lacroix, M. M., Petersen, P. C. & Buzsáki, G. Internally organized
mechanisms of the head direction sense. *Nat. Neurosci.* **18**, 569–575 (2015).

67. Leszczyński, M. & Staudigl, T. Memory-guided attention in the anterior thalamus.
Neurosci. Biobehav. Rev. **66**, 163–165 (2016).

68. Maingret, N., Girardeau, G., Todorova, R., Goutierre, M. & Zugaro, M.
Hippocampo-cortical coupling mediates memory consolidation during sleep. *Nat.*
Neurosci. **19**, 959–964 (2016).

69. Sarasso, S. *et al.* Hippocampal sleep spindles preceding neocortical sleep onset
in humans. *Neuroimage* **86**, 425–432 (2014).

70. Kim, J., Gulati, T. & Ganguly, K. Competing Roles of Slow Oscillations and Delta
Waves in Memory Consolidation versus Forgetting. *Cell* **179**, 514-526.e13
(2019).

71. Kaufmann, E., Bötzel, K., Vollmar, C., Mehrkens, J. H. & Noachtar, S. What have
we learned from 8 years of deep brain stimulation of the anterior thalamic
nucleus? Experiences and insights of a single center. *J. Neurosurg.* **135**, 619–
628 (2020).

72. Kaufmann, E. *et al.* European Expert Opinion on ANT-DBS therapy for patients
with drug-resistant epilepsy (a Delphi consensus). *Seizure* (2020).
doi:10.1016/j.seizure.2020.08.015

73. Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric
diffeomorphic image registration with cross-correlation: evaluating automated
labeling of elderly and neurodegenerative brain. *Med. Image Anal.* **12**, 26–41
(2008).

74. Ashburner, J. & Friston, K. J. Unified segmentation. *Neuroimage* **26**, 839–851
(2005).

75. Wennberg, R. A. & Lozano, A. M. Intracranial volume conduction of cortical
spikes and sleep potentials recorded with deep brain stimulating electrodes. *Clin.*
Neurophysiol. **114**, 1403–1418 (2003).

76. Iber, C., Ancoli-Israel, S., Chesson, A. & Quan, S. *The AASM Manual for the*
Scoring of Sleep and Associated Events: Rules, Terminology, and Technical
Specification. American Academy of Sleep Medicine (2007).

77. Helfrich, R. F. *et al.* Bidirectional prefrontal-hippocampal dynamics organize
information transfer during sleep in humans. *Nat. Commun.* (2019).

doi:10.1038/s41467-019-11444-x

78.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: Open source
software for advanced analysis of MEG, EEG, and invasive electrophysiological
data. *Comput. Intell. Neurosci.* **2011**, 156869 (2011).

79.   Berens, P. CircStat: A MATLAB toolbox for circular statistics. *J. Stat. Softw.* **31**,
1–21 (2009).

1011

1012   Acknowledgments

1013   This work was supported by the European Research Council (https://erc.europa.eu/,
1014   Starting Grant 802681 awarded to TSt). We are indebted to all patients who volunteered
1015   to participate in this study. We thank the staff and physicians of the at the Epilepsy Center,
1016   Department of Neurology, Ludwig Maximilians University, Munich for assistance.

1017

1018   Author Contributions

1019   Conceptualization: TS, TSt; Resources: EK, NS, JHM; Formal Analysis: TS, TSt; Funding
1020   acquisition: TSt; Writing – original draft: TS, TSt; Writing– review & editing: TS, TSs, EK,
1021   SN

1022

1023   Corresponding author

1024   Correspondence to Tobias Staudigl

1025

1026   Competing interests

1027   The authors declare no competing interests.

Supplementary Information for:

The human thalamus orchestrates neocortical oscillations during NREM sleep.

Thomas Schreiner[1], Elisabeth Kaufmann[2], Soheyl Noachtar[2], Jan-Hinnerk Mehrkens[3] & Tobias Staudigl[1]

1 Department of Psychology, Ludwig-Maximilians-Universität München, Munich, Germany

2 Epilepsy Center, Department of Neurology, Ludwig-Maximilians-Universität München, Munich, Germany

3 Department of Neurosurgery, Ludwig-Maximilians-Universität München, Munich, Germany

Supplementary tables

Supplementary Table 1 | Sleep architecture: Data are means ± s.e.m. N1, N2: NREM sleep stages N1 & N2, SWS: slow-wave sleep, REM: rapid eye movement sleep, WASO: wake after sleep onset. TST: total sleep time (in minutes).

	N1	N2	SWS	REM	WASO	TST [min]
Sleep stage [%]	6.1 ± 1.7	47.4 ± 4.2	16.5 ± 3.5	12.7 ± 2.2	15.9 ± 3.5	529.4 ± 30.1

Supplementary Table 2 | scalp electrodes: overview of scalp electrode that were selected for the respective analyses.

	ANT-SO	ANT-spindle	MD-SO	MD-spindle
P1	Fz	Fp1	Fz	P4
P2	F4	F4	FP2	Pz
P3	F8	F4	F4	F4
P4	FT10	Pz	FT10	C3
P5	F4	Pz	F3	P3
P6	Fz	C3	Fz	Pz
P7	FT10	C4	Fz	Pz
P8	-	-	F7	Cz

Supplementary Table 3 | Sleep Oscillations ANT: Data are means ± s.e.m. Number of detected events and percentage of paired events (i.e., percentage of co-occurring events across sites, relative to the overall number of events at given site) in NREM sleep at ANT contacts and scalp electrodes. Density (events/min of NREM sleep)

	ANT	Scalp	t	p
SO number	1573.8 ± 129.9	1513.5 ± 130.1	1.11	0.29
SO density	6.29 ± 0.21	5.81 ± 0.24	1.32	0.21
SO-coupling [%]	33.74 ± 3.02	31.11 ± 3.25	2.07	0.06
spindle number	2203.8 ± 213.5	1903.3 ± 211.5	3.19	0.008
spindle density	8.83 ± 0.21	6.09 ± 1.08	2.56	0.017
spindle-coupling [%]	43.8 ± 4.2	37.3 ± 4.1	3.43	0.005
SO-spindle number	431 ± 52.1	304.1 ± 69.8	3.18	0.008

1087 Supplementary Table 4 | Sleep Oscillations MD: Data are means ± s.e.m. Number of detected events
1088 and percentage of paired events (i.e., percentage of co-occurring events across sites, relative to the
1089 overall number of events at given site) in NREM sleep at MD contacts and scalp electrodes. Density
1090 (events/min of NREM sleep).

	MD	Scalp	t	p
SO number	1408.5 ± 126.2	1476.9 ± 129.1	1.06	0.31
SO density	6.21 ± 0.22	6.40 ± 0.25	-0.5	0.61
SO-coupling [%]	30.45 ± 3.02	29.51 ± 3.15	1.37	0.35
spindle number	2127.8 ± 238.2	1796.5 ± 299.8	3.35	0.005
spindle density	9.81 ± 0.98	6.14 ± 1.04	2.43	0.024
spindle-coupling [%]	45.1 ± 5.5	30.2 ± 4.6	2.18	0.046
SO-spindle number	488 ± 53.3	338.5 ± 79.7	1.78	0.095

1091
1092
1093 Supplementary Table 5 | SO features: Data are means ± s.e.m. Duration and relative occurrence during
1094 N2 and N3 sleep for scalp (ANT and MD analyses related), ANT and MD derived SOs.

SO	Duration [sec]	Slope [μV/sec]	Rel. N2 [%]	Rel. N3 [%]
scalp ANT	1.31 ± 0.01	67.55 ± 6.63	46.56 ± 3.79	53.43 ± 3.79
scalp MD	1.35 ± 0.02	80.69 ±12.93	49.21 ± 4.38	50.78 ± 4.38
ANT	1.30 ± 0.01	75.99 ± 9.61	46.05 ± 3.87	53.94 ± 3.87
MD	1.30 ± 0.02	74.82 ± 7.58	47.70 ± 4.89	52.29± 4.89

1095
1096
1097 Supplementary Table 6 | spindle features: Data are means ± s.e.m. Peak frequency, duration and relative
1098 occurrence during N2 and N3 sleep for scalp (ANT and MD analyses related), ANT and MD derived
1099 spindles

spindles	Peak-Freq [Hz]	Duration [ms]	Rel. N2 [%]	Rel. N3 [%]
scalp ANT	13.02 ± 0.15	0.81 ± 0.03	73.71 ± 3.81	26.29 ± 3.81
scalp MD	13.07 ± 0.13	0.77 ± 0.05	70.68 ± 5.14	29.32 ± 5.14
ANT	13.17 ± 0.09	0.79 ± 0.04	73.51 ± 3.76	26.49 ± 3.76
MD	13.03 ± 0.11	0.78 ± 0.03	74.02 ± 3.49	25.97 ± 3.49

1100
1101
1102 Supplementary Table 7 | drug regimen at the time of recordings.

	Anticonvulsant
P1	Levetiracetam, Phenytoin, Lamotrigine
P2	Levetiracetam, Lamotrigine
P3	Lacosamide, Oxcarbazepine, Topiramate
P4	Levetiracetam, Lamotrigine, Oxcarbazepin, Zonisamide
P5	Levetiracetam, Lamotrigine
P6	Lamotrigine, Lacosamide, Zonisamide
P7	Oxcarbazepin, Topiramate
P8	Topiramate

1108 Supplementary Table 8 | Documented effects drugs on sleep architecture.

Drug	Effect on sleep
Lacosamide[1]	none
Lamotrigine[2]	REM↑, N2↑, SWS↓
Levetiracetam[3]	N2↑, REM↓
Oxcarbazepine	unknown
Phenytoin[4]	SWS↓, REM↓
Topiramate[5]	none
Zonisamide[6]	none

1) Hudson J.D., Guptil J.T., Bynes W., et al. (2015). Assessment of the effects of lacosamide on sleep parameters in
healthy subjects. Seizure; 25: 155-159 2) Placidi F, Marciani MG, Diomedi M, et al (2000). Effects of lamotrigine on
nocturnal sleep, daytime somnolence and cognitive functions in focal epilepsy. Acta Neurol Scand;102:81–86. 3)
Cicolin A, Magliola U, Giordano A, et al. (2006). Effects of levetiracetam on nocturnal sleep and daytime vigilance in
healthy volunteers. Epilepsia; 47:82–85. 4) Benjamin Legros & Carl W Bazil (2003). Effects of antiepileptic drugs on
sleep architecture: a pilot study. Sleep Med; 4(1):51-5. 5) Bonanni E, Galli R, Maestri M, et al. (2004). Daytime
sleepiness in epilepsy patients receiving topiramate monotherapy. Epilepsia; 45:333–337. 6) Romigi A, Izzi F, Placidi
F, et al (2013). Effects of zonisamide as add-on therapy on sleep-wake cycle in focal epilepsy: a polysomnographic
study. Epilepsy Behav; 26:170–174.

1118
1119

1120 Supplementary Table 9 | Descriptives Slow and Fast spindles: Data are means ± s.e.m. Number of
1121 detected events and percentage of paired events between thalamic contacts and scalp electrodes.
1122 Density (events/min).

1123

ANT spindle	slow	fast	t	P
spindle number	1850.5 ± 132.8	1829.3 ± 155.1	0.13	0.85
spindle density	7.47 ± 0.34	7.55 ± 0.58	-0.1	0.87
spindle-coupling [%]	23.5 ± 3.1	31.7 ± 3.9	-1.8	0.08
Peak Freq. Hz	10.29 ± 0.15	14.02 ± 0.11		
MD spindle				
spindle number	1990.6 ± 132.8	1945.1 ± 222.9	0.32	0.70
spindle density	8.56 ± 0.97	8.62 ± 0.57	-0.1	0.92
spindle-coupling [%]	27.0 ± 2.7	26.4 ± 5.2	0.011	0.90
Peak Freq. Hz	10.20 ± 0.12	14.08 ± 0.12		

1124
1125

Supplementary figures

Supplementary Fig. 1 | Local and non-local ANT spindles. Non-local ANT spindles were defined as spindles identified in the ANT whose peak (maximally negative amplitude) overlapped with MD spindle peaks ± 50 ms (267.91 ± 51.57 ANT spindles out of 2203.8 ± 213.5; 11.78 ± 1.84%). All other spindles identified in the ANT were defined as local ANT spindles. The same temporal dynamics with regards to thalamo-cortical spindle coupling and SO-spindle coupling were found for local and non-local ANT spindles. **(a)** Occurrence probabilities of local ANT spindle peaks relative to neocortical spindle peaks (maximal negative amplitude, time = 0; dashed line; bin size = 50 ms), indicating that local ANT spindles precede neocortical spindles (positive clusters from -0.3 to -0.05 sec, $p < 0.0001$ & 0.2 to 0.25 sec, $p = 0.023$; time of peak: -0.1 sec). **(b)** Local ANT spindles preferential emerged around ANT SO down states (-0.15 to 0.25 sec; $p = 0.009$; peak: 0 sec). **(c)** Phases of the SO-spindle modulation derived from local ANT events. Local ANT spindles started specifically around the ANT SO down states (mean coupling direction: −168.93° ± 10.10°; Rayleigh $z = 7.93$, $p < 0.0001$). **(d)** Occurrence probabilities of non-local ANT spindle peaks relative to neocortical spindle peaks, indicating that non-local ANT spindles likewise precede neocortical spindles (positive cluster from -0.15 to 0 sec, $p < 0.017$; time of peak: -0.05 sec). **(e)** Non-local ANT spindles also preferential emerged around ANT SO down states (-0.05 to 0.1 sec; $p = 0.017$; peak: -0.05 sec). **(c)** Phases of the SO-spindle modulation derived from non-local ANT events. Non-local ANT spindles started specifically around the ANT SO down states (mean coupling direction: −157.91° ± 10.21°; Rayleigh $z = 7.86$, $p < 0.0001$).

1150
Supplementary Fig. 2 | Local and non-local MD spindles. Non-local MD spindles were defined as
spindles identified in the MD whose peak (maximally negative amplitude) overlapped with ANT spindle
peaks ± 50 ms (149.83 ± 35.69 MD spindles out of 2127.8 ± 238.2; 7.04%). All other spindles identified
in the ANT were defined as local ANT spindles. The same temporal dynamics with regards to thalamo-
cortical spindle coupling and SO-spindle coupling were found for local and non-local MD spindles. (a)
Occurrence probabilities of local MD spindle peaks relative to neocortical spindle peaks (maximal
negative amplitude, time = 0; dashed line; bin size = 50 ms), indicating that local MD spindles precede
neocortical spindles (positive cluster from -0.25 to 0 sec, p < 0.0001; time of peak: -0.05 sec). (b) Local
ANT spindles preferential emerged around ANT SO down states (-0.25 to 0.05 sec; p < 0.0001; peak: -
0.05 sec). (c) Phases of the SO-spindle modulation derived from local ANT events. Local ANT spindles
started specifically around the ANT SO down states (mean coupling direction: −172.05° ± 7.60°; Rayleigh
z = 9.59, p < 0.0001). (d) Occurrence probabilities of non-local ANT spindle peaks relative to neocortical
spindle peaks, indicating that non-local ANT spindles likewise precede neocortical spindles (positive
cluster from -0.1 to 0 sec, p < 0.0001; time of peak: -0.05 sec). (e) Non-local ANT spindles also preferential
emerged around ANT SO down states (-0.2 to 0 sec; p = 0.017; peak: -0.3 sec). (c) Phases of the SO-
spindle modulation derived from non-local ANT events. Non-local ANT spindles started specifically
around the ANT SO down states (mean coupling direction: −155.38° ± 8.68°; Rayleigh z = 8.91, p <
0.0001).
1169
1170
1171

1172
Supplementary Fig. 3 | SO coupling frontal iEEG – frontal scalp: To examine whether the skull, acting as
a low pass filter, might have delayed the picked-up signal at the scalp level relative to thalamic
recordings, an additional, independent data-set was analyzed. Specifically, SO coupling between frontal
scalp electrodes (electrode FP1) and frontal intracranial contacts in three pre-surgical epilepsy patients
(frontal iEEG contacts, N = 13) during NREM sleep was assessed. If the skull indeed delays the measured
SO negative peak amplitude at the scalp level then a comparable delay of ~50 ms with regards to SOs
should be observable in this data, were intracranial and scalp electrodes are in close distance. On average
we detected 1450 ± 137.58 SOs in frontal intracranial contacts and 1245.7 ± 161.97 SOs in frontal scalp
EEG recordings during NREM sleep, with a coupling rate of 30.15 ± 1.22 %. (a) Occurrence probabilities
of intracranial detected SO down state peaks relative to scalp detected down state peaks (bin size = 50
ms), indicated that SOs in frontal intracranial and scalp recordings emerged synchronously (positive
cluster from -0.1 to 0.1 sec; p < 0.0009; time of peak = 0 sec). (b) We then determined the phase of
intracranial SOs for all paired SO-events (i.e., all ANT SOs within ± 750 ms of neocortical SOs) at the time
of neocortical SO down state. We found a significant nonuniform distribution across contacts (Rayleigh
z = 12.82, p < 0.0001), with the phase of intracranial SOs being almost identical to the phase of scalp
detected events (the phase of neocortical SO down states corresponds to ± π; mean coupling direction:
−179.37 ± 1.84°). (c) Example of NREM sleep segment (15 sec), comprising SOs (top row: scalp recording;
bottom row: intracranial). (d) Grand average EEG trace of neocortical SOs (mean ± SEM, negative peak,
time 0). (e) Grand average iEEG trace of intracranial SOs (mean ± SEM, negative peak, time 0).

1200
Supplementary Fig. 4| Estimating the potential influence of volume conducted SOs on the timing of
thalamo-cortical SO coupling: While the bipolar montage represents the gold standard in order to
resolve localized activity in intracranial recordings [37–39], volume conducted SOs might still intrude the
measured signal. To accurately estimate the possible impact of volume conduction on our SO related
results we ran additional control analyses on the data of a sample patient. First, we identified events
where SOs are present in scalp EEG recordings (average referenced including scalp EEG electrodes only,
as in the main analyses) but not in the bipolar referenced ANT recordings (for an example event see (a);
for ERPs of all identified events (scalp in grey, ANT in blue) see (c)). The same events were then isolated
using a common average reference that included both scalp EEG and ANT recordings. The rationale of
this procedure is to isolate SOs in the ANT that were exclusively a product of volume conduction and by
that allowing to estimate potential technical delays in SO peaks between EEG and bipolar iEEG
recordings. In sum, no such delay became apparent beyond 180 degree phase shift due to the polarity
reversal of the ANT signal in the common average montage. (b) Shows the same example event as in (a)
using a common average reference (including both scalp and intracranial recordings). (d) Illustrates all
identified events used in (c) again using a common average reference (including both scalp (grey) and
intracranial recordings (blue)), highlighting that SOs in scalp recordings and volume conducted SOs in
the ANT, exhibiting a reversed polarity, peaked at the same time. The insets show in blue the phase of
volume conducted SOs in ANT contacts at the time of neocortical SO downstate, illustrating that volume
conducted SOs peaked at almost exactly the opposite phase as neocortical SOs (Rayleigh z = 282.71, p <
0.00001; mean coupling direction: 1.59°). Cortical SO down-states using a common average reference
(including iEEG as in b and d) peaked at 179.45° (Rayleigh z = 282.71; p < 0.00001).

1231
Supplementary Fig. 5 | ANT-scalp dynamics with regards to contact pairs were both contacts (n = 8; full
ANT) or just one contact of a bipolar pair (n = 4, single ANT) were localized to the ANT: (a) Occurrence
probabilities of full ANT SO down state peaks relative to scalp derived SO down state peaks (bin size = 50
ms), indicated that full ANT SOs preceded scalp derived SOs (positive cluster from -0.15 to 0 sec; p <
0.014; time of peak = -0.05 sec). (b) Occurrence probabilities of single ANT SO down state peaks relative
to scalp derived SO down state peaks (bin size = 50 ms). While single ANT SOs preceded descriptively
scalp derived SOs, no significant differences were observable when tested against event-free occurrence
probabilities (p > 0.1). (c) No significant difference became apparent when directly comparing full ANT-
scalp occurrence probabilities (green) and single ANT-scalp occurrence probabilities (orange)(d)
Occurrence probabilities of full ANT spindle peaks relative to neocortical spindle peaks (maximal negative
amplitude, time = 0; dashed line; bin size = 50 ms), indicating that full ANT spindles precede neocortical
spindles (positive clusters from -0.35 to -0.2 sec, p < 0.001 & -0.1 to 0.05 sec, p < 0.001; time of peak: -
0.05 sec). (e) Again, single ANT sleep spindles, exhibited descriptively the same dynamics as full ANT
spindle, but did not reach significance when tested against event-free occurrence probabilities (p > 0.1).
(f) No significant difference became apparent when directly comparing spindle related full ANT-scalp
occurrence probabilities (green) and single ANT-scalp occurrence probabilities (orange). (g) Occurrence
probabilities of full ANT spindle onsets with respect to full ANT SO down states (bin size = 50 ms),
illustrating that spindles preferentially emerged around ANT SO down states (-0.05 to 0 sec; p = 0.048).
(h) The same SO-spindle related dynamic became apparent with regards to single ANT contact pairs, but
again did not reach significance when tested against event-free occurrence probabilities (p > 0.1). (i)
Direct comparison of full and single ANT SO-spindle dynamics (green and orange, respectively) did not
yield any significant difference (p > 0.1).

1254
1255

1256
Supplementary Fig. 6 | Lateralization of coupling: (a) Occurrence probabilities of left ANT SO down state
peaks relative to right ANT SO down state peaks (bin size = 50 ms), indicated that SOs in the left and right
ANT emerged synchronously (positive cluster from -0.1 to 0.1 sec; p < 0.0001; time of peak = 0 sec). (b)
Occurrence probabilities for sleep spindles derived from the left ANT, relative to right ANT detected
spindles, indicated that SOs in the left and right ANT emerged synchronously (positive cluster from -0.25
to 0.25 sec; p < 0.0001; time of peak = 0 sec). (c) Occurrence probabilities of left MD SO down state
peaks relative to right MD SO down state peaks, (positive cluster from -0.15 to 0.05 sec; p < 0.0001; time
of peak = 0 sec). (d) Occurrence probabilities for sleep spindles derived from the left MD, relative to right
MD detected spindles (positive cluster from -0.2 to 0.1 sec; p < 0.0001; time of peak = 0 sec).
1266
1267

1268
Supplementary Fig. 7 | SO coupling ANT-Fz: Occurrence probabilities of ANT SO down state peaks
relative to SO down state peaks derived from scalp electrode Fz (time = 0; dashed line; bin size = 50 ms)),
supporting the main finding that ANT SOs precede neocortical SOs. The solid black line indicates
significant differences, resulting from comparison with SO-free control events (positive cluster from -
0.15 to 0.05 sec, p < 0.005; time of peak: -0.05 sec). Note that Fz was not used as generic seed electrode
in the main analysis (Fig. 1a), as it was contaminated with epileptic activity in 3/8 patients.
1275
1276
1277

1278
Supplementary Fig. 8 | SO coupling ANT-Occipital: To assess the specificity of ANT – frontal interaction
with regards to SOs (Figure 2 a + b), the occurrence probabilities of ANT SOs relative to neocortical SOs
derived from occipital electrodes (i.e., electrode O2) were evaluated. Due to missing / noisy occipital
electrodes 2 patients (4 ANT contacts) had to be removed from this analysis. Occipital SOs that did not
overlap with frontal detected SOs (± 750 ms, n = 1054.91 ± 118.38) were isolated, to ensure that the
captured SOs are not mainly products of volume conduction. (a) Occurrence probabilities revealed, that
SOs in the ANT did not precede, but peaked in parallel to occipital derived SOs (positive cluster from -
0.05 to 0.1 sec, p = 0.01; time of peak: 0.0 sec). (b) The phase of thalamic SOs for all paired SO-events at
the time of occipital SO down state (± 750 ms, n = 176.12 ± 29.71; note that only 16.31 ± 1.88 % of ANT
SOs were coupled to occipital derived SOs as compared to 33.74 ± 3.02 % in case of frontal SOs; t = -
4.37, p = 0.0003). A significant nonuniform distribution across contacts (Rayleigh z = 7.24, p < 0.001) was
detectable, with the phase of ANT SOs following their occipital counterparts (mean coupling direction:
166.56 ± 6.20°). (c) Direct comparison of occurrence probabilities for ANT-occipital and ANT-frontal SO
interactions revealed a significant difference between the two distributions from -0.25 to -0.05 sec (p =
0.013), indicating that ANT SOs preferentially emerged earlier with regards to frontal scalp SOs (light
blue) as compared to occipital scalp SOs (dark blue). (d) Comparing the phase distributions for the
coupling between thalamus and neocortex in relation to frontal and occipital SOs yielded a statistical
trend in the direction that ANT SOs coupled to earlier phases of frontal SOs as compared to occipital SOs
(Watson-Williams test: F = 4.12, p = 0.056).
1298
1299
1300
1301

Supplementary Fig. 9 | Thalamic spindles coupled to frontal and parietal scalp spindles: (a + b) Occurrence probabilities of ANT spindle peaks relative to frontal (a) and parietal (b) neocortical spindle peaks (maximal negative amplitude, time = 0; bin size = 50 ms), indicating that ANT spindles precede neocortical spindles detected from both frontal and parietal scalp electrodes (frontal: first positive cluster from -0.3 to -0.05 sec, p < 0.001; second cluster from 0.05 to 0.15 sec, p = 0.005; time of peak: -0.05 sec; parietal: positive cluster from -0.25 to 0.05 s, p < 0.001). (c + d) Occurrence probabilities of MD spindle peaks relative to frontal (c) and parietal (d) neocortical spindle peaks (maximal negative amplitude, time = 0), indicating that MD spindles precede neocortical spindles detected from both frontal and parietal scalp electrodes (frontal: positive cluster from -0.2 to 0.1 sec, p < 0.001; time of peak: -0.05 sec; parietal: first significant positive cluster from -0.2 to 0.05 sec, p < 0.001; second positive cluster from 0.15 to 0.2 sec, p = 0.01; time of peak = 0).

Supplementary Fig. 10 | ANT - cortical coupling for fast and slow spindles: Occurrence probabilities of ANT fast **(a)** and slow spindle **(b)** peaks relative to cortical spindles (same electrode sites as in the main analyses; for details see Supplementary table 2; maximal negative amplitude, time = 0; bin size = 50 ms). Both fast and slow ANT spindles significantly precede neocortical fast and slow spindles (fast spindles: first positive cluster from -0.3 to -0.1 sec, p = 0.0007; second cluster from 0.15 to 0.2 sec, p = 0.015; time of peak: -0.1 sec; slow spindles: positive cluster from -0.1 to 0.05 sec, p = 0.001). **(c)** the comparison of fast ANT-neocortical (dark blue) and slow ANT-neocortical SO interactions (light blue), did not yield significant differences (p > 0.05). **(d + e)** Occurrence probabilities for ANT fast and slow sleep spindles with regards to frontal neocortical fast and slow spindles. Fast ANT spindles significantly precede neocortical frontal fast spindles (first positive cluster from -0.4 to -0.05 sec, p < 0.0001; second cluster from 0.15 to 0.2 sec, p = 0.008; time of peak: -0.1 sec). In case of slow spindles, no significant differences were observable when tested against event-free occurrence probabilities (p = 0.06). **(f)** the comparison of frontal fast ANT-neocortical (dark blue) and slow ANT-neocortical SO interactions (light blue), did not yield significant differences (p > 0.05) **(g + h)** Occurrence probabilities for ANT fast and slow sleep spindles with regards to parietal neocortical fast and slow spindles. Fast ANT spindles significantly precede neocortical parietal fast spindles (first positive cluster from -0.25 to -0.2 sec, p = 0.008; second cluster from -0.1 to 0.05 sec, p = 0.0002; time of peak: -0.1 sec). In case of slow spindles, no significant differences were observable when tested against event-free occurrence probabilities (p = 0.09). **(i)** the comparison of parietal fast ANT-neocortical (dark blue) and slow ANT-neocortical SO interactions (light blue), did not yield significant differences (p > 0.05).

1340
Supplementary Fig. 11 | MD - cortical coupling for fast and slow spindles: Occurrence probabilities of
MD fast (a) and slow spindle (b) peaks relative to cortical spindles (same electrode sites as in the main
analyses; for details see Supplementary table 2; maximal negative amplitude, time = 0; bin size = 50 ms).
Both fast and slow MD spindles significantly precede neocortical fast and slow spindles (fast spindles:
positive cluster from -0.3 to 0.2 sec, p < 0.0001; slow spindles: positive cluster from -0.1 to -0.05 sec, p
= 0.006). (c) the comparison of fast MD-neocortical (dark red) and slow MD-neocortical SO interactions
(light red), yielded multiple significant differences (negative cluster 1 from -0.75 to -0.5 sec, p > 0.0001;
negative cluster 2 from -1.1 to -0.95 sec, p = 0.002; negative cluster 3 from 1.2 to 1.3 sec, p = 0.004). (d
+ e) Occurrence probabilities for MD fast and slow sleep spindles with regards to frontal neocortical fast
and slow spindles. Both fast and slow MD spindles significantly precede neocortical frontal fast and slow
spindles (fast spindles: positive cluster from -0.35 to 0.3 sec, p < 0.0001, time of peak: -0.05 sec; slow
spindles: first positive cluster from -0.25 to -0.15 sec, p = 0.002; second positive cluster from -0.05 to
0.15 sec, p < 0.0001; time of peak: -0.05). (f) the comparison of fast frontal MD-neocortical (dark red)
and slow MD-neocortical SO interactions (light red), yielded two negative clusters (negative cluster 1
from -0.95 to -0.7 sec, p > 0.0001; negative cluster 2 from -1.35 to -1.15 sec p > 0.0001). (g + h)
Occurrence probabilities for MD fast and slow sleep spindles with regards to parietal neocortical fast and
slow spindles. Both fast and slow MD spindles significantly precede neocortical parietal fast and slow
spindles (fast spindles: positive cluster from -0.25 to 0.15 sec, p < 0.0001, time of peak: -0.1 sec; slow
spindles: positive cluster from -0.1 to -0.05 sec, p = 0.006; time of peak: -0.05). (i) the comparison of fast
parietal MD-neocortical (dark red) and slow MD-neocortical SO interactions (light red), one two negative
clusters (negative cluster 1 from -0.7 to -0.5 sec, p > 0.0001).
1362