There and Back to the Present: An Evolutionary Tale on Biological Diversity

Leandro Duarte[1*], Gabriel Nakamura[1], Vanderlei Debastiani[1], Renan Maestri[1], Maria João Ramos Pereira[1], Marcus Cianciaruso[2], José Alexandre F. Diniz-Filho[2]

[1] *Programa de Pós-graduação em Ecologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul*

Av. Bento Gonçalves 9500 CP 15007 Porto Alegre, 91501-970, Brazil

[2] *Programa de Pós-graduação em Ecologia e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Goiás. CP 131, 74690-900, Goiânia, GO, Brazil*

*Corresponding author: duarte.ldas@gmail.com

Running head: Niche evolution predicts diversity

Keywords: Ornstein-Uhlenbeck model, adaptive rate, Approximate Bayesian Computation, eco-evolutionary dynamics

ABSTRACT: Ecologists often agree on the importance of macroevolution for niche-mediated distribution of biological diversity along environmental gradients. Yet, macroevolutionary diversification and dispersal in time and space generate uneven geographic distribution of phylogenetic pools, which affects the imprint let by macroevolution on local species pools. In this article we introduce an individual-based simulation approach coupled to Approximate Bayesian Computation (ABC) that allows to parameterize the adaptation rate of species' niche positions along the evolution of a monophyletic lineage, and the intensity of dispersal limitation, associated with the distribution of biological diversity between assemblages potentially connected by dispersal (metacommunity). The analytical tool was implemented in an R package called *mcfly*. We evaluated the statistical performance of the analytical framework using simulated datasets, which confirmed the suitability of the analysis to estimate adaptation rate and dispersal limitation parameters. Further, we evaluated the role played by niche evolution and dispersal limitation on species diversity distribution of Phyllostomidae bats across the Neotropics. The framework proposed here shed light on the links between niche evolution, dispersal limitation and the distribution of biological diversity, and thereby improved our understanding of evolutionary imprints on ecological

31   patterns. Perhaps more importantly, it offers new possibilities for solving the eco-

32   evolutionary puzzle.

Introduction

Why some places have so many species while others have so few? This is a central question that generations of ecologists after G. E. Hutchinson have been trying to answer after his seminal article (Hutchinson 1959). Indeed, biological diversity is the most widely addressed feature of ecological assemblages; yet, its interpretation may be tricky (Ricotta 2005). The mechanisms determining the striking variation of biological diversity over space and across environmental gradients have been debated for decades (Vellend 2016). Alternative (and complementary) theories have been proposed to explain biological diversity, either based on species niches (Hutchinson 1957; Levine and HilleRisLambers 2009), neutral processes (Hubbell 2001; MacArthur and Levins 1967) or biogeographic and historical factors (Gerhold et al. 2018; MacArthur 1972; Ricklefs and Schluter 1993).

Over the last decades, community phylogenetics (Cavender-Bares et al. 2009; Mouquet et al. 2012; Webb et al. 2002) has posited that phylogenetic relationships among sympatric species may link niche evolution to biogeography and, ultimately, help understanding patterns in regional and local biological diversity patterns. Accordingly, niche divergence between closely related species may be somehow prevented along evolutionary history (Felsenstein 1985; Harvey and Pagel 1991; Nosil 2012). Such tendency of species niche to vary less than expected by chance along evolution is often interpreted as evidence of phylogenetic niche conservatism (Losos 2008; Pyron et al. 2015; Wiens and Graham 2005). If so, and whether dispersal plays only a minor role in species assembly process (which is very disputable, see Vellend 2016), phylogenetic niche conservatism would provide an explicit causal link between the phylogenetic diversity of local communities and major ecological mechanisms (environmental filtering, biotic interactions, see Webb et al. 2002). Following this rationale, some ecologists have been advocating the use of phylogenetic covariation as a proxy for niche similarity among species (Cadotte et al. 2012; Tucker et al. 2018).

Nonetheless, evidence against such 'phylogeny-as-proxy' approach has been accumulated over the last years, as similar phylogenetic diversity patterns may be explained by different, and often opposing, mechanisms (Gerhold et al. 2015; Godoy et al. 2014; Mayfield and Levine 2010; Swenson 2019). As controversial conclusions on the usefulness of interpreting phylogenetic diversity in the light of ecological mechanisms promoting biological diversity proliferated, further theoretical advances in community phylogenetics have decelerated over the last years (but see Cadotte et al.

2017; Tucker et al. 2018). Tracing back plausible evolutionary scenarios underlying

species distribution patterns across current environmental gradients remains a main

challenge for community phylogenetics (Webb et al. 2002).

Unravelling evolutionary processes underlying niche divergence might help

explaining the distribution of extant species belonging to a given clade across ecological

gradients (McPeek 2017; Weber et al. 2017). Nonetheless, such enterprise faces two big

challenges. First, theoretical and empirical evidence often demonstrate that the

inference of eco-evolutionary niche dynamics based on static estimates of phylogenetic

signal in species traits is not unequivocal (Münkemüller et al. 2015; Revell et al. 2008).

This is because strong phylogenetic signal in species traits can emerge from processes

other than niche conservatism, including neutral evolution (Cooper et al. 2010; Gould

and Lewontin 1979; Harvey and Pagel 1991; Nosil 2012). Indeed, under high adaptive

rate pulling species niches to an optimum value (which might be also interpreted as

niche conservatism), phylogenetic signal in the niche is completely vanished (Butler

and King 2004; Hansen et al. 2008). Second, ecologists often measure species traits to

characterize their niches (Elton 1927; McPeek 2017), under the sound premise that

species niches are mostly molded by ecological pressures determining phenotypic

patterns of species (Ackerly 2003; Hansen and Martins 1996). In this sense, trait

dominance and/or complementarity across environmental gradients (Garnier et al. 2016)

is often taken as evidence of niche-mediated species assembly patterns. However,

species traits correlate among themselves due to their shared evolutionary history,

which results in complex trait-environment relationships. Therefore, inferring which

specific trait(s) explain the distribution of biological diversity across environmental

gradients is not trivial (Duarte et al. 2018). Indeed, searching for an evolutionary basis

of niche-based species assembly across environmental gradients is still a topic of debate

among ecologists (Cadotte et al. 2017; McPeek 2017; Tucker et al. 2018; Weber et al.

2017).

If neither phylogenetic signal in niche traits nor trait-environment correlations

allow to build a solid bridge between niche evolution and species assembly across

environmental gradients, then what? A starting point is to consider under which

circumstances we should expect a detectable imprint of evolutionary processes on

biological diversity. First, such imprint only makes sense if we consider a monophyletic

lineage with most branches well represented across the geographic space under analysis.

Otherwise, we may be investigating phylogenetic signal in biological diversity based on

species (1) whose evolution was independent for the most time lineages have been

diversifying, for example, tree ferns and angiosperms (Duarte et al. 2012) or Anura and

Gymnophiona (Loyola et al. 2013), or (2) whose phylogenetic relationships are too

incomplete or fragmented to provide a realistic evolutionary scenario (Crisp and Cook

2005). However, even when the species pool belongs to a monophyletic, well

represented lineage, endemic to the region or biome we are interested in, the imprint left

by phylogeny on regional or local diversity patterns may either persist or be completely

erased (Maestri and Duarte 2020). This is because species assembly, and therefore

biological diversity, is an instantaneous snapshot of complex eco-evolutionary

dynamics acting upon co-occurring species (Shipley 2010; Weber et al. 2017). If the

adaptation rate of species from an ancestral niche position towards an optimum niche

condition, mediated by short-term biotic interactions (Paine et al. 2018) and/or

environmental filters (Kraft et al. 2015), was sufficiently strong to break phylogenetic

signal in species niches (Hansen et al. 2008), no phylogenetic imprint on species

assembly would be detected. Thus, it is not surprising that tracing back evolutionary

processes underlying biological diversity patterns based on static phylogenetic signal

among species may lead to disparate conclusions (Gerhold et al. 2015). Appropriate

tools for the evaluation of the influence of phylogeny on biological diversity across

space are necessary for this purpose.

A possible solution to solve this puzzle implies building alternative scenarios of

niche evolution given a phylogenetic hypothesis to generate expected species assembly

patterns. This should allow us to infer the most likely scenarios of niche evolution

determining empirical species assembly patterns based on the set of expectations that

most resemble the empirical data. In this article we introduce a model-based simulation

approach that allows to estimate the strength of adaptation rate along the niche

evolution of species belonging to a monophyletic lineage and distributed across

metacommunities, defined as sets of local assemblages connected by dispersal (Leibold

et al. 2004). Moreover, the new framework also allows to estimate to what extent

dispersal limitation influences diversity distribution across space. We start by defining

the theoretical aspects of niche- and neutral-based species assembly into local

assemblages, which provide the necessary background to a further contextualization of

niche evolution as a component of niche-based species assembly. Afterwards, we

introduce a rejection-based Approximate Bayesian Computation (ABC) algorithm

(Beaumont 2010; Csilléry et al. 2010) to estimate the strength of adaptation rate and

dispersal limitation given a set of sites described by species belonging to a

monophyletic clade, a phylogenetic hypothesis for these species, spatial coordinates of

sites, and an environmental factor associated with species assembly patterns. We

demonstrate the usefulness of the simulation approach for inferring the imprint left by

niche evolution and dispersal limitation on species diversity patterns across

metacommunity and biogeographical scales by evaluating the statistical performance of

the ABC algorithm using simulated datasets. Further, we investigate the role played by

niche evolution and dispersal limitation on species diversity distribution of

Phyllostomidae bats across the Neotropics.

Parameterizing niche and neutral processes determining species assembly

Niche theory partially explains in part the distribution of biological diversity across

space. The availability of key resources and the prevalence of specific levels of

environmental conditions in a site enable populations of some species to thrive, while

others decrease their numbers for the same reason. Together with environmental

filtering, biotic interactions modulate the probability of maintenance of viable

populations of some species in the presence of others (Ackerly 2003; Hutchinson 1957;

MacArthur and Levins 1967; Vellend 2016; Violle et al. 2012, but see Kraft et al.

2015).

On the other hand, neutral dynamics relies on dispersal-based species assembly

to explain biological diversity. Neutral dynamics is modeled based on dispersal

limitation, immigration rates, local and regional species abundances, and speciation

events determining demography in local species assemblages, assuming that species are

ecologically equivalent in respect to their niches (Hubbell 2001). Even though no one

dispute the importance of niche-based processes to explain species assembly patterns,

dispersal processes also play an important role to determine species distribution across

space (Vellend 2016). Therefore, the importance of neutral processes (Bell 2000;

Hubbell 2001) cannot be underestimated. In fact, such dichotomy between niche and

neutral processes is only apparent (Gravel et al. 2006), since these processes represent

complementary mechanisms determining biological diversity (Vellend 2016). Both

niche and neutral processes provide a solid theoretical background to model species

assembly into local assemblages (Gravel et al. 2006) distributed in a metacommunity

context.

167   Consider a metacommunity M defined by sets of local assemblages (k) with
168 some degree of connection by dispersal. Let N_{*k} be the number of individuals inhabiting
169 each k assemblage, and N_{*M} be the total number of individuals in the metacommunity,
170 such that $N_{*M} = \sum_{k=1}^{M} N_{*k}$. In the absence of a major disturbance, N_{*k} is expected to
171 increase until k achieves carrying capacity (J_k), quantified as the maximum number of
172 individuals allowed to inhabit k, given the minimum amount of space and resources per
173 capita demanded by the organisms (Del Monte-Luna et al. 2004). Thus, under J_k, we
174 expect $dN_{*k}(t) = 0$, and species assembly likely starts to follow a zero-sum dynamic
175 (Hubbell 2001). This means that an individual currently inhabiting k must necessarily
176 die before a new one can be recruited, either by birth or immigration from other
177 assemblage.
178   Given there is vacant space in k to be occupied by new recruits, what processes
179 will determine their identity? As we seen before, niche processes may be the answer to
180 this question, at least partially. For a given species i belonging to the species pool q, the
181 niche position y_i can be defined as the environmental condition that maximizes the
182 probability of $N_{ik} > 0$ such that $dN_{ik}(t) \geq 0$. Further, the niche breadth s_i can be defined
183 as the amount of deviation from y_i that allows the conditions $N_{ik} > 0$ and $dN_{ik}(t) \geq 0$ to
184 be satisfied (Sexton et al. 2017). From a niche perspective, and considering only the
185 influence of environmental filtering, the probability (λ_{ik}) of species i to recruit a new
186 individual in k can be modeled as a deterministic Gaussian function based on their niche
187 position and breadth, as follows:

$$\lambda_{ik} = e^{\frac{-(x_k - y_i)^2}{2s_i^2}}. \quad (1)$$

189 Accordingly, x_k is the environmental position of k across an environmental gradient, y_i
190 is the niche position of i (measured in the same scale of x_k), and s_i is the niche breadth
191 of i. This model has been originally proposed by Minchin (1987), and has been adopted
192 in several studies to simulate species assemblages (Dray and Legendre 2008; Duarte et
193 al. 2018; Nakamura et al. 2020; Peres-Neto et al. 2017; Peres-Neto et al. 2012; Sokol et
194 al. 2017; Sokol et al. 2015).
195   We may now compute a recruitment probability ($R_{\text{Niche}_{ik}}$) that implicitly
196 incorporates the effect of biotic interactions (Gravel et al. 2006):

$$R_{\text{Niche}_{ik}} = \frac{\lambda_{ik}}{\sum_{j \neq i}^{q} \lambda_{jk}}. \quad (2)$$

Note that λ_{ik} is standardized by the probabilities of all other j species, such that j

$\in q$, which are also candidates to recruit new individuals, competing directly with i.

Since $R_{\mathrm{Niche}_{ik}}$ is only defined by niche position and breadth, q can contain not only the

species already resident in k, but also potential immigrant species occurring at any

assemblage $l,$ such that $l \in M$. The only condition to be a candidate recruiter species to k

is to show $R_{\mathrm{Niche}_{ik}} > 0$. This condition agrees with the species sorting model of

metacommunity theory (Leibold et al. 2004).

Gravel et al. (2006) improved this niche-based model of species assembly by

incorporating parameters from Hubbell's neutral model (Hubbell 2001), which provides

spatial context to species assembly in local assemblages. First, let us consider dispersal

limitation as a mechanism affecting the success of any species i belonging to the

regional species pool q, and occurring in any l assemblage, to disperse new recruits to

assemblage k. The probability δ_{lk} of any i inhabiting l to contribute to the immigrant

pool of k may be modeled as a Gaussian function of the shortest linear distance (r,

rescaled between 0 and 1, see Sokol et al. 2017) between k and l, and the slope w of the

dispersal kernel (the probability density function of the dispersal success from a source

to a sink assemblage, see Nathan and Muller-Landau 2000) for the species in l (Gravel

et al. 2006; Sokol et al. 2017; Sokol et al. 2015), as follows:

216
$$\delta_{lk} = e^{-wr_{lk}^2}. \quad (3)$$

For a slope $w = 0$, $\delta_{lk} = 1$, independently of the distance between l and k, which means

that the species in assemblage l face no barrier to disperse to k (the assemblages are

panmictic). On the other hand, for slope $w = 10$, $\delta_{lk} = 0.9$ for sites showing $r = 0.1$, but

decreases fast to $\delta_{lk} = 0.08$ if the distance between the assemblages increases to $r = 0.5$.

Over the next sections we introduce a simulation-based framework to estimate w for

empirical datasets, which enables quantifying dispersal limitation underlying diversity

patterns (see *Estimating niche and neutral parameters underlying species diversity*

gradients using Approximate Bayesian Computation).

The probability of any species i to successfully disperse new recruits from any

assemblage l ($R_{\mathrm{Dispersal}_{iM}}$), is a function of the abundance of i across M, or $N_{iM} =$

$\sum_{l \neq k}^{M} N_{il}$, and δ_{lk}, and is contingent on the probability of dispersal for all other species q in

M (Gravel et al. 2006), as follows:

229
$$R_{\text{Dispersal}_{iM}} = \frac{\sum_{l \neq k}^{M} N_{il} \delta_{lk}}{\sum_{j \neq i}^{q} \sum_{l \neq k}^{M} N_{jl} \delta_{lk}}. \quad (4)$$

It is important to observe that even for $\delta_{lk} = 1$, immigrant species arriving to site k may

not succeed as new colonizers, depending on how resistant k is to immigration.

Assemblages where resident species are strong recruiters may prevent the entrance of

new colonizers (e.g. Duarte et al. 2006), which reduces the chance of new species

become part of the assemblage, even in the absence of dispersal limitation. For a site k

under zero-sum dynamic, let d_k be the number of instantaneous deaths in k. Site k is

now open to the recruitment of $J_k - d_k$ new individuals, which may be generated either

by resident parents inhabiting k (b_k), or by individuals dispersing from any site $l \in M$

(b_l), such that $J_k = (b_k - d_k) + b_l$. The immigration coefficient m_k (Hubbell 2001) can

then be computed as

240
$$m_k = \frac{b_l}{b_l + b_k - 1}. \quad (5)$$

241   Now we can compute the recruitment probability of a species i in the

242   assemblage k given the neutral parameters $R_{\text{Dispersal}_{iM}}$, N_{ik}, N_{iM}, and m_k (Sokol et al.

243   2015), as follows:

244
$$R_{\text{Neutral}_{ik}} = m_k \cdot R_{\text{Dispersal}_{iM}} + (1 - m_k) \cdot N_{ik}. \quad (6)$$

245   Finally, we compute the recruitment probability of any species $i \in q$ in any

246   assemblage $k \in M$ that considers both niche and neutral processes, as

247
$$R_{ik} = R_{\text{Niche}_{ik}} \cdot R_{\text{Neutral}_{ik}}. \quad (7)$$

The definition of R_{ik} provides an integrative solution to model species assembly patterns

in local assemblages that allow estimating the strength of niche and neutral process

determining the dynamics of biological diversity across space. Yet, the evaluation of the

imprint left by niche evolution on species assembly, and ultimately on biological

diversity, is still lacking. Over the next section we present a solution to fill this gap.

Niche evolution and species assembly across environmental gradients

The Ornstein-Uhlenbeck (OU) model of trait evolution (Hansen 1997) allows

parameterizing the strength of the adaptation rate α of species niches evolving towards

an optimum niche condition θ, as follows:

257
$$dy_i(t) = \alpha \, [\theta - y_i(t)] \, dt + \sigma \, dB(t). \quad (8)$$

For any θ and given the niche position $y_i(t)$ of species i, adaptation rate α is a force that

deterministically decreases the fitness of $y_i(t)$ as $\theta - y_i(t) \neq 0$. As $\alpha \to 0$, the strength of

adaptation rate decreases, and trait divergence among species is increasingly modeled

by the stochastic component of the OU model ($dB[t]$), which is a stochastic white-noise

process showing Gaussian distribution ($N[0,dt)]$, Brownian motion), and standard

deviation σ, which indicates the amplitude of niche position variation among species

(Butler and King 2004; Hansen et al. 2008). Therefore, under low α values, species

niche position evolves under strong phylogenetic inertia, generating high phylogenetic

signal in species niche positions (Blomberg and Garland 2002). On the other hand, as α

$\to \infty$, species niches tend to evolve faster from the ancestral state ($y_{i\,anc.}$) towards θ, and

the imprint of phylogenetic inertia on species niches tends to be erased. The rate of

niche evolution towards θ can be estimated as a phylogenetic half-life (Hansen 1997;

Hansen et al. 2008), which indicates the amount of time units needed to move half the

distance between the ancestral state and the optimum condition, and is calculated as

272
$$t_{1/2} = \frac{\ln 2}{\alpha}. \qquad (9)$$

Note that $t_{1/2}$ is expressed in time units, which means that the effect α exerts on niche

evolution depends on the timespan between $y_{i\,anc.}$ and θ. If the timespan is long enough

(say, 200 million years), even a small α value (say, $\alpha = 0.01$) may generate a moderate

movement towards θ ($t_{1/2} = 69.3$). Nonetheless, if the evolutionary timespan is shorter

than $\cong 70$ million years, the phylogenetic half-life corresponding to $\alpha = 0.01$ will be too

slow to produce significant niche evolution towards the niche optimum, and therefore

niche evolution may show phylogenetic inertia not quite distinguishable from Brownian

motion ($\alpha = 0$, $t_{1/2} = +\infty$). Thus, the expected value of a niche y_i, evolving under OU

model during a time t, results from a deterministic process modulated by the adaptation

rate α that forces $y_i(t)$ to converge towards the optimum niche condition θ, and a

stochastic, Brownian process, as follows:

284
$$E\left[y_i(t)\right] = \theta\,(1 - e^{-\alpha t}) + y_{i\,anc.}\,e^{-\alpha t}. \qquad (10)$$

The critical step is to translate the OU parameters α and θ into quantities that can

be estimated from assemblages described by species abundances or occurrences, an

environmental variable associated with species assembly, and a phylogenetic tree. In a

metacommunity M composed by k local assemblages defined by q species belonging to

a monophyletic lineage, those local assemblages with the highest levels of biological

diversity can be considered as offering the best environmental condition x to most

species in q (Enquist et al. 2015; Violle et al. 2007), and therefore it is likely to be an

appropriate estimator of θ. We might then take x as an empirical estimate of θ, or x_θ.

293   Niche filtering is expected to be stronger in assemblages with environmental

294   condition deviating from x_θ, leading to lower species diversity. Differently from x_θ,

295   which can be estimated from data, the parameter α cannot be directly assessed, but its

296   posterior distribution can be estimated by means of rejection-based Approximate

297   Bayesian Computation (Beaumont 2010; Csilléry et al. 2010). We explain this in detail

298   in the next section.

299   For now, it is important to understand what the parameter α can inform about the

role of niche evolution on species assembly patterns. If $\alpha \to 0$, we may conclude that

niche evolution was mainly a stochastic Brownian process, which means that the

adaptation rate towards x_θ was very weak along niche evolution, insufficient to break

the phylogenetic signal of the ancestral niche of species. Under such evolutionary

scenario, niche divergence between species will be a function of the time since their

evolutionary divergence, leading to strong phylogenetic covariation in species niches.

Furthermore, the total variance in the niche position will be high, which means that

niche positions of species will be more loosely distributed across the upper and lower

limits of the environmental condition x. Niche-based assembly processes in local

communities will tend to be phylogenetically structured, while differences in diversity

values among assemblages will be not remarkable. As $\alpha \to \infty$, adaptation rate of species

niches towards x_θ becomes faster along niche evolution, so that any track of

phylogenetic signal in the niche vanishes along evolution (Cooper et al. 2016). Niche

divergence between species will be constrained by the optimum niche condition, with

values approaching x_θ as α increases, decreasing the total variance of the niche position

among species. In local assemblages showing environmental conditions close x_θ, steep

diversity peaks are expected to occur.

Estimating niche and neutral parameters underlying species diversity gradients

using Approximate Bayesian Computation

Consider that assemblages in M are distributed across an environmental gradient

defined by a single environmental condition, or multiple conditions summarized in a

single factor or principal component x. We may estimate x_θ as the environmental

condition interval that maximizes species diversity in M. In turn, whenever the niche

position y_i of a given species in M approaches x_θ, the probability of occurrence of that

species in the assemblage increases, conditioned to the niche positions of all other

species in the assemblage. The adaptation rate of y along the evolutionary history (i.e.

OU's α) is a niche-assembly parameter that can be estimated given the species diversity

variation in M, x, the spatial distribution of local assemblages in M, and the

phylogenetic relatedness among species distributed across M. By doing so, we can

reveal how strong is phylogenetic signal in niche position of species. On the other hand,

species diversity in a local assemblage may be limited by the dispersal capacity of

species. As we have seen before, this depends on the spatial distance between sites (r)

and the slope w of the dispersal kernel, in complement to niche characteristics of species

(Gravel et al. 2006). Therefore, estimating w allows quantifying the influence of

dispersal limitation on the distribution of species diversity across M.

To estimate niche (OU's α) and neutral (slope w) parameters associated with the

distribution of species diversity, we developed an Approximate Bayesian Computation

(ABC) algorithm (Beaumont et al. 2002; Csilléry et al. 2010) that works as

schematically represented in figure 1.

Figure 1: Schematic representation of *mcfly*, an Approximate Bayesian Computation framework
developed to estimate the posterior distribution of adaptation rate (OU's α) and dispersal kernel slope (*w*)
parameters driving species diversity gradients along an environmental gradient. **1**. The *real
metacommunity* provides the empirical measures of species diversity, the *real summary statistic* (*R*). **2**.
Environment is a dimension of species niches that provides the optimum niche (θ), variance (σ) and root
niche position parameters necessary to simulate species niche based on the *phylogeny,* according to
Ornstein-Uhlenbeck evolutionary model. We also define niche breadth in the simulation based on
environment. **3**. A prior distribution of OU's α is used to simulate species niche positions based on
environment parameters and *phylogeny*. **4**. *Spatial coordinates* of *k* sites are used to define the prior
distribution of slope *w*; together with immigration rate (*m*), number of individuals at *k* sites (J_{L_k}) and the
total number of individuals in *M* (J_M) provide the neutral parameters of simulations. **5**. Metacommunities
are simulated based on niche and neutral parameters (*Simulator*). **6**. Each metacommunity simulation
provides species diversity measures (*Summarizer*) taken as the simulated summary statistics (*S*). **7**.
Simulated diversities (*S*) are classified based on a pre-defined minimum correlation threshold (tolerance)
to the observed diversity (*R*). Every *S* object not discarded (*Reject*) is then *scanned* to record its OU's α
and slope *w* to provide a posterior distribution of adaptation rate (α) and dispersal limitation (*w*),
respectively.

We implemented the ABC framework in the R package *mcfly* (freely available at
(https://github.com/GabrielNakamura/mcfly), which has ´mcfly´ as the core function.

360   An additional function that helps users to define the appropriate tolerance threshold

361   ('define_tolerance') is also available in the package, as well as a working example to

362   illustrate how the user can set all the parameters needed to run ´Mcfly´ function. The

363   example can be easily accessed in R by typing *vignette("mcfly_vignette")*.

364   **The 'simulator engine'**

The simulation framework implemented in *mcfly* package derives from the individual-

based simulation protocol developed by Sokol et al. (2015, 2017) and available in the R

package *MCSim* (available at https://github.com/sokole/MCSim). Accordingly, sets of

assemblages are simulated based on lottery colonization and recruitment dynamics,

following niche and neutral mechanisms (Gravel et al. 2006), as explained in the section

Parameterizing niche and neutral processes determining species assembly.

371   *Metacommunity dimensions and species pools*

Simulated metacommunities M_{Sim} show the same number of sites of the

observed metacommunity M. If M contains information on species abundances, the

number of individuals in each simulated assemblage equals the number shown in the

respective observed local assemblage. Because the original simulation protocol of

MCSim is individual-based, when only species incidence (presence/absence) data is

available the simulation performed in *mcfly* addresses 100 individuals to each species

present in each local assemblage prior to the simulation of M_{Sim}. In the end, M_{Sim} is

transformed to an incidence matrix.

Noteworthy, the phylogenetic tree used to build M_{Sim} should be as complete as

possible to increase the quality of niche simulations, which means it should not be

pruned to keep only the species present in M. In all simulations, the species pool of M_{Sim}

will be the same defining $M,$ independently of the size of the species pool included in

the phylogeny. According to Sokol et al. (2015), recruitment patterns in M_{Sim} stabilize

after 30 simulation time steps. By default, we set simulations to stop after 50 timesteps.

386   *Niche prior and other parameters*

At each simulation run, the niche position of each species is simulated for all species

included in the phylogenetic tree using the function 'rTraitCont' of the package *ape*

(Paradis and Schliep 2019) as a continuous trait y following OU model. The adaptation

rate parameter α of niche simulation is randomly sampled from a prior distribution, with

predefined size and distribution mode. The minimum value of the α parameter is set to

correspond to the half-life ($t_{1/2}$, see eq. 9) equal to the total timespan of the phylogeny.

For a phylogeny with timespan = 30 Myr, the minimum α value will be 0.023, since

394
$$t_{1/2} = \ln(2)/0.023 = 30.$$

395   The maximum α parameter value corresponds to $t_{1/2} = 1/30 \cdot$ timespan. Thus, for that

396   same timespan = 30 Myr, the maximum α value = 0.693, since

397
$$t_{1/2} = \ln(2)/0.693 = (1/30) \cdot 30 = 1.$$

In the example above, the range of α parameters between the minimum and

maximum limits expresses a broad variation in phylogenetic signal. Indeed, we observe

an asymptotic decay in phylogenetic signal, estimated by Blomberg's K (Blomberg and

Garland 2002), with increasing OU's α values (fig. 2). Considering the α parameter

interval ranging between 0.023 and 0.693, K statistics is shown to vary between 0.983

and 0.137, respectively. Further, the lower 95% credibility limit for OU's α values

estimated from vectors randomly drawn from Gaussian distribution (white noise)

reveals that niche positions showing α values higher 0.628 cannot be distinguished from

white noise variables and show very low phylogenetic signal ≤ 0.165. Therefore, we

conclude that the range of prior α values defined in the simulation algorithm is

appropriate. Note that the simulation does not rescale the branch lengths of phylogenetic

tree. Therefore, the prior distribution of OU's α values will differ from the above

whenever the timespan differs from 30 Myr, and therefore the interpretation of results

must consider the timespan of the phylogenetic tree under analysis. By default, *mcfly*

generates a uniform distribution of the α prior, but the user may opt for generating a

prior α with half-life values uniformly distributed, instead. In the next sections we

provide a sensitivity analysis to evaluate the implications of the α prior distribution for

the posterior distribution of OU's α parameter.

Figure 2: Relationship between Blomberg's K statistic and the adaptation rate parameter of Ornstein-Uhlenbeck model (OU's α) for simulated niche positions computed for 1,000 sets of 50 species each (black circles). Gray areas indicate lower and upper credibility intervals (95%) for Blomberg's K and OU's α values estimated from white noise vectors (N[0,1]).

Another critical niche parameter to be defined is the optimum niche position

(x_θ). In our framework, x_θ is empirically defined as the values of the environmental

condition e showing the highest species diversities in M. In *mcfly*, all e_k values

corresponding to ≥ the 5% highest species diversity values will constitute a vector of x_θ

values. At each metacommunity simulation process, a single x_θ value is randomly drawn

from that vector and taken as the optimum niche. Thus, it is important to note that the

more unimodal is diversity distribution across e, the more homogeneous will x_θ among

different M_{Sim}, which is likely to decrease the credibility intervals of the posterior

distribution of OU's α and slope w parameters. Moreover, the variance of the niche

position (σ) was empirically defined as $\sigma = \sqrt{\text{standard deviation }(x)}$, and root node value

$= \overline{x}$.

433   *Neutral prior and other parameters*

The spatial coordinates of each assemblage in M are used to compute pairwise distances

r between local assemblages in M, which are rescaled to vary between 0 and 1. As we

have seen in equation 3, r is a component defining the probability of a species to

disperse from a site to another, whose effect on dispersal limitation is modulated by the

slope w, the neutral assembly parameter of interest in our framework. It is important to

note that the range of variation of w, which expresses the strength of dispersal

limitation, depends on the shape of the spatial distribution of sites. When sites are

evenly distributed across the space, as in a cell grid, *w* slopes must vary over broader

scales to capture dispersal limitation over shorter distances, and vice-versa. Thus, to

build a prior distribution of slope *w*, we first defined the maximum distance *r* between

sites enough to connect all sites in *M* by means of a minimum spanning tree (Legendre

and Legendre 2012). This threshold distance provides information on how far two sites

should be to each other to dispersal limitation to become relevant and is widely used in

spatial analysis (Borcard and Legendre 2002). Then, we simulated a uniform

distribution of dispersal probabilities between sites (δ_{lk}, see eq. 3), varying between 0

and 1, and with a predefined size. Based on equation 3, we then computed the slope *w*

corresponding to every simulated δ_{lk} for a hypothetical pair of sites located at distance *r*

to each other. By doing so, we build the prior distribution of slopes *w*, which is not

uniform itself, but decays logarithmically as *w* increases.

As we have seen in eq. 5, the probability of immigration in each assemblage (*m*)

has also to be defined prior to the simulation, which is not trivial in most cases. The

user may opt for indicating an arbitrary value; the default value in *mcfly* is 0.5, which

means that the simulation allows a half of the individuals to be replaced at each timestep

of the simulation process. Alternatively, the user may inform empirically estimated *m*

for each site, whenever these data are available.

The summary statistic

Our simulation framework summarizes species diversity in the local assemblages of *M*

and M_{Sim} using Rényi entropy (Rényi 1961), which is a flexible measure that generalizes

widely used diversity measures, such as Shannon and Simpson diversity indices (Anand

and Orlóci 1996). Rényi entropy (H_{order}) is computed as follows:

464
$$H_{order} = \frac{\log_2 \sum_{i=1}^{q} p_i^{order}}{1 - order}. \quad (11)$$

Accordingly, *q* is the number of species present in the community, p_i is the proportional

abundance of each species *i* in the community, and order is a scalar that modifies the

sensitivity of entropy to variation in species abundances (evenness). In this study, we

analyzed three entropy orders: 1, 2, and 12. For entropy order = 1, Rényi entropy is

equivalent to Shannon diversity index, while entropy order = 2 assumes values

equivalent to Simpson index (Anand and Orlóci 1996). Higher entropy orders increase

the importance of evenness in the diversity index, which tends to stabilize for entropy

order > 12 (Anand and Orlóci 1996). For datasets characterizing sites by species

incidences (presence/absence data), Rényi entropy is equivalent to species richness (or

entropy order = 0), independently of the entropy order used.

The posterior distribution of adaptive rate (α) and dispersal limitation (w)

Since species diversity is computed for M and all M_{Sim} datasets, the correlation between

observed and simulated diversities is a way to evaluate how similar are observed and

simulated metacommunities. This step involves defining a threshold correlation value

below which metacommunities are rejected. In ABC analysis, a tolerance value is

usually defined, and expresses the maximum distance tolerated between observed and

simulated summary statistics (Beaumont 2010). In *mcfly*, the tolerance expresses the

complement of the correlation (1 – correlation) and must be informed prior to the

simulations. The remaining M_{Sim} datasets surpassing the tolerance threshold provide

their corresponding OU's α and slope w parameters to a posterior distribution of niche

and neutral parameters underlying diversity patterns in M, respectively. A highest

posterior distribution (HPD) provides the credibility intervals of those parameters

(Turkkan and Pham-Gia 1993).

488   **Evaluating *mcfly* performance using simulated metacommunities**

489   To evaluate the performance of *mcfly* we simulated 2,000 metacommunities M using the

490   function 'metasim' of the package *MCSim* (Sokol et al. 2017; Sokol et al. 2015), each of

491   them showing 50 local assemblages, with known niche and neutral parameters, which

492   allowed to evaluate the efficiency of the proposed simulation approach to estimate

493   them.

494   Prior to metacommunity simulation, phylogenetic trees containing 50 species

were simulated using the function 'sim.bdtree' of package *geiger* (Harmon et al. 2008),

assuming homogeneous birth-death processes across all lineages. Speciation followed

purely stochastic birth (speciation rate = 0.1, extinction rate = 0). The number of

lineages increased exponentially with time. Evolutionary timespan was fixed as 30 Myr.

These phylogenies defined the species pools used to perform all further simulations. All

50 species in each phylogeny were allowed to occur in at least one out of 50

communities representing a metacommunity M.

502   For each phylogeny, a single OU's α value defining the real adaptive rate of

species niche positions was randomly drawn from a uniform distribution ranging

between minimum (= 0.023) and maximum (= 0.693) α values. Further, an

environmental variable x was simulated from a uniform distribution (U[0,100]), and its

mean value ($\overline{x}$) was defined as the optimum niche position (θ). OU's α, $\overline{x}$ and other

niche parameters described in the section *Niche prior and other parameters* were then

used to simulate the niche positions y_i of species ranging between 0 and 100 based on

OU method of the 'rTraitCont' function of package *ape* (Paradis and Schliep 2019). For

metacommunity simulation, species niche breadth (s_i, eq. 1) was fixed as 10, which

confers reasonable niche breadth to species (Duarte et al. 2018; Peres-Neto et al. 2012).

Spatial coordinates of assemblages in M were simulated from a uniform

distribution (U[0,1]). The number of individuals per local assemblage and the total

number of individuals in M were defined as, respectively, J_k =1000 individuals, J_M =

1000 x 50 sites = 50,000 individuals. The probability of immigration (*m,* see eq. 5) was

defined as 0.5. The frequencies of species across M were extracted from a multinormal

distribution ranging between 0.1 and 0.4 that was further normalized based on J_M.

Species colonization or recruitment across M stopped after 50 simulation time steps.

Noteworthy, in some simulations, a low number of species was capable of colonizing at

least one local assemblage in M, and therefore gamma diversity in M varied among the

simulations. We analyzed only M datasets containing, at least, 20 species distributed

across local assemblages. We divided the 2,000 M datasets into five subgroups of 400

metacommunities, each of them simulated under different dispersal limitation strengths

(*w* = 0, 1, 5, 15 or 100, see *Neutral prior and other parameters*). Thus, the real *w* slope

of each simulated M was known. After metacommunity simulation, each M dataset was

converted into an incidence matrix describing species occurrences in each local

assemblage.

After defining M datasets, we submitted each metacommunity to ABC analysis

using *mcfly* to evaluate the suitability of the framework to estimate the posterior

distribution of the adaptive rate of niche positions (OU's α) and dispersal limitation

strength (slope *w*). Prior to ABC analysis, the 2,000 metacommunities M were split into

two groups containing 1,000 metacommunities each. In the first group, the prior

distribution of OU's α used in the ABC algorithm was defined as a uniform distribution

of α values ('uniform' α prior). In the second group, α prior distribution was defined

based on a uniform distribution of α half-lives ('half-life' α prior), and the

corresponding α values were then taken as OU's α prior (see *Niche prior and other*

parameters). The maximum sample size of α and *w* priors used to analyze each

metacommunity M was defined as 14,400 simulations of M_{Sim} datasets, while the

posterior sample size was defined as 240. Since we had no previous information about

the appropriate tolerance value (see previous section), prior to ABC we computed

correlations between the summary statistics computed for each M dataset and for 480

M_{Sim} datasets to obtain a correlation distribution. The complement of the 95[th] percentile

of the respective correlation distribution was then taken as the tolerance value in ABC

performed for each metacommunity M.

At each simulation step, *mcfly* simulated M_{Sim} under predefined adaptation rate

(OU's α) and dispersal limitation (slope w), computed the summary statistics (Rényi

entropy of order 1), and correlated it to the real summary statistics computed from M.

Whenever the complement of correlation was ≤ the tolerance limit, α and w parameters

were recorded in the posterior distribution of α and w parameters. After the algorithm

reached either the posterior sample size or the maximum prior size (in cases where the

posterior sample size was not reached), the median values of each parameter in the

posterior distribution were recorded.

Statistical performance of ABC based on simulated datasets

We evaluated the statistical performance of the ABC approach using structural equation

modelling based on d-separation tests (Shipley 2000), which estimates the validity of

causal models using Fisher's C as test statistic that follows a chi-square distribution

with $2k$ degrees of freedom, where k is number of independence relationships necessary

to evaluate the model given the number of variables and the number of causal

relationships proposed in the model (Shipley 2000). Accordingly, the probability of the

null hypothesis of the fitted model to correspond to expected causal relationships

indicates the overall validity of the proposed model. Path coefficients (p) linking

variables in the causal model were estimated using ordinary least squares (OLS). The

analysis was performed using the 'psem' function of the package *piecewiseSEM*

(Lefcheck 2016).

We built the causal models considering first the effects of both real OU's α and

w on the number of species in M (which varied across simulations, as some species

were lost during the assembly), and on the median of the posterior distribution of both

OU's α and w. Furthermore, we evaluated the influence of the prior OU's α mode

('uniform' or 'half-life') on the median of the posterior distribution of OU's α. Finally,

the influence of the number of species in M on both was the median of the posterior

distribution of both OU's α and w. In a second set of structural models, we replaced real

and posterior OU's α by the respective half-life and tested similar causality

relationships. Prior to causal testing, half-life and *w* slopes were log transformed.

We also evaluated causal relationships within each prior OU's α mode. These

models were tested because we found that the effect of prior OU's α mode on the

median of the posterior distribution of OU's α was stronger than the effect of the real

OU's α (the same occurred in the half-life model). Evaluating causal relationships

within each prior OU's α mode enabled us to perform a sensitivity analysis of the ABC

approach to the prior OU's α.

In all causal models, we considered that the residuals of the median of the

posterior distribution of OU's α and *w* slope were correlated to each other.

582   **A study case: phyllostomid bats**

We analyzed the influence of niche evolution and dispersal limitation on the distribution

of species diversity of the Phyllostomidae, a species-rich and ecological diverse family

of bats. Phyllostomid bats are predominantly distributed in the Neotropics, from

California to northern Argentina and Chile (Fleming et al. 2020). The family has 222

known species (Simmons and Cirranello 2020), displaying a large array of dietary and

morphological specializations to feed on fruits, nectar, and other parts of plants, as well

as on insects, vertebrates, and blood (Baker et al. 2012; Fleming et al. 2020; Monteiro

and Nogueira 2011; Rossoni et al. 2019). The remarkable diversity of their ecological

features mirrors a strong latitudinal and environmental gradient of species richness

(Pereira and Palmeirim 2013; Villalobos and Arita 2010), which makes this family

particularly appropriate for our purpose. We gathered geographic range maps for

phyllostomid species from IUCN database (IUCN 2020), and defined species

assemblages based on species incidences in a grid composed by 110 × 110 km cells

distributed across the Neotropical region strictu sensu (see Maestri and Duarte 2020;

Morrone 2014). Phylogenetic relationships among species were taken from the

consensus phylogenetic hypothesis proposed by Shi and Rabosky (2015), the most

comprehensive bat phylogeny available to date. The complete Chiroptera phylogeny

was dated using 24 fossil calibration points distributed along the whole timespan of the

tree. Pruning geographic data given available genetic information resulted in a dataset

of 136 species (*ca.* 62% of total phyllostomid richness) distributed across 1,338

assemblages in the Neotropical region. The total timespan of the pruned phylogeny was

= 34 Myr.

We computed entropy of order 1 for sites described by phyllostomid species and

analyzed its association with each of four climatic variables (mean annual temperature,

temperature seasonality, mean annual precipitation and precipitation seasonality) and

two topographical variables (mean altitude and altitudinal range variation). Climatic

variables were taken from the WorldClim2 database (Fick and Hijmans 2017), and

topographic variables were taken from NASA (http://neo.sci.gsfc.nasa.gov/).

Phyllostomid entropy was strongly negatively related to temperature seasonality ($r = -$

0.91), corroborating results obtained elsewhere (Stevens 2011), and was taken as the

niche condition (x) in further analysis.

Then, based on the species by sites matrix (assemblages), the spatial coordinates

of the sites, the niche condition (precipitation seasonality), and phylogenetic

relationships among species, we were able to estimate the posterior distribution of the

adaptive rate of species niche positions (OU's α) and dispersal limitation strength (slope

w) underlying the distribution of species diversity across phyllostomid assemblages

using ABC analysis implemented in *mcfly*. To simulate niche positions, we defined

$\sigma = \sqrt{\text{standard deviation } (y)}$, root node value $= \bar{y}$. The probability m of new colonizers to

arrive at simulated assemblages $= 0.5$. Niche breadth (s_i) was set to 10. Species

colonization or recruitment in each set of simulated communities stopped after 50

simulation time steps. We set the maximum sample size of OU's α (uniform mode) and

w priors to 36,000, and the size of the posterior sample to 480. The tolerance in ABC

analysis ($1 - r$) was set to 0.1.

626   **Results**

627   *Simulated datasets*

The results of global causal model analyzing factors influencing the posterior

distribution of OU's α and w slope are shown in figure 3*A*. The overall model fit

(Fisher's C $= 2.67$, $P_{df=4} = 0.62$) indicated that the model was causally valid. The

median of the posterior distribution of OU's α was mostly explained by the OU's α

prior mode ($p = 0.69$), and to a lesser extent by the real OU's α ($p = 36$) and the real w

slope ($p = 0.2$). The number of species in the metacommunity did not affect the median

of the OU's α posterior distribution ($p < 0.01$). On the other hand, the median of the

posterior distribution of w was mostly explained by the real w slope ($p = 0.85$).

Replacing the median of the posterior distribution of OU's α by half-life (fig. 3*B*) did

not change the overall architecture of causal relationships. The overall model remained

valid (Fisher's C = 1.18, $P_{df=4}$ = 0.88); noteworthy, the strength of the path coefficient

linking the real half-life to the median of the posterior distribution of OU's α increased

(p = 0.42), while the latter was still mostly explained by the OU's α prior mode (p = -

0.74). All other causal relationships behaved similarly to the observed in the previous

model (fig. 3*A*).

Bivariate relationships among real OU's α, half-life and *w* slope and the

corresponding median values of the posterior distribution of OU's α, half-life and *w*

slope are shown in figure 4. In figures 4*A*, *C* and *E*, we observe relationships obtained

when the prior distribution of OU's α was uniform, whereas figures 4*B*, *D* and *F* show

the relationships obtained using the half-life derived prior distribution of OU's α.

Exploring these bivariate relationships suggests that using the uniform prior distribution

of OU's α produce stronger associations between real OU's α or half-life and the

median posterior distribution of OU's α (fig. 4*A*) or half-life (fig. 4*C*), respectively, than

the corresponding relationships based on half-life derived OU's α prior (figs. 4*B*, *D*).

But even using the uniform OU's α prior, the dispersion of the median posterior

distribution of both OU's α and half-life tended to increase towards higher real OU's α,

which should possibly lead to lower reliability of the posterior distribution of the α

parameter under higher real OU's α. Such tendency was minimized when the

association between log-transformed real half-life and median of the posterior

distribution of half-life was considered, as we also found a linear relationship between

those variables when the uniform prior distribution of OU's α was used (fig. 4*C*).

Further, the association between real *w* and the median posterior distribution of *w* slope

was not affected by OU's α prior mode, as expected (figs. 4*E*, *F*).

As the OU's α prior mode was found to be the best predictor of the median

posterior distribution of OU's α, we built the causal models considering only the tests

performed using each OU's α prior mode ('uniform' or 'half-life'). Figures 5*A*, *B* show

models built considering the uniform OU's α prior, while figures 5*C*, *D* show models

built for the half-life OU's α prior. In general, models using the half-life derived OU's α

had worse overall fit than models using the uniform OU's α prior, which had lower

Fisher's C statistics than the former. Within the models using the uniform OU's α prior,

the most adequate scenario was obtained for the scenario considering log-transformed

real half-life explaining the median posterior distribution of half-life, considering the

uniform OU's α prior (p = 0.67; fig. 5*B*). Furthermore, that model showed the lowest

correlation between the median of the posterior distribution of OU's α and *w* slope.

Figure 3: Causal models relating real OU's α or half-life, real *w* slope, OU's α prior mode and number of
species in 2,000 simulated metacommunities on the median of the posterior distribution of OU's α of
half-life and *w* slope. Causal relationships were estimated using ordinary least squares. Half-life and *w*
slopes were log-transformed prior to analysis. Validity of overall models was tested using d-separation
tests (Shipley 2000). *A*) OU's α model (Fisher's C = 2.67, $P_{df=4}$ = 0.62). *B*) Half-life model (Fisher's C =
1.18, $P_{df=4}$ = 0.88). Arrow width indicates the strength of the respective causal relationship.
679

681 **Figure 4:** Bivariate relationships between real niche (OU's α or half-life) or dispersal (*w* slope)
682 parameters of simulated metacommunities and the corresponding median of the posterior distribution of
683 OU's α, half-life and *w* slope. *A, C, E*) Uniform OU's α prior. *B, D, F*) Half-life derived OU's α prior.
684 Dashed lines delimit confidence intervals (95%) of predictions.
685

Figure 5: Causal models relating real OU's α or half-life, real *w* slope and number of species in 1,000 simulated metacommunities on the median of the posterior distribution of OU's α of half-life and *w* slope estimated using uniform (*A, B*) or half-life derived (*C, D*) OU's α prior. Causal relationships were estimated using ordinary least squares. Half-life and *w* slopes were log-transformed prior to analysis. Validity of overall models was tested using d-separation tests (Shipley 2000). *A*) OU's α model, uniform α prior (Fisher's $C = 1.59$, $P_{df=2} = 0.45$). *B*) Half-life model, uniform α prior (Fisher's $C = 0.63$, $P_{df=2} = 0.73$). *C*) OU's α model, half-life α prior (Fisher's $C = 4.13$, $P_{df=2} = 0.13$). *D*) Half-life model, half-life α prior (Fisher's $C = 4.64$, $P_{df=2} = 0.10$). Arrow width indicates the strength of the respective causal relationship.

Phyllostomid bats

ABC analysis of phyllostomid bats generated a prior distribution of OU's α and *w* slope parameters containing 16,040 simulated metacommunities. Both prior and posterior distribution of parameters are shown in figure 6. The posterior distribution of OU's α (fig. 6*A*) indicated a density peak at $\alpha = 0.05$, which means a niche half-life ($t_{1/2} =$) 13.9 Myr, or 41% of the total timespan of the phylogenetic tree of phyllostomids (34 Myr). That is to say that the distribution of entropy across the temperature seasonality gradient showed high phylogenetic signal.

Moreover, the analysis estimated a peak density for *w* slope = 58.52. The minimum spanning tree computed based on the geographic coordinates of sites indicated that the maximum distance necessary to connect all cells in the map was 263 kilometers. Based on eq. 3, at such distance the probability of a species to disperse from

a site to another was very high ($\delta_{lk} > 0.96$, see the subplot within fig. 6*B*); the

probability δ_{lk} only decreases to 0.5 at distances $\cong$ 1,000 kilometers.

Figure 7*A* shows the observed entropy distribution of bat assemblages across the

Neotropic, while figure 7*B* provides mean estimates of entropy computed from those

simulations whose OU's α and *w* slope parameters fell within the boundaries of the

most representative HPD computed for both parameters. As we can see, ABC algorithm

captured most entropy variation across sites. In figure 7*C*, we plotted the residuals of a

cubic model (subplot within fig. 7*C*) relating the observed entropy of sites (y-axis) to

ABC-estimated entropy (x-axis). As we can see, although there is a strong association

between entropies, it is not quite linear, especially at lower and upper limits of entropy

values. Interestingly, the residual plot indicates that in the southernmost regions of the

Neotropic, where entropy values were lower, the simulation framework either under or

overestimated the entropy of bat assemblages (fig. 7*C*).

Figure 6: Histograms and density plots of the posterior distribution of (*A*) OU's α and (*B*) *w* slope
parameters underlying species diversity of phyllostomid bats along a gradient in temperature seasonality
in the Neotropic. Parameters were estimated using ABC analysis implemented in *mcfly* package. Prior
sample size: 16,040 simulations. Posterior sample size: 480 simulations. Only the most representative
highest posterior density intervals (HPD$_1$ 90%) were interpreted.

Figure 7: Species diversity gradients of phyllostomid bats across the Neotropic. (*A*) Observed entropy. The subplot within shows strong negative correlation between entropy and temperature seasonality, with highest entropy values associated with lower temperature seasonality. (*B*) mean entropy estimated using ABC analysis. Mean values were computed for simulations with OU's α and *w* slopes falling within the boundaries of the most representative HPD interval (see fig. 6) computed for the corresponding posterior distribution of each estimated parameter. (*C*) Residual values of the cubic model relating estimated (x-axis) and observed entropy (y-axis) (see subplot within the main plot).

Discussion

The framework proposed in this study and implemented in the package *mcfly* successfully enabled us to infer the most plausible scenario of niche evolution determining biological diversity gradients across spatially-structured environmental gradients. We demonstrated that our model-based simulation approach, coupled to rejection-based Approximate Bayesian Computation (ABC) algorithm (Beaumont 2010; Csilléry et al. 2010), allows to estimate (i) the strength of adaptation rate along the niche evolution of species from a monophyletic lineage and distributed across metacommunities; and (ii) the role played by dispersal limitation on the distribution of biological diversity across space. Considering the initial aims of community phylogenetics proposed by Webb et al. (2002), our approach provides meaningful estimates of the role played by phylogenetic relationships on species assembly patterns, and helps to reconcile niche and neutral processes, a topic heatedly debated over the last decades (Vellend 2016, and references therein). Previous studies have already pointed out the need of integrating niche evolution to diversity gradients (Cadotte et al. 2017). This has been explored with correlative approaches relating phylogenetic and trait diversity (Tucker et al. 2018) or, indirectly, by estimating phylogenetic signal in one or more traits used to compute functional diversity (Muscarella et al. 2016). However, we demonstrated it is possible to link niche evolution models and species diversity gradients without assuming a priori phenotypic traits to define species niches. This releases the need of assuming a given trait (or group of traits) as the most appropriate for assessing niche-environment associations (why not any other[s]?). An advantage of our approach is that it allows the evaluation of datasets for which a complete characterization of niche dimensions is currently not available. Thus, it may fill an important gap in the knowledge on the relationship between phylogenetic relationships among species and ecological diversity.

The analyses performed using simulated datasets demonstrated that the ABC approach implemented in *mcfly* is suitable to estimate the adaptation rate of species niche positions and the degree of dispersal limitation across the assemblages in a metacommunity based on species diversity variation across environmental gradients. Nonetheless, the analytical framework showed to be sensitive to the mode of the prior OU's α distribution (Roos et al. 2015). Therefore, we recommend the use of the uniform distribution of OU's α to avoid biased posterior distributions of OU's α parameter. Causal models detected solid links between real and estimated niche parameters (and

for the dispersal parameter - w slope), either where OU's α or niche half-life was

considered. Thus, even though a formally defined function relating biological diversity

to evolutionary models of niche evolution is not available to date, the ABC approach

implemented in the package *mcfly* proved to be a useful way to estimate the imprint of

niche evolution on diversity gradients (Cadotte et al. 2017). Although evolutionary

analyses usually suffer from incomplete datasets (FitzJohn et al. 2009; Nee et al. 1994;

Pybus and Harvey 2000), our framework demonstrated to be robust even for

metacommunities showing incomplete phylogenetic data (simulated metacommunities

contained at least 40% of total phylogenetic pool), which is quite common in ecological

and biogeographic data. Yet, some caution may be necessary where regional assembly

contains only a small fraction of the species within the phylogenetic tree.

Phyllostomid bats showed strong dependency on niche evolution towards an

optimum of low temperature seasonality across Neotropical assemblages, corroborating

results obtained by Stevens (2011). Phyllostomids have high dispersal capacity up to

distances of *ca.* 1000 km, which is in conformity with the large geographic range of

most of its species (Willig et al. 2003). Moreover, the estimated OU'α of phyllostomid's

niche positions was low (α = 0.05), which confers a relatively high half-life to niche

positions of species (13.9 Myr). Those results suggest that those bats preferably track

habitats offering niche conditions close to their optimum over long distances, instead of

quickly adapting their niches towards an optimum condition (see also Villalobos et al.

2013). That strategy was already hypothesized to explain plant distribution (Donoghue

2008). This finding may explain evidence showing that closely related phyllostomid

species tend to live in geographic proximity (Villalobos et al. 2014).

In general, estimated entropy closely mirrored observed richness in bat

assemblages, although not quite linearly, as revealed by the model relating observed to

estimated entropy. For instance, at higher observed entropy levels in Amazonia region,

simulations tended to slightly underestimate entropy. Certainly, another relevant niche

factors might enable diversity levels to be higher than those expected by climatic niche

alone. Vertical stratification in tropical rainforests, associated with seasonal floods

affecting nutrient and roost availability, is a well-known driver of local phyllostomid

diversity in Amazonian forests (Bernard 2001; Pereira et al. 2010; Pereira et al. 2009).

Some other remarkable deviation patterns were observed in two regions: (1) Southern

Atlantic forests, where entropy was strongly underestimated. Here, phyllostomids are a

small subsample of the diversity of the family, as may be inferred from the IUCN

distribution maps (Rojas et al. 2018a), and are mostly represented by generalist

frugivores who present higher speciation rates than phyllostomids of other feeding

guilds (Rojas et al. 2018b). Such generalist diet would allow them to cope with the

unpredictability of food resources in the subtropical regions. (2) The "dry" diagonal of

South America, where entropy was in general overestimated. Neotropical rainforests

and savannas share a significant part of the generalist frugivore assemblage, but the

absence of dense forests in the more open physiognomies negatively affects the

occurrence of several frugivorous phyllostomids (Stevens 2006), particularly those

specialized in canopy fruits. Therefore, the phyllostomid assemblages of the South

American "dry" diagonal become more represented by obligatory, and specialized,

nectarivores, which show lower rates of speciation than those observed for the

generalist herbivore lineages (Rojas et al. 2018b); this might explain entropy

overestimation. In all cases, a single optimum niche condition might fail to account for

other relevant variables such as vegetation types. Indeed, the current version of *mcfly*

has this limitation, which should be fixed in further developments of the analytical

framework. Yet, in its current version, *mcfly* allows for multiple conditions to be

included as a principal component, and users may thereby incorporate climatic and

landscape data to the analysis. Finally, mismatches between observed and estimated

entropy in some areas may be, to some extent, due to inconsistencies in IUCN maps.

The framework proposed here shed light on the link between niche evolution

and the distribution of biological diversity, and thereby improved our understanding of

evolutionary imprints on ecological patterns. Yet, we believe that the main strength of

this new approach is to open avenues of investigation that may help solving the eco-

evolutionary puzzle. For instance, multiple niche optima, possibly derived from

different niche conditions, and affecting differentially distinct clades, might provide

more realistic scenarios than the single optimum scenario developed here. Further, our

framework deals with evolutionary imprints on ecological patterns, but it does not allow

evaluating how higher ecological mechanisms (sensu Vellend 2016) may ultimately

affect evolutionary dynamics (Weber et al. 2017). Further developments of the

framework proposed here should help filling these (and other) gaps in near future. This

tale is to be continued.

Acknowledgements

LD, MVC, MJRP and JAFDF research activities have been supported by CNPq

Productivity Fellowship (grants 307527/2018-2, 306590/2018-2, 311297/2018-8 and

301799/2016-4, respectively). LD, GN, MVC and JAFDF research have been

conducted in the context of the National Institutes for Science and Technology (INCT)

in Ecology, Evolution and Biodiversity Conservation, supported by MCTIC/CNPq

(proc. 465610/2014-5) and FAPEG (proc. 201810267000023). The PhyloPic

(http://phylopic.org) image of a phyllostomid bat used in figures 6 and 7 was created by

Melissa Ingala, and its use was authorized under the license available at

https://creativecommons.org/licenses/by/3.0/. Figure 1 was brilliantly designed by Clara

Heinrich (linkedin.com/in/clara-heinrich-4bb3a3112).

A ciência brasileira resiste.

References

Ackerly, D. D. 2003. Community assembly, niche conservatism, and adaptive evolution in

852   changing environments. International Journal of Plant Sciences 164:S165-S184.

853   Anand, M., and L. Orlóci. 1996. Complexity in plant communities: the notion and

854   quantification. Journal of Theoretical Biology 179:179-186.

855   Baker, R. J., O. R. P. Bininda-Emonds, H. Mantilla-Meluk, C. A. Porter, and R. A. V. D.

856   Bussche. 2012. Molecular time scale of diversification of feeding strategy and

857   morphology in NewWorld Leaf-Nosed Bats (Phyllostomidae): a phylogenetic

858   perspective, Pages 385–409. Evolutionary history of bats. Cambridge, U.K., Cambridge

859   Univ. Press.

860   Beaumont, M. A. 2010. Approximate Bayesian computation in evolution and ecology. Annual

861   review of ecology, evolution, and systematics 41:379-406.

862   Beaumont, M. A., W. Zhang, and D. J. Balding. 2002. Approximate Bayesian computation in

863   population genetics. Genetics 162:2025-2035.

864   Bell, G. 2000. The distribution of abundance in neutral communities. The American Naturalist

865   155:606-617.

866   Bernard, E. 2001. Vertical stratification of bat communities in primary forests of Central

867   Amazon, Brazil. Journal of Tropical Ecology 17:115-126.

868   Blomberg, S. P., and T. Garland. 2002. Tempo and mode in evolution: phylogenetic inertia,

869   adaptation and comparative methods. Journal of Evolutionary Biology 15:899-910.

870   Borcard, D., and P. Legendre. 2002. All-scale spatial analysis of ecological data by means of

871   principal coordinates of neighbour matrices. Ecological Modelling 153:51-68.

872   Butler, M., and A. King. 2004. Phylogenetic comparative analysis: A modeling approach for

873   adaptive evolution. The American Naturalist 164:683-695.

Cadotte, M. W., R. Dinnage, and D. Tilman. 2012. Phylogenetic diversity promotes ecosystem
stability. Ecology 93:S223-S233.

Cadotte, M. W., T. Jonathan Davies, and P. R. Peres-Neto. 2017. Why phylogenies do not

always predict ecological differences. Ecological Monographs 87:535-551.

Cavender-Bares, J., K. H. Kozak, P. V. A. Fine, and S. W. Kembel. 2009. The merging of
community ecology and phylogenetic biology. Ecology Letters 12:693-715.

Cooper, N., W. Jetz, and R. P. Freckleton. 2010. Phylogenetic comparative approaches for
studying niche conservatism. Journal of Evolutionary Biology 23:2529-2539.

Cooper, N., G. H. Thomas, C. Venditti, A. Meade, and R. P. Freckleton. 2016. A cautionary
note on the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biological
Journal of the Linnean Society 118:64-77.

Crisp, M. D., and L. G. Cook. 2005. Do early branching lineages signify ancestral traits? Trends
in Ecology & Evolution 20:122-128.

Csilléry, K., M. G. Blum, O. E. Gaggiotti, and O. François. 2010. Approximate Bayesian
computation (ABC) in practice. Trends in ecology & evolution 25:410-418.

Del Monte-Luna, P., B. W. Brook, M. J. Zetina-Rejón, and V. H. Cruz-Escalona. 2004. The

carrying capacity of ecosystems. Global ecology and biogeography 13:485-495.

Donoghue, M. J. 2008. A phylogenetic perspective on the distribution of plant diversity.
Proceedings of the National Academy of Sciences 105:11549-11555.

Dray, S., and P. Legendre. 2008. Testing the species traits-environment relationships: the
fourth-corner problem revisited. Ecology 89:3400-3412.

Duarte, L. D. S., V. J. Debastiani, M. B. Carlucci, and J. A. F. Diniz-Filho. 2018. Analyzing
community-weighted trait means across environmental gradients: should phylogeny
stay or should it go? Ecology 99:385–398.

Duarte, L. d. S., R. E. Machado, S. M. Hartz, and V. D. Pillar. 2006. What saplings can tell us
about forest expansion over natural grasslands. Journal of Vegetation Science 17:799-
808.

Duarte, L. D. S., P. V. Prieto, and V. D. Pillar. 2012. Assessing spatial and environmental
drivers of phylogenetic structure in Brazilian Araucaria forests. Ecography 35:952-960.

Elton, C. 1927, Animal Ecology. London, Sedgwick and Jackson.

Enquist, B. J., J. Norberg, S. P. Bonser, C. Violle, C. T. Webb, A. Henderson, L. L. Sloat et al.
2015. Scaling from Traits to Ecosystems: Developing a General Trait Driver Theory via
Integrating Trait-Based and Metabolic Scaling Theories. . Advances in Ecological
Research 52:249–318.

Felsenstein, J. 1985. Phylogenies and the comparative method. American Naturalist 125:1-15.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. International Journal of Climatology.

FitzJohn, R. G., W. P. Maddison, and S. P. Otto. 2009. Estimating trait-dependent speciation

and extinction rates from incompletely resolved phylogenies. Systematic biology

58:595-611.

Fleming, T. H., L. D. Dávalos, and M. A. R. Mello. 2020, Phyllostomidae bats: a unique

mammalian radiation. Chicago, The University of Chicago Press

Garnier, E., M.-L. Navas, and K. Grigulis. 2016, Plant functional diversity: organism traits,

community structure, and ecosystem properties., Oxford University Press.

Gerhold, P., J. F. Cahill, M. Winter, I. V. Bartish, and A. Prinzing. 2015. Phylogenetic patterns

are not proxies of community assembly mechanisms (they are far better). Functional

Ecology 29:600-614.

Gerhold, P., M. B. Carlucci, Ş. Proches, and A. Prinzing. 2018. The deep past controls the

phylogenetic structure of present, local communities. Annual Review of Ecology,

Evolution, and Systematics 49:477-497.

Godoy, O., N. J. B. Kraft, and J. M. Levine. 2014. Phylogenetic relatedness and the

determinants of competitive outcomes. Ecology Letters 17:836-844.

Gould, S. J., and R. C. Lewontin. 1979. The spandrels of San Marco and the Panglossian

paradigm: a critique of the adaptationist programme. Proceedings of the Royal Society

of London. Series B. Biological Sciences 205:581-598.

Gravel, D., C. D. Canham, M. Beaudet, and C. Messier. 2006. Reconciling niche and neutrality:

the continuum hypothesis. Ecology Letters 9:399-409.

Hansen, T. F. 1997. Stabilizing selection and the comparative analysis of adaptation. Evolution

51:1341-1351.

Hansen, T. F., and E. P. Martins. 1996. Translating between microevolutionary process and

macroevolutionary patterns: the correlation structure of interspecific data. Evolution

50:1404-1417.

Hansen, T. F., J. Pienaar, and S. H. Orzack. 2008. A comparative method for studying

adaptation to a randomly evolving environment. Evolution: International Journal of

Organic Evolution 62:1965-1977.

Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER:

investigating evolutionary radiations. Bioinformatics 24:129-131.

Harvey, P. H., and M. D. Pagel. 1991, The comparative method in evolutionary biology.

Oxford, Oxford University Press.

Hubbell, S. P. 2001, The Unified Neutral Theory of Biodiversity and Biogeography. Princeton,

Princeton University Press.

Hutchinson, G. E. 1957. Concluding remarks. Cold Spring Harbor Symposia on Quantitative
Biology 22:415–427.

Hutchinson, G. E. 1959. Homage to Santa Rosalia or why are there so many kinds of animals?
The American Naturalist 93:145-159.

IUCN. 2020. The IUCN Red List of Threatened Species. Version 6.2.

Kraft, N. J. B., P. B. Adler, O. Godoy, E. James, S. Fuller, and J. M. Levine. 2015. Community
assembly, coexistence, and the environmental filtering metaphor. Functional Ecology
29:592-599.

Lefcheck, J. S. 2016. piecewiseSEM: Piecewise structural equation modelling in r for ecology,
evolution, and systematics. Methods in Ecology and Evolution 7:573-579.

Legendre, P., and L. Legendre. 2012, Numerical Ecology. Amsterdam, Elsevier.

Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D.
Holt et al. 2004. The metacommunity concept: a framework for multi-scale community
ecology. Ecology Letters 7:601-613.

Levine, J. M., and J. HilleRisLambers. 2009. The importance of niches for the maintenance of
species diversity. Nature 461:254-257.

Losos, J. B. 2008. Phylogenetic niche conservatism, phylogenetic signal and the relationship
between phylogenetic relatedness and ecological similarity among species. Ecology
Letters 11:995-1003.

Loyola, R. D., P. Lemes, F. T. Brum, D. B. Provete, and L. D. S. Duarte. 2013. Clade-specific
consequences of climate change to amphibians in Atlantic Forest protected areas.
Ecography 37:65-72.

MacArthur, R., and R. Levins. 1967. The limiting similarity, convergence, and divergence of
coexisting species. The American Naturalist 101:377-385.

MacArthur, R. H. 1972, Geographical Ecology: Patterns in the Distribution of Species.
Princeton, Princeton University Press.

Maestri, R., and L. Duarte. 2020. Evolutionary imprints on species distribution patterns across
the Neotropics, Pages 103-119 Neotropical Diversification: Patterns and Processes,
Springer.

Mayfield, M. M., and J. M. Levine. 2010. Opposing effects of competitive exclusion on the
phylogenetic structure of communities. Ecology Letters 13:1085-1093.

McPeek, M. A. 2017, Evolutionary Community Ecology, v. 58, Princeton University Press.

Minchin, P. R. 1987. Simulation of multidimensional community patterns: towards a
comprehensive model. Vegetatio 71:145-156.

Monteiro, L. R., and M. R. Nogueira. 2011. Evolutionary patterns and processes in the radiation
of phyllostomid bats. BMC Evolutionary Biology 11:1.

Morrone, J. J. 2014. Biogeographical regionalisation of the Netropical region. Zootaxa
3782:001-110

Mouquet, N., V. Devictor, C. N. Meynard, F. Munoz, L.-F. Bersier, J. Chave, P. Couteron et al.
2012. Ecophylogenetics: advances and perspectives. Biological Reviews 87:769-785.

Münkemüller, T., F. C. Boucher, W. Thuiller, and S. Lavergne. 2015. Phylogenetic niche
conservatism – common pitfalls and ways forward. Functional Ecology 29:627-639.

Muscarella, R., M. Uriarte, T. M. Aide, D. L. Erickson, J. Forero-Montaña, W. J. Kress, N. G.
Swenson et al. 2016. Functional convergence and phylogenetic divergence during
secondary succession of subtropical wet forests in Puerto Rico. Journal of Vegetation
Science 27:283-294.

Nakamura, G., L. O. Gonçalves, and L. d. S. Duarte. 2020. Revisiting the dimensionality of
biological diversity. Ecography 43:539-548.

Nathan, R., and H. C. Muller-Landau. 2000. Spatial patterns of seed dispersal, their
determinants and consequences for recruitment. Trends in Ecology & Evolution 15:278-
285.

Nee, S., R. M. May, and P. H. Harvey. 1994. The reconstructed evolutionary process.
Philosophical Transactions of the Royal Society of London. Series B: Biological
Sciences 344:305-311.

Nosil, P. 2012, Ecological speciation, Oxford University Press.

Paine, C. T., A. Deasey, and A. B. Duthie. 2018. Towards the general mechanistic prediction of
community dynamics. Functional ecology 32:1681-1692.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35:526-528.

Pereira, M. J. R., J. T. Marques, and J. M. Palmeirim. 2010. Vertical stratification of bat
assemblages in flooded and unflooded Amazonian forests. Current Zoology 56:469-
478.

Pereira, M. J. R., J. T. Marques, J. Santana, C. D. Santos, J. Valsecchi, H. L. De Queiroz, P.
Beja et al. 2009. Structuring of Amazonian bat assemblages: the roles of flooding
patterns and floodwater nutrient load. Journal of Animal Ecology 78:1163-1171.

Pereira, M. J. R., and J. M. Palmeirim. 2013. Latitudinal diversity gradients in New World Bats:
are they a consequence of niche conservatism? Plos One 8:e69245.

Peres-Neto, P. R., S. Dray, and C. J. F. t. Braak. 2017. Linking trait variation to the
environment: critical issues with community-weighted mean correlation resolved by the
fourth-corner approach. Ecography 40:806-816.

Peres-Neto, P. R., M. A. Leibold, and S. Dray. 2012. Assessing the effects of spatial
contingency and environmental filtering on metacommunity phylogenetics. Ecology
93:S14-S30.

Pybus, O. G., and P. H. Harvey. 2000. Testing macro–evolutionary models using incomplete molecular phylogenies. Proceedings of the Royal Society of London. Series B: Biological Sciences 267:2267-2272.

Pyron, R. A., G. C. Costa, M. A. Patten, and F. T. Burbrink. 2015. Phylogenetic niche conservatism and the evolutionary basis of ecological speciation. Biological Reviews 90:1248-1262.

Rényi, A. 1961, On measures of entropy and information Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics.

Revell, L. J., L. J. Harmon, and D. C. Collar. 2008. Phylogenetic signal, evolutionary process, and rate. Systematic Biology 57:591-601.

Ricklefs, R. E., and D. Schluter. 1993. Species diversity in ecological communities: historical and geographical perspectives.

Ricotta, C. 2005. Through the Jungle of Biological Diversity. Acta Biotheoretica 53:29-38.

Rojas, D., M. Moreira, M. J. Ramos Pereira, C. Fonseca, and L. M. Dávalos. 2018a. Updated distribution maps for neotropical bats in the superfamily Noctilionoidea. Ecology 99:2131-2131.

Rojas, D., M. J. Ramos Pereira, C. Fonseca, and L. M. Dávalos. 2018b. Eating down the food chain: generalism is not an evolutionary dead end for herbivores. Ecology Letters 21:402-410.

Roos, M., T. G. Martins, L. Held, and H. Rue. 2015. Sensitivity Analysis for Bayesian Hierarchical Models. Bayesian Analysis 10:321-349, 329.

Rossoni, D. M., B. M. A. Costa, N. P. Giannini, and G. Marroig. 2019. A multiple peak adaptive landscape based on feeding strategies and roosting ecology shaped the evolution of cranial covariance structure and morphological differentiation in phyllostomid bats. Evolution 73:961-981.

Sexton, J. P., J. Montiel, J. E. Shay, M. R. Stephens, and R. A. Slatyer. 2017. Evolution of ecological niche breadth. Annual Review of Ecology, Evolution, and Systematics 48.

Shi, J. J., and D. L. Rabosky. 2015. Speciation dynamics during the global radiation of extant bats. Evolution 69:1528-1545.

Shipley, B. 2000, Cause and correlation in biology: a user's guide to path analysis, structural equations and causal inference, Cambridge Unversity Press.

—. 2010, From plant traits to vegetation structure: chance and selection in the assembly of ecological communities, Cambridge University Press.

Simmons, N. B., and A. L. Cirranello. 2020. Bat Species of the World: A taxonomic and geographic database.

Sokol, E. R., B. L. Brown, and J. Barrett. 2017. A simulation-based approach to understand how

metacommunity characteristics influence emergent biodiversity patterns. Oikos

126:723-737.

Sokol, E. R., B. L. Brown, C. C. Carey, B. M. Tornwall, C. M. Swan, and J. E. Barrett. 2015.

Linking management to biodiversity in built ponds using metacommunity simulations.

Ecological Modelling 296:36-45.

Stevens, R. D. 2006. Historical processes enhance patterns of diversity along latitudinal

gradients. Proceedings. Biological sciences 273:2283-2289.

—. 2011. Relative effects of time for speciation and tropical niche conservatism on the

latitudinal diversity gradient of phyllostomid bats. Proceedings of the Royal Society B:

Biological Sciences 278:2528-2536.

Swenson, N. G. 2019, Phylogenetic Ecology: A History, Critique, and Remodeling, University

of Chicago Press.

Tucker, C. M., T. J. Davies, M. W. Cadotte, and W. D. Pearse. 2018. On the relationship

between phylogenetic diversity and trait diversity. Ecology 99:1473-1479.

Turkkan, N., and T. Pham-Gia. 1993. Computation of the highest posterior density interval in

bayesian analysis. Journal of Statistical Computation and Simulation 44:243-250.

Vellend, M. 2016, The Theory of Ecological Communities, Princeton University Press.

Villalobos, F., and H. T. Arita. 2010. The diversity field of New World leaf-nosed bats

(Phyllostomidae). Global Ecology and Biogeography 19:200-211.

Villalobos, F., A. Lira-Noriega, J. Soberón, and H. T. Arita. 2014. Co-diversity and co-

distribution in phyllostomid bats: Evaluating the relative roles of climate and niche

conservatism. Basic and Applied Ecology 15:85-91.

Villalobos, F., T. F. Rangel, and J. A. F. Diniz-Filho. 2013. Phylogenetic fields of species:

cross-species patterns of phylogenetic structure and geographical coexistence.

Proceedings of the Royal Society B: Biological Sciences 280.

Violle, C., B. J. Enquist, B. J. McGill, L. Jiang, C. H. Albert, C. Hulshof, V. Jung et al. 2012.

The return of the variance: intraspecific variability in community ecology. Trends in

Ecology & Evolution 27:244-252.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007.

Let the concept of trait be functional! . Oikos 116:882–892.

Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and

community ecology. Annual Review of Ecology and Systematics 33:475-505.

Weber, M. G., C. E. Wagner, R. J. Best, L. J. Harmon, and B. Matthews. 2017. Evolution in a

community context: on integrating ecological interactions and macroevolution. Trends

in Ecology & Evolution 32:291–304.

Wiens, J. J., and C. H. Graham. 2005. Niche conservatism: integrating evolution, ecology, and
conservation biology. Annual Review of Ecology, Evolution, and Systematics 36:519-
539.
Willig, M. R., B. D. Patterson, and R. D. Stevens. 2003. Patterns of range size, richness, and
body size in the Chiroptera, Pages 580–621 *in* T. H. Kunz, and M. B. Fenton, eds. Bat
Ecology Chicago, Univ. Chicago Press.
1096

Figure 1: Schematic representation of *mcfly*, an Approximate Bayesian Computation framework developed to estimate the posterior distribution of adaptation rate (OU's α) and dispersal kernel slope (*w*) parameters driving species diversity gradients along an environmental gradient. **1**. The *real metacommunity* provides the empirical measures of species diversity, the *real summary statistic* (*R*). **2**. *Environment* is a dimension of species niches that provides the optimum niche (θ), variance (σ) and root niche position parameters necessary to simulate species niche based on the *phylogeny,* according to Ornstein-Uhlenbeck evolutionary model. We also define niche breadth in the simulation based on *environment*. **3**. A prior distribution of OU's α is used to simulate species niche positions based on *environment* parameters and *phylogeny*. **4**. *Spatial coordinates* of *k* sites are used to define the prior distribution of slope *w*; together with immigration rate (*m*), number of individuals at *k* sites (J_{L_k}) and the total number of individuals in *M* (J_M) provide the neutral parameters of simulations. **5**. Metacommunities are simulated based on niche and neutral parameters (*Simulator*). **6**. Each metacommunity simulation provides species diversity measures (*Summarizer*) taken as the simulated summary statistics (*S*). **7**. Simulated diversities (*S*) are classified based on a pre-defined minimum correlation threshold (tolerance) to the observed diversity (*R*). Every *S* object not discarded (*Reject*) is then *scanned* to record its OU's α and slope *w* to provide a posterior distribution of adaptation rate (α) and dispersal limitation (*w*), respectively.

Figure 2: Relationship between Blomberg's K statistic and the adaptation rate parameter of Ornstein-Uhlenbeck model (OU's α) for simulated niche positions computed for 1,000 sets of 50 species each (black circles). Gray areas indicate lower and upper credibility intervals (95%) for Blomberg's K and OU's α values estimated from white noise vectors ($N[0,1]$).

Figure 3: Causal models relating real OU's α or half-life, real *w* slope, OU's α prior mode and number of species in 2,000 simulated metacommunities on the median of the posterior distribution of OU's α of half-life and *w* slope. Causal relationships were estimated using ordinary least squares. Half-life and *w* slopes were log-transformed prior to analysis. Validity of overall models was tested using d-separation tests (Shipley 2000). *A*) OU's α model (Fisher's C = 2.67, $P_{df=4}$ = 0.62). *B*) Half-life model (Fisher's C = 1.18, $P_{df=4}$ = 0.88). Arrow width indicates the strength of the respective causal relationship.

Figure 4: Bivariate relationships between real niche (OU's α or half-life) or dispersal (*w* slope) parameters of simulated metacommunities and the corresponding median of the posterior distribution of OU's α, half-life and *w* slope. *A, C, E*) Uniform OU's α prior. *B, D, F*) Half-life derived OU's α prior. Dashed lines delimit confidence intervals (95%) of predictions.

Figure 5: Causal models relating real OU's α or half-life, real *w* slope and number of species in 1,000 simulated metacommunities on the median of the posterior distribution of OU's α of half-life and *w* slope estimated using uniform (a, b) or half-life derived (c, d) OU's α prior. Causal relationships were estimated using ordinary least squares. Half-life and *w* slopes were log-transformed prior to analysis. Validity of overall models was tested using d-separation tests (Shipley 2000). *A*) OU's α model, uniform α prior (Fisher's C = 1.59, $P_{df=2}$ = 0.45). *B*) Half-life model, uniform α prior (Fisher's C = 0.63, $P_{df=2}$ = 0.73). *C*) OU's α model, half-life α prior (Fisher's C = 4.13, $P_{df=2}$ = 0.13). *D*) Half-life model, half-life α prior (Fisher's C = 4.64, $P_{df=2}$ = 0.10). Arrow width indicates the strength of the respective causal relationship.

Figure 6: Histograms and density plots of the posterior distribution of (a) OU's α and (b) *w* slope parameters underlying species diversity of phyllostomid bats along a gradient in temperature seasonality in the Neotropic. Parameters were estimated using ABC analysis implemented in *mcfly* package. Prior sample size: 16,040 simulations. Posterior sample size: 480 simulations. Only the most representative highest posterior density intervals (HPD_1 90%) were interpreted.

Figure 7: Species diversity gradients of phyllostomid bats across the Neotropic. (a) Observed entropy. The subplot within shows strong negative correlation between entropy and temperature seasonality, with highest entropy values associated with lower temperature seasonality. (b) mean entropy estimated using ABC analysis. Mean values were computed for simulations with OU's α and *w* slopes falling within the boundaries of the HPD_1 90% (see fig. 6) computed for the corresponding posterior distribution of each estimated parameter. (c) Residual values of the cubic model relating estimated (x-axis) and observed entropy (y-axis) (see subplot within the main plot).