Acetyltransferases GCN5 and PCAF are required for B lymphocyte maturation in mice

Valentyn Oksenych[1,2,3,4]*, Dan Su[1], Jeremy Austin Daniel[1]*

[1] The NNF Center for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, Blegdamsvej 3B, 2200 Copenhagen, Denmark; sudan20072007@yahoo.com

[2] Department for Cancer Research and Molecular Medicine (IKOM), Norwegian University of Science and Technology, Laboratory Center, Erling Skjalgssons gate 1, 7491 Trondheim, Norway; valentyn.oksenych@ntnu.no

[3] Current address : KG Jebsen Centre for B Cell Malignancies, Institute of Clinical Medicine, University of Oslo, N-0316 Oslo, Norway;

[4] Current address : Institute of Clinical Medicine, University of Oslo, 0318 Oslo, Norway; valentyn.oksenych@medisin.uio.no

* Correspondence: Jeremy Austin Daniel, jerodanielphd@gmail.com; and Valentyn Oksenych, valentyn.oksenych@medisin.uio.no

Abstract: B lymphocyte development has two DNA recombination processes: V(D)J recombination of the immunoglobulin (*Igh*) gene variable region and class switching of the *Igh* constant regions from IgM to IgG, IgA, or IgE. V(D)J recombination is required for successful maturation of B cells from pro-B to pre-B to immature-B and then to mature B cells in the bone marrow. CSR occurs outside of the bone marrow when mature B cells migrate to peripheral lymphoid organs, such as spleen and lymph nodes. Both V(D)J recombination and CSR depend on an "open chromatin" state that makes DNA accessible to specific enzymes, recombination activating gene (RAG), and activation-induced cytidine deaminase (AID). Acetyltransferases GCN5 and PCAF possess redundant functions acetylating histone H3 lysine 9 (H3K9). Here, we generated a mouse model that lacks both GCN5 and PCAF in B cells. We found that double-deficient mice possess low levels of mature B cells in the bone marrow and peripheral organs, an accumulation of pro-B cells in bone marrow, and reduced CSR levels. We conclude that both GCN5 and PCAF are required for B cell development *in vivo*.

Keywords: KAT2A; KAT2B; mice; acetyltransferase; B cell; lymphocyte; class switching

29
30

1. Introduction

Development of B lymphocytes starts in the bone marrow where progenitor (pro)-B cells using recombination-activating genes (RAG) generate DNA double-strand breaks (DSBs) and initiate the V(D)J recombination. In maturating B cells, the V(D)J recombination process is genetic recombination of *variable (V)*, *diversity (D)*, and *joining (J)* gene segments arranging into a newly formed VDJ part of immunoglobulin gene (*Ig*) (1-4). Following the V(D)J recombination, B cells develop from pro-B cells expressing specific markers cluster of differentiation 19 (CD19), B220/CD45, and CD43 (CD19+B220+CD43+) to pre-B cells (CD19+B220+CD43-), immature B (CD19+B220+IgM+, low immunoglobulin M, IgM) and mature B (CD19+B220+IgM+, high IgM) cells in bone marrow (1). Mature B lymphocytes leave the bone marrow and through the blood migrate to periphery populating spleen and lymph nodes.

Then, mature B cells initiate another DNA recombination process to change the constant regions of immunoglobulin genes referred to as class switch recombination (CSR). During the CSR in mice, IgM is replaced by IgG3, IgG1, IgG2a, IgG2b, IgE, or IgA (1). The CSR is initiated by non-productive transcription known as a germ-line transcription (GLT)

43
44
45
46

which is needed to separate two DNA strands. Single-stranded DNA is then targeted by 47
activation-induced cytidine deaminase (AID), a B lymphocyte-specific enzyme deaminat- 48
ing cytosine to uracil (C to U). Then uracil DNA N-glycosylase (UNG) removes the uracil 49
from DNA leading to the single-strand break formation (SSB)(5). Two SSBs facing each 50
other then form a DSB and allow recombination (1-4). Both V(D)J and CSR can be regarded 51
as processes following fundamentally similar strategies of genomic recombination (6, 7). 52

The DSBs formed during the V(D)J recombination and CSR are recognized, pro- 53
cessed, and repaired by the non-homologous end-joining pathway (NHEJ) and initiate 54
more complex signaling and chromatin modification pathway known as DNA damage re- 55
sponse (DDR) (1-4). The NHEJ is initiated when Ku70 and Ku80 heterodimer (Ku) recog- 56
nizes and binds the DSBs. Ku serves as a platform for downstream factors including DNA- 57
dependent protein kinase catalytic subunit (DNA-PKcs), X-ray repair cross-complement- 58
ing protein 4 (XRCC4), XRCC4-like factor (XLF), Paralogue of XRCC4 and XLF (PAXX), a 59
modulator of retrovirus infection (MRI), and DNA ligase 4 (LIG4) (2-4). There are addi- 60
tional factors that are sometimes optional for NHEJ, including Artemis with nuclease ac- 61
tivity required for processing of hairpin-sealed DNA ends and overhangs. 62

One type of DDR pathway acts downstream of ataxia telangiectasia mutated (ATM) 63
protein kinase, which is activated by DSBs and then phosphorylates multiple substrates, 64
including NHEJ and DDR factors. ATM phosphorylates histone H2AX, which in turn re- 65
cruits mediator of DNA damage checkpoint 1 (MDC1) and facilitates the accumulation of 66
really interesting new gene (RING) finger motif (RNF) 8 and RNF168 ubiquitin-ligases, 67
and then the p53-binding protein (53BP1). Phosphorylation of H2AX is related to the acet- 68
ylation of histones, including histone H3K9. In particular, histone acetylation relies on the 69
ATM-dependent H2AX phosphorylation and SWI/SNF chromatin remodeling factors (8). 70
Acetylation of histone H3K9 is mediated by GCN5 and PCAF (8, 9). 71

There is a complex genetic interaction associated with the functional redundancy be- 72
tween the NHEJ factors (2, 3), including the following pairs: DNA-PKcs/XLF (4, 10-12), 73
PAXX/XLF (13-19), and MRI/XLF (20, 21). Moreover, there is a genetic interaction between 74
NHEJ and DDR pathway factors, for example, ATM/XLF and H2AX/XLF (22), MDC1/XLF 75
(23), RNF8/XLF and RNF168/XLF (24), 53BP1/XLF (25, 26), and more (2, 3). Moreover, 76
acetyltransferases GCN5 and PCAF are redundant in promoting histone H3 lysine K9 acet- 77
ylation (27). 78

General control non-depressible 5 (GCN5) acetyltransferase is also known as lysine 79
acetyltransferase (KAT) 2A. Germline inactivation of GCN5 in mice resulted in early em- 80
bryonic lethality due to the role of the protein in neurogenesis (27). GCN5 is functionally 81
redundant with another acetyltransferase, KAT2B, also referred to as p300/CBP-associated 82
factor (PCAF). While inactivation of *Pcaf* gene in mice has no detectable phenotype, double 83
knockout of *Gcn5/Pcaf* genes resulted in even earlier embryonic lethality than in *Gcn5*[-/-] 84
mice (27). Because histone H3K9 acetylation works downstream of ATM and H2AX in 85
DDR (8), one could propose that GCN5, PCAF, or both enzymes are required for lympho- 86
cyte development *in vivo*. However, embryonic lethality of *Gcn5*[-/-] and *Gcn5*[-/-]*Pcaf*[-/-] mice 87
(27) challenged the studies. To overcome the obstacle, we developed a complex mouse 88
model, when *Pcaf* gene was germline-inactivated (27), while floxed *Gcn5* gene (28) was 89
conditionally-inactivated in B cell lineages by CRE recombinase expressed under *Cd19* pro- 90
moter (29). To sort out CRE-positive and CRE-negative cells, we used *Rosa26-stop-YFP* 91
knockin which only expressed YFP following the CRE activation (30). 92

Here, we found that GCN5 and PCAF acetyltransferases are functionally redundant 93
during early B cell maturation, while GCN5 is required for robust CSR. 94

2. Materials and Methods 95

2.1. Mice 96

Gcn5[f/f] (28), *Pcaf*[+/-] (27), *Cd19*[Cre+] (29) (# 006785; The Jackson Laboratory, Bar Harbor, 97
ME, USA), *Rosa26-stop-YFP*[+] (30)(# 006148, The Jackson Laboratory, Bar Harbor, ME, 98
USA), and *Aid*[-/-] (31) mice were previously described. Mice used for experiments were 99

between 8-12 weeks of age. All experiments were performed in compliance with the Danish Working Environment Authority, the Danish Animal Experiment Inspectorate, and the Department of Experimental Medicine (University of Copenhagen).

2.2. Flow cytometry

Flow cytometry experiments were performed as we described earlier (20, 23, 32-34). In particular, we used the fluorescent antibodies recognizing the proteins described below. B220 (PE-CF594, FITC, Alexa Fluor 700; all clone RA3-6B2, BD Bioscience, Franklin Lakes, NJ, USA). IgM (PerCP-eFluor 710, APC-eFluor 780, APC, FITC; all clone II/41, eBioscience, Santa Clara, CA, USA). IgG1 (PE, MOPC-21, Biolegend, SanDiego, CA, USA). IgG3 (PE, MG3-35, Biolegend, San Diego, CA, USA). CD3-APC (Biolegends, USA, #100312). CD19 (Alexa Fluor 700, APC eFluor 780, both clone 1D3, eBioscience, Santa Clara, CA, USA). CD43 (APC and PE-Cy7, both clones S7, BD Bioscience, Franklin Lakes, NJ, USA).

2.3. Class switch recombination

The CSR was performed as we described earlier (10, 22, 33-37).

2.4. Western blot

The western blot procedures were performed as we described earlier (23, 32, 33, 38). Briefly, the cells were lysed for 30 minutes on ice in Radioimmunoprecipitation assay (RIPA) buffer (Sigma Aldrich, St. Louis, MO, USA, #R0278) supplemented with cOmplete™ EDTA-free Protease Inhibitor Cocktail (Sigma Aldrich, #11873580001). Proteins were analyzed by 4-12% Bis-Tris NuPAGE gels (Invitrogen, Carlsbad, CA, USA, #NP0322), transferred to PDVF membranes (GE Healthcare, Boston, MA, USA, #GE10600023), and probed with indicated antibodies. Rat anti-AID (1:500, Active Motif, Carlsbad, CA, USA, #39886); mouse anti-GCN5, clone A-11 (1:500, Santa Cruz Biotechnology, Dallas, TX, USA, #sc-365321); rabbit anti-PCAF, clone C14G9 (1:1000, Cell Signaling Technology, Leiden, The Netherlands, #3378); rabbit anti-histone H3 (1:1000, Abcam, Cambridge, UK, #ab1791); rabbit anti-histone H3 acetyl K9 (1:500, Abcam, Cambridge, UK, #ab32129).

2.5. Statistics

We performed statistical analyses with one-way ANOVA using GraphPad Prism 8.0.1.244 (San Diego, CA, USA). In the tests, p values less than 0.05 were defined as significant, i.e. $*p<0.05$; $**p<0.01$; $***p<0.001$; and $****p<0.0001$.

3. Results

3.1. Generation of mice lacking GCN5 and PCAF in B cells

Combined inactivation of *Gcn5* and *Pcaf* genes in mice results in embryonic lethality (27). To overcome this challenge, we designed a complex genetic model when floxed *Gcn5* gene is conditionally inactivated in B cell lineages by CRE enzyme under the *Cd19* promoter (*Cd19^{Cre+}*) (29). To sort out the cells with activated CRE, we used a model with knocked-in yellow fluorescent protein gene (*YFP*) into ROSA-26 locus. The YFP is inactive until CRE removes "STOP" signal (Rosa-26-YFP+) (30). Thus, we obtained *Gcn5^{f/f}Pcaf^{/-} Cd19^{+/Cre}YFP^+* mice and simpler controls. Further in the text, we will skip *Cd19^{+/Cre}* and *YFP+* for simplicity in most of the cases, and will refer to the mice based on the status of *Gcn5* and *Pcaf* genes, i.e. as *Gcn5^{f/f}Pcaf^{/-}*, *Gcn5^{f/f}*, *Pcaf^{/-}* and WT. When the CRE is active and describing sorted B cells, we indicate *Gcn5^{-/-}*, a knockout status of the gene. Lack of GCN5 and PCAF was also validated by western blot (Figure S1) and quantitative PCR.

3.2. Mice lacking GCN5 and PCAF in B cells possess small spleens

We obtained mice with germline inactivation of *Pcaf* gene and conditional inactiva- 153
tion of *Gcn5* in B cells under the *Cd19* promoter (Figure 1). We found that germline inacti- 154
vation of *Pcaf* gene alone has no detectable effect on mouse development, in line with the 155
previous observation (27). Conditional inactivation of *Gcn5* gene in B cells had no visible 156
effect on sizes of WT and *Pcaf*-deficient mice, which were 15 to 19 g on average (p>0.1433) 157
(Figure 1A). However, inactivation of *Gcn5* resulted in smaller spleens in mice (*Gcn5*$^{-/-}$, 54 158
mg), when compared to WT (69 mg) and *Pcaf*$^{/-}$ (72 mg) mice. Combined inactivation of 159
Pcaf and *Gcn5* in B cells resulted in even smaller spleens (*Gcn5*$^{-/-}$*Pcaf*$^{/-}$, 29mg, p<0.0001). 160
Spleens of mice without CRE activity with *Gcn5* gene being floxed and functional 161
(*Gcn5*$^{f/f}$*Pcaf*$^{/-}$, 67 mg), were comparable in size to the ones of WT (Figure 1 B, C). 162

Figure 1. Generation of mice with germline inactivation of *Pcaf* and conditional inactivation of 163
Gcn5 in B cell lineages. (A) Sizes of 8 weeks-old mice of indicated genotype are similar (p>0.1433). 164
(B) Size of spleens from mice of indicated genotypes. WT vs *Pcaf*$^{/-}$, n.s. p=0.9559; WT vs *Gcn5*$^{-/-}$, 165
*p=0.0192; WT vs *Gcn5*$^{-/-}$*Pcaf*$^{/-}$, ****p<0.0001; *Pcaf*$^{/-}$ vs *Gcn5*$^{-/-}$, *p=0.0119; *Pcaf*$^{/-}$ vs *Gcn5*$^{-/-}$*Pcaf*$^{/-}$, 166
****p<0.0001; *Gcn5*$^{-/-}$ vs *Gcn5*$^{-/-}$*Pcaf*$^{/-}$, ***p=0.0008. (C) Example of spleens of indicated genotypes. 167
Gcn5$^{-/-}$ indicates *Cd19*^{Cre}-dependent inactivation of *Gcn5* in B cell lineages. 168

3.3. Mice lacking GCN5 and PCAF in B cells possess delayed B lymphocyte development 169

To detect mature *Gcn5*$^{-/-}$*Pcaf*$^{/-}$ B cells, we identified B220+IgM+ cells in the spleen us- 170
ing flow cytometry (Figure 2 A, B). Inactivation of *Pcaf* gene alone did not affect B cell 171
proportions in the spleen (58%) when compared to WT mice (52%, p=0.4777) (Figure 2A). 172
Inactivation of *Gcn5* alone resulted in an insignificant reduction of mature splenocytes 173
when compared to WT (*Gcn5*$^{-/-}$, 46%, p<0.2532), although *Gcn5*$^{-/-}$*Pcaf*$^{/-}$ mice were found to 174
have a significantly less B cell frequency in the spleen (21%, p<0.0001). Similarly, the num- 175
bers of *Gcn5*$^{-/-}$*Pcaf*$^{/-}$ B splenocytes was the lowest (3,4 million), while the number of *Gcn5*$^{-/-}$ 176
B cells (27 million) was also reduced when compared to *Pcaf*$^{/-}$ (54 million, **p=0.0065) and 177
WT (44 million, *p=0.0214) controls (Figure 2B). 178

Figure 2. Reduced levels of mature B lymphocytes in spleens and blood of 8-12-weeks-old mice of indicated genotypes. (A) Proportions of B220+IgM+ mature B cells in spleen. WT vs *Pcaf*-/-, n.s. p=0.4777; WT vs *Gcn5*-/-, ns, p=0.2532; WT vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Pcaf*-/- vs *Gcn5*-/-, *p=0.0119; *Pcaf*-/- vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Gcn5*-/- vs *Gcn5*-/-*Pcaf*-/-, ***p<0.0001. (B) Summary of experiments as shown in (A), number of cells. WT vs *Pcaf*-/-, n.s., p=0.9212; WT vs *Gcn5*-/-, *p=0.0214; WT vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Pcaf*-/- vs *Gcn5*-/-, **p=0.0053; *Pcaf*-/- vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Gcn5*-/- vs *Gcn5*-/-*Pcaf*-/-, *p<0.0269. (C) Proportions of B220+ B cells and CD3+ T cells in blood. (D) Summary of several experiments from (C) reflecting proportions of B cells in blood. WT vs *Pcaf*-/-, n.s. p=0.9196; WT vs *Gcn5*-/-, n.s., p=0.1204; WT vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Pcaf*-/- vs *Gcn5*-/-, n.s., p=0.5398; *Pcaf*-/- vs *Gcn5*-/-*Pcaf*-/-, ****p<0.0001; *Gcn5*-/- vs *Gcn5*-/-*Pcaf*-/-, ***p<0.0001. *Gcn5*-/- and *Gcn5*-/-*Pcaf*-/- indicate *Cd19*^{Cre}-de-pendent inactivation of *Gcn5* in B cell lineages.

3.4. Inactivation of Gcn5 and Pcaf results in a reduced proportion of B cells in the blood

To detect mature B cells in the blood, we used B220 markers (Figure 2C, D). Inactivation of *Pcaf* alone resulted in 62% of B cells after red blood cells were lysed, which was comparable to WT mice with 65% of B cells in the blood (p=0.92). Inactivation of *Gcn5* gene alone resulted in a modest reduction of B cell proportion to 56% (p=0.12), while combined inactivation of *Gcn5* and *Pcaf* led to even lower B cell levels in blood (22%, p<0.0001). Levels of B cells in blood of control mice without CRE recombinase expression, when *Gcn5* gene was functional (*Gcn5*^{f/f}*Pcaf*-/-) were comparable to WT mice (78%). We conclude that GCN5 and PCAF are both required and functionally redundant for B cell development in mice.

One reason of low *Gcn5*-/-*Pcaf*-/- B cell count in spleen and blood in mice could be cell death following normal development of B cells in bone marrow and migration to periphery. Another option could be blocked or delayed maturation of B cells in bone marrow during the earlier developmental stages. To test the latter possibility, we analyzed B cells in bone marrow of the mice (Figure 3).

3.5. Inactivation of Gcn5 and Pcaf results in accumulation of pro-B cells in bone marrow

To characterize B cell maturation in bone marrow, we followed the expression of B220 (B220+IgM-) and IgM (B220+IgM+) on the lymphocyte surface. Inactivation of *Gcn5* or *Pcaf* resulted in an insignificant decline of B220+IgM+ population (26%, 5-6 million) when compared to WT (32%, 8 million, p=0.44)(Figure 3 A, B). Combined inactivation of *Gcn5* and *Pcaf* resulted in an additional reduction of mature B cells in bone marrow (3 million, 16%) (Figure 3 A, B).

We further focused on B220+IgM- populations by determining the CD43+ (pro-B cells) and CD43- (pre-B cells). Proportion of early stage pro-B cells increased from WT mice

(20% on average) and *Pcaf*[-/-] mice (26%, p=0.78) to *Gcn5*[-/-] (39%, **p=0.0022) to *Gcn5*[-/-]*Pcaf*[-] 215
(55%, p<0.0001) (Figure 3 C, D). 216

We conclude that GCN5 and PCAF are required for the maturation of B cells from 217
the pro-B cell stage to pre-B and later to mature B cells. 218

Figure 3. Delayed development of GCN5/PCAF-deficient B cells in bone marrow. Flow cytometry 219
analyses of developing B lymphocytes in bone marrow of 8-12-week-old mice of indicated geno- 220
types. (A) Examples of B220+IgM- (pro-B and pre-B), B220+IgM+low (immature B) and 221
B220+IgM+high (mature B) cell populations. (B) Summary of several experiments from (A) indicating 222
B220+IgM+ cells. WT vs *Pcaf*[-/-], n.s. p=0.6445; WT vs *Gcn5*[-/-], n.s., p=0.4432; WT vs *Gcn5*[-/-]*Pcaf*[-/-], 223
***p=0.0008; *Pcaf*[-/-] vs *Gcn5*[-/-], n.s., p=0.9997; *Pcaf*[-/-] vs *Gcn5*[-/-]*Pcaf*[-/-], *p<0.0138; *Gcn5*[-/-] vs *Gcn5*[-/-]*Pcaf*[-/-], 224
*p<0.0211 (C) CD43+ (pro-B cells) and CD43- (pre-B cells) gated from B220+IgM- populations in (A). 225
(D) Summary of several experiments detecting B220+IgM-CD43+ (pro-B) cells. . WT vs *Pcaf*[-/-], n.s. 226
p=0.7825; WT vs *Gcn5*[-/-], **p=0.0022; WT vs *Gcn5*[-/-]*Pcaf*[-/-], ****p<0.0001; *Pcaf*[-/-] vs *Gcn5*[-/-], n.s., p=0.0505; 227
Pcaf[-/-] vs *Gcn5*[-/-]*Pcaf*[-/-], ****p<0.0001; *Gcn5*[-/-] vs *Gcn5*[-/-]*Pcaf*[-/-], *p<0.0322 *Gcn5*[-/-] and *Gcn5*[-/-]*Pcaf*[-/-] indicate 228
Cd19[Cre]-dependent inactivation of *Gcn5* in B cell lineages. 229

3.6. GCN5 and PCAF are required for robust class switch recombination 231

The CSR relies on the ATM-dependent DDR (1, 3, 4, 22). Because H3K9 acetylation 232
works downstream of H2AX phosphorylation, and GCN5/PCAF might work downstream 233
of ATM/ATR/DNA-PKcs, we tested if the CSR depends on GCN5 and PCAF (Figure 4). 234
We purified B splenocytes from 8 to 12 weeks old mice and stimulated the CSR from IgM 235
to IgG3 using established protocols (35, 37). We focused on matched pairs of *Gcn5*[f/f] (the 236
functional equivalent of WT cells) and *Gcn5*[-/-], as well as *Gcn5*[f/f]*Pcaf*[-/-] (the functional equiv- 237
alent of *Pcaf*[-/-]) and *Gcn5*[-/-]*Pcaf*[-/-] cells. We used *Aid*[-/-] cells as a CSR-deficient control to detect 238
an experimental background (Figure 4). Inactivation of *Pcaf* alone had no effect on CSR 239
levels (WT vs *Gcn5*[f/f]*Pcaf*[-/-], p>0.96). Contrary, the inactivation of *Gcn5* resulted in a reduc- 240
tion of CSR from about 14% in WT and *Gcn5*[f/f] cells to 6% in *Gcn5*[-/-] cells, *p<0.0008 (Fig- 241
ure 4). Combined deletion of *Pcaf* and *Gcn5* resulted in a similar reduction from 12% in 242
Gcn5[f/f]*Pcaf*[-/-] cells to 6% in *Gcn5*[-/-]*Pcaf*[-/-] cells, **p=0.0080. We conclude that GCN5 is required 243
for robust CSR to IgG3, because additional inactivation of *Pcaf* did not affect CSR levels 244
when compared *Gcn5*[-/-] and *Gcn5*[-/-]*Pcaf*[-/-] B cells, p>0.9999 (Figure 4). 245

4. Discussion

Both GCN5 and PCAF are involved in chromatin modification and DDR response, which made them relevant candidates to facilitate lymphocyte development (8, 9). One challenge was the lack of a relevant *in vivo* model because GCN5 and PCAF have certain redundant functions in acetylating H3K9, and because the germline inactivation of *Gcn5* results in early embryonic lethality in mice (27). Here, we generated and analyzed a complex mouse model which allowed studying of double-deficient *Gcn5⁻/⁻Pcaf⁻/⁻* B cells development *in vivo* and *ex vivo*. We used a germline knockout of *Pcaf* (27), a conditional knockout of *Gcn5^{f/f}*, a knockin of CRE recombinase expressed under the B cell-specific *Cd19* promoter (29), and a knockin of YFP to track the activity of CRE recombinase (30).

For such a complex mouse model (*Gcn5^{f/f}Pcaf⁻/⁻Cd19^{+/cre}Rosa-26-YFP⁺*), multiple controls were used. In one line of the controls, the mice lacking PCAF, having floxed *Gcn5* gene but expressing no CRE recombinase were considered (*Gcn5^{f/f}Pcaf⁻/⁻Rosa-26-YFP⁺*, Figure S2). The GCN5-deficient and GCN5/PCAF double-deficient B cells possessed developmental delay with lower levels of mature B cells in spleen and blood, and accumulation of progenitor B cells in bone marrow (Figures 1-4). Contrary, the control mice without CRE expression demonstrated WT levels of B cell development in all the groups, i.e. WT levels of B220+IgM+ mature B cells in the spleen (Figure S2 A and B) and blood (Figure 2S C and D). In addition, these mice possessed high levels of B220+IgM+ cells (Figure S2 E, F), as well as stable and low levels of pro-B cells (B220+IgM-CD43+) in bone marrow (Figure S2, G and H).

The mice lacking PCAF and with conditional knockout of *Gcn5* in B cells were alive and resembled WT littermates (Figure 1A). One clear feature the *Gcn5^{f/f}Pcaf⁻/⁻Cd19^{+/cre}Rosa-26-YFP⁺* mice had was a small spleen (Figure 1 B and C), which was also the case in mice lacking only *Gcn5* in B cells. The small spleen could indicate a defect in B cell development, and we indeed found low numbers of mature B cells in the spleen, blood, and bone marrow. One could propose that mature B cells lacking GCN5 or both GCN5 and PCAF possess low proliferation speed or tend to trigger apoptosis. Alternatively, GCN5 and PCAF might be required for the V(D)J recombination. This option could be tested by, for example, using vAbl pre-B cell lines as we and others did before, e.g. (3, 10, 13, 18, 22-26). Another intriguing question is whether the physical presence or enzymatic activity of GCN5 and PCAF are required for the observed phenotypes, i.e. abrogated B cell maturation and reduced levels of CSR. To investigate this question one could use specific inhibitors of GCN5 and PCAF enzymes, or enzyme-dead mutations introduced to the *Gcn5* and *Pcaf* genes.

Inactivation of *Gcn5* in murine B cells also resulted in reduced lymphomagenesis in mice overexpressing MYC oncoprotein (39). Our findings further highlight this observation, suggesting that GCN5, and potentially also PCAF, enzymes are attractive targets for cancer therapy (39).

The CSR levels were reduced in B cells lacking GCN5 (Figure 4). The challenge in this set of experiments was that the mice of *Gcn5^f/f^Pcaf^-/-^Cd19^+/cre^Rosa-26-YFP^+* genotype were rather rare and possessed a very low number of suitable B splenocytes (Figures 1 and 2). Although our data on IgG3 is sufficient, one could extend the study in the future by generating knockout cell lines lacking GCN5 and PCAF and suitable for CSR. One possible model system is CH12F3 cells capable to support CSR to IgA (40), which were used in the past for this kind of experiment (19, 32, 37). The CSR itself is a complex multistage process. Generating relevant cell lines will also provide tools to determine specific stages of CSR affected in GCN5-deficient mice, i.e. germline transcription, AID recruitment, generation of DSBs, or DNA repair.

5. Conclusions

Acetyltransferases GCN5 and PCAF possess redundant functions in B cell maturation. GCN5 is required for robust class switch recombination *ex vivo*.

Author Contributions: Conceptualization, methodology, validation, formal analysis, investigation, resources, data curation – J.A.D., D.S., and V.O.; writing—original draft preparation, V.O.; writing—review and editing, V.O. and J.A.D.; supervision, J.A.D.; project administration, J.A.D. and V.O.; funding acquisition, J.A.D. and V.O. All authors have read and agreed to the published version of the manuscript.

Funding: This research was mainly supported by the Novo Nordisk Foundation Grant NNF14CC0001 to J.A.D. In addition, V.O. was supported by The Research Council of Norway (#249774, #270491 and #291217); the Norwegian Cancer Society (#182355); The Health Authority of Central Norway (#13477 and #38811); and The Outstanding Academic Fellow Program at NTNU 2017-2021. The final part of the work was supported by the Stiftelsen Kristian Gerhard Jebsen (grant no. SKGJ-MED-019).

Acknowledgements. *Gcn5*-flox and *Pcaf*-null mice were kindly provided by the Sharon Dent laboratory.

Institutional Review Board Statement: All experiments were performed in compliance with the Danish Working Environment Authority, the Danish Animal Experiment Inspectorate, and the Department of Experimental Medicine (University of Copenhagen).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. F. W. Alt, Y. Zhang, F. L. Meng, C. Guo, B. Schwer, Mechanisms of programmed DNA lesions and genomic instability in the immune system. *Cell* **152**, 417-429 (2013).

2. X. S. Wang, B. J. Lee, S. Zha, The recent advances in non-homologous end-joining through the lens of lymphocyte development. *DNA Repair (Amst)* **94**, 102874 (2020).

3. S. Castaneda-Zegarra *et al.*, Genetic interaction between the non-homologous end-joining factors during B and T lymphocyte development: In vivo mouse models. *Scand J Immunol* **92**, e12936 (2020).

4. V. Kumar, F. W. Alt, V. Oksenych, Functional overlaps between XLF and the ATM-dependent DNA double strand break response. *DNA Repair (Amst)* **16**, 11-22 (2014).

5. D. L. Bordin, L. Lirussi, H. Nilsen, Cellular response to endogenous DNA damage: DNA base modifications in gene expression regulation. *DNA Repair (Amst)* **99**, 103051 (2021).

6. X. Zhang *et al.*, Fundamental roles of chromatin loop extrusion in antibody class switching. *Nature* **575**, 385-389 (2019).

7. Y. Zhang *et al.*, The fundamental role of chromatin loop extrusion in physiological V(D)J recombination. *Nature* **573**, 600-604 (2019).

8. H. S. Lee, J. H. Park, S. J. Kim, S. J. Kwon, J. Kwon, A cooperative activation loop among SWI/SNF, gamma-H2AX and H3 acetylation for DNA double-strand break repair. *EMBO J* **29**, 1434-1445 (2010).

9. Q. Jin *et al.*, Distinct roles of GCN5/PCAF-mediated H3K9ac and CBP/p300-mediated H3K18/27ac in nuclear receptor transactivation. *EMBO J* **30**, 249-262 (2011).

10. V. Oksenych *et al.*, Functional redundancy between the XLF and DNA-PKcs DNA repair factors in V(D)J recombination and nonhomologous DNA end joining. *Proc Natl Acad Sci U S A* **110**, 2234-2239 (2013).

11. M. Xing, M. Bjoras, J. A. Daniel, F. W. Alt, V. Oksenych, Synthetic lethality between murine DNA repair factors XLF and DNA-PKcs is rescued by inactivation of Ku70. *DNA Repair (Amst)* **57**, 133-138 (2017).

12. M. Xing, V. Oksenych, Genetic interaction between DNA repair factors PAXX, XLF, XRCC4 and DNA-PKcs in human cells. *FEBS Open Bio* **9**, 1315-1326 (2019).

13. V. Abramowski *et al.*, PAXX and Xlf interplay revealed by impaired CNS development and immunodeficiency of double KO mice. *Cell Death Differ* **25**, 444-452 (2018).

14. G. Balmus *et al.*, Synthetic lethality between PAXX and XLF in mammalian development. *Genes Dev* **30**, 2152-2157 (2016).

15. S. Castaneda-Zegarra, M. Xing, R. Gago-Fuentes, S. Saeterstad, V. Oksenych, Synthetic lethality between DNA repair factors Xlf and Paxx is rescued by inactivation of Trp53. *DNA Repair (Amst)* **73**, 164-169 (2019).

16. C. Lescale *et al.*, Specific Roles of XRCC4 Paralogs PAXX and XLF during V(D)J Recombination. *Cell Rep* **16**, 2967-2979 (2016).

17. X. Liu, Z. Shao, W. Jiang, B. J. Lee, S. Zha, PAXX promotes KU accumulation at DNA breaks and is essential for end-joining in XLF-deficient mice. *Nat Commun* **8**, 13816 (2017).

18. P. J. Hung *et al.*, Deficiency of XLF and PAXX prevents DNA double-strand break repair by non-homologous end joining in lymphocytes. *Cell Cycle* **16**, 286-295 (2017).

19. V. Kumar, F. W. Alt, R. L. Frock, PAXX and XLF DNA repair factors are functionally redundant in joining DNA breaks in a G1-arrested progenitor B-cell line. *Proc Natl Acad Sci U S A* **113**, 10619-10624 (2016).

20. S. Castaneda-Zegarra *et al.*, Leaky severe combined immunodeficiency in mice lacking non-homologous end joining factors XLF and MRI. *Aging (Albany NY)* **12**, 23578-23597 (2020).

21. P. J. Hung *et al.*, MRI Is a DNA Damage Response Adaptor during Classical Non-homologous End Joining. *Mol Cell* **71**, 332-342 e338 (2018).

22. S. Zha *et al.*, ATM damage response and XLF repair factor are functionally redundant in joining DNA breaks. *Nature* **469**, 250-254 (2011).

23. C. Beck, S. Castaneda-Zegarra, C. Huse, M. Xing, V. Oksenych, Mediator of DNA Damage Checkpoint Protein 1 Facilitates V(D)J Recombination in Cells Lacking DNA Repair Factor XLF. *Biomolecules* **10** (2019).

24. B. R. Chen *et al.*, The RNF8 and RNF168 Ubiquitin Ligases Regulate Pro- and Anti-Resection Activities at Broken DNA Ends During Non-Homologous End Joining. *DNA Repair (Amst)* **108**, 103217 (2021).

25. X. Liu *et al.*, Overlapping functions between XLF repair protein and 53BP1 DNA damage response factor in end joining and lymphocyte development. *Proc Natl Acad Sci U S A* **109**, 3903-3908 (2012).

26. V. Oksenych *et al.*, Functional redundancy between repair factor XLF and damage response mediator 53BP1 in V(D)J recombination and DNA repair. *Proc Natl Acad Sci U S A* **109**, 2455-2460 (2012).

27. W. Xu *et al.*, Loss of Gcn5l2 leads to increased apoptosis and mesodermal defects during mouse development. *Nat Genet* **26**, 229-232 (2000).

28. W. Lin *et al.*, Developmental potential of Gcn5(-/-) embryonic stem cells in vivo and in vitro. *Dev Dyn* **236**, 1547-1557 (2007).

29. R. C. Rickert, J. Roes, K. Rajewsky, B lymphocyte-specific, Cre-mediated mutagenesis in mice. *Nucleic Acids Res* **25**, 1317-1318 (1997).

30. S. Srinivas *et al.*, Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol* **1**, 4 (2001).

31. M. Muramatsu *et al.*, Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* **102**, 553-563 (2000).

32. A. Dewan *et al.*, Robust DNA repair in PAXX-deficient mammalian cells. *FEBS Open Bio* **8**, 442-448 (2018).

33. R. Gago-Fuentes *et al.*, Normal development of mice lacking PAXX, the paralogue of XRCC4 and XLF. *FEBS Open Bio* **8**, 426-434 (2018).

34. S. Castaneda-Zegarra *et al.*, Generation of a Mouse Model Lacking the Non-Homologous End-Joining Factor Mri/Cyren. *Biomolecules* **9** (2019).

35. L. M. Starnes *et al.*, A PTIP-PA1 subcomplex promotes transcription for IgH class switching independently from the associated MLL3/MLL4 methyltransferase complex. *Genes Dev* **30**, 149-163 (2016).

36. D. Su *et al.*, PTIP chromatin regulator controls development and activation of B cell subsets to license humoral immunity in mice. *Proc Natl Acad Sci U S A* **114**, E9328-E9337 (2017).

37. C. Boboila *et al.*, Robust chromosomal DNA repair via alternative end-joining in the absence of X-ray repair cross-complementing protein 1 (XRCC1). *Proc Natl Acad Sci U S A* **109**, 2473-2478 (2012).

38. R. Gago-Fuentes, V. Oksenych, Non-Homologous End Joining Factors XLF, PAXX and DNA-PKcs Maintain the Neural Stem and Progenitor Cell Population. *Biomolecules* **11** (2020).

39. A. T. Farria *et al.*, Transcriptional Activation of MYC-Induced Genes by GCN5 Promotes B-cell Lymphomagenesis. *Cancer Res* **80**, 5543-5553 (2020).

40. M. Nakamura *et al.*, High frequency class switching of an IgM+ B lymphoma clone CH12F3 to IgA+ cells. *Int Immunol* **8**, 193-201 (1996).