Stressed out underground?: Transcriptomics illuminates the genome-wide

response to heat stress in surface and subterranean diving beetles

Perry G. Beasley-Hall[1,2,3]* | Terry Bertozzi[1] | Tessa M. Bradford[1] | Charles S. P. Foster[4] |

Karl Jones[1,2] | Simon M. Tierney[5] | William F. Humphreys[6,7] | Andrew D. Austin[1] | Steven

J. B. Cooper[1,2]

[1]School of Biological Sciences, Australian Centre for Evolutionary Biology and Biodiversity,

The University of Adelaide, Adelaide, South Australia 5000, Australia

[2]South Australian Museum, Adelaide, South Australia 5000, Australia

[3]School of Life and Environmental Sciences, University of Sydney, Sydney, New South

Wales 2006, Australia

[4]University of New South Wales, Sydney, New South Wales 2052, Australia

[5]Hawkesbury Institute for the Environment, Western Sydney University, Building R2 Bourke

Street, Richmond NSW 2753, Australia. ORCiD: 0000-0002-8812-6753

[6]Western Australian Museum, Locked Bag 40, Welshpool DC, WA 6986, Australia

[7]School of Biological Sciences, University of Western Australia, Crawley, WA 6009, Australia

*Correspondence

Perry Beasley-Hall, School of Biological Sciences, Darling Building, The University of

Adelaide, Adelaide, South Australia 5000, Australia

Email: perry.beasley-hall@adelaide.edu.au

Abstract

Subterranean habitats are environmentally stable with respect to temperature, humidity, and the absence of light. The transition to a subterranean lifestyle might therefore be expected to cause considerable shifts in an organism's physiology; here, we investigate how subterranean colonisation affects thermal tolerance. Subterranean organisms might be at an increased risk of decline in the face of global temperature rises, but robust data on the fauna is lacking, particularly at the molecular level. In this study we compare the heat shock response of two species of diving beetle in the genus *Paroster*: one surface-dwelling (*P. nigroadumbratus*), the other restricted to a single aquifer (*P. macrosturtensis*). *P. macrosturtensis* has been previously established as having a lower thermal tolerance compared to surface-dwelling relatives, but the genomic basis of this difference is unknown. By sequencing transcriptomes of experimentally heat-shocked individuals and performing differential expression analysis, we demonstrate both species can mount a heat shock response at high temperatures (35C), in agreement with past survival experiments. However, the genes involved in these responses differ between species, and far greater genes are differentially expressed in the surface species*,* which may explain its more robust response to heat stress. In contrast, the subterranean species significantly upregulated the heat shock protein gene *Hsp68* in the experimental setup under conditions it would likely encounter in nature (25C), suggesting it may be more sensitive to ambient stressors, e.g. handling. The results presented here contribute to an emerging narrative concerning weakened thermal tolerances in obligate subterranean organisms at the molecular level.

KEYWORDS: heat shock response, heat shock proteins, subterranean habitats, molecular evolution, transcriptomics, differential gene expression

1 | INTRODUCTION

The transition to an obligate subterranean lifestyle can cause massive shifts in a species'
biology, from the acquisition of classic troglomorphies such as the elongation of appendages
for sensing in an aphotic environment (Culver & Pipan, 2014) to changes in lesser-studied
traits including circadian rhythm (Beale et al., 2016; Merritt & Clarke, 2011), reproductive
biology (Walker & Rose, 1998), respiration (Jones *et al.* 2019) and chemosensation (Balart-
García et al., 2021; J. Yang et al., 2016). These changes have been attributed to the stark
difference between subterranean and surface (hereafter epigean) habitats. While epigean
environments can vary immensely over both time and space, subterranean environments,
such as cave systems, generally possess high environmental stability with respect to light
levels, temperature, humidity, and nutrient availability (Badino, 2010). Animals adapted to
these environments might therefore be particularly well suited for the assessment of
responses to future climate change scenarios, particularly with respect to their thermal
tolerances and responses to increasing temperatures. Indeed, such habitats have been labelled
as undervalued "laboratories for studies of global change biology" (Mammola et al., 2019).

A near-universal response to temperature-induced stress across the tree of life–and
therefore a method by which thermal tolerance can be gauged–is the heat shock response
(hereafter HSR), which involves the synthesis of heat-shock proteins (HSPs). HSPs are
expressed at base levels under non-stressful conditions and include those that are
constitutively expressed (heat-shock cognates) - only induced when organisms are exposed to
thermal extremes, during which they assist in stabilising and refolding proteins at risk of
denaturation. An inducible HSR has been observed in almost all organisms studied to date,
and the proteins involved in this response, as well as the response itself, are highly conserved
among different domains of life (Jindal et al., 1989). An estimated 50 to 200 genes are
involved in the HSR, the most significantly induced of which are HSPs (Richter et al., 2010).

However, there are exceptions to this rule: a lack of an inducible HSR has been documented

in a wide range of species, largely those that occupy very stable thermal environments such

as Antarctic marine habitats (Somero, 2020; Tomanek, 2010).

Knowledge of the HSR in organisms from thermally stable *subterranean* habitats,

including their associated inducible HSPs and at which temperatures this response might be

activated, is almost non-existent (e.g. Colson-Proch *et al.*, 2010). The vast majority of the

literature on this topic is also devoted to vertebrates, in stark contrast to their relatively small

contribution to the biodiversity of subterranean habitats, which is dominated by invertebrate

taxa (Gibert & Deharveng, 2002; Humphreys, 2012). Moreover, a better understanding of

responses to current climate change predictions for subterranean animals has been identified

as a fundamental question in subterranean biology given emerging conservation issues

associated with their respective ecosystems (Mammola et al., 2020). To address this

knowledge gap, here we make use of genomic data from Australian representatives of a

group of aquatic invertebrates containing both epigean and subterranean lineages.

The Yilgarn Craton in central Western Australia houses a diverse subterranean diving

beetle fauna belonging to two tribes, Bidessini and Hydroporini (Dytiscidae). While epigean

species can be found practically continent-wide, subterranean taxa are isolated in calcrete

aquifers (hereafter calcretes) associated with ancient palaeodrainage systems in the region.

These calcretes are completely devoid of light and animals contained within them are

assumed to have little to no access to air above the water's surface (Jones et al., 2019). Each

calcrete houses between one and three dytiscid species, but at the time of writing only around

a quarter of the ~200 known calcretes have been sampled (Cooper et al., 2002; Leijs et al.,

2012; Watts & Humphreys, 2009). Nonetheless, lineages in both tribes are known to have

made independent yet parallel, repeated transitions into underground habitats from epigean

ancestors during the late Miocene to early Pleistocene, likely in response to continental

aridification (Langille et al., 2021; Leijs et al., 2012; Leys et al., 2003; Toussaint et al., 2015).

In each case these transitions have involved the loss of eyes, pigment, and wings (Cooper *et*

al. 2002) as well as the gain of a remarkable ability to respire directly from water (Jones *et al.*

2019). Preliminary evidence suggests that subterranean members of these lineages are less

tolerant of thermal extremes compared to epigean relatives (Jones et al., 2021), but the

molecular mechanisms underlying such tolerances–and genomic changes associated with a

subterranean transition in these animals more broadly–remain unknown.

In the present study, we focus on two members of the Hydroporini: *Paroster*

nigroadumbratus (Clark), an epigean species endemic to South Australia, and the

subterranean *Paroster macrosturtensis* (Watts & Humphreys) found exclusively in a single

calcrete (hereafter the Sturt Meadows calcrete) in the Yilgarn region of Western Australia. As

subterranean dytiscids likely descended from only a handful of epigean lineages (Leijs et al.,

2012), meaningful comparisons can be made between these taxa despite their distributions

being geographically disjunct. A recent study showed that *P. macrosturtensis* has a reduced

upper thermal tolerance (or critical thermal maximum, CT_{max}) of 38.3ºC compared to other

epigean dytiscids (42–44.5ºC) (Jones et al., 2021), mirroring previous results for other cave

beetle species (Pallarés et al., 2020). These results also suggest *P. macrosturtensis* is unlikely

to reach its thermal critical maximum under current climate change predictions. However,

given the thermal stability in its environment it remains unknown as to whether *P.*

macrosturtensis might have a modified HSR compared to its epigean relatives. Building on

this previous work, here we present transcriptomic data from individuals of *P.*

macrosturtensis and *P. nigroadumbratus* previously subjected to varying degrees of heat

shock by Jones *et al*. (2021). In the present study we specifically aim 1) to generate a high-

quality, near-complete reference transcriptome for *P. nigroadumbratus* and 2) using this

dataset, characterise and compare the HSR of *P. nigroadumbratus* and *P. macrosturtensis*,

specifically with respect to which genes are differentially expressed and the conditions under

which this occurs.

2 | METHODS

2.1 | Taxon sampling and experimental design

Beetle specimens were sourced from a previous study by Jones *et al*. (2021) that subjected

specimens to heat stress. Briefly, *P. macrosturtensis* samples (n = 11) were collected from the

Sturt Meadows calcrete in WA via existing mineral exploration boreholes. Epigean *P.*

nigroadumbratus specimens were collected from water bodies in the Mount Crawford region

of South Australia (n = 18). Individuals were kept by Jones *et al*. (2021) in constant

temperature cabinets at between 24 and 26ºC either in a 12:12h light-dark cycle (epigean

species) or in complete darkness (subterranean species) for at least seven days prior to

subsequent experimentation to minimise the effects of variable thermal history and stress

during capture and transport. Beetles were then either retained in this constant temperature

environment and euthanized in liquid N_2 (control treatment) or moved to a temperature-

controlled water bath containing pure reverse osmosis (RO) water (for epigean species) or

RO water with a salinity level mimicking calcrete microhabitats (for subterranean species).

Beetles were kept at 25ºC in this environment for 4.5 hours to determine if they could survive

the experimental setup (25ºC treatment) or allowed to acclimate to 25ºC for 1 hour before the

temperature was increased by 0.05ºC every minute until it reached 35ºC (35ºC treatment). In

all cases individuals were euthanized in liquid N_2 immediately after this time period.

2.2 | cDNA synthesis and sequencing

RNA extractions of control and heat-treated individuals were performed using single whole

bodies per treatment. Extractions were performed using an AllPrep DNA/RNA Micro kit

(Qiagen, Chadstone, VIC), initially homogenizing in lysis buffer containing β-

mercaptoethanol plus a 5 mm steel bead on a Mixer Mill (Retsch, Haan, Germany). Lysates

were processed following the manufacturer's instructions and RNA was eluted from the

RNeasy MinElute spin column in 25 µl RNAse-free water. Concentration of the RNA extract

was determined using a Quantus Flourometer (Promega, Madison, Wisconsin, US) and RNA

integrity was analysed on a TapeStation (Agilent Technologies, Santa Clara, California, US).

Library preparation was performed using a Nugen Universal Plus mRNA-Seq Library

Preparation Kit (Tecan, Männedorf Switzerland), following manufacturer's instructions. All

the barcoded cDNA samples were pooled and 100 bp paired-end sequencing was outsourced

to the Australian Genome Research Facility (AGRF) on an Illumina NovaSeq 6000 S4 lane.

The reference transcriptome for *P. nigroadumbratus* presented here was derived from

previously obtained raw RNA-seq data (Tierney et al., 2015).

2.3 | Quality control of sequence data

Trim Galore v.0.4.1 (bioinformatics.babraham.ac.uk/projects/trim_galore), a wrapper that

implements Cutadapt (Martin, 2011) and FastQC

(bioinformatics.babraham.ac.uk/projects/fastqc), was used for quality control of our raw

reads and the removal of residual sequencing adapters, with default settings for paired

Illumina reads. This protocol was applied to both the raw reads from control/heat-treated

samples and those intended for reference transcriptome assembly.

2.4 | Assembly of *Paroster nigroadumbratus* reference transcriptome

Following quality control and adapter removal, Trinity v.2.5.1 (Grabherr et al., 2011) was

used for *de novo* transcriptome assembly of our raw reads using default settings. Bowtie

v.1.1.2 (Langmead et al., 2009) was used to examine RNA-Seq read representation in the

assembly by aligning raw reads to the assembled transcriptome. Reference transcriptome

completeness was validated using the gVolante implementation of BUSCO v.5 (Nishimura et

al., 2017; Simão et al., 2015) with default settings, searching against the Arthropoda

database.

Functional annotation of the transcriptome was performed using TransDecoder v.5.5.0

(Haas et al., 2013) to predict coding regions in transcripts, Trinotate v.3.2.1 (Grabherr et al.,

2011) and SQLite (http://sqlite.org/index.html) for database integration, and BLAST v.2.7.1

(Camacho et al., 2009) and HMMER v.3.2 (http://hmmer.org) for homology searches and the

identification of protein domains. BLASTX and BLASTP searches were conducted against

the Swiss-Prot database v.2021_02 (The UniProt Consortium, 2021) for Trinity transcripts

and TransDecoder-predicted proteins, respectively. These searches were performed with an e-

value of 1-e6 to capture more distantly matching proteins and -*max_hsps* and -

max_target_seqs both set to 1 to limit our results to the best hit per query by e-value. While

the use of the latter parameter has been subject to scrutiny as top hits are selected based on

sequence order in the database in the case of equivalent matches, this behaviour appears to

occur in datasets with an extremely high proportion of gaps (Madden et al., 2019; Shah et al.,

2019), which does not apply to the present study. The Pfam database v.34.0 (Mistry et al.,

2021) was used in HMMER searches to identify protein domains with default settings.

Finally, we subsetted the ORFs inferred by TransDecoder to only include the longest

isoform per gene. These transcripts were used in downstream analyses to reduce redundancy

and allowed us to perform read quantification and differential expression analysis at the

"gene" level as opposed to that of isoforms following Balart-García *et al.* (2021).

2.5 | Differential expression analysis and Gene Ontology enrichment

analysis

Salmon v.1.1.0 (Patro et al., 2017) was used to index our annotated *P. nigroadumbratus*

reference transcriptome and quasi-map our raw reads from control/heat-treated samples

against it with an automatic inference of library type and 100 bootstrap replicates. Reads

were normalised with the trimmed mean of M-values normalization method (TMM). TMM

read counts for each replicate were visualised using principal components analysis consisting

of a total of 13 *P. nigroadumbratus* replicates and 11 for *P. macrosturtensis* in our

downstream differential expression analysis (Figure S1).

The Trinity helper scripts for edgeR v.3.32.1 (Robinson et al., 2010) were used for

differential expression analyses and enrichment of Gene Ontology (hereafter GO) terms

based on annotations from the *P. nigroadumbratus* reference transcriptome. Abundance

estimates from Salmon were first transformed for use in edgeR prior to running differential

expression analysis with default settings (*abundance_estimates_to_matrix.pl,*

run_DE_analysis.pl). Differential expression analysis results were then analysed with a *p*-

value cut-off for FDR of 0.05 and a fold-change of 1. This analysis included the

enrichment/depletion of GO terms by supplying Trinity gene lengths and GO terms for each

annotation (Young et al., 2010) and compared 25ºC and 35ºC treatments against individuals

of the same species in the control treatment.

Significantly up- and downregulated genes shared between different treatments were

identified using a custom Venn diagram webtool

(bioinformatics.psb.ugent.be/webtools/Venn) and visualised with UpSetR v.1.4.0 (Conway et

al., 2017). Enrichment of GO terms was visualised with REVIGO (Supek et al., 2011), which

summarises terms by clustering them and retrieving a meaningful subset using similarity

measures. Enriched GO terms were supplied alongside FDR values associated with

enrichment/depletion and the REVIGO analysis was performed using an allowed similarity of

0.9. Resulting tree maps were produced with values set to abs(log10(p-value)) to ensure plots

reflected the significance of enrichment of each term.

2.6 | Phylogenetic analysis of putative heat shock proteins

Genes annotated as putatively encoding heat shock proteins consisted of members of three of

the four insect families of HSP, including small heat shock proteins (sHSP), HSP70, and

HSP90. Transcripts were extracted on the basis of annotations containing the terms *heat*

shock, *HSC*, *HSP*, or *chaperone*. To ensure the inclusion of as many genes as possible for

which annotation might have been incomplete (e.g. certain HSP chaperones), we also

included transcripts putatively involved in the HSR that were differentially expressed in our

dataset yet not captured in the above search. The resulting candidate set of 38 transcripts was

checked against the UniProt database to confirm the function of the proteins they encoded,

and the identities of transcripts were then validated using BLASTN searches restricted to the

Coleoptera against the NCBI nr database. These transcripts were used as a reference to filter

raw reads from each *P. macrosturtensis* 35ºC replicate using BWA v.0.7.17 (H. Li & Durbin,

2009) and SAMtools v.1.9 (H. Li et al., 2009). Filtered *P. macrosturtensis* reads (excluding

singletons) were then pooled and mapped against reference transcripts using the Read

Mapper tool in Geneious Prime (v.2021.2.2, geneious.com) with Low Sensitivity presets.

Consensus sequences for *P. macrosturtensis* were produced from these alignments with a 0%

(majority) threshold.

Coleopteran orthologs of the above genes were retrieved from OrthoDB v.10.1

(Kriventseva et al., 2019) with LOC identifiers of the top BLASTN hit as queries. mRNA

transcripts of protein sequences in question were obtained from GenBank, representing

sequences from *Aethina tumida* Murray (Polyphaga: Cucujoidea), *Agrilus planipennis*

Fairmaire (Polyphaga: Buprestoidea), *Anoplophora glabripennis* (Motschulsky) (Polyphaga:

Chrysomeloidea), *Dendroctonus ponderosae* (Hopkins) (Phytophaga: Curculionoidea),

Leptinotarsa decemlineata Say (Polyphaga: Chrysomeloidea), *Nicrophorus vespilloides*

Herbst (Polyphaga: Staphylinoidea), *Onthophagus taurus* (Schreber) (Polyphaga:

Scarabaeoidea), and *Tribolium castaneum* (Herbst) (Polyphaga: Tenebrionoidea) alongside

our two species of interest (summarised in Table S1). Alignments were computed using

MUSCLE v.3.8.425 (Edgar, 2004), checked by eye to ensure all transcripts were in frame,

and trimmed to exclude portions of GenBank sequences outside of open reading frames.

Partitions were inferred using PartitionFinder v.2.1.1 (Lanfear et al., 2017), with a resulting

partitioning scheme by codon position. Phylogenetic analysis of HSPs was performed using

maximum likelihood with RAxML v.8.2.12 (Stamatakis, 2014) via the CIPRES Science

Gateway (Miller et al., 2010) with default settings and 1000 bootstrap replicates.

3 | RESULTS

3.1 | A high-quality reference transcriptome for *Paroster nigroadumbratus*

enables the characterisation of genes upregulated during the heat shock

response

Here we present a high-quality, near-complete transcriptome for the epigean beetle *P.*

nigroadumbratus. This dataset consisted of 75,045,266 paired-end reads (72,266,264

following quality control measures), 84% of which were incorporated into 134,246 *de novo*

assembled transcripts representing 60,683 unique Trinity "genes" and 41,979 predicted

ORFs. According to assessment in BUSCO, this transcriptome was 87.66% complete with

respect to core arthropod genes and 96.74% complete when considering partial genes. Of

these, 47,810 transcripts were functionally annotated using the Trinotate pipeline. Subsetting

our predicted peptide dataset to include only the longest isoforms per putative locus, allowing

us a proxy with which to perform our downstream analyses at the gene level, resulted in

14,897 predicted ORFs (with 11,609, or ~77%, having some level of annotation).

3.2 | Differential expression, GO enrichment analysis reveal distinct

expression profiles associated with the heat shock response

Differential expression analysis in edgeR identified a total of 802 differentially expressed

(DE) genes in the epigean *P. nigroadumbratus* samples and 153 in the subterranean *P.*

macrosturtensis, with the two species exhibiting complex and markedly different expression

profiles (Figures 1, S2, S3). Different transcripts variants of the same putative homolog were

often differentially expressed across temperature treatments; these are indicated in Figure 1 as

bracketed numbers after gene names to avoid confusion. *P. macrosturtensis* upregulated

Hsp68 at both 25ºC and 35ºC, a transcript only upregulated at 35ºC in *P. nigroadumbratus*.

No differentially expressed heat shock proteins or cochaperones were shared between 25ºC

and 35ºC treatments exclusively in one species, nor were any differentially expressed

exclusively at 25ºC in *P. macrosturtensis* and not shared by any other treatments. At 35ºC,

both species upregulated the HSP genes *Hsp83* and *Hsc70-4* and the cochaperones *Samui*,

BAG2, *DnaJ, AHSA1, and Unc45a*. Gene expression profiles unique to each species

involved:1) the downregulation of two distinct transcripts each of *l(2)efl* and *Hsp68* in *P.*

nigroadumbratus at 25 ºC; 2) the downregulation of the *BAG6* cochaperone and the

upregulation of *Hsc70-3* and *l(2)efl* and *Hsp68* in *P. nigroadumbratus* at 35ºC; and finally 3)

the upregulation of *Hsc70-2*, *Hsp68* and sHSPs *l(2)efl* and *HspB1* in *P. macrosturtensis* at

35ºC. Interestingly, *Hsp68* transcripts upregulated in *P. macrosturtensis* at 25ºC were the

same as those downregulated in *P. nigroadumbratus* under the same conditions. Similarly,

l(2)efl and the aforementioned *Hsp68* transcripts downregulated at 25ºC were also

upregulated at 35ºC in *P. nigroadumbratus*. Other annotated genes thought to be involved in

the HSR, such as *Hsc70-2*, *Hsc70-5*, *TRAP1*, and *Hsp90b1*, were not differentially expressed

in either species. A full list of differentially expressed genes shared between (or unique to)

treatments is available in Table S2.

Gene Ontology enrichment analysis enables the visualisation and collapse of similar

GO terms associated with differentially expressed genes. We did not observe any consistent

depletion of GO terms associated with downregulated genes, but enriched terms were

reflective of our differential expression results above (Figure 2). In both species, terms

associated with the HSR were enriched at 35ºC (e.g. ATPase, chaperone binding, and

unfolded protein binding). At 25ºC, ADP binding was also enriched for *P. macrosturtensis*

due to the upregulation of *Hsp68* at that treatment.

3.3 | Phylogenetic analyses

Our HSP nucleotide alignments consisted of 67 sequences across 3,669bp (HSP70 family),

29 sequences across 3,741bp (HSP90 family) and 23 sequences across 867bp (sHSP family).

Three major clades were recovered in our phylogeny with strong node support (Figure 3). All

sequences from the *Paroster* species examined here were recovered as nested within these

clades, confirming the orthology of annotated HSPs, with the exception of a transcript present

in both species and inferred as sister to the *Hsp68+Hsc70-2* clades. This transcript was not

differentially expressed in response to heat shock in our dataset.

4 | DISCUSSION

Here, we have comprehensively characterised the heat-shock response at the molecular level

in the subterranean diving beetle *Paroster macrosturtensis* and one of its surface-dwelling

relatives. Using a near-complete reference transcriptome for *P. nigroadumbratus*–the first

such dataset for a member of the Hydroporinae–we performed differential expression and GO

enrichment analysis to explore genes putatively involved in the HSR. Our results demonstrate

that both the epigean *P. nigroadumbratus* and subterranean *P. macrosturtensis* have an

inducible HSR, mirroring previous survival experiments for the genus (Jones et al., 2021).

Putative orthologs of HSP cochaperones, sHSP, HSP70, and HSP90 genes were accounted

for in our differential expression analysis. However, the conditions under which this response

is activated differs between species, and *P. macrosturtensis* notably differentially expressed

less than 20% of the number of genes in response to heat stress compared to *P.*

nigroadumbratus.

4.1 | Heat shock-induced expression profiles of *Paroster* species

HSPs identified as differentially expressed in this study support past results from beetle

species and other insects more broadly. A major trend from our results concerned the

upregulation of HSP70 genes, particularly *Hsp68*; HSP70s are highly expressed in response

to heat shock in the Colorado potato beetle (Yocum, 2001), red flour beetle (Chen et al.,

2020), and the cave beetle *Neobathyscia* (Bernabò et al., 2011). These genes are upregulated

in concert with sHSPs and HSP90 genes in the rice weevil (Yuan et al., 2014) and Japanese

sawyer beetle (Cai et al., 2017) under the same conditions. HSP70s and sHSPs are also

known to be upregulated in response to heat shock in the cigarette beetle (M.-Y. Li et al.,

2021; W.-J. Yang et al., 2019).

In addition to the heat shock proteins, we also observed the upregulation of the heat

shock cognate *Hsc70-4* and cochaperones *DnaJ*, *Samui*, *BAG2*, and *AHSA1* in response to

heat shock in *Paroster*. To our knowledge there has been no documentation of a coleopteran

heat shock cognate being upregulated in response to increasing temperature, though evidence

exists for the parasitic wasp *Pteromalus*, in which *hsc70* is also induced by heavy metal

poisoning and starvation in addition to thermal extremes (H. Wang et al., 2012). Heat shock

cognates are also upregulated during diapause in silkworm eggs (Sasibhushan et al., 2012)

and young bumble bee queens (Kim et al., 2008), potentially playing a cryoprotective role in

these species. Cochaperones are less well characterised in insects, but evidence for their

upregulation in response to heat shock has been documented for *DnaJ* in hemipterans (Huang

& Kang, 2007; Hull et al., 2013) and BAG-family members, such as *Samui,* in ants (Tonione

et al., 2020). Finally, we also observed the downregulation of multiple heat shock proteins in

both species. In *P. nigroadumbratus* this was restricted to *BAG6* (a cochaperone) at 35ºC, and

l(2)efl and *Hsp68* at 25ºC. Both sHSPs and HSP70s have been documented as being

downregulated during periods of heat stress in other insects, e.g. in silk moths (Sakano et al.,

2006).

4.2 | The subterranean species *Paroster macrosturtensis* retains a heat shock response

The dataset presented here adds to a growing body of knowledge concerning the HSR in

organisms that inhabit thermally stable environments. Central to discourse on this topic is the

climatic variability hypothesis, which posits that the thermal tolerance of a taxon is positively

correlated with its temperature ranges encountered in nature (Stevens, 1989). This hypothesis

implies species from extremely stable thermal environments can no longer tolerate

temperature extremes, and has been demonstrated in a wide variety of organisms that have

either lost or possess a reduced HSR, such as cnidarians (Brennecke et al., 1998), limpets

(Sanders et al., 1991), amphipods and sea stars (Clark et al., 2008), and midges (Rinehart et

al., 2006). However, there are exceptions to this rule, with groundwater-dwelling organisms

such as those examined here displaying an inducible HSR in response to conditions far

warmer than they would encounter in nature (Colson-Proch et al., 2010; Mermillod-Blondin

et al., 2013). The HSR of these species, as well as *P. macrosturtensis*, might be retained at

such high temperatures for a variety of reasons, including the fact that HSPs play a

pleiotropic role in responding to other forms of stress and/or these species have not occupied

their respective environments for a sufficient length of time in evolutionary terms for their

HSR to be lost, e.g. via adaptive processes or a relaxation of purifying selection (Jones et al.,

2021). The latter hypothesis is plausible for *Paroster* as *P. macrosturtensis* is also known to

have retained the ability to detect light despite inhabiting an aphotic environment for over ~3

million years (Langille et al., 2019).

The present study not only demonstrates that *P. macrosturtensis* has retained an

inducible HSR, but that its HSR appears to differ markedly from its epigean relative *P.*

nigroadumbratus at the molecular level. While *P. macrosturtensis* is endemic to the Sturt

Meadows calcrete in which groundwater temperatures are relatively narrow and do not rise

above approximately 28ºC, individuals can survive at up to 38ºC before experiencing severe

locomotive impairment (Jones et al., 2021). The retention of a HSR in *P. macrosturtensis*

supports these physiological findings. However, almost half (40%) of the *P. macrosturtensis*

cohort did not survive 24 hours after heat shock compared to zero deaths with respect to *P.*

nigroadumbratus individuals. Given the reduced thermal tolerance of *P. macrosturtensis*, it

remains to be seen if the species could tolerate such extremes in the long term. Indeed, even

cave beetles considered stenothermal–those that are only capable of surviving within an

extremely narrow temperature range–have retained the HSR (Pallarés et al., 2020), but

nonetheless cannot survive at extreme temperatures for long periods (>7 days) compared to

epigean relatives (Bernabò et al., 2011; Rizzo et al., 2015).

Our molecular data reflects the previously documented reduced thermal tolerance of

subterranean dytiscids such as *P. macrosturtensis*. In keeping with these findings, *P.*

macrosturtensis exhibited significant upregulation of a transcript of *Hsp68* at 25ºC that was

also enriched in all 35ºC treatments in response to temperature extremes. These results do not

necessarily imply *Hsp68* was upregulated in response to heat stress at 25ºC, as HSPs are

known to be involved in responding to a wide range of stressors (King & MacRae, 2015;

Sang et al., 2012; Sørensen et al., 2003) 25ºC is within the range of temperatures *P.*

macrosturtensis might encounter in nature (average 23ºC–26.2ºC) (Jones et al., 2021), and

this was the same temperature (± 1ºC) as the control treatment of Jones *et al*. (2021). On the

other hand, the upregulated *Hsp68* transcript variant was also differentially expressed in the

two 35ºC treatments in both species, potentially suggesting a link to heat stress.

The above could indicate *P. macrosturtensis* 1) did experience some initial degree of heat

shock response in the control treatment and this was further exacerbated by the addition of

the experimental setup in the 25ºC treatment, or 2) the response was triggered by stress

induced by being moved to designated 25ºC vials, given *P. nigroadumbratus* also exhibits

differential expression of HSPs at 25ºC (though the downregulation of transcripts therein are

not shared with other treatments) (Figure 1). The former scenario would reflect the reduced

thermal tolerance of *P. macrosturtensis*, which is also supported by the species differentially

expressing a far lower number of genes in response to higher temperatures compared to *P.*

nigroadumbratus. Similarly reduced numbers of differentially expressed genes have also

been associated with lower thermal tolerances in other organisms such as fish (Narum &

Campbell, 2015), lizards (Akashi et al., 2016), rotifers (Paraskevopoulou et al., 2020), red

algae (W. Wang et al., 2018), and plants (Jin et al., 2020; Sun et al., 2021), though we note

the inverse (or alternatively, no clear pattern) has been observed in a number of cases,

potentially reflecting lower levels of stress as opposed to an inability to mount a HSR

(DeBiasse & Kelly, 2016; Perez et al., 2021).

Acquiring data such as these is not only necessary to better understand the HSR in

subterranean animals in the face of climate change, but also to inform future conservation

decisions. The latter is of particular relevance to members of subterranean *Paroster* as the

groundwater in the palaeovalleys containing their calcrete habitats, as well as the calcretes themselves, are heavily utilised for mining activities. As short-range endemics to the extreme, these species are not only at risk of habitat degradation via the direct intersection of shallow aquifers with mining activities, but also through the drawdown of groundwater beneath calcretes at greater depths (Hose et al., 2015). Dytiscids (and stygofauna more broadly) in these regions therefore not only face population declines or extinctions due to the increasing threat of climate change, but also potentially co-occurring stressors from mining operations such as changes in temperature, quality, and depth of groundwater. Accordingly, the calcretes these species inhabit are considered a top conservation priority for protection by the Commonwealth Scientific and Industrial Research Organisation (Healy, 2015). Such threats are of particular concern for obligate stygofaunal species given the data presented by Jones *et al.* (2021) and, more broadly, the fact that the thermal critical maximum of subterranean beetle species is known to decrease with the degree of subterranean specialisation (Pallarés et al., 2020). An increased knowledge of the assumed fragility of Australian subterranean dytiscids in the face of these stressors is therefore crucial for informing future conservation decisions.

Overall, the findings presented here demonstrate the reduced thermal tolerance of the subterranean species *P. macrosturtensis* compared to its epigean relatives is reflected in, and further clarified by, transcriptomic data. While our data are supported by past physiological evidence that demonstrated *P. macrosturtensis* could survive at high temperatures, albeit not to the limits of epigean species (Jones et al., 2021), the present study adds a new layer to this narrative. *P. macrosturtensis* might possess increased mortality in the face of high temperatures and a lower maximum temperature it can tolerate because 1) the species differentially expresses far fewer genes in response to heat shock compared to the epigean relative *P. nigroadumbratus*, suggesting it is unable to mount a robust HSR, and 2) this

response appears to be triggered at lower temperatures and/or in response to ambient

stressors, potentially subjecting individuals to stress for longer periods of time. Ultimately, it

remains unknown as to how *P. macrosturtensis* might respond to temperature extremes in the

long term, such as under a predicted increase of up to ~5ºC by 2090 in central Western

Australia (Healy, 2015). Nonetheless, the apparently weaker HSR we have characterised for

the species implies these short range endemics may not be able to thrive in such conditions,

especially given the threat of more immediate human-induced stressors such as mining

activities.

5 | CONCLUSIONS

The present study contrasted differentially expressed genes involved in the heat shock

response (HSR) of *Paroster nigroadumbratus* and *P. macrosturtensis*, epigean and

subterranean lineages of hydroporine diving beetles endemic to Australia previously

subjected to heat shock by Jones *et al.* (2021). Our results have comprehensively

characterised the genes involved in the HSR in these species and demonstrate that the

subterranean *P. macrosturtensis* retains a HSR at the molecular level, in agreement with the

physiological data of Jones *et al.* (2021) detailing that the species can survive temperatures

far above what it would encounter in nature. However, we also show that *P. macrosturtensis*

is potentially less equipped at the molecular level to respond to higher temperatures, with the

species differentially expressing less than a fifth of the number of genes in response to heat

shock in *P. nigroadumbratus* at 35ºC. Together, these results support the idea that *P.*

macrosturtensis has lost some degree of tolerance to temperature extremes compared to

epigean species such as *P. nigroadumbratus*, in agreement with the climatic variability

hypothesis. Ultimately, sampling of additional subterranean dytiscid species in the Yilgarn

region will enable a more holistic assessment of the thermal tolerances of this unique natural

experiment at the molecular level.

ACKNOWLEDGEMENTS

We thank Flora, Peter, and Paul Axford for providing access to the Sturt Meadows calcrete

and accommodation at the Sturt Meadows pastoral property. We thank Chris Watts (South

Australian Museum) for helping with identification and collecting of surface dytiscids,

providing beetle images, and for laying the foundation for the subterranean beetle research

with WFH and Remko Leijs. We thank Rae Humphreys for assistance with field collections

of *P. macrosturtensis* and for being a wonderful host to stygofauna catchers over many years.

Funding for this project was provided by an Australian Research Council Discovery grant

(DP180103851) to SJBC, WFH, ADA and TB. The authors have no conflict of interest to

disclose.

REFERENCES

Akashi, H. D., Cádiz Díaz, A., Shigenobu, S., Makino, T., & Kawata, M. (2016).

Differentially expressed genes associated with adaptation to different thermal

environments in three sympatric Cuban Anolis lizards. *Molecular Ecology*, *25*(10),

2273–2285. https://doi.org/10.1111/mec.13625

Badino, G. (2010). UNDERGROUND METEOROLOGY-"What's the weather

underground?" *Acta Carsologica*, *39*(3), Article 3. https://doi.org/10.3986/ac.v39i3.74

Balart-García, P., Cieslak, A., Escuer, P., Rozas, J., Ribera, I., & Fernández, R. (2021).

Smelling in the dark: Phylogenomic insights into the chemosensory system of a

subterranean beetle. *Molecular Ecology*. https://doi.org/10.1111/mec.15921

Beale, A. D., Whitmore, D., & Moran, D. (2016). Life in a dark biosphere: A review of

circadian physiology in "arrhythmic" environments. *Journal of Comparative*

Physiology. B, Biochemical, Systemic, and Environmental Physiology, *186*(8), 947–

968. https://doi.org/10.1007/s00360-016-1000-6

Bernabò, P., Latella, L., Jousson, O., & Lencioni, V. (2011). Cold stenothermal cave-

dwelling beetles do have an HSP70 heat shock response. *Journal of Thermal Biology*,

36(3), 206–208. https://doi.org/10.1016/j.jtherbio.2011.03.002

Brennecke, T., Gellner, K., & Bosch, T. C. (1998). The lack of a stress response in Hydra

oligactis is due to reduced hsp70 mRNA stability. *European Journal of Biochemistry*,

255(3), 703–709. https://doi.org/10.1046/j.1432-1327.1998.2550703.x

Cai, Z., Chen, J., Cheng, J., & Lin, T. (2017). Overexpression of Three Heat Shock Proteins

Protects Monochamus alternatus (Coleoptera: Cerambycidae) From Thermal Stress.

Journal of Insect Science, *17*(113). https://doi.org/10.1093/jisesa/iex082

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden,

T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1),

421. https://doi.org/10.1186/1471-2105-10-421

Chen, M., Zhang, N., Jiang, H., Meng, X., Qiang, K., & Wang, J. (2020). Transcriptional

regulation of heat shock protein 70 genes by class I histone deacetylases in the red

flour beetle, Tribolium castaneum. *Insect Molecular Biology*, *29*(2), 221–230.

https://doi.org/10.1111/imb.12627

Clark, M. S., Fraser, K. P. P., & Peck, L. S. (2008). Lack of an HSP70 heat shock response in

two Antarctic marine invertebrates. *Polar Biology*, *31*(9), 1059–1065.

https://doi.org/10.1007/s00300-008-0447-7

Colson-Proch, C., Morales, A., Hervant, F., Konecny, L., Moulin, C., & Douady, C. J.

(2010). First cellular approach of the effects of global warming on groundwater

organisms: A study of the HSP70 gene expression. *Cell Stress & Chaperones*, *15*(3),

259–270. https://doi.org/10.1007/s12192-009-0139-4

Conway, J. R., Lex, A., & Gehlenborg, N. (2017). UpSetR: An R package for the

visualization of intersecting sets and their properties. *Bioinformatics*, *33*(18), 2938–

2940. https://doi.org/10.1093/bioinformatics/btx364

Cooper, S. J. B., Hinze, S., Leys, R., Watts, C. H. S., & Humphreys, W. F. (2002). Islands

under the desert: Molecular systematics and evolutionary origins of stygobitic water

beetles (Coleoptera : Dytiscidae) from central Western Australia. *Invertebrate*

Systematics, *16*(4), 589–590. https://doi.org/10.1071/it01039

Culver, D. C., & Pipan, T. (2014). Shallow Subterranean Habitats: Ecology, Evolution, and

Conservation. In *Shallow Subterranean Habitats*. Oxford University Press.

https://oxford.universitypressscholarship.com/view/10.1093/acprof:oso/97801996461

73.001.0001/acprof-9780199646173

DeBiasse, M. B., & Kelly, M. W. (2016). Plastic and Evolved Responses to Global Change:

What Can We Learn from Comparative Transcriptomics? *Journal of Heredity*, *107*(1),

71–81. https://doi.org/10.1093/jhered/esv073

Edgar, R. C. (2004). MUSCLE: A multiple sequence alignment method with reduced time

and space complexity. *BMC Bioinformatics*, *5*(1), 113. https://doi.org/10.1186/1471-

2105-5-113

Gibert, J., & Deharveng, L. (2002). Subterranean ecosystems: A truncated functional

biodiversity. *BioScience*, *52*(6), 473–481. https://doi.org/10.1641/0006-

3568(2002)052[0473:SEATFB]2.0.CO;2

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis,

X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke,

A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev,

542 A. (2011). Trinity: Reconstructing a full-length transcriptome without a genome from

543 RNA-Seq data. *Nature Biotechnology*, *29*(7), 644–652.

544 https://doi.org/10.1038/nbt.1883

545 Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger,

546 M. B., Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N.,

547 Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C. N., … Regev, A.

548 (2013). De novo transcript sequence reconstruction from RNA-Seq: Reference

549 generation and analysis with Trinity. *Nature Protocols*, *8*(8).

550 https://doi.org/10.1038/nprot.2013.084

551 Healy, M.-A. (2015). *It's hot and getting hotter. Australian rangelands and climate change –*

552 *Reports of the Rangelands Cluster Project.* Ninti One Limited and CSIRO, Alice

553 Springs.

554 Hose, G. C., Sreekanth, J., Barron, O., & Pollino, C. (2015). *Stygofauna in Australian*

555 *Groundwater Systems: Extent of knowledge*. CSIRO, Australia.

556 Huang, L.-H., & Kang, L. (2007). Cloning and interspecific altered expression of heat shock

557 protein genes in two leafminer species in response to thermal stress. *Insect Molecular*

558 *Biology*, *16*(4), 491–500. https://doi.org/10.1111/j.1365-2583.2007.00744.x

559 Hull, J. J., Geib, S. M., Fabrick, J. A., & Brent, C. S. (2013). Sequencing and De Novo

560 Assembly of the Western Tarnished Plant Bug (Lygus hesperus) Transcriptome.

561 *PLOS ONE*, *8*(1), e55105. https://doi.org/10.1371/journal.pone.0055105

562 Humphreys, W. (2012). Diversity Patterns in Australia. In *Encyclopedia of Caves* (pp. 203–

563 219). https://doi.org/10.1016/B978-0-12-383832-2.00029-3

564 Jin, J., Yang, L., Fan, D., Liu, X., & Hao, Q. (2020). Comparative transcriptome analysis

565 uncovers different heat stress responses in heat-resistant and heat-sensitive jujube

566 cultivars. *PLOS ONE*, *15*(9), e0235763. https://doi.org/10.1371/journal.pone.0235763

Jindal, S., Dudani, A. K., Singh, B., Harley, C. B., & Gupta, R. S. (1989). Primary structure

of a human mitochondrial protein homologous to the bacterial and plant chaperonins

and to the 65-kilodalton mycobacterial antigen. *Molecular and Cellular Biology*, *9*(5),

2279–2283. https://doi.org/10.1128/mcb.9.5.2279

Jones, K. K., Cooper, S. J. B., & Seymour, R. S. (2019). Cutaneous respiration by diving

beetles from underground aquifers of Western Australia (Coleoptera: Dytiscidae).

Journal of Experimental Biology, *222*(7), jeb196659.

https://doi.org/10.1242/jeb.196659

Jones, K. K., Humphreys, W. F., Saccò, M., Bertozzi, T., Austin, A. D., & Cooper, S. J. B.

(2021). The critical thermal maximum of diving beetles (Coleoptera: Dytiscidae): a

comparison of subterranean and surface-dwelling species. *Current Research in Insect*

Science, *1*, 100019. https://doi.org/10.1016/j.cris.2021.100019

Kim, B.-G., Shim, J.-K., Kim, D.-W., Kwon, Y. J., & Lee, K.-Y. (2008). Tissue-specific

variation of heat shock protein gene expression in relation to diapause in the

bumblebee Bombus terrestris. *Entomological Research*, *38*(1), 10–16.

https://doi.org/10.1111/j.1748-5967.2008.00142.x

King, A. M., & MacRae, T. H. (2015). Insect Heat Shock Proteins During Stress and

Diapause. *Annual Review of Entomology*, *60*(1), 59–75.

https://doi.org/10.1146/annurev-ento-011613-162107

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., &

Zdobnov, E. M. (2019). OrthoDB v10: Sampling the diversity of animal, plant,

fungal, protist, bacterial and viral genomes for evolutionary and functional

annotations of orthologs. *Nucleic Acids Research*, *47*(D1), D807–D811.

https://doi.org/10.1093/nar/gky1053

Lanfear, R., Frandsen, P. B., Wright, A. M., Senfeld, T., & Calcott, B. (2017). PartitionFinder

2: New Methods for Selecting Partitioned Models of Evolution for Molecular and

Morphological Phylogenetic Analyses. *Molecular Biology and Evolution*, *34*(3), 772–

773. https://doi.org/10.1093/molbev/msw260

Langille, B. L., Hyde, J., Saint, K. M., Bradford, T. M., Stringer, D. N., Tierney, S. M.,

Humphreys, W. F., Austin, A. D., & Cooper, S. J. B. (2021). Evidence for speciation

underground in diving beetles (Dytiscidae) from a subterranean archipelago.

Evolution, *75*(1), 166–175. https://doi.org/10.1111/evo.14135

Langille, B. L., Tierney, S. M., Austin, A. D., Humphreys, W. F., & Cooper, S. J. B. (2019).

How blind are they? Phototactic responses in stygobiont diving beetles (Coleoptera:

Dytiscidae) from calcrete aquifers of Western Australia. *Austral Entomology*, *58*(2),

425–431. https://doi.org/10.1111/aen.12330

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. L. (2009). Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. *Genome Biology*,

10(3), R25. https://doi.org/10.1186/gb-2009-10-3-r25

Leijs, R., Nes, E., Watts, C., Cooper, S., Humphreys, W., & Hogendoorn, K. (2012).

Evolution of Blind Beetles in Isolated Aquifers: A Test of Alternative Modes of

Speciation. *PloS One*, *7*, e34260. https://doi.org/10.1371/journal.pone.0034260

Leys, R., Watts, C. H. S., Cooper, S. J. B., & Humphreys, W. F. (2003). Evolution of

subterranean diving beetles (Coleoptera: Dytiscidae: Hydroporini, Didessini) in the

arid zone of Australia. *Evolution*, *57*(12), 2819–2834. https://doi.org/10.1554/03-165

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics (Oxford, England)*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,

G., Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The

Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*,

25(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Li, M.-Y., Huang, Y., Lei, X., Xu, C.-T., Li, B., Chen, D.-X., & Liu, S. (2021). Identification

of six heat shock protein 70 genes in Lasioderma serricorne (Coleoptera: Anobiidae)

and their responses to temperature stress. *Journal of Asia-Pacific Entomology*.

https://doi.org/10.1016/j.aspen.2021.05.005

Madden, T. L., Busby, B., & Ye, J. (2019). Reply to the paper: Misunderstood parameters of

NCBI BLAST impacts the correctness of bioinformatics workflows. *Bioinformatics*,

35(15), 2699–2700. https://doi.org/10.1093/bioinformatics/bty1026

Mammola, S., Amorim, I. R., Bichuette, M. E., Borges, P. A. V., Cheeptham, N., Cooper, S.

J. B., Culver, D. C., Deharveng, L., Eme, D., Ferreira, R. L., Fišer, C., Fišer, Ž., Fong,

D. W., Griebler, C., Jeffery, W. R., Jugovic, J., Kowalko, J. E., Lilley, T. M., Malard,

F., … Cardoso, P. (2020). Fundamental research questions in subterranean biology.

Biological Reviews, *95*(6), 1855–1872. https://doi.org/10.1111/brv.12642

Mammola, S., Piano, E., Cardoso, P., Vernon, P., Domínguez-Villar, D., Culver, D. C.,

Pipan, T., & Isaia, M. (2019). Climate change going deep: The effects of global

climatic alterations on cave ecosystems. *The Anthropocene Review*, *6*(1–2), 98–116.

https://doi.org/10.1177/2053019619851594

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.Journal*, *17*(1), 10–12. https://doi.org/10.14806/ej.17.1.200

Mermillod-Blondin, F., Lefour, C., Lalouette, L., Renault, D., Malard, F., Simon, L., &

Douady, C. J. (2013). Thermal tolerance breadths among groundwater crustaceans

living in a thermally constant environment. *Journal of Experimental Biology*, *216*(9),

1683–1694. https://doi.org/10.1242/jeb.081232

Merritt, D. J., & Clarke, A. K. (2011). Synchronized circadian bioluminescence in cave-

dwelling Arachnocampa tasmaniensis (Glowworms). *Journal of Biological Rhythms*,

26(1), 34–43. https://doi.org/10.1177/0748730410391947

Miller, M. A., Pfeiffer, W., & Schwartz, T. (2010). Creating the CIPRES Science Gateway

for inference of large phylogenetic trees. *2010 Gateway Computing Environments*

Workshop (GCE), 1–8. https://doi.org/10.1109/GCE.2010.5676129

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L.,

Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A.

(2021). Pfam: The protein families database in 2021. *Nucleic Acids Research*, *49*(D1),

D412–D419. https://doi.org/10.1093/nar/gkaa913

Narum, S. R., & Campbell, N. R. (2015). Transcriptomic response to heat stress among

ecologically divergent populations of redband trout. *BMC Genomics*, *16*(1), 103.

https://doi.org/10.1186/s12864-015-1246-5

Nishimura, O., Hara, Y., & Kuraku, S. (2017). GVolante for standardizing completeness

assessment of genome and transcriptome assemblies. *Bioinformatics*, *33*(22), 3635–

3637. https://doi.org/10.1093/bioinformatics/btx445

Pallarés, S., Colado, R., Botella-Cruz, M., Montes, A., Balart-García, P., Bilton, D. T.,

Millán, A., Ribera, I., & Sánchez-Fernández, D. (2020). Loss of heat acclimation

capacity could leave subterranean specialists highly sensitive to climate change.

Animal Conservation. https://doi.org/10.1111/acv.12654

Paraskevopoulou, S., Dennis, A. B., Weithoff, G., & Tiedemann, R. (2020). Temperature-

dependent life history and transcriptomic responses in heat-tolerant versus heat-

sensitive Brachionus rotifers. *Scientific Reports*, *10*(1), 13281.

https://doi.org/10.1038/s41598-020-70173-0

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides

fast and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4),

417–419. https://doi.org/10.1038/nmeth.4197

Perez, R., de Souza Araujo, N., Defrance, M., & Aron, S. (2021). Molecular adaptations to

heat stress in the thermophilic ant genus Cataglyphis. *Molecular Ecology*, *n/a*(n/a).

https://doi.org/10.1111/mec.16134

Peuß, R., Eggert, H., Armitage, S. A. O., & Kurtz, J. (2015). Downregulation of the

evolutionary capacitor Hsp90 is mediated by social cues. *Proceedings of the Royal*

Society B: Biological Sciences, *282*(1819), 20152041.

https://doi.org/10.1098/rspb.2015.2041

Richter, K., Haslbeck, M., & Buchner, J. (2010). The Heat Shock Response: Life on the

Verge of Death. *Molecular Cell*, *40*(2), 253–266.

https://doi.org/10.1016/j.molcel.2010.10.006

Rinehart, J. P., Hayward, S. A. L., Elnitsky, M. A., Sandro, L. H., Lee, R. E., & Denlinger, D.

L. (2006). Continuous up-regulation of heat shock proteins in larvae, but not adults, of

a polar insect. *Proceedings of the National Academy of Sciences of the United States*

of America, *103*(38), 14223–14227. https://doi.org/10.1073/pnas.0606840103

Rizzo, V., Sánchez-Fernández, D., Fresneda, J., Cieslak, A., & Ribera, I. (2015). Lack of

evolutionary adjustment to ambient temperature in highly specialized cave beetles.

BMC Evolutionary Biology, *15*(1), 10. https://doi.org/10.1186/s12862-015-0288-2

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: A Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinformatics*,

26(1), 139–140. https://doi.org/10.1093/bioinformatics/btp616

Sakano, D., Li, B., Xia, Q., Yamamoto, K., Fuiji, H., & Aso, Y. (2006). Genes Encoding

Small Heat Shock Proteins of the Silkworm, Bombyx mori. *Bioscience,*

Biotechnology, and Biochemistry, *70*(10), 2443–2450.

https://doi.org/10.1271/bbb.60176

Sanders, B. M., Hope, C., Pascoe, V. M., & Martin, L. S. (1991). Characterization of the

Stress Protein Response in Two Species of Collisella Limpets with Different

Temperature Tolerances. *Physiological Zoology*, *64*(6), 1471–1489.

Sang, W., Ma, W.-H., Qiu, L., Zhu, Z.-H., & Lei, C.-L. (2012). The involvement of heat

shock protein and cytochrome P450 genes in response to UV-A exposure in the beetle

Tribolium castaneum. *Journal of Insect Physiology*, *58*(6), 830–836.

https://doi.org/10.1016/j.jinsphys.2012.03.007

Sasibhushan, S., Ponnuvel, K. M., & Vijayaprakash, N. B. (2012). Diapause specific gene

expression in the eggs of multivoltine silkworm Bombyx mori, identified by

suppressive subtractive hybridization. *Comparative Biochemistry and Physiology Part*

B: Biochemistry and Molecular Biology, *161*(4), 371–379.

https://doi.org/10.1016/j.cbpb.2012.01.002

Shah, N., Nute, M. G., Warnow, T., & Pop, M. (2019). Misunderstood parameter of NCBI

BLAST impacts the correctness of bioinformatics workflows. *Bioinformatics*, *35*(9),

1613–1614. https://doi.org/10.1093/bioinformatics/bty833

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M.

(2015). BUSCO: Assessing genome assembly and annotation completeness with

single-copy orthologs. *Bioinformatics (Oxford, England)*, *31*(19), 3210–3212.

https://doi.org/10.1093/bioinformatics/btv351

Somero, G. N. (2020). The cellular stress response and temperature: Function, regulation, and

evolution. *Journal of Experimental Zoology Part A: Ecological and Integrative*

Physiology, *333*(6), 379–397. https://doi.org/10.1002/jez.2344

Sørensen, J. G., Kristensen, T. N., & Loeschcke, V. (2003). The evolutionary and ecological

role of heat shock proteins. *Ecology Letters*, *6*(11), 1025–1037.

https://doi.org/10.1046/j.1461-0248.2003.00528.x

Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics*, *30*(9), 1312–1313.

https://doi.org/10.1093/bioinformatics/btu033

Stevens, G. C. (1989). The Latitudinal Gradient in Geographical Range: How so Many

Species Coexist in the Tropics. *The American Naturalist*, *133*(2), 240–256.

Sun, M., Lin, C., Zhang, A., Wang, X., Yan, H., Khan, I., Wu, B., Feng, G., Nie, G., Zhang,

X., & Huang, L. (2021). Transcriptome sequencing revealed the molecular

mechanism of response of pearl millet root to heat stress. *Journal of Agronomy and*

Crop Science, *207*(4), 768–773. https://doi.org/10.1111/jac.12496

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO Summarizes and

Visualizes Long Lists of Gene Ontology Terms. *PLOS ONE*, *6*(7), e21800.

https://doi.org/10.1371/journal.pone.0021800

The UniProt Consortium. (2021). UniProt: The universal protein knowledgebase in 2021.

Nucleic Acids Research, *49*(D1), D480–D489. https://doi.org/10.1093/nar/gkaa1100

Tierney, S. M., Cooper, S. J. B., Saint, K. M., Bertozzi, T., Hyde, J., Humphreys, W. F., &

Austin, A. D. (2015). Opsin transcripts of predatory diving beetles: A comparison of

surface and subterranean photic niches. *Royal Society Open Science*, *2*(1), 140386.

https://doi.org/10.1098/rsos.140386

Tomanek, L. (2010). Variation in the heat shock response and its implication for predicting

the effect of global climate change on species' biogeographical distribution ranges

and metabolic costs. *The Journal of Experimental Biology*, *213*(6), 971–979.

https://doi.org/10.1242/jeb.038034

Tonione, M. A., Bi, K., & Tsutsui, N. D. (2020). Transcriptomic signatures of cold adaptation

and heat stress in the winter ant (Prenolepis imparis). *PLOS ONE*, *15*(10), e0239558.

https://doi.org/10.1371/journal.pone.0239558

Toussaint, E. F. A., Condamine, F. L., Hawlitschek, O., Watts, C. H., Porch, N., Hendrich,

L., & Balke, M. (2015). Unveiling the Diversification Dynamics of Australasian

Predaceous Diving Beetles in the Cenozoic. *Systematic Biology*, *64*(1), 3–24.

https://doi.org/10.1093/sysbio/syu067

Walker, J. A., & Rose, H. A. (1998). Oothecal structure and male genitalia of the

Geoscapheinae and some Australian Panaesthia Serville (Blattodea: Blaberidae).

Australian Journal of Entomology, *37*(1), 23–26. https://doi.org/10.1111/j.1440-

6055.1998.tb01538.x

Wang, H., Li, K., Zhu, J.-Y., Fang, Q., Ye, G.-Y., Wang, H., Li, K., & Zhu, J.-Y. (2012).

CLONING AND EXPRESSION PATTERN OF HEAT SHOCK PROTEIN GENES

FROM THE ENDOPARASITOID WASP, Pteromalus puparum IN RESPONSE TO

ENVIRONMENTAL STRESSES. *Archives of Insect Biochemistry and Physiology*,

79(4–5), 247–263. https://doi.org/10.1002/arch.21013

Wang, W., Lin, Y., Teng, F., Ji, D., Xu, Y., Chen, C., & Xie, C. (2018). Comparative

transcriptome analysis between heat-tolerant and sensitive Pyropia haitanensis strains

in response to high temperature stress. *Algal Research*, *29*, 104–112.

https://doi.org/10.1016/j.algal.2017.11.026

Watts, C. H. S., & Humphreys, W. F. (2009). Fourteen New Dytiscidae (Coleoptera) of the

760   Genera Limbodessus Guignot, Paroster Sharp, and Exocelina Broun from

761   Underground Waters in Australia. *Transactions of the Royal Society of South*

762   *Australia*, *133*(1), 62–107. https://doi.org/10.1080/03721426.2009.10887112

763   Yang, J., Chen, X., Bai, J., Fang, D., Qiu, Y., Jiang, W., Yuan, H., Bian, C., Lu, J., He, S.,

764   Pan, X., Zhang, Y., Wang, X., You, X., Wang, Y., Sun, Y., Mao, D., Liu, Y., Fan, G.,

765   … Shi, Q. (2016). The Sinocyclocheilus cavefish genome provides insights into cave

766   adaptation. *BMC Biology*, *14*(1), 1. https://doi.org/10.1186/s12915-015-0223-4

767   Yang, W.-J., Xu, K.-K., Cao, Y., Meng, Y.-L., Liu, Y., & Li, C. (2019). Identification and

768   Expression Analysis of Four Small Heat Shock Protein Genes in Cigarette Beetle,

769   Lasioderma serricorne (Fabricius). *Insects*, *10*(5), 139.

770   https://doi.org/10.3390/insects10050139

771   Yocum, G. D. (2001). Differential expression of two HSP70 transcripts in response to cold

772   shock, thermoperiod, and adult diapause in the Colorado potato beetle. *Journal of*

773   *Insect Physiology*, *47*(10), 1139–1145. https://doi.org/10.1016/S0022-1910(01)00095-

774   6

775   Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology

776   analysis for RNA-seq: Accounting for selection bias. *Genome Biology*, *11*(2), R14.

777   https://doi.org/10.1186/gb-2010-11-2-r14

778   Yuan, X., Zhou, W.-W., Zhou, Y., Liu, S., Lu, F., Yang, M.-F., Cheng, J., Gurr, G. M., &

779   Zhu, Z.-R. (2014). Composition and Expression of Heat Shock Proteins in an Invasive

780   Pest, The Rice Water Weevil (Coleoptera: Curculionidae). *Florida Entomologist*,

781   *97*(2), 611–619. https://doi.org/10.1653/024.097.0237

DATA ACCESSIBILITY STATEMENT

All raw RNA-seq data for *Paroster macrosturtensis* and *P. nigroadumbratus* used in differential

expression analyses in this study are available under BioProject ID PRJNA783065 via the Sequence

Read Archive. HSP transcripts assembled from raw *P. macrosturtensis* RNA-seq data and used in

phylogenetic analysis are available via GenBank under accession numbers OL803873-OL803888

(summarised in Table S1). The reference transcriptome of *P. nigroadumbratus* (including all isoforms)

and the annotated, subsetted dataset (only including longest isoform per Trinity "gene") are available via

FigShare (DOI: 10.25909/17169191).

AUTHOR CONTRIBUTIONS

PGBH and SJBC conceived the study. PGBH analysed the data and wrote the manuscript. TB and CSPF

provided guidance regarding bioinformatic analyses. ST performed molecular laboratory work. TMB,

KJ, WFH, and ADA helped to draft the manuscript. All authors edited the manuscript and approved the

final draft.

FIGURE 1 Differentially expressed (DE) genes in *Paroster nigroadumbratus*, a surface-dwelling species, and *P. macrosturtensis*, a subterranean relative, in response to 25ºC and 35ºC treatments *vs.* control treatment (25ºC without the experimental setup) by Jones *et al.* (2021). A) DE genes shared (denoted by black lines linking circles) or unique to (unlinked circles) each temperature treatment per species. The number of shared or unique DE genes among treatments is shown in the bar graph above, including names of DE HSPs and/or HSP cochaperones. Genes are shown as either upregulated or downregulated, denoted with arrows. Bracketed numbers in the bar graph indicate different transcripts of the same gene. B) total DE genes for each temperature treatment *vs.* control, irrespective of overlap with other treatments. Photographs by Chris Watts and Howard Hamon.

FIGURE 2 REVIGO treemap showing Gene Ontology (GO) terms associated with upregulated genes at 25ºC and 35ºC compared to the control treatment. The size of squares in the figure is proportional to the *p*-value associated with differential expression of respective genes. Similar GO terms share a colour and are represented in white text by the largest square per group. Terms associated with heat shock shown here include ADP binding, chaperone binding, and ATPase, indicated by a star.

FIGURE 3 Phylogeny of heat shock transcripts inferred in RAxML to validate the identity of putative HSP orthologs in *Paroster* species. Transcript variants are represented by branches of the same colour within a clade. Scale bar is in substitutions/site; BS = bootstrap node support. Tip labels show transcript names/GenBank accession numbers. Tip names for *P. macrosturtensis* are shared with the reference *P. nigroadumbratus* transcript that reads were assembled against. Differentially expressed transcripts under a certain condition are with respect to the control.

36