Differential transcriptomic responses to heat stress in surface and subterranean diving beetles

Perry G. Beasley-Hall[1,2,3]* | Terry Bertozzi[1,2] | Tessa M. Bradford[1,2] | Charles S. P. Foster[4] | Karl Jones[1,2] | Simon M. Tierney[5] | William F. Humphreys[6,7] | Andrew D. Austin[1,2] | Steven J. B. Cooper[1,2]

[1] School of Biological Sciences, The University of Adelaide, Adelaide, South Australia 5000, Australia.

[2] South Australian Museum, Adelaide, South Australia 5000, Australia.

[3] School of Life and Environmental Sciences, University of Sydney, Sydney, New South Wales 2006, Australia.

[4] School of Medical Sciences, University of New South Wales, Sydney, New South Wales 2052, Australia.

[5] Hawkesbury Institute for the Environment, Western Sydney University, Building R2 Bourke Street, Richmond NSW 2753, Australia

[6] Western Australian Museum, Locked Bag 40, Welshpool DC, WA 6986, Australia

[7] School of Biological Sciences, University of Western Australia, Crawley, WA 6009, Australia

*Corresponding author

Perry Beasley-Hall, School of Biological Sciences, Darling Building, The University of Adelaide, Adelaide, South Australia 5000, Australia

Email: perry.beasley-hall@adelaide.edu.au

Abstract

Subterranean habitats are generally very stable environments, and as such evolutionary transitions of organisms from surface to subterranean lifestyles may cause considerable shifts in physiology, particularly with respect to thermal tolerance. In this study we compared responses to heat shock at the molecular level in a geographically widespread, surface-dwelling water beetle to a congeneric subterranean species restricted to a single aquifer (Dytiscidae: Hydroporinae). The obligate subterranean beetle *Paroster macrosturtensis* is known to have a lower thermal tolerance compared to surface lineages (CT_{max} 38ºC *cf*. 42-46 ºC), but the genetic basis of this physiological difference has not been characterized. We experimentally manipulated the thermal environment of 24 individuals to demonstrate that both species can mount a heat shock response at high temperatures (35ºC), as determined by comparative transcriptomics. However, genes involved in these responses differ between species and a far greater number were differentially expressed in the surface taxon, suggesting it can mount a more robust heat shock response; these data may underpin its higher thermal tolerance compared to subterranean relatives. In contrast, the subterranean species examined not only differentially expressed fewer genes in response to increasing temperatures, but also in the presence of the experimental setup employed here alone. Our results suggest *P. macrosturtensis* may be comparatively poorly equipped to respond to both thermally induced stress and environmental disturbances more broadly. The molecular findings presented here have conservation implications for *P. macrosturtensis* and contribute to a growing narrative concerning weakened thermal tolerances in obligate subterranean organisms at the molecular level.

KEYWORDS: heat shock response, heat shock proteins, subterranean habitats, molecular evolution, transcriptomics, differential gene expression

1 | INTRODUCTION

The transition to an obligate subterranean lifestyle can cause massive shifts in an organism's biology (1), from the acquisition of classic troglomorphies such as the elongation of appendages for sensing in an aphotic environment (2), to changes in lesser-studied traits including circadian rhythm (3,4), reproductive biology (5), respiration (6), the number of larval instars (7,8) and chemosensation (9,10). These changes have been attributed to the stark difference between subterranean and surface (hereafter epigean) habitats. While epigean environments can vary immensely over both time and space, subterranean environments, such as cave systems, generally possess high environmental stability with respect to light levels, temperature, humidity, and nutrient availability (11–13). Animals adapted to these environments might therefore be particularly well suited for the assessment of responses to future climate change scenarios, particularly with respect to their thermal tolerances and responses to increasing temperatures. Indeed, such habitats have been labelled as undervalued natural laboratories for biological studies of global change (14).

A near-universal response to temperature-induced stress across the tree of life–and therefore a method by which thermal tolerance can be gauged–is the heat shock response (hereafter HSR), which involves the synthesis of heat-shock proteins (hereafter HSPs). HSPs include those proteins that are expressed constitutively under non-stressful conditions, called heat shock cognates, or those only induced when organisms are exposed to thermal extremes, during which they assist in stabilising and refolding proteins at risk of denaturation. An inducible HSR has been observed in almost all organisms studied to date, and the proteins involved in this response, as well as the response itself, are highly conserved among different domains of life (15). An estimated 50 to 200 genes are involved in the HSR, the most significantly induced of which are HSPs (16). However, there are exceptions to this rule: a lack of an inducible HSR has been documented in a wide range of species, largely those that occupy very stable thermal environments such as Antarctic marine habitats (17,18).

Knowledge of the HSR in organisms from thermally stable *subterranean* habitats, including

their associated inducible HSPs and at which temperatures this response might be activated, is

scarce, with only a few studies devoted to invertebrate taxa (14,19–25). However, invertebrates

overwhelmingly contribute to the biodiversity of subterranean habitats compared to vertebrates

(26,27). The bulk of existing studies on the thermal tolerance of subterranean invertebrates suggest

such taxa can withstand temperatures above those they would encounter in nature and that they have

not lost the HSR. Only a small number of these studies have examined the heat shock response

directly (28) and tend to focus on thermal tolerances gauged through survival experiments.

Moreover, a better understanding of responses to current climate change predictions for subterranean

animals has been identified as a fundamental question in subterranean biology given emerging

conservation issues associated with their respective ecosystems (29). To address this knowledge gap,

here we make use of genomic data from Australian representatives of a group of aquatic

invertebrates containing both epigean and subterranean lineages.

The Yilgarn Craton in central Western Australia (WA) houses a diverse subterranean diving

beetle fauna belonging to two tribes, Bidessini and Hydroporini (Dytiscidae). While epigean species

can be found practically continent-wide, subterranean taxa are isolated in calcrete aquifers (hereafter

calcretes) associated with ancient palaeodrainage systems in the region. These calcretes are

completely devoid of light and animals contained within them are assumed to have little to no access

to air above the water's surface (6). Each calcrete houses between one and three dytiscid species, but

at the time of writing only around a quarter of the ~200 known calcretes have been sampled (30–32).

Nonetheless, lineages in both tribes are known to have made independent yet parallel, repeated

transitions into underground habitats from epigean ancestors during the late Miocene to early

Pleistocene, likely in response to continental aridification (32–35). In each case these transitions

have involved the loss of eyes, pigment, and wings (30) as well as the gain of a remarkable ability to

respire directly from water (6). Preliminary evidence suggests that subterranean members of these

lineages are less tolerant of thermal extremes compared to epigean relatives (36), but the molecular

mechanisms underlying such tolerances–and genomic changes associated with a subterranean

transition in these animals more broadly–remain unknown. An investigation into the potential link

between the above-mentioned mode of respiration, oxygen delivery, and heat tolerance is also

lacking (37).

In the present study, we focus on two members of the Hydroporini: *Paroster*

nigroadumbratus (Clark), an epigean species endemic to South Australia, and the subterranean

Paroster macrosturtensis (Watts & Humphreys) found exclusively in a single calcrete at Sturt

Meadows in the Yilgarn region of WA. As subterranean dytiscids likely descended from only a

handful of epigean lineages, meaningful comparisons can be made between these taxa despite their

distributions being geographically disjunct and their divergence *ca.* 15 Mya (32). A recent study

showed that *P. macrosturtensis* has a reduced upper critical thermal maximum (CT_{max}) of 38.3ºC

compared to other epigean dytiscids (42–44.5ºC) (36), mirroring previous results for other cave

beetle species (23). These findings suggest *P. macrosturtensis* is unlikely to reach its thermal critical

maximum under current climate change predictions. However, given the thermal stability in its

environment, it remains unknown as to whether *P. macrosturtensis* might have a modified HSR

compared to its epigean relatives and if exposure to high temperatures may nonetheless induce

significant stress in this subterranean species. Here, we present transcriptomic data from individuals

of *P. macrosturtensis* and *P. nigroadumbratus* subjected to varying degrees of heat shock following

the results of Jones *et al*. (36). In the present study we specifically aimed 1) to generate a high-

quality, near-complete reference transcriptome for *P. nigroadumbratus* and 2) using this dataset,

characterise and compare the HSR of *P. nigroadumbratus* and *P. macrosturtensis*, specifically with

respect to which genes are differentially expressed and the conditions under which this occurs.

2 | MATERIALS AND METHODS

2.1 | Taxon sampling, experimental design, and cDNA sequencing

Beetle specimens (*P. macrosturtensis*, n=11; *P. nigroadumbratus*, n=13) were sourced as described

in a previous study and subjected to heat stress using an aquarium setup described therein (36).

Specimens were placed in one of three groups: in a controlled-temperature cabinet at 25ºC (hereafter

control), in vials within the experimental setup at 25ºC, and in vials within the experimental setup

ramped to 35ºC. The control temperature of 25ºC was selected by (36) to reflect the approximate

average groundwater temperatures of the aquifer that *P. macrosturtensis* is found in. Following

treatment, the individuals were placed in liquid nitrogen for RNA sequencing immediately after

exposure to control temperatures and thermal extremes. RNA extractions were performed using

single whole bodies prior to the synthesis and sequencing of barcoded cDNA samples. Quality

control of sequence data was performed using Trim Galore with default settings v.0.4.1

(http://bioinformatics.babraham.ac.uk/projects/trim_galore). More information regarding specimen

collection and husbandry, experimental design, cDNA sequencing, and phylogenetic analysis can be

found in the electronic supplementary material.

2.2 | Assembly of *Paroster nigroadumbratus* reference transcriptome

A reference transcriptome for *P. nigroadumbratus* was *de novo* assembled using Trinity v.2.5.1

using default settings (38) and derived from unpublished raw RNA-seq data from (39). Resulting

transcriptome completeness was validated using BUSCO v.5 (40,41), and functional annotation was

performed using TransDecoder v.5.5.0 (42) and the Trinotate v.3.2.1 pipeline (38), which employs

SQLite (http://sqlite.org/index.html), BLAST v.2.7.1 (43), and HMMER v.3.2 (http://hmmer.org).

Redundancy within the transcriptome assembly was reduced by retaining only the longest isoform

for each Trinity gene identifier, following (10), ensuring differential expression analysis was

performed at the Trinity 'gene' level rather than that of isoforms. This subsetted dataset was used in

146 downstream analyses, though we note this approach limited our ability to examine alternative splice

147 variants.

2.3 | Differential expression and Gene Ontology enrichment analysis

The above reference transcriptome was indexed and raw reads generated in this study from both *P.*

macrosturtensis and *P. nigroadumbratus* were quasi-mapped to it and normalised using Salmon

v.1.1.0 (44). Differential expression analysis was performed using edgeR v.3.32.1 (45) alongside

Gene Ontology (GO) enrichment analysis, executed using Trinity helper scripts. Differential

expression and enrichment/depletion of GO terms was gauged using two comparisons per species

(Figure 1). First, for each species the group from the experimental setup at 25ºC was compared to the

control group outside of the setup at 25ºC (hereafter comparison 1). Second, for each species the

group in the experimental setup at 35ºC was compared to the group in the experimental setup at 25ºC

(hereafter comparison 2). A diagrammatic representation of these comparisons is shown in Figure 1.

These comparisons were designed to gauge the impact of the presence of the experimental setup

alone and an increase in temperature, respectively. A full description of this process can be found in

the electronic supplementary material.

2.4 | Phylogenetic analysis of putative heat shock proteins

164 Putative heat shock protein-encoding genes were aligned with coleopteran orthologs sourced from

165 the OrthoDB v.10.1 catalogue (46) and GenBank to confirm their identity (Table S1). Phylogenetic

166 inference for each HSP family was performed in RAxML v.8.2.12 (47).

FIGURE 1 *Paroster* beetles included in this study and the experimental design used to assess differential gene expression associated with thermal extremes alone, as opposed to solely the presence of the experimental setup, following (36). Only intraspecific comparisons were made when assessing differential gene expression in our analyses; interspecific comparisons were made post-hoc. Photographs by Chris Watts and Howard Hamon.

3 | RESULTS

3.1 | A high-quality reference transcriptome for *Paroster nigroadumbratus* enables the characterisation of genes involved in the heat shock response

Here we present a high-quality, near-complete transcriptome for the epigean beetle *P. nigroadumbratus*. This dataset consisted of 75,045,266 paired-end reads (72,266,264 following quality control measures), 84% of which were incorporated into 134,246 *de novo* assembled transcripts representing 60,683 unique Trinity 'genes'and 41,979 predicted ORFs. According to the assessment using BUSCO, this transcriptome was 87.66% complete with respect to complete core arthropod genes and 96.74% complete when considering partial genes. Of these transcripts, 47,810 were able to be functionally annotated using the Trinotate pipeline. Subsetting our predicted peptide dataset to include only the longest isoforms per Trinity gene identifier, allowing us a proxy with which to perform our downstream analyses at the gene level, resulted in 14,897 predicted ORFs (with 11,609, or ~77%, having some level of annotation).

3.2 | Differential expression analysis reveals distinct expression profiles associated with the heat shock response in *Paroster*

We compared expression profiles between members of the same species, subjected to different conditions, to assess genes differentially expressed in response to the presence of the experimental setup employed here (comparison 1) or an increase in temperature in that setup (comparison 2) (Figure 2). Parallels and contrasts between *P. nigroadumbratus* and *macrosturtensis* were then assessed post-hoc. Differential expression analysis using edgeR identified a total of 723 differentially expressed (DE) genes in the epigean *P. nigroadumbratus* samples and 157 in the subterranean *P. macrosturtensis*, with the two species exhibiting complex and markedly different expression profiles (Figures 2, 3, S2, S3). *P. nigroadumbratus* consistently differentially expressed a greater number of

genes than *P. macrosturtensis*: 147 and 67 genes were differentially expressed in comparison 1 and

89 and 51 genes were differentially expressed in comparison 2, respectively. The presence of the

experimental setup and an increase in water temperature caused both species to differentially express

genes involved in the heat shock response. Contrasting expression profiles with respect to the genes

involved were not only observed between the two species (as above) but also in the response of each

species to these two different stressors (Figure 2).

In comparison 1, the surface species *P. macrosturtensis* significantly upregulated a HSP gene

putatively annotated as *Hsp68*, encoding a major heat shock protein in the HSP70 family, relative to

the control. *Paroster nigroadumbratus* upregulated the HSP *Hsp68* and downregulated the sHSP

l(2)efl in the experimental setup relative to the control. None of the HSP transcripts differentially

expressed by *P. nigroadumbratus* in response to the presence of the experimental setup only were

differentially expressed by *P. macrosturtensis* or by *P. nigroadumbratus* in comparison 2.

In comparison 2, both species upregulated the HSP genes *Hsc70-4* and *Hsp83* as well as the

cochaperones *Samui*, *DnaJ*, *AHSA1,* and *Unc45a* relative to groups at the lower temperature of 25ºC.

Both *P. nigroadumbratus* and *macrosturtensis* also upregulated separate Trinity "genes" annotated as

Hsp68 each at 35 °C relative to the 25 °C treatment (Fig. 2), likely representing closely related loci

that are yet to be comprehensively characterised in the beetles. Additional differentially expressed

genes unique to each species at 35ºC relative to the 25ºC treatment included: 1) downregulation of

the HSPs *Hsp60A* and *Hsp68* in *Paroster macrosturtensis*, and 2) the upregulation of the sHSP

l(2)efl, the HSP *Hsp68*, and the HSP cognate *Hsc70-3* in *P. nigroadumbratus*. Other annotated genes

potentially involved in the HSR, such as *Hsc70-2*, *Hsc70-5*, *Trap1*, and *Hsp90b1*, were not

differentially expressed in either species (Figure 4). A full list of differentially expressed genes

shared between (or unique to) the two beetle species under different conditions is available in Table

S2. Proteins unrelated to the HSR, yet widely differentially expressed in our dataset (i.e., with

reoccurring annotations across different Trinity 'genes') included those involved in the transport of

lipids and nutrient storage, such as vitellogenin and apolipophorins.

Gene Ontology enrichment analysis enables a high-order approximation of the functional

consequences of differentially expressed genes. We did not observe any consistent depletion of GO

terms associated with downregulated genes, but enriched terms were reflective of our differential

expression results above (Figure 3). In both species, terms associated with the HSR were enriched in

association with an increased temperature at 35ºC (e.g., ATPase activator activity [GO:0001671],

chaperone binding [GO:0051087], and unfolded protein binding [GO:0051082]). The most common

remaining Gene Ontology terms included lipid transporter activity (GO:0005319), nutrient reservoir

activity (GO:0045735), metal ion binding (GO:0046872), and zinc ion binding (GO:0008270).

3.3 | Phylogenetic analyses of HSPs

Our HSP nucleotide alignments consisted of 12 sequences across 1,743 bp (HSP60 family), 67

sequences across 3,669 bp (HSP70 family), 29 sequences across 3,741 bp (HSP90 family) and 23

sequences across 867 bp (sHSP family). Sequences clustered by gene with strong node support in our

phylogeny (Figure 4). All sequences from the *Paroster* species examined here were recovered as

nested within these clades, confirming the orthology of annotated HSPs, with the exception of a

transcript present in both species and inferred as sister to the *Hsp68*+*Hsc70-2* clades. This gene was

not differentially expressed in response to heat shock in our dataset.

FIGURE 2 Differentially expressed (DE) genes in *Paroster macrosturtensis* and *nigroadumbratus* in comparison 1 (the experimental setup used in this study relative to the control, black circles) or comparison 2 (35ºC within the setup relative to 25ºC in the setup, red circles). Comparisons are ordered by total number of DE genes. A) DE genes shared (circles linked by lines) or unique to (unlinked circles) each comparison per species, summed in the bar graph above. Up- or downregulated HSPs or HSP cochaperones are shown for each group. The gene *Hsp68* being named more than once in different groups refers to separate transcripts sharing the same putative annotation (see Table S2). B) Total DE genes for each species under different conditions. HSP gene names were sourced from Trinotate annotations and orthology was validated using phylogenetic analysis (Figure 4).

FIGURE 3 REVIGO (48) treemaps showing enriched Gene Ontology (GO) terms associated with the differential

expression of genes shown in Figure 2. Treemaps are labelled as comparison 1 (the experimental setup used in this study

relative to the control) or 2 (35℃ within the setup relative to 25℃ in the setup). The size of squares is proportional to the

p-value associated with differential expression of respective genes. Similar GO terms share a colour and are represented

in white text by the largest square per group. GO terms associated with genes involved in the heat shock response are

indicated by a star.

FIGURE 4 Phylogenies of heat shock Trinity 'genes' inferred using RAxML to validate the identity of putative HSP

orthologs in *Paroster* species. Trees are as follows: A) sHSP family, B) HSP60, C) HSP90, D) HSP70. Scale bar is in

substitutions/site; *BS* = bootstrap node support. Tip labels show transcript names/GenBank accession numbers. Tip

names for *P. macrosturtensis* are shared with the reference *P. nigroadumbratus* transcript reads were assembled against.

Tip names with stars specifically refer to genes differentially expressed in the presence of the experimental setup alone

relative to the control (comparison 1) or at 35ºC relative to 25ºC in the experimental setup (comparison 2).

4 | DISCUSSION

Here, we have comprehensively characterised the heat shock response at the molecular level in the subterranean diving beetle *Paroster macrosturtensis* and one of its surface-dwelling relatives. Using a near-complete reference transcriptome for *P. nigroadumbratus*–the first such dataset for a member of the Hydroporinae–we performed differential expression and GO enrichment analysis to explore genes putatively involved in the HSR. Our results demonstrate that both the epigean *P. nigroadumbratus* and subterranean *P. macrosturtensis* have an inducible HSR, in agreement with implications of previous survival experiments for the genus (36). Putative orthologs of HSP cochaperones, sHSP, HSP60, HSP70, and HSP90 genes were accounted for in our differential expression analysis. However, the conditions under which this response is activated differs between species, and *P. macrosturtensis* notably differentially expressed just over half of the number of genes compared to *P. nigroadumbratus* in response to a rise in temperature.

4.1 | Heat shock-induced gene expression in *Paroster*

HSPs identified as differentially expressed in this study support past results for beetle species and other insects more broadly. A major trend in our results concerned the upregulation of HSP70 genes at high temperatures, particularly *Hsp68*; HSP70s are highly expressed in response to heat shock in other beetle species, including cave-adapted subterranean taxa (28,49,50), and work in concert with sHSPs and HSP90s (51–54). In addition to the heat shock proteins, we also observed the upregulation of the heat shock cognates *Hsc70-3* and *Hsc70-4* and putative cochaperones *Tsc2, Samui*, *DnaJ*, *AHSA1*, *Unc45a* at 35ºC relative to the 25ºC treatment. To our knowledge there has been no documentation of a coleopteran heat shock cognate being upregulated in response to increasing temperature, though evidence exists for the parasitic wasp *Pteromalus*, in which *hsc70* is induced by heavy metal poisoning and starvation in addition to thermal extremes (55). Heat shock cognates are also upregulated during diapause in silkworm eggs (56) and young bumble bee queens

(57), potentially playing a cryoprotective role in these species. Cochaperones are less well

characterised in insects, but evidence for their upregulation in response to heat shock has been

documented in hemipterans and ants (58–60). We observed the downregulation of several heat shock

proteins in both species in the presence of the experimental setup relative to the control, and at high

temperatures relative to the 25ºC treatment. Both sHSPs and HSP70s have been documented as being

downregulated during periods of heat stress in other insects, e.g. in silk moths (61). In *P.*

nigroadumbratus this was restricted to the sHSP *l(2)efl* and cochaperone *Tsc2* in the experimental

setup-only comparison. In contrast, the HSPs *Hsp60A* and *Hsp68* were downregulated *in P.*

macrosturtensis at high temperatures relative the 25ºC treatment.

4.2 | Expression profiles reflect differing thermal tolerances

Our molecular data mirrors previously documented reduced thermal tolerances in subterranean

insects such as *P. macrosturtensis*. The species differentially expressed far fewer genes in response

to 35ºC relative to the 25ºC treatment compared to *P. nigroadumbratus*; similarly reduced numbers

of differentially expressed genes have also been associated with lower thermal tolerances in other

organisms such as fish (62), lizards (63), rotifers (64), red algae (65), and plants (66,67), though we

note the inverse (or alternatively, no clear pattern) has been observed in a number of cases,

potentially reflecting lower levels of stress as opposed to an inability to mount a HSR (68,69).

In keeping with the above findings, *P. macrosturtensis* also differentially expressed far fewer

genes than its epigean counterpart in response to the presence of the experimental setup alone

relative to the control (Figure 2). HSPs are known to be involved in responding to a wide range of

stressors (70–72), and the involvement of such genes is not surprising in stress unrelated to

temperature; individuals being moved into the experimental setup employed here may have induced

stress from handling, for example. The greater number of genes differentially expressed by *P.*

nigroadumbratus in this scenario may suggest *P. macrosturtensis* is potentially less able to robustly

respond to ambient stressors more broadly (i.e., environmental disturbances). Such a scenario is

supported by past work showing subterranean species are sensitive to ambient stressors under

otherwise non-stressful temperatures (73), and being an epigean species, *P. nigroadumbratus* is

presumably exposed to far more dramatic environmental fluctuations (in addition to more variable

temperatures) on a regular basis than a subterranean species such as *P. macrosturtensis*. We also note

that in the presence of the experimental setup relative to the control, *P. macrosturtensis* upregulated

the same *Hsp68*-annotated Trinity 'gene' implicated in responses to heat-induced stress in both

species (Figure 2), whereas HSPs differentially expressed by *P. nigroadumbratus* under the same

conditions did not overlap with those in other groups.

4.3 | Heat shock and the climatic variability hypothesis

The dataset we present here adds to a growing body of knowledge concerning the HSR in organisms

that inhabit thermally stable environments. Central to discourse on this topic is the climatic

variability hypothesis, which posits that the thermal tolerance of a taxon is positively correlated with

its temperature ranges encountered in nature (74). This hypothesis implies species from extremely

stable thermal environments can no longer tolerate temperature extremes, and has been demonstrated

in a wide variety of organisms that have either lost or possess a reduced HSR, such as cnidarians

(75), limpets (76), amphipods and sea stars (77), and midges (78). In contrast, species that inhabit

areas with a broader range of climatic conditions would be expected to be more robust in the face of

environmental fluctuations (79). While *P. macrosturtensis* does have a lower thermal tolerance

compared to *P. nigroadumbratus*, in line with the above hypothesis, it nonetheless has retained a

HSR at high temperatures per our transcriptomic data. Similar studies have shown certain

groundwater-dwelling organisms display an inducible HSR in response to conditions far warmer than

they would encounter in nature (20,21). The HSR of these species, as well as *P. macrosturtensis*,

might be retained at such high temperatures for a variety of reasons, including the fact that the

species has not occupied its respective environments for a sufficient length of time in evolutionary

terms for their HSR to be lost, e.g. via adaptive processes or a relaxation of purifying selection (36).

The latter scenario is plausible as *P. macrosturtensis* is also known to have retained the ability to

detect light despite inhabiting an aphotic environment for over ~3 million years (80).

4.4 | Conservation implications

While the retention of a HSR in both species examined here supports the physiological findings of

Jones *et al*. (36), almost half (4 out of 10 assayed) of the *P. macrosturtensis* cohort did not survive 24

hours after heat shock in that study. It therefore remains to be seen if the species can tolerate such

extremes in the long term. Indeed, even cave beetles considered stenothermal–those that are only

capable of surviving within an extremely narrow temperature range–have retained the HSR (23), but

nonetheless cannot survive at extreme temperatures for long periods (>7 days) compared to epigean

relatives (25,28). Threatening processes that *P. nigroadumbratus* and *macrosturtensis* are both at risk

of experiencing in their fragile habitats might impact the latter species far more negatively as a

result.

447   Temperature rises of up to 5ºC by the end of the century compared to pre-industrial levels

may occur in central Western Australia per current climate change projections (81). Water

temperatures in aquifers are generally cooler and more stable than, but are nonetheless coupled with,

conditions above-ground, and are also predicted to warm as regional temperatures increase (82–84).

The subterranean habitat of these insects is therefore unlikely to shield them from the impacts of a

warming world. The fact that *P. macrosturtensis* appears to be unable to mount as robust a HSR

compared to *P. nigroadumbratus*, and therefore may experience a significantly higher amount stress

compared to epigean species in the face of high temperatures, has conservation implications for the

understudied fauna of the Australian Yilgarn and beyond.

Datasets such as these are especially pertinent for subterranean invertebrates found in the

Yilgarn region—including *P. macrosturtensis* and its subterranean relatives, in addition to

crustaceans such as isopods and amphipods—as the groundwater in their calcrete habitats is heavily

utilised for water extraction by industry (85,86). As short-range endemics to the extreme, such

species are not only at risk of habitat degradation via climate change, but from the direct intersection

of shallow aquifers with e.g. mining activities and through the drawdown of groundwater beneath

calcretes at greater depths (87). In addition to reflecting the reduced thermal tolerances of Australian

subterranean dytiscids, the molecular data we presented here for *P. macrosturtensis* also suggests a

potentially weaker response in the face of other environmental disturbances unrelated to temperature.

These factors have the potential to render *P. macrosturtensis* more vulnerable to both of the above

threatening processes compared to epigean relatives, with implications for subterranean fauna more

broadly. An increased knowledge of the assumed fragility of Australian subterranean invertebrates in

the face of these stressors is therefore crucial for informing future conservation management plans

for these animals and their fragile habitats.

5 | CONCLUSIONS

Our findings demonstrate the reduced thermal tolerance of the subterranean species *P.*

macrosturtensis compared to its epigean relatives is reflected, and further clarified by, transcriptomic

data. While our data are supported by past physiological evidence that demonstrated *P.*

macrosturtensis could survive at high temperatures, albeit not to the limits of epigean species (36),

the present study adds a new layer to this narrative. *P. macrosturtensis* might possess increased

mortality in the face of high temperatures because the species differentially expresses far fewer genes

in response to heat shock compared to the epigean relative *P. nigroadumbratus*, suggesting it may be

unable to mount as robust a heat shock response. While *P. macrosturtensis* might be able to survive

at temperatures far above those it encounters in nature for short periods, as detailed by Jones *et al*.

(36), our results suggest the species also experiences a weaker transcriptomic response to factors

unrelated to temperature (i.e. the presence of the experimental setup employed here) relative to *P.*

nigroadumbratus. As such, *P. macrosturtensis* may not be as well-equipped to survive higher

temperatures and other threatening processes, such as disturbances to surrounding groundwater, in

the long term compared to surface-dwelling members of *Paroster*. Future work in this system will

ideally assess a far greater number of dytiscid species to further explore the trends we observe here.

As the present study did not consider the role of isoforms in the heat shock response of these

animals, broader studies could examine these responses to heat stress at a finer scale by conducting

differential expression analysis on the transcript, as opposed to gene, level.

REFERENCES

1. Tierney SM, Friedrich M, Humphreys WF, Jones TM, Warrant EJ, Wcislo WT. Consequences
of evolutionary transitions in changing photic environments. Austral Entomol. 2017;56(1):23–
46.

2. Culver DC, Pipan T. Shallow Subterranean Habitats: Ecology, Evolution, and Conservation
[Internet]. Shallow Subterranean Habitats. Oxford University Press; 2014 [cited 2021 Apr 13].
Available from:
https://oxford.universitypressscholarship.com/view/10.1093/acprof:oso/9780199646173.001.00
01/acprof-9780199646173

3. Merritt DJ, Clarke AK. Synchronized circadian bioluminescence in cave-dwelling
Arachnocampa tasmaniensis (Glowworms). J Biol Rhythms. 2011 Feb;26(1):34–43.

4. Beale AD, Whitmore D, Moran D. Life in a dark biosphere: a review of circadian physiology in
"arrhythmic" environments. J Comp Physiol [B]. 2016;186(8):947–68.

5. Walker JA, Rose HA. Oothecal structure and male genitalia of the Geoscapheinae and some
Australian Panaesthia Serville (Blattodea: Blaberidae). Aust J Entomol. 1998;37(1):23–6.

6. Jones KK, Cooper SJB, Seymour RS. Cutaneous respiration by diving beetles from
underground aquifers of Western Australia (Coleoptera: Dytiscidae). J Exp Biol. 2019 Apr
1;222(7):jeb196659.

7. Cieslak A, Fresneda J, Ribera I. Life-history specialization was not an evolutionary dead-end in
Pyrenean cave beetles. Proc R Soc B Biol Sci. 2014 Apr 22;281(1781):20132978.

8. Cieslak A, Fresneda J, Ribera I. Developmental constraints in cave beetles. Biol Lett. 2014 Oct
31;10(10):20140712.

9.   Yang J, Chen X, Bai J, Fang D, Qiu Y, Jiang W, et al. The Sinocyclocheilus cavefish genome
provides insights into cave adaptation. BMC Biol. 2016 Jan 4;14(1):1.

10.   Balart-García P, Cieslak A, Escuer P, Rozas J, Ribera I, Fernández R. Smelling in the dark:
Phylogenomic insights into the chemosensory system of a subterranean beetle. Mol Ecol
[Internet]. 2021 [cited 2021 May 7]; Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.15921

11.   Badino G. UNDERGROUND METEOROLOGY - "What's the weather underground?" Acta
Carsologica [Internet]. 2010 Dec 1 [cited 2021 Apr 13];39(3). Available from: https://ojs.zrc-
sazu.si/carsologica/article/view/74

12.   Hüppop K. Adaptation to Low Food. In: White WB, Culver DC, editors. Encyclopedia of Caves
(Second Edition) [Internet]. Amsterdam: Academic Press; 2012 [cited 2022 Jun 27]. p. 1–9.
Available from: https://www.sciencedirect.com/science/article/pii/B9780123838322000013

13.   Bourges F, Genthon P, Genty D, Lorblanchet M, Mauduit E, D'Hulst D. Conservation of
prehistoric caves and stability of their inner climate: Lessons from Chauvet and other French
caves. Sci Total Environ. 2014 Sep 15;493:79–91.

14.   Mammola S, Piano E, Cardoso P, Vernon P, Domínguez-Villar D, Culver DC, et al. Climate
change going deep: The effects of global climatic alterations on cave ecosystems. Anthr Rev.
2019 Apr 1;6(1–2):98–116.

15.   Jindal S, Dudani AK, Singh B, Harley CB, Gupta RS. Primary structure of a human
mitochondrial protein homologous to the bacterial and plant chaperonins and to the 65-
kilodalton mycobacterial antigen. Mol Cell Biol. 1989 May;9(5):2279–83.

16.   Richter K, Haslbeck M, Buchner J. The Heat Shock Response: Life on the Verge of Death. Mol
Cell. 2010 Oct 22;40(2):253–66.

17.   Somero GN. The cellular stress response and temperature: Function, regulation, and evolution.
J Exp Zool Part Ecol Integr Physiol. 2020;333(6):379–97.

18.   Tomanek L. Variation in the heat shock response and its implication for predicting the effect of
global climate change on species' biogeographical distribution ranges and metabolic costs. J
Exp Biol. 2010 Mar 15;213(6):971–9.

19.   Colado R, Pallarés S, Fresneda J, Mammola S, Rizzo V, Sánchez-Fernández D. Climatic
stability, not average habitat temperature, determines thermal tolerance of subterranean beetles.
Ecology. 2022;n/a(n/a):e3629.

20.   Colson-Proch C, Morales A, Hervant F, Konecny L, Moulin C, Douady CJ. First cellular
approach of the effects of global warming on groundwater organisms: a study of the HSP70
gene expression. Cell Stress Chaperones. 2010 May;15(3):259–70.

21.   Mermillod-Blondin F, Lefour C, Lalouette L, Renault D, Malard F, Simon L, et al. Thermal
tolerance breadths among groundwater crustaceans living in a thermally constant environment.
J Exp Biol. 2013 May 1;216(9):1683–94.

22. Pallarés S, Colado R, Pérez-Fernández T, Wesener T, Ribera I, Sánchez-Fernández D. Heat
tolerance and acclimation capacity in subterranean arthropods living under common and stable
thermal conditions. Ecol Evol. 2019;9(24):13731–9.

23. Pallarés S, Colado R, Botella-Cruz M, Montes A, Balart-García P, Bilton DT, et al. Loss of heat
acclimation capacity could leave subterranean specialists highly sensitive to climate change.
Anim Conserv [Internet]. 2020 [cited 2021 Jun 7]; Available from:
https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/acv.12654

24. Raschmanová N, Šustr V, Kováč Ľ, Parimuchová A, Devetter M. Testing the climatic
variability hypothesis in edaphic and subterranean Collembola (Hexapoda). J Therm Biol. 2018
Dec;78:391–400.

25. Rizzo V, Sánchez-Fernández D, Fresneda J, Cieslak A, Ribera I. Lack of evolutionary
adjustment to ambient temperature in highly specialized cave beetles. BMC Evol Biol. 2015
Feb 4;15(1):10.

26. Gibert J, Deharveng L. Subterranean ecosystems: a truncated functional biodiversity.
BioScience. 2002 Jun 1;52(6):473–81.

27. Humphreys W. Diversity Patterns in Australia. In: Encyclopedia of Caves. 2012. p. 203–19.

28. Bernabò P, Latella L, Jousson O, Lencioni V. Cold stenothermal cave-dwelling beetles do have
an HSP70 heat shock response. J Therm Biol. 2011 Apr 1;36(3):206–8.

29. Mammola S, Amorim IR, Bichuette ME, Borges PAV, Cheeptham N, Cooper SJB, et al.
Fundamental research questions in subterranean biology. Biol Rev. 2020;95(6):1855–72.

30. Cooper SJB, Hinze S, Leys R, Watts CHS, Humphreys WF. Islands under the desert: molecular
systematics and evolutionary origins of stygobitic water beetles (Coleoptera : Dytiscidae) from
central Western Australia. Invertebr Syst. 2002;16(4):589–90.

31. Watts CHS, Humphreys WF. Fourteen New Dytiscidae (Coleoptera) of the Genera
Limbodessus Guignot, Paroster Sharp, and Exocelina Broun from Underground Waters in
Australia. Trans R Soc S Aust. 2009 Jan 1;133(1):62–107.

32. Leijs R, Nes E, Watts C, Cooper S, Humphreys W, Hogendoorn K. Evolution of Blind Beetles
in Isolated Aquifers: A Test of Alternative Modes of Speciation. PloS One. 2012 Mar
30;7:e34260.

33. Langille BL, Hyde J, Saint KM, Bradford TM, Stringer DN, Tierney SM, et al. Evidence for
speciation underground in diving beetles (Dytiscidae) from a subterranean archipelago.
Evolution. 2021;75(1):166–75.

34. Leys R, Watts CHS, Cooper SJB, Humphreys WF. Evolution of subterranean diving beetles
(Coleoptera: Dytiscidae: Hydroporini, Didessini) in the arid zone of Australia. Evolution. 2003
Dec;57(12):2819–34.

35. Toussaint EFA, Condamine FL, Hawlitschek O, Watts CH, Porch N, Hendrich L, et al.
Unveiling the Diversification Dynamics of Australasian Predaceous Diving Beetles in the
Cenozoic. Syst Biol. 2015 Jan 1;64(1):3–24.

36. Jones KK, Humphreys WF, Saccò M, Bertozzi T, Austin AD, Cooper SJB. The critical thermal
maximum of diving beetles (Coleoptera: Dytiscidae): a comparison of subterranean and
surface-dwelling species. Curr Res Insect Sci. 2021 Jan 1;1:100019.

37. Verberk WCEP, Overgaard J, Ern R, Bayley M, Wang T, Boardman L, et al. Does oxygen limit
thermal tolerance in arthropods? A critical review of current evidence. Comp Biochem Physiol
A Mol Integr Physiol. 2016 Feb 1;192:64–78.

38. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Trinity:
reconstructing a full-length transcriptome without a genome from RNA-Seq data. Nat
Biotechnol. 2011 May 15;29(7):644–52.

39. Tierney SM, Cooper SJB, Saint KM, Bertozzi T, Hyde J, Humphreys WF, et al. Opsin
transcripts of predatory diving beetles: a comparison of surface and subterranean photic niches.
R Soc Open Sci. 2015;2(1):140386.

40. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinforma Oxf
Engl. 2015 Oct 1;31(19):3210–2.

41. Nishimura O, Hara Y, Kuraku S. gVolante for standardizing completeness assessment of
genome and transcriptome assemblies. Bioinformatics. 2017 Nov 15;33(22):3635–7.

42. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo
transcript sequence reconstruction from RNA-Seq: reference generation and analysis with
Trinity. Nat Protoc [Internet]. 2013 Aug [cited 2021 May 10];8(8). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3875132/

43. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC Bioinformatics. 2009 Dec 15;10(1):421.

44. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods. 2017 Apr;14(4):417–9.

45. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010 Jan 1;26(1):139–40.

46. Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, et al. OrthoDB v10:
sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for
evolutionary and functional annotations of orthologs. Nucleic Acids Res. 2019 Jan
8;47(D1):D807–11.

47. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014 May 1;30(9):1312–3.

48. Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO Summarizes and Visualizes Long Lists of
Gene Ontology Terms. PLOS ONE. 2011 Jul 18;6(7):e21800.

49. Chen M, Zhang N, Jiang H, Meng X, Qiang K, Wang J. Transcriptional regulation of heat
shock protein 70 genes by class I histone deacetylases in the red flour beetle, Tribolium
castaneum. Insect Mol Biol. 2020;29(2):221–30.

50. Yocum GD. Differential expression of two HSP70 transcripts in response to cold shock,
thermoperiod, and adult diapause in the Colorado potato beetle. J Insect Physiol. 2001 Sep
1;47(10):1139–45.

51. Cai Z, Chen J, Cheng J, Lin T. Overexpression of Three Heat Shock Proteins Protects
Monochamus alternatus (Coleoptera: Cerambycidae) From Thermal Stress. J Insect Sci
[Internet]. 2017 Nov 1 [cited 2021 Jun 7];17(113). Available from:
https://doi.org/10.1093/jisesa/iex082

52. Li MY, Huang Y, Lei X, Xu CT, Li B, Chen DX, et al. Identification of six heat shock protein
70 genes in Lasioderma serricorne (Coleoptera: Anobiidae) and their responses to temperature
stress. J Asia-Pac Entomol [Internet]. 2021 May 9 [cited 2021 Jun 7]; Available from:
https://www.sciencedirect.com/science/article/pii/S1226861521000674

53. Yang WJ, Xu KK, Cao Y, Meng YL, Liu Y, Li C. Identification and Expression Analysis of
Four Small Heat Shock Protein Genes in Cigarette Beetle, Lasioderma serricorne (Fabricius).
Insects. 2019 May;10(5):139.

54. Yuan X, Zhou WW, Zhou Y, Liu S, Lu F, Yang MF, et al. Composition and Expression of Heat
Shock Proteins in an Invasive Pest, The Rice Water Weevil (Coleoptera: Curculionidae). Fla
Entomol. 2014 Jun;97(2):611–9.

55. Wang H, Li K, Zhu JY, Fang Q, Ye GY, Wang H, et al. CLONING AND EXPRESSION
PATTERN OF HEAT SHOCK PROTEIN GENES FROM THE ENDOPARASITOID WASP,
Pteromalus puparum IN RESPONSE TO ENVIRONMENTAL STRESSES. Arch Insect
Biochem Physiol. 2012;79(4–5):247–63.

56. Sasibhushan S, Ponnuvel KM, Vijayaprakash NB. Diapause specific gene expression in the
eggs of multivoltine silkworm Bombyx mori, identified by suppressive subtractive
hybridization. Comp Biochem Physiol B Biochem Mol Biol. 2012 Apr 1;161(4):371–9.

57. Kim BG, Shim JK, Kim DW, Kwon YJ, Lee KY. Tissue-specific variation of heat shock
protein gene expression in relation to diapause in the bumblebee Bombus terrestris. Entomol
Res. 2008;38(1):10–6.

58. Huang LH, Kang L. Cloning and interspecific altered expression of heat shock protein genes in
two leafminer species in response to thermal stress. Insect Mol Biol. 2007;16(4):491–500.

59. Hull JJ, Geib SM, Fabrick JA, Brent CS. Sequencing and De Novo Assembly of the Western
Tarnished Plant Bug (Lygus hesperus) Transcriptome. PLOS ONE. 2013 Jan 24;8(1):e55105.

60. Tonione MA, Bi K, Tsutsui ND. Transcriptomic signatures of cold adaptation and heat stress in
the winter ant (Prenolepis imparis). PLOS ONE. 2020 Oct 1;15(10):e0239558.

61. Sakano D, Li B, Xia Q, Yamamoto K, Fuiji H, Aso Y. Genes Encoding Small Heat Shock
Proteins of the Silkworm, Bombyx mori. Biosci Biotechnol Biochem. 2006 Oct
23;70(10):2443–50.

62. Narum SR, Campbell NR. Transcriptomic response to heat stress among ecologically divergent
populations of redband trout. BMC Genomics. 2015 Feb 21;16(1):103.

63. Akashi HD, Cádiz Díaz A, Shigenobu S, Makino T, Kawata M. Differentially expressed genes
associated with adaptation to different thermal environments in three sympatric Cuban Anolis
lizards. Mol Ecol. 2016;25(10):2273–85.

64. Paraskevopoulou S, Dennis AB, Weithoff G, Tiedemann R. Temperature-dependent life history
and transcriptomic responses in heat-tolerant versus heat-sensitive Brachionus rotifers. Sci Rep.
2020 Aug 6;10(1):13281.

65. Wang W, Lin Y, Teng F, Ji D, Xu Y, Chen C, et al. Comparative transcriptome analysis
between heat-tolerant and sensitive Pyropia haitanensis strains in response to high temperature
stress. Algal Res. 2018 Jan 1;29:104–12.

66. Sun M, Lin C, Zhang A, Wang X, Yan H, Khan I, et al. Transcriptome sequencing revealed the
molecular mechanism of response of pearl millet root to heat stress. J Agron Crop Sci.
2021;207(4):768–73.

67. Jin J, Yang L, Fan D, Liu X, Hao Q. Comparative transcriptome analysis uncovers different
heat stress responses in heat-resistant and heat-sensitive jujube cultivars. PLOS ONE. 2020 Sep
21;15(9):e0235763.

68. DeBiasse MB, Kelly MW. Plastic and Evolved Responses to Global Change: What Can We
Learn from Comparative Transcriptomics? J Hered. 2016 Jan 1;107(1):71–81.

69. Perez R, de Souza Araujo N, Defrance M, Aron S. Molecular adaptations to heat stress in the
thermophilic ant genus Cataglyphis. Mol Ecol [Internet]. 2021 [cited 2021 Oct 29];n/a(n/a).
Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.16134

70. Sørensen JG, Kristensen TN, Loeschcke V. The evolutionary and ecological role of heat shock
proteins. Ecol Lett. 2003;6(11):1025–37.

71. Sang W, Ma WH, Qiu L, Zhu ZH, Lei CL. The involvement of heat shock protein and
cytochrome P450 genes in response to UV-A exposure in the beetle Tribolium castaneum. J
Insect Physiol. 2012 Jun 1;58(6):830–6.

72. King AM, MacRae TH. Insect Heat Shock Proteins During Stress and Diapause. Annu Rev
Entomol. 2015;60(1):59–75.

73. Pallarés S, Sanchez-Hernandez JC, Colado R, Balart-García P, Comas J, Sánchez-Fernández D.
Beyond survival experiments: using biomarkers of oxidative stress and neurotoxicity to assess
vulnerability of subterranean fauna to climate change. Conserv Physiol. 2020 Jan
1;8(1):coaa067.

74. Stevens GC. The Latitudinal Gradient in Geographical Range: How so Many Species Coexist in
the Tropics. Am Nat. 1989;133(2):240–56.

75. Brennecke T, Gellner K, Bosch TC. The lack of a stress response in Hydra oligactis is due to
reduced hsp70 mRNA stability. Eur J Biochem. 1998 Aug 1;255(3):703–9.

76. Sanders BM, Hope C, Pascoe VM, Martin LS. Characterization of the Stress Protein Response
in Two Species of Collisella Limpets with Different Temperature Tolerances. Physiol Zool.
1991;64(6):1471–89.

77. Clark MS, Fraser KPP, Peck LS. Lack of an HSP70 heat shock response in two Antarctic
marine invertebrates. Polar Biol. 2008 Aug 1;31(9):1059–65.

78. Rinehart JP, Hayward SAL, Elnitsky MA, Sandro LH, Lee RE, Denlinger DL. Continuous up-
regulation of heat shock proteins in larvae, but not adults, of a polar insect. Proc Natl Acad Sci
U S A. 2006 Sep 19;103(38):14223–7.

79. Hidalgo-Galiana A, Monge M, Biron DG, Canals F, Ribera I, Cieslak A. Protein expression
parallels thermal tolerance and ecologic changes in the diversification of a diving beetle species
complex. Heredity. 2016 Jan;116(1):114–23.

80. Langille BL, Tierney SM, Austin AD, Humphreys WF, Cooper SJB. How blind are they?
Phototactic responses in stygobiont diving beetles (Coleoptera: Dytiscidae) from calcrete
aquifers of Western Australia. Austral Entomol. 2019;58(2):425–31.

81. Healy MA. It's hot and getting hotter. Australian rangelands and climate change – Reports of
the Rangelands Cluster Project. Ninti One Limited and CSIRO, Alice Springs; 2015.

82. Hemmerle H, Bayer P. Climate Change Yields Groundwater Warming in Bavaria, Germany.
Front Earth Sci [Internet]. 2020 [cited 2022 Aug 22];8. Available from:
https://www.frontiersin.org/articles/10.3389/feart.2020.575894

83. Taylor CA, Stefan HG. Shallow groundwater temperature response to climate change and
urbanization. J Hydrol. 2009 Sep 15;375(3):601–12.

84. Gunawardhana LN, Kazama S. Statistical and numerical analyses of the influence of climate
variability on aquifer water levels and groundwater temperatures: The impacts of climate
change on aquifer thermal regimes. Glob Planet Change. 2012 Apr 1;86–87:66–78.

85. Cooper SJB, Saint KM, Taiti S, Austin AD, Humphreys WF. Subterranean archipelago:
mitochondrial DNA phylogeography of stygobitic isopods (Oniscidea : Haloniscus) from the
Yilgarn region of Western Australia. Invertebr Syst. 2008 May 12;22(2):195–203.

86. Bradford T, Adams M, Humphreys WF, Austin AD, Cooper SJB. DNA barcoding of
stygofauna uncovers cryptic amphipod diversity in a calcrete aquifer in Western Australia's arid
zone. Mol Ecol Resour. 2010;10(1):41–50.

87. Hose GC, Sreekanth J, Barron O, Pollino C. Stygofauna in Australian Groundwater Systems:
Extent of knowledge. 2015;CSIRO, Australia.

ACKNOWLEDGEMENTS

We thank Flora, Peter, and Paul Axford for providing access to the Sturt Meadows calcrete and

accommodation at the Sturt Meadows pastoral property. We thank Chris Watts (South Australian

Museum) for helping with identification and collecting of surface dytiscids, providing beetle images,

and for laying the foundation for the subterranean beetle research with WFH and Remko Leijs. We

737 thank Rae Humphreys for assistance with field collections of *P. macrosturtensis* and for being a

738 wonderful host to stygofauna catchers over many years. Finally, we would like to thank two

739 anonymous reviewers for helping to improve the quality of the manuscript.

AUTHOR CONTRIBUTIONS AND COMPETING INTERESTS STATEMENT

PGBH, KJ, and SJBC conceived the study. KJ performed experiments exposing beetles to heat

stress. PGBH analysed the data and wrote the manuscript. TB and CSPF provided guidance

regarding bioinformatic analyses. ST performed molecular laboratory work. TMB, KJ, WFH, and

ADA helped to draft the manuscript. All authors edited the manuscript and approved the final draft.

Funding for this project was provided by an Australian Research Council Discovery grant

(DP180103851) to SJBC, WFH, ADA and TB. The authors declare that they have no competing

interests.

DATA AVAILABILITY STATEMENT

All raw RNA-seq data for *Paroster macrosturtensis* and *P. nigroadumbratus* used in differential

expression analyses in this study are available via NCBI under BioProject PRJNA783065 (individual

Sequence Read Archive accessions for samples SRR17023302-SRR17023315). HSP transcripts

assembled from raw *P. macrosturtensis* RNA-seq data and used in phylogenetic analysis are

available via GenBank (accession numbers summarized in Table S1). The reference transcriptome of

P. nigroadumbratus (including all isoforms) and the annotated, subsetted dataset (only including

longest isoform per Trinity 'gene') are available via FigShare (DOI: 10.25909/17169191).