TITLE: High temporal resolution systems profiling reveals distinct patterns of interferon

response after Covid-19 mRNA vaccination and SARS-CoV2 infection

AUTHORS

Darawan Rinchai[1#*], Sara Deola[1#], Gabriele Zoppoli[2,3§], Basirudeen Syed Ahamed Kabeer[1§],

Sara Taleb[4§], Igor Pavlovski[1±], Selma Maacha[1±], Giusy Gentilcore[1], Mohammed Toufiq[1], Lisa

Mathew[1], Li Liu[1], Fazulur Rehaman Vempalli[1], Ghada Mubarak[1], Stephan Lorenz[1], Irene

Sivieri[3,5,6], Gabriella Cirmena[2], Chiara Dentone[2], Paola Cuccarolo[3], Daniele Roberto

Giacobbe[2,5], Federico Baldi[5], Alberto Garbarino[3], Benedetta Cigolini[3], Paolo Cremonesi[7],

Michele Bedognetti[8], Alberto Ballestrero[2,5], Matteo Bassetti[2,5], Boris P. Hejblum[9], Tracy

Augustine[1], Nicholas Van Panhuys[1], Rodolphe Thiebaut[9], Ricardo Branco[1], Tracey Chew[10],

Maryam Shojaei[11,12], Kirsty Short[13,14], Carl Feng[15,16], PREDICT-19 consortium, Susu M.

Zughaier[17], Andrea De Maria[2,5], Benjamin Tang[12,18], Ali Ait Hssain[19], Davide Bedognetti[1,3&],

Jean-Charles Grivel[1&] and Damien Chaussabel[1&*]

1. Research Branch, Sidra Medicine, PO Box 26999, Doha, Qatar

2. IRCCS Ospedale Policlinico San Martino, Genoa, Italy

3. Department of Internal Medicine and Medical Specialties, University of Genoa, Genoa,

Italy

4. Division of Genomics and Translational Biomedicine, College of Health and Life Sciences,

Hamad Bin Khalifa University, Doha, Qatar

5. Division of Infectious Diseases, Department of Health Sciences, University of Genoa,

Genoa, Italy

6. Department of Experimental and Clinical Medicine, School of Internal Medicine,

University of Florence, Florence, Italy

7. Emergency Department, E.O. Ospedali Galliera Genova, Italy

8. Azienda Sanitaria Locale 3 Genovese, Genova, Liguria, Italy

9. Univ. Bordeaux, Department of Public Health, Inserm U1219 Bordeaux Population Health

Research Centre, Inria SISTM, F-33000 Bordeaux, France

10. Sydney Informatic Hub, The University of Sydney, Sydney, Australia

11. Nepean Clinical School, University of Sydney, Sydney, New South Wales, Australia;

12. Westmead Institute for Medical Research, Westmead, New South Wales, Australia.

13. The University of Queensland, School of Chemistry and Molecular Biosciences, St Lucia,

Brisbane, Queensland, Australia;

14. Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane,

Queensland, Australia.

15. School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney,

Sydney, NSW, Australia;

16. Tuberculosis Research Program, Centenary Institute, The University of Sydney, Sydney,

NSW, Australia.

17. College of Medicine, QU Health, Qatar University, PO Box 2713, Doha, Qatar

18. Westmead Institute for Medical Research, Westmead, New South Wales, Australia.

19. Medical Intensive Care Unit, Hamad General Hospital, PO BOX 3050, Doha, Qatar

#, §, ±, & These authors contributed equally to the work

* To whom correspondence may be addressed:

drinchai@sidra.org / drinchai@gmail.com

dchaussabel@sidra.org / chaussabel@gmail.com

ABSTRACT:

Knowledge of the mechanisms underpinning the development of protective immunity

conferred by mRNA vaccines is fragmentary. Here we investigated responses to COVID-19

mRNA vaccination via ultra-low-volume sampling and high-temporal-resolution

transcriptome profiling (23 subjects across 22 timepoints, and with 117 COVID-19 patients

used as comparators). There were marked differences in the timing and amplitude of the

responses to the priming and booster doses. Notably, we identified two distinct interferon

signatures. The first signature (A28/S1) was robustly induced both post-prime and post-boost

and in both cases correlated with the subsequent development of antibody responses. In

contrast, the second interferon signature (A28/S2) was robustly induced only post-boost,

where it coincided with a transient inflammation peak. In COVID19 patients, a distinct

phenotype dominated by A28/S2 was associated with longer duration of intensive care. In

summary, high-temporal-resolution transcriptomic permitted the identification of post-

vaccination phenotypes that are determinants of the course of COVID-19 disease.

INTRODUCTION

COVID-19 vaccines are critical to the ongoing efforts to control the SARS-CoV-2 coronavirus pandemic. To date, nine vaccines have received some form of approval for use in humans, and phase III trials are ongoing for an additional 11 vaccines (1). Notable differences exist among the vaccine products in terms of their design and the levels of protection they confer, as well as the type, incidence, and severity of adverse events they may elicit. Gaining a comprehensive understanding of the immunological factors underpinning the different responses to various vaccines is a major endeavor. Yet, this knowledge is necessary for guiding timely decisions to modulate vaccination protocols (e.g., the use of different types of vaccines for the priming and booster doses). This information may also assist in matching of individuals with the growing number of available vaccines based on their demographics, health status, or any other relevant clinical/molecular phenotypes.

Blood transcriptome profiling measures the abundance of transcripts in whole blood and on a system-wide scale. It was previously employed to comprehensively profile the immune responses elicited by vaccines (2,3). Notably, this approach identified innate immune signatures arising within hours after administering vaccines (4). In a recently published report, Arunachalam et al. described the blood transcriptome profiles measured following the administration of the BNT162b2 mRNA COVID-19 vaccine (5). They reported the presence of an interferon (IFN) signature one day after the priming vaccination that was no longer detectable on day 7. They further found a more comprehensive IFN/inflammatory signature to be present 1 day after administering the booster dose. However, the sampling schedule employed in this study was relatively sparse. And the sample collection time points commonly selected in systems vaccinology studies are based on kinetics established for more conventional vaccines – with sampling at days 1 and 7 often selected since they correspond

to the peaks of the innate and adaptive immune responses elicited for instance by the

influenza or pneumococcal vaccines (6). However, the precise kinetics of the immune

response elicited by mRNA vaccines remains to be established. In the present study we

endeavored to profile the blood transcriptome of individuals prior to the administration of

the first dose of COVID-19 mRNA vaccine and for the following 9 consecutive days. Subjects

also collecting samples for deep serological profiling at three time points. The same sampling

and profiling schedule was repeated to assess the response to the second dose of the vaccine.

To achieve this, we have adopted a ultra-low volume sampling procedure consisting in the

self-collection of few drops of blood (50 ul) by fingerstick (7).

Together, this work permitted the precise delineation of a well-orchestrated immune

response to COVID-19 mRNA vaccines and identified marked differences in the magnitude,

nature, and timing of the transcriptional signatures elicited by prime and boost vaccination.

Most notably, differences in temporal patterns of responsiveness revealed distinct

components of the interferon response, which is known to play a key role in controlling SARS-

CoV-2 infection (8) and was also found here to associate with the subsequent development

of the antibody response post-vaccination.

RESULTS

Study design, implementation, and serological profiling

We successfully recruited a cohort of volunteers and implemented a high-frequency sampling

protocol. This permitted to ascertain the response to the first and second dose of COVID-19

vaccines at 10 consecutive daily timepoints: immediately before vaccination and for 9 days

after. We collected samples for serological profiling at three time points: before vaccination

and on days 7 and 14 post-vaccination (**Figure 1A**). We implemented a self-sampling blood

collection protocol so that subjects could extract small volumes (50 µl) of RNA-stabilized

blood at the required frequency (the approach is described in the Methods section and an

earlier publication (7)). RNA sequencing profiles were generated using a cost-effective 3'-

biased library preparation protocol (Lexogen QuantSeq), which is optimized for low amount

of RNA input. We generated COVID-19-specific antibody profiles from capillary blood samples

collected by Volumetric Absorptive Micro Sampling analyzed using a multiplexed Bead array

established by our team (see Methods for details). Overall, 23 subjects were enrolled in the

study, and the characteristics of this cohort are reported in **Table 1**. They received either two

doses of the Pfizer/BioNTech mRNA vaccine (BNT162b2, N = 19) or two doses of the Moderna

mRNA vaccine (N = 4). Among those 23 subjects, six had recovered from COVID-19 in the

months preceding the administration of the first vaccine dose. In total 440 RNA sequencing

profiles were generated, and this extensive dataset was shared publicly in GEO with the

accession number GSE190001. The serological profiles included reactivity to a stabilized

trimer of Spike protein, the spike protein, its receptor-binding domain, the Nucleo and

Envelope proteins, of SARS-CoV2, and the subunit S1 of SARS spike protein. The data are

provided in **Supplementary File 1**. The seroreactivity to each of these antigens was dissected

by measuring the total IgG, total IgA, and IgM, as well as the finer-scale IgG and IgA subtypes.

Serological profiling data showed a rise in the levels of antibodies in the plasma of the subjects

post-vaccination (**Figure 1B**), and this included antibodies specific for the SARS-CoV-2 Spike

protein, which is targeted by COVID-19 vaccines. No responses to the Envelope protein were

detected. Some cross-reactivity was observed with the SARS Spike protein. Notably, higher

antibody levels were induced after the first dose in individuals who had been previously

infected with the virus (**Figure 1B-C**).

Altogether, the implementation of this protocol established the feasibility of obtaining

stabilized-RNA blood samples from study-subjects post-vaccination at high-temporal

frequencies. We generated a large dataset using a cost-effective RNA-sequencing protocol

that served as the basis for subsequent analyses presented in this paper and was deposited

in a public repository. A detailed map of the serological profiles of the subjects enrolled in the

study was obtained that permitted us to explore the possible associations between blood

transcriptional responses and vaccine immunogenicity.

The post-prime interferon response peaks at day-2 and correlates with the antibody

response

Vaccines can elicit innate immune responses that are detectable systemically via blood

transcriptome profiling. But not all of them do, which is for instance the case of the aluminum-

adjuvanted Hepatitis B vaccine (9). Therefore our first question was whether transcriptional

changes could be observed during the first few days following the administration of COVID-

19 mRNA vaccines.

Analyses were carried out employing a fixed repertoire of 382 transcriptional modules

(BloodGen3) that we had recently established and characterized functionally (10)(see

methods section for details). Module responses were determined across all time points. The

differential gene-set enrichment functions of the dearseq R package were run to assess

whether changes observed throughout the nine days post-prime were statistically significant

(11). This analysis identified significant temporal changes for 22 of the 382 modules

constituting the BlooGen3 repertoire (**Supplementary File 2**).

Only seven modules were found to be changed at any given time point during the first

three days following the administration of the priming dose of the vaccine (**Figure 2A**). The

abundance of four modules was consistently increased across these time points, and all four

belonged to the module aggregate A28. Each "module aggregate" regroups sets of modules

that showed consistent abundance profiles across a reference set of 16 disease cohorts that

were employed for the construction of the BloodGen3 repertoire (see methods and (10) for

details). The module aggregate in question ("Aggregate A28") comprises of six modules. As

described in detail in one of our recent publications, all six are associated with interferon

responses (10). The gene composition of the modules and the functional annotations are

provided herein (relevant information is provided in **Supplementary File 3** and can be

accessed interactively via: https://prezi.com/view/E34MhxE5uKoZLWZ3KXjG/). The

responses observed on days 1 and 2 post-prime were mapped onto fingerprint grid plots,

where modules occupy a fixed position and are arranged by aggregate. Each aggregate

occupyies a given row (**Figure 2A**). Time-course gene-set enrichment analysis confirmed that

changes observed over time in four out of six A28 modules were significant. The response

profiles of the A28 modules showed a peak on day 2 post-vaccination. This was also visible on

a heatmap showing responses at each timepoint across individual subjects (**Figure 2B**). We

next examined whether this signature correlated with antibody responses measured 14-days

post-prime as well as at 14-days post-boost (**Figure 2C**). For this, correlation analyses were

run at the module level within Aggregate A28 using, as the endpoint, fold-changes in antibody

levels on days 7 and 14 post-prime and days 7 and 14 post-boost relative to the pre-

vaccination baseline (immediately prior to the administration of the first dose of COVID-19

mRNA vaccines). "Significance hotspots" were identified when most modules within a given

aggregate reached correlation significance thresholds. In the case of the post-prime

interferon signature, we identified such significance hotspots on days 2 and 3 post-prime for

a subset of three interferon modules, M10.1, M15.127, and M83, while a fourth module,

M15.86, also displayed significant correlations across all antibody types, but only on day 2.

Thus, we found that an interferon response is induced over the first three days following

the administration of the priming dose of mRNA vaccines. Remarkably this signature

correlated with the antibody response measured several weeks later, 14 days after the

administration of the second dose of vaccine.

A decrease in inflammation is accompanied by an increase in adaptive immune response

genes on day 5 post-prime

We were next interested in characterizing the changes occurring beyond the first three days

following administration of the priming dose. In total, 18 modules displayed changes on day

4 post-prime, of which 12 showed a decrease in abundance. These modules belonged to three

aggregates that have been associated with inflammation (A31, A33, A35). Most changes were

observed on day 4, but for some modules, changes were apparent starting on day 3 and

continued beyond days 4, day 5, or even 6 (**Supplementary Figure 1**). In our earlier work,

modules within the BloodGen3 Aggregate A35 were associated with systemic inflammation

mediated by neutrophils and were found to constitute a common denominator across a wide

range of pathologies in which systemic inflammation is present (12). The association of A35

with inflammatory processes was also ascertained based on the results of the functional

profiling analyses and the restriction in transcript expression in the reference datasets (10).

Detailed functional annotations can be accessed via interactive circle packing charts:

https://prezi.com/view/7Q20FyW6Hrs5NjMaTUyW/). Module Aggregate A33 has not been

investigated as extensively in any of our prior studies but was clearly associated with

inflammation via functional profiling (https://prezi.com/view/VBqKqHuLWCra3OJOIZRR/).

The peak response post prime was on day 5, with a total of 42 modules showing

differences in comparison to the pre-vaccination baseline (**Figure 3**). At this timepoint, most

modules showed an increase in abundance (29 were increased and 13 decreased). Some of

those modules belonged to aggregates that were associated with adaptive immunity, most

notably A27, which is associated with plasmablast responses (three out of five modules were

responsive at this timepoint). This association is based on the restriction of the expression of

the genes comprising A27 modules in plasma cells observed in a reference dataset including

a wide range of cell populations (contributed by Monaco et al. (14)) and by the presence of

the plasmablast marker CD38 and other associated genes (*IGJ*, *TNFRSF17*, *TXNDC5*) in one of

the A27 modules (M12.15). Detailed annotations and expression profiles of A27 transcripts in

the reference datasets can be accessed via https://prezi.com/view/GgIiA0K9kSFHbpVj2I85/.

Other immune-relevant modules found to be increased at this timepoint are associated with

T-cells (M12.6 from aggregate A1). Detailed functional annotations for module aggregate A1

can be accessed via https://prezi.com/view/sxap39tKxkmCNTTNIlVO/). Others were mapped

to module aggregates A24 and were associated with oxidative phosphorylation which is

known to play a role for instance in T-cell activation (6 out of 11 modules were responsive)

(13). Other modules were not yet functionally annotated, including for instance the four

responsive modules, out of 15, belonging to aggregate A26. Notably, the signatures observed

on day 5 appeared to be transient, and no modules were increased on day 6 post-prime.

Taken together, we found the number of responsive modules to peak on day 5 post-

prime. A decrease in the abundance of transcripts associated with inflammation was

accompanied by an increase in the abundance of transcripts associated with adaptive

immune responses. Notably, the latter appeared earlier than seen in response to other

vaccines where plasmablast signatures are observed around day 7 post-vaccination (6,14,15).

A post-boost interferon signature peaks on day 1 and correlates with antibody responses

After delineating temporal responses post-prime, we examined changes after the second dose of COVID-19 mRNA vaccines. Time-course gene set enrichment analysis identified significant temporal changes for 311 of 382 modules comprising the BloodGen3 repertoire (**Supplementary File 4**). After the booster dose, the peak number of responsive modules occurred on day 1, with 261 responsive modules or about two-thirds of the 382 modules constituting the BloodGen3 repertoire (**Figure 4**). This number decreased sharply afterward, with 115 responsive modules on day 2 and only 9 responsive modules on day 3. The kinetic and amplitude of the post-boost response contrasted markedly with that observed post-prime, when, as described above, the number of responsive modules after the first dose instead peaked on day 5, with changes found in 42 modules at that timepoint.

As seen from the fingerprint grid plot, the day 1 post-boost response was extensive and polyfunctional (**Figure 4**). An overall decrease in abundance was observed for aggregates broadly associated with lymphocytic cells (Aggregates A1-A8) and increased for module aggregates associated with myeloid cells, inflammation, and circulating erythroid cells (Aggregates A33-A38). In addition, a marked increase in the abundance of modules associated with interferon responses was also observed (Aggregate A28). We compared the day 1 response fingerprint of the COVID-19 mRNA booster vaccine to fingerprints derived from patients with a wide range of pathologies. These included sixteen reference datasets encompassing infectious and autoimmune diseases, as well as cancer, solid organ transplant recipients, among others (these cohorts are described in our previously published work (10,16); the respective blood transcriptome fingerprint collections are accessible via a dedicated web application: https://drinchai.shinyapps.io/BloodGen3Module/). In addition, we analyzed two original COVID-19 blood transcriptome datasets: one cohort comprising 77

Covid-19 patients with disease severities ranging from mild and moderate to severe (the

"PREDICT-19 consortium Italian cohort dataset" – see methods and published study protocol

for details (17)), while the second cohort comprised 40 COVID-19 patients recruited at the

time of admission to the intensive care unit (ICU) ("IMPROVISE cohort whole blood dataset").

These high-level comparisons showed, firstly, that the extent of the changes associated with

the day 1 response to the second dose of the COVID-19 mRNA vaccine was consistent with

that observed in some patient cohorts with acute infections (**Figure 4**). More specifically, they

were found to most resemble the responses seen in a cohort of subjects with influenza

infection, with a marked interferon response (A28) and an inflammation signature (A33, A35).

At a higher level, these response patterns were also generally consistent with those observed

in patients with a COVID-19 infection. However, the changes that occurred in response to the

vaccinations were not as extreme as those found, for instance, in patients with sepsis or with

the most severe form of COVID-19 (i.e., the IMPROVISE dataset) (most notably for

inflammation [A33, A35] and erythroid cell responses [A36-A38]).

Overall, the BloodGen3 transcriptome fingerprint observed on day 1 after the second

vaccine dose contrasted markedly with the fingerprint observed on day 1 post-prime. Yet, the

interferon response signature was found to be a common denominator between the

responses to the first and second doses, as it was observed in both cases in the first few days

following administration of the vaccine. We therefore began to dissect the post-boost

response by examining this interferon response signature in more detail.

Following the administration of the booster dose, the interferon response was

noticeably sharper in comparison to the post-prime response and peaked on day 1 instead of

day 2 (**Figure 5A**). This was illustrated by the difference in the maximum average module

response, which was close to 50% of the constitutive transcripts on day 2 post-prime and

greater than 80% on day 1 post-boost.

We decided to then perform hierarchical clustering to identify subsets of modules within

the A28 aggregates that might group together based on patterns of transcript abundance

across all subjects and timepoints. Two sets of three modules each were, thus, identified

within the A28 aggregate. The first set comprised modules M8.3, M10.1, and M15.127

(referred to as A28/S1), and the second set comprised modules M16.64, M13.17, and M15.86

(referred to as A28/S2). Interestingly, we observed post-prime that, while modules in A28/S1

peaked on day 2, those belonging to A28/S2 peaked on day 1 (**Figure 5B**). Furthermore,

A28/S1 modules showed an extended peak post-boost, with day 2 levels being almost

identical to those of the day 1 peak, while A28/S1 modules peaked sharply on day 1, with

levels decreasing rapidly thereafter. These findings suggest that both sets of modules

measured distinct types of interferon response. Indeed, public datasets in which responses

to type 1 interferon were measured *in-vivo* indicated that A28/S1 modules are likely to

represent type 1 interferon responses (**Figure 5B**), while we postulated that A28/S2 modules

might represent a type 2 interferon response. Modules forming the A28/S1 set comprise

some of the better recognized "canonical" interferon response genes, such as Oligoadenylate

Synthetase family members (OAS1, OAS2, OAS3, OASL), Interferon Induced Protein family

members (IFI6, IFI27, IFI35, IFI44, IFI44L), as well as Interferon Induced Protein With

Tetratricopeptide Repeats family members (IFIT1, IFIT3, IFIT5) (10). Modules forming the

A28/S2 set comprise instead most notably members of the Nuclear Antigen family members

SP100, SP110 and SP140, which are associated with interferon gamma signaling, as well as

transcription factors IRF9 and STAT2. Composition and functional annotations for A28

modules can be explored further at: https://prezi.com/view/E34MhxE5uKoZLWZ3KXjG/.

Finally, a strong association was found between the post-boost interferon signature and

the subsequent development of an antibody response. Indeed, positive correlations were

observed for all six A28 modules that reached significance on days 1, 2, and 3 post-boost.

Notably, this differed from the post-prime interferon response, for which significance was

reached only for four of the six modules and only on days 2 and 3.

Taken together, the high temporal resolution profiling results permitted the delineation

of distinct patterns of post-prime and post-boost interferon responses. The timing of the

responses observed at the individual module level contributed to the definition of the two

distinct sets of interferon modules. One set was associated with responses to type I interferon

in-vivo and dominated the post-prime response, with a peak on day 2. The post-boost

response showed a strong induction of both sets and also peaked on day 1.

Inflammation and erythroid cell signatures peak sharply on day 1 post-boost

We continued the dissection of the day 1 post-boost signature, focusing this time on

responses associated with inflammation and circulating erythroid cell precursors.

Aggregates A33 and A35, which are associated with inflammation, tended to decrease

from day 4 through day 6 post-prime but displayed instead a sharp and transient increase in

abundance post-boost. Indeed, a well-delineated response peak was observed on day 1 post-

boost for both the A33 and A35 modules (**Figure 6**), but in contrast to the interferon response

(A28/S1), it did not extend beyond the first day. Three distinct response patterns were

identified via hierarchical clustering among the 21 modules that formed aggregate A35. The

"A35/S1" set comprised five modules, while "A35/S2" and "A35/S3" comprised ten and six

modules, respectively. The distinction between those three A35/inflammation module sets

was rather more subtle than was the case for the A28/interferon sets. Indeed, all three

module sets peaked on day 1 post-boost. Differences were rather apparent in the inflection

of changes measured on days 2 and 3 post-boost and in the "recovery phase", as abundances

appeared to dip below the baseline and progressively rise to reach pre-vaccination levels. The

underlying biological factors driving the grouping of the modules to those three distinct sets

could not be identified at this time.

Modules for three aggregates broadly associated with erythroid cell signatures also

displayed a sharp but transient increase in transcript abundance on day 1 post-boost.

However, the abundance tended to dip afterward, with a low peak on day 4 post-boost,

before recovering by day 7. Functionally, this signature was found to be most prominently

associated with immunosuppressive states, such as late-stage cancer or pharmacological

immunosuppression (16), which is consistent with published functional studies (18,19). We

also found such signatures were associated with more severe manifestations in babies

infected with Respiratory Syncytial Virus (RSV) (16). Moreover, erythroid precursors have

been recently associated with COVID most severe clinical outcomes (20). Finally, we did not

find evidence of an association between the day 1 post-boost inflammation or erythroid cell

signatures and the antibody responses.

A plasmablast signature peaks on day 4 post-administration of the booster dose and correlates with antibody responses

After the booster dose, the number of responsive modules peaked sharply on day 1,

then rapidly subsided beyond day 2, with the number of responsive modules on days 3, 4, 5,

and 6 being reduced to 8, 11, 3, and 2, respectively. Yet, changes within this later timeframe

are meaningful, as they specifically concern the set of five modules comprising aggregate A27,

which is associated with the presence of antibody-producing cells in the peripheral blood.

Three of the five A27 modules showed significant alterations after the booster dose (M16.60,

M13.32, M12.15) (**Figure 6**). The proportion of differentially expressed transcripts in each

module was relatively modest (with an average of 15% at the peak of response), especially in

comparison with the interferon signatures described above (with an average of >80% for

some modules at the response peak). Yet, the trajectories of the five A27 modules were

relatively consistent, with only one of the modules (M15.110) showing a different pattern,

i.e., a peak on day 6, slightly above the levels observed on day 4. We also examined the

association of this post-boost plasmablast signature with the antibody response and found a

significant association starting from about day 3 and lasting until day 7 post-boost (**Figure 6**).

In summary, COVID-19 mRNA vaccination induced a plasmablast response that peaked on

day 4 post-vaccination. This was unexpected since such signatures typically are measured

around day 7 post-vaccine administration (e.g., in the case of influenza or pneumococcal

vaccines (6)). We were also able to demonstrate a logical association between this post-boost

plasmablast signature and the subsequent development of humoral immunity.

Patterns of interferon induction elicited by COVID-19 mRNA vaccines are also observed

among COVID-19 patients

Our work identified the interferon response as the most upstream factor associated with the

development of humoral immunity following COVID-19 mRNA vaccination. High-temporal

resolution profiling identified distinct patterns of interferon induction post-prime and post-

boost and we next decided to determine whether similar response patterns could be

identified among patients with COVID-19 disease.

We relied for this on the original blood transcriptome data from the PREDICT-19

consortium Italian COVID-19 cohort comprising 77 patients with a wide spectrum of disease

severity. We used the response values for the six interferon modules from Aggregate A28 to

map individual COVID-19 patient samples along with post-vaccine samples on the same t-SNE

plot (**Figure 7A**). First, we confirmed that there was no apparent separation of the vaccination

and COVID-19 patient cohorts, and that batch correction was therefore not warranted before

proceeding with comparative analyses (**Supplementary Figure 2**). This is consistent with the

results of meta-analyses we have previously conducted at the module level (16). To help with

the interpretation, k-means clustering was performed using the consolidated set of samples,

resulting in the formation of eight distinct clusters. Next, we examined the distribution of

samples from the vaccine and COVID-19 cohorts across the tSNE plot and among the eight

clusters. Timepoints at which an interferon response was detectable in vaccinated subjects

were of particular interest. Indeed, day 1 and day 2 post-prime samples (P1, P2), while

preferentially found in Clusters 1 and 5, appeared to be distributed across the entire t-SNE

plot. This is in contrast with day 1 and day 2 post-boost vaccination samples (B1, B2), which

were almost exclusively found in Cluster 5. A set of COVID-19 patients also co-localized in

Cluster 5, while others were found scattered across clusters, especially Clusters 1, 2, 6, and 3.

Interferon responses were detectable in all these clusters, but with important nuances. For

one, samples from Cluster 5 showed by far the most potent responses, with responses seen

in most cases across all six interferon modules, which was consistent with the post-boost

vaccine response (**Figure 7B**). In comparison, the response was less pronounced in samples

from Cluster 1, which was dominated by modules associated with type I interferon responses

(the A28/S1 set comprising M10.1, M8.3 and M15.127 described above). This pattern of

response was more consistent with the post-prime vaccine response. Signatures for samples

forming Clusters 2 and 6 were not well-defined and were in some cases absent, yet these

clusters also included COVID-19 patients. Samples forming Cluster 3 displayed a peculiar

signature, with an increase in the abundance of modules belonging to the A28/S2 set

(M15.64, M13.17, and M15.86) concomitantly with a decrease in modules forming A28/S1.

Among the samples forming this cluster, this pattern was most apparent for the COVID-19

patients.

Thus, we employed here the interferon responses observed post COVID-19

vaccination as a benchmark for the interpretation of COVID-19 patient signature. We were

able to establish that most COVID-19 patients display responses consistent with those found

post-vaccination, which, as established in this study, were associated with the development

of potent humoral responses. However, a subset of patients displayed patterns of interferon

response that are not typically seen in vaccinated individuals. It can thus be surmised that the

later patterns of interferon response might either be suboptimal or possibly even pathogenic.

The atypical interferon response signature observed in COVID-19 patients is associated

with a worse course of disease

The fact that some COVID-19 patients failed to display robust "post-vaccine-like" interferon

responses may be due to either a defective innate immune response, which may lead to

more severe disease course, or conversely to activation thresholds not being reached in

patients presented with milder disease.

Thus, we next examined patterns of interferon response in another original COVID-19

disease cohort, comprised exclusively of patients enrolled at the time of admission in the ICU

(the IMPROVISE cohort, which was also described above). As described above, we again

mapped individual COVID-19 patient samples along with post-vaccine samples on a t-SNE plot

based on similarities in the patterns of interferon responsiveness across the six A28 interferon

modules (**Figure 8A**). COVID-19 subjects were found to again be distributed throughout

multiple clusters. Patients who co-localized with day 1 post-boost vaccine samples tended to

have relatively short ICU stays (in Cluster 5 with potent A28/S1 and A28/S2 responses), and

only a few patients co-localized with day 2 post-prime samples in Cluster 3, which was

characterized by a more prominent A28/S1 signature compared with A28/S2. Furthermore,

distinct groups of patients in Clusters 1 and 6 displayed the peculiar pattern of interferon

response dominated by A28/S2 that was identified earlier among patients enrolled in the

PREDICT-19 cohort. Notably, patients from the IMPROVISE cohort displaying this pattern of

interferon response showed significantly lengthier stays in the ICU compared to patients

displaying patterns of interferon response that are consistent with those observed post-

vaccination (**Figure 8B** comparing left and right cluster: for length of hospital stay, t-test, p =

0.006 (**), mechanical ventilation days p =0.016 (*) and ICU stay p = 0.012(*)).

Thus, in a cohort of subjects uniformly presenting with severe disease, post-prime-like

patterns of interferon response dominated by A28/S1 were less prevalent. Post-boost-like

pattern of interferon response characterized by robust A28/S1 and A28/S2 signatures were

observed instead in most patients. A notable exception were patients presenting with

patterns of response dominated by A28/S2, not observed previously following vaccination

but which were found again in this second independent COVID-19 dataset. In this context we

could also establish that such response is associated with a worse disease course. This overall

supports the notion that patients harboring this signature may fail to mount an effective

immune response against SARS-CoV-2.

The peculiar interferon response phenotype observed in COVID-19 patients is not typically found in the context of other infections

Finally, we asked whether the A28/S2-dominated interferon response pattern associated with worse disease outcomes in COVID-19 patients was also commonly found in other infectious disease.

For this we first developed a standard definition of "Interferon Response Transcriptional Phenotypes" (IRTPs): the two distinct signatures described above, A28/S1 and A28/S2, were employed as "traits" for the definition of three main phenotypes observed following vaccination and in response to SARS-CoV2 infection. 1) IRTP I encompassed A28/S1-dominated patterns of response: "A28/S1^{++}A28/S2$^+$","A28/S1^{++}A28/S2^0" and "A28/S1$^+$A28S2$^+$" (see the method section for details). 2) IRTP II corresponded to a pattern of interferon response characterized by the strong induction of both components: A28/S1^{++}A28/S2^{++}. 3) IRTP III encompassed the A28/S2-dominated patterns of interferon response: "A28/S1$^-$A28/S2^{++}", "A28/S1$^-$A28/S2$^+$", "A28/S1^{0}A28/S2^{++}", "A28/S1^{0}A28/S2$^+$" and "A28/S1$^-$A28/S2^0". These three IRTPs were in turn employed for the stratification of our vaccination cohort at early time points following administration of the priming and booster doses, as well as both of our COVID-19 cohorts and of several reference cohorts of patients which we had generated as part of one of our earlier studies (10), focusing more particularly on pathologies known to elicit robust interferon responses, including viral infections (influenza, RSV, Human Immunodeficiency Virus [HIV]), tuberculosis or systemic lupus erythematosus (SLE) (**Figure 8C**).

Interferon Response Transcriptional Phenotype I (IRTP I), that we posit corresponds to a response dominated by type 1 interferon (IFNa, IFNb), in absence of a substantial type 2 interferon (IFNg), was found in ±1/3 of the vaccinated subjects at peak response on day 2

post-prime (**Figure 8C**: P2). It was however absent at peak response post-boost (B1). Similarly,

IRTP I was found among COVID-19 patients belonging to the PREDICT-19 cohort (although in

only about 10% of patients), but not among those belonging to the IMPROVISE cohort, who

presented with more severe disease. IRTP I was otherwise also found in ±10% of subjects

across most of our reference cohorts. However, as was the case of our severe COVID-19

cohort, it was absent in the comparator cohort comprised of patients with bacterial sepsis. In

the context of mRNA vaccination, IRTP II, which is characterized by the robust induction of

both A28/S1 and A28/S2 components, was observed following the booster dose in 95% of

samples profiled on day 1, which corresponds to the peak response. The priming dose of

Covid-19 mRNA vaccines was able to induce both components robustly but in only 48% of

samples at peak (day 2 post-prime). IRTP II was otherwise also prevalent in COVID-19 patients,

which is consistent with our earlier observation. It was also found in most samples in the other

pathologies employed as comparators – except for RSV and bacterial sepsis (40% and 48%,

respectively). Interferon response transcriptional phenotype III (IRTP III), which is

characterized by an A28/S2-dominated response was observed only rarely post-COVID-19

mRNA vaccination. It was however prevalent among COVID-19 patients, with 25% and 22% of

subjects with this phenotype in the PREDICT-19 and IMPROVISE cohorts, respectively.

However, it was not observed in patient with tuberculosis, influenza virus or HIV infection.

IRTP III is on the other hand found in 13% of patients with RSV infection and reached its peak

prevalence in patients with bacterial sepsis (36%).

In summary, those results show that in most instances both components of the

transcriptional interferon response can be robustly induced following COVID-19 vaccination

or viral infection (i.e. corresponding to IRTP II). However, incomplete patterns of induction

can also be observed in some circumstances. We hypothesize that this may be due: 1) to

activation thresholds not being reached, in the case of IRTP I or 2) to subjects failing to mount

an effective interferon response, in the case of IRTP III, which in the context of SARS-CoV-2

infection appears to impact their ability to control the infection. Notably, besides COVID-19,

IRTP III phenotypes were only observed in a limited set of pathologies, including infection

caused by RSV, a virus that is known to interfere with the interferon response (21,22), and

bacterial sepsis that is characterized by a dysregulated host response to infection (23).

DISCUSSION

Relatively little is known about the types of *in-vivo* immune responses elicited by mRNA

vaccines in humans. To address this, we employed bulk blood transcriptomics to map the

immune changes taking place *in-vivo* after the administration of priming and booster doses

of COVID-19 vaccines in adult volunteers. We did so at a high-temporal resolution, collecting

small amounts of blood before and for nine consecutive days after the administration of the

priming and booster doses of COVID-19 mRNA vaccines. The use of blood transcriptomics

eliminated the need to choose a panel of analytes to measure vaccine responses, which is

one source of bias. The daily collection and profiling schemes adopted eliminated the need

to choose specific timepoints for measuring the response, thus eliminating a second source

of bias.

Profiling blood transcript abundance post-prime and -booster doses of COVID-19

mRNA vaccines at a high-temporal resolution revealed a well-orchestrated sequence of

immune events (**Figure 9**). The immune signatures elicited following the administration of the

two doses of mRNA vaccines differed drastically. Relatively modest changes were observed

post-prime that manifested primarily as the induction of interferon-response signatures that

were detectable over the first three days following the injection of the first dose. This was

followed by a more subtle response that could be attributed to the priming of the adaptive

response between days 4 and 6. Indeed, a decrease in the abundance of transcripts for

modules associated with inflammation was observed over these three days, which included

an increase in transcripts associated with plasma cells and T-cells on day 5. No further changes

were detected beyond day 6. After the booster dose, the plasmablast response was more

robust and peaked on day 4, but was not accompanied by a T-cell response peak as was the

case post-prime. Notably, in studies assessing blood transcriptional responses to vaccines, the

peak plasmablast response is typically observed on day 7, as it is, for instance, with influenza

and pneumococcal vaccines (6,14,15). As a result, sampling schedules in common use are

designed to capture changes on days 1, 7, and sometimes day 3, but would miss the peak of

the adaptive response to COVID-19 mRNA vaccines observed in our study. In addition to

eliminating potential blind spots, high-frequency sampling and profiling also permit the

precise resolution of signatures that show the complex kinetics of a response; for instance,

the erythroid cell signature peaks sharply post-boost and recedes well below baseline over

several days before recovering. The trajectory of this signature may be of significance in the

context of vaccination, as we recently described its association with immunosuppressive

states, such as late-stage cancer and maintenance therapy in liver transplant recipients (16).

In the same work, we found this signature to be strongly associated with the development of

a more severe disease in subjects with acute respiratory syncytial virus infection; and we

furthermore putatively associated this signature with populations of circulating erythroid

cells found to possess immunosuppressive properties (18).

Arunachalam et al. previously described the elicitation of qualitatively distinct innate

signatures on day 1 following the administration of priming and booster doses of COVID-19

mRNA vaccines, with the former inducing an interferon response and the latter a mixed

response that also presented an inflammatory component (5). Our findings are consistent

with these earlier observations and, employing a high-frequency sampling and profiling

protocol, permitted to further dissect those responses. Most notably, while interferon

responses appear *a priori* as the common denominator between the post-prime and post-

boost responses, the temporal pattern of response that we observed indicates that these are,

in fact, qualitatively and quantitatively distinct. This was best evidenced by the differences in

the timing of the response peak, which corresponded to day 2 post-prime and day 1 post-

boost. The kinetics of the response post-boost is, therefore, most consistent with what is

observed following injection of a single dose of influenza vaccine (6). Interestingly, a further

investigation of the patterns of response among the six modular components of the

interferon responses (module Aggregate A28) identified two distinct sets of modules. These

two sets of three modules each, A28/S1, and A28/S2, displayed distinct kinetics and

amplitude of response post-prime and post-boost. We have described, in an earlier report,

that distinct interferon modules could be employed to stratify patients with systemic lupus

erythematosus (24). Here we sought to specifically determine whether "post-prime-like"

patterns (i.e., dominated by A28/S1 – IRTP I) or "post-boost-like" patterns (i.e., with potent

induction of both components: A28/S1++, A28/S2++ - IRTP II) could be identified among

COVID-19 patients. Indeed, since those were associated with the subsequent development of

humoral immunity in the context of vaccination it may be surmised that it would also be the

case during the course of SARS-CoV-2 infection. This question was made particularly relevant

in the context of COVID-19 disease, since it has been reported that failure to induce interferon

responses is associated with worse disease outcomes (8,25–27). In the PREDICT-19 cohort,

composed by patients with predominantly mild or moderate pathology, both phenotypes

were indeed observed, along with a third "atypical" phenotype that was not observed post-

vaccination. This latter phenotype is dominated instead by A28/S2, with A28/S1 abundance

low or even decreased (IRTP IIII). Notably, in a cohort of severe patients, both A28/S1++

A28/S2++ ("post-boost-like" / IRTP II) and A28/S2>S1 ("atypical" / IRTP III) phenotypes were

also observed, with the latter being associated with extended lengths of stay in the ICU.

However, IRTP III did not appear to be preferentially associated with death in this setting,

which may be due to the supportive care provided to the patients. While, overall, our

observations support the notion that failure to mount robust interferon responses is

associated with a less favorable course of the disease, they also show that the response

elicited in these patients may be of a peculiar type, but is altogether not entirely defective

(i.e., with only one component. A28/S1, being primarily affected). One possibility is that this

peculiar response pattern may be associated with the presence of endogenously produced

autoantibodies that neutralize interferon, as has been previously described (27,28). The high

incidence of the IRTP IIII phenotype observed in patients with bacterial sepsis (about 1 in 3),

however suggests that other mechanisms may be at play. Taken together, it is not possible

for us to be conclusive on this point at this time and further investigations are thus warranted.

Other points remain to be elucidated. This includes the timing of the adaptive response

to mRNA vaccines, which appears to rise and peak several days earlier than what is normally

observed in responses to other vaccines (± 7 day peak). The priming mechanism underpinning

the robust polyfunctional response observed on day 1 post-boost remains to be determined

as well. And in particular, whether or not such a response, which would typically be

considered to be innate, is in fact antigen-specific. Interestingly, in that respect, the subjects

who were previously infected but recovered from COVID-19 did not display a noticeable day

1 inflammatory response, and their immune systems behaved like those of naïve individuals.

However, the number of recovered subjects was small, and the study was not designed to

directly address this question. Hence, further investigations will also be necessary. Notably,

the greater amplitude of responses observed post-boost and the presence of an inflammatory

component is also consistent with previous reports of the increase in the incidence of side

effects/discomfort following COVID-19 mRNA vaccine booster doses (29,30).

Thus, while this study contributes to a better understanding of drivers of mRNA vaccines

immunogenicity it can also serve as a resource to help inform the design of studies

investigating vaccine responses. Indeed, a decrease in sequencing costs provides laboratories

an opportunity to employ transcriptome profiling approaches in novel ways. One of them

being the implementation of high-temporal resolution profiling protocols. An advantage of

the delineation of transcriptome responses at high-temporal resolution is that it is doubly

unbiased, i.e., there is no need to select transcripts for inclusion in a panel because RNA

sequencing measures all transcript species present in a sample. Similarly, there is no need to

select specific timepoints for assessing the vaccine response, as all timepoints were profiled

within a given time frame. An obvious advantage of the approach is that it permits the

removal of potential blind spots and the detection of changes that may otherwise be missed

by more sparse sampling protocols. In addition to eliminating potential blind spots high-

frequency profiling data helped resolve the vaccine response more precisely. This was the

case in our study of the interferon response, with the delineation of two distinct components

having been much more difficult if not for the resolution of peaks of response over the first

three days post first and second doses of vaccines. Some of the practical elements that may

contribute to making the routine implementation of the high-temporal resolution

transcriptomics approach viable include, as mentioned earlier, a substantial decrease in the

cost of RNA sequencing, especially 3'-biased methodologies. Along the same lines recent

publications showed, through down-sampling analysis, that much fewer deep reads than

usual are adequate for biomarker discovery projects, which could lead to further reductions

in the cost of RNA sequencing assays (31), with the lower costs permitting larger sample sizes

or, as in this case, a higher sampling frequency. Another consideration is the availability of

solutions for the in-home self-collection of samples. This is the case for the collection of RNA-

stabilized blood with our custom method, which could be further improved. Novel solutions

are also being put forward that could permit the implementation of these methods at scale

(32). Finally, as we have shown, it is possible to implement the self-collection of samples for

serology profiling within a vaccinology study.

There were several limitations to our study. While the sample size was adequate for an

initial discovery phase, a larger study cohort would help to better resolve inter-individual

variations. The dataset we generated, however, has been made available for reuse, and it

should be possible to integrate and consolidate this dataset with those generated in follow-

on studies by us and others (16). Follow-on studies would need to be purposedly designed to

formally address specific questions, for instance, comparing responses in individuals who had

previously been exposed to SARS-CoV-2 with those in naïve individuals. It would also be

interesting to compare responses elicited by the Pfizer/BioNTech and Moderna vaccines,

which was not possible in our study due to the small numbers of individuals that received the

Moderna vaccine. Indeed, although we hoped it would be possible to obtain more balanced

sample sizes for a more detailed comparison, the speed at which the vaccinations were rolled

out among our target population of healthcare workers meant we had very little control over

the number of volunteers that received the different types of vaccines or their status as naïve

or previously exposed individuals. It would also have been particularly interesting to enroll

patients from different age categories, especially the elderly population, but this again proved

impossible.

In conclusion, a several COVID-19 vaccines have already been approved for use in humans,

and an even greater number of them are currently in phase III trials (>20) (33). The data

presented herein suggest that high-temporal-resolution blood transcriptomics would provide

a valuable means to precisely map and compare the types of responses elicited by the

different types of COVID-19 vaccines. Similarly, this approach could potentially be

implemented to characterize and compare vaccine response profiles in populations that do

not respond optimally to vaccines (e.g., in the elderly, immunosuppressed, and during

pregnancy). This study also contributed to a better understanding of drivers of mRNA vaccines

immunogenicity and identified interferon signatures as early indicators of the potency of the

humoral immune response elicited in individual subjects. It also led to the definition of

functional interferon response phenotypes among COVID-19 patients which were associated

with different disease trajectories. In particular, mechanisms underlying the development of

dysfunctional interferon responses remain to be elucidated, which may yield important

insights into pathogenesis of severe COVID-19 disease.

METHODS

Subject recruitment:

<u>COVAX Cohort</u>: We enrolled adult subjects eligible to receive a COVID-19 vaccine who were

willing to adhere to the sampling schedule. The protocol was approved by Sidra Hospital IRB

(IRB number 1670047-6), and all participants gave written informed consent. Inclusion criteria

matched the clinical eligibility for receiving the vaccine, and the only exclusion criterium was

to have received a first dose of any COVID-19 vaccine. Twenty-three subjects were enrolled,

and the median age was 38 years (29-57); 20 of the subjects received the Pfizer vaccine and

three the Moderna vaccine. The demographics, health status at accrual, and vaccination side

effects are shown in Table 1. Vaccination and booster intervals were typically 21 days for

Pfizer and 29 days for Moderna.

<u>IMPROVISE cohort</u>: Adult subjects with severe COVID-19 were enrolled in this cohort under

the Hamad Medical Corporation IRB approval (MRC-05-007). Blood samples were collected

at multiple timepoints during patients' ICU stay (timepoint 1 was taken at ICU admission;

timepoints 1 to 4 were seven days apart). Subjects with burn and trauma, immunological

diseases, receiving immunosuppressive treatment, with other immune-related conditions, or

with a previous COVID-19 infection were excluded. For this analysis, 40 severe COVID-19

patients were included, with a median age of 52 (range = 30 to 92). The clinical parameters

of those patients included gender, ICU and hospital stay, mechanical ventilation duration,

ECMO initiation, comorbidities, outcomes (death/recovery), nosocomial infection onset, and

plasma therapy. Samples were also collected from control subjects who were adults and did

not: 1) present with an infectious syndrome during the last 90 days, 2) experience extreme

physical stress within the last week, 3) received during the last 90 days a treatment based on:

antivirals; antibiotics; antiparasitic; antifungals; 4) received within the last 15 days, a

treatment based on non-steroidal anti-inflammatory drugs; 5) received during the last 24

months a treatment based on: immunosuppressive therapy; corticosteroids; therapeutic

antibodies; chemotherapy and 6) a person with a history of: innate or acquired immune

deficiency; hematological disease; solid tumor; severe chronic disease; surgery or

hospitalization within the last 2 years; pregnancy within the last year; participation to a phase

I clinical assay during the last year; participation to a phase I clinical assay during the last year;

pregnant or breastfeeding women; a person with restricted liberty or under legal protection.

<u>PREDICT-19 Cohort:</u>

The "Predicting disease progression in severe viral respiratory infections and COVID-19"

(PREDICT-19) Consortium is an international consortium formed by a group of researchers

who share common interests in identifying, developing and validating clinical and/or

bioinformatics tools to improve patient triage in a pandemic such as COVID-19 (17). The

PREDICT-19 Italian cohort comprises adult subjects with mild, moderate, or severe COVID-19

diagnosed by real-time PCR on nasopharyngeal swab who were consented and enrolled at

E.O. Ospedali Galliera, and IRCCS Ospedale Policlinico San Martino, Genoa, Italy (Ethics

Committee of the Liguria Region (N.CER Liguria 163/2020- ID 10475). Blood samples were

collected during hospitalization. Subjects with burn and trauma, immunological diseases,

receiving immunosuppressive treatment for underlying disorders before COVID-19 diagnosis,

with other immune-related conditions, or with a previous COVID-19 infection were excluded.

For this analysis, ten healthy subjects and 103 COVID-19 patients were included, with a

median age of 61.76 (range = 26 to 86).

Sampling protocol:

<u>COVAX Cohort:</u> For transcriptomics applications for the COVAX study, after puncturing the

skin with a finger stick, 50 µl of blood was collected in a capillary/microfuge tube assembly

supplied by KABE Labortechnik (Numbrecht, Germany) containing 100 µl of tempus RNA-

stabilizing solution aliquoted from a regular-sized tempus tube (designed for the collection of

3 ml of blood and containing 6 ml of solution; ThermoFisher, Waltham, MA, USA). This

method is described in detail in an earlier report (7), and the collection procedure is illustrated

in an uploaded video: https://www.youtube.com/watch?v=xnrXidwg83I. Blood was collected

prior to the vaccine being administered (day 0), on the same day, and daily thereafter over

the next 10 days. This protocol was followed for both the priming and booster doses.

For serology applications, 20 µl of blood was collected using a Mitra blood collection device

(Neoteryx, Torrance, CA, USA) prior to the vaccine being administered and on days 7 and 14

after vaccination with the priming and booster doses.

<u>IMPROVISE Cohort</u>: For the IMPROVISE study, samples were collected using PaxGene Blood

RNA tubes (BD Biosciences, Franklin Lakes, NJ, USA) at all timepoints and were frozen at -20C

until further processing.

<u>PREDICT-19 Cohort</u>: For the Italian cohort of the PREDICT-19 study, blood samples were

collected during hospitalization by venipuncture in tubes containing an RNA stabilizing

solution (Tempus™ Blood RNA Tube, ThermoFisher, Waltham, MA, USA, Catalog number:

4342792) and frozen at -20C until further processing.

Multiplex serological assay

The presence of antibodies against selected Human Coronaviruses proteins in the serum was

measured with a home-built bead array based on carboxymethylated beads sets with six

distinct intensities of a UV-excitable dye. Each bead set was individually coupled to 3 SARS-

CoV-2 proteins, envelope, nucleoprotein, Spike protein in its trimeric form-or its fragments,

and the S1 fragment of SARS-CoV S protein. Therefore, the complete array consisted of 6

antigens, including five SARS-CoV-2 antigens (Full Spike Trimer, Receptor Binding Domain,

Spike S1, Nucleoprotein, and Envelope), as well as the closely related SARS-CoV-S1 protein.

The binding of human antibodies to each viral antigen (bead set) is revealed with fluorescently

labeled isotype-specific mouse monoclonal or polyclonal antibodies. We measured total IgM,

total IgG, total IgA, as well as their individual isotypes, IgG1, IgG2, IgG3, IgA1, and IgA2,

reporting a total of 48 parameters per sample. The assays were performed on filter plates and

acquired on a BD-Symphony A5 using a high-throughput-sampler. An average of 300 beads

per region was acquired, and the median fluorescence intensity (MFI) for each isotype binding

was used for characterizing the antibody response. An antibody response index was

calculated as the ratio of the MFI of pooled negative blood controls collected prior to June

2018 (Sidra IRB 1609004823) to the MFI obtained for vaccinated donor samples.

RNA extraction and QC

RNA was extracted using the Tempus Spin RNA Isolation Kit (ThermoFisher), which was

adapted for the handling of small blood volumes. The methodology has been described

previously in detail (34). Contaminating DNA was removed using the TurboDNAse kit

(ThermoFisher), and RNA was quantitated on a Qubit instrument (ThermoFisher) and QCed

using an Agilent 2100 Bioanalyzer (Agilent, Santa Clara, California, USA).

RNA sequencing

<u>COVAX & IMPROVISE Cohorts:</u> mRNA-sequencing was performed using QuantSeq 3' mRNA-

Seq Library Prep Kit FWD for Illumina (75 single-end) with a read depth of 8M and average

read alignment of 79.60%. Single samples were sequenced across four lanes, and the resulting

FASTQ files were merged by sample. Quality trimming is performed to remove adapter

sequences and polyA tails. Then trimmed reads were aligned to human genome

GRCh38/hg38 (Genome Reference Consortium Human Build 38), INSDC Assembly

GCA_000001405.28, Dec 2013) using STAR 2.6.1d and featureCounts v2.0.0 was used to

generate the raw counts. Raw expression data were normalized to size factor effects using R

package DESeq2. All downstream analyses were performed using R version 4.1 unless

otherwise specified. Global transcriptional differences between samples were assessed by

principal component analysis using the "prcomp" function. Transcriptome profiling data were

deposited, along with detailed sample information, into a public repository, the NCBI Gene

Expression Omnibus (GEO), with accession ID GSE190001 and BioProject ID: PRJNA785113

PREDICT-19 Cohort: Total RNA was isolated from whole blood lysate using the Tempus Spin

Isolation kit (Applied Biosystems) according to the manufacturer's instructions. Globin mRNA

was depleted from a portion of each total RNA sample using the GLOBINclear™-Human kit

(Thermo Fisher). Following the removal of globin transcripts transcriptome profiles were

generated via mRNA sequencing. Then mRNA-sequencing was performed using Illumina

HiSeq 4000 Technology (75 paired-end) with a read depth of 60M. Single samples were

sequenced across four lanes, resulting FASTQ files were merged by sample. All FASTQ passed

QC and were aligned to reference genome GRCh38 using STAR (2.6.1d). BAM files were

converted to a raw count's expression matrix using HTSeq (https://github.com/Sydney-

Informatics-Hub/RNASeq-DE). Raw count data was normalized using DEseq2. The ensemble

IDs targeting multiple genes were collapsed (average), and a final data matrix gene was

generated for modular repertoire analysis.

Statistical Analysis

Analyses were conducted using pre-defined gene sets. Specifically, we employed a fixed

repertoire of 382 transcriptional modules that were thoroughly functionally annotated, as

described in detail in a recent publication (10). Briefly, this repertoire of transcriptional

modules ("BloodGen3") was identified based on co-expression, as measured in a collection of

16 blood transcriptome datasets encompassing 985 individual transcriptome profiles. Sets of

co-expressed transcripts were derived from the analysis of a large weighted co-clustering

network. Downstream analysis results and visualizations were generated employing a custom

R package (35). "Module response" is defined as the percentage of constitutive transcripts

with a given abundance that was determined to be different between two study groups, or

for the same individual in comparison to a given baseline (in this study, pre-vaccination

abundance levels). The values, therefore, ranged from 100% (all constitutive transcripts

increased) to −100% (all constitutive transcripts decreased). Only the dominant trend (i.e.,

increase or decrease in abundance over control/baseline) was retained for visualization

purposes on fingerprint grids or fingerprint heatmaps, with red indicating an increase and

blue a decrease in abundance. When performing group comparisons (e.g., cases vs controls

for the disease datasets used as reference), the p-value and false discovery rate cutoffs were

applied, which are mentioned in the figure legend. When performing longitudinal analyses,

the module response is determined by employing fixed fold-change and expression difference

cutoffs. Module response values obtained were used for data visualization. Significance was

determined for each module using the differential gene set enrichment function of the

dearseq R package (11).

Definition of Interferon Response Transcriptional Phenotypes

Study cohorts were stratified based on patterns of interferon response for two distinct

interferon signatures, defined as A28/S1 (comprising modules M8.3, M10.1 and M15.127)

and A28/S2 (comprising modules M13.17, M15.64, M15.86). For this, phenotypes were

defined based on levels of response observed for these two "traits", as follows:

Percentage response of the six IFN modules were scored base on degree of response (%

response >= 50; score = 2, 0 < %response < 50; score =1 and (% response <= -50; score = -2, -

50 < %response < 0 ; score =-1). Then the average scores of S1("M8.3", "M10.1" and

"M15.127") and S2 ("M13.17" , "M15.64" , "M15.86") and phenotypes were classified using

cutoff at S1/S2++ (avg score >=1), S1/S2+(1< avg score < 0.33), S1/S20(0.33 < avg score <=

0), and S1/S2 – (avg score < 0). The phenotypes were grouped as:

-     "Interferon Response Transcriptional phenotypes I" = "IRTP I" =

"A28/S1++A28/S2+","A28/S1++A28/S20", "A28/S1+A28/S2+",

-     "IRTP II" = A28/S1++A28/S2++",

-     "IRTP III" = "A28/S1-A28/S2++","A28/S1-A28/S2+","A28/S10A28/S2++",

"A28/S10A28/S2+", "A28/S1-A28/S20"

-     The "other " category encompassed the less prevalent phenotypes remaining =

"A28/S1+A28/S20", "A28/S10A8/S2+", "A8/S1+A28/S2", "A28/S10A28/S20",

"A28/S10A28/S2-", "A28/S1-A28/S2-", "A28/S1+A28/S2++"

ACKNOWLEDGMENTS:

We wish to acknowledge Michelle Esblaca and Juvilyn Gusi for their assistance with the

training of subjects and collection of blood samples, as well as the research subjects for

volunteering and making the study possible.

AUTHOR CONTRIBUTION

Conceptualization, D.R., S.D., D.B., J-C.G., and D.C.; Methodology, D.R., S.D., B.K., I.P., S.M.,

G.G., D.B., J-C.G., and D.C.; Data Generation, D.R., S.D., G.Z., B.K., S.T., I.P., S.M., G.G., L.M.,

L.L., F-R.V., G.M., S.L., T.C., M.S., R.B., S.Z., A.DM., B.T., A. AH., D.B., J-C.G., and D.C.;

Investigation, D.R., S.D., G.Z., D.B., J-C.G., and D.C.; Resources, D.R., S.D., G.Z., I.S., G.C., C.D.,

P.Cu., D.R.G., F.B., A.G., B.C., P.Cr., T.A., N.V., M. Be., A.B., M.Ba., S.Z., A.DM., B.T., A. AH.,

D.B., J-C.G., and D.C.; Formal Analysis, D.R., S.D., B.K., I.P., S.M., G.G., M.T., L.M., L.L., F-R.V.,

G.M., S.L., B. H., R.T., T.C., M.S., D.B., J-C.G., and D.C.; Patient Enrollment and Follow up, G.Z.,

S.T., I.S., G.C., C.D., P.Cu., D.R.G., F.B., A.G., B.C., P.Cr., M. Be., A.B., M.Ba., T.A., R.B., S.Z.,

A.DM., B.T., A. AH., Writing - Original Draft, D.R., S.D., G.G., D.B., J-C.G., and D.C.; Writing -

Review & Editing, all authors; Visualization, D.R., S.D., B. H., R.T., and D.C.; Supervision, D.R.,

S.D., G.Z., S.L., K.S., S.Z., A.DM., B.T., A. AH., D.B., J-C.G., and D.C.;

COMPETING INTERESTS STATEMENT

D.R.G. outside of this work, reports investigator-initiated grants from Gilead Italia and Pfizer

Inc. M.Ba Outside of this work work, reports research grants and/or advisor/consultant

and/or Speaker/chairman Bayer, Biomerieux, Cidara, Cipla, Gilead, Menarini, MSD, Pfizer,

Shionogi". The other authors have no competing interests to report.

REFERENCES

1. Tregoning JS, Flight KE, Higham SL, Wang Z, Pierce BF. Progress of the COVID-19 vaccine
effort: viruses, vaccines and variants versus efficacy, effectiveness and escape. Nat Rev
Immunol. 2021 Aug 9;

2. Gaucher D, Therrien R, Kettaf N, Angermann BR, Boucher G, Filali-Mouhim A, et al.
Yellow fever vaccine induces integrated multilineage and polyfunctional immune
responses. J Exp Med. 2008 Dec 22;205(13):3119–31.

3. Querec TD, Akondy RS, Lee EK, Cao W, Nakaya HI, Teuwen D, et al. Systems biology
approach predicts immunogenicity of the yellow fever vaccine in humans. Nat
Immunol. 2009 Jan;10(1):116–25.

4. Hagan T, Nakaya HI, Subramaniam S, Pulendran B. Systems vaccinology: Enabling
rational vaccine design with systems biological approaches. Vaccine. 2015 Sep
29;33(40):5294–301.

5. Arunachalam PS, Scott MKD, Hagan T, Li C, Feng Y, Wimmers F, et al. Systems
vaccinology of the BNT162b2 mRNA vaccine in humans. Nature. 2021 Jul 12;

6. Obermoser G, Presnell S, Domico K, Xu H, Wang Y, Anguiano E, et al. Systems scale
interactive exploration reveals quantitative and qualitative differences in response to
influenza and pneumococcal vaccines. Immunity. 2013 Apr 18;38(4):831–44.

7.   Rinchai D, Anguiano E, Nguyen P, Chaussabel D. Finger stick blood collection for gene
867   expression profiling and storage of tempus blood RNA tubes. F1000Res. 2016;5:1385.

868   8.   Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired type I
869   interferon activity and inflammatory responses in severe COVID-19 patients. Science.
870   2020 Aug 7;369(6504):718–24.

871   9.   De Mot L, Bechtold V, Bol V, Callegaro A, Coccia M, Essaghir A, et al. Transcriptional
872   profiles of adjuvanted hepatitis B vaccines display variable interindividual homogeneity
873   but a shared core signature. Sci Transl Med. 2020 Nov 11;12(569):eaay8618.

874   10.   Altman MC, Rinchai D, Baldwin N, Toufiq M, Whalen E, Garand M, et al. Development of
875   a fixed module repertoire for the analysis and interpretation of blood transcriptome
876   data. Nat Commun. 2021 Jul 19;12(1):4385.

877   11.   Agniel D, Hejblum BP. Variance component score test for time-course gene set analysis
878   of longitudinal RNA-seq data. Biostatistics. 2017 Oct 1;18(4):589–604.

879   12.   Rawat A, Rinchai D, Toufiq M, Marr AK, Kino T, Garand M, et al. A Neutrophil-Driven
880   Inflammatory Signature Characterizes the Blood Transcriptome Fingerprint of Psoriasis.
881   Front Immunol [Internet]. 2020 [cited 2020 Dec 7];11. Available from:
882   https://www.frontiersin.org/articles/10.3389/fimmu.2020.587946/full

883   13.   Sena LA, Li S, Jairaman A, Prakriya M, Ezponda T, Hildeman DA, et al. Mitochondria Are
884   Required for Antigen-Specific T Cell Activation through Reactive Oxygen Species
885   Signaling. Immunity. 2013 Feb 21;38(2):225–36.

886   14.   Nakaya HI, Wrammert J, Lee EK, Racioppi L, Marie-Kunze S, Haining WN, et al. Systems
887   biology of vaccination for seasonal influenza in humans. Nat Immunol. 2011 Jul
888   10;12(8):786–95.

889   15.   Nakaya HI, Hagan T, Duraisingham SS, Lee EK, Kwissa M, Rouphael N, et al. Systems
890   Analysis of Immunity to Influenza Vaccination across Multiple Years and in Diverse
891   Populations Reveals Shared Molecular Signatures. Immunity. 2015 Dec 15;43(6):1186–
892   98.

893   16.   Rinchai D, Altman MC, Konza O, Hässler S, Martina F, Toufiq M, et al. Definition of
894   erythroid cell-positive blood transcriptome phenotypes associated with severe
895   respiratory syncytial virus infection. Clinical and Translational Medicine.
896   2020;10(8):e244.

897   17.   Tang B, Shojaei M, Wang Y, Nalos M, Mclean A, Afrasiabi A, et al. Prospective validation
898   study of prognostic biomarkers to predict adverse outcomes in patients with COVID-
899   19: a study protocol. BMJ Open. 2021 Jan 6;11(1):e044497.

900   18.   Elahi S. Neglected Cells: Immunomodulatory Roles of CD71+ Erythroid Cells. Trends
901   Immunol. 2019 Mar;40(3):181–5.

19. Elahi S, Ertelt JM, Kinder JM, Jiang TT, Zhang X, Xin L, et al. Immunosuppressive CD71+
erythroid cells compromise neonatal host defence against infection. Nature. 2013 Dec
5;504(7478):158–62.

20. Shahbaz S, Xu L, Osman M, Sligl W, Shields J, Joyce M, et al. Erythroid precursors and
progenitors suppress adaptive immunity and get invaded by SARS-CoV-2. Stem Cell
Reports. 2021 May 11;16(5):1165–81.

21. Spann KM, Tran K-C, Chi B, Rabin RL, Collins PL. Suppression of the induction of alpha,
beta, and lambda interferons by the NS1 and NS2 proteins of human respiratory
syncytial virus in human epithelial cells and macrophages [corrected]. J Virol. 2004
Apr;78(8):4363–9.

22. Bossert B, Conzelmann K-K. Respiratory syncytial virus (RSV) nonstructural (NS) proteins
as host range determinants: a chimeric bovine RSV with NS genes from human RSV is
attenuated in interferon-competent bovine cells. J Virol. 2002 May;76(9):4287–93.

23. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The
Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). JAMA.
2016 Feb 23;315(8):801–10.

24. Chiche L, Jourde-Chiche N, Whalen E, Presnell S, Gersuk V, Dang K, et al. Modular
transcriptional repertoire analyses of adults with systemic lupus erythematosus reveal
distinct type I and type II interferon signatures. Arthritis & Rheumatology (Hoboken,
NJ). 2014 Jun;66(6):1583–95.

25. Galani I-E, Rovina N, Lampropoulou V, Triantafyllia V, Manioudaki M, Pavlos E, et al.
Untuned antiviral immunity in COVID-19 revealed by temporal type I/III interferon
patterns and flu comparison. Nat Immunol. 2021 Jan;22(1):32–40.

26. Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, et al. Inborn errors of type
I IFN immunity in patients with life-threatening COVID-19. Science. 2020 Oct
23;370(6515):eabd4570.

27. Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann H-H, Zhang Y, et al.
Autoantibodies against type I IFNs in patients with life-threatening COVID-19. Science.
2020 Oct 23;370(6515):eabd4585.

28. Bastard P, Gervais A, Le Voyer T, Rosain J, Philippot Q, Manry J, et al. Autoantibodies
neutralizing type I IFNs are present in ~4% of uninfected individuals over 70 years old
and account for ~20% of COVID-19 deaths. Sci Immunol. 2021 Aug 19;6(62):eabl4340.

29. Cheng H, Peng Z, Luo W, Si S, Mo M, Zhou H, et al. Efficacy and Safety of COVID-19
Vaccines in Phase III Trials: A Meta-Analysis. Vaccines. 2021 Jun;9(6):582.

30. Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Phase I/II study
of COVID-19 RNA vaccine BNT162b1 in adults. Nature. 2020 Oct;586(7830):589–93.

31. Milanez-Almeida P, Martins AJ, Germain RN, Tsang JS. Cancer prognosis with shallow
tumor RNA sequencing. Nat Med. 2020 Feb;26(2):188–92.

32. Haack AJ, Lim FY, Kennedy DS, Day JH, Adams KN, Lee JJ, et al. homeRNA: A self-
sampling kit for the collection of peripheral blood and stabilization of RNA [Internet].
2021 Feb [cited 2021 Oct 30] p. 2021.02.08.430337. Available from:
https://www.biorxiv.org/content/10.1101/2021.02.08.430337v1

33. Sadarangani M, Marchant A, Kollmann TR. Immunological mechanisms of vaccine-
induced protection against COVID-19 in humans. Nat Rev Immunol. 2021
Aug;21(8):475–84.

34. Syed Ahamed Kabeer B, Tomei S, Mattei V, Brummaier T, McGready R, Nosten F, et al. A
protocol for extraction of total RNA from finger stick whole blood samples preserved
with TempusTM solution. F1000Res. 2018 Nov 2;7:1739.

35. Rinchai D, Roelands J, Toufiq M, Hendrickx W, Altman MC, Bedognetti D, et al.
BloodGen3Module: Blood transcriptional module repertoire analysis and visualization
using R. Bioinformatics. 2021 Feb 24;

 Table 1: Subject characteristics

Patient ID	Vaccine name	Gender	Age	Ethnicity	Previous COVID-19	Underlying disease	Drugs	Symptoms at prime (Type)	Symptoms at prime (Grade)	Symptoms at boost (Type)	Symptoms at boost (Grade)
PZB1	Pfizer Biontech	Female	38	Asian	Yes	No	no	Myalgia	G1	Fever/Myalgia	G1
PZB2	Pfizer Biontech	Male	47	Caucasian	Yes	No	no	Myalgia	G1	Myalgia	G1
PZB3	Pfizer Biontech	Male	57	Caucasian	Yes	T2D	Metformin, Insulin	Myalgia	G1	Myalgia	G1
PZB4	Pfizer Biontech	Female	34	Indian	No	No	no	None	NA	Chills/Insomnia/Headache/Myalgia/Fatigue	G3
PZB5	Pfizer Biontech	Male	38	Indian	No	No	no	Myalgia	G1	Myalgia	G1
PZB6	Pfizer Biontech	Female	48	Caucasian	No	No	no	Myalgia/Headache	G1	Myalgia	G1
PZB7	Pfizer Biontech	Female	34	Caucasia/Arab	No	Hashimoto thyroiditis	no	None	NA	None	NA
PZB8	Pfizer Biontech	Female	29	Arab	No	No	no	Myalgia/Swelling	G1	Fever/Myalgia	G1
PZB9	Pfizer Biontech	Male	41	Arab	No	Allergic rhinitis	no	Myalgia	G1	Myalgia	G1
PZB10	Pfizer Biontech	Female	35	Arab	No	No	no	Myalgia	G1	Fever/Insomnia/Myalgia/Fatigue	G2
PZB11	Pfizer Biontech	Male	41	Caucasian	No	No	no	Myalgia	G2	Myalgia	G1
PZB12	Pfizer Biontech	Female	34	Indian	No	Hypothyroidism	Levothyroxine	None	NA	Fever/Myalgia	G1
PZB13	Pfizer Biontech	Male	29	Indian	No	No	no	Fatigue	G1	Fever/Heaviness in arm	G2
PZB14	Pfizer Biontech	Female	38	Arab	No	Allergic	no	Myalgia/Headache	G1	Fatigue	G1
PZB15	Pfizer Biontech	Male	43	Arab	No	Hypothyroidism	Levothyroxine	None	NA	Myalgia/Headache	G2
PZB16	Pfizer Biontech	Female	39	Indian	No	No	no	Heaviness in arm	G1	Fever/Myalgia/Fatigue	G2
PZB17	Pfizer Biontech	Female	42	Indian	Yes	T2D, Hypertension	Methformin, Telmisartan	Fever/Headache/Myalgia/Fatigue	G2	Fatigue/Gastritis	G2
MDA18	Moderna	Male	42	Caucasian	No	Hypertension	Amlodipine, Ramipril	Myalgia	G1	Myalgia	G1
PZB19	Pfizer Biontech	Female	39	Caucasian	No	No	no	Myalgia	G1	Headache/Myalgia/Arthralgia	G3
MDA20	Moderna	Female	36	Arab	Yes	No	no	Chills/Myalgia	G2	Chills/Myalgia	G2
MDA21	Moderna	Female	36	Caucasian	No	No	no	Myalgia	G1	Feveer/Skin rash/Myalgia	G2
PZB25	Pfizer Biontech	Female	39	Caucasian	No	No	no	Myalgia	G1	Myalgia	G1

| MDA26 | Moderna | Female | 30 | Caucasian | Yes | Asthma | Seretide, Salbutamol | Fever/Headache/Myalgia/Fatigue | G3 | Asthma attack/Fever/Myalgia/Fatigue | G3 |

Footnotes:

T2D: Type 2 Diabetes

Vaccine type and lot, and subjects' characteristics were recorded, including demographic, biometric data, blood group, underlying diseases, drugs usage, and previous COVID-19 disease.

Every subject recorded and graded the symptoms that occurred after the first and second vaccinations doses, according to the NIH "DAIDS AE Grading Table"

Figure 1: Antibody response to COVID-19 mRNA vaccination. (A) Schematic representation of the study design. (B) The heatmap represents changes in abundance of antibodies specific to several SARS-CoV-2 antigens and control antigens relative to pre-vaccination levels. Red indicates a relative increase, and green indicates a relative decrease in abundance. Columns represent subjects arranged by timepoint and have colored tracks at the top indicating whether the subjects were previously infected with SARS-CoV-2 or not. The histogram above represents the average log2 fold-change over baseline for a given column. The rows represent antibody reactivities arranged by antigen specificity. The different rows represent the isotypes of reactive antibodies, arranged according to the color legend specified below the heat map. (C) Changes in antibody levels expressed as an "antibody index" are shown on the box plots, each corresponding to a given antibody type of a given specificity. Lines indicate changes for individuals previously infected with SARS-CoV-2 and who had recovered (in pink) and for naïve individuals (in green). Centerlines, box limits, and whiskers represent the median, interquartile range, and 1.5x interquartile range, respectively. Multiple pairwise tests (paired t-test) were performed comparing antibodies levels to baseline (D0). Asterisks: * represent $p < .01$, **represent $p < .001$, *** represent $p < .0001$.

Figure 2: Characterization of the post-prime interferon response signature. (A) The bar graph shows the cumulative module response at the various timepoints following the administration of the priming dose of the vaccine (noted P1-P14). The Y-axis values and numbers on the bars indicate the number of modules meeting the 15% response threshold (out of a total of 382 modules constituting the BloodGen3 repertoire, with percentage response corresponding to the proportion of transcripts predominantly increased or decreased compared to baseline using FDR < 0.1 as the cutoff to determine significance [DESeq2]). The number of modules for which abundance was predominantly increased is shown in red, and those for which abundance was predominantly decreased are shown in blue. The fingerprint grid plots represent the overall module responses on day 1 post-prime (P1) and day 2 post-prime (P2). Modules from the BloodGen3 repertoire occupy fixed positions on the fingerprint grids. They are arranged as rows based on membership to module aggregates (rows A1 through A38). Changes compared to the pre-vaccination baseline are indicated on the grid by red and blue spots of varying color intensity, which indicate the "percentage response" for a given module. The color key at the top indicates the various functions attributed

to the modules that are represented on the grid. The response of the six modules comprising aggregate A28 is represented on a line graph that shows the proportion of responsive transcripts for each module across all the post-prime timepoints. For each module, the statistical significance of the overall response was determined by time-course gene set enrichment analysis. Four of the six A28 modules met significance thresholds FDR < 0.1 (M8.3: p-value = 1.9-e4, FDR = 0.019, M10.1: p-value = 1.9-e4, FDR = 0.019, M15.127: p-value = 1.9-e4, FDR = 0.019, 727 and M15.86: p-value = 3.9-e4, FDR = 0.031) and all six A28 modules p < 0.05 (M13.17: p-value = 1.5-e3, FDR = 0.101 and M15.64: p-value = 0.044, FDR = 0.727). We also ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). (B) Heatmaps represent proportions of transcripts that changed within the six A28 modules at different timepoints and across different individuals compared to pre-vaccination baseline values. Red indicates that transcripts were predominantly increased over the baseline, and blue indicates that transcripts were predominantly decreased. Rows represent the six A28 modules arranged within an aggregate via hierarchical clustering. Columns represent samples grouped by timepoint and show profiles of individual subjects within each timepoint. (C) The heatmaps represent associations (Spearman correlation test) between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across nine consecutive days after the priming dose. The heatmap at the top provides the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the six A28 interferon modules. The heatmap below shows the significance of the correlations shown on the heatmap directly above, with the same ordering of rows and columns.

Figure 3: Characterization of responses on day 5 post-prime. (A) The bar graph shows the cumulative number of responsive modules at each timepoint following the administration of the priming dose of the vaccine (noted P1-P14). The fingerprint grid plot shows changes observed at P5 (day 5 post-prime). The position of the modules on the grid is fixed. The percent response of individual modules is represented on the grid by red and blue spots of varying color intensity denoting a predominant increase or decrease in abundance, respectively. The percentage response of a given module corresponds to the proportion of transcripts predominantly increased or decreased compared to baseline, meeting a significance cutoff of FDR < 0.1. The color key at the top indicates the various functions attributed to the modules that are represented on the grid. **(B)** The heatmap represents Log2 average fold change in abundance of transcripts constituting sets of modules associated with given functional annotations on P5. Rows represent individual transcripts grouped according to the module aggregate they originate from, corresponding to the different rows on the fingerprint grid plot on the left. Each module aggregate is associated with a unique function, as indicated by the color key above. The columns on the heatmap represent individual subjects coded with the type of vaccine received (Pfizer BioNtech = PZB; Moderna = MDA).

Figure 4: Fingerprint grid plots mapping changes observed on day 1 post-boost and across reference datasets.
The bar graphs show the cumulative module response at the various timepoints post-priming and booster doses (noted P1-P14 and B1-B14, respectively). The Y-axis values and numbers on the bars indicate the number of modules meeting the 15% response threshold (out of a total of 382 modules constituting the BloodGen3 repertoire, with percentage response corresponding to the proportion of transcripts predominantly increased or decreased compared to baseline meeting a significance cutoff of DESeq2, FDR < 0.1. The fingerprint grid plots show changes in transcript abundance in a given study group in comparison to baseline (pre-vaccination sample or uninfected control group – with the percent response of individual modules shown by red and blue spots of varying color intensity denoting predominant increase or decrease in abundance, respectively. Changes are shown in the top grid for a group comparison of 1 day after receiving the booster dose of COVID-19 mRNA vaccines with baseline pre-vaccination samples (this study). Grids in the middle and bottom positions show changes for patients with acute infections caused by influenza virus (public dataset) or SARS-CoV-2 (this study) and for patients with bacterial sepsis (public dataset). The color key at the top indicates the various functions attributed to the modules that occupy a fixed position on the grid plot.

Figure 5: Characterization of the day 1 post-boost interferon response signature. (A) The fingerprint grid plot maps the modular response observed on day 1 post-boost (percent response is determined based on statistical cutoff: DESeq2, FRD < 0.1). The six modules forming the A28 aggregate are highlighted. The line graphs below represent the summarized percentage responses at the module level, encompassing all study subjects (one line per module). Percentage response accounts for the proportion of transcripts for a given module showing differences in abundance post-prime (left) or post-boost (right) compared to baseline pre-vaccination levels. Changes in transcript abundance post-prime and post-boost for two distinct sets of interferon response modules that received the denomination A28/S1 and A28/S2 are plotted on separate graphs. For each module, statistical significance for the overall response was determined by time course gene set enrichment analysis. Significance was reported post-prime in Figure 2. Post-boost all six A28 modules met significance thresholds p < 0.001 and FDR < 0.001 (M8.3: p-

value = 1.9-e4, FDR = 3.6-e4, M10.1: p-value = 1.9-e4, FDR = 3.6-e4, M13.17: p-value = 1.9-e4, FDR = 3.6-e4, M15.127: p-value = 1.9-e4, FDR = 3.6-e4, M15.64: p-value = 1.9-e4, FDR = 3.6-e4 and M15.86: p-value = 1.9-e4, FDR = 3.6-e4). In addition, we ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). The circle packing plots on the left show module responses at the individual transcript level for two public blood transcriptome datasets. The larger circle below indicates official symbols for the individual transcripts. It also highlights the modules included in A28/S2, shown directly on the right. The smaller circles above show changes in abundance of A28 transcripts for two public datasets. One study (GSE11342) measured blood transcriptional response in patients with Hepatitis C infection treated with alpha-interferon (23). The second study (GSE26104) measured transcriptional response in subjects with multiple sclerosis treated with beta-interferon (24). A red circle indicates a significant increase in the abundance of transcripts compared to the pre-treatment baseline (*|fold-change| > 1.5, FDR < 0.1). **(B)** Changes in abundance compared to baseline pre-vaccination levels are represented on a heatmap, with modules as rows and individual samples as columns. The modules are arranged by hierarchical clustering based on abundance patterns across samples. The samples are arranged by timepoints post-prime (top) and post-boost (bottom). **(C)** The heatmaps represent associations (Spearman correlation) between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across nine consecutive days after the booster dose. The heatmap on top provides the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the six A28 interferon modules. The heatmap below shows the significance of the correlations shown on the heatmap on top, with the same ordering of rows and columns.

Figure 6: Characterization of post-boost inflammation, erythroid cell, and plasmablast responses. (A) The bar graph at the top represents the number of response modules at any given time point post-prime and post-boost. The fingerprint grid plots show the modules that had changes compared with a fixed visualization and interpretation framework. Changes are shown for the day 1 post-boost timepoint (left) as well as day 5 (right) (percent response is determined based on statistical cutoff: DESeq2, FRD < 0.1). On the left grid, modules belonging to aggregates A35 (associated with inflammation) and A37 (associated with erythroid cells) are highlighted. The profiles of those modules are represented on the line graphs below, which show the average percentage responses of A35 and A37 modules across multiple timepoints. The percentage response for a given module is the proportion of its constitutive transcripts showing significant changes, ranging from 0% to 100% when transcripts were predominantly increased to 0% to −100% when transcripts were predominantly decreased. Each line represents the profile of the modules

constituting a given aggregate. Line graphs for A35 were split into three sets according to differences in clustering patterns (A35/S1, A35/S2, and A35/S3). On the right grid, modules belonging to aggregates A27 (associated with platelets) are highlighted. The corresponding line graph below represents the changes in abundance of A27 modules over time following administration of the second dose of vaccine. For each module, statistical significance for the overall response was determined by time course gene set enrichment analysis using the dearseq R package. For A35, 20 of 21 modules met significance thresholds (p-value < 0.05 and FDR < 0.01). It was also the case in 11 of 11 modules for A37 and 4 of 5 modules for A27 (**Supplementary file 4**). In addition, we ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). **(B).** The heatmaps represent associations between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across nine consecutive days after the booster dose. Specifically, the heatmap at the top represents the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the five A27 plasmablast modules. The heatmap below shows the significance of the correlations shown on the heatmap at the top, with the same order of rows and columns.

Figure 7: Comparing patterns of interferon response in vaccinated individuals and a cohort of COVID-19 patients. A. The tSNE plot represents similarities in patterns of interferon response induction across the six modules forming aggregate A28 and among samples comprised in our vaccination cohort and one of our COVID-19 disease cohorts (PREDICT-19 / Italy). COVID-19 samples are shown in red along with specific post-vaccination timepoints (post-prime days 1 and 2 [P1, P2], post-boost days 1 and 2 [B1, B2]). Samples from the consolidated cohorts were partitioned into 8 clusters via k-means clustering, the distribution of which is shown on the tSNE plot on the top right. **B.** Heatmaps show patterns of response for the six interferon response modules across the eight sample clusters. The red colors indicate that the abundance of transcripts for a given module is predominantly increased with the intensity representing the proportion of constitutive transcripts meeting a given threshold, which at the level of individual samples is a fixed fold change and difference cutoff (|Fold change| > 1.5, and |difference| > 10 in a given sample over its respective pre-vaccination baseline). The blue color denotes a predominant decrease in abundance of constitutive transcripts compared to the same individual's pre-vaccination baseline. Details are shown below for Clusters 3, 5, and 8 in separate heatmaps.

Figure 8: Comparison of interferon response patterns of vaccinated individuals and a cohort of COVID-19 patients with severe disease under intensive care. A. The tSNE plot represents similarities in patterns of interferon response induction across the six modules forming aggregate A28 and among samples comprised in our vaccination cohort and one of our COVID-19 disease cohorts (IMPROVISE). Specific post-vaccination timepoints (post-prime days 1 and 2 [P1, P2], post-boost days 1 and 2 [B1, B2]), as well as repeat sampling for COVID-19 patient (TP1, TP2, TP3, TP4, all collected during ICU stay) are shown on the plot on the left. Samples from the consolidated cohorts were partitioned into 8 clusters via k-means clustering, the distribution of which is shown on the tSNE plot on the center. Length of ICU stay is shown on the tSNE plot on the right. Patterns of response for the six interferon response modules across the eight sample clusters are shown on a heatmap below. The red colors indicate that the abundance of transcripts for a given module is predominantly increased with the intensity representing the proportion of constitutive transcripts meeting a given threshold, which at the level of individual samples is a fixed

fold change and difference cutoff (|Fold change |> 1.5, and |difference| > 10 in a given sample over its respective pre-vaccination baseline). The blue color denotes a predominant decrease in abundance of constitutive transcripts compared to the same individual's pre-vaccination baseline. **B.** The heatmap shows patterns of interferon responses for COVID-19 patients with severe disease upon ICU admission. Multiple clinical parameters are shown on the tracks above (ECMO [Extracorporeal Membrane Oxygenation], HTN [Hypertension], CAD [Coronary Artery Disease], CKD [Chronic Kidney Disease], CHF [Congestive Heart Failure]). The histogram represents the length of stay in the hospital, in the ICU, and under mechanical ventilation, in days. **C.** The bar graph represents for different datasets the proportion of samples corresponding to Interferon Response Transcriptional Phenotypes (IRTP) I, II or III, according to the following definition: IRTP I = (S1++S2+","S1++S20","S1+S2); IRTP II = (S1++S2++); IRTP III = ("S1-S2++","S1-S2+","S10S2++","S10S2+","S1-S20"). The datasets in question were derived from the present study: response to COVID-19 mRNA vaccination (N=23) on days 1 and 2 post-prime (P1 and P2, respectively), days 1 and 2 post-boost (B1 and B2, respectively); as well as COVID-19 disease cohorts (PREDICT-19 [N=114] and IMPROVISE [N=). Others were derived from an earlier study and include reference cohorts of patients with acute influenza infection (FLU, N=25), HIV infection (N=28), active pulmonary tuberculosis (PTB, N=23), acute RSV infection (N=70), bacterial sepsis (N=33) and SLE (N=55).

Figure 9: Summary. This diagrammatic representation summarizes the temporal trajectories of blood transcriptional signatures elicited in response to the first and second doses of mRNA vaccines.