TITLE: High temporal resolution transcriptomic profiling delineates distinct patterns of

interferon response following Covid-19 mRNA vaccination and SARS-CoV2 infection

AUTHORS

Darawan Rinchai[1#*], Sara Deola[1#], Gabriele Zoppoli[2,3§], Basirudeen Syed Ahamed Kabeer[1§],

Sara Taleb[4§], Igor Pavlovski[1±], Selma Maacha[1±], Giusy Gentilcore[1], Mohammed Toufiq[1], Lisa

Mathew[1], Li Liu[1], Fazulur Rehaman Vempalli[1], Ghada Mubarak[1], Stephan Lorenz[1], Irene

Sivieri[3,5,6], Gabriella Cirmena[2], Chiara Dentone[2], Paola Cuccarolo[3], Daniele Giacobbe[2,5],

Federico Baldi[5], Alberto Garbarino[3], Benedetta Cigolini[3], Paolo Cremonesi[7], Michele

Bedognetti[8], Alberto Ballestrero[2,5], Matteo Bassetti[2,5], Boris P. Hejblum[9], Tracy Augustine[1],

Nicholas Van Panhuys[1], Rodolphe Thiebaut[9], Ricardo Branco[1], Tracey Chew[10], Maryam

Shojaei[11,12], Kirsty Short[13,14], Carl Feng[15,16], PREDICT-19 consortium, Susu M. Zughaier[17],

Andrea De Maria[2,5], Benjamin Tang[12,18], Ali Ait Hssain[19], Davide Bedognetti[1&], Jean-Charles

Grivel[1&] and Damien Chaussabel[1&*]

1. Research Branch, Sidra Medicine, PO Box 26999, Doha, Qatar

2. IRCCS Ospedale Policlinico San Martino, Genoa, Italy

3. Department of Internal Medicine and Medical Specialties, University of Genoa, Genoa,

Italy

4. Division of Genomics and Translational Biomedicine, College of Health and Life Sciences,

Hamad Bin Khalifa University, Doha, Qatar

5. Division of Infectious Diseases, Department of Health Sciences, University of Genoa,

Genoa, Italy

6. Department of Experimental and Clinical Medicine, School of Internal Medicine,

University of Florence, Florence, Italy

7. Emergency Department, E.O. Ospedali Galliera Genova, Italy

8. Azienda Sanitaria Locale 3 Genovese, Genova, Liguria, Italy

9. Univ. Bordeaux, Department of Public Health, Inserm U1219 Bordeaux Population Health

Research Centre, Inria SISTM, F-33000 Bordeaux, France

10. Sydney Informatic Hub, The University of Sydney, Sydney, Australia

11. Nepean Clinical School, University of Sydney, Sydney, New South Wales, Australia;

12. Westmead Institute for Medical Research, Westmead, New South Wales, Australia.

13. The University of Queensland, School of Chemistry and Molecular Biosciences, St Lucia,

Brisbane, Queensland, Australia;

14. Australian Infectious Diseases Research Centre, The University of Queensland, Brisbane,

Queensland, Australia.

15. School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney,

Sydney, NSW, Australia;

16. Tuberculosis Research Program, Centenary Institute, The University of Sydney, Sydney,

NSW, Australia.

17. College of Medicine, QU Health, Qatar University, PO Box 2713, Doha, Qatar

18. Westmead Institute for Medical Research, Westmead, New South Wales, Australia.

19. Medical Intensive Care Unit, Hamad General Hospital, PO BOX 3050, Doha, Qatar

#, §, ±, & These authors contributed equally to the work

* To whom correspondence may be addressed:

drinchai@sidra.org

dchaussabel@sidra.org

ABSTRACT:

Knowledge of the factors contributing to the development of protective immunity after

vaccination with COVID-19 mRNA vaccines is fragmentary. Thus we employed high-

temporal-resolution transcriptome profiling and in-depth characterization of antibody

production approaches to investigate responses to COVID-19 mRNA vaccination. There were

marked differences in the timing and amplitude of the responses to the priming and booster

doses. Notably, two distinct interferon signatures were identified, that differed based on

their temporal patterns of induction. The first signature (S1), which was preferentially

induced by type I interferon, peaked at day 2 post-prime and at day 1 post-boost, and in

both instances was associated with subsequent development of the antibody response. In

contrast, the second interferon signature (S2) peaked at day 1 both post-prime and post-

boost but was found to be potently induced only post-boost, where it coincided with a

robust inflammation peak. Notably, we also observed "post-prime-like" ($S1^{++}, S2^{0/+}$) and

"post-boost-like" ($S1^{++}, S2^{++}$) patterns of interferon response among COVID-19 patients. A

post-boost-like signature was observed in most severely ill patients at admission to the

intensive care unit and was associated with a shorter hospital stay. Interestingly, severely ill

patients who stayed hospitalized the longest showed a peculiar pattern of interferon

induction ($S1^{-/0}, S2^{+}$), that we did not observe following the administration of mRNA

vaccines. In summary, high temporal resolution profiling revealed an elaborate array of

immune responses elicited by priming and booster doses of COVID-19 mRNA vaccines.

Furthermore, it contributed to the identification of distinct interferon-response phenotypes

underpinning vaccine immunogenicity and the course of COVID-19 disease.

INTRODUCTION

INTRODUCTION

COVID-19 vaccines are critical to the ongoing efforts to control the SARS-CoV-2 coronavirus

pandemic. To date, nine vaccines have received some form of approval for use in humans,

and phase III trials are ongoing for an additional 11 vaccines (1). Notable differences exist

among the vaccine products in terms of their design and the levels of protection they

confer, as well as the type, incidence, and severity of adverse events they may elicit. Gaining

a comprehensive understanding of the immunological factors underpinning the different

responses to diverse vaccines is a major endeavor. Yet, this knowledge is necessary for

guiding timely decisions to modulate vaccination protocols (e.g., the use of different types

of vaccines for the priming and booster doses). This information may also assist in matching

of individuals with the growing number of available vaccines based on their demographics,

health status, or any other relevant clinical/molecular phenotypes. Blood transcriptome

profiling, which measures the abundance of transcripts in whole blood and on a system-

wide scale, has previously been employed to comprehensively profile the immune

responses elicited by vaccines. Notably, this approach identified innate immune signatures

arising within hours after administering widely used, as well as experimental, vaccines (2). In

a recently published report, Arunachalam et al. described the blood transcriptome profiles

measured following BNT162b2 mRNA COVID-19 vaccination (3). They reported the presence

of an interferon (IFN) signature 1 day after the priming vaccination that was no longer

detectable on day 7. They further found that a more complex IFN/inflammatory response

was present 1 day after the administration of the booster dose.

However, as the use of mRNA vaccines in humans is new, the precise kinetics of the

immune response they elicit have not been characterized. Therefore, through this study, we

endeavored to profile the transcriptional changes occurring on a systems-wide scale prior to

the administration of the first vaccine dose and for 9 consecutive days thereafter, with

subjects also collecting samples for deep serological profiling at three time points. The same

sampling and profiling schedule being repeated to assess the response to the second dose

of the vaccine. Together, this work permitted the precise delineation of a well-orchestrated

immune response to COVID-19 mRNA vaccines and identified marked differences in the

magnitude, nature, and timing of the transcriptional signatures elicited by prime and boost

vaccination. Most notably, differences in temporal patterns of responsiveness revealed

distinct components of the interferon response, which is known to play a key role in

controlling SARS-CoV-2 infection (4) and was also found here to associate with the

subsequent development of the antibody response post-vaccination.

RESULTS

Study design, implementation and serological profiling

We successfully recruited a cohort of volunteers and implemented a high-frequency

sampling protocol following the administration of the priming and booster doses of COVID-

19 mRNA vaccines. The response to each dose was ascertained via the collection and

profiling of samples at 10 consecutive daily timepoints: immediately before vaccination and

for 9 days thereafter. We collected samples for serological profiling at three time points:

prior to vaccination and on days 7 and 14 post-vaccination. A self-sampling blood collection

protocol was implemented so that subjects could extract small volumes (50 µl) of RNA-

stabilized blood at the required frequency (the approach is described in the Methods

section and an earlier publication (5)). RNA sequencing profiles were generated using a cost-

effective 3'-biased library preparation protocol (Lexogen QuantSeq). We also established a

self-collection of capillary blood samples for measuring the levels and types of COVID-19-

specific antibodies. Antibody profiles were generated from samples collected by Volumetric

Absorptive Micro Sampling using a multiplexed Bead array established by our team (see

Methods for details). Overall, 23 subjects were enrolled in the study, and the characteristics

of this cohort are reported in **Table 1**. In total, 23 subjects received either two doses of the

Pfizer/BioNTech mRNA vaccine (BNT162b2, N = 19) or two doses of the Moderna mRNA

vaccine (N = 4). Among those 23 subjects, six had recovered from COVID-19 in the months

preceding the administration of the first vaccine dose. In total 440 RNA sequencing profiles

were generated, and this extensive dataset was shared publicly in GEO with the accession

number GSE190001. The serological profiles included reactivity to Spike proteins from seven

human coronaviruses (including SARS-CoV-2), as well the Nucleo and Envelope proteins, the

receptor-binding domain, and the stabilized S-trimer of SARS-CoV-2. The data are provided

in **Supplementary File 1**. The seroreactivity to each of these antigens was dissected by

measuring the total IgG, total IgA, and IgM, as well as the finer-scale IgG and IgA subtypes.

Serological profiling data showed a rise in the levels of antibodies in the plasma of the

subjects post-vaccination (**Figure 1A**), and this included antibodies specific for the SARS-

CoV-2 Spike protein, which is targeted by COVID-19 vaccines. No responses to the Envelope

protein were measured. Some cross-reactivity was observed with the SARS Spike protein.

Notably higher antibody levels were induced after the first dose in individuals who had been

previously infected with the virus (**Figure 1B**).

Altogether, the implementation of this protocol established the feasibility of

obtaining stabilized-RNA blood samples from study-subjects post-vaccination at high

temporal frequencies. We generated a large dataset using a cost-effective RNA-sequencing

protocol that served as the basis for subsequent analyses presented in this paper and was

deposited in a public repository. A detailed map of the serological profiles of the subjects

enrolled in the study was obtained that permitted us to explore the possible associations

between blood transcriptional responses and vaccine immunogenicity.

The post-prime interferon response peaks at day-2 and correlates with antibody response

Vaccines can elicit innate immune responses that are detectable systemically via blood

transcriptome profiling. But not all of them do, as for instance in the case of the aluminum-

adjuvanted Hepatitis B vaccine (6). Our first question, therefore, was whether

transcriptional changes could be observed during the first few days following the

administration of COVID-19 mRNA vaccines.

Analyses were carried out employing a fixed repertoire of 382 transcriptional modules

(BloodGen3) that we had recently established and characterized functionally (7)(see

methods section for details). Module responses were determined across all time points, and

the differential gene-set enrichment functions of the dearseq R package were run to assess

whether changes observed throughout the 9 days post-prime were statistically significant

(8). This analysis identified significant temporal changes for 22 of the 382 modules

constituting the BlooGen3 repertoire (**Supplementary File 3**).

Only seven modules were found to be changed at any given time point during the first 3

days following the administration of the priming dose of the vaccine (**Figure 2A**). The

abundance of four modules was consistently increased across these time points, and all four

belonged to the module aggregate A28. Each "module aggregate" regrouped sets of

modules that showed consistent abundance profiles across a reference set of 16 disease

cohorts that were employed for the construction of the BloodGen3 repertoire (see methods

for details). The module aggregate in question ("Aggregate A28") is comprised of six

modules. As described in detail in one of our recent publications, all six are associated with

interferon responses (7). The gene composition of the modules and the functional

annotations are provided herein (relevant information is provided in **Supplementary File 2**

and can be accessed interactively via: https://prezi.com/view/E34MhxE5uKoZLWZ3KXjG/).

The responses observed on days 1 and 2 post-prime were mapped onto fingerprint grid

plots, where modules occupy a fixed position and are arranged by aggregate, with each

aggregate occupying a given row (**Figure 2A**). Time-course gene-set enrichment analysis

confirmed that changes observed over time for the four A28 modules were significant. The

p-values for the two remaining modules from Aggregate A28 did not reach significance. The

response profiles of the A28 modules showed a peak on day 2 post-vaccination. This was

also visible on a heatmap showing responses at each timepoint across individual subjects

(**Figure 2B**). We next examined whether this signature correlated with antibody responses

measured 14-days post -as well as at 14-days post-boost (**Figure 2C**). For this, correlation

analyses were run at the module level within Aggregate A28 using, as the endpoint, fold-

changes in antibody levels on days 7 and 14 post-prime and days 7 and 14 post-boost

relative to the pre-vaccination baseline (immediately prior to the administration of the first

dose of COVID-19 mRNA vaccines). "Significance hotspots" were identified when most

modules within a given aggregate reached correlation significance thresholds. In the case of

the post-prime interferon signature, we identified a significance hotspot on days 2 and 3

post-prime for a subset of three interferon modules, M10.1, M15.127, and M83, while a

fourth module, M15.86, also displayed significant correlations across all antibody types, but

only on day 2.

Thus, we found that the administration of the priming dose of mRNA vaccine induces a

potent innate response characterized by an increase in the abundance of transcripts

associated with interferon responses. It was followed by more marked changes over the

subsequent days, as indicated by the rise in the number of responsive modules beyond the

third day.

A decrease in inflammation is accompanied by an increase in adaptive immune response

genes on day 5 post-prime

We were next interested in characterizing the changes occurring beyond the first 3 days

following administration of the priming dose. In total, 18 modules displayed changes on day

4 post-prime, of which 12 showed a decrease in abundance. These modules belonged to

three aggregates that have been associated with inflammation (A31, A33, A35). Most

changes were observed on day 4, but for some modules, changes were apparent starting on

day 3 and continued beyond day 4, day 5, or even day 6 (**Supplementary Figure 1**). In our

earlier work, modules within the BloodGen3 Aggregate A35 were associated with systemic

inflammation mediated by neutrophils and were found to constitute a common

denominator across a wide range of pathologies in which systemic inflammation is present

(9). The association of A35 with inflammatory processes was also ascertained based on the

results of the functional profiling analyses and the restriction in transcript expression in the

reference datasets (7). Detailed functional annotations can be accessed via interactive circle

packing charts: https://prezi.com/view/7Q20FyW6Hrs5NjMaTUyW/). Module Aggregate A33 has

not been investigated as extensively in any of our prior studies but was clearly associated

with   inflammation   via   functional   profiling

(https://prezi.com/view/VBqKqHuLWCra3OJOIZRR/).

216   The number of responsive modules post-prime peaked on day 5, with a total of 42

217 modules showing differences in comparison to the pre-vaccination baseline (**Figure 3**). At

218 this timepoint, most modules showed an increase in abundance (29 were increased and 13

219 decreased). Some of those modules belonged to aggregates that were associated with

220 adaptive immunity, most notably A27, which is associated with plasmablast responses

221 (three out of five modules were responsive at this timepoint). Others were mapped to

222 module aggregates for which immunological significance is not yet well defined; these

223 included Aggregate A24 (associated with oxidative phosphorylation, in which 6 out of 11

224 modules were responsive) and Aggregate A26 (in which 4 out of 15 modules were

225 responsive, none of which are functionally annotated). Other immune-relevant modules

226 found to be increased at this timepoint are associated with T-cells (M12.6 from aggregate

227 A1). Detailed functional annotations for module aggregate A1 can be accessed via

228 https://prezi.com/view/sxap39tKxkmCNTTNIlVO/). The five modules in Aggregate A27 are

229 functionally associated with plasmablasts, as determined from the restriction of expression

230 observed for the genes forming this aggregate in a reference dataset, including a wide range

231 of cell populations (contributed by Monaco et al. (14)) and by the presence of the

232 plasmablast marker CD38 and other associated genes (*IGJ*, *TNFRSF17*, *TXNDC5*) in one of the

233 modules (M12.15). Detailed annotations for A27 can be accessed via

234 https://prezi.com/view/GgIiA0K9kSFHbpVj2I85/. Notably, these signatures appeared to be

235 transient, and no modules were increased on day 6 post-prime.

236   Taken together, our data showed a peak in the number of responsive modules on

237 day 5 post-prime. The signature at the module repertoire-level appeared relatively modest.

238 A decrease in the abundance of transcripts associated with inflammation was accompanied

239 by an increase in the abundance of transcripts associated with adaptive immune responses.

Notably, the latter was more subdued, and it also appeared earlier than typically seen in

vaccination responses, in which plasmablast responses are usually observed around day 7

post-vaccination (10–12).

A post-boost interferon signature peaks on day 1 and correlates with antibody responses

After delineating temporal responses post-prime, we examined changes after the

second dose of COVID-19 mRNA vaccines. Time course gene set enrichment analysis

identified significant temporal changes for 311 of 382 modules comprising the BloodGen3

repertoire (**Supplementary File 4**). After the booster dose, the peak number of responsive

modules occurred on day 1, with 261 responsive modules or about two-thirds of the 382

modules constituting the BloodGen3 repertoire (**Figure 4A**). This number decreased sharply

afterwards, with 115 responsive modules on day 2 and only 9 responsive modules on day 3.

The kinetic and amplitude of the post-boost response contrasted markedly with that

observed post-prime, when, as described above, the number of responsive modules after

the first dose instead peaked on day 4, with changes found in 42 modules at that timepoint.

As seen from the fingerprint grid plot, the day 1 post-boost response was extensive

and polyfunctional (**Figure 4B**). An overall decrease in abundance was observed for

aggregates broadly associated with lymphocytic cells (Aggregates A1-A8) and increased for

module aggregates associated with myeloid cells, inflammation, and circulating erythroid

cells (Aggregates A33-A38). In addition, a marked increase in the abundance of modules

associated with interferon responses was also observed (Aggregate A28). We next

compared the day 1 response fingerprint of the COVID-19 mRNA booster vaccine to

fingerprints derived from reference datasets encompassing a wide range of infectious

diseases (these cohorts are described in previously published work (7,13); the respective

blood transcriptome fingerprint collections are accessible via a dedicated web application:

https://drinchai.shinyapps.io/BloodGen3Module/). In addition, we more recently employed

the BloodGen3 module repertoire in the analysis of two COVID-19 blood transcriptome

datasets: one cohort comprising 77 Covid-19 patients with disease severities ranging from

mild and moderate to severe (the "PREDICT-19 consortium Italian cohort dataset") (14),

while the second cohort comprised 40 COVID-19 patients recruited at the time of admission

to the intensive care unit (ICU) ("IMPROVISE cohort whole blood dataset"). These high-level

comparisons showed, firstly, that the extent of the changes associated with the day 1

response to the second dose of the COVID-19 mRNA vaccine was consistent with that

observed in some patient cohorts with acute infections (**Figure 4B**). More specifically, they

were found to most resemble the responses seen in a cohort of subjects with influenza

infection, with a marked interferon response (A28) and an inflammation signature (A33,

A35). At a higher level, these response patterns were also generally consistent with those

observed in patients with a COVID-19 infection. However, the changes that occurred in

response to the vaccinations were not as extreme as those found, for instance, in patients

with sepsis or with the most severe form of COVID-19 (i.e., the IMPROVISE dataset) (most

notably for inflammation [A33, A35] and erythroid cell responses [A36-A38]).

Overall, the BloodGen3 transcriptome fingerprint observed on day 1 after the

second dose contrasted markedly with that observed on day 1 post-prime, which was

comparatively muted. The interferon response signature was indeed the only common

denominator between the responses to the first and second doses, as it was observed in

both cases in the first few days following administration of the vaccine. We therefore began

to dissect the post-boost response by examining this interferon response signature in more

detail.

Following the administration of the booster dose, the interferon response was

noticeably sharper in comparison to the post-prime response and peaked on day 1 instead

of day 2 (**Figure 5A**). This was illustrated by the difference in the maximum average module

response, which was close to 50% of the constitutive transcripts on day 2 post-prime and

greater than 80% on day 1 post-boost.

We decided to then perform hierarchical clustering to identify the subsets of modules

within the A28 aggregates that were grouped together based on patterns of transcript

abundance across all subjects and timepoints. Two sets of three modules each were, thus,

identified within the A28 aggregate. The first set comprised modules M8.3, M10.1, and

M15.127 (referred to as A28/S1), and the second set comprised modules M16.64, M13.17,

and M15.86 (referred to as A28/S2). Interestingly, we observed post-prime that, while

modules in A28/S1 peaked on day 2, those belonging to A28/S2 peaked on day 1 (**Figure

5B**). Furthermore, A28/S1 modules showed an extended peak post-boost, with day 2 levels

being almost identical to those of the day 1 peak, while A28/S1 modules peaked sharply on

day 1, with levels decreasing rapidly thereafter. These findings suggest that both sets of

modules measured distinct types of interferon response. Indeed, the reference public

datasets in which responses to type 1 interferon were measured in vivo indicated that

A28/S1 modules are likely to represent type 1 interferon responses (**Figure 5B**), while we

postulated that A28/S2 modules represent a type 2 interferon response. Enrichment of

literature terms associated with interferon or anti-viral response was previously found to be

more pronounced in the case of modules forming the A28/S1 set (7). Indeed, these

comprise some of the better recognized "canonical" interferon response genes, such as

Oligoadenylate Synthetase family members (OAS1, OAS2, OAS3, OASL), Interferon Induced

Protein family members (IFI6, IFI27, IFI35, IFI44, IFI44L), as well as Interferon Induced

Protein With Tetratricopeptide Repeats family members (IFIT1, IFIT3, IFIT5). Modules

forming the A28/S2 set comprise instead most notably members of the Nuclear Antigen

family members SP100, SP110 and SP140, which are associated with interferon gamma

signaling, as well as transcription factors IRF9 and STAT2. Composition and functional

annotations   for   A28   modules   can   be   explored   further   at:

https://prezi.com/view/E34MhxE5uKoZLWZ3KXjG/.

Finally, a strong association was found between the post-boost interferon signature and

the subsequent development of an antibody response. Indeed, positive correlations were

observed for all six A28 modules that reached significance on days 1, 2, and 3 post-boost.

Notably, this differed from the post-prime interferon response, for which significance was

reached only for four of the six modules and only on days 2 and 3.

Taken together, the high temporal resolution profiling results permitted the delineation

of distinct patterns of post-prime and post-boost interferon responses. The timing of the

responses observed at the individual module level contributed to the definition of the two

distinct sets of interferon modules. One set was associated with responses to type I

interferon in vivo and dominated the post-prime response, with a peak on day 2. The post-

boost response showed a strong induction of both sets and also peaked on day 1.

Inflammation and erythroid cell signatures peak sharply on day 1 post-boost

We continued the dissection of the day 1 post-boost signature, focusing this time on

responses associated with inflammation and circulating erythroid cell precursors.

Aggregates A33 and A35, which are associated with inflammation, tended to

decrease from day 4 through day 6 post-prime but displayed a contrastingly sharp and

transient increase in abundance post-boost. Indeed, a well-delineated response peak was

observed on day 1 post-boost for both the A33 and A35 modules (**Figure 6**), but in contrast

to the interferon response, it did not extend beyond the first day. Three distinct response

patterns were identified via hierarchical clustering among the 21 modules that formed

aggregate A35. The "Boost A35/S1" set comprised five modules, while "Boost A35/S2" and

"Boost A35/S3" comprised ten and six modules, respectively. The distinction between those

three A35/inflammation module sets was rather more subtle than was the case for the

A28/interferon sets. Indeed, all three module sets peaked on day 1 post-boost. Differences

were rather apparent in the inflection of changes measured on days 2 and 3 post-boost and

in the "recovery phase", as abundances appeared to dip below baseline and progressively

rise to reach pre-vaccination levels. The underlying biological factors driving the grouping of

the modules to those three distinct sets could not be identified at this time.

Modules for three aggregates broadly associated with erythroid cell signatures also

displayed a sharp but transient increase in transcript abundance on day 1 post-boost.

However, the abundance tended to dip afterwards, with a lower peak on day 4 post-boost,

before recovering by day 7. Functionally, this signature was found to be most prominently

associated with immunosuppressive states, such as late-stage cancer or pharmacological

immunosuppression (13), which is consistent with published functional studies (15,16). We

also found such signatures were associated with more severe manifestations in babies

infected with Respiratory Syncytial Virus (RSV) (13). Finally, we did not find evidence of an

association between the day 1 post-boost inflammation or erythroid cell signatures and the

antibody responses.

A plasmablast signature peaks on day 4 post-administration of the booster dose and

correlates with antibody responses

After the booster dose, the number of responsive modules peaked sharply on day 1,

then rapidly subsided beyond day 2, with the number of responsive modules on days 3, 4, 5,

and 6 being reduced to 8, 11, 3, and 2, respectively. Yet, changes within this later timeframe

are meaningful, as they specifically concern the set of five modules comprising aggregate

A27, which is associated with the presence of antibody-producing cells in the peripheral

blood.

Three of the five A27 modules showed significant alterations after the booster dose

(M16.60, M13.32, M12.15) (**Figure 6**). The proportion of differentially expressed transcripts

in each module was relatively modest (with an average of 15% at the peak of response),

especially in comparison with the interferon signatures described above (with an average of

>80% for some modules at the response peak). The trajectories of the five A27 modules

appeared relatively consistent, with only one of the modules (M15.110) showing a different

pattern, i.e., a peak on day 6, slightly above the levels observed on day 4. We also examined

the association of this post-boost plasmablast signature with the antibody response and

found a significant association starting from about day 3 and lasting until day 7 post-boost

(**Figure 6**).

In summary, COVID-19 mRNA vaccination induced a plasmablast response that peaked

at a post-vaccination timepoint that is not routinely included in sampling protocols (day 4).

Instead, plasmablast responses have been reported around day 7 post-vaccine

administration (e.g., with influenza or pneumococcal vaccines (11)). However, this type of

signature could not be detected following COVID-19 mRNA vaccination at this timepoint.

We also found an association between this post-boost plasmablast signature and the

development of an antibody response.

Patterns of interferon responses consistent with those measured day 1 post-boost are found in some COVID-19 patients and associated with shorter intensive care unit stays

An important objective of this study was to identify key determinants of vaccine immunogenicity. We found that only a few of the blood transcriptional signatures that have been identified were associated with the development of antibody responses following COVID-19 mRNA vaccination. Logically, the plasmablast signature that we have just described was one of them, and the other two were the interferon signatures observed post-prime and post-boost. We surmised that if such interferon signatures were observed in COVID-19 patients, they may also be associated with the development of a protective immune response and favorable clinical outcome. This might especially be the case for the post-boost interferon response, since a significant rise in neutralizing antibodies was reportedly observed only after the second dose of the vaccine.

Therefore, we sought to next determine whether "post-prime-like" and "post-boost-like" interferon response signatures could be observed among the COVID-19 patients, and whether these were associated with a more favorable course of the disease. We relied on blood transcriptome data from the two cohorts of COVID-19 patients described above, namely the "PREDICT-19 consortium Italian cohort" comprising 77 patients with a wide spectrum of disease severity and the "IMPROVISE Whole Blood Dataset" comprised exclusively of patients admitted to the ICU.

Focusing first on the PREDICT-19 Italian cohort, we used the response values for the six interferon modules from Aggregate A28 to map individual COVID-19 patient samples along with post-vaccine samples on the same t-SNE plot (**Figure 7**). First, we confirmed that there was no apparent separation of the vaccination and COVID-19 patient cohorts and that batch correction was therefore not warranted before proceeding with comparative analyses

(**Supplementary Figure 2**). This is consistent with the results of meta-analyses we have

previously conducted at the module level (13). To help with the interpretation, k-means

clustering was performed using the consolidated set of samples, resulting in the formation

of eight distinct clusters (**Figure 8**). Next, we examined the distribution of samples from the

vaccine and COVID-19 cohorts across the tSNE plot and among the eight clusters.

Timepoints at which an interferon response was detectable in vaccinated subjects were of

particular interest. Indeed, day 1 and day 2 post-prime samples (P1, P2), while preferentially

found in Clusters 1 and 5, appeared to be distributed across the entire t-SNE plot. This is in

contrast with day 1 and day 2 post-boost vaccination samples (B1, B2), which were almost

exclusively found in Cluster 5. A set of COVID-19 patients also co-localized in Cluster 5, while

others were found scattered across clusters, especially Clusters 1, 2, 6, and 3. Interferon

responses were detectable in all of these clusters, but with important nuances. For one,

samples from Cluster 5 showed by far the most potent responses, with responses seen in

most cases across all six interferon modules, which was consistent with the post-boost

vaccine response. In comparison, the response was less pronounced in samples from Cluster

1, which was dominated by modules associated with type I interferon responses (the

A28/S1 set comprising M10.1, M8.3 and M15.127 described above). This A28/S1-high and

A28/S2-low phenotype was more pronounced in the day 2 post-prime signature because it

corresponded to the peak of the A28/S1 response, which occurred after the peak of the

A28/S2 set, and indeed, day 2 post-prime samples were preferentially found in this cluster.

Signatures for samples forming Clusters 2 and 6 were not well-defined and were in some

cases absent, yet these clusters also included COVID-19 patients. Samples forming Cluster 3

displayed a peculiar signature, with an increase in the abundance of modules belonging to

the A28/S2 set (M15.64, M13.17, and M15.86) concomitantly with a decrease in modules

forming A28/S1. Among the samples forming this cluster, this pattern was most apparent

for the COVID-19 patients.

To summarize at this juncture, we found that COVID-19 patients can present with

distinct interferon response signatures. In subjects of Cluster 5, the pattern is consistent

with the response on day 1 to booster doses of COVID-19 mRNA vaccines, which in the

vaccination setting, is correlated with the development of a potent antibody response. In

subjects from Cluster 1, the response was instead dominated by modules belonging to the

A28/S1 set, which are responsive to type I interferons and most consistent with the

response on day 2 to the priming dose of the COVID-19 mRNA vaccine. Finally, in COVID-19

patients, a third pattern of response was observed that does not have an equivalent in

vaccinated individuals or in other pathologies that we have previously examined and was

characterized by an increase in A28/S2 modules concomitant with a decrease in A28/S2

modules. This latter interferon response pattern was observed in only a small number of

COVID-19 subjects, a high proportion of whom presented with a more severe form of the

disease.

It was difficult to determine whether absent or sub-optimal interferon responses

among PREDICT-19 subjects could be due to an impaired response to the infection or simply

a milder course of the disease. Thus, we next investigated the patterns of interferon

responses among ICU patients enrolled in the IMPROVISE cohort. Employing an approach

similar to the one described above, we mapped individual COVID-19 patient samples along

with post-vaccine samples on a t-SNE plot on the basis of similarities in the patterns of

interferon responsiveness across the six interferon modules (**Figure 8A**). COVID-19 subjects

were found to again be distributed throughout multiple clusters. Patients who co-localized

with day 1 post-boost vaccine samples tended to have relatively short ICU stays (in Cluster 5

= A28/S1$^+$,A28/S2$^+$), and only a few patients co-localized with day 2 post-prime samples in

Cluster 3, which was characterized by a more prominent A28/S1 signature compared with

A28/S2 (A28/S1$^+$,A28/S2$^{+/0}$). Contrastingly, distinct patient "hostspots" were found in

Clusters 1 and 6 that were characterized by the opposite response pattern, i.e., higher

abundance levels for A28/S2 compared with A28/S1 (Cluster 6 = A28/S1$^{0/-}$, A28/S2$^{+/0}$ Cluster

1 = A28/S1$^-$, A28/S2^0). Notably, these patient clusters tended to have longer ICU stays

(**Figure 8B**).

DISCUSSION

Relatively little is known about the types of in vivo immune responses elicited by mRNA

vaccines in humans. To address this, we employed bulk blood transcriptomics to map the

immune changes taking place in vivo after the administration of priming and booster doses

of COVID-19 vaccines in adult volunteers. We did so at a high temporal resolution, collecting

small amounts of blood before and 9 consecutive days after the administration of the

priming and booster doses of COVID-19 mRNA vaccines. The use of blood transcriptomics

eliminated the need to choose a panel of analytes to measure vaccine responses, which is

one source of bias. The daily collection and profiling schemes adopted eliminated the need

to choose specific timepoints for measuring the response, thus eliminating a second source

of bias.

Profiling blood transcript abundance post-prime and -booster doses of COVID-19

mRNA vaccines at a high temporal resolution revealed a well-orchestrated sequence of

immune events (**Figure 9**). The immune signatures elicited following the administration of

the two doses of mRNA vaccines differed drastically. Relatively modest changes were

observed post-prime that manifested primarily as the induction of interferon-response

signatures that were detectable over the first 3 days following the injection of the first dose.

This was followed by a more subtle response that could be attributed to the priming of the

adaptive response between days 4 and 6. Indeed, a decrease in the abundance of

transcripts for modules associated with inflammation was observed over these 3 days,

which comprised an increase in transcripts associated with plasma cells and T-cells on day 5.

No further changes were detected beyond day 6. After the booster dose, the plasmablast

response peaked on day 4. Notably, in studies assessing blood transcriptional responses to

vaccines, the peak plasmablast response is typically observed on day 7, as it is for instance

with influenza and pneumococcal vaccines (10–12). As a result, sampling schedules in

common use are designed to capture changes on day 1, 7, and sometimes day 3, but would

miss the peak of the adaptive response to COVID-19 mRNA vaccines observed in our study.

In addition to eliminating potential blind spots, high-frequency sampling and profiling also

permit the precise resolution of signatures that show the complex kinetics of a response; for

instance, the erythroid cell signature peaks sharply post-boost and recedes well below

baseline over several days before recovering. The trajectory of this signature may be of

significance in the context of vaccination, as we recently described its association with

immunosuppressive states, such as late-stage cancer and maintenance therapy in liver

transplant recipients (13). In the same work, we found this signature to be strongly

associated with the development of more a severe disease in subjects with acute

respiratory syncytial virus infection; and we furthermore putatively associated this signature

with populations of circulating erythroid cells found to possess immunosuppressive

properties (15).

Arunachalam et al. previously described the elicitation of qualitatively distinct innate

signatures on day 1 following the administration of priming and booster doses of COVID-19

mRNA vaccines, with the former inducing an interferon response and the latter a mixed

response that also presented an inflammatory component (3). Our findings are consistent

with these earlier observations, and employing a high-frequency sampling and profiling

protocol permitted us to further dissect those responses. Most notably, while interferon

responses appear *a priori* as the common denominator between the post-prime and post-

boost responses, the temporal pattern of response that we observed indicates that these

are, in fact, qualitatively distinct. This was evidenced by the differences in the timing of the

response peak, which corresponded to day 2 post-prime and day 1 post-boost. The kinetics

of the response post-boost being therefore most consistent with what is observed following

injection of a single dose of influenza vaccine (11). Interestingly, a further investigation of

the patterns of response among the six modular components of the interferon responses

(module Aggregate A28) identified two distinct sets of modules. The first one (A28/S1),

which appeared to be preferentially associated with type 1 interferon responses, peaked on

day 2 post-prime. The post-prime response of the second set of interferon modules was

more modest and peaked on day 1 instead of day 2. Both sets showed strong responses

post-boost, with both peaking on day 1, but again showing distinct kinetics with a more

sustained response being observed in the case of A28/S1. Association with antibody

responses measured on day 14 post-boost followed a similar pattern, with a preferential

association with A28/S1 observed post-prime on days 2 and 3, and an association with both

A28/S1 and A28/S2 post-boost on days 1 and 2. We have described, in our earlier report,

that distinct interferon modules could be employed to stratify patients with systemic lupus

erythematosus (17). Here, we sought to specifically determine whether COVID-19 patients

present with "post-prime-like" (i.e., A28/S1+) or "post-boost-like" (i.e., A28/S1+, A28/S2+)

patterns of interferon response, since those were associated with the subsequent

development of antibody responses. This was even more relevant in the context of COVID-

19 disease, as it has been reported that failure to induce interferon responses is associated

with worse disease outcomes (4,18–20). In the first patient cohort, which constituted

patients with predominantly mild or moderate pathology, both phenotypes were indeed

observed, along with a third "atypical" phenotype that was not observed post-vaccination

that is dominated instead by A28/S2 (with A28/S1 abundance low or even decreased).

Notably, in a second patient cohort made up exclusively of ICU patients, both A28/S1+

A28/S2+ (post-boost-like) and A28/S2+ (atypical) phenotypes were also observed, with the

latter being clearly associated with extended lengths of stay in the ICU. However, the

atypical phenotype did not appear to be preferentially associated with death in this setting,

which may be due to the supportive care provided to the patients. While, overall, our

observations support the notion that robust interferon responses are associated with a

more favorable course of the disease, they also show that the response may be of a

different type for other patients, rather than being altogether defective (i.e., A28/S2+). One

possibility is that this peculiar response pattern is associated with the presence of

endogenously produced autoantibodies that neutralize interferon, as has been previously

described (20,21). However, it is not possible to be conclusive on this point and further

investigations are warranted.

Other points remain to be elucidated. This includes the timing of the adaptive response

to mRNA vaccines, which appears to rise and peak several days earlier than is normally

observed in responses to other vaccines (± 7 day peak). Also remaining to be determined is

the priming mechanism underpinning the robust polyfunctional response observed on day 1

post-boost. One of the questions being whether or not such a response, which would

typically be considered to be innate, is antigen-specific. Interestingly, in that respect, the

subjects who were previously infected but recovered from COVID-19 did not display a

noticeable day 1 inflammatory response, and their immune systems behaved like those of

naïve individuals. However, the number of recovered subjects was small, and the study was

not designed to directly address this question, hence further investigations will also be

necessary. Notably, the greater amplitude of responses observed post-boost and the

presence of an inflammatory component is also consistent with previous reports of the

increase in the incidence of side effects/discomfort following COVID-19 mRNA vaccine

booster doses (22,23).

The work presented here could help inform the design of studies aiming to investigate

vaccine responses. Indeed, a decrease in sequencing costs provides laboratories an

opportunity to employ transcriptome profiling approaches in novel ways, such as the

implementation of high-temporal frequency protocols. An advantage of the delineation of

transcriptome responses at high-temporal resolution is that it is doubly unbiased, i.e., there

is no need to select transcripts for inclusion in a panel because RNA sequencing measures all

transcript species present in a sample. Similarly, there is no need to select specific

timepoints for assessing the vaccine response, as all timepoints were profiled within a given

time frame. An obvious advantage of the approach is that it permits the removal of

potential blind spots, and the detection of changes that are potentially otherwise missed by

more sparse sampling protocols. We have shown here that the peak plasmablast response

following COVID-19 mRNA vaccination occurs unexpectedly at around day 4 and wanes by

day 7, when such a response would normally be expected after an influenza vaccination.

This is significant, as this signature was also shown to correlate with the extent of the

antibody response. A somewhat less obvious advantage is that, as exemplified by the

interferon signatures, high-frequency profiling data permitted us to more precisely resolve

the vaccine response. For instance, in this case, we were able to show that the interferon

response peaked on day 2 post-prime but on day 1 post-boost. To be more precise, a

component of the post-prime interferon response peaked on day 2, while the other peaked

on day 1. Some of the practical elements that may contribute to making the routine

implementation of high-temporal resolution transcriptomics approaches viable include, as

mentioned earlier, a substantial decrease in the cost of RNA sequencing, especially lower

depth, 3'-biased methodologies. Recent publications showed, through down-sampling

analysis, that much fewer deep reads than usual are adequate for biomarker discovery

projects (24), with the lower costs permitting larger sample sizes or, as in this case, a higher

sampling frequency. Another consideration is the availability of solutions for the in-home

self-collection of samples. This is the case for the collection of RNA-stabilized blood,

although our homegrown method could be improved and interesting solutions are being

put forward (25). As we have shown, it is possible to implement the self-collection of

samples for serology profiling within a vaccinology study.

There were several limitations to our study. While the sample size was adequate for an

initial discovery phase, a larger study cohort would help to better resolve inter-individual

variations. The dataset we generated, however, has been made available for reuse, and it

should be possible to integrate and consolidate this dataset with those generated in follow-

on studies by us and others (13). Follow on studies would need to be purposely designed

to formally address specific questions, for instance, comparing responses in individuals who

had previously been exposed to SARS-CoV-2 with those in naïve individuals. It would also be

interesting to compare responses elicited by the Pfizer/BioNTech and Moderna vaccines,

which was not possible in our study due to the small numbers of individuals that received

the Moderna vaccine. Indeed, although we hoped it would be possible to obtain more

balanced sample sizes for a more detailed comparison, the speed at which the vaccinations

were rolled out among our target population of healthcare workers meant we had very little

control over the number of volunteers that received the different types of vaccines or their

status as naïve or previously exposed individuals. It would also have been particularly

interesting to enroll patients from different age categories, especially the elderly

population, but this again proved impossible.

In conclusion, a considerable number of COVID-19 vaccines have already been approved

for use in humans, and others are currently in phase III trials (>20) (26). The data presented

herein suggest that high-temporal-frequency blood transcriptomics are an appropriate way

to precisely map the types of responses elicited by the different COVID-19 vaccines and,

thus, benchmark a wide range of platforms that have been employed for vaccine

development. Characterizing and comparing vaccine response profiles in populations that

may not respond optimally to vaccines should also prove particularly informative (e.g., in

the elderly, immunosuppressed, and during pregnancy). Finally, the most likely roll-out of a

systematic third injection of the booster dose should offer opportunities to further dissect

the immune responses to mRNA vaccines in individuals who have been previously

immunized.

METHODS

Subject recruitment:

COVAX Cohort: We enrolled adult subjects eligible to receive a COVID-19 vaccine and who

were willing to adhere to the sampling schedule. The protocol was approved by Sidra

Hospital IRB (IRB number 1670047-6), and all participants gave written informed consent.

Inclusion criteria matched the clinical eligibility for receiving the vaccine, and the only

exclusion criterium was subjects that already had a first dose of any COVID-19 vaccine.

Twenty-three subjects were enrolled, and the median age was 38 years (29-57); 20 of the

subjects received the Pfizer vaccine and 3 the Moderna vaccine. The demographics, health

status at accrual, and vaccination side effects are shown in Table 1. Vaccination and booster

intervals were typically 21 days for Pfizer and 29 days for Moderna.

IMPROVISE cohort: Adult subjects with severe COVID-19 were enrolled to this cohort under

the Hamad Medical Corporation IRB approval (MRC-05-007). Blood samples were collected

at multiple timepoints during patients' ICU stay (timepoint 1 was taken at ICU admission;

timepoints 1 to 4 were 7 days apart). Subjects with burn and trauma, immunological

diseases, receiving immunosuppressive treatment, with other immune-related conditions,

or with a previous COVID-19 infection were excluded. For this analysis, 40 severe COVID-19

patients were included, with a median age of 52 (range = 30 to 92). The clinical parameters

of those patients included gender, ICU and hospital stay, mechanical ventilation duration,

ECMO initiation, comorbidities, outcomes (death/recovery), nosocomial infection onset, and

plasma therapy. Samples were also collected from control subjects who were adults and did

not: 1) present with an infectious syndrome during the last 90 days, 2) experience extreme

physical stress within the last week, 3) received during the last 90 days a treatment based

on: antivirals; antibiotics; antiparasitic; antifungals; 4) received within the last 15 days, a

treatment based on non- steroidal anti-inflammatory drugs; 5) received during the last 24

months a treatment based on: immunosuppressive therapy; corticosteroids; therapeutic

antibodies; chemotherapy and 6) a person with history of: innate or acquired immune

deficiency; hematological disease; solid tumor; severe chronic disease; surgery or

hospitalization within the last 2 years; pregnancy within the last year; participation to a

phase I clinical assay during the last year; participation to a phase I clinical assay during the

last year; pregnant or breastfeeding women; person with restricted liberty or under legal

protection.

PREDICT-19 Cohort:

The "Predicting disease progression in severe viral respiratory infections and COVID-19"

(PREDICT-19) Consortium is an international consortium formed by a group of researchers

who share common interests in identifying, developing and validating clinical and/or

bioinformatic tools to improve patient triage in a pandemic such as COVID-19 (14). The

PREDICT-19 Italian cohort comprises adult subjects with mild, moderate, or severe COVID-

19 diagnosed by real-time PCR on nasopharyngeal swab who were consented and enrolled

at E.O. Ospedali Galliera, and IRCCS Ospedale Policlinico San Martino, Genoa, Italy (Ethics

Committee of the Liguria Region (N.CER Liguria 114/2020-ID 10420). Blood samples were

collected during hospitalization. Subjects with burn and trauma, immunological diseases,

receiving immunosuppressive treatment for underlying disorders before COVID-19

diagnosis, with other immune-related conditions, or with a previous COVID-19 infection

were excluded. For this analysis, 10 healthy subjects and 103 COVID-19 patients were

included, with a median age of 61.76 (range = 26 to 86).

Sampling protocol:

COVAX Cohort: For transcriptomics applications for the COVAX study, after puncturing the

skin with a finger stick, 50 µl of blood was collected in a capillary/microfuge tube assembly

supplied by KABE Labortechnik (Numbrecht, Germany) containing 100 µl of tempus RNA-

stabilizing solution aliquoted from a regular-sized tempus tube (designed for the collection

of 3 ml of blood and containing 6 ml of solution; ThermoFisher, Waltham, MA, USA). This

method is described in detail in an earlier report (5), and the collection procedure is

illustrated in an uploaded video: https://www.youtube.com/watch?v=xnrXidwg83I. Blood

was collected prior to the vaccine being administered (day 0), on the same day, and daily

thereafter over the next 10 days. This protocol was followed for both the priming and

booster doses.

For serology applications, 20 µl of blood was collected using a Mitra blood collection device

(Neoteryx, Torrance, CA, USA) prior to the vaccine being administered and on days 7 and 14

after vaccination with the priming and booster doses.

<u>IMPROVISE Cohort</u>: For the IMPROVISE study, samples were collected using PaxGene Blood

RNA tubes (BD Biosciences, Franklin Lakes, NJ, USA) at all timepoints and were frozen at -

20C until further processing.

<u>PREDICT-19 Cohort</u>: For the Italian cohort of the PREDICT-19 study, blood samples were

collected during hospitalization by venipuncture during in tubes containing an RNA

stabilizing solution (Tempus™ Blood RNA Tube, ThermoFisher, Waltham, MA, USA, Catalog

number: 4342792) and frozen at -20C until further processing.

Multiplex serological assay

The concentration of selected proteins in the serum samples was measured with a home-

built bead array based on carboxymethylated beads with 6 distinct intensities of a UV-

excitable dye. Each was individually coupled to 3 SARS-CoV-2 proteins, envelope,

nucleoprotein, S-trimer and fragments thereof and the S1 fragment of SARS-CoV S protein.

Therefore, the complete array consisted of 6 antigens, including five SARS-CoV-2 antigens

(Full Spike Trimer, RBD, Spike S1, Nucleoprotein, and Envelope), as well as the closely

related SARS-CoV-S1 protein. The readout of this assay provided the immunoglobulin

isotypic response to each of the antigens, i.e., total IgM, total IgG, total IgA, as well as their

individual isotypes, IgG1, IgG2, IgG3, IgA1, and IgA2, containing a total of 42 analytes. The

assays were performed on filter plates and acquired on a BD-Symphony A5. Ig-binding to

the spike protein antigens differed for each bead region. An average of 300 beads per

region was acquired, and the median fluorescence intensity for isotype binding was used for

the calculations. An antibody response index was calculated by ratioing form each antigen

and isotype, the MFI of polled negative blood controls collected prior to June 2018 (Sidra

IRB 1609004823) to the MFI obtained for vaccinated donor samples.

RNA extraction and QC

RNA was extracted using the Tempus Spin RNA Isolation Kit (ThermoFisher), which was

adapted for the handling of small blood volumes. The methodology has been described

previously in detail (27). Contaminating DNA was removed using the TurboDNAse kit

(ThermoFisher), and RNA was quantitated on a Qubit instrument (ThermoFisher) and QCed

using an Agilent 2100 Bioanalyzer (Agilent, Santa Clara, California, USA).

RNA sequencing

<u>COVAX & IMPROVISE Cohorts:</u> mRNA-sequencing was performed using QuantSeq 3' mRNA-

Seq Library Prep Kit FWD for Illumina (75 single-end) with a read depth of 8M and average

read alignment of 79.60%. Single samples were sequenced across four lanes, and the

resulting FASTQ files were merged by sample. Quality trimming is performed to remove

adapter sequences and polyA tails. Then trimmed reads were aligned to human genome

GRCh38/hg38 (Genome Reference Consortium Human Build 38), INSDC Assembly

GCA_000001405.28, Dec 2013) using STAR 2.6.1d and featureCounts v2.0.0 was used to

generate the raw counts. Raw expression data were normalized to size factor effects using R

package DESeq2. All downstream analyses were performed using R version 4.1, unless

otherwise specified. Global transcriptional differences between samples were assessed by

principal component analysis using the "prcomp" function. Transcriptome profiling data

were deposited, along with detailed sample information, into a public repository, the NCBI

Gene Expression Ominibus (GEO), with accession ID GSE190001 and BioProject ID:

PRJNA785113

PREDICT-19 Cohort: Total RNA was isolated from whole blood lysate using the Tempus Spin

Isolation kit (Applied Biosystems) according to the manufacturer's instructions. Globin

mRNA was depleted from a portion of each total RNA sample using the GLOBINclear™-

Human kit (Thermo fisher). Following removal of globin transcripts transcriptome profiles

were generated via mRNA sequencing. Then mRNA-sequencing was performed using

Illumina HiSeq 4000 Technology (75 paired-end) with a read depth of 60M. Single samples

were sequenced across four lanes, resulting FASTQ files were merged by sample. All FASTQ

passed QC and were aligned to reference genome GRCh38 using STAR (2.6.1d). BAM files

were converted to a raw count's expression matrix using HTSeq

(https://github.com/Sydney-Informatics-Hub/RNASeq-DE). Raw count data was normalized

using DEseq2. The ensemble ID targeting multiple genes were collapsed (average) and a

final data matrix gene was for modular repertoire analysis.

Statistical Analysis

Analyses were conducted using pre-defined gene sets. Specifically, we employed a fixed repertoire of 382 transcriptional modules that were thoroughly functionally annotated, as described in detail in a recent publication (7). Briefly, this repertoire of transcriptional modules ("BloodGen3") was identified based on co-expression, as measured in a collection of 16 blood transcriptome datasets encompassing 985 individual transcriptome profiles. Sets of co-expressed transcripts were derived from the analysis of a large weighted co-clustering network. Downstream analysis results and visualizations were generated employing a custom R package (28). "Module response" is defined as the percentage of constitutive transcripts with a given abundance that were determined to be different between two study groups, or for the same individual in comparison to a given baseline (in this study, pre-vaccination abundance levels). The values therefore ranged from 100% (all constitutive transcripts increased) to −100% (all constitutive transcripts decreased). Only the dominant trend (i.e., increase or decrease in abundance over control/baseline) was retained for visualization purposes on fingerprint grids or fingerprint heatmaps, with red indicating an increase and blue a decrease in abundance. When performing group comparisons (e.g., cases vs controls for the disease datasets used as reference), the p-value and false discovery rate cutoffs were applied, which are mentioned in the figure legend. When performing longitudinal analyses, the module response is determined by employing fixed fold-change and expression difference cutoffs. Module response values obtained were used for data visualization. Significance was determined for each module using the differential gene set enrichment function of the dearseq R package (8).

ACKNOWLEDGMENTS:

We wish to acknowledge Michelle Esblaca and Juvilyn Gusi for their assistance with the

training of subjects and collection of blood samples, as well as the research subjects for

volunteering and making the study possible.

REFERENCES

1. Tregoning JS, Flight KE, Higham SL, Wang Z, Pierce BF. Progress of the COVID-19 vaccine
effort: viruses, vaccines and variants versus efficacy, effectiveness and escape. Nat Rev
Immunol. 2021 Aug 9;

2. Hagan T, Nakaya HI, Subramaniam S, Pulendran B. Systems vaccinology: Enabling
rational vaccine design with systems biological approaches. Vaccine. 2015 Sep
29;33(40):5294–301.

3. Arunachalam PS, Scott MKD, Hagan T, Li C, Feng Y, Wimmers F, et al. Systems
vaccinology of the BNT162b2 mRNA vaccine in humans. Nature. 2021 Jul 12;

4. Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired type I
interferon activity and inflammatory responses in severe COVID-19 patients. Science.
2020 Aug 7;369(6504):718–24.

5. Rinchai D, Anguiano E, Nguyen P, Chaussabel D. Finger stick blood collection for gene
expression profiling and storage of tempus blood RNA tubes. F1000Res. 2016;5:1385.

6. De Mot L, Bechtold V, Bol V, Callegaro A, Coccia M, Essaghir A, et al. Transcriptional
profiles of adjuvanted hepatitis B vaccines display variable interindividual homogeneity
but a shared core signature. Sci Transl Med. 2020 Nov 11;12(569):eaay8618.

7. Altman MC, Rinchai D, Baldwin N, Toufiq M, Whalen E, Garand M, et al. Development of
a fixed module repertoire for the analysis and interpretation of blood transcriptome
data. Nat Commun. 2021 Jul 19;12(1):4385.

8. Agniel D, Hejblum BP. Variance component score test for time-course gene set analysis
of longitudinal RNA-seq data. Biostatistics. 2017 Oct 1;18(4):589–604.

9. Rawat A, Rinchai D, Toufiq M, Marr AK, Kino T, Garand M, et al. A Neutrophil-Driven
Inflammatory Signature Characterizes the Blood Transcriptome Fingerprint of Psoriasis.
Front Immunol [Internet]. 2020 [cited 2020 Dec 7];11. Available from:
https://www.frontiersin.org/articles/10.3389/fimmu.2020.587946/full

10. Nakaya HI, Wrammert J, Lee EK, Racioppi L, Marie-Kunze S, Haining WN, et al. Systems
biology of vaccination for seasonal influenza in humans. Nat Immunol. 2011 Jul
10;12(8):786–95.

11. Obermoser G, Presnell S, Domico K, Xu H, Wang Y, Anguiano E, et al. Systems scale
interactive exploration reveals quantitative and qualitative differences in response to
influenza and pneumococcal vaccines. Immunity. 2013 Apr 18;38(4):831–44.

12. Nakaya HI, Hagan T, Duraisingham SS, Lee EK, Kwissa M, Rouphael N, et al. Systems
Analysis of Immunity to Influenza Vaccination across Multiple Years and in Diverse
Populations Reveals Shared Molecular Signatures. Immunity. 2015 Dec 15;43(6):1186–
98.

13. Rinchai D, Altman MC, Konza O, Hässler S, Martina F, Toufiq M, et al. Definition of
erythroid cell-positive blood transcriptome phenotypes associated with severe
respiratory syncytial virus infection. Clinical and Translational Medicine.
2020;10(8):e244.

14. Tang B, Shojaei M, Wang Y, Nalos M, Mclean A, Afrasiabi A, et al. Prospective validation
study of prognostic biomarkers to predict adverse outcomes in patients with COVID-
19: a study protocol. BMJ Open. 2021 Jan 6;11(1):e044497.

15. Elahi S. Neglected Cells: Immunomodulatory Roles of CD71+ Erythroid Cells. Trends
Immunol. 2019 Mar;40(3):181–5.

16. Elahi S, Ertelt JM, Kinder JM, Jiang TT, Zhang X, Xin L, et al. Immunosuppressive CD71+
erythroid cells compromise neonatal host defence against infection. Nature. 2013 Dec
5;504(7478):158–62.

17. Chiche L, Jourde-Chiche N, Whalen E, Presnell S, Gersuk V, Dang K, et al. Modular
transcriptional repertoire analyses of adults with systemic lupus erythematosus reveal
distinct type I and type II interferon signatures. Arthritis & Rheumatology (Hoboken,
NJ). 2014 Jun;66(6):1583–95.

18. Galani I-E, Rovina N, Lampropoulou V, Triantafyllia V, Manioudaki M, Pavlos E, et al.
Untuned antiviral immunity in COVID-19 revealed by temporal type I/III interferon
patterns and flu comparison. Nat Immunol. 2021 Jan;22(1):32–40.

19. Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, et al. Inborn errors of type
I IFN immunity in patients with life-threatening COVID-19. Science. 2020 Oct
23;370(6515):eabd4570.

20. Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann H-H, Zhang Y, et al.
Autoantibodies against type I IFNs in patients with life-threatening COVID-19. Science.
2020 Oct 23;370(6515):eabd4585.

21. Bastard P, Gervais A, Le Voyer T, Rosain J, Philippot Q, Manry J, et al. Autoantibodies
neutralizing type I IFNs are present in ~4% of uninfected individuals over 70 years old
and account for ~20% of COVID-19 deaths. Sci Immunol. 2021 Aug 19;6(62):eabl4340.

22. Cheng H, Peng Z, Luo W, Si S, Mo M, Zhou H, et al. Efficacy and Safety of COVID-19
Vaccines in Phase III Trials: A Meta-Analysis. Vaccines. 2021 Jun;9(6):582.

23. Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Phase I/II study
of COVID-19 RNA vaccine BNT162b1 in adults. Nature. 2020 Oct;586(7830):589–93.

24. Milanez-Almeida P, Martins AJ, Germain RN, Tsang JS. Cancer prognosis with shallow
tumor RNA sequencing. Nat Med. 2020 Feb;26(2):188–92.

25. Haack AJ, Lim FY, Kennedy DS, Day JH, Adams KN, Lee JJ, et al. homeRNA: A self-
sampling kit for the collection of peripheral blood and stabilization of RNA [Internet].
2021 Feb [cited 2021 Oct 30] p. 2021.02.08.430337. Available from:
https://www.biorxiv.org/content/10.1101/2021.02.08.430337v1

26. Sadarangani M, Marchant A, Kollmann TR. Immunological mechanisms of vaccine-
induced protection against COVID-19 in humans. Nat Rev Immunol. 2021
Aug;21(8):475–84.

27. Syed Ahamed Kabeer B, Tomei S, Mattei V, Brummaier T, McGready R, Nosten F, et al. A
protocol for extraction of total RNA from finger stick whole blood samples preserved
with TempusTM solution. F1000Res. 2018 Nov 2;7:1739.

28. Rinchai D, Roelands J, Toufiq M, Hendrickx W, Altman MC, Bedognetti D, et al.
BloodGen3Module: Blood transcriptional module repertoire analysis and visualization
using R. Bioinformatics. 2021 Feb 24;

863   **Table 1: Subject characteristics**

Patient ID	Vaccine name	Gender	Age	Ethnicity	Previous COVID-19	Underlying disease	Drugs	Symptoms at prime (Type)	Symptoms at prime (Grade)	Symptoms at boost (Type)	Symptoms at boost (Grade)
PZB1	Pfizer Biontech	Female	38	Asian	Yes	No	no	Myalgia	G1	Fever/Myalgia	G1
PZB2	Pfizer Biontech	Male	47	Caucasian	Yes	No	no	Myalgia	G1	Myalgia	G1
PZB3	Pfizer Biontech	Male	57	Caucasian	Yes	T2D	Metformin, Insulin	Myalgia	G1	Myalgia	G1
PZB4	Pfizer Biontech	Female	34	Indian	No	No	no	None	NA	Chills/Insomnia/Headache/Myalgia/Fa	G3

tigue

ID	Vaccine	Sex	Age	Ethnicity	Previous COVID	Underlying disease	Drugs	Symptoms (dose 1)	Grade	Symptoms (dose 2)	Grade
PZB5	Pfizer Biontech	Male	38	Indian	No	No	no	Myalgia	G1	Myalgia	G1
PZB6	Pfizer Biontech	Female	48	Caucasian	No	No	no	Myalgia/Headache	G1	Myalgia	G1
PZB7	Pfizer Biontech	Female	34	Caucasia/Arab	No	Hashimoto thyroiditis	no	None	NA	None	NA
PZB8	Pfizer Biontech	Female	29	Arab	No	No	no	Myalgia/Swelling	G1	Fever/Myalgia	G1
PZB9	Pfizer Biontech	Male	41	Arab	No	Allergic rhinitis	no	Myalgia	G1	Myalgia	G1
PZB10	Pfizer Biontech	Female	35	Arab	No	No	no	Myalgia	G1	Fever/Insomnia/Myalgia/Fatigue	G2
PZB11	Pfizer Biontech	Male	41	Caucasian	No	No	no	Myalgia	G2	Myalgia	G1
PZB12	Pfizer Biontech	Female	34	Indian	No	Hypothyroidism	Levothyroxine	None	NA	Fever/Myalgia	G1
PZB13	Pfizer Biontech	Male	29	Indian	No	No	no	Fatigue	G1	Fever/Heaviness in arm	G2
PZB14	Pfizer Biontech	Female	38	Arab	No	Allergic	no	Myalgia/Headache	G1	Fatigue	G1
PZB15	Pfizer Biontech	Male	43	Arab	No	Hypothyroidism	Levothyroxine	None	NA	Myalgia/Headache	G2
PZB16	Pfizer Biontech	Female	39	Indian	No	No	no	Heaviness in arm	G1	Fever/Myalgia/Fatigue	G2
PZB17	Pfizer Biontech	Female	42	Indian	Yes	T2D, Hypertension	Methformin, Telmisartan	Fever/Headache/Myalgia/Fatigue	G2	Fatigue/Gastritis	G2
MDA18	Moderna	Male	42	Caucasian	No	Hypertension	Amlodipine, Ramipril	Myalgia	G1	Myalgia	G1
PZB19	Pfizer Biontech	Female	39	Caucasian	No	No	no	Myalgia	G1	Headache/Myalgia/Arthralgia	G3
MDA20	Moderna	Female	36	Arab	Yes	No	no	Chills/Myalgia	G2	Chills/Myalgia	G2
MDA21	Moderna	Female	36	Caucasian	No	No	no	Myalgia	G1	Feveer/Skin rash/Myalgia	G2
PZB25	Pfizer Biontech	Female	39	Caucasian	No	No	no	Myalgia	G1	Myalgia	G1
MDA26	Moderna	Female	30	Caucasian	Yes	Asthma	Seretide, Salbutamol	Fever/Headache/Myalgia/Fatigue	G3	Asthma attack/Fever/Myalgia/Fatigue	G3

Footnotes:

T2D: Type 2 Diabetes

Vaccine type and lot, and subjects' characteristics were recorded, including demographic, biometric data, blood group, underlying diseases, drugs usage, and previous COVID-19 disease.

Every subject recorded and graded the symptoms that occurred after the first and second vaccinations doses, according to the NIH "DAIDS AE Grading Table"

Figure 1: Antibody response to COVID-19 mRNA vaccination. (A) Schematic representation of the study design. (B) The heatmap represents changes in abundance of antibodies specific to several SARS-CoV-2 antigens and control antigens relative to pre-vaccination levels. Red indicates a relative increase and green indicates a relative decrease in abundance. Columns represent subjects, which are arranged by timepoint and have colored tracks at the top indicating whether the subjects were previously infected with SARS-CoV-2. The histogram above represents the average log2 fold-change over baseline for a given column. The rows represent antibodies for which abundances were profiled, which are arranged according to specificity. For each antigen, the different rows represent different types of antibodies. (C) Changes in antibody levels, expressed as an "antibody index", are shown on the box plots, each corresponding to a given antibody type of a given specificity. Lines indicate changes for individuals previously infected with SARS-CoV-2 and who had recovered (in pink) and for naïve individuals (in green). Multiple pairwise tests (paired t-test) were performed comparing antibodies levels to baseline (D0). Asterisks: * represent p < .01, **represent p < .001, *** represent p < .0001.

Figure 2: Characterization of the post-prime interferon response signature. (A) The bar graph shows the cumulative module response at the various timepoints following the administration of the priming dose of the vaccine (noted P1-P14). The Y-axis values and numbers on the bars indicate the number of modules meeting the 15% response threshold (out of a total of 382 modules constituting the BloodGen3 repertoire, with percentage response corresponding to the proportion of transcripts predominantly increased or decreased compared to baseline using FDR < 0.1 as the cutoff to determine significance [DESeq2]). The number of modules for which abundance was predominantly increased is shown in red and for those for which abundance was predominantly decreased is shown in blue. The fingerprint grid plots represent the overall module responses on day 1 post-prime (P1) and day 2 post-prime (P2). Modules from the BloodGen3 repertoire occupy fixed positions on the fingerprint grids. They are arranged as rows based on membership to module aggregates (rows A1 through A38). Changes compared to pre-vaccination baseline are indicated on the grid by red and blue spots of varying color intensity, which indicate the

"percentage response" for a given module. The color key at the top indicates the various functions attributed to the modules that are represented on the grid. The response of the six modules comprising aggregate A28 is represented on a line graph that shows the proportion of responsive transcripts for each module across all the post-prime timepoints. For each of module statistical significance for the overall response was determined by time course gene set enrichment analysis. Four of the six A28 modules met significance thresholds FDR < 0.1 (M8.3: p-value = 1.9-e4, FDR = 0.019, M10.1: p-value = 1.9-e4, FDR = 0.019, M15.127: p-value = 1.9-e4, FDR = 0.019, 727 and M15.86: p-value = 3.9-e4, FDR = 0.031) and all six A28 modules p < 0.05 (M13.17: p-value = 1.5-e3, FDR = 0.101 and M15.64: p-value = 0.044, FDR = 0.727). We also ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). (B) Heatmaps represent proportions of transcripts that changed within the six A28 modules at different timepoints and across different individuals compared to pre-vaccination baseline values. Red indicates that transcripts were predominantly increased over the baseline, and blue indicates that transcripts were predominantly decreased. Rows represent the six A28 modules, which are arranged within an aggregate via hierarchical clustering. Columns represent samples, which are grouped by timepoint and show profiles of individual subjects within each timepoint. (C) The heatmaps represent associations (spearman correlation test) between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across 9 consecutive days after the priming dose. The heatmap at the top provides the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the six A28 interferon modules. The heatmap below shows the significance of the correlations shown on the heatmap directly above, with the same ordering of rows and columns.

Figure 3: Characterization of responses on day 5 post-prime. (A) The bar graph shows the cumulative module response at the various timepoints following the administration of the priming dose of the vaccine (noted P1-P14). The fingerprint grid plot shows changes elicited by vaccination on day 5 following administration of the priming dose. The position of the modules on the grid is fixed. The percent response of individual modules is represented on the grid by red and blue spots of varying color intensity denoting predominant increase or decrease in abundance, respectively. The percentage response of a given module corresponds to the proportion of transcripts predominantly increased or decreased compared to baseline, meeting a significance cutoff of FDR < 0.1. The color key at the top indicates the various functions attributed to the modules that are represented on the grid. **(B)** The heatmap represents Log2 average fold change in abundance of transcripts constituting sets of modules associated with given functional annotations on day 5 post-vaccination. Rows represent individual transcripts grouped according to the module aggregate they originate from, corresponding to the different rows on the fingerprint grid plot on the left. Each module aggregate is associated with a different function, as indicated by the color key above. The columns on the heatmap represent individual subjects (Pfizer BioNtech = PZB; Moderna = MDA).

Figure 4: Fingerprint grid plots mapping changes observed on day 1 post-boost and across reference datasets.
The bar graphs show the cumulative module response at the various timepoints post-priming and booster doses (noted P1-P14 and B1-B14, respectively). The Y-axis values and numbers on the bars indicate the number of modules meeting the 15% response threshold (out of a total of 382 modules constituting the BloodGen3 repertoire, with percentage response corresponding to the proportion of transcripts predominantly increased or decreased compared to baseline meeting a significance cutoff of DESeq2, FDR < 0.1. The fingerprint grid plots show changes in transcript abundance in a given study group in comparison to baseline (pre-vaccination sample or uninfected control group – with the percent response of individual modules shown by red and blue spots of varying color intensity denoting predominant increase or decrease in abundance, respectively. Changes are shown in the top grid for a group comparison of 1 day after receiving the booster dose of COVID-19 mRNA vaccines with baseline pre-vaccination samples (this study). Grids in the middle and bottom positions show changes for patients with acute infections caused by influenza virus (public dataset) or SARS-CoV-2 (this study) and for patients with bacterial sepsis (public dataset). The color key at the top indicates the various functions attributed to the modules that occupy a fixed position on the grid plot.

Figure 5: Characterization of the day 1 post-boost interferon response signature. (A) The fingerprint grid plot maps the modular response observed on day 1 post-boost (percent response is determined based on statistical cutoff: DESeq2, FRD < 0.1). The six modules forming the A28 aggregate are highlighted. The line graphs below represent the summarized percentage responses at the module level, encompassing all study subjects (one line per module). Percentage response accounts for the proportion of transcripts for a given module that show differences in abundance post-prime (left) or post-boost (right) compared to baseline pre-vaccination levels. Changes in transcript abundance post-prime and post-boost for two distinct sets of interferon response modules that received the denomination A28/S1 and A28/S2 are plotted on separate graphs. For each of module statistical significance for the overall response was determined by time course gene set enrichment analysis. Significance was reported post-prime in Figure 2. Post-boost all six A28 modules met significance thresholds p < 0.001 and FDR < 0.001 (M8.3: p-

value = 1.9-e4, FDR = 3.6-e4, M10.1: p-value = 1.9-e4, FDR = 3.6-e4, M13.17: p-value = 1.9-e4, FDR = 3.6-e4, M15.127: p-value = 1.9-e4, FDR = 3.6-e4, M15.64: p-value = 1.9-e4, FDR = 3.6-e4 and M15.86: p-value = 1.9-e4, FDR = 3.6-e4). In addition, we ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). The circle packing plots on the left show module responses at the individual transcript level for two public blood transcriptome datasets. The larger circle below indicates official symbols for the individual transcripts. It also highlights the modules included in A28/S2, which are shown directly on the right. The smaller circles above show changes in abundance of A28 transcripts for two public datasets. One study (GSE11342) measured blood transcriptional response in patients with Hepatitis C infection treated with alpha-interferon (23). The second study (GSE26104) measured transcriptional response in subjects with multiple sclerosis treated with beta-interferon (24). A red circle indicates a significant increase in the abundance of transcripts compared to the pre-treatment baseline (*|fold-change| > 1.5, FDR < 0.1). **(B)** Changes in abundance compared to baseline pre-vaccination levels are represented on a heatmap, with modules as rows and individual samples as columns. The modules are arranged by hierarchical clustering based on abundance patterns across samples. The samples are arranged by timepoints post-prime (top) and post-boost (bottom). **(C)** The heatmaps represent associations (Spearman correlation) between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across 9 consecutive days after the booster dose. The heatmap on top provides the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the six A28 interferon modules. The heatmap below shows the significance of the correlations shown on the heatmap on top, with the same ordering of rows and columns.

Figure 6: Characterization of post-boost inflammation, erythroid cell, and plasmablast responses. (A) The bar graph at the top represents the number of response modules at any given time point post-prime and post-boost. The fingerprint grid plots show the modules that had changes compared with a fixed visualization and interpretation framework. Changes are shown for the day 1 post-boost timepoint (left) as well as day 5 (right) (percent response is determined based on statistical cutoff: DESeq2, FRD < 0.1). On the left grid, modules belonging to aggregates A35 (associated with inflammation) and A37 (associated with erythroid cells) are highlighted. The profiles of those modules are represented on the line graphs below, which show the average percentage responses of A35 and A37 modules across multiple timepoints. The percentage response for a given module is the proportion of its constitutive transcripts showing significant changes, which ranged from 0% to 100% when transcripts were predominantly

increased to 0% to −100% when transcripts were predominantly decreased. Each line represents the profile of the modules constituting a given aggregate. Line graphs for A35 were split into three sets according to differences in clustering patterns (A35/S1, A35/S2 and A35/S3). On the right grid, modules belonging to aggregates A27 (associated with platelets) are highlighted. The corresponding line graph below represents the changes in abundance of A27 modules over time following administration of the second dose of vaccine. For each module statistical significance for the overall response was determined by time course gene set enrichment analysis using the dearseq R package. For A35, 20 of 21 modules met significance thresholds (p-value < 0.05 and FDR < 0.01). It was also the case in 11 of 11 modules for A37 and 4 of 5 modules for A27 (**Supplementary file 4**). In addition, we ascertained the significance of changes measured post-prime at the level of this module aggregate and at each time point (paired t-test comparing module response at each time point relative to the pre-vaccination baseline; * p<0.01, ** p<0.001, *** p<0.0001). **(B).** The heatmaps represent associations between antibody responses measured 14 days after administration of COVID-19 booster doses and transcriptional responses measured across 9 consecutive days after the booster dose. Specifically, the heatmap at the top represent the correlation coefficients across multiple days and for each day across multiple subjects, with rows corresponding to the five A27 plasmablast modules. The heatmap below shows the significance of the correlations shown on the heatmap at the top, with the same order of rows and columns.

Figure 7: Comparing patterns of interferon response in vaccinated individual and a cohort of COVID-19 patients. Similarities in patterns of interferon response induction across the six modules forming aggregate A28 and among our vaccination cohort and one of our COVID-19 disease cohorts (PREDICT-19 / Italy) are represented on a tSNE plot. COVID-19 samples are shown in red along with specific post-vaccination timepoints (post-prime days 1 and 2 [P1, P2], post-boost days 1 and 2 [B1, B2]). Samples from the consolidated cohorts were partitioned into 8 clusters via k-means clustering, the distribution of which is shown on the tSNE plot on the top right. Patterns of response for the 6 interferon response modules across the 8 sample clusters are shown on a heatmap. The red colors indicate that abundance of transcripts for a given module is predominantly increased with the intensity representing the proportion of constitutive transcripts meeting a given threshold, which at the level of individual samples is a fixed fold change and difference cutoff (|Fold change| > 1.5, and |difference| > 10 in a given sample over its respective pre-vaccination baseline). The blue color denotes a predominant decrease in abundance of constitutive transcripts compared to the same individual's pre-vaccination baseline. Details are shown below for Clusters 3, 5, and 8 in separate heatmaps.

Figure 8: Comparison of interferon response patterns of vaccinated individuals and a cohort of COVID-19 patients with severe disease under intensive care. Similarities in patterns of interferon response induction across the six modules forming aggregate A28 and among our vaccination cohort and one of our COVID-19 disease cohorts (IMPROVISE) are represented in a tSNE plot, onto which disease severity information has been overlaid, along with specific post-vaccination timepoints (post-prime days 1 and 2 [P1, P2], post-boost days 1 and 2 [B1, B2]). Samples from the consolidated cohorts were partitioned into 8 clusters via k-means clustering, the distribution of which is shown on the tSNE plot on the top right. Patterns of response for the 6 interferon response modules across the 8 sample clusters are shown on a heatmap. The red colors indicate that abundance of transcripts for a given module are predominantly increased with the intensity representing the proportion of constitutive transcripts meeting a given threshold, which at the level of individual samples is a fixed fold change and difference cutoff (|Fold change |> 1.5, and |difference| > 10 in a given sample over its respective pre-vaccination baseline). The blue color denotes a predominant decrease in abundance of constitutive transcripts compared to the same individual's pre-vaccination baseline. The heatmap at the bottom shows patterns of interferon responses for COVID-19 patients with severe disease upon admission to the ICU. Multiple clinical parameters are shown on the tracks above. The histogram represents length of stay in the hospital, in the ICU, and time under mechanical ventilation.

Figure 9: Summary. This diagrammatic representation summarizes the temporal trajectories of blood transcriptional signatures elicited in response to the first and second doses of mRNA vaccines.