A core salivary microbiome shows the high prevalence of bacterial members yet

variability across human populations

Xinwei Ruan [1], Jiaqiang Luo[1], Pangzhen Zhang[1] and Kate Howell[1, *]

[1]School of Agriculture and Food, Faculty of Veterinary and Agricultural Sciences, University of

Melbourne, Parkville 3010, Australia

Xinwei Ruan: ORCID: 0000-0002-1963-3807

Jiaqiang Luo: ORCID: 0000-0001-6459-3309

Pangzhen Zhang: ORCID: 0000-0002-9794-2269

* Corresponding author. khowell@unimelb.edu.au; ORCID 0000-0001-6498-0472

Abstract

Background

Human saliva contains diverse bacterial communities, reflecting human health status, dietary patterns

and contributing to variability in the sensory perception of food. Many descriptions of salivary

microbiome diversity compare commonalities and differences with reference to a diseased state, but

the composition of healthy saliva has not been described.

Results

Here, we use a meta-analysis approach to define and explore the core membership of the human

salivary microbial community by collecting and re-analysing raw 16S rRNA amplicon sequencing

data from 47 studies with 2206 saliva samples. We found 68 core bacterial taxa that were consistently

detected. Differences induced by various host intrinsic and behaviour factors, including gender, age,

geographic location, tobacco usage, and alcohol consumption, were evident. The core of the salivary

microbiome was verified by collecting and analysing saliva in an independent study.

Conclusion

These results suggest that the methods used can effectively define a core microbial community in

human saliva with high prevalence. The core salivary microbiome demonstrated both stability and

variability among populations. Geographic location was identified as the host factor with the largest

effect in shaping salivary microbiota. The independent analysis underlined that the impact of

geographic variation is likely due to diet.

Background

Human saliva plays an essential role in influencing the sensory perception of foods and beverages and

driving the purchase decisions of consumers. When food is taken into the mouth, mixing and

mastication allow a semi-solid bolus to be formed, and at the same time, aroma and flavour to be

released from the food [1]. The variation in perceived responses can be attributed to the inter-

individual variability in salivary composition, especially salivary microorganisms. As a complex

ecosystem, the human oral cavity hosts thousands of bacterial taxa, accompanied by interactions with

other microorganisms [2]. It is an ecological system that contains many distinct sub-niches, including

saliva, dental plaques, gingival sulcus, epithelial cells on the cheek, tongue, and teeth [3]. High

heterogeneity has been reported between the composition of microbial communities that colonise on

different sites [4]. The saliva is recognised as a reservoir with microorganisms from all ecological

niches in the human mouth with long-term stability [5]. The ensemble of microorganisms and the

expressed genetic material in human saliva is known as the "salivary microbiome".

The contribution of salivary microbiome to sensory perception of foods has been described by various

studies [6-8]. However, the diverse conclusions suggest that the role of the salivary microbiome may

be confounded by inter-individual variance. Meanwhile, comparing results from different studies

introduces significant technical and bioinformatic biases [9] especially when studies have targeted

different 16S rRNA hypervariable regions for amplification [10]. Characterising the microbial

communities commonly found in most human saliva regardless of the study-specific variation could

help establish the connection between salivary composition and food preference. On this basis, the

shifts from a common salivary microbiome by diseases or host lifestyle factors will also be more

prominent. A meta-analysis can summarise the existing knowledge and identify the commonalities

and differences in salivary microbiota between people from various backgrounds.

The core oral microbiome of a healthy human has been tried to be defined for more than ten years

[11]. The core microbiome is described as the common group of microbes that are important for host

biological function [12]. Defining the core only depends on occupancy frequency does not necessarily

reflect the underlying host-microbes functional relationship. However, it provides a foundation for

prioritising members adapted to the host environment [13]. A variety of studies have been devoted to

discovering the changes in human salivary microbiota based on different conditions. The variability in

the microbial profile of human saliva has not only been associated with oral diseases [14-16] but also

various chronic diseases that do not occur in the oral cavity, such as diabetes [17] pancreatic cancer

[18] and Polycystic ovary syndrome [19]. Although the shift in human salivary composition caused by

diseases has been studied for decades, our understanding of the impact of host intrinsic and behaviour

factors is still limited.

Many host characteristics have been shown to have an impact on the composition of the salivary

microorganisms, including age [20], diet [21, 22], ethnicity [23], gender [24], smoking [25], alcohol

use [26], circadian rhythm [27], body mass index [28], and the type of stimulation [29]. Some studies

have correlated the diverse microbiome with the distinct sensory responses between consumer groups

[30, 31] It has also been reported that people from different countries are colonised with distinct

salivary bacterial communities [32]. Li et al. analysed the human oral microbiome from Africa,

Alaska, and Germany and reported differences between the human groups living in various climate

conditions [33]. However, no study to date has demonstrated a clear global pattern in salivary

microbial composition.

In this paper, we collected raw 16S rRNA sequences of human salivary microbiota from 47 publicly

available datasets spanning 15 different countries. These raw data were systematically re-analysed and

pooled together to define a core salivary microbiome. We classified all sequences into operational

taxonomic units (OTUs) at 97% identity against the Human Oral Microbiome Database (HOMD) to

minimise the technical variation induced by comparing data from different hypervariable regions. It

allowed us to make a comparison between studies and reduced the redundancy in the dataset for

defining the "core". Using the metadata acquired with raw data, we also investigated the influences of

several host factors and technical factors on human salivary microbiota. Factors that showed a

potentially strong impact on shaping microbial communities in saliva were selected, and the taxa as

potential biomarkers were identified, and linked with functional predictions. Finally, saliva samples

were collected from independent, healthy individuals and analysed for microbial composition to

confirm the results found from the global dataset. These data confirmed the composition of the core

microbiome members, but the verification of geographic origin was not possible. Our study

contributes to fundamental understandings of the stable and differential salivary microbiome across

healthy adult populations. We have identified bacteria linked to particular identities of participants

and points to salivary microbiome composition being linked to diet, rather than ethnic origin.

Methods

1.   Literature search and data collection

To acquire sufficient data from healthy human saliva, available public studies related to human

salivary microbiota were systematically reviewed. A literature search was performed using the

combination of relative terms in EMBASE, MEDLINE and Web of Science for the studies published

before November 2020 using the terms described in supplementary data (Table S1). A supplementary

dataset search in NCBI's Sequence Read Archive (SRA) was also performed using the search term

'salivary microbiome'. The included studies met the following criteria: 1) Having samples from

participants without any diagnosed disease state. For studies investigating the influence of certain

kind of disease on salivary microbiota, only samples collected from healthy controls were included in

further analysis; 2) Using whole human saliva collected by spitting, swab, mouth washing or oral

rinsing, samples exclusively extracted from any specific oral spot, like tongue surface, parotid gland,

supragingival plaque, were excluded; 3) Using 16 rRNA gene high-throughput sequencing and

sequenced with the Illumina MiSeq platform; 4) Having and sequence file with quality score and

associated metadata, information about geographic locations are required; 5) Having freely available

sequencing data; 6) Sequencing data correctly separated according to the metadata. Raw sequence

data acquired from the healthy individuals of selected studies were downloaded from SRA, European

Nucleotide Archive (ENA), using SRA Toolkit. The files were converted to the FASTQ formats if

necessary.

2. 16S rRNA gene sequence processing

Sequence data from each selected study were processed separately using QIIME2 (version.2020.2)

[34]. Sequences with primers were trimmed with "q2-cutadapt" (https://github.com/qiime2/q2-

cutadapt) to retain the targeted hypervariable regions. The demultiplexed paired-end sequences were

firstly joined by "q2-vsearch" (https://github.com/qiime2/q2-vsearch), then subjected to a quality

filter with a minimum quality of Q30. The remaining reads were then clustered into operational

taxonomic units (OTUs) at 97% similarity against the expanded Human Oral Microbiome Database

(eHOMD) version 15.1 [35] by the closed-reference OTU picking command. Reads that failed to

match a reference sequence in the HOMD database were discarded. The chimeras and features with a

frequency ≤10 or detected in a single sample were also removed. The resulting tables and sequences

from all studies were merged by QIIME2's merge and merge-seqs commands. Samples with <2000

reads were also removed. Taxonomic annotations were assigned to the representative sequences of

each OTU using the HOMD database. For the factor groups containing samples with unknown

metadata, the unknown sample was removed from the group before the downstream analyses.

3.   **Diversity measures in R**

The merged OTU table was exported into BIOM format. Further analyses were carried out in R

(version 4.1.0) with custom scripts as detailed below. Samples with >2,000 reads were retained and

processed with four normalisations: 1) Rarefying samples to 5,000 (Rarefaction, RAR); 2) Samples

were rarefied to 5,000 and converted to relative abundance (Rarefied Total-sum Scaling, RRA); 3)

Samples were converted to relative abundance directly (Total-sum scaling, TSS); 4) As described by

Romano et al. [36], zeros were added to data through the count zero multiplicative approach using

the *cmultRepl* function of the *zCompostions* package [37] in R (Centred Log-ratio, CLR).

The alpha-diversity of all samples grouped by studies were calculated in the form of Chao1, Shannon,

and Simpson's diversity indices. The beta-diversity was assessed at different taxonomic levels,

including OTU, species, genus, family, order, class, and phylum level. For the OTU level, the beta-

diversity of data processed with the first three normalisations were determined using the weighted

UniFrac distances and Bray-Curtis dissimilarities. Euclidean distances were calculated for all

normalisations. At the other levels, Bray-Curtis dissimilarity was used for the first three

normalisations, while Euclidean distances were combined with all the normalisations. Permutational

multivariate analysis of variance (PERMANOVA) using the *adonis2* function of *vegan* package [38]

with 999 permutations was conducted to investigate the statistical differences caused by different

factors, adjusting for study.

4. Defining the core microbiome

The core microbiome was determined based on the abundance-occupancy pattern. Two methods

adapted from Shade and Stopnisek [39] and Wu et al. [13] were used. For the methods adapted from

the study of Shade and Stopnisek [39], samples were rarefied to 5,000. Both Bray-Curtis similarity

and weighted Unifrac distance were used to determine the contribution in the percentage of the

prospective core set to the overall beta diversity. For the method adapted from Wu et al. [13], OTUs

were filtered out with the mean relative abundance (MRA) bigger than 0.1% and the presence in more

than 75% of samples or 100% occupancy in more than 10 studies. To investigate the bacteria-bacteria

interactions in salivary microbial communities, the co-occurrence network was constructed using

pairwise Spearman's correlation based on relative abundance. The Spearman's correlation was

calculated using the *rcorr* function in the *Hmisc* R package [40] and visualised by Cytoscape v3.8.2

[41]. A correlation with Spearman's correlation coefficient > 0.5 or < -0.5 and p-value < 0.01 is

considered as statistically robust and shown in the network.

5. Differential abundance analyses

5.1 Random Forest

The normalised abundance of taxa in the phylum, class, order, family, genus, species and OTU level

were classified against each provided metadata categories to determine which factor has the largest

effect on the salivary microbiota. A random forest classifier was created in R using the *randomForest*

package [42] with default parameters. We used the *randomForest* (importance = TRUE, proximity =

TRUE) function to generate the classification model for seven categories. For four categories that

have the random forest classifier with average error rate lower that 20%, including hypervariable

region (5.0%), geographic location (10.4%), tobacco usage (12.7%), sample type (13.8%),

differentialt taxa were defined using cross-validation. Cross-validation was performed by the *rfcv()*

function for selecting appropriate features. The *varImpPlot* function was used to show the importance

of features in the classification. The importance of features and the cross-validation curve were

visualized by using the *ggplot2* package [43] in R.

5.2 Analysis of Compositions of Microbiomes with Bias Correction (ANCOM-BC)

ANCOM-BC [44] were performed to identify the taxa with different relative abundance between

Chinese and Western samples. Function *ANCOMBC* were used with holm-bonferroni false discovery

rate correction and other default parameters. The hypervariable regions used by different studies were

used as the covariate.

6.   Functional prediction

Microbial metagenomes were inferred from 16S rRNA gene-based bacterial profiles, and the

functional prediction were conducted based on Kyoto Encyclopedia of Gene and Genomes (KEGG)

database [45] using the default pipeline in Phylogenetic Investigation of Communities by

Reconstruction of Unobserved States 2 (PICRUSt2) (Douglas et al., 2020). The ANCOM-BC analysis

was used to identify the differential abundant KEGG pathways by geographic location, adjusting for

hypervariable regions. At the same time, a random forest model was established for distinguishing

Chinese and Western samples, and the importance of pathways was measured using mean decreased

accuracy. Spearman's correlation was performed to assess the relationship between the relative

abundance of differential pathways and genera. The significant correlations were visualised using the

corrplot package [46] in R.

7.   Comparison between Chinese and Western people on an independent cohort

Saliva samples were collected from 26 participants (aged 20-60 years) recruited for a wine assessment

experiment and consisted of 13 Chinese and 13 Western wine experts (Table 1). The study was

approved by the Office for Research Ethics and Integrity of the University of Melbourne (Ethics ID:

1852616). Each group had six female panellists and seven male panellists. The Western panellists

were defined as people who have lived in Australia for more than ten years. Chinese panellists were

defined as people who were born in China and had lived in Australia for no more than 18 months.

Bacteria genomic DNA was extracted from human saliva using QIAGEN® MagAttract® PowerSoil®

DNA KF Kit [47] and subjected to 16S rRNA amplicon sequencing on the Illumina platform

following the Earth Microbiome Project protocols (https://earthmicrobiome.org/protocols-and-

standards/16s/). The raw data are available in NCBI Sequence Read Archive, with accession number

PRJNA786805.

The raw sequences were processed using the same pipeline in the meta-analysis as described in

section 2. Additionally, raw sequencing reads were denoised into zero-radios OTUs (zOTUs) by

UNOISE3 pipeline [48], and taxonomically classified by classifiers trained on the full-length 16S

rRNA gene SILVA v138 [49] database and eHOMD v15.22 [35], respectively. The affiliation

between each ZOTUs and the originating OTU was determined using a customised code adapted from

Stopnisek and Shade [50] and available at

https://github.com/XINWEIR/SalivaryMicrobiome_MetaAnalysis. The relative abundance of taxa at

the genus level in this cohort was used as the test set for the random forest model trained using the

genus-level OTU assignment information in the meta-analysis.

Results

1. Inclusion of studies and sequences

In this study, we extracted the 16S rRNA gene amplicon sequencing data of healthy human saliva

from 47 studies (Figure 1A). We abstracted data from subjects who had no diagnosed disease state,

hereafter named "healthy". Of course, subjects could have had subclinical diseases or may have

altered health status for undisclosed reasons, but we considered that this would be true of the wider

human population and therefore able to be included in our study. A total of 107,005,868 high quality

16S rRNA sequences were obtained. After removing all samples below 2,000 reads, 2206 samples

with 909 features were retained. The retained samples included studies from 15 countries (Table S2).

Most studies were conducted in three geographic regions: North America, Europe, and China (Figure

1B). Most sequences included in this meta-analysis were generated from the hypervariable region

"V3-V4" and "V4". Samples from studies targeted at "V1-V2" or the "V4-V5" region were classified

as "others" (Figure 1C; Table S3). Similarly, saliva samples collected by unconventional methods like

"swab" were also classified as "others". People who recorded a smoking habit, whether they smoke e-

cigarettes or tobacco, were categorised as smokers. Similarly, people who drink alcohol, regardless of

the frequency or the type of alcohol consumed, were classified as drinkers. The given age was treated

as a categorical factor, classifying into "18-30", "31-55", and "56+". The samples without associated

information to a particular category or classified as "others" were excluded for downstream analyses

related to the impact of this category. For example, the following analyses measuring the effect of

hypervariable region on microbial profiles included only comparisons between the V3-V4 and V4

region (Figure 1C).

2.   Intrinsic and lifestyle factors have a significant effect on the host salivary microbiome

Large variability between studies was observed in the number of reads, taxonomic profile, and alpha

diversity (Figure 2; Figure S1). Phylum *Bacteroidetes*, *Proteobacteria*, *Firmicutes*, *Fusobacteria* were

dominated among all studies, while their proportion varies (Figure 2A). When studies were grouped

by the geographic locations they originated from (coloured in Figure 2B, C, D), there is generally no

difference between their intra-community diversity, represented by Shannon, Chao1, and Simpson

indices. Only one study conducted in Qatar showed relatively lower Chao 1 index and higher

Simpson's diversity indices than studies from other locations. However, it is hard to decide whether

such variation is caused by the geographic location or other technical variations.

Because of the large disparity of methodologies amongst the studies used in our global analysis, we

applied several different strategies for normalisation as described in Methods. When investigating the

influences of different categories using permutational multivariate analysis of variance

(PERMANOVA) tests, these normalising methods were combined as appropriate with different

distance metrics, including Bray-Curtis, weighted UniFrac, and Euclidean distance. Overall, the effect

of rarefaction (RAR), total-sum scaling (TSS), and the rarefied relative abundance transformation

(RRA) were very similar in the result of PERMANOVA (Figure 3A-D; Table S4). Meanwhile, the

centred log-ratio transformation (CLR) enlarged the variance induced by an unwanted technical

factor, namely the amplified hypervariable region, at all taxonomic levels.

The beta-diversity analyses showed that all metadata categories measured have a significant (p<0.001)

effect on the bacterial profile of human saliva at all taxonomic levels, adjusted for the study effect

(Table S4). However, only limited variation among samples has been explained by these factors ($R^2 <$

10%). In contrast, "study" accounts for around 35% of the variability between samples. At the OTU

level, the combination of weighted UniFrac distance and the total-sum scaling (TSS) transformed data

best minimised the variability raised by different hypervariable regions (Figure 3D). The results of

the unconstrained principal coordinate analysis (PCoA) are in agreement with the results of

PERMANOVA. When using Bray-Curtis dissimilarity and Euclidean distance, the samples separated

distinctly according to the hypervariable regions in PCoA plots, whereas the plot constructed using

the weighted UniFrac revealed the clusters formed by the samples from different geographic locations

(Figure S2). A distinct separation of samples from three main geographic locations (Figure 3F), with

more than half (58.0%) of the variance explained by the first two dimensions, using weighted UniFrac

distance. In contrast, the differences between locations were confounded by which hypervariable

regions were sequenced in the PCoA plot for Bray-Curtis dissimilarity (Figure 3E). The results

suggest that host intrinsic and lifestyle factors significantly influence the microbial profile in human

saliva, regardless of the variation induced by technical factors.

3. A core microbiome is defined from saliva from healthy humans

Despite the large intra- and inter-study variability, many OTUs still showed a consistently high

presence and relative abundance across studies (Figure S3). These persistent OTUs detected across

studies with different protocols could be functionally important for the salivary microbiome of healthy

adults. We wanted to identify the most widespread microbial taxa within a specific population that

allows us to better understand the broad structure of microbiomes and their potential functional

consequences [12]. The abundance and occurrence frequency of taxa are two important criteria used

to define the "core salivary microbiome". Conventionally, thresholds on these two parameters filter all

taxa detected, and taxa that meet both criteria can be classified as the core salivary microbiome [13].

Recently, a more standardised procedure based on abundance-occupancy distribution was proposed

[39]. We employed both strategies to define the core salivary microbiome to identify the microbial

features with high persistence and robustness in human saliva. Considering the sequences involved in

this meta-analysis were collected from studies targeted at different hypervariable regions, close-

referenced clustering at 97% identity were used to cluster sequences into OTUs. In addition, taxa

defined at 100% sequence identity may increase the redundancy in the dataset [39]. Therefore, the

core salivary microbiome was defined using the clustered OTUs at 97% identity. To begin, the core

OTUs were determined by filtering all OTUs based on mean relative abundance and occurrence

frequency using the criteria described in the Methods [13]. In total, 11.6% of all OTUs (105 OTUs)

were included as Core 1 (Figure 4A: MRA + OCC; Table S5). Meanwhile, according to the method

proposed by Shade and Stopnisek [39], OTUs were ranked depending on their occupancy across

studies, and the contribution of top-ranked OTUs to beta-diversity was expressed by Bray-Curtis

similarity and weighted UniFrac distance. Two groups of the core microbiome were prioritised by

these two indices, to give different inclusions in the core, consisting of the top 69 OTUs (using

weighted UniFrac; Figure 4A: BC) and 94 OTUs (using Bray-Curtis; Figure 4A: wUF) OTUs (Figure

S4).

Overall, sixty-eight OTUs were shared across all three methods (Figure 4A), accounting for 7.5% of

all OTUs detected and 72.5% of all 16S rRNA gene sequences after clustering and filtering (Figure

4B, C). *Firmicutes* account for nearly half (46.4 %) of all core OTUs, while only one OTU belongs

to *Saccharibacteria*. The mean relative abundance (MRA) of each OTU in sub-groups classified by

different factors was also measured (Figure 4D). On average, the core OTUs were highly prevalent

(73.2% ± 3.4% of cumulative relative abundance) in saliva samples across different levels in

subgroups classified by age, gender, geographic locations, hypervariable regions, sample type,

smoking, and drinking habits. The core OTUs were clustered into four main groups based on their

distribution pattern in sub-groups (Figure 4E). The eight OTUs affiliated to Cluster 1 showed overall

high abundance in all sub-groups. Cluster 2 consists of core OTUs with a slightly lower mean relative

abundance than Cluster 1 and higher intra-group variability. Notably, although having a higher MRA

than some members of Cluster 1, "476_9291" was still classified as Cluster 2. The reason could be its

biased presence in sub-groups. For example, the relative abundance of "476_9291" is higher in

samples from China than other two locations. The other two clusters contain OTUs with lower MRA

than Cluster 1 and 2, while variations can still be observed within sub-groups.

Bacteria-bacteria interactions play an important role in shaping microbiota. Therefore, co-occurrence

network analysis could be a useful approach for finding the most important part of the microbial

community. We applied a network analysis built by Spearman's correlations to investigate whether

the core OTUs defined were also important to the structure of the co-occurrence pattern. A correlation

with Spearman's correlation coefficient >0.5 or < -0.5 and p-value < 0.01 is considered statistically

robust [51, 52]. The resulting co-occurrence network contains 293 nodes and 1,424 significant

correlations (edges) (Figure S5). Small modules with less than seven nodes were not displayed.

Although most core OTUs have relatively low connectivity, they tend to associate with each other

rather than with rare OTUs. Nine OTUs were identified as potential "hub" OTUs based on their

centrality and the number of links in the network (Figure S5, 6). Because of the central position in the

network, the hub taxa are regarded as the key contributor to community stability. Two core OTUs

were also identified as hub taxa in the network, which are "122BU057" (*Megasphaera*

micronuciformis) and "524_3631" (*Veillonella atypica*). Compared to other "hub" taxa, they showed

lower connectivity and relatively high betweenness centrality (Figure S6).

4.   Geographic location is the host factor with the largest impact on bacterial composition

To investigate which metadata category has the largest impact on the salivary microbiome, we

established random forest models to link the seven categories described above and a new category,

study, with the salivary microbiota data at seven taxonomic levels (OTU, phylum, class, order, family,

genus, and species). The effect of four normalisation methods was compared using the error rate

generated by random forest classification. In total, 224 random forest models were constructed

(Figure 5A). Among four normalisation methods, total-sum scaling produced the models that were, on

average, the most accurate. Generally, the random forest models built with microbial communities at

OTU levels have the lowest error rate (mean = 13.0%), while the models constructed at phylum levels

have the highest (mean =26.8%). The model built with the hypervariable region used for sequencing

was also the category that showed the lowest error rate (Figure 3G). Geographic locations

demonstrated the second important impact on the bacterial communities, with the lowest error rate

among biological factors. The random forest model constructed by the other two categories, sample

type and tobacco usage, also showed a relatively lower error rate than other categories. Study

constructed the models with high error rates at phylum (44.3% ± 2.5%) and class level (29.3% ±

2.1%). However, the error rate of models built with study rapidly dropped with the increase of

taxonomic levels, reaching 10.5% ± 5.1% at OTU level. Gender and age range led to poorly

performing models.

We wanted to determine whether the defined core microbiome could be used as biomarkers to

differentiate people categorised by intrinsic and lifestyle factors. The random forest models showed

high accuracy at OTU level were used (i.e., geographic location and smoking factors). The differential

OTUs induced by hypervariable regions and sample types were also analysed to exclude the influence

of technical factors. We further performed ten-fold cross-validation five times to measure the

importance of OTUs used to train the model. All OTUs before the point that the cross-validation error

curve starts to stabilise were defined as important OTUs. In total, we defined 59, 57, 34 and 70

important OTUs as biomarkers to differentiate samples according to geographic location, smoking

habit, hypervariable region, and sample type, respectively (Figure S7). Of these, 28, 10, 13 and 22

biomarkers were also classified as "core" (Figure 5B; Figure S8). Although 31 core OTUs showed the

importance in discriminating samples according to geographic location and smoking, nearly half of

them (15 OTUs) had the possibility of being confounded by technical factors (Figure 5B). After

excluding the OTUs that could be influenced by other factors, core OTU "322AK152" (*Bergeyella*

sp.HMT_322) was the OTU with the highest contribution to the classification of samples from three

geographic locations. Meanwhile, "122BU057" (*Megasphaera micronuciformis*) showed the highest

importance among the core differential OTUs specific to smoking, followed by "524_3631"

(*Veillonella atypica*). We were surprised to find that these two OTUs were the only two OTUs that

were defined as hub taxa in the co-occurrence network analysis (Figure S6).

5. The salivary microbiota as biomarkers to differentiate Chinese and Western

We further analysed the changes caused by geographic locations in higher taxonomic hierarchies, where many differences have been revealed. Of particular interest were taxa under phylum *Synergistetes* and *Spirochaetes*, Class *Mollicutes* and *Betaproteobacteria*, Family *Clostridiales*, and genus *Prevotella*. Interestingly, many taxa showed a higher relative abundance in the Chinese samples, both compared to North American samples and compared to European samples (Figure 6A). It suggested that the variance induced by geographic locations may be dominated by the differences between samples from Chinese and Western people. Therefore, we combined samples from North America and Europe into a single group, "Western." Compared to the Chinese grouping, the Western group has significantly lower within-sample diversity (alpha-diversity) (Wilcoxon rank-sum test, $p < 0.001$; Figure 6B, C). Next, we examined the differences between Chinese and Western in the salivary microbiota at the genus and species level (Table S6, S7). Besides establishing a random forest model, we also identified differential taxa using ANCOM-BC, adjusting for the hypervariable region. We found 48 genera identified as significantly different by both methods (Figure 6D, Table S6). Among them, *Arachnia*, *Filifactor*, *Ottowia*, *Neisseria*, *Aggregatibacter*, one genus from *Gracilibacteria*, one genus from *Clostridiales*, and three other genera belonging to the family *Peptostreptococcaceae* were strongly (standardized effect size>10, mean decreased accuracy >10) enriched in Chinese samples.

Meanwhile, *Prevotella*, *Scardovia*, *Bergeyell*, *Veillonella*, *Oribacterium*, and one genus belonging to the family *Erysipelotrichaceae* were strongly (standardized effect size>7 mean decreased accuracy >10) enriched in Western samples.

Finally, we performed the functional prediction-based 16S rRNA gene profiles to investigate whether differences in the salivary microbiota between Chinese and Western affect its function. Two methods, ANCOM-BC, and random forest model were used to identify which pathways were differential between Chinese and Western. The result of ANCOM-BC indicated that 69 pathways related to metabolism were differentially abundant between the two groups. The random forest classification

model established using KEGG pathways demonstrated an error rate of 10.01% and revealed 46

differential pathways. Among them, thirty pathways belonging to nine upper pathways (level 2) were

simultaneously defined by two methods as differing in abundance between Chinese and Western

(Figure 6E, Table S8). A variety of pathways was in higher abundance in Chinese samples. The

enrichment of these pathways in Chinese samples was mainly associated with the increased

abundance of *Neisseria* and *Lautropia* and the depleted abundance of *Prevotella*, *Veillonella*,

and *Atopobium*. Notably, three lipid metabolism pathways enriched in Chinese samples, including

"Ether lipid metabolism" (ko00565), "alpha-Linolenic acid metabolism" (ko00592), and "Linoleic

acid metabolism" (ko00591), have the highest standardised effect size (W statistics, Table S8). The

enrichment of these pathways related to lipid metabolism has been positively associated with the

higher abundance of *Neisseri*a in Chinese. *Neisseria* may have also contributed to the pathway

"Carotenoid biosynthesis" (ko00906). Another metabolic pathway related to the metabolism of

terpenoids and polyketides, "Sesquiterpenoid and triterpenoid biosynthesis" (ko00909), showed a

positive correlation with a genus belong to *Peptostreptococcaceae*. In contrast, only one pathway

named "Flavone and flavonol biosynthesis" (ko00944) was enriched in the saliva samples from

Western. A strong positive correlation has been demonstrated between this pathway and the increased

abundance of *Veillonella* in the samples from the Western grouping.

6.   Validation of the core in an independent Australian cohort

To validate the prevalence of the core OTUs in human saliva, we collected saliva samples from 13

Chinese and 13 Western participants in Melbourne and sequenced the extracted DNA with 515F-

806R primers. In total, 841,188 high-quality 16S rRNA sequences were obtained, which clustered into

397 OTUs with 97% identity to the HOMD database. Among them, the core OTUs we defined in the

meta-analysis showed high relative abundance ($78.3 \pm 6.9\%$) in all collected samples. To increase the

accuracy of the OTU assignment, we denoised the sequencing reads using the UNOISE3 pipeline and

generated ZOTUs with 100% sequence identity. After re-clustering the core OTUs defined in the

meta-analysis to ZOTUs, we observed that 59 of the identified OTUs in this independent dataset

consisted of 87 ZOTUs, and thus made up close to 80% of the relative abundance (Figure 7A).

Although some sequences belonging to the same ZOTU are clustered to different OTUs, all the core

OTUs contain at least one highly abundant ZOTU. The taxonomic profiles of the global core

annotated by the HOMD database and the ZOTUs annotated by the SILVA database were very

similar at the genus level (Figure 8A).

We wanted to verify the observed differences between the OTUs in saliva samples from Chinese and

Western people in this independent dataset. Although two groups did not differ significantly when

considering the Shannon diversity index (Wilcoxon rank-sum test, $p = 0.073$; Figure 8B), the Chinese

group showed a higher Chao 1 Index than the Western group (Wilcoxon rank-sum test, $p < 0.001$;

Figure 8C), which is in agreeance with the result of the meta-analysis. Meanwhile, no significant

differences were observed between the two groups by beta-diversity analyses (Bray-Curtis and

weighted uniFrac distance, Table S9). We used the random forest classification model constructed

using the genus level profile of the large-scale dataset to predict the Chinese and Western samples in

this independent study. This dataset's genus-level relative abundance table was prepared from both the

OTU table with 97% identity to HOMD v15.1 database (Figure 8D) and the ZOTU table annotated by

the HOMD v15.22 database (Figure 8E). The accuracy of both predictions was relatively low, with

57.7% for the OTU table and 50% for the ZOTU table. Interestingly, most Western samples were

correctly classified, while most samples from Chinese participants were classified as being 'Western'

in this analysis.

Discussion

There is ample knowledge on the disease-affected salivary microbiota, yet our perspectives to the

bacteria present in healthy humans remains limited. Our systematic selection of studies, together with

the re-analysis of the 16S rRNA amplicon sequencing data from 47 studies offers a comprehensive

description of the salivary microbiome presented in adults without diagnosed disease. Our study has

defined the core members of salivary bacterial communities across 2211 samples from 47 studies and

has used metadata captured in these studies to investigate the role of different intrinsic and extrinsic

factors on the occurrence of these core. It is clear that core members differ between geographic

locations of collected saliva, and our analysis shows that Chinese participants are different from

Western participants (encompassing European and North American studies). A prediction of the

pathways enriched in each collective indicates that bacterial metabolic pathways are likely to

influence the aroma and flavour perception of foods. These results show that despite the core

microbial members of saliva being common across humans, there are differences, likely due to diet.

We suggest that the aroma and flavour of foods and beverages are likely to be differently affected in

healthy humans across the globe, meaning that preference and consumption of different foods is likely

to be prioritised. These results have important consequences for food and beverage design,

composition, and dietary advice across the globe.

Based on the abundance-occupancy pattern, the definition of core microbiome highlighted the

persistent and conserved microbial communities in human saliva across the globe. Here, we compared

two approaches adapted from two studies (references) to defining the core. The method adapted from

the study of Wu et al. [13] is a relatively conventional strategy that has been chosen by many other

studies investigating different ecosystems, such as soil [53], compost [54], wastewater treatment

plants [55], and human's intestinal system [56]. The thresholds were simply setting on each taxon's

mean relative abundance and occupancy across all samples. For studies aimed to determine the spatial

or temporal core microbiome, additional thresholds will be added on the prevalence of taxa within

sub-groups. However, the thresholds used by different studies are usually arbitrary. Some studies have

even adopted only abundance or occupancy alone as criteria for defining the core. Therefore, a

generalised approach for defining the core microbial members from diverse datasets based on

abundance-occupancy was also applied in this study [39]. Rather than over space or time, we

determined the occupancy of OTUs according to their detection over study. When evaluating the

contribution of top-ranked OTUs in occupancy to the beta-diversity of the community, we further

used weighted Unifrac due to its effectiveness in minimising the biases induced by the selection of

hypervariable regions.

The resulting core members identified by these methods have a lot in common. A majority of the core

OTUs defined by Shade's method is also included in the cores defined by Wu's method, suggesting

the recently developed multi-step approach is effective in determining taxa with high prevalence. The

general high relative abundance of the core across different sub-groups emphasised the utility of this

pipeline in identifying the persistent members across diverse datasets. Most of the core salivary

microbiota we defined had been proposed in previous studies as prevalent bacteria in the human oral

cavity that persistently span across different individuals [11, 57-59]. The dominant genus of the core

we defined, *Streptococcus*, *Neisseria*, and *Prevotella*, were concluded as core human salivary

microbiome by a recent study based on the MG-RAST data [60]. Ten OTUs belonging to genus

Streptococcus were included in the "core", two of which were classified to the cluster with overall the

highest relative abundance across all dimensions. The prevalence of *Streptococcus* we observed is

consistent with a previous study defining the healthy core from the 454 pyrosequencing results of

three individuals [11]. The most abundant core OTU we found, *Streptococcus*

oralis subspecies *dentisani*, has been documented in previous studies as potential oral health-

promoting organisms and being highly abundant at various oral niches of healthy humans [61, 62].

497
We further conducted a co-occurrence network analysis to investigate the role of these core

microbiota in shaping the microbial community and found the co-existence between many members

of the core (Figure S5). The presence of the rare OTU that became the hub suggests that although

some taxa are not persistently detected across the community, they may still be important for the

overall structure of the salivary microbiome. It has been proposed that the oral microbiota of healthy

individuals is both homeostatic and dynamic [57]. The core microbial members consistently present in

human saliva identified here may explain the stability of oral microbiota to some extent.

Moreover, we conducted an independent study to verify the prevalence of the core defined from the

published studies. The high relative abundance and occurrence of the original core OTUs in this

independent cohort suggest that the core human salivary microbiome we defined can be applied to

different datasets. After re-clustering the core OTUs to ZOTUs with 100% sequence identity, we may

conclude that the same members constitute the core, even if different taxonomic resolution is applied.

Besides the core microbiome, there are "variable" microbiota in human microbial communities, which

vary among individuals because of unique lifestyle and genetic factors [63]. We performed analyses

for beta-diversity of samples (Figure 3) and random forest classifications (Figure 5A), demonstrating

several factors-both technical and physiological-significantly discriminated between sub-populations.

Because of the high heterogeneity between studies in their methodology, large inter-study variability

was the main factor that affected the observed salivary microbiota [64, 65]. As one of the main

technical factors that may induce the variation, the impact of chosen hypervariable regions for

sequencing on driving microbial community structures has been confirmed by our study. In addition,

the selection of different primers for the same region and DNA extraction methods may also lead to

technical variations [66]. However, studies' choices of primers and DNA extraction protocols are

more diverse than hypervariable regions (Table S2), making it difficult to group them into categories

as simple as for the variable regions. By adjusting the analyses by study, the variability caused by

study-specific technical factors other than hypervariable regions could also be covered. For future

studies, it is important to establish a standardised DNA extraction and sequencing protocol on a global

scale. During the literature search, we found that the inter-study variation may also be attributed to the

criteria of recruiting participants. Although all samples included in this meta-analysis were collected

from the control groups and population without specific diagnosed disease, the definition of 'healthy'

varies. For example, the use of antibiotics was not always considered as an exclusion criterion, and

when included, different time intervals were adopted. However, for this study, we aimed to construct

a microbial community that reflects the salivary microbiota of real-life consumers. Therefore, samples

collected from individuals without known severe systemic diseases were included.

Besides the influence of variables related to study design, this study has also revealed the importance

of various host intrinsic and lifestyle factors. We observed the change in salivary microbial

composition induced by smoking habits. As a recognised risk factor to oral health, the role of smoking

in shaping the human oral microbiome has received increasing attention [67]. Previous studies have

reported the change of *Megasphaera micronuciformis* caused by smoking in the microbial profile of

the human tongue surface [68] and upper gastrointestinal tract [69]. In agreement with these findings,

we identified a core OTU belonging to *Megasphaera micronuciformis* as a biomarker for reported

smoking. In addition, we found the differential abundance of an OTU belonging to *Veillonella atypica*

between smokers and non-smokers, which agrees with a study where this species already found to be

increased in the saliva [70]. Another strong determinant of the salivary microbiome we defined was

"sample type". As we found in this study, the difference between the mouthwash sample and the other

two collection methods is greater than the difference between the stimulated and unstimulated saliva.

Contrary to the result of Jo et al. [29], the OTU belonging to *Neisseria flava* has not been identified as

differential taxa for the type of saliva. The microbial composition of alcohol drinkers and non-

drinkers was also found to be different, while the small sample size acquired hindered our ability to

draw large-scale downstream conclusions. Although the significant differences we observed in

salivary microbiota were also attributed to the gender and age of participants, the variations they

explained are relatively low compared to other factors.

Geographic location has been identified as the host physiological factor with the largest impact on

salivary microbiota (Figure 5A). Although it only explained limited variability between samples'

microbial profiles, the observed variations were robust to the heterogeneity induced by different

hypervariable regions used (Figure 3E, F). To date, little is known regarding the influence of

geographic locations on the human salivary microbiome. A comparative study reported the

differences in saliva microbial composition in Alaskans, Africans, and Germans [33]. To our

knowledge, the geographically structured microbial communities have not been observed in healthy

human saliva based on the large-scale dataset used in this study. Our result also demonstrated several

core OTUs that may differentiate saliva samples from North America, Europe, and China. It suggests

that the global prevalent core microbiota is not necessary to be stable across populations. Given the

high abundance and occupancy frequency of the core microbiome, we would expect these taxa to be

effective indicators to predict the geographic background of saliva donors.

565 Due to the sometimes large differences in culture and lifestyle between Western and non-Western

566 populations, we further grouped our data into Western and Chinese samples. The comparison between

567 Western and non-Western populations has already been applied to the human gut microbiota, whereas

568 less is known about the saliva microbiota [71]. Our study found a difference in the abundance of

569 *Veillonella* spp between saliva from Western and Chinese people, where *Veillonella* was generally

570 higher in Western samples. Such differences may influence the flavone and flavonol biosynthesis

571 pathways in the oral cavity. Our previous study revealed the Western-born and Chinese-born wine

572 experts had different responses to the astringency of wine. Since flavonol is a well-known constituent

573 of wine-related to the bitterness and astringency perception [72], we would hypothesis that the

574 enrichment of *Veillonella* in Western may affect their sensitivity to the phenolic compounds in wine.

575 It has also been suggested that the regular consumption of flavonoid-rich foods, such as oolong tea,

576 may increase the abundance of *Veillonella* spp. in human saliva [73]. These influences can potentially

577 be the bridges that link the differences between Chinese and Western groups in sensory evaluation

578 and their salivary microbiota together. Because the amplicon sequencing data cannot speak directly to

579 the functional sequences of the observed difference, shotgun metagenome sequencing will be

580 necessary to verify the exact association. *Prevotella* abundance has previously been reported to be

581 enriched in the gut microbiota of non-Western populations [74], which is opposed to our observation

582 in saliva. However, there is substantial species-level diversity in *Prevotella* (Table S7), making it

583 plausible that different species belonging to *Prevotella* may respond differently to the geographic

584 background [75].

586 Although the independent dataset we collected did not show differences between samples taken from

587 Chinese and Western participants, the results of random forest classification may lead to some

588 interesting hypotheses. The prediction of models revealed that most of the Chinese samples in this

589 cohort were classified as Western. The donors of these samples were wine experts, of Chinese

590 ethnicity, born in China and living in Australia for no more than 18 months. Recent studies reported

591 that immigrants from Asia experience a "Westernization" of gut microbiota induced by dietary

592 acculturation [74, 75]. We hypothesise that such a phenomenon may also happen in salivary

593   microbiota. It may suggest that dietary pattern is a more important determinant than ethnicity in

594   shaping the salivary microbiota of the participants, leading to variation among different geographic

595   locations that we observed in the meta-analysis.

Conclusions

In summary, we have defined a core bacterial community in saliva from healthy humans, and this core

demonstrated both stability and variability among populations. The prevalence of the core members of

the saliva microbiome has been confirmed in an independent cohort. We have revealed the influence

of various host factors, such as geographic locations, incidence of smoking and drinking, on the

salivary microbiome. We also identified microbial and functional biomarkers to differentiate the

Chinese and Western people, underlying the potential relationship between salivary microbiota and

sensory perception. Results in this work will provide foundational information to inform future

studies to understand the similarities and differences in saliva microbial composition, potentially

associating oral to aroma and flavour perception of foods.

Availability of data and materials

The sequencing data supporting the conclusion of the meta-analysis in this article are available in

publicly accessible databases (full details can be found in Table S2). The sequencing data generated

and/or analysed during the current study are available in the NCBI Bioproject repository,

PRJNA786805 (https://www.ncbi.nlm.nih.gov/bioproject/786805). Original scripts generated during

the current study are available in Github

(https://github.com/XINWEIR/SalivaryMicrobiome_MetaAnalysis).

Acknowledgements

This study was funded by the Faculty of Veterinary and Agricultural Sciences at the University of Melbourne. JL and XR gratefully acknowledge a Melbourne Research Scholarship administered by the University of Melbourne.

Legends-Results

Figure 1. Overview of literature search procedure and metadata of included studies. a) Large-scale literature searching and data filtering process, followed by the number of samples submitted to the bioinformatic analyses; b) The locations of studies, the scale of symbols that reflect the number of samples of each study; c) Distribution of metadata categories.

Figure 2. Summary of taxonomic composition and alpha diversity of included studies. A) The mean community composition of each study at the phylum level; The alpha-diversity measured by B) Shannon index; C) Chao 1 index; D) Simpson's index, the colour of boxes stands for the geographic location of the studies. The horizontal bars within boxes represent medians. The tops and bottoms of boxes represent the 75th and 25th percentiles, respectively.

Figure 3. The variability in human salivary microbiota have been explained by different factors. Among them, hypervariable regions and geographic locations have the largest impact. The effect of the categories on the clustering of the sample was measured using PERMANOVA at four taxonomic levels: family **(A),** Genus **(B),** species **(C)** and OTU level **(D).** The colour indicates the different combinations of normalisation (TSS, Total-sum scaling; RRA, Rarefied relative abundance; CLR, Centred log ratio) and indices (BC, Bray-Curtis; EUC, Euclidean; wUF, weighted uniFrac). Because the results of rarefication (RAR) were very close to TSS and RRA, they were not displayed in the figures. Principal coordinate analysis (PCoA) with Bray-Curtis **(E)** and weighted uniFrac **(F)** showing the differences between samples from North America, Europe, and China.

Figure 4. The core OTUs defined by abundance-occupancy pattern. A) Venn diagram showing the interaction between three methods used to define the core. Sixty-eight OTUs were defined as the

core for all methods. (MRA+OCC: The thresholds were setting on mean relative abundance and

occupancy to define the core; BC: The method adapted from Shade and Stopnisek using Bray-Curtis

similarity; wUF: The method adapted from Shade and Stopnisek using weighted uniFrac distance). **B)**

Pie chart showing the number of the core (pink) versus other OTUs (blue) identified in percentage. **C)**

Pie chart showing the relative abundance of the core and other OTUs across all samples. **D)** Relative

abundance of 68 core OTUs across subgroups classified by seven categories. **E)** Heatmap showing the

log-transformed mean relative abundance of each core OTU at each level of different categories.

Figure 5. Salivary microbiome members which significantly contribute to categorisation of

metadata. Random Forest models showed the impact of categories on salivary microbiome and

the core OTUs contributing to accuracy of these models. **A)** Error rate (%) for the random forest

classifications conducted with samples grouped by eight different categories. **B)** Phylogenetic tree

indicates the taxonomic information of 68 core OTUs. The coloured squares between the tree and the

annotation of phylum indicate the OTUs that were defined by the Random Forest model as

"important" for distinguishing between different levels in each category. The bars on the outmost ring

showing the mean relative abundance of each OTU.

Figure 6. Distinct microbial profiles are evident in the saliva samples from Chinese and Western

adults. **A)** Taxonomic hierarchies show the relative enrichment of taxa in three geographic locations

at phylum through species level. Coloured nodes indicate log2-fold increase in median abundance of

the group in x-axis (pink) or y-axis (blue). Only taxa showed significant changes (false discovery rate-

adjusted Wilcoxon rank sum q < 0.05) are displayed. **B)** and **C)** Comparison of salivary microbial

alpha diversity between the Chinese and Western samples, calculated by Chao1 (B: $p< 0.001$,

Wilcoxon rank-sum test) and Shannon index (C: $p< 0.001$, Wilcoxon rank-sum test). **D)** Differential

abundant genera identified between saliva from Chinese and Western samples. The panel on the left

indicates the standardised effect sizes (W statistic) estimated via the difference on relative abundance

using ANCOM-BC (taxa enriched in Western samples have a value shifted to right, whereas taxa

enriched in Chinese samples have a value shifted to left); The panel in the middle shows the relative

abundance of selected genera; the panel on the right indicates the Mean Decrease Accuracy of the

random forest model established. **E)** Spearman's correlation coefficients were calculated between

each pairwise comparison of differential genus and KEGG pathway. Only significantly correlated

comparisons (p<0.01, FDR adjusted Spearman's rank correlation) are displayed. The only Western-

enriched pathway is marked in pink.

Figure 7. An independent cohort verifies the definition of core microbiome membership

but cannot classify based on cultural background. A) Alluvial plot showing the affiliation of

ZOTUs to their originating core OTUs defined in the meta-analysis. **B)** and **C)** Comparison of salivary

microbial alpha diversity between the Chinese and Western samples, calculated by Shannon (B: $p =$

0.073, Wilcoxon rank-sum test) and Chao1 index (C: $p< 0.001$, Wilcoxon rank-sum test). **D)** and **E)**

The prediction of the cultural backgrounds of the samples according to the random forest

classification model constructed using the genus profiles of samples in the meta-analysis. The genus

level profiles of samples processed by **D)** closed-reference clustering with 97% sequence identity and

E) UNOISE3 denoising with 100% sequence identity were used as the test set.

Legends-Supplementary Figures

Figure S1 Average sequencing depth and rarefaction curve for the complete 16S rRNA dataset.

A) Mean read number of samples from each study. The dash line indicates that all samples below this

depth (depth = 2,000) have been removed. **B)** The rarefaction curve reflects the increase of sample's

Shannon index with sequencing depth. The curve was basically stabilised at sequencing depth = 2000.

Figure S2 The effect of different combinations between normalisations and distance matrices on

reducing the impact of hypervariable regions in the PCoA plot. PCoA plots showing Bray-Curtis

dissimilarity **(A-C)**, weighted uniFrac distance **(D-F)**, Euclidean distance **(G-J)** under rarefaction

(RAR) **(A, D, G)**, total-sum scaling (TSS) **(B, E, H)**, rarefied relative abundance transformation

(RRA) **(C, F, I)** and centred log-ratio transformation **(J)**. Percentage of variances explained by the

first two principal coordinates are shown on the axes.

Figure S3 Phylogenetic tree showing the presence and absence of top 500 OTUs with highest mean relative abundance. The colour strips on the innermost ring indicates which phylum the OTUs belong to. The presence of coloured circles on the 17 rings in the middle indicate that an OTU was found in a specific level of a sub-group. The grey bars on the outermost layer represent how many studies that an OTU presented. The scale lines are used to highlight the number of 10, 20 and 40.

Figure S4 The percentage contribution of the top-ranked OTUs to the beta-diversity of the dataset. The beta-diversity is calculated for the whole dataset and for only the top-ranked OTUs using both Bray-Curtis similarity (A) and weighted uniFrac distance (B). The percentage contribution of top-ranked OTUs is calculated by dividing the beta-diversity among top-ranked OTUs using the beta-diversity of the whole dataset. The dash lines indicate the last points at which the increase on the contribution is 2%. All OTUs before this point (the point itself was also included) were defined as "core".

Figure S5 Bacterial co-occurrence network verifies the role of identified core salivary microbiome members. Small modules with less than seven nodes were not displayed. The size of nodes is proportioned to the connectivity of nodes (node degree). Core OTUs from Figure 4 are indicated as yellow, while rare OTUs are in green. The edges between nodes represent the strength of the correlation (Spearman's correlation coefficient, $\rho \geq 0.5$, $p < 0.01$). The shape of "hub" OTUs are indicated as squares with the OTU name displayed.

Figure S6 The Betweenness and Closeness centrality of OTUs involved in the network analysis. The "hub" OTUs were identified as OTUs with either high connectivity (node degree) or centrality (betweenness **(A)** and closeness **(B)** centrality). The accession number of "hub" OTUs are indicated. The core OTUs are marked in yellow, while the rare OTUs were coloured in green.

Figure S7 The optimal number for defining the biomarker OTUs of four categories. The

contribution of the OTUs used to differentiate the levels in each category on ten-fold cross-validation.

The OTUs were ranked in the order of importance. The dash lines represent the point at which the

curve starts to become overall stable.

Figure S8 The top important OTUs identified by Random Forest classification model

established by the relative abundance of human salivary microbiota. The differential OTUs

defined were ranked in descending order of their importance (Mean Decrease Accuracy). The colour

of bars reflects the phylum level information of OTUs. The Mean Decrease Accuracy bar of core

OTUs were marked with bolded black borders.

Legends-Supplementary Tables

Table S1 The terms used for searching in databases, including Medline, EMBASE, and Web of

Science.

Table S2 Studies included after large-scale literature searches that met all the inclusion criteria.

Table S3 The full metadata used in this study, including 2206 samples with unique accession

numbers.

Table S4 The influence of seven factors at seven taxonomic levels on human salivary microbial

communities, measured by PERMANOVA with adonis2 function (permutation = 999).

PERMANOVA models were adjusted for study.

Table S5 Core OTUs of adults' saliva microbiome defined by **a)** method adapted from Wu et al.

(2019) ("1"= yes, "0"=no), **b)** method adapted from Shade and Stopnisek (2019).

Table S6 Genus with differential abundance between samples from Western and Chinese people

identified by both ANCOM-BC and Random Forest model, adjusted for hypervariable regions

sequenced.

Table S7 Species with differential abundance between samples from Western and Chinese people

identified by both ANCOM-BC and Random Forest model, adjusted for hypervariable regions

sequenced.

Table S8 KEGG pathways with differential abundance between samples from Western and Chinese

people identified by both ANCOM-BC and Random Forest model, adjusted for hypervariable regions

sequenced.

Table S9 The differences between Chinese and Western participants in the independent cohort,

measured by PERMANOVA with adonis2 function (permutation = 999). The PERMANOVA model

was adjusted for the gender and age range of participants.

Table 1 Studies included after large-scale literature searches that met all the inclusion criteria.

Study	Database	Accession Number	Location	Hypervariable Region	Number of Samples	Sample Type
[76]	SRA	PRJNA323410	India	V3-V4	51	Unstimulated
[77]	SRA	PRJNA577839	Europe	V3-V4	37	Stimulated
[78]	SRA	SRP125370	India	V3-V4	12	Other
[79]	SRA	PRJNA380250	United States	V1-V2/V4-V5	4	Unstimulated
[80]	ENA	PRJEB9010	Europe	V3-V4	70	Stimulated
[81]	SRA	PRJNA438728	India	V3-V4	30	Mouthwash
[82]	SRA	PRJNA321534	United States	V4	18	Unstimulated
[83]	SRA	PRJNA361501	Europe	V3-V4	99	Other
[84]	ENA	PRJEB37445	China	V3-V4	436	Stimulated
[57]	ENA	PRJEB11529	Canada	V3-V4	96	Stimulated
[85]	SRA	PRJNA609244	Europe	V1-V2	13	Unstimulated
[86]	SRA	PRJNA503603	China	V3-V4	24	Unstimulated
[87]	SRA	PRJNA495719	China	V3-V4	22	Unstimulated
[88]	ENA	PRJEB37299	Europe	V4	11	Stimulated
[89]	SRA	PRJNA386665	China	V4	127	Unstimulated
[90]	SRA	PRJNA578492	China	V3-V4	14	Unstimulated
[91]	SRA	PRJNA326866	Europe	V1-V2	20	Unstimulated
[92]	SRA	PRJNA484857	China	V3-V4	15	Unstimulated
[93]	ENA	PRJEB21767	Europe	V3-V4	47	Stimulated

[94]	SRA	PRJNA586897	China	V3-V4	64	Unstimulated
[95]	SRA	SRP113577	China	V4-V5	71	Unstimulated
[96]	SRA	PRJNA292800	United States	V4	28	Unstimulated
[97]	SRA	PRJNA602902	United States	V3-V4	119	Stimulated
[98]	SRA	PRJNA623352	Europe	V3-V4	17	Unstimulated
[99]	SRA	PRJNA356414	Europe	V3-V4	10	Unstimulated
[100]	SRA	PRJNA602902	United States	V3-V4	40	Stimulated
[101]	SRA	PRJNA587625	Qatar	V3-V4	73	Unstimulated
[102]	SRA	PRJNA612815	Europe	V1-V3/V3-V4/V4-V5/V6-V8	44	Other
[103]	SRA	PRJNA413706	United States	V4	30	Stimulated
[104]	SRA	PRJNA601054	China	V3-V4	27	Unstimulated
[105]	SRA	PRJNA321349	United States	V3-V4	20	Stimulated
[106]	SRA	PRJNA578951	China	V3-V4	22	Unstimulated
[107]	SRA	PRJNA539937	United States	V3-V4	25	Other
[108]	SRA	PRJNA421234	New Zealand	V3-V4	7	Unstimulated
[109]	Qiita	10823	United States	V4	150	Mouthwash
[110]	SRA	PRJNA306560	China	V4	18	Stimulated
[110]	SRA	PRJNA414355	China	V3-V4	37	Unstimulated
[111]	SRA	PRJNA587078/	China	V3-V4	21	Unstimulated
[112]	SRA	PRJNA556311	China	V3-V4	20	Unstimulated

[113]	ENA	PRJEB18476	Europe	V4	11	Stimulated
[114]	SRA	PRJNA414682	China	V3-V4	20	Unstimulated
[115]	SRA	PRJNA634162	United States	V4	75	Other
[116]	SRA	PRJNA515166	Malaysia	V3-V4	72	Unstimulated
[117]	SRA	PRJNA542018	China	V4	10	Unstimulated
[118]	SRA	PRJNA586723	China	V3-V4	60	Stimulated and unstimualted
[119]	SRA	PRJNA534340	China	V3-V4	120	Stimulated and unstimualted
[120]	SRA	PRJNA598080	Europe	V3-V4	12	Unstimulated

References

1. Mosca AC, Chen J. Food-saliva interactions: Mechanisms and implications. Trends Food Sci Technol. 2017;66:125-34. https://doi.org/https://doi.org/10.1016/j.tifs.2017.06.005.

2. Liu D, Jiang X, Zheng HJ, Xie B, Wang H, He T, et al., editors. The modularity of microbial interaction network in healthy human saliva: Stability and specificity. 2017 IEEE International Conference on Bioinformatics and Biomedicine (BIBM); 2017 13-16 Nov. 2017.

3. Acharya A, Chan Y, Kheur S, Jin LJ, Watt RM, Mattheos N. Salivary microbiome in non-oral disease: A summary of evidence and commentary. Arch Oral Biol. 2017;83:169-73. https://doi.org/https://doi.org/10.1016/j.archoralbio.2017.07.019.

4. Mark Welch JL, Rossetti BJ, Rieken CW, Dewhirst FE, Borisy GG. Biogeography of a human oral microbiome at the micron scale. Proceedings of the National Academy of Sciences. 2016;113(6):E791-E800. https://doi.org/10.1073/pnas.1522149113.

5. Shaw L, Ribeiro ALR, Levine AP, Pontikos N, Balloux F, Segal AW, et al. The Human Salivary Microbiome Is Shaped by Shared Environment Rather than Genetics: Evidence from a Large Family of Closely Related Individuals. mBio. 2017;8(5):e01237-17. https://doi.org/doi:10.1128/mBio.01237-17.

6. Muñoz-González C, Cueva C, Ángeles Pozo-Bayón M, Victoria Moreno-Arribas M. Ability of human oral microbiota to produce wine odorant aglycones from odourless grape glycosidic aroma precursors. Food Chem. 2015;187:112-9. https://doi.org/https://doi.org/10.1016/j.foodchem.2015.04.068.

7. Parker M, Onetto C, Hixson J, Bilogrevic E, Schueth L, Pisaniello L, et al. Factors Contributing to Interindividual Variation in Retronasal Odor Perception from Aroma Glycosides: The Role of Odorant Sensory Detection Threshold, Oral Microbiota, and Hydrolysis in Saliva. Journal of Agricultural and Food Chemistry. 2020;68(38):10299-309. https://doi.org/10.1021/acs.jafc.9b05450.

8. Piombino P, Genovese A, Esposito S, Moio L, Cutolo PP, Chambery A, et al. Saliva from Obese Individuals Suppresses the Release of Aroma Compounds from Wine. PLoS One. 2014;9(1):e85611. https://doi.org/10.1371/journal.pone.0085611.

9. De Filippis F, Parente E, Zotta T, Ercolini D. A comparison of bioinformatic approaches for 16S rRNA gene profiling of food bacterial microbiota. Int J Food Microbiol. 2018;265:9-17. https://doi.org/https://doi.org/10.1016/j.ijfoodmicro.2017.10.028.

10. Soriano-Lerma A, Pérez-Carrasco V, Sánchez-Marañón M, Ortiz-González M, Sánchez-Martín V, Gijón J, et al. Influence of 16S rRNA target region on the outcome of microbiome studies in soil and saliva samples. Sci Rep. 2020;10(1):13637. https://doi.org/10.1038/s41598-020-70141-8.

11. Zaura E, Keijser BJF, Huse SM, Crielaard W. Defining the healthy "core microbiome" of oral microbial communities. BMC Microbiol. 2009;9(1):259. https://doi.org/10.1186/1471-2180-9-259.

12. Risely A. Applying the core microbiome to understand host–microbe systems. J Anim Ecol. 2020;89(7):1549-58. https://doi.org/https://doi.org/10.1111/1365-2656.13229.

13. Wu L, Ning D, Zhang B, Li Y, Zhang P, Shan X, et al. Global diversity and biogeography of bacterial communities in wastewater treatment plants. Nat Microbiol. 2019;4(7):1183-95. https://doi.org/10.1038/s41564-019-0426-5.

14. Guo R, Zheng Y, Zhang L, Shi J, Li W. Salivary microbiome and periodontal status of patients with periodontitis during the initial stage of orthodontic treatment. Am J Orthod Dentofacial Orthop. 2021;159(5):644-52. https://doi.org/https://doi.org/10.1016/j.ajodo.2019.11.026.

15.   Relvas M, Regueira-Iglesias A, Balsa-Castro C, Salazar F, Pacheco JJ, Cabral C, et al.
Relationship between dental and periodontal health status and the salivary microbiome:
bacterial diversity, co-occurrence networks and predictive models. Sci Rep. 2021;11(1):929.
https://doi.org/10.1038/s41598-020-79875-x.
16.   Wang K, Lu W, Tu Q, Ge Y, He J, Zhou Y, et al. Preliminary analysis of salivary
microbiome and their potential roles in oral lichen planus. Sci Rep. 2016;6(1):22943.
https://doi.org/10.1038/srep22943.
17.   Sabharwal A, Ganley K, Miecznikowski JC, Haase EM, Barnes V, Scannapieco FA.
The salivary microbiome of diabetic and non-diabetic adults with periodontal disease. J
Periodontol. 2019;90(1):26-34. https://doi.org/https://doi.org/10.1002/JPER.18-0167.
18.   Torres PJ, Fletcher EM, Gibbons SM, Bouvet M, Doran KS, Kelley ST.
Characterization of the salivary microbiome in patients with pancreatic cancer. PeerJ.
2015;3:e1373.
19.   Lindheim L, Bashir M, Münzker J, Trummer C, Zachhuber V, Pieber TR, et al. The
Salivary Microbiome in Polycystic Ovary Syndrome (PCOS) and Its Association with
Disease-Related Parameters: A Pilot Study. Front Microbiol. 2016;7(1270).
https://doi.org/10.3389/fmicb.2016.01270.
20.   Nomura Y, Kakuta E, Okada A, Otsuka R, Shimada M, Tomizawa Y, et al. Oral
Microbiome in Four Female Centenarians. Applied Sciences. 2020;10(15):5312.
21.   Hansen TH, Kern T, Bak EG, Kashani A, Allin KH, Nielsen T, et al. Impact of a
vegan diet on the human salivary microbiota. Sci Rep. 2018;8(1):5847.
https://doi.org/10.1038/s41598-018-24207-3.
22.   Kato I, Vasquez A, Moyerbrailean G, Land S, Djuric Z, Sun J, et al. Nutritional
Correlates of Human Oral Microbiome. J Am Coll Nutr. 2017;36(2):88-98.
https://doi.org/10.1080/07315724.2016.1185386.
23.   Liu K, Chen S, Huang J, Ren F, Yang T, Long D, et al. Oral Microbiota of Children Is
Conserved across Han, Tibetan and Hui Groups and Is Correlated with Diet and Gut
Microbiota. Microorganisms. 2021;9(5):1030.
24.   Renson A, Jones HE, Beghini F, Segata N, Zolnik CP, Usyk M, et al.
Sociodemographic variation in the oral microbiome. Ann Epidemiol. 2019;35:73-80.e2.
https://doi.org/https://doi.org/10.1016/j.annepidem.2019.03.006.
25.   Wu J, Peters BA, Dominianni C, Zhang Y, Pei Z, Yang L, et al. Cigarette smoking
and the oral microbiome in a large study of American adults. The ISME Journal.
2016;10(10):2435-46. https://doi.org/10.1038/ismej.2016.37.
26.   Fan X, Peters BA, Jacobs EJ, Gapstur SM, Purdue MP, Freedman ND, et al. Drinking
alcohol is associated with variation in the human oral microbiome in a large study of
American adults. Microbiome. 2018;6(1):59. https://doi.org/10.1186/s40168-018-0448-x.
27.   Takayasu L, Suda W, Takanashi K, Iioka E, Kurokawa R, Shindo C, et al. Circadian
oscillations of microbial and functional composition in the human salivary microbiome. DNA
Res. 2017;24(3):261-70. https://doi.org/10.1093/dnares/dsx001.
28.   Wu Y, Chi X, Zhang Q, Chen F, Deng X. Characterization of the salivary microbiome
in people with obesity. PeerJ. 2018;6:e4458.
29.   Jo R, Nishimoto Y, Umezawa K, Yama K, Aita Y, Ichiba Y, et al. Comparison of oral
microbiome profiles in stimulated and unstimulated saliva, tongue, and mouth-rinsed water.
Sci Rep. 2019;9(1):16124. https://doi.org/10.1038/s41598-019-52445-6.
30.   Cattaneo C, Riso P, Laureati M, Gargari G, Pagliarini E. Exploring Associations
between Interindividual Differences in Taste Perception, Oral Microbiota Composition, and
Reported Food Intake. Nutrients. 2019;11(5):1167.

31.   Gardner A, So PW, Carpenter GH. Intraoral Microbial Metabolism and Association
with Host Taste Perception. J Dent Res. 2020;99(6):739-45.
https://doi.org/10.1177/0022034520917142.
32.   Henne K, Li J, Stoneking M, Kessler O, Schilling H, Sonanini A, et al. Global
analysis of saliva as a source of bacterial genes for insights into human population structure
and migration studies. BMC Evol Biol. 2014;14(1):190. https://doi.org/10.1186/s12862-014-
0190-3.
33.   Li J, Quinque D, Horz H-P, Li M, Rzhetskaya M, Raff JA, et al. Comparative analysis
of the human saliva microbiome from different climate zones: Alaska, Germany, and Africa.
BMC Microbiol. 2014;14(1):316. https://doi.org/10.1186/s12866-014-0316-1.
34.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nat Biotechnol. 2019;37(8):852-7. https://doi.org/10.1038/s41587-019-0209-9.
35.   Chen T, Yu W-H, Izard J, Baranova OV, Lakshmanan A, Dewhirst FE. The Human
Oral Microbiome Database: a web accessible resource for investigating oral microbe
taxonomic and genomic information. Database. 2010;2010.
https://doi.org/10.1093/database/baq013.
36.   Romano S, Savva GM, Bedarf JR, Charles IG, Hildebrand F, Narbad A. Meta-
analysis of the Parkinson's disease gut microbiome suggests alterations linked to intestinal
inflammation. npj Parkinson's Disease. 2021;7(1):27. https://doi.org/10.1038/s41531-021-
00156-z.
37.   Palarea-Albaladejo J, Martín-Fernández JA. zCompositions — R package for
multivariate imputation of left-censored data under a compositional approach. Chemometrics
Intellig Lab Syst. 2015;143:85-96.
https://doi.org/https://doi.org/10.1016/j.chemolab.2015.02.019.
38.   Oksanen J, Guillaume; BF, Michael; F, Roeland; K, Pierre; L, Dan; M, et al. vegan:
Community Ecology Package. R package version 2.5-7.; 2020.
39.   Shade A, Stopnisek N. Abundance-occupancy distributions to prioritize plant core
microbiome membership. Curr Opin Microbiol. 2019;49:50-8.
https://doi.org/https://doi.org/10.1016/j.mib.2019.09.008.
40.   Frank E Harrell Jr wcfCDamo. Hmisc: Harrell Miscellaneous. R package version 4.5-
0. 2021. https://CRAN.R-project.org/package=Hmisc.
41.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome
Res. 2003;13(11):2498-504.
42.   Liaw A, Wiener M. Classification and regression by randomForest. R news.
2002;2(3):18-22.
43.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.
https://ggplot2.tidyverse.org.
44.   Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction.
Nature Communications. 2020;11(1):3514. https://doi.org/10.1038/s41467-020-17041-7.
45.   Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. Data,
information, knowledge and principle: back to metabolism in KEGG. Nucleic Acids Res.
2013;42(D1):D199-D205. https://doi.org/10.1093/nar/gkt1076.
46.   Wei T, Simko V. R package 'corrplot': Visualization of a Correlation Matrix (Version
0.90). 2021. https://github.com/taiyun/corrplot.
47.   Marotz C, Amir A, Humphrey G, Gaffney J, Gogul G, Knight R. DNA extraction for
streamlined metagenomics of diverse environmental samples. BioTechniques.
2017;62(6):290-3. https://doi.org/10.2144/000114559.

48.   Edgar RC. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon
sequencing. bioRxiv. 2016:081257. https://doi.org/10.1101/081257.
49.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res. 2013;41(Database issue):D590-D6. https://doi.org/10.1093/nar/gks1219.
50.   Stopnisek N, Shade A. Persistent microbiome members in the common bean
rhizosphere: an integrated analysis of space, time, and plant genotype. The ISME Journal.
2021;15(9):2708-22. https://doi.org/10.1038/s41396-021-00955-5.
51.   Domin H, Zurita-Gutiérrez YH, Scotti M, Buttlar J, Hentschel Humeida U, Fraune S.
Predicted Bacterial Interactions Affect in Vivo Microbial Colonization Dynamics in
Nematostella. Front Microbiol. 2018;9(728). https://doi.org/10.3389/fmicb.2018.00728.
52.   Fan P, Nelson CD, Driver JD, Elzo MA, Peñagaricano F, Jeong KC. Host genetics
exerts lifelong effects upon hindgut microbiota and its association with bovine growth and
immunity. The ISME Journal. 2021;15(8):2306-21. https://doi.org/10.1038/s41396-021-
00925-x.
53.   Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A, Eldridge DJ,
Bardgett RD, et al. A global atlas of the dominant bacteria found in soil. Science.
2018;359(6373):320-5. https://doi.org/doi:10.1126/science.aap9516.
54.   Wang Y, Gong J, Li J, Xin Y, Hao Z, Chen C, et al. Insights into bacterial diversity in
compost: Core microbiome and prevalence of potential pathogenic bacteria. Sci Total
Environ. 2020;718:137304. https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.137304.
55.   Giordano C, Boscaro V, Munz G, Mori G, Vannini C. Summer holidays as break-
point in shaping a tannery sludge microbial community around a stable core microbiota. Sci
Rep. 2016;6(1):30376. https://doi.org/10.1038/srep30376.
56.   Salonen A, Salojärvi J, Lahti L, de Vos WM. The adult intestinal core microbiota is
determined by analysis depth and health status. Clinical Microbiology and Infection.
2012;18(s4):16-20. https://doi.org/https://doi.org/10.1111/j.1469-0691.2012.03855.x.
57.   Hall MW, Singh N, Ng KF, Lam DK, Goldberg MB, Tenenbaum HC, et al. Inter-
personal diversity and temporal dynamics of dental, tongue, and salivary microbiota in the
healthy oral cavity. npj Biofilms and Microbiomes. 2017;3(1):2.
https://doi.org/10.1038/s41522-016-0011-0.
58.   Simón-Soro Á, Tomás I, Cabrera-Rubio R, Catalan MD, Nyvad B, Mira A. Microbial
Geography of the Oral Cavity. J Dent Res. 2013;92(7):616-21.
https://doi.org/10.1177/0022034513488119.
59.   Yao T, Han X, Guan T, Zhai C, Liu C, Liu C, et al. Exploration of the microbiome
community for saliva, skin, and a mixture of both from a population living in Guangdong. Int
J Legal Med. 2021;135(1):53-62. https://doi.org/10.1007/s00414-020-02329-6.
60.   Oliveira SG, Nishiyama RR, Trigo CAC, Mattos-Guaraldi AL, Dávila AMR, Jardim
R, et al. Core of the saliva microbiome: an analysis of the MG-RAST data. BMC Oral Health.
2021;21(1):351. https://doi.org/10.1186/s12903-021-01719-5.
61.   Conrads G, Westenberger J, Lürkens M, Abdelbary MMH. Isolation and Bacteriocin-
Related Typing of Streptococcus dentisani. Frontiers in Cellular and Infection Microbiology.
2019;9(110). https://doi.org/10.3389/fcimb.2019.00110.
62.   López-López A, Camelo-Castillo A, Ferrer MD, Simon-Soro Á, Mira A. Health-
Associated Niche Inhabitants as Oral Probiotics: The Case of Streptococcus dentisani. Front
Microbiol. 2017;8(379). https://doi.org/10.3389/fmicb.2017.00379.
63.   Zarco M, Vess T, Ginsburg G. The oral microbiome in health and disease and the
potential impact on personalized dental medicine. Oral Dis. 2012;18(2):109-20.
https://doi.org/https://doi.org/10.1111/j.1601-0825.2011.01851.x.

64.   Cornejo-Granados F, Gallardo-Becerra L, Leonardo-Reza M, Ochoa-Romo JP,
Ochoa-Leyva A. A meta-analysis reveals the environmental and host factors shaping the
structure and function of the shrimp microbiota. PeerJ. 2018;6:e5382.
65.   Holman DB, Brunelle BW, Trachsel J, Allen HK, Bik H. Meta-analysis To Define a
Core Microbiota in the Swine Gut. mSystems. 2017;2(3):e00004-17.
https://doi.org/doi:10.1128/mSystems.00004-17.
66.   Wright RJ, Langille MGI, Walker TR. Food or just a free ride? A meta-analysis
reveals the global diversity of the Plastisphere. ISME J. 2021;15(3):789-806.
https://doi.org/10.1038/s41396-020-00814-9.
67.   Roberts FA, Darveau RP. Microbial protection and virulence in periodontal tissue as a
function of polymicrobial communities: symbiosis and dysbiosis. Periodontol 2000.
2015;69(1):18-27. https://doi.org/https://doi.org/10.1111/prd.12087.
68.   Sato N, Kakuta M, Hasegawa T, Yamaguchi R, Uchino E, Kobayashi W, et al.
Metagenomic analysis of bacterial species in tongue microbiome of current and never
smokers. npj Biofilms and Microbiomes. 2020;6(1):11. https://doi.org/10.1038/s41522-020-
0121-6.
69.   Vogtmann E, Flores R, Yu G, Freedman ND, Shi J, Gail MH, et al. Association
between tobacco use and the upper gastrointestinal microbiome among Chinese men. Cancer
Causes Control. 2015;26(4):581-8. https://doi.org/10.1007/s10552-015-0535-2.
70.   Karabudak S, Ari O, Durmaz B, Dal T, Basyigit T, Kalcioglu MT, et al. Analysis of
the effect of smoking on the buccal microbiome using next-generation sequencing
technology. J Med Microbiol. 2019;68(8):1148-58.
https://doi.org/https://doi.org/10.1099/jmm.0.001003.
71.   Prasoodanan P. K V, Sharma AK, Mahajan S, Dhakan DB, Maji A, Scaria J, et al.
Western and non-western gut microbiomes reveal new roles of Prevotella in carbohydrate
metabolism and mouth–gut axis. npj Biofilms and Microbiomes. 2021;7(1):77.
https://doi.org/10.1038/s41522-021-00248-x.
72.   García-Estévez I, Ramos-Pineda AM, Escribano-Bailón MT. Interactions between
wine phenolic compounds and human saliva in astringency perception. Food Funct.
2018;9(3):1294-309.
73.   Liu Z, Guo H, Zhang W, Ni L. Salivary Microbiota Shifts under Sustained
Consumption of Oolong Tea in Healthy Adults. Nutrients. 2020;12(4):966.
74.   Vangay P, Johnson AJ, Ward TL, Al-Ghalith GA, Shields-Cutler RR, Hillmann BM,
et al. US Immigration Westernizes the Human Gut Microbiome. Cell. 2018;175(4):962-
72.e10. https://doi.org/https://doi.org/10.1016/j.cell.2018.10.029.
75.   Peters BA, Yi SS, Beasley JM, Cobbs EN, Choi HS, Beggs DB, et al. US nativity and
dietary acculturation impact the gut microbiome in a diverse US population. The ISME
Journal. 2020;14(7):1639-50. https://doi.org/10.1038/s41396-020-0630-6.
76.   Acharya A, Chan Y, Kheur S, Kheur M, Gopalakrishnan D, Watt R, et al. Salivary
microbiome of an urban Indian cohort and patterns linked to subclinical inflammation. Oral
diseases. 2017;23(7):926-40.
77.   Anderson A, Al-Ahmad A, Schlueter N, Frese C, Hellwig E, Binder N. Influence of
the long-term use of oral hygiene products containing stannous ions on the salivary
microbiome–a randomized controlled trial. Scientific reports. 2020;10(1):1-8.
78.   Bhushan B, Yadav A, Singh S, Ganju L. Diversity and functional analysis of salivary
microflora of Indian Antarctic expeditionaries. Journal of oral microbiology.
2019;11(1):1581513.
79.   Cabral DJ, Wurster JI, Flokas ME, Alevizakos M, Zabat M, Korry BJ, et al. The
salivary microbiome is consistent between subjects and resistant to impacts of short-term

hospitalization. Scientific Reports. 2017;7(1):11040. https://doi.org/10.1038/s41598-017-
11427-2.

80.   Cameron SJS, Huws SA, Hegarty MJ, Smith DPM, Mur LAJ. The human salivary
microbiome exhibits temporal stability in bacterial diversity. FEMS Microbiology Ecology.
2015;91(9). https://doi.org/10.1093/femsec/fiv091.

81.   Chaudhari DS, Dhotre DP, Agarwal DM, Gaike AH, Bhalerao D, Jadhav P, et al. Gut,
oral and skin microbiome of Indian patrilineal families reveal perceptible association with
age. Scientific Reports. 2020;10(1):5685. https://doi.org/10.1038/s41598-020-62195-5.

82.   Chen C, Hemme C, Beleno J, Shi ZJ, Ning D, Qin Y, et al. Oral microbiota of
periodontal health and disease and their changes after nonsurgical periodontal therapy. The
ISME Journal. 2018;12(5):1210-24. https://doi.org/10.1038/s41396-017-0037-1.

83.   Collado MC, Engen PA, Bandín C, Cabrera-Rubio R, Voigt RM, Green SJ, et al.
Timing of food intake impacts daily rhythms of human salivary microbiota: a randomized,
crossover study. The FASEB Journal. 2018;32(4):2060-72.

84.   Debelius JW, Huang T, Cai Y, Ploner A, Barrett D, Zhou X, et al. Subspecies niche
specialization in the oral microbiome is associated with nasopharyngeal carcinoma risk.
Msystems. 2020;5(4):e00065-20.

85.   Hijazi K, Morrison RW, Mukhopadhya I, Martin B, Gemmell M, Shaw S, et al. Oral
bacterial diversity is inversely correlated with mucosal inflammation. Oral Diseases.
2020;26(7):1566-75. https://doi.org/https://doi.org/10.1111/odi.13420.

86.   Ji Y, Liang X, Lu H. Analysis of by high-throughput sequencing: Helicobacter pylori
infection and salivary microbiome. BMC Oral Health. 2020;20(1):84.
https://doi.org/10.1186/s12903-020-01070-1.

87.   Jiang Q, Liu J, Chen L, Gan N, Yang D. The Oral Microbiome in the Elderly With
Dental Caries and Health. Frontiers in cellular and infection microbiology. 2019;8(442).
https://doi.org/10.3389/fcimb.2018.00442.

88.   Kumpitsch C, Moissl-Eichinger C, Pock J, Thurnher D, Wolf A. Preliminary insights
into the impact of primary radiochemotherapy on the salivary microbiome in head and neck
squamous cell carcinoma. Scientific reports. 2020;10(1):1-12.

89.   Lee W-H, Chen H-M, Yang S-F, Liang C, Peng C-Y, Lin F-M, et al. Bacterial
alterations in salivary microbiota and their association in oral cancer. Scientific reports.
2017;7(1):1-11.

90.   Lin M, Li X, Wang J, Cheng C, Zhang T, Han X, et al. Saliva microbiome changes in
patients with periodontitis with and without chronic obstructive pulmonary disease. Frontiers
in cellular infection microbiology. 2020;10:124.

91.   Lindheim L, Bashir M, Münzker J, Trummer C, Zachhuber V, Pieber TR, et al. The
salivary microbiome in polycystic ovary syndrome (pcos) and its association with disease-
related parameters: a pilot study. Frontiers in microbiology. 2016;7:1270.

92.   Liu Y, Zhang Q, Hu X, Chen F, Hua H. Characteristics of the salivary microbiota in
cheilitis granulomatosa. Med Oral Patol Oral Cir Bucal. 2019;24(6):e719-e25.
https://doi.org/10.4317/medoral.23041.

93.   Lundmark A, Hu YO, Huss M, Johannsen G, Andersson AF, Yucel-Lindberg T.
Identification of salivary microbiota and its association with host inflammatory mediators in
periodontitis. Frontiers in cellular infection microbiology. 2019;9:216.

94.   Menon R, Gomez A, Brandt B, Leung Y, Gopinath D, Watt R, et al. Long-term
impact of oral surgery with or without amoxicillin on the oral microbiome-A prospective
cohort study. Scientific reports. 2019;9(1):1-10.

95.   Niu C, Dong T, Jiang W, Gao L, Yuan K, Hu X, et al. Pregnancy-Related Ecological
Shifts of Salivary Microbiota and its Association with Salivary Sex Hormones. 2020.

96. Ozga AT, Sankaranarayanan K, Tito RY, Obregon-Tito AJ, Foster MW, Tallbull G, et al. Oral microbiome diversity among Cheyenne and Arapaho individuals from Oklahoma. American journal of physical anthropology. 2016;161(2):321-7.

97. Pushalkar S, Paul B, Li Q, Yang J, Vasconcelos R, Makwana S, et al. Electronic Cigarette Aerosol Modulates the Oral Microbiome and Increases Risk of Infection. iScience. 2020;23(3):100884. https://doi.org/https://doi.org/10.1016/j.isci.2020.100884.

98. Relvas M, Regueira-Iglesias A, Balsa-Castro C, Salazar F, Pacheco JJ, Cabral C, et al. Assessing the impact of dental and periodontal statuses on the salivary microbiome: a global oral health scale. medRxiv. 2020.

99. Russo E, Bacci G, Chiellini C, Fagorzi C, Niccolai E, Taddei A, et al. Preliminary comparison of oral and intestinal human microbiota in patients with colorectal cancer: a pilot study. Frontiers in microbiology. 2018;8:2699.

100. Sierra MA, Li Q, Pushalkar S, Paul B, Sandoval TA, Kamer AR, et al. The influences of bioinformatics tools and reference databases in analyzing the human oral microbial community. Genes. 2020;11(8):878.

101. Sohail MU, Elrayess MA, Al Thani AA, Al-Asmakh M, Yassine HM. Profiling the oral microbiome and plasma biochemistry of obese hyperglycemic subjects in Qatar. Microorganisms. 2019;7(12):645.

102. Soriano-Lerma A, Pérez-Carrasco V, Sánchez-Marañón M, Ortiz-González M, Sánchez-Martín V, Gijón J, et al. Influence of 16S rRNA target region on the outcome of microbiome studies in soil and saliva samples. Scientific reports. 2020;10(1):1-13.

103. Stewart CJ, Auchtung TA, Ajami NJ, Velasquez K, Smith DP, De La Garza II R, et al. Effects of tobacco smoke and electronic cigarette vapor exposure on the oral and gut microbiota in humans: a pilot study. PeerJ. 2018;6:e4693.

104. Sun X, Li M, Xia L, Fang Z, Yu S, Gao J, et al. Alteration of salivary microbiome in periodontitis with or without type-2 diabetes mellitus and metformin treatment. Scientific reports. 2020;10(1):1-14.

105. Tian N, Faller L, Leffler DA, Kelly CP, Hansen J, Bosch JA, et al. Salivary gluten degradation and oral microbial profiles in healthy individuals and celiac disease patients. Applied and environmental microbiology. 2017;83(6):e03330-16.

106. Tong Y, Zheng L, Qing P, Zhao H, Li Y, Su L, et al. Oral microbiota perturbations are linked to high risk for rheumatoid arthritis. Frontiers in cellular infection microbiology. 2020;9:475.

107. Urbaniak C, Lorenzi H, Thissen J, Jaing C, Crucian B, Sams C, et al. The influence of spaceflight on the astronaut salivary microbiome and the search for a microbiome biomarker for viral reactivation. Microbiome. 2020;8(1):1-14.

108. Vesty A, Gear K, Biswas K, Radcliff FJ, Taylor MW, Douglas RG. Microbial and inflammatory-based salivary biomarkers of head and neck squamous cell carcinoma. Clinical and experimental dental research. 2018;4(6):255-62. https://doi.org/https://doi.org/10.1002/cre2.139.

109. Vogtmann E, Chen J, Kibriya MG, Amir A, Shi J, Chen Y, et al. Comparison of oral collection methods for studies of microbiota. Cancer Epidemiology and Prevention Biomarkers. 2019;28(1):137-43.

110. Wang T, Yu L, Xu C, Pan K, Mo M, Duan M, et al. Chronic fatigue syndrome patients have alterations in their oral microbiome composition and function. PloS one. 2018;13(9):e0203503.

111. Wang Q, Rao Y, Guo X, Liu N, Liu S, Wen P, et al. Oral microbiome in patients with oesophageal squamous cell carcinoma. Scientific reports. 2019;9(1):1-9.

112. Wang X, Zhao Z, Tang N, Zhao Y, Xu J, Li L, et al. Microbial community analysis of
saliva and biopsies in patients with oral lichen planus. Frontiers in microbiology.
2020;11:629.
113. Wolf A, Moissl-Eichinger C, Perras A, Koskinen K, Tomazic PV, Thurnher D. The
salivary microbiome as an indicator of carcinogenesis in patients with oropharyngeal
squamous cell carcinoma: A pilot study. Scientific reports. 2017;7(1):1-10.
114. Xun Z, Zhang Q, Xu T, Chen N, Chen F. Dysbiosis and Ecotypes of the Salivary
Microbiome Associated With Inflammatory Bowel Diseases and the Assistance in Diagnosis
of Diseases Using Oral Bacterial Profiles. Frontiers in microbiology. 2018;9(1136).
https://doi.org/10.3389/fmicb.2018.01136.
115. Yano Y, Hua X, Wan Y, Suman S, Zhu B, Dagnall CL, et al. Comparison of Oral
Microbiota Collected Using Multiple Methods and Recommendations for New
Epidemiologic Studies. Msystems. 2020;5(4):e00156-20.
116. Yeo L-F, Aghakhanian FF, Tan JS, Gan HM, Phipps ME. Health and saliva
microbiomes of a semi-urbanized indigenous tribe in Peninsular Malaysia. F1000Research.
2019;8.
117. Yu FY, Wang QQ, Li M, Cheng Y-H, Cheng Y-SL, Zhou Y, et al. Dysbiosis of saliva
microbiome in patients with oral lichen planus. BMC microbiology. 2020;20(1):1-12.
118. Zhu C, Yuan C, Wei F-Q, Sun X-Y, Zheng S-G. Comparative evaluation of
peptidome and microbiota in different types of saliva samples. Ann Transl Med.
2020;8(11):686-. https://doi.org/10.21037/atm-20-393.
119. Zhu C, Yuan C, Wei FQ, Sun XY, Zheng SG. Intraindividual Variation and Personal
Specificity of Salivary Microbiota. J Dent Res. 2020;99(9):1062-71.
https://doi.org/10.1177/0022034520917155.
120. Ziganshina EE, Sagitov II, Akhmetova RF, Saleeva GT, Kiassov AP, Gogoleva NE,
et al. Comparison of the Microbiota and Inorganic Anion Content in the Saliva of Patients
with Gastroesophageal Reflux Disease and Gastroesophageal Reflux Disease-Free
Individuals. BioMed Research International. 2020;2020:2681791.
https://doi.org/10.1155/2020/2681791.
1135

Figure 1. Overview of literature search procedure and metadata of included studies. a) Large-scale literature searching and data filtering process, followed by the number of samples submitted to the bioinformatic analyses; b) The locations of studies, the scale of symbols that reflect the number of samples of each study; c) Distribution of metadata categories.

Figure 2. Summary of taxonomic composition and alpha diversity of included studies. A) The mean community composition of each study at the phylum level; The alpha-diversity measured by **B)** Shannon index; **C)** Chao 1 index; **D)** Simpson's index, the colour of boxes stands for the geographic location of the studies. The horizontal bars within boxes represent medians. The tops and bottoms of boxes represent the 75th and 25th percentiles, respectively.

Figure 3. The variability in human salivary microbiota have been explained by different factors. Among them, hypervariable regions and geographic locations have the largest impact. The effect of the categories on the clustering of the sample was measured using PERMANOVA at four taxonomic levels: family **(A),** Genus **(B),** species **(C)** and OTU level **(D).** The colour indicates the different combinations of normalisation (TSS, Total-sum scaling; RRA, Rarefied relative abundance; CLR, Centred log ratio) and indices (BC, Bray-Curtis; EUC, Euclidean; wUF, weighted uniFrac). Because the results of rarefication (RAR) were very close to TSS and RRA, they were not displayed in the figures. Principal coordinate analysis (PCoA) with Bray-Curtis **(E)** and weighted uniFrac **(F)** showing the differences between samples from North America, Europe, and China.

Figure 4. The core OTUs defined by abundance-occupancy pattern. A) Venn diagram showing the interaction between three methods used to define the core. Sixty-eight OTUs were defined as the core for all methods. (MRA+OCC: The thresholds were setting on mean relative abundance and occupancy to define the core; BC: The method adapted from Shade and Stopnisek using Bray-Curtis similarity; wUF: The method adapted from Shade and Stopnisek using weighted uniFrac distance). **B)** Pie chart showing the number of the core (pink) versus other OTUs (blue) identified in percentage. **C)** Pie chart showing the relative abundance of the core and other OTUs across all samples. **D)** Relative abundance of 68 core OTUs across subgroups classified by seven categories. **E)** Heatmap showing the log-transformed mean relative abundance of each core OTU at each level of different categories.

Figure 5. Salivary microbiome members which significantly contribute to categorisation of meta data. Random Forest models showed the impact of categories on salivary microbiome and the core OTUs contributing to accuracy of these models. A) Error rate (%) for the random forest classifications conducted with samples grouped by eight different categories. **B)** Phylogenetic tree indicates the taxonomic information of 68 core OTUs. The coloured squares between the tree and the annotation of phylum indicate the OTUs that were defined by the Random Forest model as "important" for distinguishing between different levels in each category. The bars on the outmost ring showing the mean relative abundance of each OTU.

Figure 6. Distinct microbial profiles are evident in the saliva samples from Chinese and Western adults. A) Taxonomic hierarchies show the relative enrichment of taxa in three geographic locations at phylum through species level. Coloured nodes indicate log2-fold increase in median abundance of the group in x-axis (pink) or y-axis (blue). Only taxa showed significant changes (false discovery rate-adjusted Wilcoxon rank sum q < 0.05) are displayed. **B)** and **C)** Comparison of salivary microbial alpha diversity between the Chinese and Western samples, calculated by Chao1 (B: *p*< 0.001, Wilcoxon rank-sum test) and Shannon index (C: *p*< 0.001, Wilcoxon rank-sum test). **D)** Differential abundant genera identified between saliva from Chinese and Western samples. The panel on the left indicates the standardised effect sizes (W statistic) estimated via the difference on relative abundance using ANCOM-BC (taxa enriched in Western samples have a value shifted to right, whereas taxa enriched in Chinese samples have a value shifted to left); The panel in the middle shows the relative abundance of selected genera; the panel on the right indicates the Mean Decrease Accuracy of the random forest model established. **E)** Spearman's correlation coefficients were calculated between each pairwise comparison of differential genus and KEGG pathway. Only significantly correlated comparisons (p<0.01, FDR adjusted Spearman's rank correlation) are displayed. The only Western-enriched pathway is marked in pink.

Figure 7. Validation of the findings in the meta-analysis in an independent cohort. A) Alluvial plot showing the affiliation of ZOTUs to their originating core OTUs defined in the meta-analysis. **B)** and **C)** Comparison of salivary microbial alpha diversity between the Chinese and Western samples, calculated by Shannon (B: $p = 0.073$, Wilcoxon rank-sum test) and Chao1 index (C: $p < 0.001$, Wilcoxon rank-sum test). **D)** and **E)** The prediction of the cultural backgrounds of the samples according to the random forest classification model constructed using the genus profiles of samples in the meta-analysis. The genus level profiles of samples processed by **D)** closed-reference clustering with 97% sequence identity and **E)** UNOISE3 denoising with 100% sequence identity were used as the test set.