An organism-wide atlas of hormonal signaling based on the mouse lemur single-cell transcriptome

Shixuan Liu[1, 2, 3], Camille Ezran[2, 3], Michael F. Z. Wang[4], Zhengda Li[1], The Tabula Microcebus Consortium[11], Jonathon Z. Long[5, 6], Iwijn De Vlaminck[4], Sheng Wang[7], Christin Kuo[8], Jacques Epelbaum[9,10], Jeremy Terrien[9], Mark A. Krasnow[2, 3]*, James E. Ferrell, Jr.[1, 2]*

1. Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA, USA;
2. Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA;
3. Howard Hughes Medical Institute, Stanford University School of Medicine, Stanford, CA, USA;
4. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA;
5. Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA;
6. Stanford ChEM-H, Stanford University, Stanford, Stanford, CA, USA;
7. Paul G. Allen School of Computer Science & Engineering, University of Washington, Seattle, WA, USA;
8. Department of Pediatrics, Stanford University School of Medicine, Stanford, CA, USA;
9. Unité Mécanismes Adaptatifs et Evolution (MECADEV), UMR 7179, CNRS, Museum National d'Histoire Naturelle, Brunoy, France;
10. Unité Mixte de Recherche en Santé 894 INSERM, Centre de Psychiatrie et Neurosciences, Université Paris Descartes, Sorbonne, Paris, France;
11. See the full list of the consortium members and their affiliations at the end of the manuscript.

*Corresponding authors, james.ferrell@stanford.edu, krasnow@stanford.edu

Abstract

Hormones coordinate long-range cell-cell communication in multicellular organisms and play vital roles in normal physiology, metabolism, and health. Using the newly-completed organism-wide single cell transcriptional atlas of a non-human primate, the mouse lemur (*Microcebus murinus*), we have systematically identified hormone-producing and -target cells for 87 classes of hormones, and have created a browsable atlas for hormone signaling that reveals previously unreported sites of hormone regulation and species-specific rewiring. Hormone ligands and receptors exhibited cell-type-dependent, stereotypical expression patterns, and their transcriptional profiles faithfully classified the discrete cell types defined by the full transcriptome, despite their comprising less than 1% of the transcriptome. Although individual cell types generally exhibited the same characteristic patterns of hormonal gene expression, a number of examples of similar or seemingly-identical cell types (e.g., endothelial cells of the lung versus of other organs) displayed different hormonal gene expression patterns. By linking ligand-expressing cells to the cells expressing the corresponding receptor, we constructed an organism-wide map of the hormonal cell-cell communication network. The hormonal cell-cell network was remarkably densely and robustly connected, and included classical hierarchical circuits (e.g. pituitary → peripheral endocrine gland → diverse cell types) as well as examples of highly distributed control. The network also included both well-known examples of feedback loops and a long list of potential novel feedback circuits. This primate hormone atlas provides a powerful resource to facilitate discovery of regulation on an organism-wide scale and at single-cell resolution, complementing the single-site-focused strategy of classical endocrine studies. The network nature of hormone regulation and the principles discovered here further emphasize the importance of a systems approach to understanding hormone regulation.

Introduction

Hormones are chemical messengers that circulate through the bloodstream and control and coordinate functions of both nearby and distant cells. Since the discovery of secretin over a hundred years ago[1], about 100 mammalian hormones have been identified[2]. Over the decades, these hormones have been purified, sequenced, and synthesized, their receptors have been identified, and the intracellular signaling pathways they regulate have been explored[3]. Hormones regulate diverse biological processes ranging from growth and development, reproduction, metabolism, and immune responses, to mood and behavior. Many hormones are also therapeutically significant. For example, the discovery of insulin[4] changed type I diabetes from an inevitably-fatal disease to a manageable one. Likewise, corticosteroids, a class of small molecule adrenal hormones, are used to treat diverse pathological conditions, including the inflammation attendant to COVID-19, where corticosteroids reduce the death rate in severe cases by ~30%[5,6].

The classical endocrine hormones are produced by secretory cells residing in one of the nine endocrine glands. However, it is now clear that other cell types, like fat cells and leukocytes, secrete global regulators into the circulatory system[7]. In addition, some endocrine functions, like menstrual cycles in human females, depend upon the interplay and feedback between multiple hormones and multiple cell types[8]. For these reasons, a complete understanding of endocrine physiology requires comprehensive, global approaches. Variations in hormone regulation and function in different species further complicate the situation. Although comparative studies have shown that many hormones are highly conserved across vertebrates, the hormone-producing and -receiving cells are sometimes rewired for different functions[3].

The grey mouse lemur (*Microcebus murinus*) is a small non-human primate with a lifespan of 5-13 years. Because of its small size, rapid reproduction, and close genetic proximity to humans, the mouse lemur is an appealing model organism for studies of primate biology, behavior, aging, and disease[9–13]. With the advances in single-cell RNA-sequencing (scRNAseq) techniques, we have recently created a molecular cell atlas of the mouse lemur by profiling transcripts from ~225,000 cells from 27 mouse lemur organs and tissues of 4 individuals[14] (https://tabula-microcebus.ds.czbiohub.org/). This organism-wide dataset provides an unprecedented opportunity to study global hormone regulation at single cell resolution in a non-human primate.

In this study, we present a comprehensive analysis of primate hormone signaling based on the information in the mouse lemur molecular cell atlas[14]. We systematically mapped hormone-producing and target cells for 87 hormone classes, and created a browsable atlas that depicts gene expression of hormone ligands, modulators, and receptors. This systematic analysis detected both canonical and unreported sites of hormone secretion and action, and identified several examples of species-specific rewiring. We also analyzed the global expression patterns of hormone ligands and receptors, constructed and characterized an organism-wide cell communication network for hormone signaling, and systematically searched the network for feedback circuits. This analysis revealed a few remarkable principles of hormone signaling and a long list of potentially novel feedback mechanisms.

The hormone atlas provides a useful resource and a general approach to study hormone regulation on an organism-wide scale and at single-cell resolution, complementing the more reductionistic strategies typical in classical endocrine studies. The network nature of hormone

95 regulation and principles discovered here further emphasize the importance of a systems
96 approach in understanding hormone regulation.

Results

A comprehensive table of genes involved in the biosynthesis and sensing of human and mouse lemur hormones

Although many hormones have been studied for decades, there is not a well-annotated database
of hormone ligands, synthases, processing enzymes, and receptors. Therefore, we integrated
multiple sources to compile a comprehensive gene table for human hormone signaling (**Table 1**).
For the hormone list, we mainly referenced the *Handbook of Hormones[2]*, the Wikipedia *List of*
human hormones, and the *Handbook of Biologically Active Peptides[15]*, and added newly
discovered hormones including asprosin[16] and the endocrine fibroblast growth factors FGF21,
FGF19, and FGF23[17]. To collect the genes involved in hormone production, we manually
assembled the ligand genes for peptide and protein hormones and the hormone synthesis genes
for small molecule hormones. Maturation or activation of many hormones requires specific
enzymes such as angiotensin-converting enzyme (ACE) for angiotensin, the deiodinases DIO1/2
for thyroid hormones, and prohormone processing enzymes (e.g., PCSK1, PCSK2, CPE) for
many peptide hormones[15]. In addition, hormone function may involve endocrine modulators
such as IGF binding proteins (IGFBPs) for IGF1/2. Accordingly, we added these processing
enzymes and modulators for the corresponding hormones. Next, we collected the hormone
receptor genes that are responsible for signaling in the target cells. We searched ligand-receptor
databases[18,19] for the receptor genes of protein and peptide hormones, and manually added the
receptor genes for the small molecule hormones. Most hormones have characterized ligands and
receptors with a few exceptions in which the genes remain unidentified (e.g., endomorphin
ligand and irisin receptor). In total, we collected information for 87 groups of hormones
including 106 ligand genes, 38 synthases, 24 processing enzymes/modulators, and 164 receptor
genes.

When a hormone group included multiple ligand, receptor, synthase, or processing enzyme
genes, we incorporated AND/OR logic gates among the genes (**Table 1**). Sometimes two or
more of the genes were required for signaling; this AND gate logic was designated by an
ampersand (&). For example, amylin receptors are heterodimeric complexes of CALCR and
RAMP proteins, and both subunits are required for function; thus, CALCR and RAMP were
connected by an AND symbol. Similarly, AND gates were applied among hormone synthases
that catalyze different steps, and between hormone ligands and the corresponding processing
enzymes. On the other hand, some receptor subtypes function independently. For example, there
are nine different adrenergic receptor genes, and none of these receptors require any of the others
for function. Genes like these were linked by OR gates, designated by a vertical line (|).

We next looked up the mouse lemur orthologs of the human hormone genes in the most up-to-
date mouse lemur genome assembly (Mmur_3.0) with both NCBI and Ensembl ortholog
databases (**Table 2**). Almost all (99.1%, 329/332) of the human hormone-related genes were
annotated in the mouse lemur genome. The three genes without mouse lemur orthologs (LHB,

GALP, and INSL4) may result from incomplete annotation of the mouse lemur genome. For example, LHB encodes the beta subunit of luteinizing hormone, which is thought to be expressed in the gonadotrophs of the pituitary gland across all vertebrates. The mouse lemur atlas includes a small number of gonadotrophs that might be expected to express the LHB transcripts. Therefore, we searched for transcripts expressed in the gonadotrophs and identified a candidate gene region[20], currently unannotated in NCBI and Ensembl, with high sequence similarity to human and macaque LHB. The putative LHB region is located next to the RUVBL2 gene, as in humans and mice[21]. The gene region also is selectively expressed in the expected cell type, gonadotrophs (**Fig. S1-luteinizing hormone**). We therefore conclude that the identified region is a likely candidate of the mouse lemur LHB gene. For GALP and INSL4, the atlas lacked the tissue or cell type that is known to express the genes. INSL4, a primate-only paralog of relaxin, is expressed by the placenta in humans[22]. GALP was reported to be secreted by certain hypothalamic neurons[2]. Neither the placenta nor the hypothalamic neurons were surveyed in the mouse lemur cell atlas[14]. Thus, taken together, at least 330/332 (99.4%), and possibly all, of the human hormone-related genes are present in the mouse lemur. For comparison, the mouse genome lacks five human hormone-related genes (98.5% present, 327/332), and four are known to have been lost (GNRH2, MCHR2, MLN, MLNR) (**Table 2**). The percentages of human hormone-related orthologs present in mouse lemur or mouse genomes were higher than the percentage of protein-coding genes (>99.1% vs. 89.5% in the mouse lemur genome, and 98.5% vs. 89.8% in the mouse genome). Most (86.3%) human hormone-related proteins were more similar in sequence to their mouse lemur orthologs than their mouse homologs (**Fig. S2**). These findings underscore the close relationship between human and mouse lemur hormone-related genes.

Organism-wide identification of hormone-producing and target cells

We systematically identified the source cells and target cells for the 87 known hormone groups by examining the expression of genes involved in hormone synthesis and sensing across 739 cell types profiled by 10x droplet-based single cell RNA sequencing from 27 different tissues included in the mouse lemur cell atlas[14]. Source cells were defined as cell types with positive expression of the ligand (for protein/peptide hormones) or synthase(s) (for small molecule hormones) and any necessary modulator genes. Target cells were defined as the cell types with positive expression of the hormone receptor genes. We generated a browsable set of figures that depict the global expression profiles of the hormone-related genes for each of the 87 hormone groups as well as one set of figures for prohormone processing enzymes and one for endocrine secretory proteins (**Fig. S1**). **Fig. 1** shows the cell-type specific expression of four representative examples of hormones (in red) and their receptor(s) (in blue) across the five, color-coded cell compartment types (1. epithelial/neural/germ cells; 2. endothelial cells; 3. stromal cells; 4. lymphoid cells; and 5. myeloid cells). This systematic review identified many canonical sites of hormone secretion and targets, affirming the validity of the single-cell transcriptomic measurements and analysis approach. The analysis also revealed many less-recognized or unreported hormone source/target cells and interesting global patterns of hormone signaling. Below, we highlight a few examples of well-known, lesser-known, and previously unknown sources and target cells. We also showcase examples where hormone signaling is likely rewired in the mouse lemur versus humans or mice.

Prolactin (PRL). The first example is a classical endocrine hormone, prolactin. Prolactin was one of the first hormones discovered[23] and has been studied in depth over the decades. Prolactin is secreted by the lactotrophs of the anterior pituitary gland, the master endocrine organ. Although prolactin was originally regarded as a female-specific hormone that promotes milk secretion, it is now clear that prolactin plays pleiotropic roles in both sexes[2]. These roles include regulation of water and electrolyte balance in the kidney, and control of skin cell proliferation and hair/feather growth and shedding[24]. Prolactin also plays a major role in energy homeostasis by regulating cell growth and metabolism in the liver, adipose tissues, and pancreas in both sexes[25], and may be involved in the immune response[26]. These sex-non-specific functions are conserved, not just among milk-producing mammals, but across all vertebrates.

As expected, prolactin transcripts were selectively expressed in the lactotrophs of the mouse lemur pituitary gland, and were not detected in other tissues (**Fig. 1, Fig. S1-prolactin**). The prolactin receptor (PRLR) was found in both male and female lemurs in the expected cell types, including kidney nephron tubule cells, skin epithelial cells, pancreatic ductal and acinar cells, hepatocytes, intestinal epithelial cells, adipocytes, reticular cells, plasma cells, choroid plexus cells, and the neighboring TSH-secreting thyrotrophs and FSH/LH-secreting gonadotrophs in the pituitary gland (**Fig. 1, Fig. S1-prolactin**). Thus, single-cell RNA sequencing confirms that prolactin signaling is similar in mouse lemurs and other animals.

Adiponectin (ADIPOQ) and leptin (LEP). Mouse lemurs undergo dramatic seasonal changes in body weight and fat reserves[27–29]. Adipocytes were originally regarded only as nutrient storage cells, but it is now clear that they also serve an endocrine role[30]. Adiponectin and leptin are two important metabolic hormones secreted by adipocytes[31,32]. Surprisingly, leptin transcripts were almost undetectable in the adipocytes captured in the atlas (see detailed discussion in the accompanying Tabula Microcebus manuscript 2) (**Fig. S1-leptin**).

However, adiponectin was abundantly and specifically expressed in mouse lemur adipocytes captured from multiple tissues, including four adipose depots (subcutaneous, mesenteric, interscapular brown, and perigonadal), as well as kidney, limb muscle, pancreas, and bone (**Fig. 1, Fig. S1-adiponectin**). Both UCP1-low white-like and UCP1-high brown-like adipocytes highly expressed ADIPOQ.

Previous research has characterized tissue-specific functions of adiponectin in the liver, pancreas, skeletal muscle, endothelial cells, macrophages, and adipocytes[33]. While we indeed observed expression of the adiponectin receptor ADIPOR1 in these tissues in the mouse lemur, we also found ADIPOR1 to be abundantly expressed in almost all lemur cell types (**Fig. 1**). The ubiquitous expression of this receptor suggests that adiponectin exerts an organism-wide effect through ADIPOR1.

In comparison, ADIPOR2, the other adiponectin receptor, was expressed more selectively, most notably in the male germ cells and adipocytes. Interestingly, the mouse lemur expression pattern of ADIPOR2 differs from that of the mouse in which ADIPOR2 is reported to be most highly expressed in the liver and almost undetectable in the testis[34], suggesting differential ADIPOR2 effects in the two species.

Motilin (MLN). Comparing expression patterns of the mouse lemur to that of the mouse and human, we identified interesting cases of hormone network rewiring. A telling example is

motilin, canonically considered to be a gastrointestinal hormone secreted by enteroendocrine cells to regulate gastrointestinal smooth muscle movement[15]. Mice lack both the motilin and the motilin receptor genes (**Table 2**). Indeed, genome comparative studies reported that motilin signaling is conserved in vertebrates but is lost in rodents[35]. The lack of motilin signaling in mice and rats has hampered progress on further characterizing motilin function as well as the development of motilin agonists for the treatment of gastroparesis[36]. In the mouse lemur, we observed motilin expression in enteroendocrine cells in the small intestine but not colon (**Fig. 1A**), which is consistent with the canonical view. However, there was also a broad range of other motilin-expressing cell types, including kidney nephron tubule cells, chemosensory tuft cells from both the airway (lung and trachea) and small intestine, certain brain neurons, adipocytes (higher in UCP1-high brown-like adipocytes than in white-like adipocytes), mesothelial cells, and the majority of the CD4+CD8-CCL5- helper T cells (**Fig. 1**, **Fig. S1-motilin**). The receptor was found to be most highly expressed in pancreatic delta cells and several epithelial cell types of the skin (e.g., sweat gland cells). Together, this expression pattern suggests that motilin is involved in a diverse array of mouse lemur physiological functions in addition to the regulation of gut motility.

Resistin (RETN). We also noted interesting species differences in resistin signaling. Resistin is a phylogenetically recent hormone, present only in mammals[2]. It is expressed mainly by adipocytes in mice but by macrophages, monocytes, and neutrophils in humans[37,38]. In the mouse lemur, we found that resistin is highly expressed in macrophages, monocytes, and neutrophils, and nearly absent from adipocytes (**Fig. 1, Fig. S1-resistin**), as it is in humans. Interestingly, resistin is also highly expressed in many of the vascular endothelial cell types and in metastatic uterine tumor cells, suggesting additional sources for this hormone.

Prohormone processing enzymes. Prohormone processing enzymes are required for the maturation of many peptide hormones, including insulin, adrenocorticotropic hormone (ACTH), gastric inhibitory polypeptide (GIP), and somatostatin[15]. The required processing enzymes have been identified for some but not all hormones[39,40]. In this study, we considered both the ligand and the known or suggested processing enzymes when determining if a cell type was a hormone source. Selective expression of the processing enzymes can help distinguish hormones that share the same precursor gene. For example, proglucagon (GCG) is the precursor for both glucagon and glucagon-like peptide-1 (GLP-1). GCG was expressed in both pancreatic α cells and intestinal enteroendocrine cells, but the two cell types expressed different processing enzymes: α cells expressed only PCSK2 which is required for glucagon synthesis, and enteroendocrine cells expressed only PCSK1 which is required for GLP-1 synthesis (**Fig. S1-glucagon**). Thus, as is the case in mice and humans [41], α cells are sources of glucagon and enteroendocrine cells are sources of GLP-1 in the mouse lemur.

Interestingly, we noticed that some cell types in the mouse lemur transcribed prohormone genes without expressing the respective processing enzymes. For example, proopiomelanocortin (POMC) is the precursor of multiple hormones including ACTH, melanocyte-stimulating hormone (MSH), β-endorphin, and lipotropin, and the first step in POMC processing is carried out by PCSK1[42]. As expected, ACTH-secreting corticotrophs expressed both POMC and the necessary processing enzymes, but mouse lemur male germ cells expressed abundant POMC without PCSK1 (**Fig. S1-proopiomelanocortin-derived hormones**). Similarly, male germ cells expressed GIP, which also requires PCSK1 processing. We also found renin (REN) expression in

the intestinal and airway tuft cells without the necessary PCSK1, oxytocin (OXT) expression in
kidney proximal tubule epithelium without the required carboxypeptidases E (CPE), and
neuropeptide Y (NPY) expression in bladder urothelial cells without the necessary CPE (**Fig.**
S1). These results suggest several possible hypotheses: different peptidases are involved, the
prohormone transcripts are not translated, the prohormone proteins may function without
proteolysis, or the processing enzymes may be expressed below detection levels in the animals
that were sampled.

In summary, we have depicted the organism-wide, single-cell expression patterns for hormones,
synthases, hormone modulators, and receptors from 87 hormone groups, and have systematically
identified their sources and targets (**Fig. S1**). This global analysis detected canonical cell types
underlying hormone secretion and target functions while also revealing previously unreported
sources and targets. This organism-wide single-cell hormone atlas should facilitate the
systematic study of hormone regulation in an unbiased, global manner, paving the way for the
discovery of novel aspects of hormone regulation and function.

Hormone-related gene expression faithfully classifies cell type

Next, we clustered mouse lemur cell types based on the transcriptional profiles of only the
hormone-related genes (**Fig. 2A-B**). We organized the cell types into 75 clusters and summarized
the hormones and hormone receptors that were expressed in the majority of the cell types for
each cluster in **Table 3**. We noted that cells of the same type almost always clustered closely
irrespective of their tissue of origin, and different cell types were well-separated in the cluster
dendrogram (**Fig. 2A**). For example, all lymphatic endothelial cells clustered closely and showed
characteristic IGF1 expression despite their coming from nine different tissues and four
individuals (cluster 53, **Fig. 3B**, **Fig. S1-insulin-like growth factor, Table 3**). Schwann cells
from nine tissues clustered into myelinating and non-myelinating subtypes (cluster 40 and 37).
Adipocytes sampled from four adipose depots and six other tissues formed one cluster (cluster
13, **Table 3**). Skeletal muscle stem cells from the limb muscle and tongue clustered together
(cluster 17, **Table 3**). Mesothelial cells sampled from eight visceral organs formed a cluster
distant from other stromal cells and showed distinctive expression of NPR1, CFB, PTGIS,
ITGB3, and PNMT (cluster 44, **Fig. 2A-B, Fig. S4A, Table 3**). Although the analysis only
examined the transcripts of 329 hormone-related genes, ~1% of all detected genes, these results
suggest that hormone-related gene expression is indicative of the overall transcriptome among
different cell types.

To further test this hypothesis, three comparative analyses were performed. First, we projected
the high-dimensional transcriptional profiles of the mouse lemur cell types into two-dimensional
uniform manifold approximation and projection (UMAP) representations, which illustrate cell
type similarity in the space of the selected genes. We compared the spatial relationship of cell
types in two UMAPs: one constructed based on only the 329 hormone-related genes (**Fig. 2B**),
the other based on all genes excluding the hormone genes (>30,000 genes) (**Fig. S3**). In each
case, the data points represent cell types rather than individual cells (**Fig. 2B, Fig. S3, Fig. S4**).
The two UMAPs showed remarkable similarity and cell types were similarly clustered with only
minor differences (see **Fig. S4C** for comparison). One such difference was the chemosensory tuft
cells that reside in the epithelium of both the airway (lung, trachea) and small intestine. In the

space of hormone-related genes, tuft cells from all three tissues formed a cluster (cluster 21) with
distinctive expression of renin (REN), cyclooxygenase 1 (PTGS1), arachidonate 5-lipoxygenase-
activating protein (ALOX5AP, required for leukotriene production), and motilin (MLN) (**Fig.**
2B, Fig. S1, Table 3). However, in the space of all genes, the lung, trachea, and intestinal tuft
cells were separated from each other (labeled as 21*A, 21*B, and 21*C in **Fig. S3**). Another
notable exception was skin melanocytes (cluster 38), which clustered closely with
oligodendrocytes (cluster 39) when using only the hormone-related genes (**Fig. 2B**) but clustered
with skin epithelial cell types (cluster 36) in the space of all genes (**Fig. S3**). These differences
notwithstanding, the hormone-gene and transcriptome-based cell type UMAPs were qualitatively
similar and shared an overall clustering pattern.

We then compared the transcriptional distances between pairs of cell types for either just the
hormone-related genes (hormone-based distances) or the non-hormonal transcriptome
(transcriptome-based distances) (**Fig. 2C, Fig. S5A-B**). The two distances were positively
correlated, with a Pearson correlation coefficient (r) of 0.70 ($p < 1E-15$). Similarly, we compared
hormone-based transcriptional distances between pairs of cell types to a Cell Ontology-based
distance. Cell Ontology distances represent a canonical view of the relatedness of cell types[43]
and were calculated as the mean of the Cell Ontology embedding-based distances from the Cell
Ontology graph and the text-based distances from the text descriptions of the Cell Ontology
terms[44]. These two distances were also significantly correlated, with $r=0.45$ ($p < 1E-15$) (**Fig.**
2D, Fig. S5A and C). Taken together, these analyses support the notion that the small number of
hormone-related genes faithfully classifies most of the cell types of the mouse lemur.

We then examined if the expression of hormone ligands or receptors alone could distinguish cell
types (**Fig. S4**). Interestingly, for most cell type clusters, hormone ligands or receptors alone
classified cell types similarly. This suggests that most cell types have stereotypical expression of
proteins involved in both incoming and outgoing hormonal signaling. A small fraction of clusters
were segregated from other clusters mainly by only the ligands or receptors. For example,
clusters of pituitary neuroendocrine cells (cluster 7) and tuft cells (cluster 21) are distinguished
mainly by hormone ligands, whereas clusters of the skeletal muscle stem cells (cluster 17) and
lung endothelial cells (cluster 51) are mostly based on differences in hormone receptor
expression.

In summary, we found that cells generally show stereotypical and cell-type-specific expression
of hormone ligands and receptors. Thus, hormone secretion and sensing essentially labels the
identity of the cells.

Hormonal regulation nevertheless does sometimes differ in related cell types

Although cells of the same type tended to have similar hormonal expression patterns, we
identified a number of interesting exceptions where closely related cell types differed in their
hormone signaling. Below we highlight several such examples in which we identified changes in
hormonal signaling among cells at different development stages (spermatogenesis), cells from
different tissue origins (vascular endothelial cells), cell types that are not discretely segregated
but form a continuous gradient (kidney nephron and vasa recta), cells of different molecular
subtypes (hepatocytes), and primary vs. metastatic tumor cells (**Fig. 3**).

Spermatogenesis. Spermatogenesis is a tightly regulated process that happens asynchronously and continuously in the male testis[45]. Therefore, sampling the testis at one fixed time point can capture snapshots of the entire process. The atlas includes seven sperm and sperm progenitor cell types that are at different spermatogenesis stages as indicated by the expression of known stage markers[14]. Interestingly, these cells exhibited spermatogenesis-stage-dependent hormone signaling (**Fig. 3A**). The seven cell types fall into three major stages: the undifferentiated spermatogonia, the spermatocytes (which undergo meiosis), and the haploid spermatids that elongate and form mature sperms. Comparing hormone-related gene expression among these three major stages (cluster 1, 2, 8), over 50 hormone ligands, enzymes, and receptors were differentially expressed (**Fig. 3A**), suggesting an extensive shift in hormone signaling. Previous studies on spermatogenesis have mainly focused on testosterone and on the supporting cells rather than sperms and sperm progenitors[46,47]. The present data suggest that sperms and sperm progenitor cells are not only differentially regulated by a variety of hormones but may also secrete a changing array of signaling molecules as they mature. Some of these molecules have been recently suggested to be involved in sperm functions. For example, progesterone was known to activate a sperm-specific calcium channel prior fertilization to induce Ca^{2+} influx, but the underlying progesterone receptor(s) remained unidentified[48]. Two non-classical progesterone membrane receptors (PAQR5/9) were highly expressed in the lemur spermatids and therefore may mediate the process (**Fig. 3A**). Interestingly, two additional membrane progesterone receptors (PAQR8, PGRMC1) were specifically expressed in the undifferentiated spermatogonia (**Fig. 3A**), suggesting that they may contribute to regulation of early spermatogenesis by progesterone[49]. Insulin-like factor 6 (INSL6), a member of the secretory relaxin family, was selectively expressed in the lemur spermatocytes (**Fig. 3A**). Supporting a function in these cells, knockout of INSL6 arrests spermatogenesis at late meiotic prophase in mice[50]. This analysis suggests that sperms and sperm progenitor cells may be directly engaged in hormone signaling with surrounding and potentially distant cells to affect spermatogenesis and sperm functions.

Vascular endothelium. Endothelial cells, including artery, vein, capillary, and lymphatic cells, line the surface of the circulatory system and are all derived from a developmentally uniform lineage[51]. The mouse lemur cell atlas includes a large number of endothelial cells captured from different organs, providing an opportunity for cross-organ comparison. Although vascular endothelial cells in general shared similar transcriptional profiles (**Fig. 2A-B, Fig. S3**), the lung and brain vascular endothelial cells exhibited unique hormone signaling signatures (**Fig. 3B**).

Endothelial cells of artery, vein, and capillary in the mouse lemur lung showed higher expression of hormone receptors VIPR1, CALCRL, and RAMP1 (**Fig. 3B**) compared to endothelial cells from other organs. VIPR1 is a receptor for vasoactive intestinal peptide (VIP), a strong bronchodilator and one of the most abundant signaling peptides in the lung[52]. Similar expression of VIPR1 is also observed in human but not mouse lung endothelial cells[53,54]. RAMP1 forms heterodimeric receptors with CALCR or CALCRL that are regulated by several hormone ligands including CGRP, adrenomedullin, amylin, and calcitonin, all of which have been reported to have vasodilatory effects[2]. Interestingly, RAMP1 is expressed in the lung endothelium of the mouse lemur but not in mice or humans, whereas RAMP2 is expressed in all three species and RAMP3 is only expressed in humans (**Fig. S1-adrenomedullin**)[53,54] (also see accompanying Tabula Microcebus manuscripts 1[14] and 2). These results suggest that lung vascular endothelium is a unique hormonal target in comparison to vasculature elsewhere, possibly because of the distinctive nature of the pulmonary circulation (e.g., high volume, low pressure) compared to the

systemic circulation. The lung endothelium also shows species variation in the hormone
signature, perhaps reflecting the relatively recent appearance of lungs in evolution compared to
other visceral organs.

In contrast, brain vascular endothelium was found to express higher levels of a few hormone
ligands/synthases. Endothelin-3 (ET-3), a member of the vasoconstricting endothelin family, was
selectively expressed in the brain vascular endothelium (**Fig. 3B**). ET-3 peptide has been
detected at various brain locations in rodents, but it remained unsettled what cells secreted the
peptide and what physiological role it played[55]. Our results suggest that ET-3 is synthesized by
the endothelial cells in the brain, which could target the nearby EDNRB-expressing SLC7A10+
astrocytes and the EDNRA-expressing vascular smooth muscle cells and pericytes of the brain
(**Fig. S1-endothelin**). Additionally, the brain endothelium also expressed high levels of PTGES
and PTGS2/COX-2 (**Fig. 3B**), which are the synthases for prostaglandin E2 (PGE2). PGE2 has
been suggested to regulate inflammation-induced fever in mice[56,57]. Interestingly, brain
endothelial cells also selectively lacked expression of IGFBP4, which was abundantly expressed
in the endothelium of all other tissues (**Fig. 3B**).

The expression of the detected hormone signature genes (RAMP1, CALCRL, and VIPR1 in
lung, and EDN3, PTGES, and PTGS2 in the brain) was higher in endothelial cells than other cell
types from the same organ (**Fig. S1**). Thus, it is unlikely that these organ-specific endothelial
expression patterns resulted from cross-contamination (signal spreading). Altogether, lung and
brain endothelial cells exhibited organ-specific hormone signaling which likely regulates organ-
specific physiological functions.

Kidney nephron and vasa recta. Although discrete cell types were common in the mouse lemur
cell atlas, there were clear examples of spatially continuous cell types that displayed a continuum
of gene expression. These were the kidney nephron epithelium and the vasa recta endothelium[14],
and these cells also displayed gradients of hormone signaling along their spatial axes (**Fig. 3C-
D**).

Nephrons are single-cell layered, one-directional tubules that serve as the structural and
functional units of the kidney. We identified a large number of hormone-related genes that are
specific to different sections of the nephron epithelium (**Fig. 3C**). Some of these hormones target
specific nephron sections to regulate urine formation. For example, vasopressin is well known
for its antidiuretic effect, promoting water reabsorption in the collecting duct principal cells[2].
Indeed, the vasopressin receptor AVPR2 was expressed exclusively in the collecting duct
principal cells. Interestingly, the collecting duct intercalated cells expressed another vasopressin
receptor, AVPR1A (**Fig. 3C**), which has been recently reported to mediate the regulation of acid
homeostasis by vasopressin[58]. Although the kidney epithelium is not commonly considered to be
an endocrine tissue, many hormone ligands/synthases were also expressed in different nephron
sections, including angiotensin (AGT), adrenomedullin 2 (ADM2), motilin (MLN), and
neuropeptide Y (NPY). This suggests that nephron epithelium may play multifaceted roles in
hormone production in addition to urine formation.

We also identified interesting differences in hormone signaling in the vascular endothelium of
different regions of the mouse lemur kidney. In particular, we identified a gradual shift in
hormone signaling along the arteriole-to-venule-axis of the vasa recta endothelium (**Fig. 3D**).

The vasa recta align parallel to the nephron tubules in the kidney medulla and function
collaboratively with the nephrons in urine formation[59]. **Fig. 3D** shows the hormone-related genes
that are significantly differentially expressed along the vasa recta axis. Notably, several genes
involved in IGF signaling were differentially expressed: IGF1, which was expressed in the
lymphatic endothelial cells but almost absent from other vascular endothelial types (**Fig. 3B, Fig.**
S1-IGF), was highly expressed at the venule end of the vasa recta; IGF2 and IGF1R exhibited
the opposite pattern with higher expression in the arteriole end; and four IGFBPs (IGFBP3-6),
which modulate IGF signaling, exhibited axis-dependent expression patterns. This spatial
organization of IGF signaling may be evolutionarily conserved, as a recent study on mouse
kidney vasculature development shows a similar pattern of IGF1 expression in the mouse vasa
recta[60].

Hepatocyte subtypes. The mouse lemur cell atlas identified multiple novel molecular subtypes
of known cell types[14]. For example, the liver hepatocytes of the mouse lemur clustered into two
discrete subtypes, APOB+ hepatocytes and PHYH+ hepatocytes. These two hepatocyte subtypes
are likely independent of the known zonal heterogeneity, and a similar subtype division was
observed in mouse hepatocytes[14]. Interestingly, a number of hormone-related genes were among
the most highly differentially expressed genes in the two hepatocyte subtypes in both the mouse
lemur and mouse (**Fig. 3E**)[14]. Hormones including glucagon, androgens, and prolactin appear to
selectively target the APOB+ hepatocytes as indicated by differential expression of the
corresponding receptors (GCGR, AR, PRLR). In addition, angiotensin (AGT), a canonical
hepatic hormone, was expressed at a much higher level in the APOB+ hepatocytes, whereas no
subtype difference was detected in the expression of hepcidin (HAMP) or IGF1, two other
hormones abundantly produced by the liver. Hepatocytes are known to show sex-dependent
hormone regulation[61]; however, the subtype differences in hormone gene expression were not
sex-dependent (**Fig. S6**).

Liver is a central organ of metabolism and plays important roles in hormonal regulation. The
detected heterogeneity in hormone signaling suggests potentially important functional
divergence between the hepatocyte subtypes in both the mouse lemur and mouse. A deeper
understanding of the mechanisms and regulation of hepatocyte hormone heterogeneity would be
particularly relevant to the study of mouse lemur-specific physiology, like hibernation/torpor and
seasonal body weight changes[28].

Metastatic vs. primary tumor cells. The atlas also contains cells that are pathologic, including
cells from the lung tissue of one female mouse lemur that were hypothesized by gene signature
analysis and confirmed by histology to be metastatic tumor cells of endometrial cancer origin
(also see accompanying Tabula Microcebus manuscripts 1[14] and 2). The tumor cells clustered
with the epithelial cells of the uterus from a second female individual (also, as it happened, with
confirmed endometrial cancer), particularly the MUC16+ non-ciliated uterus epithelial cells
which are likely primary tumor cells according to signature gene expression (see accompanying
Tabula Microcebus manuscript 2). The metastatic and primary tumor cell types shared
characteristic expression of oxytocin receptor (OXTR) and inhibin beta B subunit (INHBB) (**Fig.**
3F, Fig. 2A-B, Fig. S3, Table 3), suggesting that the metastatic tumor in the lung originated
from cells resembling the primary MUC16+ uterine tumor cells. However, certain aspects of
hormone signaling were likely to have been rewired in the metastatic tumor (**Fig. 3F**). Notably,
the tumor cells showed a loss of estrogen receptors (ESR1), a change associated with high grade

and advanced stage endometrial cancers in humans[62] and more often reported in high mortality
type 2 human endometrial cancers[63].

Interestingly, the metastatic tumor cells also acquired new abilities in hormone production. For
example, elevated expression of prostaglandin E2 synthase (PTGES2) was present in metastatic
cells (**Fig. 3F**). PGE2 has been suggested to be a tumorigenic factor in many cancers[64]. A recent
study identified that the PGE2 synthase PTGES2 was elevated in human endometrial cancer
tissues and found that PGE2 promotes cell proliferation and invasion in *in vitro* cell assays[65].
Resistin was also found to be highly expressed in the metastatic cells (**Fig. 3F**). Resistin has been
reported to promote invasiveness of breast cancer cells *in vitro*[66]. Thus, endometrial tumors may
utilize autocrine resistin signaling to promote metastasis. The shift in hormone signaling
identified in the lemur metastatic endometrial tumor is similar overall to that reported in the
human. This suggests that the rewiring of hormone signaling may be an important step in tumor
development and metastasis. Mouse lemurs may serve as a useful animal model to study human
cancers that are not well modeled in mice such as endometrial cancers (see more in
accompanying Tabula Microcebus manuscript 2).

In summary, we have highlighted five examples where closely related cells differ in their
hormone signaling. These closely related cells usually share the same developmental origin but
diverge during cell differentiation or development. Cells likely undergo changes in hormone
signaling during the process to interact with other cells and accommodate their specific
functional requirements.

The hormonal cell communication network is densely connected

Long-range hormonal signaling connects both nearby and distant cells through the bloodstream
and enables cross-organ communication. By linking the source cells that produce a hormone
ligand to the target cells that express the corresponding receptor, we constructed a global cell
communication network for mouse lemur hormone signaling (**Fig. 4A**). Nodes of the network are
cell types and edges are directed from hormone source to target cells.

The hormone network was remarkably densely connected; 51% of all possible directed edges
were found to be present. In fact, in the graphical depiction of the network (**Fig. 4A**), individual
edges are hardly distinguishable because the edges are so dense. The density of the hormone
network is comparable to that of the cytokine network[67], but much higher than the densities of
other characterized biological networks. For example, the network densities of the yeast, fly, and
human protein-protein interaction networks are all estimated to be ~0.1%[68–70], and the yeast
genetic interaction network density is ~3%[71]. The dense connection between cell types by
hormonal regulation is also highly robust: 67% of all existing edges are connected by more than
one ligand-receptor pair, and the network maintains high density connection when randomly
removing 10% of the ligand-receptor pairs (**Fig. S7A**).

The density of the hormone network is largely due to the fact that multiple hormone ligands and
receptors are expressed in most cell types (**Fig. 4B**). On average, cell types expressed 17.0 ± 7.9
(mean ± S.D.) receptors or receptor complexes. In comparison, fewer hormone ligands were

expressed (2.8 ± 2.5, mean ± S.D.). Only a few cell types expressed 10 or more hormone ligands
(e.g., enteroendocrine cells, pancreatic endocrine cells, certain neurons, and certain fibroblasts),
and 38.3% of cell types expressed a single ligand or no ligand. This indicates that cells in general
respond to more hormones than they produce.

The five neuroendocrine cell types of the anterior pituitary—corticotrophs, thyrotrophs,
gonadotrophs, lactotrophs, and somatotrophs—play a vital role in endocrinology. These cells are
components of regulatory cascades, and classically they are regarded as receiving regulatory
signals from one or a few hormones produced in the hypothalamus, and in turn producing one or
a few hormones to regulate a peripheral endocrine gland. Yet even these cells were found to
express numerous receptors, ranging from 17 (corticotrophs) to 29 (gonadotrophs and
somatotrophs) (**Table 4**). These cells appeared to be regulated by growth factors, neuropeptides,
peripheral and classical hypothalamic hormones, as well as neurotransmitters (e.g., dopamine)
that were not on our hormone list. Likewise, they were found to express genes for several
hormones in addition to the classical hormones (**Table 4**). For example, corticotrophs expressed
galanin (GAL), gastrin (GAS), and cortistatin (CORT), as well as the ACTH precursor (POMC)
and its processing enzymes PCSK1, PCSK2, and CPE.

Taken together, this analysis suggests that virtually all cell types across the body are extensively
engaged in cell communication via hormonal signaling and that the connections are likely
evolutionarily robust to random mutations that disturb hormone regulation. The dense connection
of the network resulted in a large number of highly connected nodes and no obvious network
cores (**Fig. 4A, Fig. S8**). This differs from the commonly studied scale-free networks, which are
dominated by a small number of highly-connected hub nodes[72]. Studies on hormone regulation
usually focus on one specific site of secretion and/or target; our analysis emphasizes the
importance of a systems view in understanding hormone regulation.

A small number of hormones are global regulators

We next examined properties of the hormone network edges. In contrast to the protein-protein
interaction network and the genetic interaction network, the edges of the hormone
communication network are directional and can be distinguished into different types according to
the specific ligand-receptor pairing. The hormone network contained 332 different edge types
that are selected pairings of 128 ligands and 166 receptors. A ligand or receptor may contain
multiple genes as required for ligand synthesis/maturation and in receptor complexes (see **Table
1**).

Some ligands and receptors were expressed in highly specific cell types while others were more
broadly expressed. We therefore quantified for each ligand or receptor a generality score to
evaluate the range of its expression across different cell types in the atlas (**Fig. 4D-E, Fig. S9,
Table 5**). Strikingly, a small number of hormone receptors were globally expressed. For
example, the top ranking receptor, adiponectin receptor 1 (ADIPOR1), was expressed in 93.4%
of the cell types. Other broadly expressed hormone receptors included the renin receptor
ATP6AP2 (90.4%), the membrane-associated progesterone receptor PGRMC1 (84.1%), the
cortisol receptor NR3C1 (73.7%), the resistin receptor CAP1 (72.9%), the IGF1 receptor IGF1R

(69.5%), the opioid growth factor receptor OGFR (65.9%), and the thyroid hormone receptor
THRA in complex with RXRB (54.3%). While it has been recognized that some of these
hormones exert organism-wide effects, our results suggest that they target not only a broad range
of tissues but likely regulate a large portion or perhaps majority of the cells of the body.

In comparison, most hormone ligands and receptors were expressed in specific cell types; ~85%
of the hormone ligands and ~60% of the hormone receptors were detected in less than 5% of the
cell types. Certain hormones like follicle stimulating hormone (FSHB and CGA) were only
detected in a single cell type. Only a few hormone ligands were relatively broadly expressed,
including adrenomedullin (ADM, 18.6%), acylation stimulating protein (C3, 18.1%), IGF2
(17.9%), and resistin (RETN, 15.4%). The generality score followed a power-law distribution
toward the tail where a small number of globally expressed receptors dominated the distribution
(**Fig. S9B-C**). This resulted in a bimodal distribution of the network node outdegree (abundance
of outgoing edges/downstream regulation) (**Fig. 4A and C, Fig. S8**). Cells expressing hormone
ligand(s) to the high-generality receptors (e.g., adipocytes which secrete adiponectin) sent
globally connected outgoing edges, and other cells that only expressed ligand(s) to specific
receptors sent a much smaller number of outgoing edges. In contrast, the nodes' indegree
(abundance of incoming edges/upstream regulation) was more homogeneously distributed
because of the higher specificity of hormone ligand expression and large number of hormone
receptors expressed in the cells (**Fig. 4A-C, Fig. S8**).

Receptors for the same hormone tend to be expressed in mutually exclusive cell types

Many hormones have more than one receptor, which may mediate different functions (**Table 1**).
We therefore examined if receptors for the same ligand tend to be co-expressed or if their
expression is mutually exclusive across different cell types. To quantitatively compare the target
cell type profiles, we calculated a Jaccard index (JI, intersection over union) for each qualifying
receptor pair, namely the ratio of cell types that co-expressed both receptors to the cell types that
expressed either of the receptors. This analysis revealed that the majority of the receptor pairs
had extremely low JIs: 25% (41/161) of the receptor pairs had a JI equaling zero, meaning no
overlap in their target cell types, and 57% (92/161) of the receptor pairs had a JI of less than 5%
(**Fig. 4F**). A telling case is the adrenoceptors; although epinephrine (adrenaline) targets many
cell types, different adrenoceptor subtypes were selectively employed in different cell types (**Fig.
4F, Fig. S1-epinephrine/norepinephrine**). For example, adrenoceptors ADRA1D and ADRB3
shared no target cell types; ADRA1D was expressed broadly in vascular smooth muscle cells and
pericytes of multiple organs, certain glial cells, and leptomeningeal cells, whereas ADRB3
mainly targeted adipocytes. Vasopressin receptors AVPR1A and AVPR2 also had less than 3%
overlap in their target cell types; AVPR2 was predominantly expressed in the kidney collecting
duct principal cells, while AVPR1A was highly expressed in the kidney collecting duct
intercalated cells and broadly expressed in many stromal cell types (**Fig. S1-vasopressin**).

A few receptor pairs showed a tendency toward co-expression, suggesting possible interactions
in cell signaling. Supporting the notion, one top ranked pair was the IGF receptors IGF1R and
IGF2R (JI=0.40), which are known to interact in IGF2 signaling. IGF2R internalizes after IGF2
binding and clears IGF2 from the cell surface, and therefore functions as a competitive inhibitor

of IGF2-IGF1R signaling[73,74]. Two other top ranked pairs were NPY1R/NPY5R (JI=0.49),
receptors for neuropeptide Y, peptide YY and pancreatic polypeptide[75], and
ADIPOR1/ADIPOR2 (0.39), receptors for adiponectin. Coexpression of NPY1R and NPY5R
has been identified in rodent brain, and recent studies indicate that the two GPCRs may form
heterodimers, resulting in modulated agonist binding properties[76,77]. Similarly, biochemical
studies on ADIPOR1 and ADIPOR2 in cell culture systems have shown that the two receptors
can form heterodimers promoting membrane expression of ADIPOR2[78,79].

Taken together, this analysis showed that the expression of distinct receptors for the same
hormone tends to be mutually exclusive, suggesting functional divergence and specification
among hormone receptor subtypes. The rare cases of co-expressed receptors appear to involve
functional interaction or cross-regulation between the receptor pairs.

Feedback circuits are widespread in the hormonal cell communication network

Hormone regulation often involves feedback control. We therefore searched the mouse lemur
hormone cell communication network to systematically identify potential feedback regulation.
For simplicity, we focused on two-node feedback circuits, namely cell type pairs that are
connected by edges in both directions (**Fig. 5A**). This systematic search revealed a vast number
of feedback circuits; 29% of all node pairs formed feedback circuits, showing that feedback
control is widespread in the hormone network (**Table 6**). We then tested if feedback circuits are
enriched in the hormone network, i.e. more prevalent than would be expected by chance. We
applied degree-preserving random rewiring[80] to randomly permute edge connections without
changing node degree (i.e., network edge number and distribution). **Fig. 5B** shows the
distribution of the feedback circuit counts from such randomly permuted networks ($n = 1000$).
The number of feedback circuits in the permuted networks was consistently lower than in the
original hormone network. However, the difference was small (on average, only ~0.2%).
Additionally, we randomly removed 10% of ligand-receptor pairs to test how perturbations to
ligand-receptor binding affect feedback distribution of the network. This analysis revealed that
the network feedback circuits remained abundant despite edge removal as cell connections
usually involved multiple ligand-receptor pairs (**Fig. S7B, Fig. 4B**). Taken together, this analysis
suggests that feedback circuits are abundantly and robustly distributed as a result of the dense
connection of the hormone network. Evolution may have further enriched feedback circuits in
hormone regulation as suggested by the random rewiring analysis.

We next reviewed selected feedback circuits between endocrine cell types (**Fig. 5C-L**) to
compare with literature reports on hormonal feedback regulation. The analysis successfully
detected well-known feedback regulation. For example, **Fig. 5C** shows the feedback loop
between gonadotrophs in the anterior pituitary and the estrogen-producing granulosa cells of the
ovary (fortuitously obtained from female perigonadal fat), a classical example of hormonal
feedback control that is evolutionarily conserved across vertebrates[81]. Consistent with this
scheme, mouse lemur gonadotrophs expressed two highly-conserved gonadotropins, follicle
stimulating hormone (FSH, a dimer of FSHB and CGA) and luteinizing hormone (LH,
LHB/CGA dimer), which target granulosa cells through the corresponding receptors FSHR and
LHCGR. Granulosa cells, in turn, expressed key synthases required for the production of sex

hormones, which feed back to gonadotrophs through androgen and estrogen receptors (AR, ESR1). Additional feedback regulation from granulosa cells to gonadotrophs was also mediated through anti-Müllerian hormone (AMH) and inhibin/activin signaling. We also detected the well-established feedback loop between somatotrophs in the anterior pituitary and hepatocytes in the liver, which was mediated through growth hormone (GH), IGF1[2,82], and potentially hepcidin and FGF21 signaling[83] (**Fig. 5D**). Additionally, in the pancreas, we detected the known feedback circuits between pancreatic α and β cells through glucagon and insulin signaling and between α and δ cells through glucagon and somatostatin signaling (**Fig. 5E-F**)[84]. The detection of these various known feedback loops supports the validity of the comprehensive scRNAseq approach.

The analysis also revealed a long list of potential endocrine feedback circuits that have not been reported, or have been reported recently but are not yet widely appreciated. For example, we detected a feedback loop between pancreatic α cells and hepatocytes with one leg from α cells to hepatocytes mediated by glucagon and the other leg from hepatocytes to α cells through IGF1 and FGF21 (**Fig. 5G**). Regulation of hepatocyte glycogenolysis and gluconeogenesis by glucagon is well-established[85]. Previous work on the fasting-induced hormone FGF21 has shown that injection of glucagon increases plasma FGF21 in humans[86] and injection of FGF21 reduces glucagon in mice[87]. A recent study also observed that plasma IGF1 and glucagon levels are negatively correlated in non-diabetic humans and found that IGF1 negatively modulates glucagon secretion in an α cell culture model[88]. Considering that the liver is a major source of circulating IGF1 and FGF21, our results and the literature reports suggest that α cells likely form a negative feedback loop with hepatocytes in maintaining metabolic homeostasis.

Additionally, adipocytes, through adiponectin, formed feedback loops with pituitary neuroendocrine cells, including the growth hormone secreting somatotrophs, the prolactin secreting lactotrophs, and the ACTH secreting corticotrophs (**Fig. 5H-J**). It has been established that growth hormone, prolactin, and ACTH reduce adiponectin secretion in adipocytes[89–91], but much less is known about the effect of adiponectin on pituitary neuroendocrine cells. Interestingly, a recent study using cultured primary pituitary cells from two non-human primate species found that adiponectin decreases growth hormone and ACTH release and increases prolactin release[92]. This suggests that adipocytes form a negative feedback loop with lactotrophs which possibly stabilizes prolactin and adiponectin levels, and form double-negative feedback loops with somatotrophs and corticotrophs, possibly functioning as bistable switches[93]. It remains to be characterized how adiponectin affects pituitary hormone secretion, what are the physiological functions of the pituitary-adipocyte feedback control, and whether this regulation is specific to primates.

We also detected feedback circuits of intestinal enteroendocrine cells with pancreatic β cells and adipocytes (**Fig. 5K-L**). It is known that enteroendocrine cells secrete GIP, which induces glucose-dependent insulin secretion in β cells after eating and exposure of the intestinal lumen to nutrients[2]. Despite the important physiological role of GIP in insulin secretion and glucose homeostasis, there is limited research on the potential feedback regulation of GIP secretion by insulin. García-Martínez et al. reported that insulin induces prolonged GIP expression in an enteroendocrine cell line[94] , suggesting a positive feedback regulation and synergistic effects between the two hormones in maintaining glucose homeostasis under a surge of blood glucose after eating.

690 Taken together, this analysis identified an extensive list of potential feedback circuits in the
691 global hormonal cell communication network (**Table 6**), providing a rich resource for generating
692 hypotheses on endocrine regulations and to compare hormonal regulatory mechanisms in
693 different species.

Exploring hormonal regulation of the circannual rhythms that control organismal biology

Lastly, the mouse lemur hormone atlas may bring new insights to the study of mouse lemur
physiology, such as circannual rhythms[29,95], social interactions[96], and aging[11–13,97]. As an
adaptation to seasonal environmental changes, mouse lemurs have evolved striking seasonal
rhythms in metabolism, behavior, and reproduction throughout their 5–10-year lifespan. Mouse
lemurs build fat reserves in preparation for the winter and enter daily torpor, a hypometabolic
state, to preserve energy, while in the summer the animals maintain a lean body and regrow their
reproductive organs for mating. The global nature of these phenotypic changes suggests that they
may be hormonally mediated. To date, only 12 circulating hormones have been measured in
mouse lemurs, but all showed dramatic seasonal changes (**Fig. 6A, Table 7**). This suggests that
hormones are likely the key messengers coordinating and driving various seasonal changes
among different organs. The hormone atlas suggests that these seasonally changing hormones
target a broad array of cells and tissues; 95% of the cell types possessed receptors for at least one
of these 12 hormones. This finding suggests that seasonal rhythms engage not only tissues that
demonstrate obvious morphological changes (e.g., adipose tissues and gonads), but involve
global shifts in cellular metabolism and physiology (**Fig. 6B**).

Discussion

Hormonal signaling is vital to multicellular life, allowing cells to communicate with both
surrounding cells and remote cells in other tissues. In this study, we systematically mapped
hormone-producing and -target cells across the 739 cell types from 27 different tissues from the
mouse lemur molecular cell atlas. We generated a browsable atlas that depicts gene expression of
ligands, modulators, and receptors for 87 hormone classes (**Fig. S1, Fig. 1**). This dataset
complements the scale and resolution of classical endocrine studies and brings a cellular and
molecular foundation for studying novel aspects of hormone regulation. The mouse lemur
hormone atlas represents a primate model of hormone signaling and should be useful in future
organism-wide cross-species comparisons.

The hormone atlas also provides an opportunity to extract global principles of hormone
signaling. The hormone-related genes exhibited cell-type-dependent, stereotypical expression
patterns, and their transcriptional profiles faithfully classified cell types, despite their constituting
less than 1% of the transcriptome (**Fig. 2, Fig. S3, Fig. S5**). Ligands alone and receptors alone
also clustered cell types well, with certain clusters mostly defined by their ligands and others by
their receptors (**Fig. S4**). It is remarkable, perhaps, how well a cell's intercellular communication
repertoire encapsulates the cell's identity.

There were, however, some exceptions to this general rule. For example, sperms and sperm
progenitor cells exhibited dramatic changes in hormone gene expression during spermatogenesis

despite their close developmental connection (**Fig. 3A**). In addition, vascular endothelial cells from lung and brain exhibited hormone signatures that were distinct from those of other vascular endothelial cells (**Fig. 3B**). Hormone signaling varied in an orderly and graded fashion along the epithelium and vasa recta endothelium of the nephron (**Fig. 3C and D**). Hepatocytes were found to fall into two molecular subtypes with differential hormonal gene expression (**Fig. 3E, Fig. S6**), suggesting different physiological roles. And finally, metastatic endometrial tumor cells exhibited different hormonal signaling from that seen in their likely cells of origin in the uterus (**Fig. 3F**), suggesting possible rewiring of hormone signaling during tumor development and metastasis.

The cell-type specific gene expression data allowed us to construct an organism-wide cell communication network for mouse lemur hormone signaling, by connecting the hormone-producing cells to the corresponding target cells (**Fig. 4**). This hormone network was exceptionally dense and robust to deletion of individual edges (**Fig. 4A, Fig. S7A**), which mainly resulted from two network features. First, most cells in the network expressed multiple hormone receptors and/or ligands, though in general cells were regulated by more hormones than they produced (**Fig. 4B**). Second, although most hormone ligands and receptors were expressed in specific cell types, a small fraction of hormone receptors were nearly ubiquitously expressed (e.g., adiponectin receptors, a membrane progesterone receptor, the IGF1 receptor, the thyroid hormone receptor, and the cortisol receptor), resulting in almost global connection of the relevant hormone-producing cells to the other cells of the body (**Fig. 4C-E, Fig. S8, Fig. S9**). Interestingly, despite the density of the hormone signaling network, subtypes of hormone receptors that bind to the same ligand were generally expressed in a mutually exclusive fashion, supporting the notion that different hormone receptor subtypes usually mediate different physiological functions through separate target cells (**Fig. 4F**). The dense connection of the network indicates extensive hormonal communication among the cells and contributes to a large number of highly connected nodes and no obvious network cores. Whereas regulatory circuits of individual hormones may be hierarchical, as classically viewed, the overall decentralized network architecture of the hormone interactome highlights the importance of taking a comprehensive approach to the understanding of global hormonal function.

Feedback regulation was widespread in hormonal signaling, and it was robust to network perturbations (**Fig. 5B, Fig. S7B**). The analysis identified well-known feedback circuits among several endocrine cell pairs and detected additional, previously unreported, hormones involved in feedback control (**Fig. 5C-F**). We also identified many novel feedback circuits (**Fig. 5G-L**) which may be involved in adaptation or homeostasis (negative feedback circuits) or in generating switch-like responses (positive and double-negative feedback circuits). This hypothesis-generating analysis provided an extensive list of potential feedback circuits (**Table 6**) to guide future functional validation and characterization studies. Our findings may also inform studies of the systems biology and network topology of hormone signaling (see for example Karin et al.[98]).

Mouse lemurs display dramatic seasonal changes in body weight, hibernation propensity, and reproductive activities[29,95,99]. Such seasonal rhythms are likely regulated, at least in part, by seasonally-varying hormones that coordinate the changes across different organs of the body (**Fig. 6A**). The hormone atlas shows that almost all (95%) of the mouse lemur cell types are regulated by seasonally varying hormones (**Fig. 6B**). This suggests that most, if not all, mouse lemur cells across all organs of the body experience seasonal changes in metabolism, growth,

and function. By systematic mapping of hormone secreting and target cells, the atlas also points
to candidate cell types and signaling to explore the upstream and downstream regulation for each
of the seasonally varying hormones. Circannual rhythms are widespread in nature[100,101] and
humans display seasonally varying hormone levels and cellular transcriptomes[102–104]. The
findings of the present study may apply to species beyond the mouse lemur, and may bring new
insights for season-associated human diseases[105–107] because of the close genetic proximity of
mouse lemurs and humans.

Single-cell RNA sequencing is a powerful tool for assaying cellular function and activity, but
there are limitations. Although the technique has improved rapidly in sensitivity, accuracy, and
efficiency, scRNAseq still faces low capture efficiency and high transcript dropout[108]. With the
large number of cells assayed in the mouse lemur molecular cell atlas, we averaged gene
expression across cells of the same cell type to smooth the noise in individual cells from dropout.
In the analysis of hormone regulation, genes involved in hormone production, such as ligands
and modulators, tend to be abundantly expressed; but certain hormone receptor proteins may be
low in copy number and slow in turnover, and therefore may result in false-negative errors. In
addition, there can be false-positive errors caused by contamination from other cells in the same
sample (i.e., signal spreading). This is most notable for abundantly expressed ligand genes and
tissues affected by endogenous digestive enzymes (e.g., pancreas). In this study, we scored
positive expression of a hormone ligand only when both the ligand gene and the necessary
modulators (such as prohormone processing enzymes) were present. This approach mitigated
false-positive detection of highly abundant ligands. Lastly, scRNAseq has low capture efficiency
for certain cells such as neurons and large sized adipocytes. Nevertheless, the mouse lemur cell
atlas included several neuronal subtypes from the brain cortex, brainstem, and retina, as well as
both UCP1-high and UCP1-low adipocytes from multiple tissues[14]. Although not yet a complete
sampling, the mouse lemur cell atlas is one of the most comprehensive and accurately-annotated
organism-wide scRNAseq datasets reported to date. Other organism-wide scRNAseq atlases
have been created or are being created for other species, including mouse[54,109,110], ascidian[111],
cynomolgus monkey[112], and human[113], bringing exciting resources to the study of the evolution
of hormone signaling, as well as the cellular and molecular mechanisms and interactions
underlying hormone-regulated biology.

Traditionally, biologists focus on one particular cell type and/or organ system. However,
hormone signaling by its very nature connects different cell types and organs. The emerging
organism-wide atlases open a new door in understanding the cross-organ communication
systems that allow organisms to function as a reliable robust whole.

Methods

Mapping and counting transcript reads: The single-cell gene expression matrix of the mouse
lemur cell atlas was obtained from The Tabula Microcebus[14] where methods and parameters to
extract the transcript counts can be found. In this study, we used only the 10x droplet-based
sequencing derived data for all analysis. In brief, the *Microcebus murinus* genome assembly

(Mmur 3.0, NCBI) was used to create a genome FASTA file. Sequencing reads were processed
with Cell Ranger (version 2.2, 10x Genomics) to generate cell barcodes and the gene expression
matrix. We additionally counted reads for three genes (MC1R, SCT, LHB) that are unannotated
in Mmur 3.0, NCBI. For MC1R and SCT, we used their annotation in the Ensembl *Microcebus*
murinus genome. For LHB, we used the uTAR-scRNA-seq pipeline described by Wang et al.[20]
to first predict its chromosome location in the mouse lemur genome. In brief, the pipeline
employs the approach of groHMM[114] to detect transcriptionally active regions (TARs) from
aligned sequencing data and annotates these TARs as unannotated (uTARs) or annotated
(aTARs) based on their overlap with the existing annotation. For the LHB_uTAR counts, we
uncovered uTARs from the mouse lemur pituitary and hypothalamus dataset and used
BLASTn[115,116] to find nucleotide sequence homology of all uTARs against human LHB. We
found one uTAR at position NC_033681.1:19143249-19145949 with high sequence homology
to human and macaque LHB. The LHB_uTAR is highly differentially expressed in the expected
cell type (gonadotroph). The LHB_uTAR is located next to gene RUVBL2, in conservation with
the co-location pattern of the two genes in humans and mice[21]. We therefore conclude that the
identified LHB_uTAR is the likely coding region of the mouse lemur LHB gene. We then used
Drop-seq tools[117] to extract transcript counts for the three genes (MC1R, SCT, LHB_uTAR).

Protein sequence analysis: We compared the sequence similarity between orthologs of human
and mouse or lemur. Protein sequences were retrieved from NCBI according to protein RefSeq
using the Matlab built-in function getgenpept. Conversion between gene ID and protein Refseq
identifier was performed using bioDBnet (https://biodbnet.abcc.ncifcrf.gov/db/db2db.php). We
aligned ortholog protein sequences and used the maximal scores in the cases of one-to-multi or
multi-to-multi ortholog mapping. Global alignment was carried out using the Needleman-
Wunsch algorithm and default parameters of the function nwalign.

Identification of hormone-producing and target cells: The functional concentration is
different for different proteins, so we applied adaptive thresholding to determine if a gene is
positively expressed in a cell type, where cell type is defined as a unique combination of the
cells' annotation and tissue of origin, as described in the accompanying Tabula Microcebus
manuscript 1[14]. For example, vein cells of the lung and brain were considered two different cell
types. Cell types that were of low quality (low gene count or transcript count) or annotated as
technical doublets were removed from the analysis. For each gene, two cell type parameters were
used, average expression level and percentage of nonzero cells. Single-cell expression levels
were first normalized by total transcript counts, multiplied by 10,000, averaged across all cells in
the cell type, then natural log transformed (i.e., log(1+AverageUMIcountPer10KTranscripts);
note that 1 is added to the average so that genes with zero transcripts will yield zero rather than
negative infinity). To determine positive expression at the cell type level, we first applied a low-
level absolute threshold on average expression level (0.037) and percent of nonzero cells (5%)
below either of which we considered the expression to be negative. Second, we calculated a
"maximal" expression level of the gene by searching the high-expression levels across the atlas.
Specifically, we considered the 99th percentile of the cells that passed the first thresholding to be
the maximum to increase the robustness of the analysis. Lastly, we applied relative thresholding
of the "maximal" level (for ligand genes at 20% and other gene types at 5%). Cell types that

passed all thresholding criteria were considered to have functional expression of the gene. We
used a higher relative threshold for the ligand genes because 1) ligands are secreted into the
circulation so a highly expressing source outshadows a low expressing source, whereas non-
ligand products of other genes (e.g., receptors, synthases, processing enzymes) function inside
individual cells; 2) we noted that ligand genes were usually expressed at higher levels and were
more easily affected by cross-contamination (signal spreading) inside a tissue. For hormone
ligands and receptors that included multiple genes, we applied AND or OR logic gates as
indicated in **Table 1** to determine if a cell type expressed the ligand or receptor. For an AND
logic gate, a functional cell type should express all its components above the functional level;
while for an OR logic gate, expression of any of its components would suffice.

Cell type distances, clustering, and visualization: We performed hierarchical clustering of the
cell types included in the mouse lemur atlas based on the transcriptional profiles of hormone
ligand, synthase, modulator, and receptor genes. Analysis was carried out at the cell type level
and relative average gene expression levels were used to calculate cell-type pairwise distances.
To test consistency of hormone-related gene expression among mouse lemur individuals, we
separated cells from different lemurs. Because circulating immune cells sampled from different
tissues were already kept as different cell types, immune cell types were not separated for
different lemurs to avoid overcrowding of the data. We found that this simplification only
slightly influenced the clustering results of individual cell types and did not affect the overall
clustering pattern. Next, cell types were filtered to remove potentially low quality cell types that
were: 1) low in cell number, 2) technical doublets, or 3) low in transcript and/or gene count. We
also manually corrected for notable cross-contamination in certain cell types. For example,
almost all non-endocrine cell types of the pancreas showed high expression of insulin (INS) and
proglucagon (GCG), probably caused by cross-contamination due to autolysis in the pancreas
during tissue processing. We therefore removed INS and GCG expression from all non-
endocrine pancreatic cell types. Similarly, we corrected contamination of GUCA2A, GUCA2B,
GIP, NTS, and SST from stromal, endothelial, and immune cells of the intestine (small intestine
and colon). Matlab built-in functions were used in the hierarchical clustering analysis. We used
cosine distances to calculate cell type pairwise distances and then applied average-linkage to
construct the dendrogram (cophenetic correlation coefficient = 0.80). Cell type clusters were then
divided at a threshold of 0.49 with minor manual inputs to further divide certain large clusters or
merge an apparently close neighboring leaf. We obtained 75 total clusters and numbered them
according to their sequence in the dendrogram shown in **Fig. 2A**. We also named the clusters
according to the shared characteristics of its cell types. Cell types included in each cluster are
listed in **Table 3** with the list of hormones, hormone receptors, and hormone-related genes that
were expressed in the majority ($\geq$ 50%) of the cell types for each cluster.

In addition to visualizing the cell type clustering via dendrogram, we also applied uniform
manifold approximation and projection (UMAP)[118] to embed the high dimensional expression
data to 2D using a Matlab UMAP package[119]. Cell type pairwise cosine distances were used as
input when generating the UMAPs.

For comparative analysis, we additionally calculated distances between cell types by either the
non-hormonal transcriptome or by Cell Ontology[43]. Transcriptome-based distances were

calculated similarly as the hormone-based distances were, with relative expression levels of non-
hormonal genes used to calculate pairwise cosine distances. For Cell Ontology distances, each
cell type was assigned a Cell Ontology term during cell type annotation as described in the
accompanying Tabula Microcebus manuscript 1[14]. Cell Ontology distances were calculated as
the mean of Cell Ontology embedding-based distances and text-based distances[44]. Cell Ontology
embedding-based distances represent distances between cell types in the Cell Ontology graph.
Text-based distances were calculated using an adapted version of Sentence-BERT[44,120].

To estimate the quality of the clustering results, we calculated silhouette values and compared
cluster average silhouette values based on only the hormone ligand genes, hormone receptor
genes, or the non-hormonal transcriptome. Silhouette values for a cell type i is from the equation
$s(i) = \frac{b(i)-a(i)}{max\{b(i),a(i)\}}$, for cell types belonging to a cluster with more than one member. $a(i)$ is the
mean in-cluster distance, namely the mean distance between cell type i and all other cell types in
the same cluster. $b(i)$ is the minimal out-cluster distance, namely the minimal mean distance
between cell type i and members of other clusters.

Vasa recta trajectory analysis: Analysis was performed independently for kidney samples from
three mouse lemur individuals using an in-house program written in Matlab. To detect the
arteriole-to-venule trajectory, data were first pre-processed following the standard procedure as
in Seurat[121]. In brief, single cell transcript counts for all sequenced genes were normalized to
total transcript counts, log transformed, and linearly scaled so that each gene has a mean
expression of 0 and standard deviation of 1. Then genes with high dispersion (variance-to-mean
ratio) compared to other genes with the same mean expression were selected as highly variable
genes for principal component analysis (PCA). Following PCA, the top 20 principal components
that were not driven by extreme outlier data points or immediate early genes were used to
generate a 2D UMAP using cell-cell Euclidean distances as input[118]. To quantify the trajectory,
we detected the probability density ridge of the data points in the UMAP using automated image
processing. The direction of the trajectory was manually assigned according to marker gene
expression. Individual cells were aligned to the trajectory by finding the shortest connecting
point to the trajectory. Data points that were too distant from the trajectory were deemed outliers
and removed from the following analysis. To detect the hormone-related genes that follow a
trajectory dependent expression pattern, we calculated the Spearman correlation coefficient
between the gene expression level and 20 unimodal patterns. These preassigned patterns have
their data smoothly changing along the trajectory and have a single peak. We let the 20 patterns
have their peaks uniformly distributed from 0 to 1. This analysis tested, for each gene, whether
its distribution along the trajectory follows one or more of the 20 unimodal patterns. All p values
were then multi-testing corrected to detect significant genes and their associated patterns. Lastly,
we compared results from the three kidney samples and kept the genes that showed consistent
patterns in all three samples.

Identification of differentially expressed genes: We used the Wilcoxon rank-sum test to detect
differentially expressed hormone genes among cell populations. Genes that were expressed
above functional level in any of the analyzed cell populations were included in the analysis. For

hormone-related genes differentially expressed during spermatogenesis, we tested whether each
gene is differentially expressed in one of the three sperm cell clusters compared to the other two
clusters. For the analysis of endothelial cells, we tested whether each gene is differentially
expressed in the lung or brain endothelial cells compared to all other endothelial cells. For the
kidney nephron analysis, we tested whether each gene is differentially expressed in one of the
nephron clusters compared to the rest of the nephron epithelial cells. For the hepatocyte analysis,
we tested whether each gene is differentially expressed in one subtype compared to the other in
both the mouse lemur and the mouse. The mouse hepatocyte data was obtained from the Tabula
Muris Senis[54] smartseq2 plate-based sequencing of mouse hepatocytes. For the tumor cell
analysis, we tested whether each gene is differentially expressed in the metastatic cells versus the
MUC16+ non-ciliated uterus epithelial cells. All p values were multi-testing corrected, and then
all genes with corrected p values less than 0.05 were examined to confirm consistent expression
of the gene in all relevant cell types and mouse lemur individuals.

Constructing and analyzing the hormonal cell communication network: Cells with the same
annotation name and tissue of origin were grouped as a cell type/node. Data from different lemur
individuals were merged in the analysis. Cell types that were likely low quality were removed.
Edge connections were determined independently for all possible cell type pairs. For example, if
cell type A expressed a hormone ligand and cell type B expressed the corresponding hormone
receptor(s), a directed edge from A to B was drawn. Note that this allows self-loops on any cell
type that expresses both the ligand and corresponding receptor. When multiple directed edges
connected two cell types, they were merged for the purpose of edge counting.

Network density is defined as the number of edges divided by the number of possible edges in
the network. Because the hormone network is directed, its network density is calculated as $\frac{N_{edges}}{N_{nodes}^2}$.
For comparison, the density of an undirected network (e.g., protein interaction network) is
calculated as $\frac{2*N_{edges}}{N_{nodes}\cdot(N_{nodes}+1)}$ if it allows self-loops, or $\frac{2*N_{edges}}{N_{nodes}\cdot(N_{nodes}-1)}$ if it does not allow self-
loops.

Outdegree and indegree are node features and defined as the relative number of outgoing edges
(node expressing a ligand) and incoming edges (node expressing a receptor). Edge counts were
normalized by cell type abundance so that node degrees range from 0-100% and represent the
fraction of the cell types that the node connects with by outgoing or incoming edges. To
normalize the node degrees, we used a weighted cell type abundance determined by the
clustering so that each cluster contributes the same weight. This is because certain cell types like
circulating immune cells are likely over-represented in the network as they appear in multiple
tissues and would be categorized as different "cell types" by our definition. We did not merge
circulating cell types from different tissues because it is challenging for some cell types like T
cells to determine whether they are circulating or potentially tissue specific or tissue-resident.
We also tried simply normalizing the node degree by the total number of nodes; similar
conclusions were reached using either method.

Calculating generality scores for hormone ligands and receptors: To calculate generality scores for the hormone ligands and receptors, we counted the cell types that express the ligand or receptor. Cell types that were likely low quality were removed from the analysis. Cell type counts were normalized by cell type abundances similarly as was the node degree, so that the generality score was not dominated by over-represented cell types. In comparison, we also counted the cell type clusters that express the hormone ligands and receptors. Here we defined that a gene was expressed in a cell type cluster if more than 50% of the cell types in the cluster positively expressed the gene. For hormone ligands and receptor complexes that contained multiple genes, we applied the AND or OR logic gates indicated in **Table 1** to determine positive expression in cell type clusters.

To analyze the distribution of generality scores, we tested if the distribution followed normal, exponential, lognormal, or gamma distributions. We first performed parametric distribution fitting and then performed Kolmogorov–Smirnov tests comparing the fitted versus data cumulative distribution functions. This analysis used Matlab built-in functions fitdist and kstest. Additionally, we tested if the data followed a power law distribution using plfit and plpva functions[122].

Calculating target cell Jaccard indices for different receptors that bind to the same ligand: We calculated Jaccard indices for pairs of receptors that 1) can bind to the same ligand as described in **Table 1**, and 2) were both expressed in at least one atlas cell type. Jaccard index here is calculated as the number of cell types that express both receptors divided by the number of cell types that express either of the receptors. Cell types that were likely low quality were removed from the analysis. Data from different mouse lemur individuals were merged in the analysis.

Network perturbation analysis: We performed two types of network perturbations: degree-preserving random rewiring and random edge removal. For degree-preserving random rewiring, we applied the algorithm developed by Maslov and Sneppen[80]. The algorithm starts with the original network and performs repeated rewiring steps. For each rewiring, it randomly identifies two edges (e.g., A→B and C→D) and swaps the connections (i.e., A→D and C→B). This allows constructing a random network while preserving the node indegree and outdegree of the original network. For the perturbation by edge removal, we randomly selected 10% of the edge types (ligand-receptor pairs) and removed these edges from the original network. Note for this analysis, multiple directed edges connecting the same cell type pairs were not merged. Therefore, the connection between two cell types would be removed only if all of its redundant edges (of different ligand-receptor pairs) were selected to be removed.

Hormone assays: Cortisol concentrations (ng/ml) were measured in duplicates from urine samples by Cortisol Urine ELISA (LDN®, catalog no. MS E-5100). Urine creatinine (mg/ml) was measured using MicrovueTM Creatinine Elisa kit (Quidel R Corporation, catalog no. 8009). Creatinine concentration was used to normalize urine measurements as an indicator of renal

1025   filtration activity. Cortisol measurements are thus expressed in ng/mg creatine. Thyroxine (T4,
1026   nmol/L) was assayed in plasma samples using T4 ELISA kit (LDN®, catalog no. TF E-2400)
1027   and the 6-point standard curve was adapted from 0 to 100 nmol/L.

1028

Acknowledgments

1030   We are grateful to Sandra Schmid, Uri Alon, Chaitan Khosla, the Krasnow lab, and the Ferrell
1031   lab for helpful discussions and comments on the manuscript. This work is supported by NIH
1032   grant R35 GM131792 to J. E. Ferrell, funding from Wall Center to M. A. Krasnow, and a
1033   fellowship from the Wu Tsai Neurosciences Institute to S. Liu. M. A. Krasnow is an investigator
1034   of the Howard Hughes Medical Institute.

1035

Author Contributions

S. Liu, M.A. Krasnow, and J.E. Ferrell conceived the project. S. Liu designed and performed all
the analysis with input from C. Ezran, Z. Li, C. Kuo, J. Epelbaum, J. Terrien, M.A. Krasnow,
and J.E. Ferrell. S. Liu, C. Kuo, and J. Long assembled the hormone list. M.F.Z. Wang and I.D.
Vlaminck performed uTAR analysis on the candidate lemur LHB gene. S. Wang calculated the
cell ontology distances. S. Liu, J. Epelbaum, and J. Terrien analyzed the seasonally oscillating
hormones. The Tabula Microcebus Consortium constructed the mouse lemur molecular cell
atlas. S. Liu and J.E. Ferrell wrote the manuscript with input from all authors.

1044

Figure legends

Fig. 1. **Expression of selected hormone ligands and receptors across the mouse lemur cell**
atlas. Four exemplary sets of hormone ligands and receptors are shown: prolactin (PRL, PRLR),
adiponectin (ADIPOQ, ADIPOR1, ADIPOR2), motilin (MLN, MLNR), and resistin (RETN,
CAP1, DCN, ROR1, TLR4). Cell types are arranged by compartment: epithelial/neural/germ
cells (**A**), stromal cells (**B**), endothelial cells (**C**), lymphoid cells (**D**), and myeloid cells (**E**).
Within each panel, rows are cell types ordered by tissue of origin. Cell type names are labeled on
the right, tissue of origin on the left, and gene symbols on the top (red for ligand genes, blue for
receptor genes). Dots represent average gene expression (circle color) and percent of cells
positive for the gene (circle size). The red/blue circle edges highlight the cell types deemed
positive for the ligand or receptor gene expression (see **Methods**). Cell types that are potentially
low quality (e.g., low cell count) are excluded from this plot. A complete set of figures that
display ligand and receptor gene expression for all 87 hormone classes can be found in **Fig. S1**.

1058

Fig. 2. Hormone-related gene expression faithfully classifies cell type. A. Clustering of mouse lemur cell types by hormone-related genes. At the top is a dendrogram of the hierarchical clustering. Clusters of cell types are color-coded, and cluster numbers are labeled as space allows. Cell types included in each cluster are listed in **Table 3**. The lower part of the panel shows the heatmap of the relative gene expression levels for the hormone-related genes. Cluster-specific hormone genes are ordered based on the cluster with the highest expression level, with the genes preferentially expressed in the left-hand cell type clusters on the top. Broadly expressed genes (expressed in more than 35% of cell types) are placed in the bottom and ordered from least to most broadly expressed. **B**. UMAP visualization of mouse lemur cell types based on the expression of hormone-related genes. Circles are cell types (unique combination of annotation name and tissue of origin) and color-coded by cell type compartment types (epithelial, neural, germ, stromal, endothelial, lymphoid, and myeloid). Dashed lines circumscribe the cell type clusters as in panel A (also see **Table 3**), and cluster IDs and names were labeled nearby. As UMAPs qualitatively display relationships among data points, the extremely distant clusters (i.e., cluster 44. mesothelial cells) were shifted up in UMAP-2 towards the center of the figure for display purposes. **C, D**. Scatter plots showing positive correlation (*p*-values < 1E-15) between cell type pairwise distances based on hormone-related gene expression versus the non-hormonal transcriptome (**C**) or Cell Ontology descriptors (**D**). Red lines show linear fitting of the data.

Fig. 3. Hormonal regulation sometimes differs in closely related cell types. Panels show expression of the hormone ligand, synthase, modulator, and receptor genes that are differentially expressed during spermatogenesis (**A**), endothelial cells of the lung and brain (**B**), epithelial cells along the spatial axis of the kidney nephron tubule (**C**), endothelial cells along the arteriole-to-venule axis of the kidney vasa recta (**D**), two molecular subtypes of hepatocytes (**E**), and between metastatic tumor cells in the lung of lemur L2 and the putative cell type of metastatic origin (MUC16+ non-ciliated uterus epithelial cells) sampled from L3 (**F**). **A-C, E and F** are dot plots showing average gene expression and percent of positive cells of the indicated genes and cell types, merging data from different mouse lemur individuals. In **A**, cell types are arranged along the course of the spermatogenesis. In **B**, endothelial cells are arranged by tissue of origin. In **C**, cell types are arranged according to their spatial distribution along the kidney nephron tubule. In **D**, on the top are a schematic representation of the kidney nephron and vasa recta and the UMAP visualization of the vasa recta cells based on expression of all genes. The black line shows the detected arteriole-to-venule trajectory with the arrow pointing in the direction of blood flow. Cells are color-coded according to the cell type annotation names in the atlas. Grey lines show the alignment of the individual cells to the trajectory. The bottom portion of the panel shows single cell scatter plots of gene expression levels along the arteriole-to-venule trajectory for the endothelial markers and hormone genes that are differentially expressed. Shown here is vasa recta data from one animal. Similar results were obtained from two other lemurs. **PCT**: proximal convoluted tubule. **PT**: proximal tubule. **PST**: proximal straight tubule (also called loop of Henle thick descending limb). **LHd**: loop of Henle thin descending limb. **LHa1**: loop of Henle thin ascending limb. **LHa2**: loop of Henle thick ascending limb. **DCT**: distal convoluted tubule. **MD**: macula densa. **CDp**: collecting duct principal cell. **CDiα**: collecting duct α intercalated cell. **CDiβ**: collecting duct β intercalated cell. **D** shows the fitted arteriole-to-venule trajectory of the kidney vasa recta capillary network (upper panel) and single-cell gene expression levels along the trajectory for the endothelial markers and differentially expressed

1104 hormone-related genes. Shown here are representative results from one of three lemur vasa recta
1105 samples. **VRD**: vasa recta descending section. **VRA**: vasa recta ascending section.

1106

Fig. 4. The hormonal cell-cell communication network. **A.** A representation of the network as
constructed by force-directed graph drawing. Nodes (cell types) are color-coded according to the
node outdegree (% of cells connected to the current node by outgoing edges) and the node circle
size shows the node indegree (% of cells connected to the current node by incoming edges).
Network edges are directed and connect nodes expressing a hormone ligand to all nodes
expressing the corresponding hormone receptor. **B.** Number of ligand and receptor types
expressed by the network nodes (cell types), ranked from most to fewest. **C.** Indegree and
outdegree of the network nodes (cell types), ranked from high to low values. Node degree is
normalized to the total number of nodes and cluster size and measures the percentage of cell
types connected from or to a network node. **D.** Generality score of all hormone ligands and
receptors ranked from the most generally expressed to most selectively expressed. Generality
score is defined as the percentage of nodes (cluster size normalized) positively expressing the
gene(s) involved in ligand synthesis or receptor binding. Also see **Fig. S9A** for generality score
defined as the number of cell type clusters positively expressing the gene(s). **E.** Scatterplot
showing generality scores of the hormone ligands and corresponding receptors. Each circle is a
unique ligand-receptor pair. **F.** Ranking (left) and distribution (right) of Jaccard indices of
receptor pairs binding to the same ligand across all hormone receptors. Jaccard index is
calculated as the number of cell types co-expressing both receptors divided by the number of cell
types expressing either of the receptors.

1126

Fig. 5. Feedback circuits in the hormonal cell-cell communication network. A. Definition of
two-node feedback circuits. **B**. Comparison of the number of 2-node feedback circuits identified
in the hormonal cell-cell communication network (red dashed line) and that of permuted
networks (n = 1000). Permutation preserved node outdegree and indegree. Note that none of the
permuted networks has as many 2-node feedback circuits as in the original network. **C-H**.
Examples of two-node feedback circuits identified in the network, focusing on the endocrine cell
types. Solid black arrows indicate known regulation, dashed orange arrows indicate predicted
regulation with partial literature support, and dotted red arrows predicted regulation without
earlier knowledge. Parentheses are used to group multiple genes that are expressed in the
relevant cell type that function either together (&) or independently (/).

1137

Fig. 6. Seasonality of hormone concentrations in the mouse lemur likely affects almost all
cells and tissues. A. Concentrations of twelve hormones measured in summer (long-
photoperiod) and winter (short-photoperiod) in the mouse lemur. Bars represent fold changes in
hormone concentration compared to average summer levels. Error bars represent 95%
confidence intervals. *P*-values were calculated by 2-sample *t*-test. Data were listed in **Table 7**,
collected from literature and supplemented with additional new measurements: estradiol[123,124],
testosterone[125–127], thyroxine[123,128] (and additional data in Table 7), DHEA[124], cortisol[129] (and

additional data in Table 7), IGF1[130], melatonin[131], and gut hormones (GIP, PPY, insulin, PYY,
and GLP-1)[132,133]. **B.** Mouse lemur cell types targeted by any of these 12 seasonal hormones. Cell
types are displayed in the same UMAP plot as in **Fig. 2B**. Circles are cell types and are color
coded according to whether they express receptors for any of the 12 hormones. Dashed lines
circumscribe the cell type clusters and are colored according to cell type compartments.

1150

Fig. S1. Ligand and receptor gene expression for 87 hormone classes across the mouse
lemur cell atlas. Each dot plot shows one of the 87 hormone groups and one of the five cell
compartment types (1. epithelial/neural/germ cells; 2. endothelial cells; 3. stromal cells; 4.
lymphoid cells; and 5. myeloid cells). The hormone group name and cell compartment type is
indicated on the top right of the figure. The dot plots are arranged in a polar coordinate style.
Radial lines represent individual cell types, with cell type names labeled inside the inner circle
and the tissue source labeled outside the outer circle. Cell types are arranged by their tissue of
origin. Circular lines represent the related genes for this hormone group (red for ligands or
synthases, light red for processing enzymes and/or other modulator genes, and blue for receptor
genes). The dark colored dots on the intersections of the radial lines and circular lines depict
mean gene expression level (by dot shade) and percent of positive cells (by dot size) for the
respective gene and cell type. The outermost red or blue dots indicate positive expression of one
of the hormone(s) (red) or hormone receptor(s) (blue), or both (purple) for the hormone group by
the respective cell type (see **Methods**). The outermost grey dots indicate potentially low-quality
cell types (e.g., low cell number (<10) and pancreatic cells that show notable signal
contamination) whether or not the cell type expresses one the hormone ligand(s) (red outline),
receptor(s) (blue outline), both (purple outline), or neither (no outline).

1168

1169 **Fig. S2. Protein sequences of hormonal genes are generally more similar between human**
1170 **and mouse lemur than human and mouse.** Each circle is a hormone-related protein. Sequences
1171 were aligned by the Needleman-Wunsch algorithm and sequence identity was calculated as the
1172 percentage of identical amino acids. Points below the red line are proteins more similar between
1173 human and mouse lemur than between human and mouse.

1174

Fig. S3. UMAP visualization of mouse lemur cell types based on the non-hormonal
transcriptome. Circles (cell types) are color-coded by cell compartment types (epithelial, neural,
germ, stromal, endothelial, lymphoid, and myeloid). Dashed lines circumscribe cell type clusters
as in **Fig. 2A-B** and cluster IDs and names are labeled next to the cell type clusters. Cell types
that were clustered by the hormone-related genes but not the non-hormonal transcriptome are
labeled with asterisks (*). Note some cell types that formed discrete clusters based on hormonal
gene expression intermingled with other clusters when mapped based on the non-hormonal
transcriptome (e.g., cluster 51-52 and 54-58), and therefore were annotated together.

1183

Fig. S4. Classification of mouse lemur cell types using only the hormone ligand genes (A) or the hormone receptor genes (B). In panels **A** and **B**, cell types are displayed by the UMAP algorithm according to the cell type distances calculated with either the hormone ligands/modulators or the hormone receptors respectively. Circles (cell types) are color-coded by the cell compartment types (epithelial, neural, germ, stromal, endothelial, lymphoid, and myeloid). Dashed lines circumscribe cell type clusters as in **Fig. 2A-B** and cluster IDs are annotated nearby. Cell types that were clustered using all the hormone-related genes but not in this analysis are labeled with asterisks (*). As UMAPs qualitatively display relationships among data points, the extremely distant clusters in panel A (i.e., cluster 44. mesothelial cells by hormone ligand genes) were shifted towards the center of the figure for display purposes. **C.** Cluster averaged silhouette values of the 68 cell type clusters by both hormone ligand and receptor genes (black), only the hormone ligand genes (red), only the hormone receptor genes (blue), or the non-hormonal transcriptome (blank). The silhouette values are a measure of how well each cell type cluster was classified (see **Methods**).

Fig. S5. Heatmap representation of cell type pairwise distances calculated by three different methods. A. Hormone-gene transcriptional cosine distances. **B.** Non-hormonal gene transcriptional cosine distances. **C.** Cell Ontology-based distances. Cell types were ordered by cluster as in **Fig. 2A**, and cluster IDs were labeled along the diagonal for large clusters.

Fig. S6. Dot plot showing the differentially expressed hormone ligands/synthases and receptors separately in the hepatocytes of male and female animals (in comparison to **Fig. 3E).**

Fig. S7. The hormonal cell-cell communication network is robust to edge removal. Distribution of network density (**A**) and number of 2-node feedback circuits (**B**) in networks where 10% of ligand-receptor connections were randomly removed.

Fig. S8. Node properties of the hormonal cell-cell communication network. Heatmap of indegree, outdegree, number of hormone ligands and receptors expressed, and six node centrality measurements (hubs, authorities, incloseness, outcloseness, pagerank, and betweenness) for all nodes (cell types) of the hormonal cell-cell communication network. Because the scales of the node properties are different, shown here are normalized values linearly scaled by median and median-absolute-deviation. Cell types (columns) are ordered by the clustering of the hormone-related genes (as in **Fig. 2A**) and cluster numbers are labeled as space allowed. Node properties (rows) are ordered by hierarchical clustering of the normalized values.

Fig. S9. Generality scores for the hormone ligands and receptors. A. Generality score, defined as the number of clusters with positive expression of the ligand or receptor (see **Fig. 4C**

1223 for comparison), ranked from most generally expressed to most selectively expressed. **B-C**.
1224 Fitting the distribution of the ligand and receptor generality scores. Normal, exponential, log-
1225 normal, and gamma distributions failed to fit the distribution (**B**). The tail of the distribution can
1226 be approximated by a power law distribution $prob(\mathrm{x}) \sim \mathrm{x}^{-\alpha}$ for $\mathrm{x} \geq \mathrm{xmin}$ (**C**).

1227

Supplemental Tables

1228

Table 1. Genes involved in the biosynthesis and sensing of 87 classes of hormones. Rows
show hormones and columns show the hormone class name, hormone name(s), type of
hormones, symbol of the genes involved in biosynthesis (Ligand/Enzyme Gene(s)) and sensing
(Receptor Gene(s)), whether the hormone synthesis or maturation requires coordination of
multiple tissues, years discovered, classical sites of secretion and targets, and approximate
plasma concentrations in humans. Genes in the table are human genes.

1235

1236 **Table 2. Mouse lemur and mouse orthologs of the human hormone-related genes in Table**
1237 **1.** Rows show genes and columns show gene type, symbols of human, lemur, and mouse genes
1238 respectively. Special cases such as unknown/missing orthologs were highlighted and labeled
1239 with *.

1240

1241 **Table 3. Clusters of mouse lemur cell types according to expression of hormone-related**
1242 **genes.** Rows show clusters and columns show cluster ID, name, cell types included in the
1243 cluster, and the hormones, receptors, and hormone-related genes that were positively expressed
1244 in the majority ($\geq 50\%$) of the cluster cell types.

1245

1246 **Table 4. Hormones that regulate or are produced by the anterior pituitary neuroendocrine**
1247 **cells.** Rows show the hormones and hormone receptors expressed in individual pituitary
1248 neuroendocrine cell types. Columns show the corresponding genes, hormone names, and
1249 upstream/downstream cell types.

1250

1251 **Table 5. Generality scores of the hormone ligands, modulators and receptors.** Rows show
1252 hormone ligands, modulators, or receptors. Columns show entry types, symbols of human and
1253 mouse lemur genes, whether the entry includes a single or multiple genes, and generality scores
1254 calculated as the percentage of positive cell types and the number of positive clusters.

1255

Table 6. Two node feedback circuits detected from the mouse lemur hormone cell-cell communication network. Rows show the detected circuits and columns show the two cell types and hormone signaling that connects the cell types.

Table 7. Individual measurements of concentrations of mouse lemur seasonally-changing hormones. Rows show individual measurements and columns show the season of measurements, concentration, and hormone names.

References

1. W. M. Bayliss, E. H. S. The mechanism of pancreatic secretion. *J. Physiol.* **28**, 325 (1902).
2. Takei, Y., Ando, H. & Tsutsui, K. *Handbook of Hormones: Comparative Endocrinology for Basic and Clinical Research*. (Academic Press, 2015).
3. Tata, J. R. One hundred years of hormones. *EMBO Rep.* **6**, 490 (2005).
4. Lewis, G. F. & Brubaker, P. L. The discovery of insulin revisited: lessons for the modern era. *J. Clin. Invest.* **131**, (2021).
5. Horby, P. *et al.* Dexamethasone in Hospitalized Patients with Covid-19. *N. Engl. J. Med.* **384**, (2021).
6. Sterne, J. A. C. *et al.* Association Between Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-19: A Meta-analysis. *JAMA* **324**, (2020).
7. Scheja, L. & Heeren, J. The endocrine function of adipose tissues in health and cardiometabolic disease. *Nat. Rev. Endocrinol.* **15**, 507–524 (2019).
8. Jabbour, H. N., Kelly, R. W., Fraser, H. M. & Critchley, H. O. D. Endocrine regulation of menstruation. *Endocr. Rev.* **27**, 17–46 (2006).
9. Ezran, C. *et al.* The Mouse Lemur, a Genetic Model Organism for Primate Biology, Behavior, and Health. *Genetics* **206**, 651–664 (2017).
10. Bons, N., Rieger, F., Prudhomme, D., Fisher, A. & Krause, K. H. Microcebus murinus: a useful primate model for human cerebral aging and Alzheimer's disease? *Genes Brain Behav.* **5**, (2006).
11. Languille, S. *et al.* The grey mouse lemur: a non-human primate model for ageing studies. *Ageing Res. Rev.* **11**, (2012).
12. Kraska, A. *et al.* Age-associated cerebral atrophy in mouse lemur primates. *Neurobiol. Aging* **32**, 894–906 (2011).
13. Pifferi, F. *et al.* Promoting healthspan and lifespan with caloric restriction in primates. *Commun Biol* **2**, 107 (2019).
14. The Tabula Microcebus Consortium. Tabula Microcebus: A transcriptomic cell atlas of mouse lemur, an emerging primate model organism. *bioRxiv* 2021.12.12.469460 (2021) doi:10.1101/2021.12.12.469460.
15. Kastin, A. *Handbook of Biologically Active Peptides*. (Academic Press, 2013).
16. Romere, C. *et al.* Asprosin, a Fasting-Induced Glucogenic Protein Hormone. *Cell* **165**, 566–579 (2016).
17. Itoh, N., Ohta, H. & Konishi, M. Endocrine FGFs: Evolution, Physiology, Pathophysiology, and Pharmacotherapy. *Front. Endocrinol.* **6**, 154 (2015).
18. Ramilowski, J. A. *et al.* A draft network of ligand–receptor-mediated multicellular signalling in human. *Nature Communications* vol. 6 (2015).
19. Armstrong, J. F. *et al.* The IUPHAR/BPS Guide to PHARMACOLOGY in 2020: extending

immunopharmacology content and introducing the IUPHAR/MMV Guide to MALARIA
PHARMACOLOGY. *Nucleic Acids Res.* **48**, D1006–D1021 (2020).

20. Wang, M. F. Z. *et al.* Uncovering transcriptional dark matter via gene annotation independent single-
cell RNA sequencing analysis. *Nat. Commun.* **12**, 1–10 (2021).

21. Parfait, B. *et al.* Human TIP49b/RUVBL2 gene: genomic structure, expression pattern, physical link
to the human CGB/LHB gene cluster on chromosome 19q13.3. *Ann. Genet.* **43**, 69–74 (2000).

22. Wilkinson, T. N., Speed, T. P., Tregear, G. W. & Bathgate, R. A. D. Evolution of the relaxin-like
peptide family. *BMC Evol. Biol.* **5**, 14 (2005).

23. Riddle, O., Bates, R. W. & Dykshorn, S. W. THE PREPARATION, IDENTIFICATION AND
ASSAY OF PROLACTIN—A HORMONE OF THE ANTERIOR PITUITARY. *American Journal*
of Physiology-Legacy Content vol. 105 191–216 (1933).

24. Bole-Feysot, C., Goffin, V., Edery, M., Binart, N. & Kelly, P. A. Prolactin (PRL) and its receptor:
actions, signal transduction pathways and phenotypes observed in PRL receptor knockout mice.
Endocr. Rev. **19**, 225–268 (1998).

25. Tovar, S. & Diéguez, C. Prolactin and energy homeostasis: pathophysiological mechanisms and
therapeutic considerations. *Endocrinology* vol. 155 659–662 (2014).

26. Paré, P. *et al.* Molecular evolutionary insights from PRLR in mammals. *Gen. Comp. Endocrinol.*
309, 113791 (2021).

27. Giroud, S. *et al.* The Grey Mouse Lemur Uses Season-Dependent Fat or Protein Sparing Strategies
to Face Chronic Food Restriction. *PLoS ONE* vol. 5 e8823 (2010).

28. Terrien, J. *et al.* Metabolic and genomic adaptations to winter fattening in a primate species, the grey
mouse lemur (Microcebus murinus). *International Journal of Obesity* vol. 42 221–230 (2018).

29. Perret, M. & Aujard, F. Regulation by Photoperiod of Seasonal Changes in Body Mass and
Reproductive Function in Gray Mouse Lemurs (Microcebus murinus): Differential Responses by
Sex. *Int. J. Primatol.* **22**, 5–24 (2001).

30. Galic, S., Oakhill, J. S. & Steinberg, G. R. Adipose tissue as an endocrine organ. *Molecular and*
Cellular Endocrinology vol. 316 129–139 (2010).

31. Maeda, K. *et al.* cDNA cloning and expression of a novel adipose specific collagen-like factor,
apM1 (AdiPose Most abundant Gene transcript 1). *Biochem. Biophys. Res. Commun.* **221**, 286–289
(1996).

32. Zhang, Y. *et al.* Positional cloning of the mouse obese gene and its human homologue. *Nature* **372**,
(1994).

33. Ruan, H. & Dong, L. Q. Adiponectin signaling and function in insulin target tissues. *Journal of*
Molecular Cell Biology vol. 8 101–109 (2016).

34. Yamauchi, T. *et al.* Cloning of adiponectin receptors that mediate antidiabetic metabolic effects.
Nature **423**, 762–769 (2003).

35. He, J., Irwin, D. M., Chen, R. & Zhang, Y.-P. Stepwise loss of motilin and its specific receptor genes
in rodents. *J. Mol. Endocrinol.* **44**, 37–44 (2010).

36. Sanger, G. J., Wang, Y., Hobson, A. & Broad, J. Motilin: towards a new understanding of the
gastrointestinal neuropharmacology and therapeutic use of motilin receptor agonists. *Br. J.*
Pharmacol. **170**, 1323–1332 (2013).

37. Hu, Q., Tan, H. & Irwin, D. M. Evolution of the Vertebrate Resistin Gene Family. *PLoS One* **10**,
e0130188 (2015).

38. Schwartz, D. R. & Lazar, M. A. Human Resistin: Found in Translation From Mouse to Man. *Trends*
Endocrinol. Metab. **22**, 259 (2011).

39. Zhang, X. *et al.* Neuropeptidomic Analysis Establishes a Major Role for Prohormone Convertase-2
in Neuropeptide Biosynthesis. *J. Neurochem.* **112**, 1168 (2010).

40. Seidah, N. G. The proprotein convertases, 20 years later. *Methods Mol. Biol.* **768**, 23–57 (2011).

41. Sandoval, D. A. & D'Alessio, D. A. Physiology of proglucagon peptides: role of glucagon and GLP-
1 in health and disease. *Physiol. Rev.* **95**, 513–548 (2015).

42. Wardlaw, S. L. Hypothalamic proopiomelanocortin processing and the regulation of energy balance.

Eur. J. Pharmacol. **660**, 213 (2011).

43. Smith, B. *et al.* The OBO Foundry: coordinated evolution of ontologies to support biomedical data
integration. *Nat. Biotechnol.* **25**, 1251–1255 (2007).

44. Wang, S. *et al.* Unifying single-cell annotations based on the Cell Ontology. *Cold Spring Harbor*
Laboratory 810234 (2020) doi:10.1101/810234.

45. Griswold, M. D. Spermatogenesis: The Commitment to Meiosis. *Physiol. Rev.* **96**, 1–17 (2016).

46. Smith, L. B. & Walker, W. H. The regulation of spermatogenesis by androgens. *Seminars in Cell &*
Developmental Biology vol. 30 2–13 (2014).

47. Ruwanpura, S. M., McLachlan, R. I. & Meachem, S. J. Hormonal regulation of male germ cell
development. *Journal of Endocrinology* vol. 205 117–131 (2010).

48. Lishko, P. V., Botchkina, I. L. & Kirichok, Y. Progesterone activates the principal Ca2+ channel of
human sperm. *Nature* **471**, 387–391 (2011).

49. Lue, Y. *et al.* Functional role of progestin and the progesterone receptor in the suppression of
spermatogenesis in rodents. *Andrology* **1**, 308–317 (2013).

50. Burnicka-Turek, O. *et al.* Inactivation of insulin-like factor 6 disrupts the progression of
spermatogenesis at late meiotic prophase. *Endocrinology* **150**, 4348–4357 (2009).

51. Dyer, L. A. & Patterson, C. Development of the endothelium: an emphasis on heterogeneity. *Semin.*
Thromb. Hemost. **36**, 227–235 (2010).

52. Groneberg, D. A., Springer, J. & Fischer, A. Vasoactive intestinal polypeptide as mediator of
asthma. *Pulm. Pharmacol. Ther.* **14**, 391–401 (2001).

53. Travaglini, K. J. *et al.* A molecular cell atlas of the human lung from single-cell RNA sequencing.
Nature **587**, 619–625 (2020).

54. Tabula Muris Consortium. A single-cell transcriptomic atlas characterizes ageing tissues in the
mouse. *Nature* **583**, 590–595 (2020).

55. Barnes, K. & Turner, A. J. The endothelin system and endothelin-converting enzyme in the brain:
molecular and cellular studies. *Neurochem. Res.* **22**, 1033–1040 (1997).

56. Wilhelms, D. B. *et al.* Deletion of prostaglandin E2 synthesizing enzymes in brain endothelial cells
attenuates inflammatory fever. *J. Neurosci.* **34**, 11684–11690 (2014).

57. Engström, L. *et al.* Lipopolysaccharide-induced fever depends on prostaglandin E2 production
specifically in brain endothelial cells. *Endocrinology* **153**, 4849–4861 (2012).

58. Giesecke, T. *et al.* Vasopressin Increases Urinary Acidification via V1a Receptors in Collecting Duct
Intercalated Cells. *Journal of the American Society of Nephrology* vol. 30 946–961 (2019).

59. Pannabecker, T. L. & Dantzler, W. H. Three-dimensional architecture of inner medullary vasa recta.
Am. J. Physiol. Renal Physiol. **290**, F1355–66 (2006).

60. Barry, D. M. *et al.* Molecular determinants of nephron vascular specialization in the kidney. *Nat.*
Commun. **10**, 5705 (2019).

61. Waxman, D. J. & O'Connor, C. Growth hormone regulation of sex-dependent liver gene expression.
Mol. Endocrinol. **20**, 2613–2629 (2006).

62. Backes, F. J. *et al.* Estrogen receptor-alpha as a predictive biomarker in endometrioid endometrial
cancer. *Gynecol. Oncol.* **141**, 312–317 (2016).

63. Shen, F., Gao, Y., Ding, J. & Chen, Q. Is the positivity of estrogen receptor or progesterone receptor
different between type 1 and type 2 endometrial cancer? *Oncotarget* vol. 8 506–511 (2017).

64. Nakanishi, M. & Rosenberg, D. W. Multifaceted roles of PGE2 in inflammation and cancer. *Semin.*
Immunopathol. **35**, 123–137 (2013).

65. Ke, J. *et al.* Prostaglandin E2 (PGE2) promotes proliferation and invasion by enhancing SUMO-1
activity via EP4 receptor in endometrial cancer. *Tumour Biol.* **37**, 12203–12211 (2016).

66. Lee, J. O. *et al.* Resistin, a fat-derived secretory factor, promotes metastasis of MDA-MB-231
human breast cancer cells through ERM activation. *Sci. Rep.* **6**, 18923 (2016).

67. Frankenstein, Z., Alon, U. & Cohen, I. R. The immune-body cytokine network defines a social
architecture of cell interactions. *Biol. Direct* **1**, (2006).

68. Giot, L. *et al.* A protein interaction map of Drosophila melanogaster. *Science* **302**, 1727–1736

(2003).

69. Yook, S. H., Oltvai, Z. N. & Barabási, A. L. Functional and topological characterization of protein interaction networks. *Proteomics* **4**, (2004).

70. Rolland, T. *et al.* A proteome-scale map of the human interactome network. *Cell* **159**, 1212–1226 (2014).

71. Costanzo, M. *et al.* The Genetic Landscape of a Cell. *Science* **327**, 425–431 (2010).

72. Barabási, A.-L. Scale-Free and Hierarchical Structures in Complex Networks. *AIP Conference Proceedings* (2003) doi:10.1063/1.1571285.

73. Iams, W. T. & Lovly, C. M. Molecular Pathways: Clinical Applications and Future Direction of Insulin-like Growth Factor-1 Receptor Pathway Blockade. *Clin. Cancer Res.* **21**, 4270–4277 (2015).

74. Ghosh, P., Dahms, N. M. & Kornfeld, S. Mannose 6-phosphate receptors: new twists in the tale. *Nat. Rev. Mol. Cell Biol.* **4**, 202–212 (2003).

75. Larhammar, D. & Salaneck, E. Molecular evolution of NPY receptor subtypes. *Neuropeptides* **38**, 141–151 (2004).

76. Kilpatrick, L. E., Humphrys, L. J. & Holliday, N. D. A G protein-coupled receptor dimer imaging assay reveals selectively modified pharmacology of neuropeptide Y Y1/Y5 receptor heterodimers. *Mol. Pharmacol.* **87**, (2015).

77. Gehlert, D. R., Schober, D. A., Morin, M. & Berglund, M. M. Co-expression of neuropeptide Y Y1 and Y5 receptors results in heterodimerization and altered functional properties. *Biochem. Pharmacol.* **74**, 1652–1664 (2007).

78. Almabouada, F. *et al.* Adiponectin Receptors Form Homomers and Heteromers Exhibiting Distinct Ligand Binding and Intracellular Signaling Properties. *J. Biol. Chem.* **288**, 3112–3125 (2013).

79. Keshvari, S. *et al.* Characterisation of the adiponectin receptors: the non-conserved N-terminal region of AdipoR2 prevents its expression at the cell-surface. *Biochem. Biophys. Res. Commun.* **432**, 28–33 (2013).

80. Maslov, S. & Sneppen, K. Specificity and stability in topology of protein networks. *Science* **296**, 910–913 (2002).

81. Dwyer, A. A. & Quinton, R. Anatomy and Physiology of the Hypothalamic-Pituitary-Gonadal (HPG) Axis. *Advanced Practice in Endocrinology Nursing* 839–852 (2019) doi:10.1007/978-3-319-99817-6_43.

82. Martín-Estal, I., de la Garza, R. G. & Castilla-Cortázar, I. Intrauterine Growth Retardation (IUGR) as a Novel Condition of Insulin-Like Growth Factor-1 (IGF-1) Deficiency. *Rev. Physiol. Biochem. Pharmacol.* **170**, 1–35 (2016).

83. Inagaki, T. *et al.* Inhibition of growth hormone signaling by the fasting-induced hormone FGF21. *Cell Metab.* **8**, 77–83 (2008).

84. Hartig, S. M. & Cox, A. R. Paracrine signaling in islet function and survival. *J. Mol. Med.* **98**, 451–467 (2020).

85. Habegger, K. M. *et al.* The metabolic actions of glucagon revisited. *Nat. Rev. Endocrinol.* **6**, 689–697 (2010).

86. Habegger, K. M. *et al.* Fibroblast growth factor 21 mediates specific glucagon actions. *Diabetes* **62**, 1453–1463 (2013).

87. Mu, J. *et al.* FGF21 analogs of sustained action enabled by orthogonal biosynthesis demonstrate enhanced antidiabetic pharmacology in rodents. *Diabetes* **61**, 505–512 (2012).

88. Mancuso, E. *et al.* Insulin-like growth factor-1 is a negative modulator of glucagon secretion. *Oncotarget* **8**, 51719–51732 (2017).

89. Kopchick, J. J., Berryman, D. E., Puri, V., Lee, K. Y. & Jorgensen, J. O. L. The effects of growth hormone on adipose tissue: old observations, new mechanisms. *Nat. Rev. Endocrinol.* **16**, 135–146 (2020).

90. Carré, N. & Binart, N. Prolactin and adipose tissue. *Biochimie* vol. 97 16–21 (2014).

91. Iwen, K. A. *et al.* Melanocortin crosstalk with adipose functions: ACTH directly induces insulin resistance, promotes a pro-inflammatory adipokine profile and stimulates UCP-1 in adipocytes. *J.*

Endocrinol. **196**, (2008).

92. Sarmento-Cabral, A. *et al.* Adipokines (Leptin, Adiponectin, Resistin) Differentially Regulate All Hormonal Cell Types in Primary Anterior Pituitary Cell Cultures from Two Primate Species. *Sci. Rep.* **7**, 43537 (2017).

93. Ferrell, J. E., Jr. Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr. Opin. Cell Biol.* **14**, 140–148 (2002).

94. García-Martínez, J. M., Chocarro-Calvo, A., De la Vieja, A. & García-Jiménez, C. Insulin drives glucose-dependent insulinotropic peptide expression via glucose-dependent regulation of FoxO1 and LEF1/β-catenin. *Biochim. Biophys. Acta* **1839**, 1141–1150 (2014).

95. Petter-Rousseaux, A. Age of Microcebus murinus at the onset of testicular development : Preliminary observations on photoperiodic effect. *Ann. Biol. anim. Bioch. Biophys.* **19**, 1801–1806 (1979).

96. Perret, M. Environmental and social determinants of sexual function in the male lesser mouse lemur (Microcebus murinus). *Folia Primatol.* **59**, 1–25 (1992).

97. Epelbaum, J. & Terrien, J. Mini-review: Aging of the neuroendocrine system: Insights from nonhuman primate models. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **100**, (2020).

98. Karin, O. *et al.* A new model for the HPA axis explains dysregulation of stress hormones on the timescale of weeks. *Mol. Syst. Biol.* **16**, e9510 (2020).

99. Perret, M. & Aujard, F. Daily hypothermia and torpor in a tropical primate: synchronization by 24-h light-dark cycle. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* vol. 281 R1925–R1933 (2001).

100. Gwinner, E. *Circannual Rhythms: Endogenous Annual Clocks in the Organization of Seasonal Processes*. (Springer, 2011).

101. Lincoln, G. A brief history of circannual time. *J. Neuroendocrinol.* **31**, (2019).

102. Tendler, A. *et al.* Hormone seasonality in medical records suggests circannual endocrine circuits. *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

103. Sailani, M. R. *et al.* Deep longitudinal multiomics profiling reveals two biological seasonal patterns in California. *Nat. Commun.* **11**, 4933 (2020).

104. Dopico, X. C. *et al.* Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat. Commun.* **6**, 7000 (2015).

105. Moriyama, M., Hugentobler, W. J. & Iwasaki, A. Seasonality of Respiratory Viral Infections. *Annual review of virology* **7**, (2020).

106. Marti-Soler, H. *et al.* Seasonal Variation of Overall and Cardiovascular Mortality: A Study in 19 Countries from Different Geographic Locations. *PLoS One* **9**, e113500 (2014).

107. Kurlansik, S. L. & Ibay, A. D. Seasonal affective disorder. *Am. Fam. Physician* **86**, (2012).

108. Chen, G., Ning, B. & Shi, T. Single-Cell RNA-Seq Technologies and Related Computational Data Analysis. *Front. Genet.* **10**, 317 (2019).

109. The Tabula Muris Consortium. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* **562**, 367–372 (2018).

110. Han, X. *et al.* Mapping the Mouse Cell Atlas by Microwell-Seq. *Cell* **172**, (2018).

111. Cao, C. *et al.* Comprehensive single cell transcriptome lineages of a proto-vertebrate. *Nature* **571**, 349 (2019).

112. Han, L. *et al.* Single-cell atlas of a non-human primate reveals new pathogenic mechanisms of COVID-19. *Cold Spring Harbor Laboratory* 2020.04.10.022103 (2020) doi:10.1101/2020.04.10.022103.

113. Tabula Sapiens Consortium* *et al.* The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. *Science* **376**, eabl4896 (2022).

114. Chae, M., Danko, C. G. & Lee Kraus, W. groHMM: a computational tool for identifying unannotated and cell type-specific transcription units from global run-on sequencing data. *BMC Bioinformatics* **16**, 1–16 (2015).

115. Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res.* **36**, (2008).

1504   116. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search
1505   tool. *J. Mol. Biol.* **215**, 403–410 (1990).
1506   117. Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
1507   Nanoliter Droplets. *Cell* **161**, 1202–1214 (2015).
1508   118. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for
1509   Dimension Reduction. (2018).
1510   119. Meehan, C., Ebrahimian, J., Moore, W. & Meehan, S. *Uniform Manifold Approximation and
1511   Projection (UMAP). MATLAB Central File Exchange.
1512   https://www.mathworks.com/matlabcentral/fileexchange/71902*. (2021).
1513   120. Reimers, N. & Gurevych, I. Sentence-BERT: Sentence Embeddings using Siamese BERT-Networks.
1514   (2019).
1515   121. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
1516   data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420 (2018).
1517   122. Clauset, A., Shalizi, C. R. & Newman, M. E. J. Power-law distributions in empirical data. *SIAM Rev.
1518   Soc. Ind. Appl. Math.* **51**, 661–703 (2009).
1519   123. Noiret, A. *et al.* Sex-Specific Response to Caloric Restriction After Reproductive Investment in
1520   Microcebus murinus: An Integrative Approach. *Front. Physiol.* **11**, (2020).
1521   124. Perret, M. & Aujard, F. Aging and season affect plasma dehydroepiandrosterone sulfate (DHEA-S)
1522   levels in a primate. *Exp. Gerontol.* **40**, 582–587 (2005).
1523   125. Dal-Pan, A. *et al.* Caloric restriction or resveratrol supplementation and ageing in a non-human
1524   primate: first-year outcome of the RESTRIKAL study in Microcebus murinus. *Age* **33**, (2011).
1525   126. Petter-Rousseaux, A. & Picon, R. Annual variation in the plasma testosterone in Microcebus
1526   murinus. *Folia Primatol.* **36**, (1981).
1527   127. Schilling, A. & Perret, M. Removal of the olfactory bulbs modifies the gonadal responses to
1528   photoperiod in the lesser mouse lemur (Microcebus murinus). *Biol. Reprod.* **49**, 58–65 (1993).
1529   128. Petter-Rousseaux, A. Annual variations in the plasma thyroxine level in Microcebus murinus. *Gen.
1530   Comp. Endocrinol.* **55**, 405–409 (1984).
1531   129. Perret, M. & Predine, J. Effects of long-term grouping on serum cortisol levels in Microcebus
1532   murinus (Prosimii). *Horm. Behav.* **18**, 346–358 (1984).
1533   130. Aujard, F., Bluet-Pajot, M. T., Zizzari, P., Perret, M. & Epelbaum, J. IGF-1: a marker of individual
1534   life-span in a primate. *Ageing Research* **1**, (2010).
1535   131. Aujard, F. *et al.* Artificially accelerated aging by shortened photoperiod alters early gene expression
1536   (Fos) in the suprachiasmatic nucleus and sulfatoxymelatonin excretion in a small primate,
1537   Microcebus murinus. *Neuroscience* vol. 105 403–412 (2001).
1538   132. Giroud, S. *et al.* Gut hormones in relation to body mass and torpor pattern changes during food
1539   restriction and re-feeding in the gray mouse lemur. *J. Comp. Physiol. B* **179**, 99–111 (2009).
1540   133. Dal-Pan, A., Blanc, S. & Aujard, F. Resveratrol suppresses body mass gain in a seasonal non-human
1541   primate model of obesity. *BMC Physiol.* **10**, (2010).

1542

Fig. 3

A Blood hormone level in summer vs. winter

	Hormone	p-value	mean conc. in summer
peak in summer	Estradiol	8.6E-07	376 pg/mg creat
	Testosterone	3.3E-07	60.7 ng/mL
	Thyroxine	3.2E-09	32.5 nmol/L
	DHEA	1.5E-14	8.07 µg/dL
	Cortisol	1.3E-03	523 ng/mg Creat
	IGF1	2.1E-08	433 ng/mL
peak in winter	Melatonin	1.4E-02	9.56 ng/mg Creat
	GIP	7.0E-04	35.1 pg/mL
	PPY	1.2E-05	59.4 pg/mL
	Insulin	1.1E-06	1.01 ng/mL
	PYY	6.9E-15	23.0 pg/mL
	GLP-1	4.6E-17	17.9 pg/mL

log2(fold change)

B

Epithelial Endothelial
Stromal Neural Germ
Lymphoid Myeloid

● Targeted by the 12 seasonal hormones
● Not targeted by the 12 seasonal hormones

UMAP 2
UMAP 1

By transcriptome (excluding hormone-related genes)

Fig. S4

A

B Fitting generality distribution

C Fitting power law distribution

The Tabula Microcebus Consortium

Name	Email	Lab	Affiliations
Snigdha Agarwal	snigdha.agarwal@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Aditi Agrawal	aditi.agrawal@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Ahmad Al-Moujahed	ahmadm@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Alina Alam	alina.ara.alam@gmail.com	Charles Chan	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Megan A. Albertelli	albertel@stanford.edu	Megan Albertelli	Department of Comparative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Paul Allegakoen	paul.allegakoen@ucsf.edu	Franklin Huang	Division of Hematology/Oncology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
Thomas Ambrosi	ambrosi@stanford.edu	Charles Chan	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Jane Antony	janetony@stanford.edu	Michael Clarke	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Steven Artandi	sartandi@stanford.edu	Steven Artandi	Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA.; Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.
Fabienne Aujard	fabienne.aujard@mnhn.fr	Fabienne Aujard	Adaptive Mechanisms and Evolution (MECADEV), UMR 7179, National Center for Scientific Research, National Museum of Natural History, Brunoy, France.
Kyle Awayan	kyle.awayan@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Ankit Baghel	abaghel@stanford.edu	Philip Beachy	Department of Biology, Stanford University, Stanford, CA, USA.
Isaac Bakerman	isaacbakerman@email.arizona.edu	Patricia Nguyen	Department of Medicine, Division of Cardiovascular Medicine, Stanford University, Stanford, CA, USA.
Trygve E. Bakken	trygveb@alleninstitute.org	Allen Institute for Brain Science	Human Cell Types Department, Allen Institute for Brain Science, Seattle, WA, USA.
Jalal Baruni	baruni@stanford.edu	Mark Krasnow; Liqun Luo	Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biology, Stanford University, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Philip Beachy	pbeachy@stanford.edu	Philip Beachy	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Urology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA, USA.
Biter Bilen	biterbilen@gmail.com	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.
Olga Botvinnik	olga.botvinnik@gmail.com	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Scott D. Boyd	sboyd1@stanford.edu	Scott Boyd	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.
Deviana Burhan	deviana.burhan@ucsf.edu	Bruce Wang	Department of Medicine and Liver Center, University of California San Francisco, San Francisco, CA, USA.
Kerriann M. Casey	kmcasey@stanford.edu	Kerriann Casey	Department of Comparative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Charles Chan	chazchan@stanford.edu	Charles Chan	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Charles A. Chang	changca@stanford.edu	Seung Kim	Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.; Stanford Diabetes Research Center, Stanford, CA, USA.
Stephen Chang	scthree@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Department of Medicine, Division of Cardiovascular Medicine, Stanford University, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Ming Chen	ming2017@stanford.edu	Douglas Vollrath	Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.
Michael F. Clarke	mfclarke@stanford.edu	Michael Clarke	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Sheela Crasta	sheelat@stanford.edu	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.

Rebecca Culver	becca.culver@gmail.com	Kerwyn Casey Huang	Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.
Jessica D'Addabbo	jcdaddab@stanford.edu	Patricia Nguyen	Division of Cardiovascular Medicine, Stanford University, Stanford, CA, USA.; Stanford Cardiovascular Institute, Stanford, CA, USA.
Spyros Darmanis	spyros.darmanis@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Roozbeh Dehghannasiri	rdehghan@stanford.edu	Julia Salzman	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biomedical Data Science, Stanford University, Stanford, CA, USA.
Song-Lin Ding	songd@alleninstitute.org	Allen Institute for Brain Science	Human Cell Types Department, Allen Institute for Brain Science, Seattle, WA, USA.
Connor V. Duffy	cvduffy@stanford.edu	Silvana Konermann	Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Jacques Epelbaum	jacques.epelbaum@inserm.fr	Jacques Epelbaum	Adaptive Mechanisms and Evolution (MECADEV), UMR 7179, National Center for Scientific Research, National Museum of Natural History, Brunoy, France.; Unité Mixte de Recherche en Santé 894 INSERM, Centre de Psychiatrie et Neurosciences, Université Paris Descartes, Sorbonne, Paris, France
F. Hernán Espinoza	hespinoz@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Camille Ezran	camillee@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Jean Farup	jean@biomed.au.dk	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biomedicine, Aarhus University, Aarhus, Denmark.
James E. Ferrell, Jr.	james.ferrell@stanford.edu	James Ferrell	Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.
Hannah K. Frank	hkfrank@tulane.edu	Scott Boyd	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA, USA.
Margaret Fuller	mtfuller@stanford.edu	Margaret Fuller	Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.
Astrid Gillich	agillich@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Elias Godoy	eligodoy@stanford.edu	Stanford Veterinary Service Center	Department of Comparative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Dita Gratzinger	ditag@stanford.edu	Dita Gratzinger	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.
Lisbeth A. Guethlein	libbyg@stanford.edu	Peter Parham	Department of Structural Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA.
Yan Hang	yanh@stanford.edu	Seung Kim	Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.; Stanford Diabetes Research Center, Stanford, CA, USA.
Kazuteru Hasegawa	kazuteru@stanford.edu	Steven Artandi	Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA.; Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.
Rebecca D. Hodge	rebeccah@alleninstitute.org	Allen Institute for Brain Science	Human Cell Types Department, Allen Institute for Brain Science, Seattle, WA, USA.
Malachia Hoover	malachia@stanford.edu	Charles Chan	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Franklin W. Huang	franklin.huang@ucsf.edu	Franklin Huang	Division of Hematology/Oncology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
Kerwyn C. Huang	kchuang@stanford.edu	Kerwyn Casey Huang	Department of Bioengineering, Stanford University, Stanford, CA, USA.; Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA.; Chan Zuckerberg Biohub, San Francisco, CA, USA.
Shelly Huynh	shelly.huynh@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.

Taichi Isobe	tisobe.kyushu@gmail.com	Michael Clarke	Department of Oncology and Social Medicine, Kyushu University, Japan.
Carly Israel	carly.israel@yale.edu	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
SoRi Jang	sorijang@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Qiuyu Jing	qjingab@connect.ust.hk	Angela Wu	Division of Life Science, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Robert C. Jones	jonesbob@stanford.edu	Stephen Quake	Department of Bioengineering, Stanford University, Stanford, CA, USA.
Jengmin Kang	jminkang@stanford.edu	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.
Caitlin J. Karanewsky	ckaranew@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Jim Karkanias	jim.karkanias@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Justus Kebschull	justus@kebschull.me	Justus Kebschull	Department of Biomedical Engineering, Johns Hopkins School of Medicine, Baltimore, MD, USA.
Aaron Kershner	kershner@stanford.edu	Philip Beachy	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Institute of Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Lily Kim	lilykim@stanford.edu	Douglas Vollrath	Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.
Seung K. Kim	seungkim@stanford.edu	Seung Kim	Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.; Stanford Diabetes Research Center, Stanford, CA, USA.; JDRF Center of Excellence, Stanford, CA, USA.; Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.
E. Christopher Kirk	eckirk@austin.utexas.edu	E. Christopher Kirk	Department of Anthropology, University of Texas at Austin, Austin, TX, USA.
Winston Koh	Winston_koh@imcb.a-star.edu.sg	Molecular Engineering Lab	Institute of Bioengineering and Bioimaging, Agency of Science Technology and Research, Singapore; Bioinformatics Institute, Agency of Science Technology and Research, Singapore
Silvana Konermann	silvanak@stanford.edu	Silvana Konermann	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
William Kong	wwkong@stanford.edu	Philip Beachy	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Mark A. Krasnow	krasnow@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Christin Kuo	ckuo@stanford.edu	Christin Kuo	Department of Pediatrics, Stanford University School of Medicine, Stanford, CA, USA.
Corinne Lautier	corinne.lautier@umontpellier.fr	Corinne Lautier	MMDN, University of Montpellier, EPHE-PSL, INSERM, Montpellier, France.
Song Eun Lee	songlee@stanford.edu	Tony Wyss-Coray	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.; Wu Tsai Neurosciences Institute, Stanford, CA, USA.
Ed S. Lein	edl@alleninstitute.org	Allen Institute for Brain Science	Human Cell Types Department, Allen Institute for Brain Science, Seattle, WA, USA.
Rebecca Lewis	rjlewis@austin.utexas.edu	Rebecca Lewis	Department of Anthropology, University of Texas at Austin, Austin, TX, USA.
Peng Li	penglium@umich.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Shengda Lin	shengda_lin@zju.edu.cn	Steven Artandi; Shengda Lin	Zhejiang Provincial Key Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang University, Hangzhou, Zhejiang, China.
Shixuan Liu	shixuan@stanford.edu	Mark Krasnow; James Ferrell	Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Yin Liu	yinl@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Gabriel Loeb	gabriel.loeb@ucsf.edu	Gabriel Loeb	Division of Nephrology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
Jonathan Z. Long	jzlong@stanford.edu	Jonathan Long	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.; Stanford ChEM-H, Stanford, CA, USA.
Wan-Jin Lu	wjlu@stanford.edu	Philip Beachy	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.

Katherine Lucot	kllucot@stanford.edu	Tom Montine	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.
Liqun Luo	lluo@stanford.edu	Liqun Luo	Department of Biology, Stanford University, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Aaron McGeever	aaron.mcgeever@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Ross Metzger	rmetzger@stanford.edu	Ross Metzger	Department of Pediatrics, Stanford University School of Medicine, Stanford, CA, USA.; Division of Cardiology, Stanford University School of Medicine, Stanford, CA, USA.
Jingsi Ming	jsming@fem.ecnu.edu.cn	Can Yang; Jingsi Ming	Academy of Statistics and Interdisciplinary Sciences, KLATASDS-MOE, East China Normal University, Shanghai, China.; Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Tom Montine	tmontine@stanford.edu	Tom Montine	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.
Antoine de Morree	demorree@stanford.edu	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biomedicine, Aarhus University, Aarhus, Denmark.
Maurizio Morri	morri@stanford.edu	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Karim Mrouj	karim.mrouj@stanford.edu	Philip Beachy	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Shravani Mukherjee	smukherj@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Ahmad Nabhan	anabhan123@gmail.com	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Saba Nafees	saba.nafees@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Norma Neff	norma.neff@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Patrick Neuhöfer	pneuhoef@stanford.edu	Steve Artandi	Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA.; Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.
Patricia Nguyen	pknguyen@stanford.edu	Patricia Nguyen	Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Stanford Cardiovascular Institute, Stanford, CA, USA.; Division of Cardiovascular Medicine, Stanford University, Stanford, CA, USA.
Jennifer Okamoto	jenokamoto@gmail.com	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Julia Olivieri	jolivier@stanford.edu	Julia Salzman	Institute for Computational and Mathematical Engineering, Stanford University, Stanford, CA, USA.
Youcef Ouadah	yzouadah@caltech.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Program in Cancer Biology, Stanford University School of Medicine, Stanford, CA, USA.
Honor Paine	honor.paine@ucsf.edu	Bruce Wang	Department of Medicine and Liver Center, University of California San Francisco, San Francisco, CA, USA.
Peter Parham	peropa@stanford.edu	Peter Parham	Department of Structural Biology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA.
Jozeph L. Pendleton	zephpen@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Lolita Penland	lpenland@calicolabs.com	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Martine Perret	martine.perret@mnhn.fr	Martine Perret	Adaptive Mechanisms and Evolution (MECADEV), UMR 7179, National Center for Scientific Research, National Museum of Natural History, Brunoy, France.
Angela Oliveira Pisco	angela.pisco@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Zhen Qi	zhenqi@stanford.edu	Michael Clarke	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Stephen R. Quake	steve@czbiohub.org; steve@quake-lab.org	Stephen Quake; CZ Biohub	Department of Bioengineering, Stanford University, Stanford, CA, USA.; Chan Zuckerberg Biohub, San Francisco, CA, USA.; Department of Applied Physics, Stanford University, Stanford, CA, USA.
Ute Radespiel	ute.radespiel@tiho-hannover.de	Ute Radespiel	Institute of Zoology, University of Veterinary Medicine Hannover, Germany.
Thomas A. Rando	rando@stanford.edu	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.

Hajanirina Noëline Ravelonjanahary	hajanravel@gmail.com	Mark Krasnow	Department of Animal Biology, Faculty of Science, University of Antananarivo, Antananarivo 101, BP 566, Madagascar.
Andriamahery Razafindrakoto	amaherix@gmail.com	Mark Krasnow	Department of Animal Biology, Faculty of Science, University of Antananarivo, Antananarivo 101, BP 566, Madagascar.
Julia Salzman	horence@stanford.edu	Julia Salzman	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Department of Biomedical Data Science, Stanford University, Stanford, CA, USA.
Nicholas Schaum	ndschaum@gmail.com	Tony Wyss-Coray	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.
Robert Schopler	bobby.schopler@duke.edu	Duke Lemur Center	Duke Lemur Center, Durham, NC, USA.
Bronwyn Scott	bronwyns@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Liza Shapiro	liza.shapiro@austin.utexas.edu	Liza Shapiro	Department of Anthropology, University of Texas at Austin, Austin, TX, USA.
Hosu Sin	hosu@stanford.edu	Margaret Fuller	Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA.
Rahul Sinha	sinhar@stanford.edu	Irving Weissman	Stanford Institute for Stem Cell Biology and Regenerative Medicine, Stanford, CA, USA.
Rene Sit	rene.sit@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Geoff Stanley	gmstanle@stanford.edu	Stephen Quake	Department of Bioengineering, Stanford University, Stanford, CA, USA.
Lubert Stryer	stryer@stanford.edu	Lubert Stryer	Department of Neurobiology, Stanford University School of Medicine, Stanford, CA, USA.
Varun Ramanan Subramaniam	varunramanan608@gmail.com	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Aditi Swarup	aswarup@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Weilun Tan	weilun.tan@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Alexander Tarashansky	tarashanst@gmail.com; alexander@tarashansky.com	Bo Wang; CZ Biohub	Department of Bioengineering, Stanford University, Stanford, CA, USA.; Chan Zuckerberg Biohub, San Francisco, CA, USA.
Aris Taychameekiatchai	aris.taychameekiatchai@ucsf.edu	Bruce Wang	Department of Medicine and Liver Center, University of California San Francisco, San Francisco, CA, USA.
Jérémy Terrien	jeremy.terrien@mnhn.fr	Jérémy Terrien	Adaptive Mechanisms and Evolution (MECADEV), UMR 7179, National Center for Scientific Research, National Museum of Natural History, Brunoy, France.
Kyle J. Travaglini	ktrav@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Andoni Urtasun	a.urtasun.arricaberri@gmail.com	Thomas Rando	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.
Sivakamasundari V	sivav@stanford.edu	Philip Beachy	Institute for Stem Cell Biology and Regenerative Medicine, Stanford University School of Medicine, Stanford, CA, USA.
Avin Veerakumar	avinv@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Department of Bioengineering, Stanford University, Stanford, CA, USA.; Medical Scientist Training Program, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Venkata N. P. Vemuri	pranathi.vemuri@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Jean-Michel Verdier	jean-michel.verdier@ephe.psl.eu	Jean-Michel Verdier	MMDN, University of Montpellier, EPHE-PSL, INSERM, Montpellier, France.
Iwijn De Vlaminck	vlaminck@cornell.edu; idevlaminck@gmail.com	Iwijn De Vlaminck	Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA.
Douglas Vollrath	vollrath@stanford.edu	Douglas Vollrath	Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.
Bo Wang	wangbo@stanford.edu	Bo Wang	Department of Bioengineering, Stanford University, Stanford, CA, USA.
Bruce Wang	bruce.wang@ucsf.edu	Bruce Wang	Department of Medicine and Liver Center, University of California San Francisco, San Francisco, CA, USA.
Gefei Wang	gwangas@connect.ust.hk	Can Yang	Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Michael F.Z. Wang	fw262@cornell.edu	Iwijn De Vlaminck	Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA.; Department of Computational Biology, Cornell University, Ithaca, NY, USA.
Sheng Wang	swang@cs.washington.edu	Sheng Wang	Paul G. Allen School of Computer Science & Engineering, University of Washington, Seattle, WA, USA.
James Webber	james.webber@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.

Hannah Weinstein	hnw2105@cumc.columbia.edu	Franklin Huang	Division of Hematology/Oncology, Department of Medicine, University of California San Francisco, San Francisco, CA, USA.; Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA, USA.; Bakar Computational Health Sciences Institute, University of California San Francisco, San Francisco, CA, USA.
Irving L. Weissman	irv@stanford.edu	Irving Weissman	Stanford Institute for Stem Cell Biology and Regenerative Medicine, Stanford, CA, USA.
Amanda L. Wiggenhorn	awiggen@stanford.edu	Jonathan Long	Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA.; Department of Chemistry, Stanford University, Stanford, CA, USA.
Cathy V. Williams	cathy.williams@duke.edu	Duke Lemur Center	Duke Lemur Center, Durham, NC, USA.
Patricia Wright	Patricia.Wright@Stonybrook.edu; patchapplewright@gmail.com	Patricia Wright	Department of Anthropology, Stony Brook University, NY, USA.
Albert Y. Wu	awu1@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Angela Wu	angelawu@ust.hk	Angela Wu	Division of Life Science, Hong Kong University of Science and Technology, Hong Kong SAR, China.; Department of Chemical and Biological Engineering, Hong Kong University of Science and Technology, Hong Kong SAR, China.; Center for Aging Science, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Tony Wyss-Coray	twc@stanford.edu	Tony Wyss-Coray	Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA.; Wu Tsai Neurosciences Institute, Stanford, CA, USA.
Bao Xiang	bxli@stanford.edu	Albert Wu	Department of Ophthalmology, Stanford University School of Medicine, Stanford, CA, USA.
Jia Yan	rose.yan@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Can Yang	macyang@ust.hk	Can Yang	Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Jinxurong Yang	jyang@diagnoa.com	Angela Wu	Division of Life Science, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Anne D. Yoder	anne.yoder@duke.edu	Anne Yoder	Department of Biology, University Program in Genetics & Genomics, Duke University, Durham, NC, USA.
Brian Yu	brian.yu@czbiohub.org	CZ Biohub	Chan Zuckerberg Biohub, San Francisco, CA, USA.
Andrea R. Yung	andreay@stanford.edu	Mark Krasnow	Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Yue Zhang	helen18@stanford.edu	Mark Krasnow	Department of Biology, Stanford University, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.
Jia Zhao	jzhaoaz@connect.ust.hk	Can Yang	Department of Mathematics, Hong Kong University of Science and Technology, Hong Kong SAR, China.
Zicheng Zhao	zczhao@stanford.edu; rezczhao@gmail.com	Megan Albertelli; Mark Krasnow	Department of Comparative Medicine, Stanford University School of Medicine, Stanford, CA, USA.; Howard Hughes Medical Institute, USA.