Stratified microbial communities are highly sensitive towards multiple combined global change factors, revealing antagonistic and synergistic effects

Marcel Suleiman, Yves Choffat, Xue Zheng, Owen L Petchey

Department of Evolutionary Biology and Environmental studies, University of Zurich

Corresponding author: Marcel Suleiman, marcel.suleiman@uzh.ch, Tel: +41 44 635 4773

Competing interests
The authors declare no competing financial interests.

Abstract

Microbial communities in many ecosystems are facing a broad range of global change scenarios, resulting in microbial changes and possibly regime shifts with unknown ecological consequences. While the influence of single stressors is already described in numerous studies, the effects of multiple stressors working simultaneously are still poorly understood. In this study, we used 240 highly replicable oxic/anoxic aquatic lab micro-ecosystems to understand the influence of four stressors (fertilizer, glyphosate, metal pollution, antibiotics) in all possible combinations at three different temperatures (20 °C, 24 °C, and 28 °C) to shed light into consequences of multiple stressors on different levels of organization, ranging from species abundance to community and ecosystem parameters. Our data reveal that (i) combination of specific stressors can change the biological consequence and direction compared to single stressors in all levels of organisation (ii), effects of stressor combinations are modified by temperature, and (iii) that the number of stressors applied also lead to significant changes. In sum, our study confirmed the need of investigating multiple stressors working simultaneously across different ecological levels of organisation.

Introduction

Microbial communities, as drivers of many ecosystems, must face increasing amounts and strengths of anthropogenic global change (Christensen et al. 2006; Jackson et al. 2016). Confrontation with fertilizer (Suleiman et al. 2017) and metal pollution (Xu et al. 2018a), pesticides like glyphosate(Solomon and Thompson 2003; Relyea 2009), antibiotics (Xu et al. 2018b) and an ongoing temperature increase (Wu et al. 2011) force populations and whole ecosystems to develop in different ways compared to less or unaffected ones. For decades, many studies analysed and proved that even slight changes in environmental conditions can lead to large taxonomic and functional microbial change, and even to regime shifts, with potential (long term) consequences for biogeochemically cycles and ecological function of the affected habitat (Gruber 2011; Suleiman et al. 2021b).

While most studies so far focused on one global change stressor, recent studies (Christensen et al. 2006; Rillig et al. 2019; Suleiman et al. 2021a) indicated that multiple stressors applied simultaneously do not always show additive effects, which would be the sum of the individual stressors. Rather, they demonstrate that interaction effects can occur. These can be synergistic (combined effect greater than the sum of the effect of individual stressors) or antagonistic (combined effect less than the sum of the effect of individual stressors). Furthermore, it was shown, that not just the character of stressor, but also the number of factors combined plays a crucial role (Rillig et al. 2019). Nevertheless, the amount of studies testing three or more stressors for an experimental system in the lab remains very low (Rillig et al. 2019), which highlights the need of investigating combined effects on experimental ecosystems, and is a critical aspect of the research field of climate change microbiology (Hutchins et al. 2019).

The limited number of studies with three or more stressors may be in part due to the

logistical demands of such experiments and the complexity of interpreting their results. For

example, to assess all interactions, one needs a fully-factorial experimental design, and so

experiments can very quickly become large. Regarding interpretation, it can be difficult to

understand the meaning of interactions among more than three stressors, such that even if

one would conduct a fully-factorial experiment, traditional methods for interpreting

interactions (such as interaction plots) may be insufficient. Some studies have instead focused

on the effect of variation in the number of stressors, and have applied only a subsample of all

possible combinations (e.g. Brennan and Collins 2015; Rillig et al. 2019).

Another gap in understanding is how stressors and combinations of stressors act

across levels of ecological organization, from individuals to ecosystem (Galic et al. 2018). For

example, Galic et al (2018) in a case study of a model of amphipod feeding behavior, showed

that responses to multiple stressors at the individual level were not consistent with those at

higher levels of organization. In their case-study, the nature of this inconsistency would lead

to underestimation of effects at population and ecosystem levels if effects at the individual

level were assumed to hold across levels of organisation.

These gaps in knowledge are problematic. One reason is that interactions can be

source of ecological surprises, i.e. if we assume additivity we can be surprised if there are

interactions (Christensen et al 2006). Also, unless there are some generalities about

interactions among drivers, we will never be able to predict the effect of a new (previously

unobserved) combination of drivers, so interactions will also then be a surprise (Christensen

et al 2006). Hence, we and others (e.g. Simmons et al 2021, Rillig et al 2018) are interested in

discovering if there are any general patterns that will allow us to predict the effects of

combinations of environmental changes. This report is about our search for signals of general

patterns in the effects of combinations of environmental drivers. For example, is the

relationship between a biological variable and the number of drivers dependent on the level

of ecological organization of the biological variable. And do the strength and nature of the

interaction effects vary with level of ecological organization.

Microbial networks of aquatic ecosystems are complex (Christensen et al. 2006; DAVIS

et al. 2010; Faust and Raes 2012), consisting of feedback reactions of numerous abiotic and

biotic interactions (Singh et al. 2009), with and within specific functional microbial groups

(Bush et al. 2017; Richardson et al. 2018). Recently, previous work indicated that these

networks are sensitive to environmental change (Christensen et al. 2006; Shade et al. 2011,

2012; Suleiman et al. 2021b, a), identifying these habitats as appropriate systems for study

and understanding influences of global changes.

In this work, we applied four different global change stressors (fertilizer, glyphosate,

metal pollution, antibiotics, and temperature) in all combinations possible with increasing

temperature (20 °C, 24 °C, and 28 °C), on a recently developed and highly replicable stratified

aquatic microbial lab system, resulting in a total number of 240 analysed experimental micro-

ecosystems. Our experiment includes the analysis of several abiotic (oxygen, total nitrogen,

total organic carbon, pH) and biotic variables (diversity richness, microbial community

composition, genera abundances), since recent studies have shown that various levels of

ecosystems can be affected (Weithoff et al. 2000; Shade et al. 2011, 2012; Suleiman et al.

2021b). We hypothesize that (i) depending on the combinations of stressors the ecosystems

will be driven in different directions and (ii) increasing temperatures can change the influence

of combined stressors (iii) increasing number of stressors goes along with significant effects.

Material and methods

Experimental set up

Incubation of stratified microbial communities was performed in micro-ecosystems consisting of standard glass test tubes (4 mL volume). These test tubes were closed with plastic lids, with a small whole (0.5 mm) that allowed gas exchange between the headspace and the atmosphere. Each micro-ecosystem consisted of sediment and water samples taken in May from a small pond in Zurich, Switzerland (47°23'51.2"N 8°32'33.3"E, the temperature of 19 °C, pH of 7) at a depth of 25 cm. Sediment was homogenized (30 min mixing) and subsequently supplemented with sterile 0.25 % crystalline cellulose, 0.25 % methyl-cellulose, 0.5 % $CaSO_4$, 0.1% $CaCO_3$, and 0.005 % $NH_4H_2PO_4$. Glass tubes were filled with a 3 mm layer of supplemented sediment and covered with 3.5 mL pond water (with 0.005 % $NH_4H_2PO_4$), resulting in 500 µL headspace. Microecosystems were incubated at room temperature for 2 hours to let sediment settle, before the different treatments were applied. Incubation took place for 23 days (about three and a half weeks) at 20 °C, 24 °C and 28 °C, respectively.

Multiple stressor treatments

In this work, there were two levels of four stressors, namely $NH_4H_2PO_4$-fertilizer (F), Glyphosate (G), metal pollution (M) and antibiotics (A). Furthermore, control micro-ecosystems (C) were used without application of any stressors. Additionally, the experiment was carried out at 20 °C, 24 °C and 28 °C under a dark-light cycle of 8:16 h, respectively. The temperature treatment and four stressor treatments were applied in a fully-factorial design. Five replicates per treatment combination resulted in 240 micro-ecosystems, according to the following design:

127   ["Control","F","G","M","A","FG","FM","FA","GM","GA","MA","FGM","FGA","GMA","FMA",

128   "FGMA"] * 5 replicates * 3 Temperatures

129   = 240 micro-ecosystems

For the stressor treatments, the following chemical compounds were added to the micro-

ecosystems 2 hours after the assembly of the microecosystems:

• $NH_4H_2PO_4$ in final concentration of 0.01 % as fertilizer (F), like used in a previous

study (Suleiman et al., 2021)

• Glyphosate in final concentration of 0.0001 % (G) like used in a recent study of (da

Costa et al. 2021)

• Trace element solution SL-10 final concentration 10x higher than recommended for

metal pollution (M)

• Mixture of penicillin and ampicillin (0,05 mg/100 mL each) in final concentration of

0.000005 % for antibiotic treatment (A), since such amounts were detected in

natural aquatic habitats (Xu et al. 2018b)

Stressors were applied to the micro-ecosystems only on day 0; there were no further press

or pulse perturbations.

Sampling and full length 16S rRNA sequencing
144
After incubation for 23 days, the whole micro-ecosystem was centrifuged for 5 min at 10.000

rpm (sediment and water column together). DNA was extracted using ZymoBIOMICS DNA

Miniprep Kit (ZymoResearch), following the manufacturer's instructions. Full-length 16S rRNA

gene amplification was performed using the primer pair 27F forward primer (5'-

AGRGTTYGATYMTGGCTCAG-3') and 1592R reverse primer (5'-RGYTACCTTGTTACGACTT-3'),

like reported previously (Suleiman et al., 2021).

PCR products were visualized on an 1 % (w/v) agarose gel and were pooled for

sequencing in equal concentrations. PCR product pools were purified using AMPure®PB beads

(PacBio). Sequencing was performed at the Functional Genomic Center Zürich, Switzerland,

and performed using SMRT Technology (PacBio) as reported previously (on three SMRT

chips). Sequencing data quality were checked using the PacBio SMRT Link software.

156
157   Bioinformatics
158
Sequencing data were filtered based on primer sequences, length (1300-1600 bp), quality,

error rates and chimeras usind *dada2* (Callahan et al. 2016). The final sequence table was

aligned using SILVA ribosomal RNA database (Quast et al. 2012), using version 138 (non-

redundant dataset 99). A phyloseq object was created using the *phyloseq* package (McMurdie

and Holmes 2013), consisting of amplicon sequence variant (ASV) table, taxonomy table and

sample data. For further analysis, the R packages *phyloseq* (McMurdie and Holmes 2013) and

vegan (Oksanen et al. 2019) were used.

In order to compare the effects of the multiple stressors, we analysed various

response variables, namely community composition based on NMDS1, NMDS2, Shannon

richness, oxygen, TN, TOC, and abundances of specific genera and functional groups.

Microbial community composition was quantified using NMDS (non-metric

multidimensional scaling) based on Bray-Curtis scores with the *metaMDS* function of the

vegan R package (Oksanen et al. 2019), with three dimensions used (k=3, try=100). Shannon

richness was calculated using *phyloseq* package (McMurdie and Holmes 2013). Functional

groups were constructed by filtering for cyanobacterial and sulfur-related species, which

resulted in a reduced dataset consisting of members of the order of *Cyanobacteriales*,

Chromatiales,   *Synechococcales*,   *Chlorobiales*,   *Leptolyngbyales*,   *Desulfobulbales*,

Desulfovibrionales, *Limnotrichales* and *Campylobacterales*. Detailed scripts and a phyloseq

with all sequences are available on github (https://github.com/Marcel2907/multiple-

stressor).

Measurement of abiotic factors
180
181 Oxygen concentration was measured 1.5 cm below the water/air interphase using an

182 oxygen dipping probe (PreSens, Germany) at the last day of incubation. pH was measured in

183 the supernatant of the centrifuged micro-ecosystem, as well as total nitrogen (TN) and total

184 organic carbon using a TOC analyser.

185 Results
186
We first results about the microbial community and genera abundances of the 240 micro-

ecosystems (Fig. 1), then, the impact of stressor combination as well as number of stressors

were calculated for diversity richness (Shannon), community structure (NMDS1), community

structure of a subpopulation, as well oxygen and pH (Fig. 2). Additional abiotic and biotic

variables were investigated (NMDS2, total nitrogen, total carbon) and can be find in detail in

the Supplementary Fig. 3. The summary results of all variables (species, community,

ecosystem) analysed based on the impact of stressor combination can be found in Fig. 3, and

the influence of number of stressors in Fig. 4.

In total, 20'319 unique ASVs were identified after running the dada2 pipeline. After

filtering to retain ASVs that appear at least in a rel. abundance of 0.1 % in at least one sample,

the total number of reads decreased to 5830 taxa. The control micro-ecosystems consisted

of mainly cyanobacteria (e.g. *Phormidaceae*) and phototrophic sulfur bacteria (specially

Chromatiaceae) (Fig. 1 a + b, Supplementary Fig. 1). Depending on the stressor combination

and temperature, the microbial communities and genera abundances showed strong

differences compared to the controls. Micro-ecosystems facing stressors of glyphosate and

metal pollution, as a single stressor or in combination, were still dominated by members of

Phormidiaceae and *Chromatiaceae* like the controls, but in all remaining treatments the

abundance of members of these families was strongly decreased (Supplementary Fig. 1).

205
Also on genus level, we observed shifts in microbial composition depending on stressor

combination and temperature (Fig. 1 b). The cyanobacterial genus *Tychonema* is highly

abundant in metal pollution treatment, and with increasing temperature also in glyphosate +

metal pollution treatments (20 % at 28 °C, 7 % at 20 °C). *Chromatium*, in contrast, appears in

several treatments and reached highest rel. abundance at 20 °C when confronted with G:M

(20 %), but vanished with increasing temperature in all treatments (< 5%). Statistical analysis

of the relative abundance dependent on stressor combinations and temperature were

performed on genus level for *Tychonema*, *Sulfuricurvum* and *Chromatium*, as representative

species for Cyanobacteria and sulfur-depending bacteria (Fig. 3). Statistical analysis confirmed

significant changes of relative species abundance depending on stressors, their combination

and the temperature, but also depended on number of stressors applied (Fig. 4).

ASV diversity of the micro-ecosystems changed depending on the combination of

stressors (Fig. 2a+b), resulting in Shannon index values between 3 and 7, with the majority of

samples between 5 and 6, indicating a high diversity in most of the micro-ecosystems. Several

significant stressor treatments and combinatios were identified, namely temperature

increase (p 1.41e-03), fertilizer (p 9.32e-05), antibiotics (p 7.85e-07), as well as combination

of fertilizer+antibiotics (p 3.82e-05) and fertilizer+antibiotics+glyphosate (p 3.02e-02) (Fig.

2a). All listed significant stressor combinations had a positive effect on diversity, except of

temperature increase to 28 °C and fertilizer:antibiotic treatment (Fig. 3). Interestingly, the

negative effect of fertilizer:antibiotic was reversed to a positive effect again by adding

glyphosate, while the treatment of glyphosate alone had no impact (Fig. 2a, Fig. 3). When

analysing the effect of number of stressors and temperature increase on diversity, just

temperature was identified to have a significant effect, but the number of stressors had no

impact (Fig. 2b, Fig. 4).

The influence of multiple stressors on microbial community composition was analysed

using NMDS, and fertilizer was identified as a key stressor for shifting microbial communities

(NMDS1 p 5.11e-77) (Fig. 2b+c, Supplementary Figure 2). Numerous stressor combinations

showed significant differences for NMDS score 1 and NMDS score 2 compared to the controls

(Fig. 2c for NMDS1, Supplementary Fig. 3 for NMDS2, Fig. 3 for both), ranging from single to

four-factorial combined stressors. Different impact directions (positive or negative score

change) of the estimates can be identified depending on the stressor combinations (Fig. 2c,

Fig. 3). The three-way interaction of fertilizer+antibiotics+temperature on NMDS1 score is a

good example how combination of stressors can change the impact direction (Fig. 5): When

no fertilizer was present, the NMDS1 score was negative, while in presence of fertilizer, the

NMDS1 score was positive. The combination of antibiotics and temperature changed

additionally the magnitude of the NMDS1 score. NMDS1 and NMDS2 scores were affected by

number of stressors (p 5.07e-17 and p 3.96e-04, respectivley), but not by total number of

stressors interacting with temperature (Fig. 2d, Fig. 4). The number of stressors did also affect

NMDS1, but not NMDS2 scores (Fig. 2f, Fig. 4).

Analysis of the subset of cyanobacteria, phototrophic sulfur bacteria and sulfate-

reducing bacteria confirmed the trends observed for NMDS of the whole data set (Fig. 2e for

NMDS1, Supplementary Fig. 3 for NMDS2, Fig. 3 for both), but showed slight differences on

three-factorial and four-factorial stressor combinations.

Oxygen concentration was highly dependent on the stressor combinations used, and

various strong effects were identified (Fig. 2g), both positive and negative in direction. All

stressors combined had a negative effect on oxygen concentration (p 4.04e-02, non-

standardised estimate - 6.1). The number of stressors applied had a strong effect on the

oxygen concentration (p 3.02e-05), as well as temperature (p 4.7e-08) and the interaction of

temperature + number of factors (p 0.04) (Fig. 2h, Fig. 4).

The pH of the micro-ecosystems was influenced by numerous treatment

combinations, but the strongest influences was identified as fertilizer addition, which

decreased the pH from 8.5 to 6.5 (p 4.76e-118). Stressor combinations that include fertilizer

addition had lower magnitude though were of the same sign (Fig. 3).Number of stressors had

a negative effect on pH, though this was mostly due to the strong influence of fertilizer and

its greater probability of occurrence in treatment combinations with more stressors applied

(Fig. 4).

Effects on total nitrogen were driven by addition of fertilizer (p 1.58e-65), and this was

not changed by addition of other stressors (Fig. 3, Supplementary Fig. 3). Furthermore,

number of stressors was highly affecting the total nitrogen concentration (p 4.09e-12), which

again can be explained by fertilizer addition alone. Total carbon concentration was

significantly negative affected by fertilizer (p 0.006), metal pollution (p 0.025) and

temperature (p 0.029) (Fig. 3, Supplementary Fig. 3).

Fig. 2 Effects of stressor combination and number of stressors on temperature applied for microbial community and ecosystem variables. Shannon diversity index in dependence on (a) temperature and stressor combination and (b) temperature and number of stressors. NMDS1 scores in dependence on (c) temperature and stressor combination and (d) temperature and number of stressors. NMDS1_subpopulation scores dependence on (e) temperature and stressor combination and (f) temperature and number of stressors. Oxygen concentration in dependence on (g) temperature and stressor combination and (h) temperature and number of stressors. pH scores in dependence on (i) temperature and stressor combination and (j) temperature and number of stressors. Standard errors are shown (n = 5 replicates).

287
Fig. 3 Summary of statistical analyses of all variables analysed.
We analysed how 12 variables (grouped by genus appearance, microbial richness and community structure and abiotic
factors) are affected by four factors applied in all combinations possible and additionally at three different temperatures.
Coefficient intervals (t-test) are shown for significant affected stressor combination (F-test p-value < 0.05). Color scheme
illustrates the direction of the influence by the stressor (based on estimate positive or negative, estimates of each variable
were standardized by dividing by highest estimate value. Grey: not significant based on f-test (p value > 0.05). tn= total
nitrogen, tc= total carbon.
295
296

297
298 **Fig. 4 Slope of effect of number of stressors over temperature for all level or organisation (species, community,**
299 **ecosystem).** 95 % confidence intervals are shown. tn=total nitrogen, tc=total carbon.

301
302
303 **Fig. 5 Visualization of the significant stressor (interactions) of temperature, fertilizer and antibiotics on NMDS1.** NMDS1 is
304 affected by the presence of fertilizer, the presence of antibiotics and temperature, but also by the interactions of fertilizer-
305 antibiotics and Fertilizer-antibiotics-temperature.

Discussion

Our study revealed strong evidence of effects of the combination and the total number of stressor on taxa abundances, community composition, and ecosystem properties, and therefore confirmed recent findings that multiple stressors are highly important to study (Rillig et al. 2019). The addition of a stressor, and/or temperature increase, could change the biological consequence of already applied stressors, even when the added stressor applied alone had no effect (Fig. 3). We observed three trends here: The addition of a further stressor can increase (e.g. F:A:T28), decrease (e.g. F:T28) or turn the whole direction of stressors which were already applied (e.g. F:A on *Tychonema* abundance, Fig. 6). This demonstrates clearly the need of analysing multiple stressors in all possible combinations. Furthermore, our study also confirmed that the number of stressors applied alone can have an effect on microbial communities and thereby provides some hope that number of stressors can be used as a predictor, though it is clear that individual stressor effects can greatly contributed to the effect of number of stressors (Rillig et al. 2019). The effect of number of stressors can be negatively (e.g. *Tychonema* abundance) or positively (e.g. total nitrogen), which highlights again the need of analysing various ecological response variables of species, microbial communities and the whole ecosystem.

While fertilizer, metal pollution and antibiotics had numerous effects when applied as a single stressor, glyphosate did not show this trend. This finding is probably concentration dependent and should be regarded with caution. Nevertheless, it is even more interesting that when adding glyphosate to more stressors it can have an effect (see e.g. diversity richness). Strong interactions were detected for antibiotics and fertilizer, and interestingly, these stressors revealed in combination different biological consequences compared to

334   applied as a single stressor, and temperature could also change the effects of the fertilizer

335   antibiotic stressor duo again (Fig. 6).

The effect of temperature is in our view highly important to notice, since it is

predicting that observed biological consequences caused by global change can change with

on-going global warming. An increase of temperature can lead to new interactions among

stressor combinations (F:A with increasing temperature led again to significant affected

variables) or can lead to significant effects that were not observed at 20 °C (G:M, G:M:24 and

G:M:28 for NMDS1).

Our study also contributes to close the gap of understanding how stressors and their

combination act across various levels of organisation, since our results revealed that

individuals can be affected differently compared to abiotic factors of the ecosystem or the

microbial community composition. Therefore, our study is in line with the findings of Galic et

al. (2018) who reported already differences of the influence of multiple stressors across levels

of organisation. Interestingly, in our study, some treatment combinations, like fertilizer and

antibiotics working simultaneously, showed significant influence across most of the different

levels of organisation (except total nitrogen and total carbon).

Stressors and how they are affecting ecosystems and communities are concentration-

dependent. In this study, we did used only two levels of stressor concentration and added the

stressor in a single pulse treatment. More information about specific concentration

thresholds and press disturbances are important to study and may clarify some observed

trends, as well as the timing of the stressor applied. In addition, responses of functional

aspects of the aquatic microbial communities should be investigated in future studies, which

could shed light into activated and deactivated pathways and enzymes used in microbial

communities to react to changing environmental conditions.

In sum, our study confirmed the need of analysing the effects of multiple stressors on

microbial communities, and indicate further that a broad range of levels of organisation

(species, microbial community, ecosystem) should be analysed due to its variation of

significant effects. Furthermore, our study showed that combination of specific stressors can

change the biological consequence and direction compared to single stressors in all levels of

organisation and that the effects of stressor combinations can be modified by temperature.

372
373

Literature

Brennan G, Collins S (2015) Growth responses of a green alga to multiple environmental drivers. Nat Clim Chang 5:892–897. https://doi.org/10.1038/nclimate2682

Bush T, Diao M, Allen RJ, et al (2017) Oxic-Anoxic regime shifts mediated by feedbacks between biogeochemical processes and microbial community dynamics. Nat Commun 8:. https://doi.org/10.1038/s41467-017-00912-x

Callahan BJ, McMurdie PJ, Rosen MJ, et al (2016) DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583. https://doi.org/10.1038/nmeth.3869

Christensen MR, Graham MD, Vinebrooke RD, et al (2006) Multiple anthropogenic stressors cause ecological surprises in boreal lakes. Glob Chang Biol 12:2316–2322. https://doi.org/https://doi.org/10.1111/j.1365-2486.2006.01257.x

da Costa N, Fugère V, Hébert M-P, et al (2021) Resistance, resilience, and functional redundancy of freshwater bacterioplankton communities facing a gradient of agricultural stressors in a mesocosm experiment. Mol Ecol 30:4771–4788. https://doi.org/https://doi.org/10.1111/mec.16100

Davis J, Sim L, Chambers J (2010) Multiple stressors and regime shifts in shallow aquatic ecosystems in antipodean landscapes. Freshw Biol 55:5–18. https://doi.org/https://doi.org/10.1111/j.1365-2427.2009.02376.x

Faust K, Raes J (2012) Microbial interactions: from networks to models. Nat Rev Microbiol 10:538–550. https://doi.org/10.1038/nrmicro2832

Galic N, Sullivan LL, Grimm V, Forbes VE (2018) When things don't add up: quantifying impacts of multiple stressors from individual metabolism to ecosystem processing. Ecol Lett 21:568–577. https://doi.org/https://doi.org/10.1111/ele.12923

Gruber N (2011) Warming up, turning sour, losing breath: ocean biogeochemistry under global change. Philos Trans R Soc A Math Phys Eng Sci 369:1980–1996. https://doi.org/10.1098/rsta.2011.0003

Hutchins DA, Jansson JK, Remais J V, et al (2019) Climate change microbiology — problems and perspectives. Nat Rev Microbiol 17:391–396. https://doi.org/10.1038/s41579-019-0178-5

Jackson MC, Loewen CJG, Vinebrooke RD, Chimimba CT (2016) Net effects of multiple stressors in freshwater ecosystems: a meta-analysis. Glob Chang Biol 22:180–189. https://doi.org/https://doi.org/10.1111/gcb.13028

421 McMurdie PJ, Holmes S (2013) Phyloseq: An R Package for Reproducible Interactive Analysis
422   and Graphics of Microbiome Census Data. PLoS One 8:.
423   https://doi.org/10.1371/journal.pone.0061217
424
425 Oksanen J, Blanchet FG, Friendly M, et al (2019) Package "vegan" Title Community Ecology
426   Package. Community Ecol Packag 2:1–297
427
428 Quast C, Pruesse E, Yilmaz P, et al (2012) The SILVA ribosomal RNA gene database project:
429   improved data processing and web-based tools. Nucleic Acids Res 41:D590–D596.
430   https://doi.org/10.1093/nar/gks1219
431
432 Relyea RA (2009) A cocktail of contaminants: How mixtures of pesticides at low
433   concentrations affect aquatic communities. Oecologia 159:363–376.
434   https://doi.org/10.1007/s00442-008-1213-9
435
436 Richardson J, Miller C, Maberly SC, et al (2018) Effects of multiple stressors on cyanobacteria
437   abundance vary with lake type. Glob Chang Biol 24:5044–5055.
438   https://doi.org/https://doi.org/10.1111/gcb.14396
439
440 Rillig MC, Ryo M, Lehmann A, et al (2019) The role of multiple global change factors in
441   driving soil functions and microbial biodiversity. Science (80-) 366:886–890.
442   https://doi.org/10.1126/science.aay2832
443
444 Shade A, Read JS, Welkie DG, et al (2011) Resistance, resilience and recovery: Aquatic
445   bacterial dynamics after water column disturbance. Environ Microbiol 13:2752–2767.
446   https://doi.org/10.1111/j.1462-2920.2011.02546.x
447
448 Shade A, Read JS, Youngblut ND, et al (2012) Lake microbial communities are resilient after a
449   whole-ecosystem disturbancefile:///Users/marcel/Downloads/10.1007_PL00008896-
450   citation.ris. ISME J 6:2153–2167. https://doi.org/10.1038/ismej.2012.56
451
452 Singh BK, Dawson LA, Macdonald CA, Buckland SM (2009) Impact of biotic and abiotic
453   interaction on soil microbial communities and functions: A field study. Appl Soil Ecol
454   41:239–248. https://doi.org/https://doi.org/10.1016/j.apsoil.2008.10.003
455
456 Solomon KR, Thompson DG (2003) Ecological risk assessment for aquatic organisms from
457   over-water uses of glyphosate. J Toxicol Environ Heal - Part B Crit Rev 6:289–324.
458   https://doi.org/10.1080/10937400306468
459
460 Suleiman M, Brandt FB, Brenzinger K, et al (2017) Potential N2O Emissions from the Tanks of
461   Bromeliads Suggest an Additional Source of N2O in the Neotropics. Microb Ecol
462   73:751–754. https://doi.org/10.1007/s00248-016-0903-9
463
Suleiman M, Choffat Y, Daugaard U, Petchey OL (2021a) Large and interacting effects of
temperature and nutrient addition on stratified microbial ecosystems in a small,
replicated, and liquid-dominated Winogradsky column approach. Microbiologyopen
10:e1189. https://doi.org/https://doi.org/10.1002/mbo3.1189

Suleiman M, Pennekamp F, Choffat Y, Petchey OL (2021b) Contrasting resistance and
resilience to light variation of the coupled oxic and anoxic components of an
experimental microbial ecosystem. bioRxiv.
https://doi.org/10.1101/2021.08.13.456201
472

473   Weithoff G, Lorke A, Walz N (2000) Effects of water-column mixing on bacteria,
474   phytoplankton, and rotifers under different levels of herbivory in a shallow eutrophic
475   lake. Oecologia 125:91–100. https://doi.org/10.1007/PL00008896
476

477   Wu Z, Dijkstra P, Koch GW, et al (2011) Responses of terrestrial ecosystems to temperature
478   and precipitation change: a meta-analysis of experimental manipulation. Glob Chang
479   Biol 17:927–942. https://doi.org/https://doi.org/10.1111/j.1365-2486.2010.02302.x
480

481   Xu J, Chen Y, Zheng L, et al (2018a) Assessment of heavy metal pollution in the sediment of
482   the main tributaries of Dongting Lake, China. Water (Switzerland) 10:1–16.
483   https://doi.org/10.3390/w10081060
484

485   Xu Z, Li T, Bi J, Wang C (2018b) Spatiotemporal heterogeneity of antibiotic pollution and
486   ecological risk assessment in Taihu Lake Basin, China. Sci Total Environ 643:12–20.
487   https://doi.org/10.1016/j.scitotenv.2018.06.175
488
489

Acknowledgements

MS was funded by Forschungskredit of the UZH (FK-20-125). OLP was supported by the

University of Zurich Research Priority Programs in Global Change and Biodiversity. OLP and

CZ were supported by Swiss National Science Foundation (Project 310030_188431). We thank

the Predictive Ecology group and especially Uriah Daugaard for valuable input and data

analysis support. We thank the Functional Genomic Center Zurich for sequencing efforts.

496
497
498 **Author contribution**
499
OLP and MS planned the experimental set-up. MS performed all experiments in the lab. MS

performed the up- and downstream bioinformatics. MS and OLP performed the data analysis.

YC provided technical support and performed the TOC analysis. XZ performed the DNA

extractions and provided technical support. OLP and MS drafted the manuscript. All authors

confirmed the final version of the manuscript.

Competing interests

The authors declare no competing financial interests.

Supplementary Information

Supplementary Fig. 1 Relative abundance of Chromatiaceae and Phormidiaceae in the micro-ecosystem depending on
stressor combination and temperature. Mean is shown by the circle, standard errors are shown, n=5)

513
514 **Supplementary Fig. 2 Beta-diversity based on NMDS analysis (Bray-Curtis) in relation to temperature and stressor**
515 **combinations.** Stress is 0.16

516
517

Supplementary Fig. 3 Effects of stressor combination and number of stressors on temperature applied for microbial community and ecosystem variables. NMDS2 scores in dependence on (a) temperature and stressor combination and (b) temperature and number of stressors. NMDS2_subpopulation scores in dependence on (c) temperature and stressor combination and (d) temperature and number of stressors. Total nitrogen concentration in dependence on (e) temperature and stressor combination and (f) temperature and number of stressors. Total carbon concentration in dependence on (g) temperature and stressor combination and (h) temperature and number of stressors. Standard errors are shown (n = 5 replicates).