Spontaneous parthenogenesis in the parasitoid wasp *Cotesia typhae*: low frequency anomaly or evolving process?

Claire Capdevielle Dulac[1, *], Romain Benoist[1], Sarah Paquet[1], Paul-André Calatayud[1], Julius Obonyo[2], Laure Kaiser[1], Florence Mougel[1]

[1] *Université Paris-Saclay, CNRS, IRD, UMR Évolution, Génomes, Comportement et Écologie, 91198, Gif-sur-Yvette, France.*

[2] *icipe, International Center of Insect Physiology and Ecology, P.O. Box 30772-00100, Nairobi, Kenya*

* *Corresponding author: claire.capdevielle-dulac@egce.cnrs-gif.fr*

Abstract

Hymenopterans are haplodiploids and unlike most other Arthropods they do not possess sexual chromosomes. Sex determination typically happens via the ploidy of individuals: haploids become males and diploids become females. Arrhenotoky is believed to be the ancestral reproduction mode in Hymenopterans, with haploid males produced parthenogenetically, and diploid females produced sexually. However, a number of transitions towards thelytoky (diploid females produced parthenogenetically) have appeared in Hymenopterans, and in most cases populations or species are either totally arrhenotokous or totally thelytokous. Here we present the case of *Cotesia typhae* (Fernandez-Triana), a Braconidae that produces parthenogenetic females at a low frequency. The phenotyping of two laboratory strains and one natural population showed that this frequency is variable, and that this rare thelytokous phenomenon also happens in the wild. Moreover, mated females from one of the laboratory strains produce a few parthenogenetic daughters among a majority of sexual daughters. The analysis of daughters of heterozygous virgin females allowed us to show that a mechanism similar to automixis with central fusion is very likely at play in *C. typhae*. This mechanism allows some parts of the genome to remain heterozygous, especially at the chromosomes' centromeres, which can be advantageous depending on the sex determination system involved. Lastly, in most species, the origin of thelytoky is either bacterial or genetic, and an antibiotic treatment as well as PCR experiments did not demonstrate a bacterial cause in *C. typhae*. The unusual case of low parthenogenetic frequency described in this species constitutes another example of the fascinating diversity of sex determination systems in Arthropods.

Introduction

Sexual reproduction is the most widespread reproductive strategy among multicellular organisms and especially in animals. In contrast with its predominance, this reproductive mode appears costly due, for instance, to the necessity to detect and attract a partner, escape sexually transmitted diseases or avoid predation during mating. Because they share parenthood with their mate, sexual individuals transmit two-fold less of their genetic material to their progeny compared to asexual counterparts. The ubiquity of sex despite such disadvantages led to the definition of the so-called "paradox of sex" (Meirmans et al., 2012; Otto, 2009).

Numerous cases of evolution toward asexual reproduction or parthenogenesis have been reported, notably within arthropod taxa (The Tree of Sex Consortium, 2014). Parthenogenesis can produce either males (arrhenotoky) or females (thelytoky) from unfertilized eggs, but only the last case strictly coincides with asexual reproduction. It is also referred to as parthenogenesis *sensu stricto*.

Thelytoky has been observed in almost all basal Hexapoda and non-holometabolous insect taxa
(Vershinina and Kuznetsova, 2016) as well as in many holometabolous insect species (Gokhman and
Kuznetsova, 2018). This wide taxonomic range illustrates the frequent transition from sexual to
asexual taxa that arose independently in various lineages. This scattered distribution hides a global
low percentage of parthenogenesis: thelytokous species represent less than 1% of the Hexapoda
(Gokhman and Kuznetsova, 2018). The proportion of asexual lineages is also highly heterogeneous
among taxa. Liegeois et al., (2021) detected frequencies between 0 and 6.7% among families of
mayflies. Van der Kooi et al. (2017) reported frequencies ranging from 0 to 38% among genera of
haplodiploid arthropods.

Transition from a sexual to asexual reproductive mode requires bypassing genetic and
developmental constraints, a challenge that may be easier to face in some taxa. In most species with
a haplodiploid sex determination system, males develop from unfertilized eggs and are haploid while
females develop from fertilized eggs leading to a diploid state. In such cases, embryonic development
is initiated independently from egg fertilization, a trait probably favouring an evolution toward
thelytoky (Vorburger, 2014). The variable frequency of asexual reproduction even among
haplodiploid lineages indicates that other factors allowing the transition toward this reproductive
mode remain to be identified (van der Kooi et al., 2017).

The multiple and independent acquisitions of asexual reproduction are associated with numerous
mechanisms that maintain or restore diploid state and produce females (Rabeling and Kronauer,
2013; Vorburger, 2014), as illustrated in Figure 1A. The figure focuses on genetic consequences in
terms of heterozygosity, but each described situation may result from different cytological
mechanisms. Apomixis induces clonal reproduction and allows the complete preservation of
heterozygosity. It may arise from mitosis, but also from endoreplication preceding meiosis with
sister chromosome pairing, resulting in recombination between identical chromosomes (Archetti,
2010; Ma and Schwander, 2017). In automixis, meiosis occurs and is followed by different diploid
restoration processes. Two meiosis products may assemble to generate a diploid cell: i) fusion of
non-sister products separated during the first *reductional* division in central fusion or ii) fusion of
sister cells produced during the second *equational* division in terminal fusion. Note that similar
patterns are obtained when one meiotic division is suppressed to ensure the maintenance of a
diploid state: the lack of first division is equivalent to central fusion while the lack of second division
equates to terminal fusion. The restoration of diploidy may also result from gamete duplication
involving either fusion of mitosis products or chromosomal replication without cellular division. In
some lineages, the restoration of diploidy may operate during embryogenesis *via* endomitosis (Little
et al., 2017; Pardo et al., 1995). Other mechanisms not illustrated here involve endomitosis followed
by meiosis with non-sister chromosomes pairing or inverted meiosis with central or terminal fusion
(Archetti, 2022, 2010). The consequences of thelytoky in terms of heterozygosity are variable
depending on the mechanism: from complete homozygosity in one generation under gamete
duplication to completely preserved heterozygosity in apomixis, with intermediate levels of
homozygosity in terminal and central fusion. According to the biology of a given species and the
degree of necessity for maintaining heterozygosity, one or other mechanism may be favoured.

Three main origins of thelytoky have been described: hybridization, bacterial endosymbiosis and
genetic mutation (Tvedte et al., 2019). Hybridization, joining genomes from two distinct species,
leads to improper chromosome pairing and dysfunctional meiosis that may promote asexuality
(Morgan-Richards and Trewick, 2005). Endosymbiotic origin is the most widely studied cause of
parthenogenesis (Ma and Schwander, 2017). To date, only bacteria have been shown to act as
parthenogenesis inducers, but it is likely that other microorganisms could be involved. Most of the

described causative agents belong to the genera *Wolbachia*, *Rickettsia* and *Cardinium*, endosymbionts also known to induce cytoplasmic incompatibility or feminization of male embryos. The particularity of endosymbiont induced parthenogenesis resides in its partial or total reversibility. Thelytokous species treated with antibiotics or heat may revert to sexual reproduction, although often performing less well than true sexual counterparts (Stouthamer et al., 1990). The genetic origin of thelytoky has often been suggested when antibiotic or heat treatment had no effect, but the precise identification of loci responsible for parthenogenesis has only been conducted in a few species (Chapman et al., 2015; Jarosch et al., 2011; Lattorff et al., 2005; Sandrock and Vorburger, 2011).

The relative frequency of thelytoky and sexual reproduction within species also varies according to taxa (Gokhman and Kuznetsova, 2018; Vershinina and Kuznetsova, 2016). Some species are described as obligate thelytokous when this mode of reproduction is the only one observed. Alternatively, thelytoky appears cyclic in some species where asexual generations alternate with sexual ones (Neiman et al., 2014). In other cases, polymorphism in the reproductive mode is observed either between populations (Foray et al., 2013; Leach et al., 2009) or within populations (Liu et al., 2019). Even in such polymorphic situations, thelytokous females usually produce female only progeny, albeit with a very low frequency of males in some cases, allowing rare events of sexual reproduction (Pijls et al., 1996).

Spontaneous occurrence of parthenogenesis has also been described in species reproducing via a sexual mode and qualified as tychoparthenogenesis (Ball, 2001; Pardo et al., 1995). Tychoparthenogenesis is characterized by a low hatching rate and a weak survival probability of the offspring (Little et al., 2017). It is typically considered as a dead-end accidental phenomenon in species adapted to sexual reproduction, although it may also correspond to an intermediate state in the evolution toward asexuality (van der Kooi and Schwander, 2015).

Cotesia typhae (Fernandez-Triana; Hymenoptera, Braconidae) is a gregarious endoparasitoid wasp native to Eastern Africa (Kaiser et al., 2017, 2015). It is specialized to one host, the corn stemborer *Sesamia nonagrioides* (Lefèbvre, Lepidoptera, Noctuidae). *Cotesia typhae* reproduces sexually and fertilized females typically lay 70-100 eggs in the first host encountered, among which about 70% develop into females and 30% into males (Benoist et al., 2020b). At least in laboratory conditions, sister-brother mating (sib-mating) frequently occurs indicating that inbreeding is not detrimental to this species. A genetic survey was conducted on this parasitoid wasp to compare two laboratory strains initiated from wild individuals sampled in two distant Kenyan localities, Kobodo and Makindu (Benoist et al., 2020a). The study led to the construction of a genetic map, based on crosses between the two strains. The phenotyping of the progenies obtained from these controlled crosses revealed an extremely variable sex-ratio, ranging from 100% to as low as 5% of females. Such a phenomenon could result from poorly mated females but also from rare thelytokous events in the progeny of unfertilized females. This last hypothesis was validated in a preliminary experiment allowing virgin females to oviposit. Among the numerous males emerging from the parasitized hosts, a few females were detected.

The aim of this study is to describe the low frequency thelytoky phenomenon in *Cotesia typhae* as a possible case of asexuality emergence. We first confirmed that this low frequency phenomenon is not restricted to artificial breeding but is also observed in natural conditions. We then tested whether thelytoky is induced by environmental conditions, that is lack of fertilisation, or observed in the progeny of mated as well as virgin females. Finally, we addressed the question of the mechanisms allowing the asexual production of females, because these mechanisms are tightly linked to the loss or conservation of heterozygosity and to the evolvability of asexual lineages.

Material and Methods

Biological material

Two separate *Cotesia typhae* parasitoid strains were obtained from adults that emerged from naturally parasitized *Sesamia nonagrioides* caterpillars collected in the field at two localities in Kenya (Kobodo: 0.679S, 34.412E; West Kenya; 3 caterpillars collected in 2013 and Makindu: 2.278S, 37.825E; South-East Kenya; 10 caterpillars collected in 2010-2011). Isofemale lines were initiated in 2016 and inbred rearings have been subsequently kept for more than 80 generations at the Evolution, Génomes, Comportement et Ecologie laboratory (EGCE, Gif-sur-Yvette, France), where cross experiments and phenotyping were performed. The phenotyping of the wild population was performed on individuals that emerged from naturally parasitized *Sesamia nonagrioides* caterpillars collected in the field in 2020 at Kobodo (see above). The *S. nonagrioides* host strain used was initiated from caterpillars collected at Makindu (see above) and Kabaa (1.24S, 37.44E). The rearing protocol of *C. typhae* and *S. nonagrioides* is detailed in Benoist et al. (2020b).

DNA extraction and genotyping methods

For our different experiments, DNA was extracted from *C. typhae* individuals using the NucleoSpin® Tissue from Macherey Nagel, following the manufacturer's instructions. For the experiment analysing the offspring of mated females, a direct PCR method was used instead of a classic DNA extraction because of the high number of individuals to be genotyped. In this case, for each individual, the abdomen was removed (because the presence of gametes could hinder the genotyping) and the rest of the body was placed in 20µL of Dilution Buffer and 0.5µL of DNA Release Additive (Thermoscientific). The tubes were kept at room temperature for 5 minutes then placed at 98°C for 2 minutes. One microliter of this mix or of DNA was then used as template for the following PCR.

Two different methods were used to genotype the chosen SNP markers, either HRM (High Resolution Melt), or allele specific PCR. HRM is based on the analysis of melt curves of DNA fragments after amplification by PCR. The melt curves are different according to the nucleotide composition of the DNA fragments, and therefore allow discrimination between homozygotes and heterozygotes at a given SNP. For each SNP marker, a 10µL mix was made with about 1ng of DNA, 0.2µM of each primer, and 5µL of Precision Melt Supermix (Bio-Rad), completed with water. The PCR protocol was 95°C for 2 minutes, followed by 40 cycles of 95°C for 10 seconds, 60°C for 30 seconds, 72°C for 30 seconds, followed by a complete denaturation of 30 seconds at 95°C before performing the melt curve. The melt curve was performed on a CFX96[TM] Real-Time System (Bio-Rad), and was started by an initial step of 1 minute at 60°C, followed by 10 seconds of every 0.2°C increment between 65°C and 95°C. The raw data resulting from the melt curves were analysed with the uAnalyze v2.1 software (Dwight et al., 2012) in order to infer the individuals' genotypes at each SNP marker.

Two SNP markers (8225nov and 21770nov) were genotyped using allele specific PCR. For this method, two parallel amplifications were performed on individuals, one of the primers of the couple being a common primer, and the other one being a specific primer, either to the Makindu, or Kobodo allele. About 1ng of DNA was mixed with 1X buffer, 3mM $MgCl_2$, 0.4mM dNTP, 0.4µM of each primer, 1U GoTaq® Flexi DNA Polymerase (Promega), and completed with water. The PCR programme was 5 minutes at 95°C, followed by 40 cycles of 95°C for 1 minute, 50°C or 55°C for 1 minute (50°C for 8225nov and 55°C for 21770nov), 72°C for one minute and a final elongation of 5 minutes at 72°C.

180 The PCR products were then run on a 2% agarose gel to check which PCR were positive and therefore
181 infer the genotypes.

182 All the primers used were designed for this study and their sequences are given in Supplementary
183 Table 2.

Phenotyping the strains/populations for the thelytokous character

In this study, the phenotyping consists of counting the number of males and females in the offspring
of virgin females, to quantify the thelytoky phenomenon. To obtain virgin females, individual
cocoons were isolated from cocoon masses and kept in tubes with a moistened cotton wool ball and
a drop of honey at 27°C until the emergence of the adults. The virgin females were then each
allowed to oviposit in one *S. nonagrioides* caterpillar, and the number of males and females in their
offspring was counted after the development of the new *C. typhae* generation.

Flow cytometry for ploidy analysis

194 Flow cytometry analysis was performed on one control female from a mixed cocoon mass (produced
195 by a fertilized female), two control males, and five parthenogenetic females (produced by a virgin
196 female), all coming from the Makindu laboratory strain, to determine their ploidy. The individuals
197 were frozen in liquid nitrogen and processed in the Imagerie-Gif Platform of Institute for Integrative
198 Biology of the Cell (I2BC), CNRS, Gif-sur-Yvette according to the protocol in (Bourge et al., 2018).

Fecundity assessment of parthenogenetic females

Parthenogenetic females were tested for their fecundity and their ability to perform thelytoky. To
test whether mated parthenogenetic females had the same fecundity as mated control females,
cocoon masses resulting from the eggs laid by *C. typhae* Makindu virgin females in *S. nonagrioides*
caterpillars were divided in smaller cocoon packs to spot the few parthenogenetic females more
easily among the males after the emergence of the adults. Adults were left together for one day with
water and honey to allow mating, and eleven parthenogenetic females were then allowed to oviposit
in *S. nonagrioides* caterpillars. After the emergence of the resulting offspring, the number of males
and females was counted for each one of them and the sex-ratio was calculated for comparison with
that obtained from fertilized females from the control Makindu laboratory strain. To test whether
virgin parthenogenetic females were able to produce parthenogenetic daughters, all the cocoons
from 5 virgin females' progenies were isolated and the virgin parthenogenetic females emerging
from these cocoons were allowed to oviposit in *S. nonagrioides* caterpillars. The number of males
and females was then counted in each resulting offspring.

Identifying the thelytoky mechanism in *Cotesia typhae*

To find out which thelytoky mechanism is at play in the *C. typhae* Makindu laboratory strain, virgin
heterozygous females are needed to analyse the recombination patterns of their offspring. Indeed,
according to the mechanism, the female offspring will be more or less heterozygous, as explained in
introduction (Figure 1A). To obtain virgin heterozygous females, six controlled crosses were
performed between the Makindu and Kobodo laboratory strains, 3 in each direction (Figure 1B). Prior

to this, cocoons had been isolated from masses of each strain, in order to obtain virgin males and
females for the crosses. Cocoons were then isolated from the masses resulting from the crosses,
leading to the emergence of virgin F1 heterozygous females. 57 of these females were allowed to
oviposit in *S. nonagrioides* caterpillars, and the offspring of the 57 resulting cocoon masses were
sexed and counted. Six females and four males from the two parental strains (including the
individuals used for the initial crosses), five F1 heterozygous females and the nine parthenogenetic
females obtained through this experiment were kept for DNA extraction and genotyping, in order to
analyse the recombination patterns resulting from parthenogenesis.

To analyse the recombination patterns of the nine parthenogenetic females, we genotyped 63 SNP
(Single Nucleotide Polymorphism) markers, having different alleles between the Makindu and
Kobodo strains, and being distributed along the 10 chromosomes of the genetic map of *Cotesia*
typhae, (Benoist et al., 2020a). Four chromosomes contained more markers than the others to
investigate the recombination patterns along chromosomes. For these four chromosomes, the
markers were chosen to have about 10cM between two successive markers when possible. The
markers were genotyped according to the protocol described above, and their genetic position is
given in Supplementary Table 1.

Identifying the thelytoky phenomenon in mated females' progenies

To see if mated females produce parthenogenetic daughters as well as sexual daughters, 40 crosses
between Makindu virgin females and Kobodo males were performed, according to the protocol
described in the previous section. By genotyping the daughters of these crosses with a SNP marker
differing between the Makindu and Kobodo strains, we can deduce if they were produced sexually (if
heterozygous at the marker) or through parthenogenesis (if homozygous for the Makindu allele at
the marker).

Out of the 40 crosses, five resulted in male only offspring and were removed from our analysis. The
35 remaining crosses lead to a mix of male and female offspring, for a total of 1861 daughters and
1803 sons. The 1861 daughters were genotyped at one SNP marker, 27068nov, according to the
protocol described in the dedicated section. All the females that had a clear Makindu homozygote
genotype and all the females presenting an uncertain genotype were then genotyped at 2 more
markers (8225nov and 21770nov) to confirm their status.

1A)

1B)

Figure 1: A) Expected genotype patterns of parthenogenetic daughters according to the main known
thelytoky mechanisms. The cytological mechanisms are simplified because the figure focuses on the

genetic consequences that can be achieved in different ways (see introduction for further details).
The patterns can vary from complete loss of heterozygosity (under gamete duplication) to complete
maintenance of heterozygosity (under apomixis). B) Crosses performed in this study in order to
obtain virgin heterozygous females. The analysis of recombination patterns of their daughters can
give clues on the thelytoky mechanism involved.

Search for a bacterial cause of thelytoky in *Cotesia typhae*

To find out if the cause of thelytoky in *C. typhae* could be bacterial, we first performed PCR with
primers designed to amplify DNA sequences from several micro-organisms known to manipulate sex
in insects. Ten virgin Makindu females that produced parthenogenetic daughters and two virgin
Makindu females that didn't produce daughters were tested with 8 primer sets, taken from Foray et
al., (2013), except for one primer set, specific to *Wolbachia*, taken from Casiraghi et al., (2005). The
primer sequences, Tm used for PCR, and their original publication are given in Supplementary Table
3. About 1ng of DNA was mixed with 1X buffer, 3mM $MgCl_2$, 0.4mM dNTP, 0.4μM of each primer, 1U
GoTaq® Flexi DNA Polymerase (Promega), and completed with water. The PCR programme was 5
minutes at 95°C, followed by 40 cycles of 95°C for 1 minute, Tm for 1 minute, 72°C for one minute
and a final elongation of 5 minutes at 72°C. The PCR products were then run on a 1% agarose gel to
check for positive amplification.

The amplified fragments obtained with the *Arsenophonus* primer set were sequenced with the
BigDye™ Terminator v1.1 Cycle Sequencing Kit (ThermoFisher Scientific), following the
manufacturer's protocol. After the identification of the bacteria *Pantoea dispersa* through
sequencing, 8 virgin Kobodo females that didn't produce any parthenogenetic daughters were also
tested with this primer set. Since our Kobodo laboratory strain doesn't undergo thelytoky, this test
was performed to check if *Pantoea dispersa* could be the causative agent of thelytoky in *C. typhae*.

To complete this experiment, we performed an antibiotic treatment on the Makindu laboratory
strain to remove any potential sex manipulating bacteria in *C. typhae* females. Rifampicin was added
in the host caterpillars' artificial diet, at a final concentration of 2g/L, for 4 *C. typhae* generations. This
treatment has previously been shown to eliminate *Wolbachia* bacteria in a close species, *Cotesia*
sesamiae (Mochiah et al., 2002). The phenotyping results after this first treatment being ambiguous,
it was continued for 7 more generations, with tetracyclin added for the 4 last generations, also at a
final concentration of 2g/L. At that point, 79 virgin *C. typhae* females were phenotyped, according to
the phenotyping protocol previously described.

Results

Phenotypes of the different strains/populations

We phenotyped two laboratory strains for the thelytokous character, Makindu and Kobodo, and a
wild population, coming from the Kobodo locality. The numbers of available virgin females obtained
for phenotyping were as follows: 99 from 13 different cocoon masses for the Makindu strain, 40 from
7 cocoon masses for the Kobodo strain, and 29 from 6 cocoon masses for the Kobodo wild
population. The results (Table 1) are very contrasted between these three populations, since the
number of parthenogenetically produced females (hereafter referred as parthenogenetic females) is
null in the Kobodo isofemale strain, intermediate in the Kobodo wild population (28% of virgin
females produced daughters), and high in the Makindu isofemale strain (68% of virgin females

produced daughters). The thelytokous phenotype is therefore present in the wild and is not a laboratory artefact, but was apparently lost in the Kobodo laboratory strain (likely due to genetic drift), or present at a frequency too low to be detected. Unfortunately, the wild Makindu population does not exist anymore and could not be tested in this study.

		N	Number of virgin females that produced female offspring (parthenogenetically) Percentage [95% confidence interval]	Number of males produced by all virgin females	Number of females produced by all virgin females Percentage [95% confidence interval]	Mean number [min ; max] of parthenogenetic females in the offspring presenting females
Makindu isofemale strain		99	67 68% [57.4-76.5]	10657	225 2% [1.81-2.36]	3.4 [1;8]
Kobodo isofemale strain		40	0	5405	0	-
Kobodo wild population		29	8 28% [13.4-47.5]	2132	8 0.4% [0.17-0.77]	1 [1;1]
Makindu strain after antibiotic treatment	After 4 generations	55	15 27% [16.5-41.2]	5886	17 0.3% [0.17-0.47]	1.13 [1;2]
	After 11 generations	79	13 16% [9.4-26.9]	8272	16 0.2% [0.11-0.32]	1.23 [1;4]

Table 1: Results of the thelytoky phenotyping of the different strains/populations. The phenotyping is based on the number and frequency of daughters produced parthenogenetically (parthenogenetic females) in the offspring of virgin *C. typhae* females. "N" is the total number of virgin females.

Ploidy of the daughters of virgin females

One daughter from the progeny of a fertilized female and 2 males, all belonging to the Makindu laboratory strain, were processed by flow cytometry as respective controls for diploid and haploid *Cotesia typhae* genomes. Five parthenogenetic daughters of virgin females were then processed, resulting in an estimated genome size identical to the control female and twice that of the control

312   males (Table 2). We can therefore conclude that *C. typhae* parthenogenetic females are diploid and
313   not the result of feminization of haploid eggs.

Sample	Size (pg)	Size (Mpb)	Ploidy
Control female	0.47	458.27	Diploid
Control male 1	0.26	251.84	Haploid
Control male 2	0.25	241.12	Haploid
Parthenogenetic female 1	0.48	467.51	Diploid
Parthenogenetic female 2	0.48	473.24	Diploid
Parthenogenetic female 3	0.49	475.28	Diploid
Parthenogenetic female 4	0.49	475.91	Diploid
Parthenogenetic female 5	0.5	484.26	Diploid

Table 2: Genome size estimated by flow cytometry. Parthenogenetic females have the same genome
size as the control female, corresponding to about twice the males' haploid genome size.

Fecundity of parthenogenetic females

Eleven parthenogenetic females (issued from virgin Makindu mothers) were randomly allowed to
mate with their brothers and were used to parasitize eleven caterpillars. Out of these eleven
females, 4 had male only offspring and 7 had a mixed offspring. The number of offspring per female
and the sex-ratio are indicated in Table 3. No significant difference of the offspring size and sex ratio
was observed between the control and parthenogenetic female datasets (p-value obtained following
Mann-Whitney rank test was 0.559 for offspring number and 0.07 for sex-ratio). The fecundity of
parthenogenetic females is therefore equivalent to that of the control females.

	N	Mean number of offspring per mated female ± Standard Error	Mean female sex-ratio (when mated) ± Standard Error
Control Makindu strain[a]	41	59 ± 4.2	0.78 ± 0.03
Parthenogenetic females	7	54.3 ± 9.2	0.65 ± 0.12

Table 3: Comparison of the fecundity between Makindu parthenogenetic females and control
females of the same laboratory strain. a: data from (Benoist et al., 2017).

All the cocoons resulting from the egg-laying of five virgin females (385 cocoons in total) were
isolated in order to obtain virgin parthenogenetic females. Fourteen females were thus obtained, out
of which ten produced an offspring. Three out of these 10 offspring (30%) contained parthenogenetic
females, for a total of 6 females and 773 males. Virgin parthenogenetic females are therefore able to
produce parthenogenetic females themselves.

Thelytoky mechanism occurring in *Cotesia typhae*

The genotyping of the 63 SNP markers first confirmed that fathers and mothers of the initial crosses between the Makindu and Kobodo laboratory strains were homozygotes for their strain's alleles. The 57 virgin F1 daughters resulting from these crosses were thus heterozygous at the SNP markers, which was confirmed for the 5 F1 daughters that were genotyped. Each of these females successfully parasitized a host larva, and from the 57 resulting offspring, six contained parthenogenetic females (originating from 4 of the initial 6 crosses, 2 in each cross direction), corresponding to a total of 9 F2 parthenogenetic females for 6653 males. These 9 females were genotyped for the 63 SNP. The genotypes and the deduced recombination events are presented in Supplementary Table 1. The recombination patterns of the 4 chromosomes genotyped with a higher density of markers are shown in Figure 2.

For six of the females, a mixture of heterozygous and homozygous markers was observed, with a surplus of heterozygotes (280 heterozygous genotypes for 94 homozygous genotypes). The number and pattern of heterozygous markers for these females indicates a mechanism similar to automixis with central fusion (it could either result from the lack of first meiotic division or from the fusion of two non-sister meiotic products). Indeed, the central parts of the chromosomes maintain a heterozygous state while there is a recombination gradient leading to more homozygous genotypes towards the extremities of the chromosomes (Fig. 2). On average, nine recombination events per genome were detected for these six females with a minimum value of five events and a maximum of 16 events detected. Based on the density of the markers characterized, these results are consistent with the genetic length measured by Benoist et al. (2020a).

Figure 2: Proportion of heterozygous females (out of nine) for each genotyped SNP marker. The results are only shown for the 4 chromosomes for which a higher number of markers were genotyped. The black segments on the x axis are indicative of the genetic position of all the markers of the genetic map (Benoist et al., 2020a) and the red segments with the dotted lines correspond to the positions of the markers genotyped in this study. The occurrence of homozygous and heterozygous states along the chromosomes is congruent with a mechanism similar to automixis. Given the metacentric nature of *C. typhae* chromosomes (C. Bressac, personal communication), the observation of 100% heterozygosity in the central part of the chromosomes suggests that diploidy is

either maintained by the suppression of the first meiotic division or restored through central fusion
and is indicative of the position of each chromosome's centromere.

For the other 3 parthenogenetic females, all the 63 markers were heterozygous, revealing no
detection of recombination event on the 10 chromosomes. We estimated the probability of such an
observation under the hypothesis that a unique mechanism such as central fusion occurs. For each
chromosome, we calculated a mean number of recombinations based on the nine parthenogenetic
females. Assuming that the number of recombinations on a chromosome follows a Poisson
distribution, we can estimate the probability of zero recombination for each chromosome based on
the mean number estimate. It varies according to the genetic length of the chromosome and to the
density of markers genotyped: it was estimated between 0.29 for chromosome 2 and 0.8 for
chromosome 9. Multiplying the probability over the ten chromosomes, we calculated a probability of
0.0025 to observe an entirely heterozygous parthenogenetic daughter. Using this individual
probability, we estimated that the probability to detect three out of nine parthenogenetic daughters
showing no recombination events on the 10 chromosomes was 1.76×10^{-6}. This hypothesis is very
unlikely, therefore we suspect another mechanism could also be at play in causing thelytoky in
Cotesia typhae. It is interesting to note that we observed both patterns (partial homozygosity and
complete heterozygosity) in the offspring resulting from initial crosses of both directions. Moreover,
one of the F1 heterozygous females displayed both patterns in her progeny. The raw data and the R
script used to estimate the given probability are available at https://zenodo.org/record/6420801.

Presence of parthenogenetic females among the daughters of mated females

Among the 40 crosses between Makindu females and Kobodo males, five gave male-only offspring
and were excluded. The 35 remaining crosses that presented offspring comprising both males and
females were kept in our analysis, leading to a total of 1861 females and 1803 males. All 1861
females were genotyped for one SNP marker. Females resulting from fecundation should be
heterozygous while parthenogenetic females should be homozygous for the Makindu allele. In total,
we found 14 homozygous females, which were confirmed by the genotyping of 2 other markers.
These 14 females correspond to 0.77% of the parthenogenetic offspring (males resulting from
arrhenotoky representing 99.33%) and originate from 10 different mothers (29% of the 35 mothers)
(Table 4). Even though the percentage of parthenogenetic females found is much smaller than in the
offspring of virgin females, this finding shows that the female progeny of mated females can come
from a mixture of parthenogenesis and sexual reproduction.

	N	Number of females that produced parthenogenetic daughters Percentage [95% confidence interval]	Number of parthenogenetic offspring (males plus parthenogenetic females)	Number of parthenogenetic females among the parthenogenetic offspring Percentage [95% confidence interval]	Mean number [min ; max] of parthenogenetic females in the offspring presenting females
Makindu virgin	99	67 68% [57.4-76.5]	10882	225 2% [1.81-2.36]	3.4 [1;8]

females					
Makindu mated females	35	10 29% [15.2-46.5]	1817	14 0.77% [0.44-1.32]	1.4 [1;3]

Table 4: Comparison of the frequency of the thelytokous character between virgin and mated Makindu females. In each case, the number of parthenogenetic daughters is presented as a percentage of the total number of parthenogenetic offspring, mainly composed of males obtained from arrhenotoky. "N" is the number of offspring analysed.

Origin of thelytoky in *Cotesia typhae*

In order to find out if thelytoky in *Cotesia typhae* has a bacterial origin, we extracted DNA from Makindu virgin mothers that produced daughters and used primers to try to amplify the DNA of six different micro-organisms known for sex manipulation in insects: *Wolbachia, Ricketssia, Cardinium, Arsenophonus, Spiroplasma* and *Microsporidia* (Foray et al., 2013). Only one primer set led to a solid amplification, the one designed to amplify *Arsenophonus* 23S. After sequencing the amplified fragment, the bacterium was identified not as *Arsenophonus* but as *Pantoea dispersa*, for which no mention in relation to thelytoky was found in the literature. We then tried to amplify this same bacterium from the DNA of Kobodo virgin mothers, who don't produce any daughters: *Pantoea dispersa* was present in all the samples tested. This makes it unlikely for this bacterium to be responsible for thelytoky in *C. typhae*.

After rearing parasitized caterpillars for four generations on a rifampicin diet, we phenotyped the Makindu strain again. 55 virgin females, coming from 5 different cocoon masses, were allowed to parasitize *Sesamia nonagrioides* caterpillars. Fifteen of these virgin females produced daughters, leading to a total of 17 daughters for 5886 sons (Table 1). Another phenotyping was performed on 79 females from 10 different cocoon masses, after 11 generations of a rifampicin diet (with tetracyclin added for the last 4 generations). Thirteen of these virgin females produced daughters, leading to a total of 16 daughters for 8272 sons (Table 1). The percentage of thelytokous females is thus smaller than the one observed before antibiotic treatment but not null.

Discussion

The phenotypic survey presented here confirms the biological reality of low frequency asexual production of females in the haplo-diploid Hymenoptera *Cotesia typhae*. The process has been observed in a significant number of progenies from both an inbred laboratory strain and a natural population. It has been shown to occur in the progeny of virgin as well as fertilized females, despite concerning only a small fraction of the individuals from a cocoon mass.

This configuration of low frequency thelytoky is poorly illustrated in the literature, except in the well-studied species *Apis mellifera* for which the phenomenon has been described for a long time (Mackensen, 1943; Tucker, 1958). In the honey bee, both workers and virgin queens are able to produce a small proportion of females among unfertilized progeny (Gloag et al., 2019). However, in a common acceptation, the expression thelytoky is rather defined as a "parthenogenetic mode where females produce only females from unfertilized eggs" (Vershinina and Kuznetsova, 2016). Among most illustrated examples of parthenogenesis *sensu stricto*, even when facultative, asexual

production of females involves the whole progeny. The case described here is somewhat closer to
what is called tychoparthenogenesis, based on the frequency of birth of parthenogenetic female eggs
(Whiting, 1945). Tychoparthenogenesis is defined as "kind of occasional thelytoky characterized by
the spontaneous hatching of a small proportion of eggs laid by virgin females" (Pardo et al., 1995). It
has been mainly described in diplodiploid species where embryonic development is induced by
sperm fertilization. In such species, developmental constraints and inbreeding depression prevent
successful hatching of unfertilized eggs in most of the cases (Little et al., 2017). In haplodiploid
species, unfertilized eggs hatch with a high frequency because they naturally produce males in
species reproducing sexually. It is thus difficult to classify *C. typhae* as a tychoparthenogenetic
species. Moreover, because the daughters produced parthenogenetically turned out to be viable and
fertile in *C. typhae*, low frequency thelytoky may be either neutral or beneficial but not
disadvantageous as in most cases of tychoparthenogenesis described.

The question remains as to whether the phenomenon is accidental or an ongoing evolutionary
process due to its adaptive benefit (van der Kooi and Schwander, 2015). In the eusocial species *Apis*
mellifera, low frequency thelytoky is clearly beneficial to workers when confronted to a queen-less
colony (Gloag et al., 2019). This advantage led to an increased frequency of workers' reproduction in
an honey bee subspecies (*Apis m. capensis*) and to the development of social parasitism where laying
workers get adopted by a colony and compete the local queen (Neumann, 2001). Studying the
occurrence of parthenogenesis among Ephemeroptera, Liegeois et al. (2021) suggested that asexual
reproduction was selectively advantageous in many species from this insect order despite its
associated low hatching success. The benefit derives from the short adult life and the low dispersal
ability that reduce the probability of encountering a reproductive partner. The fitness of sexually
reproducing individuals may consequently be reduced under certain circumstances. As in mayflies, *C.*
typhae has an adult life limited to a few days (between two and three days in laboratory conditions,
Kaiser et al., 2017). However, it is gregarious and mating between sisters and brothers emerging
from the same cocoon mass is observed in rearing conditions. Female access to male fertilization
should thus be facilitated unless sib-mating is avoided in natural conditions as demonstrated for its
relative species *Cotesia glomerata* (Gu and Dorn, 2003) or *Venturia canescens* (Collet et al., 2020).
Even in the presence of sexual partners, limited access to fertilization may also derive from
ineffective copulation in cases of highly female biased sex ratios for example (Boivin, 2013). Whether
it results from restricted access to males or to sperm itself, sperm limitation may favour expansion of
asexual reproduction. Further experiments are needed to estimate mating and fertilization success of
Cotesia typhae in natural conditions.

Beyond reproductive strategy itself, parthenogenesis has been shown to be associated with
ecological characteristics that may favour or prevent its evolution. Two opposite ecological trends
have been described co-occurring with asexual reproduction expansion: the "general purpose
genotype" (GPG) where asexual lineages are observed on broader ecological niches than their sexual
counterparts and the "frozen niche variation" (FNV) where parthenogenetic species or populations
have far more restricted niches than sexual ones (Tvedte et al., 2019). Exploring a wide dataset of
haplodiploid arthropods reproducing exclusively parthenogenetically (obligate parthenogenesis), van
der Kooi et al. (2017) concluded that GPG was the most common situation. They showed that most
parthenogenetic species have broader ecological and geographical range than close relative sexual
species but also that transition toward parthenogenesis was more likely for species exhibiting a wide
distribution. Studying the relative advantage of sexual and asexual reproduction, Song et al., (2012)
developed a model based on resource availability, in spatial and temporal heterogeneous situations.
They showed that sexual reproduction prevails in most of the cases but that asexual reproduction
may be favoured when resource diversity is low and resource remains abundant over generations.

Such a model corroborates numerous cases of ecological specialization of asexual lineages that have
been described, such as *Venturia canescens*. In this polymorphic species, two kinds of populations
live in sympatry: parthenogenetic populations found in stable anthropic habitats (bakeries and
granaries) and sexual ones associated with natural and more instable resources (Schneider et al.,
2002). Interestingly, before being characterized as a new species, *Cotesia typhae* was first identified
as a specialized clade (only one host insect, *Sesamia nonagrioides*, mainly found on one host plant,
Typha domingiensis) of the parasitoid species *Cotesia sesamiae*. According to (Branca et al., 2019),
some populations of *C. sesamiae* are less specialized than others. Studying the existence of
thelytokous reproduction in those populations would be informative about the possible link between
emerging parthenogenesis and specialization.

Regarding the mechanism involved in thelytokous reproduction, we faced an unexpected result as
data strongly suggest that two different processes may co-occur: automixis with central fusion (or a
similar cytological mechanism) and apomixis. More surprisingly, the two supposed mechanisms were
observed to co-occur in the progeny of a single female (K4M1) and independently of the cross
direction to obtain F1 virgin mothers (Kobodo female x Makindu male or Makindu female x Kobodo
male). Unfortunately, this result is supported by small sample size due to the scarcity of the
phenomenon. We may wonder whether a unique mechanism, distinct from those already described,
could explain such a result. Ma and Schwander (2017) describe for example an unusual process
where meiosis is inverted (sister chromatids separate before homologs) followed by terminal fusion.
However, the resulting progeny of such a process is 100% heterozygous, a result that does not differ
from apomixis. Another mechanism presented in the same review implies an endoreplication
preceding meiosis. Assuming such a process occurs in *C. typhae*, and hypothesizing that
recombination, and consequently segregation during the first division, may arise either between
identical or between homologous chromosomes, some intermediate situations are expected. Once
again, it does not reconcile the clear-cut figure we observe with individuals entirely heterozygous
suggesting zero recombination between homologs and individuals for which recombination is
observed for almost all homologs. If different mechanisms truly co-occur to produce females in an
asexual way, it could reflect an accidental phenomenon arising from the relaxed control of sexual
reproduction as observed in the honey bee (Aamidor et al., 2018). To better understand the
mechanism underlying thelytoky in *C. typhae*, a cytological approach of meiosis and parthenogenesis
would be necessary.

Despite the lack of a unified mechanism to explain the genotypic profile observed in the F2 progenies
obtained, we can confirm that recombination occurred in at least 6 out of 9 cases, and that these
recombination events were as frequent as those observed in sexual reproduction (Benoist et al.,
2020a). By contrast, severe reductions of recombination rates were observed associated with
parthenogenetic reproduction in the literature. For example, recombination in thelytokous workers is
reduced by up to 10-fold in comparison to their sexually reproducing mothers in the Cape bee, *Apis
mellifera capensis*, the social parasite of honeybee which reproduces parthenogenetically *via*
automixis with central fusion (Baudry et al., 2004). In the little fire ant *Wasmannia auropunctata*,
sexual populations coexist with asexual populations in which reproductive queens are produced by
automictic parthenogenesis with central fusion. In asexual populations, recombination rate is
reduced by a factor of 45 compared to the sexual populations (Rey et al., 2011). The reduction of
recombination rate is assumed to mitigate the potential deleterious impact of thelytoky: under
automixis with central fusion, heterozygosity is preserved unless recombination occurs (Figure 1). In
species affected by inbreeding depression, a homozygosity increase would be detrimental and could
be advantageously limited by a low recombination rate. As the molecular mechanisms involved in
thelytoky and recombination are probably distinct, the situation observed in the Cape bee and little

fire ant may result from a long-term evolutionary process. If the phenomenon described in *C. typhae*
is recent, it may explain the unchanged recombination rate.

Otherwise, the inbreeding impact could be meaningless in *C. typhae*. Hymenoptera are haplodiploid
and could thus be less sensitive to inbreeding because most of the deleterious alleles are purged at
the haploid state in males (Hedrick and Parker, 1997; Henter, 2003). However, their sex
determination system may be highly compelling regarding homozygosity and ability to reproduce *via*
thelytoky (Vorburger, 2014). The most common, and likely ancestral, sex determination system is
governed by the genotype at one (sl-CSD: single locus Complementary Sex Determination) or few loci
(ml-CSD: multi locus CSD) (Heimpel and de Boer, 2008). Under such a determinism, individuals that
are heterozygous at least at one of these CSD loci develop as diploid females while hemizygous or
homozygous individuals at all CSD loci develop as haploid or diploid males respectively. In most
hymenopteran species, diploid males have a low survival rate and/or are often sterile. Enhanced
homozygosity due to thelytoky may be very costly when it results in diploid male production (de Boer
et al., 2015, 2012; van Wilgenburg et al., 2006; Zhou et al., 2007). Other sex determination systems
that are less sensitive to homozygosity have been described in Hymenoptera, such as Paternal
Genome Elimination (Heimpel and de Boer, 2008) or genome imprinting, described in *Nasonia*
vitripennis (van de Zande and Verhulst, 2014). The mechanism of sex determination is unknown in *C.*
typhae. However, it has been reared for more than 80 generations in laboratory conditions starting
from an isofemale line and seems poorly sensitive to homozygosity increase.

The bacterial origin of thelytoky in *C. typhae* could not be either confirmed or completely discarded
in the present study as an intermediate state (in terms of frequency of parthenogenesis) was
observed following antibiotic treatment. The knowledge of the genetic mechanism could give some
clues about the origin of parthenogenesis as endosymbionts have been mainly shown to favour
gamete duplication. However, detailing specific interactions reveals a more complex picture.
Cardinium is able to feminize diploid males (Giorgini et al., 2009) but also to induce automixis with
central fusion (Zchori-Fein and Perlman, 2004). *Wolbachia* is mainly known to induce gamete
duplication (Leach et al., 2009; Ma and Schwander, 2017) but it has also been described to promote
apomixis (Weeks and Breeuwer, 2001). *Rickettsia* has also been shown to trigger functional apomixis
(Adachi-Hagimori et al., 2008). Furthermore, the list of endosymbionts is probably partial and in most
of the documented examples of parthenogenesis endosymbiotically determined, the cytological
mechanism remains unknown. Evidence that microorganisms can promote all processes of
parthenogenesis will probably arise from future research. The examples of demonstrated genetic
determinism of thelytoky are rare and only concern automixis with central fusion. This is the case for
the Cape honey bee (Verma and Ruttner, 1983) and for the wasp *Venturia canescens* (Beukeboom
and Pijnacker, 2000). However, Tsutsui et al., (2014) described an apomixis mechanism in the
parasitoid wasp *Meteorus pulchricornis* for which they proposed a genetic origin of thelytoky. Even
more than for endosymbiont origin, genetic determinism of parthenogenesis requires thorough
investigations to determine whether it is restricted to a few cytological mechanisms. Anyway, the
clearly evidenced mechanism of autoximis with central fusion in *C. typhae* does not allow to settle
between genetic and endosymbiont origin as this mechanism is common to both situations.

Conclusion

In this study, we described an unusual example of low frequency thelytokous reproduction within a
sexually reproducing species. We demonstrated that this asexual reproduction is likely the result of
different mechanisms and occurs even in the progeny of fertilized females, an undescribed

phenomenon to our knowledge. As most studies on asexual reproduction focus on obligate or cyclical situations, we may wonder whether such low frequency and probably accidental thelytoky is common but until now mostly undetected among sexually reproductive species. It is well known that asexuality has emerged many times in numerous lineages. If the occurrence of accidental parthenogenesis turns out to be usual, it could represent the first step of evolutionary trajectories favoured either by reproductive or ecological advantages of asexual lineages.

Acknowledgments

The present work has benefited from the I2BC Cytometry platform (Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France) with the help of Mickaël Bourge. We thank Rémi Jeannette and Sylvie Nortier for insect rearing at Gif, and Daniel Couch for proofreading.

This work was supported by the French National Research Agency (project Cotebio ANR-17-CE32-0015), and by the authors' operating grants from IRD, CNRS and *icipe*. R. Benoist was funded by the « Ecole doctorale 227 MNHN-UPMC Sciences de la Nature et de l'Homme: évolution et écologie ».

All experimentations were realized under the juridical frame of a Material Transfer Agreement signed between IRD, icipe and CNRS (CNRS 072057/IRD 302227/00) and the authorization to import Cotesia in France delivered by the DRIAAF of Ile de France (IDF 2017-OI-26-032).

We thank Christoph Haag, Jens Bast and Michael Lattorff for their constructive comments that helped us improve this paper.

Data availability

Raw data for Figure 2, and Tables 1, 3 and 4 are available at https://zenodo.org/record/6420801

References

Aamidor, S.E., Yagound, B., Ronai, I., Oldroyd, B.P., 2018. Sex mosaics in the honeybee: how haplodiploidy makes possible the evolution of novel forms of reproduction in social Hymenoptera. Biol. Lett. 14, 20180670. https://doi.org/10.1098/rsbl.2018.0670

Adachi-Hagimori, T., Miura, K., Stouthamer, R., 2008. A new cytogenetic mechanism for bacterial endosymbiont-induced parthenogenesis in Hymenoptera. Proc. R. Soc. B. 275, 2667–2673. https://doi.org/10.1098/rspb.2008.0792

Archetti, M., 2022. Evidence from automixis with inverted meiosis for the maintenance of sex by loss of complementation. J of Evolutionary Biology 35, 40–50. https://doi.org/10.1111/jeb.13975

Archetti, M., 2010. Complementation, Genetic Conflict, and the Evolution of Sex and Recombination. Journal of Heredity 101, S21–S33. https://doi.org/10.1093/jhered/esq009

Ball, S.L., 2001. Tychoparthenogenesis and mixed mating in natural populations of the mayfly Stenonema femoratum. Heredity 87, 373–380.

Baudry, E., Kryger, P., Allsopp, M., Koeniger, N., Vautrin, D., Mougel, F., Cornuet, J.M., Solignac, M.,
2004. Whole-Genome Scan in Thelytokous-Laying Workers of the Cape Honeybee (Apis mellifera
capensis): Central Fusion, Reduced Recombination Rates and Centromere Mapping Using Half-Tetrad
Analysis. Genetics 252, 243–252.

Benoist, R., Capdevielle-Dulac, C., Chantre, C., Jeannette, R., Calatayud, P., Drezen, J., Dupas, S., Le
Rouzic, A., Le Ru, B., Moreau, L., Van Dijk, E., Kaiser, L., Mougel, F., 2020a. Quantitative trait loci
involved in the reproductive success of a parasitoid wasp. Mol Ecol 29, 3476–3493.
https://doi.org/10.1111/mec.15567

Benoist, R., Chantre, C., Capdevielle-Dulac, C., Bodet, M., Mougel, F., Calatayud, P.A., Dupas, S.,
Huguet, E., Jeannette, R., Obonyo, J., Odorico, C., Silvain, J.F., Le Ru, B., Kaiser, L., 2017. Relationship
between oviposition, virulence gene expression and parasitism success in Cotesia typhae nov. sp.
parasitoid strains. Genetica 145, 469–479. https://doi.org/10.1007/s10709-017-9987-5

Benoist, R., Paquet, S., Decourcelle, F., Guez, J., Jeannette, R., Calatayud, P.-A., Le Ru, B., Mougel, F.,
Kaiser, L., 2020b. Role of egg-laying behavior, virulence and local adaptation in a parasitoid's chances
of reproducing in a new host. Journal of Insect Physiology 120, 103987.
https://doi.org/10.1016/j.jinsphys.2019.103987

Beukeboom, L.W., Pijnacker, L.P., 2000. Automictic parthenogenesis in the parasitoid Venturia
canescens (Hymenoptera: Ichneumonidae) revisited 43, 6.

Boivin, G., 2013. Sperm as a limiting factor in mating success in Hymenoptera parasitoids. Entomol
Exp Appl 146, 149–155. https://doi.org/10.1111/j.1570-7458.2012.01291.x

Bourge, M., Brown, S.C., Siljak-Yakovlev, S., 2018. Flow cytometry as tool in plant sciences, with
emphasis on genome size and ploidy level assessment. Genetics & Applications 2, 1–12.

Branca, A., Le Ru, B., Calatayud, P.-A., Obonyo, J., Musyoka, B., Capdevielle-Dulac, C., Kaiser-Arnauld,
L., Silvain, J.-F., Gauthier, J., Paillusson, C., Gayral, P., Herniou, E.A., Dupas, S., 2019. Relative
Influence of Host, Wolbachia, Geography and Climate on the Genetic Structure of the Sub-saharan
Parasitic Wasp Cotesia sesamiae. Front. Ecol. Evol. 7, 309. https://doi.org/10.3389/fevo.2019.00309

Casiraghi, M., Bordenstein, S.R., Baldo, L., Lo, N., Beninati, T., Wernegreen, J.J., Werren, J.H., Bandi,
C., 2005. Phylogeny of Wolbachia pipientis based on gltA, groEL and ftsZ gene sequences: clustering
of arthropod and nematode symbionts in the F supergroup, and evidence for further diversity in the
Wolbachia tree. Microbiology 151, 4015–4022. https://doi.org/10.1099/mic.0.28313-0

Chapman, N.C., Beekman, M., Allsopp, M.H., Rinderer, T.E., Lim, J., Oxley, P.R., Oldroyd, B.P., 2015.
Inheritance of thelytoky in the honey bee Apis mellifera capensis. Heredity 114, 584–592.
https://doi.org/10.1038/hdy.2014.127

Collet, M., Amat, I., Sauzet, S., Auguste, A., Fauvergue, X., Mouton, L., Desouhant, E., 2020. Insects
and incest: Sib-mating tolerance in natural populations of a parasitoid wasp. Mol Ecol 29, 596–609.
https://doi.org/10.1111/mec.15340

de Boer, J.G., Groenen, M.A., Pannebakker, B.A., Beukeboom, L.W., Kraus, R.H., 2015. Population-
level consequences of complementary sex determination in a solitary parasitoid. BMC Evol Biol 15,
98. https://doi.org/10.1186/s12862-015-0340-2

de Boer, J.G., Kuijper, B., Heimpel, G.E., Beukeboom, L.W., 2012. Sex determination meltdown upon
biological control introduction of the parasitoid *Cotesia rubecula?* Evol Appl 5, 444–454.
https://doi.org/10.1111/j.1752-4571.2012.00270.x

Dwight, Z.L., Palais, R., Wittwer, C.T., 2012. uAnalyze: Web-Based High-Resolution DNA Melting
Analysis with Comparison to Thermodynamic Predictions. IEEE/ACM Trans. Comput. Biol. and Bioinf.
9, 1805–1811. https://doi.org/10.1109/TCBB.2012.112

Foray, V., Helene, H., Martinez, S., Gibert, P., Desouhant, E., 2013. Occurrence of arrhenotoky and
thelytoky in a parasitic wasp Venturia canescens (Hymenoptera: Ichneumonidae): Effect of
endosymbionts or existence of two distinct reproductive modes? Eur. J. Entomol. 110, 103–107.
https://doi.org/10.14411/eje.2013.014

Giorgini, M., Monti, M.M., Caprio, E., Stouthamer, R., Hunter, M.S., 2009. Feminization and the
collapse of haplodiploidy in an asexual parasitoid wasp harboring the bacterial symbiont Cardinium.
Heredity 102, 365–371. https://doi.org/10.1038/hdy.2008.135

Gloag, R., Remnant, E.J., Oldroyd, B.P., 2019. The frequency of thelytokous parthenogenesis in
European-derived Apis mellifera virgin queens. Apidologie 50, 295–303.
https://doi.org/10.1007/s13592-019-00649-0

Gokhman, V.E., Kuznetsova, V.G., 2018. Parthenogenesis in Hexapoda: holometabolous insects. J
Zool Syst Evol Res 56, 23–34. https://doi.org/10.1111/jzs.12183

Gu, H., Dorn, S., 2003. Mating system and sex allocation in the gregarious parasitoid Cotesia
glomerata. Animal Behaviour 66, 259–264. https://doi.org/10.1006/anbe.2003.2185

Hedrick, P.W., Parker, J.D., 1997. Evolutionary Genetics and Genetic Variation of Haplodiploids and X-
Linked Genes. Annu. Rev. Ecol. Syst. 28, 55–83. https://doi.org/10.1146/annurev.ecolsys.28.1.55

Heimpel, G.E., de Boer, J.G., 2008. Sex determination in the hymenoptera. Annual review of
entomology 53, 209–30. https://doi.org/10.1146/annurev.ento.53.103106.093441

Henter, H.J., 2003. INBREEDING DEPRESSION AND HAPLODIPLOIDY: EXPERIMENTAL MEASURES IN A
PARASITOID AND COMPARISONS ACROSS DIPLOID AND HAPLODIPLOID INSECT TAXA. Evolution 57,
1793–1803. https://doi.org/10.1111/j.0014-3820.2003.tb00587.x

Jarosch, A., Stolle, E., Crewe, R.M., Moritz, R.F.A., 2011. Alternative splicing of a single transcription
factor drives selfish reproductive behavior in honeybee workers (Apis mellifera). Proceedings of the
National Academy of Sciences 108, 15282–15287. https://doi.org/10.1073/pnas.1109343108

Kaiser, L., Fernandez-Triana, J., Capdevielle-Dulac, C., Chantre, C., Bodet, M., Kaoula, F., Benoist, R.,
Calatayud, P.-A., Dupas, S., Herniou, E.A., Jeannette, R., Obonyo, J., Silvain, J.-F., Le Ru, B., 2017.
Systematics and biology of Cotesia typhae sp. n. (Hymenoptera, Braconidae, Microgastrinae), a
potential biological control agent against the noctuid Mediterranean corn borer, Sesamia
nonagrioides. ZK 682, 105–136. https://doi.org/10.3897/zookeys.682.13016

Kaiser, L., Le Ru, B.P., Kaoula, F., Paillusson, C., Capdevielle-Dulac, C., Obonyo, J.O., Herniou, E.A.,
Jancek, S., Branca, A., Calatayud, P., Silvain, J., Dupas, S., 2015. Ongoing ecological speciation in
Cotesia sesamiae , a biological control agent of cereal stem borers. Evol Appl 8, 807–820.
https://doi.org/10.1111/eva.12260

Lattorff, H.M.G., Moritz, R.F.A., Fuchs, S., 2005. A single locus determines thelytokous
parthenogenesis of laying honeybee workers (Apis mellifera capensis). Heredity 94, 533–537.
https://doi.org/10.1038/sj.hdy.6800654

Leach, I.M., Pannebakker, B.A., Schneider, M.V., Driessen, G., van de Zande, L., Beukeboom, L.W.,
2009. Thelytoky in Hymenoptera with Venturia canescens and Leptopilina clavipes as Case Studies,
in: Schön, I., Martens, K., Dijk, P. (Eds.), Lost Sex. Springer Netherlands, Dordrecht, pp. 347–375.
https://doi.org/10.1007/978-90-481-2770-2_17

Liegeois, M., Sartori, M., Schwander, T., 2021. Extremely Widespread Parthenogenesis and a Trade-
Off Between Alternative Forms of Reproduction in Mayflies (Ephemeroptera). Journal of Heredity
112, 45–57. https://doi.org/10.1093/jhered/esaa027

Little, C.J., Chapuis, M.-P., Blondin, L., Chapuis, E., Jourdan-Pineau, H., 2017. Exploring the
relationship between tychoparthenogenesis and inbreeding depression in the Desert Locust,
Schistocerca gregaria. Ecol Evol 7, 6003–6011. https://doi.org/10.1002/ece3.3103

Liu, Q., Zhou, J., Zhang, C., Ning, S., Duan, L., Dong, H., 2019. Co-occurrence of thelytokous and
bisexual Trichogramma dendrolimi Matsumura (Hymenoptera: Trichogrammatidae) in a natural
population. Sci Rep 9, 17480. https://doi.org/10.1038/s41598-019-53992-8

Ma, W.-J., Schwander, T., 2017. Patterns and mechanisms in instances of endosymbiont-induced
parthenogenesis. J. Evol. Biol. 30, 868–888. https://doi.org/10.1111/jeb.13069

Mackensen, O., 1943. The Occurrence of Parthenogenetic Females in Some Strains of Honeybees.
Journal of Economic Entomology 36, 465–467. https://doi.org/10.1093/jee/36.3.465

Meirmans, S., Meirmans, P.G., Kirkendall, L.R., 2012. The Costs Of Sex: Facing Real-world
Complexities. The Quarterly Review of Biology 87, 19–40. https://doi.org/10.1086/663945

Mochiah, M.B., Ngi-Song, A.J., Overholt, W.A., Stouthamer, R., 2002. Wolbachia infection in Cotesia
sesamiae (Hymenoptera: Braconidae) causes cytoplasmic incompatibility: implications for biological
control. Biological Control 25, 74–80. https://doi.org/10.1016/S1049-9644(02)00045-2

Morgan-Richards, M., Trewick, S.A., 2005. Hybrid origin of a parthenogenetic genus? Molecular
Ecology 14, 2133–2142. https://doi.org/10.1111/j.1365-294X.2005.02575.x

Neiman, M., Sharbel, T.F., Schwander, T., 2014. Genetic causes of transitions from sexual
reproduction to asexuality in plants and animals. J. Evol. Biol. 27, 1346–1359.
https://doi.org/10.1111/jeb.12357

Neumann, P., 2001. Social parasitism by honeybee workers (Apis mellifera capensis Escholtz): host
finding and resistance of hybrid host colonies. Behavioral Ecology 12, 419–428.
https://doi.org/10.1093/beheco/12.4.419

Otto, S.P., 2009. The Evolutionary Enigma of Sex. The American Naturalist 174, S1–S14.
https://doi.org/10.1086/599084

Pardo, M.C., López-León, M.D., Cabrero, J., Camacho, J.P.M., 1995. Cytological and developmental
analysis of tychoparthenogenesis in Locusta migratoria. Heredity 75, 485–494.
https://doi.org/10.1038/hdy.1995.165

Pijls, J.W.A.M., van Steenbergen, H.J., van Alphen, J.J.M., 1996. Asexuality cured: the relations and
differences between sexual and asexual Apoanagyrus diversicornis. Heredity 76, 506–513.
https://doi.org/10.1038/hdy.1996.73

Rabeling, C., Kronauer, D.J.C., 2013. Thelytokous Parthenogenesis in Eusocial Hymenoptera. Annu.
Rev. Entomol. 58, 273–292. https://doi.org/10.1146/annurev-ento-120811-153710

Rey, O., Loiseau, A., Facon, B., Foucaud, J., Orivel, J., Cornuet, J.-M., Robert, S., Dobigny, G., Delabie,
J.H.C., Mariano, C.D.S.F., Estoup, A., 2011. Meiotic Recombination Dramatically Decreased in
Thelytokous Queens of the Little Fire Ant and Their Sexually Produced Workers. Molecular Biology
and Evolution 28, 2591–2601. https://doi.org/10.1093/molbev/msr082

Sandrock, C., Vorburger, C., 2011. Single-Locus Recessive Inheritance of Asexual Reproduction in a
Parasitoid Wasp. Current Biology 21, 433–437. https://doi.org/10.1016/j.cub.2011.01.070

Schneider, M.V., Beukeboom, L.W., Driessen, G., Lapchin, L., Bernstein, C., Van Alphen, J.J.M., 2002.
Geographical distribution and genetic relatedness of sympatrical thelytokous and arrhenotokous
populations of the parasitoid Venturia canescens (Hymenoptera): Thelytoky and arrhenotoky in
Venturia canescens. Journal of Evolutionary Biology 15, 191–200. https://doi.org/10.1046/j.1420-
9101.2002.00394.x

Song, Y., Scheu, S., Drossel, B., 2012. The ecological advantage of sexual reproduction in multicellular
long-lived organisms: The ecological advantage of sexual reproduction. Journal of Evolutionary
Biology 25, 556–565. https://doi.org/10.1111/j.1420-9101.2012.02454.x

Stouthamer, R., Pinto, J.D., Platner, G.R., Luck, R.F., 1990. Taxonomic Status of Thelytokous Forms of
Trichogramma (Hymenoptera: Trichogrammatidae). Annals of the Entomological Society of America
83, 475–481. https://doi.org/10.1093/aesa/83.3.475

The Tree of Sex Consortium, 2014. Tree of Sex: A database of sexual systems. Sci Data 1, 140015.
https://doi.org/10.1038/sdata.2014.15

Tsutsui, Y., Maeto, K., Hamaguchi, K., Isaki, Y., Takami, Y., Naito, T., Miura, K., 2014. Apomictic
parthenogenesis in a parasitoid wasp *Meteorus pulchricornis* , uncommon in the haplodiploid order
Hymenoptera. Bull. Entomol. Res. 104, 307–313. https://doi.org/10.1017/S0007485314000017

Tucker, K.W., 1958. AUTOMICTIC PARTHENOGENESIS IN THE HONEY BEE. Genetics 43, 299–316.
https://doi.org/10.1093/genetics/43.3.299

Tvedte, E.S., Logsdon, J.M., Forbes, A.A., 2019. Sex loss in insects: causes of asexuality and
consequences for genomes. Current Opinion in Insect Science 31, 77–83.
https://doi.org/10.1016/j.cois.2018.11.007

van de Zande, L., Verhulst, E.C., 2014. Genomic Imprinting and Maternal Effect Genes in Haplodiploid
Sex Determination. Sex Dev 8, 74–82. https://doi.org/10.1159/000357146

van der Kooi, C.J., Matthey-Doret, C., Schwander, T., 2017. Evolution and comparative ecology of
parthenogenesis in haplodiploid arthropods. Evolution Letters 1, 304–316.
https://doi.org/10.1002/evl3.30

van Wilgenburg, E., Driessen, G., Beukeboom, L., 2006. Single locus complementary sex
determination in Hymenoptera: an "unintelligent" design? Front Zool 3, 1.
https://doi.org/10.1186/1742-9994-3-1

van der Kooi, C.J., Schwander, T., 2015. Parthenogenesis: Birth of a New Lineage or Reproductive
Accident? Current Biology 25, R659–R661. https://doi.org/10.1016/j.cub.2015.06.055

Verma, S., Ruttner, F., 1983. Cytological analysis of the thelytokous parthenogenesis in the Cape
honeybee (Apis mellifera capensis Escholtz). Apidologie 14, 41–57.
https://doi.org/10.1051/apido:19830104

Vershinina, A.O., Kuznetsova, V.G., 2016. Parthenogenesis in Hexapoda: Entognatha and non-
holometabolous insects. J Zoolog Syst Evol Res 54, 257–268. https://doi.org/10.1111/jzs.12141

Vorburger, C., 2014. Thelytoky and Sex Determination in the Hymenoptera: Mutual Constraints. Sex
Dev 8, 50–58. https://doi.org/10.1159/000356508

Weeks, A.R., Breeuwer, J.A.J., 2001. *Wolbachia* –induced parthenogenesis in a genus of
phytophagous mites. Proc. R. Soc. Lond. B 268, 2245–2251. https://doi.org/10.1098/rspb.2001.1797

Whiting, P.W., 1945. The Evolution of Male Haploidy. The Quarterly Review of Biology 20, 231–260.
https://doi.org/10.1086/394884

Zchori-Fein, E., Perlman, S.J., 2004. Distribution of the bacterial symbiont Cardinium in arthropods:
Molecular Ecology 13, 2009–2016. https://doi.org/10.1111/j.1365-294X.2004.02203.x

Zhou, Y., Gu, H., Dorn, S., 2007. Effects of inbreeding on fitness components of Cotesia glomerata, a
parasitoid wasp with single-locus complementary sex determination (sl-CSD). Biological Control 40,
273–279. https://doi.org/10.1016/j.biocontrol.2006.11.002

782