1
2
3
4 **Adenine Base Editing *in vivo* with a Single Adeno-Associated Virus Vector**
5
6

8   Han Zhang[a], Nathan Bamidele[a], Pengpeng Liu[b], Ogooluwa Ojelabi[a, 1], Xin D. Gao[a,2], Tomás
9   Rodriguez[a], Haoyang Cheng[a], Jun Xie[d,e,f,g], Guangping Gao[d,e,f,g], Scot A. Wolfe[b,g], Wen
10 Xue[a,b,c,g], and Erik J. Sontheimer[a,c,g,h]

11
12
13
[a]RNA Therapeutics Institute, University of Massachusetts Medical School, Worcester,
Massachusetts, 01605, USA

[b]Department of Molecular, Cell and Cancer Biology, University of Massachusetts Medical
School, Worcester, Massachusetts, 01605, USA

[c]Program in Molecular Medicine, University of Massachusetts Medical School, Worcester,
Massachusetts, 01605, USA

[d]Horae Gene Therapy Center, University of Massachusetts Medical School, Worcester, MA,
01605, USA

[e]Viral Vector Core, University of Massachusetts Medical, School, Worcester, MA, 01605, USA
[f]Department of Microbiology and Physiological Systems, University of Massachusetts Medical
School, Worcester, MA, 01605, USA

[g]Li Weibo Institute for Rare Diseases Research, University of Massachusetts Medical School,
Worcester, Massachusetts, 01605, USA

[1]Present address: LogicBio Therapeutics, Lexington, Massachusetts, 02421, USA

[2]Present address: Merkin Institute of Transformative Technologies in Healthcare, Broad Institute
of Harvard and MIT, Cambridge, Massachusetts, 02142, USA

[h]Correspondence: erik.sontheimer@umassmed.edu

31
32
33

Abstract

Base editors (BEs) have opened new avenues for the treatment of genetic diseases. However, advances in delivery approaches are needed to enable disease targeting of a broad range of tissues and cell types. Adeno-associated virus (AAV) vectors remain one of the most promising delivery vehicles for gene therapies. Currently, most BE/guide combinations and their promoters exceed the packaging limit (~5 kb) of AAVs. Dual-AAV delivery strategies often require high viral doses that impose safety concerns. In this study, we engineered an adenine base editor using a compact Cas9 from *Neisseria meningitidis* (Nme2Cas9). Compared to the well-characterized *Streptococcus pyogenes* Cas9-containing ABEs, Nme2-ABE possesses a distinct PAM (N_4CC) and editing window, exhibits fewer off-target effects, and can efficiently install therapeutically relevant mutations in both human and mouse genomes. Importantly, we show that *in vivo* delivery of Nme2-ABE and its guide RNA by a single-AAV vector can efficiently edit mouse genomic loci and revert the disease mutation and phenotype in an adult mouse model of tyrosinemia. We anticipate that Nme2-ABE, by virtue of its compact size and broad targeting range, will enable a range of therapeutic applications with improved safety and efficacy due in part to packaging in a single-vector system.

Key words: AAV, base editing, genome editing, CRISPR, sgRNA, deaminase

Introduction

Point mutations represent the largest class of known human pathogenic genetic variants [1,2]. Base editors (BEs), which comprise a single-guide RNA (sgRNA) loaded onto a Cas9 (nuclease-inactivated or nickase form) fused to a deaminase enzyme, enable precise installation of A•T to G•C (in the case of adenine base editors (ABEs) [3]) or C•G to T•A (in the case of cytidine base editors (CBEs) [4]) substitutions. In contrast to traditional nuclease-dependent genome editing approaches, base editors do not generate double-stranded DNA breaks (DSBs), do not require a DNA donor template, and are more efficient in editing non-dividing cells, making them attractive agents for *in vivo* therapeutic genome editing.

While robust editing has been achieved in many cultured mammalian cell systems, safe and effective *in vivo* delivery of base editors remains a major challenge. To date, both non-viral and viral delivery methods have shown great promise for delivering base editors for *in vivo* therapeutic purposes in rodents and primates [5,6]. For example, *in vivo* delivery via AAVs has achieved efficient editing in a wide range of tissue and cell types including liver [5–10], heart [11], muscle [12,13], retina [14,15], inner ear [16], and central nervous system (CNS) [17,18]. However, the large coding size (5.2 kb) of the best-characterized *Streptococcus pyogenes* Cas9 (SpyCas9)-containing BEs exceed the packaging limit of AAV (5 kb) [19,20]. Currently, *in vivo* delivery of base editors by AAV has been approached by splitting the SpyCas9 base editor between two AAVs and relying on the use of intein trans-splicing for the assembly of the full-length effector [13,16–18,21,22]. Although effective, this approach requires transduction of the target cell by both AAVs and successful *in trans* splicing of the two intein halves. The requirement to deliver two AAV vectors also increases the viral dosage needed for a treatment, which raises safety concerns and adds burdens to AAV manufacturing [23–25].

Compact Cas9 orthologs are ideal candidates for engineering base editors suitable for single-AAV delivery [26–28]. For example, single-AAV delivery of a domain-inlaid *Staphylococcus aureus* Cas9 (SauCas9) ABE has been reported in cultured HEK293 cells [29]. Previously, we developed a *Neisseria meningitidis* Cas9 (Nme2Cas9) as an *in vivo* genome editing platform [27,30]. Nme2Cas9 is a compact, naturally accurate genome editor with a distinct N_4CC PAM specificity. Recently, two other groups have successfully implemented Nme2-CBEs in cultured mammalian cells and in rabbits [31], as well as Nme2-ABEs in rice [32]. Here, we develop ABEs using Nme2Cas9 (Nme2-ABEs) and define their editing efficiencies, editing windows, and off-target activities in comparison to those of the widely applied Spy-ABEs in cultured mammalian cells. Next, we show that Nme2-ABE can edit multiple therapeutically significant loci, including one of the most common mutations occurring in Rett syndrome patients that cannot be targeted by other compact ABEs (e.g. ABEs derived from SauCas9 and SauCas9-KKH [33–36]) due to PAM restrictions. Lastly, by optimizing the promoter and the nuclear localization signals, we show that Nme2-ABE and its guide can be packaged into a single-AAV vector genome for *in vivo* delivery. One systematic administration of the single-AAV vector encoding both Nme2-ABE and sgRNA readily corrects the disease-causing mutation and phenotype in an adult mouse model of hereditary tyrosinemia type 1 (HT1).

Results

Development of Nme2-ABE and comparison of editing windows and off-target effects to those of Spy-ABE

First, to evaluate base editing efficiency in a streamlined manner, we developed an ABE reporter cell line in which a G-to-A mutation in the mCherry coding sequence generates a nonsense mutation. Adenine base editing can reverse the mutation and recover red fluorescence, and the editing efficiency can be readily measured by fluorescence-activated cell sorting (FACS) (**Figure 1a, Supplementary Figure 1**). Initially, we constructed Nme2-ABE7.10 by linking a TadA-TadA7.10 dimer from the Spy-ABE7.10 to the N-terminus of the Nme2Cas9 HNH nickase [3]. However, by plasmid transient transfection, Nme2-ABE7.10 showed poor activity in the ABE reporter cell line. Because the evolved TadA8e is highly active and compatible with a wide range of Cas9s [35], we next engineered Nme2-ABE8e by linking TadA8e monomer to the N-terminus of the Nme2Cas9 HNH nickase (**Figure 1b**). We found that Nme2-ABE8e supports robust editing activity in the ABE reporter cell line (**Figure 1c**). Next, to define the editing window and editing efficiency of Nme2-ABE8e, and to compare them to those parameters of Spy-ABE7.10 and Spy-ABE8e, we transfected HEK293T cells with plasmids expressing each ABE along with sgRNAs targeting 12 human genomic loci for Nme2-ABE8e [including eight dual-target sites (target sites followed by NGGNCC PAMs for both SpyCas9 and Nme2Cas9) [30] and four Nme2Cas9-specific target sites]. We found that this first-generation Nme2-ABE8e has a broad but shallow editing window that, due to modest activity, was difficult to define clearly (**Figure 1d**).

We next sought to understand the potential extent of off-target editing by Nme2-ABE8e. It has been shown that the major source of DNA off-target base editing is Cas9-dependent [37,38], caused by Cas9 binding and unwinding at near-cognate sequences. Because Nme2Cas9 is highly accurate during nuclease-driven editing in cells and *in vivo* [27,30,39], we hypothesized that Nme2-ABE8e would exhibit similar accuracy advantages relative to Spy-ABE8e. As an initial test, we systematically investigated the tolerance for nucleotide mismatches between the guide and the target sequence for the two effectors. We designed a panel of guides targeting the ABE reporter with single- and di-nucleotide mismatches with the target sequence for both Nme2-ABE8e and Spy-ABE8e and measured their activities by plasmid transfection and FACS (**Figure 1e**). Considering the differences in on-target efficiencies between the two effectors, we further normalized the activities of the mismatched guides to those of the perfectly complementary guides for each effector. We found that Nme2-ABE8e exhibited significantly lower off-target editing propensity than Spy-ABE8e: while single-nucleotide mismatches in the seed region (guide nucleotide positions 17-24 for Nme2Cas9, and 10-20 for SpCas9) and the majority of dinucleotide mismatches significantly compromised the editing efficiency of Nme2-ABE8e, these near-cognate sequences were mostly efficiently edited by Spy-ABE8e (**Figure 1e**).

Inhibition of base editing by anti-CRISPR proteins that limit DNA binding activity

The development of Nme2Cas9 base editing platforms raises the possibility that regulation strategies developed for nuclease-based editing could be similarly implemented in the case of base editing. One such strategy is the use of anti-CRISPR (Acr) proteins that limit Cas9 DNA-binding activity [40–43], which have been deployed to reduce both off-target [44] and off-tissue editing [45]. AcrIIC3 and AcrIIC4 have been reported to reduce Nme2Cas9 DNA binding activity [41,43,46] but have no effect on SpyCas9 [41,43]. To test whether such Acr proteins can

function as off-switches for Nme2-ABE8e base editing, we co-transfected the ABE reporter cell
line described above with plasmids expressing Nme2-ABE8e, sgRNA, and Acr proteins. Spy-
ABE7.10 and AcrIIA4 (an anti-CRISPR that prevents SpyCas9 DNA binding [40], nuclease
editing [40,44], and base editing [47]) was used as a positive control. Conversely, AcrE2, which is
a Type I-E Acr [43] that has no effect on SpyCas9 or Nme2Cas9 activity, was used as a negative
control. As expected, Spy-ABE7.10 base editing was reduced to background levels by AcrIIA4,
but AcrE2, AcrIIC3, and AcrIIC4 had no effect (**Supplementary Figure 3a**). By contrast,
Nme2-ABE8e editing was strongly inhibited by AcrIIC3 and AcrIIC4, but AcrE2 and AcrIIA4
had no effect. These results confirm that these anti-CRISPRs can be effective off-switches for
Nme2-ABE8e editing.
156
Tissue-specific miRNAs, in combination with miRNA response elements [MREs] in the 3'UTR
of an Acr construct, have been used to restrict editing to cell types that express such miRNAs,
both in cultured cells [41,48] and *in vivo* [45]. To determine whether such a strategy could be
used to control Nme2-ABE8e, we inserted miR-122 MREs into the 3'UTRs of our AcrIIC3 and
AcrIIC4 constructs. Mir-122 is a hepatocyte-specific miRNA that is expressed in Huh7 cells but
not HEK293 cells, and has been used to validate this strategy for nuclease-based editing. Again,
AcrIIC3 and AcrIIC4 inhibited Nme2-ABE8e activity at an endogenous site in HEK293T cells,
and inhibition largely persisted even with the MREs (**Supplementary Figure 3b**), as expected
since miR-122 is not present to silence Acr expression. In Huh7 cells, which are transfected with
lower efficiencies, AcrIIC3 and AcrIIC4 again inhibited Nme2-ABE8e, but this inhibition of
editing activity was largely relieved by the insertion of the MRE122 sites. These results indicate
that miRNA-repressible anti-CRISPRs can be used to enforce the cell-type specificity of base
editing, as it can for nuclease editing.
170
Installation of therapeutically relevant edits with Nme2-ABE8e
We next tested the potential of Nme2-ABE8e to correct pathogenic mutations or to introduce
disease-suppressing mutations. One of the most common mutations that cause Rett syndrome is
a C•G to T•A base transition that produces a nonsense mutation in the human *MeCP2* gene
(c.502 C>T; p.R168X) [49–51]. Because the target adenine is within a cytidine-rich region, this
mutation is not readily accessible for ABEs based on other compact Cas9s such as SauCas9 and
SauCas9-KKH [33–36]. To test whether Nme2-ABE8e can correct this pathogenic mutation, we
electroporated Nme2-ABE8e mRNA with a synthetic sgRNA into a Rett syndrome patient-
derived fibroblast cell line that possesses this mutation. By amplicon deep sequencing, we found
that Nme2-ABE8e successfully edited the target adenine (A10) (**Figure 2a, 2b**). A bystander edit
at an upstream adenine (A16) causes a missense mutation (c.496 T>C; p.S166P), although this
occurs with only one-fourth the frequency of intended edit at A10. Because S166 has been shown
to be subject to phosphorylation in mice and is conserved from *X. laevis* to humans [52], further
investigation will be needed to determine whether bystander editing at A16 impairs functional
rescue of edited cells. However, as with other base editors [3,29,53–58], future protein
engineering efforts adjusting the editing window promise to provide greater control over Nme2-
ABE editing outcomes.
188
Next, we sought to generate a disease-suppressing mutation that has been shown to reverse
phenotypes of a validated Duchenne muscular dystrophy (DMD) mouse model (ΔEx51) [59].
The ΔEx51 mouse model was generated by deletion of the exon 51 in the *Dmd* gene, resulting in
a downstream premature stop codon in exon 52, causing the production of a nonfunctional

truncated dystrophin protein. Previously, it has been shown that the *Dmd* reading frame can be
restored by skipping exon 50 by adenine base editing (**Figure 2c**) [13]. However, *in vivo* base
editing using ABE_{max}-SpCas9-NG delivered by dual-AAV vectors was limited to local muscle
injection due to the high viral dosage required to achieve therapeutic benefit. We identified a
guide design for Nme2-ABE8e to target the adenine (A7) within the splice donor site downstream
of exon 50 (**Figure 2c**). By plasmid transfection in the mouse N2a cell line and amplicon deep
sequencing, we found that Nme2-ABE8e can generate $17.67 \pm 4.57\%$ editing at A7 (**Figure 2d**).
The efficient editing at multiple bystander adenines is not a concern in this case as those adenines
are within the skipped exon 50 or the intron.
202
Optimization of an Nme2-ABE8e construct for single AAV delivery
Previously, we showed that Nme2Cas9 with one or two sgRNAs can be packaged into a single
AAV vector and support efficient editing *in vivo* [27,30]. We reasoned, based upon the compact
sizes of Nme2Cas9 and TadA8e, that Nme2-ABE8e with a sgRNA could be packaged into a
single AAV for *in vivo* delivery. To achieve this, we first replaced NmeCas9 with Nme2-ABE8e in
the minimized all-in-one AAV vector reported previously [27]. We attached one cMyc NLS
sequence on each terminus of Nme2-ABE8e while retaining the original promoters for effector
and sgRNA expression (**Figure 3a,** 2x cMyc). By transient transfection of vector backbone
plasmids, the single-AAV construct successfully edited the ABE reporter cell line. To further
improve Nme2-ABE editing efficiency, we then tested three different NLS configurations: 1) one
cMyc NLS on the N-terminus and two cMyc NLSs on the C-terminus (3x cMyc); 2) one Ty1
NLS, which derived from the yeast Ty1 retrotransposon that supports robust nuclear localization
in dPspCas13b fusion proteins, on the N-terminus (Ty1) [60]; and 3) one bipartite SV40 NLS
(BP_SV40) on each terminus (2x BP_SV40) (**Figure 3a**) [61]. When transfecting the vector
plasmid into the ABE reporter cell line, the 2x BP_SV40 construct showed the highest editing
efficiency (**Figure 3b**).
219
The total length of the vector constructed with the 2xBP_SV40 NLS, hereafter Nme2-ABE8e-
U6, is 4998 bp, very close to the packaging limit of AAV. To test whether we could further
reduce the vector size without significantly compromising editing efficiency, we turned to a
recently reported "miniU6" promoter that has been shown to support sgRNA expression and
achieve comparable editing efficiencies as with the complete U6 promoter [62]. Upon
replacement of the U6 promoter with miniU6 promoter, the vector, hereafter Nme2-ABE8e-
miniU6, was shortened to 4860 bp, within the packaging limit of AAV (**Figure 3c**). Both
constructs induced robust editing in the ABE reporter cell line via transient transfection of the
vector plasmids (**Figure 3c**).
229
To avoid potential ABE reporter-specific effects, we further tested both single-AAV vector
backbone plasmids at two endogenous target sites: 1) one of the human dual-target sites, DS12,
and 2) a previously reported Nme2Cas9 target site in the mouse *Rosa26* gene [45]. By plasmid
transfection in human HEK293T or mouse N2a cells, we observed significant editing at these
loci by both vectors, although the Nme2-ABE8e-miniU6 vector was somewhat less efficient
(**Figure 3c**). We thus chose both vector designs for the subsequent *in vivo* study.
236
237   **Hydrodynamic injection of single-AAV vector plasmids corrects the disease**
238   **mutation and phenotype in an adult mouse model of HT1**

To test the *in vivo* editing efficiency and therapeutic potential of the single-AAV constructs, we chose to target a pathogenic mutation associated with the liver disease HT1. HT1 is caused by mutations in fumarylacetoacetate hydrolase (FAH), which catalyzes one step of the tyrosine catabolic pathway. FAH deficiency leads to accumulations of toxic fumarylacetoacetate and succinyl acetoacetate, causing liver, kidney, and CNS damage [63]. The *Fah*[PM/PM] mouse model possesses a G•C to A•T point mutation in the last nucleotide of exon 8, which causes skipping of exon 8 and FAH deficiency (**Figure 4a**). Without treatment, FAH deficient mice will rapidly lose weight and eventually die. The *Fah*[PM/PM] mouse can be treated with 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione (NTBC), an inhibitor of an enzyme upstream within the tyrosine degradation pathway, which prevents toxin accumulation [64]. Previously, we and others have tested various *in vivo* gene-editing tools to treat the *Fah*[PM/PM] mouse model, including Cas9-directed HDR [27,65,66], base editing [67,68], microhomology-directed end joining [69], and prime editing [70]. Among these, multiple approaches including AAV, lipid nanoparticle (LNP), and plasmid hydrodynamic tail-vein injection have been used to deliver the gene-editing agents into this mouse model. However, caveats should be considered when comparing the efficiencies of different gene-editing strategies: only the initial editing efficiency (measured before NTBC withdrawal) reflects the activity of the gene-editing agents, because after NTBC withdraw, the hepatocytes in which *Fah* has been repaired will show clonal expansion over time due to their survival advantage.

We first validated a guide targeting the point mutation by electroporation of the single-AAV vector plasmids, either Nme2-ABE8e-U6 or Nme2-ABE8e-miniU6, into mouse embryonic fibroblasts (MEFs) isolated from a *Fah*[PM/PM] mouse (**Figure 4b**). We detected modest but significant editing by both vectors at the target adenine at position 13 (A13), despite low (~12%) plasmid electroporation efficiencies. We also observed higher levels of bystander editing at A16 with both vectors, and a lower level of bystander editing at A10 for Nme2-ABE8e-U6. Bystander editing at A10 (which changes an active-site-proximal serine into an alanine) has been observed previously with Spy-ABE as well [67]. The effect (if any) of the intronic A16 edit on intron 8 splice donor activity has not been defined (**Figure 4c**).

To test our single-AAV vectors of Nme2-ABE8e *in vivo*, we first performed hydrodynamic tail-vein injections of the AAV-vector plasmids into 10-week-old HT1 mice, or PBS injection as a negative control group [71]. We also injected plasmids expressing Spy-RA6.3, which is a codon-optimized Spy-ABE, as a positive control [67]. Seven days post-injection, we sacrificed 2 mice from each experimental group, and 1 mouse from each control group, to measure the editing efficiency before hepatocyte expansion. We then withdrew NTBC for the rest of the mice for long-term phenotypic study. Before NTBC withdrawal, anti-FAH immunohistochemistry (IHC) staining showed 4.58 ± 1.1% FAH+ hepatocytes from the group that was injected with the Nme2-ABE8e-U6 plasmid, and 1.71 ± 0.49% from the group injected with the Nme2-ABE8e-miniU6 plasmid (**Figure 4d**). The mouse injected with Spy-RA6.3 plasmid showed 4.5% FAH+ hepatocytes, consistent with previously reported data [67] (**Figure 4d**). After NTBC withdrawal, we monitored body weight changes. The PBS injected mice rapidly lost body weight after NTBC withdrawal and were euthanized. By contrast, mice injected with either the Nme2-ABE8e-U6 or Nme2-ABE8e-miniU6 plasmid gradually gained body weight, suggesting rescue of the pathological phenotype (**Figure 4e**). Forty days after NTBC withdrawal, we sacrificed mice from all surviving groups. To determine whether Nme2-ABE8e successfully corrects the *Fah* splicing defect, we extracted total RNA from the livers and performed reverse transcription PCR

(RT-PCR) using primers that spanned exons 5 and 9. By contrast to the PBS-injected mice,
which only showed a 305 bp PCR product corresponding to the truncated mRNA lacking exon
8, we observed that the 405 bp PCR product (containing exon 8) predominated in the Nme2-
ABE treated mice (**Figure 4f**). Sanger sequencing of the 405 bp bands further confirmed the
presence of the corrected G residue at the end of exon 8 (**Figure 4g**). When performing anti-
FAH IHC staining, we observed expansion of FAH+ hepatocytes in the groups that were injected
with either of the single-AAV vector plasmids (**Figure 4h**). Amplicon deep sequencing of
genomic DNA from the livers of treated mice again provided evidence for Nme2-ABE activity
(**Figure 4i**). By contrast to the efficiencies achieved in the MEF cells, we observed a lower
editing at A16 and no significant editing at A10, likely due to partial (A16) or complete (A10)
selection against mice harboring bystander edits at those positions. These data indicate that our
Nme2-ABE8e single-AAV vector plasmids can correct the disease genotype and phenotype of the
*Fah*PM/PM mice *in vivo*.
299
In vivo base editing by single AAV delivered Nme2-ABE8e in the *Fah*PM/PM mice
Encouraged by the initial results, we packaged AAV9 with the Nme2-ABE8e-U6 construct, as
well as the Nme2-ABE8e-miniU6 construct, considering that the relatively smaller size of the
latter may potentially benefit packaging efficiency (**Figure 5a**). However, both constructs
yielded similar vector titers. Next, to confirm the AAV genome integrity, we performed AAV
genomic DNA extraction and alkaline gel electrophoresis. We did not observe any sign of
genome truncation (**Supplemental Figure 4**). We then tail-vein injected 8-week-old *Fah*PM/PM
mice at a dosage of 4×10^{11} vg per mouse. We kept the mice on NTBC for one month before
analyzing the editing efficiency. One month after AAV injection and before NTBC withdrawal,
we sacrificed the mice and performed IHC staining using an anti-FAH antibody. The negative
control groups injected with AAV9 expressing Nme2-ABE and a sgRNA targeting the *Rosa26*
gene did not show any FAH+ hepatocytes. In contrast, we observed 6.49 ± 2.08% FAH+
hepatocytes in the AAV9-Nme2-ABE8e-U6-*Fah* treated group, and 1.62 ± 0.49% FAH+
hepatocytes in AAV9-Nme2-ABE8e-miniU6-*Fah* treated group (**Figure 5b**). Because repair of
1/100,000 hepatocytes was reported to rescue the phenotype, both AAV constructs achieved
efficiencies that were well above the therapeutic threshold [72]. Moreover, the percentage of
edited hepatocytes by AAV9-Nme2-ABE8e-U6-*Fah* was higher than what has been reported
previously by other genome editing strategies [65–67,70]. By targeted deep sequencing, the
editing efficiency at the target adenine (A13) in the AAV9-Nme2-ABE8e-U6 treated group is
0.34 ± 0.14%, while no significant editing was observed in the AAV9-Nme2-ABE8e-miniU6
treated group, possibly due to low efficiency that was below the detection limit of amplicon deep
sequencing (**Figure 5c**). The reason for the higher frequency of FAH+ hepatocytes than the
frequency of editing at the DNA level is likely due to hepatocyte polyploidy [73], as well as the
presence of genomic DNA from nonparenchymal cells. Similar distinctions in FAH+ frequencies
and genomic readouts were also observed in previous studies using this mouse model [67,70]. We
also measured the editing efficiency at the *Rosa26* locus and observed ~5% editing efficiency at
the target site, indicating that the efficiency of AAV9-delivered Nme2-ABE8e is target site
dependent, and higher-efficiency sites can be identified (**Figure 5d**). We did not detect any
above-background level of indel, indicating that single-AAV delivered Nme2-ABE8e can install
precise editing *in vivo* without generating unwanted indels (**Figure 5c, d**).
330
331 Nme2Cas9 is known to be highly accurate in nuclease based editing in cells [27,30,39], and our
332 mismatch scanning experiments (**Figure 1e**) strongly suggest that the same will be true *in vivo*.

To evaluate potential Cas9-dependent off-target effects in the AAV9-injected mice, we searched for genome-wide off-target sites for Nme2Cas9 using Cas-OFFinder [74], allowing for up to 6 mismatches. We then performed amplicon deep sequencing in the AAV9 treated livers at the two top-ranking potential off-target sites, each including 5 mismatches. We did not detect any above-background A•T-to-G•C editing at these sites (**Supplementary Figure 5**).

Optimizing effector and sgRNA arrangement improves editing efficiency by AAV delivery

Previous studies have shown that effector and sgRNA placement and orientation within the AAV genome can affect transgene expression levels and editing efficiencies [18, 26, 85]. To test if different arrangements of sgRNA and effector cassettes in the Nme2-ABE8e all-in-one AAV construct can further increase *in vivo* editing efficiency, we moved the U6-sgRNA cassette to the 3' end of the AAV genome and reversed its orientation, similar to the optimal arrangement reported by Fry *et al.* [85] (**Figure 6a**). Using the same *Rosa26* guide described above (**Figure 5d**), we packaged the rearranged construct in AAV9 capsids and performed tail-vein injections in 8-week-old mice. We observed significantly improved editing efficiency (34 ± 11.6%) with the optimized construct (**Figure 6b**), which is significantly greater than the 4.7 ± 0.94% efficiency with the original configuration, and comparable to the editing efficiency achieved previously by dual-AAV delivered split-intein SpyCas9-ABEmax targeting the *Dmmt1* gene in adult mouse liver (38 ± 2.9%) [18]. We conclude that efficient adenine base editing can be achieved *in vivo* via single-AAV delivery of the Nme2-ABE8e system.

Discussion

Rapidly evolving and precise genome editing tools such as base editors and prime editors possess great potential to address the root causes of human genetic diseases [2,75,76]. However, the safe and effective delivery of genome editors remains a major challenge. Viral delivery using AAV, which is the only FDA-approved *in vivo* gene therapy vector to date, has a limited packaging capacity. Previous studies using AAV to deliver base editors or prime editors have been reported to be effective in rodents [7,8,10–13,15–18,21], but all of them required multiple vectors and most of them required high viral dosage [7,10–13,17,18]. To date, AAV administered at high dose has been reported to relate to severe toxicity or even death in nonhuman primates, piglets, and humans [23,24,77–79]. Engineering single-AAV deliverable genome editing tools has the potential to achieve therapeutic benefits at lower dosage, which would not only ease manufacturing burdens but also reduce likelihoods of serious adverse events [80,81].

Engineering new genome editors based on compact Cas9s is an alternative approach to address this issue. Previously, single-AAV delivery of Cas9 nucleases has been reported to generate NHEJ-based editing *in vivo* [26,30,82]. Precision editing via HDR has also been achieved with single-AAV systems, albeit with low efficiencies [27,83]. The only single-AAV BE system reported previously (which was not tested *in vivo*) was based on SauCas9, which has limited targeting range due to its PAM constraints. In this study, we constructed and characterized a compact, accurate adenine base editor, Nme2-ABE8e, which can target many sites that are inaccessible to SauCas9 base editors due to their distinct PAMs. We showed that Nme2-ABE8e with a sgRNA can be packaged into a single AAV vector, and a single intravenous injection in an adult disease mouse model of tyrosinemia reversed both the disease mutation and phenotypes. Furthermore, the dosage used in this study (4×10^{11} per mouse or 2×10^{13} vg/kg) is well below the 1×10^{14} vg/kg systemic doses that have been tolerated in clinical trials [77,78].

Although the editing efficiency was modest at the *Fah* disease locus, we reached therapeutic thresholds for HT1, and the initial editing efficiency of the AAV9-delivered Nme2-ABE8e-U6 construct exceeded that reported previously at this locus using other precision genome editing tools and delivery approaches [65–67,70]. Moreover, by optimizing transgene orientation in the AAV construct, we achieved significantly improved editing efficiency (~5-fold increase) by AAV9-delivered Nme2-ABE8e targeting the *Rosa26* gene, comparable to that achieved previously by dual-AAV delivered split-intein SpyCas9-ABEmax in adult mouse liver targeting another genomic site [18]. Future optimization of this system promises to further improve efficiency. There are multiple potential explanations for the inconsistent editing efficiencies we achieved with our first-generation Nme2Cas9 ABE that suggest directions for future improvement. First, structural analyses of Nme2Cas9 [84] indicate that the position of the N-terminally fused TadA8e domain relative to the predicted path of the displaced strand is not optimal. With other effectors, domain-inlaid deaminase fusions have proven to be advantageous in some contexts [29,54], and the same is likely to be true with Nme2-ABEs. Second, the current Nme2-ABE8e has a wide editing window that could result in increased bystander editing. For example, we observed higher editing at a bystander adenine (A16) at the *Fah* locus compared to the on-target adenine at A13. Optimizations of linker composition and length [3,53], as well as domain-inlaid systems [29,54–58], may confer greater control over the editing window and improved efficiency in editing the intended position.

Our studies also suggest factors that must be considered for optimal guide expression. In contrast
to a previous study that showed no significant difference between the U6 and miniU6 promoters
in supporting Cas9 editing efficiency [62], we found that the construct with the miniU6 promoter
was consistently less efficient than that with the U6 promoter, including via AAV delivery.
Because the previous study compared these promoters in T cells by lentivirus transduction, our
observations may only apply to certain delivery strategies and cellular contexts.
408
In summary, we have engineered and characterized Nme2-ABE8e editing in mammalian cell
culture and achieved efficient *in vivo* editing by delivery of a single AAV vector. To our
knowledge, this is the first single-AAV-delivered *in vivo* base editing reported to date [29]. We
anticipate that Nme2-ABE8e, with its distinct PAM specificity, editing window, and high
accuracy, will provide additional targetability, safety, and therapeutic potential for genome
engineering applications.
415

Methods

Cell culture. HEK293T cells (ATCC CRL-3216), ABE reporter cells, MEF cells, and mouse N2a cells (ATCC CCL-131) were cultured in in Dulbecco's Modified Eagle Media (DMEM, Genesee Scientific Cat. #: 25-500) supplemented with 10% Fetal Bovine Serum (Gibco Cat. #: 26140079). Rett syndrome human patient-derived fibroblasts (hPDFs) were obtained from the Rett Syndrome Research Trust and cultured with DMEM supplemented with 15% FBS and 1x non-essential amino acids (Gibco Cat. #: 11140050). All cells were incubated in a 37°C incubator with 5% CO_2.

Molecular cloning. To generate the CMV-Nme2-ABE8e and the CMV-Nme2-ABE7.10 plasmids used in **Figure 1**, the Nme2-ABE8e, Nme2-ABE7.10, Spy-ABE8e, and Spy-ABE7.10 constructs were cloned into the pCMV-PE2 vector backbone (Addgene #132775) by Gibson assembly. Briefly, the pCMV-PE2 plasmid was digested with NotI and PmeI restriction enzymes, and the plasmid backbone was then Gibson-assembled with five fragments: N-terminal NLS, TadA8e (for Nme2-ABE8e or Spy-ABE8e), or TadA-TadA*7.10 (for Nme2-ABE7.10 or Spy-ABE7.10), the linker, Nme2Cas9-D16A or SpCas9-D10A nickase, and the C-terminal NLS. The ABE reporter construct was cloned by site-directed mutagenesis to change the 47th amino acid (Glutamine, CAG) of the mCherry coding sequence to a stop codon (TAG). The ABE reporter was further cloned into a lentiviral transfer vector backbone (Addgene #99373) by Gibson assembly. The sgRNA expression plasmids used in **Figure 1** were cloned from pBluescriptSKII (Addene #74705) in two steps. First, pBluescriptSKII was digested by NotI and XbaI, and a gene fragment that contains a U6 promoter, a type-IIS restriction cloning site (BfuAI), and a tracrRNA was assembled into the backbone by Gibson assembly. The plasmids were further digested with BfuAI and ligated to the annealed oligos to insert the guide sequences. The single-AAV vector plasmids in **Figure 3a** were cloned from the Addgene #119924 plasmid by replacing the Nme2Cas9 sequence with the Nme2-ABE8e sequence, and then subsequently re-cloned to encode different NLS configurations and the miniU6 promoter by restriction enzyme digestion and Gibson assembly. The NLS configuration of Nme2-ABE8e mRNA used in **Figure 2b** was 2x_BPSV40, and the plasmid used in **Figure 2d** was the single-AAV Nme2-ABE8e-U6 plasmid shown in **Figure 3a** (2xBP_SV40). To clone the AAV-Nme2-ABE8e_V2 plasmid in **Figure 6**, first, the 2xBP_SV40 plasmid from **Figure 3a** was digested by PmeI and NotI, the AAV backbone fragment was then Gibson-assembled with the fragment containing U1a-Nme2-ABE8e, and the fragment containing U6-sgRNA with homology sequences by overhang PCR. Most of the plasmids used in **Supplementary Figure 3a** were previously deposited in Addgene (AcrIIC3, Addgene #85713; AcrIIC4, Addgene #113434; Acr-E2, Addgene #85677). To clone the plasmid expressing AcrIIA4, Addgene #85713 plasmid was digested with XhoI and BamHI and Gibson-assembled with a gene fragment containing AcrIIA4 coding sequence [46]. In **Supplementary Figure 3b**, the AcrIIC3-MRE122 plasmid was cloned from Addgene plasmid #129531 by replacing the *Rosa26*-targeting guide with the DS12-targeting guide through restriction enzyme cloning. Subsequent replacement of the coding sequence of AcrIIC3 with AcrIIC4 generated the AcrIIC4-MRE122 plasmid. Sequences of plasmids first described in this paper can be found in the **Supplementary note** and will be made available from Addgene.

ABE reporter HEK293T cell line. Lentivirus was produced following instructions from Addgene (https://www.addgene.org/protocols/lentivirus-production/). Briefly, HEK293T cells were transfected with the transfer vector and the packaging plasmids psPAX2 (Addgene #12260)

and pMD2.G (Addgene #12259), using Lipofectamine 3000 (ThermoFisher Cat. #: L3000015).
Two days later, the medium was collected and filtered through a 0.45 µm filter (Cytiva Cat. #:
6780-25040) to remove cell debris. The viral titer was determined using Lenti-X™GoStix™
(Takara Bio Cat. #: 631280). HEK293T cells were transduced with lentivirus encoding the ABE
reporter at varying dilutions (1:10, 1:100, 1:500, and 1:1000) in the presence of 8 µg/ml
polybrene (Millipore Sigma Cat. #: TR-1003-G). Three days after transduction, the media was
removed, and fresh media was supplemented with 2 µg/ml Puromycin to select cells expressing
the full-length reporter construct. Seven days after selection, the puromycin-resistant cells were
collected. Single-cell clones were established by serial dilution in 96-well plates.
472
473 **Fluorescent reporter assay.** Forty-eight hours after transfection, cells were trypsinized and
474 harvested into microcentrifuge tubes. After centrifuging at 300 x g for 3 mins, cells were
475 resuspended into 150ul 1xPBS for flow cytometry analysis (MACSQuant VYB). For each
476 sample, 10,000 events were counted for FACS analysis. Data was analyzed using Flowjo v10.
477 Representative gating strategy can be found in **Supplementary Figure 2**.
478
***In vitro* transcription of Nme2-ABE8e mRNA.** Nme2-ABE8e mRNA was *in vitro*
transcribed using the NEB HiScribe T7 RNA synthesis kit, from 500ng of a linearized template.
Uridine was fully substituted with 1-methylpseudouridine, and mRNA was capped co-
transcriptionally using CleanCap AG analog (TriLink Biotechnologies Cat. #N-1081 and N-
7113, respectively). All enzymes were purchased from New England Biolabs. Transcription was
conducted according to the manufacturer's protocol with the following amendments:
transcriptions were completed in $1\times$ NEB HiScribe transcription buffer, and 4 mM CleanCap
AG was used during transcription. Transcription reactions were incubated at 37 °C for 2 hours
then treated with 0.4 U/µl DNase I (final concentration) for 15 min at 37 °C. mRNAs were
purified with NEB Monarch RNA purification columns and treated with pyrophosphatase for 1
hour with 0.25 U/µg Antarctic phosphatase (final concentration) in 1X Antarctic phosphatase
buffer. The final product was further purified with a NEB Monarch RNA column and eluted in
water.
492
Transfection and electroporation. For plasmid transfection, cells were seeded in 24-well
plates at 80,000 cells per well in culture media and incubated overnight. Briefly, plasmids were
transfected at 400ng per well when targeting the ABE reporter, and 1ug per well for the
endogenous target sites. Specifically, in **Figure 1c**, **1e**, **2d**, **3b**, **3c**, and **Supplementary**
Figure 3, plasmids were transfected using Lipofectamine3000 (ThermoFisher Cat #L3000001),
while in **Figure 1d** and **Supplementary Figure 2**, plasmids were transfected using
Lipofectamine2000 (ThermoFisher Cat #11668030), following the manufacturer's protocols.
Electroporation was performed using Neon TransfectionSystem 10 ul kit (ThermoFisher Cat #:
MPK 1096), with the following electroporation parameters: Pulse voltage (1650 v), Pulse width
(20 ms), Pulse number (1). Specifically, in **Figure 2a**, 264 ng Nme2-ABE8e mRNA and 100
pmole of sgRNA were electroporated into 50,000 Rett Syndrome hPDFs, while in **Figure 4c**, 1
ug plasmid DNA was electroporated into 100,000 MEF cells.
505
506 **AAV production**. AAV vector packaging was done at the Viral Vector Core of the Horae Gene
507 Therapy Center at the UMass Chan Medical School as previously described [86]. Constructs
508 were packaged in AAV9 capsids and viral titers were determined by digital droplet PCR and gel
509 electrophoresis followed by silver staining.

AAV genomic DNA extraction and alkaline agarose gel electrophoresis. Genomic DNAs were extracted from 10^{11} vg AAV by incubating with 20 units of DNase I (ThermoFisher Cat. #: EN0521) at 37 °C for 30 minutes and then with an equal volume of 2 x Pronase solution (Sigma-Aldrich, Cat. # 10165921001) for 4 hours. The genomic DNA was subsequently purified by phenol-chloroform (ThermoFisher Cat. #:15593-049) extraction and ethanol precipitation. DNA pellets were resuspended in water and analyzed by alkaline agarose gel electrophoresis and SYBR Gold staining.

Mouse experiments. All animal study protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at UMass Chan Medical School. The *Fah*$^{PM/PM}$ mice were kept on water supplemented with 10 mg/L 2-(2-nitro-4-trifluoromethylbenzoyl)-1,3-cyclohexanedione (NTBC; Sigma-Aldrich, Cat. #: PHR1731-1G). Mice with more than 20% weight loss were humanely euthanized according to IACUC guidelines. For hydrodynamic tail vein injections, plasmids were prepared by EndoFree Plasmid Maxi Kit (Qiagen Cat. #12362). Briefly, 30 ug SpCas8-RA6.3 and 30 ug sgRNA plasmids, or 60 ug Nme2-ABE8e single AAV plasmids were suspended in 2 ml saline and injected via tail vein within 5-7 seconds into 10-week-old *Fah*$^{PM/PM}$ mice. Mice were euthanized 7 days after injection and livers were collected for analysis. For AAV injection, a dosage of 4×10^{11} vg per mouse (in 200 ul saline) was tail-vein injected into 8-week-old *Fah*$^{PM/PM}$ mice.

Genomic DNA extraction from cultured cells and mouse liver. For cultured cells, genomic DNAs were extracted 72 hours after plasmid transfection, or 48 hours after mRNA and sgRNA electroporation. Briefly, cell culture media were aspirated, and cells were washed with 1x PBS. Genomic DNAs were prepared using QuickExtract DNA Extraction Solution (Lucigen) following the manufacturer's protocols. To extract genomic DNA from mouse livers, all 5 lobes of mouse liver were combined and pulverized in liquid nitrogen, and 15 mg of the tissue from each mouse liver was used for genomic DNA extraction using GenElute Mammalian Genomic DNA Miniprep Kit (Millipore Sigma Cat. #: G1N350).

Amplicon deep sequencing and data analysis. Genomic DNA was amplified by PCR using Q5 High-Fidelity 2X Master Mix (NEB Cat. #: M0492) for 20 cycles. One microliter of the unpurified PCR product was used as a template for 20 cycles of barcoding PCR. The barcoding PCR reactions were further pooled and gel-extracted using Zymo gel extraction kit and DNA clean & concentrator (Zymo research Cat. #: 11-301 and 11-303) and quantified by Qubit 1x dsDNA HS assay kits (Thermo Fisher Scientific, Cat #: Q32851). Sequencing of the pooled amplicons was performed using an Illumina MiniSeq system (300-cycles, FC-420-1004) following the manufacturer's protocol. The raw MiniSeq output was de-multiplexed using bcl2fastq2 (Illumina, version 2.20.0) with the flag --barcode-mismatches 0. To align the generated fastq files and to quantify editing efficiency, CRISPResso2 [87] (version 2.0.40) was used in batch mode with base editor output and the following flags: -w 15, -q 30. Indel frequency = (insertions reads + deletions reads)/all aligned reads x 100%.

Immunohistochemistry (IHC). Mice were euthanized by CO_2 asphyxiation and livers were fixed with 10% neutral buffered formalin (Epredia Cat. #: 5735), sectioned at 10um, and stained with hematoxylin and eosin for pathology analysis. For IHC, liver sections were dewaxed,

556 rehydrated, and stained using an anti-FAH antibody (Abcam Cat. #: ab83770) at 1:400 dilution
557 as described previously [67].
558
559 **Reverse transcription PCR**. All 5 lobes of each mouse liver were combined and pulverized in
560 liquid nitrogen. Total RNA was extracted from 50 mg of liver tissue using TRIzol reagent
561 (ThermoFisher Cat. #: 15596026) and reverse-transcribed using SuperScript III First-Strand
562 Synthesis System (ThermoFisher Cat. #:18080051). PCR was performed using primers
563 previously described [67].
564
565 **Author contributions**
566
567 H.Z., X.D.G., and E.J.S conceived the study. H.Z. designed, performed, and analyzed the *in vivo*
568 experiments. H.Z., N.B., O.O., P.L., and X.D.G., designed, performed, and analyzed the *in vitro*
569 experiments. T.R. and H.C. analyzed the deep sequencing data. E.J.S., W.X., and S.A.W.
570 supervised research. H.Z. and E.J.S. wrote the manuscript with contributions from N.B., P.L.,
571 and H.C. All authors edited the manuscript.
572

573 **Acknowledgements**
574
We thank Yueying Cao and Greg Cottle for their assistance with the mouse colony maintenance
and tail vein injections. We thank Zexiang Chen for his help with the *in vitro* mRNA
transcription, Nadia Amrani for providing the AcrIIA4 plasmid, Tingting Jiang for providing
*Fah*PM/PM MEF cells and Spy-RA6.3 plasmid, and NamKung Suk for his assistance with the
alkaline gel electrophoresis. We also thank the UMMS Viral Vector Core for AAV packaging
service, the UMMS Morphology Core for tissue sectioning and IHC staining, and the Rett
Syndrome Research Trust for patient-derived fibroblasts. We are grateful to all members of the
Gao, Wolfe, Xue and Sontheimer labs for their valuable discussions, advice, and helpful
feedback.
584
585 **Funding**
586
This work was supported by the National Institutes of Health [grant numbers R01GM125797 to
E.J.S., F31GM143879 to N.B., and R01HL150669 to S.A.W.; the Rett Syndrome Research
Trust [to S.A.W. and E.J.S.]; and the Leducq Foundation [grant number 20CVD04 to E.J.S.].
G.G. was supported by grants from National Institutes of Health (R01NS076991-01,
P01AI100263-01, P01HL131471-02, R01AI121135, UG3HL147367-01, R01HL097088, and
U19AI149646-01). W.X. was supported by grants from the National Institutes of Health
(DP2HL137167, P01HL131471 and UH3HL147367), American Cancer Society (129056-RSG-
16-093), and the Cystic Fibrosis Foundation.
595

596 **Declaration of Competing Interests**
597
598 H.Z., N.B., P.L., X.D.G., S.A.W., W.X., and E.J.S. are co-inventors on patent filings related to
599 this work. G.G. is scientific co-founder, scientific advisor, and equity holder of Voyager
600 Therapeutics, Adrenas Therapeutics, and Aspa Therapeutics. E.J.S. is a co-founder, scientific
601 advisor, and equity holder of Intellia Therapeutics.
602

References

[1] M.J. Landrum, J.M. Lee, M. Benson, G. Brown, C. Chao, S. Chitipiralla, B. Gu, J. Hart, D. Hoffman, J. Hoover, W. Jang, K. Katz, M. Ovetsky, G. Riley, A. Sethi, R. Tully, R. Villamarin-Salomon, W. Rubinstein, D.R. Maglott, ClinVar: public archive of interpretations of clinically relevant variants, Nucleic Acids Res. 44 (2016) D862–8.

[2] H.A. Rees, D.R. Liu, Base editing: precision chemistry on the genome and transcriptome of living cells, Nat. Rev. Genet. 19 (2018) 770–788.

[3] N.M. Gaudelli, A.C. Komor, H.A. Rees, M.S. Packer, A.H. Badran, D.I. Bryson, D.R. Liu, Programmable base editing of A•T to G•C in genomic DNA without DNA cleavage, Nature. 551 (2017) 464–471.

[4] A.C. Komor, Y.B. Kim, M.S. Packer, J.A. Zuris, D.R. Liu, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, Nature. 533 (2016) 420–424.

[5] T. Rothgangl, M.K. Dennis, P.J.C. Lin, R. Oka, D. Witzigmann, L. Villiger, W. Qi, M. Hruzova, L. Kissling, D. Lenggenhager, C. Borrelli, S. Egli, N. Frey, N. Bakker, J.A. Walker 2nd, A.P. Kadina, D.V. Victorov, M. Pacesa, S. Kreutzer, Z. Kontarakis, A. Moor, M. Jinek, D. Weissman, M. Stoffel, R. van Boxtel, K. Holden, N. Pardi, B. Thöny, J. Häberle, Y.K. Tam, S.C. Semple, G. Schwank, In vivo adenine base editing of PCSK9 in macaques reduces LDL cholesterol levels, Nat. Biotechnol. 39 (2021) 949–957.

[6] K. Musunuru, A.C. Chadwick, T. Mizoguchi, S.P. Garcia, J.E. DeNizio, C.W. Reiss, K. Wang, S. Iyer, C. Dutta, V. Clendaniel, M. Amaonye, A. Beach, K. Berth, S. Biswas, M.C. Braun, H.-M. Chen, T.V. Colace, J.D. Ganey, S.A. Gangopadhyay, R. Garrity, L.N. Kasiewicz, J. Lavoie, J.A. Madsen, Y. Matsumoto, A.M. Mazzola, Y.S. Nasrullah, J. Nneji, H. Ren, A. Sanjeev, M. Shay, M.R. Stahley, S.H.Y. Fan, Y.K. Tam, N.M. Gaudelli, G. Ciaramella, L.E. Stolz, P. Malyala, C.J. Cheng, K.G. Rajeev, E. Rohde, A.M. Bellinger, S. Kathiresan, In vivo CRISPR base editing of PCSK9 durably lowers cholesterol in primates, Nature. 593 (2021) 429–434.

[7] L. Wang, W. Xue, H. Zhang, R. Gao, H. Qiu, J. Wei, L. Zhou, Y.-N. Lei, X. Wu, X. Li, C. Liu, J. Wu, Q. Chen, H. Ma, X. Huang, C. Cai, Y. Zhang, B. Yang, H. Yin, L. Yang, J. Chen, Eliminating base-editor-induced genome-wide and transcriptome-wide off-target mutations, Nat. Cell Biol. 23 (2021) 552–563.

[8] P. Liu, S.-Q. Liang, C. Zheng, E. Mintzer, Y.G. Zhao, K. Ponnienselvan, A. Mir, E.J. Sontheimer, G. Gao, T.R. Flotte, S.A. Wolfe, W. Xue, Improved prime editors enable pathogenic allele correction and cancer modelling in adult mice, Nat. Commun. 12 (2021). https://doi.org/10.1038/s41467-021-22295-w.

[9] L. Villiger, T. Rothgangl, D. Witzigmann, R. Oka, P.J.C. Lin, W. Qi, S. Janjuha, C. Berk, F. Ringnalda, M.B. Beattie, M. Stoffel, B. Thöny, J. Hall, H. Rehrauer, R. van Boxtel, Y.K. Tam, G. Schwank, In vivo cytidine base editing of hepatocytes without detectable off-target mutations in RNA and DNA, Nat Biomed Eng. 5 (2021) 179–189.

[10] D. Böck, T. Rothgangl, L. Villiger, L. Schmidheini, N. Mathis, E. Ioannidi, S. Kreutzer, Z. Kontarakis, N. Rimann, H.M. Grisch-Chan, B. Thöny, G. Schwank, Treatment of a metabolic liver disease by in vivo prime editing in mice, bioRxiv. (2021) 2021.08.17.456632. https://doi.org/10.1101/2021.08.17.456632.

[11] L.W. Koblan, M.R. Erdos, C. Wilson, W.A. Cabral, J.M. Levy, Z.-M. Xiong, U.L. Tavarez, L.M. Davison, Y.G. Gete, X. Mao, G.A. Newby, S.P. Doherty, N. Narisu, Q. Sheng, C. Krilow, C.Y. Lin, L.B. Gordon, K. Cao, F.S. Collins, J.D. Brown, D.R. Liu, In vivo base

649 editing rescues Hutchinson-Gilford progeria syndrome in mice, Nature. 589 (2021) 608–614.

650 [12] S.-M. Ryu, T. Koo, K. Kim, K. Lim, G. Baek, S.-T. Kim, H.S. Kim, D.-E. Kim, H. Lee,
651 E. Chung, J.-S. Kim, Adenine base editing in mouse embryos and an adult mouse model of
652 Duchenne muscular dystrophy, (n.d.). https://doi.org/10.1038/nbt.xxxx.

653 [13] F. Chemello, A.C. Chai, H. Li, C. Rodriguez-Caycedo, E. Sanchez-Ortiz, A. Atmanli,
654 A.A. Mireault, N. Liu, R. Bassel-Duby, E.N. Olson, Precise correction of Duchenne muscular
655 dystrophy exon deletion mutations by base and prime editing, Sci Adv. 7 (2021).
656 https://doi.org/10.1126/sciadv.abg4910.

657 [14] S. Suh, E.H. Choi, H. Leinonen, A.T. Foik, G.A. Newby, W.-H. Yeh, Z. Dong, P.D.
658 Kiser, D.C. Lyon, D.R. Liu, K. Palczewski, Restoration of visual function in adult mice with an
659 inherited retinal disease via adenine base editing, Nat Biomed Eng. 5 (2021) 169–178.

660 [15] S. Zhi, Y. Chen, G. Wu, J. Wen, J. Wu, Q. Liu, Y. Li, R. Kang, S. Hu, J. Wang, P.
661 Liang, J. Huang, Dual-AAV delivering split prime editor system for in vivo genome editing, Mol.
662 Ther. (2021). https://doi.org/10.1016/j.ymthe.2021.07.011.

663 [16] W.-H. Yeh, O. Shubina-Oleinik, J.M. Levy, B. Pan, G.A. Newby, M. Wornow, R. Burt,
664 J.C. Chen, J.R. Holt, D.R. Liu, In vivo base editing restores sensory transduction and transiently
665 improves auditory function in a mouse model of recessive deafness, Sci. Transl. Med. 12 (2020).
666 https://doi.org/10.1126/scitranslmed.aay9101.

667 [17] C.K.W. Lim, M. Gapinske, A.K. Brooks, W.S. Woods, J.E. Powell, M.A. Zeballos C, J.
668 Winter, P. Perez-Pinera, T. Gaj, Treatment of a Mouse Model of ALS by In Vivo Base Editing,
669 Mol. Ther. 28 (2020) 1177–1189.

670 [18] J.M. Levy, W.-H. Yeh, N. Pendse, J.R. Davis, E. Hennessey, R. Butcher, L.W. Koblan, J.
671 Comander, Q. Liu, D.R. Liu, Cytosine and adenine base editing of the brain, liver, retina, heart
672 and skeletal muscle of mice via adeno-associated viruses, Nat Biomed Eng. 4 (2020) 97–110.

673 [19] Z. Wu, H. Yang, P. Colosi, Effect of genome size on AAV vector packaging, Mol. Ther.
674 18 (2010) 80–86.

675 [20] J.Y. Dong, P.D. Fan, R.A. Frizzell, Quantitative analysis of the packaging capacity of
676 recombinant adeno-associated virus, Hum. Gene Ther. 7 (1996) 2101–2112.

677 [21] L. Villiger, H.M. Grisch-Chan, H. Lindsay, F. Ringnalda, C.B. Pogliano, G. Allegri, R.
678 Fingerhut, J. Häberle, J. Matos, M.D. Robinson, B. Thöny, G. Schwank, Treatment of a
679 metabolic liver disease by in vivo genome base editing in adult mice, Nat. Med. 24 (2018) 1519–
680 1525.

681 [22] J. Winter, A. Luu, M. Gapinske, S. Manandhar, S. Shirguppe, W.S. Woods, J.S. Song, P.
682 Perez-Pinera, Targeted exon skipping with AAV-mediated split adenine base editors, Cell
683 Discov. 5 (2019) 41.

684 [23] C. Hinderer, N. Katz, E.L. Buza, C. Dyer, T. Goode, P. Bell, L.K. Richman, J.M.
685 Wilson, Severe Toxicity in Nonhuman Primates and Piglets Following High-Dose Intravenous
686 Administration of an Adeno-Associated Virus Vector Expressing Human SMN, Hum. Gene
687 Ther. 29 (2018) 285–298.

688 [24] J.M. Wilson, T.R. Flotte, Moving Forward After Two Deaths in a Gene Therapy Trial of
689 Myotubular Myopathy, Human Gene Therapy. 31 (2020) 695–696.
690 https://doi.org/10.1089/hum.2020.182.

691 [25] L. Wang, J. Smith, C. Breton, P. Clark, J. Zhang, L. Ying, Y. Che, J. Lape, P. Bell, R.
692 Calcedo, E.L. Buza, A. Saveliev, V.V. Bartsevich, Z. He, J. White, M. Li, D. Jantz, J.M. Wilson,
693 Meganuclease targeting of PCSK9 in macaque liver leads to stable reduction in serum
694 cholesterol, Nat. Biotechnol. 36 (2018) 717–725.

695 [26] F.A. Ran, L. Cong, W.X. Yan, D.A. Scott, J.S. Gootenberg, A.J. Kriz, B. Zetsche, O.

Shalem, X. Wu, K.S. Makarova, E.V. Koonin, P.A. Sharp, F. Zhang, In vivo genome editing using Staphylococcus aureus Cas9, Nature. 520 (2015) 186–191.

[27]   R. Ibraheim, P.W.L. Tai, A. Mir, N. Javeed, J. Wang, T.C. Rodríguez, S. Namkung, S. Nelson, E.S. Khokhar, E. Mintzer, S. Maitland, Z. Chen, Y. Cao, E. Tsagkaraki, S.A. Wolfe, D. Wang, A.A. Pai, W. Xue, G. Gao, E.J. Sontheimer, Self-inactivating, all-in-one AAV vectors for precision Cas9 genome editing via homology-directed repair in vivo, Nat. Commun. 12 (2021) 6267.

[28]   M.L. Maeder, M. Stefanidakis, C.J. Wilson, R. Baral, L.A. Barrera, G.S. Bounoutas, D. Bumcrot, H. Chao, D.M. Ciulla, J.A. DaSilva, A. Dass, V. Dhanapal, T.J. Fennell, A.E. Friedland, G. Giannoukos, S.W. Gloskowski, A. Glucksmann, G.M. Gotta, H. Jayaram, S.J. Haskett, B. Hopkins, J.E. Horng, S. Joshi, E. Marco, R. Mepani, D. Reyon, T. Ta, D.G. Tabbaa, S.J. Samuelsson, S. Shen, M.N. Skor, P. Stetkiewicz, T. Wang, C. Yudkoff, V.E. Myer, C.F. Albright, H. Jiang, Development of a gene-editing approach to restore vision loss in Leber congenital amaurosis type 10, Nat. Med. 25 (2019) 229–233.

[29]   M.T. Nguyen Tran, M.K.N. Mohd Khalid, Q. Wang, J.K.R. Walker, G.E. Lidgerwood, K.L. Dilworth, L. Lisowski, A. Pébay, A.W. Hewitt, Engineering domain-inlaid SaCas9 adenine base editors with reduced RNA off-targets and increased on-target DNA editing, Nat. Commun. 11 (2020) 4871.

[30]   A. Edraki, A. Mir, R. Ibraheim, I. Gainetdinov, Y. Yoon, C.-Q. Song, Y. Cao, J. Gallant, W. Xue, J.A. Rivera-Pérez, E.J. Sontheimer, A Compact, High-Accuracy Cas9 with a Dinucleotide PAM for In Vivo Genome Editing, Mol. Cell. 73 (2019) 714–726.e4.

[31]   Z. Liu, S. Chen, Y. Jia, H. Shan, M. Chen, Y. Song, L. Lai, Z. Li, Efficient and high-fidelity base editor with expanded PAM compatibility for cytidine dinucleotide, Sci. China Life Sci. 64 (2021) 1355–1367.

[32]   R. Xu, R. Qin, H. Xie, J. Li, X. Liu, M. Zhu, Y. Sun, Y. Yu, P. Lu, P. Wei, Genome editing with type II-C CRISPR-Cas9 systems from Neisseria meningitidis in rice, Plant Biotechnol. J. (2021). https://doi.org/10.1111/pbi.13716.

[33]   K. Hua, X. Tao, J.-K. Zhu, Expanding the base editing scope in rice by using Cas9 variants, Plant Biotechnol. J. 17 (2019) 499–504.

[34]   L. Yang, X. Zhang, L. Wang, S. Yin, B. Zhu, L. Xie, Q. Duan, H. Hu, R. Zheng, Y. Wei, L. Peng, H. Han, J. Zhang, W. Qiu, H. Geng, S. Siwko, X. Zhang, M. Liu, D. Li, Correction to: Increasing targeting scope of adenosine base editors in mouse and rat embryos through fusion of TadA deaminase with Cas9 variants, Protein Cell. 10 (2019) 700.

[35]   M.F. Richter, K.T. Zhao, E. Eton, A. Lapinaite, G.A. Newby, B.W. Thuronyi, C. Wilson, L.W. Koblan, J. Zeng, D.E. Bauer, J.A. Doudna, D.R. Liu, Phage-assisted evolution of an adenine base editor with improved Cas domain compatibility and activity, Nat. Biotechnol. 38 (2020) 883–891.

[36]   N.M. Gaudelli, D.K. Lam, H.A. Rees, N.M. Solá-Esteves, L.A. Barrera, D.A. Born, A. Edwards, J.M. Gehrke, S.-J. Lee, A.J. Liquori, R. Murray, M.S. Packer, C. Rinaldi, I.M. Slaymaker, J. Yen, L.E. Young, G. Ciaramella, Directed evolution of adenine base editors with increased activity and therapeutic application, Nat. Biotechnol. 38 (2020) 892–900.

[37]   H.A. Rees, A.C. Komor, W.-H. Yeh, J. Caetano-Lopes, M. Warman, A.S.B. Edge, D.R. Liu, Improving the DNA specificity and applicability of base editing through protein engineering and protein delivery, Nat. Commun. 8 (2017) 15790.

[38]   D. Kim, K. Lim, S.-T. Kim, S.-H. Yoon, K. Kim, S.-M. Ryu, J.-S. Kim, Genome-wide target specificities of CRISPR RNA-guided programmable deaminases, Nat. Biotechnol. 35 (2017) 475–480.

[39]   J. Wen, T. Cao, J. Wu, Y. Chen, S. Zhi, Y. Huang, P. Zhen, G. Wu, L. Aagaard, J. Zhong, P. Liang, J. Huang, Single AAV-mediated CRISPR-Nme2Cas9 efficiently reduces mutant hTTR expression in a transgenic mouse model of transthyretin amyloidosis, Mol. Ther. (2021). https://doi.org/10.1016/j.ymthe.2021.05.010.

[40]   B.J. Rauch, M.R. Silvis, J.F. Hultquist, C.S. Waters, M.J. McGregor, N.J. Krogan, J. Bondy-Denomy, Inhibition of CRISPR-Cas9 with Bacteriophage Proteins, Cell. 168 (2017) 150–158.e10.

[41]   J. Lee, A. Mir, A. Edraki, B. Garcia, N. Amrani, H.E. Lou, I. Gainetdinov, A. Pawluk, R. Ibraheim, X.D. Gao, P. Liu, A.R. Davidson, K.L. Maxwell, E.J. Sontheimer, Potent Cas9 Inhibition in Bacterial and Human Cells by AcrIIC4 and AcrIIC5 Anti-CRISPR Proteins, MBio. 9 (2018). https://doi.org/10.1128/mBio.02321-18.

[42]   S. Hwang, K.L. Maxwell, Meet the Anti-CRISPRs: Widespread Protein Inhibitors of CRISPR-Cas Systems, CRISPR J. 2 (2019) 23–30.

[43]   A. Pawluk, N. Amrani, Y. Zhang, B. Garcia, Y. Hidalgo-Reyes, J. Lee, A. Edraki, M. Shah, E.J. Sontheimer, K.L. Maxwell, A.R. Davidson, Naturally Occurring Off-Switches for CRISPR-Cas9, Cell. 167 (2016) 1829–1838.e9.

[44]   J. Shin, F. Jiang, J.-J. Liu, N.L. Bray, B.J. Rauch, S.H. Baik, E. Nogales, J. Bondy-Denomy, J.E. Corn, J.A. Doudna, Disabling Cas9 by an anti-CRISPR DNA mimic, Sci Adv. 3 (2017) e1701620.

[45]   J. Lee, H. Mou, R. Ibraheim, S.-Q. Liang, P. Liu, W. Xue, E.J. Sontheimer, Tissue-restricted genome editing in vivo specified by microRNA-repressible anti-CRISPR proteins, RNA. 25 (2019) 1421–1431.

[46]   L.B. Harrington, K.W. Doxzen, E. Ma, J.-J. Liu, G.J. Knott, A. Edraki, B. Garcia, N. Amrani, J.S. Chen, J.C. Cofsky, P.J. Kranzusch, E.J. Sontheimer, A.R. Davidson, K.L. Maxwell, J.A. Doudna, A Broad-Spectrum Inhibitor of CRISPR-Cas9, Cell. 170 (2017) 1224–1233.e15.

[47]   M. Liang, T. Sui, Z. Liu, M. Chen, H. Liu, H. Shan, L. Lai, Z. Li, AcrIIA5 Suppresses Base Editors and Reduces Their Off-Target Effects, Cells. 9 (2020). https://doi.org/10.3390/cells9081786.

[48]   M.D. Hoffmann, S. Aschenbrenner, S. Grosse, K. Rapti, C. Domenger, J. Fakhiri, M. Mastel, K. Börner, R. Eils, D. Grimm, D. Niopek, Cell-specific CRISPR–Cas9 activation by microRNA-dependent expression of anti-CRISPR proteins, Nucleic Acids Res. 47 (2019) e75–e75.

[49]   T. Bienvenu, A. Carrié, N. de Roux, M.C. Vinet, P. Jonveaux, P. Couvert, L. Villard, A. Arzimanoglou, C. Beldjord, M. Fontes, M. Tardieu, J. Chelly, MECP2 mutations account for most cases of typical forms of Rett syndrome, Hum. Mol. Genet. 9 (2000) 1377–1384.

[50]   M.J. Lyst, A. Bird, Rett syndrome: a complex disorder with simple roots, Nat. Rev. Genet. 16 (2015) 261–275.

[51]   R. Krishnaraj, G. Ho, J. Christodoulou, RettBASE: Rett syndrome database update, Hum. Mutat. 38 (2017) 922–931.

[52]   G. Stefanelli, A. Gandaglia, M. Costa, M.S. Cheema, D. Di Marino, I. Barbiero, C. Kilstrup-Nielsen, J. Ausió, N. Landsberger, Brain phosphorylation of MeCP2 at serine 164 is developmentally regulated and globally alters its chromatin association, Sci. Rep. 6 (2016) 28295.

[53]   J. Tan, F. Zhang, D. Karcher, R. Bock, Engineering of high-precision base editors for site-specific single nucleotide replacement, Nature Communications. 10 (2019). https://doi.org/10.1038/s41467-018-08034-8.

[54]   S.H. Chu, M. Packer, H. Rees, D. Lam, Y. Yu, J. Marshall, L.-I. Cheng, D. Lam, J. Olins, F.A. Ran, A. Liquori, B. Gantzer, J. Decker, D. Born, L. Barrera, A. Hartigan, N.

Gaudelli, G. Ciaramella, I.M. Slaymaker, Rationally Designed Base Editors for Precise Editing
of the Sickle Cell Disease Mutation, CRISPR J. 4 (2021) 169–177.

[55]   T.P. Huang, K.T. Zhao, S.M. Miller, N.M. Gaudelli, B.L. Oakes, C. Fellmann, D.F.
Savage, D.R. Liu, Circularly permuted and PAM-modified Cas9 variants broaden the targeting
scope of base editors, Nat. Biotechnol. 37 (2019) 626–631.

[56]   Y. Wang, L. Zhou, N. Liu, S. Yao, BE-PIGS: a base-editing tool with deaminases inlaid
into Cas9 PI domain significantly expanded the editing scope, Signal Transduct Target Ther. 4
(2019) 36.

[57]   S. Li, B. Yuan, J. Cao, J. Chen, J. Chen, J. Qiu, X.-M. Zhao, X. Wang, Z. Qiu, T.-L.
Cheng, Docking sites inside Cas9 for adenine base editing diversification and RNA off-target
elimination, Nat. Commun. 11 (2020) 5827.

[58]   Y. Liu, C. Zhou, S. Huang, L. Dang, Y. Wei, J. He, Y. Zhou, S. Mao, W. Tao, Y.
Zhang, H. Yang, X. Huang, T. Chi, A Cas-embedding strategy for minimizing off-target effects
of DNA base editors, Nat. Commun. 11 (2020) 6073.

[59]   F. Chemello, Z. Wang, H. Li, J.R. McAnally, N. Liu, R. Bassel-Duby, E.N. Olson,
Degenerative and regenerative pathways underlying Duchenne muscular dystrophy revealed by
single-nucleus RNA sequencing, Proc. Natl. Acad. Sci. U. S. A. 117 (2020) 29691–29701.

[60]   K.M. Anderson, P. Poosala, S.R. Lindley, D.M. Anderson, Targeted Cleavage and
Polyadenylation of RNA by CRISPR-Cas13, bioRxiv. (2019) 531111.
https://doi.org/10.1101/531111.

[61]   L.W. Koblan, J.L. Doman, C. Wilson, J.M. Levy, T. Tay, G.A. Newby, J.P. Maianti, A.
Raguram, D.R. Liu, Improving cytidine and adenine base editors by expression optimization
and ancestral reconstruction, Nat. Biotechnol. 36 (2018) 843–846.

[62]   R. Preece, C. Georgiadis, S.A. Gkazi, A. Etuk, A. Christi, W. Qasim, "Mini" U6 Pol III
promoter exhibits nucleosome redundancy and supports multiplexed coupling of CRISPR/Cas9
effects, Gene Ther. 27 (2020) 451–458.

[63]   J.L. Aponte, G.A. Sega, L.J. Hauser, M.S. Dhar, C.M. Withrow, D.A. Carpenter, E.M.
Rinchik, C.T. Culiat, D.K. Johnson, Point mutations in the murine fumarylacetoacetate
hydrolase gene: Animal models for the human genetic disorder hereditary tyrosinemia type 1,
Proc. Natl. Acad. Sci. U. S. A. 98 (2001) 641–645.

[64]   S. Lindstedt, Treatment of hereditary tyrosinaemia type I by inhibition of 4-
hydroxyphenylpyruvate dioxygenase, The Lancet. 340 (1992) 813–817.
https://doi.org/10.1016/0140-6736(92)92685-9.

[65]   H. Yin, W. Xue, S. Chen, R.L. Bogorad, E. Benedetti, M. Grompe, V. Koteliansky, P.A.
Sharp, T. Jacks, D.G. Anderson, Genome editing with Cas9 in adult mice corrects a disease
mutation and phenotype, Nat. Biotechnol. 32 (2014) 551–553.

[66]   H. Yin, C.-Q. Song, J.R. Dorkin, L.J. Zhu, Y. Li, Q. Wu, A. Park, J. Yang, S. Suresh, A.
Bizhanova, A. Gupta, M.F. Bolukbasi, S. Walsh, R.L. Bogorad, G. Gao, Z. Weng, Y. Dong, V.
Koteliansky, S.A. Wolfe, R. Langer, W. Xue, D.G. Anderson, Therapeutic genome editing by
combined viral and non-viral delivery of CRISPR system components in vivo, Nat. Biotechnol.
34 (2016) 328–333.

[67]   C.-Q. Song, T. Jiang, M. Richter, L.H. Rhym, L.W. Koblan, M.P. Zafra, E.M. Schatoff,
J.L. Doman, Y. Cao, L.E. Dow, L.J. Zhu, D.G. Anderson, D.R. Liu, H. Yin, W. Xue, Adenine
base editing in an adult mouse model of tyrosinaemia, Nat Biomed Eng. 4 (2020) 125–130.

[68]   T. Jiang, J.M. Henderson, K. Coote, Y. Cheng, H.C. Valley, X.-O. Zhang, Q. Wang,
L.H. Rhym, Y. Cao, G.A. Newby, H. Bihler, M. Mense, Z. Weng, D.G. Anderson, A.P.
McCaffrey, D.R. Liu, W. Xue, Chemical modifications of adenine base editor mRNA and guide

RNA expand its application scope, Nat. Commun. 11 (2020) 1979.

[69]   J.H. Shin, S. Jung, S. Ramakrishna, H.H. Kim, J. Lee, In vivo gene correction with targeted sequence substitution through microhomology-mediated end joining, Biochem. Biophys. Res. Commun. 502 (2018) 116–122.

[70]   H. Jang, D.H. Jo, C.S. Cho, J.H. Shin, J.H. Seo, G. Yu, R. Gopalappa, D. Kim, S.-R. Cho, J.H. Kim, H.H. Kim, Application of prime editing to the correction of mutations and phenotypes in adult mice with liver and eye diseases, Nat Biomed Eng. (2021). https://doi.org/10.1038/s41551-021-00788-9.

[71]   G. Zhang, V. Budker, J.A. Wolff, High levels of foreign gene expression in hepatocytes after tail vein injections of naked plasmid DNA, Hum. Gene Ther. 10 (1999) 1735–1737.

[72]   N.K. Paulk, K. Wursthorn, Z. Wang, M.J. Finegold, M.A. Kay, M. Grompe, Adeno-associated virus gene repair corrects a mouse model of hereditary tyrosinemia in vivo, Hepatology. 51 (2010) 1200–1208. https://doi.org/10.1002/hep.23481.

[73]   M.-J. Wang, F. Chen, J.T.Y. Lau, Y.-P. Hu, Hepatocyte polyploidization and its association with pathophysiological processes, Cell Death Dis. 8 (2017) e2805.

[74]   S. Bae, J. Park, J.-S. Kim, Cas-OFFinder: a fast and versatile algorithm that searches for potential off-target sites of Cas9 RNA-guided endonucleases, Bioinformatics. 30 (2014) 1473–1475.

[75]   A.V. Anzalone, L.W. Koblan, D.R. Liu, Genome editing with CRISPR-Cas nucleases, base editors, transposases and prime editors, Nat. Biotechnol. 38 (2020) 824–844.

[76]   G.A. Newby, D.R. Liu, In vivo somatic cell base editing and prime editing, Mol. Ther. (2021). https://doi.org/10.1016/j.ymthe.2021.09.002.

[77]   High-dose AAV gene therapy deaths, Nat. Biotechnol. 38 (2020) 910.

[78]   L. Morales, Y. Gambhir, J. Bennett, H.H. Stedman, Broader Implications of Progressive Liver Dysfunction and Lethal Sepsis in Two Boys following Systemic High-Dose AAV, Mol. Ther. 28 (2020) 1753–1755.

[79]   T.R. Flotte, Revisiting the "New" Inflammatory Toxicities of Adeno-Associated Virus Vectors, Hum. Gene Ther. 31 (2020) 398–399.

[80]   D. Wang, P.W.L. Tai, G. Gao, Adeno-associated virus vector as a platform for gene therapy delivery, Nat. Rev. Drug Discov. 18 (2019) 358–378.

[81]   D. Wang, F. Zhang, G. Gao, CRISPR-Based Therapeutic Genome Editing: Strategies and In Vivo Delivery by AAV Vectors, Cell. 181 (2020) 136–150.

[82]   R. Ibraheim, C.-Q. Song, A. Mir, N. Amrani, W. Xue, E.J. Sontheimer, All-in-one adeno-associated virus delivery and genome editing by Neisseria meningitidis Cas9 in vivo, Genome Biol. 19 (2018) 137.

[83]   S.A. Krooss, Z. Dai, F. Schmidt, A. Rovai, J. Fakhiri, A. Dhingra, Q. Yuan, T. Yang, A. Balakrishnan, L. Steinbrück, S. Srivaratharajan, M.P. Manns, A. Schambach, D. Grimm, J. Bohne, A.D. Sharma, H. Büning, M. Ott, Ex Vivo/In vivo Gene Editing in Hepatocytes Using "All-in-One" CRISPR-Adeno-Associated Virus Vectors with a Self-Linearizing Repair Template, iScience. 23 (2020) 100764. https://doi.org/10.1016/j.isci.2019.100764.

[84]   W. Sun, J. Yang, Z. Cheng, N. Amrani, C. Liu, K. Wang, R. Ibraheim, A. Edraki, X. Huang, M. Wang, J. Wang, L. Liu, G. Sheng, Y. Yang, J. Lou, E.J. Sontheimer, Y. Wang, Structures of Neisseria meningitidis Cas9 Complexes in Catalytically Poised and Anti-CRISPR-Inhibited States, Mol. Cell. 76 (2019) 938–952.e5.

[85]   L.E. Fry, C.F. Peddle, M. Stevanovic, A.R. Barnard, M.E. McClements, R.E. MacLaren, Promoter Orientation within an AAV-CRISPR Vector Affects Cas9 Expression and Gene Editing Efficiency, CRISPR J. 3 (2020) 276–283.

884   [86]   M. Sena-Esteves, G. Gao, Introducing Genes into Mammalian Cells: Viral Vectors, Cold
885   Spring Harb. Protoc. 2020 (2020) 095513.
886   [87]   K. Clement, H. Rees, M.C. Canver, J.M. Gehrke, R. Farouni, J.Y. Hsu, M.A. Cole, D.R.
887   Liu, J.K. Joung, D.E. Bauer, L. Pinello, CRISPResso2 provides accurate and rapid genome
888   editing sequence analysis, Nat. Biotechnol. 37 (2019) 224–226.

Figures

Figure 1
a). Schematic representation of the ABE reporter HEK293T cell line. **b).** Schematic representation of the Nme2-ABE constructs. **c).** Comparison of editing efficiencies of Nme2-ABEs to those of Spy-ABEs in the ABE reporter cell line by plasmid transient transfection (n = 3 biological replicates). **d).** Summary of editing windows and comparison of editing efficiencies for the ABEs at endogenous genomic loci. Each data point represents the mean A-to-G editing efficiency at the indicated position of the spacer across 12 Nme2Cas9 target sites and 8 SpyCas9 target sites, respectively (n = 3 biological replicates). Summary of the individual A-to-G conversion and indel efficiencies at each target site can be found in **Supplementary Figure 2**. **e).** Comparison of Nme2-ABE8e mismatch tolerance to that of Spy-ABE8e in the ABE reporter cell line. The activities of the effectors with the mismatched guides are normalized to that of the perfectly complementary (WT) guide. Red, mismatched nucleotides; green, PAM sequence (n = 3 biological replicates).

Figure 2

a). Schematic representation of a nonsense mutation in the human *MeCP2* gene (c.502 C>T; p.R168X) that causes Rett Syndrome. The black underline denotes the target sequence of an Nme2-ABE8e guide for reverting the mutant A to G (wildtype) at position 10 (red, bold). The PAM region is underlined in green. A bystander edit at position 16 (orange) can generate a missense mutation (c.496 T>C; p.S166P). **b).** Amplicon deep sequencing quantifying the editing efficiency in Rett patient-derived fibroblasts transfected with the Nme2-ABE8e mRNA and the synthetic sgRNA_*MeCP2* noted in **a** (n = 3 biological replicates). **c).** Schematic representation of the exon skipping strategy that restores the reading frame of the mouse *Dmd* transcript. Deletion of exon 51 (ΔEx51) can alter the reading frame and generate a premature stop codon in exon 52 (red). Adenine base editing at the splice site of exon 50 (red) by Nme2-ABE8e can cause exon 50 skipping (gray) and restore the *Dmd* reading frame. The PAM region is underlined in green. **d).** Amplicon deep sequencing quantifies the editing efficiency at the target site in mouse N2a cells transfected with the Nme2-ABE8e and sgRNA_*Dmd* expression plasmid (n = 3 biological replicates).

Figure 3

a). Schematic representation of single AAV constructs with different NLS configurations. **b).** Comparison of different NLS configurations by plasmid transfection in the ABE reporter cell line. **c).** Comparison of Nme2-ABE activity when sgRNA expression is driven by the U6 or miniU6 promoters in the 2xBP_SV40 NLS construct targeting the ABE reporter site (left) or endogenous human (middle) and mouse (right) genomic sites by plasmid transfection in cultured cells followed by amplicon deep sequencing (n = 3 biological replicates). Data represent mean ± SD; ns, P > 0.05; *, P < 0.05; **, P < 0.01; ***, P < 0.001; ****, P < 0.0001 (one-way ANOVA).

a

7 | 8 | 9

A

1 2   5   10   13   1617
AACGACTGGAGCAGTAATGCCTGG**TGGCCC**
                                    PAM

Nme2-ABE8e +
sgRNA_*Fah*
↓

7 | 8 | 9
       G

b

Nme2-ABE8e-U6

sgRNA — U6 — U1a — BP_SV40-NLS — Nme2-ABE8e — BP_SV40-NLS — polyA

Nme2-ABE8e-miniU6

sgRNA — miniU6 — U1a — BP_SV40-NLS — Nme2-ABE8e — BP_SV40-NLS — polyA

c

% A to G conversion

● Nme2Cas9-ABE8e-U6
● Nme2Cas9-ABE8e-miniU6
● negative control

A1G ns ns | A2G ns ns | A5G ns ns | A10G ** ns | A13G *** ** | A16G *** ** | A17G * | Indel ns ns

d

PBS | Spy-RA6.3
Nme2-ABE8e-U6 | Nme2-ABE8e-miniU6

% FAH⁺ hepatocytes

PBS | Sp-RA6.3 | Nme2-ABE8e-U6 | Nme2-ABE8e-miniU6

e

Body weight ratio

● Nme2-ABE8e-U6
■ Nme2-ABE8e-miniU6
▲ PBS

Days since NTBC withdrawal

f

Nme2-ABE8e-U6 | Nme2-ABE8e-miniU6 | PBS

400bp
300bp

400bp | 5 | 6 | 7 | 8 | 9
300bp | 5 | 6 | 7 | 9

g

A10 G13

CTGGAGCGCACGAGACATCC
Exon 8 | Exon 9

h

PBS | Nme2-ABE8e-U6 | Nme2-ABE8e-miniU6

IHC

H&E

i

% Edited

● Nme2-ABE8e-U6
● Nme2-ABE8e-miniU6
● PBS

A1G ns ns | A2G ns ns | A5G ns ns | A10G ns ns | A13G *** *** | A16G * ** | A17G * * | Indel ns ns

Figure 4

a). Illustration of the pathogenic point mutation in the *Fah*$^{PM/PM}$ mouse model that causes exon 8 skipping of the *Fah* gene, and the guide design for Nme2-ABE8e to correct the point mutation. Red and bold, target adenine; orange, other bystander adenines; green and underlined, PAM. **b).** Illustration of constructs of the single-AAV vector plasmids used in *in vivo* studies. **c).** Editing efficiencies at the *Fah* mutant site by AAV plasmid electroporation in MEF cells derived from the *Fah*$^{PM/PM}$ mouse. Data are from amplicon deep sequencing (n = 2 biological replicates). **d).** Anti-FAH immunohistochemistry (IHC) staining showing FAH$^+$ hepatocytes, before NTBC withdrawal, in the *Fah*$^{PM/PM}$ mouse hydrodynamically injected with the indicated plasmid. The bar graph quantifies the percentage of FAH$^+$ hepatocytes detected by IHC (n = 2 mice per group). Scale bars, 500 µm. **e).** Body weight plot of mice injected with the single-AAV vector plasmid showing gradual weight gain over a month after NTBC withdrawal. **f).** RT–PCR analysis of the plasmid- or PBS-injected mouse livers using primers that span exons 5 and 9. The wild-type amplicon is 405 bp and exon 8 skipped amplicon is 305 bp. **g).** Representative Sanger sequencing trace of the 405 bp RT-PCR band. **h).** Anti-FAH IHC staining showing expansion of FAH$^+$ hepatocytes 40 days post NTBC withdrawal. Scale bars, 500 µm. **i).** Quantification of the editing efficiency by amplicon deep sequencing of genomic DNA of the treated mouse livers harvested 40 days post NTBC withdrawal. **d-i**, n = 2 mice per group). Data represent mean ± SD; ns, $P > 0.05$; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ (one-way ANOVA).

Figure 5

a). Schematic representation of the AAV constructs. **b).** Anti-FAH immunohistochemistry (IHC) staining showing FAH+ hepatocytes, before NTBC withdrawal, in the $Fah^{PM/PM}$ mouse injected with AAV9 expressing Nme2-ABE8e with a sgRNA targeting either the Fah gene, or the $Rosa26$ gene that serves as a negative control. Scale bar, 500 μm. The bar graph quantifies the percentage of FAH+ hepatocytes detected by IHC (n = 4 mice in the Fah targeting group, n = 3 mice in the $Rosa26$ targeting group). **c, d).** Quantification of the editing efficiency by amplicon deep sequencing of genomic DNA from the AAV9-injected mouse livers harvested before NTBC withdrawal (n = 4 mice in the Fah targeting group, n = 3 mice in the $Rosa26$ targeting group and the PBS control group). Data represent mean ± SD; ns, $P > 0.05$; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ (one-way ANOVA).

Figure 6

a). Schematic representation of the original (left) and the optimized AAV constructs (AAV9-Nm2-ABE8e-U6_V2, right). **b).** Quantification of the editing efficiency at the *Rosa26* locus by amplicon deep sequencing using mouse livers injected with indicated AAV (n = 3 mice in the AAV injected group, n = 1 mouse in the PBS injected group). Data from the original AAV configuration (**a**, left) is the same as that from the experiment shown in **Figure 5d**. Data represent mean ± SD, ns, P > 0.05; *, P<0.1; **, P < 0.01(one-way ANOVA).

Supplementary Figure 1
Representative flow cytometry gating strategy for the ABE reporter cell line.

Supplementary Figure 2

Summary of the individual A-to-G conversion efficiency and the indel efficiency (bottom) at 12 target sites for Nme2-ABE8e, which includes 8 dual-target sites (DS 2 - 28) and 4 Nme2Cas9-specific target sites (Nme2 1 - 4), and 8 dual-target sites for Spy-ABE7.10 and Spy-ABE8e. Each data point represents the A-to-G conversion efficiency at the indicated nucleotide position measured by amplicon deep sequencing (n = 3 biological replicates).

Supplementary Figure 3
a). Anti-CRISPR proteins inhibit the activity of either Spy-ABE7.10 or Nme2-ABE8e in the ABE reporter cell line as a function of their previously defined nuclease specificity (n = 2 biological replicates). **b).** Editing efficiency at the A12 (the highest edited adenine within the target) of the DS12 site by Nme2-ABE8e controlled by microRNA-repressible anti-CRIPSR proteins. Data generated by Sanger sequencing and EditR analysis (n = 2 biological replicates). Data represent mean ± SD; ns, P > 0.05; *, P < 0.05; **, P < 0.01; ****, P < 0.0001 (one-way ANOVA).

Supplementary Figure 4
Alkaline gel electrophoresis of AAV9 genomic DNA.

1. AAV9-Nme2-ABE8e-miniU6-*Fah*
2. AAV9-Nme2-ABE8e-U6-*Fah*
3. AAV9-Nme2-ABE8e-miniU6-*Rosa26*
4. AAV9-Nme2-ABE8e-U6-*Rosa26*

a

Fah on-target site AACGACTGGAGCAGTAATGCCTGG<u>TGGCCC</u>
OT1 GGCGGCTGGAGCAGGATTGCCTGG<u>AGTTCC</u>
OT2 CAGGATTGGAACAGTAATGACTGG<u>CAAACC</u>

b

OT1

Un-injected *Fah*^{PM/PM} mouse

AAV9 injected *Fah*^{PM/PM} mouse

OT2

Un-injected *Fah*^{PM/PM} mouse

AAV9 injected *Fah*^{PM/PM} mouse

Supplementary Figure 5
a). Sequence of the *Fah* on-target site and two top-rated Cas-OFFinder predicted off-target sites for Nme2-ABE8e. Bases that are different from the on-target site are labeled in red. PAM, green, underlined. **b).** Representative amplicon deep sequencing reads at the predicted off-target sites in mice injected with AAV9 expressing Nme2-ABE8e and sgRNA-*Fah*.

Supplementary note
Nucleotide sequences of plasmids

Nme2-ABE7.10 construct in **Figure 1**
Legend: CMV enhancer, CMV promoter, NLS, **TadA**, linker, TadA7.10, Nme2Cas9-D16A, 3xHA, poly-A signal
GACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTTA
CATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCA
TAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAA
GTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGACC
TTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCA
ATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTTTTGGCAC
CAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGG
AGGTCTATATAAGCAGAGCTGGTTTAGTGAACCGTCAGATCCGCTAGAGATCCGCGGCCGCtaatacgactcactatagggcgaa
ttgggtaccgggcccGCCACCATGgctagcCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCAGCGGCGG
CGGCAGCGGCCCGGCAGCTAAGAGGGTGAAACTGGATggcggtagcggcggaggcagcggtggcggcagcggccccctcgagccaaaga
agaagcggaaagtctctgaagtcgagtttagccacgagtattggatgaggcacgcactgaccctggcaaagcgagcatgggatgaaagagaagtccccgt
gggcgccgtgctggtgcacaacaatagagtgatcggagagggatggaacaggccaatcggccgccacgaccctaccgcacacgcagagatcatggcactg
aggcagggaggcctggtcatgcagaattaccgcctgatcgatgccaccctgtatgtgacactggagccatgcgtgatgtgcgcaggagcaatgatccacagc
aggatcggaagagtggtgttcggagcacgggacgccaagaccggcgcagcaggctccctgatggatgtgctgcaccaccccggcatgaaccaccgggtgg
agatcacagaggaatcctggcagacgagtgcgccgccctgctgagcgatttctttagaatgcggagacaggagatcaaggcccagaagaaggcacagag
ctccaccgatctctggaggatctagcggaggatcctctggaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggagg
atcctctgaggtggagttttcccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggaggtgcctgtgtgggagccgtgctg
gtgctgaacaatagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggccctgagacagggcggcctg
gtcatgcagaactacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtg
tttggcgtgaggaacgcaaaaaccggcgccgcaggctccctgatggacgtgctgcactaccccggcatgaatcaccgcgtcgaaattaccgagggaatcctgg
cagatgaatgtgccgccctgctgtgctatttctttcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaccgactccggaggatctagcg
gaggctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagcggggggtcaatggccgccttcaagcctaac
ccaatcaattacatcctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgaggaggagaatcctatccggctgatcgatctggg
gcgtgagagtgtttgagagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacggagcgtgaggcgcctgacacggagaagg
gcacacaggctgctgagggcacgccggctgctgaagagagagggcgtgctgcaggcagcagacttcgatgagaatggcctgatcaagagcctgccaaacac
cccctggcagctgagagcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacctgatcaagcaccgcggctacctgagccag
cggaagaacgagggagagacagcagacaaggagctgggcgccctgctgaagggagtggccaacaatgcccacgccctgcagaccggcgatttcaggacac
ctgccgagctggccctgaataagtttgagaaggagtccggccacatcagaaaccagagggggcgactatagccacacctctcccgcaaggatctgcaggccg
agctgatcctgctgttcgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggaggggaatcgagaccctgctgatgacacagaggcctg
ccctgtccggcgacgcagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaagaacacctacacagccgagcggtttatct
ggctgacaaagctgaacaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagagggccaccctgatggatgagccttaccggaag
tctaagctgacatatgcccaggccagaaagctgctgggcctggaggacaccgccttctttaagggcctgagatacggcaaggataatgccgaggcctccacac
tgatggagatgaaggcctatcacgccatctctcgcgccctggagaaggaggggcctgaaggacaagaagtccccctgaacctgagctccgagctgcaggatg
agatcggcaccgccttctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacaggtgcagcctgagatcctggaggccctgctgaagcacat
ctctttcgataagtttgtgcagatcagcctgaaggccctgagaaggatcgtgccactgatggagcagggcaagcggtacgacgaggcctgcgccgagatctac
ggcgatcactatggcaagaagaacacagaggagaagatctatctgcccccctatccctgccgacgagatcagaaatcctgtggtgctgagggccctgtcccagg
caagaaaagtgatcaacgagtggtgcgccggtacggatctccagcccggatccacatcgagaccgccagagaagtgggcaagagcttcaaggaccggaag
gagatcgagaagagacaggaggagaatcgcaaggatcggggagaaggccgccgccaagtttaggggagtacttccctaactttgtgggcgagccaaagtctaa
ggacatcctgaagctgcgcctgtacgagcagcagcacggccaagtgtctgtatagcggcaaggagatcaatctggtgcggctgaacgagaagggctatgtgga
gatcgatcacgccctgccttctccagaacctgggacgattcttttaacaataaggtgctggtgctgggcagcgagaaccagaataagggcaatcagacaccat
acgagtatttcaatggcaaggacaactccaggggagtggcaggagttcaaggcccgcgtggagacctctagatttcccaggagcaagaagcagcggatcctgc
tgcagaagttcgacgaggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgccagtttgtggccgatcacatcctgctgacc

ggcaagggcaagagaagggtgttcgcctctaatggccagatcacaaacctgctgaggggattttggggactgaggaaggtgcgggcagagaatgacagaca
ccacgcactggatgcagtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgaggtataaggagatgaacgcctttgacggcaa
gaccatcgataaggagacaggcaaggtgctgcaccagaagacccacttcccccagccttgggagttctttgcccaggaagtgatgatccgggtgttcggcaag
ccagacggcaagcctgagtttgaggaggccgataccccagagaagctgaggacactgctggcagagaagctgtctagcaggccagaggcagtgcacgagta
cgtgaccccactgttcgtgtccagggcacccaatcggaagatgtctggcgcccacaaggacacactgagaagcgccaagagggtttgtgaagcacaacgagaa
gatctccgtgaagagagtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaacggcagggagatcgagctgtatgaggccct
gaaggcaaggctggaggcctacggaggaaatgccaagcaggccttcgacccaaaggataaccccttttataagaagggaggacagctggtgaaggccgtgc
gggtggagaagacccaggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcgatatggtgagagtggacgtgttctgtaag
gtggataagaagggcaagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatcgattgcaagggctacagaatcg
acgatagctatacattcgtgtttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtggagtttgcctactatatcaattgcgactc
ctctaacggcaggttctacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaatctggtgctgatccagaagtatcaggtga
acgagctgggcaaggagatcaggccatgtcggctgaagaagcgcccacccgtgcgggatatcGGCACCGGCGGGtctggaggatctagcggaggatcc
tctggaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggaggatccGTATACCCATACGATGTTCCTGAC
TATGCGGGCTATCCCTATGACGTCCCGGACTATGCAGGATCGTATCCTTATGACGTTCCAGATTACGCTGGATCCGCCGCT
CCGGCAGCTAAGAAAAAGAAACTGGATTTCGAATCCGGAgaattcctgcagcccggggggatccactagttctagaGGATCCGCCGCTC
CCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCGGCAGCGGCGGCGGCGGCAGCGGCGGCGGCAGCGGCggcggtagcggcggaggca
gcggtggcggcagcggcAAGCGCACCGCCGACAGCCAGCACAGCACCCCCCCCAAGACCAAGCGCAAGGTGGGCGGCAGCGG
CGGCGGCAGCGGCGGCGGCAGCGGCTCCGGACCGGCAGCTAAGAGGGTGAAACTGGATtgaTAAgcggccgccaccgcggtg
gagctccagcttttgttccctttagtgagggttaatGAATTCGAGCCCAAGAAGAAGAGGAAAGTCTAACCGGTCATCATCACCATCAC
CATTGAGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCTCCCCCGTGCCTT
CCTTGACCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGAAAATTGCATCGCATTGTCTGAGTAGGTGTC
ATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGAT
GCGGTGGGCTCTATGG

Nme2-ABE8e construct in **Figure 1 and Supplementary Figure 3a**
Legend: CMV enhancer, CMV promoter, NLS, TadA8e, linker, Nme2Cas9-D16A, 3xHA, poly-A signal
GACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTTA
CATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCA
TAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAA
GTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGACC
TTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCA
ATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTTTTGGCAC
CAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGG
AGGTCTATATAAGCAGAGCTGGTTTAGTGAACCGTCAGATCCGCTAGAGATCCGCGGCCGCtaatacgactcactatagggcgaa
ttgggtaccgggcccGCCACCATGgctagcCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCAGCGGCGG
CGGCAGCGGCCCGGCAGCTAAGAGGGTGAAACTGGATggcggtagcggcggaggcagcggtggcggcagcggccccctcgagTctgagg
tggagttttcccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggaggtgcctgtgggagccgtgctggtgctgaacaa
tagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggccctgagacagggcggcctggtcatgcagaa
ctacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtgtttggcgtgag
gaacAGCaaaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcaccgcgtcgaaattaccgagggaatcctggcagatga
atgtgccgccctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaTcAactccggaggatctagcggagg
ctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagcgggggggtcaatggccgccttcaagcctaacccaat
caattacatcctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgaggaggagaatcctatccggctgatcgatctgggcgtg
agagtgtttgagagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacggagcgtgaggcgcctgacacggagaagggcaca
caggctgctgagggcacgccggctgctgaagagagagaggcgtgctgcaggcagcagacttcgatgagaatggcctgatcaagagcctgccaaacaccccctg
gcagctgagagcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacctgatcaagcaccgcggctacctgagccagcggaa
gaacgagggagagacagcagacaaggagctgggcgccctgctgaagggagtggccaacaatgcccacgccctgcagaccggcgatttcaggacacctgccg
agctggccctgaataagtttgagaaggagtccggccacatcagaaaccagagggggcgactatagccacaccttctcccgcaaggatctgcaggccgagctga
tcctgctgttcgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggaggggaatcgagaccctgctgatgacacagaggcctgccctgt

ccggcgacgcagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaagaacacctacacagccgagcggtttatctggctga
caaagctgaacaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagagggccaccctgatggatgagccttaccggaagtctaag
ctgacatatgcccaggccagaaagctgctgggcctggaggacaccgccttctttaagggcctgagatacggcaaggataatgccgaggcctccacactgatgg
agatgaaggcctatcacgccatctctcgcgccctggagaaggagggcctgaaggacaagaagtcccccctgaacctgagctccgagctgcaggatgagatcg
gcaccgccttctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacagggtgcagcctgagatcctggaggccctgctgaagcacatctcttttc
gataagtttgtgcagatcagcctgaaggccctgagaaggatcgtgccactgatggagcagggcaagcggtacgacgaggcctgcgccgagatctacggcgat
cactatggcaagaagaacacagaggagaagatctatctgccccctatccctgccgacgagatcagaaatcctgtggtgctgagggccctgtcccaggcaaga
aaagtgatcaacggagtggtgcgccggtacggatctccagcccggatccacatcgagaccccagagaagtgggcaagagcttcaaggaccggaaggagat
cgagaagagacaggaggagaatcgcaaggatcgggagaaggccgccgccaagtttagggagtacttccctaactttgtgggcgagccaaagtctaaggaca
tcctgaagctgcgcctgtacgagcagcagcacggcaagtgtctgtatagcggcaaggagatcaatctggtgcggctgaacgagaagggctatgtggagatcg
atcacgccctgcctttctccagaacctgggacgattctttaacaataaggtgctggtgctgggcagcgagaaccagaataagggcaatcagacaccatacga
gtatttcaatggcaaggacaactccagggagtggcaggagttcaaggcccgcgtggagacctctagatttcccaggagcaagaagcagcggatcctgctgca
gaagttcgacgaggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgccagtttgtggccgatcacatcctgctgaccggca
aagggcaagagaagggtgttcgcctctaatggccagatcacaaacctgctgagggggattttggggactgaggaaggtgcgggcagagaatgacagacaccac
gcactggatgcagtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgaggtataaggagatgaacgccctttgacggcaagacc
atcgataaggagacaggcaaggtgctgcaccagaagacccacttccccccagccttgggagttctttgcccaggaagtgatgatcccgggtgttcggcaagccag
acggcaagcctgagtttgaggaggccgataccccagagaagctgaggacactgctggcagagaagctgtctagcaggccagaggcagtgcacgagtacgtg
accccactgttcgtgtccagggcacccaatcggaagatgtctggcgccccacaaggacacactgagaagcgccaagaggtttgtgaagcacaacgagaagatc
tccgtgaagagagtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaacggcagggagatcgagctgtatgaggccctgaag
gcaaggctggaggcctacggaggaaatgccaagcaggccttcgacccaaaggataaccccttttataagaagggaggacagctggtgaaggccgtgcgggt
ggagaagacccaggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcgatatggtgagagtggacgtgttctgtaaggtgg
ataagaagggcaagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatcgattgcaagggctacagaatcgacga
tagctatacattctgttttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtggagtttgcctactatatcaattgcgactcctcta
acggcaggttctacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaatctggtgctgatccagaagtatcaggtgaacga
gctgggcaaggagatcaggccatgtcggctgaagaagcgcccacccgtgcgggt gatatcGGCACCGGCGGGtctggaggatctagcggaggatcctctg
gaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggaggatccGTATACCCATACGATGTTCCTGACTAT
GCGGGCTATCCCTATGACGTCCCGGACTATGCAGGATCGTATCCTTATGACGTTCCAGATTACGCTGGATCCGCCGCTCCG
GCAGCTAAGAAAAAGAAACTGGATTTCGAATCCGGAgaattcctgcagcccgggggatccactagttctagaGGATCCGCCGCTCCCG
CCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCGGCAGCGGCGGCGGCAGCGGCGgcggtagcggcggaggcagcg
gtggcggcagcggcAAGCGCACCGCCGACAGCCAGCACAGCACCCCCCCCCAAGACCAAGCGCAAGGTGGGCGGCAGCGGCG
GCGGCAGCGGCGGCGGCAGCGGCTCCGGACCGGCAGCTAAGAGGGTGAAACTGGATtgaTAAgcggccgccaccgcggtggag
ctccagcttttgttccctttagtgagggttaatGAATTCGAGCCCAAGAAGAAGAGGAAAGTCTAACCGGTCATCATCACCATCACCAT
TGAGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCTCCCCCGTGCCTTCCT
TGACCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGAAAATTGCATCGCATTGTCTGAGTAGGTGTCATT
CTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGATGCG
GTGGGCTCTATGG

Spy-ABE7.10 construct in **Figure 1 and Supplementary Figure 3a**
Legend: CMV enhancer, CMV promoter, NLS, **TadA**, linker, TadA7.10, SpyCas9-D10A, 3xHA, poly-A signal
GACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTTA
CATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCA
TAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAA
GTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGACC
TTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCA
ATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTTTTGGCAC
CAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGG
AGGTCTATATAAGCAGAGCTGGTTTAGTGAACCGTCAGATCCGCTAGAGATCCGCGGCCGCtaatacgactcactatagggcgaa
ttgggtaccgggcccGCCACCATGGctagcCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCGGCAGCGGCGG
CGGCAGCGGCCCGGCAGCTAAGAGGGTGAAACTGGATggcggtagcggcggaggcagcggtggcggcagcggcccccctcgagccaaaga

agaagcggaaagtctctgaagtcgagtttagccacgagtattggatgaggcacgcactgaccctggcaaagcgagcatgggatgaaagagaagtccccgt
gggcgccgtgctggtgcacaacaatagagtgatcggagagggatggaacaggccaatcggccgccacgcaccctaccgcacacgcagagatcatggcactg
aggcagggaggcctggtcatgcagaattaccgcctgatcgatgccaccctgtatgtgacactggagccatgcgtgatgtgcgcaggagcaatgatccacagc
aggatcggaagagtggtgttcggagcacgggacgccaagaccggcgcagcaggctccctgatggatgtgctgcaccacccccggcatgaaccaccgggtgg
agatcacagagggaatcctggcagacgagtgcgccgccctgctgagcgatttctttagaatgcggagacaggagatcaaggcccagaagaaggcacagag
ctccaccgactctggaggatctagcggaggatcctctggaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggagg
atcctctgaggtggagtttccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggaggtgcctgtgggagccgtgctg
gtgctgaacaatagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggccctgagacagggcggcctg
gtcatgcagaactacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtg
tttggcgtgaggaacgcaaaaaccggcgccgcaggctccctgatggacgtgctgcactacccccggcatgaatcaccgcgtcgaaattaccgagggaatcctgg
cagatgaatgtgccgccctgctgtgctatttctttcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaccgactccggaggatctagcg
gaggctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagcgggggggtcaatggccgccttcaagcctaac
ccaatcaattacgatatcGACAAGAAGTACTCCATTGGGCTCGcTATCGGTACCAACAGCGTCGGCTGGGCCGTCATTACGGAC
GAGTACAAGGTGCCGAGCAAAAAATTCAAAGTTCTGGGCAATACCGATCGCCACAGCATAAAGAAGAACCTCATTGGAG
CCCTCCTGTTCGACTCCGGGGAGACGGCCGAAGCCACGCGGCTCAAAAGAACAGCACGGCGCAGATATACCCGCAGAAA
GAATCGGATCTGCTACCTGCAGGAGATCTTTAGTAATGAGATGGCTAAGGTGGATGACTCTTTCTTCCATAGGCTGGAGG
AGTCCTTTTTGGTGGAGGAGGATAAAAAGCACGAGCGCCACCCAATCTTTGGCAATATCGTGGACGAGGTGGCGTACCA
TGAAAAGTACCCAACCATATATCATCTGAGGAAGAAGCTGGTAGACAGTACTGATAAGGCTGACTTGCGGTTGATCTATC
TCGCGCTGGCGCACATGATCAAATTTCGGGGACACTTCCTCATCGAGGGGGACCTGAACCCAGACAACAGCGATGTCGA
CAAACTCTTTATCCAACTGGTTCAGACTTACAATCAGCTTTTCGAGGAGAACCCGATCAACGCATCCGGCGTTGACGCCAA
AGCAATCCTGAGCGCTAGGCTGTCCAAATCCCGGCGGCTCGAAAACCTCATCGCACAGCTCCCTGGGGAGAAGAAGAAC
GGCCTGTTTGGTAATCTTATCGCCCTGTCACTCGGGCTGACCCCCAACTTTAAATCTAACTTCGACCTGGCCGAAGATGCC
AAGCTGCAACTGAGCAAAGACACCTACGATGATGATCTCGACAATCTGCTGGCCCAGATCGGCGACCAGTACGCAGACC
TTTTTTTGGCGGCAAAGAACCTGTCAGACGCCATTCTGCTGAGTGATATTCTGCGAGTGAACACGGAGATCACCAAAGCT
CCGCTGAGCGCTAGTATGATCAAGCGCTATGATGAGCACCACCAAGACTTGACTTTGCTGAAGGCCCTTGTCAGACAGCA
ACTGCCTGAGAAGTACAAGGAAATTTTCTTCGATCAGTCTAAAAATGGCTACGCCGGATACATTGACGGCGGAGCAAGCC
AGGAGGAATTTTACAAATTTATTAAGCCCATCTTGGAAAAAATGGACGGCACCGAGGAGCTGCTGGTAAAGCTGAACAG
AGAAGATCTGTTGCGCAAACAGCGCACTTTCGACAATGGAAGCATCCCCCACCAGATTCACCTGGGCGAACTGCACGCTA
TCCTCAGGCGGCAAGAGGATTTCTACCCCTTTTTTGAAAGATAACAGGGAAAAGATTGAGAAAATCCTCACATTTCGGATA
CCCTACTATGTAGGCCCCCTCGCTCGGGGAAATTCCAGATTCGCGTGGATGACTCGCAAATCAGAAGAGACCATCACTCC
CTGGAACTTCGAGGAAGTCGTGGATAAGGGGGCCTCTGCCCAGTCCTTCATCGAAAGGATGACTAACTTTGATAAAAATC
TGCCTAACGAAAAGGTGCTTCCTAAACACTCTCTGCTGTACGAGTACTTCACAGTTTATAACGAGCTCACCAAGGTCAAAT
ACGTCACAGAAGGGATGAGAAAGCCAGCATTCCTGTCTGGAGAGCAGAAGAAAGCTATCGTGGACCTCCTCTTCAAGAC
GAACCGGAAAGTTACCGTGAAACAGCTCAAAGAAGACTATTTCAAAAAGATTGAATGTTTCGACTCTGTTGAAATCAGCG
GAGTGGAGGATCGCTTCAACGCATCCCTGGGAACGTATCACGATCTCCTGAAAATCATTAAAGACAAGGACTTCCTGGAC
AATGAGGAGAACGAGGACATTCTTGAGGACATTGTCCTCACCCTTACGTTGTTTGAAGATAGGGAGATGATTGAAGAAC
GCTTGAAAACTTACGCTCATCTCTTCGACGACAAAGTCATGAAACAGCTCAAGAGACGCCGATATACAGGATGGGGGCG
GCTGTCAAGAAAACTGATCAATGGCATCCGAGACAAGCAGAGTGGAAAGACAATCCTGGATTTTCTTAAGTCCGATGGA
TTTGCCAACCGGAACTTCATGCAGTTGATCCATGATGACTCTCTCACCTTTAAGGAGGACATCCAGAAAGCACAAGTTTCT
GGCCAGGGGGACAGTCTTCACGAGCACATCGCTAATCTTGCAGGTAGCCCAGCTATCAAAAAGGGAATACTGCAGACCG
TTAAGGTCGTGGATGAACTCGTCAAAGTAATGGGAAGGCATAAGCCCGAGAATATCGTTATCGAGATGGCCCGAGAGAA
CCAAACTACCCAGAAGGGACAGAAGAACAGTAGGGAAAGGATGAAGAGGATTGAAGAGGGTATAAAAGAACTGGGGT
CCCAAATCCTTAAGGAACACCCAGTTGAAAACACCCAGCTTCAGAATGAGAAGCTCTACCTGTACTACCTGCAGAACGGC
AGGGACATGTACGTGGATCAGGAACTGGACATCAACCGGTTGTCCGACTACGACGTGGATCATATCGTGCCCCAAAGCTT
TCTCAAAGATGATTCTATTGATAATAAAGTGTTGACAAGATCCGATAAAAATAGAGGGAAGAGTGATAACGTCCCCTCAG
AAGAAGTTGTCAAGAAAATGAAAAATTATTGGCGGCAGCTGCTGAACGCCAAACTGATCACACAACGGAAGTTCGATAA
TCTGACTAAGGCTGAACGAGGTGGCCTGTCTGAGTTGGATAAAGCCGGCTTCATCAAAAGGCAGCTTGTTGAGACACGC
CAGATCACCAAGCACGTGGCCCAAATTCTCGATTCACGCATGAACACCAAGTACGATGAAAATGACAAACTGATTCGAGA
GGTGAAAGTTATTACTCTGAAGTCTAAGCTGGTCTCAGATTTCAGAAAGGACTTTCAGTTTTATAAGGTGAGAGAGATCA

ACAATTACCACCATGCGCATGATGCCTACCTGAATGCAGTGGTAGGCACTGCACTTATCAAAAAATATCCCAAGCTGGAA
TCTGAATTTGTTTACGGAGACTATAAAGTGTACGATGTTAGGAAAATGATCGCAAAGTCTGAGCAGGAAATAGGCAAGG
CCACCGCTAAGTACTTCTTTTACAGCAATATTATGAATTTTTTCAAGACCGAGATTACACTGGCCAATGGAGAGATTCGGA
AGCGACCACTTATCGAAACAAACGGAGAAACAGGAGAAATCGTGTGGGACAAGGGTAGGGATTTCGCGACAGTCCGCA
AGGTCCTGTCCATGCCGCAGGTGAACATCGTTAAAAAGACCGAGTACAGACCGGAGGCTTCTCCAAGGAAAGTATCCT
CCCGAAAAGGAACAGCGACAAGCTGATCGCACGCAAAAAGATTGGGACCCCAAGAAATACGGCGGATTCGATTCTCCT
ACAGTCGCTTACAGTGTACTGGTTGTGGCCAAAGTGGAGAAAGGGAAGTCTAAAAAACTCAAAAGCGTCAAGGAACTGC
TGGGCATCACAATCATGGAGCGATCCAGCTTCGAGAAAAACCCCATCGACTTTCTCGAAGCGAAAGGATATAAAGAGGT
CAAAAAAGACCTCATCATTAAGCTGCCCAAGTACTCTCTCTTTGAGCTTGAAAACGGCCGGAAACGAATGCTCGCTAGTG
CGGGCGAGCTGCAGAAAGGTAACGAGCTGGCACTGCCCTCTAAATACGTTAATTTCTTGTATCTGGCCAGCCACTATGAA
AAGCTCAAAGGGTCTCCCGAAGATAATGAGCAGAAGCAGCTGTTCGTGGAACAACACAAACACTACCTTGATGAGATCA
TCGAGCAAATAAGCGAGTTCTCCAAAAGAGTGATCCTCGCCGACGCTAACCTCGATAAGGTGCTTTCTGCTTACAATAAG
CACAGGGATAAGCCCATCAGGGAGCAGGCAGAAAACATTATCCACTTGTTTACTCTGACCAACTTGGGCGCGCCTGCAGC
CTTCAAGTACTTCGACACCACCATAGACAGAAAGCGGTACACCTCTACAAAGGAGGTCCTGGACGCCACACTGATTCATC
AGTCAATTACGGGGCTCTATGAAACAAGAATCGACCTCTCTCAGCTCGGTGGAGACgatatcGGCACCGGCGGGtctggagga
tctagcggaggatcctctggaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggaggatccGTATACCCATAC
GATGTTCCTGACTATGCGGGCTATCCCTATGACGTCCCGGACTATGCAGGATCGTATCCTTATGACGTTCCAGATTACGCT
GGATCCGCCGCTCCGGCAGCTAAGAAAAAGAAACTGGATTTCGAATCCGGAgaattcctgcagcccggggggatccactagttctagaG
GATCCGCCGCTCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCGGCAGCGGCGGCGGCAGCGGCGGCGGCAGCGGCggcg
gtagcggcggaggcagcggtggcggcagcggcAAGCGCACCGCCGACAGCCAGCACAGCACCCCCCCCCAAGACCAAGCGCAAGGT
GGGCGGCAGCGGCGGCGGCAGCGGCGGCGGCAGCGGCTCCGGACCGGCAGCTAAGAGGGTGAAACTGGATtgaTAAgc
ggccgccaccgcggtggagctccagcttttgttccctttagtgagggttaatGAATTCGAGCCCAAGAAGAAGAGGAAAGTCTAACCGGTCA
TCATCACCATCACCATTGAGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCC
TCCCCCGTGCCTTCCTTGACCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGAAAATTGCATCGCATTGTC
TGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGG
CATGCTGGGGATGCGGTGGGCTCTATGG

Spy-ABE8e construct in **Figure 1**
Legend: CMV enhancer, CMV promoter, NLS, TadA8e, linker, SpyCas9-D10A, 3xHA, poly-A signal
GACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTTA
CATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCA
TAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAA
GTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGACC
TTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCA
ATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTTTTGGCAC
CAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGG
AGGTCTATATAAGCAGAGCTGGTTTAGTGAACCGTCAGATCCGCTAGAGATCCGCGGCCGCtaatacgactcactatagggcgaa
ttgggtaccgggcccGCCACCATGGctagcCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCAGCGGCGG
CGGCAGCGGCCCGGCAGCTAAGAGGGTGAAACTGGATggcggtagcggcggaggcagcggtggcggcagcggcccccctcgagTctgagg
tggagttttcccacgagtactggatgagacatgccctgacctggccaagagggcacgcgatgagagggaggtgcctgtgggagccgtgctggtgctgaacaa
tagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggccctgagacagggcggcctggtcatgcagaa
ctacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtgtttggcgtgag
gaacAGCaaaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcaccgcgtcgaaattaccgagggaatcctggcagatga
atgtgccgccctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaTcAactccggaggatctagcggagg
ctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagcgggggggtcaatggccgccttcaagcctaacccaat
caattacgatatcGACAAGAAGTACTCCATTGGGCTCGcTATCGGTACCAACAGCGTCGGCTGGGCCGTCATTACGGACGAGT
ACAAGGTGCCGAGCAAAAAAATTCAAAGTTCTGGGCAATACCGATCGCCACAGCATAAAGAAGAACCTCATTGGAGCCCT
CCTGTTCGACTCCGGGGAGACGGCCGAAGCCACGCGGCTCAAAAGAACAGCACGGCGCAGATATACCCGCAGAAGAA
TCGGATCTGCTACCTGCAGGAGATCTTTAGTAATGAGATGGCTAAGGTGGATGACTCTTTCTTCCATAGGCTGGAGGAGT

CCTTTTTGGTGGAGGAGGATAAAAAGCACGAGCGCCACCCAATCTTTGGCAATATCGTGGACGAGGTGGCGTACCATGA
AAAGTACCCAACCATATATCATCTGAGGAAGAAGCTGGTAGACAGTACTGATAAGGCTGACTTGCGGTTGATCTATCTCG
CGCTGGCGCACATGATCAAATTTCGGGGACACTTCCTCATCGAGGGGGACCTGAACCCAGACAACAGCGATGTCGACAA
ACTCTTTATCCAACTGGTTCAGACTTACAATCAGCTTTTCGAGGAGAACCCGATCAACGCATCCGGCGTTGACGCCAAAGC
AATCCTGAGCGCTAGGCTGTCCAAATCCCGGCGGCTCGAAAACCTCATCGCACAGCTCCCTGGGGAGAAGAAGAACGGC
CTGTTTGGTAATCTTATCGCCCTGTCACTCGGGCTGACCCCCAACTTTAAATCTAACTTCGACCTGGCCGAAGATGCCAAG
CTGCAACTGAGCAAAGACACCTACGATGATGATCTCGACAATCTGCTGGCCCAGATCGGCGACCAGTACGCAGACCTTTT
TTTGGCGGCAAAGAACCTGTCAGACGCCATTCTGCTGAGTGATATTCTGCGAGTGAACACGGAGATCACCCAAAGCTCCGC
TGAGCGCTAGTATGATCAAGCGCTATGATGAGCACCACCAAGACTTGACTTTGCTGAAGGCCCTTGTCAGACAGCAACTG
CCTGAGAAGTACAAGGAAATTTTCTTCGATCAGTCTAAAAATGGCTACGCCGGATACATTGACGGCGGAGCAAGCCAGG
AGGAATTTTACAAATTTATTAAGCCCATCTTGGAAAAAATGGACGGCACCGAGGAGCTGCTGGTAAAGCTGAACAGAGA
AGATCTGTTGCGCAAACAGCGCACTTTCGACAATGGAAGCATCCCCCACCAGATTCACCTGGGCGAACTGCACGCTATCC
TCAGGCGGCAAGAGGATTTCTACCCCTTTTTGAAAGATAACAGGGAAAAGATTGAGAAAATCCTCACATTTCGGATACCC
TACTATGTAGGCCCCCTCGCTCGGGGAAATTCCAGATTCGCGTGGATGACTCGCAAATCAGAAGAGACCATCACTCCCTG
GAACTTCGAGGAAGTCGTGGATAAGGGGGCCTCTGCCCAGTCCTTCATCGAAAGGATGACTAACTTTGATAAAAATCTGC
CTAACGAAAAGGTGCTTCCTAAACACTCTCTGCTGTACGAGTACTTCACAGTTTATAACGAGCTCACCAAGGTCAAATACG
TCACAGAAGGGATGAGAAAGCCAGCATTCCTGTCTGGAGAGCAGAAGAAAGCTATCGTGGACCTCCTCTTCAAGACGAA
CCGGAAAGTTACCGTGAAACAGCTCAAAGAAGACTATTTCAAAAAGATTGAATGTTTCGACTCTGTTGAAATCAGCGGAG
TGGAGGATCGCTTCAACGCATCCCTGGGAACGTATCACGATCTCCTGAAAATCATTAAAGACAAGGACTTCCTGGACAAT
GAGGAGAACGAGGACATTCTTGAGGACATTGTCCTCACCCTTACGTTGTTTGAAGATAGGGAGATGATTGAAGAACGCTT
GAAAACTTACGCTCATCTCTTCGACGACAAAGTCATGAAACAGCTCAAGAGACGCCGATATACAGGATGGGGGCGGCTG
TCAAGAAAACTGATCAATGGCATCCGAGACAAGCAGAGTGGAAAGACAATCCTGGATTTTCTTAAGTCCGATGGATTTGC
CAACCGGAACTTCATGCAGTTGATCCATGATGACTCTCTCACCTTTAAGGAGGACATCCAGAAAGCACAAGTTTCTGGCCA
GGGGGACAGTCTTCACGAGCACATCGCTAATCTTGCAGGTAGCCCAGCTATCAAAAAGGGAATACTGCAGACCGTTAAG
GTCGTGGATGAACTCGTCAAAGTAATGGGAAGGCATAAGCCCGAGAATATCGTTATCGAGATGGCCCGAGAGAACCAAA
CTACCCAGAAGGGACAGAAGAACAGTAGGGAAAGGATGAAGAGGATTGAAGAGGGTATAAAAGAACTGGGGTCCCAA
ATCCTTAAGGAACACCCAGTTGAAAACACCCAGCTTCAGAATGAGAAGCTCTACCTGTACTACCTGCAGAACGGCAGGGA
CATGTACGTGGATCAGGAACTGGACATCAACCGGTTGTCCGACTACGACGTGGATCATATCGTGCCCCAAAGCTTTCTCA
AAGATGATTCTATTGATAATAAAGTGTTGACAAGATCCGATAAAAATAGAGGGAAGAGTGATAACGTCCCCTCAGAAGA
AGTTGTCAAGAAATGAAAAATTATTGGCGGCAGCTGCTGAACGCCAAACTGATCACACAACGGAAGTTCGATAATCTGA
CTAAGGCTGAACGAGGTGGCCTGTCTGAGTTGGATAAAGCCGGCTTCATCAAAAGGCAGCTTGTTGAGACACGCCAGAT
CACCAAGCACGTGGCCCAAATTCTCGATTCACGCATGAACACCAAGTACGATGAAAATGACAAACTGATTCGAGAGGTGA
AAGTTATTACTCTGAAGTCTAAGCTGGTCTCAGATTTCAGAAAGGACTTTCAGTTTTATAAGGTGAGAGAGATCAACAATT
ACCACCATGCGCATGATGCCTACCTGAATGCAGTGGTAGGCACTGCACTTATCAAAAAATATCCCAAGCTGGAATCTGAA
TTTGTTTACGGAGACTATAAAGTGTACGATGTTAGGAAAATGATCGCAAAGTCTGAGCAGGAAATAGGCAAGGCCACCG
CTAAGTACTTCTTTTACAGCAATATTATGAATTTTTTCAAGACCGAGATTACACTGGCCAATGGAGAGATTCGGAAGCGAC
CACTTATCGAAACAAACGGAGAAACAGGAGAAATCGTGTGGGACAAGGGTAGGGATTTCGCGACAGTCCGCAAGGTCC
TGTCCATGCCGCAGGTGAACATCGTTAAAAAGACCGAAGTACAGACCGGAGGCTTCTCCAAGGAAAGTATCCTCCCGAA
AAGGAACAGCGACAAGCTGATCGCACGCAAAAAGATTGGGACCCCAAGAAATACGGCGGATTCGATTCTCCTACAGTC
GCTTACAGTGTACTGGTTGTGGCCAAAGTGGAGAAAGGGAGTCTAAAAAACTCAAAGCGTCAAGGAACTGCTGGGC
ATCACAATCATGGAGCGATCCAGCTTCGAGAAAAACCCCATCGACTTTCTCGAAGCGAAAGGATATAAAGAGGTCAAAA
AAGACCTCATCATTAAGCTGCCCAAGTACTCTCTCTTTGAGCTTGAAAACGGCCGGAAACGAATGCTCGCTAGTGCGGGC
GAGCTGCAGAAAGGTAACGAGCTGGCACTGCCCTCTAAATACGTTAATTTCTTGTATCTGGCCAGCCACTATGAAAAGCT
CAAAGGGTCTCCCGAAGATAATGAGCAGAAGCAGCTGTTCGTGGAACAACACAAACACTACCTTGATGAGATCATCGAG
CAAATAAGCGAGTTCTCCAAAAGAGTGATCCTCGCCGACGCTAACCTCGATAAGGTGCTTTCTGCTTACAATAAGCACAG
GGATAAGCCCATCAGGGAGCAGGCAGAAAACATTATCCACTTGTTTACTCTGACCAACTTGGGCGCGCCTGCAGCCTTCA
AGTACTTCGACACCACCATAGACAGAAAGCGGTACACCTCTACAAAGGAGGTCCTGGACGCCACACTGATTCATCAGTCA
ATTACGGGGCTCTATGAAACAAGAATCGACCTCTCTCAGCTCGGTGGAGACgatatcGGCACCGGCGGGtctggaggatctagc
ggaggatcctctggaagcgagacaccaggcacaagcgagtccgccacaccagagagctccggcggctcctccggaggatccGTATACCCATACGATG

TTCCTGACTATGCGGGCTATCCCTATGACGTCCCGGACTATGCAGGATCGTATCCTTATGACGTTCCAGATTACGCTGGAT
CCGCCGCTCCGGCAGCTAAGAAAAGAAACTGGATTTCGAATCCGGAgaattcctgcagcccggggggatccactagttctagaGGATC
CGCCGCTCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCGGCAGCGGCGGCGGCGGCAGCGGCGGCGGCAGCGGCggcggtagc
ggcggaggcagcggtggcggcagcggcAAGCGCACCGCCGACAGCCAGCACAGCACCCCCCCCCAAGACCAAGCGCAAGGTGGGC
GGCAGCGGCGGCGGCAGCGGCGGCGGCAGCGGCTCCGGACCGGCAGCTAAGAGGGTGAAACTGGATTtgaTAAgcggccg
ccaccgcggtggagctccagcttttgttccctttagtgagggttaatGAATTCGAGCCCAAGAAGAAGAGGGAAAGTCTAACCGGTCATCAT
CACCATCACCATTGAGTTTAAACCCGCTGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCTCCC
CCGTGCCTTCCTTGACCCTGGAAGGTGCCACTCCCACTGTCCTTTCCTAATAAAATGAGAAAATTGCATCGCATTGTCTGA
GTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCAT
GCTGGGGATGCGGTGGGCTCTATGG

U6-sgRNA construct for Nme2-ABE in **Figure 1**
Legend: U6 promoter, **guide sequence**, guide RNA scaffold, U6 termination signal

GAGGGCCTATTTCCCATGATTCCTTCATATTTGCATATACGATACAAGGCTGTTAGAGAGATAATTGGAATTAATTTGACT
GTAAACACAAAGATATTAGTACAAAATACGTGACGTAGAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGT
TTTAAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTCGATTTCTTGGCTTTATATATCTTGTGGAAAGGAC**NN
NNNNNNNNNNNNNNNNNNNNNNNN**GTTGTAGCTCCCTTTCTCATTTCGGAAACGAAATGAGAACCGTTGCTACAATAAG
GCCGTCTGAAAAGATGTGCCGCAACGCTCTGCCCCTTAAAGCTTCTGCTTTAAGGGGCATCGTTTATTTTTTT

U6-sgRNA construct for Spy-ABE in **Figure 1**
Legend: U6 promoter**, guide sequence**, guide RNA scaffold, U6 termination signal
GAGGGCCTATTTCCCATGATTCCTTCATATTTGCATATACGATACAAGGCTGTTAGAGAGATAATTGGAATTAATTTGACT
GTAAACACAAAGATATTAGTACAAAATACGTGACGTAGAAAGTAATAATTTCTTGGGTAGTTTGCAGTTTTAAAATTATGT
TTTAAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTCGATTTCTTGGCTTTATATATCTTGTGGAAAGGAC**NN
NNNNNNNNNNNNNNNNNNNNNNN**GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAG
TGGCACCGAGTCGGTGCTTTTTTTT

Single-AAV vector with 2xcMyc NLS in **Figure 3a**
Legend: ITR, U1a promoter, cMyc NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, U6 promoter,
guide sequence, guide RNA scaffold, U6 termination signal
Cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggcgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgc
gcagagagggagtggccaactccatcactaggggttcctgcggcctctagaGTTTAAACAAAAAAA
TAAACGATGCCCCTTAAAGCAGAAGCTTTAAGGGGCAGAGCGTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAAC
GTTTCGGGAGCTACAAC**NNNNNNNNNNNNNNNNNNNNNNNNNNNNNN**GGTGTTTCGTCCTTTCCACAAGATATATAAAGCCA
AGAAATCGAAATACTTTCAAGTTACGGTAAGCATATGATAGTCCATTTTAAAACATAATTTTAAAACTGCAAACTACCCAA
GAAATTATTACTTTCTACGTCACGTATTTTGTACTAATATCTTTGTGTTTACAGTCAAATTAATTCTAATTATCTCTCTAACA
GCCTTGTATCGTATATGCAAATATGAAGGAATCATGGGAAATAGGCCCTCTCTAGaatggaggcggtactatgtagatgagaattca
ggagcaaactgggaaaagcaactgcttccaaatatttgtgattttttacagtgtagttttggaaaaactcttagcctaccaattcttctaagtgttttaaaatgtggg
agccagtacacatgaagttatagagtgttttaatgaggcttaaatatttaccgtaactatgaaatgctacgcatatcatgctgttcaggctccgtggccacgcaa
ctcatactACCGGTGCCACCATGCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCTctgaggtggagttttc
ccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggaggtgcctgtgggagccgtgctggtgctgaacaatagagtgat
cggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggccctgagacagggcggcctggtcatgcagaactacagact
gattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtgtttggcgtgaggaacAGCa
aaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcaccgcgtcgaaattaccgagggaatcctggcagatgaatgtgccgcc
ctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaTcAactccggaggatctagcggaggctcctctggc
tctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagcgggggggtcaatggccgccttcaagcctaacccaatcaattacat
cctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgaggaggagaatcctatccggctgatcgatctgggcgtgagagtgtttg
agagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacggagcgtgaggcgcctgacacggagaagggcacacaggctgctg

agggcacgccggctgctgaagagagagaggcgtgctgcaggcagcagacttcgatgagaatggcctgatcaagagcctgccaaacacccctggcagctgag
agcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacctgatcaagcaccgcggctacctgagccagcggaagaacgaggg
agagacagcacaaggagctgggcgccctgctgaaggagtggccaacaatgcccacgccctgcagaccggcgatttcaggacacctgccgagctggccc
tgaataagtttgagaaggagtccggccacatcagaaaccagagggcgactatagccacaccttctcccgcaaggatctgcaggccgagctgatcctgctgtt
cgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggagggaatcgagaccctgctgatgacacagaggcctgccctgtccggcgacg
cagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaagaacacctacacagccgagcggtttatctggctgacaaagctga
acaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagagggccaccctgatggatgagccttaccggaagtctaagctgacatatg
cccaggccagaaagctgctgggcctggaggacaccgccttctttaagggcctgagatacggcaaggataatgccgaggcctccacactgatggagatgaagg
cctatcacgccatctctcgcgccctggagaaggagggcctgaaggacaagaagtccccccctgaacctgagctccgagctgcaggatgagatcggcaccgcctt
ctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacaggtgcagcctgagatcctggaggccctgctgaagcacatctctttcgataagtttgt
gcagatcagcctgaaggccctgagaaggatcgtgccactgatggagcagggcaagcggtacgacgaggcctgcgccgagatctacggcgatcactatggca
agaagaacacagaggagaagatctatctgcccccctatccctgccgacgagatcagaaatcctgtggtgctgagggccctgtcccaggcaagaaaagtgatca
acggagtggtgcgccggtacggatctccagcccggatccacatcgagaccgccagagaagtgggcaagagcttcaaggaccggaaggagatcgagaagag
acaggaggagaatcgcaaggatcgggagaaggccgccgccaagtttagggagtacttccctaactttgtgggcgagccaaagtctaaggacatcctgaagct
gcgcctgtacgagcagcagcacggcaagtgtctgtatagcggcaaggagatcaatctggtgcggctgaacgagaagggctatgtggagatcgatcacgccct
gcctttctccagaacctgggacgattcttttaacaataaggtgctggtgctgggcagcgagaaccagaataagggcaatcagacaccatacgagtatttcaatg
gcaaggacaactccagggagtggcaggagttcaaggcccgcgtggagacctctagatttcccaggagcaagaagcagcggatcctgctgcagaagttcgacg
aggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgccagtttgtggccgatcacatcctgctgaccggcaagggcaagag
aagggtgttcgcctctaatggccagatcacaaacctgctgaggggattttggggactgaggaaggtgcgggcagagaatgacagacaccacgcactggatgc
agtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgaggtataaggagatgaacgccttttgacggcaagaccatcgataagga
gacaggcaaggtgctgcaccagaagacccacttcccccagccttgggagttctttgcccaggaagtgatgatccgggtgttcggcaagccagacggcaagcct
gagtttgaggaggccgatacccccagagaagctgaggacactgctggcagagaagctgtctagcaggccagaggcagtgcacgagtacgtgaccccactgttc
gtgtccagggcacccaatcggaagatgtctggcgcccacaaggacacactgagaagcgccaagaggtttgtgaagcacaacgagaagatctccgtgaagag
agtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaacggcagggagatcgagctgtatgaggccctgaaggcaaggctgg
aggcctacggaggaaatgccaagcaggccttcgacccaaaggataacccctttataagaagggaggacagctggtgaaggccgtgcgggtggagaagacc
caggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcgatatggtgagagtggacgtgttctgtaaggtggataagaagggc
aagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatcgattgcaagggctacagaatcgacgatagctatacatt
ctgttttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtggagtttgcctactatatcaattgcgactcctctaacggcaggttc
tacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaatctggtgctgatccagaagtatcaggtgaacgagctgggcaag
gagatcaggcc**ATGTCGGCTGAAGAAGCGCCC**acccgtgcgg**GAGGAT**CCCGCCGCCAAGCGCGTGAAGCTGGACTAAAATAA
AAGATCTTTATTTTCATTAGATCTGTGTGTTGGTTTTTTGTGTAAGCgcggccgcaggaacccctagtgatggagttggccactccctctct
gcgcgctcgctcgctcactgaggccgggcgaccaaaggtcgcccgacgcccgggctttgcccgggcggcctcagtgagcgagcgagcgcgcagctgcctgca
gg

Single-AAV vector with 3xcMyc NLS in **Figure 3a**
Legend: ITR, U1a promoter, cMyc NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, U6 promoter,
guide sequence, guide RNA scaffold, U6 termination signal
Cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggcgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgc
gcagagagggagtggccaactccatcactaggggttcctgcggcctctagaGTTTAAACAAAAAAA
TAAACGATGCCCCTTAAAGCAGAAGCTTTAAGGGGCAGAGCGTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAAC
GTTTCGGGAGCTACAAC**NNNNNNNNNNNNNNNNNNNNNNNNNNNNN**GGTGTTTCGTCCTTTCCACAAGATATATAAAGCCA
AGAAATCGAAATACTTTCAAGTTACGGTAAGCATATGATAGTCCATTTTAAAACATAATTTTAAAACTGCAAACTACCCAA
GAAATTATTACTTTCTACGTCACGTATTTTGTACTAATATCTTTGTGTTTACAGTCAAATTAATTCTAATTATCTCTCTAACA
GCCTTGTATCGTATATGCAAATATGAAGGAATCATGGGAAATAGGCCCTCTCTAGAatggaggcggtactatgtagatgagaattca
ggagcaaactgggaaaagcaactgcttccaaatatttgtgattttttacagtgtagtttggaaaaactcttagcctaccaattcttctaagtgtttttaaaatgtggg
agccagtacacatgaagttatagagtgttttaatgaggcttaaatatttaccgtaactatgaaatgctacgcatatcatgctgttcaggctccgtggccacgcaa
ctcatactACCGGTGCCACCATGCCCGCCGCCAAGCGCGTGAAGCTGGACGGCGGCAGCGGCGGCGGCTctgaggtggagttttc
ccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggaggtgcctgtgggagccgtgctggtgctgaacaatagagtgat
cggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatggcccctgagacagggcggcctggtcatgcagaactacagact

gattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtgtttggcgtgaggaacAGCa
aaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcaccgcgtcgaaattaccgagggaatcctggcagatgaatgtgccgcc
ctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggcccagagctccaTcAactccggaggatctagcggaggctcctctggc
tctgagacacctggcacaagcgagagcgcaacacctgaaagcagcgggggcagcagcggggggtcaatggccgcctttcaagcctaacccaatcaattacat
cctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgaggaggagaatcctatccggctgatcgatctgggcgtgagagtgtttg
agagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacggagcgtgaggcgcctgacacggagaagggcacacaggctgctg
agggcacgccggctgctgaagagagagggcgtgctgcaggcagcagacttcgatgagaatggcctgatcaagagcctgccaaacaccccctggcagctgag
agcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacctgatcaagcaccgcggctacctgagccagcggaagaacgaggg
agagacagcagacaaggagctgggcgccctgctgaagggagtggccaacaatgcccacgccctgcagaccggcgatttcaggacacctgccgagctggccc
tgaataagtttgagaaggagtccggccacatcagaaaccagagagggcgactatagccacaccttctcccgcaaggatctgcaggccgagctgatcctgctgtt
cgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggagggaatcgagaccctgctgatgacacagaggcctgccctgtccggcgacg
cagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaagaacacctacacagccgagcggtttatctggctgacaaagctga
acaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagagggccaccctgatggatgagccttaccggaagtctaagctgacatatg
cccaggccagaaagctgctggggcctggaggacaccgccttctttaagggcctgagatacggcaaggataatgccgaggcctccacactgatggagatgaagg
cctatcacgccatctctcgcgccctggagaaggagggcctgaaggacaagaagtcccccctgaacctgagctccgagctgcaggatgagatcggcaccgcctt
ctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacagggtgcagcctgagatcctggaggccctgctgaagcacatctctttcgataagtttgt
gcagatcagcctgaaggccctgagaaggatcgtgccactgatggagcagggcaagcggtacgacgaggcctgcgccgagatctacggcgatcactatggca
agaagaacacagaggagaagatctatctgccccctatccctgccgacgagatcagaaatcctgtggtgctgagggccctgtcccaggcaagaaaagtgatca
acggagtggtgcgccggtacggatctccagcccggatccacatcgagaccgccagagaagtgggcaagagcttcaaggaccggaaggagatcgagaagag
acaggaggagaatcgcaaggatcgggagaaggccgccgccaagtttagggagtacttccctaactttgtgggcgagcccaaagtctaaggacatcctgaagct
gcgcctgtacgagcagcagcacggccaagtgtctgtatagcggcaaggagatcaatctggtgcggctgaacgagaagggctatgtggagatcgatcacgccct
gcctttctccagaacctgggacgattcttttaacaataaggtgctggtgctgggcagcgagaaccagaataagggcaatcagacaccatacgagtatttcaatg
gcaaggacaactccagggagtggcaggagttcaaggcccgcgtggagacctctagatttcccaggagcaagaagcagcggatcctgctgcagaagttcgacg
aggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgccagtttgtggccgatcacatcctgctgaccggcaagggcaagag
aagggtgttcgcctctaatggccagatcacaaacctgctgagggggattttggggactgaggaaggtgcgggcagagaatgacagacaccacgcactggatgc
agtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgaggtataaggagatgaacgcctttgacggcaagaccatcgataagga
gacaggcaaggtgctgcaccagaagacccacttcccccagccttgggagttctttgcccaggaagtgatgatccgggtgttcggcaagccagacggcaagcct
gagtttgaggaggccgataccccagagaagctgaggacactgctggcagagaagctgtctagcaggccagaggcagtgcacgagtacgtgacccactgttc
gtgtccaggcacccaatcggaagatgtctggcgcccacaaggacacactgagaagcgccaagaggtttgtgaagcacaacgagaagatctccgtgaagag
agtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaacggcagggagatcgagctgtatgaggccctgaaggcaaggctgg
aggcctacggaggaaatgccaagcaggccttcgacccaaaggataaccccttttataagaaggaggacagctggtgaaggccgtgcgggtggagaagacc
caggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcgatatggtgagagtggacgtgttctgtaaggtggataagaagggc
aagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatcgattgcaagggctacagaatcgacgatagctatacatt
ctgtttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtggagtttgcctactatatcaattgcgactcctctaacggcaggttc
tacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaatctggtgctgatccagaagtatcaggtgaacgagctgggcaag
gagatcaggccATGTCGGCTGAAGAAGCGCCCacccgtgcggGAGGATCCGGCAGCTAAGAGGGTGAAACTGGATGGCGGCA
GCGGCGGCGGCAGCGGCGGCGGCAGCGGCCCCGCCGCCAAGCGCGTGAAGCTGGACTAAAATAAAAGATCTTTATTTTC
ATTAGATCTGTGTGTTGGTTTTTTGTGTAAGCgcggccgcaggaacccctagtgatggagttggccactccctctctgcgcgctcgctcgctcact
gaggccgggcgaccaaaggtcgcccgacgcccgggctttgcccgggcggcctcagtgagcgagcgagcgcgcagctgcctgcagg

Single-AAV vector with Ty1 NLS in **Figure 3a**

Legend: ITR, U1a promoter, Ty1 NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, U6 promoter, **guide sequence**, guide RNA scaffold, U6 termination signal

Cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggcgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgc
gcagagagggagtggccaactccatcactaggggttcctgcggcctctagaGTTTAAACAAAAAAA
TAAACGATGCCCCTTAAAGCAGAAGCTTTAAGGGGCAGAGCGTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAAC
GTTTCGGGAGCTACAACNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNGGTGTTTCGTCCTTTCCACAAGATATATAAAGCCA
AGAAATCGAAATACTTTCAAGTTACGGTAAGCATATGATAGTCCATTTTAAAACATAATTTTAAAACTGCAAACTACCCAA
GAAATTATTACTTTCTACGTCACGTATTTTGTACTAATATCTTTGTGTTTACAGTCAAATTAATTCTAATTATCTCTCTAACA

GCCTTGTATCGTATATGCAAATATGAAGGAATCATGGGAAATAGGCCCTCTCTAGAatggaggcggtactatgtagatgagaattca
ggagcaaactgggaaaagcaactgcttccaaatatttgtgattttttacagtgtagtttttggaaaaactcttagcctaccaattcttctaagtgtttttaaaatgtggg
agccagtacacatgaagttatagagtgttttaatgaggcttaaatatttaccgtaactatgaaatgctacgcatatcatgctgttcaggctccgtggccacgcaa
ctcatactACCGGTGCCACCATGAACTCAAAGAAAAGGTCACTGGAGGACAACGAAACGGAAATCAAAGTCTCCAGAGACA
CCTGGAACACTAAGAACATGCGGTCCCTGGAACCGCCACGAAGCAAGAAACGGATACATGGCGGTAGCGGCGGAGGCA
GCGGTGGCGGCAGCGGCTctgaggtggagtttttcccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagggagg
tgcctgtgggagccgtgctggtgctgaacaatagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaattatgg
ccctgagacagggcggcctggtcatgcagaactacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgatccac
tctaggatcggccgcgtggtgtttggcgtgaggaacAGCaaaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcaccgcgt
cgaaattaccgagggaatcctggcagatgaatgtgccgccctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggcccagag
ctccaTcAactccggaggatcagcggaggctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggcagcagcgggg
ggtcaatggccgccttcaagcctaacccaatcaattacatcctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgaggagga
gaatcctatccggctgatcgatctgggcgtgagagtgtttgagagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacggagc
gtgaggcgcctgacacggagaagggcacacaggctgctgagggcacgccggctgctgaagagagagaggcgtgctgcaggcagcagacttcgatgagaatg
gcctgatcaagagcctgccaaacacccctggcagctgagagcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacctgat
caagcaccgcggctacctgagccagcggaagaacgagggagagacagcagacaaggagctgggcgccctgctgaagggagtggccaacaatgcccacgcc
ctgcagaccggcgatttcaggacacctgccgagctggccctgaataagtttgagaaggagtccggccacatcagaaaccagagggggcgactatagccacacc
ttctcccgcaaggatctgcaggccgagctgatcctgctgttcgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggagggaatcgag
accctgctgatgacacagaggcctgccctgtccggcgacgcagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaagaac
acctacacagccgagcggtttatctggctgacaaagctgaacaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagagagggccacc
ctgatggatgagccttaccggaagtctaagctgacatatgcccaggccagaaagctgctgggcctggaggacaccgcctttctttaagggcctgagatacggca
aggataatgccgaggcctccacactgatggagatgaaggcctatcacgccatctctcgcgccctggagaaggagaggcctgaaggacaagaagtcccccctga
acctgagctccgagctgcaggatgagatcggcaccgccttctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacagggtgcagcctgagat
cctggaggccctgctgaagcacatctctttcgataagtttgtgcagatcagcctgaaggccctgagaaggatcgtgccactgatggagcagggcaagcggtac
gacgaggcctgcgccgagatctacggcgatcactatggcaagaagaacacagaggagaagatctatctgccccctatccctgccgacgagatcagaaatcct
gtggtgctgagggccctgtcccaggcaagaaaagtgatcaacggagtggtgcgccggtacggatctccagcccggatccacatcgagaccgccagagaagtg
ggcaagagcttcaaggaccggaaggagatcgagaagagacaggaggagaatcgcaaggatcgggagaaggccgccgccaagtttagggagtactttcccta
actttgtgggcgagccaaagtctaaggacatcctgaagctgcgcctgtacgagcagcagcacggcaagtgtctgtatagcggcaaggagatcaatctggtgcg
gctgaacgagaagggctatgtggagatcgatcacgccctgcctttctccagaacctgggacgattcttttaacaataaggtgctggtgctgggcagcgagaacc
agaataagggcaatcagacaccatacgagtatttcaatggcaaggacaactccagggagtggcaggagttcaaggcccgcgtggagacctctagatttccca
ggagcaagaagcagcggatcctgctgcagaagttcgacgaggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgccagtt
tgtggccgatcacatcctgctgaccggcaagggcaagagaaagggtgttcgcctctaatggccagatcacaaacctgctgagggggattttggggactgaggaag
gtgcgggcagagaatgacagacaccacgcactggatgcagtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgaggtataa
ggagatgaacgcctttgacggcaagaccatcgataaggagacaggcaaggtgctgcaccagaagacccacttcccccagccttgggagttctttgcccagga
agtgatgatccgggtgttcggcaagccagacggcaagcctgagtttgaggaggccgatacccagagaagctgaggacactgctggcagagaagctgtctag
caggccagaggcagtgcacgagtacgtgaccccactgttcgtgtccagggcacccaatcggaagatgtctggcgcccacaaggacacactgagaagcgccaa
gaggtttgtgaagcacaacgagaagatctccgtgaagagagtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaacggcag
ggagatcgagctgtatgaggccctgaaggcaaggctggaggcctacggaggaaatgccaagcaggccttcgacccaaaggataaccccttttataagaaggg
aggacagctggtgaaggccgtgtgcgggtggagaagacccaggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcgatatgg
tgagagtggacgtgttctgtaaggtggataagaagggcaagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatc
gattgcaagggctacagaatcgacgatagctatacattctgtttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtggagtt t
gcctactatatcaattgcgactcctctaacggcaggttctacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccaccccagaatctggt
gctgatccagaagtatcaggtgaacgagctgggcaaggagatcaggccATGTCGGCTGAAGAAGCGCCCacccgtgcggTAAAATAAAAGAT
CTTTATTTTCATTAGATCTGTGTGTTGGTTTTTTGTGTAAGCgcggccgcaggaacccctagtgatggagttggccactccctctctgcgcgct
cgctcgctcactgaggccgggcgaccaaaggtcgcccgacgcccgggctttgcccgggcggcctcagtgagcgagcgagcgcgcagctgcctgcagg

Single-AAV vector with 2xBP_SV40 NLS in **Figure 2d, 3a, and Supplementary Figure 3b**
Legend: ITR, U1a promoter, BP_SV40 NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, U6 promoter,
guide sequence, guide RNA scaffold, U6 termination signal

Cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggcgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgc
gcagagagggagtggccaactccatcactagggg ttcctgcggcctctagaGTTTAAACAAAAAAA
TAAACGATGCCCCTTAAAGCAGAAGCTTTAAGGGGGCAGAGCGTTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAAC
GTTTCGGGAGCTACAACNNNNNNNNNNNNNNNNNNNNNNNNNNNNNGGTGTTTCGTCCTTTCCACAAGATATATAAAGCCA
AGAAATCGAAATACTTTCAAGTTACGGTAAGCATATGATAGTCCATTTTAAAACATAATTTTAAAACTGCAAACTACCCAA
GAAATTATTACTTTCTACGTCACGTATTTTGTACTAATATCTTTGTGTTTACAGTCAAATTAATTCTAATTATCTCTCTAACA
GCCTTGTATCGTATATGCAAATATGAAGGAATCATGGGAAATAGGCCCTCTCTAGAatggaggcggtactatgtagatgagaattca
ggagcaaactgggaaaagcaactgcttccaaatatttgtgattttttacagtgtagttttggaaaaactcttagcctaccaattcttctaagtgttttaaaatgtggg
agccagtacacatgaagttatagagtgttttaatgaggcttaaatatttaccgtaactatgaaatgctacgcatatcatgctgttcaggctccgtggccacgcaa
ctcatactACCGGTGCCACCATGaaaagaaccgccgacggcagcgaattcgagcccaagaagaagaggaaagtcGGCGGTAGCGGCGGAG
GCAGCGGTGGCGGCAGCGGCTctgaggtggagtttтcccacgagtactggatgagacatgccctgaccctggccaagagggcacgcgatgagagg
gaggtgcctgtgggagccgtgctggtgctgaacaatagagtgatcggcgagggctggaacagagccatcggcctgcacgacccaacagcccatgccgaaatt
atggccctgagacagggcggcctggtcatgcagaactacagactgattgacgccaccctgtacgtgacattcgagccttgcgtgatgtgcgccggcgccatgat
ccactctaggatcggccgcgtggtgtttggcgtgaggaacAGCaaaCGGggcgccgcaggctccctgatgAacgtgctgAactaccccggcatgaatcacc
gcgtcgaaattaccgagggaatcctggcagatgaatgtgccgccctgctgtgcGACttctACcggatgcctagacaggtgttcaatgctcagaagaaggccc
agagctccaTcAactccggaggatctagcggaggctcctctggctctgagacacctggcacaagcgagagcgcaacacctgaaagcagcggggggcagcagc
ggggggtcaatggccgccttcaagcctaacccaatcaattacatcctgggactggccatcggaatcgcatccgtgggatgggctatggtggagatcgacgagg
aggagaatcctatccggctgatcgatctgggcgtgagagtgtttgagagggccgaggtgccaaagaccggcgattctctggctatggcccggagactggcacg
gagcgtgaggcgcctgacacggagaagggcacacaggctgctgagggcacgccggctgctgaagagagagaggcgtgctgcaggcagcagacttcgatgag
aatggcctgatcaagagcctgccaaacacccctggcagctgagagcagccgccctggacaggaagctgacaccactggagtggtctgccgtgctgctgcacc
tgatcaagcaccgcggctacctgagccagcggaagaacgagggagagacagcagacaaggagctgggcgccctgctgaagggagtggccaacaatgccca
cgccctgcagaccggcgatttcaggacacctgccgagctggccctgaataagtttgagaaggagtccggccacatcagaaaccagagggggcgactatagcca
caccttctccccgcaaggatctgcaggccgagctgatcctgctgttcgagaagcagaaggagtttggcaatccacacgtgagcggaggcctgaaggagggaatc
gagaccctgctgatgacacagaggcctgccctgtccggcgacgcagtgcagaagatgctgggacactgcaccttcgagcctgcagagccaaaggccgccaag
aacacctacacagccgagcggtttatctggctgacaaagctgaacaatctgagaatcctggagcagggatccgagaggccactgaccgacacagagaggc
caccctgatggatgagccttaccggaagtctaagctgacatatgcccaggccagaaagctgctgggcctggaggacaccgccttctttaagggcctgagatac
ggcaaggataatgccgaggcctccacactgatggagatgaaggcctatcacgccatctctcgcgccctggagaaggagaggcctgaaggacaagaagtccccc
ctgaacctgagctccgagctgcaggatgagatcggcaccgccttctctctgtttaagaccgacgaggatatcacaggccgcctgaaggacagggtgcagcctg
agatcctggaggccctgctgaagcacatctctttcgataagtttgtgcagatcagcctgaaggcccctgagaaggatcgtgccactgatggagcagggcaagcg
gtacgacgaggcctgcgccgagatctacggcgatcactatggcaagaagaacacagaggagaagatctatctgcccccctatccctgccgacgagatcagaaa
tcctgtggtgctgagggccctgtcccaggcaagaaaagtgatcaacggagtggtgcgccggtacggatctccagcccggatccacatcgagaccgccagaga
agtgggcaagagcttcaaggaccggaaggagatcgagaagagacaggaggagaatcgcaaggatcgggagaaggccgccgccaagtttagggagtacttc
cctaactttgtgggcgagccaaagtctaaggacatcctgaagctgcgcctgtacgagcagcagcacggcaagtgtctgtatagcggcaaggagatcaatctgg
tgcggctgaacgagaagggctatgtggagatcgatcacgccctgcctttctccagaacctgggacgattcttttaacaataaggtgctggtgctgggcagcgag
aaccagaataagggcaatcagacaccatacgagtatttcaatggcaaggacaactccaggagtggcaggagttcaaggcccgcgtggagacctctagattt
cccaggagcaagaagcagcggatcctgctgcagaagttcgacgaggatggctttaaggagtgcaacctgaatgacaccagatacgtgaaccggttcctgtgc
cagtttgtggccgatcacatcctgctgaccggcaagggcaagagaaggtgttcgcctctaatggccagatcacaaacctgctgaggggatttttggggactga
ggaaggtgcgggcagagaatgacagacaccacgcactggatgcagtggtggtggcatgcagcaccgtggcaatgcagcagaagatcacaagattcgtgagg
tataaggagatgaacgcctttgacggcaagaccatcgataaggagacaggcaaggtgctgcaccagaagacccacttcccccagccttgggagttctttgccc
aggaagtgatgatccggggtgttcggcaagccagacggcaagcctgagtttgaggaggccgatacccagagaagctgaggacactgctggcagagaagctg
tctagcaggccagaggcagtgcacgagtacgtgaccccactgttcgtgtccagggcacccaatcggaagatgtctggcgcccacaaggacacactgagaagc
gccaagaggtttgtgaagcacaacgagaagatctccgtgaagagagtgtggctgaccgagatcaagctggccgatctggagaacatggtgaattacaagaac
ggcagggagatcgagctgtatgaggccctgaaggcaaggctggaggcctacgaggaaatgccaagcaggccttcgacccaaaggataaccccttttataag
aagggaggacagctggtgaaggccgtgcgggtggagaagacccaggagagcggcgtgctgctgaataagaagaacgcctacacaatcgccgacaatggcg
atatggtgagagtggacgtgttctgtaaggtggataagaagggcaagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgcca
gacatcgattgcaagggctacagaatcgacgatagctatacattctgtttttttccctgcacaagtatgacctgatcgccttccagaaggatgagaagtccaaggtg
gagtttgcctactatatcaattgcgactcctctaacggcaggttctacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaat
ctggtgctgatccagaagtatcaggtgaacgagctgggcaaggagatcaggccATGTCGGCTGAAGAAGCGCCCCacccgtgcggGAGGATaaa
agaaccgccgacggcagcgaattcgagcccaagaagaagaggaaagtctaaAATAAAAGATCTTTATTTTCATTAGATCTGTGTGTTGGTT

TTTTGTGTAAGCgcggccgcaggaaccccctagtgatggagttggccactccctctctgcgcgctcgctcgctcactgaggccgggcgaccaaaggtcgccc
gacgcccgggctttgcccgggcggcctcagtgagcgagcgagcgcgcagctgcctgcagg

Single-AAV vector with miniU6 promoter in **Figure 3c**
Legend: ITR, U1a promoter, BP_SV40 NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, miniU6
promoter, **guide sequence**, guide RNA scaffold, U6 termination signal
Cctgcaggcagctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggcgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgc
gcagagagggagtggccaactccatcactaggggttcctgcggcctctagaGTTTAAACAAAAAAA
TAAACGATGCCCCCTTAAAGCAGAAGCTTTTAAGGGGCAGAGCGTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAAC
GTTTCGGGAGCTACAACNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNGGTGTTTCGTCCTTTCCACAAGATATATAAAGCCA
AGAAATCGAAATACTTTCAAGTTACGGTAAGCTATCGTATATGCAAATATGAAGGAATCATGGGAAATAGGCCCTCTCTA
Gaatggaggcggtactatgtagatgagaattcaggagcaaactgggaaaagcaactgcttccaaatatttgtgattttttacagtgtagtttttggaaaaactctt
agcctaccaattcttctaagtgtttttaaaatgtgggagccagtacacatgaagttatagagtgtttttaatgaggcttaaatatttaccgtaactatgaaatgctac
gcatatcatgctgttcaggctccgtggccacgcaactcatactACCGGTGCCACCATGgaaaagaaccgccgacggcagcgaattcgagcccaagaaga
agaggaaagtcGGCGGTAGCGGCGGAGGCAGCGGTGGCGGCAGCGGCTctgaggtggagtttttcccacgagtactggatgagacatgccc
tgaccctggccaagagggcacgcgatgagaggaggtgcctgtgggagccgtgctggtgctgaacaatagagtgatcggcgagggctggaacagagccatc
ggcctgcacgacccaacagcccatgccgaaattatggcccctgagacagggcggcctggtcatgcagaactacagactgattgacgccaccctgtacgtgacat
tcgagccttgcgtgatgtgcgccggcgccatgatccactctaggatcggccgcgtggtgtttggcgtgaggaacAGCaaaCGGgcgccgcaggctccctga
tgAacgtgctgAactacccggcatgaatcaccgcgtcgaaattaccgaggaatcctggcagatgaatgtgccgccctgctgtgcGACttctACcggatgc
ctagacaggtgttcaatgctcagaagaaggcccagagctccaTcAactccgaggatctagcggaggctcctctggctctgagacacctggcacaagcgaga
gcgcaacacctgaaagcagcggggggcagcagcggggggtcaatggccgccttcaagcctaacccaatcaattacatcctgggactggccatcggaatcgcat
ccgtgggatgggctatggtggagatcgacgaggaggagaatcctatccggctgatcgatctgggcgtgagagtgtttgagagggccgaggtgccaaagaccg
gcgattctctggctatggcccggagactggcacggagcgtgaggcgcctgacacggagaagggcacacaggctgctgagggcacgccggctgctgaagaga
gagggcgtgctgcaggcagcagacttcgatgagaatggcctgatcaagagcctgccaaacacccctggcagctgagagcagccgccctggacaggaagct
gacaccactggagtggtctgccgtgctgctgcacctgatcaagcaccgcggctacctgagccagcggaagaacgagggagagacagcagacaaggagctgg
gcgccctgctgaagggagtggccaacaatgcccacgccctgcagaccggcgatttcaggacacctgccgagctggcccctgaataagtttgagaaggagtccg
gccacatcagaaaccagagggcgactatagccacaccttctcccgcaaggatctgcaggccgagctgatcctgctgttcgagaagcagaaggagtttggca
atccacacgtgagcggaggcctgaaggagggaatcgagaccctgctgatgacacagaggcctgccctgtccggcgacgcagtgcagaagatgctgggacact
gcaccttcgagcctgcagagccaaaggccgccaagaacacctacacagccgagcggtttatctggctgacaaagctgaacaatctgagaatcctggagcagg
gatccgagaggccactgaccgacacagagagggccaccctgatggatgagcccttaccggaagtctaagctgacatatgcccaggccagaaagctgctgggcc
tggaggacaccgccttctttaagggcctgagatacggcaaggataatgccgaggcctccacactgatggagatgaaggcctatcacgccatctctcgcgccctg
gagaaggagggcctgaaggacaagaagtccccccctgaacctgagctccgagctgcaggatgagatcggcaccgccttctctctgtttaagaccgacgaggat
atcacaggccgcctgaaggacagggtgcagcctgagatcctggaggccctgctgaagcacatctctcttcgataagtttgtgcagatcagcctgaaggccctgag
aaggatcgtgccactgatggagcagggcaagcggtacgacgaggcctgcgccgagatctacggcgatcactatggcaagaagaacacagaggagaagatct
atctgccccctatccctgccgacgagatcagaaatcctgtggtgctgagggccctgtcccaggcaagaaaagtgatcaacggagtggtgcgccggtacggatct
ccagcccggatccacatcgagaccgccagagaagtgggcaagagcttcaaggaccggaaggagatcgagaagagacaggaggagaatcgcaaggatcgg
gagaaggccgccgccaagtttagggagtacttccctaactttgtgggcgagccaaagtctaaggacatcctgaagctgcgcctgtacgagcagcagcacggca
agtgtctgtatagcggcaaggagatcaatctggtgcggctgaacgagaagggctatgtggagatcgatcacgccctgcctttctccagaacctgggacgattct
tttaacaataaggtgctggtgctgggcagcgagaaccagaataagggcaatcagacaccatacgagtatttcaatggcaaggacaactccaggggagtggcag
gagttcaaggcccgcgtggagacctctagatttcccaggagcaagaagcagcggatcctgctgcagaagttcgacgaggatggctttaaggagtgcaacctga
atgacaccagatacgtgaaccggttcctgtgccagtttgtggccgatcacatcctgctgaccggcaaggggcaagagaagggtgttcgcctctaatggccagatc
acaaacctgctgaggggattttggggactgaggaaggtgcgggcagagaatgacagacaccacgcactggatgcagtggtggtggcatgcagcaccgtggc
aatgcagcagaagatcacaagattcgtgaggtataaggagatgaacgcctttgacggcaagaccatcgataaggagacaggcaaggtgctgcaccagaaga
cccacttcccccagccttgggagttctttgcccaggaagtgatgatccgggtgttcggcaagccagacggcaagcctgagtttgaggaggccgatacccccagag
aagctgaggacactgctggcagagaagctgtctagcaggccagaggcagtgcacgagtacgtgaccccactgttcgtgtccagggcacccaatcggaagatg
tctggcgcccacaaggacacactgaagaagcgccaagagagagtttgtgaagcacaacgagaagatctccgtgaagagagtgttggctgaccgagatcaagctggc
cgatctggagaacatggtgaattacaagaacggcagggagatcgagctgtatgaggccctgaaggcaaggctggaggcctacggaggaaatgccaagcagg
ccttcgacccaaaggataaccccctttataagaagggaggacagctggtgaaggccgtgcgggtggagaagacccaggagagcggcgtgctgctgaataag
aagaacgcctacacaatcgccgacaatggcgatatggtgagagtggacgtgttctgtaaggtggataagaacggcaagaatcagtactttatcgtgcctatct

AAV-Nm2-ABE8e-U6_V2 in **Figure 6**
Legend: ITR, U1a promoter, BP_SV40 NLS, TadA8e, linker, Nme2Cas9-D16A, short poly-A signal, U6 promoter,
guide sequence, guide RNA scaffold, U6 termination signal

caagaatcagtactttatcgtgcctatctatgcctggcaggtggccgagaacatcctgccagacatcgattgcaagggctacagaatcgacgatagctatacat
tctgttttttccctgcacaagtatgacctgatcgcccttccagaaggatgagaagtccaaggtggagtttgcctactatatcaattgcgactcctctaacggcaggtt
ctacctggcctggcacgataagggcagcaaggagcagcagtttcgcatctccacccagaatctggtgctgatccagaagtatcaggtgaacgagctgggcaag
gagatcaggccatgtcggctgaagaagcgcccacccgtgcggGAGGATaaaagaaccgccgacggcagcgaattcgagcccaagaagaagaggaaagt
ctaaAATAAAAGATCTTTATTTTCATTAGATCTGTGTGTTGGTTTTTTGTGTaagcgctctagagagggcctatttcccatgattccttcata
tttgcatatacgatacaaggctgttagagagataattagaattaatttgactgtaaacacaaagatattagtacaaaatacgtgacgtagaaagtaataatttct
tgggtagtttgcagttttaaaattatgttttaaaatggactatcatatgcttaccgtaacttgaaagtatttcgatttcttggctttatatatcttgtggaaaggacg
aaacaccNNNNNNNNNNNNNNNNNNNNNNNgttgtagctcccgaaacgttgctacaataaggccgtctgaaaagatgtgccgcaacgctc
tgcccccttaaagcttctgctttaaggggcatcgtttattttttttgtttaaacgcaggaacccctagtgatggagttggccactccctctctgcgcgctcgctcgctca
ctgaggccgggcgaccaaaggtcgcccgacgcccgggctttgcccgggcggcctcagtgagcgagcgagcgcgcagctgcctgcagg

AcrIIA4 plasmid in **Supplementary Figure 3**
Legend: CMV IE94 promoter, AcrIIA4, SV40 polyA signal,
GACCATAGCCAATTCAATATGGCGTATATGGACTCATGCCAATTCAATATGGTGGATCTGGACCTGTGCCAATTCAATATG
GCGTATATGGACTCGTGCCAATTCAATATGGTGGATCTGGACCCCAGCCAATTCAATATGGCGGACTTGGCACCATGCCA
ATTCAATATGGCGGACTTGGCACTGTGCCAACTGGGGAGGGGTCTACTTGGCACGGTGCCAAGTTTGAGGAGGGGTCTT
GGCCCTGTGCCAAGTCCGCCATATTGAATTGGCATGGTGCCAATAATGGCGGCCATATTGGCTATATGCCAGGATCAATA
TATAGGCAATATCCAATATGGCCCTATGCCAATATGGCTATTGGCCAGGTTCAATACTATGTATTGGCCCTATGCCATATA
GTATTCCATATATGGGTTTTCCTATTGACGTAGATAGCCCCTCCCAATGGGCGGTCCCATATACCATATATGGGGCTTCCT
AATACCGCCCATAGCCACTCCCCCATTGACGTCAATGGTCTCTATATATGGTCTTTCCTATTGACGTCATATGGGCGGTCCT
ATTGACGTATATGGCGCCTCCCCCATTGACGTCAATTACGGTAAATGGCCCGCCTGGCTCAATGCCCATTGACGTCAATAG
GACCACCCACCATTGACGTCAATGGGATGGCTCATTGCCCATTCATATCCGTTCTCACGCCCCCCTATTGACGTCAATGACG
GTAAATGGCCCACTTGGCAGTACATCAATATCTATTAATAGTAACTTGGCAAGTACATTACTATTGGAAGGACGCCAGGG
TACATTGGCAGTACTCCCATTGACGTCAATGGCGGTAAATGGCCCGCGATGGCTGCCAAGTACATCCCCATTGACGTCAA
TGGGGAGGGGCAATGACGCAAATGGGCGTTCCATTGACGTAAATGGGCGGTAGGCGTGCCTAATGGGAGGTCTATATA
AGCAATGCTCGTTTAGGGAACATGAACATTAACGACCTCATACGAGAGATTAAGAACAAAGATTACACCGTCAAACTGTC
AGGAACTGATAGTAACTCAATCACCCAGCTTATTATCAGGGTAAACAATGATGGGAATGAATATGTGATATCTGAGAGCG
AAAACGAGTCTATCGTCGAGAAATTCATTTCCGCTTTTAAGAACGGGTGGAATCAGGAATATGAGGATGAAGAAGAATTT
TACAATGACATGCAGACGATCACGTTGAAAAGTGAACTGAACTAACTCGAGCCTCTAGAACTATAGTGAGTCGTATTACG
TAGATCCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACTAGAATGCAGTGAAAAAAATGCTTTATTTGT
GAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAGTT

AcrIIC4-MRE122 plasmid in **Supplementary Figure 3b**
Legend: ITR, CMV enhancer, CB promoter, SV40 interon, AcrIIC4, FLAG, SV40 NLS, 3 x MRE122, BGh polyA, U6
promoter, **guide sequence**, guide RNA scaffold, U6 termination signal
ctgcgcgctcgctcgctcactgaggccgcccgggcaaagcccgggccgtcgggcgacctttggtcgcccggcctcagtgagcgagcgagcgcgcagagaggga
gtgtagccatgctctaggaagatcaattcggtacaattcagctagcGTTTAAACAAAAAAATAAACGATGCCCCTTAAAGCAGAAGCTTTA
AGGGGCAGAGCGTTGCGGCACATCTTTTCAGACGGCCTTATTGTAGCAACGTTTCGGGAGCTACAACNNNNNNNNNNNN
NNNNNNNNNNggtgtttcgtcctttccacaagatatataaagccaagaaatcgaaatactttcaagttacggtaagcatatgatagtccattttaaaaca
taattttaaaactgcaaactacccaagaaattattactttctacgtcacgtattttgtactaatatctttgtgtttacagtcaaattaattctaattatctctctaaca
gccttgtatcgtatatgcaaatatgaaggaatcatgggaaataggccctccgcgtcgacattgattattgactctggtcgttacataactacggtaaatggccc
gcctggctgaccgcccaacgacccgcccattgacgtcaataatgacgtatgttcccatagtaacgccaataggacttttccattgacgtcaatggtgggagtat
ttacggtaaactgcccacttggcagtacatcaagtgtatcatatgccaagtacgccccctattgacgtcaatgacggtaaatggcccgcctggcattatgcccag
tacatgaccttatgggactttcctacttggcagtacatctactcgaggccacgttctgcttcactctccccatctccccccccctccccacccccaattttgtatttatt
tattttttaattattttgtgcagcgatggggcggggggggggggggggcgcgcgccaggcggggcggggcggggcgaggggcggggcggggcgagg
cggagaggtgcggcggcagccaatcagagcggcgcgctccgaaagtttccttttatggcgaggcggcggcggcggcggccctataaaaagcgaagcgcgcg
gcgggcgggagcgggatcagccaccgcggtggcggcctagagtcgacgaggaactgaaaaccagaaagttaactggtaagtttagtctttttgtcttttattt
caggtccccggatccggtggtggtgcaaatcaaagaactgctcctcagtggatgttgcctttacttctaggcctgtacggaagtgttacttctgctctaaaagctgc
ggaattgtacccgcggccgatccaCCGGTgccgccaccATGAAGATCACCAGCAGCAACTTCGCGACCATTGCGACCAGCGAGAACT

TTGCGAAGCTGAGCGTGCTGCCGAAAAACCACCGTGAGCCGATCAAGGGTCTGTTCAAAAGCGCGGTTGAACAGTTTAG
CAGCGCGCGTGACTTCTTTAAGAACGAGAACTACAGCAAAGAGCTGGCGGAAAAGTTCAACAAAGAAGCGGTGAACGA
GGCGGTTGAAAAGCTGCAAAAAGCGATCGATCTGGCGGAAAAACAGGGCATTCAATTTggcggggattataaggatgatgatgat
aagggctccCCTAAGAAGAAGAGAAAGGTGTGATAAAGCGGCCATCAAGCTtGACAAACACCATTGTCACACTCCAACAAAC
ACCATTGTCACACTCCAACAAACACCATTGTCACACTCCACCATAgAGCTcgctgatcagcctcgactgtgccttctagttgccagccatct
gttgtttgcccctcccccgtgccttccttgaccctggaaggtgccactcccactgtcctttcctaataaaatgaggaaattgcatcgcattgtctgagtaggtgtcat
tctattctggggggtggggtggggcaggacagcaaggggggaggattgggaagacaattaggtagataagtagcatggcgggttaatcattaactacaaggaa
cccctagtgatggagttggccactccctctctgcgcgctcgctcgctcactgaggccgggcgaccaaaggtcgcccgacgcccgggctttgcccgggcggcctc
agtgagcgagcgagcgcgcag