The *Blepharisma stoltei* macronuclear genome: towards the origins of whole genome reorganization

Minakshi Singh[1], Kwee Boon Brandon Seah[1], Christiane Emmerich[1], Aditi Singh[1], Christian Woehle[2], Bruno Huettel[2], Adam Byerly[3], Naomi Alexandra Stover[4], Mayumi Sugiura[5], Terue Harumoto[5], Estienne Carl Swart[1,*]

1. Max Planck Institute for Biology, Tuebingen, Germany
2. Max Planck Genome Center Cologne, Max Planck Institute for Plant Breeding, Cologne, Germany
3. Department of Computer Science and Information Systems, Bradley University, Peoria IL, USA
4. Department of Biology, Bradley University, Peoria, IL, USA
5. Nara Women's University, Nara, Japan
* Corresponding author

Abstract

Both the function and origin of genome reorganization, in which germline genomes are transformed into somatic genomes, are great biological mysteries. In ciliates, ubiquitous microbial eukaryotes with both somatic and germline nuclei in the same cell, tens of thousands of internally eliminated sequences (IESs) scattered throughout the genome are excised. *Blepharisma* represents one of the two earliest diverging ciliate classes, and has unique, dual pathways of somatic nuclear development, making it ideal for investigating the functioning, origins and evolution of these processes. Here, we report the somatic genome assembly of *Blepharisma stoltei* strain ATCC 30299 (41 Mb), arranged as numerous alternative telomere-capped minichromosomes, tens to hundreds of kilobases long. This genome encodes eight PiggyBac transposase homologs liberated from transposons. All are subject to purifying selection, but just one, the putative IES excisase, has a complete catalytic amino acid triad. We proposed that PiggyBac homologs may have been ancestral excisases that enabled extensive genome reorganization.

Abvreviations

- MIC - micronucleus
- MAC - macronucleus
- IES - interspersed eliminated sequence
- MDS - macronuclear-destined sequence
- PacBio - Pacific Biosciences
- CLR - continuous long read (PacBio)
- CCS - circular consensus sequence (PacBio)
- HiFi - High-fidelity read (PacBio)
- ATAS - alternative telomere addition site
- PBLE - PiggyBac-like element
- PGBD - PiggyBac element-derived
- Pgm - PiggyMac
- PgmL - PiggyMac-like

Introduction

Acquisition of new organelles defines and is responsible for the origins of eukaryotes [1]. Ciliates represent one of a couple of eukaryotic lineages with more than one kind of nucleus within individual cells [2]. Profound differences between the ciliate nuclei and the genomes they contain now exist, with some parallels to the germline-soma distinction in multicellular organisms [3]. Ciliate nuclear and genomic dimorphism provides a special opportunity to observe the consequences of evolution of a specialized organelle clearly derived from a general one and for studying the evolution of new, functionally differentiated genes that enable this unusual situation.

The processes responsible for large-scale developmental DNA elimination [4] in ciliate nuclei are under active study, but what the responsible ancestral molecules were remains to be determined. Knowledge of the molecules responsible for ciliate genome reorganization is dominated by *Tetrahymena* and *Paramecium* (class Oligohymenophorea), with additional input from *Oxytricha*, *Stylonychia* and *Euplotes* (class Spirotrichea) [5,6], whereas this remains to be investigated in the remaining nine or so classes. To gain fresh insights into genome reorganization and tackle questions about its origin we focused on the ciliate species *Blepharisma stoltei*. Together with its sister-class, Karyorelictea, the class Heterotrichea, to which this *B. stoltei* belongs, represent the earliest branching ciliate lineages, more distantly related to current model ciliates than those models are to each other [7]. Furthermore, *Blepharisma* exhibits distinctive alternative somatic developmental pathways, which have potential utility in disentangling processes involved in genome reorganization from preceding ones that may indirectly influence it.

Blepharisma is a distinctive genus of single-celled ciliates (Figure 1) known for the red, light-sensitive pigment, blepharismin, in the sub-pellicular membranes of representative species [8],

and unusual nuclear/developmental biology (Figure 2) [9]. To date molecular investigations and

genomics of ciliates have predominantly focused on oligohymenophoreans and spirotrichs

(Figure 3, Table S1). In recent years, publication of a draft genome for the heterotrich ciliate,

Stentor, has facilitated revival of this genus for investigations of cellular regeneration [10–12].

However, significant hurdles still need to be overcome to investigate genome reorganization in

Stentor coeruleus since requisite cell mating has not been observed in the reference somatic

genome strain (personal communication, Mark Slabodnick), and very high lethality has been

reported for other strains in which mating occurred [13]. We therefore focused on *Blepharisma*

which is amenable to such investigations, with controlled induction of mating, and, critically,

procedures for investigating cellular and nuclear development have been established from more

than a century of meticulous cytological research [8,14–25].

The *Blepharisma stoltei* strains used in the present study were originally isolated in Germany

(strain ATCC 30299) and Japan (strain HT-IV), with the former continuously cultured for over

fifty years, and the latter for over a decade. The cells are comparatively straightforward to

maintain, e.g., stable cultures can be established in a simple salt medium on a few grains of rice

passaged every few months. Due to their distinctive pigmentation and large size several

Blepharisma species are excellent subjects for introducing cell biology concepts to non-

specialists, and are thus readily available for educational purposes from commercial suppliers

like Carolina Biological Supply Company (USA). They are ideal subjects for behavioral and

developmental investigations, e.g., as voracious predators of smaller ciliates and other

unicellular species, and also exhibit pronounced phenotypic plasticity, including forming cysts

and giant, cannibal cells under suitable conditions [8].

Like all ciliates [3], *Blepharisma* cells have two types of nuclei: a macronucleus (MAC) which is

very large and transcriptionally active during vegetative growth, and a small, generally

transcriptionally inactive micronucleus (MIC), which serves as the germline. Each *Blepharisma*

cell has one MAC and several MICs (Figure 1A, B). In vegetative propagation (asexual

replication) of *Blepharisma*, cell fission results in about half of the MAC pinching off which is

then distributed to each of the resulting daughter cells together with the mitotically divided MICs.

Upon starvation, *Blepharisma* cells, like other ciliates, are also capable of conjugation and the

associated sexual processes. Essential for developmental investigations, the intricate ballet of

nuclear movements and morphological changes occurring during *Blepharisma* conjugation is

well-documented [9] (Figure 2). During this process about half of the MICs in each of the cells

undergo meiosis (meiotic MICs) and the rest do not (somatic MICs) (Figure 2B). One of the

meiotic MICs eventually gives rise to two haploid gametic nuclei. One gametic MIC (the

migratory nucleus) from each conjugating cell is exchanged with that of its partner. In parallel in

partnered cells, subsequent fusion of the migratory and stationary haploid nuclei generates a

zygotic nucleus (synkaryon), and after successive mitotic divisions gives rise to both new MICs

and new MACs (known as anlagen). The new MACs continue to mature, eventually growing in

size and DNA content [9].

Conveniently for investigations of development and genome reorganization, *Blepharisma* is one

of only two ciliate genera, along with *Euplotes* [26–29], where conjugation has been shown to be

mediated through pheromone-like substances called gamones. *Blepharisma* has two mating

types, distinguished by their gamone production. Mating type I cells release gamone 1, a ~30

kDa tyrosine-rich [30] glycoprotein [31,32]; mating type II cells release gamone 2, calcium-3-(2'-

formylamino-5'-hydroxybenzoyl) lactate, a small-molecule effector [33]. *Blepharisma* cells

commit to conjugation when complementary mating types recognize each other's gamones, with

the cells remaining paired while meiosis and fertilization occur and eventually new MACs begin

to form.

As in model ciliates, during *Blepharisma* anlagen development, MIC-specific sequences are

removed from a MIC genome copy to form a functional MAC genome (Seah, et al. in prep.). The

new MAC genomes of model ciliate species are largely free of mobile elements and other forms

of "junk" DNA contained in the MIC genome [34]. In the best studied ciliates, a form of genome

reorganization assisted by small RNAs (sRNAs) is thought to occur [5]. Specific genome

segments limited to the MIC, termed internally eliminated sequences (IESs), are excised by

domesticated transposases, and may have originated from transposons [3,5,35,36]. Large scale

genome-wide DNA amplification accompanies the genome reorganization process, producing

thousands of copies in mature MACs of larger ciliate species [3,34].

We were motivated to investigate genome reorganization in *Blepharisma*, as, unlike model

ciliates, these cells can produce two kinds of anlagen, and particularly because one of the two

pathways of their development skips the complex series of mitoses, meioses, nuclear

exchanges and fertilization [9] (Figure 2). Primary anlagen mature in the conventional manner

from zygotic nuclei. Somatic MICs which have not undergone meiosis can give rise to secondary

anlagen, which can develop into mature macronuclei [9]. This occurs frequently in strains with a

high frequency of selfing (conjugation among cells within a clonal population), in preference to

development of primary macronuclear anlagen [9]. This alternative pathway of MAC

development has also been observed experimentally after removal of primary macronuclear

anlagen by microsurgery [9]. As conjugation progresses, the old (maternal) MACs are

progressively degraded [9]. Since the *B. stoltei* MIC genome has numerous IESs which interrupt

genes (Seah et al. in prep.), in principle, to produce functional, mature MAC genomes, it is

essential that reorganization of DNA occurs in both primary and secondary anlagen.

Here we provide essential macronuclear genome and developmental, transcriptomic resources

for *B. stoltei* and present the first investigations of possible molecules involved in its genome

reorganization. Like *Stentor coeruleus* [10], *Blepharisma stoltei* genes have the shortest known

spliceosomal introns, predominantly 15 or 16 nt long. From long-read sequencing the *B. stoltei*

MAC genome appears to be organized in the form of numerous minichromosomes. Among the

MAC-encoded transposase genes we identified in *Blepharisma* were PiggyBac transposase

homologs, which, as far as we are aware, have not been reported in any ciliates other than

Paramecium and *Tetrahymena*. A few *Blepharisma* PiggyBac homologs are substantially

upregulated in MAC development, including one with a complete catalytic triad which is the main

candidate IES excisase. Consistent with ancient origins of genome reorganization in ciliates,

Blepharisma shares pronounced development-specific upregulation of homologs known to be

involved in this process, notably Dicer-like and Piwi proteins responsible for sRNA biogenesis.

Blepharisma therefore represents an invaluable outgroup for investigations of the evolution of

genome reorganization.

161 Results

162 A compact, extensively fragmented genome

The draft *Blepharisma stoltei* ATCC 30299 MAC genome is compact, at 41 Mb, and AT rich

(66%), like most sequenced ciliate MAC genomes, and relatively complete (Figure 3; Table S1,

2, Figure S1). From joint variant calling of reads from strains ATCC 30299 and HT-IV mapped to

this assembly, strain ATCC 30299 appears to be virtually homozygous, with only 1277

heterozygous single-nucleotide polymorphisms (SNPs) observed compared to 193725 in strain

HT-IV (i.e., individual heterozygosity of 3.08×10^{-5} vs. 4.67×10^{-3} respectively). Low SNP levels

were likely beneficial for the overall assembly contiguity, since heterozygosity poses significant

algorithmic challenges for genome assembly software [37]. For brevity's sake, we refer to this

draft genome as the *Blepharisma* MAC genome (and likewise "*Blepharisma*" for the associated

strain). Though the final assembly comprises 64 telomere-to-telomere sequences, it is not

possible to define conventional chromosome ends given the extensive natural fragmentation of

the *Blepharisma* MAC genome (characterized in the next section), hence we refer to them

simply as "contigs".

The basic telomere unit of *B. stoltei* is a permutation of CCCTAACA, like its heterotrich relative

Stentor coeruleus [10] (Figure S2). However, a compelling candidate for a telomerase ncRNA

(TERC) could not be found in either *Blepharisma* or *Stentor* using Infernal [38] and RFAM

models (RF00025 - ciliate TERC; RF00024 - vertebrate TERC). Thus, it was not possible to

delimit the ends of the repeat. Heterotrichs may use a different kind of ncRNA, or one which is

very divergent from known TERCs. In contrast to the extremely short (20 bp) macronuclear

telomeres of spirotrichs like *Oxytricha* with extreme MAC genome fragmentation [39],

sequenced *Blepharisma* macronuclear telomeres are moderately long (Figure S2A), with a

mode of 209 bp (i.e., 26 repeats of the 8 bp motif), extending to a few kilobases.

Approximately one in eight reads in the *Blepharisma* HiFi library were telomere-bearing and

distributed across the entire genome (Figure 4A), using a moderately strict definition of

possessing at least three consecutive telomeric repeats. In contrast, the telomere-bearing reads

of the model ciliate *Tetrahymena thermophila* predominantly map to chromosome ends (Figure

S3A), and only one in fifty nine *Tetrahymena* CLR reads are telomere-bearing, as identified by

three consecutive telomeric subunit repeats (i.e., 3×CCCCAA).

Typically, only a minority of mapped reads are telomere-bearing at individual internal contig

positions, and so we term them alternative telomere addition sites (ATASs) (Figure 4A). We

identified a total of 46705 potential ATASs in *Blepharisma* by mapping PacBio HiFi reads onto

the MAC reference, and searching for partially mapped reads that contained telomeric repeats in

the unmapped (clipped) segments without any intervening sequence between the mapped

segment and the beginning/end of the telomeric repeat. The majority of these sites (38686) were

represented by only one mapped HiFi read.

The expected distance between telomeres, and hence the average MAC DNA molecule length,

is about 130 kb. This is consistent with the raw input MAC DNA lengths, which were mostly

longer than 10 kb and as long as 1.5 Mb (Figure S3C, D), and our observation that a small

fraction (1.3%) of *Blepharisma*'s HiFi reads are bound by telomeres on both ends. Excluding the

length of the telomeres, these telomere-bound reads may be as short as 4 kb (Figure S2B).

Given the frequency of telomere-bearing reads, we expect many additional two-telomere DNA

molecules longer than 12 kb, the approximate maximum length of the HiFi reads (Figure S3C,

D).

Since the lengths of the sequenced two-telomere DNA molecules on average imply that they

encode multiple genes, we propose classifying them as "minichromosomes". This places them

between the length of "nanochromosomes" of ciliates like *Oxytricha* and *Stylonychia*, which

typically encode single genes and are on the order of a few kilobases long [39,40], and MAC

chromosomes of *Paramecium tetraurelia* and *Tetrahymena thermophila* which are hundreds of

kilobases to megabases long [41–43]. Recently it was reported that the *Paramecium bursaria*

MAC genome is considerably more fragmented than those of other previously examined

Paramecium species [44]. The DNA molecules from *P. bursaria* have thus also been classified

as minichromosomes [44].

Alternative telomere addition sites in the MAC genome tend to be intergenic in model ciliates like

Oxytricha trifallax [39]. In *Blepharisma*, we found more intergenic ATASs (28309) than intragenic

ones (18396). As intergenic regions only make up 10.1 Mb of the assembly, the intergenic

frequency of ATASs is about five-fold higher (2.81 per 1 kb) than intragenic frequency (0.562 per

1 kb). The presence of intragenic ATASs raises the question how the cell tolerates or deals with

mRNAs encoding partial proteins transcribed from 3' truncated genes. Since the sequence data

was from a clonal population, it is not possible to tell how much ATAS variability there is within

individual cells. However, it is conceivable that their positional variation in single cells reflects

that of the population. In this case, together with redundancy from massive DNA amplification

there would likely be sufficient intact copies of every gene.

Beyond the first 2-5 bp corresponding to the junction sequences, the average base composition

on the chromosome flanking ATAS junctions shows an asymmetrical bias (Figure 4B). From

position +6 onwards there is an enrichment of T to about 40% and A to 35-39%, compared to

the genome-wide frequencies of 33% each. At position +19 to +23, there is a slight decrease in

T to 37-39%. AT values gradually decline back to about 35% each by position +150.

Correspondingly, G and C are depleted downstream of ATAS junctions, dropping to a minimum

of 8.6% and 11% respectively around position +37, compared to the genome-wide average of

17% each. AT enrichment and GC depletion upstream of ATAS junctions are less pronounced.

If breakage and chromosome healing were random, we would not expect such an asymmetry.

This suggests that there is a nucleotide bias, whether in the initiation of breaks, telomere

addition, or in the processing of breaks before telomere addition. However, we have not yet

identified any conserved motif like the 15 bp chromosome breakage site (CBS) in *Tetrahymena*

[45] nor a short 10-bp sequence periodicity in base composition like in *Oxytricha trifallax* [46].

Therefore, telomere addition in *B. stoltei* appears to involve base-pairing of short segments of

about 2 bp between the telomere and chromosome, with a bias centered on the "CT" in the

telomere unit, and an asymmetrical preference for AT-rich sequences on the chromosomal side

of the junction.

Tiny spliceosomal introns

The *Blepharisma stoltei* ATCC 30299 macronuclear genome is gene-dense (25,711 predicted

genes), with short intergenic regions, tiny introns and untranslated regions (UTRs) (Figure 5A).

In contrast to *Stentor* which, unusually for ciliates, uses the standard genetic code, *B. stoltei*

uses an alternative genetic code with UGA codons reassigned from stops to tryptophan (Figure

S5).

Like *Stentor* [10], most (82%) *Blepharisma* genes have no introns. In line with genome

compactness, during our inspections we also observed numerous overlapping poly(A)-tailed

RNA-seq reads on opposite strands derived from convergently transcribed gene pairs. The

correlation of the lengths of different noncoding region classes (intergenic regions, introns and

UTRs) can be explained by them being subject to common, neutral evolutionary processes [47].

Blepharisma introns are mostly (97%) 15 or 16 nucleotides (nt) long, like those of *Stentor*

(Figure 5B). Though intron reduction (7389 introns predicted in the reference *B. stoltei* MAC

genome, i.e., 0.29 introns per gene) is not as extreme as some other microbial eukaryotes, like

Giardia lamblia [48], where almost all have been lost, both *Blepharisma* and *Stentor* have much

fewer introns relative to other ciliates (e.g., intron densities of 1.6, 2.3 and 4.8 introns per gene

in *Paramecium*, *Oxytricha* and *Tetrahymena*, respectively [49]) and to the putative, relatively

intron-rich eukaryotic common ancestor [50], along with their extreme length reduction.

Blepharisma 15 nt introns possess a characteristic branch-point "A", as would be expected in

classical models of lariat formation during mRNA splicing (Figure 5C). 16 nt introns almost

invariably have an "A" at either 10 or 11 nt downstream of the donor site (i.e., only one of 499

does not, but has "A" at 9 nt), although this is not obvious in the consensus sequence logo

because the position is variable (Figure S6D). Similarly, 17 nt introns all possess "A" at 10-12 nt

downstream of the donor site. Only a few intron bases, 5-8 and 12, of *Blepharisma*'s 15 nt

introns are relatively unconstrained (Figure 5C). This leaves little room for the presence of any

additional regulatory elements in the mRNA or underlying DNA.

279 Extensive duplications of transmembrane protein genes

A notable extended ~220 kb region encoding 53 genes belonging to a single orthologous group

(orthogroup), OG0000085 is present on Contig_1 (Figure 5A). Four additional OG0000085

genes are present at the opposite end of Contig_1, and 24 copies are found on other contigs,

often clustered together (Figure 5D). The DNA coverage across this region is lower (74×) than

the rest of Contig_1 (185×). Though there is uncertainty in the exact extent, given the sheer

volume of reads involved, the assembled sequences certainly correspond to highly repetitive

regions of the MAC genome. At the junction between the lower and higher coverage regions

more than 30 HiFi reads link the two regions of coverage, and a similar number of telomere-

bearing reads are in close proximity. At the junction we also observe at least two potential

locations of IESs, corresponding to regions that may be partially IES/partially MDS.

Large clusters of genes from particular orthogroups can be found on additional contigs (Figure

5D). In total 551 (2%) of predicted *B. stoltei* genes belong to the orthogroups with the largest

clusters per contig. Some of the largest contiguous clusters of genes from these orthogroups are

situated at the ends of contigs, suggesting they may have caused assembly breaks beyond

them. One contig, split off from other connected components in the assembly graph,

predominantly encodes genes from a single orthogroup (contig_64, 43× coverage; Figure 5D;

Figure S7). Further increases in read length and accuracy may allow assemblers to fully resolve

these in future. Curiously, all the orthogroups corresponding to the largest contiguous clusters of

genes appear to be transmembrane proteins, or decayed remnants thereof. The nature of these

proteins is described in Supplemental Text ("Properties of proteins encoded by extensive

duplications").

302 Features of gene expression during new MAC development

To gain an overview of the molecular processes during genome reorganization in *Blepharisma*

we examined gene expression trends across a developmental time series. Complementary

mating strains of *B. stoltei* were treated with gamones of the opposite mating type, before mixing

to initiate conjugation [9,51]. Samples for morphological staging and RNA-seq were taken at

intervals from the time of mixing ("0 hour" time point) up to 38 hours (Figure S8). The

progression of nuclear morphological development and the proportion of cells in each stage at

various time points is illustrated in Figure S9.

During conjugation in *Blepharisma*, meiosis begins around 2 h after conjugating cell pairs form

and continues up to around 18 h, by when gametic nuclei generated by meiosis have been

exchanged (Figure 2A, B, Figure S9). This is followed by karyogamy and mitotic multiplication of

the zygotic nucleus (22 hours). At the 26 h time point, new, developing primary MACs can be

observed in the conjugating pairs, in the form of large, irregular bodies (Figure 2A, Figure S9).

These macronuclear anlagen mature into the new MACs of the exconjugant cell by 38 h, after

which cell division generates two daughter cells. Smaller secondary MACs, derived directly from

MICs without all the intermediate nuclear stages, can also be seen from 22 hours to 30 hours,

but eventually disappear, giving sole way to the primary MACs (Figure 2A, Figure S9).

Examining gene expression at 26 hours, when the majority of cells are forming a new MAC

(Figure S9), we observe two broad trends: relatively stable constitutive gene expression (Table

S5; Data S3), e.g., an actin homolog (ENA accession: BSTOLATCC_MAC19444) and a

bacteria-like globin protein (BSTOLATCC_MAC21846), versus pronounced development-

specific upregulation (Table S6; Data S3), e.g., a histone (BSTOLATCC_MAC21995) an HMG

box protein (BSTOLATCC_MAC14030), and a translation initiation factor (eIF4E,

BSTOLATCC_MAC5291).

We eschewed a shallow Gene Ontology (GO) enrichment analysis in favour of close scrutiny of

a smaller subset of genes strongly upregulated during new MAC formation which we expect are

more relevant to molecular biologists. For this, computational gene annotations in combination

with BLASTP searches and examination of literature associated with homologs was used.

Ranking the relative gene expression at 26 hours vs. the average expression of starved,

gamone treated, and 0 hour cells, in descending order, reveals numerous genes of interest in

the context of genome reorganization, including homologs of proteins known to be involved in

genome reorganization in model ciliate species (Table S6).

Among the top 100 genes ranked this way (69× to 825× upregulation) nine contain PFAM

transposase domains: DDE_Tnp_1_7, DDE_3, MULE and DDE_Tnp_IS1595 (e.g.,

BSTOLATCC_MAC2188, BSTOLATCC_MAC14490, BSTOLATCC_MAC18054,

BSTOLATCC_MAC18052, respectively). We also observe small RNA (sRNA) biogenesis and

transport proteins, i.e., a Piwi protein (BSTOLATCC_MAC5406) and a Dicer-like protein

(BSTOLATCC_MAC1138; see also "Supplemental text", "Homologs of small RNA-related

proteins involved in ciliate genome reorganization" and Figure S12), and a POT1 telomere-

binding protein homolog (POT1.4; BSTOLATCC_MAC1496; Supplemental Text "Telomere-

binding protein paralogs"). Numerous homologs of genes involved in DNA repair and chromatin

are also present among these highly developmentally upregulated genes ("Supplemental text",

"Development-specific upregulation of proteins associated with DNA repair and chromatin" and

"Supplemental text", "Development-specific histone variant upregulation"). The presence of

proteins involved in either transcription initiation or translation initiation among these highly

upregulated genes suggests a possible manner to coordinate the regulation of development-

specific gene expression ("Supplemental text", "Development-specific upregulation of proteins

associated with initiation of transcription and translation").

354 PiggyBac homologs in *Blepharisma* are candidate IES excisases

In *Paramecium tetraurelia* and *Tetrahymena thermophila*, PiggyBac transposases are

responsible for IES excision during genome reorganization occurring in the developing new

MAC [52,53]. These transposases appear to have been domesticated, i.e., their genes are no

longer contained in transposons but are encoded in the somatic genome where they provide a

necessary function in genome development [52,53]. PiggyBac belongs to the DDE/D-

superfamily of transposases due to the presence of a protein domain containing the catalytic

triad of 3 aspartic acid residues (DDD), where the third residue can also be a glutamate (E).

PiggyBac homologs typically have a DDD catalytic triad, rather than the more common DDE

triad of other DDE/D transposases [54]. The DDD catalytic motif is present in the PiggyMac

(Pgm) of *Paramecium* and the *Tetrahymena* PiggyBac homologs Tpb1 and Tpb2 [53,55].

Among ciliates, domesticated PiggyBac transposases have so far only been reported in the

model oligohymenophorean genera *Paramecium* and *Tetrahymena*, and have not been

detected in either the MAC or MIC genomes of the spirotrich *Oxytricha trifallax* [39,56].

We detected more transposase domains (9 distinct PFAM identifiers) in *B. stoltei* than any of the

other ciliate species we examined (Figure 6A). Seven of these domains are also found in at

least one other ciliate species. Using HMMER searches with the PFAM domain characteristic of

PiggyBac homologs, DDE_Tnp_1_7 (PF13843), we found eight homologs in *B. stoltei* ATCC,

none of which were flanked by terminal repeats identified by RepeatModeler. We also found

PiggyBac homologs in *B. stoltei* HT-IV and *B. japonicum* R1072. Reminiscent of *Paramecium*

tetraurelia, which, among ten PiggyMac homologs, has just one homolog with a complete

catalytic triad [55], the DDD triad is preserved in just a single *B. stoltei* ATCC 30299 PiggyBac

homolog (Figure 6B; Contig_49.g1063, BSTOLATCC_MAC17466). This gene is strongly

upregulated during development from 22 to 38 hours after formation of heterotypic pairs, when

new MACs develop and IES excision is required (Figure 6B). Furthermore, there are significant

similarities in the basic properties of *Blepharisma* and *Paramecium* IESs, which will be reported

in detail in the subsequent *B. stoltei* MIC genome paper (Seah et al., in prep.). Consequently,

adopting the *Paramecium* nomenclature, we refer to our primary candidate IES excisase as

Blepharisma PiggyMac (BPgm) and the other homologs as BPgm-Likes (BPgmLs).

PiggyMac homologs are also present in other heterotrich ciliates but have not yet been

described because of genome assembly or annotation challenges. Using BPgm as a query

sequence, we found convincing homologs containing the conserved catalytic DDD-motif in a

genome assembly of the heterotrichous ciliate *Condylostoma magnum* (TBLASTN e-value 2e-

24 to 2e-37). While we failed to detect the DDE_Tnp_1_7 domain in predicted genes of the

heterotrich *Stentor coeruleus*, we were able to detect a relatively weak adjacent TBLASTN

matches split across two frames in its draft MAC genome (e-value 7e-15; SteCoe_contig_741

positions 6558-5475). After joining ORFs corresponding to this region and translating them, we

obtained a more convincing DDE_Tnp_1_7 match with HMMER3 (e-value 2e-24). This either

corresponds to a pseudogene or a poorly assembled genomic region. It is also possible that

additional PiggyBac homologs were missed in unassembled *Stentor* MAC genome regions.

In addition, we searched for PiggyMac homologs in the MAC genome of the pathogenic

oligohymenophorean ciliate *Ichthyophthirius multifiliis* [57]. TBLASTN searches using the *T.*

thermophila Tpb2 as a query returned no hits. A HMMER search using hmmscan with a six-

frame translation of the *I. multifiliis* MAC genome against the PFAM-A database also did not

return any matches with independent E-values (i-E-value) less than 1. We note that based on

BUSCO analyses (Figure S1) the *I. multifiliis* genome appears to be less complete than other

ciliates we examined. So, a better genome assembly will be needed to investigate the possibility

that PiggyBac homologs are encoded elsewhere in this MAC genome.

Other than the PFAM DDE_Tnp_1_7 domain, three of the *Blepharisma* PiggyBac homologs also

possess a short, characteristic cysteine-rich domain (CRD) (Figure 6C). This domain is essential

for Pgm activity and *Paramecium* IES excision [58]. PiggyBac CRDs have been classified into

three different groups [58]. In *Blepharisma*, the CRD consists of five cysteine residues arranged

as CxxC-CxxCxxxxH-Cxxx(Y)H (where C, H, Y and x respectively denote cysteine, histidine,

tyrosine and any other residue). Two *Blepharisma* homologs possess this CRD without the

penultimate tyrosine residue, while the third contains a tyrosine residue before the final histidine.

This -YH feature towards the end of the CxxC-CxxCxxxxH-Cxxx(Y)H CRD is shared by all the

PiggyBac homologs we found in *Condylostoma*, as well as the PiggyBac-like element (PBLE)

from the bat *Myotis lucifugus* (Piggybat) and the human PiggyBac element-derived (PGBD)

proteins PGBD2 and PGBD3. In contrast, PiggyBac homologs from *Paramecium* and

Tetrahymena have a CRD with six cysteine residues arranged in the variants of the motif CxxC-

CxxC-Cx{2-7}Cx{3,4}H, and group together with human PGDB4 and *Spodoptera frugiperda*

PBLE (Figure 6C).

Previous experiments involving individual or paired gene knockdowns of most of the ten

Paramecium tetraurelia PiggyMac(-like) paralogs lead to substantial IES retention, even though

only one PiggyMac gene (PGM) has the complete catalytic triad, indicating that all these

proteins are functional [55]. To examine whether PiggyMac homologs in *Paramecium* are

functionally constrained we examined non-synonymous (d_N) to synonymous substitution rate

(d_S) ratios ($\omega = d_N/d_S$) for pairwise codon sequence alignments within and between species, with

the latter using two closely related *Paramecium* species (*P. tetraurelia* and *P. octaurelia*). All the

d_N/d_S values for pairwise comparisons of each of the catalytically incomplete *P. tetraurelia*

PgmLs versus the complete Pgm, were less than 1, ranging from 0.01 to 0.25 (Table S7). All

d_N/d_S values for pairwise comparisons between *P. tetraurelia* and *P. octaurelia* PiggyBac

orthologs were also substantially less than 1, ranging from 0.02 to 0.11 (Table S8). Since $d_N/d_S=$

1 indicates genes evolving neutrally [59], none of these genes are likely pseudogenes, and all

are subject to similar strong purifying selection.

Since only one of the eight homologs has the complete DDD catalytic triad characteristic of

functional PiggyBac transposases, this is the only possible catalytically active PiggyMac in *B.*

stoltei. The *in vitro* catalytic activity of the *T. ni* PiggyBac transposase is abolished when any of

the catalytic residues are substituted by alanine [60]. Conversely, we expect all *Blepharisma*

PiggyMac homologs with an incomplete catalytic triad to be catalytically inactive. In pairwise

comparisons of each of the catalytically incomplete homologs versus the complete one d_N/d_S

ranges from 0.0076 to 0.1351 (Table S9). As these estimates fall in a similar range to those

obtained for *Paramecium* PiggyMac/PiggyMac-likes, this indicates that the *B. stoltei* PiggyMac

homologs are predominantly subject to comparable purifying selection and unlikely to be

pseudogenes, but may have differentiated and acquired new functional roles.

A common origin for PiggyBac transposases in ciliates

To determine whether the *Blepharisma* PiggyBac homologs share a common ciliate ancestor, or

whether they arose from independent acquisitions in major ciliate groups, we explored a range

of phylogenetic methods. We created a phylogeny of 155 sequences of DDE_Tnp_1_7 domains

of PiggyBac homologs which are representatives of putative domesticated transposases from

the genus *Blepharisma* (*B. stoltei* ATCC 30299, *B. stoltei* HT-IV, *B. japonicum*), *Condylostoma*

magnum, *Paramecium* spp., *Tetrahymena thermophila*, together with PiggyBac-like element

(PBLE [61]) transposases from a stramenopile species, an archaeplastid species, several

opisthokont species and an amoebozoan, homologs of the human PiggyBac-derived 1 gene

(PGBD1), PGBD3, PGBD4 and PGBD5 homologs (Figure 7; Data S1). The three BPgms from

the three *Blepharisma* strains whose MAC genomes we sequenced form a monophyletic clade,

with the BPgmLs 1 and 2 as outgroups. The BPgmLs 3,4,5,6 and 7 are grouped together with

the *Condylostoma* Pgms (each of which appears to have a complete catalytic triad). These two

clades form a heterotrich-specific group in the phylogeny. We observe that the ciliate Pgms and

PgmLs largely cluster to form a single clade. *T. thermophila* Tpb7 is an exception which appears

to group together with homologs of PGBD5 from *H. sapiens, G. gallus* and *D. rerio* but has an

aBayes support value of 0.84, much lower than the mode (1.0) and average (0.91) for this

phylogeny. The other exceptions are two fungal PBLEs and two red algal PBLEs with low

aBayes support within the predominantly monophyletic ciliate clade.

We applied tree topology tests to ascertain the likelihood of a monophyletic clade of ciliate Pgms

and PgmLs. Constrained trees were compared by different methods including bootstrap

proportion using the RELL method [62], a one-sided Kishino-Hasegawa test, a Shimodaira-

Hasegawa test [63,64], estimating confidence using expected likelihood weights [65], and an

approximately unbiased test [66] (Table S10). A tree with the constraint of monophyly for all

ciliate Pgms and PgmLs lacks support in comparison to an unconstrained tree with all the ciliate

Pgms and PgmLs and eukaryotic PBLEs. After excluding *T. thermophila* Tpb7, a tree

constrained with a clade consisting of the remaining ciliate Pgms and PgmLs is supported by all

tests. In other words, the ciliate PiggyBac homologs except Tpb7 likely originated in a ciliate

common ancestor.

475   Discussion

The genus *Blepharisma* represents one of the earliest diverging ciliate lineages, the

heterotrichs, forming an outgroup to the best-studied and deeply divergent oligohymenophorean

and spirotrich ciliates [7]. *Blepharisma* species thus provide an excellent vantage point to

compare unique processes that have accompanied the evolution of nuclear and genomic

dimorphism in ciliates, particularly the extensive genomic reorganization that occurs during the

new macronucleus (MAC) development. Comparisons between *Blepharisma* and other ciliates

also enable deeper consideration of the potential state of the ciliate common ancestor. The

annotated draft *B. stoltei* ATCC 30299 MAC genome and associated transcriptomic data provide

the basis for comparative studies of key characteristics of genome reorganization processes.

485   A minichromosomal architecture with pronounced telomere

486   addition site variation

Ciliate MAC genome architectures span from longer DNA molecules in the

oligohymenophoreans *Paramecium tetraurelia* and *Tetrahymena thermophila*, which are more

similar to conventional eukaryotic chromosomes, to numerous, predominantly single-gene

nanochromosomes of spirotrichs like *Oxytricha trifallax* and *Euplotes crassus* [3]. The *B. stoltei*

ATCC 30299 MAC genome organization (and like that of *Paramecium bursaria* [44]) lies

between these two extremes with many minichromosomes representing numerous alternative

telomere addition sites. This form of natural sequence heterogeneity creates significant new

challenges for generating single representative consensus sequences, because genome

assembly methods assume chromosome boundaries are generally consistent, as they are in

multicellular eukaryote models, down to the terminology used to describe them [78]. Fortunately,

the reference *Blepharisma* MAC genome is largely homozygous, promoting contiguity in the

final, 41 Mb assembly of 64 two-telomere-capped contigs. Nevertheless, plenty of room for

refinement remains, including development of new ways to better assemble, describe and

organize the constituents of genomes with such extreme sequence variability.

Despite the abundance of *Blepharisma* MAC genome telomeres, we did not detect a typical

ncRNA gene corresponding to the telomerase RNA component (TERC) of the ribozyme

responsible for telomere synthesis in the MAC genome. We suspect this is due to ncRNAs

presenting a far greater challenge to detect than protein-coding genes and the presence of

highly divergent ncRNA with insufficient similarity to the handful of taxonomically-restricted

TERCs identified in oligohymenophorean and spirotrich ciliates and other eukaryotes so far. On

the other hand, five homologs of POT1, the canonical telomere-binding protein, are present

(Figure S4). One of these is highly upregulated when the new MAC genome is forming, and

presumably like model ciliates the DNA is fragmented and replicated, requiring further telomere

synthesis and telomere-binding proteins.

512 A potentially catalytically active PiggyMac is present in

513 *Blepharisma*

In current models of IES excision, MIC-limited sequences are demarcated by deposition of

methylation marks on histones in an sRNA-dependent process [5]. These sequences are then

recognized by domesticated transposases whose excision is supported by additional proteins,

including those that recognize these marks [5]. Together with MIC sequencing and from the

same time course we employed in the present study for RNA-seq we have observed abundant,

development-specific sRNA production in *Blepharisma* like that in other ciliates (Seah et al. in

prep), and so we examined the expression of potential genes responsible for their biogenesis,

along with other genes highly upregulated during new MAC formation. Homologs of proteins

implicated in ciliate genome reorganization were present among the genes most highly

differentially upregulated during new MAC development, notably including Dicer-like and Piwi

proteins, and domesticated transposases.

Following this study, we will report detailed analyses of a draft *B. stoltei* ATCC 30299 MIC

genome, which possesses abundant and typically TA-bound IESs (Seah et al. in prep). In

oligohymenophorean ciliates *Tetrahymena* and *Paramecium* there is a considerable body of

evidence that PiggyBac homologs are responsible for IES excision [35,52,53,55,79]. The

responsible IES excisases in the less-studied spirotrich ciliates, *Oxytricha*, *Stylonychia* and

Euplotes, are not as evident. In *Oxytricha* the TBE transposases are considered to be involved

in IES excision, but, unlike the MAC genome-encoded primary IES excisase (Tpb2) in

Tetrahymena and all the *Paramecium* PiggyMacs and PiggyMac-likes, are encoded by full-

length transposons present in the MIC genome, but absent from the MAC [74]. The considerable

heterogeneity in *Oxytricha* IES boundaries and "unscrambling" of a large subset of them [56],

together with the observation of pronounced developmental upregulation of numerous additional

MAC- and MIC-encoded transposases raises the possibility that additional transposases other

than those of TBEs could also be involved in IES excision [39,56]. Knowledge of IESs in other

ciliates is sparse (primarily confined to the phyllopharyngean *Chilodonella uncinata* [80,81]),

and, as far as we are aware, no specific IES excisases have been proposed for them.

Since the oligohymenophorean ciliate PiggyBac homologs are clear IES excisases, we sought

and found eight homologs of these genes in the *B. stoltei* ATCC 30299 MAC genome.

Blepharisma is the first ciliate genus aside from *Tetrahymena* and *Paramecium* in which such

proteins have been reported. Additional searches revealed clear PiggyBac homologs in

Condylostoma magnum, and a weaker pair of matches in *Stentor coeruleus*, suggesting that

these are a common feature of heterotrich ciliates. Reminiscent of *Paramecium tetraurelia*, in

which just one of the nine PiggyBac homologs, PiggyMac, has a complete DDD catalytic triad

[55], a single *B. stoltei* PiggyBac homolog has a complete DDD catalytic triad. The gene

encoding this protein in *B. stoltei* ATCC 30299 is highly upregulated during MAC development,

together with two other homologs that lack a complete catalytic triad. As is characteristic of

PiggyBac homologs, each of these three homologs also has a C-terminal, cysteine-rich, zinc

finger domain, which most closely resemble those of *Condylostoma magnum* homologs. The

organization of the heterotrich PiggyBac homolog zinc finger domains appears to be more

similar to that of comparable domains of *Homo sapiens* PGBD2 and PGBD3 homologs than that

of the zinc finger domains in *Paramecium* and *Tetrahymena* PiggyBac homologs (which are

more similar to the *Homo sapiens* PGBD4 zinc fingers [58]).

In *Paramecium*, since the discovery of multiple PiggyBac homologs (PiggyMac-likes), there

have been questions about their role, since, aside from PiggyMac, they all have incomplete

catalytic triads, and are thus likely catalytically inactive, but nevertheless their gene knockdowns

lead to pronounced IES retention [55]. It has therefore been proposed that the PiggyMac-likes

may function as heteromeric multi-subunit complexes in conjunction with PiggyMac during DNA

excision [55]. On the other hand cryo-EM structures available for moth PiggyBac transposase

support a model in which these proteins function as a homodimeric complex *in vitro* [82].

Furthermore, the primary *Tetrahymena* PiggyBac, Tpb2, is able to perform cleavage *in vitro*,

without the assistance of other PiggyBac homologs [53]. In other eukaryotes, domesticated

PiggyBacs without complete catalytic triads are thought to be retained by virtue of useful DNA-

binding roles [83]. One possibility for such purely DNA-binding transposases in ciliates could be

in competitively regulating (taming) the excision of DNA by the catalytically active transposases.

Future experimental analyses of the BPgm and the BPgm-likes could aid in resolving the

conundrums and understanding of possible interactions between catalytically active and inactive

transposases.

Domesticated PiggyBac homologs are the main candidate IES excisases in *Blepharisma*

In addition to the PiggyBac homologs, we also found MAC genome-encoded transposases with

the PFAM domains "DDE_1", "DDE_3", "DDE_Tnp_IS1595" and "MULE" in *Blepharisma* (Figure

6., Supplementary results, "Additional domesticated transposases "). All the genes encoding

proteins with these domains lack flanking terminal repeats characteristic of active transposons,

suggesting these genes are further classes of domesticated transposases. In *Blepharisma* and

numerous other organisms, the DDE_1 domains co-occur with CENPB domains. Two such

proteins represent totally different proposed exaptations in mammals (centromere-binding

protein) and fission yeast (regulatory protein) [68–70]. Given the great evolutionary distances

involved, there is no specific reason to expect that the *Blepharisma* homologs have either

function. None of the three proteins with co-occurring DDE_1 and CENPB domains have a

complete catalytic triad, making it unlikely that these are active transposases or IES excisases,

though all three are noticeably upregulated during MAC development. Six proteins with the

PFAM domain DDE_3 are also encoded by *Blepharisma* MAC genes, of which five possess a

complete catalytic triad. DDE_3 domains are also characteristic of TBE transposases in

Oxytricha and Tec transposases in *Euplotes*. All the "DDE_3" protein genes are upregulated

during conjugation in *B. stoltei*, and are expressed at particularly high levels during development

of the new MAC. A number of DDE_Tnp_IS1595 and MULE domain-containing proteins have

complete catalytic triads and also show pronounced upregulation during *B. stoltei* MAC

development.

Upon excision, classical cut-and-paste transposases in eukaryotes typically leave behind

additional bases, notably including those of the target-site duplication that arose when they were

inserted, forming a "footprint" [84]. PiggyBac homologs are unique in performing precise,

"seamless" excision in eukaryotes [85], conserving the number of bases at the site of

transposon insertion after excision, a property that makes them popular for genetic engineering

[82]. *Tetrahymena* Tpb2 is the one exception among PiggyBac homologs associated with

imprecise excision in this eukaryote [53]. Since intragenic IESs are abundant in *Blepharisma*,

like *Paramecium* and unlike *Tetrahymena*, it is essential that these are excised precisely without

addition or removal of nucleotides, which, at best, could result in translation of additional amino

acids and, at worst, translation frameshifts.

Though there are clearly numerous additional domesticated transposases with complete

catalytic triads and whose genes are substantially upregulated during *Blepharisma*

development, whether they are capable of excision, and if this is precise, needs to be

established. *Tetrahymena* has distinct domesticated transposases that excise different subsets

of IESs, namely those that are predominant, imprecisely excised and intergenic (by Tpb2) [53],

versus those that are rare, precisely excised and intragenic (by Tpb1 and Tpb6) [79,86]. We

could envisage if the additional *Blepharisma* domesticated transposases are still capable of

excision, but not a precise form, an involvement in excision of a subset of the numerous

intergenic IESs. Thus, while we cannot rule out the involvement of other possible domesticated

transposases and PiggyMac-likes, currently *B. stoltei* PiggyMac is the primary candidate IES

excisase.

A possible PiggyBac homolog ciliate common ancestor

In light of the discovery of PiggyMacs in heterotrichous ciliates, we investigated whether the last

common ancestor of ciliates also possessed a PiggyMac. Phylogenetic analyses, together with

tree topology tests, indicate that the oligohymenophorean and heterotrichous PiggyMacs and

PiggyMac-likes form a monophyletic clade. The lack of PiggyBac homologs in other ciliate

classes such as the Spirotrichea, and potentially the oligohymenophorean *Ichthyophthirius*

multifiliis (with the caveat of apparent lower genome completeness), raises the question whether

the PiggyMacs were lost in these lineages or were gained independently from the same source

by both heterotrichs and a subset of oligohymenophoreans. We think the former is more likely.

However, the alternative cannot be completely dismissed, because non-model ciliates, where

the genome assembly quality allows reliable gene and domain annotations, have only been

sparsely sampled.

Future directions

The *B. stoltei* ATCC 30299 MAC genome together with a corresponding MIC genome (Seah et

al., in prep.) pave the way for future investigations of a peculiar, direct pathway to new MAC

genome development which skips the upstream complexity of meiotic and fertilization-

associated nuclear developmental processes of the standard pathway [9]. The pair of *B. stoltei*

strains we have are both now low frequency selfers, in which the conventional, indirect MAC

development pathway dominates. Comparisons with fresh, high frequency *Blepharisma* selfers

collected from the wild will facilitate comparative gene expression analyses with the direct MAC

development pathway, which will assist in distinguishing expression upregulation due to meiotic

and mitotic processes preceding indirect new MAC development. As a representative of one of

the two deepest branching ciliate lineages, the assembled somatic genome will facilitate

investigations beyond those reported here into the enigmatic origins of nuclear and genomic

dualism within single cells.

Materials and Methods

Strains and localities

The strains used and their original isolation localities were: *Blepharisma stoltei* ATCC 30299,
Lake Federsee, Germany [24]; *Blepharisma stoltei* HT-IV, Aichi prefecture, Japan; *Blepharisma*
japonicum R1072, from an isolate from Bangalore, India [22].

Cell cultivation, harvesting and cleanup

For genomic DNA isolation *B. stoltei* ATCC 30299 and HT-IV cells were cultured in Synthetic
Medium for *Blepharisma* (SMB) [87] at 27˚C. Belpharismas were fed *Chlorogonium elongatum*
grown in Tris-acetate phosphate (TAP) medium [88] at room temperature. *Chlorogonium* cells
were pelleted at 1500 g at room temperature for 3 minutes to remove most of the TAP medium,
and resuspended in 50 mL SMB. 50 ml of dense *Chlorogonium* was used to feed 1 litre of
Blepharisma culture once every three days.

Blepharisma stoltei ATCC 30299 and HT-IV cells used for RNA extraction were cultured in
Lettuce medium inoculated with *Enterbacter aerogenes* and maintained at 25˚C [89].

Blepharisma cultures were concentrated by centrifugation in pear-shaped flasks at 100 g for 2
minutes using a Hettich Rotanta 460 centrifuge with swing out buckets. Pelleted cells were
washed with SMB and centrifuged again at 100 g for 2 minutes. The washed pellet was then
transferred to a cylindrical tube capped with a 100 μm-pore nylon membrane at the base and
immersed in SMB to filter residual algal debris from the washed cells. The cells were allowed to
diffuse through the membrane overnight into the surrounding medium. The next day, the

665 cylinder with the membrane was carefully removed while attempting to minimize dislodging any

666 debris collected on the membrane. Cell density after harvesting was determined by cell counting

667 under the microscope.

668 DNA isolation, library preparation and sequencing

B. stoltei macronuclei were isolated by sucrose gradient centrifugation [4]. DNA was isolated

with a Qiagen 20/G genomic-tip kit according to the manufacturer's instructions. Purified DNA

from the isolated MACs was fragmented, size selected and used to prepare libraries according

to standard PacBio HiFi SMRTbell protocols. The libraries were sequenced in circular

consensus mode to generate HiFi reads.

Total genomic DNA from *B. stoltei* HT-IV and *B. stoltei* ATCC 30299 was isolated with the

SigmaAldrich GenElute Mammalian genomic DNA kit. A sequencing library was prepared with a

NEBnext FS DNA Library Prep Kit for Illumina and sequenced on an Illumina HiSeq 3000

sequencer, generating 150 bp paired-end reads.

Total genomic DNA from *B. japonicum* was isolated with the Qiagen MagAttract HMW DNA kit.

A long-read PacBio sequencing library was prepared using the SMRTbell® Express Template

Preparation Kit 2.0 according to the manufacturers' instructions and sequenced on an PacBio

Sequel platform with 1 SMRT cell. Independently, total genomic DNA form *B. japonicum* was

isolated with the SigmaAldrich GenElute Mammalian genomic DNA kit and an sequencing

library was prepared with the TruSeq Nano DNA Library Prep Kit (Illumina) and sequenced on

an Illumina NovaSeq6000 to generate 150 bp paired-end reads.

Gamone 1/ Cell-Free Fluid (CFF) isolation and conjugation activity assay

B. stoltei ATCC 30299 cells were cultured and harvested and concentrated to a density of 2000 cells/mL according to the procedure described in "Cell cultivation, Harvesting and Cleanup". This concentrated cell culture was incubated overnight at 27˚C. The next day, the cells were harvested, and the supernatant collected and preserved at 4˚C at all times after extraction. The supernatant was then filtered through a 0.22 µm-pore filter. BSA (10 mg/mL) was added to produce the final CFF at a final BSA concentration of 0.01%.

To assess the activity of the CFF, serial dilutions of the CFF were made to obtain the gamone activity in terms of units (U) [90].The activity of the isolated CFF was 2^{10} U.

Conjugation time course and RNA isolation for high-throughput sequencing

B. stoltei cells for the complementary strains, ATCC 30299 and HT-IV, were cultivated and harvested by gentle centrifugation to achieve a final cell concentration of 2000 cells/ml for each strain. Non-gamone treated ATCC 30299 (A1) and HT-IV cells (H1) were collected (time point: - 3 hours). Strain ATCC 30299 cells were then treated with synthetic gamone 2 (final concentration 1.5 µg/mL) and strain HT-IV cells were treated with cell-free fluid with a gamone 1 activity of ~2^{10} U/ml for three hours (Figure S8).

Homotypic pair formation in both cultures was checked after three hours. More than 75% of the cells in both cultures formed homotypic pairs. At this point the samples A2 (ATCC 30299) and H2 (HT-IV) were independently isolated for RNA extraction as gamone-treated control cells just

before mixing. For the rest of the culture, homotypic pairs in both cultures were separated by

pipetting them gently with a wide-bore pipette tip. Once all pairs had been separated, the two

cultures were mixed together. This constitutes the experiment's 0-h time point. The conjugating

culture was observed and samples collected for RNA isolation or cell fixation at 2 h, 6 h, 14 h,

18 h, 22 h, 26 h, 30 h and 38 h (Figure S8). Further details of the sample staging approach are

described in [9] and [51]. At each time point including samples A1, H1, A2 and H2, 7 mL of

culture was harvested for RNA-extraction using Trizol. The total RNA obtained was then

separated into a small RNA fraction < 200 nt and a fraction with RNA fragments > 200 nt using

the Zymo RNA Clean and Concentrator-5 kit according to the manufacturer's instructions. RNA-

seq libraries were prepared by BGI according to their standard protocols and sequenced on a

BGISeq 500 instrument.

Separate 2 mL aliquots of cells at each time point for which RNA was extracted were

concentrated by centrifuging gently at 100 rcf. 50 µL of the concentrated cells were fixed with

Carnoy's fixative (ethanol:acetic acid, 6:1), stained with DAPI and imaged to determine the state

of nuclear development [9].

Cell fixation and imaging

B. stoltei cells were harvested as above ("Cell cultivation"), and fixed with an equal volume of

"ZFAE" fixative, containing zinc sulfate (0.25 M, Sigma Aldrich), formalin, glacial acetic acid and

ethanol (Carl Roth), freshly prepared by mixing in a ratio of 10:2:2:5. Fixed cells were pelleted

(1000 g; 1 min), resuspended in 1% TritonX-100 in PHEM buffer to permeabilize (5 min; room

temperature), pelleted and resuspended in 2% (w/v) formaldehyde in PHEM buffer to fix further

(10 min; room temp.), then pelleted and washed twice with 3% (w/v) BSA in TBSTEM buffer

(~10 min; room temp.). For indirect immunofluorescence, washed cells were incubated with

primary antibody rat anti-alpha tubulin (Abcam, ab6161; 1:100 dilution in 3% w/v BSA/TBSTEM;

60 min; room temp.) then secondary antibody goat anti-rat IgG H&L labeled with AlexaFluor 488

(Abcam, ab150157, 1:500 dilution in 3% w/v BSA/TBSTEM; 20 min; room temp.). Nuclei were

counterstained with DAPI (1 µg/mL) in 3% (w/v) BSA/TBSTEM. A z-stack of images was

acquired using a confocal laser scanning microscope (Leica TCS SP8), equipped with a HC PL

APO 40× 1.30 Oil CS2 objective and a 1 photomultiplier tube and 3 HyD detectors, for DAPI

(405 nm excitation, 420-470 nm emission) and Alexa Fluor 488 (488 nm excitation, 510-530 nm

emission). Scanning was performed in sequential exposure mode. Spatial sampling was

achieved according to Nyquist criteria. ImageJ (Fiji) [91] was used to adjust image contrast and

brightness and overlay the DAPI and AlexaFluor 488 channels. The z-stack was temporally

color-coded.

For a nuclear 3D reconstruction (Figure 1B), cells were fixed in 1% (w/v) formaldehyde and

0.25% (w/v) glutaraldehyde. Nuclei were stained with Hoechst 33342 (Invitrogen) (5 µM in the

culture media), and imaged with a confocal laser scanning microscope (Zeiss, LSM780)

equipped with an LD C-Apochromat 40x/1,1 W Korr objective and a 32 channel GaAsP array

detector, with 405 nm excitation and 420-470 nm emission. Spatial sampling was achieved

according to Nyquist criteria. The IMARIS (Bitplane) software v8.0.2 was used for three-

dimensional reconstructions and contrast adjustments.

For scanning electron microscopy (SEM) cells were fixed in 0.5% osmium tetroxide/2%

formaldehyde/2.5% glutaraldehyde in 0.06x PHEM for 5-10 minutes at room temperature. Cells

were post-fixed with 2% formaldehyde/2.5% glutaraldehyde in 0.06x PHEM for 4-6 hours at

room temperature. Subsequently, samples were dehydrated in a graded ethanol series with 1

day per step followed by critical point drying (Polaron) with CO_2. Finally, the cells were sputter-

coated with a 4 nm thick layer of platinum (CCU-010, Safematic) and examined with a field

emission scanning electron microscope (Regulus 8230, Hitachi High Technologies) at an

accelerating voltage of 3 kV.

Genome assembly

Two MAC genome assemblies for *B. stoltei* ATCC 30299 (70× and 76× coverage) were

produced with Flye (version 2.7-b1585) [92] for the two separate PacBio Sequel II libraries

(independent replicates) using default parameters and the switches: --pacbio-hifi -g 45m. The

approximate genome assembly size was chosen based on preliminary Illumina genome

assemblies of approximately 40 Mb. Additional assemblies using the combined coverage (145×)

of the two libraries were produced using either Flye version 2.7-b1585 or 2.8.1-b1676, and the

same parameters. Two rounds of extension and merging were then used, first comparing the

70× and 76× assemblies to each other, then comparing the 145× assembly to the former

merged assembly. Assembly graphs were all relatively simple, with few tangles to be resolved

(Figure S7). Minimap2 [93] was used for pairwise comparison of the assemblies using the

parameters: -x asm5 --frag=yes --secondary=no, and the resultant aligned sequences were

visually inspected and manually merged or extended where possible using Geneious (version

2020.1.2) [94].

Visual inspection of read mapping to the combined assembly was then used to trim off contig

ends where there was little correspondence between the assembly consensus and the mapped

reads - which we classify as "cruft". Read mapping to cruft regions was often lower or uneven,

suggestive of repeats. Alternatively, these features could be due to trace MIC sequences, or

sites of alternative chromosome breakage during development which lead to sequences that are

neither purely MAC nor MIC. A few contigs with similar dubious mapping of reads at internal

locations, which were also clear sites of chromosome fragmentation (evident by abundant

telomere-bearing reads in the vicinity) were split apart and trimmed back as for the contig ends.

Telomere-bearing reads mapped to the non-trimmed region nearest to the trimmed site were

then used to define contig ends, adding representative telomeric repeats from one of the

underlying sequences mapped to each of the ends. The main genome assembly with gene

predictions can be obtained from the European Nucleotide Archive (ENA) (PRJEB40285;

accession GCA_905310155). "Cruft" sequences are also available from the same project

accession.

Two separate assemblies were generated for *Blepharisma japonicum.* A genome assembly for

Blepharisma japonicum strain R1072 was generated from Illumina reads, using SPAdes

genome assembler (v3.14.0) [95]. An assembly with PacBio Sequel long reads was produced

with Ra (v0.2.1) [96], which uses the Overlap-Layout-Consensus paradigm. The assembly

produced with Ra was more contiguous, with 268 contigs, in comparison to 1510 contigs in the

SPAdes assembly, and was chosen as the reference assembly for *Blepharisma japonicum*

(ENA accession: ERR6474383)*.*

Condylostoma magnum genomic reads (study accession PRJEB9019) from a previous study

[97] were reassembled to improve contiguity and remove bacterial contamination. Reads were

trimmed with bbduk.sh from the BBmap package v38.22

(https://sourceforge.net/projects/bbmap/), using minimum PHRED quality score 2 (both ends)

and k-mer trimming for Illumina adapters and Phi-X phage sequence (right end), retaining only

reads ≥25 bp. Trimmed reads were error-corrected and reassembled with SPAdes v3.13.0 [95]

using k-mer values 21, 33, 55, 77, 99. To identify potential contaminants, the unassembled

reads were screened with phyloFlash v3.3b1 [98] against SILVA v132 [99]; the coding density

under the standard genetic code and prokaryotic gene model were also estimated using

Prodigal v2.6.3 [100]. Plotting the coverage vs. GC% of the initial assembly showed that most of

the likely bacterial contigs (high prokaryotic coding density, lower coverage, presence of

bacterial SSU rRNA sequences) had >=40% GC, so we retained only contigs with <40% GC as

the final *C. magnum* genome bin. The final assembly is available from the ENA bioproject

PRJEB48875 (accession GCA_920105805).

All assemblies were inspected with the quality assessment tool QUAST [101].

818   Variant calling

Illumina total genomic DNA-seq libraries for *B. stoltei* strains ATCC 30299 (ENA accession:

ERR6061285) and HT-IV (ERR6064674) were mapped to the ATCC 30299 reference assembly

with bowtie2 v2.4.2 [102]. Alignments were tagged with the MC tag (CIGAR string for mate/next

segment) using samtools [103] fixmate. The BAM file was sorted and indexed, read groups were

added with bamaddrg (commit 9baba65, https://github.com/ekg/bamaddrg), and duplicate reads

were removed with Picard MarkDuplicates v2.25.1 (http://broadinstitute.github.io/picard/).

Variants were called from the combined BAM file with freebayes v1.3.2 [104] in diploid mode,

with maximum coverage 1000 (option -g). The resultant VCF file was combined and indexed

with bcftools v1.12 [103], then filtered to retain only SNPs with quality score > 20, and at least

one alternate allele.

Annotation of alternative telomere addition sites

Alternative telomere addition sites (ATASs) were annotated by mapping PacBio HiFi reads to

the curated reference MAC assembly described above, using minimap2 and the following flags:

-x asm20 --secondary=no --MD. We expect reads representing alternative telomere additions to

have one portion mapping to the assembly (excluding telomeric regions), with the other portion

containing telomeric repeats being soft-clipped in the BAM record. For each mapped read with a

soft-clipped segment, we extracted the clipped sequence, and the coordinates and orientation of

the clip relative to the reference. We searched for ≥ 24 bp tandem direct repeats of the telomere

unit (i.e., ≥3 repeats of the 8 bp unit) in the clipped segment with NCRF v1.01.02 [105], which

can detect tandem repeats in the presence of noise, e.g., from sequencing error. The orientation
of the telomere sequence, the distance from the end of the telomeric repeat to the clip junction
('gap'), and the number of telomere-bearing reads vs. total mapped reads at each junction were
also recorded. Junctions with zero gap between telomere repeat and clip junction were
annotated as ATASs. The above procedure was implemented in the MILTEL module of the
software package BleTIES v0.1.3 [106].
844
MILTEL output was processed with Python scripts depending on Biopython [107], pybedtools
[108], Bedtools [109], and Matplotlib [110], to summarize statistics of junction sequences and
telomere permutations at ATAS junctions, and to extract genomic sequences flanking ATASs for
sequence logos. Logos were drawn with Weblogo v3.7.5 [111], with sequences oriented such
that the telomere would be added on the 5' end of the ATAS junctions.

To calculate the expected minichromosome length, we assumed that ATASs were independent

and identically distributed in the genome following a Poisson distribution. About 47×10^3 ATASs

were annotated, supported on average by a single read. Given a genome of 42 Mbp at 145×

coverage, the expected rate of encountering an ATAS is 47×10^3 / (145 × 42 Mbp), so the

distance between ATASs (i.e., the minichromosome length) is exponentially distributed with

expectation (145 × 42 Mbp) / 47×10^3 = 130 kbp.

857 RNA-seq read mapping

To permit correct mapping of tiny introns RNA-seq data was mapped to the *B. stoltei* ATCC

30299 MAC genome using a version of HISAT2 [112] with modified source code, with the static

variable minIntronLen in hisat2.cpp lowered to 9 from 20 (change available in the HISAT2 github

fork: https://github.com/Swart-lab/hisat2/; commit hash 86527b9). HISAT2 was run with default

parameters and parameters --min-intronlen 9 --max-intronlen 500. It should be noted that RNA-

seq from timepoints in which *B. stoltei* ATCC 30299 and *B. stoltei* HT-IV cells were mixed

together were only mapped to the former genome assembly, and so reads for up to three alleles

may map to each of the genes in this assembly.

866 Genetic code prediction

We used the program PORC (Prediction Of Reassigned Codons; available from

https://github.com/Swart-lab/PORC) previously written to predict genetic codes in protist

transcriptomes [97] to predict the *B. stoltei* genetic code. This program was used to translate the

draft *B. stoltei* ATCC 30299 genome assembly in all six frames (with the standard genetic

code). Like the program FACIL [113] that inspired PORC, the frequencies of amino acids in

PFAM (version 34.0) protein domain profiles aligned to the six frame translation by HMMER

3.1b2 [114] (default search parameters; domains used for prediction with conditional E-values <

1e-20), and correspondingly also to the underlying codon, are used to infer the most likely amino

acid encoded by each codon (Figure S5).

876   Gene prediction

We created a wrapper program, Intronarrator, to predict genes in *Blepharisma* and other

heterotrichs, accommodating their tiny introns. Intronarrator can be downloaded and installed

together with dependencies via Conda from GitHub (https://github.com/Swart-lab/Intronarrator).

Intronarrator directly infers introns from spliced RNA-seq reads mapped by HISAT2 from the

entire developmental time course we generated. RNA-seq reads densely cover almost the entire

Blepharisma MAC genome, aside from intergenic regions, and most potential protein-coding

genes (Figure 5A). After predicting the introns and removing them to create an intron-minus

genome, Intronarrator runs AUGUSTUS (version 3.3.3) using its intronless model. It then adds

back the introns to the intronless gene predictions to produce the final gene predictions.

Introns are inferred from "CIGAR" string annotations in mapped RNA-seq BAM files, using the

regular expression "[0-9]+M([0-9][0-9])N[0-9]+M" to select spliced reads. For intron inference we

only used primary alignments with: MAPQ >= 10; just a single "N", indicating one potential

intron, per read; and at least 6 mapped bases flanking both the 5' and 3' intron boundaries (to

limit spurious chance matches of a few bases that might otherwise lead to incorrect intron

prediction). The most important parameters for Intronarrator are a cut-off of 0.2 for the fraction of

spliced reads covering a potential intron, and a minimum of 10 or more spliced reads to call an

intron. The splicing fraction cut-off was chosen based on the overall distribution of splicing

(Figure S6A-C). From our visual examination of mapped RNA-seq reads and gene predictions,

values less than this were typically "cryptic" excision events [115] which remove potentially

essential protein-coding sequences, rather than genuine introns. Intronarrator classifies an

intron as sense (7389 in total, excluding alternative splicing), when the majority of reads

(irrespective of splicing) mapping to the intron are the same strand, and antisense (554 in total)

when they are not. The most frequently spliced intron was chosen in rare cases of overlapping

alternative intron splicing.

To eliminate spurious prediction of protein-coding genes overlapping ncRNA genes, we also

incorporated ncRNA prediction in Intronarrator. Infernal [38] (default parameters; e-value < 1e-6)

was used to predict a restricted set of conserved ncRNAs models (i.e., tRNAs, rRNAs, SRP, and

spliceosomal RNAs) from RFAM 14.0 [116]. These ncRNAs were hard-masked (with "N"

characters) before AUGUSTUS gene prediction. Both Infernal ncRNA predictions (excluding

tRNAs) and tRNA-scan SE 2.0 [117] (default parameters) tRNA predictions are annotated in the

B. stoltei ATCC 30299 assembly deposited in the European Nucleotide Archive.

Since we found that *Blepharisma stoltei*, like *Blepharisma japonicum* [97], uses a non-standard

genetic code, with UGA codon translated as tryptophan, gene predictions use the "The Mold,

Protozoan, and Coelenterate Mitochondrial Code and the Mycoplasma/Spiroplasma Code

(transl_table=4)" from the NCBI genetic codes. The default AUGUSTUS gene prediction

parameters override alternative (mitochondrial) start codons permitted by NCBI genetic code 4,

other than ATG. So, all predicted *B. stoltei* gene coding sequences begin with ATG.

RNA-seq read mapping relative to gene predictions of Contig_1 of *B. stoltei* ATCC30299 was

visualized with PyGenomeTracks [118].

Assessment of genome completeness

921 A BUSCO (version 4.0.2) [119] analysis of the assembled MAC genomes of *B. stoltei* and *B.*

922 *japonicum* was performed on the set of predicted proteins (BUSCO mode -prot) using the

923 BUSCO Alveolata database. The completeness of the *Blepharisma* genomes was compared to

924 the protein-level BUSCO analysis of the published genome assemblies of ciliates *T.*

925 *thermophila*, *P. tetraurelia*, *S. coeruleus* and *I. multifiliis* (Figure S1).

Gene annotation

927 Pannzer2 [120] (default parameters) and EggNog (version 2.0.1) [121] were used for gene

928 annotation. Annotations were combined and are available from the Max Planck Society's Open

929 Research Repository, Edmond (https://dx.doi.org/10.17617/3.8c). Protein domain annotations

930 were performed using hmmscan from HMMER3 (version 3.3, Nov 2019) [114] vs. the PFAM

931 database (Pfam-A.full, 33.0, retrieved on June 23, 2020) with default parameters.

Gene expression analysis

Features from RNA-seq reads mapped to the *B. stoltei* ATCC 30299 genome over the

developmental time-course were extracted using featureCounts from the Subread package

[122]. Further analysis was performed using the R software environment. Genes with a total

read count of less than 50, across all timepoints, were filtered out of the dataset. The remaining

genes were passed as a DGElist object to edgeR [123]. Each time point, representing one

library, was normalized for library size using the edgeR function calcNormFactors. The

normalized read counts were transformed into TPM (transcripts per million) values [124,125].

The TPM-values for different genes were compared across timepoints to examine changes in

gene expression. Heatmaps showing log2(TPM) changes across timepoints were plotted using

the tidyverse collection of R packages (https://www.tidyverse.org/) and RColorBrewer

(https://rdrr.io/cran/RColorBrewer/). Tabulated gene expression estimates together with protein

annotations are available from Edmond (https://dx.doi.org/10.17617/3.8c).

Sequence visualization and analysis

946 Nucleotide and amino acid sequences were visualized using Geneious Prime (Biomatters Ltd.)

947 [94]. Multiple sequence alignments were performed with MAFFT version 7.450 [126,127].

948 Phylogenetic trees were constructed with PhyML version 3.3.20180621 [128].

Orthogroup inference and analysis of orthogroup clusters

OrthoFinder version 2.5.2 with default parameters (i.e., using Diamond for searching, MAFFT for

multiple alignment and FastTree for phylogenies) was used to define orthogroups, i.e., sets of

genes descended from the last common ancestor of the chosen species. Proteomes for the

following ciliate species were used: *Tetrahymena thermophila*, *Oxytricha trifallax*, *Stentor*

coeruleus (data from ciliate.org [129]); *Euplotes octocarinatus* (EOGD [130]); *Paramecium*

tetraurelia, *Paramecium caudatum* (data from ParameciumDB [131]); plus *Perkinsus marinus*

ATCC 50983 (GenBank accession: AAXJ00000000) as a non-ciliate outgroup. Orthogroup

clusters are available as Data S2, or from Edmond (https://dx.doi.org/10.17617/3.8c).

d_N/d_S estimation

We generated pairwise coding sequence alignments of PiggyMac paralog nucleotide sequences

from *P. tetraurelia* and *P. octaurelia* using MAFFT version 7.450 [127] [126] (algorithm: "auto",

scoring matrix: 200PAM/k=2, gap open penalty 1.53, offset value 0.123) using the "translation

align" panel of Geneious Prime (version 2020.1.2) [94]. PAML version 4.9 [132] was used to

estimate d_N/d_S values in pairwise mode (runmode = -2, seqtype = 1, CodonFreq = 2). For

Blepharisma stoltei, we generated pairwise coding sequence alignments of the *Blepharisma*

PiggyMac homolog, BPgm (Contig_49.g1063; BSTOLATCC_MAC17466), with the *Blepharisma*

Pgm-likes (BPgmLs) using ClustalW 2.1 [133] (Genetic code: *Blepharisma*, Protein alignment

options: ClustalW alignment, Cost matrix: BLOSUM, Gap open cost: 10, Gap open penalty: 0.1)

using "translation align" panel of Geneious Prime (version 2020.1.2). PAML version 4.9 was

used to estimate dN/dS values in pairwise mode (runmode = -2, seqtype = 1, CodonFreq = 2).

Phylogenetic analysis

Protein sequences of PBLEs and PGDBs were obtained from Bouallègue et al [61]. Protein

sequences of *Paramecium* and *Tetrahymena* Pgms and PgmLs were obtained from

ParameciumDB [131] (PGM, PGMLs1-5) and ciliate.org [129] (Tpb1, Tpb2, Tpb7, LIA5),

respectively. *Condylostoma* and *Blepharisma* Pgms and PgmLs were obtained from genome

assemblies (accessions GCA_920105805 and GCA_905310155, respectively). Other PiggyBac

protein sequences (*T. ni, D. purpureum*) were obtained from Uniprot [134]. Sequence

manipulation was done using Geneious (Biomatters Ltd.). The Geneious plug-in for

InterProScan [135] was used to identify DDE_Tnp_1_7 domains using the PFAM-A database

[136]. The DDE_Tnp_1_7 domains were extracted and aligned using the MAFFT plug-in

(v7.450) for Geneious [127] (Algorithm: Auto, Scoring matrix: BLOSUM62, Gap open penalty:

1.53, Offset value: 0.123). The best-fit model of evolution for each alignment was identified

using ModelTest-NG [137] (modeltest-ng -i <input_file>.phy -d aa -p 4 -T phyml -o

<output_file>.out). The best model according to Bayesian criteria (BIC) were used to create

phylogenetic trees using PhyML [128] [phyml -i <input_file>.phy -d aa -m LG -f m -v 0 -a e -c 4 -

o tlr]. Phylogenetic trees thus generated were visualized with FigTree (v1.4.4) (Andrew

Rambaut, http://tree.bio.ed.ac.uk/). Tree topology tests were performed using IQtree (v2.1.4-

beta) [138] (iqtree -s <input file> -m LG+G4 -z <unconstrained and constrained treefiles> -n 0 -

zb 10000 -au -nt AUTO).

Repeat annotation

Interspersed repeat element families were predicted with RepeatModeler v2.0.1 (default

settings, random number seed 12345) with the following dependencies: rmblast v2.9.0+

(http://www.repeatmasker.org/RMBlast.html), TRF 4.09 (Benson, 1999), RECON (Bao and

Eddy, 2002), RepeatScout 1.0.6 (Price et al., 2005), RepeatMasker v4.1.1

(http://www.repeatmasker.org/RMDownload.html). Repeat families were also classified in the

pipeline by RepeatClassifier v2.0.1 through comparison against RepeatMasker's repeat protein

database and the Dfam database. Consensus sequences of the predicted repeat families,

produced by RepeatModeler, were then used to annotate repeats with RepeatMasker, using

rmblast as the search engine.

Terminal inverted repeats (TIRs) of selected repeat element families were identified by aligning

the consensus sequence from RepeatModeler, and/or selected full-length elements, with their

respective reverse complements using MAFFT (Katoh and Standley, 2013) (plugin version

distributed with Geneious). TIRs from the Dfam DNA transposon termini signatures database

(v1.1, https://www.dfam.org/releases/dna_termini_1.1/dna_termini_1.1.hmm.gz) (Storer et al.,

2021) were searched with hmmsearch (HMMer v3.2.1) against the IES sequences, to identify

matches to TIR signatures of major transposon subfamilies.

1007

Data availability

The draft *Blepharisma stoltei* ATCC 30299 macronuclear genome assembly is accessible from bleph.ciliate.org and from the European Nucleotide Archive (ENA) bioproject PRJEB40285 under the accession GCA_905310155. PacBio CCS reads (ERR5873783 and ERR5873334) and subreads (ERR5962314) used to assemble the genome are also available from ENA. Illumina DNA-seq data for the *B. stoltei* ATCC 30299 and HT-IV strains is available from accessions ERR6061285 and ERR6064674, respectively. The RNA-seq developmental time course is available from the bioproject PRJEB45374 (accessions ERR6049461-ERR6049485).

Illumina and PacBio Sequel sequencing data for *Blepharisma japonicum* strain R1702 is available from the ENA bioproject PRJEB46921 (Illumina accessions: ERR6473251, ERR6474356; PacBio accession: ERR6474383).

Acknowledgements

This paper is dedicated in memory of Akio Miyake and his decades of inspirational *Blepharisma* research. We thank Federico Buonanno for the provision of *B. stoltei* ATCC 30299 cells and culturing advice, Christa Lanz and the MPI for Biology's genome center, Jürgen Berger for SEM imaging, and Sebastien Colin and the MPI for Biology's Light Microscopy Facility for the 3D nuclear reconstruction.

Bibliography

1. Martin WF, Garg S, Zimorski V. Endosymbiotic theories for eukaryote origin. Philos Trans R Soc Lond B Biol Sci. 2015;370: 20140330. doi:10.1098/rstb.2014.0330

2. Orias E. On the evolution of the karyorelict ciliate life cycle: heterophasic ciliates and the origin of ciliate binary fission. BioSystems. 1991;25: 67–73. doi:10.1016/0303-2647(91)90013-B

3. Prescott DM. The DNA of ciliated protozoa. Microbiol Rev. 1994;58: 233–267. doi:10.1128/mr.58.2.233-267.1994

4. Lauth MR, Spear BB, Heumann J, Prescott DM. DNA of ciliated protozoa: DNA sequence diminution during macronuclear development of Oxytricha. Cell. 1976;7: 67–74. doi:10.1016/0092-8674(76)90256-7

5. Chalker DL, Meyer E, Mochizuki K. Epigenetics of ciliates. Cold Spring Harb Perspect Biol. 2013;5: a017764. doi:10.1101/cshperspect.a017764

6. Vogt A, Goldman AD, Mochizuki K, Landweber LF. Transposon domestication versus mutualism in ciliate genome rearrangements. PLoS Genet. 2013;9: e1003659. doi:10.1371/journal.pgen.1003659

7. Lynn DH, editor. The Ciliated Protozoa. Dordrecht: Springer Netherlands; 2010. doi:10.1007/978-1-4020-8239-9

8. Giese AC. Blepharisma: The Biology of a Light-sensitive Protozoan. illustrated ed. Stanford University Press; 1973.

9. Miyake A, Rivola V, Harumoto T. Double paths of macronucleus differentiation at conjugation in Blepharisma japonicum. Eur J Protistol. 1991;27: 178–200. doi:10.1016/S0932-4739(11)80340-8

10. Slabodnick MM, Ruby JG, Reiff SB, Swart EC, Gosai S, Prabakaran S, et al. The

Macronuclear Genome of Stentor coeruleus Reveals Tiny Introns in a Giant Cell. Curr

Biol. 2017;27: 569–575. doi:10.1016/j.cub.2016.12.057

11. Slabodnick MM, Marshall WF. Stentor coeruleus. Curr Biol. 2014;24: R783-4.

doi:10.1016/j.cub.2014.06.044

12. Zhang KS, Blauch LR, Huang W, Marshall WF, Tang SKY. Microfluidic guillotine reveals

multiple timescales and mechanical modes of wound response in Stentor coeruleus.

BMC Biol. 2021;19: 63. doi:10.1186/s12915-021-00970-0

13. Rapport EW, Rapport DJ, Berger J, Kupers V. Induction of conjugation in Stentor

coeruleus. Trans Am Microsc Soc. 1976;95: 220–224. doi:10.2307/3225068

14. Miyake A, Harumoto T. Asymmetrical cell division in Blepharisma japonicum: difference

between daughter cells in mating-type expression. Exp Cell Res. 1990;190: 65–68.

doi:10.1016/0014-4827(90)90144-y

15. Terazima MN, Harumoto T. Defense function of pigment granules in the ciliate

Blepharisma japonicum against two predatory protists, Amoeba proteus (Rhizopodea)

and Climacostomum virens (Ciliata). Zool Sci. 2004;21: 823–828. doi:10.2108/zsj.21.823

16. Kumazawa H. Homopolar Grafting in Blepharisma japonicum. Journal of Experimental

Zoology. 1979;207: 1–16.

17. Sugiura M, Shiotani H, Suzaki T, Harumoto T. Behavioural changes induced by the

conjugation-inducing pheromones, gamone 1 and 2, in the ciliate Blepharisma

japonicum. Eur J Protistol. 2010;46: 143–149. doi:10.1016/j.ejop.2010.01.002

18. Young D. Macronuclear reorganization in Blepharisma undulans. Doctoral dissertation.

1937.

19. Friedl E, Miyake A, Heckmann K. Requirement of successive protein syntheses for the

progress of meiosis in Blepharisma. Exp Cell Res. 1983;145: 105–113.

doi:10.1016/s0014-4827(83)80013-5

20.   Miyake A, Tulli M, Nobili R. Requirement of protein synthesis in the initiation of meiosis

and other nuclear changes in conjugation of Blepharisma. Exp Cell Res. 1979;120: 87–

93. doi:10.1016/0014-4827(79)90539-1

21.   Inaba F. Conjugation between two strains of *Blepharisma*. J Protozool. 1965;12: 146–

151. doi:10.1111/j.1550-7408.1965.tb01828.x

22.   Harumoto T, Miyake A, Ishikawa N, Sugibayashi R, Zenfuku K, Iio H. Chemical defense

by means of pigmented extrusomes in the ciliate Blepharisma japonicum. Eur J Protistol.

1998;34: 458–470. doi:10.1016/S0932-4739(98)80014-X

23.   Kobayashi M, Miura M, Takusagawa M, Sugiura M, Harumoto T. Two possible barriers

blocking conjugation between different megakaryotypes of Blepharisma. Zool Sci.

2015;32: 53–61. doi:10.2108/zs140151

24.   Repak AJ. Encystment and Excystment of the Heterotrichous Ciliate Blepharisma stoltei

Isquith. Journal of Protozoology. 1968;5: 407–412.

25.   Salvini M, Durante M, Nobili R. Characterization of macronuclear DNA in Blepharisma

japonicum. Protoplasma. 1983;117: 82–88. doi:10.1007/BF01281787

26.   Kimball RF. The nature and inheritance of mating types in euplotes patella. Genetics.

1942;27: 269–285.

27.   Katashima RYO. Mating Types in *Euplotes eurystomus*. J Protozool. 1959;6: 75–83.

doi:10.1111/j.1550-7408.1959.tb03930.x

28.   Luporini P, Miceli C, Ortenzi C. Evidence that the ciliate Euplotes raikovi releases

mating-inducing factors (gamones). J Exp Zool. 1983;226: 1–9.

doi:10.1002/jez.1402260102

29.   Vallesi A, Giuli G, Bradshaw RA, Luporini P. Autocrine mitogenic activity of pheromones

produced by the protozoan ciliate Euplotes raikovi. Nature. 1995;376: 522–524.

doi:10.1038/376522a0

30.   Braun V, Miyake A. Composition of blepharmone, a conjugation-inducing glycoprotein of

the ciliate Blepharisma. FEBS Lett. 1975;53: 131–134. doi:10.1016/0014-

5793(75)80003-2

31.   Miyake A, Beyer J. Blepharmone: a conjugation-inducing glycoprotein in the ciliate

blepharisma. Science. 1974;185: 621–623. doi:10.1126/science.185.4151.621

32.   Sugiura M, Harumoto T. Identification, characterization, and complete amino acid

sequence of the conjugation-inducing glycoprotein (blepharmone) in the ciliate

Blepharisma japonicum. Proc Natl Acad Sci USA. 2001;98: 14446–14451.

doi:10.1073/pnas.221457698

33.   Kubota T, Tokoroyama T, Tsukuda Y, Koyama H, Miyake A. Isolation and structure

determination of blepharismin, a conjugation initiating gamone in the ciliate blepharisma.

Science. 1973;179: 400–402. doi:10.1126/science.179.4071.400

34.   Klobutcher LA, Herrick G. Developmental genome reorganization in ciliated protozoa:

the transposon link. Prog Nucleic Acid Res Mol Biol. 1997;56: 1–62. doi:10.1016/S0079-

6603(08)61001-6

35.   Arnaiz O, Mathy N, Baudry C, Malinsky S, Aury J-M, Denby Wilkes C, et al. The

Paramecium germline genome provides a niche for intragenic parasitic DNA:

evolutionary dynamics of internal eliminated sequences. PLoS Genet. 2012;8:

e1002984. doi:10.1371/journal.pgen.1002984

36.   Klobutcher LA, Herrick G. Consensus inverted terminal repeat sequence of Paramecium

IESs: resemblance to termini of Tc1-related and Euplotes Tec transposons. Nucleic

Acids Res. 1995;23: 2006–2013. doi:10.1093/nar/23.11.2006

37.   Chin C-S, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, et al. Phased

1125   diploid genome assembly with single-molecule real-time sequencing. Nat Methods.

1126   2016;13: 1050–1054. doi:10.1038/nmeth.4035

1127   38.   Nawrocki EP, Kolbe DL, Eddy SR. Infernal 1.0: inference of RNA alignments.

1128   Bioinformatics. 2009;25: 1335–1337. doi:10.1093/bioinformatics/btp157

1129   39.   Swart EC, Bracht JR, Magrini V, Minx P, Chen X, Zhou Y, et al. The Oxytricha trifallax

1130   macronuclear genome: a complex eukaryotic genome with 16,000 tiny chromosomes.

1131   PLoS Biol. 2013;11: e1001473. doi:10.1371/journal.pbio.1001473

1132   40.   Aeschlimann SH, Jönsson F, Postberg J, Stover NA, Petera RL, Lipps H-J, et al. The

1133   draft assembly of the radically organized Stylonychia lemnae macronuclear genome.

1134   Genome Biol Evol. 2014;6: 1707–1723. doi:10.1093/gbe/evu139

1135   41.   Zagulski M, Nowak JK, Le Mouël A, Nowacki M, Migdalski A, Gromadka R, et al. High

1136   coding density on the largest Paramecium tetraurelia somatic chromosome. Curr Biol.

1137   2004;14: 1397–1404. doi:10.1016/j.cub.2004.07.029

1138   42.   Sheng Y, Duan L, Cheng T, Qiao Y, Stover NA, Gao S. The completed macronuclear

1139   genome of a model ciliate Tetrahymena thermophila and its application in genome

1140   scrambling and copy number analyses. Sci China Life Sci. 2020;63: 1534–1542.

1141   doi:10.1007/s11427-020-1689-4

1142   43.   Aury J-M, Jaillon O, Duret L, Noel B, Jubin C, Porcel BM, et al. Global trends of whole-

1143   genome duplications revealed by the ciliate Paramecium tetraurelia. Nature. 2006;444:

1144   171–178. doi:10.1038/nature05230

1145   44.   Cheng Y-H, Liu C-FJ, Yu Y-H, Jhou Y-T, Fujishima M, Tsai IJ, et al. Genome plasticity in

1146   Paramecium bursaria revealed by population genomics. BMC Biol. 2020;18: 180.

1147   doi:10.1186/s12915-020-00912-2

1148 45. Yao MC, Yao CH, Monks B. The controlling sequence for site-specific chromosome

1149   breakage in Tetrahymena. Cell. 1990;63: 763–772. doi:10.1016/0092-8674(90)90142-2

1150 46. Cavalcanti ARO, Dunn DM, Weiss R, Herrick G, Landweber LF, Doak TG. Sequence

1151   features of Oxytricha trifallax (class Spirotrichea) macronuclear telomeric and

1152   subtelomeric sequences. Protist. 2004;155: 311–322. doi:10.1078/1434461041844196

1153 47. Lynch M. The origins of eukaryotic gene structure. Mol Biol Evol. 2006;23: 450–468.

1154   doi:10.1093/molbev/msj050

1155 48. Roy SW, Hudson AJ, Joseph J, Yee J, Russell AG. Numerous fragmented spliceosomal

1156   introns, AT-AC splicing, and an unusual dynein gene expression pathway in Giardia

1157   lamblia. Mol Biol Evol. 2012;29: 43–49. doi:10.1093/molbev/msr063

1158 49. Bondarenko VS, Gelfand MS. Evolution of the Exon-Intron Structure in Ciliate Genomes.

1159   PLoS ONE. 2016;11: e0161476. doi:10.1371/journal.pone.0161476

1160 50. Csuros M, Rogozin IB, Koonin EV. A detailed history of intron-rich eukaryotic ancestors

1161   inferred from a global survey of 100 complete genomes. PLoS Comput Biol. 2011;7:

1162   e1002150. doi:10.1371/journal.pcbi.1002150

1163 51. Sugiura M, Tanaka Y, Suzaki T, Harumoto T. Alternative gene expression in type I and

1164   type II cells may enable further nuclear changes during conjugation of Blepharisma

1165   japonicum. Protist. 2012;163: 204–216. doi:10.1016/j.protis.2011.07.007

1166 52. Baudry C, Malinsky S, Restituito M, Kapusta A, Rosa S, Meyer E, et al. PiggyMac, a

1167   domesticated piggyBac transposase involved in programmed genome rearrangements in

1168   the ciliate Paramecium tetraurelia. Genes Dev. 2009;23: 2478–2483.

1169   doi:10.1101/gad.547309

1170 53. Cheng C-Y, Vogt A, Mochizuki K, Yao M-C. A domesticated piggyBac transposase plays

key roles in heterochromatin dynamics and DNA cleavage during programmed DNA

deletion in Tetrahymena thermophila. Mol Biol Cell. 2010;21: 1753–1762.

doi:10.1091/mbc.e09-12-1079

54. Yuan Y-W, Wessler SR. The catalytic domain of all eukaryotic cut-and-paste

transposase superfamilies. Proc Natl Acad Sci USA. 2011;108: 7884–7889.

doi:10.1073/pnas.1104208108

55. Bischerour J, Bhullar S, Denby Wilkes C, Régnier V, Mathy N, Dubois E, et al. Six

domesticated PiggyBac transposases together carry out programmed DNA elimination in

Paramecium. eLife. 2018;7. doi:10.7554/eLife.37927

56. Chen X, Bracht JR, Goldman AD, Dolzhenko E, Clay DM, Swart EC, et al. The

architecture of a scrambled genome reveals massive levels of genomic rearrangement

during development. Cell. 2014;158: 1187–1198. doi:10.1016/j.cell.2014.07.034

57. Coyne RS, Hannick L, Shanmugam D, Hostetler JB, Brami D, Joardar VS, et al.

Comparative genomics of the pathogenic ciliate Ichthyophthirius multifiliis, its free-living

relatives and a host species provide insights into adoption of a parasitic lifestyle and

prospects for disease control. Genome Biol. 2011;12: R100. doi:10.1186/gb-2011-12-10-

r100

58. Guérineau M, Bessa L, Moriau S, Lescop E, Bontems F, Mathy N, et al. The unusual

structure of the PiggyMac cysteine-rich domain reveals zinc finger diversity in PiggyBac-

related transposases. Mob DNA. 2021;12: 12. doi:10.1186/s13100-021-00240-4

59. Yang Z, Nielsen R. Estimating synonymous and nonsynonymous substitution rates

under realistic evolutionary models. Mol Biol Evol. 2000;17: 32–43.

doi:10.1093/oxfordjournals.molbev.a026236

60. Mitra R, Fain-Thornton J, Craig NL. piggyBac can bypass DNA synthesis during cut and

paste transposition. EMBO J. 2008;27: 1097–1109. doi:10.1038/emboj.2008.41

61. Bouallègue M, Rouault J-D, Hua-Van A, Makni M, Capy P. Molecular Evolution of

piggyBac Superfamily: From Selfishness to Domestication. Genome Biol Evol. 2017;9:

323–339. doi:10.1093/gbe/evw292

62. Kishino H, Miyata T, Hasegawa M. Maximum likelihood inference of protein phylogeny

and the origin of chloroplasts. J Mol Evol. 1990;31: 151–160. doi:10.1007/BF02109483

63. Kishino H, Hasegawa M. Evaluation of the maximum likelihood estimate of the

evolutionary tree topologies from DNA sequence data, and the branching order in

hominoidea. J Mol Evol. 1989;29: 170–179. doi:10.1007/BF02100115

64. Ota R, Waddell PJ, Hasegawa M, Shimodaira H, Kishino H. Appropriate likelihood ratio

tests and marginal distributions for evolutionary tree models with constraints on

parameters. Mol Biol Evol. 2000;17: 798–803.

doi:10.1093/oxfordjournals.molbev.a026358

65. Strimmer K, Rambaut A. Inferring confidence sets of possibly misspecified gene trees.

Proc Biol Sci. 2002;269: 137–142. doi:10.1098/rspb.2001.1862

66. Shimodaira H. An approximately unbiased test of phylogenetic tree selection. Syst Biol.

2002;51: 492–508. doi:10.1080/10635150290069913

67. Tanaka Y, Nureki O, Kurumizaka H, Fukai S, Kawaguchi S, Ikuta M, et al. Crystal

structure of the CENP-B protein-DNA complex: the DNA-binding domains of CENP-B

induce kinks in the CENP-B box DNA. EMBO J. 2001;20: 6612–6618.

doi:10.1093/emboj/20.23.6612

68. Casola C, Hucks D, Feschotte C. Convergent domestication of pogo-like transposases

into centromere-binding proteins in fission yeast and mammals. Mol Biol Evol. 2008;25:

29–41. doi:10.1093/molbev/msm221

69.   Hohmann S. Characterisation of PDC2, a gene necessary for high level expression of

1220   pyruvate decarboxylase structural genes in Saccharomyces cerevisiae. Mol Gen Genet.

1221   1993;241: 657–666. doi:10.1007/BF00279908

1222   70.   Mojzita D, Hohmann S. Pdc2 coordinates expression of the THI regulon in the yeast

1223   Saccharomyces cerevisiae. Mol Genet Genomics. 2006;276: 147–161.

1224   doi:10.1007/s00438-006-0130-z

1225   71.   Gao B, Wang Y, Diaby M, Zong W, Shen D, Wang S, et al. Evolution of pogo, a separate

1226   superfamily of IS630-Tc1-mariner transposons, revealing recurrent domestication events

1227   in vertebrates. Mob DNA. 2020;11: 25. doi:10.1186/s13100-020-00220-0

1228   72.   Witherspoon DJ, Doak TG, Williams KR, Seegmiller A, Seger J, Herrick G. Selection on

1229   the protein-coding genes of the TBE1 family of transposable elements in the ciliates

1230   Oxytricha fallax and O. trifallax. Mol Biol Evol. 1997;14: 696–706.

1231   doi:10.1093/oxfordjournals.molbev.a025809

1232   73.   Williams K, Doak TG, Herrick G. Developmental precise excision of Oxytricha trifallax

1233   telomere-bearing elements and formation of circles closed by a copy of the flanking

1234   target duplication. EMBO J. 1993;12: 4593–4601. doi:10.1002/j.1460-

1235   2075.1993.tb06148.x

1236   74.   Nowacki M, Higgins BP, Maquilan GM, Swart EC, Doak TG, Landweber LF. A functional

1237   role for transposases in a large eukaryotic genome. Science. 2009;324: 935–938.

1238   doi:10.1126/science.1170023

1239   75.   Jahn CL, Doktor SZ, Frels JS, Jaraczewski JW, Krikau MF. Structures of the Euplotes

1240   crassus Tec1 and Tec2 elements: identification of putative transposase coding regions.

1241   Gene. 1993;133: 71–78. doi:10.1016/0378-1119(93)90226-s

1242   76.   Doak TG, Doerder FP, Jahn CL, Herrick G. A proposed superfamily of transposase

genes: transposon-like elements in ciliated protozoa and a common "D35E" motif. Proc Natl Acad Sci USA. 1994;91: 942–946. doi:10.1073/pnas.91.3.942

77. Dupeyron M, Baril T, Bass C, Hayward A. Phylogenetic analysis of the Tc1/mariner superfamily reveals the unexplored diversity of pogo-like elements. Mob DNA. 2020;11: 21. doi:10.1186/s13100-020-00212-0

78. Lewin HA, Graves JAM, Ryder OA, Graphodatsky AS, O'Brien SJ. Precision nomenclature for the new genomics. Gigascience. 2019;8. doi:10.1093/gigascience/giz086

79. Feng L, Wang G, Hamilton EP, Xiong J, Yan G, Chen K, et al. A germline-limited piggyBac transposase gene is required for precise excision in Tetrahymena genome rearrangement. Nucleic Acids Res. 2017;45: 9481–9502. doi:10.1093/nar/gkx652

80. Zufall RA, Sturm M, Mahon BC. Evolution of germline-limited sequences in two populations of the ciliate Chilodonella uncinata. J Mol Evol. 2012;74: 140–146. doi:10.1007/s00239-012-9493-4

81. Zufall RA, Katz LA. Micronuclear and macronuclear forms of beta-tubulin genes in the ciliate Chilodonella uncinata reveal insights into genome processing and protein evolution. J Eukaryot Microbiol. 2007;54: 275–282. doi:10.1111/j.1550-7408.2007.00267.x

82. Chen Q, Luo W, Veach RA, Hickman AB, Wilson MH, Dyda F. Structural basis of seamless excision and specific targeting by piggyBac transposase. Nat Commun. 2020;11: 3446. doi:10.1038/s41467-020-17128-1

83. Sarkar A, Sim C, Hong YS, Hogan JR, Fraser MJ, Robertson HM, et al. Molecular evolutionary analysis of the widespread piggyBac transposon family and related "domesticated" sequences. Mol Genet Genomics. 2003;270: 173–180.

doi:10.1007/s00438-003-0909-0

84.   van Luenen HG, Colloms SD, Plasterk RH. The mechanism of transposition of Tc3 in C.

elegans. Cell. 1994;79: 293–301. doi:10.1016/0092-8674(94)90198-8

85.   Elick TA, Bauser CA, Fraser MJ. Excision of the piggyBac transposable element in vitro

is a precise event that is enhanced by the expression of its encoded transposase.

Genetica. 1996;98: 33–41. doi:10.1007/BF00120216

86.   Cheng C-Y, Young JM, Lin C-YG, Chao J-L, Malik HS, Yao M-C. The piggyBac

transposon-derived genes TPB1 and TPB6 mediate essential transposon-like excision

during the developmental rearrangement of key genes in Tetrahymena thermophila.

Genes Dev. 2016;30: 2724–2736. doi:10.1101/gad.290460.116

87.   Miyake A, Beyer J. Cell interaction by means of soluble factors (gamones) in conjugation

of Blepharisma intermedium. Exp Cell Res. 1973;76: 15–24. doi:10.1016/0014-

4827(73)90413-8

88.   Andersen RA. Algal Culturing Techniques. 1st Edition. 2004.

89.   Miyake A, Harumoto T, Salvi B, Rivola V. Defensive function of pigment granules in

Blepharisma japonicum. Eur J Protistol. 1990;25: 310–315. doi:10.1016/S0932-

4739(11)80122-7

90.   Miyake A. Cell interaction by gamones in blepharisma. Sexual interactions in eukaryotic

microbes. Elsevier; 1981. pp. 95–129. doi:10.1016/B978-0-12-524160-1.50010-2

91.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

doi:10.1038/nmeth.2019

92.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using

repeat graphs. Nat Biotechnol. 2019;37: 540–546. doi:10.1038/s41587-019-0072-8

93.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34:

1292   3094–3100. doi:10.1093/bioinformatics/bty191

1293   94.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious

1294   Basic: an integrated and extendable desktop software platform for the organization and

1295   analysis of sequence data. Bioinformatics. 2012;28: 1647–1649.

1296   doi:10.1093/bioinformatics/bts199

1297   95.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes de novo

1298   assembler. Curr Protoc Bioinformatics. 2020;70: e102. doi:10.1002/cpbi.102

1299   96.   Vaser R, Sikic M. Yet another de novo genome assembler. BioRxiv. 2019.

1300   doi:10.1101/656306

1301   97.   Swart EC, Serra V, Petroni G, Nowacki M. Genetic Codes with No Dedicated Stop

1302   Codon: Context-Dependent Translation Termination. Cell. 2016;166: 691–702.

1303   doi:10.1016/j.cell.2016.06.020

1304   98.   Gruber-Vodicka HR, Seah BKB, Pruesse E. phyloFlash: Rapid Small-Subunit rRNA

1305   Profiling and Targeted Assembly from Metagenomes. mSystems. 2020;5.

1306   doi:10.1128/mSystems.00920-20

1307   99.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA

1308   ribosomal RNA gene database project: improved data processing and web-based tools.

1309   Nucleic Acids Res. 2013;41: D590-6. doi:10.1093/nar/gks1219

1310   100.   Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic

1311   gene recognition and translation initiation site identification. BMC Bioinformatics.

1312   2010;11: 119. doi:10.1186/1471-2105-11-119

1313   101.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

1314   genome assemblies. Bioinformatics. 2013;29: 1072–1075.

1315   doi:10.1093/bioinformatics/btt086

1316   102.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

1317   2012;9: 357–359. doi:10.1038/nmeth.1923

1318   103.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of

1319   SAMtools and BCFtools. Gigascience. 2021;10. doi:10.1093/gigascience/giab008

1320   104.   Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing. v2

1321   ed. arXiv; 2012 Jul p. arXiv:1207.3907 [q-bio.GN]. Available:

1322   https://arxiv.org/abs/1207.3907

1323   105.   Harris RS, Cechova M, Makova KD. Noise-cancelling repeat finder: uncovering tandem

1324   repeats in error-prone long-read sequencing data. Bioinformatics. 2019;35: 4809–4811.

1325   doi:10.1093/bioinformatics/btz484

1326   106.   Seah BKB, Swart EC. BleTIES: Annotation of natural genome editing in ciliates using

1327   long read sequencing. BioRxiv. 2021. doi:10.1101/2021.05.18.444610

1328   107.   Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely

1329   available Python tools for computational molecular biology and bioinformatics.

1330   Bioinformatics. 2009;25: 1422–1423. doi:10.1093/bioinformatics/btp163

1331   108.   Dale RK, Pedersen BS, Quinlan AR. Pybedtools: a flexible Python library for

1332   manipulating genomic datasets and annotations. Bioinformatics. 2011;27: 3423–3424.

1333   doi:10.1093/bioinformatics/btr539

1334   109.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

1335   features. Bioinformatics. 2010;26: 841–842. doi:10.1093/bioinformatics/btq033

1336   110.   Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007;9: 90–95.

1337   doi:10.1109/MCSE.2007.55

1338   111.   Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: a sequence logo generator.

1339   Genome Res. 2004;14: 1188–1190. doi:10.1101/gr.849004

1340   112.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and

1341   genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37: 907–915.

1342   doi:10.1038/s41587-019-0201-4

1343   113.   Dutilh BE, Jurgelenaite R, Szklarczyk R, van Hijum SAFT, Harhangi HR, Schmid M, et

1344   al. FACIL: Fast and Accurate Genetic Code Inference and Logo. Bioinformatics.

1345   2011;27: 1929–1933. doi:10.1093/bioinformatics/btr316

1346   114.   Eddy SR. Accelerated profile HMM searches. PLoS Comput Biol. 2011;7: e1002195.

1347   doi:10.1371/journal.pcbi.1002195

1348   115.   Saudemont B, Popa A, Parmley JL, Rocher V, Blugeon C, Necsulea A, et al. The fitness

1349   cost of mis-splicing is the main determinant of alternative splicing patterns. Genome Biol.

1350   2017;18: 208. doi:10.1186/s13059-017-1344-6

1351   116.   Kalvari I, Argasinska J, Quinones-Olvera N, Nawrocki EP, Rivas E, Eddy SR, et al. Rfam

1352   13.0: shifting to a genome-centric resource for non-coding RNA families. Nucleic Acids

1353   Res. 2018;46: D335–D342. doi:10.1093/nar/gkx1038

1354   117.   Chan PP, Lin BY, Mak AJ, Lowe TM. tRNAscan-SE 2.0: Improved Detection and

1355   Functional Classification of Transfer RNA Genes. BioRxiv. 2019. doi:10.1101/614032

1356   118.   Lopez-Delisle L, Rabbani L, Wolff J, Bhardwaj V, Backofen R, Grüning B, et al.

1357   pyGenomeTracks: reproducible plots for multivariate genomic datasets. Bioinformatics.

1358   2021;37: 422–423. doi:10.1093/bioinformatics/btaa692

1359   119.   Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, et al.

1360   BUSCO applications from quality assessments to gene prediction and phylogenomics.

1361   Mol Biol Evol. 2018;35: 543–548. doi:10.1093/molbev/msx319

1362 120. Törönen P, Medlar A, Holm L. PANNZER2: a rapid functional annotation web server.

1363    Nucleic Acids Res. 2018;46: W84–W88. doi:10.1093/nar/gky350

1364 121. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, et

1365    al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology

1366    resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019;47:

1367    D309–D314. doi:10.1093/nar/gky1085

1368 122. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

1369    assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–930.

1370    doi:10.1093/bioinformatics/btt656

1371 123. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

1372    expression analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140.

1373    doi:10.1093/bioinformatics/btp616

1374 124. Li B, Ruotti V, Stewart RM, Thomson JA, Dewey CN. RNA-Seq gene expression

1375    estimation with read mapping uncertainty. Bioinformatics. 2010;26: 493–500.

1376    doi:10.1093/bioinformatics/btp692

1377 125. Wagner GP, Kin K, Lynch VJ. Measurement of mRNA abundance using RNA-seq data:

1378    RPKM measure is inconsistent among samples. Theory Biosci. 2012;131: 281–285.

1379    doi:10.1007/s12064-012-0162-3

1380 126. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple

1381    sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30: 3059–

1382    3066. doi:10.1093/nar/gkf436

1383 127. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

1384    improvements in performance and usability. Mol Biol Evol. 2013;30: 772–780.

1385    doi:10.1093/molbev/mst010

128.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms

and methods to estimate maximum-likelihood phylogenies: assessing the performance

of PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

129.   Stover NA, Punia RS, Bowen MS, Dolins SB, Clark TG. Tetrahymena Genome

Database Wiki: a community-maintained model organism database. Database (Oxford).

2012;2012: bas007. doi:10.1093/database/bas007

130.   Wang R-L, Miao W, Wang W, Xiong J, Liang A-H. EOGD: the Euplotes octocarinatus

genome database. BMC Genomics. 2018;19: 63. doi:10.1186/s12864-018-4445-z

131.   Arnaiz O, Meyer E, Sperling L. ParameciumDB 2019: integrating genomic data across

the genus for functional and evolutionary biology. Nucleic Acids Res. 2020;48: D599–

D605. doi:10.1093/nar/gkz948

132.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007;24:

1586–1591. doi:10.1093/molbev/msm088

133.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.

Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23: 2947–2948.

doi:10.1093/bioinformatics/btm404

134.   UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. Nucleic

Acids Res. 2021;49: D480–D489. doi:10.1093/nar/gkaa1100

135.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:

genome-scale protein function classification. Bioinformatics. 2014;30: 1236–1240.

doi:10.1093/bioinformatics/btu031

136.   Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.

Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49: D412–D419.

doi:10.1093/nar/gkaa913

137. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. ModelTest-NG: A

New and Scalable Tool for the Selection of DNA and Protein Evolutionary Models. Mol

Biol Evol. 2020;37: 291–294. doi:10.1093/molbev/msz189

138. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et

al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era. Mol Biol Evol. 2020;37: 1530–1534. doi:10.1093/molbev/msaa015

139. Suzuki S. Parthenogenetic conjugation in Blepharisma undulans japonicus Suzuki. Bull

Yamagata Univ Natural Sci. 1957;4: 69–84.

140. Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, et al. OrthoDB

v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes

for evolutionary and functional annotations of orthologs. Nucleic Acids Res. 2019;47:

D807–D811. doi:10.1093/nar/gky1053

141. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo

genome assemblies. Bioinformatics. 2015;31: 3350–3352.

doi:10.1093/bioinformatics/btv383

1425

Figure captions

1426

1427 **Figure 1. *Blepharisma* cells and properties of ciliate MAC genomes**. **A.** *B. stoltei* ATCC

1428 30299 cell stained with anti-alpha tubulin-Alexa488 (depth-color coded red to yellow) and DAPI

1429 (cyan). **B.** Snapshot of a 3D reconstruction (Imaris, Bitplane) from CLSM images of Hoechst

1430 33342 (dsDNA dye, Invitrogen™) fluorescence (Ex405 nm / Em420-470 nm). **C.** Scanning

1431 electron microscope (SEM) image of a *B. stoltei* ATCC 30299 cell (false color).

1432

Figure 2. Nuclear development during *Blepharisma stoltei* conjugation. **A.** Cell

development cycle with inset showing the four main kinds of nuclei, highlighted with arrows.

Illustration of various cell stages adapted from [139]). Detailed staging collected for an RNA-seq

time course is provided in Figure S9. **B.** Schematic of nuclear processes occurring during

conjugation (classified according to, and modified from [9]). Nuclear events occurring before and

up to, but not including fusion of the gametic nuclei (syngamy) are classified into sixteen pre-

gamic stages where the MICs undergo meiosis and the haploid products of meiotic MICs are

exchanged between the conjugating cells, followed by karyogamy. After karyogamy, cells are

classified into 10 stages S (synkaryon), D1 (1^{st} mitosis), I1 (1^{st} interphase), D2 (2^{nd} mitosis), I2

(2^{nd} interphase), D3 (3^{rd} mitosis), I3 (3^{rd} interphase), D4 (4^{th} mitosis), E1 (1^{st} embryonic stage),

E2 (2^{nd} embryonic stage). After E2, the exconjugants divide further and are classified into 6

stages of cell division (CD1-6) which we did not follow here.

1445

1446 **Figure 3. Basic properties of ciliate MAC genomes.** In cell diagrams MACs are green and

1447 MICs are small black dots in close proximity to MACs. Citations for genome properties are in

1448 Data S1.

1449

1450 **Figure 4**. **Alternative telomere addition sites. A.** Mapping of a subset telomere-containing

1451 HiFi reads to a *B. stoltei* MAC genome contig region, with alternative telomere addition sites

1452 (ATASs) shown by blue (5') or mauve (3') arrows. Pink bars at read ends indicate soft-masking

1453 by minimap2, typically of telomeric repeats. **B.** Mean base frequencies in +/- 1 kbp flanking

1454 ATAS junctions.

1455

Figure 5. A gene-dense somatic genome with tiny introns. A. HiFi (DNA) and RNA-seq

coverage across a representative *B. stoltei* ATCC30299 MAC genome contig (Contig_1). Y

scale is linear for HiFi reads and logarithmic (base 10) for RNA-seq. Plus strand (relative to the

contig) RNA-seq coverage is green; minus strand RNA-seq coverage is blue. Between the RNA-

seq coverage graphs each arrow represents a predicted gene, colored green when transcribed

from the sense strand and blue from the antisense one. Two orthogroups classified by

OrthoFinder are shown in orange and pink, respectively. **B.** Distribution of intron lengths from

predicted genes. **C.** Sequence logos for 15 bp introns (splicing frequency > 0.5). **D.** Clustered

orthogroups (Data S2) in the *B. stoltei* MAC genome.

1465

Figure 6. MAC genome-encoded transposases in ciliates and properties of a putative

***Blepharisma* IES excisase. A.** Presence/absence matrix of PFAM transposase domains

detected in predicted MAC genome-encoded ciliate proteins. Ciliate classes are indicated before

the binomial species names. Note that *Ichthyophthirius multifiliis* is not shown because no

domains with i-E-values < 1e-4 were detected. PiggyBac homolog phylogeny with PFAM

domain architecture and gene expression heatmap for *Blepharisma*. **B**. DDE_Tnp_1_7 domain

phylogeny with PFAM domain architecture and gene expression heatmap for *Blepharisma*. **C.**

Cysteine-rich domains of PiggyBac homologs. PBLE transposases: Ago (*Aphis gossypii*); Bmo

(*Bombyx mori*); Cag (*Ctenoplusia agnata*); Har (*Helicoverpa armigera*); Hvi (*Heliothis virescens*);

PB-Tni (*Trichoplusia ni*); Mlu (PiggyBat from *Myotis lucifugus*); PLE-wu (*Spodoptera frugiperda*).

Domesticated PGBD transposases: Oni (*Oreochromis niloticus*); Pny (*Pundamilia nyererei*);

Lia5, Tpb1, Tpb2, Tpb6 and Tpb7 (*Tetrahymena thermophila*); Pgm, PgmL1, PgmL2,

PgmL3a/b/c, PgmL4a/b, PgmL5a/b (*Paramecium tetraurelia*); Tru (*Takifugu rubripes*); Pgbd2,

Pgbd3 and Pgbd4 (*Homo sapiens*).

1480

Figure 7. Phylogeny of ciliate PiggyBac homologs, eukaryotic PBLEs and PGBD5

homologs. Colors indicate eukaryotic lineages: purple=Alveolata, green=Stramenopiles,

cyan=Archaeplastida, yellow=Opisthokonta, blue=Amoebozoa. Clade highlighted in pink

represents all heterotrich Pgms and PgmLs. aBayes values generated by PhyML are

represented as line widths and branch colors. aBayes values lower than 0.9 are represented in

red-grey, with lower values appearing more red. The tree is rooted at the PiggyBac-like element

of *Entamoeba invadens.*

1488

1489

1490

1491

1492

1493

Supplementary figure captions

Figure S1. Analysis of assembly completeness by Benchmarking Universal Single-Copy

Orthologs (BUSCO). Completeness of the *B. stoltei* ATCC 30299 MAC assembly was

estimated by the percentage of BUSCOs found in the assembly with reference to the OrthoDB

v10 alveolate database [140]. The nature of the ortholog-matches is indicated by characters

followed by counts: C (complete orthologs) - light blue, D (duplicated orthologs) - dark blue, F

(fragmented orthologs) - yellow and M (missing orthologs) - red.

1501

Figure S2. Properties of minichromosomes, telomeres, and alternative telomere addition

sites. A. Length distribution of telomeres of telomere-bearing HiFi reads. **B.** Length distribution

of HiFi reads delimited by telomeres. **C**. Diagram of a telomere-bearing read mapped onto

genome reference at an ATAS. Sequence which is ambiguously chromosomal or telomeric is

"junction sequence"; junction coordinate which maximizes telomere repeat length on the read is

the "first identifiable breakpoint"; the coordinate maximizing alignment length to reference is the

"last identifiable breakpoint". The last telomeric unit permutation at the last identifiable

breakpoint is underlined (length 8 bp). **D.** Sequence logos of chromosomal sequence at ATAS

junctions, sorted by which permutation of the telomeric repeat is present (plot labels). Logos are

aligned to the "last identifiable breakpoint" between positions 20 and 21; telomeric repeats on

telomere-bearing reads begin to the left of the breakpoint. **E**. Frequencies of 2-mers in whole

genome (blue), in telomeres (green), and at ATAS junctions (chromosomal side after last

identifiable breakpoint, orange). **F.** Histogram of junction sequence lengths for ATASs in *B.*

stoltei. **G.** Counts of each telomere repeat permutation at ATAS junctions (last identifiable

breakpoint).

1517

Figure S3. **Femto Pulse analyses of *B. stoltei* MAC DNA and mapping of *T. thermophila***

telomere-bearing reads. A. Mapping of PacBio CLR reads with 3 consecutive telomeric

repeats to a representative *T. thermophila* MAC chromosome (Chr_001 from ciliate.org). **B.**

Length distribution of input MAC DNA sizes prior to fragmentation and library preparation

(Femto Pulse; LM = lower maker) - replicate 1. RFU=relative fluorescent units. **C.** Length

distribution of input MAC DNA sizes prior to fragmentation and library preparation (Femto Pulse;

LM = lower maker) - replicate 2.

1525

1526 **Figure S4. POT1 paralog phylogeny, PFAM domain architecture, and gene expression in**

1527 ***Blepharisma*.** Diagram elements as described in Figure 6B.

1528

1529 **Figure S5. *Blepharisma stoltei* genetic code.** Prediction for *B. stoltei* ATCC 30299 MAC

1530 genome by PORC; codons that are stops in the standard genetic code are highlighted in orange.

1531

Figure S6. Intron splicing. A. Distribution of intron splicing fraction of candidate sense introns

in the *B. stoltei* MAC genome. **B**. Distribution of intron splicing fractions of introns according to

intron lengths. **C.** Distribution of intron splicing fraction of candidate antisense introns. **D.**

Sequence logos for all predicted 16 nt introns, and 16 nt introns with "A" at either position -7 or -

6 (counting from the 3' end). The number of introns underlying the logos are indicated to the

right. **E.** Distribution of intron splicing fractions of introns according to intron lengths. **F.** Sample

of RNA-seq reads mapped to a GT-GG intron from gene BSTOLATCC_MAC21551

(Contig_57.g761). Translation in alternative reading frames downstream of the predicted intron

leads to premature stop codons soon after the intron.

1541

1542 **Figure S7**. ***B. stoltei* ATCC30299 MAC genome assembly graph.** Bandage [141]

1543 representation of Flye 2.8.1 assembly graph. Edges corresponding to contigs are colored by

1544 coverage (brightest pink = 160×, black=0×).

1545

Figure S8. Experimental approach for conjugation RNA-seq time series. Complementary

mating type strains of *Blepharisma stoltei* were harvested and cleaned by starving overnight.

The cleaned cultures were treated in a time-staggered format, with gamones of the

complementary mating type, where gamone 2 was a solution of the synthetic gamone 2 calcium

salt and gamone 1 was provided as the cell-free fluid (CFF) harvested from mating-type I cells.

Two sets of time-staggered gamone-treated cultures were used for the time series. Set I,

indicated by the solid line, was mixed and used to observe and collect samples at 0 hours, 2

hours, 6 hours, 26 hours and 30 hours after mixing. Set II, indicated by the dashed lines, was

mixed and used to observe and collect samples at 14 hours, 18 hours, 22 hours and 38 hours

after mixing. Test tubes indicate Trizol samples prepared for RNA-extraction which were stored

at -80 °C before processing. Cells collected for imaging were obtained shortly before the

remainder were transferred into Trizol.

1558

Figure S9. Developmental staging of *B. stoltei* for RNA-seq. A. The classification of nuclear

morphology into stages has been done according to previous descriptions [9]. Nuclear events

occurring before and up to, but not including fusion of the gametic nuclei (syngamy) are

classified into sixteen stages indicated by roman numerals. These are the pre-gamic stages of

conjugation where the MICs undergo meiosis and the haploid products of meiotic MICs are

exchanged between the conjugating cells. Stages after syngamy are classified into 10 stages,

stage S (synkaryon), D1 (1st mitosis), I1(1st interphase), D2 (2nd mitosis), I2 (2nd interphase),

D3 (3rd mitosis), I3 (3rd interphase), D4 (4th mitosis), E1 (1st embryonic stage), E2 (2nd

embryonic stage). After E2, the exconjugants divide further and are classified into 6 stages of

cell division (CD1-6) which we did not follow here. Illustration of various cell stages adapted from

[139]). Stacked bars show the proportion of cells at each time point at different stages of

development, preceded by the number of cells inspected (n).

1571

1572 **Figure S10. DDE_1 and DDE_3 domain transposases in *Blepharisma*. A.** DDE_1 domain

1573 phylogeny with PFAM domain architecture and gene expression heatmap for *Blepharisma*. **B.**

1574 DDE_3 domain phylogeny with PFAM domain architecture and gene expression heatmap for

1575 *Blepharisma*.

1576

1577 **Figure S11. DDE_Tnp_IS1595 and MULE domain transposases in *Blepharisma*. A.**

1578 DDE_Tnp_IS1595 domain phylogeny with PFAM domain architecture and gene expression

1579 heatmap for *Blepharisma*. **B.** MULE domain phylogeny with PFAM domain architecture and

1580 gene expression heatmap for *Blepharisma*.

1581

1582 **Figure S12. Small RNA-related proteins in *Blepharisma*. A.** ResIII, Helicase_c and

1583 Ribonuclease_3 domain phylogeny with PFAM domain architecture and gene expression

1584 heatmap for *Blepharisma*. **B.** PIWI domain phylogeny with PFAM domain architecture and gene

1585 expression heatmap for *B. stoltei*.

1586

1587 **Figure S13. Histones and histone-domain-containing proteins in *Blepharisma*.** Gene

1588 expression heatmaps are shown as in previous figures, are clustered according to major histone

1589 type as classified using HistoneDB domain models. Domains from PFAM and HistoneDB are

1590 shown to the right.

Figure 1.

A.

B.

C.

Figure 2.

A.

Old MAC
Secondary MAC
Developing/ new MAC
New MIC
········· Cell boundary

Genome reorganization

38 h   0 h
26 h
22 h   18 h

Meiosis

Karyogamy

Post-karyogamic stages

B.

Meiotic micronuclei   Somatic micronuclei   Macronucleus

Time (h)

18
S
D1
20
I1
D2
22
I2
D3
24
I3
26
D4
28
E1

New micronuclei   Primary macronuclear anlagen   Secondary macronuclear anlagen   Old macronucleus

Figure 3.

Species	Genome size (~Mb)	Genome architecture	Genes (zygosity)	Codon reassignments		
				UAA	UAG	UGA
Blepharisma stoltei (ATCC 302099)	**41**	Mini-chromosomes	25726 (n)	*	*	W
Stentor coeruleus	77	?	31426 (?)	*	*	*
Paramecium tetraurelia	72	Chromosomes	39642 (n)	Q	Q	*
Tetrahymena thermophila	103	Chromosomes	26258 (n)	Q	Q	*
Euplotes octocarinatus	88	Nano-chromosomes	29076 (n)	*	*	C
Stylonychia lemnae	52	Nano-chromosomes	15102 (n)	Q	Q	*
Oxytricha trifallax	50	Nano-chromosomes	18400 (n)	Q	Q	*
Perkinsus olseni	63	Chromosomes	17342 (4n)	*	*	*

Blepharisma stoltei

Stentor coeruleus

Paramecium tetraurelia

Tetrahymena thermophila

Euplotes octocarinatus

Stylonychia lemnae

Oxytricha trifallax

Perkinsus olseni (non-ciliate)

50 μm

2 μm

Figure 4.

A.

B.

Figure 5.

Figure 6.

Figure 7.

Figure S1.

Figure S2.

Figure S3.

A.

C.

D.

Figure S4.

Domain architecture

PFAM domains

Figure S5.

Figure S6.

Figure S7.

Contig_64

Figure S8.

Figure S9.

Figure S10.

A. DDE_1 domain-containing proteins

B. DDE_3 domain-containing proteins

Figure S11.

A. DDE_Tnp_IS1595 domain-containing proteins

Stentor coeruleus.1 — D _ _
Blepharisma stoltei ATCC 30299.1 — D _ _
Blepharisma stoltei ATCC 30299.2 — DD_
Blepharisma stoltei ATCC 30299.3 — DDE
Blepharisma stoltei ATCC 30299.4 — DDE
Blepharisma stoltei ATCC 30299.5 — DDE
Stentor coeruleus.2 — DDE
Stentor coeruleus.3 — DDE
Tetrahymena thermophila — DD_
Stylonychia, Oxytricha — DDE
Perkinsus marinus — DDE

0.9

Catalytic residues   **Domain architecture**

DDE_Tnp_IS1595

PFAM domains

B. MULE domain-containing proteins

Blepharisma stoltei ATCC 30299.1 — DDE
Blepharisma stoltei ATCC 30299.2 — DDE
Blepharisma stoltei ATCC 30299.3 — DDE
Blepharisma stoltei ATCC 30299.4 — DDE
Blepharisma stoltei ATCC 30299.5 — DDE
Blepharisma stoltei ATCC 30299.6 — DD_
Stylonychia lemnae.1 — DDE
Oxytricha trifallax.1 — DDE
Oxytricha trifallax.2 — DDE
Stylonychia lemnae.2 — DDE
Oxytricha trifallax.3 — DDE
Stylonychia lemnae.3 — DDE
Oxytricha trifallax.4 — DDE
Oxytricha trifallax.5 — DDE
Stylonychia lemnae.4 — DDE
Oxytricha trifallax.6 — DDE
Stylonychia lemnae.5 — DDE
Perkinsus marinus.1,2,3 — DD_, DDE, DDE
Perkinsus marinus.4 — DDE,DDE
Perkinsus marinus.5 — DDE,DDE
Perkinsus marinus.6 — DDE

0.7

Catalytic residues   **Domain architecture**

MULE
SWIM

PFAM domains

Figure S12.

A. Dicers, Dicer-likes and Dicer-derivatives

B. Piwi proteins

Figure S13.