Article Title: Limited plasticity in gene expression for providing or receiving parental care

under different temperatures

Short Running Title: Plasticity in gene expression influencing parenting

Authors: Moss, Jeanette B.[1†‡], Cunningham, Christopher B.[1], McKinney, Elizabeth C.[1], and

Moore, Allen J.[1]

Author Affiliations:

[1] Department of Entomology, University of Georgia, Athens, GA 30602, U.S.A.

[†] Current Affiliation: Department of Evolution, Ecology, and Behavior, University of Illinois,

Urbana, IL 61801, U.S.A.

[‡]Correspondence

Email: JenBMoss@gmail.com

Phone: +1 (301) 471-3649

ABSTRACT

Parenting is thought to evolve to buffer offspring from variable, unpredictable, and

challenging environmental conditions. In the subsocial carrion beetle, *Nicrophorus orbicollis*,

stressful temperatures during parenting do not affect parental behavior despite imposing steep

fitness costs to parents. Here, we ask if plasticity of gene expression underpins this behavioral

stability or facilitates independent compensation by larvae. To test this we characterized gene

expression of parents and offspring before and during active parenting under benign (20°C) and

stressful (24°C) temperatures to identify genes of parents and offspring associated with thermal

response, parenting/being parented, and gene expression plasticity associated with behavioral

stability of parental care. The main effects of thermal and social condition each shaped patterns

of gene expression in females, males, and larvae. In addition, we implicated 79 genes in females

as 'buffering' parental behavior across environments. The majority of these underwent

significant changes in expression in actively parenting mothers at the benign temperature, but not

at the stressful temperature. Our results suggest that neither genetic programs for parenting nor

their effects on offspring gene expression are fundamentally different under stressful conditions,

and that behavioral stability is associated primarily with the maintenance of existing genetic

programs rather than replacement or supplementation. Thus, while selection for compensatory

gene expression could expand the range of thermal conditions parents will tolerate, without

expanding the toolkit of genes involved selection is unlikely to lead to adaptive changes of

function.

Keywords: Insects, Parental Care, Transcriptomics, Parent-Offspring, Climate Change

INTRODUCTION

Parental care evolves to buffer offspring against environmental stress and once evolved, can be instrumental for ensuring success across a range of ecological conditions even beyond those that promoted its evolution (Clutton-Brock, 1991; Royle, Smiseth, & Kölliker, 2012; Wilson, 1975). In the face of rapidly changing environments, phenotypic plasticity in family interactions could contribute to species persistence by allowing parents to invest more when conditions are poor and for offspring to compensate when conditions exceed those that parents can or are willing to buffer. In some cases, this phenotypic plasticity may be immediately detectable as a change in behavior. For example, many birds compensate for hot breeding season conditions by extending their time and effort at the nest, sometimes risking dehydration to maintain stable incubation conditions for embryos (McClintock, Hepp, & Kennamer, 2014; Clauser & McRae, 2016; Sharpe, Bayter, & Gardner, 2021), while offspring have been shown to upregulate metabolic and immune pathways in response to the removal of a caring parent (Körner et al., 2020; Mashoodh, Westoby, & Kilner, 2021; Ziadie, Ebot-Ojong, McKinney, & Moore, 2019) or exposure to pathogens (Körner et al., 2020). However, when parents and offspring are pushed to their physiological limits, maintaining a phenotype stability (i.e., continuing to provide care at the same level) is likely just as important for ensuring offspring success as is altering behavior.

Transcriptomic approaches are useful for examining gene expression plasticity beneath the surface of phenotypes because gene expression is inherently plastic, and gene expression profiles can undergo profound changes while maintaining stable higher-level phenotypes (Eberwine & Kim, 2015; Nijhout, Best, & Reed, 2019; Rivera et al., 2021); that is, many 'genetic programs' can give rise to identical phenotypes. While most research in this area has

focused on the role of plastic gene expression in maintaining physiological performance and

stabilizing morphological traits under stress (Oleksiak, Roach, & Crawford, 2005; Cheviron,

Whitehead, & Brumfield, 2008; Cheviron & Brumfield, 2011; Peck et al. 2015), compensatory

molecular mechanisms are also expected to underpin stability of complex behaviors across

fluctuating environments (Fischer, Hauber, & Bell, 2021). This perspective is valuable to the

study of parental care because parenting evolves to minimize environmental fluctuations and

involves transformational changes of gene expression (Parker et al. 2015; Ray et al. 2016;

Bukhari et a. 2019; Fischer et al. 2019; Lopes & de Brujn), and the ultimate configuration of

expressed genes associated with care may differ depending on environment. Because parental

care is inherently a social exchange and phenotypic plasticity can stem from donors and/or

recipients of care, a comprehensive test of these predictions should profile the gene expression of

parents and offspring simultaneously (Butler & Maruska 2021).

Here, we performed RNA-seq on a subsocial burying beetle, *Nicrophorus orbicollis*, to

explore if plasticity of gene expression helps buffer families against environmental stressors.

There is a strong basis for studying the genetics of parenting and development of burying beetles

(Benowitz, McKinney, Cunningham, & Moore, 2017; Cunningham et al., 2019; Jacobs et al.,

2016; Palmer et al., 2016; Parker et al., 2015; Won et al., 2018), yet the genes responsible for

mounting responses to major environmental stressors while beetles are parenting/being parented

remain uncharacterized. Such mechanisms are particularly relevant in the context of rapid

environmental change, as they could provide targets of selection in populations unable to

behaviorally avoid stressful conditions. Even less understood is the extent to which application

of a secondary stressor may modify core parental responses; i.e., how plasticity in behavioral

gene expression buffers higher level phenotypes.

Higher temperatures during parenting for *N. orbicollis* impose steep reproductive and

survival costs; however, females and males do not change their parental care behavior, type, or

intensity (Moss and Moore, 2021). Whether behavioral stability of parents results from gene

expression changes that alter or supplement the genetic program for parental care, or whether

gene expression changes in offspring may independently compensate for thermal stress, is not

known. Here, we address these possibilities while controlling for any confounding effects of

differences in care. We examined the effects of two thermal environments to characterize

'thermal response' genes (20°C or 24°C, the same as that investigated in Moss and Moore, 2021)

and parent-offspring interactions to characterize 'parenting/being parented' genes (before or

during active parenting), comparing responses of mothers, fathers, and offspring to understand

the similarity of each's response. Our prediction was that thermal stressors and parental care will

each elicit independent gene expression responses and that these will differ between family

members (represented by the main effects of the statistical model). Further, we predicted that

some genes would show distinct patterns of expression change in response to the combination of

stressors to stabilize the behaviors across environments ("buffering genes" represented by the

interaction term of the statistical model); specifically, higher temperatures should significantly

modify expression of core genes for parenting/being parented or elicit distinct changes in gene

expression during family interactions if genetic plasticity facilitates behavioral stability across

environments.

MATERIALS AND METHODS

Study System

Nicrophorus orbicollis is a biparental carrion beetle that breeds on small vertebrate

carcasses in woodlands ranging from southern Canada to northern Texas. At the southeastern

edge of this distribution, daily mean temperatures as low as 18°C and as high as 25°C may arise

over the course of the long summer breeding season (Moss & Moore, 2021). Temperatures at the

upper end of this range (23–25°C) are highly challenging for parents and developing offspring,

with families forced to breed under these conditions suffering shortened adult and larval

lifespans and reduced clutch size larval mass (Moss & Moore, 2021). As in most members of the

genus, *N. orbicollis* expresses a large repertoire of pre- and post-hatching care behaviors (Eggert

& Müller, 1997; Scott, 1998). Carcass preparation begins with the removal of hair and liberal

application of anal excretions, which suppress microbial growth on the prepared 'brood ball.'

Females oviposit throughout the soil surrounding the brood ball, and within 2–3 days these eggs

hatch and larvae migrate through the soil to the brood ball attended by their parents. At this

stage, parents commence directly provisioning begging young *via* regurgitation. While larvae of

N. orbicollis have relatively high starvation tolerance (mean time to starvation: ~17 hours), upon

hatching their capacity to self-feed from a prepared carcass is extremely limited, and they rely on

at least three hours of post-hatching parental care for survival (Capodeanu-Nägler et al. 2018).

This post-hatching stage is likely to be the most energetically demanding stage of care for

parents as it is when attendance to offspring is highest and the stage at which we see the most sex

differences of parental behavior (Moss & Moore, 2021).

Study Design

We used *N. orbicollis* beetles that had been collected locally and reared in the laboratory

for one generation at the University of Georgia, as described in Moss and Moore (2021).

Different beetles were used to generate behavioral (Moss and Moore 2021) and transcriptomic

datasets (this study). Briefly, adults were housed individually in plastic containers (9 cm

diameter, 4 cm deep; Eco Products, Boulder, CO, USA) filled with potting soil and fed organic

ground beef twice weekly *ad libitum*. Stock breeding took place at room temperature (20 ±

0.5°C), and resulting offspring were divided evenly between two incubators starting three days

into pupation for a thermal acclimation period prior to breeding. These incubators were

programmed to long-day light conditions (reverse light:dark 14:10) on ramping temperature

cycles to simulate a diurnal range: the first fluctuating between 21 and 20°C and the second

between 25 and 24°C. Because breeding takes place underground in relatively thermostable

conditions, however, only the lower end of the diurnal range of each thermal treatment (20°C

and 24°C) was used for breeding. Virgin, non-sibling beetles (aged ≥ 14 days post-eclosion)

were paired in plastic boxes (17.2 x 12.7 x 6.4 cm; Pioneer Plastics, Dixon, KY, USA) filled

with approximately 2 cm of moistened soil and a 40–45 g thawed mouse carcass (RodentPro,

Evansville, IN, USA) and moved to either a constant dark temperature-controlled room (20°C) or

incubator (24°C) corresponding to their acclimation environment. Boxes were monitored twice

daily for the presence of eggs, and after 2–3 days eggs were collected into petri dishes containing

damp filter paper and stored at their respective treatment temperatures. Eggs were checked every

four hours between 0800 and 1700 until hatching, which occurs within two days after laying.

In addition to manipulating the thermal environments in which care took place, we also

generated within-temperature treatment controls to disentangle the effects of "active" parenting

(the stage at which parents and larvae interact socially, which we refer to as parenting or direct

parental care) from a general parenting state which includes behavior such as carcass preparation

that does not require social interactions and begins before larvae are present. Thus, we sampled

at two time points: 1) post-larval hatching but before larvae and parents interacted (≤ 16 hours

old larvae); and 2) after the first 24 hours of direct parental care. These larvae were collected

directly from the petri dish immediately when found to have hatched, with parents collected

simultaneously from the carcass they were preparing. Larvae of families for parenting samples,

by contrast, were transferred to brood balls attended by parents when found to have hatched.

These families were left undisturbed for 24 hours so that direct parenting could take place before

collecting both parents and offspring (24–40 hours old larvae) for analysis. Parents in both

treatments had initiated indirect parental care involving preparation of a brood ball for larval

feeding. Timing of sampling parents before interactions coincided with the circadian switch to

active parenting (Oldekop, Smiseth, Piggins, & Moore, 2007).

Sample Collection, Preparation, and RNA Sequencing

175   Our design yielded ten families at 20°C (five before parenting, five during parenting) and

nine families at 24°C (four before parenting, five during parenting). We collected the heads of

adults, consisting of brain and associated "fat body" tissue, as in previous studies (Parker et al

2015). This allowed us to capture gene expression in the two tissue types that influence behavior

in insects. Because of the size of larvae at hatching and because we had no *a-priori* predictions

for the specific tissues affected by interactions with adults, we collected whole bodies. Larval

samples therefore consisted of seven to ten larvae for families before parenting and two to four

larvae for families during parenting. Individual adult samples, both before and during parenting,

were age-matched. Samples were flash frozen in liquid nitrogen and stored at -80°C until RNA

extraction. Extractions were performed following the Qiagen RNAeasy Lipid kit protocols

(Qiagen, Venlo, the Netherlands). Sequencing libraries were prepared from 1.3 μg total RNA

using the Illumina TruSeq mRNA Stranded Library Kit (Illumina, Inc., San Diego, CA, USA)

according to the manufacturer's protocol with standard Illumina adapters and primers.

Sequencing was carried out on a NovaSeq 6000 platform with a 150 bp paired-end protocol,

targeting 40M read pairs per sample.

Read Mapping, Transcriptome Assembly, and Annotation

Reads were trimmed using Trimmomatic (Version 0.39; Bolger et al. 2014) to remove

adapter sequences with default parameters, except low-quality bases (Phred <15 at the leading

and trailing ends and from 4-base sliding windows), and short reads (<36 bp). Reads were

corrected using RCorrector with default parameters (Version 1.0.3.1; Song and Florea 2015).

Quality was assessed using FastQC (Version 0.11.9;

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/; Supplementary Table 1). Paired-end

reads were mapped against a draft reassembly of the WGS data of Benowitz et al. (2017) of *N.*

orbicollis genome (NCBI BioProject PRJNA371654; Supplementary Methods 1). Read mapping

was performed using HISAT2 with default parameters (Version 2.1.0; Kim et al. 2015). A

reference-guided transcriptome was assembled using StringTie (Version 2.1.1; Pertea et al. 2015)

and redundant transcripts were removed using CD-HIT (Version 4.8.1; Li and Godzik 2006)

with a sequence similarity cut-off value of 95%. Finally, we assessed the completeness of the

assembly using BUSCO (Insecta dataset; Version 5.0.0; Simão et al. 2015).

Annotation of contigs was performed using a combination of approaches. Briefly, the

longest isoform of each gene was screened against the annotated and published genome of the

closely related burying beetle *N. vespilloides* (NCBI Assembly Accession No.

GCF_001412225.1; Cunningham et al., 2015) using Magic-BLAST (Version 1.5.0; Boratyn et

al. 2019). Transcripts were also fed through the 'dammit' pipeline (http://dib-

lab.github.io/dammit/), which identifies candidate coding regions in transcripts with

Transdecoder (Version 2.0.1; https://hpc.ilri.cgiar.org/transdecoder-software) and annotates

them, drawing from multiple databases. We used the *N. vespilloides* proteome (Cunningham et

al., 2015) as a user database to search against and also retained any hits to known protein

domains in Pfam-A (Version 32.0; El-Gebali et al. 2019) and orthologs in OrthoDB (Version 10;

Kriventseva et al. 2019). Orthologous genes were retrieved from the OrthoDB database by

searching cluster IDs at the level Polyphaga. Gene ontology (GO) terms for annotated genes

were assigned by using Pfam accession numbers and GO terms of OrthoDB assignment. GO

terms were also supplemented using eggNOG-mapper (Version 5.0; Huerta-Cepas et al. 2019)

and OrthoDB searches at the Order (Insecta) level. Unique GO terms from any source were

retained for a gene.

Transcript Quantification and Filtering

We estimated transcript abundances for each sample separately using StringTie. A

transcript count matrix was extracted with the prepDE.py script provided with StringTie and

imported into R (Version 4.0.3; R Core Development Team 2019) along with sample library

sizes (calculated using Picardtools, Version 2.21.6; http://picard.sourceforge.net) and transcript

lengths. We computed the fragments per kilobase per million (FPKM) matrix with the package

countToFPKM (Version 1.0; Alhendi 2019). For each family member and social state (before or

during parenting), we retained transcripts that were expressed >2 FPKM in more than half of

samples and then combined all unique transcripts across groups in downstream analyses (Bloch

et al., 2018). Finally, we used *IsoformSwitchAnalyzeR* (Version 1.12.0; Vitting-Seerup and

Sandelin 2019) to convert transcript-level counts into gene-level counts.

Differential Expression Analysis

We performed differential expression analysis using *DESeq2* (Bioconductor Version

1.30.0; Love et al. 2014). To cluster samples, we first applied variance stabilizing transformation

using DESeq2 and performed a principal components analysis (PCA) using R's built-in 'prcomp'

function. Visualization of the clustering was done using *DESeq2*'s 'pcaplot' function and

extracted principal components were regressed on each grouping factor (family member,

temperature, parenting, and the statistical interactions among these factors) *via* analysis of

variance (ANOVA using R). Significant differences among family members in the full model

ANOVA led us to analyze males, females, and larvae separately. For each family member,

differential expression was estimated using parametric dispersion. We used the likelihood ratio

test (LRT) implemented in *DESeq2* to compare the goodness-of-fit of several models for each

family member to estimate differential gene expression. Two tests were used to capture gene

expression changes in response to main effects. The first compared a full model fitted with both

temperature and parenting (temperature + parenting) to a reduced model fitted only with

parenting to identify differentially expressed genes mutually affected by temperature across

parenting states (henceforth, 'thermal response' genes). The second compared the same full

model to a reduced model fitted only with temperature to produce a list of differentially

expressed genes mutually affected by parenting across thermal environments (henceforth,

'parenting' genes). The *p*-values that resulted from these tests were adjusted for multiple testing

using the Benjamini-Hochberg correction (Benjamini & Hochberg, 1995) and genes with

adjusted $p < 0.05$ were considered statistically significantly differentially expressed genes.

To assess similarity of gene expression responses to the main effects of temperature and

parenting across family members, we extracted lists of overlapping genes from the intersect of

male, female, and larval parenting/being parented and thermal response gene sets. To determine

whether overlap was significantly greater than expected by chance given variable input sizes, we

further carried out randomization tests. Briefly, 10,000 gene sets were randomly generated for

males, females, and larvae by sampling from the full list of expressed genes in that family

member a number equivalent to the size of the gene set being simulated. These random gene sets

were used to calculate a null distribution of overlap values, and significance was calculated as

the proportion exceeding the observed overlap for each pairwise comparison. While overlap of

the most highly significant genes between two sets lends itself to intuitive interpretation, use of

fixed thresholds can be overly stringent and lead to false negatives. Therefore, we implemented a

complementary threshold-free approach: rank-rank hypergeometric overlap (RRHO) analysis.

This analysis ranks DE genes according to their significance level along with the magnitude and

direction of change to make qualitative assessments of concordance between complete gene sets

independent of p-value cutoff. Results were visualized using the stratified method of the R

package *RRHO2* (Version 1.0; Cahill et al. 2018), with genes ranked according to the

recommended metric of -log10(p-value)*sign(log$_2$ fold change).

Finally, we assessed whether the genetic program for parenting/being parented was

altered or supplemented under the addition of thermal stress in two ways. First, we performed a

differential expression analysis using the LRT approach to identify genes showing distinct

responses to parenting/being parented (substantial differences in magnitude or different signs) in

the presence of thermal stress (henceforth, 'buffering' genes). To obtain this list, we compared a

full model fitted with a statistical interaction term (temperature x parenting) to a reduced model

fitted only with the main effects (temperature + parenting). To further characterize patterns of

differential gene expression across social and temperature groups, we performed an ANOVA

followed by post-hoc contrasts (Tukey HSD) at the level of each gene in the buffering set,

comparing the magnitude of expression and sign of the response at 24°C vs 20°C. Second, we

compared differential expression of parenting/being parented genes between the two thermal

environments. Even if changes of expression of these genes are similar enough across thermal

environments to be captured as main effects, the magnitude of the response may be enhanced or

dampened due to thermal stress. To test this possibility, we estimated log2fold changes of

expression of parenting/being parented genes separately for each temperature treatment using the

'ashr' shrinkage estimator with false discovery rate correction (Stephens 2017) and fit a

regression to these data. A slope different than one allows rejection of the null hypothesis that

responses are generally similar in magnitude across thermal treatments and provides a more

holistic comparison of the treatments.

Functional Enrichment Analysis and Candidate Genes

We queried the differentially expressed gene sets and the overlapping gene sets for

enriched Gene Ontology terms using topGO (Version 2.42.0; Alexa and Rahnenfuhrer 2020)

searching for enrichment in all three gene ontology categories: biological process, molecular

function, and cellular component. The nodeSize parameter was set to 10 to remove GO terms

with fewer than ten annotated genes and only terms with >1 significant gene were retained.

Significance was estimated using a Fisher's exact test following the program's suggested

protocol. To examine the directionality of change of enriched pathways, Z-scores were

calculated following the formula of Walter, Sánchez-Cabo, & Ricote (2015): number of

downregulated genes annotated subtracted from the number of upregulated genes annotated for

the term divided by the square-root of the total number of genes annotated for the term.

RESULTS

Our final transcriptome assembly contained 91% of single-copy conserved nucleotide

orthologs (Complete: 90.5% [Single Copy: 76.4%, Duplicated: 14.1%], Fragmented: 4.2%,

Missing: 5.3%) and a total of 12,406 genes.

Thermal and Social Conditions Shape Gene Expression of Family Members

We visualized differences in gene expression among male parents, female parents, and

offspring using principal components analysis. Parents and offspring separated strongly along

two primary axes (Fig 1A), which together explained 77.8% of total variance in gene expression

across all samples (PC1: 73.08%; PC2: 4.76%; Table 1). Family member (parent vs larvae)

clearly separated along PC1, while separation along PC2 was driven primarily by parenting

(Table 1). The interactive effect of family member and parenting and family member and

temperature were significantly associated with PC2 (Table 1; Fig 1B). Visualizing family

members separately with their own dispersions further clarified how global gene expression

differed among groups. Female gene expression displayed considerable overlap between thermal

environments but was most divided with respect to parenting, particularly in the 20°C

environment (Fig 1C). Male gene expression differentiated most strongly with temperature (Fig

1D). In larvae, behavioral groups formed distinct clusters, with parenting accounting for far

greater among-group differences than temperature (Fig 1E).

Plasticity of Gene Expression Underpins Thermal Response, Parenting/Being Parented, and

Buffering

We used differential expression analysis to characterize plastic changes in gene

expression associated with the main effects of temperature and parenting, as well as their

interaction. We identified genes showing both shared (overlapping) and distinct responses to

main effects across family members, consistent with variation in physiology and behavior

between sexes and life stages. There was also gene expression variation associated with

buffering of behavioral responses, where family members expressing the same behaviors in

different environments showed different patterns of gene expression. These included subtle

expression variation of parenting/being parented genes across temperature treatments, as well as

some changes that occurred specifically in response to the interaction of temperature and

parenting.

i.　Thermal Response Genes

Temperature induced a stronger gene expression response in males (n = 418 differentially

expressed genes; Supplementary Table 2) and larvae (n = 487 differentially expressed genes;

Supplementary Table 3) than in females (n = 284 differentially expressed genes; Supplementary

Table 4). However, male, female, and larval responses to temperature showed statistically

significant overlap in terms of gene identity (male-female: $P < 0.001$; female-larvae: $P < 0.001$;

male-larvae: $P = 0.006$; Fig 2A). Global concordance was pronounced between males and

females (Fig 2C) – both in genes that were up-regulated and down-regulated in response to high

temperature – whereas this signal was weak when comparing females to larvae (Fig 2D) and

males to larvae (Fig 2E).

ii.   Parenting Genes

With the parenting main effect, the highest number of differentially expressed genes was

associated with being parented, with 7,189 genes differentiating larvae before and during

interactions with parents (Supplementary Table 5). Adults had fewer differentially expressed

genes than larvae, and within adults there were sex differences. Fewer differentially expressed

genes were detected in males after the arrival of larvae (n = 63; Supplementary Table 6) than in

females (n =189; Supplementary Table 7). Despite this, females and males showed statistically

significant overlap of gene identities while parenting (Fig 2B; P = 0.044). When considering all

expressed genes, concordance was striking but directional, with overlap between the sexes

concentrated in genes downregulated in both during larval interaction (Fig 2F). Females

interacting with larvae and larvae interacting with parents shared more differentially expressed

genes in common (n = 115) than parenting males and parented larvae (n = 28), although this did

not reach statistical significance (P = 0.059) and neither comparison showed global signatures of

concordance of gene expression (Fig 2G–H).

iii.   Buffering Genes

To evaluate the role of gene expression plasticity in facilitating parental buffering and

offspring compensation under stressful environments, we performed a transcriptome-wide survey

for genes showing distinct responses to parenting/being parented at 24°C versus 20°C, or a

significant interaction between temperature and parenting. More genes met this criterion and

were differentially expressed in females (n = 79; Supplementary Table 8) than in males (n = 4;

Supplementary Table 9) or larvae (n = 26; Supplementary Table 10). To test whether genes

showing plasticity indeed showed stronger and broader expression changes at the more stressful

temperature, the 79 buffering genes in females were subjected to closer examination

(Supplementary Table 11). At the benign temperature (20°C), most genes (n = 57) showed

increases of expression under active parenting, whereas at the higher temperature (24°C) only

five genes showed any change in expression. Hence, most buffering genes fell into two

categories: gene expression levels before parenting were equivalent between thermal treatments

(n = 35) or gene expression levels started out significantly higher at 24°C than at 20°C (n = 32;

e.g., Apolipophorin-III; Fig 3C). Because plasticity was temperature-dependent, gene expression

levels during parenting were either significantly lower at 24°C (n = 29) or did not differ between

thermal environments (n = 45), and only five genes showed significantly higher expression levels

during parenting in the warmer treatment.

We next compared gene expression responses to parenting/being parented across

temperature treatments. Among the core genes for parenting/being parented (genes that were

differentially expressed between parenting states across temperatures), differential expression

was weaker at 24°C relative to 20°C in females (slope = 0.514; Fig3A, B) and males (slope =

0.357), whereas the slope of this regression in larvae was close to one (slope = 0.976), suggesting

that responses of larvae were generally similar in magnitude across thermal environments.

Functional Enrichment Analysis

Functional enrichment of GO terms in the within-family member temperature contrasts

revealed both shared and distinct enriched functional characteristics across females, males, and

larvae. The most significantly enriched GO terms in parents were related to chitin and

carbohydrate metabolism (Supplementary Table 12–13), whereas larvae showed the most

enrichment for cellular housekeeping functions (e.g. ribosome biogenesis and rRNA processing;

Supplementary Table 14). Compared to the 20°C group, females, males, and larvae in the 24°C

group predominantly showed downregulation of functional pathways.

Actively parenting females showed significant positive enrichment in a range of

metabolic, catabolic, and biosynthetic functions compared to their respective non-interacting

controls (Supplementary Table 15). Forty-five of the 64 GO terms that were enriched in the

female parenting gene set were also significantly enriched in the overlap between female

parenting genes and larval parented genes, particularly terms related to the metabolism of

organic, carboxylic, and fatty acids (Supplementary Table 16). Conversely, analysis of GO terms

in males revealed significant downregulation of non-overlapping GO terms (e.g. larval cuticle

patterning; Supplementary Table 17). Larval functional pathways were predominantly

upregulated in response to interactions with parents and spanned many categories, from cellular

organization and differentiation to whole organism development (Supplementary Table 18).

Finally, we examined Gene Ontology term enrichment under the interaction of parenting

and thermal environment. Buffering genes of males (n = 4) showed enrichment in three pathways

related to lipid transport and localization (Supplementary Table 20) while those in larvae (n =

26) showed enrichment in five pathways (Supplementary Table 21), the most significant of

which (synaptic target inhibition and negative chemotaxis) were upregulated in parented larvae

responding to thermal stress. In females interacting with larvae, we found 45 significantly

enriched GO term annotations for genes that were differentially expressed between 20°C and

24°C, including 26 GO terms that were also enriched in the female parenting gene set but in the

opposite direction (Supplementary Tables 12, 19; Fig 3D). Top GO terms related to the oxidation

and catabolism of fatty and organic acids and comprised genes that were significantly

downregulated, on average (Supplementary Table 19).

DISCUSSION

Parenting is expected to evolve to buffer environmental stress, but parenting involves an

interaction between the parent and the offspring. For parenting to be effective, the family

interaction needs to be predictable and the mechanisms responsible for stabilizing these

interactions across variable environments are not well understood. Plasticity of gene expression

could allow families to optimize dynamics for their current environment. If true, then parenting-

related responses to an ecologically relevant stressor should have a gene expression signature.

We test this prediction in a subsocial insect with biparental care. We found that thermal and

social conditions both shape patterns of gene expression in *N. orbicollis* females, males, and

larvae, and that many responses are distinct to specific family members. Further, we found

evidence for some plasticity of gene expression in response to the combination of social and

thermal stressors. Specifically, parenting males and females, but not parented larvae, showed

weaker gene expression responses to interactions with larvae under stressful temperatures. In

females 79 genes responded to parenting in an entirely temperature-dependent manner, showing

significant changes in expression at the benign temperature but not at the stressful temperature.

Taken together with our previous findings of limited behavioral plasticity of *N. orbicollis* parents

exposed to stressful temperatures (Moss & Moore, 2021), this implies that behavioral stability

under stress is associated primarily with maintenance of existing genetic programs, rather than

switching to an alternative or supplemented genetic program for the same complex behavior.

Parental investment and offspring development are sensitive to environmental quality as

poor conditions can impose competing physiological demands between parenting and thermal

response. To capture and compare the gene expression and hence the molecular pathways

involved in responses to thermal stress, we sampled families from a benign (20°C) and a

stressful (24°C) thermal environment across the breeding cycle. While our previous work

showed that *N. orbicollis* do not modify their behavior in response to temperature (Moss and

Moore 2021), constitutive changes of gene expression of adults and larvae exposed to high

temperatures across stages of care suggest that any phenotypic stabilization over this thermal

range is at least partly due to the buffering action of background physiological processes. In

parents, the predominant gene expression response to heat stress was downregulation,

particularly of pathways involved in cuticle formation and carbohydrate metabolism

(Supplementary Tables 12–13). Similarly, larvae showed downregulation of many pathways

involved in cellular housekeeping functions (Supplementary Table 14). These findings are

consistent with several long-term acclimation experiments reporting genome-wide

downregulation in response to extreme heat stress, which could either point to an inability to

mount systemic responses (i.e., due to insufficient evolutionary history of such extremes; Levine,

Eckert, & Begun 2011; Becker et al., 2018) or an adaptive molecular mechanism for metabolic

compensation (Yampolsky et al. 2014). The high temperature applied here was at the upper limit

but still within the range experienced in the habitat our beetles were collected (Moss and Moore,

2021), suggesting the latter as a more likely explanation. In sum, thermal stress appears to be

mitigated through the regulation of different functional pathways in parents vs larvae but follows

the same general pattern – tempering all gene expression responses at high temperatures – which

could impose carry-over effects on gene expression underlying parent-offspring and offspring-

parent interactions

With our study, we provide a glimpse into these mechanisms by comparing the gene

expression profiles of family members before and 24 hours after offspring colonize the brood

ball. Thousands of differentially expressed genes were detected in larvae spanning many

functional categories, some due to differences of sampling ages dictated by experimental design.

While we cannot disentangle the direct effects being parented *per se* on gene expression

(Capodeanu-Nägler et al., 2018; Mashoodh et al., 2021) from indirect effects associated with the

rapid development and growth occurring at this early stage (Won et al., 2018), our sampling

design was biologically realistic given that orphaned *N. orbicollis* larvae usually succumb to

starvation within a day (Capodeanu-Nägler et al. 2018). In comparison, changes of parental gene

expression associated with having larvae to parent were subtle (~100–200 differentially

expressed genes), in line with the hypothesis that the modification of behavior within a state

involves a much smaller subset of the genes required to transition between states (i.e., *N.*

vespilloides transitioning into parenting from a non-reproductive state differentially express up to

650 genes*; Parker et al. 2015, Cunningham et al. 2017, Bell, Bukhari, & Sanogo, 2016;

Benowitz, McKinney, Cunningham, et al., 2017; Cunningham et al., 2019).

As in previous work on the related *N. vespilloides* (Parker et al. 2015), males and females

parenting together showed strong concordance in overall gene expression patterns even though,

as has been found for every study of parenting in male burying beetles, we detected fewer

significantly differentially expressed genes for males. More so than males, however, actively

parenting females showed intriguing patterns of overlap (Fig 3B) and concordance (Fig 3G) with

parented larvae in terms of differentially expressed genes and pathways (i.e., males showed anti-

concordance with larvae; Fig 3H). Given that mothers in biparental pairs appear to perform a

disproportionate share of larval provisioning (Moss & Moore, 2021), we propose that these

overlapping pathways, which include lipid and carboxylic acid metabolism, could play a role in

self-feeding (i.e., from a shared resource) and/or nutrient exchange (i.e., mouth-to-mouth

regurgitation). In either case, genes with shared or reciprocal functional roles between family

members could serve as important targets of selection during adaptation to stressful

environments, which may demand more efficient performance of caregivers, offspring, or both.

Finally, we characterized patterns of gene expression in response to the combined

stressors of parenting/being parented and thermal stress. If plasticity of gene expression is

important for maintaining behavioral stability of families across environments, then genes

underlying parent-offspring interactions should be expressed differently or include

more/different genes dependent on environment. The first way we tested this was by comparing

the magnitude of differential expression of parenting/being parented genes between 24°C and

20°C. We predicted that if plasticity of gene expression underlies the ability of parents to buffer

environmental stress or for offspring to compensate independently, then responses would be

more pronounced at the higher temperature. Instead, we found that genes for receiving care (i.e.,

in larvae) were differentially expressed to the same degree regardless of temperature and that

genes for providing care (i.e., in males and females) showed less robust expression changes at

24°C than at 20°C (Fig 3A,B). Weaker differential expression at 24°C is nevertheless a form of

plasticity, and like global gene expression responses to thermal stress, could reflect adaptive

metabolic compensation by parents.

Another possibility is that plastic responses to parenting/being parented mediated by

thermal stress involve completely different genes to those captured by the main effects of

parenting. If true, then many differentially expressed genes should be captured by the interaction

between parenting and temperature, and changes in expression of these genes before and during

parenting should be more pronounced and/or change their sign at the higher temperature. Overall

few genes satisfied this condition in any family member. While not entirely surprising given that

interaction terms have lower statistical power than main effects, this suggests that individuals in

the same behavioral state but at a different temperature still rely on more-or-less the same

genetic program. The largest gene expression response occurred in heat stressed actively

parenting mothers, and closer inspection of this 'buffering' gene set (n = 79) revealed the nature

of this plasticity. Genes that were differentially expressed at 20°C (i.e., significantly increased in

expression during parenting) did not change expression or a temporal shift followed by sustained

expression (i.e., increased before parenting but then showed no further change during parenting;

Fig 3C) at 24°C. Thus, while temperature may independently influence the expression of some

parenting genes the effects are not synergistic, and rather than amplify differences between

behavioral states, thermal stress appears to have a tempering effect.

Our inferences based on differential expression analysis were further corroborated by

functional enrichment analysis of maternal buffering genes, which revealed significant

downregulation of pathways involved in fatty acid oxidation and lipid processing

(Supplementary Table 18). Lipids are the principal macronutrient used for fueling energetically

intensive behavior in insects (e.g. flight: Canavoso et al. 2003), yet previous efforts to link post-

hatching care (i.e., presumed to be the costliest form of care) to increased lipid metabolism in

burying beetles found no relationship (Benowitz, McKinney, Roy-Zokan, Cunningham, &

Moore, 2017). We suggest that females restrict the use of stored energy even more in an

energetically demanding environment, implying that the combined stressors of temperature and

parenting could directly mediate trade-offs between current and future reproduction. In line with

this, actively parenting cichlids exposed to stressful levels of noise show pronounced gene

expression changes in brain regions responsible for homeostatic functioning, suggesting that

stressful environments may trigger a neural switch from offspring-promoting to self-promoting

behaviors (Butler & Maruska 2021). Offspring of stressed mothers in our study in turn

upregulated genes involved in forming the larval serum protein complex – a structure that

sequesters amino acids early in development for the synthesis of adult proteins (Chrysanthis,

Kaliafas, & Mintzas, 1994). Whether this gene expression response by larvae may help

compensate for constrained parental investment or could simply be an indirect effect of plasticity

in mothers remains to be seen. What is clear is that parental gene expression responses to larvae

are tempered at higher temperatures but generally involve the same genes, meaning that the

adaptive function most likely fulfilled by plasticity of gene expression is metabolic

compensation. Thus, the scope for temperature adaptation to proceed *via* selection on alternative

genetic programs underlying care would appear limited.

ACKNOWLEDGEMENTS: We thank PJ Moore, EA Shelby, JT Washington, and KE Kollars

for assistance with colony maintenance and Kathryn E. Kollars for donating her talents of

illustration to create the figures in this manuscript. Robert Schmitz, Hannah Houston, James

Parker, and Tyler Earp assisted with library preparation and Illumina sequencing. Eva Fischer

provided helpful discussions and suggestions regarding statistical analyses and all members of

the Fischer lab provided valuable and constructive feedback on this manuscript. We thank

Romain Libbrecht, Tadeusz Kawecki and an anonymous reviewer for thorough and helpful peer

reviews. The authors declare no conflicts of interest.

References

Alexa, A., & Rahnenfuhrer, J. (2020). topGO: Enrichment analysis for gene ontology. *R Package*

Version 2.42.0.

Alhendi, A. (2019). countToFPKM: Convert counts to fragments per kilobase of transcript per

million (FPKM). *R Package Version 3.1.0.*

Becker, D., Reydelet, Y., Lopez, J. A., Jackson, C., Colbourne, J. K., Hawat, S., … Paul, R. J.

(2018). The transcriptomic and proteomic responses of *Daphnia pulex* to changes in

temperature and food supply comprise environment-specific and clone-specific elements.

BMC Genomics, *19*, 1–30. doi:10.1186/S12864-018-4742-6/FIGURES/15

Bell, A. M., Bukhari, S. A., & Sanogo, Y. O. (2016). Natural variation in brain gene expression

profiles of aggressive and nonaggressive individual sticklebacks. *Behaviour*, *153*, 1723–

1743. doi:10.1163/1568539X-00003393

Benjamini, Y., & Hochberg, Y. (1995). *Controlling the False Discovery Rate: a Practical and*

Powerful Approach to Multiple Testing. J. R. Statist. Soc. B (Vol. 57).

Benowitz, K. M., McKinney, E. C., Cunningham, C. B., & Moore, A. J. (2017). Relating

quantitative variation within a behavior to variation in transcription. *Evolution*, *71*, 1999–

2009. doi:10.1111/evo.13273

Benowitz, K. M., McKinney, E. C., Roy-Zokan, E. M., Cunningham, C. B., & Moore, A. J.

(2017). The role of lipid metabolism during parental care in two species of burying beetle

(*Nicrophorus* spp.). *Animal Behaviour*, *129*, 143–149. doi:10.1016/j.anbehav.2017.05.019

Benowitz, K. M., & Moore, A. J. (2016). Biparental care is predominant and beneficial to

parents in the burying beetle *Nicrophorus orbicollis* (Coleoptera: Silphidae). *Biological*

Journal of the Linnean Society, *119*, 1082–1088. doi:10.1111/bij.12830

Bloch, N. I., Corral-López, A., Buechel, S. D., Kotrschal, A., Kolm, N., & Mank, J. E. (2018).

Early neurogenomic response associated with variation in guppy female mate preference.

Nature Ecology & Evolution 2018 2, 1772–1781. doi:10.1038/s41559-018-0682-4

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*, *30*, 2114–2120. doi:10.1093/bioinformatics/btu170

Boratyn, G. M., Thierry-Mieg, J., Thierry-Mieg, D., Busby, B., & Madden, T. L. (2019). Magic-

BLAST, an accurate RNA-seq aligner for long and short reads. *BMC Bioinformatics*, *20*,

405. doi:10.1186/s12859-019-2996-x

Bukhari, S. A., Saul, M. C., James, N., Bensky, M. K., Stein, L. R., Trapp, R., & Bell, A. M.

(2019). Neurogenomic insights into paternal care and its relation to territorial aggression.

Nature Communications, 10, 4437. doi:10.1038/s41467-019-12212-7

Cahill, K. M., Huo, Z., Tseng, G. C., Logan, R. W., & Seney, M. L. (2018). Improved

identification of concordant and discordant gene expression signatures using an updated

rank-rank hypergeometric overlap approach. *Scientific Reports 8*, 1–11.

doi:10.1038/s41598-018-27903-2

Canavoso, L. E., Jouni, Z. E., Karnas, K. J., Pennington, J. E., & Wells, M. A. (2003). Fat

metabolism in insects. *Annual Review of Nutrition*, *21*, 23–46.

doi:10.1146/ANNUREV.NUTR.21.1.23

Capodeanu-Nägler, A., Prang, M. A., Trumbo, S. T., Vogel, H., Eggert, A. K., Sakaluk, S. K., &

Steiger, S. (2018). Offspring dependence on parental care and the role of parental transfer of

oral fluids in burying beetles. *Frontiers in Zoology*, *15*, 33. doi:10.1186/s12983-018-0278-5

Cheviron, Z. A., & Brumfield, R. T. (2011). Genomic insights into adaptation to high-altitude

environments. *Heredity 108*, 354–361. doi:10.1038/hdy.2011.85

Cheviron, Z. A., Whitehead, A., & Brumfield, R. T. (2008). Transcriptomic variation and

plasticity in rufous-collared sparrows (*Zonotrichia capensis*) along an altitudinal gradient.

Molecular Ecology, *17*, 4556–4569. doi:10.1111/J.1365-294X.2008.03942.X

Chrysanthis, G., Kaliafas, A. D., & Mintzas, A. C. (1994). Biosynthesis and tissue distribution of

four major larval serum proteins during development of *Ceratitis capitata* (Diptera). *Insect*

Biochemistry and Molecular Biology, *24*, 811–818. doi:10.1016/0965-1748(94)90109-0

Clauser, A. J., & McRae, S. B. (2016). Plasticity in incubation behavior and shading by king rails

Rallus elegans in response to temperature. *Journal of Avian Biology, 48,* 479–488. doi:

10.1111/jav.01056

Clutton-Brock, T. H. (1991). *The evolution of parental care*. Princeton, New Jersey, U.S.A.:

Princeton University Press.

Cunningham, C. B., Badgett, M. J., Meagher, R. B., Orlando, R., & Moore, A. J. (2017).

Ethological principles predict the neuropeptides co-opted to influence parenting. *Nature*

Communications, *81*, 4225. doi:10.1038/ncomms14225

Cunningham, C. B., Ji, L., McKinney, E. C., Benowitz, K. M., Schmitz, R. J., & Moore, A. J.

(2019). Changes of gene expression but not cytosine methylation are associated with male

parental care reflecting behavioural state, social context and individual flexibility. *Journal*

of Experimental Biology, *222*, 1–9. doi:10.1242/jeb.188649

Cunningham, C. B., Ji, L., Wiberg, R. A. W., Shelton, J., McKinney, E. C., Parker, D. J., …

Moore, A. J. (2015). The genome and methylome of a beetle with complex social behavior,

Nicrophorus vespilloides (coleoptera: Silphidae). *Genome Biology and Evolution*, *7*, 3383–

3396. doi:10.1093/gbe/evv194

Eberwine, J., & Kim, J. (2015). Cellular deconstruction: Finding meaning in individual cell

variation. *Trends in Cell Biology*, *25*, 569–578. doi:10.1016/J.TCB.2015.07.004

Eggert, A.-K., & Müller, J. K. (1997). Biparental care and social evolution in burying beetles:

lessons from the larder. In J. C. Choe & B. I. Crespi (Eds.), *The Evolution of Social*

Behavior in Insects and Arachnids (pp. 216–236). Cambridge, Massachusetts, USA:

Cambridge University Press. doi:10.1017/cbo9780511721953.011

El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C., … Finn, R. D.

(2019). The Pfam protein families database in 2019. *Nucleic Acids Research*, *47*, D427–

D432. doi:10.1093/nar/gky995

Fischer, E. K., Roland, A. B., Moskowitz, N. A., Tapia, E. E., Summers, K., Coloma, L. A., &

O'Connell, L. A. (2019). The neural basis of tadpole transport in poison frogs. *Proceedings*

of the Royal Society B: Biological Sciences, 286, 20191084. doi:10.1098/rspb.2019.1084

Fischer, E. K., Hauber, M. E., & Bell, A. M. (2021). Back to the basics? Transcriptomics offers

integrative insights into the role of space, time and the environment for gene expression and

behaviour. *Biology Letters*, *17*, 20210293. doi:10.1098/RSBL.2021.0293

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., …

Bork, P. (2019). EggNOG 5.0: A hierarchical, functionally and phylogenetically annotated

orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, *47*,

D309–D314. doi:10.1093/nar/gky1085

Jacobs, C. G. C., Steiger, S., Heckel, D. G., Wielsch, N., Vilcinskas, A., & Vogel, H. (2016).

Sex, offspring and carcass determine antimicrobial peptide expression in the burying beetle.

Scientific Reports 6, 1–8. doi:10.1038/srep25409

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low

memory requirements. *Nature Methods*, *12*, 357–360. doi:10.1038/nmeth.3317

Körner, M., Vogelweith, F., Libbrecht, R., Foitzik, S., Feldmeyer, B., & Meunier, J. (2020).

Offspring reverse transcriptome responses to maternal deprivation when reared with

pathogens in an insect with facultative family life. *Proceedings of the Royal Society B*, *287,*

20200440. doi:10.1098/RSPB.2020.0440

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., &

Zdobnov, E. M. (2019). OrthoDB v10: Sampling the diversity of animal, plant, fungal,

protist, bacterial and viral genomes for evolutionary and functional annotations of

orthologs. *Nucleic Acids Research*, *47*, D807–D811. doi:10.1093/nar/gky1053

Levine, M. T., Eckert, M. L., & Begun, D. J. (2011). Whole-genome expression plasticity across

tropical and temperatee *Drosophila melanogaster* populations from Eastern Australia.

Molecular Biology and Evolution, 28, 249–256. doi.org/10.1093/molbev/msq197

Li, W., & Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics*, *22*, 1658–1659.

doi:10.1093/bioinformatics/btl158

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*, 550. doi:10.1186/s13059-

014-0550-8

Butler, J. M., & Maruska, K. P. (2021). Noise during mouthbrooding impairs maternal care

behaviors and juvenile development and alters brain transcriptomes in the African cichlid

fish *Astatotilapia burtoni. Genes, Brain and Behavior, 20,* e12692. doi: 10.1111/gbb.12692

Mashoodh, R., Westoby, J., & Kilner, R. (2021). Evolved changes in DNA methylation in

response to the sustained loss of parental care in the burying beetle. *BioRxiv*.

doi:10.1101/2021.03.25.436923

McClintock, M. E., Hepp, G. R., & Kennamer, R. A. (2014). Plasticity of incubation behaviors

helps Wood Ducks (*Aix sponsa*) maintain an optimal thermal environment for developing

embryos. *The Auk*, *131*, 672–680. doi:10.1642/AUK-14-57.1

Moss, J. B., & Moore, A. J. (2021). Constrained flexibility of parental cooperation limits

adaptive responses to harsh conditions. *Evolution*, *75*, 1835–1849. doi:10.1111/EVO.14285

Nijhout, H. F., Best, J. A., & Reed, M. C. (2019). Systems biology of robustness and homeostatic

mechanisms. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*, *11*, e1440.

doi:10.1002/WSBM.1440

Oldekop, J. A., Smiseth, P. T., Piggins, H. D., & Moore, A. J. (2007). Adaptive switch from

infanticide to parental care: How do beetles time their behaviour? *Journal of Evolutionary*

Biology, *20*, 1998–2004. doi:10.1111/J.1420-9101.2007.01364.X

Oleksiak, M. F., Roach, J. L., & Crawford, D. L. (2005). Natural variation in cardiac metabolism

and gene expression in *Fundulus heteroclitus. Nature Genetics, 37,* 67–72. doi:

10.1038/ng1483

Palmer, W. J., Duarte, A., Schrader, M., Day, J. P., Kilner, R., & Jiggins, F. M. (2016). A gene

associated with social immunity in the burying beetle *Nicrophorus vespilloides*.

Proceedings of the Royal Society B: Biological Sciences, *283*(1823), 20152733.

doi:10.1098/rspb.2015.2733

Parker, D. J., Cunningham, C. B., Walling, C. A., Stamper, C. E., Head, M. L., Roy-Zokan, E.

M., … Moore, A. J. (2015). Transcriptomes of parents identify parenting strategies and

sexual conflict in a subsocial beetle. *Nature Communications*, *6*, 8449.

doi:10.1038/ncomms9449

Peck, L. S., Thorne, M. A. S., Hoffman, J. I., Morley, S. A., & Clark, M. S. (2015). Variability

among individuals is generated at the gene expression level. *Ecology, 96,* 2004–2014. doi:

10.1890/14-0726.1.

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T., & Salzberg, S. L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

Nature Biotechnology, *33*, 290–295. doi:10.1038/nbt.3122

R Core Development Team. (2019). R: A language and environment for statistical computing. *R*

Foundation for Statistical Computing.

Ray, S., Tzeng, R. Y., DiCarlo, L. M., Bundy, J. L., Vied, C., Tyson, G., Nowakowski, R., &

Arbeitman, M. N. (2016). An examination of dynamic gene expression changes in the

mouse brain during pregnancy and the postpartum period. *G2, 6,* 221–233. doi:

10.1534/g3.115.020982

Rivera, H. E., Aichelman, H. E., Fifer, J. E., Kriefall, N. G., Wuitchik, D. M., Wuitchik, S. J. S.,

& Davies, S. W. (2021). A framework for understanding gene expression plasticity and its

influence on stress tolerance. *Molecular Ecology*, *30*, 1381–1397. doi:10.1111/MEC.15820

Royle, N. J., Smiseth, P. T., & Kölliker, M. (2012). *The Evolution of Parental Care*. Oxford,

U.K.: Oxford University Press.

Scott, M. P. (1998). The ecology and behavior of burying beetles. *Annual Review of Entomology*,

43, 595–618. doi:10.1146/annurev.ento.43.1.595

Scott, M. P., & Traniello, J. F. A. (1990). Behavioural and ecological correlates of male and

female parental care and reproductive success in burying beetles (*Nicrophorus spp.*).

Animal Behaviour, *39*, 274–283. doi:10.1016/S0003-3472(05)80871-1

Sharpe, L. L., Bayter, C., & Gardner, J. L. (2021). Too hot to handle? Behavioural plasticity

during incubation in a small, Australian passerine. *Journal of Thermal Biology*, *98*, 102921.

doi:10.1016/J.JTHERBIO.2021.102921

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics*, *31*, 3210–3212. doi:10.1093/bioinformatics/btv351

Song, L., & Florea, L. (2015). Rcorrector: Efficient and accurate error correction for Illumina

RNA-seq reads. *GigaScience*, *4*(1), 48. doi:10.1186/s13742-015-0089-y

Trumbo, S. T. (1991). Reproductive benefits and the duration of paternal care in a biparental

burying beetle, *Nicrophorus orbicollis*. *Behaviour*, *117*, 82–105.

doi:10.1163/156853991X00139

Vitting-Seerup, K., & Sandelin, A. (2019). IsoformSwitchAnalyzeR: analysis of changes in

genome-wide patterns of alternative splicing and its functional consequences.

Bioinformatics, *35*(21), 4469–4471. doi:10.1093/bioinformatics/btz247

Walter, W., Sánchez-Cabo, F., & Ricote, M. (2015). GOplot: an R package for visually

combining expression data with functional analysis. *Bioinformatics*, *31*, 2912–2914.

doi:10.1093/BIOINFORMATICS/BTV300

Wilson, E. O. (1975). *Sociobiology: The New Synthesis*. Cambridge, U.K.: Belknap.

Won, H. I., Schulze, T. T., Clement, E. J., Watson, G. F., Watson, S. M., Warner, R. C., …

Davis, P. H. (2018). De novo assembly of the burying beetle *Nicrophorus orbicollis*

(Coleoptera: Silphidae) transcriptome across developmental stages with identification of

key immune transcripts. *Journal of Genomics*, *6*, 41. doi:10.7150/JGEN.24228

Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender,

M. E. (2014). Functional genomics of acclimation and adaptation in response to thermal

stress in *Daphnia*. *BMC Genomics, 15,* 859. doi: 10.1186/1471-2164-15-859

Ziadie, M. A., Ebot-Ojong, F., McKinney, E. C., & Moore, A. J. (2019). Evolution of personal

and social immunity in the context of parental care. *American Naturalist*, *193*, 296–308.

doi:10.1086/701122

DATA ACCESSIBILITY: All sequencing data is deposited in NCBI GenBank under the

BioProject ID PRJA71530. Raw sequence reads are deposited in the Sequence Read Archive

(SRA) under submission SUB9220525 and Accession numbers SAMN18349415–

SAMN18349472 and the assembled transcriptome is deposited in the Transcriptome Shotgun

Assembly (TSA) Sequence Database under submission SUB9312680. Supporting data and R

scripts are deposited as an Open Science Framework project: osf.io/ztyqa.

BENEFITS GENERATED: Benefits from this research accrue from the sharing of our data and

results on public databases as described above.

AUTHOR CONTRIBUTIONS: JBM and AJM conceived and designed the study. JBM, ECM,

CBC, and AJM collected and analyzed the data. JBM, CBC, and AJM wrote the paper with input

from ECM. All authors gave final approval for publication.

Table 1: ANOVA of principal components of overall gene expression.

Variable	PC1 (73.08% variance)		PC2 (4.76% variance)	
	$F_{2,55}$	*P*	$F_{2,55}$	*P*
Family Member	**5346.224**	<0.0001	0.624	0.540
Temp	**4.047**	0.050	**7.073**	0.011
Parenting	**27.238**	<0.0001	**88.311**	<0.0001
Family Member x Temperature	0.499	0.611	**11.406**	<0.0001
Family Member x Parenting	**4.852**	0.012	**92.516**	<0.0001
Temperature x Parenting	2.139	0.151	0.925	0.341
Family Member x Temperature x Parenting	1.653	0.203	1.287	0.286

Figure 1: Principal components analysis (PCA) of normalized transcriptomic profiles across the

three family members (females, males, and larvae) and four treatment groups (20°C:before

parenting, 20°C:during parenting, 24°C:before parenting, and 24°C:during parenting). (A)

Sample clustering along the first two principal components of the full model PCA (~Family

Member*Temperature*Parenting). Eclipses are drawn for 95% confidence intervals. (B–D)

Clustering of female, male, and larval samples, respectively, when analysis is repeated using

only those samples. Illustration of *N. orbicollis* family in Fig 1A by K.E. Kollars.

Figure 2: Concordance in differential gene expression between family members as estimated by

direct overlap of differentially expressed genes (A–B) and rank-rank hypergeometric overlap

(RRHO; C–H). Numbers of shared differentially expressed genes associated with (A) thermal

response and (B) parenting are shown in Venn diagrams for females (red), males (blue), and

larvae (yellow). The bottom panel shows the results of the RRHO analysis of all 12,406 genes

for each of the respective contrasts (20°C vs 24°C: C–E; and with vs without parenting: F–H).

The x (family member 1) and y (family member 2) axes of each plot correspond to the

significance of differential expression (ranked -log10(p-values)) multiplied by the signed

log2fold difference in gene expression for each family member-specific analysis. White

boundaries demarcate the switch from down- to up-regulation between behavioral groups for

each family member. Pixel color corresponds to the value of the -log10(p-value) from the

differential expression analysis, such that hot spots in the plot designate the strength and

directionality of concordance in gene expression between family members.

Figure 3: (A) Normalized expression profiles of the 189 genes that were differentially expressed

in females before and during parenting depicted as heatmaps, with each column representing an

individual sample (grouped by social and temperature treatment) and each row representing a

gene (clustered in a dendrogram by expression similarity). Cell colors represent normalized

count values, such that the visual distinctiveness of clusters of samples corresponds to the

magnitude of difference in their gene expression between groups; (B) Regression of log2fold

changes in expression of 189 genes that were differentially expressed in females before and

during parenting between 24°C and 20°C environments; (C) Distribution of normalized

expression values of Apolipophorin-III across four behavioral groups of females: before

parenting at 20°C, before parenting at 24°C, during parenting at 20°C, and during parenting at

24°C; (D) Directionality of expression (calculated as a z-score) of 26 GO terms identified as

significantly enriched in females in response to parenting only (blue bars) as well as in response

802   to the interaction of parenting and temperature (gold bars). Z-scores correspond to the

803   standardized number of genes annotated for a term that are upregulated relative to

804   downregulated, such that values less than zero correspond to overall downregulation.