1 **A salvaging strategy enables stable metabolite provisioning among free-living bacteria**

3 Sebastian Gude, Gordon J. Pherribo, and Michiko E. Taga[#]

5 Department of Plant & Microbial Biology, University of California, Berkeley, Berkeley, CA 94720

6 USA

8 [#] Correspondence to taga@berkeley.edu

Abstract

All organisms rely on complex metabolites such as amino acids, nucleotides, and cofactors for essential metabolic processes. Some microbes synthesize these fundamental ingredients of life *de novo*, while others rely on uptake to fulfill their metabolic needs. Although certain metabolic processes are inherently 'leaky', the mechanisms enabling stable metabolite provisioning among microbes in the absence of a host remain largely unclear. In particular, how can metabolite provisioning among free-living bacteria be maintained under the evolutionary pressure to economize resources? Salvaging, the process of 'recycling and reusing', can be a metabolically efficient route to obtain access to required resources. Here, we show experimentally how precursor salvaging in engineered *Escherichia coli* populations can lead to stable, long-term metabolite provisioning. We find that salvaged cobamides (vitamin B_{12} and related enzyme cofactors) are readily made available to non-productive population members, yet salvagers are strongly protected from overexploitation due to partial metabolite privatization. We also describe a previously unnoted benefit of precursor salvaging, namely the removal of the non-functional, proliferation-inhibiting precursor. As long as compatible precursors are present, any microbe possessing the terminal steps of a biosynthetic process can, in principle, forgo *de novo* biosynthesis in favor of salvaging. Consequently, precursor salvaging likely represents a potent, yet overlooked, alternative to *de novo* biosynthesis for the acquisition and provisioning of metabolites in free-living bacterial populations.

Introduction

The general mechanisms enabling stable, long-term metabolite provisioning among free-living bacteria remain poorly understood[1-3]. While host-associated bacteria may count on a consistent supply of metabolites from their host, their free-living counterparts may not. Not only do the latter have no reliable access to metabolites provided by a host, but they also lack simple means for positive assortment[4], i.e., the ability of metabolite providers to preferentially supply metabolites to population members that benefit the provider, which is typically enabled by structured environments. In the absence of positive assortment, non-productive population members may easily invade, overexploit, and displace metabolite providers[5-8], causing an irreversible reduction in community diversity and metabolic capacity.

Bacteria require a diverse pool of metabolites. These essential ingredients of life can be obtained through *de novo* biosynthesis or via uptake from the environment[9]. Metabolite provisioning among bacteria is frequently enabled by an intricate network of metabolic interactions[10]. Members of these complex communities can be broadly categorized according to their metabolic archetype for a particular metabolite as producers, degraders, salvagers, dependent consumers, and independents (Fig. 1A). While independents, by definition, do not partake in interactions involving the metabolite, dependent consumers (i.e., non-productive population members) critically rely on metabolically active producers, degraders, or salvagers to fulfill their metabolic needs. Since certain biological functions are inherently 'leaky'[11-14], some metabolites, such as degradation products of extracellularly processed polypeptides and polysaccharides[15,16], are readily supplied as 'public goods'. Yet, this type of metabolite provisioning is inherently limited to particular classes of externally processed polymeric metabolites.

An alternative mode of gaining access to metabolites is salvaging, the process of

'recycling and reusing'[17]. Bacterial salvaging is most commonly associated with iron acquisition[18-

20]. In aerobic environments, many microbes release high-affinity iron-chelating siderophores to

gain access to otherwise insoluble Fe^{3+}. Yet, bacterial salvaging is by no means limited to the

uptake of inorganic substances such as metals. Bacteria can also take up a variety of

environmental metabolites in lieu of synthesizing them *de novo*. Classes of metabolites known to

be salvaged include sugars, amino acids, nucleotides, and vitamins[21-26]. Notably, metabolite

salvaging is not restricted to complete, terminal metabolites. Chemically stable precursors, for

example intermediates of central metabolism or compounds involved in specialized metabolism,

can be salvaged as well[27,28].

While the genetic and biochemical basis of salvaging is well understood for many

metabolites due to in-depth studies in a few bacterial species, less is known about the ecological

ramifications of salvaging in multi-member bacterial populations[21-25,27]. Here, as an example, we

focus on the salvaging of the cobamide precursor, cobinamide (Cbi), and explore how its

salvaging can shape bacterial population dynamics and stability. Cobamides (vitamin B_{12} and

related enzyme cofactors) are cobalt-containing cofactors used in diverse metabolic pathways,

and cobamide precursor salvaging is widely predicted among sequenced bacterial genomes[29]

(Fig. 1B, Supp. Table 1). In this process, the incomplete and non-functional cobamide precursor

Cbi is taken up and, in the presence of the required gene products for cobamide precursor

salvaging, combined with a nucleoside containing a 'lower ligand' base of variable structure to

form a complete, functional cobamide. By studying cobamide precursor salvaging, we

demonstrate that salvaging can be an effective strategy enabling stable metabolite provisioning.

Salvagers readily release assembled complete cobamides into the environment, where they can

be utilized by non-productive population members, yet surprisingly we found that salvagers were

strongly protected from overexploitation. These findings indicate that precursor salvaging may be

a highly effective, generalizable way of enabling metabolite provisioning, even when means for

positive assortment are absent.

Results and Discussion

<u>Salvaged metabolites are readily provisioned</u>

To experimentally investigate precursor salvaging in the context of mixed free-living

bacterial populations, we generated cobamide-dependent *Escherichia coli* populations (Fig. 2A,

see Methods for details). Salvagers (Sal) only proliferated in glycerol minimal medium when either

the complete cobamide vitamin B_{12} (B_{12}) or the incomplete, non-functional cobamide precursor

cobinamide (Cbi) and the lower ligand, 5,6-dimethylbenzimidazole (DMB), which together can be

salvaged and assembled into B_{12} inside the cell, were supplied (Fig. 2B, blue symbols). As

expected, no proliferation was observed in the presence of Cbi and DMB for dependent

consumers (Dep) which have a disrupted cobamide precursor salvaging pathway (Fig. 2B, gray

symbols).

Mixed salvager-dependent populations proliferated in the presence of Cbi and DMB (Fig.

2B). While salvagers were expected to proliferate, it remained unclear whether dependents would

also be able to proliferate given that they rely on provisioning of the complete cobamide B_{12}.

Previous studies reported a need to engineer or evolve 'overproduction' strains to enable

provisioning of metabolites such as amino acids[11,30]. Yet, conversely, a recent report

demonstrated the general potency of vitamins and cofactors to promote stable metabolic

interactions[31]. To dissect individual contributions to the overall increase in population size of the

mixed salvager-dependent population, we established a calibrated fluorescence-based assay to

track the dynamics of each subpopulation in well-stirred culture conditions over time (Supp. Fig.

1). The increase in the total population size was found to be driven by proliferation of both

salvagers and dependents, indicating that the salvaged cobamides were readily released into the

environment as a 'public good' and made available (i.e., provisioned) to the dependents (Fig. 2B

and C).

Details of the proliferation dynamics markedly differed between the two subpopulations.

Salvagers proliferated with no lag and experienced a growth rate of ~0.47 h^{-1} (Fig. 2B and C),

comparable to their performance when cultured alone (Fig. 2B). Notably, dependents experienced

an extended lag before proliferation commenced at a slightly reduced growth rate of ~0.40 h^{-1},

and their final abundance, which they attained at the same time as salvagers ceased to proliferate,

was only about 1/10 of the level achieved by salvagers (Fig. 2C). These findings suggested that

the benefits of the salvaged complete cobamides were asymmetrically accessible to salvagers

and dependents, even though mechanisms for positive assortment[4], such as spatial exclusion of

dependents in structured environments[6], were inaccessible under these (well-stirred) culture

conditions.

<u>Salvaging 'preference' determines population dynamics</u>

We next explored the potential limits of the unprompted cobamide release and

provisioning observed in mixed salvager-dependent populations. We reasoned that salvaging-

based proliferation and metabolite provisioning should depend on the quantity and chemical

identity of the available precursors. To test this hypothesis, we systematically varied external

lower ligand supplementation and monitored proliferation of jointly cultured salvagers and

dependents. Proliferation dose response curves in the presence of Cbi and various levels of DMB

indicated that salvagers successfully proliferated at all tested concentrations of the externally

supplied lower ligand (Fig. 3A). Conversely, proliferation of dependents ceased completely below

100 pM of externally supplied DMB, again pointing towards a preferential access of salvagers to

the assembled cobamide. A similar trend was observed when a chemically different lower ligand,

2-methyladenine (2MA, lower ligand of the cobamide factor A, FA), was supplied (Fig. 3B). Once

more only the proliferation of the dependents ceased at low levels of external lower ligand

supplementation while salvagers continued to proliferate. Notably, a ~300-fold higher level of 2MA

compared to DMB was required to enable dependent proliferation, indicating that chemical identity

indeed played an important role. We reasoned that this difference in dependent proliferation

between the two chemically distinct lower ligands could be caused by a difference in the metabolic

'preference' for the respective complete cobamides, which has been reported for several bacterial

species[32-34]. Alternatively, a difference in the efficiency of the precursor salvaging pathway for

installation of the two lower ligands could be decisive. Proliferation dose response curves of

dependents cultured alone in the presence of B_{12} or FA showed only a minor (~2-fold) difference

in metabolic 'preference' between the two complete cobamides (Fig. 3C). This finding indicates

that the drastic difference in dependent proliferation between the supplementation with the

chemically distinct lower ligands DMB and 2MA was not due to a metabolic 'preference' for the

complete cobamides B_{12} and FA, but was instead mainly caused by a difference in lower ligand

'preference' of the precursor salvaging pathway. Similar observations of a salvaging 'preference'

in bacteria have previously been reported[35,36].

The proliferation dose response curves of jointly cultured salvagers and dependents

contained an additional noteworthy detail (Fig. 3A and B). Namely, proliferation of salvagers was

observed for levels of externally supplied lower ligand that, even if conversion to complete

cobamide was 100% efficient, were below the minimal level of externally supplied complete

cobamide required for proliferation (Fig. 3C). Thus, salvagers were able to generate proliferation-

supporting levels of complete cobamides in the absence of sufficient external lower ligand

supplementation, presumably by incorporating internally produced lower ligands. Using high-

performance liquid chromatography (HPLC) on cell extracts obtained from salvagers cultured in

the presence of Cbi with no externally supplied lower ligand, we identified these 'default'

cobamides as a mixture of FA (lower ligand: 2MA) and pseudo-B_{12} (pB_{12}, lower ligand: Adenine)

based on comparison of their respective retention times and characteristic peak spectra with

purified standards (Fig. 3D, Supp. Fig. 4). Thus, as observed previously[37], salvagers can

supplement limiting external precursor supply by tapping into their internal metabolite pool to

obtain *de novo* synthesized lower ligands (here 2MA and adenine) to generate required complete

cobamides.

<u>Salvaging can eliminate negative effects of metabolite accumulation</u>

Precursor salvaging seemed not to incur a noticeable metabolic cost under our culture

conditions, as salvagers proliferated at comparable growth rates in the presence of either the

complete cobamide B_{12} or the precursors Cbi and DMB (Fig. 2B). Nevertheless, the fact that

metabolically valuable salvaged complete cobamides were readily released and provisioned to

dependents remained puzzling, particularly because it invites overexploitation by dependents

which may cause a collapse of the entire mixed population[5-8]. In an attempt to gain some basic

understanding of the factors that may facilitate the unprompted release of the metabolically

valuable salvaged cobamides, we tried taking a more abstract perspective on the precursor

salvaging process by comparing its functioning to other similarly structured cellular activities. On

the most basic level, cobamide precursor salvagers take up metabolites, modify them

intracellularly, and finally release them back into the extracellular environment. While the

unprompted release of metabolically valuable salvaged metabolites begs an explanation, the

similarly structured process of antibiotic inactivation certainly does not. Here, toxic compounds

are taken up, internally converted into less harmful forms, and subsequently released into the

environment[38]. We wondered whether in addition to its 'constructive' characteristic, i.e., the

generation of complete, functional cobamides, cobamide precursor salvaging could also have

benefits through its 'consuming' characteristic, i.e., the assimilation of the incomplete, non-

functional cobamide precursor Cbi. The latter could be beneficial if the presence of high levels of

precursors interferes with proliferation as has been noted for the accumulation of some

metabolites[39,40].

Proliferation dose response curves of dependents cultured alone on various levels of the

complete cobamide B_{12} in the absence and presence of high levels (10-fold higher than the

highest tested level of B_{12}) of Cbi indeed revealed a clear proliferation-inhibiting effect (Fig. 4A).

The presence of high levels of environmental Cbi not only increased the EC50 of the proliferation

dose response curve by about 5-fold (Fig. 4A, black arrow) but also induced the appearance of a

drastic lag before proliferation was initiated for the lowest B_{12} levels that supported proliferation in

the presence of surplus Cbi (Fig. 4B). Thus, though Cbi is metabolically beneficial as a precursor

for salvaging, it clearly has negative effects when present in excess.

We reasoned that this proliferation-inhibition caused by surplus Cbi could potentially be

overcome, or at least be diminished, by the 'consuming' character of precursor salvaging. Indeed,

salvagers cultured under the same conditions as described above did not exhibit the lag

previously observed for dependents (Supp. Fig. 2, Fig. 4B). Yet, the inhibitory effect of Cbi seen

at higher B_{12} levels remained unchanged (Fig. 4C, black arrow). Cobamide precursor salvaging

in the absence of external lower ligand supplementation is likely limited by the internal availability

of *de novo* synthesized lower ligands, which, under our conditions, could be insufficient to

assimilate sufficient amounts of the proliferation-inhibiting Cbi. As a direct test, we supplied

salvagers with a 10-fold excess of externally supplied lower ligand. Under these conditions

salvagers were indeed able to completely relieve the detrimental effects of surplus Cbi (Fig. 4D).

Consequently, the 'consuming' character of precursor salvaging, i.e., the uptake and assimilation

of proliferation-inhibiting precursors and the subsequent export of complete cobamides, can be

beneficial to salvagers, and thus may (partially) explain the unprompted release of salvaged

metabolites.

To explore whether these Janus-like (i.e., two-sided) benefits of precursor salvaging were

reflected in its regulatory logic, we constructed a fluorescent reporter for the expression of the

major genes involved in the cobamide precursor salvaging process (i.e., cobUST) to determine

whether salvaging is regulated by precursor abundance. While gene expression changed slightly

(~4-fold) over time, we did not observe differential gene expression between conditions that

required precursor salvaging (supplementation of Cbi and DMB) and conditions that did not

(supplementation of B_{12}) (Supp. Fig. 3). Thus, our results suggested that cobamide precursor

salvaging in *E. coli* was not regulated at the level of gene expression under our culturing

conditions, and that the precursor salvaging pathway was expressed independently of the need

for cobamide precursor salvaging.

Partial privatization enables salvagers to persist stably

Unprompted metabolite release in the absence of mechanisms for positive assortment

bears the risk of overexploitation, potentially causing an associated collapse of a metabolically

interconnected population if non-productive members (i.e., dependents) possess a proliferation

advantage over their productive counterparts (i.e., salvagers)[41]. As an initial step to assess the

risk of population collapse in mixed salvager-dependent populations, we created faster-

proliferating mutant variants by moving the salvager and dependent genotypes into an *E. coli*

MG1655 GB-1 mutant background (Sal[GB-1] and Dep[GB-1])[42]. The GB-1 background confers a

growth rate advantage of ~45% compared to the original salvager and dependent in glycerol

minimal medium (compare Figs. 5A and 2B)[42]. Combinatorial culturing of all four possible

salvager-dependent pairs as 1:1 mixtures resulted in expected changes in the final population

composition (Fig. 5B). Specifically, when jointly cultured with Sal, Dep only reached about 1/10 of

the abundance of Sal after 24 hours, in agreement with our previous observations (Fig. 2C).

Jointly cultured Sal[GB-1] and Dep[GB-1] showed a similar pattern, indicating that only relative, and not

absolute, proliferation characteristics determined the population composition. Dep proliferation

was much reduced when it was cultured with the faster proliferating Sal[GB-1] mutant, whereas the

contrary effect was observed if the dependent (Dep[GB-1]) had a proliferation advantage over the

salvager (Sal). Together, these findings demonstrated how relative changes in the maximally

achievable growth rate can drastically reshape the composition of metabolically linked

populations. Moreover, it suggested that dependents may indeed have the ability to overexploit

salvagers if they have sufficiently large growth rate advantages.

To explore the consequences of growth rate differences on long-term population dynamics

and stability more rigorously, we exposed all four mixed salvager-dependent populations to a

sequence of proliferation-dilution cycles (Fig. 5C). Mixed populations were initialized with either

salvagers or dependents 100-fold in the majority, with a fixed total initial population size. The

culture medium including glycerol, Cbi, and DMB was replenished at each transfer. Interestingly,

salvagers were able to persist, and even achieved clear numerical dominance, in all four mixed

populations (Fig. 5C). These findings demonstrated that salvagers were strongly protected from

overexploitation even if dependents were able to proliferate faster by as much as 45%. Details of

long-term population dynamics varied between the four mixed populations. Differences in

maximally achievable growth rate was a dominant factor determining the rate at which population

ratios changed over time. Faster-proliferating salvagers outcompeted slower-proliferating

dependents at a higher rate (Fig. 5C, Sal^{GB-1}/Dep) compared to slower-proliferating salvagers

(Fig. 5C, Sal/Dep and Sal/Dep^{GB-1}). Similarly, faster-proliferating dependents fared better than

their slower-proliferating counterparts, as they were outcompeted at a lower rate (Fig. 5C).

Interestingly, the population ratio (from transfer 1 onwards) of Sal/Dep changed at a slightly higher

rate than that of Sal^{GB-1}/Dep^{GB-1}, despite the observation that there was no difference in maximally

achievable growth rate within each of the strain pairs (Fig. 2B, Fig. 5A and C). This finding may

indicate that, counter to a commonly held assumption and our findings in single batch cultures

(Fig. 5B), long-term competition dynamics in this metabolically interconnected microbial system

were not solely determined by relative differences in growth rate but rather absolute growth rates

may play a role as well. Alternatively, the observed difference in long-term competition dynamics

could also be explained by phenotypic variations other than differences in exponential growth rate

in the strain backgrounds. For example, differences in cobamide release during stationary phase

could shift the long-term population dynamics to the benefit of the dependents. The faster

proliferating GB-1 mutant was originally obtained through controlled laboratory evolution

experiments in which it was constantly kept in exponential phase[42]. Detailed multi-omics

characterization of the GB-1 mutant indeed indicated that its proteome shows a clear reduction in

stationary phase and stress-induced proteins[43]. The latter may cause a reduction in survival

probability when populations, as in our experimental setting and likely in many natural settings,

spent a considerable time in stationary phase. A repeated occurrence of substantial cell death,

together with its associated release of intracellular material, may enable the dependents to obtain

access to otherwise sequestered metabolites due to the disintegration of cells during stationary

phase.

Together, these findings indicate how intracellular metabolic processes like cobamide

precursor salvaging, that innately enable the (partial) retention of metabolites inside the cell prior

to release, are fundamentally distinct from extracellular processes like degradation of some

polysaccharides[15]. In the latter case, metabolites immediately become 'public goods' that have to

be taken up from the environment, thus inherently limiting the ability to privatize resources due to

the finite affinities of import systems[44,45]. Hence, our findings demonstrate the ability of intracellular

metabolic processes to establish stable metabolic interactions, with important ramifications on

community stability and the protection of productive population members.

Conclusion

Cobamide precursor salvaging has been experimentally observed and is predicted to be

widespread among bacteria[29,46]. Here, we showed that (1) salvaging of cobamide precursors does

not incur a noticeable metabolic burden, (2) salvaged cobamides are readily released and

provisioned to others, (3) cobamide precursor salvaging provides benefits through the generation

of complete functional cofactors as well as through the removal of non-functional, proliferation-

inhibiting precursors, and (4) organisms engaging in cobamide precursor salvaging are

intrinsically protected from overexploitation due to partial privatization. While likely not all of these

features are transferable to salvaging of other types of metabolites, some certainly will. For

example, removal of proliferation-inhibiting metabolites is likely widespread as high levels of many

compounds, such as cysteine[40], can lead to disturbances of optimal metabolic fluxes and thus

hinder proliferation. Furthermore, (partial) privatization, which has also recently been proposed in

the context of other microbial activities[7,47,48], is likely applicable to a broad set of metabolites,

including amino acids, nucleotides, and cofactors, that are inherently processed inside the cell.

Thus, organisms generating these intracellular metabolites, either via salvaging or *de novo*

biosynthesis, are likely naturally protected from overexploitation by non-productive population

members, and therefore may facilitate the emergence and stable maintenance of metabolite

provisioning interactions.

Salvaging, the concept of 'recycling and reusing', provides a resource-efficient way to use

environmental metabolites and inorganic substances that were released due to metabolite

overflow or cell lysis. It is important to note that metabolite salvaging is not limited to the uptake

and completion of precursors, which was the focus of this study. Salvaging can also be combined

with partial catabolism and metabolite modification to interconvert complete, yet enzymatically

unfavorable, metabolites into more useful forms[49,50]. Taken together, salvagers of various

metabolites likely play an important, yet underappreciated, role in many natural bacterial

populations by acting as hubs facilitating efficient community functioning by rerouting and

interconverting environmental metabolic fluxes. Innate protection from overexploitation by non-

productive population members positions salvaging of intracellular metabolites as an effective

mechanism to act in a broad range of environmental conditions, even those in which positive

assortment is not accessible (e.g., free-living populations).

Figures

Figure 1: Metabolic archetypes in bacterial populations

A) Illustration of distinct metabolic archetypes in bacterial populations. Producers create complex

metabolites from simple precursors such as carbon- and nitrogen-containing compounds.

Synthesized metabolites may be released and may benefit others. Dependent consumers take

up simple compounds and complex metabolites to fulfill their metabolic needs. Dependence on

complex metabolites may be facultative or obligate. Degraders break down complex compounds

such as polysaccharides or polypeptides. Degradation products are often accessible to others.

Salvagers take up and modify compounds from the environment. Salvaged compounds may be

released to the benefit of others. Independents neither create nor utilize a given metabolite. The

metabolic archetype of an organism varies for different metabolites, e.g., a producer of one

metabolite may be a consumer, degrader, salvager, or even be independent of another

metabolite. **B.** Distribution of predicted cobamide archetypes among sequenced bacteria[29]. Colors

as defined in A. Phylogenetic information is given in Supplementary Table 1.

Figure 2: Salvaged cobamides are readily provisioned

A) Illustration of cobamide archetypes. B) Exponential growth rates of salvagers (Sal, blue) and dependents (Dep, gray) in glycerol minimal medium with various additions (10 nM each). Strains were cultured alone (mono) or together (co, initial ratio: 1:1). Global means are shown as open circles and biological replicates as dots. C) Proliferation of jointly cultured salvagers (blue) and dependents (gray). Initial ratio: 1:1. Abbreviations: incomplete, non-functional cobamide precursor cobinamide (Cbi); lower ligand, a required component of a cobamide (LL); complete, functional cobamide (Cob); vitamin B_{12} (B12); 5,6-dimethylbenzimidazole, the lower ligand of B_{12} (DMB); bacterial units (b.u., see Methods for details).

Figure 3: Salvaging 'preference' shapes population dynamics

A-B) Growth rates of Sal (blue) and Dep (gray) when jointly cultured at an initial ratio of 1:1 in

glycerol minimal medium supplemented with the cobamide precursor Cbi [10 nM] and varying

concentrations of external lower ligand, DMB (A) and 2MA (B). **C)** Growth rate of Dep in glycerol

minimal medium with varying concentrations of the complete cobamides B_{12} (green, EC_{50} =

1.5×10^{-11} M) and FA (yellow, EC_{50} = 3.1×10^{-11} M). EC_{50} is the cobamide concentration at half-

maximal growth rate. In panels A-C, global means are shown as open circles, biological replicates

as dots, and sigmoidal fits as lines. **D)** HPLC analysis of cell extracts of Sal cultured in glycerol

minimal medium supplemented with Cbi [10 nM] in the absence of an externally supplied lower

ligand (black). Standards of cobinamide (Cbi, blue), factor A (FA, yellow), pseudo-B_{12} (pB12, red),

and vitamin B_{12} (B12, green) are shown. Abbreviations: a.u., arbitrary units.

Figure 4: Salvaging can relieve proliferation inhibition

A-C) Growth rate (A) and lag time (B) of Dep and growth rate of Sal (C) in glycerol minimal medium

as a function of the concentration of B_{12} in the absence (green) and presence (black) of the

cobamide precursor Cbi [10^{-7} M]. Global means are shown as open circles, biological replicates

as dots, and sigmoidal fits as lines. **D)** Same conditions as C, but with the lower ligand DMB [10^{-6} M] also added (black).

Figure 5: Partial cobamide privatization protects salvagers from overexploitation

A) Growth rates of Sal[GB-1] (blue) and Dep[GB-1] (gray) when cultured alone in glycerol minimal

medium with various additions (10 nM each). **B)** Growth yields after 24 hours of jointly cultured

salvagers (Sal or Sal[GB-1], blue) and dependents (Dep or Dep[GB-1], gray) in glycerol minimal medium

supplemented with Cbi [10 nM] and DMB [10 nM]. Initial ratio: 1:1. In panels A and B, global

means are shown as open circles and biological replicates as dots. **C)** Population ratio (green

dots) as function of transfer for jointly cultured salvagers (Sal or Sal[GB-1]) and dependents (Dep or

Dep[GB-1]) undergoing daily 1:100 [v:v] proliferation-dilution cycles in glycerol minimal medium

supplemented with Cbi [10 nM] and DMB [10 nM]. Solid green lines connect individual cultures

through subsequent transfers. Gray areas mark transfer-specific estimates of population ratio

detection limits (for details see Methods).

Supplementary Figure 1: Calibrated fluorescence as proxy for abundance in mixed populations

Examples of raw YFP and CFP fluorescence as function of raw optical density at 600 nm (OD_{600})

extracted from proliferating, fluorescently labeled salvager populations (Sal) cultured in glycerol

minimal medium supplemented with Cbi [10 nM] and DMB [10 nM]. Non-specific fluorescence

background and crosstalk between fluorescence channels were small (bottom left and top right).

YFP fluorescence underestimated abundances for raw OD_{600} >0.8. Day- and medium-specific

fluorescence-to-abundance conversion factors were extracted as slopes from linear fits in the

range $0 \leq$ raw $OD_{600} \leq 0.6$ and averaged over all biological replicates. The equivalent of a blank-

corrected OD_{600} of 1 was defined as 1 b.u. (arbitrary bacterial unit). Abbreviations: arbitrary units

(a.u.).

Supplementary Figure 2: Salvagers do not exhibit lag

A) Lag time of Sal in glycerol minimal medium as function of B_{12} concentration in presence (black)

and absence (green, hidden behind black symbols) of Cbi [10^{-7} M]. **B)** Lag time of Sal in glycerol

minimal medium as function of B_{12} concentration in presence (black) and absence (green, hidden

behind black symbols) of Cbi [10^{-7} M] and DMB [10^{-6} M]. In panels A and B, global means are

shown as open circles and biological replicates as dots.

Supplementary Figure 3: Cobamide precursor salvaging is not differentially regulated at the

level of transcription

Relative fluorescence (fluorescence in arbitrary units per bacterial abundance) of Sal carrying a

YFP transcriptional reporter for cobUST (i.e., major cobamide salvaging genes) activity as

function of time in glycerol minimal medium supplemented with Cbi [10 nM] and DMB [10 nM]

(blue) or B_{12} [10 nM] (red). Dashed vertical line marks approximate time point at which proliferation

ceases. Abbreviations: arbitrary units (a.u.) and bacterial units (b.u., see Methods for details).

Supplementary Figure 4: UV-Vis spectra of HPLC peaks

Absorbance as a function of wavelength for selected HPLC peaks labeled in Fig. 3D. **A-B)** Peak spectra of Sal extracts. **C-G)** Peak spectra of known standards. Vertical dashed lines mark 355 nm and 550 nm, respectively.

Supplementary Figure 5: Classification of flow cytometry events

A-E) AmCyan vs FITC scatter plots color-coded by event counts. AmCyan and FITC axes scales

are logicle. Dashed lines mark gating conditions for YFP-negative-CFP-positive (cyan) and YFP-

positive-CFP-negative (yellow) events, respectively. YFP-negative-CFP-negative events are

located in the lower left quadrant of each plot, i.e., below the horizontal, dashed cyan lines and

left of the vertical, dashed yellow lines. YFP-positive-CFP-positive events are located in the upper

right quadrant of each plot, i.e., all events that do not fall into any of the other three gates. Notably,

YFP-positive-CFP-positive events are very rare. Hence, no correction was applied, and YFP-

positive-CFP-positive events were simply ignored for population ratio estimations. **A/C)** Mixed

cultures of Sal/Dep at transfer 0 and transfer 1, respectively. **B/D)** Mixed cultures of Sal^{GB-1}/Dep^{GB-1}

at transfer 0 and transfer 1, respectively. A notable increase in YFP-negative-CFP-negative

events can be seen at transfer 1 only if GB-1 cells are present in the mixed culture. **E)** Blank at

transfer 1. **F)** Number of YFP-negative-CFP-negative (YNCN) events as function of the total

number of GB-1 events detected in a culture. The total number of GB-1 events was estimated

from YFP-negative-CFP-positive and YFP-positive-CFP-negative event counts and the known

strain background identity (WT or GB-1). Solid lines mark YFP-negative-CFP-negative event

counts of blanks. Both axes are linear below 10 and logarithmic thereafter. Cultures with a higher

number of GB-1 events show a tendency to contain more YFP-negative-CFP-negative events

(except for transfer 0, dark blue), suggesting that the occurrence of YFP-negative-CFP-negative

events is mainly caused by strain background identity (WT or GB-1) rather than cobamide

salvaging phenotype (salvager or dependent). No correction was applied, since adding the YFP-

negative-CFP-negative events to the counts of GB-1 events only marginally altered the population

ratio estimates.

Phylum	Producers	Salvagers	Dependents	Independents
Nitrospinae	0	2	8	0
Candidatus Azambacteria	0	0	10	0
Candidatus Curtissbacteria	0	0	9	1
Candidatus Saccharibacteria	0	0	2	8
Chlorobi	0	10	0	0
Candidatus Sungbacteria	0	0	6	5
Ignavibacteriae	0	0	12	0
Candidatus Magasanikbacteria	0	0	8	4
Candidatus Staskawiczbacteria	0	0	4	9
Candidatus Taylorbacteria	0	0	6	8
Candidatus Gottesmanbacteria	0	0	13	2
Candidatus Omnitrophica	0	2	14	0
Candidatus Doudnabacteria	0	0	15	1
Candidatus Zambryskibacteria	0	0	17	0
Elusimicrobia	1	3	12	1
Candidatus Kaiserbacteria	0	0	2	15
Candidatus Uhrbacteria	0	0	16	2
Fibrobacteres	0	1	18	0
Candidatus Peregrinibacteria	0	0	18	1
Candidatus Daviesbacteria	0	0	16	4
Synergistetes	6	9	5	0
Aquificae	0	0	13	8
Candidatus Levybacteria	0	0	17	5
Nitrospirae	8	7	6	1

Candidatus Woesebacteria	0	0	19	4
Candidatus Giovannonibacteria	0	0	19	8
Candidatus Roizmanbacteria	0	0	21	7
Verrucomicrobia	0	2	26	1
Candidatus Yanofskybacteria	0	0	34	0
Thermotogae	2	9	24	0
Planctomycetes	8	1	28	0
Acidobacteria	4	7	28	0
Chlamydiae	0	0	16	26
Chloroflexi	11	16	19	0
Fusobacteria	44	4	0	2
Deinococcus-Thermus	15	23	14	0
Spirochaetes	17	11	40	30
Tenericutes	0	1	26	115
Cyanobacteria	203	2	20	2
Bacteroidetes	6	324	689	21
Actinobacteria	1022	38	552	195
Firmicutes	646	473	744	275
Proteobacteria	2180	974	1423	258

459 **Supplementary Table 1:** Cobamide archetypes

460 Counts of cobamide archetypes per phylum containing at least 10 genomes[29]. Same data (and

461 same ordering, top-to-bottom in table corresponds to left-to-right in figure) as shown in Fig. 1B.

name	genotype	source
Sal	*Escherichia coli* MG1655 ΔmetE	Kirmiz et al[51]
Dep	*Escherichia coli* MG1655 ΔmetE ΔcobUST ΔcobC	Mok et al[50]
Sal[GB-1]	*Escherichia coli* MG1655 GB-1 ΔmetE	This study.
Dep[GB-1]	*Escherichia coli* MG1655 GB-1 ΔmetE ΔcobUST ΔcobC	This study.

463 **Supplementary Table 2:** Bacterial strains

name	backbone	insert	source
pSG013	Kan^R marker, pBR322 origin, and rop gene	constitutive promoter (BBa_J23100, iGEM), RBS (BBa_B0034, iGEM), and mCerulean	This study.
pSG015	Kan^R marker, pBR322 origin, and rop gene	constitutive promoter (BBa_J23100, iGEM), RBS (BBa_B0034, iGEM), and mCitrine	This study.
pSG033	Kan^R marker, pBR322 origin, and rop gene	cobUST regulatory region (genomic region 500 bp upstream of cobUST start codon in *E. coli* MG1655 wt) and mCitrine	This study.

465 **Supplementary Table 3:** Plasmids

Methods

<u>Strains and plasmids</u>

All strains are derivatives of *E. coli* strain MG1655. Sal[GB-1] and Dep[GB-1] were derived from *Escherichia coli* MG1655 mutant GB-1[42]. Strain Sal[GB-1] was constructed by introducing into strain GB-1 the *metE*::Kan[R] allele from donor strain JW3805-1[52] via P1 transduction and subsequently removing the Kan[R] marker by transient introduction of the plasmid pCP20 encoding the FLP recombinase[53]. To generate strain Dep[GB-1], the *cobUST* operon was deleted via lambda red recombineering[53] and the gene *cobC* was replaced via P1 transduction as described above. After each step, the selective marker was removed. Gene deletions were verified via PCR and phenotyping of growth in glycerol minimal medium supplemented with either methionine, vitamin B_{12}, cobinamide and DMB, or no addition.

Fluorescent marker and reporter plasmids are based on pETMini[50] and were assembled via isothermal cloning[54]. Insert sequences were verified via Sanger sequencing.

All bacterial strains and plasmids are listed in Supplementary Table 2 and 3.

<u>Media and culturing</u>

For proliferation and serial passaging assays, 2 mL glycerol minimal medium (50 mM KPO_4, 67 mM NaCl, 7.6 mM $(NH_4)_2SO_4$, 500 µM $MgSO_4$, 1.25 µM $Fe_2(SO_4)_3$, 0.2% [v:v] glycerol, pH 7.4) supplemented with 1 g/L methionine was inoculated with single colonies from LB plates (10 g/L tryptone, 5 g/L yeast extract, 5 g/L NaCl, 15 g/L Bacto agar), and incubated at 37 °C with aeration (200 rpm) until saturation was reached after approx. 24 hours. Cultures were transferred into 2 mL fresh glycerol minimal medium supplemented with methionine as 1:100 [v:v] dilutions, incubated at 37 °C with aeration, and harvested in mid-exponential phase (approx. OD_{600} 0.3-0.6). Cells were washed three times by centrifugation at 14000 rpm followed by resuspension in unsupplemented minimal medium. Optical density (OD_{600}) was adjusted to 10^{-1} and, if required,

cultures were mixed 1:1 [v:v], or as indicated, to obtain mixed populations. If necessary, media

were supplemented with 25 mg/L kanamycin to retain plasmids.

For proliferation assays, precultured and OD-adjusted cultures were transferred as 1:10

[v:v] dilutions into glass-bottom 96-well culture plates (CellVis) containing a final volume of 200

µL fresh glycerol minimal medium supplemented with either 10 nM vitamin B_{12}, 10 nM cobinamide

and 10 nM 5,6-dimethylbenzimidazole (DMB), or no additions, if not indicated otherwise. 96-well

culture plates were sealed with an evaporation seal (Breathe-Easy® sealing membrane) and

incubated at 37 °C with continuous shaking in a multi-well plate reader (Tecan Spark) for 36 or

60 hours. Absorbance at 600 nm and fluorescence (CFP: excitation 455/5 nm, emission 475/5

nm; YFP: excitation 514/5 nm, emission 550/30 nm) were recorded every 10 minutes.

Fluorescence measurements were converted into units equivalent to OD_{600} (see below and

Supplementary Figure 1). Fluorescence was recorded only for experiments containing two strains.

All proliferation experiments were performed in biological triplicates started on the same day.

Proliferation phenotypes showed little day-to-day variation.

For serial passaging assays, precultured and OD-adjusted mixed cultures were

transferred as 1:10 [v:v] dilutions into clear plastic 96-well culture plates (Corning®) containing a

final volume of 200 µL fresh glycerol minimal medium supplemented with 10 nM cobinamide and

10 nM DMB. Wells without cultures were filled with 200 µL dH2O. 96-well culture plates were

sealed with an evaporation seal (AeraSeal^{TM}, EXCEL Scientific) and incubated at 37 °C while

shaking at 1200 rpm in a heated plate shaker (Southwest Science). Cultures were transferred

every 24 hours as 1:100 [v:v] dilutions into fresh glycerol minimal medium supplemented with 10

nM cobinamide and 10 nM DMB. At each transfer samples were taken for composition analysis

at the Flow Cytometry Facility in the Cancer Research Laboratory at UC Berkeley. Prior to flow

cytometry analysis (BD LSRFortessa™ Cell Analyzer), samples were diluted 1:1000 [v:v] into

unsupplemented minimal medium. Forward scatter (FSC; V=350, log), side scatter (SSC; V=280,

log), FITC (V=650, log), and AmCyan (V=750, log) were recorded. Event detection was triggered

by a threshold in SSC of 500. Samples were run at approx. 300-600 events per second, 10^5

events were collected per sample, and lines were flushed with 10% bleach, rinse solution, and

dH2O between sample runs. The serial transfer experiment was performed in biological triplicate

started on the same day.

For corrinoid extractions, 2 mL glycerol minimal medium supplemented with 100 mg/L

methionine was inoculated with single colonies of strains grown on LB plates and incubated at 37

°C with aeration until saturation was reached. Cells were washed twice by centrifugation at 14000

rpm and resuspension in unsupplemented minimal medium, diluted 1:500 [v:v] into 500 mL

glycerol minimal medium supplemented with 10 nM cobinamide, incubated at 37 °C with aeration

for 24 hours, and harvested for metabolite extraction, purification, and profiling (see below).

<u>Evaluation of proliferation dynamics</u>

For proliferation assays, population dynamics were evaluated in Python[55] (version 3.8)

with the following steps. (1) Fluorescence-to-absorbance conversion (if applicable):

Fluorescence-to-absorbance conversion factors were estimated in a day-strain-medium-

dependent manner for each fluorescent channel. Conversion factors were obtained as replicate-

averaged slopes of linear fits (*NumPy*[56] (version 1.21.2) function *polyfit*) of raw fluorescence vs

raw absorbance (A_{600}) in the range $A_{600} > 0$ and $A_{600} < 0.3$ of cultures containing only a single

marker plasmid. Raw fluorescence reads were then converted into equivalent units of raw

absorbance by multiplication of the conversion factors with the raw fluorescence reads. (2)

Absorbance-to-abundance conversion: All reads were pathlength-corrected to obtain abundances

in arbitrary bacterial units (b.u.; 1 b.u. is equivalent to an OD_{600} of 1). (3) Blank subtraction: For

reads based on absorbance measurements, blanks were estimated in a well-specific manner as

the difference of the mean of the first two time points and the known initial abundance. For reads

based on fluorescence measurements, blanks were estimated in a global manner as means over

all wells containing cultures that did not carry the specific fluorescent marker plasmid. (4)

Smoothing and shortening: Blank-corrected abundance timeseries were smoothed with a

centered rolling mean filter (*pandas* (version 1.3.2) *DataFrame* function *rolling*) using a window

width of 7 data points with all data points equally weighted and a minimal window width of 1 point.

Thereafter, smoothed abundance timeseries were shortened to 24 or 60 hours. (5) Extraction of

proliferation parameters: Growth rate estimates were obtained via a Theil-Sen estimator (*SciPy*[57]

(version 1.7.1) module *stats.mstats* function *theilslopes*) which was applied to log-transformed

abundances. Estimation was restricted to data points for which the abundance was larger than

1.5×10^{-2} b.u. and smaller than 6×10^{-2} b.u. (Fig. 3C and Fig. 4ACD) or the abundance was larger

than 1.5×10^{-2} b.u. and latest data point for which the rolling-window growth rate estimate (see

below) was larger than 0.1 h^{-1} (Fig. 2B, Fig. 3AB, and Fig. 5A). Growth rate was set to zero if

fewer than 7 data points fell into the estimation interval. Growth rate estimates were generally

found to be reproducible within +/- ~0.025 h^{-1} (day-to-day and channel-to-channel variability).

Rolling-window growth rate estimates were obtained by applying a Theil-Sen estimator (*SciPy*[57]

(version 1.7.1) module *stats.mstats* function *theilslopes*) to log-transformed abundances in a

centered rolling window fashion (*pandas* (version 1.3.2) *DataFrame* function *rolling*) using a

window width of 19 data points with all data points equally weighted and a minimal window width

of 10 data points. Lag time estimates were obtained by linear interpolation of growth rates and

intercepts obtained from the Theil-Sen estimator to known log-transformed initial abundances.

Lag time was set to negative infinity for non-growing populations. (6) Extraction of dose response

curve parameters (if applicable): Dose response curve parameters were obtained via non-linear

fitting (*SciPy*[57] (version 1.7.1) module *optimize* function *curve_fit*) of extracted proliferation

parameters to a four-parameter dose response equation:

566
$$y = y_{bottom} + x^n \times \frac{y_{top} - y_{bottom}}{x^n + EC50^n}$$

with variables x (effector concentration) and y (proliferation parameter), and parameters y_{bottom}

(proliferation parameter value for $x \to 0$), y_{top} (proliferation parameter value for $x \to \infty$), n (Hill

coefficient), and $EC50$ (half maximal effective concentration).

<u>Evaluation of proliferation-dilution cycles</u>

Flow cytometry data was analyzed in custom Python[55] (version 3.8) scripts using the

FlowCal[58] (version 1.3.0) package. Briefly, data were loaded and transformed into arbitrary units

via FlowCal. Gating was directly performed in Python. Events were classified as YFP-positive-

CFP-negative if FITC > $1x10^3$ a.u. and AmCyan < $4x10^3$ a.u., as YFP-negative-CFP-positive if

FITC < $7x10^2$ a.u. and AmCyan > $2x10^3$ a.u., as YFP-negative-CFP-negative if FITC < $1x10^3$ a.u.

and AmCyan < $2x10^3$ a.u., and as YFP-positive-CFP-positive otherwise. Some gating conditions

were altered for transfer zero as cells pre-cultured in the presence of methionine and harvested

in mid-exponential phase showed slightly different fluorescence properties than cells cultured in

the presence of cobinamide and DMB and harvested in stationary phase. In particular, YFP-

negative-CFP-positive if FITC < $7x10^2$ a.u. and AmCyan > $5x10^3$ a.u. and YFP-negative-CFP-

negative if FITC < $1x10^3$ a.u. and AmCyan < $5x10^3$ a.u. All other gating was as described above.

Population ratios (Sal : Dep) were determined as the ratio of the number of YFP-negative-CFP-

positive events over the number of YFP-positive-CFP-negative events. The dynamic range of the

population ratio estimation was determined in a transfer-specific manner by the ratio of the

number of YFP-negative-CFP-positive events obtained when measuring a blank (or 1, if no such

events where detected) over the total number of events collected per mixed cultures (i.e., 10^5)

and the ratio of the total number of events collected per mixed cultures (i.e., 10^5) over the number

of YFP-positive-CFP-negative events obtained when measuring a blank (or 1, if no such events

where detected), respectively. For a discussion of YFP-negative-CFP-negative events and YFP-

positive-CFP-positive events see Supplementary Figure 5.

Corrinoid extraction and analysis

Corrinoid extraction, cyanation, and purification were performed as previously described[46,59]. Briefly, 500 mL of cultures were harvested by centrifugation and corrinoids were extracted from the pellet with methanol and cyanated with 20 mg potassium cyanide per gram (wet weight) of cells. Following cyanation, samples were desalted using C18 SepPak (Waters Associates, Milford, MA). HPLC purification and profiling were performed as previously described[46].

Contributions

S.G. and M.E.T. conceived research. S.G., G.J.P., and M.E.T. developed *E. coli*-cobamide model system. S.G. designed and performed all experiments except corrinoid extraction and analysis. G.J.P. designed and performed corrinoid extraction and analysis. S.G. analyzed all data. S.G. and M.E.T. interpreted the data and wrote the manuscript with input from G.J.P.

Data and code availability

All data and code are available from the corresponding author upon reasonable request.

Acknowledgement

This work was funded by National Institutes of Health grants R01GM114535 and R35GM139633 to M.E.T. We thank O. Sokolovskaya for initially suggesting to employ *E. coli* mutant strains. We thank K. Mok and O. Sokolovskaya for assistance with HPLC measurements and H. Nolla at the Flow Cytometry Facility in the Cancer Research Laboratory at UC Berkeley for assistance with flow cytometry measurements. We thank members of the Taga lab for helpful discussions. We also thank K. Kennedy, K. Mok, and Z. Hallberg for critical reading of the manuscript. *E. coli* strain MG1655 GB-1 was a kind gift of B.O. Palsson.

References

1   D'Souza, G. *et al.* Ecology and evolution of metabolic cross-feeding interactions in bacteria. *Natural Product Reports* **35**, 455-488 (2018).

2   Gude, S., Pherribo, G. J. & Taga, M. E. Emergence of Metabolite Provisioning as a By-Product of Evolved Biological Functions. *mSystems* **5**, e00259-00220, doi:10.1128/mSystems.00259-20 (2020).

3   Campbell, K., Herrera-Dominguez, L., Correia-Melo, C., Zelezniak, A. & Ralser, M. Biochemical principles enabling metabolic cooperativity and phenotypic heterogeneity at the single cell level. *Current Opinion in Systems Biology* **8**, 97-108, doi:https://doi.org/10.1016/j.coisb.2017.12.001 (2018).

4   Fletcher, J. A. & Doebeli, M. A simple and general explanation for the evolution of altruism. *Proceedings of the Royal Society B: Biological Sciences* **276**, 13-19, doi:doi:10.1098/rspb.2008.0829 (2009).

5   Travisano, M. & Velicer, G. J. Strategies of microbial cheater control. *Trends in Microbiology* **12**, 72-78, doi:https://doi.org/10.1016/j.tim.2003.12.009 (2004).

6   Pande, S. *et al.* Privatization of cooperative benefits stabilizes mutualistic cross-feeding interactions in spatially structured environments. *The ISME Journal* **10**, 1413-1423, doi:10.1038/ismej.2015.212 (2016).

7   Smith, R. P. *et al.* The public and private benefit of an impure public good determines the sensitivity of bacteria to population collapse in a snowdrift game. *Environmental Microbiology* **21**, 4330-4342, doi:https://doi.org/10.1111/1462-2920.14796 (2019).

8   Rankin, D. J., Bargum, K. & Kokko, H. The tragedy of the commons in evolutionary biology.

Trends in Ecology & Evolution **22**, 643-651, doi:https://doi.org/10.1016/j.tree.2007.07.009

(2007).

9   D'Souza, G. *et al.* LESS IS MORE: SELECTIVE ADVANTAGES CAN EXPLAIN THE PREVALENT LOSS OF

BIOSYNTHETIC GENES IN BACTERIA. *Evolution* **68**, 2559-2570, doi:10.1111/evo.12468 (2014).

10   Freilich, S. *et al.* Competitive and cooperative metabolic interactions in bacterial communities.

Nature Communications **2**, 589, doi:10.1038/ncomms1597 (2011).

11   Harcombe, W. NOVEL COOPERATION EXPERIMENTALLY EVOLVED BETWEEN SPECIES. *Evolution*

64, 2166-2172, doi:10.1111/j.1558-5646.2010.00959.x (2010).

12   Mee, M. T., Collins, J. J., Church, G. M. & Wang, H. H. Syntrophic exchange in synthetic microbial

communities. *Proceedings of the National Academy of Sciences* **111**, E2149-E2156,

doi:10.1073/pnas.1405641111 (2014).

13   Pacheco, A. R., Moel, M. & Segrè, D. Costless metabolic secretions as drivers of interspecies

interactions in microbial ecosystems. *Nature Communications* **10**, 103, doi:10.1038/s41467-018-

07946-9 (2019).

14   Morris, J. J., Lenski, R. E. & Zinser, E. R. The Black Queen Hypothesis: Evolution of Dependencies

through Adaptive Gene Loss. *mBio* **3**, e00036-00012, doi:10.1128/mBio.00036-12 (2012).

15   Gore, J., Youk, H. & van Oudenaarden, A. Snowdrift game dynamics and facultative cheating in

yeast. *Nature* **459**, 253-256, doi:10.1038/nature07921 (2009).

16   Gur, E., Biran, D. & Ron, E. Z. Regulated proteolysis in Gram-negative bacteria — how and when?

Nature Reviews Microbiology **9**, 839-848, doi:10.1038/nrmicro2669 (2011).

17   Villalba, G., Segarra, M., Fernández, A. I., Chimenos, J. M. & Espiell, F. A proposal for quantifying

the recyclability of materials. *Resources, Conservation and Recycling* **37**, 39-53,

doi:https://doi.org/10.1016/S0921-3449(02)00056-3 (2002).

18   Neilands, J. B. Siderophores: Structure and Function of Microbial Iron Transport Compounds (*).

Journal of Biological Chemistry **270**, 26723-26726, doi:10.1074/jbc.270.45.26723 (1995).

19   Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. Siderophore cheating and cheating

resistance shape competition for iron in soil and freshwater Pseudomonas communities. *Nature*

Communications **8**, 414, doi:10.1038/s41467-017-00509-4 (2017).

20   Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host

interactions. *Nature Reviews Microbiology* **18**, 152-163, doi:10.1038/s41579-019-0284-4 (2020).

21   Johnson, M. D. L., Echlin, H., Dao, T. H. & Rosch, J. W. Characterization of NAD salvage pathways

and their role in virulence in Streptococcus pneumoniae. *Microbiology* **161**, 2127-2136,

doi:10.1099/mic.0.000164 (2015).

22   Melnick, J. *et al.* Identification of the Two Missing Bacterial Genes Involved in Thiamine Salvage:

Thiamine Pyrophosphokinase and Thiamine Kinase. *J Bacteriol* **186**, 3660-3662,

doi:10.1128/jb.186.11.3660-3662.2004 (2004).

23   Nygaard, P. in *Bacillus subtilis and Other Gram-Positive Bacteria*   (American Society of

Microbiology, 1993).

24   Sekowska, A. *et al.* Bacterial variations on the methionine salvage pathway. *BMC Microbiology* **4**,

9, doi:10.1186/1471-2180-4-9 (2004).

25   Yuan, Y. *et al.* Discovery of novel bacterial queuine salvage enzymes and pathways in human

pathogens. *Proceedings of the National Academy of Sciences* **116**, 19126-19135,

doi:10.1073/pnas.1909604116 (2019).

26   Datta, M. S., Sliwerska, E., Gore, J., Polz, M. F. & Cordero, O. X. Microbial interactions lead to

rapid micro-scale successions on model marine particles. *Nature Communications* **7**, 11965,

doi:10.1038/ncomms11965 (2016).

27 Escalante-Semerena, J. C. Conversion of Cobinamide into Adenosylcobamide in Bacteria and

Archaea. *J Bacteriol* **189**, 4555-4560, doi:10.1128/jb.00503-07 (2007).

28 Contreras, H., Chim, N., Credali, A. & Goulding, C. W. Heme uptake in bacterial pathogens.

Current Opinion in Chemical Biology **19**, 34-41, doi:https://doi.org/10.1016/j.cbpa.2013.12.014

(2014).

29 Shelton, A. N. *et al.* Uneven distribution of cobamide biosynthesis and dependence in bacteria

predicted by comparative genomics. *The ISME Journal* **13**, 789-804, doi:10.1038/s41396-018-

0304-9 (2019).

30 Pande, S. *et al.* Fitness and stability of obligate cross-feeding interactions that emerge upon

gene loss in bacteria. *The ISME Journal* **8**, 953-962, doi:10.1038/ismej.2013.211 (2014).

31 Noto Guillen, M., Rosener, B., Sayin, S. & Mitchell, A. Assembling stable syntrophic Escherichia

coli communities by comprehensively identifying beneficiaries of secreted goods. *Cell Systems*,

doi:https://doi.org/10.1016/j.cels.2021.08.002 (2021).

32 Anderson, P. J. *et al.* One Pathway Can Incorporate either Adenine or Dimethylbenzimidazole as

an α-Axial Ligand of B12 Cofactors in Salmonella enterica. *J Bacteriol* **190**, 1160-1171,

doi:doi:10.1128/JB.01386-07 (2008).

33 Chan, C. H. & Escalante-Semerena, J. C. ArsAB, a novel enzyme from Sporomusa ovata activates

phenolic bases for adenosylcobamide biosynthesis. *Molecular Microbiology* **81**, 952-967,

doi:https://doi.org/10.1111/j.1365-2958.2011.07741.x (2011).

34 Degnan, Patrick H., Barry, Natasha A., Mok, Kenny C., Taga, Michiko E. & Goodman, Andrew L.

Human Gut Microbes Use Multiple Transporters to Distinguish Vitamin B12 Analogs and

Compete in the Gut. *Cell Host & Microbe* **15**, 47-57,

doi:https://doi.org/10.1016/j.chom.2013.12.007 (2014).

35　Crofts, Terence S., Seth, Erica C., Hazra, Amrita B. & Taga, Michiko E. Cobamide Structure

Depends on Both Lower Ligand Availability and CobT Substrate Specificity. *Chemistry & Biology*

20, 1265-1274, doi:https://doi.org/10.1016/j.chembiol.2013.08.006 (2013).

36　Hazra, Amrita B., Tran, Jennifer L. A., Crofts, Terence S. & Taga, Michiko E. Analysis of Substrate

Specificity in CobT Homologs Reveals Widespread Preference for DMB, the Lower Axial Ligand of

Vitamin B12. *Chemistry & Biology* **20**, 1275-1285,

doi:https://doi.org/10.1016/j.chembiol.2013.08.007 (2013).

37　Hazra, A. B. *et al.* Anaerobic biosynthesis of the lower ligand of vitamin B$_{12}$.

Proceedings of the National Academy of Sciences **112**, 10792-10797,

doi:10.1073/pnas.1509132112 (2015).

38　Davies, J. Inactivation of Antibiotics and the Dissemination of Resistance Genes. *Science* **264**,

375-382, doi:doi:10.1126/science.8153624 (1994).

39　Wolfe, A. J. The Acetate Switch. *Microbiology and Molecular Biology Reviews* **69**, 12-50,

doi:10.1128/mmbr.69.1.12-50.2005 (2005).

40　Jones, C. M., Hernández Lozada, N. J. & Pfleger, B. F. Efflux systems in bacteria and their

metabolic engineering applications. *Applied Microbiology and Biotechnology* **99**, 9381-9393,

doi:10.1007/s00253-015-6963-9 (2015).

41　Smith, P. & Schuster, M. Public goods and cheating in microbes. *Current Biology* **29**, R442-R447,

doi:https://doi.org/10.1016/j.cub.2019.03.001 (2019).

42　Herring, C. D. *et al.* Comparative genome sequencing of Escherichia coli allows observation of

bacterial evolution on a laboratory timescale. *Nature Genetics* **38**, 1406-1412,

doi:10.1038/ng1906 (2006).

43   Cheng, K.-K. *et al.* Global metabolic network reorganization by adaptive mutations allows fast

growth of Escherichia coli on glycerol. *Nature Communications* **5**, 3233,

doi:10.1038/ncomms4233 (2014).

44   Burkovski, A. & Krämer, R. Bacterial amino acid transport proteins: occurrence, functions, and

significance for biotechnological applications. *Applied Microbiology and Biotechnology* **58**, 265-

274, doi:10.1007/s00253-001-0869-4 (2002).

45   Jojima, T., Omumasaba, C. A., Inui, M. & Yukawa, H. Sugar transporters in efficient utilization of

mixed sugar substrates: current knowledge and outlook. *Applied Microbiology and*

Biotechnology **85**, 471-480, doi:10.1007/s00253-009-2292-1 (2010).

46   Yi, S. *et al.* Versatility in Corrinoid Salvaging and Remodeling Pathways Supports Corrinoid-

Dependent Metabolism in Dehalococcoides mccartyi. *Applied and Environmental Microbiology*

78, 7745-7752, doi:doi:10.1128/AEM.02150-12 (2012).

47   Estrela, S., Morris, J. J. & Kerr, B. Private benefits and metabolic conflicts shape the emergence

of microbial interdependencies. *Environmental Microbiology* **18**, 1415-1427, doi:10.1111/1462-

2920.13028 (2016).

48   Lindsay, R. J., Pawlowska, B. J. & Gudelj, I. Privatization of public goods can cause population

decline. *Nature Ecology & Evolution* **3**, 1206-1216, doi:10.1038/s41559-019-0944-9 (2019).

49   Gray, M. J. & Escalante-Semerena, J. C. The cobinamide amidohydrolase (cobyric acid-forming)

CbiZ enzyme: a critical activity of the cobamide remodelling system of Rhodobacter sphaeroides.

Molecular Microbiology **74**, 1198-1210, doi:10.1111/j.1365-2958.2009.06928.x (2009).

50   Mok, K. C. *et al.* Identification of a Novel Cobamide Remodeling Enzyme in the Beneficial Human

Gut Bacterium Akkermansia muciniphila. *mBio* **11**, e02507-02520, doi:doi:10.1128/mBio.02507-

20 (2020).

51   Kirmiz, N. *et al.* Comparative Genomics Guides Elucidation of Vitamin B$_{12}$

Biosynthesis in Novel Human-Associated *Akkermansia* Strains. *Applied and Environmental*

Microbiology **86**, e02117-02119, doi:doi:10.1128/AEM.02117-19 (2020).

52   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the

Keio collection. *Molecular Systems Biology* **2**, 2006.0008,

doi:https://doi.org/10.1038/msb4100050 (2006).

53   Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia coli

K-12 using PCR products. *Proceedings of the National Academy of Sciences* **97**, 6640-6645,

doi:10.1073/pnas.120163297 (2000).

54   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases.

Nature Methods **6**, 343-345, doi:10.1038/nmeth.1318 (2009).

55   *Python Software Foundation, Python v.3.8*, <http://www.python.org> (2019).

56   Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357-362, doi:10.1038/s41586-

020-2649-2 (2020).

57   Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature*

Methods **17**, 261-272, doi:10.1038/s41592-019-0686-2 (2020).

58   Castillo-Hair, S. M. *et al.* FlowCal: A User-Friendly, Open Source Software Tool for Automatically

Converting Flow Cytometry Data from Arbitrary to Calibrated Units. *ACS Synthetic Biology* **5**,

774-780, doi:10.1021/acssynbio.5b00284 (2016).

59   Sokolovskaya, O. M. *et al.* Cofactor Selectivity in Methylmalonyl Coenzyme A Mutase, a Model

Cobamide-Dependent Enzyme. *mBio* **10**, e01303-01319, doi:doi:10.1128/mBio.01303-19 (2019).

777