Timescale of degradation-driven protein level fluctuation in the yeast *Saccharomyces cerevisiae*

Bahareh Mahrou[1, 2,]*, Azady Pirhanov[1], Moluk Hadi Alijanvand[3], Yong Ku Cho[1, 4,]*, Yong-Jun Shin[1]

[1] Biomedical Engineering Department, University of Connecticut, Storrs, CT 06269, USA

[2] Electrical Engineering Department, University of Connecticut, Storrs, CT 06069, USA

[3] Department of Epidemiology and Biostatistics, School of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

[4] Chemical and Biomolecular Engineering Department, University of Connecticut, Storrs, CT 06269, USA

* Corresponding authors: Bahareh Mahrou, E-mail: bahareh.mahrou@uconn.edu

Yong Ku Cho, E-mail: cho@uconn.edu

Abstract

Generating robust, predictable perturbations in cellular protein levels will advance our understanding of protein function and enable control of physiological outcomes in biotechnology applications. Previous studies have shown that controlling RNA transcription achieves perturbations in protein levels over a timescale of several hours. Here, we demonstrate the potential for harnessing the protein degradation machinery to achieve robust, rapid control of a specific protein level in the yeast *Saccharomyces cerevisiae*. Using a light-driven protein degradation machinery and red fluorescent proteins as reporters, we show that under constant transcriptional induction, repeated triangular fluctuations in protein levels can be generated by controlling the protein degradation rate. Consistent with previous results using transcriptional control, we observed a continuous decrease in the magnitude of fluctuations as the modulation frequency increased, indicating low-pass filtering of input perturbation. However, compared to hour-scale fluctuations observed using transcriptional control, modulating the protein degradation rate enabled five to ten minute-scale fluctuations. Our study demonstrates the potential for repeated control of protein levels by controlling protein degradation rate, at timescales much shorter than that achieved by transcriptional control.

Introduction

Extensive research led to insights on how the dynamics and complexity of intracellular regulatory networks are organized to maintain a cellular response (Huang and Ingber 2000, Goldberg 2007, Huang and Ingber 2007, Vogel and Marcotte 2012, Ciechanover 2017). One form of regulation is protein turnover, which refers to the continual renewal or replacement of intracellular proteins. The output of this process is the steady-state intracellular protein concentration, maintained dynamically by continuous production and degradation. The steady-state shifts in response to extracellular stimuli or pathophysiologic conditions, which cause changes in production or degradation rates (Baracos 2000, Pratt, Petty et al. 2002, Mizushima and Klionsky 2007, Hinkson and Elias 2011, Ciechanover 2017, Kneppers, Langen et al. 2017). A powerful approach for gaining fundamental insights for such complex systems would be a bottom-up approach, in which the protein turnover is controlled using external stimuli.

Although production and degradation both contribute to protein turnover, control strategies have largely focused on gene transcription. Multitude of factors involved in transcriptional regulation and its impact on biology and disease have been well characterized (Hill and Treisman 1995, Tautz 2000, Bintu, Buchler et al. 2005, Lee and Young 2013). Numerous previous studies have demonstrated control of transcription using optical and chemical approaches to modulate protein levels (Bennett, Pang et al. 2008, Milias-Argeitis, Summers et al. 2011, Toettcher, Gong et al. 2011, Uhlendorf, Miermont et al. 2012, Melendez, Patel et al. 2014, Milias-Argeitis, Rullan et al. 2016). On the other hand, the effect of degradation process on protein turnover control has been far less characterized. In nature, regulation of protein levels by modulating the degradation rate have been discovered in several proteins with high importance in physiology and drug

development (Hellmann and Estelle 2002, Burger and Seth 2004, Varshavsky 2008). For instance,

the tumor suppressor protein p53 is kept on at a low steady-state level by Mdm2 that mediates its

turnover through protein degradation (Adams 2003, Zilfou and Lowe 2009, Brady and Attardi

2010). Upon DNA damage by γ radiation, the steady-state is perturbed, and p53 turnover exhibits

an oscillatory behavior indicating that state transitions are occurring repeatedly (Geva-Zatorsky,

Rosenfeld et al. 2006). These examples demonstrate the robustness of regulating degradation rates

to control protein levels.

A major finding from controlling transcription was the experimental observation of low-

pass filtering of transcriptional induction in the yeast *S. cerevisiae* (Bennett, Pang et al. 2008). In

this previous study, Bennett and colleagues modulated the transcription rate of a fluorescent

protein by controlling the extracellular concentration of glucose (transcriptional repressor) over a

constant level of galactose (transcriptional inducer) using a microfluidic system (Bennett, Pang et

al. 2008). Protein turnover faithfully responded to slow glucose fluctuations, while being

insensitive to high-frequency fluctuations. Such low-pass filtering behavior may be important

in preventing distortion of cellular response by high-frequency noise signals from the stochastic

intracellular environment.

Here we applied a light-activated protein degradation machinery (Renicke, Schuster et al.

2013) in *S. cerevisiae* to study the dynamics of protein level fluctuations driven by changes in the

degradation rate. In this approach, a reporter protein is fused to a blue light-activated degradation

signal (photo-sensitive degron, psd), which represents a protein with controllable degradation rates

(**Fig. 1**). We take advantage of the light-controlled degradation machinery to analyze the

frequency-dependent fluctuations in protein turnover, induced by varying frequency of light inputs

that modulate the protein degradation. Square-wave blue light with varying frequencies was

delivered to *S. cerevisiae* cells constitutively expressing psd and Δpsd constructs (**Fig. 1**). Using

this system, we observed that upon application of low-frequency square waves of light, the protein

concentration oscillates with the same corresponding frequency. When the blue light input

frequency increases, the amplitude of oscillation decreases, characteristic of a low-pass filter

response. However, compared to previously reported transcriptionally controlled fluctuations, the

response to degradation-driven fluctuations was faster, enabling protein fluctuations on the order

of 10 minutes. This suggests that compared to controlling gene expression, regulating protein

degradation allows a much faster response in protein levels.

We applied the system identification approach to model the impact of light-activated

degradation on protein turnover. System identification is widely used to study the response of a

system under time-varying stimulations to construct models of dynamical systems from the

measured input and output data (Ljung 1998). We defined a model for the protein turnover based

on system identification to better understand its low-pass filtering characteristics. We modeled the

dynamic of protein turnover with protein production and degradation as inputs and protein level

as the output. Based on our experimental observation on how the protein level changes under

different input fluctuations, we adaptively estimated the model parameters. The estimated model

parameters were used for the frequency-domain analysis, which allowed quantitative comparison

of degradation driven low-pass filtering to previous transcriptional control results.

Figure 1.

a   **psd construct**

b   **Δpsd construct**

Figure 1. Schematic of photo-sensitive degron (psd) constructs used in this study. In the psd construct
(**a**), a protein degradation signal sequence derived from the ornithine decarboxylase is fused to the blue-
light sensing light-oxygen-voltage (LOV) domain from *Arabidopsis thaliana*. Upon blue light activation,
the degradation signal accessibility increases, resulting in faster protein degradation. The Δpsd construct
(**b**), contains all components of psd except the degradation signal sequence.

Results and Discussion

Low-pass filtering of degradation-driven protein level fluctuations in the yeast *S. cerevisiae*

The psd module (**Fig. 1**) comprises of light oxygen voltage 2 domain from *Arabidopsis*

thaliana (*At*LOV2), fused with a derivative of the carboxy-terminal degron of murine ornithine

decarboxylase (ODC), enabling the control of ubiquitin-independent proteasomal degradation rate

upon blue light illumination (Renicke, Schuster et al. 2013). While *At*LOV2 is sensitive and

responsive to blue light, the ODC degradation signal is responsible for increased protein

degradation (**Fig. 1**). To observe the effect of fluctuations in protein degradation rate on protein

turnover, we applied pulses of blue light and monitored the fluorescence emission from the red

fluorescent protein tagRFP fused to psd (**Fig. 1a**). To validate protein degradation due to the ODC

degradation signal, we generated a control construct that lacks the signal sequence, termed Δpsd

(**Fig. 1b**). The expression of these constructs in yeast *S. cerevisiae* was continuously driven using

the constitutive GAL_{1-10} promoter under galactose. To validate continuous expression, we

measured the *tagRFP* transcript level by extracting the mRNA from yeast cells and performing a

quantitative polymerase chain reaction (qPCR) of the transcript (**Fig. 2a,** see **Supp. Fig. 1** and

Supp. Table 1 for RNA quality control and primers used). We illuminated the cells with maximum

blue light intensity used in this study (light power density of 0.027 mW/mm^2 (106.02 μmol/m^{2}s))

and the longest duration ($1/f = 100$ min where f indicates illumination frequency) to assess the

effect of illumination on gene expression. The mRNA was extracted from cells at the start of the

experiment and at the end of each illumination cycle (see inset in **Fig. 2a**). Two primer sets that

bind to different regions of *tagRFP* were used for amplification, along with primers for two

previously validated housekeeping genes, *TFC1* and *UBC6* (Teste, Duquenne et al. 2009), as a

reference. Expression of *tagRFP* remained unchanged for 200 minutes and a significant increase

was detected between 200 to 300 minutes (**Fig. 2a**, $p = 0.014$ and 0.023, student's t-test comparing

ΔCt at 200 min. to that at 300 min. for the two *tagRFP* primer sets, respectively (n = 3 independent

culture preparations)). This result indicates that the GAL_{1-10} promoter allows continuous

expression of the tagRFP constructs, unhindered by light illumination.

Figure 2.

Figure 2. Experimental validation and observation of degradation-driven protein level fluctuations in yeast expressing tagRFP fused to psd or Δpsd. (**a**) qPCR analysis of *tagRFP* transcripts during blue light illumination (inset indicates illumination timepoints). Two sets of primers (primer sets 1 and 2) that bind to different regions of *tagRFP* were used, and previously validated housekeeping genes *TFC1* and *UBC6* were used as references. ΔCt values were calculated by subtracting the Ct of *tagRFP* from the average of the Ct of the two reference genes. *: $p < 0.05$, student's t-test comparing ΔCt at 200 min. to that at 300 min. Error bars indicate standard deviation from n = 3 independent culture preparations. (**b**) Normalized fluorescence level of tagRFP-psd and tagRFP-Δpsd under dark (top panels); tagRFP-Δpsd illuminated with blue light cycles (bottom panels). Solid lines indicate the mean fluorescence intensities and lighter lines indicate standard deviation from n = 10 to 15 colonies of cells. (**c**) Fluorescence of tagRFP-psd under varying frequencies of blue light illumination. *f* denotes the temporal frequency and 1/*f* indicates cycle period. Solid lines indicate the mean fluorescence intensities and lighter lines indicate standard deviation from n = 10 to 15 colonies. (**d**) Experiment for 1/*f* = 20 min. was repeated with 25 colonies of cells. Same intensity of blue light (power density of 0.027 mW/mm^2 (106.02 μmol/m^{2}s)) was used in all panels. In panels (b) and (c), data for each illumination frequency was obtained in separate experiments using independent cultures of cells.

Since the cell cycle of *S. cerevisiae* is on the order of 100 min. (Herskowitz 1988), we

observed cell division during the repeated illumination cycles (**Supp. Fig. 2**). Proteins in each cell

are divided into daughter cells, resulting in dilution of proteins. One approach to quantify the entire

amount of proteins throughout cell division is to sum the fluorescence intensity from a colony of

cells originating from mother cells (Talia, Skotheim et al. 2007). In agreement with previous

studies, we found that summing the fluorescence intensity of a colony of cells results in continuous

increase (red line in **Supp. Fig. 2**), while the fluorescence intensity divided by the area of cells

fluctuates as the cells divide (blue line in **Supp. Fig. 2**). To capture the change in protein level

decoupled from cell division, we used the sum of fluorescence intensities of a colony of cells as a

measure of protein levels throughout this study.

We then tested the effect of pulses of blue light on yeast cells expressing tagRFP-psd or

tagRFP-Δpsd. In the absence of blue light, the normalized fluorescence level of tagRFP-psd and

tagRFP-Δpsd (see **Materials and Methods** section for calculation) showed a continuous increase

over 300 min. (**Fig. 2b**). The fluorescence level of tagRFP-Δpsd was generally higher than that of

tagRFP-psd (**Fig. 2b**), likely due to the fact that the LOV domain conformation is in equilibrium

between active and inactive states even under dark (Strickland, Yao et al. 2010), resulting in slight

background degradation even without blue illumination. As anticipated, the level of tagRFP-Δpsd

was unresponsive to two distinct frequencies of illumination (**Fig. 2b**, $f = 1/100$ mins and $f = 1/80$

mins, respectively in experiments 1 and 2). In contrast, fluorescence level of tagRFP-psd showed

distinct fluctuations with varying frequency of blue light (**Fig. 2c**). We repeated the experiment 5

times, each at a distinct illumination frequency (**Fig. 2c**). Fluctuation in protein level was clearly

visible up to illumination frequency of $1/f = 40$ mins, but not at the highest illumination frequency

tested ($1/f = 20$ mins) (**Fig. 2c**). When the experiment for $1/f = 20$ mins was repeated with a larger

number of colonies cells, the fluctuation was discernable albeit markedly reduced (**Fig. 2d**). Based

on these findings, we conclude that the output fluctuations observed in the experiment are the

result of light-induced increase in protein degradation rate mediated by the psd.

The reduction in fluctuations in protein levels at higher input illumination frequency is

characteristic of low-pass filtering (Antoniou 2016), and indicate that the input perturbation

driving protein degradation has been filtered out at these higher frequencies. To analyze the low-

pass behavior of the degradation driven protein turnover, we determined the time delay between

the input blue light illumination frequency and output protein fluorescence intensity using cross-

correlation. We calculated the cross-correlation (see **Supplementary Information** for calculation)

for experiments using tagRFP-psd shown in **Fig. 2c**. Since the protein level decreases upon

illumination, a minimum in the cross-correlation occurs (**Fig. 3**, indicated by arrows). Based on

the minimum value for each experiment, the time delay between the input and output signals

ranged from 1 to 15 minutes (see **Supplementary Information** for calculation). We find that the

time at which minima occur shortens as the input frequency fluctuations increases, characteristic

of a low-pass filtering system.

Figure 3.

Figure 3. Cross-correlation between input blue light illumination and protein fluorescence intensity.
The curves are cross-correlations resulted from four experiments with varying input frequencies indicated
in the inset. The arrows show the minimum (after time 0) of each curve. Note that the minimums do not
directly indicate the time delay, but allow the calculation of the time delay as shown in the Supplementary
Information.

Degradation-driven fluctuations in a faster maturing fluorescent protein.

Fluorescent proteins contain a chromophore formed through autocatalytic maturation
(Wachter 2007, Craggs 2009, Subach and Verkhusha 2012). The fluorescence is only detected
after maturation, effectively causing a slowdown of apparent protein production rate. Although
not tested, a shift in protein maturation time is expected to impact the low-pass filtering behavior

when transcription rate is modulated as in Bennett *et al.*'s work (Bennett, Pang et al. 2008). In

contrast, when protein degradation rate is modulated, as in our experiments, a shift in protein

maturation time should not affect the low-pass filtering behavior.

To test this, we replaced tagRFP in tagRFP-psd with the fluorescent protein mCherry

(mCherry-psd), which has a significantly shorter maturation time compared to that of tagRFP

(maturation half-time of ~40 minutes for mCherry vs. ~100 minutes for TagRFP (Merzlyak,

Goedhart et al. 2007)). We then repeated the experiment shown in **Fig. 2c** using mCherry-psd and

monitored the mCherry fluorescence in colony numbers similar to that in **Fig. 2d** (25 to 30 colonies

having 4 to 8 cells at the beginning of the experiment). Clear fluctuations in mCherry fluorescence

were visible, down to illumination frequency of $1/f = 20$ min (**Fig. 4a**). We also conducted more

experiments with higher frequencies of input ($1/f = 10$ min and $1/f = 5$ min) and detected small but

distinguishable fluctuations (**Fig. 4b** and see **Supp. Fig. 4** for extended experiments). We observed

consistent decrease in the amplitude of output fluctuations upon increasing the frequency of

degradation fluctuation, characteristic of a low-pass filtering behavior.

Figure 4.

Figure 4. Experimental observation of degradation-driven protein level fluctuations in yeast expressing mCherry fused to psd. (**a**) Fluorescence of mCherry-psd under varying frequencies of blue light illumination. (**b**) Fluorescence of mCherry-psd under higher frequency illumination. For frequency $1/f$ = 20 min., experiment in panel (a) was repeated. For all panels, solid lines indicate the mean fluorescence intensities and lighter lines indicate standard deviation from n = 25 to 30 colonies. Blue light intensity was 0.027 mW/mm² (106.02 μmol/m²s)) for all panels.

System identification and frequency-domain analysis.

To quantitatively assess the low-pass filtering, we constructed a mathematical model for degradation-driven protein turnover using the difference equation approach we previously described (Shin 2016). The model equation can be expressed as:

$$237 \qquad y(n) = p_y y(n-1) + p_{pro} u_{pro}(n-1) + p_{deg} u_{deg}(n-1) \qquad (1)$$

where n is the discrete-time index, y(n) is the normalized fluorescence intensity at time n, $u_{pro}(n)$

is the normalized amount of galactose (transcription inducer) at time n, $u_{deg}(n)$ is the normalized

intensity of blue light (degradation rate controller) at time n. Parameters p_y, p_{pro}, and p_{deg} indicate

endogenous degradation, production, and light-induced degradation, respectively. These

parameters can be adaptively estimated from the experimental data using the normalized least

mean square (NLMS) algorithm, one of commonly used adaptive system identification algorithms

(Sayed 2011). Assuming constant protein production, as indicated by relatively consistent

transcription (**Fig. 2a**), eq. 1 can be simplified by approximating $u_{pro}(n)$ to 1:

$$246 \qquad y(n) = p_y y(n-1) + p_{pro} + p_{deg} u_{deg}(n-1) \qquad (2)$$

247   The above equation represents a multiple-input single-output (MISO) system that can be

248   transformed into the frequency domain using the z–transform.

$$249 \qquad Y(z) = \frac{p_{deg}}{z-p_y} U_{deg}(z) + \frac{p_{pro}z^2}{(z-1)(z-p_y)} \qquad (3)$$

250   Eq. 3 illustrates the relationship between the single output Y(z) and two independent inputs (blue

251   light intensity, $U_{deg}(z)$, and the normalized galactose concentration which is assumed to be constant

252   at 1). The equation can be re-expressed in matrix form:

$$253 \qquad Y(z) = \begin{bmatrix} \frac{p_{deg}}{z-p_y} & \frac{p_{pro}z^2}{(z-1)(z-p_y)} \end{bmatrix} \begin{bmatrix} U_{deg}(z) \\ 1 \end{bmatrix} = [H_1(z) \quad H_2(z)] \begin{bmatrix} U(z) \\ 1 \end{bmatrix} \qquad (4)$$

$$254 \qquad H_1(z) = \frac{p_{deg}}{z-p_y} \qquad (5)$$

$$255 \qquad H_2(z) = \frac{p_{pro}z^2}{(z-1)(z-p_y)} \qquad (6)$$

where $H_1(z)$ shows the relationship between the output $Y(z)$ (protein fluorescence intensity) and

input $U_{deg}(z)$ (blue light intensity). The unknown parameter values p_y and p_{deg} in $H_1(z)$ were

estimated using the NLMS adaptive system identification algorithm (See **Supplementary**

Information and **Supp. Fig. 5**). The Bode plot in **Fig. 5** (indicated as 'Model') shows the

amplitude and phase shift response of $H_1(z)$, calculated based on the mean and standard deviation

of estimated parameter values acquired from each experiment (See **Supplementary Information**).

The amplitude responses clearly show low-pass filtering, indicated by a steady decrease in the

amplitude of the fluctuations as input blue light illumination frequency increases (**Fig. 5a**). We

performed a statistical comparison of experimental results (tagRFP-psd vs. mCherry-psd) as well

as experiments to the model (tagRFP-psd vs. model, mCherry-psd vs. model). The summary of

statistical comparisons in **Table 1** show that the model captured the amplitude and phase shift

responses well. As for amplitude response, no significant difference was found when comparing

model vs. tagRFP-psd as well as model vs. mCherry-psd (**Fig. 5a** and **Table 1**). When comparing

phase shift response of model vs. tagRFP-psd, significant difference was found only for at $1/f =$

100 min and $1/f = 80$ min, but not at other frequencies (**Fig. 5b** and **Table 1**). Phase shift response

of model vs. mCherry-psd was not significantly different at all illumination frequencies (**Fig. 5b**

and **Table 1**). When comparing tagRFP-psd vs. mCherry-psd, mCherry-psd showed significantly

smaller amplitude response at frequencies $1/f = 80$ min, $1/f = 60$ min, and $1/f = 40$ min (**Fig. 5a**

and **Table 1**), but no significant difference was found for phase shift (**Fig. 5b** and **Table 1**). These

results show that both tagRFP-psd and mCherry-psd showed low-pass filtering with similar time

delay, and confirm that the difference in maturation time between tagRFP and mCherry do not

cause a change in the low-pass filtering behavior. Comparing the Bode plot of our experiments

with the previous study using transcriptional control (Bennett, Pang et al. 2008), we found that the

same amplitude of fluctuations (e.g., an amplitude of around 0.3 which is equivalent to -10.5 dB)

was achieved at approximately 3-fold higher frequencies (0.06 rad/min) in degradation-driven

control (achieved at frequency of 0.02 rad/min), indicating a much faster timescale. Taken

together, these results demonstrate that degradation driven control is a robust approach for

controlling protein levels at minutes timescale.

Although we demonstrated this capability in the yeast *S. cerevisiae*, we anticipate the

timescale determined here will apply to many other cell types, considering the UPS is widely

conserved among eukaryotes. In fact, light-sensitive degradation using the psd has been

successfully demonstrated in cell types including human, and other mammalian cells (Sun, Zhang

et al. 2017, Baaske, Gonschorek et al. 2018). Moreover, light-activation is highly suitable for

feedback control since the intensity, frequency, and location of illumination can be instantly

modulated. Since we demonstrated that controlling the degradation rate is sufficient to drive

protein levels under constant transcription, we anticipate this approach to allow generating protein

oscillations previously not achievable using transcriptional control.

Figure 5.

a

b

Figure 5. Model and experimental comparison of low-pass filtering using Bode plots. (**a**) Amplitude response and (**b**) phase shift response from tagRFP-psd, mCherry-psd, and model calculations as a function of input frequency. Data points indicate median and error bars indicate interquartile range (IQR) from four colonies of cells. Statistical analysis of the results is found in Table 1.

295
296
297
298

300 **Table 1.**

Set / Group	f = 2π/100 (1/f = 100 min)		f = 2π/80 (1/f = 80 min)		f = 2π/60 (1/f = 60 min)		f = 2π/40 (1/f = 40 min)		f = 2π/20 (1/f = 20 min)	
	Amplitude	Phase shift	Amplitude	Phase shift	Amplitude	Phase shift	Amplitude	Phase shift	Amplitude	Phase shift
	Median (IQR)									
Model	-12.691 (3.352)	2.586 (0.027)	-14.579 (4.359)	2.554 (0.030)	-20.007 (4.805)	2.510 (0.008)	-23.572 (4.840)	2.444 (0.010)	-25.096 (4.847)	2.300 (0.016)
tagRFP-psd	-11.537 (0.642)	3.016 (0.204)	-11.293 (0.288)	2.906 (0.236)	-16.055 (0.440)	2.985 (0.157)	-17.9937 (0.446)	2.435 (0.550)	-35.654 (0.446)	2.199 (0.471)
mCherry-psd	-15.686 (2.608)	2.953 (0.173)	-15.935 (1.590)	2.710 (0.196)	-22.007 (3.538)	2.251 (0.209)	-25.2071 (4.433)	2.435 (0.314)	-34.437 (4.433)	2.042 (0.314)
P-value	0.095	0.024*	0.023*	0.030*	0.024*	0.115	0.023*	0.981	0.351	0.701

302 *Difference in median value of model, tagRFP-psd, and mCherry-psd assessed using Kruskal-Wallis test, P-value < 0.05

303 Gray highlighted cells in each column indicate pairs with significant difference as assessed using NEMENYI test for multiple pairwise comparisons between the

304 groups, P-value < 0.05

Conclusions

Here we demonstrated robust generation of controlled protein level fluctuations in the yeast *S. cerevisiae* using a blue-light induced protein degradation machinery. We found that by monitoring the total fluorescence of colonies of cells, protein fluctuations with minutes timescale can be reproducibly generated. By using two red fluorescent proteins with distinct maturation time, we show that the delay in protein production rate does not change the protein level fluctuation generated by modulating the protein degradation rate. The results show that controlling protein degradation rates under constant transcription as a means for modulating protein levels, demonstrating a clear pathway for feedback control of protein levels with minutes timescale.

Materials and Methods

Cloning and expression of a photo-sensitive degron in yeast

The photo-sensitive degron (psd) module used in this study has been previously described by Renicke and colleagues (Renicke, Schuster et al. 2013). A red fluorescent reporter protein (tagRFP) was fused to the N-terminus of *At*LOV2. The tagRFP-psd construct was cloned into the pRS316 plasmid (Huang and Shusta 2005) (a gift from Dr. Eric Shusta) using restriction enzymes NotI and BsrGI (New England Biolabs) under the galactose inducible GAL_{1-10} promoter. Psd without the ODC degron (Δpsd) was generated using polymerase chain reaction and cloned using the same restriction sites. The resulting plasmids were named pRS316-tagRFP-psd and pRS316-tagRFP-Δpsd respectively. The plasmids were transformed into the yeast *Saccharomyces*

cerevisiae strain BJ5464 (BJα, α *ura3-52 trp1 leu2Δ1 his3Δ200 pep4:: HIS3 prb1Δ1.6R can1*

GAL) cells, using a commercially available kit (Zymo Research). Transformed cells were grown

in SD-CAA medium (0.1 M sodium phosphate, pH 6.0, 6.7 g/L yeast nitrogen base, 5.0 g/L

casamino acids, 20.0 g/L glucose), supplemented with tryptophan (0.04 g/L) at 30 °C on an orbital

shaker (250 rpm).

qPCR gene expression analysis

Briefly, cells containing pRS316-tagRFP-Δpsd plasmid were grown in SD-CAA medium

supplemented with tryptophan (0.04 g/L) at 30 °C on an orbital shaker (250 rpm) overnight. Yeast

cells grown overnight were diluted to OD_{600} of 1, and the medium was replaced with the induction

medium, SG-CAA (0.1 M sodium phosphate, pH 6.0, 6.7 g/L yeast nitrogen base, 5.0 g/L casamino

acids, 20.0 g/L galactose) supplemented with tryptophan (0.04 g/L) and incubated at 16 °C on an

orbital shaker (250 rpm) for 20 hrs. Then cells were diluted 5-fold in SG-CAA (supplemented with

tryptophan) and grown at 30 °C on an orbital shaker (250 rpm) for period of 400 mins with blue

light illumination with period of $1/f$ = 100 mins. Cells (10 mL of culture) were collected at the

end of each 100 min illumination cycle (blue light power density of 0.027 mW/mm^2 (106.02

μmol/m^{2}s)), and stored at 4 °C until analysis. RNA extraction was performed right after samples

collecting samplesusing RNeasy Mini Kit (Qiagen cat# 74104) according to manufacturer's

protocols. Quality of purified RNAs were checked by measuring the ratio of absorbance at 260 nm

to 280 nm (A260/A280) by using the NanoDrop ND-1000 (Nanodrop Technologies). Integrity and

concentration of RNA were assessed by using the automated electrophoresis station Agilent

Tapestation 4200 (Bio-Rad).

All RNA samples were diluted in RNase/DNase free water (Fisher Scientific cat# BP28191)) at

the same concentration and a volume corresponding to 500 ng was added to iScript™ cDNA

synthesis kit reagents (Bio-Rad cat# 1708890) for cDNA synthesis, per kit instructions. The

reaction mix was incubated for 5 min at 25 °C, followed by 30 min at 46 °C and 5 min at 95 °C

and hold at 4 °C. The cDNA samples were then stored at –80 °C.

qPCR was performed by using SsoAdvanced^{TM} Universal SYBR Green Supermix (Bio-Rad cat#

1725270). 2 µl of cDNA was used per total of 20 µl reaction mixture as a template. *TFC1* and

UBC6 genes were used as housekeeping genes. See Supplementary Fig. 1 and Table 1 for RNA

quality control and primers used. Cycling conditions included a polymerase activation step of 98°C

for 2 min and 40 cycles of 98°C for 2 sec and annealing /extension at 60°C for 5 sec followed by

melt curve analysis from 65 to 95°C in 0.5°C increments. Size of amplicons for each gene were

verified on a 2% agarose gel. ΔCt values were calculated by subtracting the threshold cycle (Ct)

of *tagRFP* from the average of the Ct of the two reference genes.

Light-induced degradation and imaging

Yeast cells grown overnight were diluted to OD_{600} of 1, and the medium was replaced with the

induction medium, SG-CAA (0.1 M sodium phosphate, pH 6.0, 6.7 g/L yeast nitrogen base, 5.0

g/L casamino acids, 20.0 g/L galactose) supplemented with tryptophan (0.04 g/L) at 16 ºC on an

orbital shaker (250 rpm) for 20 hrs. Cells were kept in the dark during the entire induction period

to prevent unwanted degradation. To activate degradation, blue light (470 nm) was delivered at a

power density of 0.027 mW/mm^2 (106.02 µmol/m^{2}s) using a light emitting diode (LED, Thorlabs).

For imaging tagRFP, green light (530 nm) was delivered at a power density of 0.69 mW/mm^2.

Extra precautions were taken to prevent exposure to light after induction until the

experiment was completed. For imaging, cells were put into 96 wells plates coated with

Concanavalin A (2 mg/mL, Sigma Aldrich) to attach the cells to the bottom of plates. Unbound

cells were washed with SG-CAA medium.

To quantify the protein level, the sum of fluorescence intensity of each colony was

measured over the experiment. To identify regions with colonies, 'Otsu dark' auto threshold

method in ImageJ library was used. The mean fluorescence intensity over all colonies of all 5

experiments is defined as the normalization factor. The sum for each colony was then normalized

to the normalization factor. Additionally, an offset specific to each experiment is used in all

experiments' dataset so that the first value starts at a normalized value of 1. In each experiment,

measurements from at least 10 colonies were recorded and averaged.

LED intensities were controlled by an Arduino (See **Supplementary Information**)

controlled using the µManager (Edelstein, Tsuchida et al. 2014). The system snaps images from

all samples every minute while exposing psd and Δpsd samples to blue LED controlled via the

Java programming language (See Supplementary Information). The image processing/analysis

was done using the imageJ/µManager libraries (See **Supplementary Information** for more details

and the codes are available in Github (Mahrou 2021)).

Statistical analysis

To generate the Bode plot, the median phase and amplitude values from each of the 5

frequencies from 4 colonies of cells and median of 4 parameter values estimated for the model

were calculated. The median values between the three datasets (Model, tagRFP-psd, and mCherry-

psd) were compared using the Kruskal-Wallis test. When the median differences between the

groups were statistically significant (P-value<0.05), the NEMENYI test was also used to compare

the median differences between groups of two. Statistical analyses were performed using real

statistics in excel (Charles Zaiontz, www.real-statistics.com).

Acknowledgements

396 This work was funded by the University of Connecticut, the Alzheimer's Association (2019-

397 AARG-NTF-640971), and NIH (1R21NS111358-01A1). We thank Bo Reese and Lu Li of Center

398 for Genome Innovation within the Institute for Systems Genomics at the University of Connecticut

399 for their help in the RNA quality assessment and qPCR analysis. The authors declare that they

400 have no conflicts of interest with the contents of this article.

Author Contributions

402 B.M. designed and performed the experiments, derived the model and analyzed the data. A.P.

403 cloned the psd constructs and conducted the qPCR gene expression analysis. M.H. conducted the

404 statistical analysis. Y.C. and Y.S. supervised the project and assisted with the analysis and the

405 interpretation of data and model. All authors discussed the results and contributed to the

406 manuscript.

References

Adams, J. (2003). "Potential for proteasome inhibition in the treatment of cancer." <u>Drug discovery today</u> **8**(7): 307-315

Antoniou, A. (2016). <u>Digital signal processing</u>, McGraw-Hill.

Baaske, J., P. Gonschorek, R. Engesser, A. Dominguez-Monedero, K. Raute, P. Fischbach, K. Muller, E. Cachat, W. W. A. Schamel, S. Minguet, J. A. Davies, J. Timmer, W. Weber and M. D. Zurbriggen (2018). "Dual-controlled optogenetic system for the rapid down-regulation of protein levels in mammalian cells." <u>Sci Rep</u> **8**(1): 15024.10.1038/s41598-018-32929-7

Baracos, V. E. (2000). "Regulation of skeletal-muscle–protein turnover in cancer-associated cachexia." <u>Nutrition</u> **16**(10): 1015-1018

Bennett, M. R., W. L. Pang, N. A. Ostroff, B. L. Baumgartner, S. Nayak, L. S. Tsimring and J. Hasty (2008). "Metabolic gene regulation in a dynamically changing environment." <u>Nature</u> **454**(7208): 1119

Bintu, L., N. E. Buchler, H. G. Garcia, U. Gerland, T. Hwa, J. Kondev and R. Phillips (2005). "Transcriptional regulation by the numbers: models." <u>Current opinion in genetics & development</u> **15**(2): 116-124

Brady, C. A. and L. D. Attardi (2010). "p53 at a glance." <u>J Cell Sci</u> **123**(15): 2527-2532

Burger, A. M. and A. K. Seth (2004). "The ubiquitin-mediated protein degradation pathway in cancer: therapeutic implications." <u>European journal of cancer</u> **40**(15): 2217-2229

Ciechanover, A. (2017). "Intracellular protein degradation: from a vague idea thru the lysosome and the ubiquitin-proteasome system and onto human diseases and drug targeting." <u>Best Practice & Research Clinical Haematology</u> **30**(4): 341-355

Craggs, T. D. (2009). "Green fluorescent protein: structure, folding and chromophore maturation." <u>Chemical Society Reviews</u> **38**(10): 2865-2875

Edelstein, A. D., M. A. Tsuchida, N. Amodaj, H. Pinkard, R. D. Vale and N. Stuurman (2014). "Advanced methods of microscope control using μManager software." <u>Journal of Biological Methods</u> **1**(2): e10.10.14440/jbm.2014.36

Geva-Zatorsky, N., N. Rosenfeld, S. Itzkovitz, R. Milo, A. Sigal, E. Dekel, T. Yarnitzky, Y. Liron, P. Polak and G. Lahav (2006). "Oscillations and variability in the p53 system." <u>Molecular systems biology</u> **2**(1)

Goldberg, A. (2007). "Functions of the proteasome: from protein degradation and immune surveillance to cancer therapy." <u>Biochemical Society Transactions</u> **35**(1): 12-17

Hellmann, H. and M. Estelle (2002). "Plant development: regulation by protein degradation." <u>science</u> **297**(5582): 793-797

Herskowitz, I. (1988). "Life cycle of the budding yeast Saccharomyces cerevisiae." <u>Microbiol Rev</u> **52**(4): 536-553.10.1128/mr.52.4.536-553.1988

Hill, C. S. and R. Treisman (1995). "Transcriptional Regulation by Extracellular signals: Mechanisms and
Specificity." Cell **80**(2): 199-211.https://doi.org/10.1016/0092-8674(95)90403-4

Hinkson, I. V. and J. E. Elias (2011). "The dynamic state of protein turnover: It's about time." Trends in
cell biology **21**(5): 293-303

Huang, D. and E. V. Shusta (2005). "Secretion and surface display of green fluorescent protein using the
yeast Saccharomyces cerevisiae." Biotechnol Prog **21**(2): 349-357.10.1021/bp0497482

Huang, S. and D. E. Ingber (2000). "Shape-dependent control of cell growth, differentiation, and apoptosis:
switching between attractors in cell regulatory networks." Experimental cell research **261**(1): 91-103

Huang, S. and D. E. Ingber (2007). "A non-genetic basis for cancer progression and metastasis: self-
organizing attractors in cell regulatory networks." Breast disease **26**(1): 27-54

Kneppers, A. E., R. C. Langen, H. R. Gosker, L. B. Verdijk, N. C. Lipovec, P. A. Leermakers, M. C.
Kelders, C. C. de Theije, D. Omersa and M. Lainscak (2017). "Increased Myogenic and Protein Turnover
Signaling in Skeletal Muscle of Chronic Obstructive Pulmonary Disease Patients With Sarcopenia." Journal
of the American Medical Directors Association **18**(7): 637. e631-637. e611

Lee, T. I. and R. A. Young (2013). "Transcriptional regulation and its misregulation in disease." Cell
152(6): 1237-1251

Ljung, L. (1998). System identification. Signal analysis and prediction, Springer**:** 163-173.

Mahrou, B. (2021). "Degradation driven protein turnover." from https://github.com/Bam12006/The-low-
pass-filtering-effect-of-degradation-driven-protein-turnover.

Melendez, J., M. Patel, B. L. Oakes, P. Xu, P. Morton and M. N. McClean (2014). "Real-time optogenetic
control of intracellular protein concentration in microbial cell cultures." Integrative Biology **6**(3): 366-372

Merzlyak, E. M., J. Goedhart, D. Shcherbo, M. E. Bulina, A. S. Shcheglov, A. F. Fradkov, A. Gaintzeva,
K. A. Lukyanov, S. Lukyanov and T. W. Gadella (2007). "Bright monomeric red fluorescent protein with
an extended fluorescence lifetime." Nature methods **4**(7): 555

Milias-Argeitis, A., M. Rullan, S. K. Aoki, P. Buchmann and M. Khammash (2016). "Automated
optogenetic feedback control for precise and robust regulation of gene expression and cell growth." Nature
communications **7**: 12546

Milias-Argeitis, A., S. Summers, J. Stewart-Ornstein, I. Zuleta, D. Pincus, H. El-Samad, M. Khammash
and J. Lygeros (2011). "In silico feedback for in vivo regulation of a gene expression circuit." Nature
biotechnology **29**(12): 1114

Mizushima, N. and D. J. Klionsky (2007). "Protein turnover via autophagy: implications for metabolism."
Annu. Rev. Nutr. **27**: 19-40

Pratt, J. M., J. Petty, I. Riba-Garcia, D. H. Robertson, S. J. Gaskell, S. G. Oliver and R. J. Beynon (2002).
"Dynamics of protein turnover, a missing dimension in proteomics." Molecular & Cellular Proteomics **1**(8):
579-591

Renicke, C., D. Schuster, S. Usherenko, L.-O. Essen and C. Taxis (2013). "A LOV2 domain-based
optogenetic tool to control protein degradation and cellular function." Chemistry & biology **20**(4): 619-626

Sayed, A. H. (2011). Adaptive filters, John Wiley & Sons.

Shin, Y.-J. (2016). "Digital Signal Processing and Control for the Study of Gene Networks." Scientific
reports **6**: 24733

Strickland, D., X. Yao, G. Gawlak, M. K. Rosen, K. H. Gardner and T. R. Sosnick (2010). "Rationally
improving LOV domain-based photoswitches." Nat Methods **7**(8): 623-626.10.1038/nmeth.1473

Subach, F. V. and V. V. Verkhusha (2012). "Chromophore transformations in red fluorescent proteins."
Chem Rev **112**(7): 4308-4327.10.1021/cr2001965

Sun, W., W. Zhang, C. Zhang, M. Mao, Y. Zhao, X. Chen and Y. Yang (2017). "Light-induced protein
degradation in human-derived cells." Biochem Biophys Res Commun **487**(2): 241-
246.10.1016/j.bbrc.2017.04.041

Talia, S. D., J. M. Skotheim, J. M. Bean, E. D. Siggia and F. R. Cross (2007). "The effects of molecular
noise and size control on variability in the budding yeast cell cycle." Nature **448**(7156): 947-
951.10.1038/nature06072

Tautz, D. (2000). "Evolution of transcriptional regulation." Current opinion in genetics & development
10(5): 575-579

Teste, M.-A., M. Duquenne, J. M. François and J.-L. Parrou (2009). "Validation of reference genes for
quantitative expression analysis by real-time RT-PCR in Saccharomyces cerevisiae." BMC molecular
biology **10**(1): 99

Toettcher, J. E., D. Gong, W. A. Lim and O. D. Weiner (2011). "Light-based feedback for controlling
intracellular signaling dynamics." Nature methods **8**(10): 837

Uhlendorf, J., A. Miermont, T. Delaveau, G. Charvin, F. Fages, S. Bottani, G. Batt and P. Hersen (2012).
"Long-term model predictive control of gene expression at the population and single-cell levels."
Proceedings of the National Academy of Sciences **109**(35): 14271-14276

Varshavsky, A. (2008). "Discovery of cellular regulation by protein degradation." Journal of Biological
Chemistry **283**(50): 34469-34489

Vogel, C. and E. M. Marcotte (2012). "Insights into the regulation of protein abundance from proteomic
and transcriptomic analyses." Nature Reviews Genetics **13**(4): 227

Wachter, R. M. (2007). "Chromogenic cross-link formation in green fluorescent protein." Accounts of
chemical research **40**(2): 120-127

Zilfou, J. T. and S. W. Lowe (2009). "Tumor suppressive functions of p53." Cold Spring Harbor
perspectives in biology **1**(5): a001883