TITLE
***pop-1/TCF*, *ref-2/ZIC* and T-box factors regulate the development of anterior cells in the *C. elegans* embryo**

AUTHORS
Jonathan D. Rumley[1], Elicia A. Preston[1], Dylan Cook[1], Felicia L. Peng[1], Amanda L. Zacharias[2, 3], Lucy Wu[1], Ilona Jileaeva[1], John Isaac Murray[1, 4]

1. **Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104**
2. **Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229**
3. **Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH 45267**
4. **Corresponding author - jmurr@pennmedicine.upenn.edu**

ABSTRACT

Patterning of the anterior-posterior axis is fundamental to animal development. The Wnt pathway plays a major role in this process by activating the expression of posterior genes in animals from worms to humans. This observation raises the question of whether the Wnt pathway or other regulators control the expression of the many anterior-expressed genes. We found that the expression of five anterior-specific genes in *Caenorhabditis elegans* embryos depends on the Wnt pathway effectors *pop-1/TCF* and *sys-1/*β-catenin. We focused further on one of these anterior genes, *ref-2/ZIC,* a conserved transcription factor expressed in multiple anterior lineages. Live imaging of *ref-2* mutant embryos identified defects in cell division timing and position in anterior lineages. *Cis*-regulatory dissection identified three *ref-2* transcriptional enhancers, one of which is necessary and sufficient for anterior-specific expression. This enhancer is activated by the T-box transcription factors TBX-37 and TBX-38, and surprisingly, concatemerized TBX-37/38 binding sites are sufficient to drive anterior-biased expression alone, despite the broad expression of TBX-37 and TBX-38. Taken together, our results highlight the diverse mechanisms used to regulate anterior expression patterns in the embryo.

INTRODUCTION

Anterior-posterior patterning regulation

Proper anterior-posterior patterning is critical for animal embryonic development and requires the Wnt pathway across bilaterian animals from worms to humans. Typically, posterior cell identities

are induced by posteriorly-produced Wnt ligands signaling through the canonical Wnt pathway

[1]. In this pathway, signal transduction in response to Wnt activates posterior-expressed gene

transcription through the transcription factor TCF and its co-activator β-catenin [2]. Although this

conserved role for the Wnt pathway in regulating the expression of posterior genes is well

documented [3], much less is known about how genes expressed in anterior cells are regulated.

Caenorhabditis elegans has been widely used to study anterior-posterior patterning because it is

genetically tractable, its embryonic lineage is invariant, and most of its developmental regulators

are highly conserved in vertebrates and other animals [4–10]. Also, the majority of embryonic cell

divisions are oriented along the anterior-posterior axis, and require the Wnt pathway to

differentiate the two daughter fates [11–13]. Following each anterior-posterior cell division, the

Wnt pathway is activated in the more posterior daughter cell, leading to high nuclear SYS-1/β-

catenin and low nuclear POP-1/TCF, whereas the anterior daughter cell has low nuclear SYS-1

and high nuclear POP-1[11–13]. The combination of high nuclear SYS-1 and low nuclear POP-1

in posterior cells makes it stoichiometrically favorable for POP-1 to be bound to posterior-

expressed targets as a complex with SYS-1, thereby driving their expression [14–16]. In contrast,

anterior cells have low nuclear SYS-1 and high nuclear POP-1; for posterior-expressed targets,

this results in POP-1 being bound in the absence of SYS-1, allowing it to recruit co-repressors,

including UNC-37/groucho [17] (Figure 1A). Loss of POP-1/TCF results in posterior genes having

either reduced expression in posterior cells or increased expression in anterior cells, indicative of

its dual roles as an activator and repressor. Loss of SYS-1/β-catenin results only in down-

regulation of posterior genes in posterior cells, consistent with an exclusive role as a posterior

activator [13]. These observations leave open the question of how anterior genes are regulated in

C. elegans.

Figure 1: How are anterior genes regulated? A) The Wnt/β-catenin asymmetry pathway
regulates anterior-posterior patterning by producing asymmetric nuclear concentrations of POP-
1/TCF and SYS-1/β-catenin in anterior and posterior sister cells. Low nuclear concentrations of
POP-1 and high nuclear concentrations of SYS-1 in the posterior cell activate expression of
posterior genes and are associated with the lack of expression of anterior genes. Conversely, high
nuclear concentrations of POP-1 and low nuclear concentrations of SYS-1 in the anterior cell
repress posterior genes and are associated with expression of anterior genes. Effects of *pop-1* and
sys-1 RNAi on posterior genes are shown; effects on most anterior genes are unknown. B) In this
work, we test whether *pop-1*/TCF and *sys-1*/β-catenin regulate anterior genes in the *C. elegans*
embryo, and ask what other factors also regulate anterior expression.

Previous work suggests at least three possible mechanisms by which anterior genes could be

regulated. 1) They may be regulated by the Wnt pathway indirectly, with posteriorly-expressed

canonical Wnt targets repressing the expression of anterior genes in posterior cells. 2) They may

be directly regulated by Wnt pathway components acting in a modified (or "opposite") manner. 3)

They may be regulated by a non-Wnt-related mechanism (Figure 1B).

An example of indirect Wnt pathway regulation through a posterior repressor is seen in the *C.*

elegans embryonic EMS lineage, in which cells derived from the posterior daughter of EMS, E,

express the POP-1-activated, gut-specifying transcription factor *end-1*. In turn, *end-1* represses the

transcription factor *ceh-51*, which is normally expressed in and specifies the lineage derived from

the anterior daughter of EMS, MS [18,19].

Conversely, the *C. elegans* anterior gene *ttx-3* appears to be directly regulated by Wnt pathway

components. *ttx-3* helps specify the AIY neuron class and is regulated by POP-1 and SYS-1 in a

manner dependent on the sole *C. elegans* ZIC family transcription factor REF-2 [20]. After the

AIY grandmother divides, *ttx-3* is expressed in the anterior daughter, the AIY mother, but not in

the posterior daughter. In the AIY mother, the Wnt pathway is inactive, with low nuclear SYS-1

and high nuclear POP-1, and both *pop-1* and *ref-2* are required to activate *ttx-3* expression. In the

posterior sister, the Wnt pathway is active, with high nuclear SYS-1 levels and low nuclear POP-

1 levels, and *sys-1* is required to repress expression of *ttx-3*. POP-1 and REF-2 can directly interact,

suggesting that REF-2 and POP-1 bind as a complex to activate *ttx-3* expression in the AIY mother.

The role of SYS-1 is less clear but some evidence suggests SYS-1 may bind to the POP-1/REF-2

complex to repress expression, or may sequester away the limited POP-1, such that POP-1 cannot

interact with REF-2 [20,21]. Similar regulation has also been seen in other species. For example,

Drosophila TCF/Pangolin can act in a "reverse transcriptional switch" such that when TCF is

bound to a non-optimal site without binding to β-catenin/Armadillo it activates expression, and

when it binds along with β-catenin it represses expression [22].

Because of the role of POP-1 and SYS-1 in regulating the anterior-specific expression of *ttx-3*, we

hypothesized that these Wnt pathway components may also regulate other anterior genes. POP-1

and SYS-1 could act as general anterior expression regulators, interacting with lineage-restricted

co-regulators to ensure appropriate expression of anterior genes in the correct lineages [20,21]. We

tested this by mining single-cell-resolution expression data to identify transcription factors

expressed in anterior-specific patterns [23–25], and show that of five we tested, all require *pop-1*

and/or *sys-1* for either anterior expression or posterior repression. We focus in more detail on a

single anterior-specific gene, *ref-2/ZIC*. By automated lineage analysis of mutants, we find that

ref-2 is required for normal cell division timing and cell position in *ref-2* expressing lineages.

Embryonic expression of *ref-2* is driven by at least three developmental enhancers, two of which

drive early embryonic expression and one of which drives expression in later-stage embryos. The

most distal enhancer (3.9 kb upstream of the transcription start site) drives highly anterior-biased

expression. Functional dissection revealed striking redundancy within this enhancer, reminiscent

of enhancer redundancy that facilitates robust gene expression in other species [26]. We also

identified a role for the T-box transcription factors *tbx-37* and *tbx-38* in driving anterior-specific

expression. Surprisingly, concatemers of a single T-box binding site reiterate much of the anterior-

specific expression pattern of *ref-2* in early embryos, suggesting a key role for these genes in

driving anterior expression.

RESULTS

Many early embryonic transcription factors have anterior-specific expression

We used large-scale single-cell expression databases derived from time-lapse 4D imaging of

embryos expressing reporter genes to identify anterior-expressed genes (Figure 2A). We define

"anterior" or "posterior" lineages as sets of related cells derived from either the more-anterior or

more-posterior daughter after a specific cell division (Figure 2B). Notably, this is distinct from

physical position; for example, a cell that divides near the posterior pole of the embryo gives rise

to both an anterior and posterior daughter lineage, both of which are located in the posterior half

of the embryo (Figure 2B). Following convention [5,27], all lineage trees in this manuscript are

drawn with the more anterior daughters on the left and the posterior daughters on the right of each

bifurcation. Earlier studies identified a strong tendency for genes to be expressed in either multiple

anterior or multiple posterior lineages during early and mid-embryogenesis; these frequently

appear as "lineally repetitive" patterns, with multiple related anterior or posterior cousin lineages

all expressing the same gene (e.g. *nob-1* and *ref-2* in Figure 2D, E) [23,24]. Both anterior-specific

and posterior-specific gene expression were similarly common. While posterior specific genes

(e.g. Figure 2D) appear to be largely regulated by canonical Wnt signaling [13], it is less clear how

anterior-specific patterns are regulated. We mined existing literature and large-scale databases of

fluorescent reporter expression patterns (Figure 2A) [23–25] to identify 19 conserved genes

expressed in anterior lineage-specific patterns (Figure 2C).

Figure 2: Identification of genes with anterior-biased expression. A) Automated lineaging traces lineages and quantifies expression from 4D confocal images. In the lineage trees, vertical lines represent cells, and horizontal lines represent cell divisions. Most cells divide along the anterior-posterior axis, with anterior cells depicted on the left branch and posterior cells on the right branch. B) Posterior and anterior genes are denoted as such based on their expression in cells descended from either a posterior or anterior sister cell, respectively, following an anterior-posterior cell division. Anterior and posterior genes can generally be expressed in cells from any part of the embryo. Anterior and posterior founder cells are labeled with orange asterisks/green triangles respectively. C) List of anterior-biased genes in the EPiC database along with their predicted human homologs. D) Expression pattern of a posterior gene, *nob-1/Hox9-13*. E) Expression patterns of seven anterior genes; most are expressed in unique combinations of mostly anterior lineages.

We measured cellular resolution expression profiles of seven of these by live confocal imaging
followed by automated cell tracking and lineage tracing (Figure 2A, E, Figure S1, Figure S2,
Figure S3) [25,28]. For each gene, we collected 4D time-lapse images of transgenic embryos
expressing a histone-mCherry (*pha-4*, *ref-2*, *elt-1*, *tbx-35*, *irx-1*, and *pes-1*), or GFP (*ceh-51*)
reporter under the control of upstream regulatory sequences of each gene (referred to here as
"promoter reporters"). For five genes we also collected images for fosmid transgenes (*irx-1, elt-1,*
pha-4 and *ref-2)* or CRISPR knock-ins (*ceh-51*) in which the protein is fused to GFP and
surrounded by its normal genomic context ("protein reporters"). Each embryo also contained a
ubiquitously expressed second-color, histone-GFP or histone-mCherry fusion, used for cell
tracking. We identified each nucleus at each time point and traced them across time using
StarryNite [28,29] cell tracking software, and used AceTree [30,31] for subsequent manual error
correction and validation that the extracted lineages were correct. Finally, we quantified the
expression of the reporter in each nucleus at each time point (Figure 2A) [25]. The results confirm
the anterior-specific expression of each gene, and, in some cases, identify previously unknown
expression patterns (Figure S1, Figure S2, Figure S3). For each promoter reporter, the anterior-
specific patterns were consistent, but the protein fusion reporters all had additional expressing
lineages and dynamics indicative of distal enhancers and post-transcriptional regulation (Figure
S1, Figure S2, Figure S3). The anterior-biased expression of these genes is also seen in embryonic
single-cell RNA-seq data (Figures S4 and S5) [32].
177
178   **The Wnt pathway effectors *pop-1* and *sys-1* are required for normal anterior-specific**
179   **expression**
180
Most genes expressed preferentially in posterior lineages require the Wnt effector transcription
factor POP-1/TCF either for activation in posterior cells (together with its coactivator SYS-1/β-
catenin) or for repression in anterior cells, and at least some are direct targets [12,13,33–36].
Although *pop-1* is required for the normal expression of some anterior genes [20,33,37], it is
unclear whether *pop-1* and *sys-1* are generally required for anterior-specific expression, as they
are for posterior-specific expression.
187

To test this, we measured the expression of five anterior gene reporters (*tbx-35*, *ceh-51*, *elt-1*, *irx-*

1, and *ref-2*) after *pop-1* and *sys-1* RNAi by live 4D imaging and lineage tracing as described

above. We used promoter reporters for *tbx-35*, *ceh-51*, *irx-1*, and *ref-2* because our wild-type

expression data indicated these reporters are sufficient for anterior regulation (Figure S1, Figure

S2, Figure S3). For *elt-1*, anterior-specific expression was more robust for the protein reporter so

we tested that reporter's dependence on *pop-1* and *sys-1* (Figure 3; Figure S6).

Reporters of two of the genes we tested, *tbx-35* and *ceh-51*, are expressed solely in the anterior

lineage MS (Figure S2; Figure S6A, B), and previous qualitative expression analyses indicated

that expression persists in the absence of *pop-1* [19,37,38]. Consistent with this, we detected

expression of both genes in the MS lineage after *pop-1* RNAi, however this expression was

significantly decreased. We also observed increased expression of both genes in the E lineage

(posterior sister of MS), although this expression remained lower than both the wild-type and *pop-*

1 RNAi MS levels (Figure 3A, B; Figure S6A, B; Table S4). We also observe low-level ectopic

expression of *tbx-35* in the E lineage under *sys-1* RNAi. These results indicate that both *tbx-35* and

ceh-51 require *pop-1* for full expression in the MS lineage and for repression in the E lineage (as

shown previously [19,37]), and that *sys-1* is required to repress *tbx-35* in the E lineage.

elt-1 reporters are expressed in several early anterior lineages that primarily give rise to ectodermal

fates (ABpla, ABpra, Caa and Cpa), and in one posterior lineage (ABarp) (Figure S1; Figure S6C).

In the Caa and Cpa lineages, anterior lineages that express both the promoter and protein reporters,

pop-1 RNAi results in decreased *elt-1* reporter expression. However, RNAi for neither *pop-1* nor

sys-1 affects expression in their posterior sister lineages Cap and Cpp (Figure 3C; Figure S6C;

Table S4). Similarly, *pop-1* RNAi results in reduced expression of the protein reporter in the ABpla

and ABpra lineages, while RNAi knockdown of neither *pop-1* nor *sys-1* has much effect on their

posterior sister lineages ABplp and ABprp (Figure S6C; Table S4). Thus, in both the ABp and C

lineages, *pop-1* is required for anterior expression of *elt-1*, but neither *pop-1* nor *sys-1* is required

for posterior repression.

Figure 3: Regulation of anterior genes by Wnt effectors *pop-1* and *sys-1*. A-E) Expression pattern (left) and quantification (right) of reporters for the anterior genes *tbx-35* (A), *ceh-51* (B), ELT-1 (C), *irx-1* (D), and *ref-2* (E) in specific anterior lineages and their posterior sisters. Expression is shown under WT conditions and following *pop-1* or *sys-1* RNAi. Expression quantification (right) is the mean expression across all measurements within that lineage from when the reporter is normally first detected until the last time point indicated in the lineage (details in Methods).

irx-1 reporters are expressed early in four related anterior sublineages of ABp, and in three anterior

sublineages of MS, while in later embryos they are expressed in both anterior and posterior

lineages (Figure S3; Figure S6D). Like for *elt-1, pop-1* RNAi reduces *irx-1* reporter expression in

anterior ABp-derived sublineages, and two of the MS-derived expressing sublineages lose

expression nearly completely. After both *pop-1* and *sys-1* RNAi, the *irx-1* reporter expression

expands into several posterior sublineages of ABp whose anterior sisters normally express *irx-1*

(Figure 3D; Figure S6D; Table S4). These results indicate that *irx-1* requires *pop-1* for expression

in anterior lineages and both *pop-1* and *sys-1* for repression in posterior lineages.

Reporters for *ref-2* are expressed in the early embryo in all MS descendants and in several related

anterior sublineages of ABa that arise at the 50-cell stage (Figure S1; Figure S6E). In later

embryos, expression occurs in two anterior sublineages of MS, in several anterior ABp-derived

sublineages, and in the Pn ventral epidermal blast cells, consistent with previous studies [39]. After

pop-1 RNAi, expression of the *ref-2* promoter reporter is significantly reduced in anterior

sublineages of ABala (Figure 3E). As for other genes, *sys-1* RNAi caused de-repression of *ref-2*

in several posterior sisters of the normally expressing sublineages, but we did not detect de-

repression in these sublineages after *pop-1* RNAi (Figure 3E; Figure S6E; Table S4). Thus, *ref-2*

requires *pop-1* for expression in anterior lineages and *sys-1* for repression in posterior lineages.

To summarize these perturbations, each anterior gene tested requires *pop-1* for full expression in

anterior lineages. Several genes also require *pop-1* (*tbx-35*, *ceh-51* and *irx-1*) or *sys-1* (*tbx-35, irx-*

1 and *ref-2*) for repression in some posterior lineages. Thus, it appears that the expression of most

anterior genes requires the Wnt pathway components POP-1 and SYS-1, but this dependency is

complex as was seen previously for posterior genes [13]. Another layer of complexity is that in

some cases, we cannot rule out that the loss of expression after *pop-1* RNAi is, at least in part, due

to indirect effects of *pop-1* loss, such as an earlier anterior to posterior fate conversion. For

example, for *irx-1* expression in descendants of the MS lineage, an MS to E fate conversion could

explain the loss of expression after *pop-1* RNAi. However, in this case, we observe consistent

reduction in *irx-1* reporter expression after *pop-1* RNAi even in embryos without a strong MS to

E conversion (as assayed by cell cycle timing). Expression changes of *irx-1* in the AB lineage

cannot be explained by known fate conversions, indicating that the expression changes are at least

in part due to a direct effect of *pop-1*.

The anterior-specific transcription factor *ref-2/ZIC* is required for normal patterns of cell

division and cell position in expressing lineages.

The striking anterior-specific expression patterns observed for early embryonic TFs raises the

question of whether these factors regulate the development of anterior lineages. To address this

question, we focused on one anterior gene, *ref-2*. *ref-2* encodes a zinc finger transcription factor

homologous to mammalian ZIC genes and *Drosophila* Odd-Paired. Previous work showed that

ref-2 is required to prevent progeny of "Pn" epidermal blast cells from fusing with the major

hypodermal syncytium in larval stages [39] and a predicted null allele of *ref-2*, *ref-2(gk178)*,

causes larval lethality [40]. Recently, REF-2 was shown to activate the anterior-specific expression

of the neural regulator *ttx-3* in the AIY mother cell by binding the *ttx-3 cis*-regulatory region and

recruiting POP-1 independent of POP-1 DNA binding activity [20]. Since *ref-2* is expressed in

many other anterior lineages in early embryos, we investigated whether *ref-2* mutants have

developmental defects in those lineages.

We tested for embryonic lineage phenotypes at single-cell resolution by analyzing 4D time-lapse

images of eight embryos homozygous for the deletion allele *ref-2(gk178)* and expressing a

ubiquitous histone-mCherry fusion for cell tracking. We used StarryNite to track the lineage of

each embryo, and used AceTree to curate the lineages through the >550-cell stage (six embryos)

or >350-cell stage (two embryos). We compared the position and cell cycle length of each cell in

each embryo to those from a reference of 17 wild-type embryos [41,42] to identify outliers

("defects") (Figure 4A).

Figure 4: *ref-2* is required for proper division timing and positioning for many embryonic cells that are descended from cells that express *ref-2*. A) Lineage tree depicting the REF-2 protein expression pattern (top), and chart indicating the number of embryos out of eight homozygous *ref-2* mutant embryos that exhibit cell cycle defects and position defects in indicated cell lineages. Yellow indicates lineages that express *ref-2* early, blue indicates lineages that express *ref-2* late, and the overlap (green) indicates both. A corresponding analysis of wild-type embryos gives 0-1 defects per cell out of all 17 embryos tested [41]. B, C) Scatter plots of mean neighbor score vs mean position deviation of cells of 17 wild-type (B) and eight *ref-2* mutant (C) embryos. Neighbor score is the ratio of each cell's distances to its ten closest wild-type neighbors between mutant and wild-type embryos. Points representing cells are colored based on WT REF-2::GFP expression levels [41]. D) Plot of cell position deviations with arrows starting at the average WT position of cells and pointing to the average position in *ref-2* mutants. Arrows are colored by the WT expression levels of REF-2::GFP for each cell. The labeled cells (ABaraaaapa and ABaraaaapp) have the greatest mean cell position deviations. E) ABaraa lineages of WT and *ref-2* mutant embryos, with cells expressing REF-2::GFP outlined on the WT tree. Two delayed and three missed cell divisions are highlighted on the *ref-2* mutant tree. F) ABaraaa lineage with cells expressing REF-2::GFP outlined. The average position deviations in microns of terminal sister cells are indicated on the terminal branches of the lineage tree. Three dimensional projections of a WT and a *ref-2* mutant embryo are shown with the positions of the terminal ABaraaa lineage cells highlighted. ABaraaaapa and ABaraaaapp are outlined in the WT embryo projection as the cells with the greatest mean position deviation in the *ref-2* mutant.

We identified a total of 55 cells with early, late or missing divisions, and 54 of these were derived

from REF-2::GFP expressing lineages ($P < 10^{-15}$). Overall 95% of cell cycle defects were delays

or missed divisions, indicating a role for *ref-2* in promoting cell division. Similarly, we identified

68 position defects, of which 56 were in *ref-2* expressing lineages (Figure 4B, C, $P = 1.1 * 10^{-14}$,

chi-squared test). Ten *ref-2*-expressing cells, and no non-expressing cells, had position defects in

two or more embryos. The cells with the strongest position and cell cycle length defects were from

the *ref-2*-expressing anterior ABaraaa lineage, which produces primarily anterior pharyngeal cells

(e1, m1 and arcade cells). The most frequent cell cycle length defects were in ABaraaaaa and

ABaraaaap (both defective in 8 of 8 embryos), which were delayed by 21.8 minutes and 14.2

minutes, respectively (Figure 4E). The largest position defects were in ABaraaaapa and

ABaraaaapp, which were mispositioned on average 6 microns posterior of their wild-type position

(Figure 4D, F). However, other defects were broadly distributed across *ref-2*-expressing lineages,

in particular MS- and ABa-derived anterior sublineages (Figure 4A). In summary, the anterior

lineage-expressed TF *ref-2* is required for the normal development of anterior lineages, although

the low penetrance of many defects suggests that *ref-2* may act with other partially redundant

regulators, similar to other early zygotic TFs [41,43–48].

Modular enhancers control *ref-2* embryonic expression

The anterior expression and phenotypes of *ref-2* raise the question of what sequences and

regulators control this expression. To identify genomic sequences responsible for *ref-2* anterior

expression, we first compared expression driven by the 31.9 kb genomic fosmid protein reporter

to the shorter 4.1kb upstream ("promoter") reporter (Figure 5A, B, C; Figure S8). In general, the

patterns were similar; both are expressed broadly in the MS lineage and in multiple anterior ABa-

derived sublineages. In the ABa lineage, the promoter reporter shows background expression in

some posterior sister sublineages suggesting the existence of repressive sequences outside of this

promoter region. Additionally, the promoter fusion reporter is detected persistently in both the MS

and ABa lineages, whereas protein reporter expression is detected transiently, likely reflecting the

use of a stable histone-mCherry reporter for the promoter reporter. Importantly, both the REF-

2::GFP protein and *ref-2* promoter reporters drive expression in progenitors of most cells defective

in *ref-2* mutants.

Figure 5: Identification of *ref-2* enhancers. A) Genome browser screenshot showing the *ref-2* locus on the X chromosome. Displayed tracks include the gene model, sequence conservation among 26 *Caenorhabditis* species [49], ChIP-seq data for TBX-37 and TBX-38 binding, and ATAC-seq data for cells in the ABa and ABp lineages [50]. Boxes indicate candidate enhancer regions as determined by conservation. Below the genome browser screenshot are models of the fluorescent reporter expression constructs we examined. B-H) Lineage trees showing the expression patterns of *ref-2* protein (B), promoter (C), -3.9 kb enhancer reporter (D), promoter lacking the -3.9 kb enhancer (E), -2.8 kb enhancer reporter (F), -1.6 kb enhancer reporter (G), and promoter in the *tbx-37 tbx-38* double mutant (H) as determined by time-lapse confocal microscopy. Sublineages used in anteriority analyses are labeled in B and underlined in black (anterior) and red (posterior) under each lineage tree. Sublineages with transient early REF-2::EGFP (protein reporter) expression are underlined in yellow under the lineage tree in B. Pn ventral epidermal cells are indicated with asterisks in insets of panels B, C, and F. I) Example lineage showing which cells were included in anteriority score analyses. Reporter expression levels were averaged across cells in the anterior or posterior lineages for the indicated cell generations. J) Anteriority scores of the *ref-2* promoter, enhancers, and promoter lacking the -3.9 kb enhancer. Lineages used for analysis are ABala, ABalaa, ABalap, ABalpa, ABalpp, ABaraa, ABarap, ABarpa, and ABarpp (see Figure 5B). K) Full somatic lineage correlation (Spearman's ρ) of each reporter's expression pattern with the mean expression pattern across embryos expressing the promoter reporter.

To identify the sequences in this region that regulate expression in these lineages, we sought to

identify minimal portions of the region with enhancer activity. We identified three portions of the

ref-2 promoter conserved among *Caenorhabditis* species, suggesting they encode important

functions (Figure 5A) [49,51,52]. The most distal of these regions is in open chromatin in ABa-

derived cells as measured in a recently published ATAC-seq data set [50]. Also, this region is

bound by the broadly-expressed ABa lineage transcription factors TBX-37 and TBX-38, as

determined by a recently published ChIP-seq data set (Figure 5A) [50]. We fused each putative

enhancer to a reporter cassette consisting of a minimal *pes-10* promoter and a *his-24*::mCherry

reporter. The *pes-10* promoter is widely used, drives no consistent embryonic expression on its

own [13,53,54] and is compatible with a wide variety of enhancers [13,53]. We introduced each

reporter into worms and used StarryNite to determine in which cells each putative enhancer drives

expression.

The most distal enhancer, located 3.9 kb upstream of the likely transcription start site, (-3.9 kb

enhancer; 449 base pairs) drives early expression (during gastrulation) in several anterior

sublineages of ABa and in anterior sublineages of MS (Figure 5D; Figure S8). This expression

occurs in nearly identical lineages to the early portion of the protein reporter expression pattern,

but while the protein reporter expression in these lineages is transient, the enhancer reporter

persists, again likely because of the use of a stable histone-mCherry reporter. The -3.9 kb enhancer

expression is also well correlated with the pattern driven by the full promoter of *ref-2* (Spearman's

$\rho > 0.6$, Figure 5K). Deleting this enhancer from the *ref-2* promoter reporter results in loss of the

strong anterior-biased expression in the ABalaa, ABalap, and ABarpa lineages, and a loss in the

weak anterior-biased expression in the ABalpa, ABalpp, ABaraa, and ABarap lineages. All

remaining consistent expression occurs in the lineages derived from the Notch-signaled cells

ABalp and ABara and has little anterior bias (Figure 5E, J; Figure; S7; Figure S8) [48]. The

significance of this expression is unclear; it can be seen in the promoter reporter but not the protein

reporter, suggesting that it is normally repressed by additional sequences outside the promoter. We

conclude that the -3.9 kb enhancer is necessary and sufficient for anterior-biased expression in the

ABa lineage.

A second 745 bp enhancer, located at -2.8 kb, drives expression in later embryos (bean stage) in

some anterior sublineages of MS and in the ABp-derived Pn ventral epidermal blast cells that were

previously reported to require *ref-2* [39]. These cells also robustly express the *ref-2* protein reporter

and weakly express the *ref-2* promoter reporter (Figure 5B, C, F, J, K). Finally, the most proximal

enhancer candidate (at -1.6 kb; 206 base pairs) drives variable, weakly anterior-biased, expression

in multiple lineages, most consistently in ABala and ABara (Figure 5G, J, K; Figure S9) and also

drives late expression in some ABp sublineages (Figure 5G; Figure S9). Much of this expression

is in cells or at stages that do not express the full-length reporters, suggesting that the activity of

this sequence differs in its normal genomic context.

To measure anterior expression bias, we developed an "anteriority score" based on the log2 ratio

between anterior and posterior sister lineage expression. Because expression tends to accumulate

over several cell cycles in each expressing lineage, we calculated this score based on the mean

expression of the progeny 2-3 cell divisions after the birth of each anterior lineage (Figure 5I).

Using this metric, we detected robust and significant anterior-biased expression for both the *ref-2*

promoter and -3.9 kb enhancer reporters in ABa-derived sublineages, and reduced or no anterior-

biased expression for the -2.8 kb and -1.6 kb enhancer reporters and for the promoter Δ -3.9 kb

enhancer reporter in the same sublineages (Figure 5J). Comparing the expression driven by each

enhancer to the full promoter showed each is positively, but imperfectly, correlated, consistent

with each driving a subset of the full pattern (Figure 5K).

ref-2* promoter expression requires the T-box transcription factors *tbx-37* and *tbx-38

A dominant feature of the early *ref-2* expression pattern is reiterated expression in six of the eight

anterior sister cells of the ectodermal ABa lineage at the 50-cell stage (AB32 stage--when there

are 16 ABa descendants). This expression is first observed in the fluorescent reporter strains in

pairs of sister cells in the following cell generation (AB64 stage--when there are 32 ABa

descendants). scRNAseq data are consistent with mRNA being expressed in the parent cell, with

reporter detection presumably delayed by the time required for the fluorophores of mCherry or

GFP to mature (Figure S5). This raises the question of whether *ref-2* expression requires the ABa-

specific transcription factors *tbx-37* and *tbx-38*. These genes encode redundant paralogous T-box

family transcription factors which are expressed throughout the ABa lineage and are required for
multiple cell fate decisions within ABa [46,50,55]. Confirming previous reports [46,50], a TBX-
38::GFP knock-in reporter made by CRISPR [50] is expressed throughout the ABa lineage and is
detectable at least from the AB16 to AB128 stages (Figure S10) (*tbx-37* and *tbx-38* RNA are
detectable by the AB4 stage [56]). To test whether *tbx-37* and *tbx-38* are necessary for *ref-2*
expression, we measured expression of the *ref-2* promoter reporter in embryos carrying
homozygous deletions of both *tbx-37* and *tbx-38*. We found that in the absence of *tbx-37* and *tbx-*
38, nearly all *ref-2* promoter expression in the ABa lineage is lost (Figure 5H; Figure S11), while
expression in the MS lineage was maintained. Thus, *tbx-37* and *tbx-38* are necessary for the
anterior-biased expression of *ref-2* in ABa.

The -3.9 kb enhancer contains three non-overlapping sub-enhancers that independently
drive anterior expression in the ABa lineage

Since the *ref-2* -3.9 kb enhancer drives expression in a similar pattern to the early expression of
the full promoter and protein reporters, and its deletion from the *ref-2* promoter results in a
reduction of anterior-biased expression in portions of the ABa lineage, it is a primary enhancer
responsible for driving the early anterior-biased expression of *ref-2*. To map features necessary or
sufficient for anterior expression, we tested the activity of eight truncated versions of the *ref-2* -
3.9 kb enhancer to identify minimal regions of the enhancer sufficient to drive anterior expression
in the ABa lineage (Figure 6, Figure S12). Instead of a single minimal region, we found three non-
overlapping regions within the -3.9 kb enhancer that each are sufficient to drive anterior-biased
enhancer activity in the ABa lineage: a promoter-distal region (bases 1 to 200), a medial region
(bases 227 to 304) and a promoter-proximal region (bases 320 to 449) (Figure 6B, C). Of these
regions, the proximal region drives the most consistent expression and this expression is most
similar to the full length -3.9 kb enhancer construct (Figure 6D), whereas the medial and distal
fragments drive more variable expression and in a subset of these lineages. Thus, although
integration of information between these sub-enhancers is likely necessary for robustness of
expression pattern, the 130 bp proximal fragment provides a model system for further dissection
to identify anterior expression regulators.

Figure 6: Three non-overlapping fragments of the -3.9 kb enhancer are each sufficient to
drive anterior-biased expression in the ABa lineage of the early embryo. A) Model of the *ref-*
2 -3.9 kb enhancer with predicted TBX-37/38 sites (cyan, predicted high affinity sites indicated by
thick line and predicted low-affinity by a thin line), and predicted high affinity POP-1 sites
(magenta). B) Expression patterns driven by full-length *ref-2* -3.9 kb enhancer and by minimal
fragments. C-D) Box plots displaying the anteriority scores (C) and Spearman's ρ (D) for the full-
length *ref-2* -3.9 kb enhancer and minimal fragments. Lineages used to determine anteriority scores
are the same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage and was calculated
relative to the mean expression of the full-length -3.9 kb enhancer. A complete set of enhancer
truncations tested is displayed in Figure S12.

456
457

TBX-37/38 binding sites are required for anterior expression of the proximal region

The proximal fragment exhibits broad sequence conservation with related nematodes, suggesting

it contains multiple important TF binding sites (Figure 5A). To identify sequences necessary for

anterior expression, we scrambled each 15 base-pair region along the length of the enhancer

fragment and tested the resulting sequences for enhancer activity (Figure S13B, C, D). Scrambling

a single region comprising base pairs 410-424 (coordinates relative to the full-length enhancer)

resulted in a complete loss of the anterior-biased expression in the ABa lineage, indicating that

this region is necessary for this expression (Figure 7B; Figure S13B). Scrambling the most

proximal regions 425-439 and 435-449 resulted in a loss of expression in subsets of the ABa

lineage (in ABar and ABalp) but maintained anterior expression in the ABala lineage (Figure

S13B). Expression was largely maintained when other regions were mutated, suggesting sequences

in these regions are not individually necessary for anterior-specific expression despite their

conservation.

The proximal region includes five predicted TBX-37/38 sites, two of which are predicted to have

high affinity for TBX-37 and TBX-38. Two of the sites overlap with either the 410-424 region that

is essential for ABa-specific expression or the 425-449 region that is required for a subset of ABa-

specific expression (Figure 7A, B; Figure S13A, B). To determine whether this region is bound by

TBX-37 and TBX-38 *in vivo,* we mined a recently published dataset that measured genome-wide

TBX-37 and TBX-38 binding by ChIP-seq. Both factors show binding in the proximal region of

the -3.9 kb enhancer (Figure 5A) [50].

To determine whether the TBX-37/38 sites are required for anterior expression, we measured the

activity of the proximal fragment after mutating the two central nucleotides of all TBX-37/38 sites.

The mutated enhancer fragment did not drive anterior-specific expression in the ABa lineage

(Figure 7B, C, D; Figure S13B, C, D; Figure S14), confirming the importance of these sites.

Additionally, the loss of anterior-biased expression following the mutation of these sites indicates

that the two predicted high affinity POP-1 sites in this fragment are not sufficient to drive anterior-

biased expression. Also, these sites are not necessary for the anterior-biased expression driven by

this enhancer fragment, as the loss of these sites does not result in a loss of the anterior-biased

expression (Figure S13B, C, D). In fact, concatemers of POP-1 sites are sufficient to drive

posterior-biased expression (Figure S13F) [13,53].

Since the high affinity site beginning at position 418 (TBX_{418}) overlaps the region (410-424)

required for expression, we hypothesized that this specific TBX-37/38 site may be required for

anterior expression in the ABa lineage. To test this, we tested the activity of the proximal fragment

with only the central two nucleotides of TBX_{418} mutated (Figure 7B, Figure S13B, E). This

mutation resulted in a loss of robust anterior-biased expression driven by the proximal fragment.

Therefore, TBX-37 and TBX-38 bind to the *ref-2* -3.9 kb enhancer proximal fragment and TBX_{418}

is necessary for it to drive robust anterior-biased expression (Figure 5A; Figure 7B, C, D; Figure

S13B, C, D; Figure S14), suggesting that TBX-37 and TBX-38 directly regulate *ref-2* expression.

Figure 7: The ABa-expressed transcription factors *tbx-37* and *tbx-38* are required for the anterior-biased expression of *ref-2* in the ABa lineage. A) Model of the *ref-2* -3.9 kb enhancer with proximal fragment (base pairs 320-449) highlighted. Predicted TBX-37/38 and POP-1 sites are indicated as in Figure 6. B) Expression patterns driven by the proximal fragment of the *ref-2* -3.9 kb enhancer with its WT sequence, with base pairs 410-424 scrambled, with all predicted TBX-37/38 sites mutated, and with TBX_{418} mutated. Expression driven by the proximal fragment with additional regions scrambled or with two predicted POP-1 sites mutated are displayed in Figure S13. C-D) Anteriority scores (C) and Spearman's ρ (D) for the proximal fragment with its WT sequence, with base pairs 410-424 scrambled, with all predicted TBX-37/38 sites mutated, and with TBX_{418} mutated. Lineages used to determine anteriority scores are the same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage and was calculated relative to the mean expression of the WT -3.9 kb enhancer proximal fragment.

Concatemerized TBX sites are sufficient to drive anterior-specific expression in the ABa lineage

Given that *tbx-37* and *tbx-38* are expressed in both anterior and posterior cells within the ABa lineage, we hypothesized that site TBX_{418} would be sufficient to drive broad expression in ABa, with other sequences in the enhancer required for anterior specificity. To test this, we tested the activity of synthetic enhancers comprising concatemerized copies of TBX_{418} separated by either short (6-7 bp) or long (11-15 bp) flanking sequences from its endogenous context (Figure 8A). Surprisingly, both of these reporters drove anterior-specific expression in the ABa lineage similar to that driven by the full proximal fragment (Figure 8B, C, D). This indicates that either TBX_{418} or its flanking sequences are sufficient both for ABa lineage expression, and for anterior-specific expression within this lineage. To test the possibility that the flanking sequences are responsible

for anterior-specificity, we scrambled those sequences while leaving the central *TBX₄₁₈* site intact

in the context of the long-flanking-sequence construct. The resulting reporter was again expressed

only in anterior sublineages similar to the full proximal fragment (Figure 8A, B, C, D). Thus, when

multimerized, *TBX₄₁₈* is sufficient both for ABa expression and for anterior specificity without

additional non-overlapping sequences.

Anteriorly-biased expression of concatemerized TBX sites is regulated by *pop-1*

ref-2 anterior-specific expression is dependent on *pop-1* and *sys-1*, and a concatemerized TBX-

37/38 binding site recapitulates part of this expression. This raises the question of whether the

TBX-37/38 binding site expression is *pop-1*-dependent. To test this, we examined the expression

of the *TBX₄₁₈* concatemer with scrambled flanking regions after *pop-1* RNAi. Knockdown of *pop-*

1 results in changes in the expression pattern, including a significant decrease in anterior

expression bias in the ABalap lineage and a significant decrease in expression level in the ABalaa

lineage (Figure 8E, F, G). We also tested the expression effects of *lit-1* RNAi, which causes an

increase in nuclear POP-1 levels in posterior cells [16,57,58]. Knockdown of *lit-1* results in a

significant decrease in the anterior bias of the *TBX₄₁₈* concatemer expression across several

sublineages of ABa, including a significant increase in expression in the ABalap lineage (Figure

8E, F, G). Some of these changes mirror the fate transformations known to occur after *pop-1* or

lit-1 RNAi, but others, including the increase in ABalap expression after *lit-1* RNAi, do not

[14,59]. Thus, our results indicate that either POP-1 or its targets influence both the lineage

specificity and anterior specificity of expression driven by *TBX₄₁₈*.

Figure 8: TBX_{418} in the *ref-2* -3.9 kb enhancer is sufficient when concatemerized to drive anterior-biased expression in the ABa lineage. A) Models of the proximal fragment of the *ref-2* -3.9 kb enhancer, the TBX_{418} seven-site concatemer with 6-7 base pair flanking regions, the TBX_{418} five-site concatemer with 11-15 base pair flanking regions, and the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions. B) Expression patterns driven by the proximal fragment of the *ref-2* -3.9 kb enhancer and each TBX-37/38 site concatemer. C-D) Anteriority scores (C) and Spearman's ρ (D) for the proximal fragment of the *ref-2* -3.9 kb enhancer and each TBX-37/38 binding site concatemer. E) Expression patterns driven by the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions after *pop-1* RNAi or *lit-1* RNAi. F) Anteriority scores for the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions in indicated lineages following indicated RNAi treatments. G) Quantification of reporter expression for the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions in the ABala lineage. Expression is shown under WT conditions and following *pop-1* or *lit-1* RNAi. Lineages used to determine global anteriority scores are the same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage expression pattern and was calculated relative to the WT proximal fragment.

DISCUSSION

Previous work has definitively shown that the Wnt pathway components POP-1 and SYS-1 regulate posteriorly expressed genes in the *C. elegans* embryo via the canonical Wnt asymmetry pathway [12,13,36]. What has been less clear is how the similar number of anteriorly expressed genes are regulated. We have demonstrated that a set of anteriorly expressed genes, *tbx-35*, *ceh-51*, *elt-1*, *irx-1*, and *ref-2*, require POP-1 and SYS-1 for their anterior-biased expression. Therefore, POP-1 and SYS-1 regulate the expression of anterior-biased genes. The mechanism by which POP-1 and SYS-1 mediate this regulation remains to be determined. Previous work demonstrated that posterior genes can be regulated by POP-1 binding directly to POP-1 binding sites in their upstream regulatory regions [53]. Thus, a simple model of POP-1 regulation of anterior genes would be indirect, with POP-1 activating the expression of repressors in posterior cells. For example, *pop-1* RNAi is known to cause the anterior MS cell to adopt the fate of its posterior sister E due in part to derepression in MS of the E-specific classic Wnt target *end-1*. *end-1* (or its targets) could then act to repress expression of MS-specific genes. This mechanism would predict that anterior genes would be expressed later than posterior genes, or that anterior genes would be expressed early in both anterior and posterior cells, but maintained only in anterior cells. This can happen. For example, in the C lineage *elt-1* is expressed in both anterior and posterior sister cells at the 4C stage, but is only maintained in the anterior granddaughters of C (Caa and Cpa), perhaps due to its repression by the Wnt target *hlh-1* in the descendants of the posterior granddaughters of C (Cap and Cpp) [33]. What we observe in most cases, however, is that anterior gene expression is activated at about the same time as posterior gene expression in sister lineages (Figures S1, S2, S3, S4, S5, and S6, Zacharias et al. 2015 [13]). This observation suggests that POP-1 could regulate some anterior genes directly. Some *pop-1* dependence of anterior genes certainly could be explained by upstream anterior-to-posterior cell fate conversions following *pop-1* RNAi; however, our results as a whole cannot be explained by the known pattern of fate transformations [46].

Indirect *pop-1*-dependent regulation of anterior genes does not explain all of its role in anterior gene expression regulation. For instance, previous work showed that loss of muscle-specifying TFs (MRFs) expressed in Cap and Cpp (*hlh-1*, *unc-120*, and *hnd-1*) is not sufficient to permit ectopic expression of *elt-1* in these cells or to convert them to an epidermal fate [33,60]. Therefore,

another factor, perhaps *pop-1* acting directly, must be responsible for restricting *elt-1* to the

anterior C granddaughters. Our results identified a requirement for *pop-1* in anterior expression of

elt-1 but not posterior repression in the C lineage. The proposed role of the MRFs in repressing

elt-1 expression [33,60,61] suggests that this effect could partly be due to misexpression of MRFs

in the anterior C granddaughters, which could be tested in the future.

Direct activation of genes by POP-1/TCF when Wnt is absent occurs in other systems. For

example, in *Drosophila* TCF/Pangolin can act in a reverse transcriptional switch such that when it

is bound directly to DNA at non-optimal sites it can activate transcription when unbound by β-

catenin, and repress transcription when bound to β-catenin. This difference in the regulatory

activity of TCF is mediated by an allosteric conformational change in the structure of TCF [22].

In *C. elegans*, POP-1 bound together with REF-2 at the *ttx-3 cis*-regulatory region, may similarly

undergo allosteric changes, leading to the opposite regulation of target genes, such that POP-1

activates the expression of *ttx-3* when bound with REF-2 in the absence of SYS-1 and represses

the expression of *ttx-3* when bound with REF-2 and SYS-1 [20,21]. Our findings are consistent

with a similar mechanism regulating *ref-2* itself.

Our observation that five copies of a single 10 base-pair TBX-37/38 binding site are sufficient to

drive anterior-biased expression in the ABa lineage suggests that a POP-1 site is likely unnecessary

for anterior expression. However, *tbx-37* and *tbx-38* cannot explain this anterior bias alone, as

these factors are expressed throughout the ABa lineage. We showed that *pop-1* is required for both

anterior expression and posterior repression of the *ref-2* promoter and for anterior expression of a

TBX-37/38 binding site concatemer. If this role of POP-1 is direct, it likely acts by binding to other

factors, possibly TBX-37 and TBX-38, rather than to DNA directly since the concatemer contains

no sequences with high affinity for POP-1. Consistent with this, TBX-38 and POP-1 were

previously found to physically interact with each other in a yeast 2-hybrid assay [62]. Intriguingly,

several other T-box factors are expressed early in the AB lineage [50,56]; future work should

determine if these play a role in restricting the activity of these sites to anterior lineages. For

instance, TBX_{418} is predicted to be bound by several other *C. elegans* T-box factors. Other T-box

factors expressed in the early ABa lineage could compete for binding at this site, with POP-1

altering the relative activity of these factors between anterior and posterior cells. Additionally, it

is unclear whether *tbx-37* and *tbx-38* also regulate posterior gene expression in the ABa lineage;

intriguingly, the expression of *pha-4*, another anterior gene, in the ABa lineage is also dependent

on *tbx-37* and *tbx-38* [46].

In contrast with our observation that TBX_{418} concatemers are sufficient to drive anterior-biased

expression, Murgan et al. 2015 found that a REF-2 binding site concatemer is not sufficient to

drive expression in the anterior AIY mother cell. Instead, Murgan et al. identified binding sites for

helix-loop-helix family transcription factors that are sufficient to drive expression in the AIY

mother and its posterior sister cell, and that combining these with REF-2 binding sites restricts

expression to the AIY mother. Thus, REF-2 acting through ZIC binding sites is not sufficient to

activate expression without cooperating factors binding other sites whereas TBX-37 and TBX-38

may be sufficient to do so. Other factors could also play a role, for example we used a *pes-10*

minimal promoter, whereas Murgan et al. did not include a minimal promoter in their constructs,

and there could be differences in sensitivity between the microscopy techniques [20].

Classic work in many species, including flies and vertebrates, has found that developmental genes

are often regulated both by partially redundant enhancers and modular enhancers that are

responsible for regulating distinct portions of the genes' expression patterns [63,64]. The extent to

which *C. elegans* genes are regulated by such distal enhancers vs promoter proximal elements has

been unclear. We found that the predicted enhancers in the promoter region of *ref-2* function

modularly, such that each enhancer drives different expression patterns, presumably providing

multiple inputs to fine-tune the expression pattern of *ref-2*. Combined with other recent studies

this adds to evidence that enhancer-mediated regulation is widespread during *C. elegans*

embryonic development [54,65–69]. Additionally, we observe modularity within the -3.9 kb

enhancer, with three non-overlapping regions of the enhancer each sufficient to drive anterior-

biased expression. Since the protein reporter is expressed in more lineages than the promoter

reporter, there are likely even more enhancers outside the promoter region that regulate expression

of *ref-2*.

Several studies have identified evidence for multiple functions of *ref-2*. *ref-2* was first identified

as being required for the production of the Pn.p ventral epidermal cells and for the inhibition of

their fusion to the epidermal syncytium hyp7 [39]. *ref-2* is also required for the initiation of the

differentiation of the cholinergic neuron AIY [40]. Also, *ref-2* has been found to be required for

female fate during sexual development [70]. We have demonstrated a role for *ref-2* in embryonic

development. *ref-2* is required for robust WT cell positioning and cell division timing in anterior

lineages. Intriguingly, an insect ortholog of *ref-2*, *odd-paired*, acts as a regulator of anterior-

posterior expression and is an anterior determinant in several species [71,72], suggesting the

function of *ref-2* in anterior fate regulation may be ancestral. Because the defects in cell position

and division timing are only partially penetrant, there are likely other, partially redundant

regulators of cell position and cell division timing that act with the sole *C. elegans* ZIC homolog

ref-2 during embryonic development. Further experiments, including suppressor and enhancer

screens and co-mutation of other transcription factors will need to be performed to identify other

factors that regulate cell position and division timing in conjunction with *ref-2*.

METHODS

***C. elegans* culture and strain generation**

C. elegans strains (Table S1) were maintained at standard growth temperatures on OP50 *E. coli*

on NGM plates (Table S2). RNAi knockdown was performed by feeding, as previously described

[73,74]. We validated the efficacy of *pop-1* RNAi by measuring the cell cycle delay resulting from

the transformation of the anterior MS lineage into an E-like lineage, the efficacy of *sys-1* RNAi by

failure of morphogenesis and embryonic death, and the efficacy of *lit-1* RNAi by measuring the

cell cycle shortening resulting from the transformation of the posterior E lineage into an MS-like

lineage (Table S2) [13,34,59]. Enhancer reporter strains were generated by microinjection into

RW10029, the GFP histone strain used for lineage tracing. Injection cocktails used to create

extrachromosomal array transgenes consisted of a linear reporter DNA construct at 10 ng/μL, with

5 ng/μL *myo-2*p::GFP, and 135 ng/μL pBluescript vector (or highest concentration possible

whenever 135 ng/μL was impossible to make) and were injected using a Narishige MN-151

micromanipulator with Tritech microinjector system. Enhancer reporters were maintained in

extrachromosomal arrays by screening for worms expressing the *myo-2*p::GFP reporter, which

have a green fluorescent pharynx. Most enhancer reporters, including all for which loss of

expression was observed, were analyzed in at least two strains to control for differences in copy

number and extrachromosomal array artifacts. Other strains were created through crosses using

standard approaches (Table S1).

Generation of transgenes

Candidate enhancers were amplified by PCR from *C. elegans* N2 strain genomic DNA with

Phusion HF polymerase (New England Biosciences) and gel purified (Qiagen). Enhancer

fragments, mutated enhancers, and binding site concatemers were ordered as either gBlocks or

Ultramers from Integrated DNA Technologies (Coralville, Iowa) and were amplified with Phusion

HF polymerase (New England Biosciences) with overhangs for stitching and either gel or PCR

purified (Qiagen). Enhancer reporters were produced by fusing via PCR stitching these constructs

to a *pes-10* minimal promoter::HIS-24::mCherry::let-858 3'UTR fragment amplified from the

POPTOP plasmid [53] (Addgene #34848). The enhancer reporters were purified with a PureLink

PCR purification kit (ThermoFisher) and/or gel purified. The desired product was determined by

its size based on gel electrophoresis. We identified putative transcription factor binding sites using

CIS-BP (http://cisbp.ccbr.utoronto.ca) [75,76].To mutate the TBX-37/38 sites, the central two

nucleotides were altered (e.g. ATAG**TG**TGAA changed to ATAG**GT**TGAA for TBX_{418}). For the

construct with all TBX-37/38 sites mutated, after mutating the five primary sites, subsequent

analysis revealed two weaker sites. All of these predicted TBX-37/38 sites were mutated. The

reporter construct for the *ref-2* promoter lacking the -3.9 kb enhancer was amplified from N2

genomic DNA by PCR and PCR stitched to the *pes-10* minimal promoter::HIS-24::mCherry

reporter, and, thus, drives expression through the *pes-10* minimal promoter. The *pes-10* promoter

drives no consistent embryonic expression on its own prior to elongation [13,53,54] and is

compatible with a wide variety of enhancers [13,53] (Table S5).

Quantitative comparisons

All quantitative comparisons were performed using R version 4.0.3 (The R Foundation for

Statistical Computing), Microsoft Excel version 16.16.27, or Python version 2.7.16. For the main

figures, a minimum N of 6 was used for all experiments, except for ELT-1::GFP expression after

sys-1 RNAi, for which N = 3. For the supplemental figures, a minimum N of 2 was used (largely

for perturbations which appeared similar to wild-type at this level). Code and raw data for these

analyses are available at https://github.com/jisaacmurray/ref-2_paper.

Expression and phenotypic analysis by 4D imaging

We obtained confocal micrographs using a Leica TCS SP5 or Stellaris scanning confocal

microscope (67 z planes at 0.5 μm spacing and 1.5 minute time spacing, with laser power

increasing by 4-fold through the embryo depth to account for attenuation of signal with depth).

Embryos obtained from self-fertilized hermaphrodites were mounted in egg buffer/methyl

cellulose with 20μm beads used as spacers and imaged at 22°C using a stage temperature controller

(Brook Industries, Lake Villa, IL) [77]. We used StarryNite software to automatically annotate

nuclei and trace lineages [29]. We corrected errors from the automated analysis and quantified

reporter expression in each nucleus relative to the local background (using the "blot" background

correction technique) with AceTree software as previously described (Table S3) [23,25].

Anterior gene WT to RNAi within lineage comparisons

To compare the expression levels of the anterior genes *tbx-35*, *ceh-51*, ELT-1, *irx-1*, and *ref-2* and

the TBX_{418} concatemer with 11-15 base pair scrambled flanking regions between WT and RNAi

embryos within the expressing lineages and their posterior sisters, we calculated for each embryo

the mean reporter expression across all measurements within each lineage (all descendant cells

and time points) starting with the cell stage that expresses the reporter at the time cells are born or

shortly afterwards to the last time point in the indicated lineage. For expression level comparisons

of extrachromosomal array transgenes, we normalized expression values to the 75th percentile

expression value of the strain with the higher 75th percentile expression value across all measured

cells in the indicated lineage within each RNAi treatment. The expression analyzed for the MS

lineage starts at an earlier stage than that for other lineages, since following *pop-1* RNAi the cell

divisions are so dramatically delayed that reporter expression comes on in earlier cell generations

in these embryos. The cell generations used in each anterior gene comparison are noted in Table

S4.

Anteriority Scores

To measure the magnitude of anterior expression bias, we developed a measure that we refer to as the anteriority score. To determine the anteriority scores, first we chose a minimum expression cutoff and set all values below this cutoff to 0 and subtracted the minimum cutoff from all expression values greater than or equal to the cutoff (an expression value of 200 was used as the cutoff for all anteriority score analyses, except a value of zero was used for the TBX-38::GFP reporter, which has very low fluorescence intensity compared to the other reporters analyzed). Next, we averaged the expression levels of the reporter of interest in the cells of each anterior and each posterior sister lineage of interest. For the *ref-2* promoter and enhancer constructs, as well as the TBX-37/38 site concatemers, we analyzed anterior and posterior lineages descended from the cells ABala, ABalaa, ABalap, ABalpa ABalpp ABaraa, ABarap, ABarpa, and ABarpp. We used expression values starting at the generation with ABaxxxxxx cells (AB128 stage) until the time point of the last division of AB128 stage cells in the ABa lineage. Next, we calculated the means of the average expression levels of all of the anterior lineages and all of the posterior lineages analyzed for each embryo. We then added a pseudocount of 1 to each of the anterior and posterior embryo means. For each experimental group, the adjusted average anterior expression is divided by the adjusted average posterior expression, and the base 2 logarithm of the result is reported-- Anteriority Score = $\log_2((\text{anterior mean} + 1)/(\text{posterior mean} +1))$.

Correlation analysis

Correlation analysis was done by Spearman's ρ, comparing each embryo's expression to the average expression of the control group. Spearman's ρ was calculated in R using the "cor" function. For Spearman's ρ we used all the data from the lineages of interest without excluding any cell generations or having a minimum value cutoff. We included cells born before the onset of expression here to include information about the cell generation of expression onset in the analysis. For the analysis comparing the *ref-2* promoter with its enhancers we used the full somatic lineage starting with the blastomeres AB, EMS, C, and D. For all other analyses we used only the ABa lineage.

Mutant cell position and cell division analysis

We identified cell position and cell cycle defects as previously described [41]. We corrected for

differences in global division rates, and considered divisions as defective if they deviated from the

wild-type cell cycle length by at least five minutes and had a z-score greater than three. Cell

positions were corrected for differences in embryo size and rotation, and considered defective if

they deviated from the expected wild type position by at least five microns, had a z-score greater

than five, and a nearest neighbor score greater than 0.8.

Statistical analyses

Statistical significance of differences in anterior or posterior expression between untreated and

RNAi-treated embryos was tested using the two-tailed Wilcoxon Ranked Sum test. Statistical

significance of differences in Spearman's ρ or Anteriority Score were tested using the one-tailed

Wilcoxon Ranked Sum test to test for differences from either the control group or from 0, as

indicated in each figure. The p-values calculated by the Wilcoxon Ranked Sum test were corrected

for multiple comparisons using the false discovery rate method in R. Significance of overlap

between positional or cell division defects and expression status were assessed using a Chi-squared

test.

FIGURE LEGENDS

Figure 1: How are anterior genes regulated? A) The Wnt/β-catenin asymmetry pathway
regulates anterior-posterior patterning by producing asymmetric nuclear concentrations of POP-
1/TCF and SYS-1/β-catenin in anterior and posterior sister cells. Low nuclear concentrations of
POP-1 and high nuclear concentrations of SYS-1 in the posterior cell activate expression of
posterior genes and are associated with the lack of expression of anterior genes. Conversely, high
nuclear concentrations of POP-1 and low nuclear concentrations of SYS-1 in the anterior cell
repress posterior genes and are associated with expression of anterior genes. Effects of *pop-1* and
sys-1 RNAi on posterior genes are shown; effects on most anterior genes are unknown. B) In this
work, we test whether *pop-1*/TCF and *sys-1*/β-catenin regulate anterior genes in the *C. elegans*
embryo, and ask what other factors also regulate anterior expression.

820
821
Figure 2: Identification of genes with anterior-biased expression. A) Automated lineaging
traces lineages and quantifies expression from 4D confocal images. In the lineage trees, vertical
lines represent cells, and horizontal lines represent cell divisions. Most cells divide along the
anterior-posterior axis, with anterior cells depicted on the left branch and posterior cells on the
right branch. B) Posterior and anterior genes are denoted as such based on their expression in cells
descended from either a posterior or anterior sister cell, respectively, following an anterior-

posterior cell division. Anterior and posterior genes can generally be expressed in cells from any
part of the embryo. Anterior and posterior founder cells are labeled with orange asterisks/green
triangles respectively. C) List of anterior-biased genes in the EPiC database along with their
predicted human homologs. D) Expression pattern of a posterior gene, *nob-1/Hox9-13*. E)
Expression patterns of seven anterior genes; most are expressed in unique combinations of mostly
anterior lineages.
834

Figure 3: Regulation of anterior genes by Wnt effectors *pop-1* and *sys-1*. A-E) Expression pattern (left) and quantification (right) of reporters for the anterior genes *tbx-35* (A), *ceh-51* (B), ELT-1 (C), *irx-1* (D), and *ref-2* (E) in specific anterior lineages and their posterior sisters. Expression is shown under WT conditions and following *pop-1* or *sys-1* RNAi. Expression

841   quantification (right) is the mean expression across all measurements within that lineage from
842   when the reporter is normally first detected until the last time point indicated in the lineage (details
843   in Methods).
844

845
846
Figure 4: *ref-2* is required for proper division timing and positioning for many embryonic
cells that are descended from cells that express *ref-2*. A) Lineage tree depicting the REF-2
protein expression pattern (top), and chart indicating the number of embryos out of eight
homozygous *ref-2* mutant embryos that exhibit cell cycle defects and position defects in indicated
cell lineages. Yellow indicates lineages that express *ref-2* early, blue indicates lineages that express
ref-2 late, and the overlap (green) indicates both. A corresponding analysis of wild-type embryos
gives 0-1 defects per cell out of all 17 embryos tested [41]. B, C) Scatter plot of mean neighbor
score vs mean position deviation of cells of 17 wild-type (B) and eight *ref-2* mutant (C) embryos.
Neighbor score is the ratio of each cell's distances to its ten closest wild-type neighbors between

mutant and wild-type embryos. Points representing cells are colored based on WT REF-2::GFP
expression levels [41]. D) Plot of cell position deviations with arrows starting at the average WT
position of cells and pointing to the average position in *ref-2* mutants. Arrows are colored by the
WT expression levels of REF-2::GFP for each cell. The labeled cells (ABaraaaapa and
ABaraaaapp) have the greatest mean cell position deviations. E) ABaraa lineages of WT and *ref-
2* mutant embryos, with cells expressing REF-2::GFP outlined on the WT tree. Two delayed and
three missed cell divisions are highlighted on the *ref-2* mutant tree. F) ABaraaa lineage with cells
expressing REF-2::GFP outlined. The average position deviations in microns of terminal sister
cells are indicated on the terminal branches of the lineage tree. Three dimensional projections of a
WT and a *ref-2* mutant embryo are shown with the positions of the terminal ABaraaa lineage cells
highlighted. ABaraaaapa and ABaraaaapp are outlined in the WT embryo projection as the cells
with the greatest mean position deviation in the *ref-2* mutant.
868

869
870

Figure 5: Identification of *ref-2* enhancers. A) Genome browser screenshot showing the *ref-2*
locus on the X chromosome. Displayed tracks include the gene model, sequence conservation
among 26 *Caenorhabditis* species [49], ChIP-seq data for TBX-37 and TBX-38 binding, and
ATAC-seq data for cells in the ABa and ABp lineages [50]. Boxes indicate candidate enhancer
regions as determined by conservation. Below the genome browser screenshot are models of the

fluorescent reporter expression constructs we examined. B-H) Lineage trees showing the
expression patterns of *ref-2* protein (B), promoter (C), -3.9 kb enhancer reporter (D), promoter
lacking the -3.9 kb enhancer (E), -2.8 kb enhancer reporter (F), -1.6 kb enhancer reporter (G), and
promoter in the *tbx-37 tbx-38* double mutant (H) as determined by time-lapse confocal microscopy.
Sublineages used in anteriority analyses are labeled in B and underlined in black (anterior) and red
(posterior) under each lineage tree. Sublineages with transient early REF-2::EGFP (protein
reporter) expression are underlined in yellow under the lineage tree in B. Pn ventral epidermal
cells are indicated with asterisks in insets of panels B, C, and F. I) Example lineage showing which
cells were included in anteriority score analyses. Reporter expression levels were averaged across
cells in the anterior or posterior lineages for the indicated cell generations. J) Anteriority scores of
the *ref-2* promoter, enhancers, and promoter lacking the -3.9 kb enhancer. Lineages used for
analysis are ABala, ABalaa, ABalap, ABalpa, ABalpp, ABaraa, ABarap, ABarpa, and ABarpp
(see Figure 5B). K) Full somatic lineage correlation (Spearman's ρ) of each reporter's expression
pattern with the mean expression pattern across embryos expressing the promoter reporter.
890

891
892
Figure 6: Three non-overlapping fragments of the -3.9 kb enhancer are each sufficient to
drive anterior-biased expression in the ABa lineage of the early embryo. A) Model of the *ref-*
2 -3.9 kb enhancer with predicted TBX-37/38 sites (cyan, predicted high affinity sites indicated by
thick line and predicted low-affinity by a thin line), and predicted high affinity POP-1 sites
(magenta). B) Expression patterns driven by full-length *ref-2* -3.9 kb enhancer and by minimal
fragments. C-D) Box plots displaying the anteriority scores (C) and Spearman's ρ (D) for the full-
length *ref-2* -3.9 kb enhancer and minimal fragments. Lineages used to determine anteriority scores
are the same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage and was calculated
relative to the mean expression of the full-length -3.9 kb enhancer. A complete set of enhancer
truncations tested is displayed in Figure S12.
903

904
905
Figure 7: The ABa-expressed transcription factors *tbx-37* and *tbx-38* are required for the
anterior-biased expression of *ref-2* in the ABa lineage. A) Model of the *ref-2* -3.9 kb enhancer
with proximal fragment (base pairs 320-449) highlighted. Predicted TBX-37/38 and POP-1 sites
are indicated as in Figure 6. B) Expression patterns driven by the proximal fragment of the *ref-2* -
3.9 kb enhancer with its WT sequence, with base pairs 410-424 scrambled, with all predicted TBX-
37/38 sites mutated, and with TBX_{418} mutated. Expression driven by the proximal fragment with
additional regions scrambled or with two predicted POP-1 sites mutated are displayed in Figure
S13. C-D) Anteriority scores (C) and Spearman's ρ (D) for the proximal fragment with its WT
sequence, with base pairs 410-424 scrambled, with all predicted TBX-37/38 sites mutated, and
with TBX_{418} mutated. Lineages used to determine anteriority scores are the same as in Figure 5J.
Spearman's ρ analysis uses the full ABa lineage and was calculated relative to the mean expression
of the WT -3.9 kb enhancer proximal fragment.
918

Figure 8: *TBX₄₁₈* in the *ref-2* -3.9 kb enhancer is sufficient when concatemerized to drive anterior-biased expression in the ABa lineage. A) Models of the proximal fragment of the *ref-2* -3.9 kb enhancer, the TBX_{418} seven-site concatemer with 6-7 base pair flanking regions, the TBX_{418} five-site concatemer with 11-15 base pair flanking regions, and the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions. B) Expression patterns driven by the proximal fragment of the *ref-2* -3.9 kb enhancer and each TBX-37/38 site concatemer. C-D) Anteriority scores (C) and Spearman's ρ (D) for the proximal fragment of the *ref-2* -3.9 kb enhancer and each TBX-37/38 binding site concatemer. E) Expression patterns driven by the TBX_{418} five-site concatemer with 11-15 base pair scrambled flanking regions after *pop-1* RNAi or

lit-1 RNAi. F) Anteriority scores for the *TBX$_{418}$* five-site concatemer with 11-15 base pair
scrambled flanking regions in indicated lineages following indicated RNAi treatments. G)
Quantification of reporter expression for the *TBX$_{418}$* five-site concatemer with 11-15 base pair
scrambled flanking regions in the ABala lineage. Expression is shown under WT conditions and
following *pop-1* or *lit-1* RNAi. Lineages used to determine global anteriority scores are the same
as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage expression pattern and was
calculated relative to the WT proximal fragment.
937

938
939

940   **Figure S1: Several early embryonic genes are expressed with anterior bias.** Full lineage
941   expression patterns are shown for integrated transgenic promoter reporters of *ref-2* and *elt-1*; and
942   for integrated transgenic protein reporters of REF-2 and ELT-1.

943
944

945 **Figure S2: Several early embryonic genes are expressed with anterior bias.**
946 Full lineage expression patterns are shown for integrated transgenic promoter reporters of *tbx-35*
947 and *ceh-51*; for an integrated transgenic protein reporter of PHA-4; and for a CRISPR knock-in
948 protein reporter of CEH-51.
949

950
951

952 **Figure S3: Several early embryonic genes are expressed with anterior bias.**
953 Full lineage expression patterns are shown for integrated transgenic promoter reporters of *irx-1*
954 and *pes-1*; and for an integrated transgenic protein reporter of IRX-1.
955

tbx-35

ceh-51

Figure S4: scRNA-seq data confirm imaging-based expression data. scRNA-seq expression data for *tbx-35* and *ceh-51*. *tbx-35* expression lineage derived from Packer et al. 2019 indicates expression in the daughters of MS (MSa and MSp) [32]. Early *C. elegans* embryo diagram derived from Tintori et al. 2016 indicates *tbx-35* expression in MS and its daughters, as well as low-level expression in E and ABpl [56]. *ceh-51* expression lineage derived from Packer et al. 2019 indicates expression in the descendants of MS from the MS2 through the MS16 generation (MS daughters through MS great-great granddaughters) [32]. Early *C. elegans* embryo diagram derived from Tintori et al. 2016 indicates *ceh-51* expression in the daughters of MS [56]. In both cases scRNA-seq expression data agree with imaging-based expression data, albeit with expression being detected earlier and persisting for fewer cell generations than for the fluorescent reporters (Figure 3 and Figure S6). Expressing cells in early embryo diagrams are labeled with their names. Gray branches on lineage trees indicate cells for which distinct transcriptomes could not be determined.

Figure S5: scRNA-seq data confirm imaging-based expression data. scRNA-seq expression data for *ref-2*, *irx-1*, and *elt-1* derived from Packer et al. 2019 [32]. scRNA-seq expression data largely agree with imaging-based expression data, albeit with expression being detected earlier and persisting for fewer cell generations than for the fluorescent reporters (Figures 3 and 5 and Figure S6). Note that while most anterior expression events are resolved in these data, a few of the ABa lineage *ref-2* expression domains are not (because the anterior vs posterior sister cells were not separately annotated at the stage when *ref-2* is expressed). However, the data that are resolved for *ref-2* are consistent with the promoter and fosmid reporter expression patterns. Gray branches on lineage trees indicate cells for which distinct transcriptomes could not be determined.

981
982
983 **Figure S6: Anterior genes are regulated by Wnt effectors *pop-1* and *sys-1*.** Full example WT,
984 *pop-1* RNAi, and *sys-1* RNAi lineages are shown for integrated transgenic promoter reporters of
985 *tbx-35* (A), *ceh-51* (B), *irx-1* (D), and *ref-2* (E) and for an integrated transgenic protein reporter of
986 ELT-1 (C).
987

988
989 **Figure S7: Deletion of the -3.9 kb enhancer from the *ref-2* promoter causes a reduction in**
990 **the anterior bias in the expression driven by the promoter in most ABa sublineages in which**
991 **the full-length promoter drives anterior-biased expression.** Box plots display anteriority scores
992 of lineages descended from indicated cells.
993

994
995 **Figure S8: Expression patterns driven by the *ref-2* promoter, the -3.9 kb enhancer, and the**
996 **promoter lacking the -3.9 kb enhancer are somewhat variable.** Displayed here are partial
997 lineages for all analyzed embryos bearing these reporters to show the variation in their expression.
998

REF-2::EGFP

ref-2 -1.6 kb enhancer

Expression

None   Low   High

1001   **Figure S9: Expression patterns driven by the *ref-2* -1.6 kb enhancer are variable.** Displayed
1002   here are full lineages for all analyzed embryos bearing this reporter to show the variation in its
1003   expression. The full lineage expression pattern of the REF-2::GFP protein reporter is shown for
1004   comparison.
1005

ref-2 Promoter and TBX-38 Protein Anteriority Scores

ref-2 promoter

ABa

A

P

TBX-38 protein

A

P

Expression

None   Low   High

ref-2 promoter

TBX-38 protein

p < 0.001

-2   0   2   4   6   8   10

Figure S10: A TBX-38::GFP CRISPR knock-in protein reporter has minimally anterior-
biased expression in the lineages in which the *ref-2* promoter drives anterior-biased
expression. Displayed are partial lineages of the P*ref-2*::*his-24*::mCherry promoter reporter
expression and the TBX-38::GFP CRISPR knock-in expression. The box plot shows the mean
anteriority scores for the *ref-2* promoter reporter and the TBX-38::GFP CRISPR knock-in for the
same lineages as in Figure 5J.
1014

1015
1016
Figure S11: *tbx-37* and *tbx-38* are required for *ref-2* promoter-driven anterior-biased
expression in the ABa lineage. Displayed are partial lineages for the expression driven by the *ref-*
2 promoter in a WT background and in a *tbx-37(tm314) tbx-38(tm581)* double mutant background.
Also shown are box plots of anteriority scores and Spearman's ρ for each of these groups,
indicating a loss of anterior bias and correlation to WT for the *ref-2* promoter expression pattern
in *tbx-37 tbx-38* double mutant embryos. Lineages used to determine anteriority scores are the
same as in Figure 5J Spearman's ρ analysis uses the full ABa lineage expression pattern and was
calculated relative to the *ref-2* promoter in the WT background.
1025

Figure S12: Three non-overlapping fragments of the -3.9 kb enhancer are each sufficient to drive anterior-biased expression in the ABa lineage of the early embryo. A) Model of the *ref-2* -3.9 kb enhancer with predicted TBX-37/38 and POP-1 sites indicated as in Figure 6. B) Expression patterns driven by full-length *ref-2* -3.9 kb enhancer and by all tested fragments. C-D) Box plots displaying the anteriority scores (C) and Spearman's ρ (D) for the full-length *ref-2* -3.9 kb enhancer and all tested fragments. Lineages used to determine anteriority scores are the same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage and was calculated relative to the full-length -3.9 kb enhancer.

Figure S13: The broadly-expressed transcription factors *tbx-37* and *tbx-38* are required for the anterior-biased expression of *ref-2* in the ABa lineage. A) Model of the *ref-2* -3.9 kb enhancer with proximal fragment (base pairs 320-449) highlighted. Predicted TBX-37/38 and POP-1 sites are indicated as in Figure 6. B) Expression patterns driven by the proximal fragment of the *ref-2* -3.9 kb enhancer with its WT sequence, with each 15 base-pair segment of the fragment scrambled, with all predicted TBX-37/38 sites mutated, with *TBX_418* mutated, and with the two

predicted POP-1 sites mutated. C-D) Box plots displaying the anteriority scores (C) and
Spearman's ρ (D) for the proximal fragment with its WT sequence, with each 15 base-pair segment
of the fragment scrambled, with all predicted TBX-37/38 sites mutated, with TBX_{418} mutated, and
with the two predicted POP-1 sites mutated. Lineages used to determine anteriority scores are the
same as in Figure 5J. Spearman's ρ analysis uses the full ABa lineage and was calculated relative
to the WT proximal fragment. E) Models of the proximal fragment of the *ref-2* -3.9 kb enhancer
with the WT sequence of TBX_{418} and the mutated sequence used in the TBX_{418} mutant reporters.
F) Expression pattern driven by the seven-copy POP-1 site concatemer POPTOP [13,53].
1053

1054

1055

Figure S14: *ref-2* -3.9 kb enhancer WT proximal fragment, proximal fragment with all TBX-37/38 sites mutated, and proximal fragment with *TBX₄₁₈* mutated exhibit variation in their expression patterns. Displayed are partial lineages for all analyzed embryos expressing reporters for the *ref-2* -3.9 kb enhancer WT proximal fragment, the *ref-2* -3.9 kb enhancer proximal fragment with base pairs 410-424 scrambled, the *ref-2* -3.9 kb enhancer proximal fragment with all TBX-37/38 sites mutated, and the *ref-2* -3.9 kb enhancer proximal fragment with *TBX₄₁₈* mutated to show the variation in the expression pattern for each of these reporters.

1063

TABLES

Table S1:
Caenorhabditis elegans strain list

Table S2:
Escherichia coli strain list

Table S3:
Quantitative analyses raw data

Table S4:
Anterior gene summary expression data

Table S5:
Recombinant DNA list

ACKNOWLEDGMENTS

We thank J. Archibald Millar, Julia Richards and Shaili Patel for technical contributions to data collection and analysis. We thank Meera Sundaram for use of her injection equipment. We thank Christopher Brown, Meera Sundaram, Michael Atchison, Peter Klein, and Mary Mullins for helpful discussions. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We also thank the Waterston lab, the Maduro lab, and the Cochella lab for kindly providing *C. elegans* strains. We thank the *C. elegans* Reverse Genetics Core Facility at the University of British Columbia, which as part of the international *C. elegans* Gene Knockout Consortium produced the *ref-2* mutant allele (*ref-2(gk178)*) used in this study. This work was funded by T32GM008216, F31GM123737, and R35GM130357.

References Cited

1. Hikasa H, Sokol SY. Wnt signaling in vertebrate axis specification. Cold Spring Harb
Perspect Biol. 2013;5: a007955. doi:10.1101/cshperspect.a007955

2. Archbold HC, Yang YX, Chen L, Cadigan KM. How do they do Wnt they do?: regulation of
transcription by the Wnt/β-catenin pathway. Acta Physiol Oxf Engl. 2012;204: 74–109.
doi:10.1111/j.1748-1716.2011.02293.x

3. Martin BL, Kimelman D. Wnt signaling and the evolution of embryonic posterior
development. Curr Biol CB. 2009;19: R215–R219. doi:10.1016/j.cub.2009.01.052

4. Zacharias AL, Murray JI. Combinatorial decoding of the invariant C. elegans embryonic
lineage in space and time. genesis. 2016;54: 182–197. doi:10.1002/dvg.22928

5. Sulston JE, Schierenberg E, White JG, Thomson JN. The embryonic cell lineage of the
nematode Caenorhabditis elegans. Dev Biol. 1983;100: 64–119. doi:10.1016/0012-
1606(83)90201-4

6. Bürglin TR, Finney M, Coulson A, Ruvkun G. Caenorhabditis elegans has scores of
homoeobox-containing genes. Nature. 1989;341: 239–243. doi:10.1038/341239a0

7. Chen W, Lim HH, Lim L. A new member of the ras superfamily, the rac1 homologue from
Caenorhabditis elegans. Cloning and sequence analysis of cDNA, pattern of developmental
expression, and biochemical characterization of the protein. J Biol Chem. 1993;268: 320–
324. doi:10.1016/S0021-9258(18)54152-1

8. van den Heuvel S. Cell-cycle regulation. WormBook. 2005. pp. 1–16.
doi:10.1895/wormbook.1.28.1

9. Shaye DD, Greenwald I. OrthoList: A Compendium of C. elegans Genes with Human
Orthologs. PLOS ONE. 2011;6: e20085. doi:10.1371/journal.pone.0020085

10. Kim W, Underwood RS, Greenwald I, Shaye DD. OrthoList 2: A New Comparative
Genomic Analysis of Human and Caenorhabditis elegans Genes. Genetics. 2018;210: 445–
461. doi:10.1534/genetics.118.301307

11. Herman M. C. elegans POP-1/TCF functions in a canonical Wnt pathway that controls cell
migration and in a noncanonical Wnt pathway that controls cell polarity. Development.
2001;128: 581–590. doi:10.1242/dev.128.4.581

12. Mizumoto K, Sawa H. Two βs or not two βs: regulation of asymmetric division by β-catenin.
Trends Cell Biol. 2007;17: 465–473. doi:10.1016/j.tcb.2007.08.004

13. Zacharias AL, Walton T, Preston E, Murray JI. Quantitative Differences in Nuclear β-catenin
and TCF Pattern Embryonic Cells in C. elegans. Sternberg PW, editor. PLOS Genet.
2015;11: e1005585. doi:10.1371/journal.pgen.1005585

14. Lin R, Hill RJ, Priess JR. POP-1 and anterior-posterior fate decisions in C. elegans embryos.
Cell. 1998;92: 229–239.

15. Maduro MF, Lin R, Rothman JH. Dynamics of a developmental switch: recursive
intracellular and intranuclear redistribution of Caenorhabditis elegans POP-1 parallels Wnt-
inhibited transcriptional repression. Dev Biol. 2002;248: 128–142.
doi:10.1006/dbio.2002.0721

16. Huang S, Shetty P, Robertson SM, Lin R. Binary cell fate specification during C. elegans
embryogenesis driven by reiterated reciprocal asymmetry of TCF POP-1 and its
coactivatorβ-catenin SYS-1. Development. 2007;134: 2685–2695. doi:10.1242/dev.008268

17. Calvo D, Victor M, Gay F, Sui G, Po-Shan Luke M, Dufourcq P, et al. A POP-1 repressor
complex restricts inappropriate cell type-specific gene transcription during Caenorhabditis
elegans embryogenesis. EMBO J. 2001;20: 7197–7208. doi:10.1093/emboj/20.24.7197

18. Maduro MF, Kasmir JJ, Zhu J, Rothman JH. The Wnt effector POP-1 and the PAL-1/Caudal
homeoprotein collaborate with SKN-1 to activate C. elegans endoderm development. Dev
Biol. 2005;285: 510–523. doi:10.1016/j.ydbio.2005.06.022

19. Owraghi M, Broitman-Maduro G, Luu T, Roberson H, Maduro MF. Roles of the Wnt
effector POP-1/TCF in the C. elegans endomesoderm specification gene network. Dev Biol.
2010;340: 209–221. doi:10.1016/j.ydbio.2009.09.042

20. Murgan S, Kari W, Rothbächer U, Iché-Torres M, Mélénec P, Hobert O, et al. Atypical
Transcriptional Activation by TCF via a Zic Transcription Factor in C. elegans Neuronal
Precursors. Dev Cell. 2015;33: 737–745. doi:10.1016/j.devcel.2015.04.018

21. Murgan S, Bertrand V. How targets select activation or repression in response to Wnt.
Worm. 2015;4: e1086869. doi:10.1080/21624054.2015.1086869

22. Zhang CU, Blauwkamp TA, Burby PE, Cadigan KM. Wnt-Mediated Repression via
Bipartite DNA Recognition by TCF in the Drosophila Hematopoietic System. Desplan C,
editor. PLoS Genet. 2014;10: e1004509. doi:10.1371/journal.pgen.1004509

23. Murray JI, Boyle TJ, Preston E, Vafeados D, Mericle B, Weisdepp P, et al. Multidimensional
regulation of gene expression in the C. elegans embryo. Genome Res. 2012;22: 1282–1294.
doi:10.1101/gr.131920.111

24. Ma X, Zhao Z, Xiao L, Xu W, Kou Y, Zhang Y, et al. A 4D single-cell protein atlas of
transcription factors delineates spatiotemporal patterning during embryogenesis. Nat
Methods. 2021;18: 893–902. doi:10.1038/s41592-021-01216-1

25. Murray JI, Bao Z, Boyle TJ, Boeck ME, Mericle BL, Nicholas TJ, et al. Automated analysis
of embryonic gene expression with cellular resolution in C. elegans. Nat Methods. 2008;5:
703–709. doi:10.1038/nmeth.1228

26. Kvon EZ, Waymack R, Gad M, Wunderlich Z. Enhancer redundancy in development and
disease. Nat Rev Genet. 2021;22: 324–336. doi:10.1038/s41576-020-00311-x

27. Sulston JE, Horvitz HR. Post-embryonic cell lineages of the nematode, Caenorhabditis
elegans. Dev Biol. 1977;56: 110–156. doi:10.1016/0012-1606(77)90158-0

28. Bao Z, Murray JI, Boyle T, Ooi SL, Sandel MJ, Waterston RH. Automated cell lineage
tracing in Caenorhabditis elegans. Proc Natl Acad Sci U S A. 2006;103: 2707.
doi:10.1073/pnas.0511111103

29. Santella A, Du Z, Nowotschin S, Hadjantonakis A-K, Bao Z. A hybrid blob-slice model for
accurate and efficient detection of fluorescence labeled nuclei in 3D. BMC Bioinformatics.
2010;11: 580. doi:10.1186/1471-2105-11-580

30. Boyle TJ, Bao Z, Murray JI, Araya CL, Waterston RH. AceTree: a tool for visual analysis of
Caenorhabditis elegans embryogenesis. BMC Bioinformatics. 2006;7: 275.
doi:10.1186/1471-2105-7-275

31. Katzman B, Tang D, Santella A, Bao Z. AceTree: a major update and case study in the long
term maintenance of open-source scientific software. BMC Bioinformatics. 2018;19: 121.
doi:10.1186/s12859-018-2127-0

32. Packer Jonathan S., Zhu Qin, Huynh Chau, Sivaramakrishnan Priya, Preston Elicia, Dueck
Hannah, et al. A lineage-resolved molecular atlas of C. elegans embryogenesis at single-cell
resolution. Science. 2019;365: eaax1971. doi:10.1126/science.aax1971

33. Yanai I, Baugh LR, Smith JJ, Roehrig C, Shen-Orr SS, Claggett JM, et al. Pairing of
competitive and topologically distinct regulatory modules enhances patterned gene
expression. Mol Syst Biol. 2008;4. doi:10.1038/msb.2008.6

34. Lin R, Thompson S, Priess JR. pop-1 encodes an HMG box protein required for the
specification of a mesoderm precursor in early C. elegans embryos. Cell. 1995;83: 599–609.
doi:10.1016/0092-8674(95)90100-0

35. Kidd AR, Miskowski JA, Siegfried KR, Sawa H, Kimble J. A β-Catenin Identified by
Functional Rather Than Sequence Criteria and Its Role in Wnt/MAPK Signaling. Cell.
2005;121: 761–772. doi:10.1016/j.cell.2005.03.029

36. Phillips BT, Kidd AR, King R, Hardin J, Kimble J. Reciprocal asymmetry of SYS-1/beta-
catenin and POP-1/TCF controls asymmetric divisions in Caenorhabditis elegans. Proc Natl
Acad Sci. 2007;104: 3231–3236. doi:10.1073/pnas.0611507104

37. Broitman-Maduro G, Owraghi M, Hung WWK, Kuntz S, Sternberg PW, Maduro MF. The
NK-2 class homeodomain factor CEH-51 and the T-box factor TBX-35 have overlapping
function in C. elegans mesoderm development. Dev Camb Engl. 2009;136: 2735–2746.
doi:10.1242/dev.038307

38. Broitman-Maduro G, Lin KT-H, Hung WWK, Maduro MF. Specification of the C. elegans
MS blastomere by the T-box factor TBX-35. Dev Camb Engl. 2006;133: 3097–3106.
doi:10.1242/dev.02475

39. Alper S, Kenyon C. The zinc finger protein REF-2 functions with the Hox genes to inhibit
cell fusion in the ventral epidermis of C. elegans. Development. 2002;129: 3335–3348.
doi:10.1242/dev.129.14.3335

40. Bertrand V, Hobert O. Linking asymmetric cell division to the terminal differentiation
program of postmitotic neurons in C. elegans. Dev Cell. 2009;16: 563–575.
doi:10.1016/j.devcel.2009.02.011

41. Walton T, Preston E, Nair G, Zacharias AL, Raj A, Murray JI. The Bicoid Class
Homeodomain Factors ceh-36/OTX and unc-30/PITX Cooperate in C. elegans Embryonic
Progenitor Cells to Regulate Robust Development. PLOS Genet. 2015;11: e1005003.
doi:10.1371/journal.pgen.1005003

42. Richards JL, Zacharias AL, Walton T, Burdick JT, Murray JI. A quantitative model of
normal Caenorhabditis elegans embryogenesis and its disruption after stress. Dev Biol.
2013;374: 12–23. doi:10.1016/j.ydbio.2012.11.034

43. Maduro MF, Meneghini MD, Bowerman B, Broitman-Maduro G, Rothman JH. Restriction
of Mesendoderm to a Single Blastomere by the Combined Action of SKN-1 and a GSK-3β
Homolog Is Mediated by MED-1 and -2 in C. elegans. Mol Cell. 2001;7: 475–485.
doi:10.1016/S1097-2765(01)00195-2

44. Maduro MF, Hill RJ, Heid PJ, Newman-Smith ED, Zhu J, Priess JR, et al. Genetic
redundancy in endoderm specification within the genus Caenorhabditis. Dev Biol. 2005;284:
509–522. doi:10.1016/j.ydbio.2005.05.016

45. Boeck ME, Boyle T, Bao Z, Murray J, Mericle B, Waterston R. Specific roles for the GATA
transcription factors end-1 and end-3 during C. elegans E-lineage development. Spec Sect
Hist Dev Biol. 2011;358: 345–355. doi:10.1016/j.ydbio.2011.08.002

46. Good K, Ciosk R, Nance J, Neves A, Hill RJ, Priess JR. The T-box transcription factors
TBX-37 and TBX-38 link GLP-1/Notch signaling to mesoderm induction in C. elegans
embryos. Development. 2004;131: 1967–1978. doi:10.1242/dev.01088

47. Andachi Y. Caenorhabditis elegans T-box genes tbx-9 and tbx-8 are required for formation
of hypodermis and body-wall muscle in embryogenesis. Genes Cells. 2004;9: 331–344.
doi:10.1111/j.1356-9597.2004.00725.x

48. Neves A, Priess JR. The REF-1 Family of bHLH Transcription Factors Pattern C. elegans
Embryos through Notch-Dependent and Notch-Independent Pathways. Dev Cell. 2005;8:
867–879. doi:10.1016/j.devcel.2005.03.012

49. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The Human
Genome Browser at UCSC. Genome Res. 2002;12: 996–1006. doi:10.1101/gr.229102

50. Charest J, Daniele T, Wang J, Bykov A, Mandlbauer A, Asparuhova M, et al. Combinatorial
Action of Temporally Segregated Transcription Factors. Dev Cell. 2020;55: 483-499.e7.
doi:10.1016/j.devcel.2020.09.002

51. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al.
Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome
Res. 2005;15: 1034–1050. doi:10.1101/gr.3715005

52. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution
rates on mammalian phylogenies. Genome Res. 2010;20: 110–121.
doi:10.1101/gr.097857.109

53. LaBonty M, Szmygiel C, Byrnes LE, Hughes S, Woollard A, Cram EJ. CACN-1/Cactin
plays a role in Wnt signaling in C. elegans. PloS One. 2014;9.
doi:10.1371/journal.pone.0101945

54. Murray JI, Preston E, Crawford JP, Rumley JD, Amom P, Anderson BD, et al. The anterior
Hox gene *ceh-13* and *elt-1/GATA* activate the posterior Hox genes
nob-1 and *php-3* to specify posterior lineages in the *C.
elegans* embryo. bioRxiv. 2021; 2021.02.09.430385. doi:10.1101/2021.02.09.430385

55. Poole RJ, Hobert O. Early Embryonic Programming of Neuronal Left/Right Asymmetry in
C. elegans. Curr Biol. 2006;16: 2279–2292. doi:10.1016/j.cub.2006.09.041

56. Tintori SC, Osborne Nishimura E, Golden P, Lieb JD, Goldstein B. A Transcriptional
Lineage of the Early C. elegans Embryo. Dev Cell. 2016;38: 430–444.
doi:10.1016/j.devcel.2016.07.025

57. Rocheleau CE, Yasuda J, Shin TH, Lin R, Sawa H, Okano H, et al. WRM-1 Activates the
LIT-1 Protein Kinase to Transduce Anterior/Posterior Polarity Signals in C. elegans. Cell.
1999;97: 717–726. doi:10.1016/S0092-8674(00)80784-9

58. Lo M-C, Gay F, Odom R, Shi Y, Lin R. Phosphorylation by the β-Catenin/MAPK Complex
Promotes 14-3-3-Mediated Nuclear Export of TCF/POP-1 in Signal-Responsive Cells in C.
elegans. Cell. 2004;117: 95–106. doi:10.1016/S0092-8674(04)00203-X

59. Kaletta T, Schnabel H, Schnabel R. Binary specification of the embryonic lineage in
Caenorhabditis elegans. Nature. 1997;390: 294–298. doi:10.1038/36869

60. Fukushige T, Brodigan TM, Schriefer LA, Waterston RH, Krause M. Defining the
transcriptional redundancy of early bodywall muscle development in C. elegans: evidence
for a unified theory of animal muscle development. Genes Dev. 2006;20: 3395–3406.
doi:10.1101/gad.1481706

61. Fukushige T, Krause M. The myogenic potency of HLH-1 reveals wide-spread
developmental plasticity in early C. elegans embryos. Dev Camb Engl. 2005;132: 1795–
1805. doi:10.1242/dev.01774

62. Simonis N, Rual J-F, Carvunis A-R, Tasan M, Lemmens I, Hirozane-Kishikawa T, et al.
Empirically controlled mapping of the Caenorhabditis elegans protein-protein interactome
network. Nat Methods. 2009;6.

63. Whiting J, Marshall H, Cook M, Krumlauf R, Rigby PW, Stott D, et al. Multiple spatially
specific enhancers are required to reconstruct the pattern of Hox-2.6 gene expression. Genes
Dev. 1991;5: 2048–2059. doi:10.1101/gad.5.11.2048

64. Epstein DJ, McMahon AP, Joyner AL. Regionalization of Sonic hedgehog transcription
along the anteroposterior axis of the mouse central nervous system is regulated by Hnf3-
dependent and -independent mechanisms. Development. 1999;126: 281–292.
doi:10.1242/dev.126.2.281

65. Teng Y, Girard L, Ferreira HB, Sternberg PW, Emmons SW. Dissection of cis-regulatory
elements in the C. elegans Hox gene egl-5 promoter. Dev Biol. 2004;276: 476–492.
doi:10.1016/j.ydbio.2004.09.012

66. Streit A, Kohler R, Marty T, Belfiore M, Takacs-Vellai K, Vigano M-A, et al. Conserved
Regulation of the Caenorhabditis elegans labial/Hox1 Gene ceh-13. Dev Biol. 2002;242: 96–
108. doi:10.1006/dbio.2001.0544

67. Landmann F, Quintin S, Labouesse M. Multiple regulatory elements with spatially and
temporally distinct activities control the expression of the epithelial differentiation gene lin-
26 in C. elegans. Dev Biol. 2004;265: 478–490. doi:10.1016/j.ydbio.2003.09.009

68. Chen RA-J, Down TA, Stempor P, Chen QB, Egelhofer TA, Hillier LW, et al. The landscape
of RNA polymerase II transcription initiation in C. elegans reveals promoter and enhancer
architectures. Genome Res. 2013;23: 1339–1347. doi:10.1101/gr.153668.112

69. Serizay J, Dong Y, Jänes J, Chesney M, Cerrato C, Ahringer J. Distinctive regulatory
architectures of germline-active and somatic genes in C. elegans. Genome Res. 2020;30:
1752–1765. doi:10.1101/gr.265934.120

70. Kuersten S, Segal SP, Verheyden J, LaMartina SM, Goodwin EB. NXF-2, REF-1, and REF-
2 Affect the Choice of Nuclear Export Pathway for tra-2 mRNA in C. elegans. Mol Cell.
2004;14: 599–610. doi:10.1016/j.molcel.2004.05.004

71. Clark E, Akam M. Odd-paired controls frequency doubling in Drosophila segmentation by
altering the pair-rule gene regulatory network. Wittkopp PJ, editor. eLife. 2016;5: e18215.
doi:10.7554/eLife.18215

72. Yoon Y, Klomp J, Martin-Martin I, Criscione F, Calvo E, Ribeiro J, et al. Embryo polarity in
moth flies and mosquitoes relies on distinct old genes with localized transcript isoforms.
Wittkopp PJ, Pick L, Popadic A, Akam M, editors. eLife. 2019;8: e46711.
doi:10.7554/eLife.46711

73. Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, et al. Systematic functional
analysis of the Caenorhabditis elegans genome using RNAi. Nature. 2003;421: 231–237.
doi:10.1038/nature01278

74. Vora S, Phillips BT. Centrosome-Associated Degradation Limits β-Catenin Inheritance by
Daughter Cells after Asymmetric Division. Curr Biol. 2015;25: 1005–1016.
doi:10.1016/j.cub.2015.02.020

75. Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, et al.
Determination and inference of eukaryotic transcription factor sequence specificity. Cell.
2014;158: 1431–1443. doi:10.1016/j.cell.2014.08.009

76. Narasimhan K, Lambert SA, Yang AWH, Riddell J, Mnaimneh S, Zheng H, et al. Mapping
and analysis of transcription factor sequence specificities. eLife. 2015;4.
doi:10.7554/eLife.06967

77. Bao Z, Murray JI. Mounting Caenorhabditis elegans Embryos for Live Imaging of
Embryogenesis. Cold Spring Harb Protoc. 2011;2011: pdb.prot065599.
doi:10.1101/pdb.prot065599

1323